Variance-Reduced Accelerated First-order Methods: Central Limit Theorems and Confidence Statements

Jinlong Lei and Uday V. Shanbhag *†

Abstract

In this paper, we consider a strongly convex stochastic optimization problem and propose three classes of variable sample-size stochastic first-order methods: (i) the standard stochastic gradient descent method; (ii) its accelerated variant; and (iii) the stochastic heavy ball method. In each scheme, the exact gradients are approximated by averaging across an increasing batch size of sampled gradients. We prove that when the sample-size increases geometrically, the generated estimates converge in mean to the optimal solution at a geometric rate for schemes (i) – (iii). Based on this result, we provide central limit statements whereby it is shown that the rescaled estimation errors converge in distribution to a normal distribution with the associated covariance matrix dependent on the Hessian matrix, covariance of the gradient noise, and the steplength. If the sample-size increases at a polynomial rate, we show that theorems (CLTs). Under certain conditions, we discuss how both the algorithms and the associated limit theorems may be extended to constrained and nonsmooth regimes. Finally, we provide an avenue to construct confidence regions for the optimal solution based on the established CLTs and test the theoretical findings on a stochastic parameter estimation problem.

1 Introduction

In this paper, we consider the strongly convex optimization problem (1):

$$\min_{x \in \mathbb{R}^m} f(x) \triangleq \mathbb{E}[F(x, \boldsymbol{\xi})],\tag{1}$$

where $\boldsymbol{\xi}: \Omega \to \mathbb{R}^d$ is a random variable defined on the probability space $(\Omega, \mathcal{F}, \mathbb{P}), F: \mathbb{R}^m \times \mathbb{R}^d \to \mathbb{R}$, and the expectation is taken over the distribution of the random vector. Stochastic optimization problems have been extensively studied, given the wide applicability of such models in almost all areas of science and engineering, ranging from communication and queueing systems to finance (cf. [4, 55]). However, in most situations, this expectation and its derivative are unavailable in closed form requiring the development

^{*}Email: leijinlong@tongji.edu.cn (Jinlong Lei), udaybag@psu.edu (Uday V. Shanbhag).

[†]Lei is the Department of Control Science and Engineering, Tongji University; She is also with the Shanghai Institute of Intelligent Science and Technology, Tongji University, Shanghai 200092, China. Shanbhag is with the Department of Industrial and Manufacturing Engineering, Pennsylvania State University, University Park, PA 16802, USA

of sampling approaches. Sample average approximation (SAA) [55] and stochastic approximation (SA) represent two commonly used approaches for contending with stochastic programs. Our focus is on SA schemes, first considered by Robbins and Monro [51] for seeking roots of a regression function with noisy observations. The standard SA algorithm $x_{k+1} = x_k - \alpha_k \nabla F(x_k, \xi_k)$, also known as the stochastic gradient descent (SGD) algorithm, updates the estimate x_{k+1} based on a sampled gradient $\nabla F(x_k, \xi_k)$. The convergence analysis usually requires suitable properties on the gradient map (such as Lipschitzian requirements) and the steplength sequence (such as non-summable but square summable). The almost sure convergence of x_k to x^* , the unique optimal solution of (1), was established in [5, 8, 11] on the basis of the Robbins-Siegmund theorem [52] while ordinary differential equation (ODE) techniques have also been employed for claiming similar statements in [7, 25, 32]. In addition, in [10], the authors developed a statistical diagnostic test to detect the phase transition, during which the iterative procedure converges towards a region of interest in the context of SGD with constant learning rate. Tight bounds on the rate of convergence can be obtained by establishing the asymptotic distribution for the iterates (cf. [7, 11, 18, 25, 32]). An instructive review of results up to around 2010 was provided in [47], while asymptotic normality of the suitably scaled iterates for SA with decreasing step-sizes has been proven in [18, 25, 32]. To be specific, the rescaled error process $\sqrt{\alpha_k}(x_k - x^*)$ asymptotically converges in distribution to a normal distribution with zero mean and with covariance depending on the Hessian matrix, the covariance of the gradient noise, and the steplength. Prior work on CLTs for standard stochastic approximation for smooth convex optimization can be traced to the seminal averaging paper by Polyak and Juditsky [50]. The asymptotic normality was further investigated in [11] for SA with expanding truncations, an avenue that does not require Lipschitz continuity of the gradients. The sequence of iterates generated by the constant steplength SA scheme has been shown to be a homogeneous Markov chain with a unique stationary distribution; see [32, Chapter 9], [39, Chapter 17], and [17].

CLTs for SA schemes are significant from the standpoint of algorithm design as well as inference.

(i) Algorithm design. Indeed, the optimal selection of the steplength depends on the Hessian matrix at x^* , see e.g., [11, Chapter 3.4]. Since the Hessian at x^* is unavailable, the optimal value of α_k can only be estimated, which has led to the development of adaptive SA methods (cf. [57, 63, 64]). Motivated by the heavy dependence of SA schemes on steplength choices, the authors in [66] developed a self-tuned rule that adapts the steplength sequence to problem parameters. In a similar vein, there has been an effort to develop optimal constant steplengths. Specifically, it was shown in [40] that with suitably selected constant stepsizes, the expected function values at the averaged iterate converge to the optimum with rate $O(1/\sqrt{k})$ in merely convex case and rate O(1/k) in strongly convex case, matching the lower bounds [41]. It is further shown in [38] that the constant SGD simulates a Markov chain with a stationary distribution, which might be used to adjust the tuning parameters of constant SGD so as to improve the convergence rate, e.g., SGD for fitting the generalized linear models [61] and approximating the Bayesian posterior inference [38].

(*ii*) Confidence statements. Furthermore, CLTs for SA schemes might allow for the possibility of constructing confidence regions for the optimal solution (e.g. [3,12,27]). In particular, Hsieh and Glynn [27] designed an approach to rigorously characterize confidence regions without explicitly estimating the covariance of the limiting normal distribution, while Chen et al. [12] proposed a plug-in estimator and a batch-means estimator for the asymptotic covariance of the average iterate from SGD. In addition, Su and Zhu [58] advocated performing SGD updates for a while and then splitting the single thread into several threads; notably, they proceeded to construct *t*-based confidence interval that can attain asymptotically exact coverage probability. The monograph [32] has extensively investigated CLTs for SA schemes under both constant and decreasing steplengths. In the context of SAA schemes, there has been some recent work on developing confidence statements for stochastic optimization [55] and stochastic variational inequality problems [33, 37].

Unfortunately, SA schemes with diminishing steps cannot recover the deterministic convergence rates seen in exact gradient methods while constant steplength SA schemes are only characterized by convergence guarantees to a neighborhood of the optimal solution. Variance-reduction schemes employing an increasing batch-size of sampled gradients (instead of the unavailable true gradient) appear to have been first alluded to in [2, 16, 26, 45] and analyzed in smooth and strongly convex [9, 19, 53, 54], smooth convex regimes [22], nonsmooth (but smoothable) convex [28], nonconvex [34], and game-theoretic [35] regimes. Notably, the linear rates in mean-squared error were derived for strongly convex smooth [54] and a subclass of nonsmooth objectives [28], while a rate $O(1/k^2)$ and O(1/k) was obtained for expected sub-optimality in convex smooth [22, 29] and nonsmooth [28], respectively. In each instance, the schemes achieve the corresponding optimal deterministic rates of convergence under suitable growth in sample-size sequences while in almost all cases, the optimal sample-complexity bounds were obtained. An excellent discussion relating sampling rates and the canonical rate (i.e. the Monte-Carlo rate) in stochastic gradient-based schemes has been provided by Pasupathy et al. [46].

Gaps and motivation. This paper is motivated by the following gaps. (i) Despite a surge of interest in variance-reduced schemes, both with and without acceleration, no limit theorems are available for claiming asymptotic normality of the scaled sequence in either unaccelerated or Nesterov accelerated regimes coupled with variance reduction. In particular, we have little understanding regarding whether such avenues have detrimental impacts in terms of the limiting behavior. Such statements are also unavailable for the related heavy-ball method. (ii) The geometric growth in sample-size required for achieving linear rates of convergence may prove onerous in some settings. Are CLTs available in settings where the sample-size grows at a polynomial rate? (iii) Finally, given a CLT, there is little by way of availability of rigorous confidence statements, barring the work by Hsieh and Glynn [27]. Can such statements be developed for the proposed class of variance-reduced first-order methods?

Justification and relationship to other variance-reduced schemes.

(i) *Terminology and applicability.* The moniker "variance-reduced" has been loosely used in stochastic optimization to capture approaches that admit deterministic convergence rates. For instance, for minimizing finite-sum expected-residual minimization problems in machine learning, techniques such as stochastic variance reduced gradient (SVRG) [30] and a fast incremental gradient method [14] achieve deterministic rates of convergence. In general probability spaces, using increasing batch-sizes of gradients represent progressively accurate approximations of the true gradient, as opposed to noisy sampled variants employed in single sample schemes [19]. The resulting schemes, often referred to as *mini-batch* SA schemes, may

achieve deterministic rates of convergence under suitable growth rates in sample-sizes.

(ii) Weaker assumptions, and stronger statements. The proposed variance-reduced framework has several key benefits often unavailable in single-sample regimes: 1) Under suitable assumptions, such schemes often achieve optimal deterministic rates in terms of iteration complexity (e.g. in constrained regimes, complexity of an iteration is essentially that of computing a projection onto a convex set) while achieving near-optimal sample complexity. This assumes profound importance when iterations are expensive as in high-dimensional and nonlinear settings; 2) In addition, the benefits of acceleration are less clear in stochastic single-sample regimes, absent such approaches (to the best of our knowledge). While the present treatment is in an unconstrained regime, we also discuss how extensions to constrained and nonsmooth regimes may be developed. (iii) Sampling requirements. Variance-reduced schemes have obvious benefits when sampling is relatively cheap compared to computational effort of an iteration. Though such schemes are often characterized by near-optimal sample complexity, one might question how to contend with batch-sizes (denoted by N_k) tending to $+\infty$. This issue is somewhat of a red herring since most SA schemes are meant to provide ϵ -approximations; (see the detailed discussions in Remark 3). With the ubiquity of multi-core architecture, such requirements are not terribly onerous. In addition, we also establish the polynomial rates and associated sample complexities for polynomially increasing sample size. They are particularly important when sampling is expensive and one would like to modulate the sampling rate to ensure that the scheme is practical.

(iv) *CLTs and confidence statements*. Apart from rate statements, we additionally establish CLT results for the proposed variance-reduced framework. In particular, we provide a clean characterization of the limiting distribution and show the dependence of the Hessian matrix, the condition number, the noise covariance, etc in the prescription of the CLT. The confidence statements are established as well; we should note that we are unaware of such statements in the context of accelerated and heavy-ball settings in a variance-reduced regime.

Outline and contributions. To address these gaps, we present CLTs and confidence statements for first-order stochastic variance-reduced algorithms for resolving (1), including the classical SGD [51], the stochastic variants of the Nesterov's accelerated method [43], and the heavy ball method [48]. We provide statements when batch-sizes increases at either a geometric or a polynomial rate. Our main contributions are summarized next.

(I) CLTs for variable sample-size gradient methods. In Section 2.1, we recall the variance-reduced (VR) stochastic gradient algorithm with a constant steplength (see Algorithm 1), where the gradient is estimated by the average of an increasing batch of sampled gradients. In Section 3, when the batch-size increases at a geometric rate, we observe that the mean-squared error diminishes at a geometric rate (see Proposition 1) and provide a preliminary Lemma 7 for establishing a CLT for a noised-corrupted linear recursion. Based on this Lemma and the linear approximation of the gradient function at x^* , we may derive a CLT (see Theorem 1) in this setting. We proceed to show that the covariance of the limiting normal distribution depends on the Hessian matrix at the solution, the condition number, the covariance of gradient noise, etc and the steplength. Additionally, we show in Section 4 that when the batch-size is increased at a polynomial rate, the sequence

of iterates converges at a corresponding polynomial rate (see Proposition 5). Then based on the CLT shown in Lemma 9 for the time-varying linear recursion, the CLT for Algorithm 1 with polynomially increasing batch-size is established in Theorem 5.

(II) CLTs for VR-accelerated gradient method. In Section 2.2, we consider a VR-accelerated gradient algorithm with constant steplengths (see Algorithm 2). Then by leveraging the geometric rate of convergence (Proposition 2), we establish amongst the first CLTs in accelerated regimes (Theorem 2) when the batch-size increases at a geometric rate. It is well-known that the accelerated variant has better constants in the iteration complexity in deterministic regimes and this is also seen in stochastic settings. Akin to earlier, when the batch-size increases at a polynomial rate, a polynomial convergence rate and the corresponding CLT are in Proposition 6 and Theorem 6.

III. CLTs for VR-heavy-ball schemes. In Section 2.3, we design a VR-heavy ball scheme with constant steplengths (see Algorithm 3). In contrast with Algorithms 1 and 2, the heavy ball method is analyzed for both quadratic and non-quadratic objective functions. When the batch-size increases geometrically, the geometric rate of convergence are respectively shown in Proposition 3 and 4 for quadratic and non-quadratic objective functions, while the corresponding CLT is established in Theorem 3 and Theorem 4. Similarly, for polynomially increasing batch-size, a polynomial rate of convergence is shown in Proposition 7 and 8 for quadratic and non-quadratic objective functions, whereas the corresponding CLTs are respectively established in Theorem 7 and Theorem 8.

IV. Confidence statements. In Section 5, inspired by [27], we provide rigorous confidence regions for the optimal solution and function value. Then in Section 6, we implement some simulations on a parameter estimation problem in the stochastic environment to validate the theoretical findings.

Notations. Let \mathbf{I}_m denote the identity matrix of dimension m and $\mathbf{0}_m \in \mathbb{R}^{m \times m}$ denote the matrix with all entries equal zero. Let $\{X_k\}$ be a sequence of random variables. $X_k \xrightarrow[k \to \infty]{d} N(0, \mathbf{S})$ denotes that X_k converges in distribution to a normal distribution $N(0, \mathbf{S})$ with mean zero and covariance \mathbf{S} , and $X_k \xrightarrow[k \to \infty]{P} X$ denotes that X_k converges in probability to X. By utilizing the definitions of the "small-o in probability" and "big-O in probability" notations $o_P(\cdot)$ and $O_P(\cdot)$ provided in [62, Chapter 2.2], the expression $e_k = o_P(1)$ implies that a sequence of random variables $\{e_k\}$ converges to zero in probability. Similarly, the expression $e_k = O_P(1)$ implies that a sequence of random variables $\{e_k\}$ is bounded in probability. For a square matrix \mathbf{P} , we denote by $\rho(\mathbf{P})$ and $\|\mathbf{P}\|$ its spectral radius and matrix two-norm, respectively.

2 First-Order Variable Sample-size Stochastic Algorithms.

Since the exact gradient $\nabla f(x)$ is expectation-valued and unavailable in a closed form, we assume that there exists a *stochastic first-order oracle* such that for any x and ξ , a sampled gradient $\nabla F(x,\xi)$ is returned and is assumed to be an unbiased estimator of $\nabla f(x)$. In this section, we present three first-order stochastic algorithms to find the optimal solution to (1). Throughout the paper, time is slotted at k = 0, 1, 2, ... and an iterate at time k is denoted by $x_k \in \mathbb{R}^m$.

2.1 Variance-reduced Gradient Method.

We present a variable sample-size stochastic gradient algorithm (Algorithm 1) to solve (1), where at iteration k, the unavailable exact gradient $\nabla f(x_k)$ is estimated via the average of an increasing batch-size of sampled gradients.

Algorithm 1 Variance reduced SGD

Given an arbitrary initial value $x_0 \in \mathbb{R}^m$, a positive constant $\alpha > 0$, and a positive integer sequence $\{N_k\}_{k>0}$. Then iterate the following equation for $k \ge 0$.

$$x_{k+1} = x_k - \alpha \frac{\sum_{j=1}^{N_k} \nabla F(x_k, \xi_{j,k})}{N_k},$$
(2)

where $\alpha > 0$ is the constant steplength, N_k is the number of sampled gradients used at time k, and $\xi_{j,k}$, $j = 1, \dots, N_k$, denote the independent and identically distributed (i.i.d.) realizations of $\boldsymbol{\xi}$.

If the gradient observation noise w_{k,N_k} is defined as

$$w_{k,N_k} \triangleq \frac{\sum_{j=1}^{N_k} \nabla F(x_k,\xi_{j,k})}{N_k} - \nabla f(x_k), \tag{3}$$

then the update (2) can be rewritten as

$$x_{k+1} = x_k - \alpha (\nabla f(x_k) + w_{k,N_k}).$$
(4)

For any k, define \mathcal{F}_k as $\mathcal{F}_k \triangleq \sigma\{x_0, \xi_{j,t}, 1 \leq j \leq N_t, 0 \leq t \leq k-1\}$. Then x_k is adapted to \mathcal{F}_k by Algorithm 1. We impose the following conditions on the objective function, the conditional expectation and the second moments of the sampled gradients produced by the stochastic first-order oracle.

Assumption 1 (i) f is continuously differentiable on \mathbb{R}^m with a Lipschitz continuous gradient, i.e., there exists a constant L > 0 such that $\|\nabla f(x) - \nabla f(x')\| \le L \|x - x'\|$ for any $x, x' \in \mathbb{R}^m$. (ii) f is η -strongly convex, i.e., $(\nabla f(x) - \nabla f(x'))^T (x - x') \ge \eta \|x - x'\|^2$ for any $x, x' \in \mathbb{R}^m$.

(iii) There exists a constant $\nu > 0$ such that for any $k \ge 0$ and $j = 1, \dots, N_k$, $\mathbb{E}[\nabla F(x_k, \boldsymbol{\xi}) | \mathcal{F}_k] = \nabla f(x_k)$ almost surely and $\mathbb{E}[\|\nabla F(x_k, \boldsymbol{\xi}) - \nabla f(x_k)\|^2 | \mathcal{F}_k] \le \nu^2$ almost surely (a.s.).

Since x_k is adapted to \mathcal{F}_k and the samples $\xi_{j,k}$, $j = 1, \dots, N_k$, are independent, we obtain from (3) and Assumption 1(iii) that for any $k \ge 0$,

$$\mathbb{E}[w_{k,N_k} \mid \mathcal{F}_k] = \frac{\sum_{j=1}^{N_k} \mathbb{E}\left[\nabla F(x_k, \boldsymbol{\xi}) - \nabla f(x_k) \mid \mathcal{F}_k\right]}{N_k} = 0, \text{ and}$$

$$\mathbb{E}[\|w_{k,N_k}\|^2 \mid \mathcal{F}_k] = \frac{\sum_{j=1}^{N_k} \mathbb{E}[\|\nabla F(x_k, \boldsymbol{\xi}) - \nabla f(x_k)\|^2 \mid \mathcal{F}_k]}{N_k^2} \le \frac{\nu^2}{N_k}.$$
(5)

hold almost surely.

Since f is strongly convex, it has a unique optimal solution denoted by x^* . Then by the optimality condition, $\nabla f(x^*) = 0$. We now introduce an inequality [3, Eqn. (2.1.24)] on f satisfying Assumptions 1(i) and 1(ii):

$$(x-y)^T (\nabla f(x) - \nabla f(y)) \ge \frac{\eta L \|x-y\|^2}{\eta + L} + \frac{\|\nabla f(x) - \nabla f(y)\|^2}{\eta + L}, \quad \forall x, y \in \mathbb{R}^m.$$
(6)

We then establish a recursion on the mean-squared estimation error, which can be proved by making a simple modification to the proof of [3, Theorem 2.1.15].

Lemma 1 Let Assumption 1 hold. Consider Algorithm 1 with $\alpha \in (\frac{0,2}{\eta+L}]$. Then

$$\mathbb{E}[\|x_{k+1} - x^*\|^2] \le \left(1 - \frac{2\alpha\eta L}{\eta + L}\right) \mathbb{E}[\|x_k - x^*\|^2] + \frac{\alpha^2 \nu^2}{N_k}, \quad \forall k \ge 0.$$
(7)

Proof. From (4) it follows that

$$||x_{k+1} - x^*||^2 = ||x_k - x^*||^2 - 2\alpha (x_k - x^*)^T (\nabla f(x_k) + w_{k,N_k}) + \alpha^2 (||\nabla f(x_k)||^2 + 2\nabla f(x_k)^T w_{k,N_k} + ||w_{k,N_k}||^2).$$
(8)

Since $\nabla f(x^*) = 0$, by using (6), we obtain that

$$(x_k - x^*)^T \nabla f(x_k) \ge \frac{\eta L}{\eta + L} \|x_k - x^*\|^2 + \frac{1}{\eta + L} \|\nabla f(x_k)\|^2.$$
(9)

Since x_k is adapted to \mathcal{F}_k , by taking expectations conditioned on \mathcal{F}_k on both sides of (8), and using (5) from Assumption 1(iii), we obtain that

$$\mathbb{E}[\|x_{k+1} - x^*\|^2 |\mathcal{F}_k] \le \|x_k - x^*\|^2 - 2\alpha (x_k - x^*)^T \nabla f(x_k) + \alpha^2 \|\nabla f(x_k)\|^2 + \frac{\alpha^2 \nu^2}{N_k} \\ \stackrel{(9)}{\le} \left(1 - \frac{2\alpha\eta L}{\eta + L}\right) \|x_k - x^*\|^2 - \alpha \left(\frac{2}{\eta + L} - \alpha\right) \|\nabla f(x_k)\|^2 + \frac{\alpha^2 \nu^2}{N_k}, \quad a.s. .$$

Then by choosing $\alpha \in (0, \frac{2}{n+L}]$ and taking unconditional expectations, we achieve (7).

2.2 Accelerated Gradient Method.

Nesterov's accelerated gradient descent method generates a sequence that converges to the solution at a rate $O(q^k)$ where $q \triangleq 1 - \sqrt{\eta/L}$ for η -strongly convex and *L*-smooth functions [3], and with a rate $O(1/k^2)$ for merely convex functions [43]. Nesterov proved that this is the best possible rate for any first-order method. As such, we combine the Nestrov's accelerated method with Algorithm 1 and propose an accelerated variable sample-size stochastic gradient descent algorithm (Algorithm 2) so as to improve the rate of convergence. Such a scheme has been employed for smooth strongly convex [28], smooth convex [22], and nonsmooth (but smoothable) convex [28] stochastic optimization problems with associated rates of convergence given by $O(q^k)$, $O(1/k^2)$, and O(1/k), respectively. The present paper takes a crucial step towards developing CLTs and confidence regions for the optimal solution.

Algorithm 2 Variance-reduced Accelerated SGD

Given arbitrary initial values $x_0 = y_0 \in \mathbb{R}^m$, positive constants α, β , and a positive integer sequence $\{N_k\}_{k\geq 0}$. Then iterate the following equations for $k \geq 0$.

$$y_{k+1} = x_k - \alpha(\nabla f(x_k) + w_{k,N_k}),$$
 (10a)

$$x_{k+1} = y_{k+1} + \beta(y_{k+1} - y_k), \tag{10b}$$

where w_{k,N_k} is defined as in (3), $\alpha > 0$ and $\beta > 0$ are constant steplengths.

Next, we establish an upper bound on the expected sub-optimality gap of the iterates generated by Algorithm 2. Its proof is similar to that in [3]; hence it is omitted here but included in the supplementary material for completeness. This is an important preliminary result to be used in the rate analysis of Algorithm 2.

Lemma 2 Let Algorithm 2 be applied to the problem (1). Suppose Assumption 1 holds and $\alpha \in (0, \frac{1}{L}]$. Set $\beta = \frac{1-\gamma}{1+\gamma}$ with $\gamma = \sqrt{\alpha \eta}$. Then for any $k \ge 1$,

$$\mathbb{E}[f(y_k) - f^*] \le \frac{(\eta + L)(1 - \gamma)^k}{2} \mathbb{E}[\|x_0 - x^*\|^2] + \nu^2 \left(\alpha + \frac{(1 - \gamma)\gamma}{2\eta}\right) \sum_{i=0}^{k-1} \frac{(1 - \gamma)^i}{N_{k-1-i}}.$$
(11)

2.3 The Heavy-Ball Method.

The classical heavy-ball method of Polyak [48] takes the form $x_{k+1} = x_k - \alpha \nabla f(x_k) + \beta(x_k - x_{k-1})$ with a steplength $\alpha > 0$ and a momentum parameter $\beta > 0$. For the class of quadratic and strongly convex functions, the author in [48] derives optimal step-size parameters and achieves the linear convergence rate $(\sqrt{L} - \sqrt{\eta})/(\sqrt{L} + \sqrt{\eta})$. However for the class of non-quadratic but *L*-smooth and η -strongly convex functions, the authors in [2] provide the global linear rate of convergence; however, this rate does not lead to an acceleration compared with unaccelerated gradient descent method (cf. [23, Lemma 2.5]). Stochastic variants of the heavy-ball method have been employed widely in practice (cf. [31,59,60] for applications to machine learning). Recent efforts [20,65] have analyzed the rate in stochastic settings and [65] derived a sublinear rate $\mathcal{O}(1/\sqrt{k})$ for general Lipschitz continuous convex objectives with bounded variance, while a rate $\mathcal{O}(1/k^{\beta})$ with $\beta \in (0, 1)$ was provided in [20] for the case of the strongly convex quadratic functions. Convergence properties remain unaddressed in variance-reduced regimes. Therefore, we consider a variable sample-size variant of the stochastic heavy ball method assuming constant steplengths, in contrast with the diminishing steplengths utilized in [20, 65] and prove the global linear convergence of the iterates. In the developed Algorithm 3, we add a momentum term $\beta(x_k - x_{k-1})$ to the variable sample-size stochastic gradient step (2) and obtain the update (12).

Algorithm 3 Variance-reduced heavy-ball SGD

Given an arbitrary initial value $x_0 \in \mathbb{R}^m$, two positive constants $\alpha, \beta > 0$, and a positive integer sequence $\{N_k\}_{k>0}$. Set $x_{-1} = x_0$. Then iterate the following equation for $k \ge 0$.

$$x_{k+1} = x_k - \alpha(\nabla f(x_k) + w_{k,N_k}) + \beta(x_k - x_{k-1}),$$
(12)

where w_{k,N_k} is defined as in (3), $\alpha > 0$ and $\beta > 0$ are constant steplengths.

In the following lemma, we first give an upper bound on the expected mean-squared error of $\{x_k\}$ for quadratic cost functions based on [49]. The proof can be found in Appendix A

Lemma 3 Suppose that Assumption 1 holds and f is a quadratic function with $\nabla^2 f(x) \equiv \mathbf{H}$ for any $x \in \mathbb{R}^m$. Consider Algorithm 3, where $\alpha \in (0, 4/L)$ and $\beta \triangleq \max\{|1 - \sqrt{\alpha \eta}|^2, |1 - \sqrt{\alpha L}|^2\} < 1$. Then

for any $\iota \in (0, 1 - \sqrt{\beta})$, there exist a constant $c(\iota)$ such that the following holds for any $k \ge 0$.

$$\mathbb{E}\left[\left\| \begin{pmatrix} x_{k+1} - x^* \\ x_k - x^* \end{pmatrix} \right\|^2 \right] \le 2(c(\iota))^2 (\sqrt{\beta} + \iota)^{2(k+1)} \left[\|x_0 - x^*\|^2 \right] + \alpha^2 \nu^2 (c(\iota))^2 \sum_{t=0}^k \frac{(\sqrt{\beta} + \iota)^{2(k-t)}}{N_t}.$$
 (13)

Additionally, we follow the idea of [2] in establishing the convergence rate of Algorithm 3 for L-smooth and η -strongly convex functions that are non-quadratic. The proof is similar to that of [2], hence it is omitted here but included in the supplementary material for completeness.

Lemma 4 Suppose that Assumption 1 holds. Consider Algorithm 3, where $\beta \in (0, 1)$ and $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$. Then for any $k \ge 0$.

$$\mathbb{E}[\|x_k - x^*\|^2] \le \frac{1}{2\hat{m}\eta} \mathbb{E}\left[\|x_0 - x^*\|^2 + \hat{m}(f(x_0) - f(x^*))\right] q^{k+1} + \frac{\alpha^2 \nu^2}{2(1-\beta)^2 \hat{m}\eta} \sum_{i=0}^k q^i / N_{k-i}.$$
 (14)

where $\hat{m} \triangleq \frac{2\alpha}{1-\beta} \left(\frac{\beta - \alpha \eta}{1-\beta} + \frac{2\eta}{\eta+L} \right)$ and $q = \max\{q_1, q_2\}$ with

$$q_1 \triangleq \frac{\beta}{\beta + \eta \left(\frac{2(1-\beta)}{L+\eta} - \alpha\right)} < 1 \text{ and } q_2 \triangleq \max\left\{0, 1 - \frac{2\alpha\eta L}{(1-\beta)(L+\eta) + 2\beta L}\right\}.$$
(15)

Based on the above two lemmas, for both the quadratic and non-quadratic strongly convex functions, we will separately derive both a geometric rate and asymptotic normality statements in Section 3 for Algorithm 3 under geometrically increasing batch-sizes, whereas the corresponding results of Algorithm 3 under polynomially increasing batch-sizes will be established in Section 4.

2.4 Pathway for addressing constrained and nonsmooth regimes.

We believe that the presented avenues hold promise for contending with constrained and nonsmooth regimes. *(i) Constrained problems.* Consider the constrained problem

$$\min_{x \in \mathcal{X}} f(x) \triangleq \mathbb{E}[F(x, \boldsymbol{\xi})], \tag{16}$$

where \mathcal{X} is a closed and convex set with a nonempty interior. Algorithm 1 can be extended to a constrained regime with an additional projection onto the convex set \mathcal{X} . Under suitable conditions, Lemma 1 holds and x_k converges almost surely to the optimal solution x^* . For the case when the optimal solution x^* lies in the interior of \mathcal{X} , the sequence $\{x_k\}_{k\geq K}$ will lie in the interior of the constrained set in an almost sure sense for sufficiently large K. Then by proceeding in a similar fashion as in Section 3, CLTs may be developed for constrained problems. If however, solutions are on the boundary of the set, a possible resolution may lie in computing CLTs of approximate minimizers of an unconstrained reformulation via penalization and barrier methods [44].

(*ii*) Nonsmooth problems. When the f is not necessarily smooth, then one avenue lies in employing smoothing approaches. Under some conditions [1], one can construct an (a, b)-smoothed convex approximation of

f, denoted by f_{η} where $f_{\eta}(x) \leq f(x) \leq f_{\eta}(x) + \eta b$ and $\|\nabla f_{\eta}(x) - \nabla f_{\eta}(y)\| \leq \frac{a}{\eta} \|x - y\|$ for all x, y. One may then integrate iterative smoothing within the variance-reduced accelerated gradient scheme as follows.

$$y_{k+1} = x_k - \alpha_k (\nabla f_{\eta_k}(x_k) + w_{k,N_k}),$$

$$x_{k+1} = y_{k+1} + \beta_k (y_{k+1} - y_k).$$

Under suitable assumptions on α_k , β_k , η_k , N_k , the sequence $\{y_k\}$ converges to a unique solution of the original problem. Under an assumption that f is smooth in a neighborhood of x^* , one can again develop CLTs in this setting. If however, f is not necessarily smooth at x^* , then one avenue might lie in developing C^2 smoothing-based approximations and provide CLTs for ϵ -solutions.

A comprehensive examination of (i) and (ii) is beyond the scope of this paper but we believe the smooth and unconstrained analysis presented in this paper is a crucial building block.

3 Central Limit Theorems under Geometrically Increasing Batch-size.

In this section, we establish CLTs for Algorithms 1, 2, and 3 when the number of sampled gradients, denoted by N_k , increases at a geometric rate.

3.1 Rate and Oracle Complexities.

Based on Lemmas 1–4, we can establish the geometric rate of convergence along with the iteration and oracle complexity guarantees for Algorithms 1-3. Related results regarding linear convergence for stochastic gradient methods can be found in [9, 19, 29, 46, 54], while a linear rate of the accelerated variants has been provided in [28, 53]. The proofs of Propositions 1-3 are similar to those in our prior work [34, Theorem 4.2 and Corollary 4.7].

Proposition 1 (Rate and Oracle Complexity for Algorithm 1) Let Assumption 1 hold and $\alpha \in (0, \frac{2}{\eta+L}]$. Consider Algorithm 1, where $N_k \triangleq \lceil \rho_1^{-(k+1)} \rceil$ for some $\rho_1 \in (q, 1)$ with $q \triangleq 1 - \frac{2\alpha\eta L}{\eta+L}$. Then

$$\mathbb{E}[\|x_k - x^*\|^2] \le \rho_1^k \left(\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2 \nu^2}{1 - q/\rho_1} \right), \quad \forall k \ge 1.$$
(17)

The iteration and oracle complexity for computing an ϵ -solution, defined as $\mathbb{E}[||x_k - x^*||^2] \leq \epsilon$, are $\mathcal{O}(\kappa \ln(1/\epsilon))$ and $\mathcal{O}(\kappa/\epsilon)$, where $\kappa \triangleq \frac{L}{\eta}$ denotes the condition number.

Proof. By substituting $N_k = \lceil \rho_1^{-(k+1)} \rceil$ into (7), using $q = 1 - \frac{2\alpha\eta L}{\eta + L}$ and $\rho_1 \in (q, 1)$, one obtains

$$\mathbb{E}[\|x_{k+1} - x^*\|^2] \le q\mathbb{E}[\|x_k - x^*\|^2] + \alpha^2 \nu^2 \rho_1^{k+1} \le q^{k+1}\mathbb{E}[\|x_0 - x^*\|^2] + \alpha^2 \nu^2 \sum_{t=0}^k q^t \rho_1^{k+1-t}$$
$$= q^{k+1}\mathbb{E}[\|x_0 - x^*\|^2] + \alpha^2 \nu^2 \rho_1^{k+1} \sum_{t=0}^k (q/\rho_1) \le \left(\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2 \nu^2}{1 - q/\rho_1}\right) \rho_1^{k+1}.$$

Hence, we derive (17). Suppose we set $\alpha \triangleq \frac{2}{\eta+L}$ and $\rho_1 \triangleq \left(\frac{\kappa}{\kappa+1}\right)^2 > q = \left(\frac{\kappa-1}{\kappa+1}\right)^2$. From (17), it follows that $\mathbb{E}[\|x_k - x^*\|^2] \leq \epsilon$ for any $k \geq K(\epsilon)$, where

$$K(\epsilon) \triangleq \frac{\ln\left(\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2 \nu^2}{1 - q/\rho_1}\right) + \ln(1/\epsilon)}{2\ln(1 + \frac{1}{\kappa})}.$$
(18)

It is noticed that $\ln(1 + 1/\kappa) \ge (1 - \frac{\kappa}{\kappa+1}) = \frac{1}{(\kappa+1)}$. Consequently, the number of iterations required to obtain an ϵ -optimal solution in a mean-squared sense, i.e. $\mathbb{E}[||x - x^*||^2] \le \epsilon$, is $\mathcal{O}(\kappa \ln(1/\epsilon))$. Finally, based on the iteration complexity bound (18), we achieve the following oracle complexity bound, measured by the number of sampled gradients, for obtaining an ϵ -optimal solution:

$$\sum_{k=0}^{K(\epsilon)-1} N_k \le \sum_{k=1}^{K(\epsilon)} \rho_1^{-k} \le \int_1^{K(\epsilon)+1} \rho_1^{-t} dt \le \frac{\rho_1^{-(K(\epsilon)+1)}}{\ln(1/\rho_1)} = \frac{\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2 \nu^2}{1 - q/\rho_1}}{2\epsilon \rho_1 \ln(1 + 1/\kappa)} = \mathcal{O}\left(\frac{\kappa}{\epsilon}\right).$$

In the following lemma, we show that the result of Proposition 1 holds as well when the noise condition Assumption 1(iii) is replaced by some state-dependent noise condition.

Lemma 5 Consider Algorithm 1, where $N_k \triangleq \lceil \rho_1^{-(k+1)} \rceil$ for some $\rho_1 \in (q, 1)$ with $q \triangleq 1 - \frac{2\alpha\eta L}{\eta+L}$. Let $\alpha \in (0, \frac{2}{\eta+L}]$ and Assumptions 1(i)-(ii) hold. In addition, suppose that there exist constants $\nu_1, \nu_2 > 0$ such that

$$\mathbb{E}[w_{k,N_k}|\mathcal{F}_k] = 0 \text{ and } \mathbb{E}[\|w_{k,N_k}\|^2|\mathcal{F}_k] \le \frac{\nu_1^2 + \nu_2^2 \|x_k\|^2}{N_k}, \ a.s., \ \forall k \ge 0.$$
(19)

Then there exists a constant c > 0 such that the following holds.

$$\mathbb{E}[\|x_{k+1} - x^*\|^2] \le \rho_1^k \left(\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2 c}{1 - q/\rho_1} \right), \quad \forall k \ge 0.$$
(20)

Further, the iteration and oracle complexity for computing an ϵ *-solution are bounded by* $\mathcal{O}(\kappa \ln(1/\epsilon))$ *and* $\mathcal{O}(\kappa/\epsilon)$ *, respectively.*

Proof. Since x_k is adapted to \mathcal{F}_k , by taking expectations conditioned on \mathcal{F}_k on both sides of (8), using $\alpha \in (0, \frac{2}{n+L}]$, (9) and (19), we obtain the following.

$$\mathbb{E}[\|x_{k+1} - x^*\|^2 | \mathcal{F}_k] \le \left(1 - \frac{2\alpha\eta L}{\eta + L}\right) \|x_k - x^*\|^2 + \frac{\alpha^2(\nu_1^2 + \nu_2^2 \|x_k\|^2)}{N_k}, \quad a.s.$$

By taking unconditional expectations, using $N_k \triangleq \lceil \rho_1^{-(k+1)} \rceil$ and $||x_k||^2 \leq 2(||x_k - x^*||^2 + ||x^*||^2)$, we obtain that

$$\mathbb{E}[\|x_{k+1} - x^*\|^2] \le \left(1 - \frac{2\alpha\eta L}{\eta + L} + 2\alpha^2\nu_2^2\rho_1^{k+1}\right)\mathbb{E}[\|x_k - x^*\|^2] + \alpha^2\rho_1^{k+1}(\nu_1^2 + 2\nu_2^2\|x^*\|^2).$$
(21)

Next, we show that $\mathbb{E}[||x_k - x^*||^2]$ is uniformly bounded by some constants.

Case 1: If $2\alpha^2\nu_2^2 \leq \rho_1 - q$, then $1 - \frac{2\alpha\eta L}{\eta + L} + 2\alpha^2\nu_2^2\rho_1^{k+1} \leq q + 2\alpha^2\nu_2^2\rho_1 \triangleq \tilde{q} < q + 2\alpha^2\nu_2^2 \leq \rho_1$. Hence, we conclude that $\tilde{q} < \rho_1$. Similarly to the derivation of (17), we obtain from (21) that for any $k \geq 0$,

$$\mathbb{E}[\|x_k - x^*\|^2] \le \tilde{q}\mathbb{E}[\|x_k - x^*\|^2] + \alpha^2 \rho_1^k(\nu_1^2 + 2\nu_2^2\|x^*\|^2) \le \rho_1^k\left(\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2(\nu_1^2 + 2\nu_2^2\|x^*\|^2)}{1 - \tilde{q}/\rho_1}\right)$$

$$\leq \mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2(\nu_1^2 + 2\nu_2^2 \|x^*\|^2)}{1 - \tilde{q}/\rho_1} \triangleq c_1.$$

Case 2: If $2\alpha^2\nu_2^2 > \rho_1 - q$, we define $\tilde{k} \triangleq \left[\frac{\ln\left(\frac{2\alpha^2\nu_2^2}{\rho_1 - q}\right)}{\ln(\rho_1^{-1})}\right]$. Then for any $k \ge \tilde{k}$, $\rho_1^{-(k+1)} > \frac{2\alpha^2\nu_2^2}{\rho_1 - q}$ and hence $2\alpha^2\nu_2^2\rho_1^{k+1} < \rho_1 - q$. Note from (21) that for any $k \le \tilde{k}$,

$$\mathbb{E}[\|x_k - x^*\|^2] \le (1 + 2\alpha^2 \nu_2^2 \rho_1) \mathbb{E}[\|x_{k-1} - x^*\|^2] + \alpha^2 (\nu_1^2 + 2\nu_2^2 \|x^*\|^2)$$

$$\le (1 + 2\alpha^2 \nu_2^2 \rho_1)^{\tilde{k}} \Big(\mathbb{E}[\|x_0 - x^*\|^2] + \alpha^2 (\nu_1^2 + 2\nu_2^2 \|x^*\|^2) \frac{(1 + 2\alpha^2 \nu_2^2 \rho_1)^{\tilde{k}} - 1}{(1 + 2\alpha^2 \nu_2^2 \rho_1) - 1} \Big) \triangleq c_2.$$

By defining $\hat{q} \triangleq q + 2\alpha^2 \nu_2^2 \rho_1^{\tilde{k}+1}$, we have $1 - \frac{2\alpha\eta L}{\eta+L} + 2\alpha^2 \nu_2^2 \rho_1^{k+1} \le \hat{q} < \rho_1$ for any $k \ge \hat{k}$. Then it follows from (21) that for any $k \ge \tilde{k}$,

$$\mathbb{E}[\|x_{k+1} - x^*\|^2] \le \hat{q}\mathbb{E}[\|x_k - x^*\|^2] + \alpha^2 \rho_1^{k+1}(\nu_1^2 + 2\nu_2^2 \|x^*\|^2) \\ \le \hat{q}^{k+1-\tilde{k}}\mathbb{E}[\|x_{\tilde{k}} - x^*\|^2] + \alpha^2(\nu_1^2 + 2\nu_2^2 \|x^*\|^2) \sum_{t=0}^{k-\tilde{k}} \rho_1^{k+1-t} \hat{q}^t \le c_2 + \frac{\alpha^2(\nu_1^2 + 2\nu_2^2 \|x^*\|^2)}{1-\hat{q}/\rho_1}.$$

Thus, for the case $2\alpha^2\nu_2^2 > \rho_1 - q$, we have that $\mathbb{E}[\|x_k - x^*\|^2] \le c_2 + \frac{\alpha^2(\nu_1^2 + 2\nu_2^2\|x^*\|^2)}{1 - \tilde{q}/\rho_1}$ for any $k \ge 1$. By combing $2\alpha^2\nu_2^2 \le \rho_1 - q$ (**Case 1**) and $2\alpha^2\nu_2^2 > \rho_1 - q$ (**Case 2**), we conclude that there exits some

By combing $2\alpha^2\nu_2^2 \le \rho_1 - q$ (Case 1) and $2\alpha^2\nu_2^2 > \rho_1 - q$ (Case 2), we conclude that there exits some constant $c_3 > 0$ such that $\mathbb{E}[||x_k - x^*||^2] \le c_3$ for any $k \ge 1$. This combines with (21) produces

$$\mathbb{E}[\|x_{k+1} - x^*\|^2] \le \left(1 - \frac{2\alpha\eta L}{\eta + L}\right) \mathbb{E}[\|x_k - x^*\|^2] + \alpha^2 \rho_1^{k+1} \underbrace{(\nu_1^2 + 2\nu_2^2 \|x^*\|^2 + 2\nu_2^2 c_3)}_{\triangleq_c}]$$
$$= q \mathbb{E}[\|x_k - x^*\|^2] + \alpha^2 c \rho_1^{k+1}.$$

The rest of the proof is the same as that of Proposition 1.

Next, we provide similar statements for Algorithm 2, and Algorithm 3 (for both quadratic and nonquadratic objective functions), for which the proofs are omitted here but included in the supplementary material for completeness.

Proposition 2 (Rate and Oracle Complexity for Algorithm 2) Let Assumption 1 hold. Consider Algorithm 2, where $\alpha \in (0, \frac{1}{L}]$, $\gamma \triangleq \sqrt{\alpha \eta}$, $\beta \triangleq \frac{1-\gamma}{1+\gamma}$, and $N_k \triangleq \lceil \rho_2^{-(k+1)} \rceil$ with $\rho_2 \in (1-\gamma, 1)$. Then for any $k \ge 0$,

$$\mathbb{E}[f(y_k)] - f^* \le \rho_2^k \left(\frac{\eta + L}{2} \mathbb{E}[\|x_0 - x^*\|^2] + \frac{\rho_2 \nu^2}{\rho_2 - (1 - \gamma)} \left(\alpha + \frac{(1 - \gamma)\gamma}{2\eta} \right) \right),$$
(22)

and $\mathbb{E}[\|x_k - x^*\|^2] \leq c\rho_2^k$ for some constant c > 0. In addition, the iteration and oracle complexity for obtaining an ϵ -solution are $\mathcal{O}(\sqrt{\kappa}\ln(1/\epsilon))$ and $\mathcal{O}(\sqrt{\kappa}/\epsilon)$, respectively.

Proposition 3 (Rate and Oracle Complexity for Algorithm 3 on Quadratic Functions) Suppose that Assumption 1 holds and $f(\cdot)$ is a quadratic function with $\nabla^2 f(x) \equiv \mathbf{H}$ for any $x \in \mathbb{R}^m$. Consider Algorithm 3, where $\alpha \triangleq \frac{4}{(\sqrt{\eta}+\sqrt{L})^2}$, $\beta \triangleq \left(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}\right)^2$, and $N_k \triangleq \lceil \rho_3^{-(k+1)} \rceil$ with $\rho_3 \in (\beta, 1)$. Then for any $\iota \in (0, \sqrt{\rho_3} - \sqrt{\beta})$, there exist a constant $c(\iota)$ such that

$$\mathbb{E}\left[\left\| \begin{pmatrix} x_{k+1} - x^* \\ x_k - x^* \end{pmatrix} \right\|^2 \right] \le c(\iota)^2 \left(2\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2 \nu^2}{1 - (\sqrt{\beta} + \iota)^2/\rho_3} \right) \rho_3^{k+1}.$$
(23)

In addition, the iteration and oracle complexity required for obtaining an ϵ -solution in a mean-squared sense are respectively $\mathcal{O}(\sqrt{\kappa}\ln(1/\epsilon))$ and $\mathcal{O}(\sqrt{\kappa}/\epsilon)$.

Proposition 4 (Rate for Algorithm 3 on Non-Quadratic Functions) Let Assumption 1 hold. Consider Algorithm 3, where $\beta \in (0, 1)$, $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$, and $N_k \triangleq \lceil \rho_4^{-(k+1)} \rceil$ with $\rho_4 \in (q, 1)$. Here $q = \max\{q_1, q_2\}$ with q_1 and q_2 defined in (15). Then for any $k \ge 0$.

$$\mathbb{E}[\|x_k - x^*\|^2] \le \frac{1}{2\hat{m}\eta} \left(\mathbb{E}\left[\|x_0 - x^*\|^2 + \hat{m}(f(x_0) - f(x^*))\right] + \frac{\alpha^2 \nu^2}{(1-\beta)^2 (1-q/\rho_4)} \right) \rho_4^{k+1},$$
(24)

where $\hat{m} \triangleq \frac{2\alpha}{1-\beta} \left(\frac{\beta-\alpha\eta}{1-\beta} + \frac{2\eta}{\eta+L} \right)$.

Remark 1 From the complexity statements in Propositions 1 and 2, we see that the dependence on the condition number is improved from κ (in Algorithm 1) to $\sqrt{\kappa}$ by the accelerated Algorithm 2. The rate and oracle complexity for the heavy-ball method (Algorithm 3) of quadratic function are similar to that observed in Algorithm 2, while Algorithm 3 for non-quadratic functions might not lead an acceleration (in the sense of the dependence on κ in the constant) of Algorithm 1 (see also Remark 9). In addition, Propositions 1-3 imply that a smaller κ leads to a smaller constant in the oracle complexity.

Remark 2 (Convergence in Probability) Because mean-squared convergence implies convergence in probability, on the basis of Propositions 1-4, we may conclude that the sequences $\{x_k\}$ and $\{y_k\}$ generated by Algorithms 1-3 converge in probability to the optimal solution x^* .

Remark 3 We discuss the choices employed in Algorithm 2 for different settings of ϵ and κ in the following table. In particularly, we observe that the sample size at the last iterate is relatively modest.

κ	ϵ	K	$\beta = \frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1}$	$\rho\in(\beta,1)$	$N_K = \lceil \rho^{-K} \rceil$
10	10^{-3}	$\sqrt{\kappa}\ln(1/\epsilon) \approx 22$	pprox 0.52	$\rho = 0.55$	$\approx 5.1e5$
10	10^{-3}	$\sqrt{\kappa}\ln(1/\epsilon) \approx 22$	pprox 0.52	$\rho = 0.85$	≈ 36
10^{3}	10^{-3}	$\sqrt{\kappa}\ln(1/\epsilon) \approx 219$	≈ 0.939	$\rho = 0.97$	≈ 789
10^{3}	10^{-4}	$\sqrt{\kappa}\ln(1/\epsilon) \approx 292$	pprox 0.939	$\rho = 0.97$	$\approx 7.3e3$

It can be seen that for well-conditioned problems (κ close to 1), the sampling rate at the last iteration reaches 10^5 when ρ is chosen closer to β and $\epsilon = 10^{-3}$. However, it is possible to choose β closer to 1 to

obtain far more reasonable sampling rates. In more practical settings with higher κ (e.g., $\kappa = 10^3$), it is seen that sampling rates are relatively modest. In addition, by leveraging the parallelizable structure, current multi-core and multi-processing techniques can contend with the challenge of computing gradient estimators with large sample sizes. Finally, it is noticed that most large-scale machine learning problems defined on finite sample spaces with cardinalities of 10^6 to 10^9 or even more. Application of variance-reduced gradient methods (such as SVRG) necessitate taking full gradients intermittently, implying that computing a gradient estimate with this sample-size is well within the reach of current computational constraints.

3.2 Preliminary Lemmas.

Before establishing CLTs for Algorithms 1-3, we first introduce a preliminary CLT on doubly-indexed random variables [11, Lemma 3.3.1]. We state it as Lemma 6, whose proof is found in [24, Chapter 12].

Lemma 6 Let ξ_{kt} , $t = 1, \dots, k$ be *m*-dimensional random vectors. Define

$$\mathbf{S}_{kt} \triangleq \mathbb{E}[\boldsymbol{\xi}_{kt} \boldsymbol{\xi}_{kt}^{T}], \quad \mathbf{R}_{kt} \triangleq \mathbb{E}[\boldsymbol{\xi}_{kt} \boldsymbol{\xi}_{kt}^{T} | \boldsymbol{\xi}_{k1}, \cdots, \boldsymbol{\xi}_{k,t-1}], \text{ and } \mathbf{S}_{k} \triangleq \sum_{t=1}^{k} \mathbf{S}_{kt}$$
(25)

Assume that
$$\mathbb{E}[\boldsymbol{\xi}_{kt}|\boldsymbol{\xi}_{k1},\cdots,\boldsymbol{\xi}_{k,t-1}]=0, \ \sup_{k\geq 1}\sum_{t=1}^{n}\mathbb{E}[\|\boldsymbol{\xi}_{kt}\|^{2}]<\infty,$$
 (26)

$$\lim_{k \to \infty} \mathbf{S}_k = \mathbf{S}, \quad \lim_{k \to \infty} \sum_{t=1}^k \mathbb{E}[\|\mathbf{S}_{kt} - \mathbf{R}_{kt}\|] = 0,$$
(27)

and
$$\lim_{k \to \infty} \sum_{t=1}^{k} \mathbb{E}[\|\boldsymbol{\xi}_{kt}\|^2 I_{[\|\boldsymbol{\xi}_{kt}\| \ge \epsilon]}] = 0, \quad \forall \epsilon > 0.$$

$$(28)$$

Then $\sum_{t=1}^{k} \boldsymbol{\xi}_{kt} \xrightarrow{d} N(0, \mathbf{S}).$

To establish the CLTs for Algorithms 1–3, we further require the following conditions.

Assumption 2 (i) $\nabla f : \mathbb{R}^m \to \mathbb{R}^m$ is differentiable at x^* with Hessian matrix $\mathbf{H} \in \mathbb{R}^{m \times m}$, and

$$\nabla f(x) = \mathbf{H}(x - x^*) + \mathbf{D}(x)(x - x^*) \text{ with } \|\mathbf{D}(x)\| \le R_D \|x - x^*\| \text{ for some } R_D > 0.$$
(29)

(ii) The noise sequence $\{w_{k,N_k}\}$ further satisfies

$$\lim_{k \to \infty} \left(N_k \mathbb{E} \left[w_{k,N_k} w_{k,N_k}^T | \mathcal{F}_k \right] \right) = \lim_{k \to \infty} \left(N_k \mathbb{E} \left[w_{k,N_k} w_{k,N_k}^T \right] \right) = \mathbf{S}_0, \quad a.s., \tag{30}$$

and
$$\lim_{r \to \infty} \sup_{k} \mathbb{E}\left[\left\| \sqrt{N_k} w_{k,N_k} \right\|^2 I_{\left[\| \sqrt{N_k} w_{k,N_k} \| > r \right]} \right] = 0,$$
(31)

where $I_{[a>b]} = 1$ if a > b, and $I_{[a>b]} = 0$, otherwise.

Remark 4 We now provide two easily understood sufficient conditions to guarantee that Assumption 2(*i*) holds. In (A), we assume that the gradient ∇f is twice differentiable with bounded second derivatives, i.e.

f is assumed to be thrice differentiable while in (B), we require that f is twice differentiable with Lipschitz continuous Hessians. Define F as $F(x) \triangleq \begin{pmatrix} F_1(x) \\ \vdots \\ F_m(x) \end{pmatrix} = \nabla f(x)$, where $F_j : \mathbb{R}^m \to \mathbb{R}$ for each $j = 1, \dots, m$.

(A) Suppose F is twice continuously differentiable and that $\nabla^2 F_j(x)$ is bounded in $x \in \mathbb{R}^m$ for $j = 1, \dots, m$. Then for any jth $(j = 1, \dots, m)$ component of F(x), by the second-order mean-value theorem, we know there exists $\tilde{x}^j \in \mathbb{R}^m$ on the line segment connecting x and x^* such that

$$F_{j}(x) = F_{j}(x^{*}) + \nabla F_{j}(x^{*})^{T}(x - x^{*}) + \frac{1}{2}(x - x^{*})^{T}\nabla^{2}F_{j}(\tilde{x}^{j})(x - x^{*})$$

$$= F_{j}(x^{*}) + \nabla F_{j}(x^{*})(x - x^{*}) + \underbrace{\left(\frac{1}{2}\nabla^{2}F_{j}(\tilde{x}^{j})(x - x^{*})\right)^{T}}_{\triangleq \mathbf{D}_{j}(x)}(x - x^{*}).$$
(32)

Since $\|\nabla^2 F_j(x)\| \leq R_{D_j}$ for all x, we have that $\|\mathbf{D}_j(x)\| \leq R_{D_j}\|x - x^*\|$. Thus, we conclude that $\mathbf{D}(x) \triangleq \begin{pmatrix} \mathbf{D}_1(x) \\ \vdots \\ \mathbf{D}_n(x) \end{pmatrix}$ satisfies $\|\mathbf{D}(x)\| = \sqrt{\sum_{j=1}^m \|\mathbf{D}_j(x)\|^2} \leq R_D \|x - x^*\|$, where $R_D = \sqrt{\sum_{j=1}^m R_{D_j}^2}$.

Since $F(x^*) = 0$ and $\nabla F(x^*) = \mathbf{H}$ is a symmetric matrix, we derive (29) by stacking (32) for $j = 1, \dots, m$.

(B). Suppose f is twice continuously differentiable with Lipschitz continuous Hessians with constant L_H . By the integral mean-value theorem for the gradient, we have that

$$\nabla f(x) = \nabla f(x^*) + \nabla^2 f(x^*)(x - x^*) + \int_0^1 \left(\nabla^2 f(x^* + \zeta(x - x^*)) - \nabla^2 f(x^*) \right) (x - x^*) d\zeta$$

= $\mathbf{H}(x - x^*) + \mathbf{D}(x)(x - x^*)$, where $\mathbf{D}(x) \triangleq \int_0^1 \left(\nabla^2 f(x^* + \zeta(x - x^*)) - \nabla^2 f(x^*) \right) d\zeta$.

We now derive a bound on $\mathbf{D}(x)$ as follows.

$$\|\mathbf{D}(x)\| = \left\| \int_0^1 \left(\nabla^2 f(x^* + \zeta(x - x^*)) - \nabla^2 f(x^*) \right) d\zeta \right\|$$

$$\leq \max_{0 \le \zeta \le 1} \left\| \left(\nabla^2 f(x^* + \zeta(x - x^*)) - \nabla^2 f(x^*) \right) \right\| \le L_H \|x - x^*\|.$$

Again, we see that (29) holds.

Remark 5 Since the Lindeberg condition condition (31) is less easily verified, we provide a sufficient condition for the ease of understanding and verification in practice. Suppose there exists a constant $\delta > 0$ and a finite value b > 0 such that $\mathbb{E}\left[\|\sqrt{N_k}w_{k,N_k}\|^{2+\delta}\right] \leq b$ for any $k \geq 0$. Therefore,

$$\mathbb{E}\left[\left\|\sqrt{N_k}w_{k,N_k}\right\|^2 I_{\left[\|\sqrt{N_k}w_{k,N_k}\|>r\right]}\right]$$

$$\leq \frac{1}{r^{\delta}} \mathbb{E}\left[\left\| \sqrt{N_k} w_{k,N_k} \right\|^{2+\delta} I_{\left[\| \sqrt{N_k} w_{k,N_k} \| > r \right]} \right] \leq \frac{1}{r^{\delta}} \mathbb{E}\left[\left\| \sqrt{N_k} w_{k,N_k} \right\|^{2+\delta} \right] \leq \frac{b}{r^{\delta}}.$$

This implies that $\sup_k \mathbb{E}\left[\left\|\sqrt{N_k}w_{k,N_k}\right\|^2 I_{\left[\left\|\sqrt{N_k}w_{k,N_k}\right\|>r\right]}\right] \leq \frac{b}{r^{\delta}}$. Hence (31) holds.

In the following, we establish the central limit theorem of a linear recursion, for which the proof is provided in Appendix B. This result will be applied in the proof of Theorems 1-4.

Lemma 7 Suppose that **P** is a square matrix with spectral radius, denoted by $\rho(\mathbf{P})$, strictly smaller than l (i.e., $\rho(\mathbf{P}) < 1$), $N_k = \lceil \rho^{-(k+1)} \rceil$ with $\rho \in (0,1)$ for any $k \ge 0$, and that $\{w_{k,N_k}\}$ satisfies Assumption 2(ii). Let $\{e_k\}$ be generated by

$$e_{k+1} = \mathbf{P}e_k - \alpha \rho^{-(k+1)/2} \mathbf{G} w_{k,N_k} + \zeta_{k+1},$$
(33)

where $\mathbb{E}[\|e_0\|^2] < \infty$ and $\zeta_k \xrightarrow{P}{k \to \infty} 0$. Then

$$\alpha^{-1}e_k \xrightarrow[k \to \infty]{} N(0, \Sigma) \text{ with } \Sigma \triangleq \sum_{t=0}^{\infty} \mathbf{P}^t \mathbf{GS}_0 \mathbf{G}^T (\mathbf{P}^t)^T.$$

3.3 Central Limit Theorems.

Based on Proposition 1 and Lemma 7, using Assumption 2, we present the central limit theorem for Algorithm 1 with geometrically increasing batch-sizes.

Theorem 1 (CLT of Algorithm 1 with Geometrically increasing N_k) Let Assumptions 1 and 2 hold. Consider Algorithm 1, where $\alpha \in (0, \frac{2}{\eta+L}]$. Set $q \triangleq 1 - \frac{2\alpha\eta L}{\eta+L}$, $N_k \triangleq \lceil \rho_1^{-(k+1)} \rceil$ with $\rho_1 \in (q, 1)$, and $\mathbf{P}_1 \triangleq \rho_1^{-1/2}(\mathbf{I}_m - \alpha \mathbf{H})$. Then $\rho(\mathbf{P}_1) < 1$ and

$$\alpha^{-1}\rho_1^{-k/2}(x_k - x^*) \xrightarrow[k \to \infty]{d} N(0, \Sigma_1), \text{ where } \Sigma_1 \triangleq \sum_{t=0}^{\infty} \mathbf{P}_1^t \mathbf{S}_0 \mathbf{P}_1^t.$$
(34)

Proof. By using (29), we can rewrite (4) as

$$x_{k+1} - x^* = x_k - x^* - \alpha \mathbf{H}(x_k - x^*) - \alpha (\nabla f(x_k) - \mathbf{H}(x_k - x^*) + w_{k,N_k})$$

= $(\mathbf{I}_m - \alpha \mathbf{H})(x_k - x^*) - \alpha (\mathbf{D}(x_k)(x_k - x^*) + w_{k,N_k}).$ (35)

Define $e_k \triangleq \rho_1^{-k/2}(x_k - x^*)$. Then by multiplying both sides of (35) by $\rho_1^{-(k+1)/2}$ and by using the definition $\mathbf{P}_1 \triangleq \rho_1^{-1/2}(\mathbf{I}_m - \alpha \mathbf{H})$, we obtain

$$e_{k+1} = \rho_1^{-1/2} (\mathbf{I}_m - \alpha \mathbf{H}) \rho_1^{-k/2} (x_k - x^*) - \alpha \rho_1^{-(k+1)/2} (\mathbf{D}(x_k) (x_k - x^*) + w_{k,N_k})$$

= $\mathbf{P}_1 e_k - \alpha \rho_1^{-(k+1)/2} w_{k,N_k} + \zeta_{k+1} \text{ with } \zeta_{k+1} \triangleq -\alpha \rho_1^{-1/2} \mathbf{D}(x_k) e_k.$ (36)

In the following, we prove that $\zeta_k \xrightarrow{P} 0$. From (17), it follows that for any $k \ge 0$,

$$\operatorname{var}(e_k) \le \mathbb{E}[\|e_k\|^2] = \mathbb{E}[\|x_k - x^*\|^2 / \rho_1^k] \le \mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2 \nu^2}{1 - q/\rho_1} \triangleq v_e^2 \text{ with } v_e > 0.$$
(37)

Chebyshev's inequality asserts that if X is a random variable with mean μ and variance σ^2 , then for any real number h > 0,

$$\mathbb{P}(\|X-\mu\| \le h\sigma) \ge 1 - h^{-2}.$$
(38)

By setting $h = \chi^{-1/2}$ for any $\chi \in (0, 1)$ and applying (38) where $X = e_k$, we have

$$\mathbb{P}\big(\|e_k - \mathbb{E}[e_k]\| \le \chi^{-1/2} \operatorname{var}(e_k)\big) \ge 1 - \chi.$$
(39)

Define the events $\mathcal{A}_1 \triangleq \{ \|e_k - \mathbb{E}[e_k] \| \leq \chi^{-1/2} \operatorname{var}(e_k) \}$, $\mathcal{A}_2 \triangleq \{ \|e_k - \mathbb{E}[e_k] \| \leq \chi^{-1/2} v_e^2 \}$, $\mathcal{A}_3 \triangleq \{ \|e_k\| \leq \|\mathbb{E}[e_k]\| + \chi^{-1/2} v_e^2 \}$, $\mathcal{A}_4 \triangleq \{ \|e_k\| \leq \mathbb{E}[\|e_k\|] + \chi^{-1/2} v_e^2 \}$, and $\mathcal{A}_5 \triangleq \{ \|e_k\| \leq v_e + \chi^{-1/2} v_e^2 \}$. Note by (37) that $\mathcal{A}_1 \subseteq \mathcal{A}_2$. We observe that $\mathcal{A}_2 \subseteq \mathcal{A}_3$ by the inequality $\|x_1 - x_2\| \geq \|x_1\| - \|x_2\|$. Since $\| \bullet \|$ is convex in x, by the Jensen's inequality we have $\|\mathbb{E}[X]\| \leq \mathbb{E}[\|X\|]$, and hence $\mathcal{A}_3 \subseteq \mathcal{A}_4$. Since $\| \bullet \|^2$ is a convex function, by using (37) and Jensen's inequality, we have $\mathbb{E}[\|e_k\|] \leq \sqrt{\mathbb{E}[\|e_k\|^2]} \leq v_e$ and hence $\mathcal{A}_4 \subseteq \mathcal{A}_5$. Thus, we have $\mathcal{A}_1 \subseteq \mathcal{A}_5$. This together with (39) implies that for any $k \geq 0$,

$$\mathbb{P}(\|e_k\| \le v_e + \chi^{-1/2} v_e^2) = \mathbb{P}(\mathcal{A}_5) \ge \mathbb{P}(\mathcal{A}_1) = \mathbb{P}(\|e_k - \mathbb{E}[e_k]\| \le \chi^{-1/2} \operatorname{var}(e_k)) \ge 1 - \chi.$$

That is to say that for any $\chi \in (0, 1)$,

$$\mathbb{P}(\|e_k\| > v_e + \chi^{-1/2} v_e^2) < \chi, \ \forall k \ge 0.$$
(40)

Therefore, we conclude that $||e_k||$ is bounded in probability (i.e., $||e_k|| = O_P(1)$).

Note by Proposition 1 and the Markov's inequality that $||x_k - x^*|| \xrightarrow{P} 0$, i.e., $||x_k - x^*|| = o_P(1)$. Then by invoking the bound that $||\mathbf{D}(x)|| \le R_D ||x - x^*||$, we conclude that $||\mathbf{D}(x_k)|| = o_P(1)$. Recall from [62, p.12] that symbols $o_P(\cdot)$ and $O_P(\cdot)$ satisfy

$$O_P(1)o_P(1) = o_P(1).$$
 (41)

This together with $\|\mathbf{D}(x_k)\| = o_P(1)$ and $\|e_k\| = O_P(1)$ implies $\|\zeta_{k+1}\| \le \alpha \rho_1^{-1/2} \|\mathbf{D}(x_k)\| \|e_k\| = O_P(1)o_P(1) \stackrel{(41)}{=} o_P(1)$. Thus, we conclude that $\zeta_k \xrightarrow[k \to \infty]{P} 0$.

Since **H** is the Hessian matrix of f(x) at $x = x^*$, by Assumptions 1(i) and 1(ii), we conclude that **H** has eigenvalues $\lambda_1, \dots, \lambda_m$ that satisfy $0 < \eta \le \lambda_m \le \lambda_{m-1} \le \dots \le \lambda_2 \le \lambda_1 \le L$. Then $\|\mathbf{I}_m - \alpha \mathbf{H}\|_2 \le \max\{|1 - \alpha \eta|, |1 - \alpha L|\}$. We first show that $|1 - \alpha \eta| \ge |1 - \alpha L|$. It can be seen that $|1 - \alpha \eta| \ge |1 - \alpha L|$ when $\alpha \in (0, 1/L]$. While for any $\alpha \in [1/L, \frac{2}{\eta+L}]$, $|1 - \alpha \eta| = 1 - \alpha \eta$, $|1 - \alpha L| = \alpha L - 1$, and hence $|1 - \alpha \eta| - |1 - \alpha L| = 2 - \alpha(\eta + L) \ge 0$. Then for any $\alpha \in (0, \frac{2}{\eta+L}]$,

$$\|\mathbf{I}_m - \alpha \mathbf{H}\| \le |1 - \alpha \eta| = 1 - \alpha \eta \le \sqrt{1 - \frac{2\alpha \eta L}{\eta + L}}.$$

The last inequality holds because $\alpha \leq \frac{2}{\eta+L} \Rightarrow \frac{2L}{\eta+L} + \alpha\eta - 2 \leq 0 \Rightarrow \frac{2\alpha\eta L}{\eta+L} + (\alpha\eta)^2 - 2\alpha\eta \leq 0 \Rightarrow (1 - \alpha\eta)^2 \leq 1 - \frac{2\alpha\eta L}{\eta+L}$. Since $q = 1 - \frac{2\alpha\eta L}{\eta+L}$ and $\rho_1 \in (q, 1)$, we have that

$$\|\mathbf{P}_1\| = \rho_1^{-1/2} \|(\mathbf{I}_m - \alpha \mathbf{H})\| \le (q/\rho_1)^{1/2} < 1.$$
(42)

Since the spectral radius of a symmetric matrix \mathbf{P}_1 equals its two norm, i.e., $\rho(\mathbf{P}_1) = \|\mathbf{P}_1\| < 1$, by invoking Lemma 7 and setting $\mathbf{G} = \mathbf{I}_m$ and $\mathbf{P} = \mathbf{P}_1$ (symmetric), we conclude that the sequence $\alpha^{-1}e_k$ generated by (36) converges in distribution to a normally distributed random variable with zero mean and covariance Σ_1 defined as in (34). Therefore, the result follows by recalling that $e_k = \rho_1^{-k/2}(x_k - x^*)$.

Based on Lemma 4, by employing similar proof arguments as in Theorem 1, we conclude that the central limit result established in Theorem 1 holds as well under Assumptions 1(i)-(ii), Assumption 2, and the state-dependent noise condition (19).

Remark 6 Suppose we set $\alpha = \frac{2}{\eta+L}$ and $\rho_1 \triangleq \left(\frac{\kappa}{\kappa+1}\right)^2$ in Algorithm 1. If $V \sim N(0, \mathbf{I}_m)$, it follows from (34) that $\frac{\eta+L}{2} \left(\frac{\kappa+1}{\kappa}\right)^k (x_k - x^*) \xrightarrow{d} N(0, \mathbf{\Sigma}_1)$. Namely,

$$x_k \stackrel{D}{\approx} x^* + \frac{2}{\eta + L} \left(1 - \frac{1}{\kappa + 1} \right)^k \boldsymbol{\Sigma}_1^{1/2} V \text{ for large } k.$$

This result implies that the sequence $\{x_k\}$ converges in distribution to the optimal solution x^* at rate $(\frac{\kappa}{\kappa+1})^k$, and $\{x_k\}$ is asymptotically normally distributed for large k. This provides the possibility of assessing confidence regions of the estimate from the normal distribution. In addition, the estimation error for large k depends on the structure of the studied problem (including η , L, and the Hessian matrix **H**) and probability distribution of the gradient noise, measured through the coupling matrix Σ_1 . Thus, the problem's difficulty is largely characterized by the covariance matrix Σ_1 .

Based on the CLT established in Theorem 1, we proceed to use the Delta method [55] to derive the asymptotic distribution of the gradient and the objective function based on the estimation sequence $\{x(k)\}$.

Corollary 1 Consider Algorithm 1 and suppose all conditions of Theorem 1 hold. Then
(i)
$$\alpha^{-1}\rho_1^{-k/2} \nabla f(x_k) \xrightarrow[k \to \infty]{d} N(0, \mathbf{H}\Sigma_1\mathbf{H}).$$

(ii) $\alpha^{-2}\rho_1^{-k}(f(x_k) - f(x^*)) \xrightarrow[k \to \infty]{d} \frac{1}{2}N(0, \Sigma_1)^T\mathbf{H}N(0, \Sigma_1).$

Proof. (i) Note by Assumption 2(i) that the gradient mapping $\nabla f(x) : \mathbb{R}^m \to \mathbb{R}^m$ is differentiable at x^* with Hessian matrix **H**. By using (34) and the Delta theorem [55, Eqn. (7.182)], we have that

$$\alpha^{-1}\rho_1^{-k/2} \left(\nabla f(x_k) - \nabla f(x^*) \right) \xrightarrow[k \to \infty]{d} \mathbf{H} N(0, \Sigma_1).$$

Then by the fact that $\nabla f(x^*) = 0$ and **H** is symmetric, the assertion (i) holds.

(ii) By using (34), $\nabla f(x^*) = 0$, and the second-order Delta theorem [55, Theorem 7.70], one obtains

$$\alpha^{-2}\rho_1^{-k}\big(f(x_k) - f(x^*)\big) \xrightarrow[k \to \infty]{d} \frac{1}{2}f''_u(x^*) \text{ with } u = N(0, \Sigma_1),$$

where $f''_u(x^*)$ denotes the second order directional derivative at x^* along the direction u. Since $f : \mathbb{R}^m \to \mathbb{R}$ is twice continuously differentiable, $f''_u(x^*) = u^T \mathbf{H} u$. Then result (ii) follows from $u = N(0, \Sigma_1)$.

Akin to Theorem 1, based on Proposition 2 and Lemma 7, we can also establish the CLT for Algorithm 2 with geometrically increasing batch-sizes.

Theorem 2 (CLT for Algorithm 2 with Geometrically Increasing N_k) Suppose Assumptions 1 and 2 hold. Consider Algorithm 2, where $\alpha \in (0, \frac{1}{L}]$, $\gamma \triangleq \sqrt{\alpha \eta}$, $\beta \triangleq \frac{1-\gamma}{1+\gamma}$, and $N_k \triangleq \lceil \rho_2^{-(k+1)} \rceil$ with $\rho_2 \in (1 - \gamma, 1)$. Then

$$\alpha^{-1}\rho_2^{-k/2} \begin{pmatrix} y_k - x^* \\ y_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \boldsymbol{\Sigma}_2) \text{ with } \boldsymbol{\Sigma}_2 \triangleq \sum_{t=0}^{\infty} \mathbf{P}_2^t \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix} (\mathbf{P}_2^t)^T,$$
where $\mathbf{P}_2 \triangleq \rho_2^{-1/2} \begin{pmatrix} (1+\beta)(\mathbf{I}_m - \alpha \mathbf{H}) & -\beta(\mathbf{I}_m - \alpha \mathbf{H}) \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix} \text{ and } \rho(\mathbf{P}_2) < 1.$

Proof. By using (29), we can rewrite (10a) as

$$y_{k+1} - x^* = (\mathbf{I}_m - \alpha \mathbf{H})(x_k - x^*) - \alpha (\mathbf{D}(x_k)(x_k - x^*) + w_{k,N_k}).$$
(43)

Note by (10b) that $x_k - x^* = (1 + \beta)(y_k - x^*) - \beta(y_{k-1} - x^*)$. This together with (43) produces

$$y_{k+1} - x^* = (\mathbf{I}_m - \alpha \mathbf{H})((1+\beta)(y_k - x^*) - \beta(y_{k-1} - x^*)) - \alpha(\mathbf{D}(x_k)(x_k - x^*) + w_{k,N_k})$$

= $(1+\beta)(\mathbf{I}_m - \alpha \mathbf{H})(y_k - x^*) - \beta(\mathbf{I}_m - \alpha \mathbf{H})(y_{k-1} - x^*) - \alpha(\mathbf{D}(x_k)(x_k - x^*) + w_{k,N_k}).$

Define $z_{k+1} \triangleq \begin{pmatrix} y_{k+1} - x^* \\ y_k - x^* \end{pmatrix}$ and $\mathbf{H}_2 \triangleq \begin{pmatrix} (1+\beta)(\mathbf{I}_m - \alpha \mathbf{H}) & -\beta(\mathbf{I}_m - \alpha \mathbf{H}) \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}$. Then based on the above equation, we obtain the following recursion.

$$z_{k+1} = \mathbf{H}_2 z_k - \alpha \begin{pmatrix} \mathbf{D}(x_k)(x_k - x^*) + w_{k,N_k} \\ 0 \end{pmatrix}.$$
(44)

The eigenvalue decomposition of **H** is given by $\mathbf{H} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$, where $\mathbf{\Lambda} \triangleq \text{diag}\{\lambda_1, \lambda_2, \cdots, \lambda_m\}$ and **U** is orthogonal. This allows us to rewrite \mathbf{H}_2 as

$$\mathbf{H}_{2} = \begin{pmatrix} \mathbf{U} & \mathbf{0}_{m} \\ \mathbf{0}_{m} & \mathbf{U} \end{pmatrix} \begin{pmatrix} (1+\beta)(\mathbf{I}_{m} - \alpha \mathbf{\Lambda}) & -\beta(\mathbf{I}_{m} - \alpha \mathbf{\Lambda}) \\ \mathbf{I}_{m} & \mathbf{0}_{m} \end{pmatrix} \begin{pmatrix} \mathbf{U} & \mathbf{0}_{m} \\ \mathbf{0}_{m} & \mathbf{U} \end{pmatrix}^{T}.$$

Then the eigenvalues of the matrix \mathbf{H}_2 are given by the roots of the equation $det(vI_{2m} - \mathbf{H}_2) = 0$, i.e.,

$$det \begin{pmatrix} v\mathbf{I}_m - (1+\beta)(\mathbf{I}_m - \alpha \mathbf{\Lambda}) & \beta(\mathbf{I}_m - \alpha \mathbf{\Lambda}) \\ -\mathbf{I}_m & v\mathbf{I}_m \end{pmatrix} = 0$$

By the property $det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = det(AD - BC)$ when the blocks A, B, C, D are square matrices of the same size and CD = DC [56], we have

$$det\left(v\left(v\mathbf{I}_m - (1+\beta)(\mathbf{I}_m - \alpha\mathbf{\Lambda})\right) + \beta(\mathbf{I}_m - \alpha\mathbf{\Lambda})\right) = 0$$

Since the matrix in the above determinant is a diagonal matrix, it can be described by the following characteristic equations for $i = 1, \dots, m$.

$$v(v - (1 + \beta)(1 - \alpha\lambda_i)) + \beta(1 - \alpha\lambda_i) = v^2 - (1 + \beta)(1 - \alpha\lambda_i)v + \beta(1 - \alpha\lambda_i) = 0.$$
(45)

By $\gamma^2 = \alpha \eta$ and $\beta = \frac{1-\gamma}{1+\gamma}$, we will show that the discriminant of the above quadratic equation is nonpositive for each $i = 1, \dots, m$.

$$\Delta_{i} = (1+\beta)^{2}(1-\alpha\lambda_{i})^{2} - 4\beta(1-\alpha\lambda_{i}) = 4(1-\alpha\lambda_{i})\left(\frac{(1-\alpha\lambda_{i})}{(1+\gamma)^{2}} - \frac{1-\gamma}{1+\gamma}\right)$$
$$= \frac{4(1-\alpha\lambda_{i})}{(1+\gamma)^{2}}\left(\gamma^{2} - \alpha\lambda_{i}\right) = \frac{4\alpha(1-\alpha\lambda_{i})}{(1+\gamma)^{2}}\left(\eta - \lambda_{i}\right) \le 0,$$

where the last inequality holds by the fact that for any $i = 1, \ldots, m, 1 - \alpha \lambda_i \ge 1 - \lambda_i/L \ge 0$ and $\eta - \lambda_i \le 0$. Consequently, (45) has two complex roots $\frac{(1+\beta)(1-\alpha\lambda_i)\pm i\sqrt{-\Delta_i}}{2}$, where i denotes the imaginary part. Thus, the magnitude of the roots is $\sqrt{\left(\frac{(1+\beta)(1-\alpha\lambda_i)}{2}\right)^2 - \frac{\Delta_i}{4}} = \sqrt{\beta(1-\alpha\lambda_i)}$. Since $\gamma^2 = \alpha \eta$, $\beta = \frac{1-\gamma}{1+\gamma}, \rho_2 \in (1-\gamma, 1)$, and $\lambda_i \in [\eta, L]$, we obtain that

$$\rho(\mathbf{H}_2) = \max_i \sqrt{\beta(1 - \alpha\lambda_i)} \le \sqrt{\beta(1 - \alpha\eta)} = \sqrt{\frac{(1 - \gamma)(1 - \gamma^2)}{1 + \gamma}} = 1 - \gamma < \rho_2 < \rho_2^{1/2}.$$
 (46)

Define $\varepsilon_k \triangleq \rho_2^{-k/2} z_k$. By multiplying both sides of (44) by $\rho_2^{-(k+1)/2}$, we obtain the following recursion with $\varsigma_{k+1} \triangleq -\alpha \rho_2^{-(k+1)/2} \mathbf{D}(x_k)(x_k - x^*)$:

$$\varepsilon_{k+1} = \rho_2^{-1/2} \mathbf{H}_2 \varepsilon_k - \alpha \rho_2^{-(k+1)/2} \begin{pmatrix} \mathbf{D}(x_k)(x_k - x^*) + w_{k,N_k} \\ 0 \end{pmatrix} = \mathbf{P}_2 \varepsilon_k - \alpha \rho_2^{-(k+1)/2} \begin{pmatrix} w_{k,N_k} \\ 0 \end{pmatrix} + \begin{pmatrix} \varsigma_{k+1} \\ 0 \end{pmatrix} + \begin{pmatrix} \varphi_{k+1} \\ \varphi_{k+1} \end{pmatrix} + \begin{pmatrix} \varphi_{k+1} \end{pmatrix} + \begin{pmatrix} \varphi_{k+1} \end{pmatrix} + \begin{pmatrix} \varphi_{k+1} \\ \varphi_{k+1} \end{pmatrix} + \begin{pmatrix} \varphi_{k+1} \end{pmatrix} + \begin{pmatrix}$$

We proceed to show that $\varsigma_k \xrightarrow[k \to \infty]{k \to \infty} 0$. By Proposition 2, we see that $\mathbb{E}[\rho_2^{-k} || x_k - x^* ||^2] \leq c$ for some constant c > 0 and $|| x_k - x^* || = o_P(1)$ by the Markov's inequality. Then by invoking the bound that $|| \mathbf{D}(x) || \leq R_D || x - x^* ||$, we achieve $|| \mathbf{D}(x_k) || = o_P(1)$. Similarly to the procedures for proving (40), we can also show that $|| \rho_2^{-k/2}(x_k - x^*) || = O_P(1)$. Therefore, we conclude that

$$\|\varsigma_{k+1}\| \le \alpha \rho_2^{-1/2} \|\mathbf{D}(x_k)\| \|\rho_2^{-k/2}(x_k - x^*)\| = o_P(1)O_P(1) \stackrel{(41)}{=} o_P(1) \Rightarrow \varsigma_{k+1} \xrightarrow{P}_{k \to \infty} 0.$$

Because $\rho(\mathbf{P}_2) = \rho_2^{-1/2} \rho(\mathbf{H}_2) < 1$ by (46), by setting $\mathbf{P} = \mathbf{P}_2$, $\mathbf{G} = \begin{pmatrix} \mathbf{I}_m \\ \mathbf{0}_m \end{pmatrix}$ and using Lemma 7, we obtain that $\alpha^{-1} \varepsilon_k \xrightarrow{d} N(0, \mathbf{\Sigma}_2)$ with $\mathbf{\Sigma}_2 \triangleq \lim_{k \to \infty} \sum_{t=0}^k \mathbf{P}_2^t \mathbf{G} \mathbf{S}_0 \mathbf{G}^T (\mathbf{P}_2^t)^T$. Then by $\mathbf{G} \mathbf{S}_0 \mathbf{G}^T = \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix}$ and $\varepsilon_k = \rho_2^{-k/2} z_k = \rho_2^{-k/2} \begin{pmatrix} y_k - x^* \\ y_{k-1} - x^* \end{pmatrix}$, we prove the result.

Remark 7 By setting $\alpha = \frac{1}{L}$ and $\rho_2 = 1 - \frac{1}{a\sqrt{\kappa}}$ for some a > 1 in Algorithm 2, we have that $1 - \gamma = 1 - \frac{1}{\sqrt{\kappa}} < \rho_2$, and Theorem 2 holds. Thus, we obtain $L\left(1 - \frac{1}{a\sqrt{\kappa}}\right)^{-k/2} \begin{pmatrix} y_k - x^* \\ y_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \Sigma_2)$.

Hence the following holds with $V \sim N(0, \mathbf{I}_{2m})$.

where

$$\begin{pmatrix} y_k \\ y_{k-1} \end{pmatrix} \stackrel{D}{\approx} \begin{pmatrix} x^* \\ x^* \end{pmatrix} + \frac{1}{L} \left(1 - \frac{1}{a\sqrt{\kappa}} \right)^{k/2} \Sigma_2^{1/2} V \text{ for large } k$$

In a similar fashion, we establish the central limit theorem for the stochastic heavy ball method (Algorithm 3) on quadratic objective functions with geometrically increasing batch-sizes based on Proposition 3 and Lemma 7.

Theorem 3 (CLT for Alg. 3 with Geometrically Increasing N_k on Quadratic Functions) Suppose that Assumptions 1 and 2(*ii*) hold, and f is a quadratic function with $\nabla^2 f(x) \equiv \mathbf{H}$ for any $x \in \mathbb{R}^m$. Consider Algorithm 3, where $\alpha \triangleq \frac{4}{(\sqrt{\eta}+\sqrt{L})^2}$, $\beta \triangleq \left(1-\frac{2}{\sqrt{\kappa}+1}\right)^2$, and $N_k \triangleq \lceil \rho_3^{-(k+1)} \rceil$ with $\rho_3 \in (\beta, 1)$ for any $k \ge 0$. Then

$$\alpha^{-1} \rho_3^{-k/2} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \boldsymbol{\Sigma}_3) \text{ with } \boldsymbol{\Sigma}_3 \triangleq \sum_{t=0}^{\infty} \mathbf{P}_3^t \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix} (\mathbf{P}_3^t)^T,$$
$$\mathbf{P}_3 \triangleq \rho_3^{-1/2} \begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha \mathbf{H} & -\beta \mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix} \text{ and } \rho(\mathbf{P}_3) < 1.$$

Proof. As has been shown in the proof of Lemma 3 (Equation (58)), we can rewrite the recursion (12) in a matrix form as follows:

$$\begin{pmatrix} x_{k+1} - x^* \\ x_k - x^* \end{pmatrix} = \underbrace{\begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha \mathbf{H} & -\beta \mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}}_{\triangleq \mathbf{H}_3} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix} - \alpha \begin{pmatrix} w_{k,N_k} \\ 0 \end{pmatrix}.$$

Define $\varepsilon_k \triangleq \rho_3^{-k/2} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix}$. By multiplying the above equation with $\rho_3^{-(k+1)/2}$, one obtains

$$\varepsilon_{k+1} = \mathbf{P}_3 \varepsilon_k - \alpha \rho_3^{-(k+1)/2} \begin{pmatrix} \mathbf{I}_m \\ \mathbf{0}_m \end{pmatrix} w_{k,N_k},$$

where $\mathbf{P}_3 = \rho_3^{-1/2} \mathbf{H}_3$. Thus, $\rho(\mathbf{P}_3) = \rho_3^{-1/2} \rho(\mathbf{H}_3) < 1$ by recalling $\rho(\mathbf{H}_3) = \sqrt{\beta}$ from (60). This together with Assumption 2(ii) proves the result by using Lemma 7.

Remark 8 Suppose we set
$$\rho_3 = \left(1 - \frac{1}{\sqrt{\kappa}+1}\right)^2$$
 in Algorithm 3, then $\rho_3 > \beta$ and Theorem 3 holds; i.e.,

$$\frac{(\sqrt{\eta}+\sqrt{L})^2}{4} \left(1 - \frac{1}{\sqrt{\kappa}+1}\right)^{-k} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \Sigma_3).$$
 If we denote $V \sim N(0, \mathbf{I}_{2m})$, then

$$\begin{pmatrix} x_k \\ x_{k-1} \end{pmatrix} \stackrel{D}{\approx} \begin{pmatrix} x^* \\ x^* \end{pmatrix} + \frac{4}{(\sqrt{\eta}+\sqrt{L})^2} \left(1 - \frac{1}{\sqrt{\kappa}+1}\right)^k \Sigma_3^{1/2} V \text{ for large } k.$$

Next, based on Proposition 4 and Lemma 7, we also show the CLT of Algorithm 3 on non-quadratic objective functions.

Theorem 4 (CLT for Alg. 3 with Geometrically Increasing N_k **on Non-Quadratic Functions)** Let Assumption 1 and 2 hold. Consider Algorithm 3, where $\beta = |1 - \sqrt{\alpha \eta}|^2$ and $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$. Let $q = \max\{q_1, q_2\}$ with q_1 and q_2 defined in (15). Set $N_k \triangleq \lceil \rho_4^{-(k+1)} \rceil$ with $\rho_4 \in (q, 1)$. Then

$$\alpha^{-1}\rho_4^{-k/2} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \hat{\boldsymbol{\Sigma}}_3) \text{ with } \hat{\boldsymbol{\Sigma}}_3 \triangleq \sum_{t=0}^{\infty} \mathbf{P}_4^t \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix} \begin{pmatrix} \mathbf{P}_4^t \end{pmatrix}^T$$

where $\mathbf{P}_4 \triangleq \rho_4^{-1/2} \begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha \mathbf{H} & -\beta \mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}$ and $\rho(\mathbf{P}_4) < 1$.

Proof. From (12) and Assumption 2(a) it follows that

$$x_{k+1} - x^* = x_k - x^* + \beta(x_k - x_{k-1}) - \alpha \mathbf{H}(x_k - x^*) - \alpha \mathbf{D}(x_k)(x_k - x^*) - \alpha w_{k,N_k}.$$

Then we may rewrite the recursion in a matrix form as follows:

$$\begin{pmatrix} x_{k+1} - x^* \\ x_k - x^* \end{pmatrix} = \underbrace{\begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha \mathbf{H} & -\beta \mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}}_{\triangleq \mathbf{H}_4} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix} - \alpha \begin{pmatrix} \mathbf{D}(x_k)(x_k - x^*) + w_{k,N_k} \\ 0 \end{pmatrix}.$$

Define $\varepsilon_k \triangleq \rho_4^{-k/2} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix}$. By multiplying the above equation with $\rho_4^{-(k+1)/2}$, one obtains

$$\varepsilon_{k+1} = \mathbf{P}_4 \varepsilon_k - \alpha \rho_4^{-(k+1)/2} \begin{pmatrix} \mathbf{I}_m \\ \mathbf{0}_m \end{pmatrix} w_{k,N_k} + \begin{pmatrix} \varsigma_{k+1} \\ 0 \end{pmatrix} \text{ with } \varsigma_{k+1} \triangleq -\alpha \rho_4^{-(k+1)/2} \mathbf{D}(x_k) (x_k - x^*),$$

where $\mathbf{P}_{4} = \rho_{4}^{-1/2} \mathbf{H}_{4}$.

Next, we show that $\rho(\mathbf{H}_4) \leq \sqrt{q}$ similar to how we showed tha $\rho(\mathbf{H}_2) \leq 1 - \gamma$ in Theorem 2. Recall the eigenvalue decomposition $\mathbf{H} = \mathbf{U}\mathbf{\Lambda}\mathbf{U}^T$, where \mathbf{U} is orthogonal and $\mathbf{\Lambda} \triangleq \operatorname{diag}\{\lambda_1, \lambda_2, \cdots, \lambda_m\}$ with satisfy $0 < \eta \leq \lambda_m \leq \lambda_{m-1} \leq \cdots \leq \lambda_2 \leq \lambda_1 \leq L$. Then we can rewrite

$$\mathbf{H}_4 = \begin{pmatrix} \mathbf{U} & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{U} \end{pmatrix} \begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha\mathbf{\Lambda} & -\beta\mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix} \begin{pmatrix} \mathbf{U} & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{U} \end{pmatrix}^T.$$

Similarly to the derivation of (45), the eigenvalues of H_4 can be described by the following characteristic equations.

$$p_i(v) = (v - (1 + \beta - \alpha \lambda_i))v + \beta = v^2 - (1 + \beta - \alpha \lambda_i)v + \beta = 0, \quad i = 1, \cdots, m.$$

Since $0 < \eta \le \lambda_m \le \lambda_{m-1} \le \dots \le \lambda_2 \le \lambda_1 \le L$, we have $|1 - \sqrt{\alpha \lambda_i}| \le \max\{|1 - \sqrt{\alpha \eta}|, |1 - \sqrt{\alpha L}|\}$. Note from $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$ that $\alpha - \left(\frac{2}{\sqrt{L}+\sqrt{\eta}}\right)^2 < \frac{2}{L+\eta} - \frac{4}{L+\eta+2\sqrt{\eta L}} \le 0$. Thus, $|1 - \sqrt{\alpha \eta}| \ge |1 - \sqrt{\alpha L}|$ and $|1 - \sqrt{\alpha \lambda_i}| \le |1 - \sqrt{\alpha \eta}| = \beta$. Then similar to (59), the discriminant of the equation $p_i(v) = 0$ is

$$\Delta_i = (1 + \beta - \alpha \lambda_i)^2 - 4\beta = \beta^2 - 2\beta(1 + \alpha \lambda_i) + (1 - \alpha \lambda_i)^2 = (\beta - (1 + \alpha \lambda_i))^2 - 4\alpha \lambda_i \le 0.$$

Then the spectral radius of \mathbf{H}_4 is

$$\rho(\mathbf{H}_4) = \sqrt{\beta} \ge |1 - \sqrt{\alpha \eta}| = 1 - \sqrt{\alpha \eta} \text{ since } \alpha \eta < 1.$$
(47)

Since $\kappa > 1$ and $\beta \in (0, 1)$, we have $(1 - \beta)(\kappa + 1) + 2\kappa\beta = \kappa + 1 + \beta(\kappa - 1) > \kappa + 1 > \frac{4\kappa}{\kappa+1}$. Therefore,

$$(1-\beta)^{2}(\kappa+1) + 2\kappa\beta(1-\beta) > \frac{4\kappa(1-\beta)}{\kappa+1} = \frac{4(1-\beta)L}{L+\eta} > 2\alpha L.$$

$$\Rightarrow 1-\beta > \frac{2\alpha L}{(1-\beta)(\kappa+1)+2\kappa\beta} = \frac{2\alpha L\eta}{(1-\beta)(L+\eta)+2L\beta}$$

$$\Rightarrow \beta < 1 - \frac{2\alpha L\eta}{(1-\beta)(L+\eta)+2L\beta} \stackrel{(15)}{\leq} q_{2}.$$

This together with (47) implies that $\rho(\mathbf{H}_4) < \sqrt{q_2} \le \sqrt{q}$ since $q = \max\{q_1, q_2\}$.

From Proposition 4 it follows that $\mathbb{E}[\rho_4^{-k} || x_k - x^* ||^2] \leq C$ for some constant C > 0 and hence $|| x_k - x^* || = o_P(1)$. Then by invoking the bound that $|| \mathbf{D}(x) || \leq R_D || x - x^* ||$, we obtain that $|| \mathbf{D}(x_k) || = o_P(1)$. Similarly to the procedures for proving (40), we can show that $|| \rho_4^{-k/2}(x_k - x^*) || = O_P(1)$. Therefore,

$$\|\varsigma_{k+1}\| \le \alpha \rho_3^{-1/2} \|\mathbf{D}(x_k)\| \|\rho_3^{-k/2}(x_k - x^*)\| = o_P(1)O_P(1) \stackrel{(41)}{=} o_P(1) \Rightarrow \varsigma_{k+1} \xrightarrow{P}_{k \to \infty} 0.$$

Since $\rho(\mathbf{P}_4) = \rho_4^{-1/2} \rho(\mathbf{H}_4) = \rho_4^{-1/2} \sqrt{q} < 1$ by $\rho(\mathbf{H}_4) < \sqrt{q}$ and $\rho_4 > q$, the result follows by invoking Lemma 7.

Since $\beta = |1 - \sqrt{\alpha \eta}|^2$ and $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$ imply that β is defined in an implicit sense, we add the following remark to clarify their existence and show how does the spectral radius depends on the condition number κ .

Remark 9 By substituting $\beta = |1 - \sqrt{\alpha \eta}|^2$ into the upper bound of α , it requires that

$$\alpha < \frac{2(1-|1-\sqrt{\alpha\eta}|^2)}{L+\eta} \Leftrightarrow \alpha(L+\eta) < 4\sqrt{\alpha\eta} - 2\alpha\eta \Leftrightarrow \alpha(L+3\eta) < 4\sqrt{\alpha\eta} \Leftrightarrow \alpha < \frac{16\eta}{(L+3\eta)^2}.$$

Then by (47), we achieve $\rho(\mathbf{H}_4) \geq 1 - \sqrt{\alpha \eta} > 1 - \sqrt{\frac{16\eta^2}{(L+3\eta)^2}} = 1 - \sqrt{\frac{16}{(\kappa+3)^2}} = 1 - \frac{4}{\kappa+3} \geq 1 - \frac{4}{\kappa}$. In comparison with the geometric rate $1 - 1/\sqrt{\kappa+1}$ of Algorithm 3 for quadratic functions, the geometric parameter of Algorithm 3 non-quadratic functions will not exceed $1 - \frac{4}{\kappa}$. Therefore, it does not lead to an acceleration compared with variable sample-size stochastic gradient algorithm (Algorithm 1), which is consistent with the statement in [23, Lemma 2.5].

4 Central Limit Theorems on Polynomial Batch-size.

There are many settings where a geometric increase in N_k is impractical. For instance, the generation of a sampled gradient is computationally expensive; an example of this commonly arises in simulation optimization problems in the context of large manufacturing or queueing simulations. To this end, we consider the use of polynomial increases in sample-size, an avenue that allows for more gentle growth, and proceed to investigate the rate of convergence and the associated central limit statements for Algorithms 1– 3. Polynomial rates have been studied in [34] as well as [46] but remain unaware of rate and complexity statements as well as the ensuing CLTs in accelerated and heavy-ball regimes.

4.1 Rate of Convergence.

We first recall some preliminary results that find utility in the rate analysis of the proposed algorithms with the polynomially increasing batch-sizes.

Lemma 8 (*i*) [35, Eqn. (17)] For any $q \in (0, 1)$ and v > 0, there holds

$$\sum_{t=1}^{k} q^{k-t} t^{-v} \le q^k \frac{e^{2v} q^{-1} - 1}{1-q} + \frac{2k^{-v}}{q \ln(1/q)}$$

(*ii*) [35, Lemma 4] For any $q \in (0,1)$ and v > 0, $q^x \le c_{q,v} x^{-v}$ for all x > 0 where $c_{q,v} \triangleq e^{-v} \left(\frac{v}{\ln(1/q)}\right)^v$.

Based on Lemmas 1-4 in Section 2 and Lemma 8, we can establish polynomial rates of convergence of the iterates generated by the three proposed methods. Omitted proofs are included in the supplementary material for purposes of completeness.

Proposition 5 (Rate statement for Algorithm 1 under polynomially increasing N_k) Suppose Assumption 1 holds and that $N_k \triangleq \lceil (k+1)^v \rceil$ for some v > 0. Consider Algorithm 1 with $\alpha \in (0, \frac{2}{\eta+L}]$. Define $q \triangleq 1 - \frac{2\alpha\eta L}{\eta+L}$ and $c_{q,v} \triangleq e^{-v} \left(\frac{v}{\ln(1/q)}\right)^v$. Then $\mathbb{E}[\|x_k - x^*\|^2] \le \left(\underbrace{c_{q,v}\mathbb{E}[\|x_0 - x^*\|^2] + \frac{\alpha^2\nu^2 c_{q,v}(e^{2v}q^{-1} - 1)}{1 - q} + \frac{2\alpha^2\nu^2}{q\ln(1/q)}}_{Q}\right)k^{-v}, \quad \forall k \ge 1.$ (48)

Then the number of iterations and sampled gradients required to obtain an ϵ -optimal solution in the meansquared sense (i.e. $\mathbb{E}[||x - x^*||^2] \leq \epsilon$) are $\mathcal{O}(v(1/\epsilon)^{1/v})$ and $\mathcal{O}(e^v v^v(1/\epsilon)^{1+1/v})$, respectively.

Proposition 6 (Rate statement for Algorithm 2 under polynomially increasing N_k) Let Algorithm 2 be applied to (1), where $N_k \triangleq \lceil (k+1)^v \rceil$ with some v > 0. Suppose Assumption 1 holds and $\alpha \in (0, \frac{1}{L}]$. Define $\gamma \triangleq \sqrt{\alpha \eta}$ and $\beta \triangleq \frac{1-\gamma}{1+\gamma}$. Then there exists a constant C(v) > 0 such that

$$\mathbb{E}[f(y_k)] - f^* \le C(v)k^{-v}, \quad \forall k \ge 1.$$

Remark 10 Since $f(x) - f(x^*) \ge \frac{\eta}{2} ||x - x^*||^2$, we have $\mathbb{E}[||y_k - x^*||^2] \le \frac{2C_v}{\eta} k^{-v}$. Then from (10b) it follows that $\mathbb{E}[||x_k - x^*||^2] \le 2(1 + \beta)^2 \mathbb{E}[||y_k - x^*||^2] + 2\beta^2 \mathbb{E}[||y_{k-1} - x^*||^2] \le ck^{-v}$ for some c > 0. Thus, the mean-squared error of Algorithm 2 also displays the polynomial rate of convergence similar to that shown in Proposition 5 for Algorithm 1. Because the mean-squared convergence implies converges in probability, the sequences $\{x_k\}$ and $\{y_k\}$ generated by Algorithm 2 satisfy $x_k \xrightarrow{P}{k \to \infty} x^*$ and $y_k \xrightarrow{P}{k \to \infty} x^*$ when the conditions of Proposition 6 hold.

Proposition 7 (Rate of Alg. 3 on quadratic Functions with polynomially increasing N_k) Suppose that Assumption 1 holds and that f is a quadratic function with $\nabla^2 f(x) \equiv \mathbf{H}$ for any $x \in \mathbb{R}^m$. Consider Algorithm 3, where $\alpha \triangleq \frac{4}{(\sqrt{\eta} + \sqrt{L})^2}$, $\beta \triangleq (\frac{\sqrt{\kappa} - 1}{\sqrt{\kappa} + 1})^2$, and $N_k \triangleq \lceil (k+1)^v \rceil$ with v > 0. Then there exists a constant C(v) > 0 such that $\mathbb{E}\left[\left\| \begin{pmatrix} x_{k+1} - x^* \\ x_k - x^* \end{pmatrix} \right\|^2 \right] \le C(v)(k+1)^{-v}, \forall k \ge 0.$

Proposition 8 (Rate of Alg. 3 on Non-Quadratic Functions with polynomially increasing N_k) Let Assumption 1 hold. Consider Algorithm 3, where $\beta \in (0, 1)$, $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$, and $N_k \triangleq \lceil (k+1)^v \rceil$ with v > 0. Then there exists a constant C(v) > 0 such that

$$\mathbb{E}[\|x_k - x^*\|^2] \le C(v)(k+1)^{-v}, \ \forall k \ge 0.$$

4.2 Central Limit Theorems under Polynomially Increasing N_k.

In this part, we first establish the asymptotic normality of a time-varying linear recursion and provide the proof in Appendix C. This result will be applied in proving Theorems 5-7.

Lemma 9 Suppose that the square matrix **A** satisfies $\rho(\mathbf{A}) < 1$, $N_k = \lceil (k+1)^v \rceil$ with v > 0, and $\{w_{k,N_k}\}$ satisfies Assumption 2(ii). Let the sequence $\{e_k\}$ be generated by

$$e_{k+1} = \mathbf{A}_k e_k - \alpha (k+1)^{\nu/2} \mathbf{G} w_{k,N_k} + \zeta_{k+1}, \quad \mathbb{E}[\|e_0\|^2] < \infty,$$
(49)

where $\mathbf{A}_0 = \mathbf{A}$, $\mathbf{A}_k \triangleq \left(\frac{k+1}{k}\right)^{\nu/2} \mathbf{A}$ for any $k \ge 1$, and $\zeta_k \xrightarrow{P}{k \to \infty} 0$. Then

$$\alpha^{-1}e_k \xrightarrow{d} N(0, \mathbf{\Sigma}) \text{ with } \mathbf{\Sigma} \triangleq \lim_{k \to \infty} \sum_{t=1}^k \left(\frac{k}{t}\right)^v \mathbf{A}^{k-t} \mathbf{GS}_0 \mathbf{G}^T \left(\mathbf{A}^T\right)^{k-t}$$

Based on Proposition 5 and Lemma 9, by using Assumption 2, we are now ready to derive the associated central limit theorem for Algorithm 1 with polynomially increasing batch-sizes.

Theorem 5 (CLT of Algorithm 1 with Polynomially increasing Batch-sizes) Suppose that Assumptions 1 and 2 hold. Consider Algorithm 1, where $\alpha \in (0, \frac{2}{\eta+L}]$ and $N_k \triangleq \lceil (k+1)^v \rceil$ with some v > 0. Define $\mathbf{A} \triangleq \mathbf{I}_m - \alpha \mathbf{H}$. Then $\rho(\mathbf{A}) < 1$ and

$$\alpha^{-1}k^{\nu/2}(x_k - x^*) \xrightarrow[k \to \infty]{d} N(0, \Sigma_4) \text{ with } \Sigma_4 \triangleq \lim_{k \to \infty} \sum_{t=1}^k \left(\frac{k}{t}\right)^{\nu} \mathbf{A}^{k-t} \mathbf{S}_0 \mathbf{A}^{k-t}.$$

Proof. We begin by noting that $x_k \xrightarrow[k \to \infty]{P} x^*$ and (35) holds. Define $e_0 \triangleq x_0 - x^*$, $e_k \triangleq k^{\nu/2}(x_k - x^*)$ for any $k \ge 1$, and $\zeta_{k+1} \triangleq -\alpha(k+1)^{\nu/2} \mathbf{D}(x_k)(x_k - x^*)$ for any $k \ge 0$. By multiplying both sides of (35) with $(k+1)^{\nu/2}$, and using $\mathbf{A}_k = \left(\frac{k+1}{k}\right)^{\nu/2} (\mathbf{I}_m - \alpha \mathbf{H})$, we achieve

$$e_{k+1} = \left(\frac{k+1}{k}\right)^{\nu/2} (\mathbf{I}_m - \alpha \mathbf{H}) k^{\nu/2} (x_k - x^*) - \alpha (k+1)^{\nu/2} (\mathbf{D}(x_k) (x_k - x^*) + w_{k,N_k}) = \mathbf{A}_k e_k - \alpha (k+1)^{\nu/2} w_{k,N_k} + \zeta_{k+1}, \quad \forall k \ge 1.$$
(50)

By setting k = 0 in (35), and using $A_0 = A = I_m - \alpha H$, we see that

$$e_1 = x_1 - x^* = \mathbf{A}_0(x_0 - x^*) - \alpha w_{1,N_1} - \alpha \mathbf{D}(x_0)(x_0 - x^*)$$

Then by $\zeta_1 = -\alpha \mathbf{D}(x_0)(x_0 - x^*)$, we see that the (50) also holds for k = 0. Hence the recursion (50) holds for any $k \ge 0$. From (42) it follows that the symmetric matrix **A** satisfies

$$\rho(\mathbf{A}) = \|\mathbf{A}\|_2 = \|\mathbf{I}_m - \alpha \mathbf{H}\|_2 = \rho^{1/2} \|\mathbf{P}_1\|_2 \le q^{1/2} < 1.$$

We conclude from Proposition 5 that $\mathbb{E}[k^v || x_k - x^* ||^2] \le c$ for some constant c > 0, and $|| x_k - x^* || = o_P(1)$. Then by invoking the bound that $|| \mathbf{D}(x) || \le R_D || x - x^* ||$, we obtain $|| \mathbf{D}(x_k) || = o_P(1)$. Similarly to the procedures for proving (40), we can show that $|| k^{v/2} (x_k - x^*) || = O_P(1)$. Therefore,

$$\|\zeta_{k+1}\| \le \alpha \left(1 + \frac{1}{k}\right)^{\nu/2} \|\mathbf{D}(x_k)\| \|k^{\nu/2} (x_k - x^*)\| = o_P(1)O_P(1) \stackrel{(41)}{=} o_P(1) \Rightarrow \zeta_{k+1} \xrightarrow{P} 0.$$
(51)

Therefore, by using Lemma 9 with $\mathbf{G} = \mathbf{I}_m$, we conclude that $\alpha^{-1}e_k \xrightarrow{d} N(0, \Sigma_4)$. By the definition $e_k = k^{v/2}(x_k - x^*)$, we obtain the result.

Similarly to Theorem 5, based on Proposition 6 and Lemma 9, we now establish the central limit theorem for Algorithm 2 under the assumption of polynomially increasing batch-sizes.

Theorem 6 (CLT of Algorithm 2 under Polynomially increasing Batch-sizes) Suppose Assumptions 1 and 2 hold. Consider Algorithm 2, where $\alpha \in (0, \frac{1}{L}]$ and $N_k = \lceil (k+1)^v \rceil$ with some v > 0. Define $\gamma \triangleq \sqrt{\alpha \eta}$, $\beta \triangleq \frac{1-\gamma}{1+\gamma}$, and $\mathbf{H}_2 \triangleq \begin{pmatrix} (1+\beta)(\mathbf{I}_m - \alpha \mathbf{H}) & -\beta(\mathbf{I}_m - \alpha \mathbf{H}) \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}$. Then $\rho(\mathbf{H}_2) < 1$ and

$$\alpha^{-1}k^{\nu/2}\begin{pmatrix} y_k - x^*\\ y_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \mathbf{\Sigma}_5) \text{ with } \mathbf{\Sigma}_5 \triangleq \lim_{k \to \infty} \sum_{t=1}^k \left(\frac{k}{t}\right)^{\nu} \mathbf{H}_2^{k-t} \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m\\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix} \left(\mathbf{H}_2^T\right)^{k-t}$$

Proof. Define $z_{k+1} \triangleq \begin{pmatrix} y_{k+1} - x^* \\ y_k - x^* \end{pmatrix}$, $\varepsilon_0 \triangleq z_0$, and $\varepsilon_k \triangleq k^{v/2} z_k$ for any $k \ge 1$. Therefore, by defining $\varsigma_{k+1} \triangleq -\alpha(k+1)^{v/2} \mathbf{D}(x_k)(x_k - x^*)$, and multiplying both sides of (44) by $(k+1)^{v/2}$, we obtain that

$$\varepsilon_{k+1} = \left(\frac{k+1}{k}\right)^{\nu/2} \mathbf{H}_2 \varepsilon_k - \alpha (k+1)^{\nu/2} \begin{pmatrix} \mathbf{I}_m \\ \mathbf{0}_m \end{pmatrix} w_{k,N_k} + \begin{pmatrix} \varsigma_{k+1} \\ 0 \end{pmatrix}.$$

By defining $\mathbf{G} \triangleq \begin{pmatrix} \mathbf{I}_m \\ \mathbf{0}_m \end{pmatrix}$, $\mathbf{A}_0 \triangleq \mathbf{H}_2$, and $\mathbf{A}_k \triangleq \left(\frac{k+1}{k}\right)^{v/2} \mathbf{H}_2$, there holds

$$\varepsilon_{k+1} = \mathbf{A}_k \varepsilon_k - \alpha (k+1)^{\nu/2} \mathbf{G} w_{k,N_k} + \begin{pmatrix} \varsigma_{k+1} \\ 0 \end{pmatrix}, \quad \forall k \ge 1.$$
(52)

From (44) it is seen that (52) holds for k = 0 as well. Thus, the recursion (52) holds for any $k \ge 0$.

Note from Remark 10 that $\mathbb{E}[k^v || x_k - x^* ||^2] \le c$ for some constant c > 0, and $|| x_k - x^* || = o_P(1)$. Then by invoking the bound that $|| \mathbf{D}(x) || \le R_D || x - x^* ||$, we derive $|| \mathbf{D}(x_k) || = o_P(1)$. Similarly to the procedures for proving (40), we can show that $|| k^{v/2} (x_k - x^*) || = O_P(1)$. Therefore,

$$\|\varsigma_{k+1}\| \le \alpha \left(1 + \frac{1}{k}\right)^{\nu/2} \|\mathbf{D}(x_k)\| \|k^{\nu/2} (x_k - x^*)\| = o_P(1)O_P(1) \stackrel{(41)}{=} o_P(1).$$

Hence $\varsigma_k \xrightarrow{P}{k \to \infty} 0$. Then by using Lemma 9 and (46), we obtain that

$$\alpha^{-1}\varepsilon_k \xrightarrow[k \to \infty]{} N(0, \Sigma_5), \text{ where } \Sigma_5 \triangleq \lim_{k \to \infty} \sum_{t=0}^k \left(\frac{k}{t}\right)^v \mathbf{H}_2^{k-t} \mathbf{GS}_0 \mathbf{G}^T \left(\mathbf{H}_2^T\right)^{k-t}.$$

Then by the fact that $\mathbf{GS}_0\mathbf{G}^T = \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix}$ and $\varepsilon_k = k^{\nu/2} \begin{pmatrix} y_k - x^* \\ y_{k-1} - x^* \end{pmatrix}$, we prove the result.

Remark 11 Suppose we set $\alpha = \frac{2}{L+\eta}$ in Algorithm 1. Then $q = 1 - \frac{2\alpha\eta L}{\eta+L} = \left(\frac{\kappa-1}{\kappa+1}\right)^2$. By (42), it is seen that the matrix **A**, defined in Theorem 5, satisfies $\|\mathbf{A}\|_2 \leq q^{1/2} = \frac{\kappa-1}{\kappa+1} = 1 - \frac{2}{\kappa+1}$. Suppose $\alpha = \frac{1}{L}$ in Algorithm 2. Then $\gamma = 1/\sqrt{\kappa}$ and by (46), we know that \mathbf{H}_2 , defined in Theorem 2, satisfies $\|\mathbf{H}_2\|_2 = 1 - \gamma = 1 - \frac{1}{\sqrt{\kappa}}$. Thus, we conclude from Theorems 5 and 6 that both the unaccelerated gradient method (Algorithm 1) and its accelerated counterpart (Algorithm 2) have convergence rates with the same order $\|x_k - x^*\| = \mathcal{O}(k^{-\nu/2})$; however the accelerated scheme has a smaller constant than its unaccelerated counterpart since $\|\mathbf{H}_2\|_2 < \|\mathbf{A}\|_2$ due to the fact that $\frac{2}{\kappa+1} \leq \frac{1}{\sqrt{\kappa}}$.

Similar to Theorem 5, based on Proposition 7 for quadratic objective functions (resp. Proposition 8 for non-quadratic objective functions) and Lemma 9, we now state the central limit theorem for Algorithm 3 with polynomially increasing batch-size (proof omitted).

Theorem 7 (CLT of Alg. 3 on Quadratic Functions with polynomially increasing N_k) Let Assumptions 1 and 2(ii) hold. Suppose that f is a quadratic function with $\nabla^2 f(x) \equiv \mathbf{H}$ for any $x \in \mathbb{R}^m$. Consider Algorithm 3, where $\alpha \triangleq \frac{4}{(\sqrt{\eta}+\sqrt{L})^2}$, $\beta \triangleq \left(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}\right)^2$, and $N_k = \lceil (k+1)^v \rceil$, v > 0. Set $\mathbf{H}_3 \triangleq \left(\begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha\mathbf{H} & -\beta\mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}$. Then $\rho(\mathbf{H}_3) < 1$ and

$$\alpha^{-1}k^{\nu/2} \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \mathbf{\Sigma}_6) \text{ with } \mathbf{\Sigma}_6 \triangleq \lim_{k \to \infty} \sum_{t=1}^k \left(\frac{k}{t}\right)^{\nu} \mathbf{H}_3^{k-t} \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix} \left(\mathbf{H}_3^T\right)^{k-t}$$

Theorem 8 (CLT of Alg. 3 on Non-Quadratic Functions with polynomially increasing N_k) Let Assumptions 1 and 2 hold. Consider Algorithm 3, where $\beta = |1 - \sqrt{\alpha \eta}|^2$, $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$, and $N_k = \lceil (k+1)^v \rceil$, v > 0. Set $\mathbf{H}_4 \triangleq \begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha \mathbf{H} & -\beta \mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}$. Then $\rho(\mathbf{H}_4) < 1$ and $\alpha^{-1}k^{v/2}\begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix} \xrightarrow{d} N(0, \hat{\boldsymbol{\Sigma}}_6)$ with $\hat{\boldsymbol{\Sigma}}_6 \triangleq \lim_{k \to \infty} \sum_{t=1}^k \begin{pmatrix} k \\ t \end{pmatrix}^v \mathbf{H}_4^{k-t} \begin{pmatrix} \mathbf{S}_0 & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{0}_m \end{pmatrix} (\mathbf{H}_4^T)^{k-t}$.

5 Confidence Regions of the Optimal Solution.

A crucial motivation for developing CLTs lies in developing confidence statements. In this section, we proceed to construct the confidence regions for the optimal solution x^* . Note that the limiting covariance

matrix is dependent on the Hessian at the solution, which is unavailable. Furthermore, we do not have an consistent estimator for the covariance matrix. Yet, in the absence of such an estimate, we proceed to develop rigorous confidence statements, adopting an approach developed in [27].

Since Algorithms 1, 2, and 3 lead to similar central limit results, we show how to construct confidence regions merely for the sequence $\{x_k\}$ generated by Algorithm 1. The simulation framework in [27] lies in generating *n* independent replications of Algorithm 1, leading to *n* copies of the random iterate x_k , denoted by x_{1k}, \dots, x_{nk} . Then the sample mean and the covariance estimator are respectively defined as

$$\bar{\mathbf{x}}_k = \frac{1}{n} \sum_{i=1}^n x_{ik} \text{ and } \mathbf{S}_k = \frac{1}{n-1} \sum_{i=1}^n (x_{ik} - \bar{\mathbf{x}}_k) (x_{ik} - \bar{\mathbf{x}}_k)^T.$$
 (53)

Based on [6, Theorem 4 and Corollary 1] and [27, Theorem 2], we may achieve the following result. The proof is given in Appendix D for completeness.

Proposition 9 Consider Algorithm 1. Suppose that the conditions of Theorem 1 hold and $n \ge m+1$. Then (i) $\sqrt{n}\alpha^{-1}\rho_1^{-k/2}(\bar{\mathbf{x}}_k - x^*) \xrightarrow[k \to \infty]{d} N(0, \Sigma_1)$. (ii) $n(\bar{\mathbf{x}}_k - x^*)^T \mathbf{S}_k^{-1}(\bar{\mathbf{x}}_k - x^*) \xrightarrow[k \to \infty]{d} \xrightarrow[k \to \infty]{m-m} F(m, n-m)$, where F(m, n-m) denotes the F-distribution with (m, n-m) degrees of freedom.

Proposition 9 can be used to construct the confidence region of the optimal solution. Define

$$X_{mk}(z) \triangleq \left\{ x \mid n(\bar{\mathbf{x}}_k - x)^T \mathbf{S}_k^{-1}(\bar{\mathbf{x}}_k - x) \le \frac{m(n-1)}{n-m} z \right\},\tag{54}$$

where z is selected such that $\mathbb{P}(F(m, n - m) \leq z) \geq 1 - \delta$ with some $\delta \in (0, 1)$. Then we have the following corollary.

Corollary 2 ([27, Proposition 3]) Consider Algorithm 1. Suppose that $n \ge m + 1$ and the conditions of Theorem 1 hold. Then the confidence region $X_{mk}(z)$ defined in (54) is asymptotically correct, i.e., $\lim_{k\to\infty} \mathbb{P}(x^* \in X_{mk}(z)) = 1 - \delta.$

The above result asserts that the estimated confidence region $X_{mk}(z)$ asymptotically covers the optimal solution x^* with probability $100(1 - \delta)\%$. The approach is easily implementable because it merely requires n independent replications of Algorithm 1, while without requiring a consistent estimator of the covariance matrix of the stationary normal distribution. The confidence regions of Algorithm 2 and Algorithm 3 can be constructed in a similar way. When it is expensive to run multiple independent trials, an alternative avenue for reducing the complexity requirements can be found in [67], where the authors employ a batch-means method for constructing the confidence region. This framework is reliant on a cancellation approach to "cancel" out the covariance matrix that is hard to estimate. This extension will be considered in future work.

In the following, we show that the sequence $\sqrt{\frac{n(n-m)}{n-1}}S_k^{-1/2}(\bar{x}_k - x^*)$ converges to a suitably defined multivariate t distribution, the proof of which is given in Appendix E.

Proposition 10 Consider Algorithm 1. Suppose that the conditions of Theorem 1 hold and n > m + 2. Then

$$\sqrt{\frac{n(n-m)}{n-1}} \mathbf{S}_k^{-1/2} (\bar{\mathbf{x}}_k - x^*) \xrightarrow{d} T_{n-m}(0, \mathbf{I}_m, m),$$
(55)

. . .

where a m-variate random vector $X \sim T_v(\mu, \mathbf{\Lambda}, m)$ (i.e., X is a t distribution with mean μ , covariance matrix $v(v-2)^{-1}\mathbf{\Lambda}, v > 2$) if it has a probability density function h given by

$$h(u) = \frac{\Gamma((v+m)/2)}{(\pi v)^{v/2} \Gamma(v/2) |\mathbf{A}|^{1/2}} \left\{ 1 + \frac{(u-\mu)^T \mathbf{A}^{-1}(u-\mu)}{v} \right\}^{-\frac{v+m}{2}}, v > 2.$$
(56)

The above result can also be adopted to construct the confidence regions. Define

$$\widetilde{X}_{mk}(\mathbf{U}) \triangleq \left\{ x \mid \sqrt{\frac{n(n-m)}{n-1}} \mathbf{S}_k^{-1/2}(\bar{\mathbf{x}}_k - x) \in \mathbf{U} \right\},$$

where the region U is selected such that $\mathbb{P}(T_{n-m}(0, \mathbf{I}_m, m) \in \mathbf{U}) \ge 1 - \delta$ with some $\delta \in (0, 1)$. Similarly to Corollary 2, the confidence region $\tilde{X}_{mk}(\mathbf{U})$ defined above is asymptotically correct, i.e., $\lim_{k\to\infty} \mathbb{P}(x^* \in \tilde{X}_{mk}(\mathbf{U}))) = 1 - \delta$. However, it might be harder to construct the confidence regions from the multivariate t distribution than that from the F-distribution since the region U might not be easily obtained.

6 Numerical Simulations.

In this section, we carry out simulations for the parameter estimation problem. We aim to estimate the unknown *m*-dimensional parameter x^* based on the gathered scalar measurements $\{d_k\}_{k\geq 1}$ given by $d_k = u_k^T x^* + \nu_k$, where $u_k \in \mathbb{R}^m$ denotes the regression vector and $\nu_k \in \mathbb{R}$ denotes the local observation noise. Assume that $\{u_k\}$ and $\{\nu_k\}$ are mutually independent i.i.d. Gaussian sequences with distributions $N(\mathbf{0}, R_u)$ and $N(0, \sigma_{\nu}^2)$, respectively. Suppose the covariance matrix R_u is positive definite. Then we might model the parameter estimation problem as the following stochastic optimization problem:

$$\min_{x \in \mathbb{R}^m} f(x) \triangleq \mathbb{E}\left[\|d_k - u_k^T x\|^2 \right].$$
(57)

Thus, $f(x) = (x - x^*)^T R_u(x - x^*) + \sigma_\nu^2$ and $\nabla f(x) = R_u(x - x^*)$. Because the Hessian matrix R_u of the objective function f is positive definite, x^* is the unique optimal solution to (57). Suppose that we can observe the regressor u_k and the measurement d_k , then the noisy observation of the gradient $\nabla f(x)$ can be constructed as $u_k u_k^T x - d_k u_k$. Set the dimension of x^* as m = 5. We run Algorithm 1 with $\alpha = \frac{2}{L+\eta}$, Algorithm 2 with $\alpha = \frac{1}{L}$ and $\beta = \frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}$, and Algorithm 3 with $\alpha = \frac{4}{(\sqrt{\eta}+\sqrt{L})^2}$ and $\beta = \left(\frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}\right)^2$, where $x_0 = y_0 = 0$ and the batch-size $N_k = \lceil \rho^{-k} \rceil$ with $\rho = \frac{\kappa^2}{(\kappa+1)^2}$. In the remainder of this section, VSS-SGD, VSS-ACC, VSS-HB represent the abbreviations of Variance-Reduced SGD (Algorithm 3), respectively.

Convergence rate, iteration and oracle complexity. We run Algorithms 1, 2, 3, and the standard SGD algorithm $x_{k+1} = x_k - \alpha_k \nabla f(x_k, \xi_{j,k})$ with $\alpha_k = R_u^{-1}/k$ setting to be the optimal tuning steplength,







Figure 1: Linear Rate

Figure 2: Iteration Complexity

Figure 3: Oracle Complexity

and terminate the schemes when $\frac{\mathbb{E}[||x_k - x^*||_2]}{||x^*||_2} \leq 10^{-3}$. We then examine their empirical rate of convergence, iteration and oracle complexity. Here the empirical mean is calculated by averaging across 100 trajectories. The convergence rate of the relative error $\frac{\mathbb{E}[||x_k - x^*||_2]}{||x^*||_2}$ is shown in Figure 1, which demonstrates that the iterates generated by Algorithms 1-3 converge in mean to the optimal solution at a linear rate. We see that the accelerated scheme (Algorithm 2) has the fastest empirical rate, while the heavy ball method (Alg. 3) tends to stabilize later in the process. The empirical relationship between the accuracy ϵ and $K(\epsilon)$ is shown in Figure 2, where $K(\epsilon)$ denotes the number of iterations required to make $\frac{\mathbb{E}[||x_k - x^*||_2]}{||x^*||_2} < \epsilon$. It is seen that the standard SGD algorithm requires far more iterations than the proposed variance reduced schemes for obtaining an approximate solution with the same accuracy. The empirical relationship between ϵ and $O(\epsilon)$ is shown in Figure 3, where $O(\epsilon)$ denotes the number of sampled gradients required to make $\frac{\mathbb{E}[||x_k - x^*||_2]}{||x^*||_2} < \epsilon$. We observe that for obtaining an estimate with the same accuracy, the accelerated scheme (Alg. 3) requires the smallest number of sampled gradients, while the variable sample-size SGD method (Alg. 1) requires more sampled gradients than standard SGD.



Figure 4: Histograms of $\rho^{-k/2}(x_k^5 - x^*)$ at k = 50 along fitted normal distributions

Limiting distributions. We run 1000 independent sample paths of Algorithms 1, 2, and 3 and terminate at k = 50. The empirically obtained largest eigenvalue of the covariance matrix S_k (estimated by (53)) at k = 50 are 1.648×10^{-5} , 1.649×10^{-6} , 1.825×10^{-6} , respectively. This might imply that the accelerated scheme (Algorithm 2) has the best performance measured by the covariance of the stationary distribution. Since the unknown parameter x^* is multi-dimensional, we merely show the limiting distribution of one component of the rescaled error $\rho^{-k/2}(x_k - x^*)$. The histograms of $\rho^{-k/2}(x_k^5 - x^*)$ at k = 50 are shown in Figure 4 along with the fitted normal distribution (the red curve), where x_k^5 denotes the fifth component



Figure 5: Histograms of $\rho^{-k}(f(x_k) - f(x^*))$ at k = 50

of x_k . It is also seen that the rescaled error is (approximately) normally distributed and among these, the accelerated scheme has the smallest variance. In addition, the histograms of the rescaled suboptimality gap $\rho^{-k}(f(x_k) - f(x^*))$ at k = 50 are shown in Figure 5. We can further conclude that the empirical sub-optimality gap of the accelerated scheme is the best among the proposed variable sample-size schemes. We further run the SGD method $x_{k+1} = x_k - R_u^{-1} \nabla f(x_k, \xi_{j,k})/k$. The histogram along with the fitted normal distribution of $\sqrt{N}(x_N^5 - x^*)$ is displayed in Figure 6, while the empirical sub-optimality gap $N(f(x_N) - f(x^*))$ is shown in Figure 7, where $N \triangleq \sum_{k=1}^{50} N_k$ denotes the number of sampled gradients utilized by Algorithms 1-3.





Figure 6: Hist. along fitted normal for $\sqrt{N}(x_N^5 - x^*)$

Figure 7: Hist. of $N(f(x_N) - f(x^*))$

Coverage probability of the constructed confidence region. In a single replication, we generate n independent sample paths for Algorithms 1, 2 and 3 and terminate each sample path when the total number of sampled gradients used reaches N_{max} . Then we can construct the 95% confidence region by (54) and check whether the true parameter lies in the constructed confidence region. We estimate the coverage probability (i.e., the proportion of replications that the confidence region contains the true value) by conducting 1000 replications. The estimated coverage probability for Algorithms 1-3 with different selection of sample paths n and simulation budget N_{max} is shown in Table 1. It shows the impact of the number of independent runs on the quality of the confidence intervals. It can be seen that the coverage probabilities are getting closer to the nominal level of 95% when the number of sample paths n grows larger. In particular, we see that $n \ge 10$ seems to produce relatively accurate confidence bands but for $n \le 8$, the intervals did drop in quality (measured in terms of coverage probability). Thus, we conclude from empirical simulations that a modest number of independent runs can generate relatively high quality confidence intervals. While it is

seen from Table 1 that the simulation budget N_{max} does not significantly impact the coverage probability; however, a larger N_{max} does lead to narrowing of the confidence region.

		VSS-SGD	VSS-ACC	VSS-HB
	n = 6	0.671 ± 0.221	0.652 ± 0.2259	0.665 ± 0.223
$N_{\rm max} = 10^3$	n = 8	0.904 ± 0.0869	0.872 ± 0.1117	0.893 ± 0.0995
	n = 10	0.946 ± 0.0511	0.923 ± 0.0591	0.951 ± 0.0466
	n = 15	0.973 ± 0.0263	0.966 ± 0.0329	0.961 ± 0.0375
	n = 6	0.664 ± 0.2233	0.62 ± 0.2358	0.646 ± 0.2289
$N_{\rm max} = 10^4$	n = 8	0.872 ± 0.117	0.877 ± 0.108	0.896 ± 0.0933
	n = 10	0.929 ± 0.066	0.944 ± 0.0592	0.949 ± 0.0484
	n = 15	0.965 ± 0.0338	0.946 ± 0.0511	0.961 ± 0.0375
	n = 6	0.651 ± 0.2274	0.656 ± 0.2259	0.664 ± 0.2233
$N_{\rm max} = 10^5$	n = 8	0.87 ± 0.1132	0.871 ± 0.1125	0.896 ± 0.0933
	n = 10	0.912 ± 0.083	0.937 ± 0.0591	0.919 ± 0.0745
	n = 15	0.952 ± 0.0457	0.964 ± 0.0347	0.956 ± 0.0421

Table 1: The estimated coverage probability of Algorithms 1-3. The ideal coverage probability is 0.95.

Polynomially increasing batch-sizes. We run Algorithms 1-3 with the batch-size increasing at a polynomial rate $N_k = \lceil k^v \rceil$, v = 2. The convergence rate of the relative error $\frac{\mathbb{E}[\|x_k - x^*\|_2]}{\|x^*\|_2}$ is shown in Figure 8, while the histograms of $k^{v/2}(x_k^5 - x^*)$ at k = 100 along with the fitted normal distribution are shown in Figure 10. It can be seen that the accelerated scheme has the best performance because it displays fastest convergence rate and the rescaled error has smallest variance.



Figure 8: Poly. Batch-size

Figure 9: Convergence rate of VSS-ACC (Alg. 2) for different problem dimension m

The effect of the problem dimension. We take Algorithm 2 as an example. By adding further numerical simulations, we examine the impact of m on algorithm performance. In particular, we implement Algorithm 2 for m = 5, 10, 20, where $\alpha = \frac{1}{L}$, $\beta = \frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}$, $N_k = \max\{\lceil m^{3/4} \rceil, \lceil \rho^{-k} \rceil\}$ with $\rho = \frac{\kappa^2}{(\kappa+1)^2}$,



Figure 10: Histograms of $k^{\nu/2}(x_k^5 - x^*)$ at k = 100

and the initial values $x_0 = y_0 = 0$. The empirical counterpart of the convergence rate of the relative error $\frac{\mathbb{E}[||x_k - x^*||_2]}{||x^*||_2}$ is shown in Figure 9, demonstrating that the larger m leads to a worse rate of convergence. We additionally test the coverage probability of the constructed confidence region for Algorithm 2 for different choices of m. In a single replication, we run Algorithm 2 for n = 30 independent sample paths and terminate them when the total number of sampled gradients used reaches 5000. Then we can construct the 95% confidence region by (54) and check whether the true parameter lies in the constructed confidence region. Finally, we estimate the coverage probability by 1000 replications. The estimated coverage probabilities for m = 5, 10, 20 are respectively $0.956 \pm 0.0421, 0.953 \pm 0.0448$, and 0.945 ± 0.052 .

7 Conclusions

In this work, we considered the strongly convex stochastic optimization problem and proposed three classes of variance reduced stochastic gradient algorithms (unaccelerated, accelerated, and heavy ball methods), where the unavailable exact gradient is approximated by an increasing batch of sampled gradients. We then establish rate and complexity guarantees. Further, we establish amongst the first formal central limit theorems for all the three schemes when the batch-size increased at either a geometric or a polynomial rate. The covariance matrix specifies how problem structure (including the strong convexity parameter, the Lipschitz constant, and the Hessian matrix) and distribution of gradient noise influences the algorithm performance. In addition, we provide an avenue to construct the confidence region of the optimal solution based on the central limit theorems. The paper concludes with an application of the proposed schemes to the stochastic parameter estimation problem to validate our theoretical findings. Yet much remains to be understood about the how one may develop analogs of such statements in constrained and nonsmooth regimes. One avenue for addressing such challenges is via a smoothing framework [1] whereby a stochastic gradient update is taken with respect to a smoothed objective and the smoothing parameter is progressively reduced (cf. [28]). It is also of interest to explore whether the central limit result on the function value can be obtained under weaker conditions. Finally, we intend to examine if confidence regions can be designed

based on the batch-means method [67] when generating independent trials is expensive.

References

- [1] Beck, A.: First-Order Methods in Optimization. SIAM, Philadelphia, PA (2017)
- [2] Bertsekas, D., Tsitsiklis, J.: Parallel and Distributed Computation: Numerical Methods. Prentice-Hall International editions. Prentice-Hall International (1989)
- [3] Bickel, P., Choi, D., Chang, X., Zhang, H., et al.: Asymptotic normality of maximum likelihood and its variational approximation for stochastic blockmodels. The Annals of Statistics 41(4), 1922–1943 (2013)
- [4] Birge, J.R., Louveaux, F.: Introduction to Stochastic Programming: Springer Series in Operations Research. Springer (1997)
- [5] Blum, J.R.: Multidimensional stochastic approximation methods. The Annals of Mathematical Statistics pp. 737–744 (1954)
- [6] Bodnar, T., Mazur, S., Podgórski, K.: Singular inverse wishart distribution and its application to portfolio theory. Journal of Multivariate Analysis 143, 314–326 (2016)
- [7] Borkar, V.S.: Stochastic approximation: a dynamical systems viewpoint, vol. 48. Springer (2009)
- [8] Bottou, L.: Online learning and stochastic approximations. On-line Learning in Neural Networks 17(9), 142 (1998)
- [9] Byrd, R.H., Chin, G.M., Nocedal, J., Wu, Y.: Sample size selection in optimization methods for machine learning. Math. Program. 134(1), 127–155 (2012)
- [10] Chee, J., Toulis, P.: Convergence diagnostics for stochastic gradient descent with constant learning rate. In: International Conference on Artificial Intelligence and Statistics, pp. 1476–1485. PMLR (2018)
- [11] Chen, H.F.: Stochastic approximation and its applications, vol. 64. Springer Science & Business Media (2006)
- [12] Chen, X., Lee, J.D., Tong, X.T., Zhang, Y.: Statistical inference for model parameters in stochastic gradient descent. The Annals of Statistics 48(1), 251–273 (2020)
- [13] DasGupta, A.: Asymptotic theory of statistics and probability. Springer Science & Business Media (2008)
- [14] Defazio, A., Bach, F., Lacoste-Julien, S.: Saga: A fast incremental gradient method with support for non-strongly convex composite objectives. In: Advances in Neural Information Processing Systems, pp. 1646–1654 (2014)

- [15] DeGroot, M.H., Schervish, M.J.: Probability and statistics. Pearson Education (2012)
- [16] Deng, G., Ferris, M.C.: Variable-number sample-path optimization. Math. Program. 117(1-2), 81–109 (2009)
- [17] Dieuleveut, A., Durmus, A., Bach, F.: Bridging the gap between constant step size stochastic gradient descent and markov chains. The Annals of Statistics 48(3), 1348–1382 (2020)
- [18] Fabian, V., et al.: On asymptotic normality in stochastic approximation. The Annals of Mathematical Statistics 39(4), 1327–1332 (1968)
- [19] Friedlander, M.P., Schmidt, M.: Hybrid deterministic-stochastic methods for data fitting. SIAM J. Scientific Computing 34(3), A1380–A1405 (2012)
- [20] Gadat, S., Panloup, F., Saadane, S., et al.: Stochastic heavy ball. Electronic Journal of Statistics 12(1), 461–529 (2018)
- [21] Ghadimi, E., Feyzmahdavian, H.R., Johansson, M.: Global convergence of the heavy-ball method for convex optimization. In: 2015 European control conference (ECC), pp. 310–315. IEEE (2015)
- [22] Ghadimi, S., Lan, G.: Accelerated gradient methods for nonconvex nonlinear and stochastic programming. Mathematical Programming 156(1-2), 59–99 (2016)
- [23] Goujaud, B., Taylor, A., Dieuleveut, A.: Provable non-accelerations of the heavy-ball method. arXiv preprint arXiv:2307.11291 (2023)
- [24] Hall, P., Heyde, C.C.: Martingale limit theory and its application. Academic press (2014)
- [25] Hasminskii, R.Z., Silver, B.: Stochastic approximation and recursive estimation, vol. 47. American Mathematical Soc. (1973)
- [26] Homem-de-Mello, T.: Variable-sample methods for stochastic optimization. ACM Trans. Model. Comput. Simul. 13(2), 108–133 (2003)
- [27] Hsieh, M.H., Glynn, P.W.: Recent advances in simulation optimization: confidence regions for stochastic approximation algorithms. In: Proceedings of the 34th Conference on Winter simulation: exploring new frontiers, pp. 370–376. Winter Simulation Conference (2002)
- [28] Jalilzadeh, A., Shanbhag, U., Blanchet, J., Glynn, P.W.: Smoothed variable sample-size accelerated proximal methods for nonsmooth stochastic convex programs. Stochastic Systems 12(4), 373–410 (2022)
- [29] Jofré, A., Thompson, P.: On variance reduction for stochastic smooth convex optimization with multiplicative noise. Math. Program. 174(1-2), 253–292 (2019)
- [30] Johnson, R., Zhang, T.: Accelerating stochastic gradient descent using predictive variance reduction. In: Advances in neural information processing systems, pp. 315–323 (2013)

- [31] Krizhevsky, A., Sutskever, I., Hinton, G.E.: Imagenet classification with deep convolutional neural networks. In: Advances in neural information processing systems, pp. 1097–1105 (2012)
- [32] Kushner, H., Yin, G.G.: Stochastic approximation and recursive algorithms and applications, vol. 35. Springer Science & Business Media (2003)
- [33] Lamm, M., Lu, S., Budhiraja, A.: Individual confidence intervals for solutions to expected value formulations of stochastic variational inequalities. Math. Program. 165(1), 151–196 (2017)
- [34] Lei, J., Shanbhag, U.V.: Asynchronous variance-reduced block schemes for composite non-convex stochastic optimization: block-specific steplengths and adapted batch-sizes. Optimization Methods and Software 37(1), 264–294 (2022)
- [35] Lei, J., Shanbhag, U.V.: Distributed variable sample-size gradient-response and best-response schemes for stochastic Nash equilibrium problems. SIAM J. Optim. 32(2), 573–603 (2022)
- [36] Lin, P.E.: Some characterizations of the multivariate t distribution. Journal of Multivariate Analysis 2(3), 339–344 (1972)
- [37] Lu, S.: A new method to build confidence regions for solutions of stochastic variational inequalities. Optimization 63(9), 1431–1443 (2014)
- [38] Mandt, S., Hoffman, M.D., Blei, D.M.: Stochastic gradient descent as approximate Bayesian inference. The Journal of Machine Learning Research 18(1), 4873–4907 (2017)
- [39] Meyn, S.P., Tweedie, R.L.: Markov chains and stochastic stability. Springer Science & Business Media (2012)
- [40] Nemirovski, A., Juditsky, A., Lan, G., Shapiro, A.: Robust stochastic approximation approach to stochastic programming. SIAM Journal on optimization 19(4), 1574–1609 (2009)
- [41] Nemirovskiĭ, A., Yudin, D.: Problem Complexity and Method Efficiency in Optimization. A Wiley-Interscience publication. Wiley (1983)
- [42] Nesterov, Y.: Introductory lectures on convex optimization: A basic course, vol. 87. Springer Science & Business Media (2013)
- [43] Nesterov, Y.E.: A method for solving the convex programming problem with convergence rate $O(1/k^2)$. In: Dokl. akad. nauk Sssr, vol. 269, pp. 543–547 (1983)
- [44] Nocedal, J., Wright, S.J.: Numerical Optimization. Springer Series in Operations Research. Springer-Verlag, New York (1999)
- [45] Pasupathy, R.: On choosing parameters in retrospective-approximation algorithms for stochastic root finding and simulation optimization. Oper. Res. 58(4-Part-1), 889–901 (2010)

- [46] Pasupathy, R., Glynn, P.W., Ghosh, S., Hashemi, F.S.: On sampling rates in simulation-based recursions. SIAM J. Optimization 28(1), 45–73 (2018)
- [47] Pasupathy, R., Kim, S.: The stochastic root-finding problem: Overview, solutions, and open questions. ACM Transactions on Modeling and Computer Simulation (TOMACS) 21(3), 19 (2011)
- [48] Polyak, B.T.: Some methods of speeding up the convergence of iteration methods. Ussr computational mathematics and mathematical physics 4(5), 1–17 (1964)
- [49] Polyak, B.T.: Introduction to optimization. Optimization Software, Inc., Publications Division, NewYork (1987)
- [50] Polyak, B.T., Juditsky, A.B.: Acceleration of stochastic approximation by averaging. SIAM Journal on Control and Optimization 30(4), 838–855 (1992)
- [51] Robbins, H., Monro, S.: A stochastic approximation method. The Annals of Mathematical Statistics pp. 400–407 (1951)
- [52] Robbins, H., Siegmund, D.: A convergence theorem for non negative almost supermartingales and some applications. In: Optimizing methods in statistics, pp. 233–257. Elsevier (1971)
- [53] Schmidt, M., Roux, N.L., Bach, F.R.: Convergence rates of inexact proximal-gradient methods for convex optimization. In: Advances in Neural Information Processing Systems, pp. 1458–1466 (2011)
- [54] Shanbhag, U.V., Blanchet, J.H.: Budget-constrained stochastic approximation. In: Proceedings of the 2015 Winter Simulation Conference, pp. 368–379. IEEE Press (2015)
- [55] Shapiro, A., Dentcheva, D., Ruszczyński, A.: Lectures on stochastic programming: modeling and theory. SIAM (2009)
- [56] Silvester, J.R.: Determinants of block matrices. The Mathematical Gazette 84(501), 460-467 (2000)
- [57] Spall, J.C.: Adaptive stochastic approximation by the simultaneous perturbation method. IEEE Transactions on Automatic Control 45(10), 1839–1853 (2000)
- [58] Su, W.J., Zhu, Y.: Uncertainty quantification for online learning and stochastic approximation via hierarchical incremental gradient descent. arXiv preprint arXiv:1802.04876 (2018)
- [59] Sutskever, I., Martens, J., Dahl, G., Hinton, G.: On the importance of initialization and momentum in deep learning. In: International Conference on Machine Learning, pp. 1139–1147 (2013)
- [60] Szegedy, C., Liu, W., Jia, Y., Sermanet, P., Reed, S., Anguelov, D., Erhan, D., Vanhoucke, V., Rabinovich, A.: Going deeper with convolutions. In: Proceedings of the IEEE Conference on Computer Vision and Pattern Recognition, pp. 1–9 (2015)
- [61] Toulis, P., Airoldi, E., Rennie, J.: Statistical analysis of stochastic gradient methods for generalized linear models. In: International Conference on Machine Learning, pp. 667–675 (2014)

- [62] Van der Vaart, A.W.: Asymptotic statistics, vol. 3. Cambridge university press (2000)
- [63] Venter, J., et al.: An extension of the robbins-monro procedure. The Annals of Mathematical Statistics 38(1), 181–190 (1967)
- [64] Wei, C., et al.: Multivariate adaptive stochastic approximation. The Annals of Statistics 15(3), 1115– 1130 (1987)
- [65] Yang, T., Lin, Q., Li, Z.: Unified convergence analysis of stochastic momentum methods for convex and non-convex optimization. arXiv preprint arXiv:1604.03257 (2016)
- [66] Yousefian, F., Nedic, A., Shanbhag, U.V.: On stochastic gradient and subgradient methods with adaptive steplength sequences. Automatica 48(1), 56–67 (2012)
- [67] Zhu, Y., Dong, J.: On constructing confidence region for model parameters in stochastic gradient descent via batch means. In: Proceedings of the 2021 Winter Simulation Conference (2021)

A Proof of Lemma 3

First, we introduce an important relationship between the matrix norm and the spectral radius of the matrix, originally proven by Gelfand and also shown in [49, Lemma 1 in Section 2.1].

Lemma 10 (Gelfand's formula). It holds that $\rho(\mathbf{P}) = \lim_{k\to\infty} \|\mathbf{P}^k\|^{1/k}$, i.e., the spectral radius of \mathbf{P} gives the asymptotic growth rate of $\|\mathbf{P}^k\|$. Then for any $\iota > 0$, there exists a constant $c = c(\iota)$ such that $\|\mathbf{P}^k\| \le c(\iota)(\rho(\mathbf{P}) + \iota)^k$, $\forall k \ge 1$.

Proof of Lemma 3. Note that $\nabla f(x) - \nabla f(x^*) = \mathbf{H}(x - x^*)$ since f(x) is quadratic and $\nabla^2 f(x) \equiv \mathbf{H}$. Then by using (12) and $\nabla f(x^*) = 0$, we have

$$\begin{aligned} x_{k+1} - x^* &= x_k - x^* + \beta(x_k - x_{k-1}) - \alpha(\nabla f(x_k) - \nabla f(x^*)) - \alpha w_{k,N_k} \\ &= x_k - x^* + \beta(x_k - x_{k-1}) - \alpha \mathbf{H}(x_k - x^*) - \alpha w_{k,N_k} \\ &= ((1+\beta)\mathbf{I}_m - \alpha \mathbf{H})(x_k - x^*) - \beta(x_{k-1} - x^*) - \alpha w_{k,N_k}. \end{aligned}$$

Denote by $z_k = \begin{pmatrix} x_k - x^* \\ x_{k-1} - x^* \end{pmatrix}$. We then write this recursion in matrix form. $z_{k+1} = \begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha \mathbf{H} & -\beta \mathbf{I}_m \\ \mathbf{I} & \mathbf{0} \end{pmatrix} z_k + \begin{pmatrix} -\alpha w_{k,N_k} \\ 0 \end{pmatrix}.$

$$z_{k+1} = \underbrace{\begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha \mathbf{H} & -\beta \mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix}}_{\triangleq \mathbf{T}} z_k + \begin{pmatrix} -\alpha w_{k,N_k} \\ 0 \end{pmatrix}.$$
(58)

By the eigenvalue decomposition, $\mathbf{H} = \mathbf{U} \mathbf{\Lambda} \mathbf{U}^T$, where \mathbf{U} is orthogonal and $\mathbf{\Lambda} \triangleq \operatorname{diag}\{\lambda_1, \lambda_2, \cdots, \lambda_m\}$ with $\lambda_i \in [\eta, L], i = 1, \cdots, m$ being the eigenvalues of \mathbf{H} . Then we can rewrite \mathbf{T} as

$$\begin{pmatrix} \mathbf{U} & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{U} \end{pmatrix} \begin{pmatrix} (1+\beta)\mathbf{I}_m - \alpha\mathbf{\Lambda} & -\beta\mathbf{I}_m \\ \mathbf{I}_m & \mathbf{0}_m \end{pmatrix} \begin{pmatrix} \mathbf{U} & \mathbf{0}_m \\ \mathbf{0}_m & \mathbf{U} \end{pmatrix}^T$$

Then the eigenvalues of the matrix T are the roots of the equation $det(vI_{2m} - T) = 0$, i.e.,

$$det \begin{pmatrix} v\mathbf{I}_m - (1+\beta)\mathbf{I}_m + \alpha \mathbf{\Lambda} & \beta \mathbf{I}_m \\ -\mathbf{I}_m & v\mathbf{I}_m \end{pmatrix} = 0.$$

By the property $det \begin{pmatrix} A & B \\ C & D \end{pmatrix} = det(AD - BC)$ when the blocks A, B, C, D are square matrices of the

same size and CD = DC [56], we have $det \left(v \left(v \mathbf{I}_m - (1 + \beta) \mathbf{I}_m + \alpha \mathbf{\Lambda} \right) + \beta \mathbf{I}_m \right) = 0$. Since the matrix in the determinant is diagonal, it can be described by the following characteristic equations.

$$p_i(v) = (v - (1 + \beta - \alpha \lambda_i))v + \beta = v^2 - (1 + \beta - \alpha \lambda_i)v + \beta = 0, \quad i = 1, \cdots, m$$

From $\eta \leq \lambda_i \leq L$ and $\alpha < 4/L$ it follows that for any $i = 1, \dots, m : 1 - \sqrt{\alpha L} \leq 1 - \sqrt{\alpha \lambda_i} \leq 1 - \sqrt{\alpha \eta}$, hence $|1 - \sqrt{\alpha \lambda_i}|^2 \leq \max\{|1 - \sqrt{\alpha \eta}|^2, |1 - \sqrt{\alpha L}|^2\} = \beta < 1$. Thus, the discriminant of the equation $p_i(v)$, denoted by Δ_i , is nonpositive.

$$\Delta_{i} = (1 + \beta - \alpha\lambda_{i})^{2} - 4\beta = (1 - \beta)^{2} - 2(1 + \beta)\alpha\lambda_{i} + (\alpha\lambda_{i})^{2}$$

$$\leq (1 - |1 - \sqrt{\alpha\lambda_{i}}|^{2})^{2} - 2(1 + |1 - \sqrt{\alpha\lambda_{i}}|^{2})\alpha\lambda_{i} + (\alpha\lambda_{i})^{2}$$

$$= (1 + |1 - \sqrt{\alpha\lambda_{i}}|^{2})^{2} - 4|1 - \sqrt{\alpha\lambda_{i}}|^{2} - 2(1 + |1 - \sqrt{\alpha\lambda_{i}}|^{2})\alpha\lambda_{i} + (\alpha\lambda_{i})^{2}$$

$$= (1 - \alpha\lambda_{i} + |1 - \sqrt{\alpha\lambda_{i}}|^{2})^{2} - 4|1 - \sqrt{\alpha\lambda_{i}}|^{2} = 0.$$
(59)

Hence $p_i(v) = 0$ has two complex roots $\frac{1+\beta-\alpha\lambda_i\pm i\sqrt{-\Delta_i}}{2}$, where $\mathbf{i} = \sqrt{-1}$. Thus, the magnitude of the roots is $\sqrt{\left(\frac{1+\beta-\alpha\lambda_i}{2}\right)^2 - \frac{\Delta_i}{4}} = \sqrt{\beta}$. Then

$$\rho(\mathbf{T}) = \sqrt{\beta}, \quad \forall k \ge 1.$$
(60)

This together with Lemma 10 implies that for any $\iota \in (0, 1 - \sqrt{\beta})$, there exists a constant $c = c(\iota)$ such that

$$\|\mathbf{T}^k\| \le c(\iota)(\sqrt{\beta} + \iota)^k, \quad \forall k \ge 1.$$
(61)

We can recursively write (58) as

$$z_{k+1} = \mathbf{T}^{k+1} z_0 - \sum_{t=0}^k \mathbf{T}^{k-t} \alpha w_{t,N_t}.$$

$$\Rightarrow ||z_{k+1}||^2 = z_0^T (\mathbf{T}^{k+1})^T \mathbf{T}^{k+1} z_0 + \alpha^2 \left\| \sum_{t=0}^k \mathbf{T}^{k-t} w_{t,N_t} \right\|^2 - 2\alpha (\mathbf{T}^{k+1} z_0)^T \sum_{t=0}^k \mathbf{T}^{k-t} w_{t,N_t}.$$
 (62)

Note from (5) that

$$\mathbb{E}[w_{i,N_{i}}^{T}w_{j,N_{j}}] = \mathbb{E}\left[\mathbb{E}[w_{i,N_{i}}^{T}w_{j,N_{j}} \mid \mathcal{F}_{j}]\right] = \mathbb{E}\left[w_{i,N_{i}}^{T}\mathbb{E}[w_{j,N_{j}}|\mathcal{F}_{j}]\right] = 0 \text{ for any } i < j,$$
$$\mathbb{E}[z_{0}^{T}(\mathbf{T}^{k+1})^{T}\sum_{t=0}^{k}\mathbf{T}^{k-t}w_{t,N_{t}}] = \sum_{t=0}^{k}\mathbb{E}[z_{0}^{T}(\mathbf{T}^{k+1})^{T}\mathbf{T}^{k-t}w_{t,N_{t}}] = \sum_{t=0}^{k}\mathbb{E}\left[\mathbb{E}[z_{0}^{T}(\mathbf{T}^{k+1})^{T}\mathbf{T}^{k-t}w_{t,N_{t}} \mid \mathcal{F}_{t}]\right] = 0$$

Then by using (5), we derive

$$\mathbb{E}\left[\left\|\sum_{t=0}^{k}\mathbf{T}^{k-t}w_{t,N_{t}}\right\|^{2}\right] = \sum_{t=0}^{k}\mathbb{E}\left[\left\|\mathbf{T}^{k-t}w_{t,N_{t}}\right\|^{2}\right] + 2\sum_{i
$$= \sum_{t=0}^{k}\mathbb{E}\left[\left\|\mathbf{T}^{k-t}w_{t,N_{t}}\right\|^{2}\right] \le \sum_{t=0}^{k}\|\mathbf{T}^{k-t}\|^{2}\frac{\nu^{2}}{N_{t}} \stackrel{(61)}{\le} \sum_{t=0}^{k}c(\iota)^{2}(\sqrt{\beta}+\iota)^{2(k-t)}\frac{\nu^{2}}{N_{t}}$$$$

Then by using (62) and (61), we obtain that

$$\mathbb{E}[\|z_{k+1}\|^{2}] \leq \|\mathbf{T}^{k+1}\|^{2} \mathbb{E}[\|z_{0}\|^{2}] + \alpha^{2} \mathbb{E}\left[\left\|\sum_{t=0}^{k} \mathbf{T}^{k-t} w_{t,N_{t}}\right\|^{2}\right]$$
$$\leq c(\iota)^{2} (\sqrt{\beta} + \iota)^{2(k+1)} \mathbb{E}\left[\|z_{0}\|^{2}\right] + \alpha^{2} \nu^{2} c(\iota)^{2} \sum_{t=0}^{k} (\sqrt{\beta} + \iota)^{2(k-t)} \frac{1}{N_{t}}.$$
$$\left[\left\|\left(x_{0} - x^{*}\right)\right\|^{2}\right]$$

By recalling $\mathbb{E}[||z_0||^2] = \left[\left\| \begin{pmatrix} x_0 - x^* \\ x_{-1} - x^* \end{pmatrix} \right\|^2 \right] = 2\mathbb{E}[||x_0 - x^*||^2]$ from $x_{-1} = x_0$, the result follows.

B Proof of Lemma 7

Define the sequence $\{u_k\}$ as follows:

$$u_{k+1} = \mathbf{P}u_k - \alpha \rho^{-(k+1)/2} \mathbf{G} w_{k,N_k}, \quad u_0 = 0.$$
(63)

By combining (63) with (33), we obtain the following recursion:

$$e_{k+1} - u_{k+1} = \mathbf{P}(e_k - u_k) + \zeta_{k+1} = \mathbf{P}^{k+1}(e_0 - u_0) + \sum_{t=0}^k \mathbf{P}^{k-t}\zeta_{t+1}.$$
 (64)

Lemma 10 together with $\rho(\mathbf{P}) < 1$ implies that for any $\varrho \in (\rho(\mathbf{P}), 1)$, there exists a constant $c(\varrho)$ such that

$$\|\mathbf{P}^k\| \le c(\varrho)\varrho^k, \quad \forall k \ge 1.$$
(65)

Then by using (65) and $\zeta_k \xrightarrow{P} 0$, we obtain from (64) that

$$\|e_{k+1} - u_{k+1}\| \le \|\mathbf{P}^{k+1}\| \|e_0 - u_0\| + \sum_{t=0}^k \|\mathbf{P}^{k-t}\| \|\zeta_{t+1}\| \le c(\varrho)\varrho^{k+1} \|e_0 - u_0\| + c(\varrho)\sum_{t=0}^k \varrho^{k-t}o(1) \xrightarrow{P}{k \to \infty} 0$$

Thus, e_k defined by (33) and u_k produced by (63) converge to the same distribution asymptotically, if indeed the limit exists. This follows from the fact that $X_k \xrightarrow{d} X$, $||X_k - Y_k|| \xrightarrow{P} 0 \Rightarrow Y_k \xrightarrow{d} X$ (see [62, Theorem 2.7(iv)]).

From (63) and $u_0 = 0$ it follows that

$$u_{k+1} = \mathbf{P}^{k+1}u_0 - \alpha \sum_{t=0}^k \mathbf{P}^{k-t}\mathbf{G}\rho^{-(t+1)/2}w_{t,N_t} = -\alpha \sum_{t=0}^k \mathbf{P}^{k-t}\mathbf{G}\rho^{-(t+1)/2}w_{t,N_t}.$$

This implies that

$$\alpha^{-1}u_k = -\sum_{t=0}^{k-1} \mathbf{P}^{k-1-t} \mathbf{G} \rho^{-(t+1)/2} w_{t,N_t} = -\sum_{t=1}^k \mathbf{P}^{k-t} \mathbf{G} \rho^{-t/2} w_{t-1,N_{t-1}}.$$
 (66)

Define $\xi_{kt} \triangleq -\mathbf{P}^{k-t}\mathbf{G}\rho^{-t/2}w_{t-1,N_{t-1}}$ for any $t: 1 \le t \le k$. We intend to apply Lemma 6. Therefore, we have to check conditions (26)-(28). By using Assumption 1(iii), $N_t \ge \rho^{-(t+1)}$, (5), and (65), we obtain that $\mathbb{E}[\boldsymbol{\xi}_{kt}|\boldsymbol{\xi}_{k1},\cdots,\boldsymbol{\xi}_{k,t-1}] = 0$ and

$$\begin{split} \mathbb{E}[\|\boldsymbol{\xi}_{kt}\|^{2}|\boldsymbol{\xi}_{k1},\cdots,\boldsymbol{\xi}_{k,t-1}] &\leq \|\mathbf{P}^{k-t}\|^{2}\|\mathbf{G}\|^{2}\rho^{-t}\mathbb{E}[\|w_{t-1,N_{t-1}}\|^{2}|\mathcal{F}_{t-1}] \\ &\leq c(\varrho)^{2}\varrho^{2(k-t)}\|\mathbf{G}\|^{2}\rho^{-t}\frac{\nu^{2}}{N_{t-1}} \leq c(\varrho)^{2}\nu^{2}\|\mathbf{G}\|^{2}\varrho^{2(k-t)} \ a.s.. \\ &\Longrightarrow \sum_{t=1}^{k}\mathbb{E}[\|\boldsymbol{\xi}_{kt}\|^{2}] \leq c(\varrho)^{2}\nu^{2}\|\mathbf{G}\|^{2}\sum_{t=1}^{k}\varrho^{2(k-t)} = \frac{c(\varrho)^{2}\nu^{2}\varrho^{2}\|\mathbf{G}\|^{2}}{1-\varrho^{2}} < \infty. \end{split}$$

Thus, (26) holds. By recalling the definitions of \mathbf{S}_{kt} and \mathbf{R}_{kt} in (25), using $\sum_{t=1}^{k} \|\mathbf{P}^{k-t}\|^2 \leq \sum_{t=1}^{k} c(\varrho)^2 \varrho^{2(k-t)} = \frac{c(\varrho)^2}{1-\varrho^2}$ from (65) and (30), and $\rho^{-t}/N_{t-1} \xrightarrow[t \to \infty]{} 1$, we have

$$\sum_{t=1}^{k} \mathbb{E}[\|\mathbf{R}_{kt} - \mathbf{S}_{kt}\|] \leq \sum_{t=1}^{k} \|\mathbf{P}^{k-t}\mathbf{G}\|^{2} \mathbb{E}\left[\left\|N_{t-1}\mathbb{E}\left[w_{t-1,N_{t-1}}w_{t-1,N_{t-1}}^{T}|\mathcal{F}_{t-1}\right] - N_{t-1}\mathbb{E}\left[w_{t-1,N_{t-1}}w_{t-1,N_{t-1}}^{T}\right]\right\|\right] \\ + \sum_{t=1}^{k} \|\mathbf{P}^{k-t}\mathbf{G}\|^{2} \mathbb{E}\left[\left\|\left(N_{t-1}\mathbb{E}\left[w_{t-1,N_{t-1}}w_{t-1,N_{t-1}}^{T}|\mathcal{F}_{t-1}\right] - N_{t-1}\mathbb{E}\left[w_{t-1,N_{t-1}}w_{t-1,N_{t-1}}^{T}\right]\right)\left(\rho^{-t}/N_{t-1} - 1\right)\right\|\right] \\ \leq \|\mathbf{G}\|^{2} \sum_{t=1}^{k} c(\varrho)^{2} \varrho^{2(k-t)} o(1) \xrightarrow[k \to \infty]{} 0.$$

This verifies the second equality in (27). We now verify the first equality in (27).

$$\mathbf{S}_{k} = \sum_{t=1}^{k} \mathbf{S}_{kt} = \sum_{t=1}^{k} \mathbf{P}^{k-t} \mathbf{G} \mathbb{E} \left[\rho^{-t} w_{t-1,N_{t-1}} w_{t-1,N_{t-1}}^{T} \right] (\mathbf{P}^{k-t} \mathbf{G})^{T}$$

$$= \sum_{t=1}^{k} \mathbf{P}^{k-t} \mathbf{G} \mathbf{S}_{0} (\mathbf{P}^{k-t} \mathbf{G})^{T} + \sum_{t=1}^{k} \mathbf{P}^{k-t} \mathbf{G} \mathbf{S}_{0} (\mathbf{P}^{k-t} \mathbf{G})^{T} (\rho^{-t}/N_{t-1} - 1)$$

$$+ \sum_{t=1}^{k} \mathbf{P}^{k-t} \mathbf{G} \left(N_{t-1} \mathbb{E} \left[w_{t-1,N_{t-1}} w_{t-1,N_{t-1}}^{T} \right] - \mathbf{S}_{0} \right) (\mathbf{P}^{k-t} \mathbf{G})^{T} \rho^{-t}/N_{t-1},$$
(67)

where the second and last terms tend to zero by using $\sum_{t=1}^{k} \|\mathbf{P}^{k-t}\|^2 \leq \frac{c(\varrho)^2}{1-\varrho^2}$, $\rho^{-t}/N_{t-1} \xrightarrow[t \to \infty]{} 1$, and (30). The first term on the right-hand side of (67) can be written as

$$\tilde{\mathbf{S}}_{k} \triangleq \sum_{t=1}^{k} \mathbf{P}^{k-t} \mathbf{G} \mathbf{S}_{0} (\mathbf{P}^{k-t} \mathbf{G})^{T} = \sum_{t=0}^{k-1} \mathbf{P}^{t} \mathbf{G} \mathbf{S}_{0} \mathbf{G}^{T} (\mathbf{P}^{t})^{T}.$$
(68)

By (65), $\sum_{t=0}^{\infty} \mathbf{P}^t \mathbf{GS}_0 \mathbf{G}^T (\mathbf{P}^t)^T$ is well defined and denoted by $\boldsymbol{\Sigma}$. Then $\boldsymbol{\Sigma} - \tilde{\mathbf{S}}_k = \sum_{t=k}^{\infty} \mathbf{P}^t \mathbf{GS}_0 \mathbf{G}^T (\mathbf{P}^t)^T$ and hence we derive from (65) that

$$\begin{split} \|\boldsymbol{\Sigma} - \tilde{\mathbf{S}}_k\| &\leq \sum_{t=k}^{\infty} \|\mathbf{P}^t \mathbf{G} \mathbf{S}_0 \mathbf{G}^T (\mathbf{P}^t)^T\| \leq \|\mathbf{G} \mathbf{S}_0 \mathbf{G}^T\| \sum_{t=k}^{\infty} \|\mathbf{P}^t\|^2 \\ &\leq \|\mathbf{G} \mathbf{S}_0 \mathbf{G}^T\| c(\varrho)^2 \sum_{t=k}^{\infty} \varrho^{2t} = \|\mathbf{G} \mathbf{S}_0 \mathbf{G}^T\| \frac{c(\varrho)^2}{1-\varrho^2} \varrho^{2k} \to 0 \text{ when } k \to \infty \end{split}$$

Therefore, the limit of (68) exists and equals Σ . This together with (67) and that the second and last terms on the right hand side of (67) tend to zero, we achieve

$$\lim_{k \to \infty} \mathbf{S}_k \triangleq \mathbf{\Sigma} = \lim_{k \to \infty} \sum_{t=0}^{\infty} \mathbf{P}^t \mathbf{G} \mathbf{S}_0 \mathbf{G}^T \left(\mathbf{P}^t \right)^T.$$

Finally, we have to verify the Lindeberg condition (28). By using $\xi_{kt} = -\mathbf{P}^{k-t}\mathbf{G}\rho^{-t/2}w_{t-1,N_{t-1}}$, $N_{t-1} = \lceil \rho^{-t} \rceil$, and (65), we obtain that

$$\|\xi_{kt}\| \le \|\mathbf{P}^{k-t}\| \|\mathbf{G}\| \rho^{-t/2} \|w_{t-1,N_{t-1}}\| \le c(\varrho) \varrho^{k-t} \|\mathbf{G}\| \sqrt{N_{t-1}} \|w_{t-1,N_{t-1}}\|.$$
(69)

Hence for any $\epsilon > 0$,

$$\left\{ \|\xi_{kt}\| > \epsilon \right\} \subset \left\{ \sqrt{N_{t-1}} \|w_{t-1,N_{t-1}}\| > \epsilon \|\mathbf{G}\|^{-1} c(\varrho)^{-1} \varrho^{-(k-t)} \right\}.$$
(70)

Because for any $t \ge 1$, $\varrho^{-(k-t)} \xrightarrow[k \to \infty]{} \infty$. Then using (31), we obtain that

$$\sup_{t\geq 1} \mathbb{E}\left[N_{t-1} \|w_{t-1,N_{t-1}}\|^2 I_{[\sqrt{N_{t-1}}} \|w_{t-1,N_{t-1}}\| > \epsilon \|\mathbf{G}\|^{-1} c(\varrho)^{-1} \varrho^{-(k-t)}]\right] \xrightarrow[k\to\infty]{} 0.$$

Consequently, for any $\epsilon > 0$, by using (69) and (70), the following holds:

$$\sum_{t=1}^{k} \mathbb{E}[\|\boldsymbol{\xi}_{kt}\|^{2} I_{[\|\boldsymbol{\xi}_{kt}\| \geq \epsilon]}] \leq \sum_{t=1}^{k} \|\mathbf{G}\|^{2} c(\varrho)^{2} \varrho^{2(k-t)} \mathbb{E}\left[N_{t-1} \|w_{t-1,N_{t-1}}\|^{2} I_{\left[\sqrt{N_{t-1}}\|w_{t-1,N_{t-1}}\| > \epsilon \|\mathbf{G}\|^{-1} c(\varrho)^{-1} \varrho^{-(k-t)}\right]\right]$$
$$= \sum_{t=1}^{k} \|\mathbf{G}\|^{2} c(\varrho)^{2} \varrho^{2(k-t)} o(1) \xrightarrow[k \to \infty]{} 0.$$

Thus, the conditions (26)-(28) hold. Then by using Lemma 6, the fact that e_k and u_k have the same limit distribution, and $\alpha^{-1}u_k = -\sum_{t=1}^k \xi_{kt}$, we proves Lemma 7.

C Proof of Lemma 9.

We observe that $\rho(\mathbf{A}) < 1$ together with Lemma 10 implies that for any $\varrho \in (\rho(\mathbf{A}), 1)$, there exists a constant $c(\varrho)$ such that

$$\|\mathbf{A}^k\| \le c(\varrho)\varrho^k, \quad \forall k \ge 1.$$
(71)

Define $\Phi_{k,j} \triangleq \mathbf{A}_k \cdots \mathbf{A}_j$ with $\Phi_{j,j+1} \triangleq \mathbf{I}_m$. Recall from the definitions $\mathbf{A}_0 = \mathbf{A}$ and $\mathbf{A}_k \triangleq \left(\frac{k+1}{k}\right)^{\nu/2} \mathbf{A}$, we obtain that $\Phi_{k,0} = (k+1)^{\nu/2} \mathbf{A}^k$ and $\Phi_{k,t+1} = \left(\frac{k+1}{t+1}\right)^{\nu/2} \mathbf{A}^{k-t}$. This combined with (71) produces

$$\|\boldsymbol{\Phi}_{k,0}\| \le c(\varrho)\varrho^k \, (k+1)^{\nu/2}, \qquad \forall k \ge 0, \tag{72}$$

and
$$\|\mathbf{\Phi}_{k,t+1}\| \le c(\varrho) \left(\frac{k+1}{t+1}\right)^{\nu/2} \varrho^{k-t}, \quad \forall k \ge t \ge 0.$$
 (73)

For any given a > 0, define $\tilde{q} \triangleq \varrho^a$ and $\tilde{v} \triangleq av/2$. Then by using (73) and Lemma 8, we derive

$$\sum_{t=0}^{k} \| \boldsymbol{\Phi}_{k,t+1} \|^{a} \leq c(\varrho)^{a} \sum_{t=0}^{k} \left(\frac{k+1}{t+1} \right)^{av/2} \varrho^{a(k+1-(t+1))} \leq c(\varrho)^{a} (k+1)^{\tilde{\nu}} \sum_{t=1}^{k+1} t^{-\tilde{\nu}} \tilde{q}^{k+1-t}$$

$$\leq c(\varrho)^{a} (k+1)^{\tilde{\nu}} \left(\tilde{q}^{k+1} \frac{e^{2\tilde{\nu}} \tilde{q}^{-1}-1}{1-\tilde{q}} + \frac{2(k+1)^{-\tilde{\nu}}}{\tilde{q} \ln(1/\tilde{q})} \right)$$

$$\leq c(\varrho)^{a} \left((k+1)^{\tilde{\nu}} \tilde{q}^{k+1} \frac{e^{2\tilde{\nu}} \tilde{q}^{-1}-1}{1-\tilde{q}} + \frac{2}{\tilde{q} \ln(1/\tilde{q})} \right) \leq c(\varrho)^{a} \left(c_{\tilde{q},\tilde{\nu}} \frac{e^{2\tilde{\nu}} \tilde{q}^{-1}-1}{1-\tilde{q}} + \frac{2}{\tilde{q} \ln(1/\tilde{q})} \right), \quad \forall k \geq 1.$$
(74)

Define an auxiliary sequence $\{u_k\}$ by

$$u_{k+1} = \mathbf{A}_k u_k - \alpha (k+1)^{\nu/2} \mathbf{G} w_{k,N_k}, \quad u_0 = 0.$$
(75)

This combined with (49) produces the following recursion:

$$e_{k+1} - u_{k+1} = \mathbf{A}_k(e_k - u_k) + \zeta_{k+1}$$

= $\mathbf{\Phi}_{k,0}(e_0 - u_0) + \sum_{t=0}^k \mathbf{\Phi}_{k,t+1} \mathbf{G} \zeta_{t+1} = \mathbf{\Phi}_{k,0} e_0 + \sum_{t=0}^k \mathbf{\Phi}_{k,t+1} \zeta_{t+1}.$

Then by using (72), (74), $\mathbb{E}[||e_0||^2] < \infty$, and $\zeta_{k+1} \xrightarrow{P} 0$, we conclude that

$$\|e_{k+1} - u_{k+1}\| = \|\Phi_{k,0}\| \|e_0\| + \sum_{t=0}^k \|\Phi_{k,t+1}\| \|\zeta_{t+1}\| \xrightarrow{P}{k \to \infty} 0.$$

This implies that e_k defined by (49) and u_k defined as in (75) have the same limit distribution if exists. Thus, it remains to find the stationary distribution of u_k .

From (75) it follows that

$$u_{k+1} = \mathbf{\Phi}_{k,0}u_0 - \alpha \sum_{t=0}^k \mathbf{\Phi}_{k,t+1}\mathbf{G}(t+1)^{\nu/2}w_{t,N_t} = -\alpha \sum_{t=0}^k \mathbf{\Phi}_{k,t+1}\mathbf{G}(t+1)^{\nu/2}w_{t,N_t}$$

This implies that

$$\alpha^{-1}u_k = -\alpha \sum_{t=0}^{k-1} \mathbf{\Phi}_{k-1,t+1} \mathbf{G}(t+1)^{\nu/2} w_{t,N_t} = -\sum_{t=1}^k \mathbf{\Phi}_{k-1,t} \mathbf{G} \ t^{\nu/2} w_{t-1,N_{t-1}}.$$
 (76)

We intend to apply Lemma 6 by defining $\xi_{kt} \triangleq -\Phi_{k-1,t}\mathbf{G}t^{v/2}w_{t-1,N_{t-1}}$ and check conditions (26), (27), and (28). Using $N_{t-1} \triangleq \lceil t^v \rceil \ge t^v$, (74), and Assumption 1(iii), we can verify (26). Also, using (30) and (74), the definitions of \mathbf{S}_{kt} and \mathbf{R}_{kt} in (25), the second equality of (27) holds. We now verify the first equality in (27).

$$\mathbf{S}_{k} = \sum_{t=1}^{k} \mathbf{S}_{kt} = \sum_{t=1}^{k} \mathbf{\Phi}_{k-1,t} \mathbf{G} \mathbb{E} \left[t^{v} w_{t-1,N_{t-1}} w_{t-1,N_{t-1}}^{T} \right] \mathbf{G}^{T} \mathbf{\Phi}_{k-1,t}^{T}$$

$$= \sum_{t=1}^{k} \mathbf{\Phi}_{k-1,t} \mathbf{G} \mathbf{S}_{0} \mathbf{G}^{T} \mathbf{\Phi}_{k-1,t}^{T} + \sum_{t=1}^{k} \mathbf{\Phi}_{k-1,t} \mathbf{G} \mathbf{S}_{0} \mathbf{G}^{T} \mathbf{\Phi}_{k-1,t}^{T} \left(t^{v}/N_{t} - 1 \right)$$

$$+ \sum_{t=1}^{k} \mathbf{\Phi}_{k-1,t} \mathbf{G} \left(N_{t-1} \mathbb{E} \left[w_{t-1,N_{t-1}} w_{t-1,N_{t-1}}^{T} \right] - \mathbf{S}_{0} \right) \mathbf{G}^{T} \mathbf{\Phi}_{k-1,t}^{T} t^{v}/N_{t-1},$$
(77)

where the second and last terms tend to zero by (30), $t^v/N_{t-1} \xrightarrow[t \to \infty]{t \to \infty} 1$, and $\sum_{t=1}^k \|\Phi_{k-1,t}\|^2 < \infty$ from (74). While the first term on the right-hand side of (77), by using (74) one obtains

$$\sum_{t=1}^{k} \| \mathbf{\Phi}_{k-1,t} \mathbf{GS}_0 \mathbf{G}^T \mathbf{\Phi}_{k-1,t}^T \| \le \| \mathbf{GS}_0 \mathbf{G}^T \| \sum_{t=1}^{k} \| \mathbf{\Phi}_{k-1,t} \|^2 < \infty.$$

Because $\sum_{t=1}^{k} \| \mathbf{\Phi}_{k-1,t} \mathbf{G} \mathbf{S}_0 \mathbf{G}^T \mathbf{\Phi}_{k-1,t}^T \|$ is monotonically increasing and bounded, its limit exists. Thus, the limit of $\sum_{t=1}^{k} \mathbf{\Phi}_{k-1,t} \mathbf{G} \mathbf{S}_0 \mathbf{G}^T \mathbf{\Phi}_{k-1,t}^T = \sum_{t=1}^{k} \left(\frac{k}{t}\right)^v \mathbf{A}^{k-t} \mathbf{G} \mathbf{S}_0 \mathbf{G}^T (\mathbf{A}^{k-t})^T$ exists and is denoted by $\boldsymbol{\Sigma}$. Then \mathbf{S}_k as defined as in (77) satisfies that

$$\lim_{k \to \infty} \mathbf{S}_k = \lim_{k \to \infty} \sum_{t=1}^k \left(\frac{k}{t}\right)^v \mathbf{A}^{k-t} \mathbf{G} \mathbf{S}_0 \mathbf{G}^T (\mathbf{A}^T)^{k-t} \triangleq \boldsymbol{\Sigma}.$$

Thus, the first equality in (27) holds.

Finally, the Lindeberg condition (28) can similarly validated as that of Lemma 7. Then all conditions of Lemma 6 hold. Thus, by Lemma 6 and (76), we conclude that $\alpha^{-1}u_k \xrightarrow{d} N(0, \Sigma)$. Because e_k defined by (49) and u_k defined as in (75) have the same limit distribution, Lemma 9 is then proved.

D Proof of Proposition 9

Based on Theorem 1 it is seen that

$$e_{ik} \triangleq \alpha^{-1} \rho_1^{-k/2} (x_{ik} - x^*) \xrightarrow[k \to \infty]{d} Y_i \sim N(0, \Sigma_1), \quad i = 1, \cdots, n,$$
(78)

where Y_1, \dots, Y_n are n i.i.d. random vectors with distribution $N(0, \Sigma_1)$. Define

$$\bar{\mathbf{e}} \triangleq \frac{1}{n} \sum_{i=1}^{n} Y_i \text{ and } \mathcal{S} \triangleq \frac{1}{n-1} \sum_{i=1}^{n} \left(Y_i - \bar{\mathbf{e}} \right) \left(Y_i - \bar{\mathbf{e}} \right)^T.$$
 (79)

Recall from [6, Theorem 4] that \bar{e} and S are independently distributed, and

$$\bar{\mathbf{e}} \sim N(0, \Sigma_1/n) \text{ and } (n-1)\mathcal{S} \sim \mathcal{W}_m(\Sigma_1, n-1),$$
(80)

where $\mathcal{W}_m(\Sigma_1, n-1)$ denotes the *m*-dimensional Wishart distribution with n-1 degrees of freedom and the matrix parameter Σ_1 . Because Σ_1 is invertible and $n \ge m+1$, the random matrix S is almost surely invertible.

(i) Denote by $\mathbf{e} = col\{e_1, \dots, e_n\} \triangleq (e_1^T, \dots, e_n^T)^T \in \mathbb{R}^{mn}$ with $e_i \in \mathbb{R}^m$, $i = 1, \dots, n$. Note that $g_1(\mathbf{e}) \triangleq \frac{1}{n} \sum_{i=1}^n e_i$ is a continuous function. Since e_{ik} , $i = 1, \dots, n$ are mutually independent, from (78) it follows that

$$col\{e_{1k},\cdots,e_{nk}\} \xrightarrow[k \to \infty]{d} col\{Y_1,\cdots,Y_n\}.$$
 (81)

Then by the continuous mapping theorem [13, Theorem 1.14] and (80), we derive

$$\sqrt{n}\alpha^{-1}\rho_1^{-k/2}(\bar{\mathbf{x}}_k - x^*) = \sqrt{n}\frac{1}{n}\sum_{i=1}^n e_{ik} \xrightarrow{d} \sqrt{n}\frac{1}{n}\sum_{i=1}^n Y_i = \sqrt{n}\bar{\mathbf{e}} \sim N(0, \boldsymbol{\Sigma}_1).$$
(82)

(ii) Since $rank(\Sigma_1) = m$ and $\sqrt{n}\bar{\mathbf{e}} \sim N(0, \Sigma_1)$ by (82), we obtain

$$U_1 \triangleq (\sqrt{n}\bar{\mathbf{e}})^T \boldsymbol{\Sigma}_1^{-1} (\sqrt{n}\bar{\mathbf{e}}) \sim \chi^2(m),$$
(83)

where $\chi^2(m)$ denotes the chi-squared distribution with m degrees of freedom. Because $rank(\Sigma_1) = m$, $\bar{\mathbf{e}} \in \mathbb{R}^m$ and S are independently distributed with $\bar{\mathbf{e}}^T \Sigma_1$ being non-zero with probability one, from [6, Corollary 1] it follows that

$$U_2 \triangleq \frac{\bar{\mathbf{e}}^T \boldsymbol{\Sigma}_1^{-1} \bar{\mathbf{e}}}{\bar{\mathbf{e}}^T \left((n-1) \boldsymbol{\mathcal{S}} \right)^{-1} \bar{\mathbf{e}}} \sim \chi^2 (n-m)$$
(84)

is independent of $\bar{\mathbf{e}}$. Hence U_1 and U_2 are independent.

From (78) and (53) it follows that

$$\bar{\mathbf{x}}_{k} - x^{*} = \frac{1}{n} \sum_{i=1}^{n} (x_{ik} - x^{*}) = \alpha \rho_{1}^{k/2} \sum_{i=1}^{n} e_{ik},$$

$$\mathbf{S}_{k} = \frac{1}{n-1} \sum_{i=1}^{n} (x_{ik} - \bar{\mathbf{x}}_{k}) (x_{ik} - \bar{\mathbf{x}}_{k})^{T} = \frac{\alpha^{2} \rho_{1}^{k}}{n-1} \sum_{i=1}^{n} (e_{ik} - \frac{1}{n} \sum_{i=1}^{n} e_{ik}) (e_{ik} - \frac{1}{n} \sum_{i=1}^{n} e_{ik}).$$
(85)

Note that $g_2(\mathbf{e}) \triangleq \frac{1}{n-1} \sum_{i=1}^n \left(e_i - \frac{1}{n} \sum_{i=1}^n e_i \right) \left(e_i - \frac{1}{n} \sum_{i=1}^n e_i \right)^T$ is a continuous function. Because the matrix inverse functional is continuous in a neighborhood of any non-singular matrix, and

$$\sum_{i=1}^{n} \left(Y_i - \frac{1}{n} \sum_{i=1}^{n} Y_i \right) \left(Y_i - \frac{1}{n} \sum_{i=1}^{n} Y_i \right)^T$$

is almost surely invertible from (80), we conclude that $(g_2(\mathbf{e}))^{-1}$ is almost surely continuous in a neighborhood of $col\{Y_1, \dots, Y_n\}$. Hence, $g(\mathbf{e}) = g_1(\mathbf{e})^T (g_2(\mathbf{e}))^{-1} g_1(\mathbf{e})$ is almost surely continuous in a neighborhood of $col\{Y_1, \dots, Y_n\}$. Therefore, by the continuous mapping theorem [13, Theorem 1.14], and (81), we have

$$n(\bar{\mathbf{x}}_{k} - x^{*})^{T} \mathbf{S}_{k}^{-1}(\bar{\mathbf{x}}_{k} - x^{*})$$

$$\stackrel{(85)}{=} n\left(\frac{\sum_{i=1}^{n} e_{ik}}{n}\right)^{T} \left(\frac{1}{n-1} \sum_{i=1}^{n} \left(e_{ik} - \frac{1}{n} \sum_{i=1}^{n} e_{ik}\right) \left(e_{ik} - \frac{1}{n} \sum_{i=1}^{n} e_{ik}\right)^{T}\right)^{-1} \frac{\sum_{i=1}^{n} e_{ik}}{n}$$

$$\stackrel{d}{\longrightarrow} n \frac{\sum_{i=1}^{n} Y_{i}}{n} \left(\frac{1}{n-1} \sum_{i=1}^{n} \left(Y_{i} - \frac{\sum_{i=1}^{n} Y_{i}}{n}\right) \left(Y_{i} - \frac{\sum_{i=1}^{n} Y_{i}}{n}\right)^{T}\right)^{-1} \frac{\sum_{i=1}^{n} Y_{i}}{n}$$

$$\stackrel{(79)}{=} n(n-1)\bar{\mathbf{e}}^{T} ((n-1)\mathcal{S})^{-1} \bar{\mathbf{e}} = (n-1) \frac{n\bar{\mathbf{e}}^{T} \mathbf{\Sigma}_{1}^{-1} \bar{\mathbf{e}}}{\bar{\mathbf{e}}^{T} ((n-1)\mathcal{S})^{-1} \bar{\mathbf{e}}}$$

$$= (n-1) \frac{U_{1}}{U_{2}} \sim \frac{m(n-1)}{n-m} F(m,n-m),$$
(86)

where the last one holds because $U_1 = n\bar{\mathbf{e}}^T \boldsymbol{\Sigma}_1^{-1} \bar{\mathbf{e}} \sim \chi^2(m)$ by (83), $U_2 = \frac{\bar{\mathbf{e}}^T \boldsymbol{\Sigma}_1^{-1} \bar{\mathbf{e}}}{\bar{\mathbf{e}}^T ((n-1)S)^{-1} \bar{\mathbf{e}}} \sim \chi^2(n-m)$ by (84), U_1 and U_2 are independent. Thus, $F(d_1, d_2)$ arises as the ratio of two appropriately scaled chi-squared variates [15].¹

E Proof of Proposition 10

Since Σ_1 is symmetric and invertible, by the property of Wishart distribution, we have that

$$\boldsymbol{\Sigma}_{1}^{-1}(n-1)\mathcal{S}\boldsymbol{\Sigma}_{1}^{-1} \overset{(80)}{\sim} \boldsymbol{\Sigma}_{1}^{-1}\mathcal{W}_{m}(\boldsymbol{\Sigma}_{1}, m+(n-m)-1)\boldsymbol{\Sigma}_{1}^{-1} \sim \mathcal{W}_{m}(\boldsymbol{\Sigma}_{1}^{-1}, m+(n-m)-1).$$

Note by (82) that $\sqrt{n(n-m)}\Sigma_1^{-1/2}\bar{\mathbf{e}} \sim N(0, (n-m)\mathbf{I}_m)$. Recall from [6, Theorem 4] that $\bar{\mathbf{e}}$ and S are independent. Then by [36, Representation B], we conclude that

$$\left((n-1)\boldsymbol{\Sigma}_{1}^{-1}\boldsymbol{\mathcal{S}}\boldsymbol{\Sigma}_{1}^{-1}\right)^{-1/2}\sqrt{n(n-m)}\boldsymbol{\Sigma}_{1}^{-1/2}\bar{\mathbf{e}}\sim T_{n-m}(0,\boldsymbol{\Sigma}_{1},m).$$

Thus, by the probability density function of $T_v(\mu, \Lambda, m)$ defined in (56), we see that

$$\boldsymbol{\Sigma}_{1}^{-1/2} \left((n-1)\boldsymbol{\Sigma}_{1}^{-1} \boldsymbol{\mathcal{S}} \boldsymbol{\Sigma}_{1}^{-1} \right)^{-1/2} \sqrt{n(n-m)} \boldsymbol{\Sigma}_{1}^{-1/2} \bar{\mathbf{e}} \sim T_{n-m}(0, \mathbf{I}_{m}, m)$$

Hence

$$((n-1)\mathcal{S})^{-1/2}\sqrt{n(n-m)}\bar{\mathbf{e}} \sim T_{n-m}(0,\mathbf{I}_m,m).$$
(87)

Similarly to the derivation of (86), we can show that $(g_2(\mathbf{e}))^{-1/2}g_1(\mathbf{e})$ is almost surely continuous in a neighborhood of $col\{Y_1, \dots, Y_n\}$. Then by the continuous mapping theorem [13, Theorem 1.14] and (81), we achieve (55) by the following.

$$\frac{((n-1)\mathbf{S}_k)^{-1/2}\sqrt{n(n-m)}(\bar{\mathbf{x}}_k - x^*)}{2(1)^{-1/2}\sqrt{n(n-m)}(\bar{\mathbf{x}}_k - x^*)}$$

¹If the random variables $U_1 \sim \chi^2(d_1)$ and $U_2 \sim \chi^2(d_2)$ are independent, then $\frac{U_1/d_1}{U_2/d_2} \sim F(d_1, d_2)$.

$$\stackrel{(85)}{=} \left((n-1) \frac{1}{n-1} \sum_{i=1}^{n} \left(e_{ik} - \frac{1}{n} \sum_{i=1}^{n} e_{ik} \right) \left(e_{ik} - \frac{1}{n} \sum_{i=1}^{n} e_{ik} \right)^{T} \right)^{-1/2} \sqrt{n(n-m)} \frac{\sum_{i=1}^{n} e_{ik}}{n}$$

$$\stackrel{d}{\longrightarrow} \left((n-1) \frac{1}{n-1} \sum_{i=1}^{n} \left(Y_{i} - \frac{\sum_{i=1}^{n} Y_{i}}{n} \right) \left(Y_{i} - \frac{\sum_{i=1}^{n} Y_{i}}{n} \right)^{T} \right)^{-1/2} \sqrt{n(n-m)} \frac{\sum_{i=1}^{n} Y_{i}}{n}$$

$$\stackrel{(79)}{=} ((n-1)\mathcal{S})^{-1/2} \sqrt{n(n-m)} \bar{\mathbf{e}} \stackrel{(87)}{\sim} T_{n-m}(0, \mathbf{I}_{m}, m).$$

SUPPLEMENTARY MATERIAL.

The proof of Lemma 2 is motivated by [3, Section 2.2] and [1, Section 3.6.2].

F Proof of Lemma 2

We define $\phi_k(x)$ and p_k as follows.

$$\phi_0(x) = f(x_0) + \frac{\eta}{2} \|x - x_0\|^2, \tag{F.1}$$

$$\phi_{k+1}(x) = (1-\gamma)\phi_k(x) + \gamma \left(f(x_k) + (x-x_k)^T h(x_k) + \frac{\eta}{2} \|x-x_k\|^2 \right),$$
(F.2)

$$p_{k+1} = (1-\gamma)p_k + \left(\alpha + \frac{(1-\gamma)\gamma}{2\eta}\right) \|w_{k,N_k}\|^2 + \alpha w_{k,N_k}^T \nabla f(x_k), \ p_0 = 0,$$
(F.3)

where $h(x_k) \triangleq \frac{x_k - y_{k+1}}{\alpha} = \nabla f(x_k) + w_{k,N_k}$.

We first show by induction that for any $k \ge 0$, $\nabla^2 \phi_k(x) = \eta \mathbf{I}_m$. By (F.1) it is seen that $\nabla^2 \phi_0(x) = \eta \mathbf{I}_m$. Suppose $\nabla^2 \phi_k(x) = \eta \mathbf{I}_m$, then by (F.2), we obtain that

$$\nabla^2 \phi_{k+1}(x) = (1-\gamma) \nabla^2 \phi_k(x) + \gamma \eta \mathbf{I}_m = \eta \mathbf{I}_m.$$

Thus, $\nabla^2 \phi_k(x) = \eta \mathbf{I}_m$ for any $k \ge 0$. Because ϕ_k is a quadratic function, we can rewrite $\phi_k(x)$ as

$$\phi_k(x) = \phi_k^* + \frac{\eta}{2} \|x - v_k\|^2 \text{ with } v_k \triangleq \operatorname*{argmin}_x \phi_k(x), \quad \forall k \ge 0.$$
(F.4)

We proceed to give a recursive form for v_{k+1} and ϕ_{k+1}^* . Noting from (F.4) that $\nabla \phi_k(x) = \eta(x - v_k)$. Then by using the first-order optimality condition $\nabla \phi_{k+1}(x) = 0$ of the unconstrained convex optimization $\min_x \phi_{k+1}(x)$, and the definition of $\phi_{k+1}(x)$ in (F.2), we obtain that

$$\nabla \phi_{k+1}(x) = (1-\gamma)\eta(x-v_k) + \gamma h(x_k) + \gamma \eta(x-x_k) = 0.$$

By rearranging the previous equation, we have that

$$v_{k+1} = (1 - \gamma)v_k + \gamma x_k - \gamma h(x_k)/\eta.$$
 (F.5)

By using (F.2) and (F.4), evaluating $\phi_{k+1}(x)$ at $x = x_k$ we obtain that

$$\phi_{k+1}^* = \phi_{k+1}(x_k) - \frac{\eta}{2} \|x_k - v_{k+1}\|^2 = (1 - \gamma)\phi_k(x_k) + \gamma f(x_k) - \frac{\eta}{2} \|x_k - v_{k+1}\|^2$$

= $(1 - \gamma)\phi_k^* + \frac{\eta(1 - \gamma)}{2} \|x_k - v_k\|^2 + \gamma f(x_k) - \frac{\eta}{2} \|v_{k+1} - x_k\|^2.$ (F.6)

Note by (F.5) that

$$\|v_{k+1} - x_k\|^2 = \|(1 - \gamma)(v_k - x_k) - \gamma h(x_k)/\eta\|^2$$

= $(1 - \gamma)^2 \|v_k - x_k\|^2 + \frac{\gamma^2}{\eta^2} \|h(x_k)\|^2 - \frac{2\gamma(1 - \gamma)}{\eta} (v_k - x_k)^T h(x_k).$

This together with (F.6) leads to

$$\phi_{k+1}^* = (1-\gamma)\phi_k^* + \gamma f(x_k) + \frac{\eta\gamma(1-\gamma)}{2} \|x_k - v_k\|^2 - \frac{\gamma^2}{2\eta} \|h(x_k)\|^2 + \gamma(1-\gamma)(v_k - x_k)^T h(x_k).$$
(F.7)

We then show by induction that the following holds for any $k \ge 0$.

$$v_k - x_k = \frac{1}{\gamma} (x_k - y_k).$$
 (F.8)

From (F.1) it is seen that the minimizer of ϕ_0 is $v_0 = x_0$. Then by the initial condition $x_0 = y_0$, we conclude that (F.8) holds for k = 0. We inductively assume that (F.8) holds for k, and proceed to prove that (F.8) holds for k + 1. By substituting $v_k = x_k + (x_k - y_k)/\gamma$ into (F.5), one obtains $v_{k+1} = (1 - \gamma)(x_k + (x_k - y_k)/\gamma) + \gamma x_k - \gamma h(x_k)/\eta$. Hence

$$v_{k+1} - x_{k+1} = \frac{1}{\gamma} \left(x_k - \gamma^2 h(x_k) / \eta \right) - \left(\frac{1}{\gamma} - 1 \right) y_k - x_{k+1}.$$

This together with $h(x_k) = \frac{x_k - y_{k+1}}{\alpha}$ and $\gamma = \sqrt{\alpha \eta}$ produces

$$v_{k+1} - x_{k+1} = \frac{1}{\gamma} \left(x_k - \alpha h(x_k) \right) - \left(\frac{1}{\gamma} - 1 \right) y_k - x_{k+1}$$

$$\stackrel{(10b)}{=} \frac{1}{\gamma} y_{k+1} - \left(\frac{1}{\gamma} - 1 \right) \frac{(1+\beta)y_{k+1} - x_{k+1}}{\beta} - x_{k+1} = \frac{1}{\gamma} (x_{k+1} - y_{k+1}),$$

where the last equality holds by $\beta = \frac{1-\gamma}{1+\gamma}$. Thus, we have shows that (F.8) holds for any $k \ge 0$.

Then by substituting (F.8) into (F.7), and using $h(x_k) = \nabla f(x_k) + w_{k,N_k}$, we obtain that

$$\phi_{k+1}^{*} = (1-\gamma)\phi_{k}^{*} + \gamma f(x_{k}) + \frac{\eta(1-\gamma)}{2\gamma} \|x_{k} - y_{k}\|^{2} - \frac{\gamma^{2}}{2\eta} \|h(x_{k})\|^{2} + (1-\gamma)(x_{k} - y_{k})^{T} h(x_{k})$$

$$= (1-\gamma)\phi_{k}^{*} + \gamma f(x_{k}) - \frac{\gamma^{2}}{2\eta} \|h(x_{k})\|^{2} + (1-\gamma)(x_{k} - y_{k})^{T} \nabla f(x_{k})$$

$$+ \frac{\eta(1-\gamma)}{2\gamma} \|x_{k} - y_{k}\|^{2} + (1-\gamma)(x_{k} - y_{k})^{T} w_{k,N_{k}}$$

$$\geq (1-\gamma)\phi_{k}^{*} + \gamma f(x_{k}) - \frac{\gamma^{2}}{2\eta} \|h(x_{k})\|^{2} + (1-\gamma)(x_{k} - y_{k})^{T} \nabla f(x_{k}) - \frac{(1-\gamma)\gamma}{2\eta} \|w_{k,N_{k}}\|^{2},$$
(F.9)

where the last inequality follows by $\frac{\eta(1-\gamma)}{2\gamma} \|a\|^2 + (1-\gamma)a^Tb \ge -\frac{(1-\gamma)\gamma}{2\eta} \|b\|^2$. We proceed to show that $\phi_k^* \ge f(y_k) - p_k$ for any $k \ge 0$. By the definitions (F.1) and (F.3), we see

We proceed to show that $\phi_k^* \ge f(y_k) - p_k$ for any $k \ge 0$. By the definitions (F.1) and (F.3), we see that $\phi_0^* = f(x_0) = f(y_0)$ and $p_0 = 0$. Hence $\phi_k^* \ge f(y_k) - p_k$ holds for k = 0. We inductively assume that $f(y_k) \le \phi_k^* + p_k$, and aim to show that $f(y_{k+1}) - \phi_{k+1}^* \le p_{k+1}$. Since f is L-smooth, by using $h(x_k) = \frac{x_k - y_{k+1}}{\alpha} = \nabla f(x_k) + w_{k,N_k}$, we obtain that

$$f(y_{k+1}) \le f(x_k) + (y_{k+1} - x_k)^T \nabla f(x_k) + \frac{L}{2} \|y_{k+1} - x_k\|^2$$

$$\leq f(x_k) - \alpha h(x_k)^T (h(x_k) - w_{k,N_k}) + \frac{L\alpha^2}{2} \|h(x_k)\|^2$$

$$\leq f(x_k) + \left(\frac{L\alpha^2}{2} - \alpha\right) \|h(x_k)\|^2 + \alpha \|w_{k,N_k}\|^2 + \alpha w_{k,N_k}^T \nabla f(x_k),$$
(F.10)

where the last inequality holds because $h(x_k)^T w_{k,N_k} = ||w_{k,N_k}||^2 + w_{k,N_k}^T \nabla f(x_k)$. By the induction assumption $f(y_k) \le \phi_k^* + p_k$ and the convexity of f, we obtain that

$$f(x_k) = (1 - \gamma)f(y_k) + (1 - \gamma)(f(x_k) - f(y_k)) + \gamma f(x_k)$$

$$\leq (1 - \gamma)\phi_k^* + (1 - \gamma)p_k + (1 - \gamma)(x_k - y_k)^T \nabla f(x_k) + \gamma f(x_k).$$

The above bound combined with (F.10) produces

$$f(y_{k+1}) \leq (1-\gamma)\phi_k^* + \gamma f(x_k) + (1-\gamma)(x_k - y_k)^T \nabla f(x_k) + (1-\gamma)p_k + \left(\frac{L\alpha^2}{2} - \alpha\right) \|h(x_k)\|^2 + \alpha \|w_{k,N_k}\|^2 + \alpha w_{k,N_k}^T \nabla f(x_k).$$

It incorporated with (F.9) leads to the following relation:

$$f(y_{k+1}) - \phi_{k+1}^* \leq \left(\frac{L\alpha^2}{2} - \alpha + \frac{\gamma^2}{2\eta}\right) \|h(x_k)\|^2 + (1 - \gamma)p_k + \alpha w_{k,N_k}^T \nabla f(x_k) + \left(\alpha + \frac{(1 - \gamma)\gamma}{2\eta}\right) \|w_{k,N_k}\|^2 \leq p_{k+1},$$
(F.11)

where the last inequality holds by the definition of p_{k+1} in (F.3) and $\frac{L\alpha^2}{2} - \alpha + \frac{\gamma^2}{2\eta} = \frac{\alpha(\alpha L - 1)}{2} \leq 0$ from $\alpha \leq 1/L$ and $\gamma^2 = \alpha \eta < 1$. Therefore, we conclude that $f(y_k) \leq \phi_k^* + p_k$ for any $k \geq 0$.

Because f is η -strongly convex, from (F.2) and $\mathbb{E}[h(x_k)|\mathcal{F}_k] = \nabla f(x_k)$ it follows that

$$\mathbb{E}[\phi_{k+1}(x)|\mathcal{F}_k] = (1-\gamma)\phi_k(x) + \gamma \left(f(x_k) + (x-x_k)^T \nabla f(x_k) + \frac{\eta}{2} \|x-x_k\|^2\right)$$
$$\leq (1-\gamma)\phi_k(x) + \gamma f(x), \quad \forall x \in \mathbb{R}^m.$$

By taking unconditional expectations, we obtain that $\mathbb{E}[\phi_{k+1}(x)] \leq (1-\gamma)\mathbb{E}[\phi_k(x)] + \gamma f(x)$ for any $x \in \mathbb{R}^m$. Therefore, by rearranging terms and setting $x = x^*$ in the above inequality, we have

$$\mathbb{E}[\phi_{k+1}(x^*)] - f(x^*) \le (1 - \gamma)(\mathbb{E}[\phi_k(x^*)] - f(x^*)) \le (1 - \gamma)^{k+1}(\mathbb{E}[\phi_0(x^*)] - f(x^*)).$$

Then by the fact that $f(y_k) \leq \phi_k^* + p_k$, there holds

$$\mathbb{E}[f(y_k)] - f(x^*) \le \mathbb{E}[\phi_k(x^*)] - f(x^*) + \mathbb{E}[p_k] \le (1 - \gamma)^k (\mathbb{E}[\phi_0(x^*)] - f(x^*)) + \mathbb{E}[p_k] \\ \le \frac{\eta + L}{2} (1 - \gamma)^k \mathbb{E}[\|x_0 - x^*\|^2] + \mathbb{E}[p_k],$$
(F.12)

where the last inequality holds because $\phi_0(x^*) - f(x^*) = f(x_0) - f(x^*) + \frac{\eta}{2} ||x^* - x_0||^2 \le \frac{\eta + L}{2} ||x^* - x_0||^2$ by the *L*-smoothness of *f* and $\nabla f(x^*) = 0$. Next, we derive a bound on $\mathbb{E}[p_k]$. By taking expectations on both sides of (F.3), using $p_0 = 0$ and (5), we obtain that

$$\mathbb{E}[p_k] = (1-\gamma)\mathbb{E}[p_{k-1}] + \left(\alpha + \frac{(1-\gamma)\gamma}{2\eta}\right)\mathbb{E}[\|w_{k-1,N_{k-1}}\|^2]$$

$$= \left(\alpha + \frac{(1-\gamma)\gamma}{2\eta}\right) \sum_{i=0}^{k-1} (1-\gamma)^{i} \mathbb{E}[\|w_{k-i-1,N_{k-i-1}}\|^{2}] \le \nu^{2} \left(\alpha + \frac{(1-\gamma)\gamma}{2\eta}\right) \sum_{i=0}^{k-1} (1-\gamma)^{i} / N_{k-1-i}.$$

This together with (F.12) proves Lemma 2.

G Proof of Lemma 4

The proof of Lemma 4 is motivated by [2, Theorem 2].

Define

$$p_k = \frac{\beta}{1 - \beta} (x_k - x_{k-1}).$$
 (G.1)

Then by (12), we have

$$\begin{aligned} x_{k+1} + p_{k+1} &= x_{k+1} + \frac{\beta}{1-\beta} (x_{k+1} - x_k) = \frac{1}{1-\beta} x_{k+1} - \frac{\beta}{1-\beta} x_k \\ &= \frac{x_k - \alpha (\nabla f(x_k) + w_{k,N_k}) + \beta (x_k - x_{k-1})}{1-\beta} x_{k+1} - \frac{\beta}{1-\beta} x_k \\ &= x_k + p_k - \frac{\alpha}{1-\beta} (\nabla f(x_k) + w_{k,N_k}). \end{aligned}$$

This implies that

$$\|x_{k+1} + p_{k+1} - x^*\|^2 = \|x_k + p_k - x^*\|^2 - \frac{2\alpha}{1-\beta} (\nabla f(x_k) + w_{k,N_k})^T (x_k + p_k - x^*) + \frac{\alpha^2}{(1-\beta)^2} \|\nabla f(x_k) + w_{k,N_k}\|^2.$$
(G.2)

With the definition of \mathcal{F}_k and the update rule (12), we see that x_k is adapted to \mathcal{F}_k , and hence p_k is adapted to \mathcal{F}_k . Then by taking conditional expectations on both sides of (G.2) on \mathcal{F}_k , we obtain

$$\mathbb{E}[\|x_{k+1} + p_{k+1} - x^*\|^2 |\mathcal{F}_k] = \|x_k + p_k - x^*\|^2 - \frac{2\alpha}{1-\beta} (\nabla f(x_k) + \mathbb{E}[w_{k,N_k}|\mathcal{F}_k])^T (x_k + p_k - x^*) \\ + \frac{\alpha^2}{(1-\beta)^2} \left(\|\nabla f(x_k)\|^2 + 2\nabla f(x_k)^T \mathbb{E}[w_{k,N_k}|\mathcal{F}_k] + \mathbb{E}[\|w_{k,N_k}\|^2 |\mathcal{F}_k] \right) \\ \stackrel{(5)}{\leq} \|x_k + p_k - x^*\|^2 - \frac{2\alpha}{1-\beta} \nabla f(x_k)^T (x_k + p_k - x^*) + \frac{\alpha^2}{(1-\beta)^2} \left(\|\nabla f(x_k)\|^2 + \frac{\nu^2}{N_k} \right)$$
(G.3)
$$\stackrel{(G.1)}{=} \|x_k + p_k - x^*\|^2 - \frac{2\alpha}{1-\beta} \nabla f(x_k)^T (x_k - x^*) - \frac{2\alpha\beta}{(1-\beta)^2} \nabla f(x_k)^T (x_k - x_{k-1}) + \frac{\alpha^2}{(1-\beta)^2} \left(\|\nabla f(x_k)\|^2 + \frac{\nu^2}{N_k} \right)$$

Since $\nabla f(x^*) = 0$, Assumption 1(i) and (ii) hold, we recall from [3, (2.1.24)] that

$$\nabla f(x_k)^T(x_k - x^*) \ge \frac{\eta L}{L + \eta} \|x_k - x^*\|^2 + \frac{1}{L + \eta} \|\nabla f(x_k)\|^2.$$
(G.4)

Since f is η -strongly convex from Assumption 1(ii), we have

$$f(x_{k-1}) - f(x_k) \ge \nabla f(x_k)^T (x_{k-1} - x_k) + \frac{\eta}{2} \|x_{k-1} - x_k\|^2.$$

$$\Rightarrow \nabla f(x_k)^T (x_k - x_{k-1}) \ge f(x_k) - f(x_{k-1}) + \frac{\eta}{2} \|x_{k-1} - x_k\|^2.$$
(G.5)

Then by substituting (G.4) and (G.5) into (G.3), we obtain

$$\mathbb{E}[\|x_{k+1} + p_{k+1} - x^*\|^2 |\mathcal{F}_k] \le \|x_k + p_k - x^*\|^2 + \frac{\alpha^2}{(1-\beta)^2} \left(\|\nabla f(x_k)\|^2 + \frac{\nu^2}{N_k} \right) \\ - \frac{2\alpha}{1-\beta} \left(\frac{\eta L}{L+\eta} \|x_k - x^*\|^2 + \frac{1}{L+\eta} \|\nabla f(x_k)\|^2 \right) - \frac{2\alpha\beta}{(1-\beta)^2} \left(f(x_k) - f(x_{k-1}) + \frac{\eta}{2} \|x_{k-1} - x_k\|^2 \right).$$

By taking unconditional expectations on both sides of the above equation, and rearranging the terms, we get

$$\mathbb{E}[\|x_{k+1} + p_{k+1} - x^*\|^2 |\mathcal{F}_k] + \frac{2\alpha\beta}{(1-\beta)^2} (f(x_k) - f(x^*)) \le \|x_k + p_k - x^*\|^2 + \frac{2\alpha\beta}{(1-\beta)^2} (f(x_{k-1}) - f(x^*)) + \frac{\alpha^2\nu^2}{(1-\beta)^2N_k} - \frac{2\alpha\eta L}{(1-\beta)(L+\eta)} \|x_k - x^*\|^2 - \frac{\alpha\beta\eta}{(1-\beta)^2} \|x_{k-1} - x_k\|^2 + \frac{\alpha}{1-\beta} \left(\frac{\alpha}{1-\beta} - \frac{2}{L+\eta}\right) \|\nabla f(x_k)\|^2.$$
(G.6)

Since $\nabla f(x^*) = 0$ and Assumption 1(ii) holds, we recall from [3, (2.1.19)] that

$$2\eta(f(x_k) - f(x^*)) \le \|\nabla f(x_k)\|^2.$$
(G.7)

By recalling that $\beta \in (0,1)$ and $\alpha \in (0, \frac{2(1-\beta)}{L+\eta})$, the last term on the right-hand side of (G.6) becomes negative. Thus, by substituting (G.7) into (G.6), we derive

$$\mathbb{E}[\|x_{k+1} + p_{k+1} - x^*\|^2 |\mathcal{F}_k] + \frac{2\alpha}{1-\beta} \left(\frac{\beta - \alpha\eta}{1-\beta} + \frac{2\eta}{\eta+L} \right) (f(x_k) - f(x^*)) \le \|x_k + p_k - x^*\|^2 + \frac{2\alpha\beta}{(1-\beta)^2} (f(x_{k-1}) - f(x^*)) - \frac{2\alpha\eta L}{(1-\beta)(L+\eta)} \|x_k - x^*\|^2 - \frac{\alpha\beta\eta}{(1-\beta)^2} \|x_{k-1} - x_k\|^2 + \frac{\alpha^2\nu^2}{(1-\beta)^2N_k}.$$
 (G.8)

Define $\hat{z}_k = \begin{pmatrix} x_k - x^* \\ x_k - x_{k-1} \end{pmatrix}$. From (G.1) it can be seen that

$$\|x_k + p_k - x^*\|^2 = \hat{z}_k^T \mathbf{M} \hat{z}_k \text{ with } \mathbf{M} \triangleq \begin{pmatrix} \mathbf{I}_m & \frac{\beta}{1-\beta} \mathbf{I}_m \\ \frac{\beta}{1-\beta} \mathbf{I}_m & \left(\frac{\beta}{(1-\beta)^2}\right)^2 \mathbf{I}_m \end{pmatrix}.$$

Then (G.8) can be rewritten as

$$\mathbb{E}[\hat{z}_{k+1}^T \mathbf{M} \hat{z}_{k+1} | \mathcal{F}_k] + \hat{m}(f(x_k) - f(x^*)) \le \hat{z}_k^T \mathbf{N} \hat{z}_k + \hat{n}(f(x_{k-1}) - f(x^*)) + \frac{\alpha^2 \nu^2}{(1-\beta)^2 N_k}, \tag{G.9}$$

where

$$\hat{m} \triangleq \frac{2\alpha}{1-\beta} \left(\frac{\beta-\alpha\eta}{1-\beta} + \frac{2\eta}{\eta+L} \right), \ \hat{n} \triangleq \frac{2\alpha\beta}{(1-\beta)^2}, \ \text{and} \ \mathbf{N} \triangleq \left(\begin{array}{cc} \left(1 - \frac{2\alpha\eta L}{(1-\beta)(L+\eta)} \right) \mathbf{I}_m & \frac{\beta}{1-\beta} \mathbf{I}_m \\ \frac{\beta}{1-\beta} \mathbf{I}_m & \frac{\beta(\beta-\alpha\eta)}{(1-\beta)^2} \mathbf{I}_m \end{array} \right).$$

With the definitions of q_1 and q_2 in (15), it has been shown in [2, Appendix] that $\hat{n} \leq q_1 \hat{m}$ and $q_2 \mathbf{M} - \mathbf{N}$ is positive semidefinite. Then from (G.9) and $q = \max\{q_1, q_2\} \in (0, 1)$, we obtain that

$$\mathbb{E}[\hat{z}_{k+1}^T \mathbf{M} \hat{z}_{k+1} | \mathcal{F}_k] + \hat{m}(f(x_k) - f(x^*)) \le q \left(\hat{z}_k^T \mathbf{M} \hat{z}_k + \hat{m}(f(x_{k-1}) - f(x^*))\right) + \frac{\alpha^2 \nu^2}{(1-\beta)^2 N_k}$$

By taking unconditional expectations on both sides of the above inequality, we achieve

$$\mathbb{E}[\hat{z}_{k+1}^T \mathbf{M} \hat{z}_{k+1} + \hat{m}(f(x_k) - f(x^*))] \le q \mathbb{E}\left[\hat{z}_k^T \mathbf{M} \hat{z}_k + \hat{m}(f(x_{k-1}) - f(x^*))\right] + \frac{\alpha^2 \nu^2}{(1-\beta)^2 N_k}$$

$$\leq q^{k+1} \mathbb{E} \left[\hat{z}_0^T \mathbf{M} \hat{z}_0 + \hat{m} (f(x_{-1}) - f(x^*)) \right] + \frac{\alpha^2 \nu^2}{(1-\beta)^2} \sum_{i=0}^k q^i / N_{k-i}$$
$$= q^{k+1} \mathbb{E} \left[\|x_0 - x^*\|^2 + \hat{m} (f(x_0) - f(x^*)) \right] + \frac{\alpha^2 \nu^2}{(1-\beta)^2} \sum_{i=0}^k q^i / N_{k-i}, \tag{G.10}$$

where the last inequality holds since $x_{-1} = x_0$ and $\hat{z}_0 = \begin{pmatrix} x_0 - x^* \\ 0 \end{pmatrix}$.

Since $\nabla f(x^*) = 0$ and Assumption 1(ii) holds, we have $f(x_k) - f(x^*) \ge 2\eta ||x_k - x^*||^2$. Then by using (G.10) and recalling the positive semidefiniteness of **M**, we obtain

$$2\hat{m}\eta\mathbb{E}[\|x_k - x^*\|^2] \le q^{k+1}\mathbb{E}\left[\|x_0 - x^*\|^2 + \hat{m}(f(x_0) - f(x^*))\right] + \frac{\alpha^2\nu^2}{(1-\beta)^2}\sum_{i=0}^k q^i/N_{k-i}.$$

Thus the required result holds.

H Proof of Proposition 2

By using $\rho_2 \in (1 - \gamma, 1)$ and substituting $N_k = \lceil \rho_2^{-(k+1)} \rceil$ into (11), we obtain that

$$\mathbb{E}[f(y_k)] - f^* \le \rho_2^k \frac{\eta + L}{2} \mathbb{E}[\|x_0 - x^*\|^2] + \nu^2 \left(\alpha + \frac{(1 - \gamma)\gamma}{2\eta}\right) \rho_2^k \sum_{i=0}^{k-1} \left(\frac{1 - \gamma}{\rho_2}\right)^i,$$

which leads to (22) by using the bound $\sum_{i=0}^{k-1} \left(\frac{1-\gamma}{\rho_2}\right)^i \leq \frac{1}{1-\frac{1-\gamma}{\rho_2}} = \frac{\rho_2}{\rho_2-(1-\gamma)}.$

Because f is η -strongly convex and $\nabla f(x^*) = 0$, we have $f(x) - f(x^*) \ge \frac{\eta}{2} ||x - x^*||^2$. Thus, y_k generated by Algorithm 2 satisfies $||y_k - x^*||^2 \le \frac{2}{\eta} (f(y_k) - f^*)$. Then from (22) it follows that

$$\mathbb{E}[\|y_k - x^*\|^2] \le c\rho_2^k \text{ with } c \triangleq \frac{2}{\eta} \left(\frac{\eta + L}{2} \mathbb{E}[\|x_0 - x^*\|^2] + \frac{\rho_2 \nu^2}{\rho_2 - (1 - \gamma)} \left(\alpha + \frac{(1 - \gamma)\gamma}{2\eta} \right) \right).$$

Note by (10b) that $||x_k - x^*|| \le ||(1 + \beta)(y_k - x^*)|| + ||\beta(y_{k-1} - x^*)||$, and hence

$$\mathbb{E}[\|x_k - x^*\|^2] \le 2(1+\beta)^2 \mathbb{E}[\|y_k - x^*\|^2] + 2\beta^2 \mathbb{E}[\|y_{k-1} - x^*\|^2] \\ \le c \left(2(1+\beta)^2 + 2\beta^2 \rho_2^{-1}\right) \rho_2^k.$$

Therefore, sequences $\{x_k\}$ and $\{y_k\}$ converge to the optimal solution x^* at a geometric rate $\mathcal{O}(\rho_2^k)$ in the mean-squared sense.

Suppose we set $\alpha \triangleq \frac{1}{L}$. Then $\gamma = \sqrt{\frac{n}{L}} = \frac{1}{\sqrt{\kappa}}$ and $\beta = \frac{\sqrt{\kappa}-1}{\sqrt{\kappa}+1}$. Select $\rho_2 \triangleq 1 - \frac{1}{2\sqrt{\kappa}}$ such that $\rho_2 > 1 - \gamma$. We can show that the number of iterations required to obtain an ϵ -optimal solution in a mean-squared sense is $\mathcal{O}\left(\frac{\ln(1/\epsilon)}{\ln(1/\rho_2)}\right) = \mathcal{O}\left(\sqrt{\kappa}\ln(1/\epsilon)\right)$ since $\ln\left(\frac{1}{1-1/(2\sqrt{\kappa})}\right) \approx \frac{1}{2\sqrt{\kappa}}$ for large κ , and hence the oracle complexity $\sum_{k=0}^{K(\epsilon)-1} N_k = \mathcal{O}\left(\sqrt{\kappa}/\epsilon\right)$.

I Proof of Proposition 3

By substituting $\alpha = \frac{4}{(\sqrt{\eta} + \sqrt{L})^2}$ into $\beta = \max\{|1 - \sqrt{\alpha \eta}|, |1 - \sqrt{\alpha L}|\}^2$, there holds $\beta = \left(1 - \frac{2}{\sqrt{\kappa} + 1}\right)^2 < 1$. Then Lemma 3 holds. Therefore, by using (13), $N_k = \lceil \rho_3^{-(k+1)} \rceil$, $x_{-1} = x_0$, $\rho_3 \in (\beta, 1)$, and $\iota \in (0, \sqrt{\rho_3} - \sqrt{\beta})$, there exists a constant $c(\iota)$ such that

$$\mathbb{E}\left[\left\| \begin{pmatrix} x_{k+1} - x^* \\ x_k - x^* \end{pmatrix} \right\|^2 \right] \le 2c(\iota)^2 (\sqrt{\beta} + \iota)^{2(k+1)} \mathbb{E}[\|x_0 - x^*\|^2] + \alpha^2 \nu^2 c(\iota)^2 \sum_{t=0}^k (\sqrt{\beta} + \iota)^{2(k-t)} \rho_3^{t+1}, \quad \forall k \ge 0.$$

Since $\iota \in (0, \sqrt{\rho_3} - \sqrt{\beta})$, we have $\sqrt{\beta} + \iota \in (\sqrt{\beta}, \rho_3)$. This together with $\sum_{t=0}^k (\sqrt{\beta} + \iota)^{2(k-t)} \rho_3^{t+1} = \sum_{t=0}^k (\sqrt{\beta} + \iota)^{2t} \rho_3^{k+1-t} = \rho_3^{k+1} \sum_{t=0}^k \left(\frac{(\sqrt{\beta} + \iota)^2}{\rho_3} \right)^t \le \frac{\rho_3^{k+1}}{1 - (\sqrt{\beta} + \iota)^2/\rho_3}$, proving (23).

By (23), $\mathbb{E}[\|x_k - x^*\|] \leq c\rho_3^k$ for some constant c > 0. Suppose $\rho_3 = \left(1 - \frac{1}{\sqrt{\kappa}+1}\right)^2 > \beta$. Akin to the proof of Proposition 1(ii), we can show that the number of iterations required to obtain an ϵ -optimal solution satisfying $\mathbb{E}[\|x_k - x^*\|^2] \leq \epsilon$ is $\mathcal{O}\left(\frac{\ln(1/\epsilon)}{\ln(1/\rho_3)}\right) = \mathcal{O}\left(\sqrt{\kappa}\ln(1/\epsilon)\right)$ since $\ln\left(\frac{1}{1-1/(\sqrt{\kappa}+1)}\right) \approx \frac{1}{\sqrt{\kappa}+1}$ for large κ , and the oracle complexity $\sum_{k=0}^{K(\epsilon)-1} N_k = \mathcal{O}\left(\sqrt{\kappa}/\epsilon\right)$.

J Proof of Proposition 4

By substituting $N_k = \lceil \rho_4^{-(k+1)} \rceil$ into (14), we obtain

$$\mathbb{E}[\|x_k - x^*\|^2] \le \frac{1}{2\hat{m}\eta} \mathbb{E}\left[\|x_0 - x^*\|^2 + \hat{m}(f(x_0) - f(x^*))\right] q^{k+1} + \frac{\alpha^2 \nu^2}{2(1-\beta)^2 \hat{m}\eta} \rho_4^{k+1} \sum_{i=0}^n (q/\rho_4)^i \\ \le \frac{1}{2\hat{m}\eta} \mathbb{E}\left[\|x_0 - x^*\|^2 + \hat{m}(f(x_0) - f(x^*))\right] \rho_4^{k+1} + \frac{\alpha^2 \nu^2}{2(1-\beta)^2 \hat{m}\eta(1-q/\rho_4)} \rho_4^{k+1},$$

ı

where the last inequality holds by $\rho_4 \in (q, 1)$. Then (24) holds.

K Proof of Proposition 5

By substituting $N_k \triangleq \lceil (k+1)^v \rceil$ into (7), using $q = 1 - \frac{2\alpha\eta L}{\eta + L}$ and Lemma 8(i), one obtains

$$\mathbb{E}[\|x_k - x^*\|^2] \le q\mathbb{E}[\|x_{k-1} - x^*\|^2] + \alpha^2 \nu^2 k^{-\nu} = q^k \mathbb{E}[\|x_0 - x^*\|^2] + \alpha^2 \nu^2 \sum_{m=1}^k q^{k-m} m^{-\nu} = q^k \mathbb{E}[\|x_0 - x^*\|^2] + \alpha^2 \nu^2 \left(q^k \frac{e^{2\nu} q^{-1} - 1}{1 - q} + \frac{2k^{-\nu}}{q \ln(1/q)}\right),$$

which together with Lemma 8(ii) proves (48). Then $\mathbb{E}[||x_k - x^*||^2] \leq \epsilon$ for any $k \geq K(\epsilon) \triangleq \left(\frac{C_v}{\epsilon}\right)^{1/v}$. By noting that $C_v = \mathcal{O}(e^v v^v)$, the iteration complexity is $\mathcal{O}(v(1/\epsilon)^{1/v})$. Therefore, the number of sampled gradients required to obtain an ϵ -NE is bounded by

$$\begin{split} \sum_{k=0}^{K(\epsilon)-1} \lceil (k+1)^v \rceil &\leq K(\epsilon) + (K(\epsilon))^v + \sum_{k=1}^{K(\epsilon)-1} k^v \leq \left(\frac{C_v}{\epsilon}\right)^{1/v} + \frac{C_v}{\epsilon} + \int_1^{K(\epsilon)} t^v dt \\ &= \left(\frac{C_v}{\epsilon}\right)^{1/v} + \frac{C_v}{\epsilon} + \frac{t^{v+1}}{v+1} \Big|_1^{K(\epsilon)} = \left(\frac{C_v}{\epsilon}\right)^{1/v} + \frac{C_v}{\epsilon} + (v+1)^{-1} \left(\frac{C_v}{\epsilon}\right)^{1+\frac{1}{v}} - (v+1)^{-1}. \end{split}$$

Therefore, the oracle complexity is $\mathcal{O}\left(e^{v}v^{v}\left(1/\epsilon\right)^{1+\frac{1}{v}}\right)$.

Proof of Proposition 6 L

It is noticed by t = k - i and Lemma 8(i),

$$\sum_{i=0}^{k-1} (1-\gamma)^i (k-i)^{-\nu} = \sum_{t=1}^k (1-\gamma)^{k-t} t^{-\nu} \le \frac{(1-\gamma)^k (e^{2\nu}(1-\gamma)^{-1}-1)}{\gamma} + \frac{2k^{-\nu}}{(1-\gamma)\ln(1/(1-\gamma))}.$$
 (L.1)

By substituting $N_k = \lceil (k+1)^v \rceil$ into (11), we obtain that

$$\mathbb{E}[f(y_k)] - f^* \le \frac{(1-\gamma)^k (\eta+L)\mathbb{E}[\|x_0-x^*\|^2]}{2} + \nu^2 \left(\alpha + \frac{(1-\gamma)\gamma}{2\eta}\right) \sum_{i=0}^{k-1} (1-\gamma)^i (k-i)^{-\nu}.$$

This together with Lemma 8(ii) and (L.1) proves the required result.

Proof of Proposition 7 Μ

By substituting $N_k = \lfloor (k+1)^v \rfloor$ into (13), we obtain that for any $\iota \in (0, 1 - \sqrt{\beta})$, there exists a constant $c(\iota)$ such that for any $k \ge 0$,

$$\mathbb{E}\left[\left\| \begin{pmatrix} x_{k+1} - x^* \\ x_k - x^* \end{pmatrix} \right\|^2 \right] \le 2(c(\iota))^2 (\sqrt{\beta} + \iota)^{2(k+1)} \mathbb{E}[\|x_0 - x^*\|^2] + \alpha^2 \nu^2 (c(\iota))^2 \sum_{t=1}^{k+1} (\sqrt{\beta} + \iota)^{2(k+1-t)} t^{-v} \\ \le 2(c(\iota))^2 (\sqrt{\beta} + \iota)^{2(k+1)} \mathbb{E}[\|x_0 - x^*\|^2] \\ + \alpha^2 \nu^2 (c(\iota))^2 \left((\sqrt{\beta} + \iota)^{2(k+1)} \frac{e^{2v} (\sqrt{\beta} + \iota)^{-2} - 1}{1 - (\sqrt{\beta} + \iota)^2} + \frac{2(k+1)^{-v}}{2(\sqrt{\beta} + \iota)^2 \ln(1/(\sqrt{\beta} + \iota))} \right)$$

where the last inequality holds by using Lemma 8(i) and recalling that $\sqrt{\beta} + \iota \in (\sqrt{\beta}, 1)$. This together with Lemma 8(ii) yields the result.

Ν **Proof of Proposition 8**

By substituting $N_k = \lceil (k+1)^v \rceil$ into (14), we obtain that

$$\mathbb{E}[\|x_k - x^*\|^2] \le \frac{1}{2\hat{m}\eta} \mathbb{E}\left[\|x_0 - x^*\|^2 + \hat{m}(f(x_0) - f(x^*))\right] q^{k+1} + \frac{\alpha^2 \nu^2}{2(1-\beta)^2 \hat{m}\eta} \sum_{i=0}^k q^i (k+1-i)^{-\nu}, \forall k \ge 0.$$

By using Lemma 8(i), we derive $\sum_{i=0}^{k} q^i (k+1-i)^{-v} = \sum_{t=1}^{k+1} q^{k+1-t} t^{-v} \le q^{k+1} \frac{e^{2v}q^{-1}-1}{1-q} + \frac{2(k+1)^{-v}}{q\ln(1/q)}.$ Thus,

$$\mathbb{E}[\|x_k - x^*\|^2] \le \frac{1}{2\hat{m}\eta} \mathbb{E}\left[\|x_0 - x^*\|^2 + \hat{m}(f(x_0) - f(x^*))\right] q^{k+1} + \frac{\alpha^2 \nu^2}{2(1-\beta)^2 \hat{m}\eta} \left(q^{k+1} \frac{e^{2\nu} q^{-1} - 1}{1-q} + \frac{2(k+1)^{-\nu}}{q\ln(1/q)}\right)$$

This together with Lemma 8(ii) yields the result.

This together with Lemma 8(ii) yields the result.

References

- Bubeck, S.: Theory of convex optimization for machine learning. arXiv preprint arXiv:1405.4980 15 (2014)
- [2] Ghadimi, E., Feyzmahdavian, H.R., Johansson, M.: Global convergence of the heavy-ball method for convex optimization. In: 2015 European control conference (ECC), pp. 310–315. IEEE (2015)
- [3] Nesterov, Y.: Introductory lectures on convex optimization: A basic course, vol. 87. Springer Science & Business Media (2013)