

ON THE POWER OF GRAPH NEURAL NETWORKS AND THE ROLE OF THE ACTIVATION FUNCTION

SAMMY KHALIFE AND AMITABH BASU

ABSTRACT. In this article we present new results about the expressivity of Graph Neural Networks (GNNs). We prove that for any GNN with piecewise polynomial activations, whose architecture size does not grow with the graph input sizes, there exists a pair of non-isomorphic rooted trees of depth two such that the GNN cannot distinguish their root vertex up to an arbitrary number of iterations. In contrast, it was already known that unbounded GNNs (those whose size is allowed to change with the graph sizes) with piecewise polynomial activations can distinguish these vertices in only two iterations. It was also known prior to our work that with ReLU (piecewise linear) activations, bounded GNNs are weaker than unbounded GNNs [ACI⁺22]. Our approach adds to this result by extending it to handle any piecewise polynomial activation function, which goes towards answering an open question formulated by Grohe [Gro21] more completely. Our second result states that if one allows activations that are not piecewise polynomial, then in two iterations a single neuron perceptron can distinguish the root vertices of any pair of nonisomorphic trees of depth two (our results hold for activations like the sigmoid, hyperbolic tan and others). This shows how the power of graph neural networks can change drastically if one changes the activation function of the neural networks. The proof of this result utilizes the Lindemann-Weierstrauss theorem from transcendental number theory.

1. INTRODUCTION

Graph Neural Networks (GNNs) form a popular framework for a variety of computational tasks involving network data, with applications ranging from analysis of social networks, structure and functionality of molecules in chemistry and biological applications, computational linguistics, simulations of physical systems, techniques to enhance optimization algorithms, to name a few. The interested reader can look at [ZAL18, BBL⁺17, DZC⁺19, DBV16, SGT⁺08, Ham20, DMI⁺15, KDZ⁺17, SYS⁺20, ZCH⁺20, BPL⁺16, CCK⁺21, SGGP⁺20], which is a small sample of a large and actively growing body of work.

Given the rise in importance of inference and learning problems involving graphs and the use of GNNs for these tasks, significant progress has been made in recent years to understand their computational capabilities. See the excellent recent survey [Jeg22] for an exposition of some aspects of this research. One direction of investigation is on their so-called *separation power* which is the ability of GNNs to distinguish graphs with different structures. In this context, it becomes natural to compare their separation power to other standard computation models on graphs, such

JOHNS HOPKINS UNIVERSITY, DEPARTMENT OF APPLIED MATHEMATICS AND STATISTICS

E-mail addresses: khalife.sammy@jhu.edu, basu.amitabh@jhu.edu.

2020 *Mathematics Subject Classification.* 68T07, 68Q19, 05D10, 11J85.

as different variants of the *Wesfeiler-Lehman algorithm* [CFI92, XHLJ18, HV21], and the very closely related *color refinement algorithm* [Gro21]. These investigations are naturally connected with *descriptive complexity theory*, especially to characterizations in terms of certain logics; see [Gro17, Gro21] for introductions to these different connections. A closely related line of work is to investigate how well general functions on the space of graphs can be approximated using functions represented by GNNs; see [AL21, KP19, MFSL19, GR22] for a sample of work along these lines. Our work in this paper focuses on improving our understanding of the separation power of GNNs.

At a high level, the computational models of GNNs, Wesfeiler-Lehman/color refinement type algorithms and certain logics in descriptive complexity are intimately connected because they all fall under the paradigm of trying to discern something about the global structure of a graph from local neighborhood computations. Informally, these algorithms iteratively maintain a state (a.k.a. “color”) for each vertex of the graph and in every iteration, the state of a vertex is updated by performing some predetermined set of operations on the set of current states of its neighbors (including itself). The different kinds of allowed states and allowed operations determine the computational paradigm. For instance, in GNNs, the states are typically vectors in some Euclidean space and the operations for updating the state are functions that can be represented by deep neural networks. As another example, in the color refinement algorithm, the states are multisets of some predetermined finite class of labels and the operations are set operations on these multisets. A natural question then arises: Given two of these models, which one is more powerful, or equivalently, can one of the models always simulate the other? Several mathematically precise answers to such questions have already been obtained. For instance, it has been proved independently by [MRF⁺19] and [XHLJ18] that the color refinement algorithm precisely captures the expressiveness of GNNs in the sense that there is a GNN distinguishing two nodes of a graph (by assigning them different state vectors) if and only if color refinement assigns different multisets to these nodes. Such a characterization holds for *unbounded* GNNs, i.e. GNNs for which the underlying neural network sizes can grow with the size of the input graphs. This implies a characterization of the distinguishability of nodes by GNNs as being equivalent to what is known as *Graded Modal Counting Logic* (GC2); see [BKM⁺20] for some recent, quantitatively precise results in this direction.

Reviewing these equivalences in a recent survey [Gro21], Grohe emphasizes the fact that the above mentioned equivalence between the separation power of GNNs and the color refinement algorithm has only been established for unbounded GNNs whose neural network sizes are allowed to grow as a function of the size of the input graphs. Question 1 on his list of important open questions in this topic asks what happens if one considers *bounded* GNNs, i.e., the the size of the neural networks is fixed a priori and cannot change as a function of the size of the input graphs. *Do bounded GNNs have the same separation power as unbounded GNNs and color refinement?* In [ACI⁺22], the authors provide a negative answer to this question in the case of Rectified Linear Unit (ReLU) activations, by proving lower bounds on the size of the GNNs with ReLU activations that are needed to simulate color refinement. Their approach combines results from communication complexity and properties of ReLU neural networks, including upper bounds on the number of their linear regions. In this article, we answer this question in the negative in the more

general case of *piecewise polynomial activations*. Given any bounded GNN with such activations, we explicitly construct two non isomorphic rooted trees of depth two such that their root nodes cannot be distinguished by the GNN. Interestingly, only the sizes of the trees depend on the GNN, but their depth does not. This result is stated formally in Theorem 3 and it holds for bounded GNNs with piecewise polynomial activations (this includes, e.g., ReLU activations). We prove a second result that shows how the activation function dramatically impacts the expressivity of bounded size GNNs: if one allows activation functions that are not piecewise polynomial, all root nodes of rooted trees of depth two can be distinguished by a single neuron perceptron. This result is formally stated in Theorem 4. [AGA⁺24] provide a more general result for analytic non polynomial activation functions: it is possible to simulate color refinement *on any graph* if one allows analytic non polynomial functions. [BKR24] also showed that among GNNs with analytic non polynomial activations, those with one dimensional embeddings at every iteration are sufficient to simulate color-refinement. In contrast with approaches presented in [AGA⁺24, BKR24], our proof does not use the analytic properties of the activation function, but linear independence over the integers of the activation function evaluated at distinct integers.

The rest of this article is organized as follows. In Section 2 we present the main definitions and formal statement of our results. In Section 3 we give an overview of the proofs. Sections 4 and 5 fill in the technical details.

2. FORMAL STATEMENT OF RESULTS

We assume graphs to be finite, undirected, simple, and vertex-labelled: a graph is a tuple $G = (V(G), E(G), P_1(G), \dots, P_\ell(G))$ consisting of a finite vertex set $V(G)$, a binary edge relation $E(G) \subset V(G)^2$ that is symmetric and irreflexive, and unary relations $P_1(G), \dots, P_\ell(G) \subset V(G)$ representing $\ell > 0$ vertex labels. In the following, we suppose that the $P_i(G)$'s form a partition of the set of vertices of G , i.e. each vertex has a unique label. Also, the number ℓ of labels, which we will also call *colors*, is supposed to be fixed and does not grow with the size of the input graphs. When there is no ambiguity about which graph G is being considered, $N(v)$ refers to the set of neighbors of v in G not including v . $|G|$ will denote the number of vertices of G . We use simple curly brackets for a set $X = \{x \in X\}$ and double curly brackets for a multiset $Y = \{\{y \in Y\}\}$. For a set X , $|X|$ is the cardinal of X . When m is a positive integer, \mathfrak{S}_m is the set of permutations of $\{1, \dots, m\}$. We will also need the following basic notions from algebra. Given any ring $(R, +, \times)$, $R[X_1, \dots, X_m]$ will denote the polynomial ring in m indeterminates with coefficients in R . An (additive) subgroup generated by $\sigma_1, \dots, \sigma_q \in R$ is the smallest (inclusion-wise) additive subgroup of R that contains $\sigma_1, \dots, \sigma_q$. Equivalently, it corresponds to the set $\{\lambda_1\sigma_1 + \dots + \lambda_q\sigma_q : \lambda \in \mathbb{Z}^q\}$.

Definition 1 (Piecewise polynomial). *Let m be a positive integer. A function $f : \mathbb{R}^m \rightarrow \mathbb{R}$ is piecewise polynomial iff there exist multivariate polynomials $P_1, \dots, P_r \in \mathbb{R}[X_1, \dots, X_m]$ such that for any $x \in \mathbb{R}^m$, there exists $i \in \{1, \dots, r\}$ such that $f(x) = P_i(x)$. A polynomial region A of f is a set such that there exists $i \in [r]$ such that $A \subseteq \{x \in \mathbb{R}^m : f(x) = P_i(x)\}$. The degree of a piecewise polynomial function f is $\deg(f) := \max\{\deg(P_1), \dots, \deg(P_r)\}$. The number of polynomial pieces of a piecewise polynomial f is the smallest r such that f can be represented as above.*

Definition 2 (Finitely generated polynomial). *Let q be a positive integer. A multivariate polynomial P is q -generated provided there exist reals $\sigma_1, \dots, \sigma_q$ such that $P = \sum_{\alpha \in S} \gamma_\alpha X^\alpha$, where γ_α belongs to the additive subgroup of \mathbb{R} generated by $\sigma_1, \dots, \sigma_q$, and S is the set of exponents of P . Under these conditions, we say that P is q -generated by the reals $\sigma_1, \dots, \sigma_q$.*

Definition 3 (Embedding, equivariance, and refinement). *Given a set X , an embedding ξ is a function that takes as input a graph G and a vertex $v \in V(G)$, and returns an element $\xi(G, v) \in X$. An embedding is equivariant if and only if for any pair of isomorphic graphs G, G' , and any isomorphism f from G to G' , it holds that $\xi(G, v) = \xi(G', f(v))$. We say that ξ refines ξ' if and only if for any graph G and any $v \in V(G)$, $\xi(G, v) = \xi(G, v') \implies \xi'(G, v) = \xi'(G, v')$. When the graph G is clear from context, we use $\xi(v)$ as shorthand for $\xi(G, v)$.*

Definition 4 (Color refinement). *Given a graph G , and $v \in V(G)$, let $(G, v) \mapsto \text{col}(G, v)$ be the function which returns the color of the node v . The color refinement refers to a procedure that returns a sequence of equivariant embeddings cr^t , computed recursively as follows:*

- $\text{cr}^0(G, v) = \text{col}(G, v)$
- For $t \geq 0$, $\text{cr}^{t+1}(G, v) := (\text{cr}^t(G, v), \{\{\text{cr}^t(G, w) : w \in N(v)\}\})$

In each round, the algorithm computes a coloring that is finer than the one computed in the previous round, that is, cr^{t+1} refines cr^t . For some $t \leq n := |G|$, this procedure stabilises: the coloring does not become strictly finer anymore.

Remark 1. *We refer the reader to the seminal work [CFI92, Sections 2 and 5] for comments about the history and connections between the color refinement and Weisfeiler-Lehman algorithms.*

Definition 5 (Graph Neural Network (GNN)). *A GNN is a recursive embedding of vertices of a labelled graph by relying on the underlying adjacency information and node features. Each vertex v is attributed an indicator vector $\xi^0(v)$ of size ℓ , encoding the color of the node v : the colors being indexed by the palette $\{1, \dots, \ell\}$, $\xi^0(v) = e_i$ (the i -th canonical vector) if the color of the vertex v is i . The GNN is fully characterized by:*

- A combination function $\text{comb} : \mathbb{R}^{2\ell} \rightarrow \mathbb{R}^\ell$ which is a feedforward neural network with given activation function $\sigma : \mathbb{R} \rightarrow \mathbb{R}$.
- The update rule of the GNN at iteration $t \in \mathbb{N}$ for any labelled graph G and vertex $v \in V(G)$, is given as follows:

$$\xi^0(v) \text{ is the indicator vector of the color of } v, \quad \xi^{t+1}(v) = \text{comb}(\xi^t(v), \sum_{w \in N(v)} \xi^t(w))$$

Remark 2. *This type of GNN is sometimes referred to as a **recurrent GNN**. The general definition (cf. for instance [Gro21]) usually considers a sequence of combine and aggregation functions which may depend on the iteration t . The aggregation functions replace the sum over the neighborhood, i.e. at each iteration $\text{comb}(\xi^t(v), \text{agg}(\{\{\xi^t(w) : w \in N(v)\}\}))$ is the new embedding for vertex v . It has been proved in [XHLJ18] that for any agg function, there is a GNN (of potentially larger size) whose aggregation function is the summation and which refines any GNN with this aggregation function. The results of this article extend to GNNs whose combination and aggregation functions are allowed to be different in different*

iterations, but are multivariate piecewise polynomials. For ease of presentation, we restrict to recurrent GNNs.

Given these definitions, we can now formally state the previously known results about the expressivity of unbounded GNNs (Theorems 1 and 2). Namely, in Theorem 2, the size of the GNN is allowed to grow with n .

Theorem 1. [Gro21, XHLJ18, MRF⁺19] *Let $d \geq 1$, and let ξ^d be an embedding computed by a GNN after d iterations. Then cr^d refines ξ , that is, for all graphs G, G' and vertices $v \in V(G)$, $v' \in V(G')$, $\text{cr}^d(v) = \text{cr}^d(v') \implies \xi^d(G, v) = \xi^d(G', v')$.*

Theorem 2. [Gro21, Theorem VIII.4][XHLJ18, MRF⁺19] *Let $n \in \mathbb{N}$. Then there is a recurrent GNN such that for all $t = 0, \dots, n$, the vertex invariant ξ^t computed in the t -th iteration of the GNN refines cr^t on all graphs of order at most n .*

In contrast, we prove Theorems 3 and 4 for bounded GNNs:

Theorem 3. *For any GNN, i.e., choice of combination function, represented by a feedforward neural network with piecewise polynomial activation, and any natural number $I \in \mathbb{N}$, there exists a pair of rooted trees T and T' (unicolored, i.e. $\ell = 1$) of depth two with root nodes s and s' respectively such that:*

- $\text{cr}^2(T, s) \neq \text{cr}^2(T', s')$, i.e. s and s' can be distinguished with color refinement in two iterations.
- $\xi^t(T, s) = \xi^t(T', s')$ for all $t \leq I$, i.e., s and s' cannot be distinguished by the GNN until iteration $I + 1$.

Theorem 4. *In two iterations, a single neuron perceptron with an activation $\sigma \in \{\exp, \text{sigmoid}, \cosh, \sinh, \tanh\}$ can distinguish the root nodes of any pair of non-isomorphic rooted trees of depth two.*

3. OVERVIEW OF THE PROOFS

To establish our first result, we will use rooted trees of the form shown in Figure 1 which is a tree of depth two whose depth one vertices have prescribed degrees k_1, \dots, k_m , with $k_1, \dots, k_m \geq 1$. Given a GNN with piecewise polynomial activation and a natural number $I \in \mathbb{N}$, we will show that there exist two sets of integers k_1, \dots, k_m and k'_1, \dots, k'_m that are not the same up to permutations, such that for the corresponding rooted trees $T[k_1, \dots, k_m]$ and $T[k'_1, \dots, k'_m]$, the GNN cannot distinguish s and s' for the first I iterations, i.e. $\xi^t(T, s) = \xi^t(T', s')$ for any $t \in \{1, \dots, I\}$. Note that the natural numbers m , and k_1, \dots, k_m and k'_1, \dots, k'_m will depend on I , the activation and the size of the neural network considered.

The proof of the first result is structured as follows. Since the trees are parameterized by m -tuples of integers k_1, \dots, k_m , the embedding of the root node computed by the GNN at any iteration is a function of these m integers. Since the activations are piecewise polynomials, these embeddings of the root node are also piecewise multivariate polynomial functions of k_1, \dots, k_m . We further prove that the *complexity* of its polynomial pieces can be controlled uniformly in the sense of Definition 2: the number of generators of such polynomial is independent of m , but depends only on the number of iterations of the GNN and the underlying neural network (Lemma 3 and 4). Then, we show that there exists a large enough region of \mathbb{R}^m on which this piecewise polynomial function is evaluated by the *same*

polynomial. This region is large enough in the following sense: we prove that for a dimension m that is sufficiently large, the region contains more integral vectors than the number of possible values a q -generated polynomial with degree at most q can take on these vectors, even after identifying vectors up to permutations of the coordinates (Lemmas 1 and 2). This implies that the polynomial piece will take the same value on two distinct integral vectors whose coordinates are not identical up to permutations. When translating this result back to the world of GNNs, this means that the two embeddings of the root nodes of the trees corresponding to these two vectors will coincide. To conclude a separation between bounded and unbounded GNNs, we justify that the unbounded ones can separate these two vertices. This is based on the previous result (Theorem 2) stating that unbounded GNNs refine color refinement.

Our second result states that for activations that are not piecewise polynomial, a one neuron perceptron GNN can distinguish the root nodes of any pair of non-isomorphic trees of depth two. In particular, we prove this when the activation function is the exponential, the sigmoid or the hyperbolic sine, cosine or tangent functions. This is done by showing that the condition $\xi^2(s) = \xi^2(s)$ corresponds to a relation between the exponentials of the integers k_1, \dots, k_m and k'_1, \dots, k'_m . Applying the Lindemann-Weirstrass Theorem in transcendental number theory (Lemma 5 and Theorem 5) leads to the conclusion that k'_1, \dots, k'_m must be a permutation of k_1, \dots, k_m , showing that the trees are isomorphic.




FIGURE 1. $T[k_1, \dots, k_m]$

4. COLLISION WITH PIECEWISE POLYNOMIAL ACTIVATIONS

Lemma 1. *Let m , q and T be positive integers. For any natural number M , let F_M be the box $\{(k_1, \dots, k_m) \in \mathbb{Z}^m : \forall i \ 1 \leq k_i \leq M\}$ and let $\sigma_1, \dots, \sigma_q$ be q reals. Let P_1, \dots, P_T be T polynomials of total degree less than q , and q -generated by $\sigma_1, \dots, \sigma_q$. Then the number of values taken by the function $x \mapsto (P_1(x), \dots, P_T(x))$ on F_M is at most*

$$\left(2 \left(\max_{i \in [q], j \in [T], \alpha \in S} |\lambda_{\alpha, i}^j| \right) M^q \binom{m+q-1}{q} + 1 \right)^{qT}$$

where S is the union of the set of exponents of the P_j 's and the $\lambda_{\alpha, i}^j$ are the coefficients in the decomposition of their coefficients over the generators $\sigma_1, \dots, \sigma_q$.

Proof. Note that if every P_j is a polynomial with integer coefficients, i.e., each P_j is generated by 1, and has degree less than q , then the proof follows by considering the maximum and the minimum of the P_j 's over F_M , as each of them takes only integer values. To deal with the finitely generated case, we reduce it to the integer case. If S is the union of the set of exponents of P_1, \dots, P_T , by assumption there exist integers $(\lambda_{\alpha,i}^j)_{(\alpha,i) \in S \times \{1, \dots, q\}}$ such that

$$\begin{aligned} P_j &= \sum_{\alpha \in S} \gamma_{\alpha,j} X^\alpha = \sum_{\alpha \in S} \left(\sum_{i=1}^q \lambda_{\alpha,i}^j \sigma_i \right) X^\alpha \\ &= \sum_{i=1}^q \left(\sum_{\alpha \in S} \lambda_{\alpha,i}^j X^\alpha \right) \\ &= \sum_{i=1}^q \left(\underbrace{\sum_{\alpha \in S} \lambda_{\alpha,i}^j X^\alpha}_{:=f_{i,j}} \right) \sigma_i \end{aligned}$$

$f_{i,j}$ is a polynomial of degree less than q with integer coefficients. As we wish to upper-bound the number of values of $\phi : x \mapsto (P_1(x), \dots, P_T(x))$ on F_M , we consider

$$\begin{aligned} \Phi : F_M &\longrightarrow \mathbb{Z}^{qT} \\ x &\mapsto (f_{i,j}(x))_{i \in [q], j \in [T]} \end{aligned}$$

The number of values of Φ on F_M controls the number of values of ϕ on F_M as each of the coordinates of ϕ is a linear combination of the coordinates of Φ with fixed linear coefficients $\sigma_1, \dots, \sigma_q$. We use the fact that each coordinate of $\Phi(x)$ is a multivariate polynomial with *integer* coefficients:

$$\begin{aligned} |\phi(F_M)| &\leq |\Phi(F_M)| \leq \prod_{i \in [q], j \in [T]} (|\max_{x \in F_M} f_{i,j}(x) - \min_{x \in F_M} f_{i,j}(x)| + 1) \\ &\leq \prod_{i \in [q], j \in [T]} (2 \max_{x \in F_M} |f_{i,j}(x)| + 1) \\ &\leq \left(2 \left(\max_{i \in [q], j \in [T], \alpha \in S} |\lambda_{\alpha,i}^j| \right) M^q \binom{m+q-1}{q} + 1 \right)^{qT} \end{aligned}$$

where the last inequality follows from the fact that $\binom{m+q-1}{q}$ is an upper-bound on the number of monomials of $f_{i,j}$. \square

Lemma 2. *Let q be a positive integer, let I be a finite subset of \mathbb{N} , and let $(f_{t,m} : \mathbb{R}^m \rightarrow \mathbb{R})_{(t,m) \in I \times \mathbb{N}}$ be a double sequence of piecewise polynomial functions satisfying:*

- i) $\deg(f_{t,m}) \leq q$ for all $(t,m) \in I \times \mathbb{N}$ (bounded degree condition).
- ii) for any $m \in \mathbb{N}$ and $t \in I$, $f_{t,m}$ is q -generated.

Then, there exists $m \in \mathbb{N}$ and two integral vectors $(k_1, \dots, k_m) \in \mathbb{N}^m$ and $(k'_1, \dots, k'_m) \in \mathbb{N}^m$ that are not equal up to a permutation such that for any $t \in I$, $f_{t,m}(k_1, \dots, k_m) = f_{t,m}(k'_1, \dots, k'_m)$.

Proof. Let q be a positive integer, I be a finite subset of \mathbb{N} and $(f_{t,m} : \mathbb{R}^m \rightarrow \mathbb{R})_{(t,m) \in I \times \mathbb{N}}$ be a double sequence of piecewise polynomial satisfying the assumptions of the lemma. Let m be any natural number such that $m > q^2|I|$. For any positive integer M , let F_M be the box $\{(k_1, \dots, k_m) \in \mathbb{Z}^m : \forall i \ 1 \leq k_i \leq M\}$.

There are $\binom{M+m-1}{m}$ multisets of size m that can be formed from $\{1, \dots, M\}$. One can find that many elements of F_M that are not equal up to a permutation. Let r be an upperbound on the number of pieces of $f_{t,m}$ for every t , i.e. $r := \max\{\text{number of pieces of } f_{t,m} : t \in I\}$: note that r exists since I is finite, but may depend on m . Then, there exists a subset of F_M with at least $\frac{1}{r|I|} \binom{M+m-1}{m}$ integral vectors that are not equal up to a permutation where for every t , $f_{t,m}$ is a polynomial $P_{t,m}$ of degree at most q , and $P_{t,m}$ is q -generated. This is true as the collection of piecewise polynomial functions $(f_{i,m})_{i \in I}$ divide F_M in at most $r^{|I|}$ regions, where in each region, for every $i \in I$, $f_{i,m}$ is polynomial.

Due to Lemma 1, the pigeonhole principle gives us that all polynomials will be equal on some vectors (k_1, \dots, k_m) and (k'_1, \dots, k'_m) of F_M not equal up to a permutation as soon as:

$$(1) \quad \underbrace{\frac{1}{r^{|I|}} \binom{M+m-1}{m}}_{\substack{\text{number of multisets formed} \\ \text{from a region of } F_M \text{ where} \\ \text{each } f_{t,m} \text{ is polynomial}}} > \underbrace{\left(2 \max_{i \in [q], t \in I} \left(\max_{\alpha \in S_{m,t}} |\lambda_{\alpha,i}^{m,t}| \right) M^q \binom{m+q-1}{q} + 1 \right)^{q|I|}}_{\substack{\text{number of values } (P_{1,m}, P_{2,m}, \dots, P_{|I|,m}) \\ \text{can take at most on } F_M}}$$

where the $\lambda_{\alpha,i}^{m,t}$ are integers in the decomposition of the polynomials $f_{t,m}$ over generators $\sigma_1^m, \dots, \sigma_q^m \in \mathbb{R}$ for the monomial X^α .

Such a value of M can be found by noticing that $\binom{M+m-1}{m}$ is a polynomial of M of degree m whereas the right hand side is a polynomial of M of degree $q^2|I|$. Since we chose m to be greater than $q^2|I|$, there exists $M \in \mathbb{N}$ such that Equation (1) holds. Hence there exist k and k' whose coordinates are not equal up any permutation and such that $f_{i,m}(k_1, \dots, k_m) = f_{i,m}(k'_1, \dots, k'_m)$ for any $i \in I$. \square

Lemma 3. *Let q be a positive integer. Let $F : \mathbb{R}^{d_2} \rightarrow \mathbb{R}$ be a piecewise multivariate polynomial whose pieces are all q generated. Let $G : \mathbb{R}^{d_1} \rightarrow \mathbb{R}^{d_2}$ be a function whose coordinates are piecewise polynomial functions whose pieces are all q -generated.*

Then the pieces of the piecewise polynomial $F \circ G : \mathbb{R}^{d_1} \rightarrow \mathbb{R}$ are all \tilde{q} -generated, where \tilde{q} depends only on q and the degree of the pieces of F and G .

Proof. We first make the following observation. Let $A_1, \dots, A_{r_1} \subseteq \mathbb{R}^{d_2}$ be polynomial regions of F and $B_1, \dots, B_{r_2} \subseteq \mathbb{R}^{d_1}$ be polynomial regions of G that union to \mathbb{R}^{d_2} and \mathbb{R}^{d_1} respectively (such always exist, cf. Definition 1). For each $i \in [r_1]$, and $j \in [r_2]$, let $Z_{ij} := (B_i \cap G^{-1}(A_j))$. Then:

- G and $F \circ G$ are polynomial on each Z_{ij} , as $Z_{ij} \subseteq B_i$, and $G(Z_{ij}) \subseteq A_j$.
- The Z_{ij} union to \mathbb{R}^{d_1} , as $\bigcup_{i \in [r_2], j \in [r_1]} Z_{ij} = \bigcup_{j \in [r_1]} \bigcup_{i \in [r_2]} B_i \cap G^{-1}(A_j) =$

$$\bigcup_{j \in [r_1]} \left(G^{-1}(A_j) \cap \underbrace{\left(\bigcup_{i \in [r_2]} B_i \right)}_{\mathbb{R}^{d_1}} \right) = \bigcup_{j \in [r_1]} G^{-1}(A_j) = \mathbb{R}^{d_1}. \quad \text{The last}$$

equality follows from $\bigcup_{i \in [r_1]} A_i = \mathbb{R}^{d_2} \implies \bigcup_{i \in [r_1]} G^{-1}(A_i) = \mathbb{R}^{d_1}$.

Hence, a polynomial piece of $F \circ G$ is of the form $F_i \circ G_j$ where F_1, \dots, F_{r_1} are the polynomial pieces of F , and G_1, \dots, G_{r_2} are the polynomial pieces of G .

If $F_i = \sum_{\alpha \in S_{F_i}} \gamma_\alpha X^\alpha$ and $G_j = \sum_{\beta \in S_{G_j}} \nu_\beta X^\beta$ with γ_α and ν_β integers then

$$\begin{aligned} (F_i \circ G_j)(X) &= \sum_{\alpha \in S_{F_i}} \gamma_\alpha \left(\sum_{\beta \in S_{G_j}} \nu_\beta X^\beta \right)^\alpha \\ &= \sum_{\alpha \in S_{F_i}} \left(\sum_{k=1}^q \lambda_{k,\alpha} \sigma_k \right) \left(\sum_{\beta \in S_{G_j}} \left(\sum_{k=1}^q \eta_{k,\alpha} \sigma_k \right) X^\beta \right)^\alpha \end{aligned}$$

After expansion, the polynomial $(F_i \circ G_j)$ has coefficients that are linear combinations of multivariate monomials of $\sigma_1, \dots, \sigma_q$ (of total degree at most the degree of $F_j + 1$ in this case) with integer coefficients, as the λ and η coefficients are supposed to be integers. This shows that $F_i \circ G_j$ is \tilde{q} -generated, where \tilde{q} depends only on p , the degree of F_i and G_j . \square

Lemma 4. *For positive integers m, k_1, \dots, k_m , let $T[k_1, \dots, k_m]$ designate the rooted tree illustrated in Figure 1. For any vertex $v \in V(T[k_1, \dots, k_m])$, Let $\xi^t(T[k_1, \dots, k_m], v)$ be the embedding of the vertex $v \in V$ obtained via a GNN with piecewise activation functions after t iterations (where $\xi^0(u) = 1$ for any vertex u of $T[k_1, \dots, k_m]$). Then, for any iteration t , there exists an integer q such that for any integer m , and any vertex $v \in V$, there exists a symmetric multivariate piecewise polynomial function F_m such that*

- $\xi^t(T[k_1, \dots, k_m], v) = F_m(k_1, \dots, k_m)$.
- $\deg(F_m) \leq q$
- each piece of F_m is q -generated

Proof. We first prove all properties of $\xi^t(T[k_1, \dots, k_m], v)$ by induction on t .

Base case: for $t = 0$ this is trivial since all vertices are initialised with the constant polynomial 1, whose degree does not depend on m , and is finitely generated.

Induction step: Suppose the property is true at iteration t , i.e there exists an integer q_t such that for each vertex w , and for every integer m , $\xi^t(T[k_1, \dots, k_m], w)$ is a multivariate polynomial of the k_i 's whose degree is upperbounded by q_t , and is q_t generated. For every vertex v of $T[k_1, \dots, k_m]$

$$\xi^{t+1}(T[k_1, \dots, k_m], v) = \phi(\xi^t(T[k_1, \dots, k_m], v), \sum_{w \in N(v)} \xi^t(T[k_1, \dots, k_m], w))$$

where ϕ is a piecewise bivariate polynomial (as a Neural Network $\mathbb{R}^2 \rightarrow \mathbb{R}$ with piecewise polynomial activation). By composition $\xi^{t+1}(T[k_1, \dots, k_m], v)$ is a piecewise multivariate polynomial of k_1, \dots, k_m such that for every integer m , $\deg(\xi^{t+1}(T[k_1, \dots, k_m], v)) \leq q_t \cdot \deg(\phi)$. Furthermore, since for every m and every vertex u , $\xi^t(T[k_1, \dots, k_m], u)$ is supposed to be q_t generated, then using the above update rule, Lemma 3 gives us that by composition, $\xi^{t+1}(T[k_1, \dots, k_m], v)$ is \tilde{q}_t generated, where \tilde{q}_t depends only on the degree of ϕ and q_t .

Setting $q_{t+1} := \max(q_t \cdot \deg(\phi), \tilde{q}_t)$ gives us the desired properties (for every vertex v and every integer m , $\xi^{t+1}(T[k_1, \dots, k_m], v)$ has degree at most q_{t+1} and is q_{t+1} generated), and ends the induction on t . \square

Proof of Theorem 3. We already know [Gro21] that color refinement refines any recurrent GNN (even with an architecture of unbounded size). We prove the existence of pairs of graphs that can be separated by the color refinement algorithm, but cannot be separated by a recurrent GNN of fixed (but arbitrary) size. We use $T[k_1, \dots, k_m]$ to refer to the tree illustrated in Figure 1. This tree has depth two, a root node s , and contains m nodes at depth one. Each vertex i at depth 1 has exactly $k_i - 1$ “children” at depth two (and therefore k_i neighbors, where k_i is a positive integer). In the following, all vertices have color label 1.

Claim: Let $T[k_1, \dots, k_m]$ and $T'[k'_1, \dots, k'_m]$ be two rooted trees given by Figure 1. If the k_i 's and k'_i 's are not equal up to a permutation, the color refinement distinguishes s and s' after two iterations, i.e. $\text{cr}^2(s) \neq \text{cr}^2(s')$.

Proof of claim. Simply note that

$$\text{cr}^2(s) = (\text{cr}^1(s), \{\{\text{cr}^1(x_1), \dots, \text{cr}^1(x_m)\}\})$$

$$\text{where } \text{cr}^1(s) = (\underbrace{1}_{\text{cr}^0(s)}, \underbrace{\{\{1, \dots, 1\}\}}_{m \text{ times}})$$

$$\text{and } \forall i \in \{1, \dots, m\} \quad \text{cr}^1(x_i) = (\underbrace{1}_{\text{cr}^0(x_i)}, \underbrace{\{\{1, \dots, 1\}\}}_{k_i \text{ times}})$$

hence $\text{cr}^2(s)$ is uniquely determined by the multiset $\{\{k_1, \dots, k_m\}\}$. \square

Let $T > 0$ be a positive integer, and for $0 \leq t \leq T$, let $f_{t,m}(k_1, \dots, k_m) := \xi^t(T[k_1, \dots, k_m], s)$ be the value returned by a GNN with piecewise polynomial activation after t iterations (note that the embeddings are one-dimensional because only one color is used). Using Lemma 4, there exists an integer q such that the double sequence $(f_{t,m})_{t \in \{0, \dots, T\}, m \in \mathbb{N}}$ of piecewise multivariate polynomials has degree at most q and such that every $f_{t,m}$ is q -generated. Lemma 2 with $I = \{0, \dots, T\}$ tells us that there exists $m \in \mathbb{N}$, and two vectors $k \in \mathbb{N}^m$ and $k' \in \mathbb{N}^m$ whose coordinates are not equal up to permutations, such that for any $t \in \{0, \dots, T\}$, $f_{t,m}(k_1, \dots, k_m) = f_{t,m}(k'_1, \dots, k'_m)$. \square

Remark 3. Note that in Theorem 3, depth two is minimal: for any pair of non isomorphic rooted trees of depth one, any GNN with one neuron perceptron, an injective activation function, weights set to one, and zero bias can distinguish their root vertex in one iteration. Indeed, in that case, $\xi^1(s) = \sigma(1 + \deg(s))$ if the GNN is recurrent with a combine function given by $\phi : \mathbb{R}^2 \rightarrow \mathbb{R}, (x_1, x_2) \mapsto \sigma(x_1 + x_2)$. Hence, $\xi^1(s) \neq \xi^1(s')$ as soon as σ is injective and s and s' have distinct degree.

5. ACTIVATIONS THAT ARE NOT PIECEWISE POLYNOMIAL

In this Section we present a proof of Theorem 4. We prove that for any pair of non isomorphic rooted trees of depth two, i.e. trees of the form $T[k_1, \dots, k_m]$ and $T'[k'_1, \dots, k'_m]$ (here the k_i 's and k'_i 's are all greater than or equal to 1, cf. Figure 1) can be distinguished by a bounded GNN with any of the following activation functions: exponential, sigmoid, or a hyperbolic sine, cosine or tangent function. Consider the following 1-neuron perceptron ϕ with activation function $\sigma, \phi : \mathbb{R}^2 \rightarrow$

\mathbb{R} , $\phi(x_1, x_2) = \sigma(x_1 + x_2)$. Then it is easy to see that:

$$\begin{aligned} \forall v \in V(T[k_1, \dots, k_m]) \quad \xi^1(v) &= \sigma(\xi^0(v) + \sum_{w \in N(v)} \xi^0(w)) = \sigma(1 + \deg(v)) \\ \xi^2(v) &= \sigma(\sigma(1 + \deg(v)) + \sum_{w \in N(v)} \sigma(1 + \deg(w))) \end{aligned}$$

In particular $\xi^2(s) = \sigma(\sigma(1+m) + \sum_{i=1}^m \sigma(k_i+1))$. Now suppose σ is either injective on \mathbb{R} , or nonnegative and injective on \mathbb{R}^+ (this is the case for the exponential, the sigmoid, the hyperbolic tan, and the hyperbolic cosine and sine), s and s' are vertices of two trees with potentially different number of leaves m and n , then

$$(2) \quad \xi^2(s) = \xi^2(s') \iff \sum_{i=0}^m \sigma(k_i + 1) = \sum_{i=0}^n \sigma(k'_i + 1)$$

where $k_0 := 1 + m$ and $k'_0 := 1 + n$. The goal of the remainder of this section is to prove that the right hand side equality of (2) implies $m = n$ and k_i 's are the same as k'_i 's, up to a permutation, for the activation functions σ of Theorem 4.

Theorem 5 (Lindemann-Weierstrass Theorem, 1885). *If $\alpha_1, \dots, \alpha_n$ are distinct algebraic numbers, then the exponentials $e^{\alpha_1}, \dots, e^{\alpha_n}$ are linearly independent over the algebraic numbers.*

Lemma 5. *Let n and m be positive integers, and $\alpha_1, \dots, \alpha_n$ and $\alpha'_1, \dots, \alpha'_m$ be algebraic numbers. Then $\sum_{i=1}^n e^{\alpha_i} = \sum_{i=1}^m e^{\alpha'_i}$ if and only if $m = n$ and the α_i 's and α'_i 's are equal up to a permutation.*

Proof. (\Leftarrow) is clear. For (\Rightarrow), by contradiction suppose that the α_i 's and α'_i 's are not equal up to a permutation. First, if the α_i 's (resp. α'_i 's) are not distinct one can group them by their number of occurrences in both sums. Then, we would have a linear dependence with integer coefficients of exponentials of integers. This contradicts Theorem 5 (Linderman-Weirstrass). \square

Proof of Theorem 4. Without loss of generality, suppose the k_i 's and k'_i 's are ordered in increasing order. For ease of notation, let α and α' be the vectors defined as $\alpha_i = k_i + 1$ for all $i \in \{1, \dots, m\}$ and $\alpha'_i = k'_i + 1$ for all $i \in \{1, \dots, n\}$. We will now prove that (2) implies $\alpha = \alpha'$ in each case.

- $\sigma \in \{\text{sigmoid}, \text{tanh}\}$. In the case of the sigmoid, (2) yields the following equation after multiplication by the product of the denominators:

$$\left(\sum_{i=1}^m e^{\alpha_i} \left(\prod_{j \neq i}^m (1 + e^{\alpha_j}) \right) \right) \prod_{i=1}^n (1 + e^{\alpha'_i}) = \left(\sum_{i=1}^n e^{\alpha'_i} \left(\prod_{j \neq i}^n (1 + e^{\alpha'_j}) \right) \right) \prod_{i=1}^m (1 + e^{\alpha_i})$$

After developing and grouping each hand side into linear combinations of exponentials we obtain an equation of the form:

$$(3) \quad \sum_{\substack{S \subseteq \{1, \dots, m\} \\ T \subseteq \{1, \dots, n\}}} \gamma_{S,T} \exp(\alpha_S + \alpha'_T) = \sum_{\substack{S \subseteq \{1, \dots, m\} \\ T \subseteq \{1, \dots, n\}}} \gamma_{S,T} \exp(\alpha'_S + \alpha_T)$$

where for $S \subseteq \{1, \dots, m\}$, $\alpha_S := \sum_{i \in S} \alpha_i$ (resp. for $T \subseteq \{1, \dots, n\}$, $\alpha'_T := \sum_{i \in T} \alpha'_i$). All $\gamma_{S,T}$ are integers, hence algebraic and $\gamma_{\emptyset, T} = 0$ for all subsets $T \subseteq \{1, \dots, n\}$.

We will prove by strong induction on $\max(m, n)$ that $\sum_{i=1}^m \sigma(\alpha_i) = \sum_{i=1}^n \sigma(\alpha'_i) \implies m = n$ and $\alpha = \alpha'$.

Base case: If $\max(m, n) = 1$ and $\sum_{i=1}^m \sigma(\alpha_i) = \sum_{i=1}^n \sigma(\alpha_1)$, then, either $m = n = 0$, or $m = n = 1$. In the first case, this is vacuously true. In the second case, we have that $\sigma(\alpha_1) = \sigma(\alpha'_1)$, and then $\alpha_1 = \alpha'_1$ follows from the injectivity of the sigmoid.

Induction step: We suppose that for some given positive integer p , and any nonnegative integers $\alpha_1, \dots, \alpha_m$ and $\alpha'_1, \dots, \alpha'_n$, such that $\max(m, n) \leq p$, then $\sum_{i=1}^m \sigma(\alpha_i) = \sum_{i=1}^n \sigma(\alpha'_i) \implies m = n$ and $\alpha = \alpha'$.

Let α and α' be vector of integers of size m and n such that $\max(m, n) = p + 1$, and $\sum_{i=1}^m \sigma(\alpha_i) = \sum_{i=1}^n \sigma(\alpha'_i)$. We saw that (3) can be derived from this equality, where $\gamma_{S,T}$ are algebraic numbers satisfying $\gamma_{\emptyset, T} = 0$ for all $T \subseteq \{1, \dots, n\}$. Moreover, the coordinates of α and α' are ordered, hence the smallest term on the left hand side is $\exp(\alpha_1)$ and the smallest term on the right hand side is $\exp(\alpha'_1)$. Using Lemma 5, this implies that $\alpha_1 = \alpha'_1$. Therefore, $\sigma(\alpha_1) = \sigma(\alpha'_1)$. In turn, this implies $\sum_{i=2}^m \sigma(\alpha_i) = \sum_{i=2}^n \sigma(\alpha'_i)$. We can apply the induction hypothesis on the vectors $(\alpha_2, \dots, \alpha_m)$ and $(\alpha'_2, \dots, \alpha'_n)$ which both have size $\leq p$. Hence, we obtain that $m - 1 = n - 1$ and $(\alpha_2, \dots, \alpha_m) = (\alpha'_2, \dots, \alpha'_m)$. This in turn proves that $m = n$, and $\alpha = \alpha'$, which ends the induction.

If $\sigma = \tanh = \frac{\exp(2\cdot) + 1}{\exp(2\cdot) - 1}$. After multiplication by the product of the denominators, (2) yields:

$$\left(\sum_{i=1}^n (e^{2\alpha_i} - 1) \prod_{j=1, j \neq i}^n (e^{2\alpha_j} + 1) \right) \prod_{j=1}^m (1 + e^{2\alpha'_j}) = \left(\sum_{i=1}^m (e^{2\alpha'_i} - 1) \prod_{j=1, j \neq i}^m (e^{2\alpha'_j} + 1) \right) \prod_{j=1}^n (1 + e^{2\alpha_j})$$

After developing into a linear combination of exponentials on each side, the arguments containing α_T with $T \neq \emptyset$ on the left hand side and α'_T with $T \neq \emptyset$ on the right hand side have positive algebraic coefficients. There are also arguments of the form α'_T on the left hand side and α_T on the right hand side (in other words, $\gamma_{\emptyset, T} \neq 0$, unlike the sigmoid case). However, note that the coefficients corresponding to these terms are (algebraic and) negative. Hence, as a consequence of Lemma 5, the arguments with negative coefficients in front of the exponentials must match up on each side, and we are left with an equation similar to (3) (the arguments have a factor 2), where again $\gamma_{\emptyset, T} = 0$. We can apply the same reasoning by induction as for the sigmoid case, to prove that $\alpha = \alpha'$.

- $\sigma \in \{\sinh, \cosh\}$. If $\sigma = \cosh$, then (2) yields:

$$\left(\sum_{j=1}^n \exp(\alpha_j) - \sum_{j=1}^m \exp(\alpha'_j) \right) + \left(\sum_{j=1}^n \exp(-\alpha_j) - \sum_{j=1}^m \exp(-\alpha'_j) \right) = 0$$

Due to Lemma 5, this can only happen if $m = n$ and for all $j \in \{1, \dots, n\}$, $\alpha_j = \alpha'_j$, because α_j, α'_j are algebraic for any $j \in \{1, \dots, n\}$, and the α_j 's and α'_j 's are ordered and positive. We conclude that $\alpha = \alpha'$. The case $\sigma \in \{\sinh\}$ can be treated similarly. \square

6. ACKNOWLEDGEMENTS

The authors gratefully acknowledge support from Air Force Office of Scientific Research (AFOSR) grant FA95502010341 and National Science Foundation (NSF) grant CCF2006587. The authors are also very grateful to insightful comments from reviewers that helped to improve the paper significantly. In particular, we learnt of the reference [ACI⁺22] from one of the reviewers. We also thank Eran Rosenbluth for pointing out an error in our first proof of Theorem 3.

REFERENCES

- [ACI⁺22] Anders Aamand, Justin Chen, Piotr Indyk, Shyam Narayanan, Ronitt Rubinfeld, Nicholas Schiefer, Sandeep Silwal, and Tal Wagner. Exponentially improving the complexity of simulating the weisfeiler-lehman test with graph neural networks. *Advances in Neural Information Processing Systems*, 35:27333–27346, 2022.
- [AGA⁺24] Tal Amir, Steven Gortler, Ilai Avni, Ravina Ravina, and Nadav Dym. Neural injective functions for multisets, measures and graphs via a finite witness theorem. *Advances in Neural Information Processing Systems*, 36, 2024.
- [AL21] Waïss Azizian and Marc Lelarge. Expressive power of invariant and equivariant graph neural networks. In *ICLR 2021-International Conference on Learning Representations*, 2021.
- [BBL⁺17] Michael M Bronstein, Joan Bruna, Yann LeCun, Arthur Szlam, and Pierre Vandergheynst. Geometric deep learning: going beyond euclidean data. *IEEE Signal Processing Magazine*, 34(4):18–42, 2017.
- [BKM⁺20] Pablo Barceló, Egor V Kostylev, Mikael Monet, Jorge Pérez, Juan Reutter, and Juan Pablo Silva. The logical expressiveness of graph neural networks. In *8th International Conference on Learning Representations (ICLR 2020)*, 2020.
- [BKR24] César Bravo, Alexander Kozachinskiy, and Cristóbal Rojas. On dimensionality of feature vectors in mpnns. *arXiv preprint arXiv:2402.03966*, 2024.
- [BPL⁺16] Peter Battaglia, Razvan Pascanu, Matthew Lai, Danilo Jimenez Rezende, et al. Interaction networks for learning about objects, relations and physics. *Advances in neural information processing systems*, 29, 2016.
- [CCK⁺21] Quentin Cappart, Didier Chételat, Elias Khalil, Andrea Lodi, Christopher Morris, and Petar Veličković. Combinatorial optimization and reasoning with graph neural networks. *arXiv preprint arXiv:2102.09544*, 2021.
- [CFI92] Jin-Yi Cai, Martin Fürer, and Neil Immerman. An optimal lower bound on the number of variables for graph identifications. *Combinatorica*, 12(4):389–410, 1992.
- [DBV16] Michaël Defferrard, Xavier Bresson, and Pierre Vandergheynst. Convolutional neural networks on graphs with fast localized spectral filtering. *Advances in neural information processing systems*, 29, 2016.
- [DMI⁺15] David K Duvenaud, Dougal Maclaurin, Jorge Iparraguirre, Rafael Bombarell, Timothy Hirzel, Alán Aspuru-Guzik, and Ryan P Adams. Convolutional networks on graphs for learning molecular fingerprints. *Advances in neural information processing systems*, 28, 2015.
- [DZC⁺19] Ming Ding, Chang Zhou, Qibin Chen, Hongxia Yang, and Jie Tang. Cognitive graph for multi-hop reading comprehension at scale. *arXiv preprint arXiv:1905.05460*, 2019.
- [GR22] Floris Geerts and Juan L Reutter. Expressiveness and approximation properties of graph neural networks. *arXiv preprint arXiv:2204.04661*, 2022.
- [Gro17] Martin Grohe. *Descriptive complexity, canonisation, and definable graph structure theory*, volume 47. Cambridge University Press, 2017.
- [Gro21] Martin Grohe. The logic of graph neural networks. In *2021 36th Annual ACM/IEEE Symposium on Logic in Computer Science (LICS)*, pages 1–17. IEEE, 2021.
- [Ham20] William L Hamilton. Graph representation learning. *Synthesis Lectures on Artificial Intelligence and Machine Learning*, 14(3):1–159, 2020.
- [HV21] Ningyuan Teresa Huang and Soledad Villar. A short tutorial on the weisfeiler-lehman test and its variants. In *ICASSP 2021-2021 IEEE International Conference on Acoustics, Speech and Signal Processing (ICASSP)*, pages 8533–8537. IEEE, 2021.
- [Jeg22] Stefanie Jegelka. Theory of graph neural networks: Representation and learning. *arXiv preprint arXiv:2204.07697*, 2022.
- [KDZ⁺17] Elias Khalil, Hanjun Dai, Yuyu Zhang, Bistra Dilkina, and Le Song. Learning combinatorial optimization algorithms over graphs. *Advances in neural information processing systems*, 30, 2017.
- [KP19] Nicolas Keriven and Gabriel Peyré. Universal invariant and equivariant graph neural networks. *Advances in Neural Information Processing Systems*, 32, 2019.
- [MFSL19] Haggai Maron, Ethan Fetaya, Nimrod Segol, and Yaron Lipman. On the universality of invariant networks. In *International conference on machine learning*, pages 4363–4371. PMLR, 2019.

- [MRF⁺19] Christopher Morris, Martin Ritzert, Matthias Fey, William L Hamilton, Jan Eric Lenssen, Gaurav Rattan, and Martin Grohe. Weisfeiler and leman go neural: Higher-order graph neural networks. In *Proceedings of the AAAI conference on artificial intelligence*, volume 33, pages 4602–4609, 2019.
- [SGGP⁺20] Alvaro Sanchez-Gonzalez, Jonathan Godwin, Tobias Pfaff, Rex Ying, Jure Leskovec, and Peter Battaglia. Learning to simulate complex physics with graph networks. In *International conference on machine learning*, pages 8459–8468. PMLR, 2020.
- [SGT⁺08] Franco Scarselli, Marco Gori, Ah Chung Tsoi, Markus Hagenbuchner, and Gabriele Monfardini. The graph neural network model. *IEEE transactions on neural networks*, 20(1):61–80, 2008.
- [SYS⁺20] Jonathan M Stokes, Kevin Yang, Kyle Swanson, Wengong Jin, Andres Cubillos-Ruiz, Nina M Donghia, Craig R MacNair, Shawn French, Lindsey A Carfrae, Zohar Bloom-Ackermann, et al. A deep learning approach to antibiotic discovery. *Cell*, 180(4):688–702, 2020.
- [XHLJ18] Keyulu Xu, Weihua Hu, Jure Leskovec, and Stefanie Jegelka. How powerful are graph neural networks? *arXiv preprint arXiv:1810.00826*, 2018.
- [ZAL18] Marinka Zitnik, Monica Agrawal, and Jure Leskovec. Modeling polypharmacy side effects with graph convolutional networks. *Bioinformatics*, 34(13):i457–i466, 2018.
- [ZCH⁺20] Jie Zhou, Ganqu Cui, Shengding Hu, Zhengyan Zhang, Cheng Yang, Zhiyuan Liu, Lifeng Wang, Changcheng Li, and Maosong Sun. Graph neural networks: A review of methods and applications. *AI open*, 1:57–81, 2020.