INTERPOLATION OF FAT POINTS ON K3 AND ABELIAN SURFACES

ADRIAN ZAHARIUC

ABSTRACT. We prove that any number of general fat points of any multiplicities impose the expected number of conditions on a linear system on a smooth projective surface, in several cases including primitive linear systems on very general K3 and abelian surfaces, 'Du Val' linear systems on blowups of \mathbb{P}^2 at 9 very general points, and certain linear systems on some ruled surfaces over elliptic curves. This is done by answering a question of the author about the case of only one fat point on a certain ruled surface, which follows from a circle of results due to Treibich–Verdier, Segal–Wilson, and others.

1. INTRODUCTION

The Segre–Harbourne–Gimigliano–Hirschowitz (SHGH) Conjecture [\[19,](#page-8-0) [12,](#page-7-0) [11,](#page-7-1) [14\]](#page-8-1) is a well-known open problem in algebraic geometry. Please see [\[2\]](#page-7-2) for a survey of the history and various approaches to the conjecture. Questions similar to the SHGH Conjecture on projective surfaces other than \mathbb{P}^2 have been studied, for instance, in [\[26,](#page-8-2) [27,](#page-8-3) [7,](#page-7-3) [15,](#page-8-4) [16,](#page-8-5) [17,](#page-8-6) [28\]](#page-8-7). Such questions also relate to Seshadri constants, though in general they require even more specific (non-asymptotic) information.

This note is a follow-up to [\[28\]](#page-8-7). We will show that, over the complex numbers, [\[28,](#page-8-7) Conjecture 2.5] can be deduced from certain facts (of a rather analytic nature), thereby establishing the results proved conditionally on this claim in [\[28\]](#page-8-7). First, we state these results clearly and extend slightly the range of situations covered.

Theorem 1.1. Let S be a smooth complex projective surface and \mathcal{L} an invertible \mathcal{O}_S -module, such that (S, \mathcal{L}) is one of the following:

- (1) a very general primitively polarized K3 surface of any degree;
- (2) a very general primitively polarized abelian surface of any degree (i.e. a very general $(1, d)$ -polarized abelian surface);
- (3) the blowup of \mathbb{P}^2 at 9 very general points, and a Du Val linear system, that is, $c_1(\mathcal{L}) = -kK_S + E$, where $E \simeq \mathbb{P}^1 \subset S$ is a (-1) -curve and k is a positive integer;
- (4) $S = \mathbb{P}{\mathcal{E}}$, where $\mathcal E$ is a rank 2 locally free sheaf on a smooth projective curve E of genus 1, such that either
	- (a) $\mathcal{E} = \mathcal{O}_E \oplus \mathcal{J}$ with $\mathcal{J} \in \text{Pic}^0(E)$ very general; or

recherches en sciences naturelles et en génie du Canada (CRSNG), RGPIN-2020-05497.

CRSNG We acknowledge the support of the Natural Sciences and Engineering Research Council of Canada (NSERC), RGPIN-2020-05497. Cette recherche a été financée par le Conseil de

²⁰²⁰ Mathematics Subject Classification. 14C20, 14H42, 14J26, 14J27, 14J28.

Key words and phrases. SHGH Conjecture, tangential cover, K3 surface, abelian surface, ruled surface, Du Val curve.

(b) $\mathcal E$ is the unique nontrivial extension of $\mathcal O_E$ by $\mathcal O_E$,

and $\mathcal L$ is any ample line bundle with intersection number 1 with the section of S corresponding to the natural surjection $\mathcal{E} \rightarrow \mathcal{O}_E$ (in either case).

Let $p_1, \ldots, p_n \in S$ be n general points, m_1, \ldots, m_n positive integers, and

(1)
$$
|\mathcal{L}(m_1,\ldots,m_n)| = \mathbb{P}H^0(S,\mathcal{L}\otimes \mathcal{I}_{p_1,S}^{m_1}\otimes\cdots\otimes \mathcal{I}_{p_n,S}^{m_n}) \subset |\mathcal{L}|
$$

the linear system of divisors in $|\mathcal{L}|$ which have multiplicity at least m_i at p_i , for all $i=1,2\ldots,n$. Then

$$
\dim |\mathcal{L}(m_1,\ldots,m_n)| = \max \left\{-1, \dim |\mathcal{L}| - \sum_{i=1}^n \frac{m_i(m_i+1)}{2}\right\},\,
$$

with the convention that empty linear systems have dimension -1 .

To clarify, in cases [1](#page-0-0) and [2,](#page-0-1) $|\mathcal{L}|$ is the *primitive* linear system. For K3 surfaces, we confirm the conjectures of De Volder and Laface (Conjectures 2.1 and 2.3 in [\[27\]](#page-8-3)) in the primitive case $(d = 1$ with the notation in [27]), please see Remark [3.4.](#page-7-4) Case [3](#page-0-2) can equally well be stated in terms of \mathbb{P}^2 , so we also obtain some cases of the SHGH Conjecture. Cases [2](#page-0-1) and [4a](#page-0-3) were not considered explicitly in [\[28\]](#page-8-7), but they can be dealt with using the same idea, and seem natural enough to include. We have also removed the generality assumption on E in [4b.](#page-1-0) To the author's knowledge, Theorem [1.1](#page-0-4) is currently the only SHGH-type result with no restrictions on the number and weights of fat points.

Using the arguments in [\[28\]](#page-8-7), Theorem [1.1](#page-0-4) boils down to case [4b](#page-1-0) of Theorem [1.1](#page-0-4) and $n = 1$, or specifically, the characteristic 0 part of [\[28,](#page-8-7) Conjecture 2.5]. The reduction is discussed in §[3.](#page-4-0)

The conjecture in [\[28\]](#page-8-7) is stated as Proposition [2.4](#page-2-0) (in the form which removes the generality assumption on the j-invariant), and $\S2$ $\S2$ is devoted to proving it. It turns out that the claim can be deduced from facts about tangential covers of elliptic curves, that were studied by Treibich, Verdier, and other authors [\[21,](#page-8-8) [22,](#page-8-9) [23,](#page-8-10) [24,](#page-8-11) [25,](#page-8-12) [20,](#page-8-13) [9\]](#page-7-5). From the point of view of SHGH-type problems, the situation is quite unexpected, since the key fact comes from analytic work of Segal and Wilson [\[18\]](#page-8-14), and it seems that no algebraic proofs are currently available.

Acknowledgements. I would like to thank Xi Chen, Brian Osserman, Edoardo Sernesi, and Armando Treibich for useful discussions. I am especially grateful to Sernesi and Treibich for invaluable help with navigating the literature.

2. Proof of the conjecture from [\[28\]](#page-8-7)

2.1. Finiteness of θ vanishing in the KP direction. Let C be an integral (possibly singular) complex projective curve of arithmetic genus $g \geq 1$, and $p \in C$ a smooth point. There is a canonical identification of the tangent line T_pC to C at p with $\mathcal{O}_C(p) \otimes \mathcal{O}_p$. The exact sequence

$$
0 \longrightarrow \mathcal{O}_C \longrightarrow \mathcal{O}_C(p) \longrightarrow \mathcal{O}_C(p) \otimes \mathcal{O}_p \longrightarrow 0
$$

gives a natural embedding $T_p C \hookrightarrow H^1(\mathcal{O}_C)$ as the first coboundary map in the associated long exact sequence, whose image thus coincides with the kernel of the map $H^1(\mathcal{O}_C) \to H^1(\mathcal{O}_C(p))$. Since $\mathcal{T}_{\text{Pic}(C)} \cong H^1(C, \mathcal{O}_C) \otimes \mathcal{O}_{\text{Pic}(C)}$, the point p induces a natural direction at any point in $Pic(C)$, which is sometimes called the Kadomtsev-Petviashvili (KP) direction.

The essential ingredient in our arguments is the following.

Theorem 2.1. Let $\Delta \hookrightarrow Pic^{g-1}(C)$ be a holomorphic map from an open disk, such that the tangent line to any point of Δ coincides with the KP direction $T_pC \subset$ $H^1(\mathcal{O}_C)$. Then there exists $\mathcal{L} \in \Delta$ such that $H^0(C, \mathcal{L}) = 0$.

Proof. Follows from [\[18,](#page-8-14) Proposition 8.6] and the 'Krichever dictionary', as stated and explained in [\[22,](#page-8-9) §3.4]. Indeed, after compactifying the Jacobian in order to have the same setup, the first two lines on page 45 of [\[22\]](#page-8-9) imply our claim, since we are claiming precisely that Δ is not contained in the theta divisor. Note also that Proposition 3.7 on the same page provides the Segal–Wilson formula for the vanishing/contact order.

Remark 2.2. We will only require the case when C has at worst planar singularities, in which the theory is somewhat simpler [\[18,](#page-8-14) p. 38]. In fact, one can even make a certain plausible conjecture later, which would ensure that we only invoke Theorem [2.1](#page-2-1) for C smooth. When C is smooth, Theorem [2.1](#page-2-1) also follows from results of Fay [\[10\]](#page-7-6). However, it is stated in [\[1,](#page-7-7) footnote to p. 326] that a direct geometric proof is not known even this case.

2.2. Conjecture 2.5 in [\[28\]](#page-8-7) in characteristic 0. Let E be a smooth complex projective curve of genus 1, $\rho : S = \mathbb{P} \mathcal{V} \to E$ the ruled surface over E, where \mathcal{V} is the unique rank 2 vector bundle that fits in a non-split short exact sequence

$$
0 \longrightarrow \mathcal{O}_E \longrightarrow \mathcal{V} \longrightarrow \mathcal{O}_E \longrightarrow 0,
$$

 $E_{\infty} \subset S$ the section corresponding to $\mathcal{V} \to \mathcal{O}_E \to 0$, $q \in E$, and $F_s = \rho^{-1}(s)$, for any $s \in E$. There is a simple way to characterize curves which occur in $|F_q + kE_{\infty}|$ due to Treibich and Verdier $[25]$: the projection map from the curve to E must be so-called tangential. We state the result in a form closer to [\[29,](#page-8-15) Lemma 3.3] (the author was unfortunately unaware of the literature when [\[29\]](#page-8-15) was written).

Lemma 2.3 (essentially [\[25,](#page-8-12) Corollaire 3.10]). Let C a projective integral curve, and $f: C \to S$ a morphism such that $f^{-1}(E_{\infty}) = \{q\}$ scheme-theoretically, and q is a nonsingular point of C. Then the composition

$$
H^1(E, \mathcal{O}_E) \xrightarrow{(\rho f)^*} H^1(C, \mathcal{O}_C) \longrightarrow H^1(C, \mathcal{O}_C(q))
$$

is equal to 0.

Proof. As in the proof of [\[29,](#page-8-15) Lemma 3.3], the map $H^1(S, \mathcal{O}_S) \to H^1(S, \mathcal{O}_S(E_{\infty}))$ induced by the inclusion $\mathcal{O}_S \subset \mathcal{O}_S(E_\infty)$ is equal to 0. The commutativity of

$$
H^{1}(E, \mathcal{O}_{E}) \xrightarrow{\rho^{*}} H^{1}(S, \mathcal{O}_{S}) \xrightarrow{f^{*}} H^{1}(C, \mathcal{O}_{C})
$$

$$
\downarrow 0 \qquad \qquad \downarrow
$$

$$
H^{1}(S, \mathcal{O}_{S}(E_{\infty})) \longrightarrow H^{1}(C, \mathcal{O}_{C}(q))
$$

completes the proof. \Box

Proposition 2.4 ([\[28,](#page-8-7) Conjecture 2.5]). Let $x \in S$ be a general (closed) point and m a positive integer. Let k be the minimum positive integer for which the linear system $|F_q + kE_\infty|$ contains a curve of multiplicity at least m at x. Then

$$
k = \frac{m(m+1)}{2}
$$

.

Here is a casual summary of the proof. Instead of having q fixed and x variable, we may equally well consider x fixed and q variable, thanks to the automorphisms of S. Imagine a hypothetical family of high multiplicity curves at x intersecting E_{∞} at the variable point q, which we partially normalize by blowing up at x. The family of effective divisors cut by this family of curves on one of the curves in the family has certain features that contradict properties of tangential covers obtained by combining Lemma [2.3](#page-2-2) and Theorem [2.1.](#page-2-1)

Proof. It is obvious that $k \leq {m+1 \choose 2}$, since $h^0(\mathcal{O}_S(F_q + kE_\infty)) = k+1$ by e.g. [\[28,](#page-8-7) Proposition 2.3]. Indeed, the map

$$
H^{0}(S, \mathcal{O}_{S}(F_{q} + kE_{\infty})) \longrightarrow H^{0}(S, \mathcal{O}_{S}(F_{q} + kE_{\infty}) \otimes \mathcal{O}_{S}/\mathcal{I}_{x, S}^{m})
$$

connot be injective as soon as $k \geq \binom{m+1}{2}$ for obvious dimension reasons.

Assume by way of contradiction that the inequality was strict, and let

$$
r = \binom{m+1}{2} - k - 1 \ge 0.
$$

Let $Y \in |F_q + kE_\infty|$ of multiplicity at least m at x. Since all divisors in $|F_q + kE_\infty|$ are sums of an integral curve with a multiple of E_{∞} (in characteristic 0, e.g. [\[28,](#page-8-7) $\S2.2$), Y must be integral by the minimality assumption on k. Moreover, by the genus-degree formula,

(2)
$$
k = p_a(Y) \ge p_g(Y) + {\text{mult}_{x} Y \choose 2} \ge 1 + {\text{mult}_{x} Y \choose 2},
$$

so the multiplicity of Y at x must be precisely m .

Let $\beta: S' \to S$ be the blowup of S at x with exceptional curve $W \subset S'$, $C \subset S'$ the proper transform of Y, and $f: C \to S$ the restriction of β to C. Let $g = p_a(C)$. We have $C \sim \beta^* Y - mW$ on S' , so

(3)
$$
C^2 = (\beta^* F_q + k\beta^* E_{\infty} - mW)^2 = -m^2 + 2k
$$

and

$$
C \cdot K_{S'} = (\beta^* F_q + k\beta^* E_{\infty} - mW) \cdot (-2\beta^* E_{\infty} + W) = m - 2.
$$

Hence, by the genus-degree formula,

(4)
$$
g = 1 + \frac{C^2 + C \cdot K_{S'}}{2} = k - {m \choose 2} = m - 1 - r.
$$

It is also clear that $g \geq 1$ since C maps non-constantly to E.

We claim that for any $s \in E$, the linear system $|F_s + kE_{\infty}|$ contains a divisor $Y(s, x)$ of multiplicity m at x. Let $y \in S \backslash E_{\infty}$ such that

(5)
$$
\rho(y) = \rho(x) + s - q \in A_0(E).
$$

Recall that the subgroup of $Aut(S)$ consisting of automorphisms which lie above translation automorphisms of E acts transitively on $S\backslash E_{\infty}$, e.g. [\[28,](#page-8-7) §2.1], and choose such an automorphism ψ such that $\psi(y) = x$. For y general, and therefore for y arbitrary by semi-continuity, there exists a divisor $Y(q, y) \in |F_q + kE_{\infty}|$ with multiplicity at least m at y by assumption. Then we define

$$
Y(s,x) = \psi(Y(q,y)) \subset S,
$$

and note that [\(5\)](#page-3-0) implies that $Y(s, x) \in |F_s + kE_{\infty}|$. Moreover, the multiplicity of $Y(q, y)$ at y, and therefore of $Y(s, x)$ at x, must be precisely m by arguments similar to the ones above, cf. (2) . (For special s, it could happen in principle that

 $Y(q, y)$ contains E_{∞} , but it doesn't really matter; we may consider the analogue of [\(2\)](#page-3-1) for the irreducible component different from E_{∞} .) Then, if $Y'(s, x) \subset S'$ is the proper transform of $Y(s, x)$, we have

(6)
$$
Y'(s,x) \sim \beta^* Y(s,x) - mW \sim \beta^* F_s + k\beta^* E_\infty - mW.
$$

Let $p_1, \ldots, p_r \in C$ be r arbitrary nonsingular points on C, and let

(7)
$$
\mathcal{L}_s = \phi^* \mathcal{O}_E(s) \otimes \mathcal{O}_C(kq + p_1 + \cdots + p_r) \otimes (\mathcal{O}_{S'}(-mW)|_C),
$$

where $\phi = \rho \circ f$. This definition makes clear that $\{\mathcal{L}_s : s \in E\}$ is a curve in Pic(C), though we prefer to think of it as

(8)
$$
\mathcal{L}_s = \mathcal{O}_C(p_1 + \cdots + p_r) \otimes f^* \mathcal{O}_{S'}(Y'(s,x)),
$$

which is equivalent by [\(6\)](#page-4-1). Then $h^0(\mathcal{L}_s) > 0$ by [\(8\)](#page-4-2) (for $s \neq q$, but in particular also for $s = q$ by semi-continuity), and

$$
\deg \mathcal{L}_s = C \cdot Y'(s, x) + r = C^2 + r = 2k - m^2 + r = m - 2 - r = g - 1
$$

by (8) , (3) and (4) .

To summarize, the curve $Z := \{ \mathcal{L}_s : s \in E \} \subset \text{Pic}^{g-1}(C)$ is a family of *effective* degree $g-1$ Cartier divisors on C, and at the same time a coset of $\phi^*(Pic^0(E))$ in $Pic(C)$ by [\(7\)](#page-4-3). It follows that

$$
T_{[\mathcal{L}_s]}Z \subset T_{[\mathcal{L}_s]} \mathrm{Pic}^{g-1}(C) = H^1(C, \mathcal{O}_C)
$$

is the image of $H^1(E, \mathcal{O}_E) \to H^1(C, \mathcal{O}_C)$, which is in fact the KP direction

$$
T_qC = H^0(C, \mathcal{O}_C(q) \otimes \mathcal{O}_q) \hookrightarrow H^1(C, \mathcal{O}_C)
$$

by Lemma [2.3.](#page-2-2) This contradicts Theorem [2.1,](#page-2-1) since $h^0(\mathcal{L}_s) > 0$ for all $s \in E$. \Box

Note that Proposition [2.4](#page-2-0) not only proves [\[28,](#page-8-7) Conjecture 2.5] in characteristic 0, but strengthens it by removing the generality assumption on the j -invariant.

3. Proof of the main result

With [\[28,](#page-8-7) Conjecture 2.5] proven, cases [1](#page-0-0) and [4b](#page-1-0) of Theorem [1.1](#page-0-4) follow directly from [\[28,](#page-8-7) Theorem 3.3], and in essence so does case [3,](#page-0-2) since the formulation in terms of \mathbb{P}^2 given in [\[28,](#page-8-7) Theorem 3.3] is equivalent to the formulation in terms of blowups of \mathbb{P}^2 at 9 points in Theorem [1.1.](#page-0-4) Indeed,

$$
\mathcal{L} = \mathcal{O}_S(3kH - kE_1 - \cdots - kE_8 - (k-1)E_9),
$$

where H is the pullback of the class of a line in \mathbb{P}^2 , and $E_1, \ldots, E_9 \subset S$ are the 9 exceptional curves, since we may assume $E = E_9$ by Cremona transformations.

This leaves cases [2](#page-0-1) and [4a.](#page-0-3) If we construct suitable specializations of (S, \mathcal{L}) in situations [2](#page-0-1) and [4a](#page-0-3) satisfying the conditions in [\[28,](#page-8-7) Proposition 3.1], then we are done. Such specializations exist, with the ultimately minor caveat that, in the abelian surface case, the base of the elliptic fibration as in loc. cit. is not \mathbb{P}^1 , but an elliptic curve instead. Thus, we need a minor extension of [\[28,](#page-8-7) Proposition 3.1].

Proposition 3.1. Let π : $X \to B$ be a smooth projective family of surfaces over a smooth quasi-projective curve B, and let $\mathcal{L}_X \in \text{Pic}(X)$ relatively ample over B. Let $X_t = \pi^{-1}(t)$ and $\mathcal{L}_t = \mathcal{L}_X|_{X_t}$ for any $t \in B$. Fix $b \in B$ closed. Assume that

(9)
$$
H^{1}(X_{b}, \mathcal{L}_{b}) = H^{2}(X_{b}, \mathcal{L}_{b}) = 0,
$$

and that the central fibre X_b has an elliptic fibration $f : X_b \to Y$, where Y is a smooth projective curve, with the following properties:

(1) For general $s \in Y$, the natural short exact sequence

$$
0 \longrightarrow N_{f^{-1}(s)/X_b} \longrightarrow N_{f^{-1}(s)/X} \longrightarrow T_b B \otimes \mathcal{O}_{f^{-1}(s)} \longrightarrow 0
$$

on $f^{-1}(s)$ is not split;

(2) There exists a fixed section $G \subset X_b$ of f, such that any divisor $D \in |\mathcal{L}_b|$ is the sum of G and dim $|\mathcal{L}_b| + p_q(Y)$ (mobile) fibres of f.

Let
$$
t \in B
$$
 general and $(S, \mathcal{L}) = (X_t, \mathcal{L}_t)$. If $m_1, m_2, ..., m_n \ge 1$, then

(10)
$$
\dim |\mathcal{L}(m_1,...,m_n)| = \max \left\{-1, \dim |\mathcal{L}| - \sum_{i=1}^n \frac{m_i(m_i+1)}{2}\right\},\,
$$

with notation as in [\(1\)](#page-1-2) for $|\mathcal{L}(m_1, \ldots, m_n)|$.

Proof. The fact that the left hand side is at least the right hand side in [\(10\)](#page-5-0) (the dimension is at least the expected dimension) is well-known and trivial, so we only need to prove the reverse inequality. When $Y \simeq \mathbb{P}^1$, this is precisely the combination of [\[28,](#page-8-7) Proposition 3.1] and Proposition [2.4.](#page-2-0) In general, most of the argument in [\[28\]](#page-8-7) applies verbatim, though the conclusion requires an additional ingredient, which is precisely the 1-dimensional interpolation ('SHGH') problem.

Lemma 3.2. Let $M \in Pic(Y)$, and $p_1, \ldots, p_n \in Y$ general points. Then

$$
\dim |\mathcal{M}(-D)| = \max\{-1, \dim |\mathcal{M}| - \deg D\}
$$

for any effective divisor D such that $\text{Supp}(D) \subseteq \{p_1, \ldots, p_n\}.$

Proof. Generalizing to non-complete linear systems, the statement boils down to the case $n = 1$, which is the well-known fact that, in characteristic 0, a linear system on a curve has only finitely many inflection points, e.g. [\[8,](#page-7-8) Proposition 1.1]. \Box

Let us explain how 1D interpolation must be applied at the end of the argument in [\[28\]](#page-8-7) to generalize the proof. Condition [2](#page-5-1) implies that there exists $\mathcal{M} \in \text{Pic}(Y)$ such that $|\mathcal{M}| \simeq |\mathcal{L}_b|$ by $D \mapsto G + f^{-1}(D)$ since $\pi^* : Pic(Y) \to Pic(X_b)$ is injective by [\[13,](#page-7-9) III, Exercise 12.4] and the projection formula. We use the notation in the proof of [\[28,](#page-8-7) Proposition 3.1]. The next-to-last centred formula states that

(11)
$$
P_b \subseteq \left\{ [D] \in |\mathcal{L}_b| : \mathrm{coeff}_{E_i} D \ge \lambda(m_i) = \frac{m_i(m_i+1)}{2} \right\} =: T,
$$

that is, the limit in X_b of divisors with high multiplicities at the chosen points must contain the elliptic fibre E_i with multiplicity at least $\lambda(m_i)$, for $i = 1, \ldots, n;$ please see also Remark [3.3](#page-6-0) below. Indeed, the argument in loc. cit. up to that point requires no changes, and the formula for $\lambda(m_i)$ is our Proposition [2.4.](#page-2-0) Under $|\mathcal{M}| \simeq |\mathcal{L}_b|$, T corresponds to $T_Y = \{D \in |\mathcal{M}| : \text{coeff}_{p_i} D \geq \lambda(m_i)\}.$ Then

$$
\dim P_b \le \dim T = \dim T_Y = \max \left\{-1, \dim |\mathcal{M}| - \sum_{i=1}^n \frac{m_i(m_i+1)}{2}\right\}
$$

$$
= \max \left\{-1, \dim |\mathcal{L}_t| - \sum_{i=1}^n \frac{m_i(m_i+1)}{2}\right\}
$$

by Lemma [3.2](#page-5-2) and dim $|\mathcal{M}| = \dim |\mathcal{L}_b| = \dim |\mathcal{L}_t|$, and we may conclude as in [\[28\]](#page-8-7). (The situation is thus slightly different when $Y \not\cong \mathbb{P}^1$, in that the limit divisors are forced to contain some elliptic fibres other than E_1, \ldots, E_n , which plays a role in obtaining the correct bound). \Box

Remark 3.3. For the reader's convenience, we review that the argument in [\[28\]](#page-8-7) entails blowing up the elliptic fibres which contain the (reduced) limits of the fat points in the central fibre. This is reminiscent of the approach of Ciliberto and Miranda to the SHGH Conjecture using degenerations [\[5,](#page-7-10) [6\]](#page-7-11). In our situation, the exceptional divisors of such blowups are isomorphic to the ruled surface in §[2.2.](#page-2-3)

It remains to construct specializations with the desired properties.

Ruled surfaces $(4a)$. It is clear that the group of automorphisms of the surface acts transitively on the set of linear systems we are considering, so all such linear systems behave identically. Let $B = Pic^0(E)$. Let P be the universal (Poincaré) line bundle over $B \times E = Pic^0(E) \times E$. Then we define $X = \mathbb{P}(\mathcal{O}_{B \times E} \oplus \mathcal{P})$ and $o = [\mathcal{O}_E] \in B$. There are two distinguished sections $\Sigma_0, \Sigma_\infty \subset X$ corresponding to the projections to the two terms of $\mathcal{O}_{B \times E} \oplus \mathcal{P}$. Let $q \in E$ and $k \geq 1$ integer, and

$$
\mathcal{L}_X = \mathcal{O}_X(k\Sigma_\infty + \pi^{-1}(B \times \{q\})).
$$

Then $X_o = E \times \mathbb{P}^1$, and condition [2](#page-5-1) follows easily from the fact that any effective divisor on $E \times \mathbb{P}^1$ with intersection number 1 with $E \times \{\text{point}\}\$ is the sum of fibres of projections to the two factors, which is elementary.

Condition [\(9\)](#page-4-4) follows from the Kodaira vanishing theorem.

To check the condition [1,](#page-5-3) let $R \simeq \text{Spec } \mathbb{C}[\epsilon]/(\epsilon^2)$ be the first order thickening of $o = [\mathcal{O}_E]$ in B. Let $y \in \mathbb{P}^1$ such that $E \times \{y\} \not\subset \Sigma_0, \Sigma_\infty$, and let's assume by way of contradiction that the sequence

(12)
$$
0 \longrightarrow N_{E \times \{y\}, X_o} \longrightarrow N_{E \times \{y\}, X} \longrightarrow N_{X_o, X}|_{E \times \{y\}} \longrightarrow 0
$$

was split. Then $H^0(N_{E\times \{y\},X_o}) \hookrightarrow H^0(N_{E\times \{y\},X})$ is not surjective, and the global sections of $H^0(N_{E\times \{y\},X})$ not coming from global sections of $H^0(N_{E\times \{y\},X_o})$ give a first order deformation of $E \times \{y\} \subset X$ flat over R. This deformation is a section of $X|_{R\times E}$, so corresponds to a surjective map ν : $\mathcal{O}_{R\times E} \oplus \mathcal{P}|_{R\times E} \to \mathcal{Q}$ to a line bundle Q on $R \times E$. Clearly, deg $Q_o = 0$. Since we are assuming $E \times \{y\} \not\subset \Sigma_0, \Sigma_{\infty}$, the restrictions of ν to $\mathcal{O}_{R\times E} \oplus 0$ and $0 \oplus \mathcal{P}|_{R\times E}$ are nonzero on $\{o\} \times E$, and are therefore isomorphisms for degree reasons. Hence, $\mathcal{Q} \simeq \mathcal{O}_{R \times E}$ and $\mathcal{Q} \simeq \mathcal{P}|_{R \times E}$. However, $\mathcal{P}|_{R\times E} \not\cong \mathcal{O}_{R\times E}$ by definition, which is a contradiction.

Abelian surfaces. Let $X \to B$ be a one-parameter family of $(1, d)$ -polarized abelian surfaces specializing to a product $E \times F$ of two elliptic curves, as in [\[29\]](#page-8-15) (or [\[3,](#page-7-12) §4] implicitly). Let us be more specific. If $E \times F$ is polarized by $\mathcal{J} = \mathcal{J}_E \boxtimes \mathcal{J}_F$, with deg $\mathcal{J}_E = 1$ and deg $\mathcal{J}_F = d$, then $(E \times F, \mathcal{J})$ is a $(1, d)$ -polarized abelian surface. Let \mathcal{A}_d be the Deligne-Mumford moduli stack of $(1, d)$ -polarized abelian surfaces, and $(U, u) \rightarrow \mathcal{A}_d$ an affine (in particular, quasi-projective) étale neighbourhood of $(E \times F, \mathcal{J})$. By the usual Kodaira-Spencer theory, there is a natural injective map

$$
T_{u}U = T_{(E \times F, \mathcal{J})} A_{d} \hookrightarrow H^{1}(E \times F, \mathcal{T}_{E \times F}) \simeq \mathbb{C}^{4}.
$$

By [\[4,](#page-7-13) Lemma 2.2], the image is contained in the annihilator of $c_1(\mathcal{J}) \in H^{1,1}(E \times F)$ relative to the natural pairing

$$
H^1(E \times F, \mathcal{T}_{E \times F}) \times H^1(E \times F, \Omega_{E \times F}) \longrightarrow \mathbb{C},
$$

and hence must coincide with it since dim $T_u U \ge \dim U = 3$. (To apply [\[4,](#page-7-13) Lemma 2.2, we need to consider a curve through u in U.) Then we take $(B, o) \rightarrow (U, u)$ to be a smooth curve through u corresponding to a *general* direction $T_oB \hookrightarrow T_uU$,

this is possible since U is quasi-projective. (In [\[29\]](#page-8-15), this *first order* generality was not really used and thus not imposed, but here it is needed.)

Condition [\(9\)](#page-4-4) again follows from the Kodaira vanishing theorem. Condition [2](#page-5-1) follows from the fact that any divisor $D \in |\mathcal{J}|$ is of the form

 $D = E \times D' + z \times F$, where $|D'| \in |\mathcal{J}_F|$ and $\{z\} = |\mathcal{J}_E|$,

which is elementary (e.g. $[3]$ or $[29]$). It remains to check condition [1,](#page-5-3) namely, that the short exact sequence notationally identical to [\(12\)](#page-6-1) is not split. Here, $y \in F$ is completely arbitrary. The proof is very similar to the proof of [\[4,](#page-7-13) Proposition 2.1], so we will be brief. It suffices to check that $H^0(N_{X_o,X}|_{E\times\{y\}}) \to H^1(N_{E\times\{y\},X_o})$ is nonzero. As in $[4, (2.10)]$ $[4, (2.10)]$, this map factors as

(13)
$$
H^0(N_{X_o,X}|_{E\times\{y\}}) \simeq T_oB \xrightarrow{\text{ks}} H^1(\mathcal{T}_{E\times F}) \longrightarrow H^1(\mathcal{T}_{E\times F}|_{E\times\{y\}}) \longrightarrow H^1(N_{E\times\{y\},X_o}),
$$

where ks is the Kodaira-Spencer map. It is clear that the composition of the last two maps in [\(13\)](#page-7-14) is surjective, and its kernel is different from the annihilator of $c_1(\mathcal{J})$ considered above – for instance, under the natural identification $H^1(\mathcal{T}_{E\times F}) \simeq \mathbb{C}^4$, the former is clearly a 'coordinate hyperplane', which the latter is not. Then the generality of the Kodaira-Spencer map (i.e. of the tangent direction in U) implies that [\(13\)](#page-7-14) is nonzero, as desired.

This completes the proof of Theorem [1.1.](#page-0-4)

Remark 3.4. Theorem [1.1](#page-0-4) rather vacuously implies the primitive case $(d = 1)$ of [\[27,](#page-8-3) Conjecture 2.1]. Then the primitive case of [\[27,](#page-8-3) Conjecture 2.3] follows from [\[27,](#page-8-3) Theorem 3.7]. To be pedantic, the equivalence (since one implicitation is trivial) of the two conjectures proved in loc. cit. does not logically imply their equivalence for $d = 1$, though the proof clarifies that there is absolutely no issue.

REFERENCES

- [1] C. Birkenhake and P. Vanhaecke. The vanishing of the theta function in the kp direction: A geometric approach. Compos. Math., 135:323–330, 2003.
- [2] C. Bocci and R. Miranda. Topics on interpolation problems in algebraic geometry. Rend. Sem. Mat. Univ. Politec. Torino, 62:279–334, 2004.
- [3] J. Bryan and C. Leung. Generating functions for the number of curves on abelian surfaces. Duke Math. J., 99:311–328, 1999.
- [4] X. Chen. A simple proof that rational curves on k3 are nodal. Math. Ann., 324:71–104, 2002.
- [5] C. Ciliberto and R. Miranda. Degenerations of planar linear systems. J. Reine Angew. Math., 501:191–220, 1998.
- [6] C. Ciliberto and R. Miranda. Linear systems of plane curves with base points of equal multiplicity. Trans. Amer. Math. Soc., 352:4037–4050, 2000.
- [7] M. Dumnicki. Special homogeneous linear systems on hirzebruch surfaces. Geom. Dedicata, 147:283–311, 2010.
- [8] D. Eisenbud and J. Harris. Divisors on general curves and cuspidal rational curves. Invent. Math., 74:371–418, 1983.
- [9] G. Farkas and N. Tarasca. Du val curves and the pointed brill–noether theorem. Selecta Math. (N.S.), 23:2243–2259, 2017.
- [10] J. Fay. On the even-order vanishing of jacobian theta functions. Duke Math. J., 51:109–132, 1984.
- [11] A. Gimigliano. On linear systmes of plane curves. PhD thesis, Queen's University, 1987.
- [12] B. Harbourne. The geometry of rational surfaces and hilbert functions of points in the plane. In Proceedings of the 1984 Vancouver conference in algebraic geometry, volume 6 of CMS Conf. Proc., pages 95–111, Providence, RI, Amer. Math. Soc., 1986.
- [13] R. Hartshorne. Algebraic geometry. Number 52 in Grad. Texts in Math. Springer, 1977.
- [14] A. Hirschowitz. Une conjecture pour la cohomologie des diviseurs sur les surfaces rationnelles génériques. J. Reine Angew. Math., 397:208-213, 1989.
- [15] J. Huizenga. Interpolation on surfaces in p^3 . Trans. Amer. Math. Soc., 365:623-644, 2013.
- [16] A. Laface. On linear systems of curves on rational scrolls. Geom. Dedicata, 90:127–144, 2002.
- [17] A. Laface. Special systems through double points on an algebraic surface. Proc. Amer. Math. Soc., 139:1971–1981, 2011.
- [18] G. Segal and G. Wilson. Loop groups and equations of kdv type. Publ. Math. IHES, 61:5–65, 1985.
- [19] B. Segre. Alcune questioni su insiemi finiti di punti in geometria algebrica. Univ. e Politec. Torino Rend. Sem. Mat., 20:67–85, 1960/1961.
- [20] E. Sernesi. On treibich-verdier curves. Pure Appl. Math. Q., 2023 (in press).
- [21] A. Treibich. Tangential polynomials and elliptic solitons. Duke Math. J., 59:611–627, 1989.
- [22] A. Treibich. Compactified jacobians of tangential covers. In D. Bennequin, editor, Colloque Systémes Intégrables en hommage à J.-L. Verdier, Boston, 1993. Birkhauser.
- [23] A. Treibich. Revetements tangentiels et condition de brill-noether. CRAS Paris, 316:815–817, 1993.
- [24] A. Treibich and J.-L. Verdier. Variétés de kritchever des solitons elliptiques. In A. Beauville and S. Ramanan, editors, Actes du Colloque Franco-Indien, Bombay, 1989.
- [25] A. Treibich and J.-L. Verdier. Solitons elliptiques. In The Grothendieck Festschrift, Volume III, number 88 in Progress in Math., pages 437–480. Birkhauser, Boston, 1991.
- [26] C. De Volder and A. Laface. Degeneration of linear systems through fat points on k3 surfaces. Trans. Amer. Math. Soc., 357:3673–3682, 2005.
- [27] C. De Volder and A. Laface. Linear systems on generic k3 surfaces. Bull. Belg. Math. Soc. Simon Stevin, 12:481–489, 2005.
- [28] A. Zahariuc. Elliptic surfaces and linear systems with fat points. Math. Z., 293:647–660, 2019.
- [29] A. Zahariuc. The severi problem for abelian surfaces in the primitive case. J. Math. Pures Appl., 158:320–349, 2022.

Department of Mathematics and Statistics, University of Windsor, 401 Sunset Ave, Windsor, ON, N9B 3P4, Canada

Email address: adrian.zahariuc@uwindsor.ca