

FINITENESS OF THE NUMBER OF IRREDUCIBLE λ -QUIDDITIES OVER A FINITE COMMUTATIVE AND UNITARY RING

FLAVIEN MABILAT

RÉSUMÉ. Les λ -quiddités de taille n sont des n -uplets d'éléments d'un ensemble fixé, solutions d'une équation matricielle apparaissant lors de l'étude des frises de Coxeter. L'étude de ces solutions passe notamment par l'utilisation d'une notion d'irréductibilité. L'objectif principal de ce texte est de montrer qu'il y a un nombre fini de λ -quiddités irréductibles lorsque l'on se place sur un anneau commutatif unitaire fini et d'obtenir dans ce cas une majoration de la taille maximale de ces dernières.

ABSTRACT. λ -quiddities of size n are n -tuples of elements of a fixed set, solutions of a matrix equation appearing in the study of Coxeter's friezes. The study of these solutions involves in particular the use of a notion of irreducibility. The main objective of this text is to show that there is a finite number of irreducible λ -quiddities over a finite unitary commutative ring and to obtain in this case an upper bound of the maximal size of them.

« Les choses n'arrivent quasi jamais comme on se les imagine. »
Madame de Sévigné, *Lettres*.

1. INTRODUCTION

Since its apparition at the begining of the 1970s, the concept of Coxeter's friezes ([3]) has drawn great attention. Indeed, these objects have been the source of a lot of beautiful results and are closely linked to many topics (see for example [13]). One of the most important elements of the study of Coxeter's friezes over a subset R of a commutative and unitary ring A is the resolution of the following equation over R :

$$M_n(a_1, \dots, a_n) := \begin{pmatrix} a_n & -1_A \\ 1_A & 0_A \end{pmatrix} \begin{pmatrix} a_{n-1} & -1_A \\ 1_A & 0_A \end{pmatrix} \cdots \begin{pmatrix} a_1 & -1_A \\ 1_A & 0_A \end{pmatrix} = -Id.$$

Besides, the presence of the matrices $M_n(a_1, \dots, a_n)$ is very interesting since they also appear in the study of a lot of other mathematical objects, such as Hirzebruch-Jung continued fractions or discrete Sturm-Liouville equations (see for instance the introduction of [15]).

Moreover, the study of the previous equation naturally leads to consider the generalised equation below over a subset R of a commutative and unitary ring $A \neq \{0_A\}$:

$$(E_R) \quad M_n(a_1, \dots, a_n) = \pm Id.$$

We will say that a solution (a_1, \dots, a_n) of (E_R) is a λ -quiddity of size n over R . Our objective is to obtain information about λ -quiddities over different sets. There are several ways to achieve this goal. The first one, and the most natural, is to look for a general construction of all the solutions of (E_R) . In this direction, we have, for example, a recursive construction and a combinatorial description of all the solutions of $(E_{\mathbb{N}^*})$ (see [15] Theorem 1 and Theorem 2). Another possibility is to find some general information, such as the number of fixed-size λ -quiddities. Several results of that kind are known (see [2, 6, 12, 14]). However, here, we will consider a third way. Indeed, to study the solutions of (E_R) it is convenient to introduce a notion of irreducible λ -quiddity which allow us to restrict our attention to these solutions only (see [4] and the next section).

2020 *Mathematics Subject Classification*. 05A05, 05E16, 05A20.

Key words and phrases. λ -quiddity; modular group; finite rings, Coxeter's friezes.

A lot of results concerning irreducible λ -quiddities are already known. In particular, we have a complete list of the irreducible solutions of (E_R) over several sets (see section 2.2). However, in most cases, we do not have such a list and we try in a more modest way to find information about irreducible solutions. In particular, two main questions arise : Is the number of irreducible λ -quiddities over R finite ? Is the size of irreducible solutions of (E_R) bounded ? If R is finite then the two questions are strictly the same and this is precisely the case we want to deal with here. In particular, we have conjectured in a previous text that, for all $N \geq 2$, $(E_{\mathbb{Z}/N\mathbb{Z}})$ has a finite number of irreducible solutions ([8] Conjecture 1). Note that the resolution of (E_R) with $R = \mathbb{Z}/N\mathbb{Z}$ is closely linked to the different writings of the elements of the congruence subgroup $\hat{\Gamma}(N) := \{C \in SL_2(\mathbb{Z}), C = \pm Id [N]\}$. Indeed, all the matrices of $SL_2(\mathbb{Z})$ can be written in the form $M_n(a_1, \dots, a_n)$, with $a_i \in \mathbb{N}^*$ (note that in this text \mathbb{N} is the set of all non negative integers). Since this expression is not unique, it is natural to look for all the writings of this form for a given matrix, or a set of matrices.

Our main objective in this text is to show that the number of irreducible λ -quiddities over a finite commutative and unitary ring A is finite, which in particular will give us a solution to the conjecture mentionned in the previous paragraph. Besides, we want to have an upper bound of the size of irreducible λ -quiddities over A . For a unitary ring A , we set $\text{car}(A)$ the characteristic of A and we recall the classical following property: if A is finite then $\text{car}(A) \neq 0$. In this article, we will prove the following results :

Theorem 1.1. *Let $(A, +, \times)$ be a finite commutative and unitary ring (different from $\{0_A\}$) and $R \subset A$ a submagma of $(A, +)$, that is to say a subpart of A closed under $+$.*

i) *There is a finite number of irreducible λ -quiddities over R .*

ii) *Let ℓ_A be the maximum size of an irreducible λ -quiddity over A .*

— *If $\text{car}(A) = 2$ then $4 \leq \ell_A \leq \frac{|SL_2(A)|}{|A|} + 2$.*

— *If $\text{car}(A) \neq 2$ then $\max(4, \text{car}(A)) \leq \ell_A \leq \frac{|SL_2(A)|}{2|A|} + 2$.*

Corollary 1.2. i) *Let $N \geq 5$ and p_1, \dots, p_r the distinct prime factors of N .*

$$N \leq \ell_{\mathbb{Z}/N\mathbb{Z}} \leq \frac{N^2}{2} \prod_{i=1}^r \left(1 - \frac{1}{p_i^2}\right) + 2.$$

ii) *Let q the power of a prime number p .*

— *If $p = 2$ then $4 \leq \ell_{\mathbb{F}_q} \leq q^2 + 1$.*

— *If $p \neq 2$ then $\max(4, p) \leq \ell_{\mathbb{F}_q} \leq \frac{q^2+3}{2}$.*

Theorem 1.1 is proved in section 3.1 while the proof of corollary 1.2 is given in section 3.2.

2. DEFINITIONS AND PRELIMINARY RESULTS

The aim of this section is to give the precise definitions of the notions mentionned in the introduction and to provide some elements which will be useful in the proofs of our main results. To have a complete view on irreducible λ -quiddities, we will also recall some known results obtained for several precise rings. In all this section, $(A, +, \times)$ is a commutative and unitary rings different from $\{0_A\}$ and R is a submagma of $(A, +)$. 0_A is the identity element of $+$ and 1_A is the identity element of \times . If $N \geq 2$ and $a \in \mathbb{Z}$, we set $\bar{a} := a + N\mathbb{Z}$.

2.1. Quiddities. We begin by the precise definition of the concept of λ -quiddity.

Definition 2.1 ([4] definition 2.2). *Let $n \in \mathbb{N}^*$. The n -tuple $(a_1, \dots, a_n) \in R^n$ is a λ -quiddity over R if (a_1, \dots, a_n) is a solution of (E_R) , that is to say if it exists $\epsilon \in \{\pm 1_A\}$ such that $M_n(a_1, \dots, a_n) = \epsilon Id$. If there is no ambiguity, we will omit to precise the ring.*

Throughout the rest of this text, we will speak indiscriminately of the solutions of (E_R) or of the λ -quiddities over R . To define the notion of irreducibility, we need the two following definitions.

Definition 2.2 ([4], lemma 2.7). Let $(a_1, \dots, a_n) \in R^n$ and $(b_1, \dots, b_m) \in R^m$. We define the following operation :

$$(a_1, \dots, a_n) \oplus (b_1, \dots, b_m) = (a_1 + b_m, a_2, \dots, a_{n-1}, a_n + b_1, b_2, \dots, b_{m-1}).$$

The $(n + m - 2)$ -tuple obtained is the sum of (a_1, \dots, a_n) with (b_1, \dots, b_m) .

Examples. We give here some examples of sums over $A = R = \mathbb{Z}$:

- $(1, 2, 3) \oplus (1, 0, -2, 4) = (5, 2, 4, 0, -2)$;
- $(-1, 1, 0, 2) \oplus (2, 2, 2) = (1, 1, 0, 4, 2)$;
- $(2, 1, -, 1, 0, -3) \oplus (2, 3, 1, 1) = (3, 1, -1, 0, -1, 3, 1)$.

The operation \oplus is very useful since it has the interesting following property. Let (b_1, \dots, b_m) a solution of (E_R) . $(a_1, \dots, a_n) \oplus (b_1, \dots, b_m)$ is a λ -quiddity over R if and only if (a_1, \dots, a_n) is a solution of (E_R) (see [4, 16]). Besides, if $0_A \in R$, we have the following equality :

$$(a_1, \dots, a_n) \oplus (0_A, 0_A) = (0_A, 0_A) \oplus (a_1, \dots, a_n) = (a_1, \dots, a_n).$$

However, \oplus is neither commutative nor associative (see [16] example 2.1).

Definition 2.3 ([4], définition 2.5). Let $(a_1, \dots, a_n) \in R^n$ and $(b_1, \dots, b_n) \in R^n$. $(a_1, \dots, a_n) \sim (b_1, \dots, b_n)$ if (b_1, \dots, b_n) can be obtained by cyclic rotations of (a_1, \dots, a_n) or of (a_n, \dots, a_1) .

\sim is an equivalence relation on the set R^n (see [16] lemme 1.7). Moreover, if $(a_1, \dots, a_n) \sim (b_1, \dots, b_n)$ then (a_1, \dots, a_n) is a solution of (E_R) if and only if (b_1, \dots, b_n) is a λ -quiddity over R (see [4] proposition 2.6).

We can now defined the notion of irreducibility used in the introduction.

Definition 2.4 ([4], définition 2.9). A λ -quiddity over R (c_1, \dots, c_n) ($n \geq 3$) is said to be reducible if there exists a λ -quiddity over R (b_1, \dots, b_l) and an m -tuple $(a_1, \dots, a_m) \in R^m$ such that

- $(c_1, \dots, c_n) \sim (a_1, \dots, a_m) \oplus (b_1, \dots, b_l)$;
- $m \geq 3$ and $l \geq 3$.

A λ -quiddity over R is said to be irreducible if it is not reducible.

Remark. We suppose $0_A \in R$. $(0_A, 0_A)$ is never considered as an irreducible solution of (E_R) .

We set $\ell_R \in \mathbb{N} \cup \{+\infty\}$ the maximum size of an irreducible λ -quiddity over R . In the next two subparts, we will give some preliminary information about ℓ_R .

2.2. Some classification results. The objective of this subsection is to gather some classification results already known concerning irreducible λ -quiddities. We begin by the case $A = \mathbb{Z}$.

Theorem 2.5 ([4] Theorem 3.1 and 3.2). *i) The set of irreducible λ -quiddities over \mathbb{Z} is*

$$\{(1, 1, 1), (-1, -1, -1), (a, 0, -a, 0), (0, a, 0, -a), a \in \mathbb{Z} - \{-1, 1\}\}.$$

ii) The only irreducible λ -quiddities over \mathbb{N} are $(1, 1, 1)$ and $(0, 0, 0, 0)$.

Theorem 2.6. *The set of irreducible λ -quiddities over \mathbb{N}^* is $\{(1, \dots, 1) \in (\mathbb{N}^*)^{3n}, n \in \mathbb{N}^*\}$.*

Proof. This is an easy consequence of the combinatorial description of the solutions of $(E_{\mathbb{N}^*})$ given in Theorem 1 of [15]. For a detailed proof see [10] proposition 4.2.1. □

These two theorems give us interesting information about irreducible λ -quiddities over infinite sets :

- It exists infinite rings such that $\ell_A \neq +\infty$.
- We can find an infinite set R such that the number of irreducible λ -quiddities over R is finite.
- It exists sets $R' \subset R$ such that $\ell_R \neq +\infty$ and $\ell_{R'} = +\infty$.

Theorem 2.7 ([11] Theorem 2.8). *Let I a set containing at least two elements and (A_i) a family of commutative and unitary rings. We suppose that at least two different A_i have zero characteristic. Let $n \geq 3$. It exists an irreducible λ -quiddity of size n over $\prod_{i \in I} A_i$.*

We also have information about finite rings. In particular, we have a precise classification of the irreducible solutions of $(E_{\mathbb{Z}/N\mathbb{Z}})$ in the cases $2 \leq N \leq 6$ (see [8] Theorem 2.5).

Theorem 2.8. *i) ([8] Theorem 2.5) The irreducible λ -quiddities over $\mathbb{Z}/4\mathbb{Z}$ are : $(\bar{1}, \bar{1}, \bar{1}), (\bar{-1}, \bar{-1}, \bar{-1}), (\bar{0}, \bar{0}, \bar{0}, \bar{0}), (\bar{0}, \bar{2}, \bar{0}, \bar{2}), (\bar{2}, \bar{0}, \bar{2}, \bar{0})$ and $(\bar{2}, \bar{2}, \bar{2}, \bar{2})$.*

ii) ([11] Theorem 2.6) The irreducible λ -quiddities over $(\mathbb{Z}/2\mathbb{Z}) \times (\mathbb{Z}/2\mathbb{Z})$ are (up to cyclic rotations):

- $((\bar{1}, \bar{1}), (\bar{1}, \bar{1}), (\bar{1}, \bar{1}))$;
- $((\bar{0}, \bar{0}), (\bar{0}, \bar{0}), (\bar{0}, \bar{0}), (\bar{0}, \bar{0})), ((\bar{0}, \bar{0}), (\bar{0}, \bar{1}), (\bar{0}, \bar{0}), (\bar{0}, \bar{1})), ((\bar{0}, \bar{0}), (\bar{1}, \bar{0}), (\bar{0}, \bar{0}), (\bar{1}, \bar{0})), ((\bar{1}, \bar{0}), (\bar{0}, \bar{1}), (\bar{1}, \bar{0}), (\bar{0}, \bar{1}))$;
- $((\bar{1}, \bar{0}), (\bar{1}, \bar{0}), (\bar{1}, \bar{0}), (\bar{1}, \bar{0})), ((\bar{0}, \bar{1}), (\bar{0}, \bar{1}), (\bar{0}, \bar{1}), (\bar{0}, \bar{1}), (\bar{0}, \bar{1}), (\bar{0}, \bar{1}))$.

iii) ([11] Theorem 2.7) The irreducible λ -quiddities over $\mathbb{F}_4 := \frac{(\mathbb{Z}/2\mathbb{Z})[X]}{\langle X^2+X+1 \rangle}$ are (up to cyclic rotations) :

- $(\bar{1}, \bar{1}, \bar{1})$;
- $(\bar{0}, \bar{0}, \bar{0}, \bar{0}), (\bar{0}, X, \bar{0}, X), (\bar{0}, X + \bar{1}, \bar{0}, X + \bar{1})$;
- $(X, X, X, X, X), (X + \bar{1}, X + \bar{1}, X + \bar{1}, X + \bar{1}, X + \bar{1})$;
- $(X, X + \bar{1}, X, X + \bar{1}, X, X + \bar{1})$;
- $(X, X, X + \bar{1}, X + \bar{1}, X, X, X + \bar{1}, X + \bar{1})$;
- $(X, X, X + \bar{1}, X, X, X + \bar{1}, X, X, X + \bar{1}), (X + \bar{1}, X + \bar{1}, X, X + \bar{1}, X + \bar{1}, X, X + \bar{1}, X + \bar{1}, X)$.

This result show us that finite rings having the same cardinality can have a very different list of irreducible λ -quiddities. In particular, ℓ_A does not only depend of $|A|$.

2.3. Preliminary results. The aim of this subpart is to collect several results which will be useful in the next section. First, we give the solutions of (E_R) of small size (see for example [9] section 3.1) :

Lemma 2.9. — (E_R) has no solution of size 1.

- $(0_A, 0_A)$ is the only λ -quiddity over A of size 2.
- $(1_A, 1_A, 1_A)$ and $(-1_A, -1_A, -1_A)$ are the only λ -quiddities over A of size 3. Besides, they are irreducible.
- The set of solutions of (E_R) of size 4 is $\{(-a, b, a, -b) \in R^4, ab = 0_A\} \cup \{(a, b, a, b) \in R^4, ab = 2 \times 1_A\}$.
- Solutions of (E_A) of size greater than 4 containing $\pm 1_A$ are reducible.
- A solution of (E_A) of size 4 is irreducible if and only if it does not contain $\pm 1_A$.
- Solutions of (E_A) of size greater than 5 containing 0_A are reducible.

The following result is an easy consequence of lemma 2.9.

Proposition 2.10. Let $(G, +)$ an infinite subgroup of an infinite ring $(A, +, \times)$. There is infinitely many irreducible λ -quiddities over G .

Proof. Since $(G, +)$ is a group, $0_A \in G$. For all $g \in G - \{-1_A, 1_A\}$, $(0_A, g, 0_A, -g)$ is an irreducible λ -quiddity over G . □

Now, we will give some elements about the cardinality of the special linear group over a finite ring. We begin by recall the two classical following results.

Theorem 2.11. Let q a power of a prime number. $|SL_2(\mathbb{F}_q)| = q(q^2 - 1)$.

Proposition 2.12. Let $n \in \mathbb{N}^*$ and A_1, \dots, A_n n commutative and unitary rings. $SL_2(A_1 \times \dots \times A_n)$ is a group isomorphic to $SL_2(A_1) \times \dots \times SL_2(A_n)$.

Proof. We consider the following application :

$$\begin{aligned} \psi : \quad & SL_2(A_1 \times \dots \times A_n) \longrightarrow SL_2(A_1) \times \dots \times SL_2(A_n) \\ & \begin{pmatrix} (a_i)_{1 \leq i \leq n} & (b_i)_{1 \leq i \leq n} \\ (c_i)_{1 \leq i \leq n} & (d_i)_{1 \leq i \leq n} \end{pmatrix} \longmapsto \begin{pmatrix} \begin{pmatrix} a_i & b_i \\ c_i & d_i \end{pmatrix} \end{pmatrix}_{1 \leq i \leq n}. \end{aligned}$$

ψ is a group isomorphism. □

Theorem 2.13 ([7] 3.24, pages 211-215). *Let $N \geq 2$ and p_1, \dots, p_r the distinct prime factors of N .*

$$|SL_2(\mathbb{Z}/N\mathbb{Z})| = N^3 \prod_{i=1}^r \left(1 - \frac{1}{p_i^2}\right).$$

We will also use an expression of the coefficients of $M_n(a_1, \dots, a_n)$. For this, we set $K_{-1} = 0$, $K_0 = 1$ and for $i \geq 1$ and $(a_1, \dots, a_i) \in A^i$:

$$K_i(a_1, \dots, a_i) = \begin{vmatrix} a_1 & 1 & & & \\ & 1 & a_2 & & \\ & & \ddots & \ddots & \ddots \\ & & & 1 & a_{i-1} & 1 \\ & & & & 1 & a_i \end{vmatrix}.$$

$K_i(a_1, \dots, a_i)$ is the continuant of a_1, \dots, a_i . By induction, we have the following equality (see for example [10] proposition 4.3.2) :

$$\textbf{Proposition 2.14. } (a_1, \dots, a_n) \in A^n. \quad M_n(a_1, \dots, a_n) = \begin{pmatrix} K_n(a_1, \dots, a_n) & -K_{n-1}(a_2, \dots, a_n) \\ K_{n-1}(a_1, \dots, a_{n-1}) & -K_{n-2}(a_2, \dots, a_{n-1}) \end{pmatrix}.$$

Lemma 2.15. *Let A a commutative and unitary ring. Let $n \geq 1$ and $(a_1, \dots, a_n) \in A^n$ such that $K_n(a_1, \dots, a_n) = \epsilon \in \{\pm 1_A\}$. We set $x = \epsilon K_{n-1}(a_2, \dots, a_n)$ and $y = \epsilon K_{n-1}(a_1, \dots, a_{n-1})$.*

$$M_{n+2}(x, a_1, \dots, a_n, y) = -\epsilon Id.$$

$$\textit{Proof. } M_{n+2}(x, a_1, \dots, a_n, y) = \begin{pmatrix} K_{n+2}(x, a_1, \dots, a_n, y) & -K_{n+1}(a_1, \dots, a_n, y) \\ K_{n+1}(x, a_1, \dots, a_n) & -K_n(a_1, \dots, a_n) \end{pmatrix}.$$

We have the two following equalities :

- $K_{n+1}(x, a_1, \dots, a_n) = xK_n(a_1, \dots, a_n) - K_{n-1}(a_2, \dots, a_n) = \epsilon x - \epsilon x = 0_A$;
- $K_{n+1}(a_1, \dots, a_n, y) = yK_n(a_1, \dots, a_n) - K_{n-1}(a_1, \dots, a_{n-1}) = \epsilon y - \epsilon y = 0_A$.

Since $\det(M_{n+2}(x, a_1, \dots, a_n, y)) = 1_A$, $K_{n+2}(x, a_1, \dots, a_n, y) = -\epsilon$. So, $M_{n+2}(x, a_1, \dots, a_n, y) = -\epsilon Id$. \square

We have seen that $R' \subset R$ does not imply in general $\ell_{R'} \leq \ell_R$. However, this is true if we only consider commutative and unitary rings.

Proposition 2.16. *Let A, B two commutative and unitary rings such that A is a subring of B . $\ell_A \leq \ell_B$.*

Proof. Let $n := \ell_A$. It exists $(a_1, \dots, a_n) \in A^n$ an irreducible λ -quiddity over A . Since $A \subset B$, (a_1, \dots, a_n) is a solution of (E_B) . Suppose it is reducible over B . It exists $m, l \geq 3$, $(b_1, \dots, b_m) \in B^m$ and (c_1, \dots, c_l) a λ -quiddity over B such that

$$(a_1, \dots, a_n) \sim (b_1, \dots, b_m) \oplus (c_1, \dots, c_l) = (b_1 + c_l, b_2, \dots, b_{m-1}, b_m + c_1, c_2, \dots, c_{l-1}).$$

In particular, $(c_2, \dots, c_{l-1}) \in A^{l-2}$ and $(b_2, \dots, b_{m-1}) \in A^{m-2}$. Besides, it exists $\alpha \in \{\pm 1_B\} = \{\pm 1_A\}$ such that $M_l(c_1, \dots, c_l) = \alpha Id$. By proposition 2.14,

$$0_B = K_{l-1}(c_1, \dots, c_{l-1}) = c_1 K_{l-2}(c_2, \dots, c_{l-1}) - K_{l-3}(c_3, \dots, c_{l-1}) = -\alpha c_1 - K_{l-3}(c_3, \dots, c_{l-1}).$$

Hence, $c_1 = -\alpha K_{l-3}(c_3, \dots, c_{l-1})$. Since $(c_3, \dots, c_{l-1}) \in A^{l-3}$ and A is a ring, $c_1 \in A$. Similar arguments show $c_l \in A$. Hence, (c_1, \dots, c_l) is a λ -quiddity over A . Besides, $(b_1 + c_l, b_m + c_1) \in A^2$. Thus, $(b_1, b_m) \in A^2$. So, (a_1, \dots, a_n) is reducible over A . Since this is not the case, (a_1, \dots, a_n) is irreducible over B and $\ell_B \geq n$. \square

Remark. It exists rings A, B verifying $A \subset B$, $A \neq B$ and $\ell_A = \ell_B$. For instance, $\mathbb{Z} \subsetneq \mathbb{Z}[\pi]$ and $\ell_{\mathbb{Z}} = \ell_{\mathbb{Z}[\pi]} = 4$ ([9] Theorem 2.7).

3. LOWER AND UPPER BOUNDS FOR ℓ_A

In a section dedicated to the well-known “pigeon-hole principle”, M. Aigner and G. Ziegler say : “Some mathematical principles, [...], are so obvious that you might think they would only produce equally obvious results.” and, immediately after, they precise “It ain’t necessarily so” (see [1] page 131). Reading these few lines gave us the idea used in what follows and the proof developed below provides an example which illustrates Aigner and Ziegler’s words.

3.1. Proof of theorem 1.1. Let $(A, +, \times)$ a finite commutative and unitary ring different from $\{0_A\}$ and R a submagma of $(A, +)$.

i) Since A is finite, $\text{car}(A) = p \neq 0$. Let $x \in R$ ($R \neq \emptyset$). Since R is closed under $+$, $0_A = \sum_{i=1}^p x \in R$.

We set $l := |SL_2(A)|$, l is well defined since A is finite. Besides, $l \geq 4$. Let $n \in \mathbb{N}^*$ such that $n \geq l + 1 \geq 5$ and $(a_1, \dots, a_n) \in R^n$. We consider the matrices $M_k(a_1, \dots, a_k)$, with $1 \leq k \leq n$. We have $n > l$ objects belonging to a set that contains only l elements. Hence, by the pigeon-hole principle, it exists $1 \leq i < j \leq n$ such that :

$$M_i(a_1, \dots, a_i) = M_j(a_1, \dots, a_j) = M_{j-i}(a_{i+1}, \dots, a_j) M_i(a_1, \dots, a_i).$$

Since $M_i(a_1, \dots, a_i)$ is invertible ($\det(M_i(a_1, \dots, a_i)) = 1_A$), we have :

$$M_{j-i}(a_{i+1}, \dots, a_j) = Id.$$

So, (a_{i+1}, \dots, a_j) is a solution of (E_R) whose size is $1 \leq j - i \leq n - 1$. We consider three cases :

— If $j - i \geq 3$ then we have following equality :

$$(a_1, \dots, a_n) \sim (a_j, \dots, a_n, a_1, \dots, a_{j-1}) = \underbrace{(0_A, a_{j+1}, \dots, a_n, a_1, \dots, a_i, 0_A)}_{n-(j-i)+2 \geq 3} \oplus (a_{i+1}, \dots, a_j).$$

$$(\text{if } j = n \text{ then } (a_1, \dots, a_n) \sim (a_n, a_1, \dots, a_{n-1}) = (0_A, a_1, \dots, a_i, 0_A) \oplus (a_{i+1}, \dots, a_n).)$$

— If $j - i = 2$ then, by lemma 2.9, $a_{i+1} = a_{i+2} = 0_A$ and we have :

$$(a_1, \dots, a_n) \sim (a_{i+3}, \dots, a_n, a_1, \dots, a_{i+2}) = \underbrace{(a_{i+3}, \dots, a_n, a_1, \dots, a_i)}_{n-2 \geq 3} \oplus (0_A, 0_A, 0_A, 0_A).$$

$$(\text{if } i = n - 2 \text{ then } (a_1, \dots, a_n) = (a_1, \dots, a_{n-2}) \oplus (0_A, 0_A, 0_A, 0_A).)$$

— By lemma 2.9, the case $j - i = 1$ is impossible.

Hence, we can written (a_1, \dots, a_n) as a sum of an u -tuple with a solution of (E_R) of size v with $u, v \geq 3$. So, a λ -quiddity whose size is greater or equal to $l + 1$ is reducible. Since R is finite (as a subpart of a finite set), we conclude that the number of irreducible λ -quiddities over R is finite.

ii) We set $H := \left\{ \begin{pmatrix} 1_A & x \\ 0_A & 1_A \end{pmatrix}, \begin{pmatrix} -1_A & x \\ 0_A & -1_A \end{pmatrix}, x \in A \right\}$. H is a subgroup of $SL_2(A)$. We set $(SL_2(A)/H)_d$ the set of the right cosets of H and

$$l := |(SL_2(A)/H)_d| = \frac{|SL_2(A)|}{|H|} = \begin{cases} \frac{|SL_2(A)|}{|A|} & \text{if } \text{car}(A) = 2; \\ \frac{|SL_2(A)|}{2|A|} & \text{if } \text{car}(A) \neq 2. \end{cases}$$

Let $n \in \mathbb{N}^*$ such that $n \geq l + 3$ and $(a_1, \dots, a_n) \in A^n$. We consider the right cosets $H M_k(a_1, \dots, a_k)$, with $1 \leq k \leq n - 2$. We have $n - 2 > l$ objects belonging to a set that contains only l elements. Hence, by the pigeon-hole principle, it exists $1 \leq i < j \leq n - 2$ such that :

$$H M_i(a_1, \dots, a_i) = H M_j(a_1, \dots, a_j).$$

So, $M = M_j(a_1, \dots, a_j)(M_i(a_1, \dots, a_i))^{-1} \in H$. However, we have the following equalities :

$$\begin{aligned} M &= M_j(a_1, \dots, a_j)(M_i(a_1, \dots, a_i))^{-1} \\ &= M_{j-i}(a_{i+1}, \dots, a_j) M_i(a_1, \dots, a_i)(M_i(a_1, \dots, a_i))^{-1} \\ &= M_{j-i}(a_{i+1}, \dots, a_j). \end{aligned}$$

Hence, $M_{j-i}(a_{i+1}, \dots, a_j) \in H$. Thus, by proposition 2.14, $K_{j-i}(a_{i+1}, \dots, a_j) = \epsilon = \pm 1_A$. We set $x = \epsilon K_{j-i-1}(a_{i+2}, \dots, a_j) \in A$ and $y = \epsilon K_{j-i-1}(a_{i+1}, \dots, a_{j-1}) \in A$. By lemma 2.15, $(x, a_{i+1}, \dots, a_j, y)$ is a solution of (E_A) . Its size is equal to $3 \leq j - i + 2 \leq n - 1$. We have the following equality :

$$(a_1, \dots, a_n) \sim (a_{j+1}, \dots, a_n, a_1, \dots, a_j) = (a_{j+1} - y, \dots, a_n, a_1, \dots, a_i - x) \oplus (x, a_{i+1}, \dots, a_j, y).$$

Hence, we can written (a_1, \dots, a_n) as a sum of an u -tuple with a solution of (E_A) of size v with $u, v \geq 3$. So, a λ -quiddity over A whose size is greater or equal to $l + 3$ is reducible. Hence, $\ell_A \leq l + 2$.

By lemma 2.9, $(0_A, 0_A, 0_A, 0_A)$ is an irreducible λ -quiddity over A . So $\ell_A \geq 4$. Since A is finite, $\text{car}(A) > 0$. We suppose $\text{car}(A) \geq 4$. To obtain the majoration $\ell_A \geq \text{car}(A)$, we will adapt the proof of

theorem 2.6 of [8]. For all $m \in \mathbb{N}$, we set $m_A = m \times 1_A$.

By induction, we have, for all $m \in \mathbb{N}$, the following equality : $K_m(2_A, \dots, 2_A) = (m+1)_A$. Indeed, $K_0 = 1_A$ and $K_1(2_A) = 2_A$ and if we suppose it exists $m \geq 1$ such that $K_m(2_A, \dots, 2_A) = (m+1)_A$ and $K_{m-1}(2_A, \dots, 2_A) = m_A$ then

$$K_{m+1}(2_A, \dots, 2_A) = 2_A K_m(2_A, \dots, 2_A) - K_{m-1}(2_A, \dots, 2_A) = (2m+2)_A - m_A = (m+2)_A.$$

By induction, we obtain the desired formula.

So, by proposition 2.14, $M_{\text{car}(A)}(2_A, \dots, 2_A) = Id$. Suppose this λ -quiddity is reducible. It exists $u, v \geq 3$, $(b_1, \dots, b_u) \in A^u$ and (c_1, \dots, c_v) a λ -quiddity over A such that :

$$(2_A, \dots, 2_A) = (b_1, \dots, b_u) \oplus (c_1, \dots, c_v) = (b_1 + c_v, b_2, \dots, b_{u-1}, b_u + c_1, c_2, \dots, c_{v-1}).$$

Hence, $c_2 = \dots = c_{v-1} = 2_A$. So, by proposition 2.14, we have :

$$\pm 1_A = K_{v-2}(c_2, \dots, c_{v-1}) = K_{v-2}(2_A, \dots, 2_A) = v_A - 1_A.$$

Thus, $\text{car}(A)$ divides v or $(v-2)$. These two cases are impossible since $3 \leq v \leq \text{car}(A) + 2 - u$.

Hence, $(2_A, \dots, 2_A) \in A^{\text{car}(A)}$ is an irreducible solution of (E_A) and so $\ell_A \geq \text{car}(A)$. □

3.2. Applications. If we compile the results of theorem 1.1 and the cardinalities recalled in theorems 2.11 and 2.13, we obtain corollary 1.2. Besides, we can modify the proof of theorem 1.1 i) to have the following result :

Proposition 3.1. *Let A a finite commutative and unitary ring and $x \in \{-1_A, 0_A, 1_A\}$. It exists $k_x \in \mathbb{N}^*$ such that for all $n \geq k_x$ and for all n -tuple $(a_1, \dots, a_n) \in A^n$, it exists $1 \leq i \leq n$ and $1 \leq j \leq n - i + 1$ such that $K_i(a_j, \dots, a_{j+i-1}) = x$.*

Proof. We set $l := |SL_2(A)|$, l is well defined since A is finite. Let $n \in \mathbb{N}^*$ such that $n \geq 3l + 1$ and $(a_1, \dots, a_n) \in A^n$. We consider the matrices $M_k(a_1, \dots, a_{3k+1})$, with $0 \leq k \leq l$. We have $l + 1 > l$ objects belonging to a set that contains only l elements. Hence, by the pigeon-hole principle, it exists $0 \leq i < j \leq l$ such that :

$$M_{3i+1}(a_1, \dots, a_{3i+1}) = M_{3j+1}(a_1, \dots, a_{3j+1}) = M_{3(j-i)}(a_{3i+2}, \dots, a_{3j+1}) M_{3i+1}(a_1, \dots, a_{3i+1}).$$

Since, $M_{3i+1}(a_1, \dots, a_{3i+1})$ is invertible, we have :

$$M_{3(j-i)}(a_{3i+2}, \dots, a_{3j+1}) = Id.$$

So, by proposition 2.14, we have :

- $K_{3(j-i)}(a_{3i+2}, \dots, a_{3j+1}) = 1_A$;
- $K_{3(j-i)-2}(a_{3i+3}, \dots, a_{3j}) = -1_A$;
- $K_{3(j-i)-1}(a_{3i+2}, \dots, a_{3j}) = 0_A$.

Besides, $1 \leq 3(j-i) - 2$ and $3(j-i) \leq n$. □

Remark. This result cannot be extended to another element of a finite commutative and unitary ring A . For instance, we can consider, for all $n \in \mathbb{N}^*$, $(0_A, \dots, 0_A) \in A^n$. Indeed, we have :

- if $i \equiv \pm 1[4]$, $K_i(0_A, \dots, 0_A) = 0_A$;
- if $i \equiv 2[4]$, $K_i(0_A, \dots, 0_A) = -1_A$;
- if $i \equiv 0[4]$, $K_i(0_A, \dots, 0_A) = 1_A$.

In [11] section 3.3, we have introduced the following open problem :

Problem. *Let I a set. Find necessary and sufficient conditions on I such that there is a finite number of irreducible λ -quiddities over $\prod_{i \in I} \mathbb{Z}/2\mathbb{Z}$.*

Theorem 1.1 give us a sufficient condition. Indeed, if I is a finite set, $A = \prod_{i \in I} \mathbb{Z}/2\mathbb{Z}$ is a finite ring and so theorem 1.1 guarantees that the number of irreducible λ -quiddities over A is finite. Besides, thanks to proposition 2.12, we have the following inequality :

$$\ell_{\prod_{i=1}^r \mathbb{Z}/2\mathbb{Z}} \leq \frac{1}{2^r} \left(\prod_{i=1}^r 6 \right) + 2 = 3^r + 2.$$

We now give some numerical applications of theorem 1.1. In the table below, we give the exact value of ℓ_A obtained with a computer (see [6] section 5.2) and the value of $\theta_A = \begin{cases} \frac{|SL_2(A)|}{|A|} + 2 & \text{if } \text{car}(A) = 2; \\ \frac{|SL_2(A)|}{2|A|} + 2 & \text{if } \text{car}(A) \neq 2. \end{cases}$

A	$\mathbb{Z}/7\mathbb{Z}$	$\mathbb{Z}/8\mathbb{Z}$	$\mathbb{Z}/9\mathbb{Z}$	$\mathbb{Z}/10\mathbb{Z}$	$\mathbb{Z}/11\mathbb{Z}$	$\mathbb{Z}/12\mathbb{Z}$	$\mathbb{Z}/13\mathbb{Z}$	$\mathbb{Z}/14\mathbb{Z}$	$\mathbb{Z}/15\mathbb{Z}$	$\mathbb{Z}/16\mathbb{Z}$
ℓ_A	9	8	12	12	19	15	25	20	26	24
θ_A	26	26	38	38	62	50	86	74	98	98

A	\mathbb{F}_8	$\mathbb{Z}/2\mathbb{Z} \times \mathbb{F}_4$	$\mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$	$\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}$	\mathbb{F}_9	\mathbb{F}_{16}	$\mathbb{Z}/17\mathbb{Z}$	$\mathbb{Z}/18\mathbb{Z}$	\mathbb{F}_{25}	\mathbb{F}_{32}
θ_A	65	47	29	34	42	257	146	110	314	1025

Note that $\ell_{\mathbb{Z}/3\mathbb{Z} \times \mathbb{Z}/3\mathbb{Z}} = 12$ ([11] section 4.1.3).

3.3. A refinement of theorem 1.1 in the case $A = \mathbb{F}_9$. The aim of this section is to find a better upper bound for $\ell_{\mathbb{F}_9}$. In the previous subpart, we have shown that $\ell_{\mathbb{F}_9} \leq 42$. Now, we will prove $\ell_{\mathbb{F}_9} \leq 32$. To do this, we will follow the proof of theorem 1.1 ii) but we will choose another subgroup.

Proposition 3.2. $\ell_{\mathbb{F}_9} \leq 32$.

Proof. $H := SL_2(\mathbb{F}_3)$ is a subgroup of $SL_2(\mathbb{F}_9)$. We set $(SL_2(\mathbb{F}_9)/H)_d$ the set of the right cosets of H and $l := |(SL_2(\mathbb{F}_9)/SL_2(\mathbb{F}_3))_d| = 30$.

Let $n \in \mathbb{N}^*$ and $(a_1, \dots, a_n) \in \mathbb{F}_9^n$ such that $n \geq l + 3$ and (a_1, \dots, a_n) is a λ -quiddity. We consider the right cosets $H M_k(a_1, \dots, a_k)$, with $1 \leq k \leq n - 2$. We have $n - 2 > l$ objects belonging to a set that contains only l elements. Hence, by the pigeon-hole principle, it exists $1 \leq i < j \leq n - 2$ such that :

$$H M_i(a_1, \dots, a_i) = H M_j(a_1, \dots, a_j).$$

So, $M = M_j(a_1, \dots, a_j)(M_i(a_1, \dots, a_i))^{-1} \in H$, that is to say $M_{j-i}(a_{i+1}, \dots, a_j) \in H$.

If $j - i = 1$ then $a_{i+1} \in \mathbb{F}_3 = \{\overline{0}, \overline{1}, \overline{-1}\}$. By lemma 2.9, (a_1, \dots, a_n) is reducible.

Now, we suppose $j - i \geq 2$. The elements of H cannot have three coefficients equal to $\overline{0}$. Thus $K_{j-i}(a_{i+1}, \dots, a_j) = \pm \overline{1}$ or $K_{j-i-1}(a_{i+2}, \dots, a_j) = \pm \overline{1}$ or $K_{j-i-1}(a_{i+1}, \dots, a_{j-1}) = \pm \overline{1}$.

We suppose $K_{j-i}(a_{i+1}, \dots, a_j) = \epsilon = \pm \overline{1}$.

We set $x = \epsilon K_{j-i-1}(a_{i+2}, \dots, a_j) \in \mathbb{F}_9$ and $y = \epsilon K_{j-i-1}(a_{i+1}, \dots, a_{j-1}) \in \mathbb{F}_9$. By lemma 2.15, $(x, a_{i+1}, \dots, a_j, y)$ is a solution of $(E_{\mathbb{F}_9})$. Its size is equal to $4 \leq j - i + 2 \leq n - 1$. We have the following equality :

$$(a_1, \dots, a_n) \sim (a_{j+1}, \dots, a_n, a_1, \dots, a_j) = \underbrace{(a_{j+1} - y, \dots, a_n, a_1, \dots, a_i - x)}_{n-(j-i) \geq 3} \oplus (x, a_{i+1}, \dots, a_j, y).$$

Hence, (a_1, \dots, a_n) is reducible.

We suppose $K_{j-i-1}(a_{i+1}, \dots, a_{j-1}) = \epsilon = \pm \overline{1}$.

We set $x = \epsilon K_{j-i-2}(a_{i+2}, \dots, a_{j-1}) \in \mathbb{F}_9$ and $y = \epsilon K_{j-i-2}(a_{i+1}, \dots, a_{j-2}) \in \mathbb{F}_9$. By lemma 2.15, $(x, a_{i+1}, \dots, a_{j-1}, y)$ is a solution of $(E_{\mathbb{F}_9})$. Its size is equal to $3 \leq j - i + 1 \leq n - 2$. We have the following equality :

$$(a_1, \dots, a_n) \sim (a_j, \dots, a_n, a_1, \dots, a_{j-1}) = \underbrace{(a_j - y, \dots, a_n, a_1, \dots, a_i - x)}_{n-(j-i)+1 \geq 4} \oplus (x, a_{i+1}, \dots, a_{j-1}, y).$$

Hence, (a_1, \dots, a_n) is reducible.

If $K_{j-i-1}(a_{i+2}, \dots, a_j) = \epsilon = \pm 1$, the proof is the same as above.

Hence, $\ell_A \leq l + 2 = 32$.

□

3.4. A still open problem. In most situations, we consider irreducible λ -quiddities over infinite subgroups (and even more precisely subgroups of $(\mathbb{C}, +)$) or over finite rings. For the first case, there are a lot of possible situations and generally we have to consider each subgroup separately. For the second case, theorem 1.1 gives us a precise answer to the problem of the finiteness of the number of irreducible solutions. Hence, we could consider that the general question of the finiteness of the number of irreducible λ -quiddities over a fixed set is closed. However, there is still a case that deserves our attention. Indeed, theorem 1.1 does not solve the problem of the finiteness of the number of irreducible λ -quiddities over a finite submagma $R \neq \{0_A\}$ of an infinite ring A (note that this case implies that $\text{car}(A) \neq 0$). Obviously, if it exists a finite subring B of A containing R then theorem 1.1 give us the solution, but we can find several situations in which it is impossible to do such a reduction.

For instance, we choose two positive integers N and k , with $N \geq 2$. We consider the two classical following sets $A := (\mathbb{Z}/N\mathbb{Z})[X]$ and $R := (\mathbb{Z}/N\mathbb{Z})_k[X] := \{P \in A, \deg(P) \leq k\}$. A is an infinite ring and R is a finite submagma of A . Besides, it does not exist a finite subring B of A containing R . Indeed, a ring containing R necessarily contains X^l for all $l \in \mathbb{N}$.

In some cases, we can easily conclude. Indeed, consider for instance the sets $A := (\mathbb{Z}/2\mathbb{Z})[X]$ and $R := \{\bar{0}, X\}$. A and R verify the desired conditions. For all $n \in \mathbb{N}^*$, $(X, \dots, X) \in R^n$ is not a λ -quiddity over R (since $\deg(K_n(X, \dots, X)) = n$). Hence, a λ -quiddity over R necessarily contains $\bar{0}$. Thus, the only irreducible λ -quiddities over R are $(\bar{0}, \bar{0}, \bar{0}, \bar{0})$, $(\bar{0}, X, \bar{0}, X)$ and $(X, \bar{0}, X, \bar{0})$. However, the general following problem remains open.

Problem. *Let R a finite submagma of an infinite ring. Do we have $\ell_R < +\infty$? If the answer is no, can we characterise the submagma R for which $\ell_R < +\infty$?*

REFERENCES

- [1] M. Aigner, G. Ziegler, *Proofs from THE BOOK*, Second edition, Springer-Verlag Berlin Heidelberg GmbH, 2001.
- [2] C. Conley, V. Ovsienko, *Quiddities of polygon dissections and the Conway-Coxeter frieze equation*, Annali della Scuola Normale Superiore di Pisa, Vol. 24 no. 4, (2023), pp 2125-2170.
- [3] H. S. M. Coxeter, *Frieze patterns*, Acta Arithmetica, Vol. 18, (1971), pp 297-310.
- [4] M. Cuntz, *A combinatorial model for tame frieze patterns*, Munster J. Math., Vol. 12 no. 1, (2019), pp 49-56.
- [5] M. Cuntz, T. Holm, *Frieze patterns over integers and other subsets of the complex numbers*, J. Comb. Algebra., Vol. 3 no. 2, (2019), pp 153-188.
- [6] M. Cuntz, F. Mabilat, *Comptage des quiddités sur les corps finis et sur quelques anneaux $\mathbb{Z}/N\mathbb{Z}$* , Annales de la Faculté des Sciences de Toulouse, to appear, arXiv:2304.03071, hal-04057675.
- [7] S. Francinou, H. Gianella, S. Nicolas, *Exercices de mathématiques-Oraux X-ENS, algèbre 2*, Cassini, Enseignement Des Mathématiques, 2009.
- [8] F. Mabilat, *Combinatoire des sous-groupes de congruence du groupe modulaire*, Annales Mathématiques Blaise Pascal, Vol. 28 no. 1, (2021), pp. 7-43. doi : 10.5802/ambp.398. <https://ambp.centre-mersenne.org/articles/10.5802/ambp.398/>.
- [9] F. Mabilat, *λ -quiddité sur $\mathbb{Z}[\alpha]$ avec α transcendant*, Mathematica Scandinavica, Vol. 128 no. 1, (2022), pp 5-13, <https://doi.org/10.7146/math.scand.a-128972>.
- [10] F. Mabilat, *Combinatoire des sous-groupes de congruence : des fractions continues à la description des sous-groupes de congruence*, Phd thesis, Université de Reims Champagne-Ardenne, (2022), <https://theses.fr/2022REIMS052>.
- [11] F. Mabilat, *λ -quiddités sur des produits directs d'anneaux*, hal-04190487, arXiv:2308.15848.

- [12] F. Mabilat, *Some counting formulas for λ -quiddities over the rings $\mathbb{Z}/2^m\mathbb{Z}$* , Bulletin of the Australian Mathematical Society, to appear, arXiv:2402.09968.
- [13] S. Morier-Genoud, *Coxeter's frieze patterns at the crossroad of algebra, geometry and combinatorics*, Bull. Lond. Math. Soc., Vol. 47 no. 6, (2015), pp 895-938.
- [14] S. Morier-Genoud, *Counting Coxeter's friezes over a finite field via moduli spaces*, Algebraic combinatoric, Vol. 4 no. 2, (2021), pp 225-240.
- [15] V. Ovsienko, *Partitions of unity in $SL(2, \mathbb{Z})$, negative continued fractions, and dissections of polygons*, Res. Math. Sci., Vol. 5 no. 2, (2018), Article 21, 25 pp.
- [16] M. Weber, M. Zhao, *Factorization of frieze patterns*, Revista de la Unión Matemática Argentina, Vol. 60 no. 2, (2019), pp 407-415.

Email address: `flavien.mabilat@univ-reims.fr`