

# Stable properties under weakly geometrically flat maps

Daniel Barlet\* and Jón Magnússon†

May 1, 2024

**ABSTRACT.** In this note we show that a weakly geometrically flat map  $\pi: M \rightarrow N$  between pure dimensional complex spaces has the local lifting property for cycles. From this result we also deduce that, under these hypotheses, several properties of  $M$  are transferred to  $N$ .

## 1 Lifting cycles locally

An equidimensional holomorphic map  $\pi: M \rightarrow N$  between two pure dimensional complex spaces with  $q := \dim M - \dim N$ , is called **weakly geometrically flat** (or geometrically flat in the weak sense) if it is surjective and there exists an analytic family  $(F_y)_{y \in N}$  of  $q$ -cycles in  $M$  such that, for any  $y \in N$  we have the equality  $|F_y| = \pi^{-1}(y)$ .

If, moreover, for very general  $y \in N$  the cycle  $F_y$  is a reduced cycle we say that  $\pi$  is **geometrically flat** (or geometrically flat in the strong sense if we want to avoid confusion).

**REMARKS.**

- (i) Weak geometric flatness is stable by base change  $f: P \rightarrow N$ , where  $f$  is a holomorphic map and  $P$  is a pure dimensional reduced complex space. On the contrary, this is not true, in general, for strong geometric flatness when  $P$  has at least one irreducible component whose image has empty interior in  $N$ .
- (ii) When  $N$  is normal and  $M, N$  are pure dimensional, every equidimensional holomorphic map  $\pi: M \rightarrow N$  is geometrically flat (in the strong sense).

---

\*Institut Elie Cartan, Géométrie,  
Université de Lorraine, CNRS UMR 7502 and Institut Universitaire de France.

†Department of Mathematics, School of Engineering and Physical Sciences, University of Iceland.

**Proposition 1.0.1** *Let  $\pi: M \rightarrow N$  be a geometrically flat holomorphic map in the weak sense between reduced complex spaces of pure dimensions  $m$  and  $n$  and  $z_0$  be a point in  $M$ . Then there exists an open connected neighborhood  $N_0$  of  $\pi(z_0)$  in  $N$  and a closed analytic subset  $Z$  of  $\pi^{-1}(N_0)$  such that  $\pi^{-1}(\pi(z_0)) \cap Z = \{z_0\}$  and  $\pi$  induces a proper and finite map  $\pi_0: Z \rightarrow N_0$ , which is geometrically flat in the weak sense.*

PROOF. Put  $y_0 := \pi(z_0)$ ,  $q := m - n$  and let  $(F_y)_{y \in N}$  be the analytic family of fiber-cycles of  $\pi$ . Let  $E = (U, B, j)$  be a  $q$ -scale on  $M$ , adapted to  $F_{y_0}$  and centered at  $z_0$ . More precisely,  $j$  is a proper holomorphic embedding of an open neighborhood  $M_0$  of  $z_0$  in  $M$  into an open neighborhood  $W$  of the origin in  $\mathbb{C}^l$  and  $U, B$  are open relatively compact polydisks in  $\mathbb{C}^q, \mathbb{C}^{l-q}$  such that  $\bar{U} \times \bar{B} \subset W$  and

$$j(F_{y_0} \cap M_0) \cap (\bar{U} \times \partial B) = \emptyset.$$

Moreover,  $E$  can be chosen in such a way that  $j(F_{y_0} \cap M_0) \cap (\{0\} \times \bar{B}) = \{(0, 0)\}$ . Then there exists an open neighborhood  $N_0$  of  $y_0$  in  $\pi(M_0)$  such that, for all  $y$  in  $N_0$ , the scale  $E$  is adapted to  $F_y$  and  $\deg_E(F_y) = \deg_E(F_{y_0})$ .

Put  $M' := M_0 \cap \pi^{-1}(N_0)$ . Then  $j(M') = (\pi \circ j^{-1})^{-1}(N_0)$  is an analytic subset of  $W$  which satisfies  $j(M') \cap (\bar{U} \times \partial B) = \emptyset$ . Hence, for all  $x$  in  $U$ ,  $j(M') \cap (\{x\} \times B)$  is an analytic subset of  $W$  and consequently  $Z_x := j^{-1}((\{x\} \times B) \cap j(M'))$  is an analytic subset of  $\pi^{-1}(N_0)$ . Furthermore, the induced map  $\pi_x: Z_x \rightarrow N_0$  is proper, surjective and with finite fibers and we may assume that  $Z_{x_0} \cap \pi^{-1}(y_0) = \{z_0\}$ . For every  $y$  in  $N_0$ , denote  $X_y$  the multigraph that  $F_y$  determines in  $U \times B$  and put  $A_y := j^*(X_y \cap U \times B) \cap (\{x\} \times B)$ .

Then  $(A_y)_{y \in N_0}$  is an analytic family of 0-cycles in  $Z_x$  such that  $|A_y| = \pi_x^{-1}(y)$  and it follows that  $Z := Z_0$  satisfies the required properties.  $\blacksquare$

REMARKS.

- (i) Under the assumptions of the proposition assume that  $\pi$  is geometrically flat in the strong sense. Then for very general  $y$  in  $N_0$ , the cycle  $F_y$  is reduced. So the map  $\pi_0: Z_0 \rightarrow N_0$  is geometrically flat in the strong sense (after shrinking  $N_0$  if necessary).
- (ii) Let  $\Sigma$  be an analytic subset of codimension  $r$  in  $M$ . Then  $j(\Sigma \cap M')$  is an analytic subset of codimension at most equal to  $r$  in  $j(M')$  and it follows that, for very general  $x$  in  $U$ , we have  $\text{codim}_{Z_x}(\Sigma \cap Z_x) = r$  (or empty). So we may choose  $x_0 \in U$  such that  $Z_{x_0} \cap \Sigma$  has codimension at most equal to  $r$  in  $Z_{x_0}$ .  $\square$

**Corollary 1.0.2** *Let  $\pi: M \rightarrow N$  be a geometrically flat map in the weak sense between pure dimensional complex spaces,  $Y$  be a rational  $p$ -cycle in  $N$  and  $y_0$  be a point in  $|Y|$ . Then, for every  $x$  in  $\pi^{-1}(y_0)$ , there exists an open neighborhood  $N_1$  of  $y_0$  in  $N$  and a rational  $p$ -cycle  $X$  in  $\pi^{-1}(N_1)$  such that  $x \in |X|$  and  $\pi_* X = Y \cap N_1$ .*

PROOF. We may assume that  $Y$  is an irreducible analytic subset of  $N$ . Then the induced map  $\pi^{-1}(Y) \rightarrow Y$  is geometrically flat in the weak sense so it is enough to consider the case where  $Y = N$ . Let  $z_0$  be a fixed point in  $\pi^{-1}(y_0)$ .

Then, by Proposition 1.0.1, there exists an open connected neighborhood  $N_0$  of  $y_0$  in  $N$  and an analytic subset  $Z$  of  $\pi^{-1}(N_0)$  such that  $\pi$  induces a proper and finite map  $\pi_0: Z \rightarrow N_0$  which is geometrically flat in the weak sense and such that (set-theoretically)  $\pi^{-1}(z_0) \cap Z = \{z_0\}$ . As the map  $\pi_0$  is both proper and open and  $N_0$  is connected,  $\pi_0$  is surjective and induces a surjective map from every irreducible component of  $Z_0$  onto an irreducible component of  $N_0$ . Let  $N'_0$  be the union of those irreducible components of  $N_0$  which do not contain  $y_0$ . Thus  $N_1 = N_0 \setminus N'_0$  is an open connected neighborhood of  $y_0$  which has only finitely many irreducible components  $C_1, \dots, C_l$ . For every  $j \in \{1, \dots, l\}$ , we choose an irreducible component  $Z_j$  of  $\pi^{-1}(N_1)$  such that  $\pi(Z_j) = C_j$  and denote  $k_j$  the degree of the induced map  $Z_j \rightarrow C_j$ . Then the rational cycle

$$Y := \frac{1}{k_1} Z_1 + \dots + \frac{1}{k_l} Z_l$$

has the desired properties. ■

## 2 The sheaf $L^\bullet$

We recall here some classical equivalent definitions of  $L^2$ -holomorphic forms on a normal complex space  $M$ .

Remember that  $M$  has **weakly rational singularities** if and only if  $\omega_M^n = L_M^n$  where  $M$  has pure dimension  $n$ . Thanks to [4] (see also [5]) this implies that  $\omega_M^p = L_M^p$  for each  $p \in [0, n]$ . If, moreover,  $M$  is Cohen-Macaulay, then  $M$  has rational singularities.

The following well known lemma gives two equivalent definitions of these (coherent) sheaves.

**Lemma 2.0.1** *Let  $M$  be a normal complex space and denote  $S(M)$  its singular locus. Then the following properties are equivalent for a holomorphic  $p$ -form  $\alpha$  defined on  $M \setminus S(M)$ :*

1. *Let  $\Sigma$  is a closed analytic subset of codimension  $\geq 2$  in  $M$  containing the singular set  $S(M)$  in  $M$ . Then for every open set  $M'$  in  $M$  and any proper holomorphic embedding  $f: X \rightarrow M'$  of a  $p$ -dimensional reduced complex space such that  $f(X) \not\subset \Sigma$ , the integral  $\int_{K \setminus f^{-1}(\Sigma)} f^*(\alpha) \wedge \overline{f^*(\alpha)}$  is finite for any compact set  $K$  in  $X$ .*
2. *There exists a desingularization  $\tau: \tilde{M} \rightarrow M$  such that  $\tau^*(\alpha)$  extends as a holomorphic  $p$ -form on  $\tilde{M}$ .*

PROOF. First assume that property 2. holds true and consider a proper holomorphic map  $f: X \rightarrow M'$  such that  $f(X) \not\subset \Sigma$ . Remark first that we may assume that  $\Sigma = S(M)$  since on the smooth part of  $M$  the sheaf  $\Omega_M^p$  has the extension property in codimension 2. Let  $\tilde{X} \rightarrow \tilde{M}'$  be the strict transform of  $\tau$  by  $f$  (so  $\tilde{X} := X \times_{M, \text{strict}} \tilde{M}$ ) and let  $\tilde{f}$  be the canonical projection. Then we have

$$\int_{pr^{-1}(K)} \tilde{f}^*(\tau^*(\alpha)) \wedge \overline{\tilde{f}^*(\tau^*(\alpha))} < +\infty$$

and this integral is equal to  $\int_K f^*(\alpha) \wedge \overline{f^*(\alpha)}$  since the projection  $pr: \tilde{X} \rightarrow X$  is a (proper) modification with center in  $f^{-1}(S(M))$ , for any compact  $K$  in  $X$ .

Conversely, if the property 1. holds true, consider a desingularization  $\tau: \tilde{M} \rightarrow M$  and take a smooth generic point  $x_0$  of an irreducible component of the exceptional set  $E$  in  $\tilde{M}$  (the preimage of the center of the modification  $\tau$ ). Then let  $g: X \rightarrow \tilde{M}$  be a  $p$ -dimensional complex submanifold of a neighborhood of  $x_0$  which is transversal to  $E^1$  at the point  $x_0$ . Then let  $f := g \circ \tau$ . The property 1. gives the fact that  $g^*(\tau^*(\alpha))$  cannot have a pole along  $E$ , since  $x_0$  is generic, and then  $\tau^*(\alpha)$  extends holomorphically across  $E$ , using again the codimension 2 extension property of the sheaf  $\Omega_M^p$  since  $\tilde{M}$  is a complex manifold.  $\blacksquare$

REMARK. Let  $Z$  be an analytic subset of an open subset  $M'$  of  $M$ , such that  $Z \cap S(M)$  is of codimension at least 2 in  $Z$ , and  $j: Z \hookrightarrow M'$  be the canonical injection. Then, due to characterisation 1. above, we have a restriction map

$$j^*(L_{M'}^p) \rightarrow L_Z^p$$

extending the pull-back morphism of holomorphic  $p$ -forms, for any  $p \geq 0$ .

### 3 The Theorem

**Theorem 3.0.1** *Let  $\pi: M \rightarrow N$  be a geometrically flat map in the weak sense between irreducible complex spaces. Assume that  $M$  has one of the following properties:*

*P.1 The complex space  $M$  is normal.*

*P.2 The complex space  $M$  is normal and every codimension 1 cycle in  $M$  is  $\mathbb{Q}$ -Cartier.*

*P.3 The complex space  $M$  is normal and every holomorphic  $p$ -form on  $M \setminus S(M)$  extends as a section of the  $L_M^p$ -sheaf (so  $L_M^p = \omega_M^p$ ).*

---

<sup>1</sup>Note that if  $E$  has codimension  $\geq 2$  the proof is complete since  $\tau^*(\alpha)$  extends holomorphically across  $E$ .

*P.4 The complex space  $M$  is normal and the de Rham complex  $0 \rightarrow \mathbb{C} \rightarrow (\omega_M^\bullet, d^\bullet)$  is exact in degree  $p$ .*

*P.5 The complex space  $M$  is normal and the de Rham complex  $0 \rightarrow \mathbb{C} \rightarrow (L_M^\bullet, d^\bullet)$  is locally exact in degree  $p$  on  $M$ .*

*Then the same property is also true for  $N$ .*

PROOF. As all five properties are local we may assume, due to Proposition 1.0.1 and Remark (ii) following it, that there exists an analytic subset  $Z$  of dimension  $n$  in  $M$  satisfying the following conditions:

- The restriction  $\pi|_Z$  is surjective, proper and geometrically flat in the weak sense.
- $S(M) \cap Z$  is of codimension at least 2. in  $Z$

Denote  $k$  the degree of  $\pi' := \pi|_Z$ .

Let us begin by property *P.1*. Take a locally bounded meromorphic function  $f$  in an open neighborhood  $N'$  of  $y_0 \in N$ . Then  $\pi^*(f)$  is holomorphic on  $\pi^{-1}(N')$  and induces a holomorphic function  $g$  on  $\pi'^{-1}(N')$ . Then the trace of  $g$  by the induced map  $\pi'^{-1}(N') \rightarrow N'$  is holomorphic and equal to  $kf$ .

Proof of *P.2* for  $N$ . Assume that  $X$  is a cycle of codimension 1 in  $N$  and  $y$  a point in  $N$ . Then  $\pi^*(X)$  is a cycle of codimension 1 in  $M$  and we may find an open neighborhood  $U$  in  $M$  of  $\pi^{-1}(y) \cap Z$  and a holomorphic function  $f$  defining the cycle  $\pi^*(X)$  in  $U$ . As  $\pi'$  is a proper map, there exists an open neighborhood  $N'$  of  $y$  such that  $\pi'^{-1}(N') \subset U \cap Z$ . Put  $Z' := \pi'^{-1}(N')$  and denote  $\pi'': Z' \rightarrow N'$  the restriction of  $\pi'$ . Then  $f|_{Z'}$  defines the cycle  $(\pi'')^*(X)$ . Since the norm of the function  $f|_{Z'}$  is meromorphic and locally bounded on  $N'$ , which is normal, it is holomorphic on  $N'$  and defines the cycle  $k(X \cap N')$ . Hence  $X$  is locally  $\mathbb{Q}$ -Cartier in  $N$ .

To prove property *P.3* for  $N$ , consider a  $p$ -form  $\alpha$  on  $N \setminus S(N)$ . Then  $\pi^*(\alpha)$  extends as a global section of the sheaf  $L_M^p$ . Since  $Z$  meets  $S(M)$  in codimension at least 2 the form  $\pi'^*(\alpha)$  is holomorphic on  $Z \setminus S(M)$  and is a section of  $L_Z^p$  due to the characterization 1. in Lemma 2.0.1. Then, since the map  $\pi'$  is a proper map with finite fibers, using again the characterization 1 above with  $\Sigma = \pi(S(M) \cap Z) \cap S(N)$  we conclude that  $\alpha$  is a section of  $L_N^p$  because the normality of  $M$  implies the normality of  $N$  (see P.1), and  $S(N)$  has codimension at least 2 in  $N$ . ■

To prove property *P.4* for  $N$  we take an open neighborhood  $N'$  of a point  $y$  in  $N$  and we consider  $\alpha \in \Gamma(N', \omega_N^p)$  such that  $d\alpha = 0$ . Then the restriction of  $\alpha$  to  $N' \setminus S(N)$  is a closed holomorphic  $p$ -form. Then  $\pi^*(\alpha)$  is also a closed holomorphic  $p$ -form on  $\pi^{-1}(N' \setminus S(N))$ . Since  $\pi^{-1}(S(N))$  and  $S(M)$  have codimension at least equal to 2,  $\pi^*(\alpha)$  extends as a  $d$ -closed section  $\beta$  of  $\omega_M^p$  on  $\pi^{-1}(N')$ . Then we may

find an open neighborhood  $U$  in  $M$  of  $\pi^{-1}(y) \cap Z$  (which is a finite set) and a section  $\beta \in \Gamma(U, \omega_M^{p-1})$  such that  $d\beta = \pi^*(\alpha)$  on  $U$ . Then  $\beta$  is a  $(p-1)$ -holomorphic form on  $U \setminus S(M)$  and then induces a  $(p-1)$ -holomorphic form on  $Z \setminus (Z \cap S(M))$ . Since  $(Z \cap S(M))$  has codimension at least 2 in  $Z$  we obtain a section on  $Z$  of  $\omega_Z^{p-1}$  (the sheaf  $\omega_Z^\bullet$  has the extension property in codimension  $\geq 2$ ) and we have  $d\beta|_Z = \pi^*(\alpha)|_Z$  since the sheaf  $\omega_Z^\bullet$  has no torsion. The direct image by  $\pi'$  gives a section  $\gamma := \pi'_*(\beta|_Z)$  on  $N'$  which satisfies  $d\gamma = k\alpha$  where  $k$  is the degree of  $\pi'$ . This proves property  $P.4$  for  $N$ .

The proof of  $P.5$  is analogous. ■

REMARK. In the paper [4] it is proved (see Theorem 1.4<sup>2</sup>), generalizing the result of [5], that  $\omega_M^n = L_M^n$  for a normal complex space implies  $\omega_M^p = L_M^p$  for any  $p \in [0, n]$ . An interesting characterization of  $L^p$  is also given in [4] (see Theorem 1.1): A section of  $\omega^p$  is in  $L^p$  if for any  $\beta \in \Omega_M^{n-p}$  and any  $\gamma \in \Omega_M^{n-p-1}$  the forms  $\alpha \wedge \beta$  and  $d\alpha \wedge \gamma$  are in  $L^n$ . □

## 4 Bibliography

### References

- [1] Barlet, D. *Le faisceau  $\omega_X^\bullet$  sur un espace analytique  $X$  de dimension pure*, in Sem. F. Norguet III, Lecture Notes 670 Springer Verlag (1978) pp. 187-204
- [2] Barlet, D et Magnússon, J. *Cycles analytiques complexes II: l'espace des cycles*, Cours Spécialisés SMF 27 2020. Chapitre VIII.
- [3] Barlet, D et Magnússon, J. *On nearly smooth complex spaces* Math. Z. 292. (2019), pp. 317-341.
- [4] Kebekus, S. and Schnell, C. *extending holomorphic forms from the regular locus of a complex space to a resolution of singularities*, arXiv: 1811.03644 [math.AG] 13 Jul 2020.
- [5] Daniel Greb, Stefan Kebekus, Sandor J. Kovacs, and Thomas Peternell. *Differential forms on log canonical spaces*. Inst. Hautes Etudes Sci. Publ. Math., 114 (1) pp. 87–169, November 2011.

---

<sup>2</sup>In fact they prove more: if  $\omega_M^k = L_M^k$  for some  $k \leq n$  then  $\omega_M^p = L_M^p$  for any  $p \in [0, k]$ .