DOES THERE EXIST THE APPLICABILITY LIMIT OF PDE TO DESCRIBE PHYSICAL PHENOMENA?

— A PERSONAL SURVEY OF QUANTIZATION, QED, TURBULENCE —

ATSUSHI INOUE

Dedication to Masuda san for his warm consideration

ABSTRACT. What does it mean to study PDE(=Partial Differential Equation)? How and what to do "to claim proudly that I'm studying a certain PDE"? Newton mechanic uses mainly ODE(=Ordinary Differential Equation) and describes nicely movements of Sun, Moon and Earth etc. Now, so-called quantum phenomenum is described by, say Schrödinger equation, PDE which explains both wave and particle characters after quantization of ODE. The coupled Maxwell-Dirac equation is also "quantized" and QED(=Quantum Electro-Dynamics) theory is invented by physicists. Though it is said this QED gives very good coincidence between theoretical¹ and experimental observed quantities, but what is the equation corresponding to QED? Or, is it possible to describe QED by "equation" in naive sense?

Contents

2
4
4
5
6
7
9
9
9
12
16
17
17

Date: May 2, 2024, based on a talk on 19 March 2023.

Author: alias atlom=a tiny little old mathematician, or suman=superman-"per", "per" pronounces "pa-" in Japanese, means "bone head" or "stupid", therefore deleted and finally "suman" stands for "sorry" in Japanese, curious!

¹By the way, without "solving" equation mathematically, how they obtain theoretical value? After H. Lewy's example, we can't believe so naively physicist claim that there must exist a solution of that PDE if we may derive it nicely from physical phenomena.

ATSUSHI INOUE

5.2. Hopf equ	lation	20
Appendix A.	Some questions on Euler or Navier-Stokes equations	23
References		24

1. INTRODUCTION WITH A BRIEF PERSONAL HISTORY OF ATLOM

Though I attended with the TV course of collegial physics² in my high school time, as I felt the score of Shingaku-Furiwake³ was insufficient to go to physics, I had no alternative but to proceed to applied mathematics because for me at that time, two branches seem to treat analogous objects. On the other hand, my career as a mathematician has been started rather accidentally. Despite I couldn't imagine I might be an "ordinary" salaried man⁴, but fortunately perhaps, just after finished my master course, I was proposed a research assistant post of department of mathematics. Such posts made increased suddenly because of "Science and Engineering Faculty Expansion Plan" by the Japanese government which began after the Sputnik shock suffered by USA. By the way, as is said "The teacher of the university is not resigned as once if I do it" or "Beggars and monks can't quit after three days", I couldn't only resign from research assistant post but also had an opportunity to study abroad as a Boursier.

But after several years at around age thirties, I was offered a new position. At that time, continuously getting salary as researcher, I should do mathematics more seriously, so I felt, and I started to study not only linear PDE but also non-linear PDE. Surely, Hörmander's works on linear PDE are overwhelming at that time.

So to restart with a little mathematician, I questioned naively why the Navier-Stokes equation is so famous and I wondered if this equation is genuinely good enough or worth studying? If it is so good as equation, it should be invariant under change of variables? At that time, the initial and boundary value problem in the time dependent domain for the Navier-Stokes equation is studied by Fujita and Sauer [40] by penalty method. Since I couldn't appreciate their method fully⁵, I find the change of variable formula for vector field or differential 1-form with the help of Wakimoto [71], under the condition that "gold fishes in the bowl don't allow even kissing⁶"! Not only this, even turbulent phenomena are believed to be governed by this same equation, why so? Though this equation is derived from conservations of momentum and mass by observing "laminar flow", why so easily believed the totally different looking phenomena, "turbulent flows", are also governed by the same equation? Not only this, the viscosity occurs after or before viewing the discrete molecular structure of water as a continuum⁷? Here it is also appropriate quoting Hopf saying:

 $^{^{2}}$ A part of NBC's educational program "Continental Classroom" in 1958–1963, which is broadcasted from NHK in 1959–1961 as "physics in nuclear age" with explanation in Japanese

 $^{^{3}}$ After 18months of general education, each student should decide to proceed in which department

⁴Seemingly, I'm not good at temporarily going along with "respectable person" as an amenable member of organization. ⁵Whether their penalty method works when $\bigcup_{0 \le t \le T} \Omega(t) \times \{t\} \subset \mathbb{R}^{d+1}$ when $\Omega(t)$ and $\Omega(t')$ is not necessarily diffeomorphic each other but with the same volume?

⁶That is, assuming that the volume of $\Omega(t)$ is constant in t and for any t, t', $\Omega(t)$ and $\Omega(t')$ are diffeomorphic each other ⁷A physicist says naively that viscosity comes from particle structure of water, they don't bother when they, to treat fluids, started to regard particles as if continuum.

Statistical mechanics constructs certain "relevant" phase distributions which characterize the "typical" phase motions and which must be used for all statistical predictions. \cdots \cdots , in statistical hydromechanics – the theory of highly turbulent fluid flow – the small scale on which the "fluid elements" interact seems to be of decisive importance. The relevant distributions based on this scale – the hypothetical Kolmogoroff distributions – must be mathematically very different from canonical distribution. So far all attempts to determine the relevant hydrodynamical phase distributions, at least to a sufficient degree of approximation, have met with considerable mathematical difficulties.

From these consideration, I wonder whether "is there applicability limit for describing physical phenomena by PDE"? and what is the limit of regarding huge number of water molecules as continuum?

As is well-known, now called "quantum phenomena" are not so well described directly by ODE(Newton Mechanics), therefore we need to use PDE, Schrödinger equation(in some sense, a quantized version of Newton Mechanics). Therefore, at least logically, there will exist some phenomena assumed to be not well describable using PDE.

For such phenomena, Gelfand not only proposed to use FDE^8 (=Functional Derivative Equation) for understanding turbulence and QED, but also questioned whether our existed mathematics tools sufficient to describe these phenomena? I completely empathize with his opinion, and I feel we need to develop theory of FDE⁹.

Apart from above, to treat initial value problem for linear hyperbolic systems of PDE at that time, there is a trend to diagonalize that system with posing conditions on properties of characteristic roots, as mathematical technique. But I feel strange why we need to diagonalize system of PDE. I feel curious to such treatise because not only until when such efforts continue but also there exists not diagonalizable system of PDE. From my point of view, the necessity of diagnalization is to apply existing theory of pseudo-differential equations or more precisely, standard symbol calculus is only confined to scalar case. Therefore, we need to treat matrix structure as it is, this is one motivation to construct superanalysis(=analysis on superspace $\Re^{m|n}$ not on \mathbb{R}^m).

Even almost 30 years passed after I started to concern with FDE or superanalysis, these subjects never belong to the main stream of researchs in mathematical society, at least in Japan. Therefore, it seems natural, at that time, judges of KAKENHI(=Grants-in-Aid for Scientific Research) disregarded my proposal concerning FDE and superanalysis almost completely, so I think. But any way, none of them, even personally, asked me what are them? Finally, I proposed to "make a room to accept appeals to the judgement of KAKENHI examiners¹⁰" which is also neglected completely. Under these situations, Kŷuya Masuda supported me behind the scenes.

⁸Someone wrote somewhere that, "if there doesn't exist Riemannian geometry or invariant theory, theory of relativity, if there doesn't exist boundary value theory, wave mechanics, and if there doesn't exist matrix theory, quantum mechanics may not be invented? New physical theory is stimulated precedent mathematical theory"

⁹But in general, we can't give meaning higher order functional derivatives at each point. Moreover, in infinite dimensional topological spaces, there doesn't exist Lebesgue-like measure which permit integration by parts [91]. This means any trial to extend "A study of PDE by functional analytic method" seems breakdown from the outset. In spite of this, physicists, using tools not yet mathematically justified, get certain theoretical values and experimental values with complicated and expensive experiments, and astonishingly these values coincide many digits. This fact seems to imply something exist which is not yet appreciated mathematically.

 $^{^{10}}$ whose names are announced after one year later

In this note, mathematics which is disregarded by judges of KAKENHI, are mentioned, because without doing so, these trials are in vain with my physical lifespan¹¹.

Since originally, the descriptive ability of PDE and its reproducibility of physical phenomena are main concern, it seems natural we wonder the difference between the derivation of classical field equation and quantum field mechanics.

In any way, it seems worth mentioning the following description which is given in Chapter 9, Functional Methods, of Itzykson and Zuber: "Quantum Field Theory" 1979:

"The path integral formalism of Feynman and Kac provides a unified view of quantum mechanics, field theory, and statistical models. Starting from the case of finitely many degrees of freedom it is generalized to include fermion systems and then extended to infinite systems. The steepest-descent method of integration exhibits the close relationship with classical mechanics and allows us to recover ordinary perturbation theory."

2. QUANTIZATION AND PATH INTEGRAL METHOD

2.1. The beginning of Path Integral Method. Following explanation is due to Feynman and Hibbs [35] but I cited here from Albeverio and Hoegh-Krohn [2].

Let consider the representation formula for the solution of Schrödinger equation on \mathbb{R}^d

$$i\hbar\frac{\partial}{\partial t}\psi(q,t) = -\frac{\hbar^2}{2m}\Delta\psi(q,t) + V(q)\psi(q,t),$$

with the initial data $\psi(q,0) = \underline{\psi}(q)$. Decomposing

$$H = H_0 + V$$
 with $H_0 = -\frac{\hbar^2}{2m}\Delta$,

and assuming that H is selfadjoint in $L^2(\mathbb{R}^d)$, we have the solution $\psi(q,t) = e^{-i\hbar^{-1}tH}\underline{\psi}(q)$ by Stone's theorem. On the other hand, Lie-Trotter-Kato's product formula asserts that even though $[H_0, V] \neq 0$, we have

$$e^{-i\hbar^{-1}tH} = e^{-i\hbar^{-1}t(H_0+V)} = \lim_{n \to \infty} (e^{-i\hbar^{-1}(t/n)V}e^{-i\hbar^{-1}(t/n)H_0})^n.$$

Since

$$e^{-i\hbar^{-1}tH_0}u(q) = (2\pi i\hbar t/m)^{-d/2} \int_{\mathbb{R}^d} dq' \, e^{im(q-q')^2/(2\hbar t)}u(q')$$

we have

$$e^{-i\hbar^{-1}(t/n)V}e^{-i\hbar^{-1}(t/n)H_0}u(q) = e^{-i\hbar^{-1}(t/n)V(q)}\int_{\mathbb{R}^d} dq' \, e^{im(q-q')^2/(2\hbar t/n)}u(q'),$$

we get,

$$(2\pi i\hbar t/m)^{-d/2} \int_{\mathbb{R}^d} dq' \, e^{im(q-q')^2/(2\hbar t)} u(q') = (2\pi i\hbar t/m)^{-dn/2} \int_{\mathbb{R}^{dn}} dq_0 \cdots dq_{n-1} e^{i\hbar S_t^*(q_n, \cdots, q_0)} u(q_0)$$

with $q' = q_0$ and $q_n = q$, where

$$S_t^*(q_n, \cdots, q_0) = \sum_{j=1}^n \left[\frac{m}{2} \frac{(q_j - q_{j-1})^2}{(t/n)^2} - V(q_j)\right] \frac{t}{n}.$$

 $^{^{11}\}mathrm{Sorry}$ to use arXiv as such personal sense

Taking $\gamma^*(\tau)$ on [0,t] as a zigzag path from q_0 at time 0, passing through $\gamma(\tau_j) = q_j \ j = 0, \dots, n$ where $\tau_j = j\frac{t}{n}$ and q_0, \dots, q_n are given points in \mathbb{R}^d . Feynman regards this $S_t^*(q_n, \dots, q_0)$ as a Riemann approximation for the classical action $S_t(\gamma)$ along the path $\gamma(\tau)$:

$$S_t(\gamma) = \int_0^t d\tau \left[\frac{m}{2} \left(\frac{d\gamma}{d\tau}\right)^2 - V(\gamma(\tau))\right]$$

Assuming $\gamma^*(\tau)$ is an approximation for any classical path belonging to the path space

$$\Gamma_{(t,\underline{q},\overline{q})} = \{ \gamma \in AC([0,t]: \mathbb{R}^d); \gamma(0) = \underline{q}, \gamma(t) = \overline{q} \}$$

and using assumed "Lebesgue-like measure" $d_F \gamma$ on $\Gamma_{(t,q,\bar{q})}$, we express its solution as

$$\psi(\bar{q},t) = \int_{\gamma(t)=\bar{q}} d_F \gamma \, e^{i\hbar^{-1}S_t(\gamma)} \underline{\psi}(\gamma(0)),$$

called Feynman's path integral expression for the solution of Schrödinger equation by Feynman's time sclicing method.

If we permit this integral representation of the solution with operations under integral sign admitted, when making $\hbar \to 0$, we have the main contribution stems from the stationary point $\gamma_c(\cdot)$ that is, $\frac{\delta S_t(\gamma)}{\delta \gamma}\Big|_{\gamma=\gamma_c} = 0$. This expression with above interpretation is persuasive to claim that classical mechanical equation appeared from quantum one when making $\hbar \to 0$.

2.2. Quantization à la Fujiwara. Using Feynman's idea rather conversely, Fujiwara [41, 42] constructed a fundamental solution of Schrödinger equation. From a given Lagrangian $L(q, \dot{q})$, he found a classical orbit $\gamma_c(\cdot) \in \Gamma_{(t,q,\bar{q})}$ of that Lagrangian mechanics:

$$\ddot{q}(s) + L(q(s), \dot{q}(s)) = 0 \quad \text{with} \quad q(0) = \underline{q}, \ \dot{q}(0) = \underline{p}^*$$

such that $\bar{q} = q(t, \underline{q}, \underline{p}^*)$, that is, $\gamma_c(0) = \underline{q}, \gamma_c(t) = \bar{q}$. Then, the action integral corresponding to γ_c is given

$$S_t(\gamma_c) = S_t(\bar{q}, \underline{q}) = \int_0^t ds \, L(\gamma_c(s), \dot{\gamma}_c(s)),$$

moreover defining van Vleck determinant as

$$D(t, \bar{q}, \underline{q}) = \det\left(\frac{\partial^2}{\partial \underline{q} \partial \bar{q}} S_t(\bar{q}, \underline{q})\right),$$

he finally define a short time propagator as a FIOp(=Fourier Integral Operator)

$$T_t \underline{\psi}(\bar{q}) = \int_{\mathbb{R}^n} d\underline{q} A(t, \bar{q}, \underline{q}) e^{i\hbar^{-1}S_t(\bar{q}, \underline{q})} \underline{\psi}(\underline{q})$$

where

$$A(t, \bar{q}, \underline{q}) = \sqrt{D(t, \bar{q}, \underline{q})}.$$

Above Fujiwara's process is justified when $\sup_q |\partial_q^{\alpha} V(q)| \leq C, (|\alpha| \geq 2)$. That is, under this condition, not only there exists a unique classical orbit, and above quantities are well-defined, but also T_t defines a bounded linear operator on $L^2(\mathbb{R}^n)$ with the property

$$||T_{t+s}u - T_sT_tu|| \le C(t^2 + s^2)||u||.$$

Moreover, following operator E_t is defined by

$$\lim_{n \to \infty} \| (T_{t/n})^n u - E_t u \| = 0$$

which gives a parametrix of the given initial value problem of Schrödinger equation [41]. The kernel representation is the fundamental solution such that

$$(E_t u)(\bar{q}) = \int_{\mathbb{R}^n} d\underline{q} K(t, \bar{q}, \underline{q}) u(\underline{q}).$$

Finally, not only how to characterize the long-time behavior¹² of $K(t, \bar{q}, \underline{q})$ [42], but also how to use the # products of FIOps by Kumano-go group, are the problems reconsidered.

Remark 2.1. (1)[Feynman-Kac formula]: Sitimulated by Feynman's idea, M. Kac represents the solution of

$$\frac{\partial}{\partial t}u(q,t) = \sigma\Delta u(q,t) - V(q)u(q,t),$$

using Wiener measure $dW(\gamma)$ by

$$u(q,t) = \int e^{-\int_0^t dW(\gamma) V(\gamma(s)+q)ds} \underline{u}(\gamma(0)+\underline{q}).$$

(2)[Problem of R/12]: For a given Riemann manifold (M, g_{jk}) , applying Fujiwara's idea to the heat equation, that is, some type of quantization, Inoue and Maeda [68] got the notorious term R/12 where R is the scalar curvature of the Riemann metric $g_{jk}(q)dq^{j}dq^{k}$. Curiously, this term was also derived as the most probable path calculation in probability theory [39, 94].

Rather recently the term R/12 appeared in S. Fukushima [43], but this seems not the direct consequence of quantization of $g_{jk}(q)dq^jdq^k$ like physicists arguments, for example, B. DeWitt [29] or F. Bastianelli et al. [14]. Fukushima tries to construct a fundamental solution of $i\frac{\partial}{\partial t} - \frac{1}{2}\Delta_g + \frac{R}{12}$. (3)[Hamiltonian formulation on manifolds?] Above mentioned works containing physicist papers, are formulated in Lagrangian case. Since it needs works to formulate Fourier transformations on manifolds, physicist papers such as Field [36, 37] claim something interesting without any mathematical estimates.

2.3. Hamilton formulation of PIM. For a given Lagrangian $L(q, \dot{q})$, above process is denoted formally as

$$K = \int d_F \gamma \, e^{i\hbar^{-1}S(\gamma)} \quad \text{or} \quad K(t, \bar{q}, \underline{q}) = \int_{\Gamma_{(t, \bar{q}, \underline{q})}} d_F \gamma \, e^{i\hbar^{-1}S(\gamma)}$$

where

$$S(\gamma) = \int_0^t ds \, L(\gamma(s), \dot{\gamma}(s)), \text{ for each } \gamma \in \Gamma_{(t, \bar{q}, \underline{q})}.$$

What does it become when Hamiltonian H(q, p) is given? Here, as is well-known, H(q, p) is related to $L(q, \dot{q})$ by Legendre transformation. How to give meaning to the following formal expression?:

$$K = \int d_F p \, d_F q \, e^{i\hbar^{-1} \int ds(\dot{q}p - H(q,p))}$$

Under same assumption as above, there exists a classical orbit $(q(t, \underline{q}, \underline{p}), p(t, \underline{q}, \underline{p}))$ to the Hamilton equation for the given H(q, p) with initial data $(\underline{q}, \underline{p})$. For any fixed \underline{p} , for the map $\underline{q} \to \overline{q} = q(t, \underline{q}, \underline{p})$, there exists an inverse map $\underline{q} = x(t, \overline{q}, \underline{p})$ for sufficiently small |t|:

$$\bar{q} = q(t, x(t, \bar{q}, \underline{p}), \underline{p}) \text{ and } \underline{q} = x(t, q(t, \underline{q}, \underline{p}), \underline{p})$$

Using these, we put

$$S(t, \bar{q}, \underline{p}) = (\underline{qp} - S_0(t, \underline{q}, \underline{p})) \Big|_{\underline{q} = x(t, \bar{q}, \underline{p})}$$

 $^{^{12}}$ For example, how Maslov index appeared in PIM?

and define

$$D(t,\bar{q},\underline{p}) = \det \left(\partial_{\bar{q}} \partial_{\underline{p}} S(t,\bar{q},\underline{p}) \right).$$

Finally we define

$$T_t u(\bar{q}) = (2\pi\hbar)^{-n/2} \int_{\mathbb{R}^n} d\underline{p} D(t, \bar{q}, \underline{p})^{1/2} e^{i\hbar^{-1}S(t, \bar{q}, \underline{p})} \hat{u}(\underline{p})$$

This gives a parametrix for Schrödinger equation [58].

Remark 2.2. In order to consider a classical orbit connecting 2 points \bar{q} and \underline{q} in t, we need 2nd order time derivatives in corresponding classical mechanical equation. Then, how one treats Weyl or Dirac equation? Hint is given in the next section.

3. Another interpretation of m.o.c., explained through the simplest case

Let check the difference between Lagarangian or Hamiltonian methods, by taking the simplest example. This is important to treat first order system of PDE such as Dirac or Weyl equation by applying Feynman's time slicing method.

Recall m.o.c.(=method of characteristics). On region $\Omega \subset \mathbb{R}^{d+1}$, we consider the following initial value problem: Let solve

(3.1)
$$\begin{cases} \frac{\partial}{\partial t}u(t,q) + \sum_{j=1}^{d} a_j(t,q) \frac{\partial}{\partial q_j}u(t,q) = b(t,q)u(t,q) + f(t,q), \\ u(\underline{t},q) = \underline{u}(q) \end{cases}$$

The characterisitic equation of (3.1) is given by

$$\begin{cases} \frac{d}{dt}q_j(t) = a_j(t, q(t)), \\ q_j(\underline{t}) = \underline{q}_j \quad \text{for} \quad j = 1, \cdots, d, \end{cases}$$

with solution denoted by

$$q(t) = q(t, \underline{t}; \underline{q}) = (q_1(t), \cdots, q_d(t)) \in \mathbb{R}^d, \underline{q} = (\underline{q}_1, \cdots, \underline{q}_d).$$

Using this,

Theorem 3.1 (method of characteristics). In (3.1), assume coefficients $a_j \in C^1(\Omega : \mathbb{R})$, $b, f \in C(\Omega : \mathbb{R})$. Taking any point $(\underline{t}, \underline{q}) \in \Omega$, assume \underline{u} is C^1 in a neighbourhood of \underline{q} . Then, there exists a unique solution u(t,q) near $(\underline{t},\underline{q})$. Moreover, putting $B(t,\underline{q}) = b(t,q(t,\underline{t};\underline{q}))$, $F(t,\underline{q}) = f(t,q(t,\underline{t};\underline{q}))$ and defining

(3.2)
$$U(t,\underline{q}) = e^{\int_{\underline{t}}^{t} d\tau \ B(\tau,\underline{q})} \left\{ \int_{\underline{t}}^{t} ds \ e^{-\int_{\underline{t}}^{s} d\tau \ B(\tau,\underline{q})} F(s,\underline{q}) + \underline{u}(\underline{q}) \right\}$$

we have a solution (3.1) given by

(3.3)
$$u(t,\bar{q}) = U(t,x(t,\bar{q})).$$

Here, $\underline{q} = x(t, \underline{t}; \overline{q})$ is the inverse function of $\overline{q} = q(t, \underline{t}; \underline{q})$.

Remark 3.1. (3.2) satisfies

$$\frac{d}{dt}U(t,\underline{q}) = B(t,\underline{q})U(t,\underline{q}) + F(t,\underline{q}).$$

An example: With this theorem in mind, we consider the following simplest case:

(3.4)
$$\begin{cases} i\hbar\frac{\partial}{\partial t}u(t,q) = a\frac{\hbar}{i}\frac{\partial}{\partial q}u(t,q) + bqu(t,q), \\ u(0,q) = \underline{u}(q). \end{cases}$$

The symbol of the righthand side of above equation is derived by

$$H(q,p) = e^{-i\hbar^{-1}qp} \left(a\frac{\hbar}{i}\frac{\partial}{\partial q} + bq \right) e^{i\hbar^{-1}qp} = ap + bq.$$

From this, we have the characteristic

$$\dot{q}(t) = \frac{\partial H(q, p)}{\partial p} = a \text{ with } q(0) = \underline{q},$$

whose solution is given

$$q(s) = \underline{q} + as$$
 with $\underline{q} = x(t, \overline{q}) = \overline{q} - at$.

Since above representation (3.2) gives

$$U(t,\underline{q}) = \underline{u}(\underline{q})e^{-i\hbar^{-1}(b\underline{q}t+2^{-1}abt^2)},$$

from (3.3), we get

$$u(t,\overline{q}) = \underline{u}(\overline{q} - at)e^{-i\hbar^{-1}(b\overline{q}t - 2^{-1}abt^2)}$$

By this procedure, the information of p(t) isn't used!

Now, we give another interpretation of this representation by Hamilton path integral method.

For $(t, \underline{q}, \underline{p})$, putting

$$S_0(t,\underline{q},\underline{p}) = \int_0^t ds [\dot{q}(s)p(s) - H(q(s),p(s))] = -b\underline{q}t - 2^{-1}abt^2$$

we define the action integral for Hamilton function H(q, p)

$$S(t,\overline{q},\underline{p}) = \underline{q}\,\underline{p} + S_0(t,\underline{q},\underline{p})\big|_{\underline{q}=x(t,\overline{q})} = \overline{q}\underline{p} - a\underline{p}t - b\overline{q}t + 2^{-1}abt^2.$$

This $S = S(t, \overline{q}, \underline{p})$ satisfies the following Hamilton-Jacobi equation:

$$\frac{\partial}{\partial t}S + H(\overline{q}, \partial_{\overline{q}}S) = 0 \quad \text{with} \quad S(0, \overline{q}, \underline{p}) = \overline{q}\underline{p}$$

In this case, van Vleck determinant is a scalar

$$\frac{\partial^2 S(t,\overline{q},\underline{p})}{\partial \overline{q} \partial p} = 1$$

and it satisfies the following continuity equation:

$$\frac{\partial}{\partial t}D + \frac{1}{2}\partial_{\overline{q}}(DH_p) = 0 \quad \text{with} \quad D(0, \overline{q}, \underline{p}) = 1 \quad \text{where} \quad H_p = \frac{\partial H}{\partial p}(\overline{q}, \partial_{\overline{q}}S)$$

Using these classical quantities S and D, modifying Feynman's method slightly¹³, we define

$$u(t,\overline{q}) = (2\pi\hbar)^{-1/2} \int d\underline{p} D^{1/2}(t,\overline{q},\underline{p}) \cdot e^{i\hbar^{-1}S(t,\overline{q},\underline{p})} \underline{\hat{u}}(\underline{p}),$$

Using $\delta(\overline{q} - at - \underline{q}) = (2\pi\hbar)^{-1} \int d\underline{p} e^{i\hbar^{-1}(\overline{q} - at - \underline{q})\underline{p}}$, we have

$$\begin{split} u(t,\overline{q}) &= (2\pi\hbar)^{-1/2} \int d\underline{p} \, e^{i\hbar^{-1}S(t,\overline{q},\underline{p})} \underline{\hat{u}}(\underline{p}) \\ &= (2\pi\hbar)^{-1} \iint d\underline{p} d\underline{q} \, e^{i\hbar^{-1}(S(t,\overline{q},\underline{p})-\underline{q}\underline{p})} \underline{u}(\underline{q}) = \int d\underline{q} \delta(\overline{q}-at-\underline{q}) \underline{u}(\underline{q}) e^{i\hbar^{-1}(-b\overline{q}t+2^{-1}abt^2)} \\ &= \underline{u}(\overline{q}-at) e^{i\hbar^{-1}(-b\overline{q}t+2^{-1}abt^2)}. \end{split}$$

 $^{^{13}}$ Though Feynman's procedure based on Lagrangian, here we changed to Hamiltonian formulation

4. Necessity of superanalysis

4.1. **Reasons of necessity.** There are several reasons at least for me: (i) As is written in p.355 of R.Feynman & A.R. Hibbs [35], 'spin' has been the object outside Feynman's procedures at that time (below, underlined by atlom):

... <u>path integrals suffer grievously from a serious defect</u>. They do not permit a discussion of spin operators or other such operators in a simple and lucid way. They find their greatest use in systems for which coordinates and their conjugate momenta are adequate. Nevertheless, spin is a simple and vital part of real quantum-mechanical systems. It is a serious limitation that the half-integral spin of the electron does not find a simple and ready representation. It can be handled if the amplitudes and quantities are considered as quarternions instead of ordinary complex numbers, but the lack of commutativity of such numbers is a serious complication.

(ii) E. Berezin [18], who invented the second quantization, proposed to "treat boson and fermion on equal footing". That is, instead of \mathbb{R} or \mathbb{C} , he claims the necessity to construct a scalar-like, non-commutative ground field, where lives electron and photon equally.

(iii) E. Witten [99] explained the meaning of supersymmetric idea in physics to mathematician by new derivation of Morse inequalities using exterior differential operations. But be careful, in many case, physicists usage of supersymmetry only means to use fermion or odd variables, not exactly treating some "symmetry" in $\Re^{m|n}$, for example K.B. Efetov's works [32] and Y.V. Fyodorov [44].

4.2. Regarding matrices as differential operators! A claim "any Clifford algebra has a representation on Grassmann algebra"¹⁵ is straight forwardly explained for 2×2 matrices case. Though analogous argument works for a set of $2^d \times 2^d$ matrices because it has Clifford relation, but for a set of general $N \times N$ matrices, we need to find suitable algebra on which we need foundation of analysis. I imagine not only the work of J.L. Martin [78, 79] but also the one by R. Campoamor-Stursberg et al. [22] give some hint on this. Especially N = 3 will be interesting when we consider vorticity equation for Euler or Navier-Stokes equation.

Recall Pauli matrices $\{\boldsymbol{\sigma}_k\}_{k=1}^3$:

$$\boldsymbol{\sigma}_1 = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \boldsymbol{\sigma}_2 = -i \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}, \quad \boldsymbol{\sigma}_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

These matrices satisfy not only Clifford relation

(4.1)
$$\boldsymbol{\sigma}_{j}\boldsymbol{\sigma}_{k} + \boldsymbol{\sigma}_{k}\boldsymbol{\sigma}_{j} = 2\delta_{jk}, \text{ where } j,k=1,2,3,$$

but also the following for any (j, k, ℓ) , an even permutation of (1, 2, 3):

(4.2)
$$\boldsymbol{\sigma}_{i}\boldsymbol{\sigma}_{k} = i\boldsymbol{\sigma}_{\ell}$$

 $^{^{14}}$ Inoue[57, 59]

¹⁵Though I don't appreciate fully Chevalley' theorem "Any Clifford algebra has a representation on Grassmann algebra", but following consideration may be sufficient of my intuitive arguments.

Decompose any 2×2 matrix \mathbb{A} by $\{\boldsymbol{\sigma}_j\}$,

(4.3)
$$\mathbb{A} = \begin{pmatrix} a & c \\ d & b \end{pmatrix} = \frac{a+b}{2} \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} + \frac{a-b}{2} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix} + \frac{c+d}{2} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} = \frac{a+b}{2} \mathbb{I}_2 + \frac{a-b}{2} \sigma_3 + \frac{c+d}{2} \sigma_1 + i\frac{c-d}{2} \sigma_2.$$

That is, a set of all 2×2 matrices has the Clifford algebra structure with Pauli matrices $\{\boldsymbol{\sigma}_k\}$ as a basis.

We identify a vector as a function of Grassmann variables. Then, a matrix is regarded as a differential operator acting on a set of functions composed with Grassmann variables, which forms something-like field similar to real or complex number field but non-commutative.

Rather abruptly we prepare¹⁶ two odd variables θ_1, θ_2 having the following relations:

(4.4)
$$\theta_1 \cdot \theta_2 + \theta_2 \cdot \theta_1 = 0, \ \theta_j \cdot \theta_j = 0, \ (j = 1, 2)$$

That is, they have product \cdot having Grassmann relations and differentiation such as

$$\frac{\partial}{\partial \theta_1} 1 = 0, \quad \frac{\partial}{\partial \theta_1} \theta_1 = 1, \quad etc.$$

Here, when we differentiate, bringing that variable in front, i.e.

$$\frac{\partial}{\partial \theta_2} \theta_1 \cdot \theta_2 = \frac{\partial}{\partial \theta_2} (-\theta_2 \cdot \theta_1) = \frac{\partial}{\partial \theta_2} (\theta_2 \cdot (-\theta_1)) = -\theta_1.$$

More concretely, for example, taking two variables z_1 and z_2 in \mathbb{R}^2 , construct differential forms dz_1 , dz_2 with exterior product \wedge and interior product \mid such that

$$dz_1 \wedge dz_2 = -dz_2 \wedge dz_1, \quad dz_j \wedge dz_j = 0, \quad \frac{\partial}{\partial z_j} \lfloor dz_k = \delta_{jk}.$$

Here, identifying dz_j as θ_j for j = 1.2, \wedge as \cdot and $\frac{\partial}{\partial z_j} \lfloor$ as $\frac{\partial}{\partial \theta_j}$, and abbreviating \cdot , we continue to explain.

Using odd variables θ_1 , θ_2 , we give the identifying maps #, \flat with Grassmann algebras as follows:

(4.5)
$$\Gamma_{0} = \{u(\theta) = u_{0} + u_{1}\theta_{1}\theta_{2} \mid u_{0}, u_{1} \in \mathbb{C}\} \stackrel{\flat}{\underset{\sharp}{\leftrightarrow}} \mathbb{C}^{2} = \left\{\mathbf{u} = \begin{pmatrix}u_{0}\\u_{1}\end{pmatrix} \mid u_{0}, u_{1} \in \mathbb{C}\right\}$$
$$\text{with} \quad (\#\mathbf{u})(\theta) = \left(\#\begin{pmatrix}u_{0}\\u_{1}\end{pmatrix}\right)(\theta) = u_{0} + u_{1}\theta_{1}\theta_{2},$$
$$\flat(u_{0} + u_{1}\theta_{1}\theta_{2}) = \flat(u(\theta)) = \begin{pmatrix}u_{0}\\u_{1}\end{pmatrix}.$$

Here, we have $u(0) = u(\theta)|_{\theta=0} = u_0$, $\partial_{\theta_2}\partial_{\theta_1}u(\theta)|_{\theta=0} = \partial_{\theta_2}(\partial_{\theta_1}u(\theta))|_{\theta=0} = u_1$, which relates a vector **u** and a function $u(\theta)$.

Now, we define differential operators w.r.t. odd variables:

(4.6)

$$\sigma_{1}(\theta,\partial_{\theta}) = \theta_{1}\theta_{2} - \frac{\partial^{2}}{\partial\theta_{1}\partial\theta_{2}},$$

$$\sigma_{2}(\theta,\partial_{\theta}) = i\left(\theta_{1}\theta_{2} + \frac{\partial^{2}}{\partial\theta_{1}\partial\theta_{2}}\right),$$

$$\sigma_{3}(\theta,\partial_{\theta}) = 1 - \theta_{1}\frac{\partial}{\partial\theta_{1}} - \theta_{2}\frac{\partial}{\partial\theta_{2}}.$$

 $^{^{16}\}text{usage}$ "preparing" odd variables $\theta_1,\,\theta_2$ is not frequently used in PDE group

Thes operators act on $u(\theta) = u_0 + u_1 \theta_1 \theta_2 \in \Gamma_0$ as follows:

$$\sigma_1(\theta, \partial_\theta)(u_0 + u_1\theta_1\theta_2) = u_0\theta_1\theta_2 + u_1,$$

$$\sigma_2(\theta, \partial_\theta)(u_0 + u_1\theta_1\theta_2) = i(u_0\theta_1\theta_2 - u_1),$$

$$\sigma_3(\theta, \partial_\theta)(u_0 + u_1\theta_1\theta_2) = u_0 - u_1\theta_1\theta_2.$$

Here,

$$\begin{split} \frac{\partial^2}{\partial \theta_1 \partial \theta_2} \theta_1 \theta_2 &= \partial_{\theta_1} \partial_{\theta_2} \theta_1 \theta_2 \\ &= \partial_{\theta_1} (\partial_{\theta_2} \theta_1 \theta_2) = \partial_{\theta_1} (\partial_{\theta_2} (-\theta_2 \theta_1)) = \partial_{\theta_1} (-\theta_1) = -1 \end{split}$$

They act on Γ_0 as,

$$\sigma_1(\theta,\partial_\theta)\sigma_2(\theta,\partial_\theta) = i\sigma_3(\theta,\partial_\theta), \quad \sigma_2(\theta,\partial_\theta)\sigma_3(\theta,\partial_\theta) = i\sigma_1(\theta,\partial_\theta), \quad \sigma_3(\theta,\partial_\theta)\sigma_1(\theta,\partial_\theta) = i\sigma_2(\theta,\partial_\theta).$$

Remark 4.1. (1) The action $\sigma_1(\theta, \partial_{\theta})$ on Γ_0 has a matrix representation

$$\flat \sigma_1(\theta, \partial_\theta) \sharp \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} = \begin{pmatrix} u_1 \\ u_0 \end{pmatrix}, \quad \flat \sigma_1(\theta, \partial_\theta) \sharp = \boldsymbol{\sigma}_1, \quad etc.$$

(2) Moreover, putting

$$\Gamma_1 = \left\{ v(\theta) = v_1 \theta_1 + v_2 \theta_2 \mid v_1, v_2 \in \mathbb{C} \right\} \underset{\sharp}{\stackrel{\flat}{\leftrightarrow}} \mathbb{C}^2 = \left\{ \mathbf{v} = \begin{pmatrix} v_1 \\ v_2 \end{pmatrix} \mid v_1, v_2 \in \mathbb{C} \right\}$$

we have

$$\sigma_j(\theta, \partial_\theta)v(\theta) = 0$$
 for any $v \in \Gamma_1$ and $j = 1, 2, 3$.

We define Fourier transformation¹⁷ for odd variables. Taking number \hbar in $\mathbb{R}^{\times} = \mathbb{R} - \{0\}$ or $i\mathbb{R}^{\times}$ called it as spin constant, we define

$$\hat{u}(\pi) = \hbar \int_{\Re^{0|2}} d\theta \, e^{-i\hbar^{-1}\langle \theta | \pi \rangle} u(\theta), \quad u(\theta) = \hbar \int_{\Re^{0|2}} d\pi \, e^{i\hbar^{-1}\langle \theta | \pi \rangle} \hat{u}(\pi).$$

In the above, the integral interval $\Re^{0|2}$ may be considered only as a symbol, at least for the time being. Here, $\langle \theta | \pi \rangle = \theta_1 \pi_1 + \theta_2 \pi_2$ and

$$\langle \theta | \pi \rangle^2 = 2\theta_1 \pi_1 \theta_2 \pi_2 = -2\theta_1 \theta_2 \pi_1 \pi_2, \quad \langle \theta | \pi \rangle^j = \overbrace{\langle \theta | \pi \rangle \langle \theta | \pi \rangle \cdots \langle \theta | \pi \rangle}^{j \text{ times}} = 0 \text{ if } j \ge 3$$

we have

$$e^{a\langle\theta|\pi\rangle} = \sum_{\ell=1}^{\infty} \frac{a^{\ell} \langle\theta|\pi\rangle^{\ell}}{\ell!} = 1 + a\langle\theta|\pi\rangle - a^{2} \theta_{1} \theta_{2} \pi_{1} \pi_{2}.$$

Remark 4.2. "Integration" should be considered preferable such that (i) all polynomials are integrable, (ii) linear with integrand and (iii) translation invariant, Berezin integration satisfies these properties.

By Fourier transformation, differential operator is regarded as multiplication in dual space¹⁸, and theory of pseudo-differential operators treats PDE with variable coefficients controlling error estimates.

Defining Weyl symbols of differential operators $\{\sigma_j(\theta, \partial_\theta)\}_{j=1}^3$ as

(4.7)
$$\begin{cases} \sigma_1(\theta, \pi) = \theta_1 \theta_2 + \hbar^{-2} \pi_1 \pi_2, \\ \sigma_2(\theta, \pi) = i(\theta_1 \theta_2 - \hbar^{-2} \pi_1 \pi_2), \\ \sigma_3(\theta, \pi) = -i\hbar^{-1} \langle \theta | \pi \rangle = -i\hbar^{-1}(\theta_1 \pi_1 + \theta_2 \pi_2), \end{cases}$$

¹⁷The integration here is very algebraic, called Berezin integral, which has properties as integral. See, more precisely, V.S. Vladimirov and I.V. Volovich [98]

 $^{^{18}}$ This means, must we recognize objects algebraically? Or four arithmetic operations is the only core to recognize mathematically?

we have, for example,

$$(\hat{\sigma}_{3}^{\mathsf{w}}(\theta,\partial_{\theta})u)(\theta) = -i\hbar \iint_{\mathfrak{R}^{0|2}\times\mathfrak{R}^{0|2}} d\pi d\theta' \, e^{i\hbar^{-1}\langle\theta-\theta'|\pi\rangle} \sigma_{3}(\frac{\theta+\theta'}{2},\pi)(u_{0}+u_{1}\theta'_{1}\theta'_{2}) = u_{0}-u_{1}\theta_{1}\theta_{2} = \sigma_{3}(\theta,\partial_{\theta})(u_{0}+u_{1}\theta_{1}\theta_{2}). \ etc.$$

Here, we used

$$\int_{\Re^{0|2}} d\pi \, e^{i k^{-1} \langle \theta - \theta' | \pi \rangle} \sigma_3(\frac{\theta + \theta'}{2}, \pi) = -i k^{-1} (\theta_1 \theta_2 - \theta'_1 \theta'_2).$$

Remark 4.3. (i) It takes many time to perceive the meaning of 1 in $\sigma_3(\theta, \partial_{\theta})$ appeared in (4.6), that is, 1 stems from Weyl quantization!

(ii) Artificially introduced spin constant k gives nice result only when $k\hbar^{-1} = 1$, that is, the action integral obtained by Jacobi method satisfies Hamilton-Jacobi equation in case $k\hbar^{-1} = 1$, therefore we assume $k = \hbar!$

Therefor, from (4.3), we have

(4.8)
$$(\sharp \mathbb{A}\mathbf{u})(\theta) = \left[\frac{a+b}{2} + \frac{a-b}{2}\sigma_3(\theta,\partial_\theta) + \frac{c+d}{2}\sigma_1(\theta,\partial_\theta) + i\frac{c-d}{2}\sigma_2(\theta,\partial_\theta)\right]u(\theta)$$

whose Weyl symbol is

(4.9)
$$\sigma^{w}(\sharp \mathbb{A}b) = \left[\frac{a+b}{2} + \frac{a-b}{2}\sigma_{3}^{w}(\theta,\pi) + \frac{c+d}{2}\sigma_{1}(\theta,\pi) + i\frac{c-d}{2}\sigma_{2}(\theta,\pi)\right].$$

Remark 4.4. In the theory of pseudo-differential operators, symbol calculus is the main ingredient where algebra is essentially \mathbb{R} . Whether analogous procedure works on superspace $\mathfrak{R}^{m|n}$? More explicitly, though the symbol of $\sigma_1(\theta, \partial_{\theta})$ is given as $\theta_1 \theta_2 + k^{-2} \pi_1 \pi_2$, but whether the inverse of $\sigma_1(\theta, \partial_{\theta})$ is calculated directly from $\theta_1 \theta_2 + k^{-2} \pi_1 \pi_2$?

4.3. Chi's example. The following result is a part of Chi [23] in 1958:

Theorem 4.1 (An example of weakly hyperbolic equation). Let

$$L(t) = L(t, \partial_q) = t^2 \partial_q^2 + b \partial_q.$$

Solving the initial value problem

(4.10)
$$u_{tt} - L(t, \partial_q)u = 0 \quad with \quad u(0,q) = \underline{u}_0(q), \quad u_t(0,q) = 0,$$

when b = 4k + 1, $k \in \mathbb{Z}_+$, we have the solution given by

$$u(t,q) = \sum_{\ell=0}^{k} \frac{2^{2\ell} k!}{(2\ell)!(k-\ell)!} t^{2\ell} \underline{u}_{0}^{(\ell)} \left(q + \frac{t^{2}}{2}\right).$$

In this section, we derive this result by completely different method from Chi.

4.3.1. Chi's proof. Applying change of variables,

$$\xi = x + \frac{t^2}{2}, \quad \eta = x - \frac{t^2}{2}$$

to u(t,x), Chi reduces above problem to the Euler-Darboux equation for $\tilde{u}(\xi,\eta) = u\left(\sqrt{\xi-\eta}, \frac{\xi+\eta}{2}\right)$ yielding,

(4.11)
$$\frac{\partial^2 \tilde{u}}{\partial \xi \partial \eta} - \frac{1-b}{4(\xi-\eta)} \frac{\partial \tilde{u}}{\partial \xi} + \frac{1+b}{4(\xi-\eta)} \frac{\partial \tilde{u}}{\partial \eta} = 0$$

with initial condition

$$\tilde{u}(\xi,\xi) = \underline{u}_0(\xi), \quad \lim_{\xi = \eta \to 0} (\xi - \eta)^{1/2} \left(\frac{\partial \tilde{u}}{\partial \xi} - \frac{\partial \tilde{u}}{\partial \eta} \right) = \underline{u}_1(\xi).$$

Remark 4.5. For

$$E(\alpha,\beta) = \partial_{\xi}\partial_{\eta} - \frac{\beta}{\xi - \eta}\partial_{\xi} + \frac{\alpha}{\xi - \eta}\partial_{\eta},$$

with $\alpha = \frac{1+b}{4}$, $\beta = \frac{1-b}{4}$, it gives (4.11). He uses the fact such that if $0 < \alpha, \beta < 1$ then the solution of $E(\alpha, \beta)u = 0$ is expressed as the convergent Euler-Darboux integral

$$u(\xi,\eta) = \frac{\Gamma(\alpha+\beta)}{\Gamma(\alpha)\Gamma(\beta)} \int_0^1 dt \, u_0(\xi+(\eta-\xi)t)t^{\beta-1}(1-t)^{\alpha-1} + \frac{\Gamma(1-\alpha-\beta)}{2\Gamma(1-\alpha)\Gamma(1-\beta)} (\eta-\xi)^{t-\alpha-\beta} \int_0^1 dt \, u_1(\xi+(\eta-\xi)t)t^{-\alpha}(1-t)^{-\beta}.$$

Moreover, for other α, β , this integral diverges as it stands, he gives these as Riemann-Liouville integral.

Seemingly, Euler-Darboux equation has some relations to special functions or Painleve's functions, therefore proceeding inversely, we might dream another perspective to these functions by using superanalysis?

4.3.2. Direct construction of a solution of Hamilton-Jacobi equation. Without reducing to Euler-Darboux equation, we give a simple minded proof (at least as a story) of Chi's equation applying superanalysis.

Putting $u_0 = u$, $u_1 = u_t$, we make (4.10) to a system:

$$i\frac{\partial}{\partial t} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix} = i \begin{pmatrix} 0 & 1 \\ L(t, \partial_q) & 0 \end{pmatrix} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}.$$

Here, we multiply *i* to both sides, rather artificially. Changing a vector representation to non-commutative scalar one by putting $u(t, x, \theta) = u_0(t, x) + u_1(t, x)\theta_1\theta_2$, we have

$$(L(t,\partial_x)\theta_1\theta_2 - \partial_{\theta_1}\partial_{\theta_2})(u_0 + u_1\theta_1\theta_2) = u_1 + Lu_0\theta_1\theta_2 \sim \begin{pmatrix} 0 & 1 \\ L(t,\partial_x) & 0 \end{pmatrix} \begin{pmatrix} u_0 \\ u_1 \end{pmatrix}.$$

Here, $q \in \mathbb{R}$ is imbedded in $x \in \mathfrak{R}^{1|0}$ such that $X_{\mathrm{B}} = q$. Defining

$$\mathcal{H}(t,\partial_x,\theta,\partial_\theta) = iL(t,\partial_x)\theta_1\theta_2 - i\partial_{\theta_1}\partial_{\theta_2}$$

whose symbol is given by

$$\mathcal{H}(t,\xi,\theta,\pi) = i(-t^2\xi^2 + ib\xi)\theta_1\theta_2 + i\pi_1\pi_2$$

and corresponding Hamilton-Jacobi equation is given by

(4.12)
$$\mathcal{S}_t + \mathcal{H}(t, \mathcal{S}_x, \theta, \mathcal{S}_\theta) = 0$$

which is solved with the initial condition $\mathcal{S}(0, x, \xi, \theta, \pi) = \langle x | \xi \rangle + \langle \theta | \pi \rangle$.

In the supersmooth category, since we have

(4.13)
$$S(t, x, \xi, \theta, \pi) = S(t, x, \xi) + X(t, x, \xi)\theta_1\theta_2 + Y(t, x, \xi)\theta_1\pi_1 + Y(t, x, \xi)\theta_2\pi_2 + V(t, x, \xi)\theta_1\pi_2 + \tilde{V}(t, x, \xi)\theta_2\pi_1 + Z(t, x, \xi)\pi_1\pi_2 + W(t, x, \xi)\theta_1\theta_2\pi_1\pi_2,$$

where

$$\begin{split} S(t,x,\xi) &= \mathcal{S}(t,x,\xi,0,0), \\ X(t,x,\xi) &= \partial_{\theta_2}\partial_{\theta_1}\mathcal{S}(t,x,\xi,\theta,\pi)\big|_{\theta=\pi=0}, \\ Y(t,x,\xi) &= \partial_{\pi_1}\partial_{\theta_1}\mathcal{S}(t,x,\xi,\theta,\pi)\big|_{\theta=\pi=0}, \quad \tilde{Y}(t,x,\xi) = \partial_{\pi_2}\partial_{\theta_2}\mathcal{S}(t,x,\xi,\theta,\pi)\big|_{\theta=\pi=0}, \\ V(t,x,\xi) &= \partial_{\pi_2}\partial_{\theta_1}\mathcal{S}(t,x,\xi,\theta,\pi)\big|_{\theta=\pi=0}, \quad \tilde{V}(t,x,\xi) = \partial_{\pi_1}\partial_{\theta_2}\mathcal{S}(t,x,\xi,\theta,\pi)\big|_{\theta=\pi=0}, \\ Z(t,x,\xi) &= \partial_{\pi_2}\partial_{\pi_1}\mathcal{S}(t,x,\xi,\theta,\pi)\big|_{\theta=\pi=0}, \quad W(t,x,\xi) = \partial_{\pi_2}\partial_{\pi_1}\partial_{\theta_2}\partial_{\theta_1}\mathcal{S}(t,x,\xi,\theta,\pi)\big|_{\theta=\pi=0}, \end{split}$$

we should seek solution S of (4.12) with this form. Since

$$\mathcal{H}(t, \mathcal{S}_x, \theta, \mathcal{S}_\theta) = i(-t^2 \mathcal{S}_x^2 + ib \mathcal{S}_x)\theta_1 \theta_2 + i \mathcal{S}_{\theta_1} \mathcal{S}_{\theta_2},$$

putting $\theta = \pi = 0$ in (4.12), we have readily

$$S_t(t, x, \xi) = 0$$
 with $S(0, x, \xi) = \langle x|\xi \rangle$.

This gives $S(t, x, \xi) = \langle x | \xi \rangle$.

Differentiating (4.12) w.r.t. θ_1 and θ_2 then restricting to $\theta = \pi = 0$, we get

$$X_t + i(-t^2\xi^2 + ib\xi) + iX^2 = 0$$

Since this is the Riccati type ODE, using $\varphi(t)$, we find $X = -i\frac{\dot{\varphi}}{\varphi}$, that is,

(4.14)
$$\ddot{\varphi} + (t^2 \xi^2 - ib\xi)\varphi = 0 \quad \text{with} \quad \dot{\varphi}(0) = 0$$

Regarding (x,ξ) as parameter, we may solve this equation by power series in t, but this procedure is postponed until explaining our construction of S and quantization.

Putting $Y(t, x, \xi) = \partial_{\pi_1} \partial_{\theta_1} \mathcal{S}(t, x, \xi, \theta, \pi) \Big|_{\theta = \pi = 0}$, we have from (4.12),

$$Y_t + iXY = 0 \quad \text{with} \quad Y(0) = 1.$$

From above structure of X, we have $Y\varphi = 1$. Analogously, we have $\tilde{Y}(t, x, \xi) = \partial_{\pi_2} \partial_{\theta_2} \mathcal{S}(t, x, \xi, \theta, \pi) \Big|_{\theta = \pi = 0}$ which equals to Y.

Calculating V and \tilde{V} analogously, we get both equal to 0.

As
$$Z(t, x, \xi) = \partial_{\pi_2} \partial_{\pi_1} S(t, x, \xi, \theta, \pi) \Big|_{\theta = \pi = 0}$$
 satisfies
 $Z_t + iY^2 = 0$ with $Z(0) = 0$

we have

$$Z(t, x, \xi) = -i \int_0^t ds \, Y^2(s) = -i \int_0^t ds \, \varphi(s)^{-2}.$$

Analogously, we get W = 0.

Therefore using φ , we have

(4.15)
$$\mathcal{S} = \langle x|\xi \rangle + X\theta_1\theta_2 + Y\langle \theta|\pi \rangle + Z\pi_1\pi_2 \text{ with } X = -i\frac{\dot{\varphi}}{\varphi}, \quad Y = \frac{1}{\varphi}, \quad Z = -i\int_0^t ds\,\varphi(s)^{-2}.$$

4.3.3. Continuity equation. Define van Vleck determinant as

$$\mathcal{D}(t, x, \theta, \xi, \pi) = \operatorname{sdet} \begin{pmatrix} \frac{\partial^2 \mathcal{S}(t, x, \theta, \xi, \pi)}{\partial x \partial \xi} & \frac{\partial^2 \mathcal{S}(t, x, \theta, \xi, \pi)}{\partial x \partial \pi} \\ \frac{\partial^2 \mathcal{S}(t, x, \theta, \xi, \pi)}{\partial \theta \partial \xi} & \frac{\partial^2 \mathcal{S}(t, x, \theta, \xi, \pi)}{\partial \theta \partial \pi} \end{pmatrix},$$

and abbreviated as $\mathcal{D} = \mathcal{D}(t, x, \theta, \xi, \pi)$. Using (4.15), we have

$$\mathcal{D} = \varphi^2.$$

4.3.4. Quantization. Using S and $\mathcal{A} = \mathcal{D}^{1/2}$, we define

(4.16)
$$\mathcal{T}_{t}\underline{u}(\theta) = (2\pi)^{-1/2} \int d\xi \, d\pi \, \mathcal{A}e^{i\mathcal{S}}\underline{\hat{u}}(\pi) \\ = (2\pi)^{-1/2} \int d\xi \, e^{i\mathcal{S}_{1}} \left[\int d\pi \, Y^{-1}e^{i\mathcal{S}_{2}}(\underline{\hat{u}}_{1} + \underline{\hat{u}}_{0}\pi_{1}\pi_{2}) \right]$$

where for the sake of notational simplicity, we put

$$S = S_1 + S_2,$$

$$S_1 = \langle x | \xi \rangle + X \theta_1 \theta_2, \quad S_2 = Y \langle \theta | \pi \rangle + Z \pi_1 \pi_2.$$

Since

$$\int d\pi Y^{-1} e^{i\mathcal{S}_2}(\underline{\hat{u}}_1 + \underline{\hat{u}}_2 \pi_1 \pi_2) = Y^{-1} \underline{\hat{u}}_0 + Y^{-1} (iZ + Y^2 \theta_1 \theta_2) \underline{\hat{u}}_1,$$

therefore, remarking $e^{iX\theta_1\theta_2} = 1 + iX\theta_1\theta_2$, we have

$$(1+iX\theta_1\theta_2)(Y^{-1}\underline{\hat{u}}_0+Y^{-1}(iZ+Y^2\theta_1\theta_2)\underline{\hat{u}}_1) = Y^{-1}\underline{\hat{u}}_0+iY^{-1}Z\underline{\hat{u}}_1+[iXY^{-1}\underline{\hat{u}}_0-(Y+XY^{-1}Z)\underline{\hat{u}}_1]\theta_1\theta_2.$$

Finally, we have

(4.17)
$$(\mathcal{T}_{t}\underline{u})(\theta) = u_{0}(t,x) + u_{1}(t,x)\theta_{1}\theta_{2} \quad \text{with}$$
$$u_{0}(t,x) = (2\pi)^{-1/2} \int d\xi \, e^{i\langle x|\xi\rangle} (Y^{-1}\underline{\hat{u}}_{0} + iY^{-1}Z\underline{\hat{u}}_{1}),$$
$$u_{1}(t,x) = (2\pi)^{-1/2} \int d\xi \, e^{i\langle x|\xi\rangle} [iXY^{-1}\underline{\hat{u}}_{0} - (Y + XY^{-1}Z)\underline{\hat{u}}_{1}].$$

Therefore, when $\underline{u}_1 = 0$, we have

(4.18)
$$u_0(t,x) = (2\pi)^{-1/2} \int d\xi \, e^{i\langle x|\xi\rangle} \varphi(t,x,\xi) \underline{\hat{u}}_0(\xi).$$

4.3.5. Calculation of φ . We try to find a power series solution w.r.t. t of (4.14). Decomposing¹⁹ $\varphi(t) = \phi(t)e^{it^2\xi/2}$, ϕ satisfies below from (4.14):

(4.19)
$$\ddot{\phi} + 2it\xi\dot{\phi} + i(1-b)\xi\phi = 0 \text{ with } \phi(0) = 1, \ \dot{\phi}(0) = 0.$$

Putting $\alpha = 2i\xi$, $\beta = i(1-b)\xi$, we rewrite (4.19) as

(4.20)
$$\ddot{\phi} + \alpha \dot{\phi} + \beta \phi = 0 \quad \text{with} \quad \phi(0) = 1, \ \dot{\phi}(0) = 0.$$

Putting $\phi(t) = \sum_{j=0}^{\infty} c_j t^j$ into above, comparing coefficients of each t^{ℓ} , we have, for any ℓ , $c_{2\ell+1} = 0$ and

$$c_{2\ell} = \frac{(-1)^{\ell} (2\alpha)^{\ell}}{(2\ell)!} \left(\frac{\beta}{2\alpha}\right)_{\ell} \quad \text{where} \quad (x)_{\ell} = x(x+1)\cdots(x+\ell-1) = \frac{\Gamma(x+\ell)}{\Gamma(x)}.$$

Remark 4.6. When $\beta = b - 1 = 4k$ with $k = 0, 1, 2, \dots$, since

$$\left(\frac{\beta}{2\alpha}\right)_{\ell} = (-k)_{\ell} = (-k)(-k+1)\cdots(-k+\ell-1) = (-1)^{\ell}(k-1)\cdots(k-\ell+1) = \frac{(-1)^{\ell}k!}{(k-\ell)!}$$

above power series becomes a finite sum, we have

(4.21)
$$\phi(t) = \sum_{j=0}^{k} \frac{4^{j} k!}{(2j)! (k-j)!} (i\xi)^{j} t^{2j}.$$

¹⁹How to find factor $e^{it^2\xi/2}$?

At last, putting (4.21) into (4.18), we get

$$\begin{aligned} u_0(t,q) &= (2\pi)^{-1/2} \int dp \, e^{iqp} \bigg(\sum_{j=0}^k \frac{4^j k!}{(2j)! (k-j)!} (ip)^j t^{2j} \bigg) e^{ipt^2/2} \underline{\hat{u}}_0(p) \\ &= (2\pi)^{-1/2} \int dp \, e^{ip(q+t^2/2)} \bigg(\sum_{j=0}^k \frac{2^{2j} k!}{(2j)! (k-j)!} (ip)^j t^{2j} \bigg) \underline{\hat{u}}_0(p) \\ &= \sum_{j=0}^k \frac{2^{2j} k!}{(2j)! (k-j)!} t^{2j} \underline{u}_0^{(j)} \bigg(q + \frac{t^2}{2} \bigg). \end{aligned}$$

4.4. An application to Random Matrix Theory. In 1983, K.B. Efetov [32] wrote a paper entitled "Supersymmetry and theory of disordered metals" where he derived Wigner's semi-circle law by applying "his superanalysis".

Let \mathfrak{H}_N be a set of $N \times N$ Hermite matrices. Identifying this topologically with \mathbb{R}^{N^2} , we introduce probability measure $d\mu_N(H)$ on \mathfrak{H}_N .

(4.22)
$$d\mu_N(H) = \prod_{k=1}^N d(\Re H_{kk}) \prod_{j
$$P_{N,J}(H) = Z_{N,J}^{-1} \exp\left[-\frac{N}{2J^2} \operatorname{tr} H^* H\right].$$$$

Here, $H = (H_{jk}), H^* = (H_{jk}^*) = (\overline{H}_{kj}) = {}^t\overline{H}, \prod_{k=1}^N d(\Re H_{kk}) \prod_{j < k}^N d(\Re H_{jk}) d(\Im H_{jk})$ is the Lebesgue measure on $\mathbb{R}^{N^2}, Z_{N,J}^{-1}$ is the normalized constant given by $Z_{N,J} = 2^{N/2} (J^2 \pi / N)^{3N/2}$.

Let $E_{\alpha} = E_{\alpha}(H)$ ($\alpha = 1, \dots, N$) be real eigenvalues of a given matrix $H \in \mathfrak{H}_N$. For Dirac's delta δ , we put

(4.23)
$$\rho_N(\lambda) = \rho_N(\lambda; H) = N^{-1} \sum_{\alpha=1}^N \delta(\lambda - E_\alpha(H)),$$

and for any function f on \mathfrak{H}_N , we consider

$$\langle f \rangle_N = \langle f(\cdot) \rangle_N = \int_{\mathfrak{H}_N} d\mu_N(H) f(H).$$

Theorem 4.2 (Wigner's semi-circle law).

(4.24)
$$\lim_{N \to \infty} \left\langle \rho_N(\lambda) \right\rangle_N = w_{sc}(\lambda) = \begin{cases} (2\pi J^2)^{-1} \sqrt{4J^2 - \lambda^2} & \text{for } |\lambda| < 2J, \\ 0 & \text{for } |\lambda| > 2J. \end{cases}$$

This expression is derived by introducing auxiliary odd variables ρ_1, ρ_2 :

(4.25)
$$\left\langle \rho_N(\lambda) \right\rangle_N = \pi^{-1} \Im \int_{\mathfrak{Q}} dQ \left(\left\{ (\lambda - i0)I_2 - Q \right\}^{-1} \right)_{bb} \exp\left[-N\mathcal{L}(Q) \right]$$

Here I_n is the $n \times n$ identity matrix, and

(4.26)

$$\mathcal{L}(Q) = \operatorname{str} [(2J^2)^{-1}Q^2 + \log((\lambda - i0)I_2 - Q)],$$

$$\mathfrak{Q} = \left\{ Q = \begin{pmatrix} x_1 & \rho_1 \\ \rho_2 & ix_2 \end{pmatrix} \mid x_1, x_2 \in \mathfrak{R}_{ev}, \ \rho_1, \rho_2 \in \mathfrak{R}_{od} \right\} \cong \mathfrak{R}^{2|2},$$

$$dQ = \frac{dx_1 dx_2}{2\pi} d\rho_1 d\rho_2,$$

$$((\lambda - i0 - x_1)(\lambda - i0 - ix_2) + \rho_1 \rho_2)$$

$$\left(((\lambda - i0)I_2 - Q)^{-1} \right)_{bb} = \frac{(\lambda - i0 - x_1)(\lambda - i0 - ix_2) + \rho_1 \rho_2}{(\lambda - i0 - x_1)^2 (\lambda - i0 - ix_2)}$$

For 2×2 -supermatrix A, $(A)_{bb}$ is boson-boson(or even-even) part, in this case A_{11} .

$$\left\langle \frac{\delta \mathcal{L}(Q)}{\delta Q}, \tilde{Q} \right\rangle = \frac{d}{d\epsilon} \mathcal{L}(Q + \epsilon \tilde{Q}) \bigg|_{\epsilon=0}$$

critical point is given by

$$\frac{\delta \mathcal{L}(Q)}{\delta Q} = \operatorname{str} \left(\frac{Q}{J^2} - \frac{1}{\lambda - Q}\right) = 0.$$

Defining these as effective saddle points, putting

$$Q_c = \left(\frac{1}{2}\lambda + \frac{1}{2}\sqrt{\lambda^2 - 4J^2}\right)I_2,$$

they have

$$\lim_{N \to \infty} \left\langle \rho_N(\lambda) \right\rangle_N = \pi^{-1} \Im(\lambda - Q_c)_{bb}^{-1} = w_{sc}(\lambda). \qquad \Box$$

Remark 4.7. Though the derivation of expression (4.25) is justified in [70], we can't justify the usage of saddle point method in this setting²¹, at that time. Since I feel this problem has relation to "Laplace's method in function space" in B. Simon [90], so I have an interest.

5. Some examples of FDE

5.1. A Schwinger-Dyson equation.

5.1.1. A derivation of A Schwinger-Dyson equation of first order. A formally given quadratic Lagrangian, we try to give meaning to the first order Schwinger-Dyson equation [53]: Let

(5.1)
$$L(q,v) = \frac{1}{2} \int_{\mathbb{R}} dt (\dot{q}(t)^2 - \omega_0^2 q(t)^2) + \frac{1}{2} \int_{\mathbb{R}^4} dx dt (|v_t(x,t)|^2 - |\nabla v(x,t)|^2) - \lambda \int_{\mathbb{R}^4} dx dt \delta(x) q(t) v(x,t)$$

be given, then critical point of L is calculated by

(5.2)
$$\frac{d}{d\epsilon}L(q+\epsilon\rho,v)\Big|_{\epsilon=0} = \int_{\mathbb{R}} dt(\dot{q}(t)\dot{\rho}(t) - \omega_0^2 q(t)\rho(t)) - \lambda \int_{\mathbb{R}^4} dx dt\delta(x)\rho(t)v(x,t) \\ = \int_{\mathbb{R}} dt \left(-\frac{d^2}{dt^2}q(t) - \omega_0^2 q(t) - \lambda \int_{\mathbb{R}^3} dx\delta(x)v(x,t) \right)\rho(t) \\ = \langle \frac{\delta L(q,v)}{\delta q(t)}, \rho(t) \rangle = 0,$$

and

(5.3)

$$\frac{d}{d\epsilon}L(q, v + \epsilon\varphi)\Big|_{\epsilon=0} = \int_{\mathbb{R}^4} dx dt (v_t(x, t)\varphi_t(x, t) - \nabla v(x, t)\nabla\varphi(x, t)) - \lambda \int_{\mathbb{R}^4} dx dt \delta(x)q(t)\varphi(x, t) = -\int_{\mathbb{R}^4} dx dt (v_{tt}(x, t) - \Delta v(x, t) + \lambda\delta(x)q(t))\varphi(x, t)) = \langle \frac{\delta L(q, v)}{\delta v(x, t)}, \varphi(x, t) \rangle = 0.$$

 $^{20}\mathrm{This}$ fact makes us remind the appearance of \hbar at one place in Feynman Path Integral

 $^{^{21}}$ Therefore, we give a proof of a very small portion which is explained in Y.V.Fyodorov [44]

From these, at least formally, we may suppose the corresponding classical orbit (q(t), v(x, t)) satisfy below:

(5.4)
$$\begin{cases} (\frac{d^2}{dt^2} + \omega_0^2)q(t) = -\lambda \int_{\mathbb{R}^3} dx \delta(x)q(t)v(x,t) \\ \Box v(x,t) = \lambda \delta(x)q(t). \end{cases}$$

Putting the generating functional as

(5.5)
$$\hat{Z} = \hat{Z}(q, v) = e^{i\hbar^{-1}L(q, v)},$$

we define its characteristic functional

(5.6)
$$Z(p,u) = Z_0^{-1} \int d_F q \, d_F v \, \hat{Z}(q,v) e^{-i\hbar^{-1}(\langle q,p \rangle + \langle v,u \rangle)}$$
with $Z_0 = \int d_F q \, d_F v \, \hat{Z}(q,v).$

This Z(p, u) satisfies formally the following equation:

(5.7)
$$\begin{cases} (\frac{d^2}{dt^2} + \omega_0^2) \frac{\delta Z(p, u)}{\delta p(t)} = -i\hbar^{-1} p(t) Z(p, u) - \lambda \int_{\mathbb{R}^3} dx \delta(x) \frac{\delta Z(p, u)}{\delta u(x, t)}, \\ \Box \frac{\delta Z(p, u)}{\delta u(x, t)} = -i\hbar^{-1} u(x, t) Z(p, u) - \lambda \delta(x) \frac{\delta Z(p, u)}{\delta p(t)}. \end{cases}$$

In fact, since

$$\frac{\delta \hat{Z}(q,v)}{\delta q(t)} = -i\hbar^{-1}\left(\frac{d^2}{dt^2} + \omega_0^2\right)q(t)\hat{Z} - i\hbar^{-1}\lambda \int_{\mathbb{R}^3} dx\delta(x)v(x,t)\hat{Z},$$
$$\frac{\delta \hat{Z}(q,v)}{\delta v(x,t)} = -i\hbar^{-1}\Box v(x,t)\hat{Z} - i\hbar^{-1}\lambda\delta(x)q(t)\hat{Z},$$

assuming that functional integration permit integration by parts, we have

$$Z_0^{-1} \int d_F q \, d_F v \, \frac{\delta \hat{Z}(q,v)}{\delta q(t)} e^{-i\hbar^{-1}(\langle q,p \rangle + \langle v,u \rangle)} = i\hbar^{-1}p(t)Z(p,u),$$

$$Z_0^{-1} \int d_F q \, d_F v \, \frac{\delta \hat{Z}(q,v)}{\delta v(x,t)} e^{-i\hbar^{-1}(\langle q,p \rangle + \langle v,u \rangle)} = i\hbar^{-1}u(x,t)Z(p,u),$$

Interchanging integration and differentiation formally, we get

$$\frac{\delta Z(p,u)}{\delta p(t)} = -i\hbar^{-1}Z_0^{-1} \int d_F q \, d_F v \, q(t)\hat{Z}(q,v)e^{-i\hbar^{-1}(\langle q,p \rangle + \langle v,u \rangle)},$$

$$\frac{\delta Z(p,u)}{\delta u(x,t)} = -i\hbar^{-1}Z_0^{-1} \int d_F q \, d_F v \, v(x,t)\hat{Z}(q,v)e^{-i\hbar^{-1}(\langle q,p \rangle + \langle v,u \rangle)}.$$

From these, we have

$$Z_0^{-1} \int d_F q \, d_F v \left(-i\hbar^{-1} \left(\frac{d^2}{dt^2} + \omega_0^2 \right) q(t) \hat{Z} - i\hbar^{-1} \lambda \int_{\mathbb{R}^3} dx \delta(x) v(x,t) \hat{Z} \right) e^{-i\hbar^{-1} \left(\langle q, p \rangle + \langle v, u \rangle \right)}$$
$$= \left(\frac{d^2}{dt^2} + \omega_0^2 \right) \frac{\delta Z(p,u)}{\delta p(t)} + \lambda \int_{\mathbb{R}^3} dx \delta(x) \frac{\delta Z(p,u)}{\delta u(x,t)} = i\hbar^{-1} p(t) Z(p,u). //$$

5.1.2. Reformulation and calculation.

Feynman propagator $E(x,t) = \Box_F^{-1}(x,t)$. As a formal solution of division problem $\Box E(x,t) = \delta(x,t)$ in $\mathcal{D}'(\mathbb{R}^4)$, we have

$$E(x,t) = \lim_{\epsilon \to 0} (2\pi)^{-4} \int_{\mathbb{R}^4} d\tau d\xi \frac{e^{-it\tau + ix\xi}}{-\tau^2 + |\xi|^2 - i\epsilon}$$

whose precise meanings are given in, for example, Gelfand and Shilov [46].

Remark 5.1. Approximate $\delta(x)$ by $\rho_{\epsilon}(x) = \epsilon^{-2}\rho(x/\epsilon)$, i.e. for $\rho(x) = \rho(|x|) \in C_0^{\infty}(\mathbb{R}^3)$, $\rho(x) \ge 0$ and $\int_{\mathbb{R}^3} dx \rho(x) = 1$, making $\epsilon \to 0$ then $\rho_{\epsilon}(x) \to \delta(x)$ in $\mathcal{D}'(\mathbb{R}^3)$.

Lemma 5.1. For any $u \in \mathcal{S}(\mathbb{R}^4)$, $\Box_F^{-1}u \in \mathcal{S}'(\mathbb{R}^4)$. Moreover, when $u \in C_0^{\infty}(\mathbb{R}^4)$, $\lim_{\epsilon \to 0} \langle \rho_{\epsilon}(\cdot), \Box_F^{-1}u(\cdot, t) \rangle$ exists in $H^{-1}(\mathbb{R})$ and formally represented as $\langle \delta, \Box_F^{-1}u \rangle(t)$ or $\langle \delta(\cdot), \Box_F^{-1}u(\cdot, t) \rangle$.

Renormalization. Applying \Box_F^{-1} to the second equation of (5.7), and putting this into the first equation of (5.7), we get

(5.8)
$$(\frac{d^2}{dt^2} + \omega_0^2 - \lambda^2 \int_{\mathbb{R}^3} dx \,\delta(x) \Box_F^{-1}(x,t)) \frac{\delta Z(p,u)}{\delta p(t)} \\ = -i\hbar^{-1} p(t) Z(p,u) + i\hbar^{-1}\lambda \int_{\mathbb{R}^3} dx \,\delta(x) \Box_F^{-1} u(x,t) Z.$$

Here, we approximate $\delta(x)$ by $\rho_{\epsilon}(x)$. In this case, putting

$$(A_{\lambda}^{\epsilon}f)(t) = \left(\frac{d^2}{dt^2} + \omega_0^2\right)f(t) - \lambda^2 \langle \rho_{\epsilon}(\cdot), (\Box_F^{-1}(\rho_{\epsilon}f))(\cdot, t) \rangle$$

using Fourier transformation and Plancherel theorem,

$$\begin{split} (\widehat{A_{\lambda}^{\epsilon}f})(\tau) &= \frac{1}{\sqrt{2\pi}} \int d\tau e^{it\tau} (A_{\lambda}^{\epsilon}f)(t) \\ &= (-\tau^2 + \omega_0^2) \widehat{f}(\tau) + \frac{\lambda^2}{4\pi^2} \int_{\mathbb{R}^3} d\xi \frac{|\widehat{\rho}_{\epsilon}(\xi)|^2}{\tau^2 - |\xi|^2 + i0} \widehat{f}(\tau) \\ &= (-\tau^2 + \omega_0^2) \widehat{f}(\tau) \\ &\quad + \frac{\lambda^2}{4\pi^2} \bigg[\int_{\mathbb{R}^3} d\xi |\widehat{\rho}_{\epsilon}(\xi)|^2 \bigg(\frac{1}{\tau^2 - |\xi|^2 + i0} + \frac{1}{|\xi|^2} \bigg) - \int_{\mathbb{R}^3} d\xi \frac{|\widehat{\rho}_{\epsilon}(\xi)|^2}{|\xi|^2} \bigg] \widehat{f}(\tau). \end{split}$$

Simply disregarding the last term which is ∞ , making $\epsilon \to 0$, then we have

$$\begin{split} \lim_{\epsilon \to 0} \int_{\mathbb{R}^3} d\xi |\hat{\rho}_{\epsilon}(\xi)|^2 \left(\frac{1}{\tau^2 - |\xi|^2 + i0} + \frac{1}{|\xi|^2} \right) \\ &= \int_{\mathbb{R}^3} d\xi \left(\frac{\tau^2 + i0}{(\tau^2 - |\xi|^2 + i0)|\xi|^2} \right) = \int_{S^2} d\omega \int_0^\infty \rho^2 d\rho \frac{\tau^2 + i0}{(\tau^2 - \rho^2 + i0)\rho^2} \\ &= 4\pi \int_0^\infty d\rho \frac{\tau^2 + i0}{\tau^2 - \rho^2 + i0} = -4\pi (2\pi i) \frac{\tau^2}{2|\tau|} = -4i\pi^2 |\tau| \end{split}$$

Therefore

$$\lim_{\epsilon \to 0} (\widehat{A_{\lambda}^{\epsilon}} f)(\tau) = \{ (-\tau^2 + \omega_0^2) - i\lambda^2 |\tau| \} \hat{f}(\tau)$$

so we define

$$A_{\lambda}^{R}f)(t) = \left(\frac{d^{2}}{dt^{2}} + \omega_{0}^{2} - i\lambda^{2}|\frac{d}{dt}|\right)f(t).$$

When $\lambda > 0$, for any $\tau \in \mathbb{R}$, since $-\tau^2 - i\lambda^2 |\tau| + \omega_0^2 \neq 0$, we have

Lemma 5.2. Operator $A_{\lambda}^{R}(\lambda > 0)$ is invertible, and $(A_{\lambda}^{R})^{-1}$ maps $H^{-1}(\mathbb{R})$ to $H^{1}(\mathbb{R})$ as bounded operator. Moreover, for any $p \in H^{1}(\mathbb{R})$, the quantity $\langle (A_{\lambda}^{R})^{-1}p, p \rangle$ is well-defined.

Problem 5.1. Though I wrote this paper, it isn't clear the sentence "Simply disregarding the last term". For example, whether we may find analogous regularization, such as zeta regularization for divergent series or divergent integral in [1] or other summability methods?

Renormalized FDE equation and a result. Now, we have the renormalized FDE equation

(5.9)
$$A_{\lambda}^{R} \frac{\delta Z(p,u)}{\delta p(t)} = -i\hbar^{-1}p(t)Z(p,u) + i\hbar^{-1}\lambda \int_{\mathbb{R}^{3}} dx \delta(x) \Box_{F}^{-1}u(x,t)Z(p,u),$$
$$\Box \frac{\delta Z(p,u)}{\delta u(x,t)} = [i\hbar^{-1}u(x,t) + i\hbar\lambda^{2}(\delta(x)(A_{\lambda}^{R})^{-1}\langle\delta(\cdot), \Box_{F}^{-1}u(\cdot,t)\rangle) - i\hbar^{-1}\lambda(\delta(A_{\lambda}^{R})^{-1}p)(x,t)]Z(p,u)$$

whose solution is obtained explicitly as

(5.10)

$$Z(p,u) = \exp[i(2\hbar)^{-1}\langle (A^R_{\lambda})^{-1}p,p\rangle + i(2\hbar)^{-1}\langle \Box^{-1}_{F}u,u\rangle + i(2\hbar)^{-1}\lambda^2\langle (A^R_{\lambda})^{-1}\langle \delta, \Box^{-1}_{F}u\rangle(t), \langle \delta, \Box^{-1}_{F}u\rangle(t)\rangle - i\hbar^{-1}\lambda\langle (A^R_{\lambda})^{-1}p, \langle \delta, \Box^{-1}_{F}u\rangle(t)\rangle]$$

and all Green function is obtained as bellow: $G^{(n,m)}(t_1, \cdots, t_m, (u_1, s_1))$

$$^{(n,m)}(t_1,\cdots,t_n,(y_1,s_1),\cdots,(y_m,s_m)))$$

= $(i\hbar)^{n+m}\frac{\delta^{n+m}Z(0,0)}{\delta p(t_1)\cdots\delta p(t_n)\delta u(y_1,s_1)\cdots\delta u(y_m,s_m)}.$

Small goals: Enlarging Feynman's idea slightly, following afore mentioned Itzykson-Zuber's claim, we dare to imagine "Quantum mechanics(Quantum field theory) stands for obtaining properties of generating function²² represented by Lagrange or Hamilton function using Feynman measure". To relate these quantities to classical mechanics(classical field theory) and to get rid of the use of non-existing Feynman measure, we take formally the characteristic function which satisfies certain FDE^{23} and give meaning to it after renormalization and solve it!

As is known, to study PDE without getting explicit solution of it, we fully use Lebesgue measure to have the existence proof using integration by parts, change of variables formula under integral sign and Fourier transformations with the method of functional analysis²⁴. As mentioned before, not only there doesn't exist a translational invariant, completely additive measure in function spaces, but also giving meaning to the trace of higher order functional derivatives is hard. For a simple quadratic interaction, we give meaning to some devices eliminating ∞ . This suggests that as is used by physicists, expand w.r.t. interaction parameter and performing Gaussian type integration, after summation method, we get something-like a solution of FDE.

Moreover, in the next paragraph, I give a trial to understand the trace of 2nd order functional derivatives.

5.2. Hopf equation. Let (M, g_{jk}) be a compact Riemannian manifold of dimension d with or without boundary ∂M . We denote by $\overset{\circ}{\mathbf{X}}_{\sigma}(M)$ and $\overset{\circ}{\Lambda}_{\sigma}^{1}(M)$, the space of all solenoidal vector fields on M which vanish near the boundary and that of all divergence free 1-forms on M which vanish near the boundary, respectively. $\tilde{\mathbf{H}}$ (resp. \mathbf{H}) stands for the completion of the space $\overset{\circ}{\Lambda}_{\sigma}^{1}(M)$ (resp. $\overset{\circ}{\mathbf{X}}_{\sigma}(M)$) w.r.t. $\tilde{\mathbf{L}}^{2}$ -norm (resp. \mathbf{L}^{2} -norm).

5.2.1. Differential geometrical expression of Navier-Stokes equation. For a vector field $u = u^j \frac{\partial}{\partial x^j}$, we have the Navier-Stokes equation

(5.11)
$$\frac{\partial u^i}{\partial t} - \nu (\Delta u)^i + (\nabla_u u)^i + p^i = f^i,$$

with

$$\delta u = \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^i} (\sqrt{g} u^i) = 0, \quad u(0, x) = \underline{u}(x) \text{ and } u(t, x) \big|_{x \in \partial M} = 0.$$

Here

$$(\Delta u)^i = \nabla^k \nabla_k u^i - R^i_k u^k, etc.$$

 $^{^{22}}$ For the time being, we don't concern with how these functions relate to the physical quantities observed by experiments 23 Since in general, it contains higher order derivatives, we need new device contrivance such as Colombeau's generalized functions

²⁴Project called "A study of PDE by functional analytic method" initiated by K. Yosida in Japan.

5.2.2. Hopf's motivation. Following Hopf's introduction in [52], we quote

The differential law which governs the motion of a deterministic mechanical system has the symbolic form

$$\frac{du}{dt} = \mathfrak{F}(u)$$

where u is an instantaneous phase of the system and where the right jand side is completely determined by the phase.

There are mechanical systems the phases of which are characterized by a very large number of independent parameters and the phase motions of which are tremendously complicated. Two examples are the classical model of a gas with its very large number of degrees of freedom and the flow of viscous incompressible fluid at a very large value of overall Reynolds number.

In both cases the important task is not the determination of the exact phase motion with an exactly given initial phases but the determination of the statistical properties of the "typical" phase motion. In order to achieve this goal statistical mechanics studies probability distributions of simultaneous phases and their evolution in time resulting from the individual phase motions.

Moreover, he claims to find statistical equilibrium of the system, we need to seek stationary phase distribution. This idea is comparable to find equilibrium state of Hamilton flow, we try to find a stable solution of Liouville equation.

5.2.3. Derivation of Hopf equation. Let $T_t u_0$ be a (some kind of) solution of Navier-Stokes equation (NS) with the initial data u_0 , for a Borel measure $\mu(\omega)$ on **H**, we put $\mu_t(\omega) = \mu(T_t^{-1}(\omega))$. Here, ω is a Borel set on **H**. The characteristic function of this measure $\mu_t(\omega)$ should satisfy the equation, called Hopf equation. More precisely,

(I) the characteristic function of this measure $d\mu_t(\cdot)$ is given by

$$W(t,\eta) = \int_{\mathbf{H}} d\mu_t(u) \, e^{i\langle u,\eta\rangle} = \int_{\mathbf{H}} d\mu(u) \, e^{i\langle T_t u,\eta\rangle}$$

which satisfies

(5.12)
$$\frac{\partial}{\partial t}W(t,\eta) = \int_{M} d_{g}x \left[-i\{\frac{\partial}{\partial x^{k}}\eta_{j}(x) - \Gamma_{jk}^{\ell}(x)\eta_{\ell}\}\frac{\delta^{2}W(t,\eta)}{\delta\eta_{j}(x)\delta\eta_{k}(x)} + \nu(\Delta\eta)_{j}(x)\frac{\delta W(t,\eta)}{\delta\eta_{j}(x)} + i\eta_{j}(x)f^{j}(t,x)W(t,\eta) \right]$$

One of the additional conditions to this equation is

(5.13)
$$\frac{1}{\sqrt{g(x)}}\frac{\partial}{\partial x^j}\left\{\sqrt{g(x)}\frac{\delta W(t,\eta)}{\delta \eta_j(x)}\right\} = 0$$

(5.14)
$$W(0,\eta) = W_0(\eta) \text{ and } W(t,0) = 1.$$

Here, $t \in (0, \infty)$ and

$$\eta = \eta(x) = \eta_j(x)dx^j \in \overset{\circ}{\Lambda}{}^1_{\sigma}(M), \quad f = f(t,x) = f^j(t,x)\partial/\partial x^j \in \overset{\circ}{\mathbf{X}}{}_{\sigma}(M).$$

Given positive definite functional $W_0(\eta)$ on **H** satisfies

(5.15)
$$W_0(\eta) = 1 \text{ and } \frac{1}{\sqrt{g(x)}} \frac{\partial}{\partial x^j} \{ \sqrt{g(x)} \frac{\delta W_0(\eta)}{\delta \eta_j(x)} \} = 0.$$

This equation is something like Liouville equation corresponding to Navier-Stokes equation. In fact, let $u(t,x) = T_t \underline{u}(x)$ be a "solution" for Navier-Stokes equation with initial data $\underline{u} = \underline{u}^j \partial / \partial x^j \in \mathbf{H}$. For Borel measure $\mu(\omega)$ on $\mathbf{H} = L^2_{\sigma}(\Omega)$, put $\mu_t(\omega) = \mu(T^{-1}_t(\omega))$ for any Borel set ω in \mathbf{H} . More explicitly, put

(5.16)
$$\Phi(t,v) = \int e^{i(v,T_t u)} \mu(du) = \int e^{i(v,u)} \mu_t(du),$$

Then, this functional satisfies Hopf equation formally. In the above, for u, we put \tilde{u} as the solenoidal part of u.

5.2.4. How to give the meaning to the trace of 2nd order functional derivative.

Problem 5.2. Our problems here are, whether does there exist a functional $W(t,\eta)$ satisfying (5.12), and how to give meaning to the 2nd order functional derivatives $\frac{\delta^2 W(t,\eta)}{\delta \eta_j(x) \delta \eta_k(x)}$?

Above problem (I) is restated as

(II) Find a Borel measure $\mu(t, \cdot)$ on **H** which satisfies

$$(5.17) \qquad -\int_{0}^{\infty} \int_{\mathbf{H}} d\mu(t,u) \frac{\partial \Phi(t,u)}{\partial t} - \int_{\mathbf{H}} d\mu_{0}(u) \Phi(0,u)$$
$$= \int_{0}^{\infty} \int_{\mathbf{H}} \int_{M} d_{g} x d\mu(t,u) dt \left[-u^{k}(x)u^{j}(x) \frac{\partial}{\partial x^{k}} \frac{\delta \Phi(t,u)}{\delta u^{j}(x)} + \Gamma_{k\ell}^{j}(x)u^{k}(x)u^{\ell}(x) \frac{\delta \Phi(t,u)}{\delta u^{j}(x)} + \nu \nabla_{k} u^{j}(x) \cdot \nabla^{k} \frac{\delta \Phi(t,u)}{\delta u^{j}(x)} - f^{j}(t,x) \frac{\delta \Phi(t,u)}{\delta u^{j}(x)} \right]$$

for any suitable test functionals $\Phi(t, u)$.

Remark 5.2. There appeared only 1st order Functional Derivatives in (II)!

Theorem 5.3 (A part of Theorem A' of Inoue [54]). Let $f(\cdot) \in L^2((0,\infty); \mathbf{V}^{-1})$ be given and suppose a Borel measure μ_0 on \mathbf{H} satisfies

$$\int_{\mathbf{H}} d\mu_0(u)(1+|u|^2) < \infty.$$

Then, there exists a basic family $\{\mu(t,\cdot)\}_{0 < t < \infty}$ of Borel measures on **H** such that

$$\begin{split} \int_0^\infty dt \, \int_{\mathbf{H}} d\mu(t, u) \Phi_t(t, u) + \int_{\mathbf{H}} d\mu_0(u) \Phi_t(0, u) \\ &= \int_0^\infty dt \left[\int_{\mathbf{H}} d\mu(t, u) \left\{ \nu a(u, \tilde{\Phi}_u(t, u)) + b(u, u, \tilde{\Phi}_u(t, u)) - \langle f(t), \Phi_u(t, u) \rangle \right\} \right] \end{split}$$

for any $\Phi \in TF$ with compact support in t, i.e. there exists a constant T depending on Φ such that $\phi(t, \cdot) = 0$ when $t \ge T$.

In the above, forms a and b are defined by

$$\begin{aligned} a(u,v) &= \int_M d_g x \, g_{ij} \nabla_k u^i \nabla^k v^j, \\ b(u,v,w) &= (\nabla_u v,w) = \int_M d_g x \, g_{ij} \{ u^k \frac{\partial}{\partial x^k} v^i + \Gamma^i_{k\ell} u^k v^\ell \} w^j \end{aligned}$$

for $u, v, w \in \mathbf{X}(M)$ with $u = u^j \frac{\partial}{\partial x^j}$ etc.

Definition 5.1 (Definition 2.7 of [54]). A real functional $\Phi(\cdot, \cdot)$ on $[0, \infty) \times \mathbf{V}$ is called TF(=test function) denoted by $\Phi \in TF$ if it satisfies the following.

(1) $\Phi(\cdot, \cdot)$ is continuous on $[0, \infty) \times \mathbf{V}$ and verifies

$$|\Phi_u(t,u)| \le c \text{ and } |\Phi_t(t,u)| \le c + c|u|$$

where $\Phi_u(t,\cdot)$ is regarded as an element in $\tilde{\mathbf{H}}$.

- (2) $\Phi(\cdot, \cdot)$ is Fréchet **H**-differentiable in the direction **V**.
- (3) $\Phi_u(\cdot, \cdot)$ is continuous on $[0, \infty) \times \mathbf{V}$ to $\tilde{\mathbf{V}}^s$ and is bounded, i.e. there exists a constant c depending on Φ such that

$$\|\Phi_u(t,u)\|_s \le c \text{ and } |\Phi_t(t,u)| \le c + c|u| \quad \text{for any} \quad (t,u) \in [0,\infty) \times \mathbf{V}.$$

Appendix A. Some questions on Euler or Navier-Stokes equations

Though the following results are rather recent one which make me not only astonish, but also such impression is criticized because "These results don't match physical reality" by physicists and rather bypassed [83, 84] saying, Euler equation admits various unphysical weak solutions. Even though, one may construct a weak solution with compact support in a torus dimension 2, a gravitational equation which is derived from classical idea

On the other hand, by the theory of general relativity, the following Einstein's field equation

$$G_{\mu\nu} + \Lambda g_{\mu\nu} = \kappa T_{\mu\nu}$$

is derived where $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}$ is the Einstein tensor, $g_{\mu\nu}$ is the metric tensor, Λ is the cosmological constant and κ is the Einstein gravitational constant. Since from this equation, it is claimed the universe begins abruptly by Big Bang, I feel some analogy with above result.

We investigate the initial-boundary value problem

(NS)
$$u_t - \nu \Delta u + u \nabla u - \nabla p = f$$
, div $u = 0$, $u(0, x) = u_0(x)$, $u|_{\partial\Omega} = 0$,

or

(E)
$$u_t + u\nabla u - \nabla p = 0$$
, div $u = f$, $u(0, x) = u_0(x)$, $u \cdot \mathbf{n} = 0$

If we introduce the idea of weak solution for the Euler equation by integration by parts with suitable test functions, we have a result for Euler equation [27, 28].

Theorem A.1 (Theorem 1.7 of Isett and Oh [72]). (Onsager's conjecture on manifolds, sharp version). Let (M, g_{jk}) be a compact Riemannian manifold and $I \subseteq \mathbb{R}$ an open interval. Let (u^{ℓ}, p) be a weak solution to the Euler equations on $I \times M$ such that $u^{\ell} \in L^3_t(I; B^{1/3}_{3,c_0}(M)) \cap C_t(I; L^2(M))$. Then, conservation of energy (E) holds.

Analogously, we have

Theorem A.2 (Theorem1.2(Non-uniqueness of weak solutions for (NS))[20]). There exists $\beta > 0$, such that nonnegative smooth function $e(t) : [0,T] \to \mathbb{R}_{\geq}$, there exists $v \in C_t^0([0,T]; H_x^\beta(\mathbb{T}^3))$ a weak solution of Navier-Stokes equation (NS), such that $\int_{\mathbb{T}^3} dx |v(x,t)|^2 = e(t)$ for all $t \in [0,T]$. Moreover, the associated vorticity $\nabla \times v$ lies in $C_t^0([0,T]; L_x^1(\mathbb{T}^3))$.

Theorem A.3 (\mathbb{R}^2 : 1993 Scheffer [86]). There exists a weak solution, not identically zero, of (E) in \mathbb{R}^2 , having compact support in $(x, t) \in \mathbb{R}^2 \times (0, T)$.

Theorem A.4 (\mathbb{T}^2 : 1997 Shnirelman [88]). There exists a weak solution, not identically zero, of (E) in \mathbb{T}^2 , having compact support in $t \in (0, T)$.

Theorem A.5 (Theorem 1.3 of Buckmaster et al [19]). For any $\epsilon >$, there exists a non-trivial continuous weak solution $v : \mathbb{T}^3 \times \mathbb{R} \to \mathbb{R}^3$ of (E), with $v \in L^1(C^{1/3-\epsilon})$ with compact support in time.

This stands for the collapse of the uniqueness, that is, if there exists very rapidly oscillating exterior force, then there exists a weak solution having space-time compact support.

Theorem A.6 ([86, 88, 89]). There exists a weak solution $u(x,t) \in L^2(\mathbb{R}^2 \times \mathbb{R})$ such that $u(x,t) \equiv 0$ for $|x|^2 + |t|^2 > 1$.

Concerning this, the following paragraph suggests us we need to use new test functions to recognize anomalous phenomena.

[Shnirelman [88]] The weak solution constructed by [86, 88] is not in fact a solution; very strong external forces are present, but they are infinitely-fast oscillating in space, and therefore are indistinguishable from zero in the sense of distributions. The smooth test-functions are not "sensitive" enough to "feel" these forces. This is the fault of sensors, not of forces.

Theorem A.7 (Theorem1.1 of [100]). Let Ω be a bounded domain in \mathbb{R}^3 with C^2 -boundary $\partial\Omega$. Let $u \in L^{\infty}(0,T;L^2(\Omega)) \cap L^2(0,T;H^1(\Omega))$ be a weak solution of the Navier-Stokes equation for any smooth test function $\varphi \in C^{\infty}(\mathbb{R}^+ \times \Omega)$ with compact support, and div $\varphi = 0$. In addition, if

 $u \in L^p(0,T;L^q(\Omega)) \cap L^s(0,T;B^{\alpha,\infty}_s(\Omega))$

for any $\frac{1}{p} + \frac{1}{q} \leq \frac{1}{2}$, $q \geq 4$, s > 2 and for any $\frac{1}{2} + \frac{1}{s} < \alpha < 1$. Then, for any $t \in [0,T]$,

$$\int_{\Omega} dx |u(t,x)|^2 + 2\nu \int_0^t \int_{\Omega} dt dx |\nabla u|^2 = \int_{\Omega} dx |u_0|^2.$$

References

- F. Aghili and S. Tafazoli, Analytical Solution to Improper Integral of Divergent Power Functions Using The Riemann Zeta Function, https://arxiv.org/pdf/1805.10480.pdf
- [2] S. Albeverio and R.J.Hoegh-Krohn, Mathematical Theory of Feynman path integrals, Lect. Notes in Mathematics, 523, Springer, 1976
- [3] S. Albeverio and S. Mazzucchi, Feynman path integrals for polynomially growing potentials, Journal of Functional Analysis 221(2005), pp. 83-121.
- [4] —, The time-dependent quartic oscillator a Feynman path integral approach, Journal of Functional Analysis 238(2006), pp. 471-488.

https://core.ac.uk/download/pdf/82515059.pdf

- [5] —, A survey on mathematical Feynman path integrals: Construction, Asymptotics, Applications, in Quantum Field Theory, eds. B. Fauser, J. Tolksdorf and E. Zeidler (2009), pp. 49-66.
- [6] V.I. Arnold, Sur la geometric differentielle des groupes de Lie de dimension infinite et ses applications à hydrodynamique des fluids parfaits, Ann. Inst. Fourier 16(1966), pp. 319-361.
- [7] A. Ascanelli and M. Cicognani, The fundamental solution for a second order weakly hyperbolic Cauchy problem, Hyperbolic Problems and Regularity Questions, Trends in Mathematics, pp. 17-25, Birkhäuser Verlag, 2006
- [8] C. Bardos, L. Szekelyhidi and E. Wiedemann, Non-uniqueness for the Euler equations: the effect of the boundary, arXiv: 1305.0773

- [9] C. Bardos and E.S. Titi, Euler equations for incompressible ideal fluids, Russian Math. Surveys 62(2007), pp. 409-451.
- [10] —, Loss of smoothness and energy conserving rough weak solutions for the 3d Euler equations, Discrete and continuous dynamical systems series S 3(2010), pp. 185-197.
 - https://www.mccme.ru/ ansobol/otarie/slides/Bardos-Titi-DCDS-Jour-June-2010.pdf
- [11] ——, Mathematics and Turbulence: where do we stand?, J. turbulence 14(2013), pp. 42-76.
- [12] —, Onsager's Conjecture for incompressible Euler equations in bounded domain, Arch.Ration. Mech.Anal. 228(2018), pp. 197-207.
- [13] T. Barker and C. Prange, Localized smoothing for the Navier-Stokes equations and concentration of critical norms near singularities, math-arXiv:1812.09115
- [14] F. Bastianelli, O. Corradini, E. Vassura, Quantum mechanical path integrals in curved spaces and the type A trace anomaly, arXiv: 1702.04247v2, 2017
- [15] E.T. Bell, Mathematics: Queen and Servant of Science, McGraw-Hill, 1951.
- [16] ——Men of mathematics, Touchstone, 1986.
- [17] E. Brezin, Grassmann variables and supersymmetry in the theory of disordered systems, in Applications of field theory to statistical mechanics, ed. L. Garrido, Lecture Notes in Physics 216(1985), pp. 115-123.
- [18] F.A. Berezin and M.S. Marinov, Particle spin dynamics as the Grassmann variant of classical mechanics, Annals of Physics 104(1977), pp. 336-362.
- [19] T. Buckmaster, C. DeLellis and L. Szekelyhidi Jr, Dissipative Euler flows with Onsager-critical spatial regularity, to appear CPAM,
- [20] T. Buckmaster and V. Vicol, Nonuniqueness of weak solutions to the Navier-Stokes equation arXiv: 1709.10033v4, 2018
- [21] E. Caliceti. Distributional Borel Summability of Odd Anharmonic Oscillators, arXiv:math-ph/9910001v1, https://arxiv.org/pdf/math-ph/9910001.pdf
- [22] R. Campoamor-Stursberg and M. Rausch de Traubenberg, Parafermions, tenary algebras and their associated superspace, arXiv:0910.3857 v1, 20 Oct 2009
- [23] M-y. Chi, On the Cauchy problem for a class of hyperbolic equations with the initial data on the parabolic degenerating line, Acta Math. Sinica 8(1958), pp. 521-528.
- [24] C. de Lellis and L. Szekelyhidi, The Euler equations as differential inclusion, Ann.of Math. 170(2009), pp. 1417-1436, arXiv: math/0702079
- [25] —, On admissibility criteria for weak solutions of the Euler equations, Arch.Ration.Mech. Anal.195(2010), 225-260, arXiv:0712.3288
- [26] —, The h-principle and the equations of fluid dynamics, Bull. Amer. Math. Soc. 49(2012), pp. 347-375.
- [27] —, Dissipative Euler flows and Onsager's conjecture, J.Eur.Math.Soc(JEMS) 16(2014), pp. 1467-1505. arxiv: 2012
- [28] —, The h-principle and Onsager's conjecture, Eur. Math. Soc. Newsl. 95(2015), pp. 19-24.
- [29] B. DeWitt, Dynamical theory in curved spaces I. A review of the classical and quantum action principles, Reviews of modern physics 29(1984), pp. 377-397.
- [30] R.J. DiPerna and P.L. Lions, Ordinary differential equations, transport theory and Sobolev spaces, Invent.math. 98(1989), pp. 511-547.
- [31] R.J. DiPerna and A.J. Majda, Oscillations and concentrations in weak solutions of the incompressible fluid equations, Commun.Math.Phys.108(1987), pp. 667-689.
- https://projecteuclid.org/download/pdf_1/euclid.cmp/1104116630
- [32] K.B. Efetov, Supersymmetry and theory of disordered metals, Advances in Physics 32(1983), pp. 53-127.
- [33] —, Random matrices and supersymmetry in disordered systems, arXiv:comd-math/0502322v1, 2005.
- [34] M. Farge, N. Kewlahan, V. Perreier and E. Goirand, Wavelets and Turbulence, Proceedings of the IEEE, 84(1996), pp. 639-669. http://ms.mcmaster.ca/kevla/pdf/articles/Proc_IEEE_v84.pdf
- [35] R. Feynman and A.R. Hibbs, Quantum Mechanics and Path Integrals, McGraw-Hill Book Co., New York, 1965.
- [36] J. Field, Quantum mechanics in space-time: The Feynman path amplitude description of physical optics, de Broglie matter waves and quark and neutrino flavour oscillations, Annals Phys. 321 (2006) 627-707. arXiv:quant-ph/0503026v1
- [37] ——, Derivation of the Schrödinger equation from the Hamilton-Jacobi equation in Feynman's path integral formulation of quantum mechanics, arXiv:1204.0653v1
- [38] C. Foias, Statistical study of Navier-Stokes equations I, II, Rend. Sem. Mat. Padova 48(1973), pp. 219-349, 49(1973), pp. 9-123.
- [39] T. Fujita and S. Kotani, The Onsager-Machlup function for diffusion processes, J. Math. Kyoto Univ. 22(1982), pp. 115-130.
- [40] H. Fujita and N. Sauer, On existence of weak solutions of the Navier-Stokes equations in regions with moving boundaries, J. Fac. Univ. Tokyo Sect IA 17(1970), pp. 403-420.
- [41] D. Fujiwara, A construction of the fundamental solution for the Schrödinger equation, J. D'Analyse Math.35(1979), pp. 41-96.

ATSUSHI INOUE

- [42] —, Remarks on convergence of the Feynman path integrals, Duke Math. J.47(1980), pp. 559-600.
- [43] S. Fukushima, Time-slicing approximation of Feynman path integrals on compact manifolds, arXiv:2009.13281v3
- [44] Y.V. Fyodorov, Basic features of Efetov's supersymmetry approach, in Mesoscopic Quantum Physics, ed E.Akkermans, G. Montambaux, J.-L. Picard and J. Zinn-Justin, 1995, Elsevier Science B.V.
- [45] I.M. Gelfand, Some aspects of functional analysis and algebra, in "International Congress of Mathematicians in Amsterdam", (1954) pp. 253-276,
- https://www.mathunion.org/fileadmin/ICM/Proceedings/ICM1954.1/ICM1954.1.ocr.pdf
- [46] I.M. Gelfand and G.E. Shilov, Generalized functions, vol1: Properties and Operations, Academic Press, New York. 1964.
- [47] E. Getzler, Pseudo-differential operators on supermanifolds and the Atiyah-Singer index theorem, Commun. Math. Phys. 92(1983), pp. 163-178.
- [48] S. Graffi, V. Grecchi and B. Simon, Borel summability: Application to the anharmonic oscillator, Physics Letters 32b(1970), pp. 631-634.
- [49] M. Gromov, Local and global in geometry, http://www.ihes.fr/~gromov/PDF/1[107].pdf
- [50] S.W. Hawking, Zeta function regularization of path integrals in curved spacetime, Commun. Math. Phys. 55 (1977) pp. 133-148.
- [51] T.I. Hesla, Collisions of smooth bodies in viscous fluids: A mathematical investigation, ProQuest Dissertations And Theses; Thesis (Ph.D.)–University of Minnesota, 2004.
- [52] E. Hopf, Statistical hydrodynamics and functional calculus, J.Rational Mech.Anal.1 (1952), pp. 87-123.
- [53] A. Inoue, Some examples exhibiting the procedures of renormalization and gauge fixing –Schwinger-dyson equations of first order, Kodai Math. J. 9(1986), pp.134-160.
- [54] —, Strong and classical solutions of the Hopf equation -an example of Functional Derivative Equation of second order, Tôhoku Math.J. 39(1987), pp. 115-144
- [55] —, On functional derivative equations invitation to analysis on functional spaces I, II(in Japanese), suugaku 42(1990) pp. 170-177, pp.261-265
- [56] —, On a construction of the fundamental solution for the free Weyl equation by Hamiltonian path-integral method an exactly solvable case with "odd variable coefficients", Tôhoku J.Math.50(1998), pp. 91-118.
- [57] —, On a construction of the fundamental solution for the free Dirac equation by Hamiltonian path-integral method –another interpretation of Zitterbewegung, Japanese J.Math.24(1998), pp. 297-334.
- [58] —, On a "Hamiltonian path-integral" derivation of the Schrödinger equation, Osaka J. Math. 36(1999), pp. 861-904.
- [59] —, A partial solution for Feynman's problem –a new derivation of the Weyl equation, Mathematical Physics and Quantum Field Theory, Electron.J.Diff.Eqns., Conf.04, 2000, pp. 121-145.
- [60] ——, Definition and characterization of supersmooth functions on superspace based on Fréchet-Grassmann algebra, https://arxiv.org/pdf/0910.3831.pdf
- [61] ——, What is Onsager conjecture? -has that some relations with the beginning and end of the universe, OPseminar, 3 September 2016, at Yonago.
- [62] ——, Byproduct of studying Onsager conjecture, can we have a little problem touching ourselves? what is the very weak solution of Euler equation?, OPseminar, 15 July 2017, at Ise
- [63] ——, Personal byproduct of studying Onsager conjecture, can we find a quick problem?, OPseminar, 1 September 2018, at Tokushima
- [64] ——, Introduction to Superanalysis and its Applications –as an example of non-commutative analysis, 2018, in preparation
- [65] —, Anharmonic oscillator by superanalysis why complex number representation is used in quantum mechanics even though observed quantities are real?, OPseminar, Zoom, 28 August 2022.
- [66] —, A parametrix of FIO type for the Schrödinger equation with quartic potentials, in preparation
- [67] A. Inoue and T. Funaki, On a new derivation of Navier-Stokes equation, Commun.Math.Phys. 65(1979), pp. 83-90.
- [68] A. Inoue and Y. Maeda, On integral transformations associated with a certain Lagrangian- as a prototype of quantization, J.Math.Soc.Japan 37(1985), pp. 219-244.
- [69] —, On a construction of a good parametrix for the Pauli equation by Hamiltonian path-integral method an application of superanalysis, Japanese J. Math.29(2003) pp. 27-107.
- [70] A. Inoue and Y. Nomura, Some refinements of Wigner's semi-circle law for Gaussian random matrices using superanalysis, Asymptotic Analysis 23(2000), pp. 329-375.
- [71] A. Inoue and M. Wakimoto, On existence of solutions of the Navier-Stokes equation in a time dependent domain, J. Fac. Sci. Univ. Tokyo Sect. IA Math. 24 (1977), pp. 303-319.
- [72] P. Isett and S-J.Oh, A heat flow approach to Onsager's conjecture for the Euler equations on manifolds, arXiv: 1310.7947v2
- [73] C. Itzykson and J-B. Zuber, Quantum Field Theory, 1979, MaGRAW-HILL INTERNATIONAL EDITIONS,

- [74] A. Júngel, J.L. López and J.Montejo-Gámez, A new derivation of the quantum Navier-Stokes equations in the Wigner-Fokker-Planck approach, Journal of Statistical Physics 145 (2011) pp. 1661-1673.
- [75] T. Kato, Remarks on zero viscosity limit for nonstationary Navier-Stokes flows with boundary, in seminar on nonlinear differential equations, ss. Chern (ed) Springer-Verlag (1984), pp 85-98.
- [76] H. Kumano-go, Pseudo-Differential Operators, Cambridge-Massachusettes-London, The MIT Press, 1981.
- [77] Y.I. Manin, New dimensions in geometry, Lecture Notes in Mathematics 1111, pp. 59-101, Berlin-New York-Tokyo-Heidelberg, Springer-Verlag, 1985.
- [78] J.L. Martin, Generalized classical dynamics, and the 'classical analogue' of a Fermi oscillator, Proc. Roy.Soc. A251(1959), pp. 536-542.
- [79] —, The Feynman principle for a Fermi system, Proc. Roy.Soc. A251(1959), pp. 543-549.
- [80] T. Matsumoto, H. Nawa and T. Sakajo, Turbulence as stochastic process, preprint.
- [81] T. Miyakawa and Y. Teramoto, Existence and periodicity of weak solutions of the Navier-Stokes equations in a time dependent domain, Hiroshima Math. J. 12(1982), pp. 523-528.
- [82] F. Rindler, Directional oscillations, concentrations, and compensated compactness via microlocal compactness forms, ArXiv: 1212.1430v3.
- [83] T. Sakajo, A singular solution for the equation of fluid mechanics concerning analysis of turbulent phenomena(in Japanese), sugaku 71(2019), pp. 178-200.
- [84] —, Singular solutions of nonlinear hydrodynamic equations arising in turbulence theory, Sugaku Expositions vol. 36 No. 1 pp. 93–117 (2023) (doi:10.1090/suga/478)
- [85] R. Saldi, On the existence of weak solutions of a non-linear mixed problem for the Navier-Stokes equations in a time dependent domain, J. Fac. Sci. Univ. Tokyo Sect.IA 32(1985), pp. 213-221.
- [86] V. Scheffer, An inviscid flow with compact support in space-time, The Journal of Geometric Analysis, 3(1973), pp. 343-401.
- [87] A. Shnirelman, On the non-uniqueness of weak solution of the Euler equations, Journées Équations aux dérivées partielles (1996), pp. 1-10.
- [88] —, On the nonuniqueness of weak solution of the Euler equation, Comm. Pure Appl. Math.50(1997), pp. 1261-1286.
- [89] ——, Weak solutions with decreasing energy of incompressible Euler equations, Commun.Math.Phys. 210(2000). pp. 541-603.
- [90] B. Simon, Functional Integration and Quantum Mechanics, 1979, Academic Press, New York San Francisco London.
- [91] O.G. Smolyanov and S.V. Fomin, Measures on linear topological spaces, Russian Math.Surveys 31(1976), pp. 1-53.
- [92] E.S. Swanson, A primer on functional methods and the Schwinger-Dyson equations, arXiv:1008.4337v2
- [93] N. Takayama, Propagation of singularities of solutions of the Euler-Darboux equation and a global structure of the space of holonomic solutions I. Funkcialj Ekvacioj, 35(1992), pp.343-403.
- [94] Y. Takahashi and S. Watanabe, The probability functionals (Onsager-Machlup functions) of diffusion processes, In: Williams, D. (eds) Stochastic Integrals. Lecture Notes in Mathematics, vol 851. pp. 433-463. Springer, Berlin, Heidelberg.
- [95] J.G. Tormo, Asymptotic series in quantum mechanics: anharmonic oscillator, http://diposit.ub.edu/dspace/bitstream/2445/59765/1/TFG-Garcia-Tormo-Joan.pdf
- [96] M. Vishik, Incompressible flows of an ideal fluid with vorticity in borderline spaces of Besov type, Ann. scient.Ec.Norm.Sup 32(1999), pp. 769-812.
- [97] V.S. Vladimirov and I.V. Volovich, Superanalysis I. Differential calculus, Theor. Math. Phys. 59(1983), pp. 317-335.
- [98] —, Superanalysis II. Integral calculus, Theor. Math. Phys. 60(1984), pp. 743-765.
- [99] E. Witten, Supersymmetry and Morse theory, J.Diff.Geom.17(1982), pp. 661-692.
- [100] C.Yu, The energy equality for the Navier-Stokes equations in bounded domains, arXiv:1802.07661v1