

A CLASS OF NUMERICAL SEMIGROUPS DEFINED BY KUNZ AND WALDI - THEIR PRINCIPAL MATRICES AND STRUCTURE

SRISHTI SINGH AND HEMA SRINIVASAN

ABSTRACT. In this paper, we explore a class of numerical semigroups initiated by Kunz and Waldi containing two coprime numbers $p < q$, which we call KW semigroups. We characterize KW numerical semigroups by their principal matrices. We present a necessary and sufficient criterion for a matrix to be the principal matrix of a KW semigroup. An explicit description of the minimal resolutions of numerical semigroups in the same class with small embedding dimensions 3 and 4 is given. We give a generalization of this notion to three dimensions using lattice paths under a plane and present some preliminary results and questions.

1. INTRODUCTION

A numerical semigroup $\langle \mathbf{a} \rangle$ is a submonoid of \mathbb{N} minimally generated by $\mathbf{a} = \{a_1, \dots, a_n\} \subseteq \mathbb{N}$ where $\gcd(a_1, \dots, a_n) = 1$. Such a semigroup will contain all but finitely many positive integers, called the gaps. The largest gap is called the Frobenius number, $F(\mathbf{a})$.

This paper is inspired by recent remarkable works of Kunz and Waldi in [5] where they build numerical semigroups of the same multiplicity p by filling in gaps. Any semigroup of embedding dimension 2 is symmetric, i.e., the number of gaps is exactly half the number of elements less than the Frobenius number. Thus, a semigroup generated by two relatively prime positive integers $p < q$, is symmetric and of multiplicity p . Kunz and Waldi build extensions of these semigroups by filling in gaps larger than p to create semigroups of higher embedding dimensions with the same multiplicity p . All the gaps of $\langle p, q \rangle$ are of the form $pq - xp - yq$ for some positive integers $x, y \in \mathbb{N}$, with $pq - p - q$ being the largest, the Frobenius number.

In [5], Kunz and Waldi study numerical semigroups of the form $\langle p, q, h_1, \dots, h_{n-2} \rangle$ where $p < q$ are relatively prime positive integers. Further, they analyze a special class of

Date: May 7, 2024.

MSC Primary: 13c05, 13c70; *Secondary:* 13D02.

Keywords: Semigroup rings, monoids, critical binomials.

semigroups with embedding dimension n , which we denote by $KW(p, q)$, by filling in gaps where $x \leq p/2, y \leq q/2$. In particular, they prove a structure theorem for this class:

Theorem 1.1. (*Corollary 3.1, Theorem (Appendix) [5]*) *Let $A = \langle p, q, h_1, \dots, h_{n-2} \rangle$ with $h_i = pq - x_i p - y_i q$, for some $0 < 2x_i \leq q, 0 < 2y_i \leq p, 1 \leq i \leq n-2$. Then the semigroup ring $k[A]$ is of type $n-1$ and its relation ideal is generated by $\binom{n}{2}$ elements.*

In this paper, we begin by studying the principal matrices of these semigroups. A “principal matrix” is an $n \times n$ integer matrix associated with a numerical semigroup of embedding dimension n . If $\mathbf{a} = \{a_1, \dots, a_n\}$ minimally generates a numerical semigroup, then its principal matrix $P(\mathbf{a}) = (a_{ij})$ is an $n \times n$ matrix, $a_{ij} \geq 0, i \neq j$, where $-a_{ii}$ is the smallest positive integer such that $a_{ii}x_i + \sum_{i \neq j} a_{ij}x_j = 0$. The diagonal entries of $P(\mathbf{a})$ are uniquely determined for a given \mathbf{a} but the matrix itself need not be unique.

Indeed for a general numerical semigroup $\langle \mathbf{a} \rangle$, the rank of $P(\mathbf{a}) \leq n-1$ and when it is $n-1$, one can recover \mathbf{a} from the $n-1$ order minors of any of the $n-1$ rows. Further, the rank of $P(\mathbf{a}) \geq n/2$, ([2]). As such, this matrix holds vital information about the semigroup itself.

For instance, a criterion is established for Gorenstein monomial curves using principal matrices in [1] and [3]. Additionally, [2] presents a characterization for principal matrices of numerical semigroups that are gluing of numerical semigroups.

In this paper, we study the principal matrices of the semigroups in $KW(p, q)$. We establish a necessary and sufficient criterion for a matrix to be the principal matrix of a numerical semigroup within Kunz and Waldi’s class in Theorem 3.1. As a consequence, we get

Corollary 1.2. *Let $3 \leq p < q$ be prime numbers. Then the principal matrix of a numerical semigroup $A = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW((p, q))$ is of the form*

$$P(A) = T(\alpha, \beta) = \begin{bmatrix} -\left(\frac{q+\alpha_{n-2}}{2}\right) & \frac{p-\beta_{n-2}}{2} & 0 & 0 & \cdots & 0 & 1 \\ \frac{q-\alpha_1}{2} & -\left(\frac{p+\beta_1}{2}\right) & 1 & 0 & \cdots & 0 & 0 \\ \alpha_1 & \beta_1 & -2 & 0 & \cdots & 0 & 0 \\ \alpha_2 & \beta_2 & 0 & -2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \alpha_{n-2} & \beta_{n-2} & 0 & 0 & \cdots & 0 & -2 \end{bmatrix},$$

for some odd positive integers α_i, β_i satisfying $q > \alpha_1 \geq \dots \geq \alpha_{n-2} \geq 0$ and $0 < \beta_1 \leq \dots \leq \beta_{n-2} < p$. Conversely, any $n \times n$ matrix of the form $P(A) = T(\alpha, \beta)$ above is a principal matrix of a $KW(p, q)$ numerical semigroup.

As an interesting fact, we observe that the defining ideals of semigroup rings $k[H]$ for $H \in KW(p, q)$ can be written as sums of 2×2 minors of $2 + \binom{n-2}{2}$ matrices of size 2×3 .

The paper is organized as follows: In section 2, we define the notion of principal matrices, the class $KW(p, q)$ of numerical semigroups and some notation. The proof of our main result (Theorem 3.1) is in section 3. In section 4, various auxiliary results pertaining to the minimal resolutions and relation ideal of semigroup rings of $KW(p, q)$ are given. We end this paper by extending the class of KW-semigroups in section 5, and conclude with some open questions.

2. PRELIMINARIES

If a set of relatively prime positive integers $\mathbf{a} = \{a_1, \dots, a_n\}$ minimally generates a semigroup $\langle \mathbf{a} \rangle$, then, there are equations

$$c_i a_i = \sum_{j \neq i, 1 \leq j \leq n} a_{ij} a_i, \quad 1 \leq i \leq n.$$

In the associated semigroup ring $k[\mathbf{a}] = k[t^{a_1}, \dots, t^{a_n}] = k[x_1, \dots, x_n]/I_{\mathbf{a}}$,

$$f_i = x^{c_i} - \prod_{j \neq i, 1 \leq j \leq n} x_j^{a_{ij}}$$

are among a set of minimal generators of $I_{\mathbf{a}}$. Further, $\sqrt{(f_1, \dots, f_n)} = \sqrt{I_{\mathbf{a}}}$. These f_i are therefore called critical binomials of the numerical semigroup $\langle \mathbf{a} \rangle$ or the binomial toric ideal $I_{\mathbf{a}}$. This is also contained in the following $n \times n$ matrix, which is called the Principal Matrix of the numerical semigroup.

Definition 2.1. (Definition 1, [2]) Let $\mathbf{a} = \{a_1, \dots, a_n\}$ be a set of positive integers min-

imally generating a semigroup $\langle \mathbf{a} \rangle$. $A = \begin{bmatrix} -c_1 & a_{12} & \cdots & a_{1n} \\ a_{21} & -c_2 & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n1} & a_{n2} & \cdots & -c_n \end{bmatrix}$ is called a principal

matrix of $\langle \mathbf{a} \rangle$ if $A\mathbf{a} = 0$, and c_i is the smallest positive integer such that $c_i a_i \in \langle \mathbf{a} - \{a_i\} \rangle$ for all $1 \leq i \leq n$.

Although the diagonal entries $-c_i$ are uniquely determined, a_{ij} are not always unique. The sequence of positive integers \mathbf{a} can be recovered from a given principal matrix A of rank $n - 1$ by factoring out the greatest common divisor of the entries of any nonzero column of the adjoint of A and taking its absolute value.

Example 2.2. *The principal matrix of the numerical semigroup $\mathbf{a} = \langle 5, 7, 11, 13 \rangle$ is*

$$P(\mathbf{a}) = \begin{pmatrix} -4 & 1 & 0 & 1 \\ 2 & -3 & 1 & 0 \\ 3 & 1 & -2 & 0 \\ 1 & 3 & 0 & -2 \end{pmatrix}$$

We can recover \mathbf{a} simply from the 3×3 minors of the first three rows.

In this paper, we study the principal matrices of numerical semigroups in $KW(p, q)$, constructed in [4] as follows:

Definition 2.3. *Let $p, q \in \mathbb{N}$ be relatively prime with $3 \leq p < q$. The set of Kunz-Waldi Semigroups associated to $p < q$, denoted by $KW(p, q)$, is the set of all numerical semigroups H with $\langle p, q \rangle \subset H \subset \langle p, q, r \rangle$, where $r = \begin{cases} \frac{p}{2}, & p \text{ even} \\ \frac{q}{2}, & q \text{ even} \\ \frac{p+q}{2}, & p \text{ and } q \text{ odd.} \end{cases}$*

All $H \in KW(p, q)$ are in one-to-one correspondence to the lattice paths in the rectangle $R \subseteq \mathbb{R}^2$ with the corners $(0, 0)$, $(0, p' - 1)$, $(q' - 1, p' - 1)$, and $(q' - 1, 0)$, where $p' = \lfloor \frac{p}{2} \rfloor$ and $q' = \lfloor \frac{q}{2} \rfloor$. The only $H \in KW(p, q)$ with embedding dimension $e(H) = 2$ are $\langle p, q \rangle$, $\langle p/2, q \rangle$, and $\langle p, q/2 \rangle$. For any other $H \in KW(p, q)$, the minimal generating set is $\{p, q, h_1, \dots, h_{e(H)-2}\}$, where $h_i = pq - x_i p - y_i q$, if (x_i, y_i) for $1 \leq i \leq e(H) - 2$ are the corners of the lattice path defining H . This gives a sufficient condition for H to be in $KW(p, q)$ and we incorporate it in the definition:

$H \in KW(p, q)$ if

$$(1) \quad 2x_i \leq q \quad \text{and} \quad 2y_i \leq p \quad \forall 1 \leq i \leq e(H) - 2$$

Remark 1. *Thus, $H = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW(p, q)$ if and only if $h_i = pq - x_i p - y_i q$ satisfying*

$$2x_i \leq q \quad \text{and} \quad 2y_i \leq p \quad \forall 1 \leq i \leq e(H) - 2,$$

$$0 < x_1 < \dots < x_{n-2}, \text{ and } y_1 > \dots > y_{n-2} > 0.$$

3. PRINCIPAL MATRIX OF KW SEMIGROUPS

In this section, we prove one of our main theorems (3.1), which provides an explicit structure of the principal matrix for any $H \in KW(p, q)$.

Theorem 3.1. *Let $p < q$ be two relatively prime positive integers. Choose positive integers $q > \alpha_1 > \alpha_2 > \dots > \alpha_{n-2} \geq 0$, of the same parity and $0 \leq \beta_1 < \beta_2 < \dots < \beta_{n-2} < p$ of the same parity. Let $h_i = \frac{p\alpha_i + q\beta_i}{2}$, $1 \leq i \leq n-2$.*

Then the matrix

$$(2) \quad T := T(\alpha, \beta) = \begin{bmatrix} -\left(\frac{q+\alpha_{n-2}}{2}\right) & \frac{p-\beta_{n-2}}{2} & 0 & 0 & \dots & 1 \\ \frac{q-\alpha_1}{2} & -\left(\frac{p+\beta_1}{2}\right) & 1 & 0 & \dots & 0 \\ \alpha_1 & \beta_1 & -2 & 0 & \dots & 0 \\ \alpha_2 & \beta_2 & 0 & -2 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_{n-2} & \beta_{n-2} & 0 & 0 & \dots & -2 \end{bmatrix}$$

is a principal matrix for the semigroup $H = \{p, q, h_1, \dots, h_{n-2}\}$ if and only if $H \in KW(p, q)$ provided any of the following is true:

- (i) p and q are both odd
- (ii) p is even and $2h_1 \neq p\alpha_1$
- (iii) p is even, $2h_1 = p\alpha_1$ but $q \leq 2\alpha_1 - \alpha_{n-2}$
- (iv) q is even and $2h_{n-2} \neq q\beta_{n-2}$
- (v) q is even and $p \leq 2\beta_{n-2} - \beta_1$.

If none of the above conditions is true and if p is even, then the principal matrix of H is

$$\begin{bmatrix} -\alpha_1 & 0 & 2 & 0 & \dots & 0 \\ \frac{q-\alpha_1}{2} & -\left(\frac{p+\beta_1}{2}\right) & 1 & 0 & \dots & 0 \\ \alpha_1 & \beta_1 & -2 & 0 & \dots & 0 \\ \alpha_2 & \beta_2 & 0 & -2 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_{n-2} & \beta_{n-2} & 0 & 0 & \dots & -2 \end{bmatrix}$$

and if q is even, it is

$$\begin{bmatrix} -\left(\frac{q+\alpha_{n-2}}{2}\right) & \frac{p-\beta_{n-2}}{2} & 0 & 0 & \dots & 1 \\ 0 & -\beta_{n-2} & 0 & 0 & \dots & 2 \\ \alpha_1 & \beta_1 & -2 & 0 & \dots & 0 \\ \alpha_2 & \beta_2 & 0 & -2 & \dots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_{n-2} & \beta_{n-2} & 0 & 0 & \dots & -2 \end{bmatrix}.$$

Proof. First, suppose that one of the conditions (i)-(v) are satisfied. Set $\mathbf{h} = \{p, q, h_1, \dots, h_{n-2}\}$. By construction, h_i is a minimal generator and $2h_i = p\alpha_i + q\beta_i$. Next, we note that

$$(3) \quad p \left(\frac{q + \alpha_{n-2}}{2} \right) = \frac{pq}{2} + h_{n-2} - \frac{q\beta_{n-2}}{2} = q \left(\frac{p - \beta_{n-2}}{2} \right) + h_{n-2} \in \langle \mathbf{h} \setminus \{q\} \rangle$$

since p and β_n have the same parity, and

$$(4) \quad q \left(\frac{p + \beta_1}{2} \right) = \frac{pq}{2} + h_1 - \frac{p\alpha_1}{2} = p \left(\frac{q - \alpha_1}{2} \right) + h_1 \in \langle \mathbf{h} \setminus \{p\} \rangle,$$

since q and α_1 are of same parity.

It remains to show that $\frac{q + \alpha_{n-2}}{2}$ and $\frac{p + \beta_1}{2}$ are the smallest integers that make (3) and (4) true, respectively.

Suppose $\delta q = \gamma p + \mu_1 h_1 + \dots + \mu_{n-2} h_{n-2}$, for some $\delta, \gamma, \mu_i (1 \leq i \leq n-2) \in \mathbb{N}$. Not all μ_i are zero as p and q are coprime. Then

$$q \left(\delta - \frac{\mu_1 \beta_1 + \dots + \mu_{n-2} \beta_{n-2}}{2} \right) = p \left(\gamma + \frac{\mu_1 \alpha_1 + \dots + \mu_{n-2} \alpha_{n-2}}{2} \right)$$

From the above equation, it follows that p divides $2\delta - (\mu_1 \beta_1 + \dots + \mu_{n-2} \beta_{n-2})$, so

$$(5) \quad 0 \leq px = 2\delta - (\mu_1 \beta_1 + \dots + \mu_{n-2} \beta_{n-2})$$

for some $x \in \mathbb{N}$. Now, if q is odd, then none of the α_i are zero and hence $x > 0$.

Observe that necessarily some $\mu_j \geq 1$, so in (5),

$$\begin{aligned} \delta &= \frac{px}{2} + \frac{\sum_{i \neq j, 1 \leq i \leq n} \mu_i \beta_i}{2} + \frac{p + \beta_j}{2} - \frac{p + \beta_j}{2} \\ &= \frac{p + \beta_j}{2} + \frac{p}{2} (x - 1) + \frac{\beta_j}{2} (\mu_j - 1) + \sum_{i \neq j} \mu_i \beta_i \\ &\geq \frac{p + \beta_j}{2} \\ &\geq \frac{p + \beta_1}{2} \end{aligned}$$

We have shown that $\frac{p + \beta_1}{2}$ is the smallest positive integer such that $(\frac{p + \beta_1}{2})p \in \langle \mathbf{h} \setminus \{p\} \rangle$.

Even if q is even, $x > 0$ unless $\alpha_{n-2} = 0$ and $2h_{n-2} = p\beta_{n-2}$. In that case, T will still be principal as long as $p \leq 2\beta_{n-2} - \beta_1$.

Similarly, suppose

$$\delta' p = \gamma' q + \nu_1 h_1 + \dots + \nu_{n-2} h_{n-2},$$

for some $\delta', \gamma', \nu_i (1 \leq i \leq n-2) \in \mathbb{N}$, necessarily some $\nu_j \geq 1$. Isolating p and q as above, we obtain that

$$qy = 2\delta' - (\nu_1\alpha_1 + \dots + \nu_{n-2}\alpha_{n-2})$$

for some $y \geq 1$ if p is not even so that none of the β_i are zero.

Then

$$\begin{aligned} \delta' &= \frac{qy}{2} + \frac{\nu_1\alpha_1 + \dots + \nu_{n-2}\alpha_{n-2}}{2} + \frac{q + \alpha_j}{2} - \frac{q + \alpha_j}{2} \\ &\geq \frac{q + \alpha_j}{2} \\ &\geq \frac{q + \alpha_{n-2}}{2} \end{aligned}$$

Again, if p happens to be even, T will still be principal as long as one of the conditions (i)-(v) is satisfied. Thus, T is a principal matrix of H . Moreover, each h_i , $1 \leq i \leq n-2$ can be expressed as follows:

$$(6) \quad h_i = \frac{p\alpha_i + q\beta_i}{2} = pq - \left(\frac{q - \alpha_i}{2}\right)p - \left(\frac{p - \beta_i}{2}\right)q$$

By hypotheses, $\alpha_i > -1$ and $\beta_i > -1$, so that $2\left(\frac{q - \alpha_i}{2}\right) = q - \alpha_i < q + 1$, and $2\left(\frac{p - \beta_i}{2}\right) < p + 1$. In view of the sufficient condition to be a KW-semigroup (1), it now follows that $H \in KW(p, q)$.

Finally, say p is even, and $2h_1 = p\alpha_1$ with $q > 2\alpha_1 - \alpha_{n-2}$. Then q is odd. And the minimal relation on p is $\alpha_1 p = 2h_1$. Hence the principal matrix is

$$\begin{bmatrix} -\alpha_1 & 0 & 2 & 0 & \cdots & 0 \\ \frac{q - \alpha_1}{2} & -\left(\frac{p + \beta_1}{2}\right) & 1 & 0 & \cdots & 0 \\ \alpha_1 & \beta_1 & -2 & 0 & \cdots & 0 \\ \alpha_2 & \beta_2 & 0 & -2 & \cdots & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\ \alpha_{n-2} & \beta_{n-2} & 0 & 0 & \cdots & -2 \end{bmatrix}$$

The case when q is even can be proved similarly. □

Consequently, we obtain a characterization of the principal matrix of any $H \in KW(p, q)$ when $3 < p < q$ are prime. This is Corollary 1.2. Recall from the Introduction:

Corollary (1.2). *Let p be odd and $q > p$ be relatively prime to p . Then the principal matrix of a numerical semigroup $A = (p, q, h_1, \dots, h_{n-2}) \in KW((p, q))$ is of the form*

$$P(A) = T(\alpha, \beta) = \begin{bmatrix} -\left(\frac{q+\alpha_{n-2}}{2}\right) & \frac{p-\beta_{n-2}}{2} & 0 & 0 & \cdots & 0 & 1 \\ \frac{q-\alpha_1}{2} & -\left(\frac{p+\beta_1}{2}\right) & 1 & 0 & \cdots & 0 & 0 \\ \alpha_1 & \beta_1 & -2 & 0 & \cdots & 0 & 0 \\ \alpha_2 & \beta_2 & 0 & -2 & \cdots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \alpha_{n-2} & \beta_{n-2} & 0 & 0 & \cdots & 0 & -2 \end{bmatrix},$$

for some odd positive integers α_i, β_i satisfying $q > \alpha_1 \geq \dots \geq \alpha_{n-2} \geq 0$ and $0 < \beta_1 \leq \dots \leq \beta_{n-2} < p$. Conversely, any $n \times n$ matrix of the form $P(A) = T(\alpha, \beta)$ above is a principal matrix of a $KW(p, q)$ numerical semigroup.

Proof. Note that $h_i = pq - px_i - qy_i$, for all $1 \leq i \leq n-2$, with $0 < 2x_i \leq q, 0 < 2y_i \leq p$. Since $H \in KW(p, q)$, it satisfies the hypotheses of Theorem 3.1. Next, since $p \geq 3$ is prime, it must be odd so necessarily $y_1 \neq p/2$. Thus, we get that the principal matrix of H is of the form $T(\alpha, \beta)$ from equation (2).

The converse is a direct consequence of Theorem 3.1. \square

Remark 2. *The maximal embedding dimension for a numerical semigroup H in $KW(p, q)$ is precisely $2 + \lfloor \frac{p}{2} \rfloor$. This maximum is achieved when the set of minimal generators of H is a part of an arithmetic sequence $(p, p+d=q, \dots, q+(p-1)d)$. However, the building of this extremal semigroup is via the gaps $h_i = q + (p-1)d = pq - p\lfloor \frac{q}{2} \rfloor - q$ and continue. For example, with $p = 7, q = 11$ we see that $h_1 = 77 - 35 - 11 = 31, h_2 = 27, h_3 = 23$ which is contained in $(7, 11, 15, 19, 23, 27, 31)$.*

4. AUXILIARY RESULTS

Let $H \in KW(p, q)$ be a numerical semigroup of embedding dimension $n \geq 3$. Thus, H is minimally generated by the set $\{p, q, h_1, \dots, h_{n-2}\}$ for some relatively prime positive integers p and q , and $h_i = pq - px_i - qy_i$, $1 \leq i \leq n-2$. Here, $q > x_{n-2} > \dots > x_1$ and $p > y_1 > \dots > y_{n-2}$ are positive integers with the same parity with p and q respectively. Let k be a field. Set $S = k[u, v, u_1, \dots, u_{n-2}]$ to be a polynomial ring and $\phi : S \rightarrow k[t]$ to be the map defined by $u \mapsto t^p, v \mapsto t^q, u_i \mapsto t^{h_i}, 1 \leq i \leq n-2$. Then the semigroup ring $k[H] = k[t^p, t^q, t^{h_1}, \dots, t^{h_{n-2}}]$ is the image of ϕ and it is isomorphic to S/I_H where $I_H = \ker \phi$, called the ideal of H , is the prime ideal defining the affine monomial curve whose coordinate ring is $k[H]$.

4.1. **A Classification of I_H .** Using results from the appendix in [5], we can express the defining ideal of the semigroup ring $k[H]$ where $H \in KW(p, q)$ for any embedding dimension $n \geq 3$ as the sum of determinantal ideals.

Theorem 4.1. *Let $H = \langle p, q, h_1, \dots, h_{n-2} \rangle \in KW(p, q)$, so that $h_i = pq - x_i p - y_i q$, $1 \leq i \leq n - 2$ with positive integers $x_1 < x_2 < \dots < x_{n-2}$, $y_1 > y_2 > \dots > y_{n-2}$, each satisfying (1): $2x_i \leq q, 2y_i \leq p$. Then the defining ideal I_H of the semigroup ring $k[H]$ is the ideal of the 2×2 minors of the following 2×3 matrices:*

$$A_{ij} = \begin{bmatrix} u_i & u^{q-2x_j} v^{p-y_i-y_j} & u_j \\ u^{x_j-x_i} & u_j & v^{y_i-y_j} \end{bmatrix}, \quad 1 \leq i < j \leq n-2$$

$$B = \begin{bmatrix} v^{y_{n-2}} & u^{q-x_1-x_{n-2}} & u_1 \\ u^{x_1} & u_{n-2} & v^{p-y_1-y_{n-2}} \end{bmatrix}$$

$$C = \begin{bmatrix} u^{q-x_1-x_{n-2}} v^{p-2y_1} & u_1 & u_{n-2} \\ u_1 & u^{x_{n-2}-x_1} & v^{y_1-y_{n-2}} \end{bmatrix}$$

Proof. According to the theorem in Appendix A. of [5], the defining ideal of H consists of the following binomials:

$$f_{ij} = u_i u_j - u^{q-x_i-x_j} v^{p-y_i-y_j}, \quad 1 \leq i < j \leq n-2$$

$$g_i = v^{y_i-y_{i+1}} u_i - u^{x_{i+1}-x_i} u_{i+1}, \quad 1 \leq i \leq n-3$$

$$h_1 = v^{p-y_1} - u^{x_1} u_1$$

$$h_2 = v^{y_{n-2}} u_{n-2} - u^{q-x_{n-2}}$$

Let M^l denote column l of a matrix M . Observe that

$$f_{ij} = \det \begin{bmatrix} A_{ij}^1 & A_{ij}^2 \end{bmatrix}, \quad 1 \leq i < j \leq n-2,$$

$$f_{jj} = \det \begin{bmatrix} A_{ij}^2 & A_{ij}^3 \end{bmatrix}, \quad 2 \leq j \leq n-2,$$

$$f_{11} = \det \begin{bmatrix} C^1 & C^2 \end{bmatrix},$$

$$g_i = \det \begin{bmatrix} A_{ij}^1 & A_{ij}^3 \end{bmatrix}, \quad 1 \leq i \leq n-3, \quad j = i+1,$$

$$h_1 = \det \begin{bmatrix} B^1 & B^3 \end{bmatrix},$$

$$h_2 = \det \begin{bmatrix} B^1 & B^2 \end{bmatrix}$$

□

4.2. Minimal Free Resolutions. In this section, we explicitly describe minimal free resolutions of the semigroup ring $k[H]$ where H is a numerical semigroup of small embedding dimension 3 or 4 in the class $KW(p, q)$, and give their Betti numbers. We express the maps in the resolution in terms of the x_i, y_i where $px_i + qy_i$ are the complements of the gaps h_i of $\langle p, q \rangle$, i.e., $h_i = pq - (px_i + qy_i)$ for $1 \leq i \leq n - 2$.

4.2.1. Embedding Dimension of H is 3. In this instance, we have $n = 3$, so $H = \langle p, q, h_1 \rangle \in KW(p, q)$, $h_1 = pq - px_1 - qy_1$. Here, $x_1, y_1 \in \mathbb{N}$ with $(x_1, y_1) \leq (q/2, p/2)$. Furthermore, $k[H] = k[t^p, t^q, t^{h_1}]$ is the semigroup ring associated to H , and $\phi : S \rightarrow R$ is the map defined by $u \mapsto t^p, v \mapsto t^q, u_1 \mapsto t^{h_1}$. Following is the resolution of $k[H]$:

$$0 \rightarrow S^2 \xrightarrow{\begin{pmatrix} u_1 & -u^{q-2x_1} \\ -u^{x_1} & v^{y_1} \\ v^{p-2y_1} & -u_1 \end{pmatrix}} S^3 \xrightarrow{\begin{pmatrix} v^{p-y_1} - u^{x_1} & u^{q-2x_1}v^{p-2y_1} - u_1^2 & u^{q-x_1} - v^{y_1}u_1 \end{pmatrix}} S^1$$

4.2.2. Embedding Dimension of H is 4. In this setting, $H = \langle p, q, h_1, h_2 \rangle \in KW(p, q)$, $h_i = pq - px_i - qy_i, i = 1, 2$. Here, $x_1 < x_2 \leq q/2, y_2 < y_1 \leq p/2$ are in \mathbb{N} . Furthermore, $k[H] = k[t^p, t^q, t^{h_1}, t^{h_2}]$ is the semigroup ring associated to H . Following is the resolution of R/I , where I is the relation ideal of $k[H]$:

$$0 \rightarrow S^3 \xrightarrow{A_1} S^8 \xrightarrow{A_2} S^6 \xrightarrow{A_3} S^1$$

The maps are given by matrices below.

$$A_1 = \begin{bmatrix} u_2 & -u^{q-x_2}v^{y_1-y_2} & 0 \\ -u_1 & u^{q-x_1-x_2} & 0 \\ u^{x_1}v^{p-2y_1} & -x_1 & 0 \\ 0 & v^{y_2} & u^{x_2-x_1} \\ -v^{p-y_1-y_2} & u_2 & 0 \\ -u^{2x_1-x_2} & 0 & -v^{y_1-y_2} \\ u^{x_2-x_2}u_1 & 0 & u_2 \\ 0 & v^{2y_2-y_1}u_2 & -u_1 \end{bmatrix},$$

$$A_2 =$$

$$\begin{bmatrix} -u^{x_1} & -u^{2x_1-x_2}v^{p-y_1-y_2} & 0 & 0 & 0 & -u_2 & -v^{p-y_1-y_2} & 0 \\ 0 & 0 & 0 & -u_2 & 0 & u_1 & u^{x_2-x_1} & -v^{\frac{y_1-y_2}{2}} \\ 0 & 0 & v^{y_2} & u_1 & u^{x_2-x_1}v^{2y_2-y_1} & 0 & 0 & u^{x_2-x_1} \\ -v^{p+y_2-2y_1} & -u^{2x_1-x_2}u_1 & u^{q+x_1-2x_2} & -u^{q-2x_2}v^{p-y_1-y_2} & -v^{2y_2-y_1}u_2 & -u^{q-x_1-x_2}v^{p-2y_1} & -u_1 & -u_2 \\ -u^{x_2-x_1}v^{p-2y_1} & -v^{p-y_1-y_2} & u_2 & 0 & u_1 & 0 & 0 & 0 \\ -u_1 & -u_2 & u^{q-2x_2}v^{y_1-y_2} & 0 & u^{q-x_1-x_2} & 0 & 0 & 0 \end{bmatrix}$$

$A_3 =$

$$\left[u^{q-2x_1}v^{p-2y_1} - u_1^2, u^{q-x_1-x_2}v^{p-y_1-y_2} - u_1u_2, u^{q-2x_2}v^{p-2y_2} - u_2^2, v^{y_1-y_2}u_1 - u^{x_2-x_1}u_2, v^{y_2}u_2 - u^{q-x_2}, u^{x_1}u_1 - v^{p-y_1} \right]$$

5. GENERALIZING KW TYPE MONOIDS TO DIMENSION THREE

In this section, we will expand this class $KW(p, q)$. Let p and q be two coprime numbers with $1 < p < q$. Let $w = r_1p + r_2q$ for some $r_1, r_2 \in \mathbb{N}$. Then the semigroup $S = \langle sp, sq, r_1p + r_2q \rangle$ with $\gcd(s, w) = 1$, is again symmetric and with Frobenius number $s(pq - p - q) + w(s - 1)$. Consider the class $R(p, q, r_1, r_2, s)$ of numerical semigroups obtained by adding to S gaps corresponding to points $(a, b, c) \in \mathbb{N}^3$ falling on a lattice path under the plane $G_0 : spx + sqy + wz = s(pq + w) - sp - sq - w$.

From here on, S is the numerical semigroup $\langle sp, sq, w \rangle$ where $p < q$ are relatively prime and s, w are relatively prime and $w = r_1p + r_2q, r_i \geq 1$.

Proposition 5.1. *Every gap of S can be written as $s(pq+w) - sp(a+1) - sp(b+1) - w(c+1)$ for some $(a, b, c) \in \mathbb{N}^3$. We may always take $c \leq s - 1$. In particular, each gap of S is of the form $s(pq + w) - sp(a + 1) - sp(b + 1) - w(c + 1)$ with $(a, b, c) < (q, p, s)$ alphabetically, and corresponds to a unique lattice point (a, b, c) below the hyperplane*

$$G_0 : spx + sqy + wz = s(pq + w) - sp - sq - w.$$


Proof. Let t be a gap of S . Since S is symmetric, $F(S) - t \in S$ so that $F(S) - t = spx + sqy + wz$ for some $(x, y, z) \in \mathbb{N}^3$. In other words,

$$(7) \quad t = s(pq + w) - sp(x + 1) - sq(y + 1) - w(z + 1)$$

Now assume $x < q, y < p, z < s$. Suppose

$$\begin{aligned} t &= s(pq + w) - sp(x + 1) - sq(y + 1) - w(z + 1) \\ &= s(pq + w) - sp(x' + 1) - sq(y' + 1) - w(z' + 1) \end{aligned}$$

FIGURE 1. Gaps of $\langle 15, 21, 17 \rangle$ are in 1-1 correspondence with lattice points under the plane $15x + 21y + 17z = 103$.



for some $(x, y, z), (x', y', z') \in \mathbb{N}^3$. Then

$$s(px + qy - px' - qy') = w(z' - z)$$

Since $(s, w) = 1$, $s \mid z' - z$, i.e., $z' = z + ms$ for some $m \in \mathbb{Z}$. But $0 \leq z, z' \leq s - 1$ so $m = 0$.

Therefore,

$$p(x - x') = q(y' - y).$$


Since p and q are relatively prime, we get $x = x' + mq$ for some $m \in \mathbb{Z}$. But $0 \leq x, x' \leq q - 1$ so $m = 0$.

Thus, we must have $x = x', y = y'$, and $z = z'$.

□

Definition 5.2. We denote by $KW(p, q, r_1, r_2, s)$ the numerical semigroups H in $R(p, q, r_1, r_2, s)$ generated by $\{sp, sq, w = r_1p + r_2q, h_1, \dots, h_{n-3}\}$ where $h_i = s(pq + w) - (x_i + 1)sp - (y_i + 1)sq - (z_i + 1)w$, where $x_i \leq \frac{q-4}{2}, y_i \leq \frac{p-4}{2}$ and $z_i \leq s - 2$.

Set $\Gamma(x, y, z) = F(S) - xsp - ysq - zw$ for $(x, y, z) \in \mathbb{N}^3$. The Apéry set of a numerical semigroup \mathbf{a} with respect to an element $a \in \mathbf{a}$, named in honour of R. Apéry, is

FIGURE 2. H obtained by adding 55, 51, 63 to S


$Ap(\mathbf{a}, a) = \{b \in \mathbf{a} \mid b - a \notin \mathbf{a}\}$. Let $H = \langle sp, sq, w, h_2, \dots, h_{n-2} \rangle \in KW(p, q, r_1, r_2, s)$. We give a characterization of the elements h_2, \dots, h_{n-2} in terms of the Apéry sets of H . To understand the motivation, we go through the following geometric illustration.




Example 5.3. Let $S = \langle sp = 36, sq = 44, w = 29 \rangle$. Obtain H by adding the gaps corresponding to a lattice point within one of the cuboids in Figure 2. Here, $H = \langle 36, 44, 29, 206, 221, 222 \rangle \in KW(9, 11, 2, 1, 4)$, where $h_2 = 221, h_3 = 222, h_4 = 206$ correspond to the lattice points $A = (1, 2, 2), B = (3, 1, 1)$ and $E = (1, 3, 1)$, respectively.

First, we compute $Ap(H, sp = 36)$. Fix $z = c$, with $c \leq z_1$, the maximum height of any cuboid in the image. This plane passes through at least one of the three cuboids, say C_i which is obtained from h_i . For each $b \leq b_i$, there is a line $y = b, z = c$ on the plane $z = c$ such that each point $(a_j, b, c), a_j \leq a_2$ is in H . Moreover, whenever $i, j \leq 2$, we have

$$\Gamma(a_i, b, c) \equiv \Gamma(a_j, b, c) \pmod{sp}$$

Out of all these points on the line $z = c, y = b$, the one with the largest x -coordinate, say P_l , corresponds to the smallest gap congruent to $sw - sqb - wc$ modulo sp . The equivalence classes are distinct for each distinct b and c , so $\Gamma(P_l) \in Ap(H, sp)$. Visually, $Ap(H, sp) \setminus S$ is contained in the set of gaps corresponding to the lattice points seen in Figure 3.

Similarly, we can find elements in $Ap(H, sq = 44)$ and $Ap(H, w = 29)$. $Ap(H, sq) \setminus S$ and $Ap(H, w) \setminus S$ can be visualized in Figures 4 and 5 respectively. One can see that the only lattice points common to all three figures 3, 4 and 5 are exactly those corresponding to h_2, h_3 and h_4 .

FIGURE 3. $Ap(H, sp) \setminus S$ FIGURE 4. $Ap(H, sq) \setminus S$ FIGURE 5. $Ap(H, w) \setminus S$

This gives rise to the following result:

Proposition 5.4. *Let $H \in KW(p, q, r_1, r_2, s)$. Then $\{\text{minimal generators of } H\} \setminus \{sp, sq, w\} = Ap(H, sp) \cap Ap(H, sq) \cap Ap(H, w)$.*

Proof. Denote by \mathbf{h} the minimal generating set of H . Let $h_i \in \mathbf{h}$. If $h_i - sp \in H$, then $h_i = \sum_j r_j h_j + asp + bsq + cw$. If $r_i = 0$, h_i is not a minimal generator. If $r_i \geq 1$, then we have $\sum_{j \neq i} r_j h_j + (r_i - 1)h_i + (a+1)sp + bsq + cw = 0$ which is impossible. Hence $h_i - sp \notin H$ and hence $h_i \in Ap(H, sp)$. Similarly, $h_i \in Ap(H, sq)$ and $h_i \in Ap(H, w)$. Thus, $\mathbf{h} \setminus \{p, sq, w\} \subset Ap(H, sp) \cap Ap(H, sq) \cap Ap(H, w)$ since $x \notin Ap(H, x)$ for any $x \in H$.

If $h \in Ap(H, sp) \cap Ap(H, sq) \cap Ap(H, w)$ then $h = \sum_{i=1}^{n-2} t_i h_i$ for $h - sp, h - sq, h - w$ cannot be in H . Further since $2h_i \in \langle sp, sq, w \rangle$ for all i , $t_i \leq 1$. If two of the t_i are not zero,

say, $i = 1, 2$, then we have $h = 2h_1 - sp(x_2 - x_1) - sq(y_2 - y_1) - w(z_2 - z_1) + \sum_{i=3}^{n-2} h_i$. By the fact that $h_i \in H$ are minimal generators, we have that one of $(x_2 - x_1), (y_2 - y_1), (z_2 - z_1)$ must be negative. So, one of $h - sp, h - sq, h - w$ must be in H which is impossible. so, only one of $t_i \neq 0$. Hence, $h = h_i$ for some i and so we get the other inclusion: $\mathbf{h} \setminus \{sp, sq, w\} = \text{Ap}(H, sp) \cap \text{Ap}(H, sq) \cap \text{Ap}(H, w)$. \square

The proposition above establishes a close relationship between the lattice path of a numerical semigroup $H \in KW(p, q, r_1, r_2, s)$ and its Apéry set. In fact, the lattice path of H can be exclusively determined by its Apéry sets $\text{Ap}(H, sp)$, $\text{Ap}(H, sq)$, and $\text{Ap}(H, w)$. Consequently, this observation could facilitate deriving an upper bound for the embedding dimension of any $H \in KW(p, q, r_1, r_2, s)$, as in Remark 2.

Kunz and Waldi show that for $H \in KW(p, q)$, the type of the semigroup ring $k[H]$ equals $e(H) - 1$. Contrasting this with the dimension three, consider $H \in R(5, 7, 2, 1, 3)$ obtained by adding a gap h of $\langle 15, 21, 17 \rangle$. In the following table. the first column consists of gaps h such that $H = \langle 15, 21, 17, h \rangle$, with embedding dimension 4 has the corresponding type in column 2.

Choice of h	$t(H)$
50,56,65,67, 71,73,82,86,88,103	3
35, 52	4


However, we show that for a class of the $h \in H$ in $KW(p, q, r_1, r_2, s)$, we prove that the type is always 3.

Example 5.5. Consider Figure 6 below. Take $H = \langle 36, 44, 29, 221 \rangle \in KW(9, 11, 2, 1, 4)$ where 221 corresponds to the lattice point $A = (1, 2, 2)$. Then $PF(H) = \{271, 316, 331\}$ where 331, 271, and 316 correspond to the points $I = (2, 0, 0), J = (0, 3, 0)$, and $K = (0, 0, 3)$, respectively.

Theorem 5.6. Let $H = \langle sp, sq, w, h \rangle \in KW(p, q, r_1, r_2, s)$ with $h = F(S) - spx - sqy - wz$. If $p_z := F(S) - (z + 1)w \notin H$, then $t(H) = 3$.

Proof. We will show that

$$PF(H) = \{p_x := F(S) - (x + 1)sp, p_y := F(S) - (y + 1)sq, p_z\}.$$

FIGURE 6. $PF(\langle 36, 44, 29, 221 \rangle)$

Claim 1. $p_x, p_y \notin H$. Suppose first that $p_x \in H$. Then $p_x = \alpha sp + \beta sq + \gamma w + \delta h$ for some $\alpha, \beta, \gamma, \delta \in \mathbb{N}$. By the choice of h to be in Kunz-Waldi class, $2h \in S$. pX is a gap of S . So, $\delta = 1$. So, we get $(\alpha + 1)sp + (\beta - \gamma)sq + (\gamma - z)w = 0$. This means either $\beta < \gamma$ or $\gamma < z$. If indeed $\gamma < z$, then $(z - \gamma)w = (\alpha + 1)sp + (\beta - \gamma)sq$. So, s must divide $z - \gamma$ but this is impossible since $z < s$. Now if $\gamma \geq z$, $0 \leq (z - \gamma)w = ((y - \beta)q - (\alpha + 1)p)s$. So, there exists $u \in \mathbb{N}$ such that $(y - \beta)q - (\alpha + 1)p = wu = r_1pu + r_2qu \implies (y - \beta - r_2u)q = (\alpha + 1 + r_1u)p > 0 \implies y > p$ a contradiction. So, $p_x \notin H$. The argument for $p_y \notin H$ is similar.

Claim 2. $p_i + sp, p_i + sq, p_i + w, p_i + h \in H$ for all $1 \leq i \leq 3$.

To see that $p_1 + sp \in H$, we have the following

$$\begin{aligned} p_1 + sp &= F(S) - (x + 1)sp + sp \\ &= F(S) - xsp \\ &= h + ysq + zw \in H \end{aligned}$$

Next, to see that $p_1 + sq \in H$, we have

$$\begin{aligned} p_1 + sq &= F(S) - (x + 1)sp + sq \\ &= spq + sw - sp - sq - w - (x + 1)sp + sq \\ &= sp(q - x - 2) + w(s - 1) \in H \end{aligned}$$

since $x \leq \lfloor q/2 \rfloor - 2 \implies q - x - 2 \geq 0$. Thirdly, for $p_1 + w \in H$, we see

$$\begin{aligned} p_1 + w &= F(S) - (x + 1)sp + w \\ &= spq + sw - sp - sq - w - (x + 1)sp + w \\ &= sp(q - x - 2) + s(r_1p + r_2q) - sq \\ &= sp(q - x - 2 + r_1) + sq(r_2 - 1) \in H \end{aligned}$$

since $r_2 \geq 1$.

Similarly, $p_2 + sp = sq(p - y - 2) + w(s - 1) \in H$ since $y \leq \lfloor p/2 \rfloor - 2 \implies p - y - 2 \geq 0$, $p_2 + sq = h + xsp + zw \in H$, and $p_2 + w = sq(p - y - 2 + r_2) + sp(r_1 - 1) \in H$ as $r_1 \geq 1$.

Next, $p_3 + sp = w(s - z - 2) + sq(p - 1) \in H$ as $z \leq s - 2$, $p_3 + sq = w(s - z - 2) + sp(q - 1) \in H$, and $p_3 + w = h + xsp + ysq \in H$.

It remains to show that $p_i + h \in H$ for all $1 \leq i \leq 3$. Well, recall that $x \leq \frac{q-4}{2}$, $y \leq \frac{p-4}{2}$, and $z \leq s - 2$ so that

$$p_1 + h = sp(q - 2x - 3) + sq(p - y - 2) + w(2s - z - 2) \in H,$$

$$p_2 + h = sp(q - x - 2) + sq(p - 2y - 3) + w(2s - z - 2) \in H,$$

and

$$p_3 + h = sp(q - x - 2) + sq(p - y - 2) + w(2s - 2z - 2) \in H.$$

Claim 3. No other gap of H is in $PF(H)$. Any gap of H is of the form $g = F(S) - spa - sqb - wc$ for some $(a, b, c) \in \mathbb{N}^3$. Suppose g is different from p_1, p_2, p_3 . It is not of the form $F(S) - (x + 1 + i)sp$ for $i > 0$ because $F(S) - (x + 1 + i)sp + sp = p_1 - (i - 1)sp \in H \implies p_1 \in H$, a contradiction to Claim 1. Similarly, g is not of the form $F(S) - (y + 1 + i)sq$ or $F(S) - (z + 1 + i)w$ for $i \geq 0$. Further, one of the following must happen: $a > x, b > y, c > w$. Otherwise, if $a \leq x, b \leq y, c \leq z$, then $g = h + (x - a)sp + (y - b)sq + (z - c)w \in H$.

Suppose $g = F(S) - spa - sqb - wc \in PF(H)$ with $a = x + j$ for some $j \in \mathbb{N} \setminus \{0\}$, and one of b, c is non-zero. Then $p_1 = g + sp(j - 1) + sqb + wc \in H$. If $b = y + j$ for some $j \geq 1$, then $p_2 = g + spa + sq(j - 1) + wc \in H$. Finally, if $c = z + j$ for some $j \geq 1$, then $p_3 = g + spa + sqb + w(j - 1) \in H$. Each of these cases is a contradiction since $p_1, p_2, p_3 \notin H$. So g cannot be different from p_1, p_2 or p_3 .

Thus, $PF(H) = \{p_1, p_2, p_3\}$ proving that the type of H is 3.

□

The condition $p_3 \notin H$ is necessary as seen in the example below.

Example 5.7. Take $H = \langle 27, 33, 29, 152 \rangle \in KW(9, 11, 2, 1, 3)$. Here, $152 = s(pq + w) - (x + 1)sp - (y + 1)sqy - (z + 1)w$ for $(x, y, z) = (3, 1, 1)$ but $t(H) = 4$. In this case, $p_3 = 237 \in H$ and $PF(H) = \{p_1, p_2, 183, 204\}$.

Further, the theorem still may not hold if $H \in R(p, q, r_1, r_2, s) \setminus KW(p, q, r_1, r_2, s)$.

Example 5.8. Take $H = \langle 15, 21, 17, 37 \rangle \in R(5, 7, 2, 1, 3)$. Here, $37 = s(pq + w) - (x + 1)sp - (y + 1)sqy - (z + 1)w$ for $(x, y, z) = (3, 1, 0)$, so that $3 > \frac{7-4}{2} = 1.5$, and $t(H) = 4$.

With this in mind, we ask the following questions:

Question 1. What are the conditions on the gaps h_i that will yield an explicit formula for the type of H ?

Question 2. Let $h_i = spx + sqy + wz = s(pq + w) - spx_i - sqy_i - wz_i$, where $x_i < \frac{q}{2}, y_i < \frac{p}{2}, z < s$. In particular, what is the type of $H = (sp, sq, w, h_i, 1 \leq i \leq n - 2) \in KW(p, q, s, w)$?

Theorem 1.1 gives the number of generators for the relation ideal of any $H \in KW(p, q)$. Consider the following tables, In the left one, the first column consists of gaps h such that $H = \langle 15, 21, 17, h \rangle$, with embedding dimension 4, and the second column has the corresponding number of minimal generators for its relation ideal. In the next table, the first column consists of h' such that $H' = \langle 15, 21, 17, 73, h' \rangle$, and the second column is $\mu(I_{H'})$, the number of minimal generators of $I_{H'}$.

Choice of h	$\mu(I_H)$	Choice of h'	$\mu(I_{H'})$
35, 50, 52, 56, 65, 67, 71, 73, 82, 86, 88	6	67	9
103	7	50, 71, 82, 86	10
		65	11

This prompts the question:

Question 3. What is the number of minimal generators for the relation ideal of $H \in KW(p, q, r_1, r_2, s)$?

Even when the embedding dimension of H is 4, we may have many different numbers of generators for its ideal as seen in the table above.

REFERENCES

- [1] Bresinsky, H., Monomial Gorenstein ideals, *Manuscripta Math* **429** (1979). 159–181. DOI: <https://doi.org/10.1007/BF01303625>.

- [2] P. Dey, H. Srinivasan, Principal matrices of numerical semigroups, *Women in commutative algebra Assoc. Women Math. Ser.* **29** (2021), 233–256. DOI: https://doi.org/10.1007/978-3-030-91986-3_9.
- [3] P. Gimenez, H. Srinivasan, A note on Gorenstein monomial curves, *Bulletin of the Brazilian Mathematical Society, New Series Springer Science and Business Media LLC* **45** (2014), 671–678. DOI: <http://dx.doi.org/10.1007/s00574-014-0068-4>.
- [4] E. Kunz, R. Waldi, Geometrical illustration of numerical semigroups and of some of their invariants, *Semigroup Forum* **89** (2014), 664–691. DOI: <http://dx.doi.org/10.1007/s00233-014-9599-7>.
- [5] E. Kunz, R. Waldi, On the deviation and the type of certain local Cohen–Macaulay rings and numerical semigroups, *Journal of Algebra* **478** (2017), 397–409. DOI: <https://doi.org/10.1016/j.jalgebra.2017.01.041>.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MISSOURI, COLUMBIA MO 65211, USA.