Optimal Lagrange Interpolation Projectors and Legendre Polynomials

Mikhail Nevskii

Department of Mathematics, P. G. Demidov Yaroslavl State University, Sovetskaya str., 14, Yaroslavl, 150003, Russia

May 2, 2024

Abstract

Let *K* be a convex body in \mathbb{R}^n , and let $\Pi_1(\mathbb{R}^n)$ be the space of polynomials in *n* variables of degree at most 1. Given an (n + 1)-element set $Y \subset K$ in general position, we let P_Y denote the Lagrange interpolation projector $P_Y : C(K) \to \Pi_1(\mathbb{R}^n)$ with nodes in *Y*. In this paper, we study upper and lower bounds for the norm of the optimal Lagrange interpolation projector, i.e., the projector with minimal operator norm where the minimum is taken over all (n + 1)-element sets of interpolation nodes in *K*. We denote this minimal norm by $\theta_n(K)$.

Our main result, Theorem 5.2, provides an explicit lower bound for the constant $\theta_n(K)$ for an arbitrary convex body $K \subset \mathbb{R}^n$ and an arbitrary $n \ge 1$. We prove that $\theta_n(K) \ge \chi_n^{-1}(\operatorname{vol}(K)/\operatorname{simp}_n(K))$ where χ_n is the Legendre polynomial of degree n and $\operatorname{simp}_n(K)$ is the maximum volume of a simplex contained in K. The proof of this result relies on a geometric characterization of the Legendre polynomials in terms of the volumes of certain convex polyhedra. More specifically, we show that for every $\gamma \ge 1$ the volume of the set $\{x = (x_1, ..., x_n) \in \mathbb{R}^n : \sum |x_j| + |1 - \sum x_j| \le \gamma\}$ is equal to $\chi_n(\gamma)/n!$.

Furthermore, if *K* is an *n*-dimensional ball, this approach leads us to the equivalence $\theta_n(K) \approx \sqrt{n}$ which is complemented by the exact formula for $\theta_n(K)$. If *K* is an *n*-dimensional cube, we obtain explicit efficient formulae for upper and lower bounds of the constant $\theta_n(K)$; moreover, for small *n*, these estimates enable us to compute the exact values of this constant.

MSC: 41A05, 52B55, 52C07

Keywords: polynomial interpolation, projector, norm, estimate, simplex, homothety, absorption index.

1. Introduction

Let *K* be a convex body in \mathbb{R}^n . Given a function $f \in C(K)$, we let $E_1(f; K)_{C(K)}$ denote the best approximation of *f* (in the *C*(*K*)-norm) by polynomials of degree at most 1 defined on \mathbb{R}^n . (We denote the space of these polynomials by $\Pi_1(\mathbb{R}^n)$.)

Let *P* be a polynomial projector on *K*, i.e., a linear operator from C(K) into $\Pi_1(\mathbb{R}^n)$ such that $P(Pf|_K) = Pf$ for every function $f \in C(K)$. Then the following well known inequality

$$E_1(f;K)_{C(K)} \le \|f - Pf\|_{C(K)} \le (1 + \|P\|_K)E_1(f;K)_{C(K)}.$$
(1.1)

E-mail address: mnevsk55@yandex.ru

holds. Here $||P||_{K} = \sup\{||Pf|_{K}||_{C(K)} : ||f||_{C(K)} \le 1\}$ is the *C*(*K*) operator norm of *P*.

Clearly, the first inequality in (1.1) is trivial because $Pf \in \Pi_1(\mathbb{R}^n)$. The second inequality is known in the literature as Lebesgue's Lemma. See, e.g., [6, p.30]. This lemma shows that, if the norm $||P||_K$ is not large, there is no significant loss of accuracy if we replace the best C(K)-approximant of fby the value Pf which linearly depends on the function f.

This approach provides a (fairly simple) tool for "linearizing" the best approximation element in various problems of Analysis involving the approximation of continuous functions defined on subsets of \mathbb{R}^n . It also motivates us to study the following problem.

Main Problem 1.1 Let *K* be a convex body in \mathbb{R}^n , and let *Y* be an (n + 1)-element subset of *K* in "general position" (i.e., *Y* is the family of vertices of a nondegenerate simplex in \mathbb{R}^n). Let P_Y be the first order Lagrange interpolation projector with nodes in *Y* (i.e., $P_Y(f) \in \Pi_1(\mathbb{R}^n)$ and $P_Y(f) = f$ on *Y* for every $f \in C(K)$).

(i) Find an (n + 1)-element set $Y_{opt} \subset K$ such that $P_{Y_{opt}}$ has the smallest operator norm among all Lagrange interpolation projectors P_Y where Y is an arbitrary (n + 1)-element subset of K.

We refer to the Lagrange interpolation projector $P_{Y_{opt}}$ as the optimal interpolation projector, and we call the set Y_{opt} as an optimal set of interpolation nodes.

(ii) Calculate or give efficient estimates for the upper and lower bounds of the constant

$$\theta_n(K) = \|P_{Y_{opt}}\|_K,$$

i.e., the minimum of the operator norm $||P_Y||_K$ where the minimum is taken over all (n + 1)-element sets *Y* of interpolation nodes in *K*.

Let us note that both upper and lower bounds of the operator norms of interpolation projectors are important in applications. Usually, upper estimates of minimal norms of projectors are obtained by considering projectors of some special type. The technique of obtaining lower estimates of minimal norms is fundamentally different – in this case it is necessary to prove the lower estimate for an arbitrary projector.

The construction and evaluation of interpolation projectors is a classical topic in Approximation Theory. These problems have been treated in many papers and monographs, see, e.g., [1, 3, 5, 9, 10, 15, 38, 40].

In this paper, we show that there exists a Lagrange interpolation projector whose operator norm does not exceed n + 1 (Theorem 3.1). On the other hand, we prove that for any Lagrange interpolation projector $P : C(K) \to \Pi_1(\mathbb{R}^n)$ the following inequality

$$||P|| \ge \chi_n^{-1} \left(\frac{\operatorname{vol}(K)}{\operatorname{simp}_n(K)} \right)$$
(1.2)

holds. Here χ_n is the standardized Legendre polynomial of degree *n* and simp_n(*K*) is the maximum volume of a simplex contained in *K*. See Theorem 5.2.

The main point of the proof of inequality (1.2) is the following geometrical characterization of Legendre polynomials given in Theorem 4.1. This theorem states that for $\gamma \ge 1$ the volume of the set

$$\left\{x \in \mathbb{R}^{n} : \sum_{j=1}^{n} |x_{j}| + \left|1 - \sum_{j=1}^{n} x_{j}\right| \le \gamma\right\}$$

is equal to $\chi_n(\gamma)/n!$.

In the cases when K coincides with an n-dimensional cube or ball, the above inequality yields the estimate $||P|| \ge c \sqrt{n}$.

Let us describe the content of the paper in more detail. In Section 2 we give basic notation, definitions, and preliminary information. Section 3 contains upper estimates of the minimum absorption coefficient of a convex body K by a simplex and also of the minimum projector norm for linear interpolation on K. In Section 4 we prove Theorem 4.1 mentioned above. In Section 5 we prove inequality (1.2). In Section 6 we present several explicit lower bounds for the constant $\theta_n(K)$ provided K is an n-dimensional ball or an n-dimensional cube. Section 7 contains inequalities of the considered type for interpolation by linear functions on an arbitrary compact set in \mathbb{R}^n . Finally, Section 8 contains concluding remarks, a review of some results on this topic, and a description of open questions.

2. Notation and preliminaries

Let us fix some notation. Throughout the paper $n \in \mathbb{N}$ is a positive integer. Given $x = (x_1, ..., x_n) \in \mathbb{R}^n$, by ||x|| we denote its standard Euclidean norm

$$||x|| = \sqrt{\langle x, x \rangle} = \left(\sum_{i=1}^{n} x_i^2\right)^{1/2}$$

Hereafter, for $x = (x_1, ..., x_n), y = (y_1, ..., y_n) \in \mathbb{R}^n$ by $\langle x, y \rangle$ we denote the standard inner product in \mathbb{R}^n : $\langle x, y \rangle = x_1y_1 + ... + x_ny_n$

Given $x^{(0)} \in \mathbb{R}^n$ and R > 0, we let $B(x^{(0)}; R)$ denote the Euclidean ball with center $x^{(0)}$ and radius R:

$$B(x^{(0)}; R) = \{x \in \mathbb{R}^n : ||x - x^{(0)}|| \le R\}.$$

We also set

$$B_n = B(0; 1)$$
 and $Q_n = [0, 1]^n$.

The notation $L(n) \approx M(n)$ means that there exist absolute constants $c_1, c_2 > 0$ such that $c_1M(n) \leq L(n) \leq c_2M(n)$.

Let *K* be a convex body in \mathbb{R}^n , i.e., a compact convex subset of \mathbb{R}^n with nonempty interior. The symbol σK denotes a homothetic copy of *K* with homothety center at the center of gravity of *K* and homothety coefficient σ .

We let vol(K) denote the volume of K. If K is a convex polytope, then by ver(K) we denote the set of all vertices of K. By *a translate* we mean the result of a parallel shift.

We say that an *n*-dimensional simplex S is *circumscribed around a convex body* K if $K \subset S$ and each (n - 1)-dimensional face of S contains a point of K. A convex polytope is *inscribed into* K if every vertex of this polytope belongs to the boundary of K.

Definition 2.1 Let $i \in \{1, ..., n\}$ and let $d_i(K)$ be the maximal length of a segment contained in *K* and parallel to the x_i -axis. We refer to $d_i(K)$ as the *i*th axial diameter of *K*.

The notion of axial diameter of a convex body was introduced by Scott [41,42].

Definition 2.2 Given convex bodies K_1 , K_2 , by $\xi(K_1; K_2)$ we denote the minimal $\sigma \ge 1$ with the property $K_1 \subset \sigma K_2$. We call $\xi(K_1, K_2)$ the absorption index of K_1 by K_2 .

By $\alpha(K_1, K_2)$ we denote the minimal $\sigma > 0$ such that K_1 is a subset of a translate of σK_2 .

Note that the equality $\xi(K_1; K_2) = 1$ is equivalent to the inclusion $K_1 \subset K_2$. Clearly, $\alpha(K_1, K_2) \leq \xi(K_1, K_2)$.

Definition 2.3 By $\xi_n(K)$ we denote the minimal absorption index of a convex body *K* by an inner nondegenerate simplex. In other words,

 $\xi_n(K) = \min\{\xi(K; S) : S \text{ is an } n \text{-dimensional simplex}, S \subset K, \operatorname{vol}(S) \neq 0\}.$

By C(K) we denote the space of all continuous functions $f: K \to \mathbb{R}$ with the uniform norm

$$||f||_{C(K)} = \max_{x \in K} |f(x)|.$$

We let $\Pi_1(\mathbb{R}^n)$ denote the space of polynomials in *n* variables of degree at most 1.

Let *S* be a nondegenerate simplex in \mathbb{R}^n with vertices $x^{(j)} = (x_1^{(j)}, \dots, x_n^{(j)}), 1 \le j \le n+1$. We define *the vertex matrix* of this simplex by

$$\mathbf{A} = \begin{pmatrix} x_1^{(1)} & \dots & x_n^{(1)} & 1 \\ x_1^{(2)} & \dots & x_n^{(2)} & 1 \\ \vdots & \vdots & \vdots & \vdots \\ x_1^{(n+1)} & \dots & x_n^{(n+1)} & 1 \end{pmatrix}.$$

Clearly, matrix A is nondegenerate and

$$\operatorname{vol}(S) = \frac{|\operatorname{det}(\mathbf{A})|}{n!}.$$
(2.1)

Let

$$\mathbf{A}^{-1} = \left(l_{ij}\right)_{i,j=1}^{n+1}.$$
(2.2)

Definition 2.4 Linear polynomials

$$\lambda_j(x) = l_{1j}x_1 + \ldots + l_{nj}x_n + l_{n+1,j}, \quad j = 1, ..., n+1,$$

are called the basic Lagrange polynomials corresponding to S.

These polynomials have the following property:

$$\lambda_j(x^{(k)}) = \delta_j^k \text{ for all } j, k = 1, ..., n + 1.$$

Here δ_i^k is the Kronecker delta. For an arbitrary $x \in \mathbb{R}^n$, we have

$$x = \sum_{j=1}^{n+1} \lambda_j(x) x^{(j)}, \quad \sum_{j=1}^{n+1} \lambda_j(x) = 1.$$

Thus, $\lambda_j(x)$ are *the barycentric coordinates of x* with respect to the simplex *S*. In turn, equations $\lambda_j(x) = 0$, j = 1, ..., n + 1, define the (n - 1)-dimensional hyperplanes containing the faces of *S*. Therefore,

$$S = \left\{ x \in \mathbb{R}^n : \lambda_j(x) \ge 0, \ j = 1, \dots, n+1 \right\}.$$

Also let us note that for every j = 1, ..., n + 1 we have

$$\lambda_j(x) = \frac{\Delta_j(x)}{\Delta}.$$
(2.3)

Here $\Delta = \det(\mathbf{A})$ and $\Delta_j(x)$ is the determinant that appears from Δ after replacing the *j*th row by the row $(x_1 \dots x_n 1)$. For more information on λ_j , see [25], [36].

In [24] we show that

$$\xi(K;S) = (n+1) \max_{1 \le k \le n+1} \max_{x \in K} (-\lambda_k(x)) + 1 \quad (K \not\subset S),$$
(2.4)

$$\alpha(K;S) = \sum_{j=1}^{n+1} \max_{x \in K} (-\lambda_j(x)) + 1$$
(2.5)

(see also [25]). The equality $\xi(K; S) = \alpha(K; S)$ holds true if and only if the simplex $\xi(S)S$ is circumscribed around *K*. This is also equivalent to the relation

$$\max_{x \in K} (-\lambda_1(x)) = \dots = \max_{x \in K} (-\lambda_{n+1}(x)).$$
(2.6)

If *K* is a convex polytope, then the maxima on *K* in (2.4)–(2.6) can also be taken over $x \in ver(K)$. Note that for $K = Q_n$ formula (2.4) is proved in [22].

For an arbitrary *n*-dimensional nondegenerate simplex *S*,

$$\alpha(Q_n; S) = \sum_{i=1}^n \frac{1}{d_i(S)}$$
(2.7)

Two various approaches to (2.7) are given in [24].

Definition 2.5 Let $x^{(j)} \in K$, $1 \le j \le n + 1$, be the vertices of a nondegenerate simplex *S*. The interpolation projector $P : C(K) \to \Pi_1(\mathbb{R}^n)$ with nodes $x^{(j)}$ is determined by the equalities $Pf(x^{(j)}) = f_j = f(x^{(j)})$, $1 \le j \le n + 1$. We say that an interpolation projector $P : C(K) \to \Pi_1(\mathbb{R}^n)$ and a nondegenerate simplex $S \subset K$ correspond to each other if the nodes of *P* coincide with the vertices of *S*. We use notation P_S and S_P .

For an interpolation projector $P = P_S$, the analogue of Lagrange interpolation formula holds:

$$Pf(x) = \sum_{j=1}^{n+1} f(x^{(j)}) \lambda_j(x).$$
 (2.8)

Here λ_i are the basic Lagrange polynomials of the simplex S_P (see Definition 1.4).

We let $||P||_K$ denote the norm of *P* as an operator from *C*(*K*) into *C*(*K*). Thanks to (2.8),

$$\begin{split} \|P\|_{K} &= \sup_{\|f\|_{C(K)}=1} \|Pf\|_{C(K)} = \sup_{-1 \le f_{j} \le 1} \max_{x \in K} \left| \sum_{j=1}^{n+1} f_{j} \lambda_{j}(x) \right| \\ &= \max_{x \in K} \sup_{-1 \le f_{j} \le 1} \left| \sum_{j=1}^{n+1} f_{j} \lambda_{j}(x) \right| = \max_{x \in K} \sup_{-1 \le f_{j} \le 1} \sum_{j=1}^{n+1} f_{j} \lambda_{j}(x). \end{split}$$

Because $\sum f_j \lambda_j(x)$ is linear in x and f_1, \ldots, f_{n+1} , we have

$$||P||_{K} = \max_{x \in K} \max_{f_{j}=\pm 1} \sum_{j=1}^{n+1} f_{j}\lambda_{j}(x) = \max_{x \in K} \sum_{j=1}^{n+1} |\lambda_{j}(x)|.$$
(2.9)

If *K* is a convex polytope in \mathbb{R}^n (e. g., *K* is a cube), a simpler equality

$$||P||_{K} = \max_{x \in \operatorname{ver}(K)} \sum_{j=1}^{n+1} |\lambda_{j}(x)|$$
(2.10)

holds.

Definition 2.6 We let $\theta_n(K)$ denote the minimal value of $||P_S||_K$ where *S* runs over all nondegenerate simplices with vertices in *K*. An interpolation projector $P : C(K) \to \Pi_1(\mathbb{R}^n)$ is called minimal if $||P||_K = \theta_n(K)$.

It was shown in [20] that for any interpolation projector $P : C(K) \to \Pi_1(\mathbb{R}^n)$ and the corresponding simplex *S* we have

$$\frac{n+1}{2n} \left(\|P\|_{K} - 1 \right) + 1 \le \xi(K; S) \le \frac{n+1}{2} \left(\|P\|_{K} - 1 \right) + 1.$$
(2.11)

Thanks to (2.11),

$$\frac{n+1}{2n} \Big(\theta_n(K) - 1 \Big) + 1 \le \xi_n(K) \le \frac{n+1}{2} \Big(\theta_n(K) - 1 \Big) + 1.$$
(2.12)

Obviously, if a projector *P* satisfies the equality

$$\xi_n(K) = \frac{n+1}{2} (||P||_K - 1) + 1, \qquad (2.13)$$

then *P* is minimal and the right-hand relation in (2.12) becomes an equality.

Occasionally, we will consider the case when n + 1 is an Hadamard number, i.e., there exists an Hadamard matrix of order n + 1.

Definition 2.7 An Hadamard matrix of order *m* is a square binary matrix **H** with entries either 1 or -1 satisfying the equality

$$\mathbf{H}^{-1} = \frac{1}{m} \mathbf{H}^T.$$

An integer *m*, for which an Hadamard matrix of order *m* exists, is called an Hadamard number.

Thus, the rows of **H** are pairwise orthogonal with respect to the standard scalar product on \mathbb{R}^{m} .

The order of an Hadamard matrix is 1 or 2 or some multiple of 4 (see [13]). But it is still unknown whether an Hadamard matrix exists for every order of the form m = 4k. This is one of the longest lasting open problems in Mathematics called *the Hadamard matrix conjecture*. The orders below 1500 for which Hadamard matrices are not known are 668, 716, 892, 956, 1132, 1244, 1388, and 1436 (see, e.g., [11, 17]).

Definition 2.8 By h_n we denote the maximum value of a determinant of order *n* with entries 0 or 1. Let v_n be the maximum volume of an *n*-dimensional simplex contained in Q_n . The numbers h_n and v_n satisfy the equality $h_n = n!v_n$ (see [14]). For any *n*, there exists in Q_n a maximum volume simplex with some vertex coinciding with a vertex of the cube. For n > 1, the following inequalities hold

$$\frac{1}{2} \left(1 - \frac{\log(4/3)}{\log n} \right) n \log n < \log(2^{n-1}h_{n-1}) \le \frac{1}{2} n \log n.$$
(2.14)

The right-hand inequality in (2.14) was proved by Hadamard [12] and the left-hand one by Clements and Lindström [4]. Consequently, for all $n \in \mathbb{N}$

$$\left(\frac{3}{4}\right)^{(n+1)/2} \frac{(n+1)^{(n+1)/2}}{2^n} < h_n \le \frac{(n+1)^{(n+1)/2}}{2^n},\tag{2.15}$$

$$\left(\frac{3}{4}\right)^{(n+1)/2} \frac{(n+1)^{(n+1)/2}}{2^n n!} < \nu_n \le \frac{(n+1)^{(n+1)/2}}{2^n n!}.$$
(2.16)

The right-hand equality in each relation holds if and only if n + 1 is an Hadamard number [14]. In some cases the right-hand inequality in (2.16) has been improved. For instance, if n is even, then

$$v_n \le \frac{n^{n/2} \sqrt{2n+1}}{2^n n!}.$$
(2.17)

If n > 1 and $n \equiv 1 \pmod{4}$, then

$$\nu_n \le \frac{(n-1)^{(n-1)/2}}{2^{n-1}(n-1)!} \tag{2.18}$$

(see [14]). For many *n*, the values of v_n and h_n are known exactly. The first 12 numbers v_n are

$$v_1 = 1, \quad v_2 = \frac{1}{2}, \quad v_3 = \frac{1}{3}, \quad v_4 = \frac{1}{8}, \quad v_5 = \frac{1}{24}, \quad v_6 = \frac{1}{80}, \quad v_7 = \frac{2}{315},$$

 $v_8 = \frac{1}{720}, \quad v_9 = \frac{1}{2520}, \quad v_{10} = \frac{1}{11340}, \quad v_{11} = \frac{9}{246400}, \quad v_{12} = \frac{3}{394240}.$

Definition 2.9 Let \varkappa_n be the volume of the ball B_n , and let σ_n be the volume of a regular simplex inscribed into B_n .

The numbers \varkappa_n and σ_n are known exactly. Namely,

$$\varkappa_n = \frac{\pi^{n/2}}{\Gamma(n/2+1)}, \qquad \sigma_n = \frac{1}{n!} \sqrt{n+1} \left(\frac{n+1}{n}\right)^{n/2}, \tag{2.19}$$

$$\varkappa_{2k} = \frac{\pi^k}{k!}, \qquad \varkappa_{2k+1} = \frac{2^{k+1}\pi^k}{(2k+1)!!} = \frac{2(k!)(4\pi)^k}{(2k+1)!}.$$
(2.20)

Definition 2.10 We let $simp_n(K)$ denote the maximum volume of a nondegenerate simplex *S* with vertices in *K*.

Obviously, $simp_n(Q_n) = v_n$. A regular simplex inscribed into an *n*-dimensional ball has the maximum volume among all simplices contained in this ball. There are no other simplices with this property (see [7,43,46]). Thus, $simp_n(B_n) = \sigma_n$.

3. Inequalities $\xi_n(K) \le n+2$ and $\theta_n(K) \le n+1$

Theorem 3.1 Let K be a convex body in \mathbb{R}^n , and let S be an arbitrary maximum volume simplex contained in K. Then

$$\xi(K;S) \le n+2$$
 and $||P_S||_K \le n+1.$ (3.1)

Proof. Using a purely geometric approach, M. Lassak [16] have proved that for every maximum volume simplex S in K the following inclusions

$$S \subset K \subset (n+2)S \tag{3.2}$$

hold. From this and Definition 2.2, we have $\xi(K; S) \le n + 2$.

Let us note that (3.2) also follows from formula (2.4). In fact, because simplex $S \subset K$ has maximum volume, $|\Delta_j(x)| \leq |\Delta|$ for any j = 1, ..., n + 1 and $x \in K$. (This inequality is immediate from (2.1).) Thanks to (2.3),

$$-\lambda_j(x) \le |\lambda(x)| = \frac{|\Delta_j(x)|}{|\Delta|} \le 1, \quad x \in K.$$
(3.3)

(Recall that λ_j are the basic Lagrange polynomials for the simplex S. See Definition 2.4.) By (2.4),

$$\xi(K;S) = (n+1) \max_{1 \le k \le n+1} \max_{x \in K} (-\lambda_k(x)) + 1 \le n+2$$

proving the left-hand inequality in (3.1).

The right-hand inequality in (3.1) follows from (2.9). Indeed, because $|\lambda(x)| \le 1$, we have

$$||P_S||_K = \max_{x \in K} \sum_{j=1}^{n+1} |\lambda_j(x)| \le n+1.$$

The proof of the theorem is complete.

The following corollary is immediate from Theorem 3.1.

Corollary 3.2 *For any covex body* $K \subset \mathbb{R}^n$ *,*

$$\xi_n(K) \le n+2, \qquad \theta_n(K) \le n+1. \tag{3.4}$$

4. Legendre polynomials and the measure of $E_{n,\gamma}$

The standardized Legendre polynomial of degree n is the function

$$\chi_n(t) = \frac{1}{2^n n!} \left[(t^2 - 1)^n \right]^{(n)}, \quad t \in \mathbb{R}$$

(Rodrigues' formula). For the properties of χ_n see, e.g., [44,45]. Legendre polynomials are orthogonal on the segment [-1, 1] with the weight $w(t) \equiv 1$. The first standardized Legendre polynomials are

$$\chi_0(t) = 1, \quad \chi_1(t) = t, \quad \chi_2(t) = \frac{1}{2} \left(3t^2 - 1 \right), \quad \chi_3(t) = \frac{1}{2} \left(5t^3 - 3t \right),$$
$$\chi_4(t) = \frac{1}{8} \left(35t^4 - 30t^2 + 3 \right), \quad \chi_5(t) = \frac{1}{8} \left(63t^5 - 70t^3 + 15t \right).$$

For these polynomials the following recurrent relations

$$\chi_{n+1}(t) = \frac{2n+1}{n+1} t \cdot \chi_n(t) - \frac{n}{n+1} \chi_{n-1}(t)$$
(4.1)

hold. In particular, this implies $\chi_n(1) = 1$. Let us also note that, if $n \ge 1$ then $\chi_n(t)$ increases as $t \ge 1$. (This property also easily follows from formula (4.3) given below.)

We let χ_n^{-1} denote the function inverse to χ_n on the semi-axis $[1, +\infty)$.

One of the key statements of our approach to the problems of optimal Lagrangian interpolation is Theorem 4.1 formulated and proven in this section. This theorem reveals some rather surprising connections between Legendre polynomials with the problems of optimal Lagrangian interpolation.

Given $\gamma \ge 1$, we let $E_{n,\gamma}$ define a set

$$E_{n,\gamma} = \left\{ x \in \mathbb{R}^n : \sum_{j=1}^n |x_j| + \left| 1 - \sum_{j=1}^n x_j \right| \le \gamma \right\}.$$
 (4.2)

Theorem 4.1 The following equalities hold:

$$\operatorname{mes}_{n}(E_{n,\gamma}) = \frac{1}{2^{n}n!} \sum_{i=0}^{n} {\binom{n}{i}}^{2} (\gamma - 1)^{n-i} (\gamma + 1)^{i} = \frac{\chi_{n}(\gamma)}{n!}.$$
(4.3)

This result was obtained in [18]. Unfortunately, this paper is practically inaccessible to a wide audience, so for the convenience of the reader we present it here.

Proof. First, let us prove the left-hand equality in (4.3). Let

$$E^{(1)} = \{x \in E_{n,\gamma} : \sum x_i > 1\}$$
 and $E^{(2)} = \{x \in E_{n,\gamma} : \sum x_i \le 1\}.$

Let us give explicit formulae for the volumes $m_1 = \text{mes}_n(E^{(1)})$ and $m_2 = \text{mes}_n(E^{(2)})$.

Let us fix $k, 1 \le k \le n$, and consider a non-empty subset $G \subset E^{(1)}$ consisting of all points $x = (x_1, ..., x_n)$ such that $x_1, ..., x_k \ge 0$ and $x_{k+1}, ..., x_n < 0$. Let $y_i = x_i$ for i = 1, ..., k, $y_i = -x_i$ for i = k + 1, ..., n, and let $y = (y_1, ..., y_n)$. Then

$$G = \left\{ y : 1 + y_{k+1} + \ldots + y_n \le y_1 + \ldots + y_k \le \frac{\gamma + 1}{2}, \ y_i \ge 0 \right\},\$$

hence

$$\operatorname{mes}_{n}(G) = \int_{1}^{\alpha} dy_{1} \int_{1}^{\alpha-y_{1}} dy_{2} \dots \int_{1}^{\alpha-y_{1}-\dots-y_{k-1}} dy_{k}$$
$$\int_{0}^{y_{1}+\dots+y_{k}-1} dy_{k+1} \int_{0}^{y_{1}+\dots+y_{k}-1-y_{k+1}} dy_{k+2} \dots \int_{0}^{y_{1}+\dots+y_{k}-1-y_{k+1}-\dots-y_{n-1}} dy_{n}.$$

Throughout the proof, $\alpha = (\gamma + 1)/2$. If b > 0, then

$$\int_{0}^{b} dz_{1} \int_{0}^{b-z_{1}} dz_{2} \dots \int_{0}^{b-z_{1}-\dots-z_{l-1}} dz_{l} = \frac{b^{l}}{l!},$$

$$\operatorname{mes}_{n}(G) = \int_{1}^{\alpha} dy_{1} \int_{1}^{\alpha-y_{1}} dy_{2} \dots \int_{1}^{\alpha-y_{1}-\dots-y_{k-1}} \frac{1}{(n-k)!} (y_{1}+\dots+y_{k}-1)^{n-k} dy_{k}$$
$$= \left(\int_{y_{1}+\dots+y_{k}\leq\alpha} -\int_{y_{1}+\dots+y_{k}\leq1}\right) \frac{1}{(n-k)!} (y_{1}+\dots+y_{k}-1)^{n-k} dy_{1} \dots dy_{k}$$
$$= J_{1}-J_{2}.$$

The first integral equals

$$J_1 = \sum_{j=1}^k (-1)^{j+1} \frac{(\alpha-1)^{n-k+j}}{(n-k+j)!} \frac{\alpha^{k-j}}{(k-j)!} + \frac{(-1)^{n+k}}{n!}.$$

The value of J_2 appears from this expression if instead of α we take 1. Consequently,

$$\operatorname{mes}_{n}(G) = J_{1} - J_{2}$$

$$= \sum_{j=1}^{k} (-1)^{j+1} \frac{(\alpha - 1)^{n-k+j}}{(n-k+j)!} \frac{\alpha^{k-j}}{(k-j)!}$$

$$= \frac{(-1)^{k+1}}{n!} \sum_{i=0}^{k-1} {n \choose i} (\alpha - 1)^{n-i} (-\alpha)^{i}.$$
(4.4)

Clearly, the set $E^{(1)}$ is the union of all pairwise disjoint sets *G* with various k = 1, ..., n. Therefore, the measure of $E^{(1)}$ is equal to

$$m_1 = \sum_{k=1}^n \binom{n}{k} \frac{(-1)^{k+1}}{n!} \sum_{i=0}^{k-1} \binom{n}{i} (\alpha - 1)^{n-i} (-\alpha)^i.$$

Changing the order of summation and using the identity

$$\sum_{k=0}^{i} (-1)^k \binom{n}{k} = (-1)^i \binom{n-1}{i}$$
(4.5)

(see, e. g., [39]) we get

$$m_{1} = \frac{1}{n!} \sum_{i=0}^{n-1} {n \choose i} (\alpha - 1)^{n-i} (-\alpha)^{i} \sum_{k=0}^{i} (-1)^{k} {n \choose k}$$
$$= \frac{1}{n!} \sum_{i=0}^{n-1} {n \choose i} (\alpha - 1)^{n-i} \alpha^{i}.$$
(4.6)

Now, let us turn to $E^{(2)}$. First, note that $E^{(2)}$ contains the domain $S = \{x_i \ge 0, \sum x_i \le 1\}$ with the measure 1/n!. Next, fix $k \in \{1, ..., n\}$ and consider the subset $G' \subset E^{(2)}$ corresponding to the

inequalities $x_1, \ldots, x_k < 0$; $x_{k+1}, \ldots, x_n \ge 0$. Put $y_1 = -x_1, \ldots, y_k = -x_k$; $y_{k+1} = x_{k+1}, \ldots, y_n = x_n$. Then

$$G' = \{y : y_{k+1} + \ldots + y_n \le 1 + y_1 + \ldots + y_k \le \frac{\gamma - 1}{2}, y_i \ge 0\}.$$

Let $\beta = (\gamma - 1)/2$. Then the following equalities hold:

$$\begin{aligned} \operatorname{mes}_{n}(G') &= \int_{0}^{\beta} dy_{1} \int_{0}^{\beta-y_{1}} dy_{2} \dots \int_{0}^{\beta-y_{1}-\dots-y_{k-1}} dy_{k} \\ & \int_{0}^{1+y_{1}+\dots+y_{k}} dy_{k+1} \int_{0}^{1+y_{1}+\dots+y_{k}-y_{k+1}} dy_{k+2} \dots \int_{0}^{1+y_{1}+\dots+y_{k}-y_{k+1}-\dots-y_{n-1}} dy_{n} \\ &= \int_{0}^{\beta} dy_{1} \int_{0}^{\beta-y_{1}} dy_{2} \dots \int_{0}^{\beta-y_{1}-\dots-y_{k-1}} \frac{(1+y_{1}+\dots+y_{k})^{n-k}}{(n-k)!} dy_{k} \\ &= \left[\sum_{j=0}^{k-1} (-1)^{k-1-j} \frac{(1+\beta)^{n-j}\beta^{j}}{(n-j)! j!} \right] + \frac{(-1)^{k}}{n!} = \frac{(-1)^{k+1}}{n!} \left(\left[\sum_{j=0}^{k-1} \binom{n}{j} (1+\beta)^{n-j} (-\beta)^{j} \right] - 1 \right). \end{aligned}$$

Clearly, the set $E^{(2)}$ is the union of all such sets G' corresponding to various k = 1, ..., n, and the simplex S. Therefore,

$$m_2 = \operatorname{mes}_{n}(E^{(2)}) = \frac{1}{n!} \left(\left\{ \sum_{k=1}^{n} (-1)^{k+1} \binom{n}{k} \left(\left[\sum_{j=0}^{k-1} \binom{n}{j} (1+\beta)^{n-j} (-\beta)^{j} \right] - 1 \right) \right\} + 1 \right).$$

Note that $1 + \beta = (\gamma + 1)/2 = \alpha$ and $\beta = (\gamma - 1)/2 = \alpha - 1$. Let us make the substitution i = n - j in the internal sum in the right hand side of this equality, and make use of the formula

$$\sum_{k=0}^{n} (-1)^{k} \binom{n}{k} = \sum_{k=1}^{n} (-1)^{k} \binom{n}{k} + 1 = 0.$$

We obtain the following:

$$m_{2} = \frac{1}{n!} \left(1 + \sum_{k=1}^{n} (-1)^{k+1} \binom{n}{k} \left((-1)^{n} \sum_{i=n-k+1}^{n} \binom{n}{i} (\alpha - 1)^{n-i} (-\alpha)^{i} - 1 \right) \right)$$

$$= \frac{(-1)^{n}}{n!} \sum_{k=1}^{n} (-1)^{k+1} \binom{n}{k} \sum_{i=n-k+1}^{n} \binom{n}{i} (\alpha - 1)^{n-i} (-\alpha)^{i}.$$

Changing the order of summation, we obtain

$$m_2 = \frac{(-1)^n}{n!} \sum_{i=1}^n \binom{n}{i} (\alpha - 1)^{n-i} (-\alpha)^i \sum_{k=n+1-i}^n (-1)^{k+1} \binom{n}{k}.$$

Using (4.5), we can write

$$\sum_{k=n+1-i}^{n} (-1)^{k+1} \binom{n}{k} = \sum_{k=n+1-i}^{n} (-1)^{k+1} \binom{n}{n-k} = \sum_{j=0}^{i-1} (-1)^{n-j+1} \binom{n}{j} = (-1)^{n+i} \binom{n-1}{i-1}$$

Therefore,

$$m_2 = \frac{1}{n!} \sum_{i=1}^n \binom{n}{i} \binom{n-1}{i-1} (\alpha - 1)^{n-i} \alpha^i.$$
(4.7)

From this, (4.6) and (4.7), we have

$$\begin{aligned} \operatorname{mes}_{n}(E_{n,\gamma}) &= m_{1} + m_{2} \\ &= \frac{1}{n!} \sum_{i=0}^{n-1} \binom{n}{i} \binom{n-1}{i} (\alpha - 1)^{n-i} \alpha^{i} + \frac{1}{n!} \sum_{i=1}^{n} \binom{n}{i} \binom{n-1}{i-1} (\alpha - 1)^{n-i} \alpha^{i} \\ &= \frac{1}{n!} \sum_{i=1}^{n-1} \binom{n}{i} \binom{n-1}{i} + \binom{n-1}{i-1} (\alpha - 1)^{n-i} \alpha^{i} + \frac{1}{n!} ((\alpha - 1)^{n} + \alpha^{n}) \\ &= \frac{1}{n!} \sum_{i=0}^{n} \binom{n}{i}^{2} (\alpha - 1)^{n-i} \alpha^{i} = \frac{1}{2^{n} n!} \sum_{i=0}^{n} \binom{n}{i}^{2} (\gamma - 1)^{n-i} (\gamma + 1)^{i} \end{aligned}$$

completing the proof of the left-hand equality in (4.3). We took into account that

$$\binom{n-1}{i} + \binom{n-1}{i-1} = \binom{n}{i}.$$

The right-hand equality in (4.3) follows from the identity

$$\sum_{i=0}^{n} {\binom{n}{i}}^{2} t^{i} = (1-t)^{n} \chi_{n} \left(\frac{1+t}{1-t}\right). \quad \text{See [39]}.$$

Let us set $t = (\gamma - 1)/(\gamma + 1)$. Then $(1 - t)^n = 2^n(\gamma + 1)^{-n}$ and $(1 + t)/(1 - t) = \gamma$. Hence,

$$\operatorname{mes}_{n}(G) = \frac{1}{2^{n}n!} \sum_{i=0}^{n} {\binom{n}{i}}^{2} (\gamma - 1)^{n-i} (\gamma + 1)^{i} = \frac{1}{2^{n}n!} \sum_{i=0}^{n} {\binom{n}{i}}^{2} (\gamma + 1)^{n-i}$$
$$= \frac{1}{2^{n}n!} (\gamma + 1)^{n} \sum_{i=0}^{n} {\binom{n}{i}}^{2} (\frac{\gamma - 1}{\gamma + 1})^{i} = \frac{\chi_{n}(\gamma)}{n!}.$$

The proof of Theorem 4.1 is complete.

Let us give some simple examples. The set $E_{1,2} = \{x \in \mathbb{R} : |x|+|1-x| \le 2\}$ is the segment [-1/2, 3/2] with the length mes₁($E_{1,2}$) = $\chi_1(2)/1! = 2$. The set $E_{2,2} = \{x \in \mathbb{R}^2 : |x_1| + |x_2| + |1 - x_1 - x_2| \le 2\}$ is a hexagon on the plane with the area mes₂($E_{2,2}$) = $\chi_2(2)/2! = 11/4$.

The three-dimensional domain

$$E_{3,2} = \{x \in \mathbb{R}^3 : |x_1| + |x_2| + |x_3| + |1 - x_1 - x_2 - x_3| \le 2\}$$

is shown in Fig. 1. It has the volume $mes_3(E_{3,2}) = \chi_3(2)/3! = 17/6$.

There is an interesting open problem related to the above-mentioned equality (4.3). Essentially, this equality along with Rodrigues' formula and other well-known relations gives a characterization of Legendre polynomials. This characterization is written via the volumes of some convex polyhedra. Namely, for $t \ge 1$

$$\chi_n(t) = n! \operatorname{mes}_n(E_{n,t}), \tag{4.8}$$

where $E_{n,t}$ is defined by (4.2).



The question arises about analogues of (4.8) for other classes of orthogonal polynomials, such as Chebyshev polynomials or, more generally, Jacobi polynomials: *Is the equality* (4.8) *a particular case of a more general pattern?* We would be grateful for any information on this matter.

Let us note that, from (4.8) and (4.1), we have

$$\operatorname{mes}_{n+1}(E_{n+1,t}) = \frac{2n+1}{(n+1)^2} t \operatorname{mes}_n(E_{n,t}) - \frac{1}{(n+1)^2} \operatorname{mes}_{n-1}(E_{n-1,t})$$

The direct establishing this recurrence relation for the measures of $E_{n,t}$ could provide a proof of Theorem 4.1 different from the above.

5. Inequality
$$\theta_n(K) \ge \chi_n^{-1} \left(\frac{\operatorname{vol}(K)}{\operatorname{simp}_n(K)} \right)$$

Based on Theorem 4.1, in this section we obtain lower bounds for the norm of the projection operator associated with linear interpolation on an arbitrary convex body K.

Theorem 5.1 Suppose $P : C(K) \to \Pi_1(\mathbb{R}^n)$ is an arbitrary interpolation projector. Then for the corresponding simplex $S \subset K$ and the node matrix **A** we have

$$||P||_{K} \ge \chi_{n}^{-1} \left(\frac{n! \operatorname{vol}(K)}{|\det(\mathbf{A})|} \right) = \chi_{n}^{-1} \left(\frac{\operatorname{vol}(K)}{\operatorname{vol}(S)} \right).$$
(5.1)

Proof. For each i = 1, ..., n, let us subtract from the *i*th row of **A** its (n + 1)th row. Denote by **B** the submatrix of order *n* which stands in the first *n* rows and columns of the result matrix. Then

 $|\det(\mathbf{B})| = |\det(\mathbf{A})| = n! \operatorname{vol}(S) \le n! \operatorname{vol}(K)$

so that

$$\frac{|\det(\mathbf{B})|}{n!\operatorname{vol}(K)} \le 1.$$
(5.2)

Let $x^{(j)}$ be the vertices and λ_j be the basic Lagrange polynomials of simplex S. Since $\lambda_1(x), \ldots, \lambda_{n+1}(x)$ are the barycentric coordinates of a point x with respect to S, we have

$$||P||_{K} = \max_{x \in K} \sum_{j=1}^{n+1} |\lambda_{j}(x)| = \max\left\{\sum_{j=1}^{n+1} |\beta_{j}| : \sum_{j=1}^{n+1} \beta_{j} = 1, \sum_{j=1}^{n+1} \beta_{j} x^{(j)} \in K\right\}.$$

Let us replace β_{n+1} with the equal value $1 - \sum_{j=1}^{n} \beta_j$. The condition $\sum_{j=1}^{n+1} \beta_j x^{(j)} \in K$ is equivalent to

$$\sum_{j=1}^{n} \beta_j (x^{(j)} - x^{(n+1)}) \in K' = K - x^{(n+1)}.$$

Hence,

$$\|P\|_{K} = \max\left\{\sum_{j=1}^{n} |\beta_{j}| + \left|1 - \sum_{j=1}^{n} \beta_{j}\right|\right\}$$
(5.3)

where the maximum is taken over all β_j such that $\sum_{j=1}^n \beta_j (x^{(j)} - x^{(n+1)}) \in K'$. Clearly, vol(K') = vol(K).

Let us consider the nondegenerate linear operator $F : \mathbb{R}^n \to \mathbb{R}^n$, which maps the point $\beta = (\beta_1, \dots, \beta_n)$ to the point $x = F(\beta)$ according to the rule

$$x = \sum_{j=1}^{n} \beta_j \left(x^{(j)} - x^{(n+1)} \right)$$

We have the matrix equality $F(\beta) = (\beta_1, ..., \beta_n)\mathbf{B}$, where **B** is the above introduced matrix of order *n* with the entries $b_{ij} = x_j^{(i)} - x_j^{(n+1)}$. Let us put

$$\gamma^* = \chi_n^{-1} \left(\frac{n! \operatorname{vol}(K)}{|\det \mathbf{B}|} \right).$$

Note that, thanks to (5.2), $(n!\operatorname{vol}(K))/|\det(\mathbf{B})| \ge 1$ so that the constant γ^* is well defined. Let us also note that $\chi_n(\gamma^*) = n!\operatorname{vol}(K)/|\det(\mathbf{B})|$.

Given $\gamma \ge 1$, let us introduce a set

$$E_{n,\gamma} = \left\{ \beta = (\beta_1, \ldots, \beta_n) \in \mathbb{R}^n : \sum_{j=1}^n |\beta_j| + \left| 1 - \sum_{j=1}^n \beta_j \right| \le \gamma \right\}.$$

Let us show that $K' \not\subset F(E_{n,\gamma})$ provided $\gamma < \gamma^*$. Indeed, this is immediate from Theorem 4.1 because, thanks to this theorem,

$$\operatorname{mes}_{n}(F(E_{n,\gamma})) < \operatorname{mes}_{n}(F(E_{n,\gamma^{*}})) = |\det \mathbf{B}| \cdot \operatorname{mes}_{n}(E_{n,\gamma^{*}})$$
$$= |\det \mathbf{B}| \cdot \frac{\chi_{n}(\gamma^{*})}{n!} = \operatorname{vol}(K) = \operatorname{vol}(K').$$

Thus, for every $\varepsilon > 0$ there exists a point $x^{(\varepsilon)}$ with the properties:

$$x^{(\varepsilon)} = \sum \beta_j^{(\varepsilon)} \left(x^{(j)} - x^{(n+1)} \right) \in K' \text{ and } \left| \sum \beta_j^{(\varepsilon)} \right| + \left| 1 - \sum \beta_j^{(\varepsilon)} \right| \ge \gamma^* - \varepsilon.$$

In view of (5.3) this gives $||P||_K \ge \gamma^* - \varepsilon$. Since $\varepsilon > 0$ is an arbitrary, we obtain

$$||P||_{K} \ge \gamma^{*} = \chi_{n}^{-1} \left(\frac{n! \operatorname{vol}(K)}{|\det(\mathbf{B})|} \right) = \chi_{n}^{-1} \left(\frac{n! \operatorname{vol}(K)}{|\det(\mathbf{A})|} \right) = \chi_{n}^{-1} \left(\frac{\operatorname{vol}(K)}{\operatorname{vol}(S)} \right)$$

The theorem is proved.

Recall that $simp_n(K)$ denotes the maximum volume of a nondegenerate simplex S with vertices in K (see Definition 2.10).

Theorem 5.2 Let K be an arbitrary convex body in \mathbb{R}^n . Then

$$\theta_n(K) \ge \chi_n^{-1} \left(\frac{\operatorname{vol}(K)}{\operatorname{simp}_n(K)} \right).$$
(5.4)

Proof. By (5.1), for any interpolation projector $P : C(K) \to \Pi_1(\mathbb{R}^n)$,

$$||P||_{K} \ge \chi_{n}^{-1}\left(\frac{\operatorname{vol}(K)}{\operatorname{vol}(S)}\right) \ge \chi_{n}^{-1}\left(\frac{\operatorname{vol}(K)}{\operatorname{simp}_{n}(K)}\right),$$

where $S = S_P$. This immediately gives (5.4).

Remark 5.3 If simplex $S \subset K$ has maximum volume, then $K \subset (n + 2)S$, see (3.2). Therefore, the ratio $vol(K)/simp_n(K)$ in (5.4) is bounded from above by $(n + 2)^n$.

In the case $K = Q_n$ we have vol(K) = 1, $simp_n(K) = v_n$ (see Definition 2.8). If $K = B_n$, then $vol(K) = \varkappa_n$, $simp_n(K) = \sigma_n$ (see Definition 2.9). Thus, in these cases, (5.4) imply the following two inequalities:

$$\theta_n(Q_n) \ge \chi_n^{-1}\left(\frac{1}{\nu_n}\right) \quad \text{and} \quad \theta_n(B_n) \ge \chi_n^{-1}\left(\frac{\varkappa_n}{\sigma_n}\right).$$
(5.5)

In the next section we will give two important consequences of these inequalities. \triangleleft

Remark 5.4 Inequalities (3.4) and (5.4) show that for each convex body $K \subset \mathbb{R}^n$

$$\chi_n^{-1}\left(\frac{\operatorname{vol}(K)}{\operatorname{simp}_n(K)}\right) \le \theta_n(K) \le n+1.$$

In Section 6 we prove analogues of these inequalities for an arbitrary (not necessarily convex) compact set $E \subset \mathbb{R}^n$ with vol(conv(E)) > 0.

6. Inequalities
$$\theta_n(Q_n) > c \sqrt{n}$$
 and $\theta_n(B_n) > c \sqrt{n}$

The Stirling formula (see, e. g., [8]) $n! = \sqrt{2\pi n} (n/e)^n e^{\frac{\zeta_n}{12n}}, 0 < \zeta_n < 1$, yields

$$\sqrt{2\pi n} \left(\frac{n}{e}\right)^n < n! < \sqrt{2\pi n} \left(\frac{n}{e}\right)^n e^{\frac{1}{12n}}.$$
(6.1)

Let us also see that for every n > 1 the following inequality

$$\chi_n^{-1}(s) > \left(\frac{s}{\binom{n}{\lfloor n/2 \rfloor}}\right)^{1/n}$$
(6.2)

holds. In fact, if $t \ge 1$ and n > 1, then, according to (4.3),

$$\chi_{n}(t) = \frac{1}{2^{n}} \sum_{i=0}^{n} {\binom{n}{i}}^{2} (t-1)^{n-i} (t+1)^{i} < {\binom{n}{\lfloor n/2 \rfloor}} \cdot \frac{1}{2^{n}} \sum_{i=0}^{n} {\binom{n}{i}} (t-1)^{n-i} (t+1)^{i}$$
$$= {\binom{n}{\lfloor n/2 \rfloor}} \cdot (2t)^{n} \cdot 2^{-n} = {\binom{n}{\lfloor n/2 \rfloor}} t^{n}$$

proving (6.2). For even *n*, we have

$$\binom{n}{\lfloor n/2 \rfloor} = \binom{n}{n/2} = \frac{n!}{\left((n/2)!\right)^2},$$

therefore,

$$\chi_n^{-1}(s) > \left(\frac{s\left((n/2)!\right)^2}{n!}\right)^{1/n}.$$
(6.3)

If *n* is odd, then

$$\binom{n}{\lfloor n/2 \rfloor} = \frac{n!}{\frac{n+1}{2}!\frac{n-1}{2}!}$$

so that, thanks to (6.2),

$$\chi_n^{-1}(s) > \left(\frac{s\frac{n+1}{2}!\frac{n-1}{2}!}{n!}\right)^{1/n}.$$
(6.4)

Theorem 6.1 *For all* $n \in \mathbb{N}$ *,*

$$\theta_n(Q_n) > \frac{\sqrt{n-1}}{e}.$$
(6.5)

Proof. The case n = 1 is trivial since $\theta_1(Q_1) = 1$. If n > 1, from (6.3), (6.4) and the Hadamard inequality (2.16), we have $\nu_n \le (n+1)^{(n+1)/2} / 2^n n!$.

For even n, thanks to the first inequality in (5.5), the left-hand inequality in (6.1), and (6.3), we get

$$\begin{aligned} \theta_n(Q_n) &\geq \chi_n^{-1}\left(\frac{1}{\nu_n}\right) \geq \chi_n^{-1}\left(\frac{2^n n!}{(n+1)^{(n+1)/2}}\right) > 2\left(\frac{[(n/2)!]^2}{(n+1)^{(n+1)/2}}\right)^{1/n} \\ &> \frac{2}{(n+1)^{1/2+1/(2n)}}\left(\sqrt{\pi n}\left(\frac{n}{2e}\right)^{n/2}\right)^{2/n} = \frac{(\pi n)^{1/n} n}{e(n+1)^{1/2+1/(2n)}} > \frac{\sqrt{n-1}}{e}. \end{aligned}$$

If n is odd, then, thanks to (6.4),

$$\begin{aligned} \theta_n(Q_n) &\geq \chi_n^{-1}\left(\frac{1}{\nu_n}\right) \geq \chi_n^{-1}\left(\frac{2^n n!}{(n+1)^{(n+1)/2}}\right) > \left(\frac{2^n \frac{n+1}{2}! \frac{n-1}{2}!}{(n+1)^{(n+1)/2}}\right)^{1/n} \\ &> 2\left(\frac{\pi \sqrt{n^2 - 1} \left(n^2 - 1\right)^{(n-1)/2} (n+1)}{(2e)^n}\right)^{1/n} \\ &= \frac{1}{e} \pi^{1/n} \sqrt{n-1} (n+1)^{1/(2n)} > \frac{\sqrt{n-1}}{e} \end{aligned}$$

proving that (6.5) holds for all n.

In some situations, the estimates of Theorem 2.4 can be improved [25]. In particular, we prove that $\theta_n > \sqrt{n/e}$ if *n* is even, and $\theta_n > n/(e\sqrt{n-1})$ provided n > 1 and $n \equiv 1 \pmod{4}$.

Let us also note that, thanks to (6.5), the inequality $\theta_n(Q_n) > c \sqrt{n}$ holds with some c, 0 < c < 1. A suitable estimate is

$$\theta_n(Q_n) > \frac{2\sqrt{2}}{3e} \sqrt{n}. \tag{6.6}$$

Indeed, if $n \le 8$, then the right-hand side of (6.6) is less than 1, while for $n \ge 9$

$$\theta_n(Q_n) > \frac{\sqrt{n-1}}{e} \ge \frac{2\sqrt{2}}{3e} \sqrt{n}.$$

Notice that $\frac{2\sqrt{2}}{3e} = 0.3468\dots$

This approach was extended to linear interpolation on the unit ball B_n , see [27].

Corollary 6.2 For every n,

$$\theta_n(B_n) \ge \chi_n^{-1} \left(\frac{\pi^{\frac{n}{2}} n!}{\Gamma(n/2+1) \sqrt{n+1} ((n+1)/n)^{n/2}} \right).$$
(6.7)

If n = 2k, then (6.7) is equivalent to the inequality

$$\theta_{2k}(B_{2k}) \ge \chi_{2k}^{-1} \left(\frac{\pi^k (2k)!}{k! \sqrt{2k+1} \left((2k+1)/2k \right)^k} \right).$$
(6.8)

For n = 2k + 1 we have

$$\theta_{2k+1}(B_{2k+1}) \ge \chi_{2k+1}^{-1} \left(\frac{2(k!)(4\pi)^k}{\sqrt{2k+2} \left((2k+2)/(2k+1) \right)^{(2k+1)/2}} \right).$$
(6.9)

Proof. It is sufficient to apply the second inequality in (5.5), (2.19), and (2.20).

Thanks to (6.3) and (6.4), inequalities (6.8) and (6.9) imply the following low bound for the constant $\theta_n(B_n)$ (see [28]).

Theorem 6.3 *There exists an absolute constant* c > 0 *such that*

$$\theta_n(B_n) > c \sqrt{n}. \tag{6.10}$$

Inequality (6.10) takes place with the constant

$$c = \frac{\sqrt[3]{\pi}}{\sqrt{12e} \cdot \sqrt[6]{3}} = 0.2135....$$
(6.11)

7. Estimates for linear interpolation on an arbitrary compact set

In this section we show how the above approach leads to estimates of the minimum projector norm under linear interpolation on a compact set.

Let *E* be an arbitrary compact in \mathbb{R}^n . Everywhere in this section, we let *K* denote *the convex hull of the set E*. We will assume that vol(K) > 0. By $||P||_E$ we denote the norm of the interpolation projector $P : C(E) \to \Pi_1(\mathbb{R}^n)$ as an operator from C(E) to C(E). We let $\theta_n(E)$ denote the minimal norm $||P||_E$ over all projectors *P* with nodes in *E*. By $simp_n(E)$ we denote the maximum volume of a simplex with vertices in *E*.

We will need the following elementary lemma.

Lemma 7.1 If $\varphi : \mathbb{R}^n \to \mathbb{R}$ is a convex continuous function, then $\max_{K} \varphi = \max_{F} \varphi$.

Proof. The maxima from the condition exist since φ is a continuous function. As K = conv(E), for any $y \in K$ there exist a number *m*, points $y^{(1)}, \ldots, y^{(m)} \in E$ and numbers μ_1, \ldots, μ_m such that

$$y = \sum_{i=1}^{m} \mu_i y^{(i)}, \quad \mu_i \ge 0, \text{ and } \sum_{i=1}^{m} \mu_i = 1.$$

Clearly, $\varphi(y^{(i)}) \leq \max_{E} \varphi$. The convexity of φ implies

$$\varphi(\mathbf{y}) = \varphi\left(\sum_{i=1}^{m} \mu_i \mathbf{y}^{(i)}\right) \le \sum_{i=1}^{m} \mu_i \varphi(\mathbf{y}^{(i)}) \le \left(\sum_{i=1}^{m} \mu_i\right) \max_{E} \varphi = \max_{E} \varphi.$$

Therefore, $\max_{K} \varphi \leq \max_{E} \varphi$. The inverse inequality is obvious.

Clearly, the result of Lemma 7.1 is well-known. It is immediate from the following Bauer's maximum principle [2]. Any function that is convex and continuous, and defined on a convex and compact set, attains its maximum at some extreme point of that set. Consequently, maximum φ on K = conv(E) is attained at some extreme point of E.

Theorem 7.2 We have

$$\theta_n(E) \ge \chi_n^{-1} \left(\frac{\operatorname{vol}(K)}{\operatorname{simp}_n(E)} \right).$$
(7.1)

Proof. First, let us note that for an arbitrary polynomial $p \in \Pi_1(\mathbb{R}^n)$ the following inequality

$$\|p\|_E = \|p\|_K \tag{7.2}$$

holds. (Recall that K = conv(E).) This is immediate from Lemma 7.1 for the convex continuous function $\varphi(x) = |p(x)|$.

Let $P : C(E) \to \Pi_1(\mathbb{R}^n)$ be an arbitrary interpolation projector with nodes in *E*. We will consider it also as a projector from C(K). Thanks to (7.2), it follows that $||P||_E = ||P||_K$. Therefore, thanks to inequality (5.4) of Theorem 5.2,

$$\theta_n(E) \ge \theta_n(K) \ge \chi_n^{-1} \left(\frac{\operatorname{vol}(K)}{\operatorname{simp}_n(K)} \right).$$

Finally, let us see that $simp_n(K) = simp_n(E)$. Indeed, let us suppose $S \subset K$ is any simplex with some vertex $x \notin E$. Without changing other vertices, we can replace vertex x by vertex x' so that the volume of the resulting simplex does not decrease. Indeed, this volume increases with $dist(x; \Gamma)$, where Γ is the (n - 1)-dimensional hyperplane containing all the vertices of the simplex except x. Let Γ be given by the equation $q(z) = \langle a, z \rangle + a_0 = 0$, $a = (a_1, \dots, a_n) \in \mathbb{R}^n$, $a_0 \in \mathbb{R}$. Then

$$\operatorname{dist}(x;\Gamma) = \frac{|q(x)|}{||a|||}$$

is obviously a convex continuous function. By Lemma 7.1, the maximum of $dist(x; \Gamma)$ on *K* is attained at some point $x' \in E$. Applying this procedure sequentially to all vertices of the simplex not belonging to *E*, we construct a new simplex with vertices on *E* without reducing the volume.

Thus, $simp_n(K) = simp_n(E)$, and the proof is complete.

Theorem 7.3 Let $E \subset \mathbb{R}^n$ be a compact set such that vol(conv(E)) > 0. Then the following inequality

$$\theta_n(E) \le n+1 \tag{7.3}$$

holds.

Proof. For an arbitrary interpolation projector $P: C(E) \to \Pi_1(\mathbb{R}^n)$, we have the following equality

$$||P||_E = \max_{x \in E} \sum_{j=1}^{n+1} |\lambda_j(x)|.$$

This can be proved in the same way as in the case E = K (see (2.9)).

Let *S* be a simplex with maximum volume over all family of simplices with vertices in *E*. For this simplex, $|\lambda_j(x)| \le 1$ (see the proof of Theorem 3.1, inequality (3.3)). By the above formula, the corresponding projector $P = P_S$ satisfies $||P_S||_E \le n + 1$. Consequently, $\theta_n(E) \le n + 1$.

Let us combine inequalities (7.1) and (7.3). If $E \subset \mathbb{R}^n$ is an arbitrary compact set satisfying the condition vol(conv(E)) > 0, then

$$\chi_n^{-1}\left(\frac{\operatorname{vol}(\operatorname{conv}(E))}{\operatorname{simp}_n(E)}\right) \le \theta_n(E) \le n+1.$$

8. Concluding remarks and open questions

Despite the apparent simplicity of formulation, the problem of finding exact values of $\theta_n(Q_n)$ is very difficult. Since 2006, these values are known only for n = 1, 2, 3, and 7 (see [19], [25]). Namely,

$$\theta_1(Q_1) = 1, \quad \theta_2(Q_2) = \frac{2\sqrt{5}}{5} + 1 = 1.8944..., \quad \theta_3(Q_3) = 2, \quad \theta_7(Q_7) = \frac{5}{2}.$$

The corresponding numbers $\xi(Q_n)$ are

$$\xi_1(Q_1) = 1, \quad \xi_2(Q_2) = \frac{3\sqrt{5}}{5} + 1 = 2.3416..., \quad \xi_3(Q_3) = 3, \quad \xi_7(Q_7) = 7.$$

Hence, for each n = 1, 2, 3, 7 the right-hand inequality in (2.12) becomes an equality:

$$\xi_n(Q_n) = \frac{n+1}{2} \left(\theta_n(Q_n) - 1 \right) + 1.$$
(8.1)

If a nondegenerate simplex S is contained in Q_n , then $d_i(S) \le 1$. Using (2.7) we get

$$\xi(Q_n; S) \ge \alpha(Q_n; S) = \sum_{i=1}^n \frac{1}{d_i(S)} \ge n$$

Therefore, always $\xi_n(Q_n) \ge n$. If n + 1 is an Hadamard number, then $\xi_n(Q_n) = n$ (see [24, 25]).

Thanks to (2.12), inequality $\xi_n(Q_n) \ge n$ implies

$$\theta_n(Q_n) \ge 3 - \frac{4}{n+1}.\tag{8.2}$$

If n = 1, 3, or 7, then in (8.2) we have an equality.

For $1 \le n \le 3$ the simplices corresponding to minimal projectors are just the same that the extremal simplices with respect to $\xi_n(Q_n)$. The proofs are given in [21, 25] (see also the recent survey [31]). Let us mention here these results.

The case n = 1 is trivial: $\theta_1(Q_1) = 1$ and a unique extremal simplex is the segment [0, 1] coinciding with Q_1 . Suppose n = 2. Up to rotations, the only simplex extremal with respect both to $\theta_2(Q_2)$ and $\xi_2(Q_2)$ is the triangle with vertices (0,0), $(1,\tau)$, $(\tau, 1)$, where $\tau = (3 - \sqrt{5})/2 = 0.3819...$ This number satisfies $\tau^2 - 3\tau + 1 = 0$ or

$$\frac{\tau}{1-\tau} = \frac{1-\tau}{1}.$$

Hence, τ delivers the golden section of the segment [0,1]. Sharp inequality $||P||_{Q_2} \ge 2\sqrt{5}/5 + 1$ for projectors corresponding to simplices $S \subset Q_2$ gives the new characterization of this classical notion.

Up to coordinate substitution, each extremal simplex in Q_3 coincides with the tetrahedron S' with vertices

$$(1, 1, 0), (1, 0, 1), (0, 1, 1), (0, 0, 0)$$

or the tetrahedron S'' with vertices

$$\left(\frac{1}{2}, 0, 0\right), \left(\frac{1}{2}, 1, 0\right), \left(0, \frac{1}{2}, 1\right), \left(1, \frac{1}{2}, 1\right).$$

In other words, if $||P_S||_{Q_3} = 3$, then either vertices of *S* coincide with vertices of the cube and form a regular tetrahedron or coincide with the centers of opposite edges of two opposite faces of the cube and does not belong to a common plane.

Let us turn to the case n = 7. Since 8 is an Hadamard number, there exists a seven-dimensional regular simplex having vertices at vertices of the cube. We can take the simplex with the vertices

$$(1, 1, 1, 1, 1, 1), (0, 1, 0, 1, 0, 1, 0), (0, 0, 1, 1, 0, 0, 1), (1, 0, 0, 1, 1, 0, 0),$$

$$(0, 0, 0, 0, 1, 1, 1), (1, 0, 1, 0, 0, 1, 0), (1, 1, 0, 0, 0, 0, 1), (0, 1, 1, 0, 1, 0, 0).$$

The equality $\xi_7(Q_7) = 7$ and inequality

$$\xi_n(Q_n) \le \frac{n+1}{2} (\theta_n(Q_n) - 1) + 1$$

imply $\theta_7(Q_7) \ge 5/2$. But for the corresponding projector, $||P||_{Q_7} = 5/2$. Therefore, $\theta_7 = 5/2$, and this projector is minimal.

Let n+1 be an Hadamard number, and let *S* be an *n*-dimensional regular simplex having the vertices at vertices of Q_n Then, for the corresponding projector $P_S : C(Q_n) \to \Pi_1(\mathbb{R}^n)$, we have

$$\|P_S\|_{Q_n} \le \sqrt{n+1}.$$
 (8.3)

Various proofs of (8.3) are given in [18] and [25]. Paper [29] contains the proof essentially making use of the structure of an Hadamard matrix. The equality $||P_S||_{Q_n} = \sqrt{n+1}$ may hold as for all regular simplices *S* having vertices at vertices of the cube (n = 1, n = 3), as for some of them (n = 15), or may not be executed at all.

As it is shown in Section 5, for each n, $\theta_n(Q_n) \ge \sqrt{n-1}/e$. Therefore, if n + 1 is an Hadamard number, then

$$\frac{\sqrt{n-1}}{e} \le \theta_n(Q_n) \le \sqrt{n+1}.$$

In other words, the above estimate $\theta_n(Q_n) \ge c \sqrt{n}$ is sharp at least when n + 1 is an Hadamard number. For these dimensions, $\theta_n(Q_n) \asymp \sqrt{n}$.

The upper bounds of $\theta_n(Q_n)$ for special *n* were improved by A. Ukhalov and his students with the help of computer methods. In particular, simplices of maximum volume in the cube were considered. In all situations where n + 1 is an Hadamard number, the full set of Hadamard matrices of the corresponding order was used. In particular, to obtain an estimate for θ_{23} , all existing 60 Hadamard matrices of order 24 were considered. To estimate θ_{27} , we have to consider 487 Hadamard matrices of order 28. Known nowaday upper estimates for $1 \le n \le 27$ are given in [37]. Here they are (for brevity, we write $\theta_n = \theta_n(Q_n)$):

$$\begin{aligned} \theta_{1} &= 1, \quad \theta_{2} = \frac{2\sqrt{5}}{5} + 1, \quad \theta_{3} = 2, \quad \theta_{4} \leq \frac{3(4 + \sqrt{2})}{7}, \quad \theta_{5} \leq 2.448804, \\ \theta_{6} \leq 2.6000 \dots, \quad \theta_{7} = \frac{5}{2}, \quad \theta_{8} \leq \frac{22}{7}, \quad \theta_{9} \leq 3, \quad \theta_{10} \leq \frac{19}{5}, \quad \theta_{11} \leq 3, \\ \theta_{12} \leq \frac{17}{5}, \quad \theta_{13} \leq \frac{49}{13}, \quad \theta_{14} \leq \frac{21}{5}, \quad \theta_{15} \leq \frac{7}{2}, \quad \theta_{16} \leq \frac{21}{5}, \quad \theta_{17} \leq \frac{139}{34}, \\ \theta_{18} \leq 5.1400 \dots, \quad \theta_{19} \leq 4, \quad \theta_{20} \leq 4.68879 \dots, \quad \theta_{21} \leq \frac{251}{50}, \quad \theta_{22} \leq \frac{1817}{335}, \\ \theta_{23} \leq \frac{9}{2}, \quad \theta_{24} \leq \frac{103}{21}, \quad \theta_{25} \leq 5, \quad \theta_{26} \leq \frac{474}{91}, \quad \theta_{27} \leq 5. \end{aligned}$$

The best nowaday known lower bound of $\theta_n(Q_n)$ has the form

$$\theta_n(Q_n) \ge \max\left[3 - \frac{4}{n+1}, \chi_n^{-1}\left(\frac{1}{\nu_n}\right)\right].$$
(8.4)

Here χ_n is the standardized Legendre polynomial of degree *n*, see Section 3. The values of the righthand side of (8.4) for $1 \le n \le 54$ are given in [37].

As noted in Section 1 (see (2.11)), the inequality

$$\xi_n(Q_n) \le \frac{n+1}{2} \left(\theta_n(Q_n) - 1 \right) + 1 \tag{8.5}$$

is true for any *n*. So far, we know only four values of *n* for which this relation becomes an equality: n = 1, 2, 3, and 7. These are exactly the cases in which we know the exact values both of $\theta_n(Q_n)$ and $\xi_n(Q_n)$. In [32] the authors conjectured that the minimum of *n* for which inequality (8.5) is strict is 4. This is still an open problem.

The above given estimate $\xi_n(Q_n) \ge n$ occurs to be exact in order of *n*. If n > 2, then

$$\xi_n(Q_n) \le \frac{n^2 - 3}{n - 1} \tag{8.6}$$

(see [23,25]). If n > 1, the right-hand side of (8.6) is strictly smaller than n + 1. Inequality $\xi_n < n + 1$ holds true also for n = 1, 2. Thus, always $n \le \xi_n(Q_n) < n + 1$, i.e., $\xi_n(Q_n) - n \in [0, 1)$. However, the exact values of the constant $\xi_n(Q_n)$ are currently only known for n = 2, n = 5 and n = 9, as well as for an infinite set of n for which there exists an Hadamard matrix of order n + 1.

In all these cases, except n = 2, the equality $\xi_n(Q_n) = n$ holds. In the noted Hadamard's case, one can give the proof using the structure of Hadamard matrix of order n + 1, see [33]. In [33], we have

also discovered the exact values of $\xi_n(Q_n)$ for n = 5 and n = 9 and constructed several infinite families of extremal simplices for n = 5, 7, 9.

Thanks to the equivalence $\xi_n(Q_n) \asymp n$ and inequality $\theta_n \ge c \sqrt{n}$, for all sufficiently large *n*, we have

$$\xi_n(Q_n) < \frac{n+1}{2} \left(\theta_n(Q_n) - 1 \right) + 1.$$
(8.7)

Let n_0 be the minimal natural number such that for all $n \ge n_0$ inequality (8.7) holds. The problem about the exact value of n_0 is very difficult. The known lower and upper bounds differ quite significantly. From the preceding, we have the estimate $n_0 \ge 8$. In 2009 we proved that $n_0 \le 57$ (see [21,25]). A sufficient condition for the validity of (8.7) for n > 2 is the inequality

$$\chi_n\left(\frac{3n-5}{n-1}\right) \cdot \nu_n < 1. \tag{8.8}$$

It was proved in [21] that (8.8) is satisfied for $n \le 57$. Later calculations allowed the upper bound of n_0 to be slightly lowered. In [34] it is noted that $n_0 \le 53$. In other words, inequality (8.7) is satisfied at least starting from n = 53. A better estimate from above for n_0 is an open problem.

Let us now proceed to interpolation on the unit ball B_n .

It was shown in [26] that $\xi_n(B_n) = n$ for all *n*. Moreover, for a simplex $S \subset B_n$, the equality $\xi(B_n; S) = n$ is equivalent to the fact that *S* is a regular simplex inscribed in the ball. By (2.11), for any projector $P : C(B_n) \to \prod_1 (\mathbb{R}^n)$ we have

$$||P||_{B_n} \ge 3 - \frac{4}{n+1}.$$
(8.9)

The right-hand equality in (2.12), i. e., the equality

$$\xi_n(B_n) = \frac{n+1}{2} \Big(\theta_n(B_n) - 1 \Big) + 1, \tag{8.10}$$

is equivalent to

$$\theta_n(B_n) = 3 - \frac{4}{n+1}.$$
(8.11)

As it shown in [35], equalities (8.10) - (8.11) take place for $1 \le n \le 4$, while starting from n = 5 we have the strict inequality

$$\xi_n(B_n) < \frac{n+1}{2} \Big(\theta_n(B_n) - 1 \Big) + 1.$$

For $1 \le n \le 4$, equality in (8.9) holds only if S_P is a regular inscribed simplex.

The fact that the lower estimate $\theta_n(B_n) > c \sqrt{n}$ obtained in Section 5 is exact in dimension *n* was first established in [35]: *the equivalence* $\theta_n(B_n) \approx \sqrt{n}$ *is valid.*

A complete solution to the problem about the values of the numbers $\theta_n(B_n)$ was given in [30]. Let us briefly describe these results. Let $\psi : [0, n + 1] \rightarrow \mathbb{R}$ be a function defined by

$$\psi(t) = \frac{2\sqrt{n}}{n+1} \left(t(n+1-t) \right)^{1/2} + \left| 1 - \frac{2t}{n+1} \right|$$

and let

$$a = a_n = \left\lfloor \frac{n+1}{2} - \frac{\sqrt{n+1}}{2} \right\rfloor.$$

Let S be a regular simplex inscribed in B_n , and let p_n be the norm of P_S . It was proved in [35] that

$$p_n = \max\{\psi(a), \psi(a+1)\}$$
 and $\sqrt{n} \le p_n \le \sqrt{n+1}$.

Moreover, $p_n = \sqrt{n}$ only for n = 1 and $p_n = \sqrt{n+1}$ holds if and only if $\sqrt{n+1}$ is an integer.

The equality $\theta_n(B_n) = p_n$ was obtained first for $1 \le n \le 4$ (different proofs are given in [35] and [28]). As a conjecture for all *n*, this statement was formulated in [28]. In [30] we developed a certain new geometric approach to the problem which enabled us to prove this conjecture, i.e., the equality $\theta_n(B_n) = p_n$ for an arbitrary positive integer *n*.

Thus, the following ineuality $\sqrt{n} \le \theta_n(B_n) \le \sqrt{n+1}$ holds for all *n*. Furthermore, we proved that the minimal is any projector corresponding to a regular simplex inscribed into the boundary sphere, and there are no other minimal projections in this case.

Let k_n coincide with that of the numbers a_n and $a_n + 1$ on which $\psi(t)$ takes a larger value. The numbers k_n increase with n, but not strictly monotonically. If $n \ge 2$, then $k_n \le n/2$. As an example, we give below the numbers k_n for $1 \le n \le 15$, n = 50, n = 100, and n = 1000 ([35]):

$$k_1 = k_2 = k_3 = k_4 = 1$$
, $k_5 = k_6 = 2$, $k_7 = k_8 = k_9 = 3$, $k_{10} = k_{11} = 4$,
 $k_{12} = k_{13} = 5$, $k_{14} = k_{15} = 6$, $k_{50} = 22$, $k_{100} = 45$, $k_{1000} = 485$.

Finally, let us note that equality (8.10) holds for those and only those dimensions n when $k_n = 1$.

Acknowledgements

I am very thankful to Pavel Shvartsman for useful suggestions and remarks.

References

- V. BARTHELMANN, E. NOVAK, AND K. RITTER. High dimensional polynomial interpolation on sparse grids, Adv. Comput. Math., 12:4 (2000), 273–288.
- [2] H. BAUER. Minimalstellen von Funktionen und Extremal punkte, Archiv der Mathematik, 9:4 (1958), 389–393.
- [3] C. DE BOOR. *Polynomial Interpolation in Several Variables*, In: R. de Millo and J. R. Rice (editors), Studies in Computer Science, 87–119, Plenum Press, 1994.
- [4] G. F. CLEMENTS AND B. LINDSTRÖM. A sequence of (±1) determinants with large values, Proc. Amer. Math. Soc., 16 (1965), 548–550.
- [5] S. DE MARCHI. Lectures on Multivariate Polynomial Interpolation, Göttingen Padova, 2015.
- [6] R. A. DEVORE AND G. G. LORENTZ. *Constructive Approximation*, Springer-Verlag: Berlin-Heidelberg, 1993.
- [7] L. FEJES TÓT. Regular Figures, New York: Macmillan/Pergamon, 1964.
- [8] G. M. FIKHTENGOL'TS. *The Course in Differential and Integral Calculation. Vol.3*, Fizmatlit, Moscow, 2001 (in Russian).

- [9] M. GASCA AND T. SAUER. Polynomial interpolation in several variables, *Adv. Comput. Math.*, **12**:4 (2000), 377–410.
- [10] M. GUNZBURGER AND A. TECKENTRUP. Optimal point sets for total degree polynomial interpolation in moderate dimensions, arXiv:1407.3291v1 [math.NA] 11 Jul 2014.
- [11] K. J. HORADAM. Hadamard Matrices and Their Applications, Princeton University Press, 2007.
- [12] J. HADAMARD. Résolution d'une question relative aux déterminants, *Bull. Sciences Math.* (2), 17 (1893), 240–246.
- [13] M. HALL, JR. *Combinatorial Theory*, Blaisdall Publishing Company, Waltham (Massachusetts)-Toronto-London, 1967.
- [14] M. HUDELSON, V. KLEE, AND D. LARMAN. Largest *j*-simplices in *d*-cubes: some relatives of the Hadamard maximum determinant problem, *Linear Algebra Appl.*, **241–243** (2019), 519–598.
- [15] V. KAARNIOJA. Multivariate polynomial interpolation over arbitrary grids, arXiv:1512.07424v1 [math.NA] 23 Dec 2015.
- [16] M. LASSAK. Approximation of convex bodies by inscribed simplices of maximum volume, *Discr. Comput. Geom.*, 21 (1999), 449–462.
- [17] P. K. MANJHI AND M. K. RAMA. Some new examples of circulant partial Hadamard matrices of type $4 H(k \times n)$, Adv. Appl. Math. Sci., **21**:5 (2022), 2559–2564.
- [18] M. V. NEVSKII. Estimates for the minimum norm of a projection in linear interpolation over the vertices of an *n*-dimensional cube, *Model. Anal. Inform. Sist.*, **10**:1 (2003), 9–19 (in Russian).
- [19] M. V. NEVSKII. Geometric methods in the minimal projection problem, *Model. Anal. Inform. Sist.*, **13**:2 (2006), 16–29 (in Russian).
- [20] M. V. NEVSKII. Inequalities for the norms of interpolating projections, *Model. Anal. Inform. Sist.*, **15**:3 (2008), 28–37 (in Russian).
- [21] M. V. NEVSKII. On a certain relation for the minimal norm of an interpolational projection, *Model. Anal. Inform. Sist.*, **16**:1 (2009), 24–43 (in Russian).
- [22] M. V. Nevskii, On a property of *n*-dimensional simplices, *Mat. Zametki*, 87:4 (2010), 580–593 (in Russian). English translation: *Math. Notes*, 87:4 (2010), 543–555.
- [23] M. V. NEVSKII. On geometric characteristics of an *n*-dimensional simplex, *Model. Anal. Inform. Sist.*, **18**:2 (2011), 52–64 (in Russian).
- [24] M. NEVSKII. Properties of axial diameters of a simplex, *Discr. Comput. Geom.*, **46**:2 (2011), 301–312.
- [25] M. V. NEVSKII. *Geometric Estimates in Polynomial Interpolation*, P. G. Demidov Yaroslavl' State University, Yaroslavl', 2012 (in Russian).

- [26] M. V. NEVSKII. On some problems for a simplex and a ball in \mathbb{R}^n , *Model. Anal. Inform. Sist.*, **25**:6 (2018), 680–691 (in Russian). English translation: *Aut. Control Comp. Sci.*, **53**:7 (2019), 644–652.
- [27] M. V. NEVSKII. Geometric estimates in interpolation on an n-dimensional ball, *Model. Anal. Inform. Sist.*, 26:3 (2019), 441–449 (in Russian). English translation: *Aut. Control Comp. Sci.*, 54:7 (2020), 712–718.
- [28] M. V. NEVSKII. On properties of a regular simplex inscribed into a ball, *Model. Anal. Inform. Sist.*, 28:2 (2021), 186–197 (in Russian). English translation: *Aut. Control Comp. Sci.*, 56:7 (2022), 778–787.
- [29] M. V. NEVSKIJ. On some estimate for the norm of an interpolation projector, *Model. Anal. Inform. Sist.*, 29:2 (2022), 92–103 (in Russian). English translation: *Aut. Control Comp. Sci.*, 57:7 (2023), 718–726.
- [30] M. V. NEVSKII. On the minimal norm of the projection operator for linear interpolation on an *n*-dimensional ball, *Mat. Zametki*, **114**:3 (2023), 477–480 (in Russian). English translation: *Math. Notes*, **114**:3 (2023), 415–418.
- [31] M. NEVSKII. Geometric estimates in linear interpolation on a cube and a ball, arXiv:2402.11611v1 [math.CA] 18 Feb 2024.
- [32] M. V. NEVSKII AND A. YU. UKHALOV. On numerical characteristics of a simplex and their estimates, *Model. Anal. Inform. Sist.*, 23:5 (2016), 602–618 (in Russian). English translation: *Aut. Control Comp. Sci.*, 51:7 (2017), 757–769.
- [33] M. NEVSKII AND A. UKHALOV. Perfect Simplices in \mathbb{R}^5 , *Beitr. Algebra Geom.*, **59**:3 (2018), 501–521.
- [34] M. V. NEVSKII AND A. YU. UKHALOV. On optimal interpolation by linear functions on an *n*dimensional cube, *Model. Anal. Inform. Sist.*, 25:3 (2018), 291–311 (in Russian). English translation: *Aut. Control Comp. Sci.*, 52:7 (2018), 828–842.
- [35] M. V. NEVSKII AND A. YU. UKHALOV. Linear interpolation on a Euclidean ball in ℝⁿ, Model. Anal. Inform. Sist., 26:2 (2019), 279–296 (in Russian). English translation: Aut. Control Comp. Sci., 54:7 (2020), 601–614.
- [36] M. V. NEVSKII AND A. YU. UKHALOV. Selected Problems in Analysis and Computational Geometry. Part 1, P. G. Demidov Yaroslavl' State University, Yaroslavl', 2020 (in Russian).
- [37] M. V. NEVSKII AND A. YU. UKHALOV. Selected Problems in Analysis and Computational Geometry. Part 2, P. G. Demidov Yaroslavl' State University, Yaroslavl', 2022 (in Russian).
- [38] S. PASHKOVSKIJ. Vychislitel'nye Primeneniya Mnogochlenov i Ryadov Chebysheva, Nauka, Moskva, 1983 (in Russian).
- [39] A. P. PRUDNIKOV, YU. A. BRYCHKOV, AND O. I. MARICHEV. *Integraly i Ryady*, Nauka, Moskva, 1981 (in Russian).
- [40] T. J. RIVLIN. *The Chebyshev Polynomials*, John Wiley & Sons, New York-Chichester-Brisbane-Toronto, 1974.

- [41] P.R. Scott. Lattices and convex sets in space, *Quart. J. Math. Oxford Ser.* (2), **36** (1985), 359–362.
- [42] P.R. Scott. Properties of axial diameters, Bull. Austral. Math. Soc., 39 (1989), 329-333.
- [43] D. SLEPIAN. The content of some extreme simplices, *Pacific J. Math.*, **31** (1969), 795–808.
- [44] P. K. SUETIN. *Klassicheskie ortogonal'nye mnogochleny*, Nauka, Moskva, 1979 (in Russian).
- [45] G. Szegö. *Orthogonal Polynomials*, Providence, American Mathematical Society, New York, 1975.
- [46] D. VANDEV. A minimal volume ellipsoid around a simplex, *C. R. Acad. Bulg. Sci.*, **45**:6 (1992), 37–40.