

Endpoints of smooth plane dendroids

In memory of professors Gary Gruenhage and Piotr Minc

David S. Lipham

ABSTRACT. Let X be a smooth dendroid in the plane. We prove that the endpoints of X are arcwise accessible from $\mathbb{R}^2 \setminus X$, and that the space of endpoints $E(X)$ has the property of a circle. Also if $E(X)$ is totally disconnected and 1-dimensional, then X must contain a Cantor set of arcs. An example is constructed to show that this is false outside the plane.

1. Introduction

Dendroids form an important class of uniquely arcwise connected continua. They can be defined as hereditarily arcwise connected continua without simple closed curves. Every contractible 1-dimensional continuum is a dendroid [3], and all dendroids are tree-like [5]. Plane dendroids are non-separating and have the fixed point property [15, 2]. For a nice survey with open problems, see [16].

The focus of this paper is on smooth plane dendroids; equivalently, plane dendroids which admit radially convex metrics [4]. We will show that each endpoint of a smooth plane dendroid X is accessible from $\mathbb{R}^2 \setminus X$, with at most one exception (§3). From here we find that the space of endpoints $E(X)$ is circle-like, in that every two of its points are separated by two other points (§4). Bellamy constructed in [1] a smooth plane dendroid with connected endpoint set which he observed to have this property.

We also examine the dimension of endpoints. It is well-known that the endpoints of a smooth plane dendroid can be 1-dimensional even when they are totally disconnected. The endpoints of the Lelek fan have this property [11]. Apparently, the Lelek fan has an uncountable collection of pairwise-disjoint arcs, meaning that it is non-Suslinian. Using the result of §4, we will show that if X is any smooth plane dendroid, and $E(X)$ is totally disconnected and 1-dimensional, then X must be non-Suslinian (§5). Here planarity is critical: We construct in §6 a (non-planar) Suslinian smooth dendroid D such that $E(D)$ is homeomorphic to the endpoints of the Lelek fan.

Acknowledgements. The example presented in §6 is attributed to Ed Tymchatyn and Piotr Minc, although it was not published until now. I am grateful to Ed Tymchatyn for sharing with me the ideas for its construction.

I also wish to thank Logan Hoehn for suggesting the use of colocal connectedness to prove accessibility of endpoints.

2020 *Mathematics Subject Classification.* 54F15, 54F45, 54F50.

Key words and phrases. dendroid, smooth, endpoint, totally disconnected, zero-dimensional, Suslinian, plane, accessible.

2. Basic notions

A **continuum** is a compact connected metric space.

A continuum is **hereditarily unicoherent** if every two of its subcontinua have connected intersection. A **dendroid** is a hereditarily unicoherent, arcwise connected continuum. If X is a dendroid and $x, y \in X$ then we let $\alpha(x, y)$ denote the unique arc in X with endpoints x and y .

A dendroid X is **smooth** if there exists $p \in X$ such that if $x_n \rightarrow x$ in X , then $\alpha(x_n, p) \rightarrow \alpha(x, p)$ in the Hausdorff distance. The point p is called an **initial point** of X ; alternatively we say that X is **smooth at p** . If X is a dendroid smooth at p , then there exists an equivalent metric d on X such that $d(x, p) = d(x, y) + d(y, p)$ whenever $y \in \alpha(x, p)$; see [4, 9]. Such a metric is called **radially convex** with respect to p .

A point $e \in X$ is an **endpoint** if e is an endpoint of every arc in X that contains it. We let $E(X)$ denote the set of all endpoints of X . Note that by [19, Lemma 3], each arc in X is contained in a maximal arc in X , each of whose endpoints must belong to $E(X)$. The endpoint set of a smooth dendroid is always G_δ [20].

A **fan** is a dendroid X with a point p such that each component of $X \setminus \{p\}$ is homeomorphic to $(0, 1]$. The **Cantor fan** is the quotient of $C \times [0, 1]$ that takes $C \times \{0\}$ to a point. The **Lelek fan** is a smooth fan with dense set of endpoints [11].

A continuum is **Suslinian** if X contains no uncountable collection of non-degenerate, pairwise-disjoint subcontinua [12]. The following helps to understand the Suslinian structure of a smooth dendroid.

PROPOSITION 1. *Let X be a smooth dendroid. The following are equivalent.*

- (1) X is Suslinian;
- (2) $X \setminus E(X)$ is the union of an increasing sequence of dendroids $X_1 \subset X_2 \subset \dots$ such that $E(X_n)$ is countable;
- (3) $X \setminus E(X)$ is a countable union of arcs.

PROOF. (1) \Rightarrow (2): Suppose that X is Suslinian. Let d be a radially convex metric on X . Define $f_n : X \rightarrow X$ by $f_n(x)$ is the unique point y of $\alpha(x, p)$ such that

$$d(y, p) = (1 - \frac{1}{n}) \cdot d(x, p).$$

By continuity of f_n , the image $X_n = f_n[X]$ is a dendroid. If $e \in E(X_n)$, then there is an arc $\alpha_e \subset X$ such that $\alpha_e \cap X_n = \{e\}$. Note that if $e \neq e'$ then $\alpha_e \cap \alpha_{e'} = \emptyset$ or $\alpha_e \cap \alpha_{e'} = \{e\}$. Since X is Suslinian this means that $E(X_n)$ must be countable.

(2) \Rightarrow (3): It suffices to show that each X_n is a countable union of arcs. Let n be given. Assuming that $E(X_n)$ is countable, $\mathcal{M} = \{\alpha(e, e') : e, e' \in E(X_n)\}$ is also countable. Given any $x \in X_n$, there is a maximal arc $M \subset X_n$ containing x . Then $x \in M \in \mathcal{M}$. Therefore $X_n = \bigcup \mathcal{M}$.

(3) \Rightarrow (1): Suppose that $X \setminus E(X) = \bigcup_{n=1}^{\infty} \alpha_n$ is a countable union of arcs. Let Q_n be a countable dense subset of α_n . We claim that the countable set $Q = \bigcup_{n=1}^{\infty} Q_n$ intersects each subcontinuum of X . Clearly this will imply that X is Suslinian. Let K be any subcontinuum of X . Then K contains an arc $\beta \subset X \setminus E(X)$. By Baire's theorem, there exists n such that α_n contains an arc of β . Then $Q_n \cap \beta \neq \emptyset$. So $Q \cap K \neq \emptyset$. \square

3. Accessibility of endpoints

If U is an open subset of \mathbb{R}^2 then a point $z \in \mathbb{R}^2 \setminus U$ is **accessible from U** if there is an arc $\alpha \subset \mathbb{R}^2$ such that $\alpha \setminus U = \{z\}$. When a continuum X is understood from context, we will say that $x \in X$ is **accessible** if x is accessible from $\mathbb{R}^2 \setminus X$.

A continuum X is **colocally connected** at $x \in X$ if for every open set V containing x there is an open $U \subset V$ such that $x \in U$ and $X \setminus U$ is connected.

A **simply connected domain** is a bounded open subset of the plane which is connected and simply connected (and therefore homeomorphic to \mathbb{R}^2).

LEMMA 2. *Let X be a dendroid in the plane. Let U be an open set in the plane such that $X \setminus U$ is connected. If W is any connected component of U , then $X \setminus W$ is connected.*

PROOF. Let $x, y \in X \setminus W$. Let $f_x : [0, 1] \hookrightarrow X$ such that $f_x(0) = x$ and $f_x(1) \in W$. Let $a = \inf\{t \in [0, 1] : f_x(t) \notin U\}$. Define f_y and b similarly for y . Let α be an arc in $X \setminus U$ from $f_x(a)$ to $f_y(b)$. Then $\beta = f_x[0, a] \cup \alpha \cup f_y[0, b]$ is an arc in $X \setminus W$ from x to y . This shows that $X \setminus W$ is connected. \square

LEMMA 3. *Let X be a dendroid in the plane. Let $S \subset \mathbb{R}^2$ be a circle with complementary components U bounded and V unbounded, such that X meets V . Suppose that W is a connected open subset of U such that $X \setminus W$ is connected. Then there is a simply connected domain $W' \subset U$ such that $W \subset W'$ and $X \setminus W'$ is connected.*

PROOF. Let W' be the union of W with all of its bounded complementary components. Clearly $W \subset W' \subset U$. By [18, Theorem 5.4], each complementary component of W meets ∂W . So W' is connected. Also W' is open and has connected complement ($\mathbb{R}^2 \setminus W'$ is just the unbounded component of $\mathbb{R}^2 \setminus W$). Therefore W' is simply connected. Note that $X \setminus W$ lies wholly in the unbounded component of $\mathbb{R}^2 \setminus W$ because $X \setminus W$ is connected and meets V . Thus $X \setminus W' = X \setminus W$ is connected. \square

LEMMA 4. *Let X be a dendroid in the plane. Let $W \subset \mathbb{R}^2$ be simply connected domain. If $X \setminus W$ is connected, then $W \setminus X$ is connected.*

PROOF. Suppose that $X \setminus W$ is connected but $W \setminus X$ is not. Then $X/(X \setminus W)$ is a dendroid that separates the 2-sphere $\mathbb{R}^2/(\mathbb{R}^2 \setminus W)$, a contradiction. \square

THEOREM 5. *Let X be a plane dendroid. If X is colocally connected at x , then x is accessible.*

PROOF. Let $\delta = \text{diam}(X)$ and for each $n = 1, 2, 3, \dots$ let S_n be the circle of radius $\delta/2n$ centered at e . Note that X meets the unbounded component of each $\mathbb{R}^2 \setminus S_n$. Let U_n be an open set in the bounded component of $\mathbb{R}^2 \setminus S_n$ such that $e \in U_n$ and $X \setminus U_n$ is connected. Let W_n be the connected component of e in U_n . By Lemma 2, $X \setminus W_n$ is connected. By Lemma 3 there is a simply connected domain $W'_n \subset U_n$ such that $e \in W'_n$ and $X \setminus W'_n$ is connected. By Lemma 4, $W'_n \setminus X$ is (path-)connected.

Now let $y_0 \in \mathbb{R}^2 \setminus X$. Let $\alpha_0 \subset \mathbb{R}^2 \setminus X$ be an arc from y_0 to some $y_1 \in W'_1 \setminus X$. Let $y_2 \in W'_2 \cap W'_1 \setminus X$. Let $\alpha_1 \subset W'_1 \setminus X$ be an arc from y_1 to y_2 . Continue this process, letting α_n be an arc in $W'_n \setminus X$ from y_n to a new point $y_{n+1} \in W'_n \cap W'_{n+1} \setminus X$. The arcs α_n form a null sequence, and their only limit point is e . Hence the (locally connected) continuum $\overline{\bigcup_{n=0}^{\infty} \alpha_n}$ contains an arc α from y_0 to e , with $\alpha \cap X = \{e\}$. \square

COROLLARY 6. *Let X be a smooth plane dendroid with initial point p . Then every endpoint $e \in E(X) \setminus \{p\}$ is accessible.*

PROOF. By [10, Theorems 3.1 and 3.5], X is colocally connected at e . By Theorem 5, e is accessible. \square

REMARK 1. The point p could be an inaccessible endpoint. See Figure 1.

REMARK 2. In Bellamy's dendroid [1], the endpoints are the only accessible points.




FIGURE 1. A smooth plane dendroid in which p is an inaccessible endpoint.

4. Separation of endpoints

We show that the space of endpoints of a smooth plane dendroid has the circle-like property described in §1.

The following is based on the θ -curve theorem.

PROPOSITION 7. *Let O be a simple closed curve in the plane. Let $a, b, c, d \in O$ in cyclic order. Let U and V be the components of $\mathbb{R}^2 \setminus O$. Let α and β be arcs in $\overline{U} \setminus \{b, d\}$ and $\overline{V} \setminus \{b, d\}$, respectively, from a to c . Then $\alpha \cup \beta$ separates b and d .*

THEOREM 8. *Let X be a plane dendroid, smooth at p . Let $e \in E(X) \setminus \{p\}$. For every $x \in X \setminus \alpha(e, p)$ there is a simple closed curve $\sigma \subset \mathbb{R}^2$ which separates e from x and has the property $|\sigma \cap E(X)| \leq 2$.*

PROOF. Let $e' \neq p$ be an endpoint of a maximal arc in X extending $\alpha(x, p)$. Let γ be an arc from e to e' such that $\gamma \cap X = \{e, e'\}$, as provided by Corollary 6 and the fact that $\mathbb{R}^2 \setminus X$ is path-connected. Let $q \in D$ such that $\alpha(q, p) = \alpha(e, p) \cap \alpha(x, p)$. The simple closed curve $\gamma \cup \alpha(e, q) \cup \alpha(e', q)$ separates the plane into two components U and V . Assume $p \in \overline{U}$. Let $b \in \gamma$.

Claim 1: *There is an arc α in \overline{U} from p to b which misses e and x , and contains at most one endpoint.* We may assume that there exists a sequence of points $x_n \in U \cap X \setminus \alpha(p, q)$ such that $x_n \rightarrow p$. Otherwise p is accessible from $U \setminus X$ and the claim becomes trivial. Since X is smooth at p we know $\alpha(x_n, p) \rightarrow p$ and so eventually $\alpha(x_n, p)$ misses e and x . Let α_1 be a maximal arc in X containing $\alpha(x_n, p)$. Then α_1 has an endpoint $e_1 \in U \setminus \{p\}$. By Corollary 6, there is an arc α_2 from e_1 to b such that $\alpha_2 \cap X = \{e\}$. Put $\alpha = \alpha(e_1, p) \cup \alpha_2$.

Claim 2: *There is an arc β in \overline{V} from q to b which misses e and x , and contains at most one endpoint.* The proof is similar to that of Claim 1. Note that $X \cap \overline{V}$ is a dendroid smooth at q . So in the non-trivial case that there exists $x_n \in V \cap X$ such that $x_n \rightarrow q$, eventually $\alpha(x_n, q)$ misses e and x . The arc $\alpha(x_n, q)$ extends to an arc β_1 with an endpoint $e_2 \in V \setminus \{p\}$. By Corollary 6, there is an arc β_2 from b to e_2 such that $\beta_2 \cap X = \{e_2\}$. Put $\beta = \alpha(e_2, q) \cup \beta_2$.

By Proposition 7, $\sigma = \alpha \cup \beta \cup \alpha(p, q)$ separates e and x . See Figure 2. \square

COROLLARY 9. *Let X be a smooth dendroid in the plane with initial point p . Within the space $E(X) \setminus \{p\}$, every two points are separated by two other points.*

5. Dimension of endpoints

A space X is **totally disconnected** if for every two points of X are contained in disjoint clopen sets. A space X is **zero-dimensional** if X has a basis of clopen sets. A space X is **almost zero-dimensional** if X has a basis of neighborhoods which are intersections of clopen sets [7]. Observe that:




FIGURE 2. Proof of Theorem 8.

zero-dimensional \Rightarrow almost zero-dimensional \Rightarrow totally disconnected.

A function $\varphi : Z \rightarrow [0, \infty)$ is **upper semi-continuous (USC)** if $\varphi^{-1}[0, t)$ is open for every $t > 0$. If Z is first countable, then this is equivalent to saying that if $z_n \rightarrow z$ and $\varphi(z_n) \geq \varphi(z)$ for all n , then $\varphi(z_n) \rightarrow \varphi(z)$.

PROPOSITION 10 ([7, Theorem 4.15]). *A space X is almost zero-dimensional if and only if X is homeomorphic to the graph of a USC function with zero-dimensional domain.*

LEMMA 11. *Let X be a smooth dendroid in the plane and $e \in E(X) \setminus \{p\}$. Let \mathcal{U}_e be the collection of all open subsets U of X such that $e \in U$ and $\partial U \cap E(X) = \emptyset$. Put*

$$F(e) = \bigcap_{U \in \mathcal{U}_e} \overline{U}.$$

If $E(X)$ is totally disconnected, then $F(e) \subset \alpha(e, p)$.

PROOF. Suppose $x \in X \setminus \alpha(e, p)$. By Theorem 8 there is an open set $U \ni e$ such that $x \notin \overline{U}$ and $F = \partial U \cap E(X)$ has at most two points. Since $E(X)$ is totally disconnected, there is an open set $V \ni e$ such that $\partial V \cap E(X) = \emptyset$ and $V \cap F = \emptyset$. Then $U \cap V \in \mathcal{U}_e$ and $x \notin \overline{U \cap V}$. Therefore $x \notin F(e)$. \square

THEOREM 12. *Let X be a smooth plane dendroid with initial point p . If $E(X)$ is totally disconnected, then $E(X) \setminus \{p\}$ is almost zero-dimensional.*

PROOF. Suppose that $Y = E(X) \setminus \{p\}$ is totally disconnected. For each $e \in Y$ there exists $U_1(e), U_2(e), \dots \in \mathcal{U}_e$ such that

$$F(e) = \bigcap_{n=1}^{\infty} \overline{U_n(e)}.$$

The collection $\{U_n(e) : e \in Y, n \geq 1\}$ is a basis for a first countable, zero-dimensional topology \mathcal{W} on Y . Let $Z = (Y, \mathcal{W})$. Let d be a radially convex metric on X , and define $\varphi : Z \rightarrow [0, \infty)$ by

$$\varphi(x) = d(x, p).$$

Claim 1: φ is USC. Suppose $e_n \rightarrow e$ in Z , and $\varphi(e_n) \geq \varphi(e)$ for each n . Let x be any accumulation point of (e_n) in X . Then $x \in F(e)$. By Lemma 11, $x \in \alpha(e, p)$. And $d(e_n, p) \geq d(e, p)$ implies $d(x, p) \geq d(e, p)$ by continuity of d . Since d is radially convex $d(e, p) = d(e, x) + d(x, p)$. So $d(e, x) = 0$ and $x = e$. Therefore $e_n \rightarrow e$ in Y .

Claim 2: Y is homeomorphic to the graph of φ . Consider the graph

$$\text{Gr}(\varphi) = \{\langle e, \varphi(e) \rangle : e \in Y\}$$

as a subspace of $Z \times [0, \infty)$. Apparently, $e \mapsto \langle e, \varphi(e) \rangle$ defines a continuous one-to-one mapping of Y onto $\text{Gr}(\varphi)$. We show that its inverse is continuous. To that end, suppose that $e_n \rightarrow e$ in Z and $\varphi(e_n) \rightarrow \varphi(e)$. If x is any accumulation point of (e_n) in X , then $x \in \alpha(e, p)$ by Lemma 11. Also $\varphi(x) = \varphi(e)$. So $x = e$. Therefore $e_n \rightarrow e$ in Y .

By Proposition 10, Y is almost zero-dimensional. \square

A space X is **zero-dimensional at** $x \in X$ if the point x has a neighborhood basis of clopen sets. Put

$$\Omega(X) = \{x \in X : X \text{ is zero-dimensional at } x\}$$

and $\Lambda(X) = X \setminus \Omega(X)$.

PROPOSITION 13 ([14, Theorem A]). *Let X be a Suslinian plane continuum. If Y is any totally disconnected Borel set in X , then $\Lambda(Y)$ is countable.*

PROPOSITION 14 ([13, Theorem 1]). *Let Y be an almost zero-dimensional space. If $\Lambda(Y)$ is countable, then Y is zero-dimensional.*

THEOREM 15. *Let X be a Suslinian smooth plane dendroid. If $E(X)$ is totally disconnected, then $E(X)$ is zero-dimensional.*

PROOF. Suppose that $E(X)$ is totally disconnected. Let p be an initial point of X . By Theorem 12, $Y = E(X) \setminus \{p\}$ is almost zero-dimensional. By Proposition 13, $\Lambda(Y)$ is countable. By Proposition 14, Y is zero-dimensional. By [8, Corollary 1.3.5], $E(X)$ is zero-dimensional. \square

COROLLARY 16. *Let X be a smooth plane dendroid. If $E(X)$ is totally disconnected and 1-dimensional, then X contains an uncountable collection of pairwise-disjoint arcs.*

6. Example

We construct a Suslinian smooth dendroid D as a quotient of the Lelek fan L , such that $E(D)$ is homeomorphic to $E(L)$. The example shows that Theorem 15 and Corollary 16 are false outside the plane.

Let's begin by understanding what a Suslinian quotient of the Cantor fan might involve. Let C be the middle-thirds Cantor set. For each

$$n = 0, 1, 2, \dots$$

let \mathcal{C}^n be the natural partition of C into 2^n disjoint closed sets of diameter 3^{-n} . In $C \times [0, 1]$, put $\langle c, 0 \rangle \sim \langle d, 0 \rangle$ for all $c, d \in C$. For each $n \geq 1$ define $\langle c, t \rangle \sim \langle d, t \rangle$ if c and d are in the same member of \mathcal{C}^n and

$$t \in [1 - 2^{1-n}, 1 - 2^{-n}].$$

The equivalence classes under the relation \sim form an upper semi-continuous decomposition of the Cantor fan, and the quotient space is the Gehman dendrite (see Figure 3). Notice that the endpoints are untouched by the identification of arcs; the endpoint set of the Gehman dendrite is the Cantor set.




FIGURE 3. A quotient of the Cantor fan.

In order to apply a similar construction to the Lelek fan, we will require a few more definitions. If \mathcal{C} and \mathcal{D} are partitions of a set X , then \mathcal{D} **refines** \mathcal{C} if for every $D \in \mathcal{D}$ there exists $C \in \mathcal{C}$ such that $D \subset C$. If \mathcal{A} is a collection of pairwise disjoint closed sets, then \mathcal{C} **respects** \mathcal{A} if for every $C \in \mathcal{C}$ and $A \in \mathcal{A}$, either $C \subset A$ or $C \cap A = \emptyset$. A **null partition** of X is a countable partition of X whose elements form a null sequence (i.e. their diameters converge to 0).

LEMMA 17. *Let $\mathcal{C} = \{C_n : n < \omega\}$ be a closed partition of the Cantor set C , and suppose that A_n is a closed subset of C_n for each $n < \omega$. Then for any $\varepsilon > 0$ there exists a null partition of C of mesh $< \varepsilon$, which refines \mathcal{C} and respects $\mathcal{A} = \{A_n : n < \omega\}$.*

PROOF. Each A_n is an intersection of clopen sets in C , so $C_n \setminus A_n$ can be written as a countable union of pairwise disjoint closed sets B_0^n, B_1^n, \dots and so forth. We now have that $\{A_n : n < \omega\} \cup \{B_i^n : n, i < \omega\}$ is a countable partition of C which can be enumerated D_0, D_1, \dots and so on. Each D_k is partitioned by a finite collection \mathcal{D}_k of pairwise disjoint closed sets of diameter $< \varepsilon/k$. The desired partition of C is formed by the members of all \mathcal{D}_k 's. \square

EXAMPLE 1. *There exists a Suslinian smooth dendroid D such that $E(D) \simeq E(L)$.*

PROOF. Given a function $\varphi : C \rightarrow [0, 1]$ with Cantor set domain, we define

$$L_0^\varphi = \bigcup_{c \in C} \{c\} \times [0, \varphi(c)].$$

We consider the Lelek fan L as the quotient of some L_0^φ that is obtained by shrinking $C \times \{0\}$ to a single point (the vertex of the fan). Thus $E(L) = \{\langle c, \varphi(c) \rangle : \varphi(c) > 0\}$.

In L_0^φ , put $\langle c, 0 \rangle \sim \langle d, 0 \rangle$ for all $c, d \in C$. Let $A_1 = \{c \in C : \varphi(c) \geq \frac{3}{4}\}$. Let \mathcal{C}^1 be a null partition of C that respects A_1 . For each $c \in C$ let

$$\varphi_1(c) = \begin{cases} \frac{1}{2} & \text{if } c \in A_1 \\ 0 & \text{otherwise.} \end{cases}$$

If c and d belong to the same member of \mathcal{C}^1 , then put $\langle c, t \rangle \sim \langle d, t \rangle$ for each $t \in [0, \varphi_1(c)]$. We continue this procedure as follows.

Suppose $n \geq 2$, and $\mathcal{C}^{n-1} = \{C_k^{n-1} : k < \omega\}$ is a partition of C . Let

$$A_k^n = \{c \in C_k^{n-1} : \varphi(c) \geq \varphi_{n-1}(c) + \frac{3}{4n}\}.$$

Put

$$\varphi_n(c) = \begin{cases} \varphi_{n-1}(c) + \frac{1}{2n} & \text{if } c \in \bigcup_{k=0}^{\infty} A_k^n \\ \varphi_{n-1}(c) & \text{otherwise.} \end{cases}$$

Let \mathcal{C}^n be a null partition of mesh $< \frac{1}{n}$ that refines \mathcal{C}^{n-1} and respects \mathcal{A}^n . If c and d belong to the same member of \mathcal{C}^n , put $\langle c, t \rangle \sim \langle d, t \rangle$ for each $t \in [\varphi_{n-1}(c), \varphi_n(c)]$.

It is easily checked that the equivalence classes under \sim form an upper semi-continuous decomposition of L (note that a sequence of equivalence classes must converge to an entire equivalence class, or to a single point of L), and moreover that D is a Suslinian smooth dendroid with endpoint set $E(L)$. \square

REMARK 3. The dendroid D not planable by Theorem 15. In fact, since $E(L)$ is uncountable and 1-dimensional at each of its points, by Proposition 13 every plane continuum that homeomorphically contains $E(L)$ is non-Suslinian.

REMARK 4. A space is called **rational** if it has a basis of open sets with countable boundaries. It is known that every rational continuum is Suslinian, and that the converse is false. The dendroid D is another example of a Suslinian but not rational continuum, owing to the fact that $E(L)$ is not rational [6, Corollary 4.8].

REMARK 5. Lelek constructed a Suslinian smooth plane dendroid which is not rational [12, Example 3.1]. Its endpoints are dense and zero-dimensional.

It is unknown whether a Bellamy dendroid (as in [1]) can be Suslinian.

QUESTION 1. *Let X be a smooth plane dendroid. If X is Suslinian, then is $E(X)$ not connected?*

References

- [1] D. Bellamy, An interesting plane dendroid, *Fund. Math.* 110 (1980), 191–207.
- [2] K. Borsuk, A theorem on fixed points, *Bull. de l'Academie Polonaise des Sciences, Sériés des Sciences Astronomiques, Math. et Phys.* (1954), 17–20.
- [3] J. Charatonik, Problems and remarks on contractibility of curves. In: Novák, Josef (ed.): *General topology and its relations to modern analysis and algebra IV*, Proceedings of the fourth Prague topological symposium, 1976. Part B: Contributed Papers. Society of Czechoslovak Mathematicians and Physicist, Praha, 1977. pp. 72–76.
- [4] J. Charatonik and C. Eberhart, On smooth dendroids, *Fund. Math.* 67 (1970), 297–322.
J. J. Charatonik, On fans, *Diss. Math. (Rozprawy Mat.)* 54 (1967).
W. J. Charatonik, The Lelek fan is unique, *Houston J. Math.* 15 (1989), no. 1, 27–34.
- [5] H. Cook, Tree-likeness of dendroids and λ -dendroids. *Fund. Math.* 68 (1970), 19–22.
- [6] J. Dijkstra and D. Lipham, On cohesive almost zero-dimensional spaces, *Can. Math. Bull.* 64 (2021), 429–441.
- [7] J. Dijkstra and J. van Mill, Erdős space and homeomorphism groups of manifolds, *Mem. Amer. Math. Soc.* 208 (2010), no. 979.
- [8] R. Engelking, *Dimension Theory*, Volume 19 of Mathematical Studies, North-Holland, Amsterdam, 1978.
- [9] B. Fugate, G. R. Gordh Jr. and Lewis Lum, Arc-Smooth Continua, *Trans. Amer. Math. Soc.* 265 (1981), 545–561.
- [10] J. Krasinkiewicz and P. Minc, Dendroids and their endpoints, *Fund. Math.* 99 (1978), 227–244.
- [11] A. Lelek, On plane dendroids and their end points in the classical sense, *Fund. Math.* 49 (1961), 301–319.
- [12] A. Lelek, On the topology of curves II, *Fund. Math.* 70 (1971), 131–138.
- [13] D. Lipham, A dichotomy for spaces near dimension zero, *Topology Appl.*, Volume 336 (2023), 108611.
- [14] D. Lipham, J. van Mill, M. Tuncali, E. Tymchatyn, K. Valkenburg, Buried points of plane continua. Preprint; <https://arxiv.org/abs/2401.10206> (2024).
- [15] R. Manka, Results and problems in fixed point theory for tree-like continua. *Topology Proc.* 39 (2012), 113–130.
- [16] V. Martínez-de-la-Vega, J. Martínez-Montejano, Open Problems on Dendroids, *Open Problems in Topology II* Chapter 35, 2007 p.319–334.
- [17] J. van Mill and M. Tuncali, Plane continua and totally disconnected sets of buried points. *Proc. Amer. Math. Soc.* 140 (2012), no. 1, 351–356.

- [18] S. B. Nadler Jr., Continuum Theory: An Introduction, Pure Appl. Math., vol. 158, Marcel Dekker, Inc., New York, 1992.
- [19] J. Nikiel, A characterization of dendroids with uncountably many endpoints in the classical sense, Houston J. Math. 9 (1983) 421–432.
- [20] J. Nikiel and E. Tymchatyn, Sets of end-points and ramification points in dendroids, Fund. Math. 138 (1991), 139–146.
- [21] G. T. Whyburn, Analytic topology, AMS Colloquium Publications, Volume 28. American Mathematical Society, New York, 1942.

DEPARTMENT OF MATHEMATICS AND DATA SCIENCE, COLLEGE OF COASTAL GEORGIA, BRUNSWICK
GA 31520, UNITED STATES OF AMERICA
Email address: `dlipham@ccga.edu`