Thermodynamic constraints on polar active matter hydrodynamics

Andrea Amoretti,^{1,2,*} Daniel K. Brattan,^{1,2,†} and Luca Martinoia^{1,2,‡}

¹Dipartimento di Fisica, Università di Genova, via Dodecaneso 33, I-16146, Genova, Italy

²I.N.F.N. - Sezione di Genova, via Dodecaneso 33, I-16146, Genova, Italy

(Dated: May 6, 2024)

In this letter we use the thermodynamics of a passive fluid to constrain transport in the corresponding active fluid which is subsequently described by the Toner-Tu equations. Acknowledging that the system fundamentally breaks boost symmetry we compel what were previously entirely phenomenological parameters in the Toner-Tu model to satisfy precise relationships among themselves. Consequently, we determine exact scalings for the transport coefficients under dynamical renormalisation group flow that reproduce the results of recent simulations.

Polar active systems [1] are composed of self-propelled particles that align their movement with their closest neighbours. These systems demonstrate collective behaviors such as flocking and dynamic pattern formation, providing valuable insights into biological phenomena and serving as a framework for understanding complex systems.

Among the various theoretical approaches aimed at describing these systems, special attention is given to the model first proposed in [2–5]. There, Toner and Tu put forward a hydrodynamic description of flocking behavior, drawing inspiration from the equations governing liquid crystals. They formulate a set of equations for a dynamic velocity field \vec{v} and a conserved number density n, resembling the Navier-Stokes equation for passive fluids. However, in this case, boost symmetry is not conserved due to the system's polar nature. The system's preferred background velocity is attained by coupling these equations to an external potential U for the velocity field, akin to the well-known Mexican hat potential. The system's active nature is further accounted for by incorporating a random noise self-driving force \vec{f} .

Despite the considerable success in elucidating the largescale dynamics of polar active systems, the Toner and Tu model lacks a clear microscopic derivation, a gap only partially addressed by kinetic theory [6–10]. These approaches rely on the assumption that momentum equals mass current, secretly imposing that Galilean boosts are a system symmetry. However, the phenomenological equations proposed by Toner and Tu break Galilean boosts, indicating a disconnect between momentum and mass current in active matter systems.

In this letter, we demonstrate that polar active fluids are akin to passive fluids lacking boost symmetry, provided they are supplemented with a non-thermal noise term and a potential inducing spontaneous symmetry breaking. For hydrodynamics to offer a sensible description [11], there must be a scale separation within the system, ensuring rapid local equilibration compared to long-scale collective dynamics. The necessity of a local Gibbs-like distribution heavily constrains the constitutive relations of the ideal-fluid model. Notably, this implies that the quantity akin to pressure in the Toner-Tu equations can be treated as a thermodynamic pressure in many respects. Consequently, we find that a subset of the derived transport coefficients is thermodynamically constrained. Our argument centers on the requirement that an active fluid maintains consistency with basic thermodynamics, particularly in its disordered phase [12]. While tuning the symmetry-breaking potential restricts the space of steadystate configurations and consequently affects collective dynamics and response to perturbations [2–4], it does not alter the necessity for constitutive relations to adhere to thermodynamic principles. This argument is also supported by recent works, suggesting that standard hydrodynamics is a good description of systems that are in non-equilibrium steady states [13, 14].

Notably, these constraints enable us to derive analytical expressions for the critical exponents of the system in arbitrary spatial dimension d, aligning precisely with numerical computations in the Vicsek model [15] and differing from those derived by [3].

Thermodynamically consistent polar active fluids - We begin by deriving the ideal constitutive relations for any thermodynamic system dependent on number density nand velocity \vec{v} . To obtain a generating functional for the relevant consistutive relations one promotes translation invariance to a local symmetry in a system without boosts [16, 17]. The leading term in a small derivative expansion for the effective action is then just the integral of a scalar, $P(\mu, \vec{v}^2)$, where μ will be the chemical potential for n. In the passive fluid, P is the usual thermodynamic pressure. Variations of the background on which the fluid sits then produce the desired constitutive relations. In particular, the ideal-fluid constitutive relations take the form [13, 16– 18]

$$g^{i} = \rho v^{i}, \ \tau^{ij} = \rho v^{i} v^{j} + P \delta^{ij}, \ j^{i} = n v^{i},$$
 (1a)

where $\vec{g} = \partial P / \partial \vec{v}$ is the momentum conjugate to the velocity, $\rho = 2\partial P / \partial \vec{v}^2$ is the kinetic mass density and $n = \partial P / \partial \mu$, and satisfy the following conservation equations

$$\partial_t g^i + \partial_j \tau^{ji} = 0$$
, $\partial_t n + \partial_i j^i = 0$. (1b)

Equations (1b) follow respectively from translation invariance and particle number conservation. Notice in particular that $\vec{j} \neq \vec{g}$ [19], which is a consequence of the breaking of the Milne or Galilean Ward identity that holds only for systems with an exact boost symmetry [20, 21].

To the ideal constitutive relations (1a) we proceed as usual and add dissipative corrections at order-one in derivatives, correcting τ^{ij} and j^i in (1). In fact, order-one hydrodynamics for fluids without boost symmetry has been studied thoroughly in [16, 17, 22]. For ease of reference, we summarise the expressions in the supplemental material (SM) [23].

With the order one consistitutive relations determined, we turn on non-thermal noise and the external potential U in the normative manner - by adding them to (1b) by hand. In this regard we treat U no differently to the noise \vec{f} in assuming that the form of the consistutive relations are unchanged by these external forces. Although \vec{g} is no longer a conserved quantity, it can certainly be a relevant hydrodynamic variable if the external potential, which breaks the symmetries, is weak enough [24, 25]. In our case U should be understood as imposing extra constraints on the system which reduce the allowed space of stationary states from those permitted to the passive fluid i.e. the stationary state velocity will be fixed to a particular value.

With these considerations we can now state the Toner-Tu equations which are

$$\partial_t \vec{v} + \lambda_1 (\vec{v} \cdot \vec{\nabla}) \vec{v} + \lambda_2 (\vec{\nabla} \cdot \vec{v}) \vec{v} + \lambda_3 \vec{\nabla} |\vec{v}|^2 + \vec{\nabla} P_1 + \vec{v} (\vec{v} \cdot \vec{\nabla} P_2) = U \vec{v} + D_B \vec{\nabla} (\vec{\nabla} \cdot \vec{v}) + D_T \nabla^2 \vec{v} + D_2 (\vec{v} \cdot \vec{\nabla})^2 \vec{v} + \vec{f} , \quad (2a)$$

$$\partial_t n + \lambda_n \nabla \cdot (n\vec{v}) = 0 , \qquad (2b)$$

where all transport coefficients above are functions of nand \vec{v}^2 , U is bounded below and has a zero at some \vec{v}^2 and we assume that \vec{f} is a Gaussian white non-thermal noise

$$\langle f_i(\vec{x},t)f_j(\vec{x}',t')\rangle = \Delta\delta_{ij}\delta^d(\vec{x}-\vec{x}')\delta(t-t') \quad (2c)$$

with Δ some constant. As was done in the original paper [3], we have set derivatives of the particle density in (2a) to zero for legibility. These are easily restored and do not affect the physics we are interested in.

The Toner-Tu equations above (2) include λ_1 , λ_2 , λ_3 and λ_n as phenomenological parameters. The equivalent terms in our model are fixed by the thermodynamics of the passive fluid,

$$\lambda_1 = \lambda_n = 1 , \ \lambda_2 = \frac{1 - \frac{\partial \rho}{\partial n} \frac{n}{\rho}}{1 + 2 \frac{\partial \rho}{\partial v^2} \frac{v^2}{\rho}} , \ \lambda_3 = 0 , \qquad (3a)$$

with

$$\vec{\nabla}P_1 = \frac{1}{\rho}\vec{\nabla}P \;, \; \vec{v}\cdot\vec{\nabla}P_2 = \frac{\frac{2}{\rho}\frac{\partial\rho}{\partial v^2}}{1+2\frac{\partial\rho}{\partial v^2}\frac{v^2}{\rho}}\vec{v}\cdot\vec{\nabla}P \;, \quad (3b)$$

where again, all thermodynamic derivatives are taken with respect to constant \vec{v}^2 or n. It should be noted that λ_3 can be shifted to an arbitrary function by redefining the pressure $P \rightarrow P + f(\vec{v}^2, n)$, as in the original Toner-Tu model. We have made the logical choice for its value and assume that we are supplied with the form of the pressure P to fix any ambiguity.

Restoring the boost symmetry amounts to setting $\vec{g} = n\vec{v}$ in (3) as is done in some works [6–10]. This choice would force upon us the constraint $\lambda_1 = 1$ and $\lambda_2 = 0$, as can be seen by direct substitution into (3a). Subsequently, one can redefine the chemical potential to remove the anisotropic pressure and general dependence of thermodynamics on \vec{v}^2 . This conclusion is independent of the precise nature of the microscopic theory, following only from symmetries. Moreover, we see that an anisotropic pressure term is a natural consequence of boost agnosticity even in equilibrium.

Without the imposition $\vec{g} = \vec{j}$, the requirement of having a local Gibbs-like distribution has heavily constrainted the ideal-fluid constitutive relations such that the four independent phenomenological parameters, λ_1 , λ_2 and λ_3 and λ_n in the original work [3] (see also (2)) are reduced to one (P) and its derivatives.

The ordered phase - Only for specific flows and choices of U in (2) would we expect the system of equations to arrive at a state with zero background velocity $\vec{v} = \vec{0}$ at late times. For a \vec{v}^2 -dependent generic potential, bounded below by zero, the system will flow to a zero of U and break rotations spontaneously. It is typical in the literature to take $U = \alpha - \beta \vec{v}^2$, so that $U\vec{v}$ behaves like a derivative of the Mexican hat potential. Around the new steady state we can then linearise in the velocity

$$\vec{v} = (v_0 + \delta v_{\parallel})\hat{x}_{\parallel} + \vec{v}_{\perp} \tag{4}$$

with δv_{\parallel} and \vec{v}_{\perp} small fluctuations which are respectively longitudinal or orthogonal to the background velocity $v_0 \hat{x}_{\parallel}$. v_0 is defined such that it obeys $U(n_0, v_0^2) = 0$, with n_0 the equilibrium particle number density. In the ordered phase the longitudinal mode associated with δv_{\parallel} is gapped, therefore it can be integrated out to find a set of reduced equations for the Goldstone-like gapless mode \vec{v}_{\perp} and $\delta n = n - n_0$. The details are rather long and we refer to [23] for specifics, here we outline the main steps, which however follow closely the analysis of [5].

We solve the equation for δv_{\parallel} iteratively in terms of \vec{v}_{\perp} and δn treating derivatives and linearisation as small in the same parameter ϵ . Eliminating δv_{\parallel} , the charge conservation equation expanded up to $\mathcal{O}(\epsilon^3)$ is

$$\partial_t \delta n + v_2 \partial_{\parallel} \delta n + n_0 \vec{\nabla}_{\perp} \cdot \vec{v}_{\perp} + w_1 \vec{\nabla}_{\perp} (\vec{v}_{\perp} \delta n) - w_2 \partial_{\parallel} (\delta n)^2 - w_3 \partial_{\parallel} |\vec{v}_{\perp}|^2 - \xi_n \partial_{\parallel} \left(\vec{\nabla}_{\perp} \cdot \vec{v}_{\perp} \right) - D_{\parallel} \partial_{\parallel}^2 \delta n - D_{\perp} \nabla_{\perp}^2 \delta n = 0$$
(5a)

where the expressions of the coefficients can be found in the SM [23]. Similarly, the equation of motion for the transverse velocity fluctuations \vec{v}_{\perp} becomes

$$\begin{aligned} \partial_t \vec{v}_{\perp} + v_0 \partial_{\parallel} \vec{v}_{\perp} + \kappa \vec{\nabla}_{\perp} \delta n + g_1 \delta n \partial_{\parallel} \vec{v}_{\perp} + g_2 \vec{v}_{\perp} \partial_{\parallel} \delta n \\ &+ g_3 \vec{\nabla}_{\perp} (\delta n)^2 + g_4 \vec{v}_{\perp} (\vec{\nabla}_{\perp} \cdot \vec{v}_{\perp}) + g_5 (\vec{v}_{\perp} \cdot \vec{\nabla}_{\perp}) \vec{v}_{\perp} \\ &- D_v \vec{\nabla}_{\perp} \partial_{\parallel} \delta n - \xi_v \vec{\nabla}_{\perp} (\vec{\nabla}_{\perp} \cdot \vec{v}_{\perp}) \\ &- \xi_{\perp} \nabla^2_{\perp} \vec{v}_{\perp} - \xi_{\parallel} \partial^2_{\parallel} \vec{v}_{\perp} = \vec{f}_{\perp} \end{aligned}$$
(5b)

The v_2 and κ terms are linear in fluctuations and derivatives, the g_i and w_i terms are the non-linear corrections, while the generalized diffusivities D_i and viscosities ξ_i are related to first-order transport coefficients [23].

The form of these reduced equations (5a) and (5b) matches exactly the expressions in [5], with a few minor differences. First, we observe that we have a new non-linear term g_4 , which is missing in [5], secondly we find that the bare coefficient D_{\perp} is non-zero, while it appears to be zero in [5]. Finally, $g_5 = 1$ and v_0 appears in front of $\partial_{\parallel} \vec{v}_{\perp}$, despite the lack of Galilean invariance. These modifications aside, we have fully reproduced the effective equations (5) of Toner and Tu for the hydrodynamics. Thus the non-equilibrium features characterizing active matter are not relevant for a hydrodynamic description.

As the linearized equations are the same as in [5], the modes and correlators take the same form, with different values of the coefficients. In particular, we can compute the modes from the full theory (2) or from the reduced equations (5a), (5b) obtaining the same expressions.

One of the key signatures of the Toner and Tu model is the fact that there are two speeds of sound $v_{\pm}(\theta)$, where θ is the angle between the wave vector \vec{q} and the background velocity. Specifically at $\theta = 0$ one finds $v_{+}(\theta = 0) = v_{0}$ and $v_{-}(\theta = 0) = v_{0}\lambda_{1}$ [3]. This effect is observed in simulations [26] and experiments [27] and used to provide an estimate of $\lambda_{1} \approx 0.75$, which was interpreted as a signature of the breaking of Galilean boost symmetry. As we showed, from our boost-agnostic theory we find $(g_{5} =)\lambda_{1} = 1$. It would naively seem that our model fails this simple test - however this is not correct. Indeed, at $\theta = 0$ we find that the sound speed once again has two values [5]

$$v_{+}(\theta = 0) = v_{0}$$
 $v_{-}(\theta = 0) = v_{0} \left(1 - \frac{n_{0}U_{n}}{2v_{0}^{2}U_{v}}\right)$ (6)

and the difference vanishes when the $U_n = \left(\frac{\partial U}{\partial n}\right)_{v^2} = 0$. The fact that $U_n \neq 0$ is what allows us to have two different speeds of sound at $\theta = 0$, despite having $\lambda_1 = 1$.

Interestingly, the parameter U_n in (6) is key to the breaking of time-reversal invariance [28, 29]. Notice that the Toner and Tu equations [3] break Onsager relations [30] not only in the coefficient U_n , but also through the λ_i (see also [31] for a discussion on time-reversal invariance in active-matter systems). This is very unusual from the point of view of hydrodynamics: usually in hydrodynamics the ideal fluid (which follows from local equilibrium) is always time-reversal invariant. Breaking of the Onsager relations only appears through viscous corrections or external forces. Contrary to this, the Toner and Tu model violates Onsager relations at lowest order in derivatives, therefore equilibrium (even in the disordered phase!) is not time-reversal invariant.

To conclude, we are able to reproduce the observed phenomenology of the Toner-Tu model from the hydrodynamic of passive fluids, suggesting that the active character does not modify the constitutive relations.

Exact scaling exponents - Having derived the Toner-Tu equations from first principles we use the fact that several of our transport coefficients are pure functions of thermodynamic parameters (3) to our advantage: namely, we constrain the scaling behaviour of the transport coefficients. Let's study how the various terms change under the dynamical renormalization group [32, 33]. Using standard notations, we rescale fields and coordinates as $(\vec{v}_{\perp}, \delta n, \vec{x}_{\perp}, x_{\parallel}, t) = (b^{\chi} \vec{v}'_{\perp}, b^{\chi} \delta n', b \vec{x}'_{\perp}, b^{\zeta} x'_{\parallel}, b^{z} t')$. With this choice of scalings the equations of motion take the same form as before, but with rescaled transport coefficients given by:

$$(w_1, g_{3,4,5})' = b^{z-1+\chi} ((w_1, g_{3,4,5}) + \text{graphs}) ,$$
 (7a)

$$(w_{2,3}, g_{1,2})' = b^{z-\zeta+\chi} \left((w_{2,3}, g_{1,2}) + \text{graphs} \right) , \quad (7b)$$

(

(

$$(\xi_n, D_v)' = b^{z-1-\zeta} \left((\xi_n, D_v) + \text{graphs} \right) , \qquad (7c)$$

$$(\xi_{\parallel}, D_{\parallel})' = b^{z-2\zeta} \left((\xi_{\parallel}, D_{\parallel}) + \text{graphs} \right) , \qquad (7d)$$

$$(\xi_{v,\perp}, D_{\perp})' = b^{z-2} \left((\xi_{v,\perp}, D_{\perp}) + \text{graphs} \right)$$
 (7e)

$$\Delta' = b^{z-\zeta+1-d-2\chi} \left(\Delta + \text{graphs}\right) , \qquad (7f)$$

where "graphs" represents perturbative corrections that are obtained from integrating out the fast modes with wave vector $b^{-1}\Lambda \leq |\vec{q}_{\perp}| \leq \Lambda$ and Λ is the UV cutoff. As was shown in [3, 33], the asymptotic $\vec{q} \to 0$ behaviour of the correlators is determined by the fixed point values of the scaling exponents ζ , z and χ . Therefore, a lot of effort has been devoted to identifying the critical points of these β -functions using both clever symmetry arguments [3, 34–36] or through more direct computation [37, 38].

The simplest such example is the linear fixed point, where non-linear terms are ignored, and it is given by $(z, \zeta, \chi) = (2, 1, \frac{2-d}{2})$. However, the linear fixed point is unstable for d < 4, as non-linearities become relevant in the IR. Another example is found in [3], where Toner and Tu obtained exact exponents in d = 2 taking advantage of an emergent pseudo-Galilean symmetry. They obtain $(z, \zeta, \chi) = (6/5, 3/5, -1/5)$. These results however are invalidated by the more accurate analysis of [5], which agrees with our expressions (5a) and (5b), due to the presence of new non-linearities ignored in the original work. One of the advantages of our derivation is that it provides a new argument to fix the values of the critical exponents exactly. Looking at the non-linear terms, the associated bare transport coefficients g_i and w_i all depend on global equilibrium thermodynamic properties (like ρ_0 , n_0 , v_0 and P), their thermodynamic derivatives (such as susceptibilities, compressibilities, ...) and derivatives of the external potential $\left(\frac{\partial U}{\partial n}\right)_{v^2}$ and $\left(\frac{\partial U}{\partial v^2}\right)_n$. Our claim is that because of the thermodynamic origin of these quantities, they should not receive perturbative corrections: after all, in global equilibrium (where these quantities are defined) there are no hydrodynamic modes that can renormalize these values.

This is indeed what happens in Navier-Stokes hydrodynamics [33]. Consider the equations in (2) and restore the Galilean boost symmetry, setting $\lambda_1 = 1$ and $\lambda_2 = \lambda_3 = 0$. Then, the usual argument goes that λ_1 cannot receive perturbative corrections to its scaling because it is protected by the Galilean symmetry and must remain equal to 1. Indeed, various forms of this (pseudo-)Galilean symmetry are at the core of the exact results obtained in the past. The same argument, however, cannot be used to fix the λ_n term in the equations, which is not fixed by Galilean invariance, but rather by thermodynamics (which determines the ideal-fluid part of the equations).

This argument could not be used in [3–5] as the λ_i in (2) were understood to be non-equilibrium transport coefficients. As they are not derived from equilibrium thermodynamics, and were treated as closer to generalized viscosities in spirit, it was not possible to fix their scalings uniquely in the past. On the other hand our derivation, which highlights the thermodynamic origin of the coefficients, indicates that these non-linear (ideal-fluid) terms should never receive perturbative corrections.

Similar conclusions apply to the derivatives of the external potential which, strictly speaking, are not obtained from equilibrium thermodynamics (in the sense that they are something more than just the equation of state). Nonetheless, they too do not receive perturbative corrections. In particular, it is well known that the positions of the peaks in the correlators (the speeds of sound) do not renormalize under dynamical renormalization group [3, 33]. Then, we can use these expressions to uniquely fix the values of the thermodynamic derivatives of the potential U. As a constraint it too determines the global hydrostatic properties of the fluid, which cannot be affected by fast modes.

Consequently, imposing that the non-linear terms g_i and w_i do not receive graphical corrections, we find two equations for the fixed point

$$z - 1 + \chi = 0$$
 $z - \zeta + \chi = 0$ (8)

To completely fix the exponents we need a third equation. To do that, we will employ the hyperscaling relation, which follows from assuming that the scaling of Δ is exact. The reason for this is very practical, and comes from the

	d = 2			d = 3		
	TT98	Vicsek	this work	TT98	Vicsek	this work
χ ζ z α	-1/5 3/5 6/5 8/5	$\begin{array}{c} -0.31(2) \\ 0.95(2) \\ 1.33(2) \\ 1.67(2) \end{array}$	-1/3 1 4/3 5/3	$ -3/5 \\ 4/5 \\ 8/5 \\ 14/9$	-0.62 1 1.77 1.59(3)	-2/3 1 5/3 23/15

TABLE I. Critical exponents for Toner an Tu [3] and Vicsek [39], compared to the exact exponent computed in this work.

fact that recent simulations suggest it is a universal law [39]. Furthermore, from a field-theoretical perspective, a constant noise correlator Δ should in general receive contributions from at least two non-linearities, each carrying one power of \vec{q} , thus contributing to $\mathcal{O}(q^2)$ corrections, which are irrelevant compared to Δ in the hydrodynamic limit, up to unlikely cancellations. Imposing this last constraint

$$z - \zeta + 1 - d - 2\chi = 0 \tag{9}$$

we find the exact scaling exponents

$$z = \frac{2+d}{3}$$
 $\zeta = 1$ $\chi = \frac{1-d}{3}$ (10)

Notice that these expressions correctly interpolate between the d < 4 theory and the mean field theory results. In the specific case of d = 2 our argument is supported by recent studies based on symmetry arguments [40].

As expected, $\chi < 0$ in d = 2, implying that velocity fluctuations are weak and do not destroy the ordered phase. The Mermin-Wagner theorem would forbid continuous phase transitions below 2 dimesions, however active matter systems defy this expectation due to their "nonequilibrium nature", typically attributed to the λ_i . In our approach however the λ_i are a consequence of local equilibrium, therefore it is unexpected that they can allow for an ordered phase below 2 dimensions.

These values appear to be in perfect agreement with recent state-of-the-art numerics on the Vicsek model [39], which found discrepancies when comparing to the standard Toner and Tu expressions, as outlined in Table I. We also included in the table the large number fluctuation exponents $\langle \delta N \rangle^2 \sim \langle N \rangle^{\alpha}$. These exponents should be 1 for thermal phases, while polar active matter hydrodynamics predicts $\alpha = 1 + (d + \zeta + 2\chi - 1)/d$ [1, 39, 41].

Outlook - In this letter, we have demonstrated that polar active fluids exhibit similarities to passive fluids lacking boost symmetry when supplemented with a nonthermal noise term and a potential inducing spontaneous symmetry breaking. Consequently, we precisely derived expressions for the critical exponent. These expressions align excellently with numerical simulations of the Vicsek model and also correlate well with experimental observations on epithelial cells [42], self-propelled rollers [27], bacterial colonies [43], and earlier simulations of the Vicsek model [44, 45]. This paves the way for more precise computations, where the constraint derived in our approach can aid in computing radiative corrections and achieving better agreement with experimental results.

On the theoretical front, we expect our argument to be valid for generic systems, like Malthusian flocks [34, 40] or incompressible phases [2]. Moreover, we omitted the consideration of temperature, as is customary in the Toner-Tu model. However, it is straightforward to introduce an additional scalar quantity, T, into our formalism, leading to a conservation equation associated with time translation. Naturally, we anticipate this equation to undergo relaxation similar to that of the equation for \vec{g} . Understanding the implications of this relaxed equation could provide insight into the nature of entropy conservation/production within the effective active matter systems.

ACKNOWLEDGEMENTS

A.A., D.B. acknowledge support from the project PRIN 2022A8CJP3 by the Italian Ministry of University and Research (MUR). L.M. acknowledge support from the project PRIN 2022ZTPK4E by the Italian Ministry of University and Research (MUR). This project has also received funding from the European Union's Horizon 2020 research and innovation programme under the Marie Sklodowska-Curie grant agreement No. 101030915.

- M. C. Marchetti, J. F. Joanny, S. Ramaswamy, T. B. Liverpool, J. Prost, M. Rao, and R. A. Simha, Hydrodynamics of soft active matter, Rev. Mod. Phys. 85, 1143 (2013).
- [2] J. Toner and Y. Tu, Long-range order in a twodimensional dynamical XY model: How birds fly together, Phys. Rev. Lett. **75**, 4326 (1995).
- [3] J. Toner and Y. Tu, Flocks, herds, and schools: A quantitative theory of flocking, Phys. Rev. E 58, 4828 (1998).
- [4] J. Toner, Y. Tu, and S. Ramaswamy, Hydrodynamics and phases of flocks, Annals of Physics **318**, 170 (2005).
- [5] J. Toner, Reanalysis of the hydrodynamic theory of fluid, polar-ordered flocks, Phys. Rev. E 86, 031918 (2012).
- [6] E. Bertin, M. Droz, and G. Grégoire, Boltzmann and hydrodynamic description for self-propelled particles, Phys. Rev. E 74, 022101 (2006).
- [7] E. Bertin, M. Droz, and G. Grégoire, Hydrodynamic equations for self-propelled particles: microscopic derivation and stability analysis, Journal of Physics A: Mathematical and Theoretical 42, 445001 (2009).
- [8] T. Ihle, Kinetic theory of flocking: Derivation of hydrodynamic equations, Phys. Rev. E 83, 030901 (2011).
- [9] A. Baskaran and M. C. Marchetti, Enhanced diffusion and ordering of self-propelled rods, Phys. Rev. Lett. 101, 268101 (2008).

- [10] A. Patelli, Landau kinetic equation for dry aligning active models, Journal of Statistical Mechanics: Theory and Experiment **2021**, 033210 (2021).
- [11] D. Forster, Hydrodynamic Fluctuations, Broken Symmetry, and Correlation Functions, Frontiers in Physics : a lecture note and reprint series (W. A. Benjamin, Advanced Book Program, 1975).
- [12] T. Agranov, R. L. Jack, M. E. Cates, and Étienne Fodor, Thermodynamically consistent flocking: From discontinuous to continuous transitions (2024), arXiv:2401.09901 [cond-mat.stat-mech].
- [13] A. Amoretti, D. K. Brattan, L. Martinoia, and I. Matthaiakakis, Non-dissipative electrically driven fluids, JHEP 2023, 218, arXiv:2211.05791 [hep-th].
- [14] D. K. Brattan, M. Matsumoto, M. Baggioli, and A. Amoretti, Relaxed hydrodynamic theory of electrically driven non-equilibrium steady states (2024), arXiv:2404.05568 [cond-mat.stat-mech].
- [15] T. Vicsek, A. Czirók, E. Ben-Jacob, I. Cohen, and O. Shochet, Novel type of phase transition in a system of self-driven particles, Phys. Rev. Lett. **75**, 1226 (1995).
- [16] J. Armas and A. Jain, Effective field theory for hydrodynamics without boosts, SciPost Phys. 11, 054 (2021), arXiv:2010.15782 [hep-th].
- [17] J. de Boer, J. Hartong, E. Have, N. A. Obers, and W. Sybesma, Non-Boost Invariant Fluid Dynamics, Sci-Post Phys. 9, 018 (2020), arXiv:2004.10759 [hep-th].
- [18] J. de Boer, J. Hartong, N. A. Obers, W. Sybesma, and S. Vandoren, Perfect Fluids, SciPost Phys. 5, 003 (2018), arXiv:1710.04708 [hep-th].
- [19] We set the particles' mass m = 1 without loss of generality.
- [20] K. Jensen, On the coupling of Galilean-invariant field theories to curved spacetime, SciPost Phys. 5, 011 (2018), arXiv:1408.6855 [hep-th].
- [21] M. H. Christensen, J. Hartong, N. A. Obers, and B. Rollier, Boundary Stress-Energy Tensor and Newton-Cartan Geometry in Lifshitz Holography, JHEP 01, 057, arXiv:1311.6471 [hep-th].
- [22] I. Novak, J. Sonner, and B. Withers, Hydrodynamics without boosts, JHEP 07, 165, arXiv:1911.02578 [hepth].
- [23] L. Martinoia, D. K. Brattan, and A. Amoretti, Supplemental information (2024).
- [24] S. Grozdanov, A. Lucas, and N. Poovuttikul, Holography and hydrodynamics with weakly broken symmetries, Phys. Rev. D 99, 086012 (2019), arXiv:1810.10016 [hep-th].
- [25] L. Martinoia, Developments in quasihydrodynamics, Ph.D. thesis, Genoa U. (2024), arXiv:2403.14254 [hep-th].
- [26] Y. Tu, J. Toner, and M. Ulm, Sound waves and the absence of galilean invariance in flocks, Phys. Rev. Lett. 80, 4819 (1998).
- [27] D. Geyer, A. Morin, and D. Bartolo, Sounds and hydrodynamics of polar active fluids, Nature Materials 17, 789–793 (2018).
- [28] A. Amoretti, D. K. Brattan, L. Martinoia, I. Matthaiakakis, and J. Rongen, Relaxation terms for anomalous hydrodynamic transport in Weyl semimetals from kinetic theory, JHEP 02, 071, arXiv:2309.05692 [hep-th].
- [29] A. Amoretti, D. K. Brattan, L. Martinoia, and I. Matthaiakakis, Restoring time-reversal covariance in relaxed hydrodynamics, Phys. Rev. D 108, 056003 (2023), arXiv:2304.01248 [hep-th].

- [30] M. J. Bowick, N. Fakhri, M. C. Marchetti, and S. Ramaswamy, Symmetry, thermodynamics, and topology in active matter, Phys. Rev. X 12, 010501 (2022).
- [31] E. Fodor, C. Nardini, M. E. Cates, J. Tailleur, P. Visco, and F. van Wijland, How far from equilibrium is active matter?, Phys. Rev. Lett. **117**, 038103 (2016).
- [32] S.-k. Ma and G. F. Mazenko, Critical dynamics of ferromagnets in 6ε dimensions: General discussion and detailed calculation, Phys. Rev. B **11**, 4077 (1975).
- [33] D. Forster, D. R. Nelson, and M. J. Stephen, Largedistance and long-time properties of a randomly stirred fluid, Phys. Rev. A 16, 732 (1977).
- [34] J. Toner, Birth, death, and flight: A theory of malthusian flocks, Phys. Rev. Lett. 108, 088102 (2012).
- [35] H. Ikeda, Scaling theory of continuous symmetry breaking under advection (2024), arXiv:2401.01603 [cond-mat.statmech].
- [36] H. Ikeda, Minimum scaling model and exact exponents for the nambu-goldstone modes in the vicsek model (2024), arXiv:2403.02086 [cond-mat.soft].
- [37] L. Chen, C. F. Lee, and J. Toner, Universality class for a nonequilibrium state of matter: A $d = 4 \varepsilon$ expansion study of malthusian flocks, Phys. Rev. E **102**, 022610 (2020).
- [38] P. Jentsch and C. F. Lee, A new universality class describes vicsek's flocking phase in physical dimensions (2024), arXiv:2402.01316 [cond-mat.soft].

- [39] B. Mahault, F. Ginelli, and H. Chaté, Quantitative assessment of the toner and tu theory of polar flocks, Phys. Rev. Lett. **123**, 218001 (2019).
- [40] H. Chaté and A. Solon, Dynamic scaling of twodimensional polar flocks (2024), arXiv:2403.03804 [condmat.stat-mech].
- [41] S. Ramaswamy, R. A. Simha, and J. Toner, Active nematics on a substrate: Giant number fluctuations and long-time tails, Europhysics Letters (EPL) 62, 196–202 (2003).
- [42] F. Giavazzi, C. Malinverno, S. Corallino, F. Ginelli, G. Scita, and R. Cerbino, Giant fluctuations and structural effects in a flocking epithelium, Journal of Physics D: Applied Physics 50, 384003 (2017).
- [43] H. P. Zhang, A. Be'er, E.-L. Florin, and H. L. Swinney, Collective motion and density fluctuations in bacterial colonies, Proceedings of the National Academy of Sciences 107, 13626 (2010), https://www.pnas.org/doi/pdf/10.1073/pnas.1001651107.
- [44] H. Chaté, F. Ginelli, G. Grégoire, and F. Raynaud, Collective motion of self-propelled particles interacting without cohesion, Physical Review E 77, 10.1103/physreve.77.046113 (2008).
- [45] S. Ngo, A. Peshkov, I. S. Aranson, E. Bertin, F. Ginelli, and H. Chaté, Large-scale chaos and fluctuations in active nematics, Physical Review Letters **113**, 10.1103/physrevlett.113.038302 (2014)

Supplemental Information

ORDER-ONE BOOST-AGNOSTIC HYDRODYNAMICS

In the main text we show that the hydrodynamics of polar active matter can be described as a passive fluid without boost symmetry in the presence of an external potential $U(n, \vec{v}^2)$ and a non-thermal noise. Therefore, it is important to study the order-one constitutive relations of boost-agnostic fluids, which have been discussed thoroughly in [S1, S2]. In particular, we can follow the comprehensive results in [S2].

Consider a normal fluid with momentum and particle number conservation

$$\partial_t g^i + \partial_j \tau^{ji} = 0 , \qquad \qquad \partial_t n + \partial_i j^i = 0 . \qquad (S1)$$

The ideal-fluid constitutive relations are given in the main text:

$$g^i = \rho v^i , \qquad (S2a)$$

$$\tau^{ij} = \rho v^i v^j + P \delta^{ij} , \qquad (S2b)$$

$$j^i = nv^i av{(S2c)}$$

and follow from the requirement of local equilibrium. Because active matter systems do not conserve energy, we must decouple the energy conservation equation that is considered in [S2] by setting the temperature to be non-dynamical. Furthermore, we will only consider fluids which obey time-reversal invariance of the microscopic theory. This leads us to impose the Onsager reciprocal relations [S3–S5]. However, we impose the Onsager relations only on the constitutive relations, while still allowing the potential U to break the time-reversal symmetry. This is necessary to account for certain results obtained in simulations. Physically, this represents the fact that our fluid only starts violating the Onsager conditions when the active part (the external potential) is turned on. It must be noted that our choice is simply dictated by a matter of convenience and simplicity, while still allowing interesting results. Including Onsager-violating terms in the constitutive relations would make the expressions more complicated, but would not change the conclusions. The constitutive relations for j^i and τ^{ij} , up to order one in derivatives and in d spatial dimensions, are [S2]:

$$j^{i} = nv^{i} - \left(\gamma_{00}P^{ij}\partial_{j}\mu + \zeta_{00}\frac{v^{i}v^{j}}{|v|^{2}}\partial_{j}\mu + \gamma_{01}\frac{P^{i(j}v^{k)}}{|v|}\sigma_{jk} + \zeta_{01}\frac{v^{i}v^{j}v^{k}}{2|v|^{3}}\sigma_{jk} + \frac{\zeta_{02}}{2}\frac{v^{i}}{|v|}P^{jk}\sigma_{jk}\right),$$
(S3a)

$$\tau^{ij} = \rho v^{i}v^{j} + P\delta^{ij} - \left[\eta \left(P^{k(i}P^{j)l} - \frac{1}{d-1}P^{ij}P^{kl}\right)\sigma_{kl} + 2\gamma_{01}\frac{P^{k(i}v^{j)}}{|v|}\partial_{k}\mu + \zeta_{01}\frac{v^{i}v^{j}v^{k}}{|v|^{3}}\partial_{k}\mu + \zeta_{02}\frac{v^{k}}{|v|}P^{ij}\partial_{k}\mu + 2\gamma_{11}\frac{v^{(i}P^{j)(k}v^{l)}}{|v|^{2}}\sigma_{kl} + \frac{1}{2}\zeta_{11}\frac{v^{i}v^{j}v^{k}v^{l}}{|v|^{4}}\sigma_{kl} + \frac{\zeta_{12}}{2}\left(P^{ij}\frac{v^{k}v^{l}}{|v|^{2}} + \frac{v^{i}v^{j}}{|v|^{2}}P^{kl}\right)\sigma_{kl} + \frac{\zeta_{22}}{2}P^{ij}P^{kl}\sigma_{kl}\right],$$
(S3b)

where $\sigma_{ij} = \partial_i v_j + \partial_j v_i$, $P^{ij} = \delta^{ij} - \frac{v^i v^j}{|v|^2}$ is the projector orthogonal to the velocity, and μ is the chemical potential. The momentum instead remains fixed at its ideal-fluid value $\vec{g} = \rho \vec{v}$ (S2). Positivity of entropy production constraints the matrices of coefficients

$$\begin{pmatrix} \zeta_{00} & \zeta_{01} & \zeta_{02} \\ \zeta_{01} & \zeta_{11} & \zeta_{12} \\ \zeta_{02} & \zeta_{12} & \zeta_{22} \end{pmatrix} \ge 0 , \qquad \begin{pmatrix} \gamma_{00} & \gamma_{01} \\ \gamma_{01} & \gamma_{11} \end{pmatrix} \ge 0 \qquad \eta \ge 0 ,$$
(S4)

i.e. the ζ and γ matrices are positive semi-definite. We can rewrite the constitutive relations using n instead of μ as thermodynamic variable, which is more natural if we want to compare our results with [S6]

$$j^{i} = nv^{i} - \left(\bar{\gamma}_{00}P^{ij}\partial_{j}n + \bar{\zeta}_{00}\frac{v^{i}v^{j}}{|v|^{2}}\partial_{j}n + \bar{\gamma}_{01}\frac{P^{i(j}v^{k})}{|v|}\sigma_{jk} + \bar{\zeta}_{01}\frac{v^{i}v^{j}v^{k}}{2|v|^{3}}\sigma_{jk} + \frac{\bar{\zeta}_{02}}{2}\frac{v^{i}}{|v|}P^{jk}\sigma_{jk} + \mathfrak{m}P^{ij}\partial_{j}v^{2}\right) , \quad (S5a)$$

$$\tau^{ij} = \rho v^{i}v^{j} + P\delta^{ij} - \left[\eta \left(P^{k(i}P^{j)l} - \frac{1}{d-1}P^{ij}P^{kl}\right)\sigma_{kl} + 2\tilde{\gamma}_{01}\frac{P^{k(i}v^{j)}}{|v|}\partial_{k}n + \tilde{\zeta}_{01}\frac{v^{i}v^{j}v^{k}}{|v|^{3}}\partial_{k}n + \tilde{\zeta}$$

where we redefined the transport coefficients by absorbing the susceptibilities

$$\bar{\gamma}_{00} = \gamma_{00} \left(\frac{\partial \mu}{\partial n}\right)_{v^2} , \qquad \bar{\gamma}_{01} = \gamma_{01} , \qquad \bar{\zeta}_{00} = \zeta_{00} \left(\frac{\partial \mu}{\partial n}\right)_{v^2} , \qquad (S6a)$$

$$\bar{\zeta}_{01} = \zeta_{01} + 2\zeta_{00} |\vec{v}| \left(\frac{\partial \mu}{\partial v^2}\right)_n, \qquad \bar{\zeta}_{02} = \zeta_{02}, \qquad \mathfrak{m} = \gamma_{00} \left(\frac{\partial \mu}{\partial v^2}\right)_n, \qquad (S6b)$$

$$\tilde{\gamma}_{01} = \gamma_{01} \left(\frac{\partial \mu}{\partial n}\right)_{v^2} , \qquad \tilde{\gamma}_{11} = \gamma_{11} , \qquad \tilde{\zeta}_{01} = \zeta_{01} \left(\frac{\partial \mu}{\partial n}\right)_{v^2} , \qquad (S6c)$$

$$\tilde{\zeta}_{11} = \zeta_{11} + 2\zeta_{01} |\vec{v}| \left(\frac{\partial \mu}{\partial v^2}\right)_n , \qquad \qquad \tilde{\zeta}_{02} = \zeta_{02} \left(\frac{\partial \mu}{\partial n}\right)_{v^2} \qquad \qquad \tilde{\zeta}_{12} = \zeta_{12} \qquad (S6d)$$

$$\tilde{\zeta}_{22} = \zeta_{22} , \qquad \qquad \mathfrak{t} = \gamma_{01} \left(\frac{\partial \mu}{\partial v^2} \right)_n , \qquad \qquad \mathfrak{n} = \zeta_{02} \left(\frac{\partial \mu}{\partial v^2} \right)_n . \tag{S6e}$$

In spite of the fact that there are 10 independent order-one transport coefficients, the equations of motion linearized around a constant background velocity $\vec{v} = v_0 \hat{x}_{\parallel}$ will be such that we can absorb all of them into three order-one diffusive coefficients in the charge conservation equation and four transport coefficients in the momentum equations.

MATCHING TONER AND TU

Consider now the Toner and Tu equations [S6, S7], and focus on the ideal fluid part while setting to zero the noise term \vec{f} and the external potential U:

$$0 = \partial_t n + \vec{\nabla} \cdot (\vec{v}n) \quad , \tag{S7a}$$

$$0 = \partial_t \vec{v} + \lambda_1 (\vec{v} \cdot \vec{\nabla}) \vec{v} + \lambda_2 \vec{v} (\vec{\nabla} \cdot \vec{v}) + \lambda_3 \vec{\nabla} |\vec{v}|^2 + \vec{\nabla} P_1 + \vec{v} (\vec{v} \cdot \vec{\nabla} P_2) .$$
(S7b)

We now ask: is it possible to obtain these equations from the hydrodynamics of passive fluids without boost symmetry?

Consider the ideal-fluid constitutive relations for boost-agnostic hydrodynamics (S2) and the conservation laws (S1). The charge conservation equation naturally agrees with the Toner and Tu model (S7). However, the same cannot be said about the momentum equation, which we write as

$$\partial_t \vec{v} + \vec{v} (\vec{\nabla} \cdot \vec{v}) + (\vec{v} \cdot \vec{\nabla}) \vec{v} + \frac{\vec{\nabla} P}{\rho} + \frac{\vec{v}}{\rho} \partial_t \rho + \frac{\vec{v}}{\rho} \vec{v} \cdot \vec{\nabla} \rho = 0 .$$
(S8)

Using the number conservation equation we can express the last two terms as

$$\partial_t \rho + \vec{v} \cdot \vec{\nabla} \rho = \left(\frac{\partial \rho}{\partial n}\right)_{v^2} \left(\partial_t n + \vec{v} \cdot \vec{\nabla} n\right) + \left(\frac{\partial \rho}{\partial \vec{v}^2}\right)_n \left(\partial_t \vec{v}^2 + \vec{v} \cdot \vec{\nabla} \vec{v}^2\right)$$
$$= -\left(\frac{\partial \rho}{\partial n}\right)_{v^2} n \vec{\nabla} \cdot \vec{v} + \left(\frac{\partial \rho}{\partial \vec{v}^2}\right)_n \left(\partial_t \vec{v}^2 + \vec{v} \cdot \vec{\nabla} \vec{v}^2\right) . \tag{S9}$$

Plugging this expression back into (S8) and projecting the momentum equation along \vec{v} we can solve for $\partial_t \vec{v}^2$. Finally, substituting the solution for $\partial_t \vec{v}^2$ again into the momentum conservation equation we arrive at the final result

$$\partial_t \vec{v} + \lambda_1 (\vec{v} \cdot \vec{\nabla}) \vec{v} + \lambda_2 \vec{v} (\vec{\nabla} \cdot \vec{v}) + \lambda_3 \vec{\nabla} |\vec{v}|^2 + \frac{\vec{\nabla}P}{\rho} - k \frac{\vec{v}}{\rho^2} (\vec{v} \cdot \vec{\nabla}P) = 0 , \qquad (S10)$$

where the values of the parameters are

$$\lambda_1 = 1 , \qquad \qquad \lambda_2 = \frac{1 - \frac{\partial \rho}{\partial n} \frac{n}{\rho}}{1 + 2 \frac{\partial \rho}{\partial v^2} \frac{v^2}{\rho}} , \qquad (S11a)$$

$$\lambda_3 = 0 , \qquad \qquad k = 2 \frac{\partial \rho}{\partial v^2} \frac{1}{1 + 2 \frac{\partial \rho}{\partial v^2} \frac{v^2}{q}} , \qquad (S11b)$$

and the thermodynamic derivatives $\frac{\partial \rho}{\partial \vec{v}^2}$ and $\frac{\partial \rho}{\partial n}$ are to be understood at constant n or \vec{v}^2 respectively. The last two terms in this equation can be identified as the transverse and longitudinal pressures P_1 and P_2 in (S7). However, compared to (S7), we find that boost-agnostic fluids should always have $\lambda_1 = 1$ and $\lambda_3 = 0$.

Even if the Galilean symmetry is broken, the ideal-fluid transport coefficients are mostly constrained to specific values. Physically, this follows from the fact that the ideal-fluid constitutive relations are not just the most general expression compatible with symmetries without derivatives, but are also constrained by the requirement of equilibrium thermodynamics. Then, the constitutive relations for a boost-agnostic fluid are a consequence of having a Gibbs-like equilibrium distribution or, equivalently, they follow from the requirement that the ideal fluid must obey the first law of thermodynamics and not produce entropy.

To conclude, notice that the value of λ_3 is rather arbitrary, both in our equation (S10) and in (S7). This happens because the pressure is a function of \vec{v}^2 . Therefore we can expand the pressure as

$$\frac{\vec{\nabla}P}{\rho} = \frac{1}{\rho} \left(\frac{\partial P}{\partial n}\right)_{v^2} \vec{\nabla}n + \frac{1}{\rho} \left(\frac{\partial P}{\partial \vec{v}^2}\right)_n \vec{\nabla} |\vec{v}|^2 , \qquad (S12)$$

where the first term can be interpreted as a new pressure gradient $\vec{\nabla}\tilde{P}$, while the second term defines $\lambda_3 = \frac{1}{\rho} \left(\frac{\partial P}{\partial \vec{n}^2}\right)_n$.

INTEGRATING OUT THE LONGITUDINAL MODE

We now consider the equations of motion for polar active matter

$$\partial_t g^i + \partial_j \tau^{ji} = U(n, \vec{v}^2) + f^i , \qquad \qquad \partial_t n + \partial_i j^i = 0 , \qquad (S13)$$

with the constitutive relations (S5). We focus on the ordered phase, in which the external potential U causes the fluid to pick a uniform velocity in the non-equilibrium background steady state and thus breaks rotations spontaneously. We expand the velocity as

$$\vec{v} = (v_0 + \delta v_{\parallel})\hat{x}_{\parallel} + \vec{v}_{\perp} , \qquad (S14)$$

4

with δv_{\parallel} and \vec{v}_{\perp} small fluctuations which are respectively longitudinal or orthogonal to the background velocity $v_0 \hat{x}_{\parallel}$. v_0 is defined such that $U(n_0, v_0^2) = 0$, with n_0 the equilibrium particle number density.

In the ordered phase the longitudinal mode associated with δv_{\parallel} is gapped, therefore we can freeze longitudinal fluctuations to find a set of reduced equations only for the Goldstone-like gapless modes \vec{v}_{\perp} and $\delta n = n - n_0$. As argued in the main text, we are interested in linear and non-linear ideal-fluid terms, or linear viscous terms [S6]. On a practical level, this amounts to counting fluctuations and derivative on a similar footing $\mathcal{O}(\delta) \sim \mathcal{O}(\partial) \sim \mathcal{O}(\epsilon)$, with ϵ a counting parameter.

Consider now the equation of motion for the gapped mode δv_{\parallel} . In this equation δv_{\parallel} appears at $\mathcal{O}(1)$, $\mathcal{O}(\partial)$ and $\mathcal{O}(\partial^2)$, however because it only appears at $\mathcal{O}(\partial)$ or higher in the equations of motion for \vec{v}_{\perp} and δn we can focus only on the ideal fluid and external potential part of the equations, which are $\mathcal{O}(1) + \mathcal{O}(\partial)$. The error we are making not including viscous correction to the equation for the heavy mode are at least $\mathcal{O}(\partial^3)$ and can be neglected in our derivative counting.

We introduce a notation for the thermodynamic derivatives of the external potential U, the pressure P and the kinetic mass density ρ

$$\left(\frac{\partial F}{\partial n}\right)_{v^2} = F_n , \qquad \left(\frac{\partial F}{\partial \vec{v}^2}\right)_n = F_v , \qquad \left(\frac{\partial^2 F}{\partial n^2}\right)_{v^2} = F_{n^2} , \qquad (S15)$$

with $F = \{U, P, \rho\}$. These thermodynamic derivatives are evaluated on the background, at $n = n_0$ and $\vec{v}^2 = v_0^2$. Then the equation of motion for the heavy mode δv_{\parallel} , up to $\mathcal{O}(\epsilon^2)$, reads

$$\rho_{0}\partial_{t}\delta v_{\parallel} + \rho_{0}v_{0}\left(2\partial_{\parallel}\delta v_{\parallel} + \vec{\nabla}_{\perp}\cdot\vec{v}_{\perp}\right) + P_{n}\partial_{\parallel}\delta n + 2v_{0}P_{v}\partial_{\parallel}\delta v_{\parallel} + v_{0}\rho_{n}\partial_{t}\delta n + v_{0}^{2}\rho_{n}\partial_{\parallel}\delta n + 2v_{0}^{2}\rho_{v}\partial_{t}\delta v_{\parallel} + 2v_{0}^{3}\rho_{v}\partial_{\parallel}\delta v_{\parallel} = \delta U(v_{0} + \delta v_{\parallel}) + \mathcal{O}(\epsilon^{3}) , \quad (S16)$$

where $\rho_0 = \rho(n_0, v_0^2)$ is the kinetic mass density computed on the background. Everything is linear in fluctuations to this point, because all the terms are at least order one in derivatives. The non-linearities appear in the fluctuations of the external potential, which we expand as

$$\delta U = U_n \delta n + U_v \left(2v_0 \delta v_{\parallel} + (\delta v_{\parallel})^2 + |\vec{v}_{\perp}|^2 \right) + U_{n^2} (\delta n)^2 + \mathcal{O}(\delta^3) .$$
(S17)

In this expansion, notice in particular that the $(\delta v_{\parallel})^2$ and $(\delta n)^2$ terms are missing from the analysis of [S7]. However, as we will see, this only adjusts the values of certain bare coefficients in the final equation.

We can now solve equation (S16) iteratively in δv_{\parallel} in terms of \vec{v}_{\perp} and δn . At lowest order in fluctuations and derivatives the solution is simply given by

$$\delta v_{\parallel} \approx -\frac{U_n}{U_v} \frac{\delta n}{2v_0} + \mathcal{O}(\epsilon^2) .$$
(S18)

We can subsequently plug this approximate solution back into the derivative and non-linear terms in (S16) and solve again for δv_{\parallel} . The complete solution is

$$\begin{split} \delta v_{\parallel} &\approx -\frac{U_n}{U_v} \frac{\delta n}{2v_0} - \left(\frac{\rho_0}{2v_0} \frac{U_n}{U_v} - v_0 \rho_n + v_0 \rho_v \frac{U_n}{U_v}\right) \frac{\partial_t \delta n}{2v_0^2 U_v} \\ &- \left(\rho_0 \frac{U_n}{U_v} - v_0^2 \rho_n - P_n + P_v \frac{U_n}{U_v} + v_0^2 \rho_v \frac{U_n}{U_v}\right) \frac{\partial_{\parallel} \delta n}{2v_0^2 U_v} \\ &+ \frac{\rho_0}{2v_0 U_v} \vec{\nabla}_{\perp} \cdot \vec{v}_{\perp} - \frac{|\vec{v}_{\perp}|^2}{2v_0} - \left(\frac{1}{8v_0^3} \frac{U_n^2}{U_v^2} + \frac{U_{n^2}}{2v_0}\right) (\delta n)^2 + \mathcal{O}(\epsilon^3) \,. \end{split}$$
(S19)

Finally, we can use the linearized ideal-fluid charge conservation equation

$$\partial_t \delta n + v_0 \partial_{\parallel} \delta n + n_0 \vec{\nabla}_{\perp} \cdot \vec{v}_{\perp} + n_0 \partial_{\parallel} \delta v_{\parallel} = 0 , \qquad (S20)$$

together with the order-zero solution (S18), to eliminate the time derivative in favour of spatial gradients. The final solution for δv_{\parallel} in terms of the other fields is

$$\delta v_{\parallel} = -\alpha_1 \delta n - \alpha_2 (\delta n)^2 - \alpha_3 \vec{\nabla}_{\perp} \cdot \vec{v}_{\perp} - \alpha_4 \partial_{\parallel} \delta n - \frac{|\vec{v}_{\perp}|^2}{2v_0} + \mathcal{O}(\epsilon^3) , \qquad (S21)$$

where we introduced the thermodynamic quantities

$$\alpha_1 = \frac{1}{2v_0} \frac{U_n}{U_v} , \qquad \qquad \alpha_3 = \frac{\rho_n n_0}{2v_0 U_v} - \frac{\rho_0}{2v_0 U_v} - \frac{\rho_v n_0}{2v_0} \frac{U_n}{U_v^2} - \frac{n_0 \rho_0}{4v_0^3} \frac{U_n}{U_v^2} , \qquad (S22a)$$

$$\alpha_2 = \frac{1}{8v_0^3} \frac{U_n^2}{U_v^2} + \frac{U_{n^2}}{2v_0} , \qquad \alpha_4 = \frac{\rho_0}{4v_0^2} \frac{U_n}{U_v^2} - \frac{U_n}{U_v^2} \frac{\rho_n n_0}{4v_0^2} - \frac{P_n}{2v_0^2 U_{v^2}} + \frac{U_n}{U_v^2} \frac{P_v}{2v_0^2} + \frac{U_n^2}{U_v^3} \frac{\rho_v n_0}{4v_0^2} + \frac{U_n^2}{U_v^3} \frac{n_0 \rho_0}{8v_0^4} .$$
 (S22b)

With this result, we can now expand the charge conservation equation up to $\mathcal{O}(\epsilon^3)$ and substitute the above expression for the heavy mode. The final equation becomes

$$\partial_t \delta n + v_2 \partial_{\parallel} \delta n + n_0 \vec{\nabla}_{\perp} \cdot \vec{v}_{\perp} + w_1 \vec{\nabla}_{\perp} (\vec{v}_{\perp} \delta n) - w_2 \partial_{\parallel} (\delta n)^2 - w_3 \partial_{\parallel} |\vec{v}_{\perp}|^2 - \xi_n \partial_{\parallel} \left(\vec{\nabla}_{\perp} \cdot \vec{v}_{\perp} \right) - D_{\parallel} \partial_{\parallel}^2 \delta n - D_{\perp} \nabla_{\perp}^2 \delta n = 0 ,$$
(S23)

with the following definitions

$$v_2 = v_0 - \alpha_1 n_0 , \qquad \qquad w_1 = 1 , \qquad (S24)$$

$$w_2 = \alpha_1 + \alpha_2 n_0 , \qquad \qquad w_3 = \frac{n_0}{2v_0} , \qquad (S25)$$

$$\xi_n = \alpha_3 n_0 + \bar{\gamma}_{01} + \bar{\zeta}_{02} , \qquad \qquad D_{\parallel} = n_0 \alpha_4 + \bar{\zeta}_{00} - \alpha_1 \bar{\zeta}_{01} , \qquad (S26)$$

$$D_{\perp} = \bar{\gamma}_{00} - \alpha_1 (2v_0 \mathfrak{m} + \bar{\gamma}_{01}) . \tag{S27}$$

Similarly, we can substitute in the equation of motion for the transverse velocity fluctuations \vec{v}_{\perp} . Doing so one finds a non-linear term of the form $\delta n \partial_t \vec{v}_{\perp}$, which can again be written in terms of spatial gradients using the ideal-fluid linearized equations of motion for \vec{v}_{\perp} . The final result is

$$\partial_{t}\vec{v}_{\perp} + v_{0}\partial_{\parallel}\vec{v}_{\perp} + \kappa\vec{\nabla}_{\perp}\delta n + g_{1}\delta n\partial_{\parallel}\vec{v}_{\perp} + g_{2}\vec{v}_{\perp}\partial_{\parallel}\delta n + g_{3}\vec{\nabla}_{\perp}(\delta n)^{2} + g_{4}\vec{v}_{\perp}\left(\vec{\nabla}_{\perp}\cdot\vec{v}_{\perp}\right) + g_{5}\left(\vec{v}_{\perp}\cdot\vec{\nabla}_{\perp}\right)\vec{v}_{\perp} - D_{v}\vec{\nabla}_{\perp}\partial_{\parallel}\delta n - \xi_{v}\vec{\nabla}_{\perp}\left(\vec{\nabla}_{\perp}\cdot\vec{v}_{\perp}\right) - \xi_{\perp}\nabla_{\perp}^{2}\vec{v}_{\perp} - \xi_{\parallel}\partial_{\parallel}^{2}\vec{v}_{\perp} = \vec{f}_{\perp} , \quad (S28)$$

with the coefficients given by

$$g_4 = 1 - \frac{1}{\rho_0} \left(n_0 \rho_n - 2v_0 n_0 \alpha_1 \rho_v - 2v_0 \alpha_3 U_v \right) , \qquad g_5 = 1 , \qquad (S29c)$$

$$D_{v} = \frac{1}{\rho_{0}} \left(2v_{0}\alpha_{4}P_{v} + \tilde{\gamma}_{01} + \tilde{\zeta}_{02} - \alpha_{1}(2v_{0}(\mathfrak{t} + \mathfrak{n}) + \tilde{\gamma}_{11} + \tilde{\zeta}_{12}) \right) , \quad \xi_{v} = \frac{1}{\rho_{0}} \left(2v_{0}\alpha_{3}P_{v} + \tilde{\zeta}_{22} \right) , \quad (S29d)$$

$$\xi_{\parallel} = \frac{\tilde{\gamma}_{11}}{\rho_0} , \qquad \qquad \xi_{\perp} = \frac{\eta}{\rho_0} . \tag{S29e}$$

The overall form of these reduced equations (S28) and (S23) agrees with the result found in [S7], even if the values of the coefficients are different, with a few minor differences. First, observe that we have a new non-linear term g_4 , which is missing in [S7], secondly we find that the bare coefficient D_{\perp} is non-zero, while it appears to be zero in [S7], finally, $g_5 = 1$ despite the lack of Galilean invariance.

- [S3] L. Onsager, Phys. Rev. **37**, 405 (1931).
- [S4] L. Onsager, Phys. Rev. 38, 2265 (1931).
- [S5] P. Kovtun, J. Phys. A 45, 473001 (2012), arXiv:1205.5040 [hep-th].
- [S6] J. Toner and Y. Tu, Phys. Rev. E 58, 4828 (1998).
- [S7] J. Toner, Phys. Rev. E 86, 031918 (2012)

[[]S1] J. de Boer, J. Hartong, E. Have, N. A. Obers, and W. Sybesma, SciPost Phys. 9, 018 (2020), arXiv:2004.10759 [hep-th].

[[]S2] J. Armas and A. Jain, SciPost Phys. 11, 054 (2021), arXiv:2010.15782 [hep-th].