ON POINTWISE CONVERGENCE OF CONE MULTIPLIERS

PENG CHEN, DANQING HE, XIAOCHUN LI, AND LIXIN YAN

ABSTRACT. For $p \ge 2$, and $\lambda > \max\{n|\frac{1}{p} - \frac{1}{2}| - \frac{1}{2}, 0\}$, we prove the pointwise convergence of cone multipliers, i.e.

$$\lim_{t\to\infty}T_t^{\lambda}(f)\to f \text{ a.e.},$$

where $f \in L^p(\mathbb{R}^n)$ satisfies supp $\widehat{f} \subset \{\xi \in \mathbb{R}^n : 1 < |\xi_n| < 2\}$. Our main tools are weighted estimates for maximal cone operators, which are consequences of trace inequalities for cones.

1. INTRODUCTION

The Bochner-Riesz operator is defined by

$$B^{\lambda}(f)(x) = \int_{\mathbb{R}^n} (1 - |\xi|^2)^{\lambda}_+ \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$

for $\lambda > 0$. It has been conjectured that B^{λ} is bounded on $L^{p}(\mathbb{R}^{n})$ when $\lambda > \max\{|\frac{n}{p} - \frac{n}{2}| - \frac{1}{2}, 0\}$. The conjecture was resolved when n = 2 by [11]; later [35] gave another proof. Due to its significant connection with the restriction theorem and other conjectures in harmonic analysis, the Bochner-Riesz conjecture has been extensively studied over the past decades; see [2], [16], [17], [18], [4], [44], [37], [25] and references therein.

A cone multiplier in \mathbb{R}^n for $\lambda > 0$ is defined as

(1.1)
$$\tilde{m}^{\lambda}(\xi) = (1 - \frac{|\xi'|^2}{\xi_n^2})^{\lambda}_+,$$

where $\xi = (\xi', \xi_n) \in \mathbb{R}^n$, and the related operator is defined by

$$\widetilde{T}^{\lambda}(f)(x) := \int_{\mathbb{R}^n} \widetilde{m}^{\lambda}(\xi) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi.$$

D. He is supported by National Key R&D Program of China (No. 2021YFA1002500), NNSF of China (No. 12322105), the New Cornerstone Science Foundation, and Natural Science Foundation of Shanghai (No. 22ZR1404900 and 23QA1400300). P. Chen was supported by NNSF of China 12171489 and Guangdong Natural Science Foundation 2022A1515011157. X. Li is partially supported by Simons foundation. L. Yan was supported by National Key R&D Program of China 2022YFA1005700.

It was Stein [43] who introduced this operator. He conjectured that the range of boundedness of \tilde{T}^{λ} in \mathbb{R}^{n} coincides with that of B^{λ} is \mathbb{R}^{n-1} , as suggested by de Leeuw's theorem [13].

On the study of Stein's conjecture in \mathbb{R}^3 , Mockenhaupt [40] reduced it to a reverse square function estimate, initially obtaining a non-optimal result using a geometric observation on the distribution of plates. Bourgain [3] later improved Mockenhaupt's result by introducing the so-called bilinear method. This method was systematically developed to study the restriction theorem and related conjectures; for further details, see [47], [48], [1], [7], [38], [28], and [42]. The reverse square function estimate also implies the well-known local smoothing estimates; see [41]. To tackle these problems, Wolff [49] introduced decoupling inequalities, which have become a crucial topic in modern harmonic analysis, appearing naturally in various fields from PDEs to number theory; see [36], [22], [33], [5], [6], [14], [15], [29], [27] and references therein. The sharp reverse square function estimate in \mathbb{R}^3 was recently obtained by Guth, Wang and Zhang [30], thereby resolving the cone multiplier conjecture and the local smoothing conjecture in \mathbb{R}^3 , which was generalized to the variable coefficients setting in [21]. More results and methods on the boundedness of cone multipliers can be found in [34], [31], and [32].

In this exposition, we shift our focus to the pointwise convergence of cone multipliers, which is connected to maximal estimates of cone multipliers. Pointwise convergence of Fourier series and Fourier integrals is a central issue in harmonic analysis. For classical results, one may refer to [50]. According to Stein's maximal principle [45], pointwise convergence problems are consequences of the boundedness problems of corresponding maximal operators. For instance, Carleson [10] demonstrated the boundedness of the Carleson operator, resolving the long-standing Lusin's conjecture regarding the pointwise convergence of Dirichlet means in \mathbb{R} ; a different proof was given by Fefferman [19]. Carbery [8] established the $L^4(\mathbb{R}^2)$ boundedness of maximal Bochner-Riesz operator. For recent progress, interested readers are directed to [46], [37], and [39].

We formulate our problem as follows. Since $\tilde{m}^{\lambda}(t^{-1}\xi) = \tilde{m}^{\lambda}(\xi)$, we dilate \tilde{m}^{λ} anisotropically by considering $\tilde{m}(t^{-1}\xi', \xi_n)$ for t > 0, which converges to $\chi_{\mathbb{R}^n}(\xi)$ almost everywhere as $t \to \infty$. This observation, combined with the Lebesgue dominated convergence theorem, implies that

(1.2)
$$\widetilde{T}_t^{\lambda}(f)(x) := \int_{\mathbb{R}^n} \widetilde{m}^{\lambda}(t^{-1}\xi',\xi_n) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$

converges to f(x) for all $x \in \mathbb{R}^n$ as $t \to \infty$ when f is a Schwartz function. It is natural to inquire about the sharp range of p such that

(1.3)
$$\lim_{t \to \infty} \tilde{T}_t^{\lambda}(f)(x) = f(x) \quad \text{a.e.} \quad \forall f \in L^p(\mathbb{R}^n)$$

By Stein's maximal principle, this result can be reduced to the study of the maximal cone operator

$$\tilde{T}_*^{\lambda}(f)(x) := \sup_{t>0} |\tilde{T}_t^{\lambda}(f)(x)|.$$

Since $\tilde{T}^{\lambda}(f) \leq \tilde{T}^{\lambda}_{*}(f)$, concerning the pointwise convergence and maximal estimates of cone multipliers, we pose the following conjecture.

Conjeture 1. Let $n \ge 3$, and $p \ge 2$. The range of p such that (1.3) holds is the same as the range of p such that $\|\tilde{T}^{\lambda}\|_{L^p(\mathbb{R}^n)\to L^p(\mathbb{R}^n)} < \infty$, i.e. $\lambda > \max\{(n-1)|\frac{1}{p}-\frac{1}{2}|-\frac{1}{2},0\}$. Moreover, $\|\tilde{T}^{\lambda}_*\|_{L^p(\mathbb{R}^n)\to L^p(\mathbb{R}^n)} < \infty$ when $\lambda > \max\{(n-1)|\frac{1}{p}-\frac{1}{2}|-\frac{1}{2},0\}$.

A corresponding conjecture for p < 2 could also be proposed, but we will not delve into it here as the boundedness of the maximal Bochner-Riesz operator is not clear even in the plane. For the pointwise convergence of \tilde{T}_{t}^{λ} , we derive the following restricted result.

Theorem 1.1. Let $n \ge 3$, $p \in [2, \infty)$, and $\lambda > \max\{|\frac{n}{p} - \frac{n}{2}| - \frac{1}{2}, 0\}$. Suppose that $f \in L^p(\mathbb{R}^n)$ satisfies that

(1.4)
$$supp \ \widehat{f} \subset \{ \xi \in \mathbb{R}^n : \ a < |\xi_n| < b \}$$

for some $0 < a < b < \infty$. Then $\tilde{T}_t^{\lambda}(f) \to f a.e.$ as $t \to \infty$.

Remark 1. This result is non-optimal compared with the conjectured range $\lambda > \max\{(n-1)|\frac{1}{p} - \frac{1}{2}| - \frac{1}{2}, 0\}$. In particular, when λ is greater than the conjectured critical index (n-2)/2, whether the pointwise convergence holds for all $p \in [2, \infty)$ is still open.

Theorem 1.1 follows obviously from the case a = 1 and b = 2. Therefore, we introduce the following notation:

$$m^{\lambda}(\xi) := (1 - \frac{|\xi'|^2}{\xi_n^2})^{\lambda}_+ \psi(2^{-1}\xi_n),$$

where ψ is a nonnegative smooth function supported in $[\frac{1}{4}, 1]$ such that

$$\sum_{\gamma \in \mathbb{Z}} \psi(2^{\gamma} t) = 1$$

for t > 0. We can define T_t^{λ} and T_*^{λ} correspondingly. For $f \in \mathcal{S}(\mathbb{R}^n)$ we define

$$T_t^{\lambda}(f)(x) := \int_{\mathbb{R}^n} m^{\lambda}(t^{-1}\xi',\xi_n)\widehat{f}(\xi)e^{2\pi i x \cdot \xi}d\xi,$$

$$T_*^{\lambda}(f)(x) := \sup_{t>0} |T_t^{\lambda}(f)(x)|,$$

and

$$L(f)(x) = \int_{\mathbb{R}^n} \Psi(2^{-1}\xi_n)\widehat{f}(\xi)e^{2\pi i x \cdot \xi}d\xi.$$

We can actually prove the following result.

Theorem 1.2. Let $n \ge 3$, $p \in [2, \infty)$, and $\lambda > \max\{|\frac{n}{p} - \frac{n}{2}| - \frac{1}{2}, 0\}$. Then, for any $f \in L^p(\mathbb{R}^n)$, $T_t^{\lambda}(f) \to Lf$ a.e. as $t \to \infty$.

Instead of focusing on the L^p boundedness of maximal Bochner-Riesz operators, Carbery, Rubio de Francia, and Vega [9] demonstrated the pointwise convergence of Bochner-Riesz means in all higher dimensions when $p \ge 2$ by establishing weighted L^2 estimates of maximal Bochner-Riesz operators. This approach proves to be effective in various problems; for instance, Chen, Duong, He, Lee, and Yan [12] adopt this strategy to establish the pointwise convergence of Bochner-Riesz means for Hermit operators. We will employ it to demonstrate that T_*^{λ} is bounded on some appropriately weighted L^2 spaces. Specifically, due to the anisotropic dilation in the definition of T_*^{λ} , we utilize the homogeneous weights given by

(1.5)
$$w_{\alpha,\beta}(x) := |x'|^{-\alpha} |x_n|^{-\beta}$$

where $x = (x', x_n) \in \mathbb{R}^{n-1} \times \mathbb{R}$. However, these weights introduce some challenging technical complexities. For example, we need to make a careful decompose wherein an L^p function is dissembled into four pieces in Proposition 2.1, and the associated trace inequalities become more intricate. Nevertheless, we can establish the following result.

Theorem 1.3. *Let* $n \ge 3$, $\alpha \in [0, n-1)$, $\beta \in [0, 1)$, *and* $\lambda > \max\{\frac{\alpha+\beta-1}{2}, 0\}$. *Then*

(1.6)
$$\|T_*^{\lambda}(f)\|_{L^2(w_{\alpha,\beta})} \le C \|f\|_{L^2(w_{\alpha,\beta})}$$

for any Schwartz function f. As a result, for any $f \in L^2(w_{\alpha,\beta})$, we have $T_t^{\lambda}(f) \to Lf$ a.e. as $t \to \infty$.

This result constitutes the primary technical contribution of this study. Since the trace inequalities represent weighted restriction estimates, which play a crucial role in establishing weighted estimates of Bochner-Riesz means as demonstrated in [9], we aim to transform the inequality (1.6) into analogous trace inequalities of cones. However, this transformation necessitates a nuanced argument that relies on the spatial distribution of f, owing to the anisotropic dilation inherent in the definition of maximal cone operators; please refer to Section 5 for a detailed explanation. To establish

the desired trace inequalities, we make use of geometric properties of cones with the help of an argument inspired by the work in [9].

This paper is organized as follows. In Section 2, we decompose an arbitrary L^p function into $L^2(w_{\alpha,\beta})$ pieces, where we have to keep the Fourier support properties for technical reasons. Section 3 discusses the reduction to square functions. In Section 4 and Section 5, we establish necessary trace inequalities for cones, which lead to desired weighted inequalities for T_*^{λ} via a delicate argument in term of the physical position of f. The proof of main theorems are given in Section 6.

Notations

By $A \lesssim B$ we mean that there exists an absolute constant *C* such that $A \leq CB$. $A \sim B$ if $A \lesssim B$ and $B \lesssim A$.

 $E + O(\delta) := \{x \in \mathbb{R}^n : dist(x, E) < C\delta\}\}$ is the *C* δ -neighborhood of *E* in \mathbb{R}^n , where *C* is an unimportant absolute constant.

 $Q(x,\ell)$ is the cube centered at $x \in \mathbb{R}^n$ with length ℓ . We denote by $NQ(x,\ell)$ the cube $Q(x,N\ell)$.

Given a multiplier $m \in L^{\infty}(\mathbb{R}^n)$, the associated linear operator T_m is defined as

$$T_m(f)(x) = \int_{\mathbb{R}^n} m(\xi) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi.$$

 $\mathcal{S}(\mathbb{R}^n)$ is the class of Schwartz functions.

2. DECOMPOSING FUNCTIONS

We decompose an arbitrary function $f \in L^p(\mathbb{R}^n)$ in terms of the weights $w_{\alpha,\beta}$ defined in (1.5) with the Fourier support remained, therefore, to obtain the pointwise convergence of cone multiplies, it is natural to consider weighted L^2 estimates of T_*^{λ} . The proof is standard, but we include it for the sake of completeness.

Proposition 2.1. Let $p \ge 2$, $\alpha_1 = \beta_1 = 0$, $0 \le \alpha_3 < (n-1)(1-\frac{2}{p}) < \alpha_2, \alpha_4 < 1 + (n-1)(1-\frac{2}{p})$, and $0 \le \beta_2 < 1-\frac{2}{p} < \beta_3, \beta_4 < 2-\frac{2}{p}$. For any $f \in L^p(\mathbb{R}^n)$, we can decompose it into

$$f = f_1 + f_2 + f_3 + f_4$$

such that $f_i \in L^2(w_i)$, where $w_i(x) = |x'|^{-\alpha_i} |x_n|^{-\beta_i}$ for i = 1, 2, 3, 4. Moreover, if supp $\widehat{f} \subset S = \{\xi \in \mathbb{R}^n : \xi_n \in (\frac{1}{2}, 2)\}$, then

(2.1)
$$supp \ \widehat{f_i} \subset S' = \{ \xi \in \mathbb{R}^n : \ \xi_n \in (\frac{1}{10}, 10) \}$$

for i = 1, 2, 3, 4.

Proof. We take a function $\varphi_k \in S(\mathbb{R}^k)$ such that supp $\widehat{\varphi}_k \subset B(0, 10^{-10})$ and $\varphi_k(0) = 1$.

We define

$$f_1(x) = f(x)\varphi_{n-1}(x')\varphi_1(x_n)$$

$$f_2(x) = f(x)(1 - \varphi_{n-1})(x')\varphi_1(x_n)$$

$$f_3(x) = f(x)\varphi_{n-1}(x')(1 - \varphi_1)(x_n)$$

$$f_4(x) = f(x)(1 - \varphi_{n-1})(x')(1 - \varphi_1)(x_n).$$

Obviously $f = f_1 + f_2 + f_3 + f_4$, and, when supp $\hat{f} \subset S$, (2.1) can be verified easily using that supp $\hat{\varphi}_k \subset B(0, 10^{-10})$.

Since $p \ge 2$, by Hölder's inequality, we obtain

$$\int_{\mathbb{R}^n} |f_1(x)|^2 w_1(x) dx = \int_{\mathbb{R}^n} |f(x)\varphi_{n-1}(x')\varphi_1(x_n)|^2 dx$$
$$\leq C ||f||_{L^p(\mathbb{R}^n)}^2.$$

Let us analyze f_4 below. By Hölder's inequality, we control $||f_4||^2_{L^2(w_4)}$ by

(2.2)
$$||f||_{L^p}^2 \left(\int_{\mathbb{R}^n} ((1-\varphi_{n-1})(x')|x'|^{-\alpha_4}(1-\varphi_1)(x_n)|x_n|^{-\beta_4})^{\frac{p}{p-2}} dx \right)^{\frac{p-2}{p}}.$$

As $\varphi_{n-1}(0) = 1$, we have

$$|1-\varphi_{n-1}(x')| \leq C|x'|.$$

This implies

$$\begin{split} &\int_{\mathbb{R}^{n-1}} ((1-\varphi_{n-1})(x')|x'|^{-\alpha_4})^{p/(p-2)} dx' \\ &\leq \int_{|x'|\leq 1} |x'|^{(1-\alpha_4)\frac{p}{p-2}} dx' + \int_{|x'|\geq 1} |x'|^{-\alpha_4\frac{p}{p-2}} dx' \\ &\lesssim \int_0^1 r^{(1-\alpha_4)\frac{p}{p-2}+n-2} dr + \int_1^\infty r^{n-2-\alpha_4\frac{p}{p-2}} dr, \end{split}$$

which is finite since $(n-1)(1-\frac{2}{p}) < \alpha_4 < 1 + (n-1)(1-\frac{2}{p})$. Similarly

$$\int_{\mathbb{R}} ((1-\varphi_1)(x_n)|x_n|^{-\beta_4})^{p/(p-2)} dx_n < \infty$$

when $\beta_4 \in (1 - \frac{2}{p}, 2 - \frac{2}{p})$. These estimates and (2.2) show that $\|f_4\|_{2^2} \leq C \|f\|_{L^p}$

$$||f_4||_{L^2(w_4)} \le C ||f||_{L^p}$$

One can also obtain similarly that

$$||f_2||_{L^2(w_2)}^2 = \int_{\mathbb{R}^n} |f_2(x)|^2 |x'|^{-\alpha_2} |x_n|^{-\beta_2} dx$$

$$\leq C \|f\|_{L^{p}}^{2} \left(\int_{\mathbb{R}^{n}} ((1 - \varphi_{n-1})(x')|x'|^{-\alpha_{2}} \varphi_{1}(x_{n})|x_{n}|^{-\beta_{2}})^{\frac{p}{p-2}} dx \right)^{\frac{p-2}{p}} \\ < C \|f\|_{L^{p}}^{2}$$

when $(n-1)(1-\frac{2}{p}) < \alpha_2 < 1 + (n-1)(1-\frac{2}{p})$ and $\beta_2 \frac{p}{p-2} < 1$. The proof for $f_3 \in L^2(w_3)$ is similar.

3. REDUCTION TO SQUARE FUNCTIONS

We recall that ψ is a smooth bump supported in $[\frac{1}{4}, 1]$ such that $\sum_{\gamma \in \mathbb{Z}} \psi(2^{\gamma}t) =$ 1 for *t* > 0,

$$m^{\lambda}(\xi) = \left(1 - \frac{|\xi'|^2}{\xi_n^2}\right)^{\lambda}_+ \Psi(2^{-1}\xi_n),$$
$$T_t^{\lambda}(f)(x) = \int_{\mathbb{R}^n} m^{\lambda}(t^{-1}\xi',\xi_n)\widehat{f}(\xi)e^{2\pi ix\cdot\xi}d\xi,$$

and

$$T^{\lambda}_*(f)(x) = \sup_{t>0} |T^{\lambda}_t(f)(x)|.$$

For given $\alpha, \beta \ge 0$ and $w_{\alpha,\beta}(x) = |x'|^{-\alpha} |x_n|^{-\beta}$, we will explore the range of λ such that T_*^{λ} is bounded on $L^2(w_{\alpha,\beta})$. We define, for $\gamma \ge 1$,

$$m_{\gamma}(\xi) = \psi(2^{\gamma}(1-\frac{|\xi'|^2}{\xi_n^2}))m^{\lambda}(\xi)2^{\gamma\lambda},$$

and

$$m_0(\xi) = m^{\lambda}(\xi) (1 - \sum_{\gamma \ge 1} \psi(2^{\gamma}(1 - \frac{|\xi'|^2}{\xi_n^2}))).$$

We remark that m_0 is supported in

$$\{\xi \in \mathbb{R}^n : |\xi'| / \xi_n \in [0,1], \, \xi_n \in [\frac{1}{2},2]\},\$$

and m_{γ} is supported in

{
$$\xi \in \mathbb{R}^n : |\xi'| / \xi_n \in [1 - 2^{-\gamma}, 1 - 2^{-\gamma-1}], \xi_n \in [\frac{1}{2}, 2]$$
}

for $\gamma \geq 1$. Obviously

(3.1)
$$T_*^{\lambda}(f)(x) \le \sum_{\gamma \ge 0} 2^{-\gamma \lambda} M_{\gamma}(f)(x),$$

where

$$M_{\gamma}(f)(x) = \sup_{t>0} \left| \int_{\mathbb{R}^n} m_{\gamma}(t^{-1}\xi',\xi_n) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi \right|$$

Concerning the weighted boundedness of M_{γ} , we can easily prove a nonoptimal result.

Proposition 3.1. *For* $\gamma \ge 0$, $\alpha \in (-(n-1), (n-1))$, *and* $\beta \in (-1, 1)$,

(3.2)
$$\|M_{\gamma}(f)\|_{L^2(w_{\alpha,\beta})} \leq C_{\gamma}\|f\|_{L^2(w_{\alpha,\beta})}.$$

To prove this proposition, we recall first some definitions and results.

We denote by Q_k a cube in \mathbb{R}^k , and M_k the standard Hardy-Littlewood maximal function in \mathbb{R}^k . A strong maximal function (on $\mathbb{R}^{n-1} \times \mathbb{R}$):

$$M_S(f)(x) = \sup_{R \ni x} \frac{1}{|R|} \int_R |f(y)| dy,$$

where the supremum runs over all *R* of the form $Q_{n-1} \times Q_1$. It is well known that M_S is bounded on $L^p(\mathbb{R}^n)$ for 1 since

(3.3)
$$M_{\mathcal{S}}(f)(x',x_n) \leq (M_1 \circ M_{k-1})(f)(x',x_n).$$

A weight *w* belongs to the (product) $A_p(\mathbb{R}^{n-1} \times \mathbb{R})$ if there exists a finite constant C_w such that for all $R = Q_{n-1} \times Q_1$,

(3.4)
$$\left(\frac{1}{|R|}\int_{R}w(x',x_n)dx'dx_n\right)\left(\frac{1}{|R|}\int_{R}w(x',x_n)^{-1/(p-1)}dx'dx_n\right)^{p-1} \leq C_w.$$

One can easily verify that $w_{\alpha,\beta}$ is an $A_2(\mathbb{R}^{n-1} \times \mathbb{R})$ weight if and only if $\alpha \in (-(n-1), n-1)$ and $\beta \in (-1, 1)$. Since M_k is bounded on $L^2(w)$ when w is an $A_2(\mathbb{R}^k)$ weight, it follows from (3.3) that

(3.5)
$$\|M_{\mathcal{S}}(f)\|_{L^{2}(w_{\alpha,\beta})} \leq C \|f\|_{L^{2}(w_{\alpha,\beta})}$$

when $\alpha \in (-(n-1), n-1)$ and $\beta \in (-1, 1)$.

Proof of Propositio 3.1. As $m_{\gamma}(\xi)$ is a compactly supported smooth function, it is routine to verify that

$$M_{\gamma}(f)(x) \leq C_{\gamma} M_S(f)(x),$$

which implies further (3.2) by (3.5).

The boundedness of T^{λ}_* on $L^2(w_{\alpha,\beta})$ is now reduced to estimating the norm $||M_{\gamma}||_{L^2(w_{\alpha,\beta})\to L^2(w_{\alpha,\beta})}$ as $\gamma\to\infty$, since we have proved that M_{γ} is bounded on $L^2(w_{\alpha,\beta})$ for all $\gamma \ge 0$.

We will focus on the case $\gamma \ge 1$ from now on.

By the Sobolev embedding, we obtain further that

$$M_{\gamma}(f)(x)^2 \le 2^{\gamma} \mathcal{G}_{\gamma}(f)(x) \tilde{\mathcal{G}}_{\gamma}(f)(x),$$

where

$$\mathcal{G}_{\gamma}(f)(x) := \left(\int_0^\infty \left|\int_{\mathbb{R}^n} m_{\gamma}(t^{-1}\xi',\xi_n)\widehat{f}(\xi)e^{2\pi ix\cdot\xi}d\xi\right|^2 \frac{dt}{t}\right)^{1/2},$$

and

$$\tilde{\mathcal{G}}_{\gamma}(f)(x) := \left(\int_0^\infty \left|\int_{\mathbb{R}^n} \tilde{m}_{\gamma}(t^{-1}\xi',\xi_n)\widehat{f}(\xi)e^{2\pi i x \cdot \xi}d\xi\right|^2 \frac{dt}{t}\right)^{1/2}$$

Here $\tilde{m}_{\gamma}(\xi',\xi_n) = (2^{-\gamma}\xi') \cdot (\nabla_{n-1}m_{\gamma})(\xi',\xi_n)$, and $\nabla_{n-1}m_{\gamma}(\cdot,\xi_n)$ is the gradient of $m_{\gamma}(\cdot,\xi_n)$ as a function defined on \mathbb{R}^{n-1} . In summary we obtain

(3.6)
$$T_*^{\lambda}(f)(x) \le M_0(f)(x) + \sum_{\gamma \ge 1} 2^{-\gamma\lambda} 2^{\gamma/2} [\mathcal{G}_{\gamma}(f)(x) \tilde{\mathcal{G}}_{\gamma}(f)(x)]^{1/2}.$$

For $0 < \delta \ll 1$, we fix a smooth function $\mu_{\delta}(\xi')$ such that

(3.7)
$$\operatorname{supp} \mu_{\delta} \subset \{ \xi' \in \mathbb{R}^{n-1} : |\xi'| \in [1-\delta, 1] \}$$

and

$$(3.8) |\partial^l \mu_{\delta}(\xi')| \le C_l \delta^{-|l|}$$

for any $l = (l_1, ..., l_{n-1}) \in \mathbb{N}^{n-1}$. Associated with μ_{δ} , we define the multiplier

$$m_{\delta}(\xi',\xi_n)=\mu_{\delta}(\xi'/\xi_n)\psi(2^{-1}\xi_n),$$

and

$$\mathbf{G}_{\delta}(f)(x) := \left(\int_0^\infty \left| \int_{\mathbb{R}^n} m_{\delta}(t^{-1}\xi',\xi_n) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi \right|^2 \frac{dt}{t} \right)^{1/2}$$

As the multipliers m_{γ} and \tilde{m}_{γ} are multipliers of this form with $\delta = 2^{-\gamma}$, the problem is now reduced to studying the boundedness of \mathbf{G}_{δ} .

Let us examine a simple case first.

Proposition 3.2. Let $\lambda > 0$. Then

(3.9)
$$\|\mathbf{G}_{\delta}(f)\|_{L^{2}(\mathbb{R}^{n})} \leq \delta^{1/2} \|f\|_{L^{2}(\mathbb{R}^{n})}$$

and T^{λ}_* is bounded on $L^2(\mathbb{R}^n)$.

Proof. By Plancherel's identity, $\|\mathbf{G}_{\delta}(f)\|_{L^{2}(\mathbb{R}^{n})}^{2}$ is equal to

$$\int_{\mathbb{R}^n} \int_0^\infty |m_\delta(t^{-1}\xi',\xi_n)|^2 \frac{dt}{t} |\widehat{f}(\xi)|^2 d\xi.$$

For a fixed ξ , the *t* such that $m_{\delta}(t^{-1}\xi', \xi_n) \neq 0$ is contained in $\frac{|\xi'|}{\xi_n} + O(\frac{|\xi'|}{\xi_n}\delta)$, hence the inner integral is bounded by $C\delta$, which implies (3.9).

Recalling (3.6) and Proposition 3.1, we obtain from (3.9) that

$$\|T_*^{\lambda}(f)\|_{L^2(\mathbb{R}^n)} \le \sum_{\gamma \ge 0} 2^{-\gamma\lambda} 2^{\gamma/2} 2^{-\gamma/2} \|f\|_{L^2(\mathbb{R}^n)} \le C \|f\|_{L^2(\mathbb{R}^n)}$$

when $\lambda > 0$.

We study the boundedness of \mathbf{G}_{δ} on $L^2(w_{\alpha,\beta})$ below. Let us fix $\delta \ll 1$, and decompose \mathbf{G}_{δ} further. Let $S_t(f)$ be the linear operator with multiplier $m_{\delta}(t^{-1}\xi',\xi_n)$, then we can write $[\mathbf{G}_{\delta}(f)(x)]^2$ as $\sum_{k\in\mathbb{Z}}|G_k(f)(x)|^2$ with

$$|G_k(f)(x)|^2 := \int_{2^k}^{2^{k+1}} |S_t(f)(x)|^2 \frac{dt}{t}$$

The study of G_{δ} can be reduced to the boundedness of G_k , as we will see in the next result.

Proposition 3.3. Let $\alpha \in (-(n-1), n-1)$ and $\beta \in (-1, 1)$. If (3.10) $\|G_0(f)\|_{L^2(w_{\alpha,\beta})} \leq A \|f\|_{L^2(w_{\alpha,\beta})}$

for any $f \in L^2(w_{\alpha,\beta})$, then

$$\|\mathbf{G}_{\boldsymbol{\delta}}(f)\|_{L^{2}(w_{\boldsymbol{\alpha},\boldsymbol{\beta}})} \lesssim A\|f\|_{L^{2}(w_{\boldsymbol{\alpha},\boldsymbol{\beta}})}$$

To prove this proposition, we establish first the following lemma.

Lemma 3.4. *Fix* $\alpha, \beta \in \mathbb{R}$ *. If*

(3.11)
$$\|G_0(f)\|_{L^2(w_{\alpha,\beta})} \le A \|f\|_{L^2(w_{\alpha,\beta})}$$

for any $f \in L^2(w_{\alpha,\beta})$, then for any $k \in \mathbb{Z}$ it holds

$$||G_k(f)||_{L^2(w_{\alpha,\beta})} \le A ||f||_{L^2(w_{\alpha,\beta})}.$$

Proof. We need to estimate

$$\int_{\mathbb{R}^n} |G_k(f)(x)|^2 w_{\alpha,\beta}(x) dx = \int_{\mathbb{R}^n} \int_{2^k}^{2^{k+1}} \left| S_t(f)(x) \right|^2 \frac{dt}{t} |x'|^{-\alpha} |x_n|^{-\beta} dx.$$

By change of variables, the inner integral is equal to

$$\int_{1}^{2} \left| \left[m_{\delta}(2^{-k}t^{-1}\xi',\xi_{n})\widehat{f}(\xi) \right]^{\vee}(x) \right|^{2} \frac{dt}{t} \\ = \int_{1}^{2} \left| 2^{k(n-1)} \left[m_{\delta}(t^{-1}\xi',\xi_{n})\widehat{f}(2^{k}\xi',\xi_{n}) \right]^{\vee}(2^{k}x',x_{n}) \right|^{2} \frac{dt}{t} \right|^{2}$$

Let $\widehat{f}_k(\xi) = \widehat{f}(2^k \xi', \xi_n)$. It follows from (3.11) that

$$\int_{\mathbb{R}^{n}} |G_{k}(f)(x)|^{2} w_{\alpha,\beta}(x) dx$$

=2^{k(n-1)}2^{k\alpha} $\int_{\mathbb{R}^{n}} \int_{1}^{2} \left| S_{t}(f_{k})(x) \right|^{2} \frac{dt}{t} |x'|^{-\alpha} |x_{n}|^{-\beta} dx$
 $\leq A^{2} 2^{k(n-1)} 2^{k\alpha} \int_{\mathbb{R}^{n}} |f_{k}(x)|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx$

$$=A^{2}2^{k(n-1)}2^{k\alpha}\int_{\mathbb{R}^{n}}|2^{-k(n-1)}f(2^{-k}x',x_{n})|^{2}|x'|^{-\alpha}|x_{n}|^{-\beta}dx$$
$$\leq A^{2}||f||^{2}_{L^{2}(w_{\alpha,\beta})}.$$

To finish the proof of Proposition 3.3, we need the following lemma to paste G_k . Let $\hat{\zeta}$ be a smooth bump supported in $\{\xi' \in \mathbb{R}^{n-1} : |\xi'| \sim 1\}$, and $\hat{\psi}$ a smooth bump supported in $\{\xi_n \in \mathbb{R} : \xi_n \sim 1\}$. We define $\widehat{f_{k,L}}(\xi) := \widehat{f}(\xi)\zeta(2^{-(k+L)}\xi')\widehat{\psi}(2^{-L}\xi_n)$. We will explore the orthogonality of $\{f_{k,L}\}$ below.

Lemma 3.5. Let $\alpha \in (-(n-1), n-1)$ and $\beta \in (-1, 1)$. Then $\sum_{k \in \mathbb{Z}} \sum_{L \in \mathbb{Z}} \int_{\mathbb{R}^n} |f_{k,L}(x)|^2 w_{\alpha,\beta}(x) dx \lesssim \int_{\mathbb{R}^n} |f(x)|^2 w_{\alpha,\beta}(x) dx.$

Proof. Let $\widehat{F_{k,L}}(\xi) = \widehat{f}(\xi)\widehat{\zeta}(2^{-k}\xi')\widehat{\psi}(2^{-L}\xi_n)$, which is different from $\widehat{f_{k,L}}(\xi)$ in that they have different locations in ξ' . By a simple change of variables on (k,L), we have

$$\begin{split} &\sum_{k\in\mathbb{Z}}\sum_{L\in\mathbb{Z}}\int_{\mathbb{R}^n}|f_{k,L}|^2w_{\alpha,\beta}(x)dx\\ &=\sum_{k\in\mathbb{Z}}\sum_{L\in\mathbb{Z}}\int_{\mathbb{R}^n}|F_{k,L}|^2w_{\alpha,\beta}(x)dx\\ &\sim\int_{\mathbb{R}^n}\int_0^1\int_0^1|\sum_k\sum_L r_k(s_1)r_L(s_2)F_{k,L}(x)|^2ds_1ds_2w_{\alpha,\beta}(x)dx, \end{split}$$

where $r_k(s_1)$ and $r_L(s_2)$ are Rademacher functions, and we use Khintchine's inequality of several variables (see [23, Appendix C.5]).

Let us write $\sum_{k} \sum_{L} r_k(s_1) r_L(s_2) F_{k,L}(x) = K_{s_1,s_2} * f$ with

$$\widehat{K}_{s_1,s_2}(\xi) = \left(\sum_k r_k(s_1)\widehat{\zeta}(2^{-k}\xi')\right)\left(\sum_L r_L(s_2)\widehat{\psi}(2^{-L}\xi_n)\right) =: \widehat{K}_{s_1}(\xi')\widehat{K}_{s_2}(\xi_n).$$

This is a bi-parameter singular integral with associated constants independent of s_1 and s_2 . Applying [20, Theorem] we obtain

$$\int_{\mathbb{R}^n} |K_{s_1,s_2} * f(x)|^2 w_{\alpha,\beta}(x) dx \lesssim \int_{\mathbb{R}^n} |f(x)|^2 w_{\alpha,\beta}(x) dx$$

uniformly in $s_1, s_2 \in [0, 1]$ as $w_{\alpha, \beta}$ is a product $A_2(\mathbb{R}^{n-1} \times \mathbb{R})$ weight in the sense of (3.4). This concludes the proof by integrating over s_1 and s_2 . \Box

Proof of Proposition 3.3. We write $\|\mathbf{G}_{\delta}(f)\|_{L^{2}(w_{\alpha,\beta})}^{2}$ as

$$\sum_{k}\int_{2^{k}}^{2^{k+1}}\int_{\mathbb{R}^{n}}\left|\left[m_{\delta}(t^{-1}\xi',\xi_{n})\widehat{f}(\xi)\right]^{\vee}(x)\right|^{2}w_{\alpha,\beta}(x)dx\frac{dt}{t},$$

which, by the support of m_{δ} , is bounded by

$$\begin{split} \sum_{k} \int_{2^{k}}^{2^{k+1}} \int_{\mathbb{R}^{n}} \left| \left[m_{\delta}(t^{-1}\xi',\xi_{n})\widehat{f}_{k,0}(\xi) \right]^{\vee}(x) \right|^{2} w_{\alpha,\beta}(x) dx \frac{dt}{t} \\ &= \sum_{k} \int_{\mathbb{R}^{n}} \int_{2^{k}}^{2^{k+1}} \left| \left[m_{\delta}(t^{-1}\xi',\xi_{n})\widehat{f}_{k,0}(\xi) \right]^{\vee}(x) \right|^{2} \frac{dt}{t} w_{\alpha,\beta}(x) dx \\ &= \sum_{k} \int_{\mathbb{R}^{n}} |G_{k}(f_{k,0})(x)|^{2} w_{\alpha,\beta}(x) dx \\ &\lesssim A^{2} \sum_{k} \int_{\mathbb{R}^{n}} |f_{k,0}(x)|^{2} w_{\alpha,\beta}(x) dx \\ &\lesssim A^{2} \int_{\mathbb{R}^{n}} |f(x)|^{2} w_{\alpha,\beta}(x) dx. \end{split}$$

In the last two steps we use Lemma 3.4 and Lemma 3.5

4. Some trace lemmas

We prove in this section some trace lemmas, whose dual forms, weighted restriction-type estimates, are good substitutes of classical restriction estimates in our setting.

Let $\alpha \ge 0, \beta \ge 0$, and

$$\Gamma_{\delta} := \{ x \in \mathbb{R}^n : \text{ dist } (x, \Gamma) < \delta \},\$$

where $\Gamma = \{\xi \in \mathbb{R}^n : |\xi'| = \xi_n \in (1,2)\}$. We are interested in determining $C_{\delta}(\alpha,\beta)$ in the following inequality.

(4.1)
$$\int_{\Gamma_{\delta}} |\widehat{f}(\xi)|^2 \leq C_{\alpha,\beta} C_{\delta}(\alpha,\beta) \int_{\mathbb{R}^n} |f(x)|^2 w_{\alpha,\beta}^{-1}(x) dx$$

where $w_{\alpha,\beta}^{-1} = |x'|^{\alpha} |x_n|^{\beta}$.

Our main result in this section is as follows.

Proposition 4.1. Let $\alpha \in [0, n-1)$ and $\beta \in [0, 1)$. Then (4.1) holds for

$$C_{\delta}(\alpha,\beta) \sim \left\{ egin{array}{ll} \delta^{lpha+eta}, & lpha+eta < 1 \ \delta\lograc{1}{\delta}, & lpha+eta = 1 \ \delta, & lpha+eta > 1. \end{array}
ight.$$

The proof of Proposition 4.1 is contained in the following results, as the case $\alpha = \beta = 0$ is trivial.

Lemma 4.2. If $\beta = 0$, then (4.1) holds for

$$C_{\delta}(\alpha, 0) \sim \left\{ egin{array}{ll} \delta^{lpha}, & 0 < lpha < 1 \ \delta \log rac{1}{\delta}, & lpha = 1 \ \delta, & 1 < lpha < n-1. \end{array}
ight.$$

This result is a direct consequence of the following classical trace inequalities for spheres.

Lemma 4.3 ([9, Lemma 3]). *Let* $\alpha \in (0, n-1)$. *It holds*

(4.2)
$$\int_{\mathbb{S}^{n-2}+O(\delta)} |\widehat{g}(\xi)|^2 d(\xi) \lesssim C_{\delta}(\alpha,0) \int_{\mathbb{R}^{n-1}} |g(x)|^2 |x|^{\alpha} dx.$$

Proof of Lemma 4.2. Let

$$f_{\xi_n}(x') = \int_{\mathbb{R}^{n-1}} \widehat{f}(\xi',\xi_n) e^{2\pi i x' \cdot \xi'} d\xi' = \int_{\mathbb{R}} f(x',x_n) e^{-2\pi i x_n \xi_n} dx_n.$$

Applying Lemma 4.3 we obtain

$$\begin{split} \int_{\Gamma_{\delta}} |\widehat{f}(\xi)|^2 d\xi &\leq \int_1^2 \int_{\mathbb{S}_{\xi_n}^{n-2} + O(\delta)} |\widehat{f}(\xi',\xi_n)|^2 d\xi' d\xi_n \\ &\lesssim C_{\delta}(\alpha,0) \int_1^2 \int_{\mathbb{R}^{n-1}} |f_{\xi_n}(x')|^2 |x'|^{\alpha} dx' d\xi_n, \end{split}$$

where $\mathbb{S}_{\xi_n}^{n-2}$ is the sphere centered at the origin of radius ξ_n . By Plancherel's identity, this is bounded by

$$C_{\delta}(\alpha,0) \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} \left| \int_{\mathbb{R}} f(x',x_n) e^{-2\pi i x_n \xi_n} dx_n \right|^2 d\xi_n |x'|^{\alpha} dx'$$

= $C_{\delta}(\alpha,0) \int_{\mathbb{R}^n} |f(x)|^2 |x'|^{\alpha} dx.$

Lemma 4.4. If $\alpha = 0$ and $\beta \in (0, 1)$, then (4.1) holds for $C_{\delta}(\alpha, \beta) \sim \delta^{\beta}$.

Proof. We want to prove

(4.3)
$$\int_{\Gamma_{\delta}} |\widehat{f}(\xi)|^2 d\xi \leq C \delta^{\beta} \int_{\mathbb{R}^n} |f(x)|^2 |x_n|^{\beta} dx,$$

which by duality is equivalent to

(4.4)
$$\int_{\mathbb{R}^n} |\widehat{g}(\xi)|^2 \frac{d\xi}{|\xi_n|^\beta} \lesssim \delta^\beta \int_{\Gamma_\delta} |g(x)|^2 dx$$

for any $g \in L^2(\Gamma_{\delta})$.

We first claim that

(4.5)
$$\int_{\mathbb{R}} |\widehat{F}(\xi_n)|^2 \frac{d\xi_n}{|\xi_n|^{\beta}} \lesssim \delta^{\beta} \int_{I_{\delta}} |F(x_n)|^2 dx_n$$

for any $F \in L^2(I_{\delta})$, where $I_{\delta} = [a - \delta, a + \delta]$ is an interval of length 2δ . Take $F(x_n) = g(x', x_n)$, which is supported in $(|x'| - \delta, |x'| + \delta)$ as $x \in \Gamma_{\delta}$,

Take $F(x_n) = g(x', x_n)$, which is supported in $(|x'| - \delta, |x'| + \delta)$ as $x \in \Gamma_{\delta}$ then

$$\int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} |\widehat{F}(\xi_n)|^2 \frac{d\xi_n}{|\xi_n|^\beta} dx' \lesssim \delta^\beta \int_{\mathbb{R}^n} |g(x)|^2 dx$$

by (4.5). The left hand side, by Plancherel's identity, is equal to

$$\int_{\mathbb{R}^n} |\widehat{g}(\xi)|^2 \frac{d\xi}{|\xi_n|^{\beta}},$$

which yields (4.4).

It remains to verify (4.5). Actually $\int_{\mathbb{R}} |\widehat{F}(\xi_n)|^2 \frac{d\xi_n}{|\xi_n|^{\beta}}$ equals

$$C_{\beta} \int_{\mathbb{R}} \int_{\mathbb{R}} F(x_n) \overline{F}(y_n) \frac{1}{|x_n - y_n|^{1 - \beta}} dx_n dy_n$$

as $\beta \in (0,1)$. It follows from this expression and the Schur test that (4.5) is a consequence of

$$\sup_{x_n\in[a-\delta,a+\delta]}\int_{a-\delta}^{a+\delta}\frac{1}{|x_n-y_n|^{1-\beta}}dy_n\lesssim\delta^{\beta},$$

which is straightforward to check using $\beta > 0$.

Lemma 4.5. If $\alpha \in (0, n-1)$ and $\beta \in (0, 1)$, then (4.1) holds for

$$C_{\delta}(\alpha,\beta)\sim \left\{ egin{array}{ll} \delta^{lpha+eta},&lpha+eta<1\ \delta\lograc{1}{\delta},&lpha+eta=1\ \delta,&lpha+eta>1. \end{array}
ight.$$

Proof. By duality (4.1) is equivalent to

(4.6)
$$\int_{\mathbb{R}^n} |\widehat{g}(\xi)|^2 |\xi'|^{-\alpha} |\xi_n|^{-\beta} d\xi \lesssim C_{\delta}(\alpha,\beta) \int_{\Gamma_{\delta}} |g(x)|^2 dx$$

By an argument similar to the proof of Lemma 4.4, it suffices to show that

(4.7)
$$\sup_{x\in\Gamma_{\delta}}\int_{\Gamma_{\delta}}|x'-y'|^{\alpha-(n-1)}|x_n-y_n|^{\beta-1}dy \lesssim C_{\delta}(\alpha,\beta).$$

The left hand side is

(4.8)
$$\sup_{x\in\Gamma_{\delta}}\int_{\Gamma_{\delta}-x}|z'|^{\alpha-(n-1)}|z_{n}|^{\beta-1}dz.$$

-	-		

By rotation in the x'-plane, we may assume that $x' = (|x'|, 0, ..., 0) \in \mathbb{R}^{n-1}$ with $|x'| \in [1, 2]$.

To analyze the integral in (4.8), we define

9

$$S_0 = \{ z \in \Gamma_{\delta} - x : |z'| \le \delta \},$$
$$S_{\ell} = \{ z \in \Gamma_{\delta} - x : \ell \delta \le |z'| \le (\ell + 1)\delta \} \text{ for } 1 \le \ell \le \delta^{-1}/1000 \},$$

and

$$S_{\infty} = \{ z \in \Gamma_{\delta} - x : |z'| \ge 1/1000 \}.$$

Obviously $\Gamma_{\delta} - x = \left[\bigcup_{0 \le \ell \le \delta^{-1}/1000} S_{\ell} \right] \bigcup S_{\infty}$. S_0 is essentially a ball of radius $\sim \delta$, and S_{ℓ} is contained in an annular cylinder for $\ell \ge 1$.

Let us calculate the integrals over S_0 and S_{∞} first.

We observe that $0 \in \Gamma_{\delta} - x$, and, for $z \in \Gamma_{\delta} - x$, we have $|z_n| \le 2\delta$ when $|z'| \le \delta$, therefore (4.9)

$$\int_{S_0} |z'|^{\alpha-(n-1)} |z_n|^{\beta-1} dz \lesssim \int_{|z'| \lesssim \delta} |z'|^{\alpha-(n-1)} dz' \int_{|z_n| \lesssim \delta} |z_n|^{\beta-1} dz_n \lesssim \delta^{\alpha+\beta}$$

as $\alpha, \beta > 0$.

To estimate the integral over S_{∞} , we denote the intersection of S_{∞} and the hyperplane $\{y \in \mathbb{R}^n : y_n = z_n\}$ by

$$E_{z_n} := \{ z' \in \mathbb{R}^{n-1} : ||z'+x'| - (z_n+x_n)| \lesssim \delta, |z'| \ge 10^{-3} \},\$$

whose (n-1)-dimensional volume is bounded by δ as E_{z_n} is contained in an annulus of radius ~ 1 and width $\sim \delta$. This yields (4.10)

$$\int_{S_{\infty}} |z'|^{\alpha - (n-1)} |z_n|^{\beta - 1} dz \lesssim \int_{|z_n| \lesssim 1} |z_n|^{\beta - 1} \int_{E_{z_n}} |z'|^{\alpha - (n-1)} dz' dz_n \lesssim \delta$$

since $|z'| \sim 1$ in E_{z_n} and $\beta > 0$.

We estimate the integral over S_{ℓ} with $1 \le \ell \le \delta^{-1}/1000$ below, where we have $\ell \delta \le |z'| \le (\ell+1)\delta$. It is easy to see that $|z_n| \le (\ell+5)\delta$ in S_{ℓ} . Fix $|z_n| \le (\ell+5)\delta$, then the set $\{z' : (z', z_n) \in S_{\ell}\}$ is contained in

$$E'_{z_n} := (\mathbb{S}^{n-2}_{\ell\delta} + O(\delta)) \cap (\mathbb{S}^{n-2}_{x_n+z_n} - x' + O(\delta)).$$

For $|z_n| \in I_k := [(k-1)\delta, k\delta]$ with $1 \le k \le \ell + 5$, we have

(4.11)
$$|E'_{z_n}| \lesssim \ell^{\frac{n-2}{2}} (\ell+10-k)^{\frac{n-4}{2}} \delta^{n-1}$$

Let

$$A_{z_n} := (\mathbb{S}_{\ell\delta}^{n-2} + O(\delta)) \cap (\mathbb{S}_{x_n+z_n}^{n-2} - x').$$

As $x_n + z_n \sim 1$, (4.12) $|E'_{z_n}| \lesssim |A_{z_n}| \delta$. Let $\alpha_{z_n,\ell}$ be the smallest angle of the triangle with side lengths $x_n + z_n$, x_n , and $\ell \delta$, i.e. the opposite angle of the side with length $\ell \delta$. Then

$$|A_{z_n}| \lesssim [\alpha_{z_n,\ell+1}]^{n-2} - [\alpha_{z_n,\ell}]^{n-2} \lesssim \ell^{\frac{n-2}{2}} (\ell+10-k)^{\frac{n-4}{2}} \delta^{n-2},$$

which implies (4.11) recalling (4.12).

It follows from (4.11) that

$$\begin{split} & \int_{S_{\ell}} |z'|^{\alpha - (n-1)} |z_{n}|^{\beta - 1} dz \\ \lesssim & \sum_{1 \le k \le \ell + 5} \int_{|z_{n}| \in I_{k}} \int_{\{z': \ (z', z_{n}) \ \in S_{\ell}\}} |z'|^{\alpha - (n-1)} dz'|z_{n}|^{\beta - 1} dz_{n} \\ \lesssim & \sum_{1 \le k \le \ell + 1} \ell^{\frac{n-2}{2}} (\ell + 10 - k)^{\frac{n-4}{2}} \delta^{n-1} (\ell \delta)^{\alpha - (n-1)} (k\delta)^{\beta - 1} \delta \\ & \sim & \delta^{\alpha + \beta} \ell^{\alpha - (n-1)} \ell^{\frac{n-2}{2}} \sum_{1 \le k \le \ell + 1} (\ell + 10 - k)^{\frac{n-4}{2}} k^{\beta - 1} \\ \lesssim & \delta^{\alpha + \beta} \ell^{\alpha + \beta - 2}. \end{split}$$

Summing over ℓ , we obtain the control

$$\begin{split} \sum_{\ell} \int_{S_{\ell}} |z'|^{\alpha - (n-1)} |z_n|^{\beta - 1} dz \lesssim &\delta^{\alpha + \beta} + \sum_{1 \leq \ell \lesssim \delta^{-1}} \delta^{\alpha + \beta} \ell^{\alpha + \beta - 2} + \delta \\ \lesssim \begin{cases} \delta^{\alpha + \beta}, & \alpha + \beta < 1 \\ \delta \log \frac{1}{\delta}, & \alpha + \beta = 1 \\ \delta, & \alpha + \beta > 1. \end{cases} \end{split}$$

This estimate finishes the proof.

5. WEIGHTED ESTIMATES

The main goal of this section is to prove the following result via trace inequalities obtained in Section 4.

Theorem 5.1. *For* $\alpha \in [0, n-1)$ *,* $\beta \in [0, 1)$ *, and* $0 < \delta \ll 1$ *, let*

$$A_{\delta}(\alpha,\beta) = \delta^{\frac{1-(\alpha+\beta)}{2}} C_{\delta}(\alpha,\beta)^{1/2} \sim \begin{cases} \delta^{1/2}, & \alpha+\beta < 1\\ \delta^{1/2}(\log\frac{1}{\delta})^{1/2}, & \alpha+\beta = 1\\ \delta^{(2-\alpha-\beta)/2}, & \alpha+\beta > 1. \end{cases}$$

Then

(5.1)
$$\|G_0(f)\|_{L^2(w_{\alpha,\beta})} \leq CA_{\delta}(\alpha,\beta) \|f\|_{L^2(w_{\alpha,\beta})}.$$

We recall that $w_{\alpha,\beta}(x) = |x'|^{-\alpha} |x_n|^{-\beta}$,

$$m_{\delta}(\xi) = \mu_{\delta}(\xi'/\xi_n)\psi(2^{-1}\xi_n)$$

with μ_{δ} satisfying (3.7) and (3.8), and

$$G_0(f)(x) = \left(\int_1^2 \left| \left[m_{\delta}(t^{-1}\xi',\xi_n) \widehat{f}(\xi) \right]^{\vee}(x) \right|^2 \frac{dt}{t} \right)^{1/2}.$$

To study G_0 , we decompose $K_{\delta} := (m_{\delta})^{\vee}$ dyadically. More precisely, we take a smooth function Ψ supported in the unit annulus such that

$$1=\sum_{j\geq j_0}\Psi_j,$$

where $\Psi_j = \Psi(2^{-j} \cdot)$ is supported in $B(0, 2^{j+1}) \setminus B(0, 2^{j-1})$ for $j \ge j_0 + 1$, and $\Psi_{j_0} = 1 - \sum_{j \ge j_0+1} \Psi_j$ is supported in $B(0, 2^{j_0})$ with $2^{j_0} \sim \delta^{-1}$. Correspondingly we decompose $K_{\delta} = \sum_{j \ge j_0} K_j$ with $K_j := K_{\delta} \Psi_j$.

As m_{δ} is supported in Γ_{δ} and $\widehat{\Psi}_j$ is essentially supported in a ball of radius $2^{-j} \leq \delta$, we see that $\widehat{K}_j = m_{\delta} * \widehat{\Psi}_j$ is essentially supported in Γ_{δ} , which we describe precisely below.

Proposition 5.2. Let $j \ge j_0$. For any $M \in \mathbb{N}$, there exists a constant $C_M > 0$ such that the following statements hold.

(*i*) For any $\xi \in \mathbb{R}^n$, we have

(5.2)
$$|\widehat{K_j}(\xi)| \le C_M (2^j \delta)^{-M}.$$

(ii) For dist $(\xi, \Gamma) \sim 2^{\ell} \delta$ with $\ell \geq 1$, we have

(5.3)
$$|\widehat{K_j}(\xi)| \le C_M 2^{-\ell M} (2^j \delta)^{-M}$$

(*iii*) For dist $(\xi, \Gamma) \geq \frac{1}{2}$,

(5.4)
$$|\widehat{K_j}(\xi)| \le C_M 2^{-jM} (1+|\xi|)^{-M}$$

Proof. (i) For $j \ge j_0 + 1$, we may rewrite \widehat{K}_j as

$$\int m_{\delta}(\xi-2^{-j}\eta)\widehat{\Psi}(\eta)d\eta$$

As one can easily show $\partial^{\alpha} m_{\delta}(\xi) \leq \delta^{-|\alpha|}$ from (3.7) and (3.8), applying Taylor's expansion to $m_{\delta}(\xi - 2^{-j} \cdot)$, and using the vanishing moments of Ψ , we obtain the claimed estimate. When $j = j_0$, we have

$$|\widehat{K}_{j_0}(\xi)| \leq C ||m_{\delta}||_{L^{\infty}} ||\widehat{\Psi}_{j_0}||_{L^1} \leq C.$$

(ii) When ξ is $2^{\ell}\delta$ away from Γ , we obtain from the condition $\xi - 2^{-j}\eta \in \Gamma_{\delta}$ that $2^{-j}|\eta| \gtrsim 2^{\ell}\delta$. This implies

$$\begin{split} \left| \int m_{\delta}(\xi - 2^{-j}\eta) \widehat{\Psi}(\eta) d\eta \right| \leq & C_{M,\Psi} \int_{|\eta| \gtrsim 2^{j+\ell} \delta} (1 + |\eta|)^{-2M} d\eta \\ \leq & C_{M,\Psi} (1 + 2^{j+\ell} \delta)^{-M} \\ \leq & C_{M,\Psi} 2^{-\ell M} (2^{j} \delta)^{-M}. \end{split}$$

(iii) (5.4) could be proved similarly via the observation that, for $\xi - 2^{-j}\eta \in \Gamma_{\delta}$, we have

$$2^{-j}|\boldsymbol{\eta}| \ge \frac{1}{10}(1+|\boldsymbol{\xi}|)$$

$$(\boldsymbol{\xi} \ \boldsymbol{\Gamma}) > 1$$

due to the assumption dist $(\xi, \Gamma) \ge \frac{1}{2}$.

We define $\widehat{K_{\delta,t}}(\xi) = m_{\delta}(t^{-1}\xi',\xi_n)$, which implies that $K_{\delta,t} = t^{n-1}K_{\delta}(tx',x_n)$. Let $K_{j,t}(x) = t^{n-1}K_j(tx',x_n)$, then

(5.5)
$$K_{\delta,t}(x) = \sum_{j \ge j_0} K_{j,t}(x)$$

Obviously

(5.6)
$$G_0(f)(x) \le \sum_{j \ge j_0} \left(\int_1^2 |(K_{j,t} * f)(x)|^2 \frac{dt}{t} \right)^{1/2}$$

We remark that $K_{j,t}$ is supported in a ball of radius $c_1 2^j$ with $c_1 \sim 1$ as $t \in [1,2]$. Therefore we can localize f by decomposing it into pieces supported in cubes of length $c_1 2^j$. For $i \in \mathbb{Z}^n$, we define $Q_i = Q(c_1 2^j i, c_1 2^j)$, then $\{Q_i\}_{i \in \mathbb{Z}^n}$ is a partition of \mathbb{R}^n . Define $f_i = f \chi_{Q_i}$, then

(5.7)
$$f = \sum_{i \in \mathbb{Z}^n} f_i.$$

To prove (5.1), it follows from (5.6) and (5.7) that we should estimate

$$\int_1^2 \int_{\mathbb{R}^n} |K_{j,t} * f_i(x)|^2 w_{\alpha,\beta}(x) dx \frac{dt}{t},$$

which relies heavily on the position of *i*. Therefore we decompose \mathbb{Z}^n into four subsets:

$$E_{1} := \{i \in \mathbb{Z}^{n} : |i'| \ge 10n, |i_{n}| \ge 10n\}$$

$$E_{2} := \{i \in \mathbb{Z}^{n} : |i'| \ge 10n, |i_{n}| \le 10n\}$$

$$E_{3} := \{i \in \mathbb{Z}^{n} : |i'| \le 10n, |i_{n}| \le 10n\}$$

$$E_{4} := \{i \in \mathbb{Z}^{n} : |i'| \le 10n, |i_{n}| \ge 10n\},$$

and discuss them separately.

Case 1: $i \in E_1$

To handle this case, we need the following lemma obtained from the Plancherel identity.

Lemma 5.3. Suppose $0 < A \le w(x) \le B$ for all $x \in 3Q_i$, where $3Q_i = Q(c_1 2^j i, 3c_1 2^j)$. Then

(5.8)
$$\left\| \left(\int_{1}^{2} |K_{j,t} * f_{i}(x)|^{2} \frac{dt}{t} \right)^{1/2} \right\|_{L^{2}(w)} \lesssim \left(\frac{B}{A}\right)^{1/2} \delta^{1/2} (2^{j} \delta)^{-M} \|f_{i}\|_{L^{2}(w)}.$$

Proof. Noticing that $K_{j,t} * f_i$ is supported in $3Q_i$, the left hand side of (5.8) is bounded by

(5.9)
$$B^{1/2} \left(\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} dx \frac{dt}{t} \right)^{1/2}.$$

As $\widehat{K_{j,t}}$ is essentially supported¹ in

$$\Gamma_{\delta,t} := \{ \xi \in \mathbb{R}^n : (t^{-1}\xi', \xi_n) \in \Gamma_{\delta} \},\$$

we may decompose the range of *t* as

$$[1,2] = \cup_{\beta=\delta^{-1}}^{2\delta^{-1}} I_{\beta}$$

with $I_{\beta} = [\beta \delta, (\beta + 1)\delta]$. Moreover, we define $\widehat{P_{\beta}g}$ as the restriction of \widehat{g} to the set

$$\{\boldsymbol{\xi}\in\mathbb{R}^n:\frac{|\boldsymbol{\xi}'|}{\xi_n}\in[\boldsymbol{\beta}\boldsymbol{\delta},(\boldsymbol{\beta}+1)\boldsymbol{\delta}],\boldsymbol{\xi}_n\in[\frac{1}{2},4]\}.$$

It follows from Proposition 5.2 that, for $t \in I_{\beta}$, the essential support of $\widehat{F_{\beta'}g}$ and the support of $\widehat{P_{\beta'}g}$ are disjoint when $|\beta - \beta'| \ge 10$. This implies that $K_{j,t} * P_{\beta'}f_i$ is essentially 0 when $t \in I_{\beta}$ with $|\beta - \beta'| \ge 10$. Hence

$$\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} dx \frac{dt}{t}$$

$$= \sum_{\beta} \int_{I_{\beta}} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} dx \frac{dt}{t}$$

$$\sim \sum_{\beta} \int_{I_{\beta}} \int \Big| \sum_{\beta'=\beta-10}^{\beta+10} (K_{j,t} * (P_{\beta'}f_{i}))(x) \Big|^{2} dx \frac{dt}{t}$$

$$\lesssim \sum_{\beta} \sum_{\beta'=\beta-10}^{\beta+10} \int_{I_{\beta}} \int |(K_{j,t} * (P_{\beta'}f_{i}))(x)|^{2} dx \frac{dt}{t}$$

$$\sim \sum_{\beta} \sum_{\beta'=\beta-10}^{\beta+10} \int_{I_{\beta}} \int |(\widehat{K_{j,t}} \widehat{P_{\beta'}f_{i}})(\xi)|^{2} d\xi \frac{dt}{t}$$

¹We present here an intuitive argument, while a strict one is provided in the appendix.

$$\begin{split} \lesssim &(2^{j}\delta)^{-M} \sum_{\beta} \sum_{\beta'=\beta-10}^{\beta+10} \int_{I_{\beta}} \int |(\widehat{P_{\beta'}f_{i}})(\xi)|^{2} d\xi \frac{dt}{t} \\ \lesssim &\delta(2^{j}\delta)^{-M} \sum_{\beta} \sum_{\beta'=\beta-10}^{\beta+10} \int |(\widehat{P_{\beta'}f_{i}})(\xi)|^{2} d\xi \\ \lesssim &\delta(2^{j}\delta)^{-M} \|f_{i}\|_{L^{2}}^{2} \\ \lesssim &A^{-1}\delta(2^{j}\delta)^{-M} \|f_{i}\|_{L^{2}(w)}^{2}, \end{split}$$

where we use (5.2). This combined with (5.9) yields (5.8).

For $i \in E_1$ and $w(x) = w_{\alpha,\beta}(x)$, we have $A \sim B$. Applying Lemma 5.3, we obtain (5.10)

$$\left(\int_{1}^{2}\int_{\mathbb{R}^{n}}|K_{j,t}*f_{i}(x)|^{2}w_{\alpha,\beta}(x)dx\frac{dt}{t}\right)^{1/2}\leq C\delta^{1/2}(2^{j}\delta)^{-M}\|f_{i}\|_{L^{2}(w_{\alpha,\beta})}.$$

Case 2: $i \in E_2$

In the remaining three cases, as $w_{\alpha,\beta}$ may not be bounded, Lemma 5.3 is not applicable. We need the following estimate to make use of trace inequalities obtained in Section 4.

Lemma 5.4. *Suppose that* $w_1(x) \ge 0$ *satisfies*

(5.11)
$$w_1(x) \ge A > 0 \qquad \forall x \in Q_i,$$

(5.12)
$$w_1(x) \sim w_1(sx', x_n) \text{ for } s \sim 1,$$

and

(5.13)
$$\int_{\Gamma_{\delta}} |\widehat{g}(\xi)|^2 d\xi \leq C_{\delta}(w_1) \int_{\mathbb{R}^n} |g(x)|^2 w_1^{-1}(x) dx.$$

Then

$$\left(\int_{1}^{2}\int_{\mathbb{R}^{n}}|K_{j,t}*f_{i}(x)|^{2}w_{1}(x)dx\frac{dt}{t}\right)^{1/2}\leq CA^{-1/2}\delta^{1/2}(2^{j}\delta)^{-M}C_{\delta}(w_{1})^{1/2}\|f_{i}\|_{L^{2}(w_{1})}.$$

Proof. As in the proof of Lemma 5.3, we write $[1,2] = \bigcup_{\beta=\delta^{-1}}^{2\delta^{-1}} I_{\beta}$ with $I_{\beta} = [\beta \delta, (\beta+1)\delta]$. Then

$$\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} w_{1}(x) dx \frac{dt}{t}$$

$$\leq C \sum_{\beta} \int_{I_{\beta}} \int_{\mathbb{R}^{n}} |\sum_{\beta'=\beta-10}^{\beta+10} (K_{j,t} * (P_{\beta'}f_{i}))(x)|^{2} w_{1}(x) dx \frac{dt}{t}$$

(5.14)
$$\leq C \sum_{\beta} \sum_{\beta'=\beta-10}^{\beta+10} \int_{I_{\beta}} \int_{\mathbb{R}^{n}} |(K_{j,t} * (P_{\beta'}f_{i}))(x)|^{2} w_{1}(x) dx \frac{dt}{t}.$$

Applying (5.13) and Proposition 5.2, we obtain essentially

$$\begin{split} \int_{\mathbb{R}^n} |K_{j,t} * g(x)|^2 dx &= \int_{\mathbb{R}^n} |\widehat{K_{j,t}}(\xi)\widehat{g}(\xi)|^2 d\xi \\ &\lesssim (2^j \delta)^{-M} \int_{\Gamma_{\delta,t}} |\widehat{g}(\xi)|^2 d\xi \\ &\lesssim (2^j \delta)^{-M} C_{\delta}(w_1) \int_{\mathbb{R}^n} |g(x)|^2 w_1^{-1}(x) dx, \end{split}$$

where in the last step we use (5.12). By duality

$$\int_{\mathbb{R}^n} |K_{j,t} * g(x)|^2 w_1(x) dx \lesssim (2^j \delta)^{-M} C_{\delta}(w_1) \int_{\mathbb{R}^n} |g(x)|^2 dx.$$

Hence (5.14) is bounded by

$$(2^{j}\delta)^{-M}C_{\delta}(w_{1})\sum_{\beta}\sum_{\beta'=\beta-10}^{\beta+10}\int_{I_{\beta}}\int_{\mathbb{R}^{n}}|P_{\beta'}f_{i}(x)|^{2}dx\frac{dt}{t}$$

$$\lesssim (2^{j}\delta)^{-M}\delta C_{\delta}(w_{1})\|f_{i}\|_{L^{2}}^{2}$$

$$\lesssim (2^{j}\delta)^{-M}\delta C_{\delta}(w_{1})A^{-1}\|f_{i}\|_{L^{2}(w_{1})}^{2}.$$

This completes the proof.

We observe that when $x \in Q_i$ with $i \in E_2$, we have $|x_n| \leq 2^j$ and $|x'|^{-\alpha} \sim |x'_i|^{-\alpha}$, where $x_i = c_1 2^j i$ is the center of Q_i . Take $w_1(x) = |x_n|^{-\beta}$, and accordingly $A = 2^{-j\beta}$, then $C_{\delta}(w_1) = \delta^{\beta}$ by Proposition 4.1 with $\alpha = 0$. Applying Lemma 5.4 with $M_1 \geq M + \beta$, we obtain

$$\begin{split} &\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} w_{\alpha,\beta}(x) dx \frac{dt}{t} \\ &\lesssim |x_{i}'|^{-\alpha} \int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} w_{1}(x) dx \frac{dt}{t} \\ &\lesssim |x_{i}'|^{-\alpha} \delta 2^{j\beta} (2^{j} \delta)^{-M_{1}} \delta^{\beta} \int_{\mathbb{R}^{n}} |f_{i}(x)|^{2} w_{1}(x) dx \\ &\lesssim \delta (2^{j} \delta)^{-M} \int_{\mathbb{R}^{n}} |f_{i}(x)|^{2} w_{\alpha,\beta}(x) dx, \end{split}$$

as $w_{\alpha,\beta}(x) \sim |x'_i|^{-\alpha} w_1(x)$ in Q_i . In summary, we have (5.15) $\left(\int_1^2 \int_{\mathbb{R}^n} |K_{j,t} * f_i(x)|^2 w_{\alpha,\beta}(x) dx \frac{dt}{t}\right)^{1/2} \leq C \delta^{1/2} (2^j \delta)^{-M} ||f_i||_{L^2(w_{\alpha,\beta})}.$

Case 3: $i \in E_3$

In this case, to apply Lemma 5.4, we take $w_1(x) = w_{\alpha,\beta}(x)$ and $A = 2^{-j(\alpha+\beta)}$ such that $A \le w_1(x)$ in Q_i . We recall that by Proposition 4.1

$$C_{\delta}(w_{lpha,eta})\lesssim \left\{egin{array}{ll} \delta^{lpha+eta},&lpha+eta<1\ \delta\lograc{1}{\delta},&lpha+eta=1\ \delta,&lpha+eta>1, \end{array}
ight.$$

therefore it follows from Lemma 5.4 that

$$(5.16) \qquad \begin{pmatrix} \int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} w_{\alpha,\beta}(x) dx \frac{dt}{t} \end{pmatrix}^{1/2} \\ \leq C2^{j(\alpha+\beta)/2} \delta^{1/2} (2^{j}\delta)^{-M} C_{\delta}(w_{\alpha,\beta})^{1/2} ||f_{i}||_{L^{2}(w_{\alpha,\beta})} \\ \lesssim ||f_{i}||_{L^{2}(w_{\alpha,\beta})} \begin{cases} \delta^{1/2} (2^{j}\delta)^{-M}, & \alpha+\beta < 1 \\ \delta^{1/2} (\log \frac{1}{\delta})^{1/2} (2^{j}\delta)^{-M}, & \alpha+\beta = 1 \\ \delta^{1-\frac{\alpha+\beta}{2}} (2^{j}\delta)^{-M}, & \alpha+\beta > 1. \end{cases}$$

Case 4: $i \in E_4$

We recall that $E_4 := \{i \in \mathbb{Z}^n : |i'| \le 10n, |i_n| \ge 10n\}$, therefore in Q_i with $i \in E_4$ we have $|x'| \le 2^j$ and $|x_n|^{-\beta} \sim (2^j |i_n|)^{-\beta}$. Taking $w_1(x) = |x'|^{-\alpha} = w_{\alpha,0}$ and $A = 2^{-j\alpha}$, we can apply Lemma 5.4 to obtain

$$\begin{split} &\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} w_{\alpha,\beta}(x) dx \frac{dt}{t} \\ \lesssim (2^{j}|i_{n}|)^{-\beta} \int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}(x)|^{2} w_{1}(x) dx \frac{dt}{t} \\ \lesssim (2^{j}|i_{n}|)^{-\beta} \delta 2^{j\alpha} (2^{j}\delta)^{-M} C_{\delta}(w_{1}) \int_{\mathbb{R}^{n}} |f_{i}(x)|^{2} w_{1}(x) dx \\ \lesssim \delta 2^{j\alpha} (2^{j}\delta)^{-M} C_{\delta}(\alpha,0) \int_{\mathbb{R}^{n}} |f_{i}(x)|^{2} w_{\alpha,\beta}(x) dx. \end{split}$$

By Proposition 4.1 with $\beta = 0$ we obtain

$$\left(\int_{1}^{2}\int_{\mathbb{R}^{n}}|K_{j,t}*f_{i}(x)|^{2}w_{\alpha,\beta}(x)dx\frac{dt}{t}\right)^{1/2}$$

(5.17)
$$\lesssim \|f_i\|_{L^2(w_{\alpha,\beta})} \begin{cases} \delta^{1/2} (2^j \delta)^{-M}, & 0 < \alpha < 1\\ \delta^{1/2} (\log \frac{1}{\delta})^{1/2} (2^j \delta)^{-M}, & \alpha = 1\\ \delta^{1-\frac{\alpha}{2}} (2^j \delta)^{-M}, & 1 < \alpha < n-1. \end{cases}$$

We are now ready to prove Theorem 5.1.

Proof of Theorem 5.1. When $\alpha = \beta = 0$, (5.1) is verified by Proposition 3.2. We discuss below the case $(\alpha, \beta) \neq (0, 0)$.

Recalling that f_i is supported in Q_i and $K_{j,t} * f_i$ is supported in $3Q_i$, we have $K_{j,t} * f = \sum_i (K_{j,t} * f_i) \chi_{3Q_i} \lesssim (\sum_i |K_{j,t} * f_i|^2)^{1/2}$ as $\{3Q_i\}_{i \in \mathbb{Z}^n}$ are finitely overlapping. It follows from the definition of f_i that $|\sum_i f_i|^2 = \sum_i |f_i|^2$. Hence

$$\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx \frac{dt}{t}$$

$$\lesssim \sum_{i} \int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx \frac{dt}{t}$$

$$\sim \left(\sum_{i \in E_{1}} + \sum_{i \in E_{2}} + \sum_{i \in E_{3}} + \sum_{i \in E_{4}}\right) \int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx \frac{dt}{t}$$

When $\alpha + \beta > 1$, by (5.16), we have, for $i \in E_3$,

$$\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f_{i}|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx \frac{dt}{t} \leq \delta^{2-\alpha-\beta} (2^{j}\delta)^{-M} ||f_{i}||_{L^{2}(w_{\alpha,\beta})}^{2}.$$

Moreover, noticing that $1 \le C\delta^{-\beta}$, the same bound holds for all $\alpha \in (0, n-1)$ when $i \in E_4$ by (5.17). These estimates and (5.10), (5.15) yield

$$\begin{split} &\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,t} * f|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx \frac{dt}{t} \\ &\lesssim \left[\delta(2^{j}\delta)^{-M} + \delta^{2-\alpha-\beta} (2^{j}\delta)^{-M} \right] \left(\sum_{i \in E_{1}} + \sum_{i \in E_{2}} + \sum_{i \in E_{3}} + \sum_{i \in E_{4}} \right) \int_{\mathbb{R}^{n}} |f_{i}|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx \\ &\lesssim \left[\delta^{2-\alpha-\beta} (2^{j}\delta)^{-M} \right] \int_{\mathbb{R}^{n}} |f|^{2} |x'|^{-\alpha} |x_{n}|^{-\beta} dx. \end{split}$$

This combined with (5.6) implies that

$$\|G_0(f)\|_{L^2(w_{\alpha,\beta})} \lesssim \delta^{(2-\alpha-\beta)/2} \|f\|_{L^2(w_{\alpha,\beta})}$$

since $j \ge j_0$ with $2^{j_0} \sim \delta^{-1}$.

The case when $\alpha + \beta \leq 1$ can be proved similarly.

6. PROOF OF MAIN RESULTS

It is necessary to show that L is well defined in $L^2(w_{\alpha,\beta})$ and $L^p(\mathbb{R}^n)$.

Lemma 6.1. Let $\alpha \in [0, n-1)$, $\beta \in [0, 1)$, and $p \in (1, \infty)$. The linear operator L initially defined on Schwartz functions is bounded on $L^2(w_{\alpha,\beta})$ and bounded on $L^p(\mathbb{R}^n)$.

Proof. We recall that

$$L(f)(x) = \int_{\mathbb{R}^n} \Psi(2^{-1}\xi_n) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi$$
$$= \int_{\mathbb{R}^{n-1}} \int_{\mathbb{R}} f(x', y_n) 2 \Psi^{\vee}(2(x_n - y_n)) dy_n dx'$$

The L^p boundedness of L follows from Minkowski's inequality directly as $\psi^{\vee} \in L^1(\mathbb{R}).$

Observing that $|g * \psi^{\vee}(x_n)| \leq CM(g)(x_n)$, which is bounded on $L^2(|x_n|^{-\beta})$ for $\beta \in (-1, 1)$, the boundedness of *L* on $L^2(w_{\alpha, \beta})$ follows.

Proof of Theorem 1.3. When $\alpha + \beta > 1$, we obtain from Theorem 5.1 and Proposition 3.3 that

$$\|\mathbf{G}_{\boldsymbol{\delta}}(f)\|_{L^{2}(w_{\boldsymbol{\alpha},\boldsymbol{\beta}})} \lesssim \boldsymbol{\delta}^{(2-\boldsymbol{\alpha}-\boldsymbol{\beta})/2} \|f\|_{L^{2}(w_{\boldsymbol{\alpha},\boldsymbol{\beta}})}.$$

It follows from this, Proposition 3.1, and (3.6) that

$$\|T_*^{\lambda}(f)\|_{L^2(w_{\alpha,\beta})} \lesssim \sum_{\gamma \ge 0} 2^{-\gamma\lambda} 2^{\gamma/2} 2^{-\gamma(2-\alpha-\beta)/2} \|f\|_{L^2(w_{\alpha,\beta})} \lesssim C \|f\|_{L^2(w_{\alpha,\beta})}$$

as $\lambda > \frac{\alpha + \beta - 1}{2}$. The case when $\alpha + \beta \le 1$ can be proved similarly with the help of The-

We have finished the proof of (1.6). Now we turn to the proof of the second statement, which follows from the boundedness of T^{λ}_* on $L^2(\mathbb{R}^n, w_{\alpha,\beta})$ and the following two lemmas.

Lemma 6.2. For any measurable set E, |E| = 0 is equivalent to $w_{\alpha,\beta}(E) =$ 0.

This lemma is trivial.

Lemma 6.3 ([23, Exercise 7.4.1]). Let $\alpha \in [0, n-1)$ and $\beta \in [0, 1)$. The Schwartz functions are dense in $L^2(w_{\alpha,\beta})$.

By these lemmas, Lemma 6.1 and (1.6), we apply Stein's maximal principle ([45], see also [23, Theorem 2.1.14.]) to obtain that

$$\lim_{t \to \infty} T_t^{\lambda}(f)(x) = L(f)(x) \qquad a.e$$

whenever $f \in L^2(w_{\alpha,\beta})$.

Next we prove Theorem 1.2.

Proof of Theorem 1.2. For a given $\lambda > 0$, we fix $p \in [2, \frac{2n}{n-2\lambda-1})$. For any $f \in L^p(\mathbb{R}^n)$, by Proposition 2.1 we can write $f = \sum_{i=1}^4 f_i$ with $f \in L^2(w_i)$. In the decomposition, for any $\varepsilon > 0$, we can select $\alpha_i, \beta_i > 0$ appropriately such that $\alpha_i + \beta_i < n(1 - \frac{2}{p}) + \varepsilon$, i = 1, 2, 3, 4. In particular, we have $\lambda > (\alpha_i + \beta_i - 1)/2$ for i = 1, 2, 3, 4.

By Theorem 1.3 we obtain

$$\lim_{t \to \infty} T_t^{\lambda}(f_i)(x) = L(f_i)(x) \qquad a.e$$

for i = 1, 2, 3, 4. This complete the proof of Theorem 1.2 by summing over *i*.

Finally we prove Theorem 1.1.

Proof of Theorem 1.1. Fix k_1 and k_2 in \mathbb{Z} such that $2^{k_1+10} < a < b < 2^{k_2-10}$. We define

$$L_k(f)(x) = \int_{\mathbb{R}^n} \psi(2^{-k-1}\xi_n) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi.$$

Then for $f \in L^p(\mathbb{R}^n)$ satisfying (1.4), L_k is well defined, and

(6.1)
$$f = \sum_{k_1 \le k \le k_2} L_k(f).$$

Let

$$T_{k,t}^{\lambda}(f)(x) := \int_{\mathbb{R}^n} (1 - \frac{|\xi'|^2}{\xi_n^2})_+^{\lambda} \psi(2^{-k-1}\xi_n) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi,$$

then a dilation argument shows that

(6.2)
$$\lim_{t \to \infty} T_{k,t}^{\lambda}(f) = L_k(f) \qquad a.e.$$

Noticing that, for $f \in L^p(\mathbb{R}^n)$ satisfying (1.4),

$$\tilde{T}_t^{\lambda}(f) = \sum_{k_1 \le k \le k_2} T_{k,t}^{\lambda}(f),$$

the conclusion follows by (6.1) and (6.2).

7. REMARKS

7.1. By a standard duality argument, we obtain from Lemma 3.5 the following orthogonality result.

Lemma 7.1. Let $k, L \in \mathbb{Z}$, $\alpha \in (-(n-1), n-1)$ and $\beta \in (-1, 1)$. There exists a constant *C* such that, for any sequence of functions $\{H_{k,L}\}$ satisfying

supp
$$\widehat{H_{k,L}} \subset \{\xi \in \mathbb{R}^n : |\xi'| \sim 2^{L+k}, \xi_n \sim 2^L\},\$$

we have

$$\int_{\mathbb{R}^n} |\sum_{L \in \mathbb{Z}} H_{k,L}(x)|^2 w_{\alpha,\beta}(x) dx \le C \sum_{L \in \mathbb{Z}} \int_{\mathbb{R}^n} |H_{k,L}(x)|^2 w_{\alpha,\beta}(x) dx.$$

Let

$$\widetilde{\mathbf{G}}_{\delta}(f)(x) := \left(\int_0^\infty \left| \int_{\mathbb{R}^n} \mu_{\delta}(\xi'/t\xi_n) \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi \right|^2 \frac{dt}{t} \right)^{1/2}$$

With the help of Lemma 7.1, we can actually obtain from (3.10) the estimate

(7.1)
$$\|\mathbf{G}_{\boldsymbol{\delta}}(f)\|_{L^{2}(w_{\boldsymbol{\alpha},\boldsymbol{\beta}})} \lesssim A \|f\|_{L^{2}(w_{\boldsymbol{\alpha},\boldsymbol{\beta}})}$$

by imitating the proof of Proposition 3.3.

By (7.1), to obtain the (weighted) boundedness of

$$\sup_{t>0} \left| \int_{\mathbb{R}^n} (1 - \frac{|\xi'|^2}{t^2 \xi_n^2})^{\lambda}_+ \widehat{f}(\xi) e^{2\pi i x \cdot \xi} d\xi \right|,$$

it remains to study the maximal function defined by

(7.2)
$$\widetilde{M}_{\varphi}(f)(x) := \sup_{t>0} \left| \left[\varphi(\xi'/t\xi_n) \widehat{f}(\xi) \right]^{\vee} \right|,$$

where $\varphi \in C_0^{\infty}(\mathbb{R}^{n-1})$ is supported in B(0,1) with $\varphi(0) = 1$. We conjecture that \widetilde{M}_{φ} is bounded on $L^2(\mathbb{R}^n)$. But we cannot prove it. By a standard argument using square functions, we can assume further that $\varphi(\xi') = 1$ for $|\xi'| \leq 1/2$.

7.2. Let

$$K_{\lambda}(x) = \int_{\mathbb{R}^n} (1 - \frac{|\xi'|^2}{\xi_n^2})^{\lambda}_+ \psi(\xi_n) e^{2\pi i x \cdot \xi} d\xi.$$

It has been proved in [32] that

$$K^{\lambda} \in L^1(\mathbb{R}^n)$$

for $\lambda > \frac{n-2}{2}$, which implies that

(7.3)
$$||T^{\lambda}(f)||_{L^{p}(\mathbb{R}^{n})} \leq C||f||_{L^{p}(\mathbb{R}^{n})}$$

for $1 . It is natural to conjecture that, when <math>\lambda > \frac{n-2}{2}$,

(7.4)
$$||T_*^{\lambda}(f)||_{L^p(\mathbb{R}^n)} \le C ||f||_{L^p(\mathbb{R}^n)}, \quad 1$$

This can be verified for $\lambda > \frac{n}{2}$

Proposition 7.2. Let $\lambda > \frac{n}{2}$. Then

$$\|T_*^{\lambda}(f)\|_{L^p(\mathbb{R}^n)} \le C \|f\|_{L^p(\mathbb{R}^n)}, \quad 1$$

This result follows from the pointwise estimate

(7.5)
$$|K_{\lambda}(x)| \le C(1+|x|)^{-\frac{n}{2}-\lambda}$$

obtained by [32]. We remark that this bound is worse than the corresponding pointwise estimate of $[(1 - |\xi|^2)_+]^{\vee}$. We are curious if we can fill the gap between Proposition 7.2 and the

We are curious if we can fill the gap between Proposition 7.2 and the conjectured range $\lambda > \frac{n-2}{2}$. This kind of gap does appear in the classical Bochner-Riesz problem.

8. Appendix

We present a strict proof of Lemma 5.3 in this section, while a strict proof of Lemma 5.4 could be obtained similarly.

Let us recall Lemma 5.3 first.

Lemma 8.1. Suppose $0 < A \le w(x) \le B$ for all $x \in 3Q_i$, where $3Q_i = Q(c_1 2^j i, 3c_1 2^j)$. Then

$$\left\| \left(\int_{1}^{2} |K_{j,t} * f_{i}(x)|^{2} \frac{dt}{t} \right)^{1/2} \right\|_{L^{2}(w)} \lesssim \left(\frac{B}{A} \right)^{1/2} \delta^{1/2} (2^{j} \delta)^{-M} \|f_{i}\|_{L^{2}(w)}.$$

Proof. Let ℓ_0 be an integer such that $2^{\ell_0} \sim (10\delta)^{-1}$. We need to decompose \widehat{K}_i further such that

$$\widehat{K}_j(\xi) = \sum_{10 \leq \ell \leq \ell_0} \widehat{K}_j(\xi) \widehat{arphi}_\ell(\xi) + \widehat{K}_j(\xi) \widehat{arphi}_\infty(\xi) =: \sum_\ell \widehat{K_{j,\ell}}(\xi),$$

where $\widehat{\varphi}_{10}(\xi)$ is supported in $\{\xi : dist(\xi, \Gamma) \leq 2^{20}\delta\}$, $\widehat{\varphi}_{\ell}$ is supported in $\{\xi : dist(\xi, \Gamma) \sim 2^{\ell}\delta\}$ for $11 \leq \ell \leq \ell_0$, and

$$\widehat{arphi}_{\infty}(\xi) = 1 - \sum_{10 \leq \ell \leq \ell_0} \widehat{arphi}_{\ell}(\xi).$$

Correspondingly we define $\widehat{K_{j,\ell,t}}(\xi) = \widehat{K_{j,\ell}}(t^{-1}\xi',\xi_n)$, and we have

$$K_{j,t} = \sum_{\ell \ge 10} K_{j,\ell,t}.$$

Using the above decomposition and that $w(x) \leq B$ we conrol $(\int_1^2 \int_{\mathbb{R}^n} |K_{j,t} * f_i(x)|^2 w(x) dx \frac{dt}{t})^{1/2}$ by

(8.1)
$$B^{1/2} \sum_{\ell \ge 10} \left(\int_{1}^{2} \int_{\mathbb{R}^{n}} |K_{j,\ell,t} * f_{i}(x)|^{2} dx \frac{dt}{t} \right)^{1/2}$$

We will consider two cases: $10 \le \ell \le \ell_0$, and $\ell \ge \infty$.

Case 1: $10 \le \ell \le \ell_0$.

Since $\widehat{K_{j,\ell,t} * f_i}(\xi) = \widehat{K_{j,\ell}}(t^{-1}\xi', \xi_n)\widehat{f_i}(\xi)$, in its support we have $|\xi'| = t\xi_n + [-2^{\ell+2}\delta, 2^{\ell+2}\delta]$

and $\xi_n \in [\frac{1}{2}, 4]$. We decompose $[1, 2] = \bigcup_{\beta=\delta^{-1}}^{2\delta^{-1}} I_{\beta}$ with $I_{\beta} = [\beta \delta, (\beta + 1)\delta]$, and the support of $\widehat{K_{j,\ell,t}}$ is

$$\{\xi \in \mathbb{R}^n : |\xi'| \in \xi_n \beta \delta + [-2^{\ell+2}\delta, 2^{\ell+2}\delta], \xi_n \in [\frac{1}{2}, 4]\}.$$

Define $\widehat{P_{\beta}g}$ as the restriction of \widehat{g} to the set

$$\{\boldsymbol{\xi}\in\mathbb{R}^n: |\boldsymbol{\xi}'|\in[\boldsymbol{\xi}_n\boldsymbol{\beta}\boldsymbol{\delta},\boldsymbol{\xi}_n(\boldsymbol{\beta}+1)\boldsymbol{\delta}], \ \boldsymbol{\xi}_n\in[\frac{1}{2},4]\}.$$

Then for $t \in I_{\beta}$, the support of $\widehat{K_{j,\ell,t}}$ and the support of $\widehat{P_{\beta'g}}$ are disjoint if $|\beta - \beta'| \ge 10 \cdot 2^{\ell}$. Hence

(8.2)
$$\int_{1}^{2} \int |K_{j,\ell,t} * f_{i}(x)|^{2} dx \frac{dt}{t}$$
$$= \sum_{\beta} \int_{I_{\beta}} \int |\sum_{\beta'=\beta-10\cdot2^{\ell}}^{\beta+10\cdot2^{\ell}} (K_{j,\ell,t} * P_{\beta'}f_{i})(x)|^{2} dx \frac{dt}{t}$$
$$\leq C2^{\ell} \sum_{\beta} \sum_{\beta'=\beta-10\cdot2^{\ell}}^{\beta+10\cdot2^{\ell}} \int_{I_{\beta}} \int |(K_{j,\ell,t} * P_{\beta'}f_{i})(x)|^{2} dx \frac{dt}{t},$$

where in the last step we use Hölder's inequality. By Proposition 5.2 (i) and (ii) we have

$$\int \left| (K_{j,\ell,t} * P_{\beta'}f_i)(x) \right|^2 dx = \int \left| \widehat{K_{j,\ell,t}}(\xi) \widehat{P_{\beta'}f_i}(\xi) \right|^2 d\xi$$
$$\lesssim 2^{-\ell M} (2^j \delta)^{-M} \int \left| \widehat{P_{\beta'}f_i}(\xi) \right|^2 d\xi.$$

Hence we control (8.2) by

$$2^{\ell}2^{-\ell M}(2^{j}\delta)^{-M}\sum_{\beta}\sum_{\beta'=\beta-10\cdot 2^{\ell}}^{\beta+10\cdot 2^{\ell}}\int_{I_{\beta}}\int \left|\widehat{P_{\beta'}f_{i}}(\xi)\right|^{2}d\xi\frac{dt}{t}$$

$$\lesssim 2^{\ell} 2^{-\ell M} (2^{j} \delta)^{-M} 2^{\ell} \delta \sum_{\beta'} \int \left| \widehat{P_{\beta'} f_{i}}(\xi) \right|^{2} d\xi$$
$$\lesssim 2^{-\ell (M-2)} \delta (2^{j} \delta)^{-M} \|f_{i}\|_{L^{2}}^{2}.$$

Case 2: $\ell = \infty$

In this case we apply Proposition 5.2 (iii) directly to obtain the control

$$\begin{split} & \int_{1}^{2} \int |K_{j,\ell,t} * f_{i}(x)|^{2} dx \frac{dt}{t} \\ \lesssim & 2^{-jM} \|f_{i}\|_{L^{2}}^{2} \\ \lesssim & \delta(2^{j}\delta)^{-M} \|f_{i}\|_{L^{2}}^{2}, \end{split}$$

where in the last step we use that $\delta \leq 1$.

By the estimates obtained in the above two cases, we can control (8.1) by

$$B^{1/2} (\delta(2^{j}\delta)^{-M})^{1/2} \sum_{\ell \ge 10} 2^{-\ell M} \|f_{i}\|_{2} \lesssim (B/A)^{1/2} \delta^{1/2} (2^{j}\delta)^{-M/2} \|f_{i}\|_{L^{2}(w)}$$

since $w(x) \ge A$ in the support of f_i .

References

- J. Bennett, A. Carbery, and T. Tao. On the multilinear restriction and Kakeya conjectures. *Acta Math.*, 196(2):261–302, 2006.
- [2] S. Bochner. Summation of multiple Fourier series by spherical means. *Trans. Amer. Math. Soc.*, 40(2):175–207, 1936.
- [3] J. Bourgain. Estimates for cone multipliers. *Geometric aspects of functional analysis* (*Israel, 1992–1994*), 77:41–60, 1995.
- [4] J. Bourgain. Some new estimates on oscillatory integrals. *Essays on Fourier analysis in honor of Elias M. Stein*, 83:83–112, 1995.
- [5] J. Bourgain and C. Demeter. The proof of the ℓ^2 decoupling conjecture. *Ann. of Math.*, 182(2):351–389, 2015.
- [6] J. Bourgain, C. Demeter, and L. Guth. Proof of the main conjecture in Vinogradov's mean value theorem for degrees higher than three. *Ann. of Math.*, 184(2):633–682, 2016.
- [7] J. Bourgain and L. Guth. Bounds on oscillatory integral operators based on multilinear estimates. *Geom. Funct. Anal.*, 21(6):1239–1295, 2011.
- [8] A. Carbery. The boundedness of the maximal Bochner-Riesz operator on $L^4(\mathbb{R}^2)$. Duke Math. J., 50(2):409–416, 1983.
- [9] A. Carbery, J. L. Rubio de Francia, and L. Vega. Almost everywhere summability of Fourier integrals. *J. London Math. Soc.*, 38(3):513–524, 1988.
- [10] L. Carleson. On convergence and growth of partial sums of Fourier series. Acta Math., 116(1):135–157, 1966.
- [11] L. Carleson and P. Sjölin. Oscillatory integrals and a multiplier problem for the disc. *Studia Math.*, 3(44):287–299, 1972.
- [12] P. Chen, X. T. Duong, D. He, S. Lee, and L. Yan. Almost everywhere convergence of Bochner-Riesz means for the Hermite operators. *Adv. Math.*, 392:108042, 2021.

- [13] K. de Leeuw. On L^p multipliers. Ann. of Math., 81(2):364–379, 1965.
- [14] X. Du, L. Guth, and X. Li. A sharp Schrödinger maximal estimate in \mathbb{R}^2 . Ann. of *Math.*, 186(2):607–640, 2017.
- [15] X. Du and R. Zhang. Sharp L² estimate of Schrödinger maximal function in higher dimensions. Ann. of Math., 189(3):837–861, 2019.
- [16] C. Fefferman. Inequalities for strongly singular convolution operators. Acta Math., 124(1):9–36, 1970.
- [17] C. Fefferman. The multiplier problem for the ball. Ann. of Math., 94:330–336, 1971.
- [18] C. Fefferman. A note on spherical summation multipliers. *Israel J. Math.*, 15(1):44– 52, 1973.
- [19] C. Fefferman. Pointwise convergence of Fourier series. Ann. of Math., 98:551–571, 1973.
- [20] R. Fefferman. A^p weights and singular integrals. Amer. J. Math., 110(5):975–987, 1988.
- [21] C. Gao, B. Liu, C. Miao, and Y. Xi. Square function estimates and local smoothing for Fourier integral operators. *Proc. Lond. Math. Soc.*, 126(6):1923–1960, 2023.
- [22] G. Garrigós and A. Seeger. On plate decompositions of cone multipliers. Proc. Edinb. Math. Soc., 52(3):631–651, 2009.
- [23] L. Grafakos. Classical Fourier analysis, volume 249. Springer, 2014.
- [24] L. Grafakos. *Modern Fourier analysis*, volume 250. Springer, 2014.
- [25] S. Guo, C. Oh, H. Wang, S. Wu, and R. Zhang. The Bochner-Riesz problem: an old approach revisited. arXiv preprint arXiv:2104.11188, 2021.
- [26] S. Guo, J. Roos, and P.-L. Yung. Sharp variation-norm estimates for oscillatory integrals related to Carleson's theorem. *Anal. PDE*, 13(5):1457–1500, 2020.
- [27] S. Guo and R. Zhang. On integer solutions of Parsell–Vinogradov systems. *Invent. Math.*, 218:1–81, 2019.
- [28] L. Guth. A restriction estimate using polynomial partitioning. J. Amer. Math. Soc., 29(2):371–413, 2016.
- [29] L. Guth, A. Iosevich, Y. Ou, and H. Wang. On Falconer's distance set problem in the plane. *Invent. Math.*, 219(3):779–830, 2020.
- [30] L. Guth, H. Wang, and R. Zhang. A sharp square function estimate for the cone in \mathbb{R}^3 . *Ann. of Math.*, 192(2):551–581, 2020.
- [31] Y. Heo. Improved bounds for high dimensional cone multipliers. *Indiana Univ. Math. J.*, 58(3):1187–1202, 2009.
- [32] Y. Heo, S. Hong, and C. Yang. An endpoint estimate for the cone multiplier. *Proc. Amer. Math. Soc.*, 138(4):1333–1347, 2010.
- [33] Y. Heo, F. Nazarov, and A. Seeger. On radial and conical Fourier multipliers. J. Geom. Anal., 21(1):96–117, 2011.
- [34] S. Hong. Weak type estimates for cone multipliers on H^p spaces, p < 1. *Proc. Amer. Math. Soc.*, 128(12):3529–3539, 2000.
- [35] L. Hörmander. Oscillatory integrals and multipliers on *FL^p*. *Ark. Mat.*, 11(1):1–11, 1973.
- [36] I. Łaba and T. Wolff. A local smoothing estimate in higher dimensions. J. Anal. Math., 88(1):149–171, 2002.
- [37] S. Lee. Improved bounds for Bochner-Riesz and maximal Bochner-Riesz operators. Duke Math. J., 122(1):205, 2004.
- [38] S. Lee and A. Vargas. On the cone multiplier in \mathbb{R}^3 . J. Funct. Anal., 263(4):925–940, 2012.

- [39] X. Li and S. Wu. New estimates of maximal Bochner-Riesz operator in the plane. *Math. Ann.*, 378(3-4):873–890, 2020.
- [40] G. Mockenhaupt. A note on the cone multiplier. Proc. Amer. Math. Soc., 117(1):145– 152, 1993.
- [41] G. Mockenhaupt, A. Seeger, and C. D. Sogge. Wave front sets, local smoothing and Bourgain's circular maximal theorem. *Ann. of Math.*, 136(1):207–218, 1992.
- [42] Y. Ou and H. Wang. A cone restriction estimate using polynomial partitioning. J. Eur. Math. Soc., 24(10):3557–3595, 2022.
- [43] E. Stein. Some problems in harmonic analysis. In Harmonic analysis in Euclidean spaces, Proceedings of the Symposium in Pure Mathematics of the Amer. Math. Soc., Williams College, Mass, Proc. Sympos. Pure Math., XXXV Part I, 1979, pages 3–20. Amer. Math. Soc., 1979.
- [44] E. Stein. Harmonic Analysis: Real-Variable Methods, Orthogonality, and Oscillatory Integrals. (PMS-43). Princeton Mathematical Series. Princeton University Press, 1993.
- [45] E. M. Stein. On limits of sequences of operators. Ann. of Math., 74(1):140–170, 1961.
- [46] T. Tao. On the maximal Bochner-Riesz conjecture in the plane for p < 2. *Trans. Amer. Math. Soc.*, 354(5):1947–1959, 2002.
- [47] T. Tao and A. Vargas. A bilinear approach to cone multipliers I. Restriction estimates. *Geom. Funct. Anal.*, 10(1):185–215, 2000.
- [48] T. Tao and A. Vargas. A bilinear approach to cone multipliers II. Applications. *Geom. Funct. Anal.*, 10(1):216–258, 2000.
- [49] T. Wolff. Local smoothing type estimates on L^p for large *p. Geom. Funct. Anal.*, 10(5):1237–1288,2000.
- [50] A. Zygmund. Trigonometric series, volume 1. Cambridge university press, 2002.

PENG CHEN, DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, 510275, P.R. CHINA

Email address: chenpeng3@mail.sysu.edu.cn

DANQING HE, SCHOOL OF MATHEMATICAL SCIENCES, FUDAN UNIVERSITY, PEO-PLE'S REPUBLIC OF CHINA

Email address: hedanqing@fudan.edu.cn

XIAOCHUN LI, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ILLINOIS AT URBANA-CHAMPAIGN, URBANA, IL, 61801, USA *Email address*: xcli@math.uiuc.edu

LIXIN YAN, DEPARTMENT OF MATHEMATICS, SUN YAT-SEN UNIVERSITY, GUANGZHOU, 510275, P.R. CHINA

Email address: mcsylx@mail.sysu.edu.cn