Emergence of a Poisson process in weakly interacting particle systems

David Padilla-Garza, Luke Peilen, Eric Thoma

Abstract

We consider the Gibbs measure of a general interacting particle system for a certain class of "weakly interacting" kernels. In particular, we show that the local point process converges to a Poisson point process as long as the inverse temperature β satisfies $N^{-1} \ll \beta \ll N^{-\frac{1}{2}}$, where N is the number of particles. This expands the temperature regime for which convergence to a Poisson point process has been proved.

1 Introduction and motivation

In this paper, we will be interested in a system of particles that interact via a pairwise interaction, and are confined by an external potential. We will consider a system of N particles that lie in d-dimensional Euclidean space. This is modeled by the Hamiltonian

$$\mathcal{H}_N(X_N) = \sum_{i \neq j} g(x_i - x_j) + N \sum_{i=1}^N V(x_i),$$
(1.1)

where $g : \mathbf{R}^d \to \mathbf{R}$ is the pair-wise interaction, $X_N := (x_1, ..., x_N) \in (\mathbf{R}^d)^N$, and $V : \mathbf{R}^d \to \mathbf{R}$ is the confining potential. We will call this an interacting particle system. We will be interested in the behaviour of such a system for large but finite N.

We are particularly interested in limiting behavior of the law of the local point process, which is defined as

$$\Xi := \sum_{i=1}^{N} \delta_{N^{1/d}(x_i - x^*)} \tag{1.2}$$

for an artibrary centering point x^* . The scaling of $N^{1/d}$ will reflect that the particles $\{x_i\}_{i=1}^N$ typically occupy a volume of order 1, so the local point process (in the weak topology) captures microscopic behavior of the system near x^* as $N \to \infty$.

Our main goal will be to prove that the law of Ξ is asymptotically Poissonian in certain temperature scalings $\beta = \beta_N$ and for a broad class of "weak interaction" kernels g. Before moving to precise statements in Section 2, we discuss more broadly the study of interacting particle systems with Hamiltonian of the form (1.1).

A very frequent form of the interaction g for $d \ge 2$ is given by the Coulomb kernel:

$$g(x) = \begin{cases} \frac{1}{|x|^{d-2}} & \text{if } d \ge 3, \\ -\log(|x|) & \text{if } d = 2. \end{cases}$$
(1.3)

Note that g satisfies that

$$-\Delta g = c_d \delta_0, \tag{1.4}$$

where δ_x denotes a Dirac delta at x, for all $d \ge 2$ and for some constant c_d depending only on d.

In d = 1, a very frequent form of g is given by

$$g(x) = -\log(|x|)$$
 if $d = 1.$ (1.5)

This is the same formula as in d = 2, but in dimension 1 g is not the fundamental solution of Laplacian (i.e. it does not satisfy equation (1.4)); instead it is a solution kernel for a fractional Laplacian.

In dimension $d \ge 1$, a generalization of the Coulomb kernel is given by the Riesz kernel:

$$g(x) = \frac{1}{|x|^{d-2s}},\tag{1.6}$$

with $0 < s < \min\{\frac{d}{2}, 1\}$. In this case g satisfies that

$$(-\Delta)^s g = c_{d,s} \delta_0. \tag{1.7}$$

For some constant $c_{d,s}$ depending only on d, s. We will refer to this setting as the Riesz case.

The study of minimizers of (1.1) when g is either a Coulomb or Riesz interaction is an active field of research, with possible applications in approximation theory [5, 11, 12, 15, 16, 21, 22, 29, 36]. However, in this paper we will take a different approach, and focus on the positive-temperature regime. In other words, we will look at the Gibbs measure associated to Hamiltonian (1.1), instead of focusing only on minimizers. This Gibbs measure is given by:

$$d\mathbb{P}_{N,\beta}(X_N) = \frac{1}{Z_{N,\beta}^V} \exp\left(-\beta \mathcal{H}_N(X_N)\right) \, dX_N,\tag{1.8}$$

where

$$Z_{N,\beta}^{V} = \int_{\mathbf{R}^{d \times N}} \exp\left(-\beta \mathcal{H}_{N}(X_{N})\right) \, \mathrm{d}X_{N}$$
(1.9)

is the partition function, and $\beta > 0$ is the inverse temperature which may depend on N.

The Gibbs measure (1.8) is linked to Random Matrix Theory if g is either a Coulomb, or a 1d log interaction. In d = 1 and if $\beta = 1, 2$ or 4 and if V is quadratic, (1.8) gives the density of eigenvalues of matrices in the Gaussian Orthogonal, Unitary or Symplectic ensembles, respectively. These ensembles are well-studied because they are determinantal ($\beta = 2$) or Pfaffian ($\beta = 1, 4$), which allows one to explicitly write down the correlation kernels. In the specific case $d = 2, \beta = 2, V(x) = |x|^2$, equation (1.8) corresponds to the Ginibre ensemble, which also has a determinantal structure. Higher dimensional Coulomb gases, and Riesz gases are also an active field of research [2, 7, 8, 26, 27, 33, 34, 40, 48, 50].

At a macroscopic level, an interacting particle system is well-described by its empirical measure, defined as

$$\operatorname{emp}_{N} := \frac{1}{N} \sum_{i=1}^{N} \delta_{x_{i}}.$$
(1.10)

The behaviour of the empirical measure is very different depending on the scaling of the temperature. If the temperature is not too large $(\frac{1}{N} \ll \beta)$, the empirical measure converges

to the equilibrium measure, denoted μ_V , and defined as the minimizer of the mean-field functional:

$$\mu_V := \operatorname{argmin}_{\mu} \mathcal{E}(\mu) + \int_{\mathbf{R}^d} V \,\mathrm{d}\mu, \qquad (1.11)$$

where

$$\mathcal{E}(\mu) := \iint_{\mathbf{R}^d \times \mathbf{R}^d} g(x - y) \,\mathrm{d}\mu_x \,\mathrm{d}\mu_y.$$
(1.12)

This convergence happens almost surely with respect to the Gibbs measure [17]. This measure has compact support as long as V grows fast enough at infinity (see section 3).

On the other hand, for large temperatures $(\beta \simeq \frac{1}{N})$, the empirical measure does not converge to the equilibrium measure. Instead, the effect of temperature and entropy is large enough that particles are not confined to a compact set. In this case, the empirical measure converges to the thermal equilibrium measure, denoted μ_{θ} and defined as

$$\mu_{\theta} := \operatorname{argmin}_{\mu} \mathcal{E}(\mu) + \int_{\mathbf{R}^d} V \,\mathrm{d}\mu + \frac{1}{\theta} \operatorname{ent}[\mu], \qquad (1.13)$$

where $\theta := N\beta$ and $ent[\cdot]$ is the Boltzmann-Gibbs entropy:

$$\operatorname{ent}[\mu] = \begin{cases} \int_{\mathbb{R}^d} \log(\frac{\mathrm{d}\mu}{\mathrm{d}x}) \,\mathrm{d}\mu & \text{if } \mu \text{ is absolutely continuous w.r.t. Lebesgue measure } \mathrm{d}x \\ \infty & \text{o.w.} \end{cases}$$

(1.14)

Unlike the empirical measure, the thermal equilibrium measure is everywhere positive (see section 3). In the case g = 0 and $V(x) = |x|^2$, which corresponds to independent Gaussian particles, the thermal equilibrium measure is a Gaussian, with variance determined by θ . Thus, the thermal equilibrium measure can be seen as an interpolation between a Gaussian probability density, and the equilibrium measure.

The empirical measure (1.10) and local point process (1.2) exhibit very different behavior in the large N limit. Due to equally weighting the N particles, the empirical measure typically exhibits a law of large numbers effect and converges (weakly) to a deterministic limit. The local point process is sensitive to (and, in the weak topology, effectively determined by) each particle in a large microscopic neighborhood of the centering point. It exhibits random, temperature dependent behavior even in the $N \to \infty$ limit, except at very low temperature.

At these low temperatures, the local point process is expected to be well approximated by minimizers of energies associated to the Gibbs measure Hamiltonian, an extremely delicate family of variational problems. For example, in certain dimensions, the local point process is conjectured to converge to a periodic lattice minimizing a next-order (renormalized) energy [35, 45] in some generality. In dimensions 8 and 24, this conjecture has also been proved, in this case due to a link to the Cohn-Kumar conjecture [35]. Given the difficulty in working with the local point process, it is often convenient to work instead with the empirical and tagged empirical fields, which are less refined but more manageable observables describing the microscopic behaviour of the particle system [34, 40, 46].

On the other extreme, in the large temperature regime $(\beta \simeq \frac{1}{N})$, the law of the local point process completely thermalizes and is given by a Poisson process in the large N limit. This was proved in the context of general integrable interactions in [24], generalizing the work of [4,28] for β -ensembles, which correspond to interacting particle systems with a Coulomb interaction. The temperature regime for Poissonian behavior is expected to be typically much wider, an expectation we confirm for our class of interactions. The microscopic behaviour of a particle system isn't necessarily asymptotically deterministic or Poissonian. For example, in the Coulomb case, if $\beta \simeq N^{1-\frac{2}{d}}$ then the microscopic behaviour is stochastic, but governed by a probability measure that favours low-renormalizedenergy configurations [25, 26, 40]. There are also interesting rigidity and tolerance properties that emerge in the $N \to \infty$ limit [13, 14, 20, 49].

This paper will be about an interacting particle system at high temperature, with a general pair-wise interaction. The study of interacting particle systems with a general interaction is a classical subject in statistical mechanics [19, 42, 43], and also an active field of study [10, 17, 18, 24]. Particle systems at high temperature have also recently drawn attention, and this subject occasionally overlaps with the analysis of general interactions [1, 10, 17, 23, 30, 31].

The analysis of interacting particle systems with a general pair-wise interaction is also linked to AI and machine learning, more specifically to neural networks. Neural networks can be used to accurately represent high-dimensional functions: given a high-dimensional function f, it may be represented as

$$f(x) := \lim_{n \to \infty} \frac{1}{n} \sum_{i=1}^{n} \varphi_i(x, \theta_i), \qquad (1.15)$$

where $\varphi_i(x, \theta_i)$ are given functions, depending on the parameter θ_i . One of the most frequent algorithms used to determine the parameters θ_i is stochastic gradient descent (SGD). Despite its ubiquity, very few rigorous results for convergence existed for SGD until recently. The approach in [37–39,51] is to model the evolution of the parameters θ_i under the SGD as a particle system with an evolution given by an SPDE. By modelling parameters as a particle system, the energy landscape for the empirical measure becomes convex, and the authors are able to show a convergence rate of $O(n^{-1})$ to the mean-field limit. In this model, the interaction between the particles depends on the error between the measurements and the approximating function, and on the functions φ_i . In general, however, it is not a Coulomb or Riesz interaction. If the φ_i are radial basis function networks, the interaction between the particles may be given by a Gaussian kernel, which is weakly interacting (see Definition 2.1). Given that the function f is high dimensional, this approach is linked to particle systems for large d as well.

The main result in this paper is that, for a specific class of interaction kernels, which we call weakly interacting, the local point process converges to a Poisson process for $N^{-1} \ll \beta \ll N^{-\frac{1}{2}}$, a wider temperature than the one considered in [24] (which, to our knowledge is currently the most general result in that direction), namely $\beta \simeq N^{-1}$. Hence, we prove that the hypothesis on the temperature scaling may be weakened for this specific class of interaction kernels. Our "weakly interacting" kernels are essentially kernels with an integrable positive Fourier transform, see Definition 2.1.

2 Main results

We will now state precisely the main results of the paper. We begin by defining exactly the class of confining potentials and interaction kernels that we deal with.

Definition 2.1. An interaction kernel g is called weakly interacting if g satisfies:

1. g(x) = g(-x). 2. $\hat{g} > 0$ a.e. 3. $\hat{g} \in L^1$. Note that, as a consequence of item 3, g is bounded below (and above). Without loss of generality, we assume that this lower bound is 0. Also as a consequence of item 3, g is continuous. Hence, we may assume without loss of generality the following:

4. g > 0 a.e.
5. g is continuous.

A confining potential V is called admissible if V satisfies:

1. V is l.s.c. 2. $\lim_{x\to\infty} V(x) = \infty$. 3. $\int_{|x|\ge 1} e^{-\alpha V(x)} dx < +\infty$ for large enough $\alpha \ge \alpha_0$.

Definition 2.2. A key quantity throughout the paper will be the potential field $h^{\mu} := g * \mu$ associated to a finite measure μ . We often take $\mu = \exp_N$.

We will now state our results for weakly interacting kernels and admissible confining potentials. As mentioned in the introduction, the main goal of the paper is to prove that the local point process converges to a Poisson point process. Along the way, we prove concentration bounds for the potential field of emp_N at given points, and asymptotes for the Laplace transform of fluctuations. These are interesting results in their own right, and so we state them in this section.

We recall the definition of the equilibrium measure μ_V in (1.11) and thermal equilibrium measure μ_{θ} in (1.13). Throughout the paper, we assume $\theta \ge \alpha_0$ from Definition 2.1.

Proposition 2.3 (Concentration bounds for the potential field). Assume that g is weakly interacting, and V is admissible. For any $k \in \mathbf{N}$, $\epsilon > 0$, and points $y_1, y_2, ..., y_k \in \mathbf{R}^d$, we have

$$\mathbb{P}_{N,\beta}\left(\left|\sum_{i=1}^{k} h^{\mathrm{emp}_{\mathrm{N}}-\mu_{\theta}}(y_{i})\right| > \epsilon\right) \le \exp\left(-N^{2}\beta\left[\frac{\Lambda\epsilon^{2}}{k^{2}} - \frac{1}{N}g(0)\right]\right),\tag{2.1}$$

for some $\Lambda > 0$ depending only on g.

The proof is found in Section 4.

Proposition 2.4 (Asymptotics of the Laplace transform of field fluctuations). Assume that g is weakly interacting, and V is admissible. Let $k \in \mathbb{N}$ and $y_1, y_2, ..., y_k \in \mathbb{R}^d$. Then

$$\mathbb{E}_{N,\beta}\left[\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mathrm{emp}_{N}}(y_{i})\right)\right)\right] = M_{N}\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})\right)\right) + A_{N},\qquad(2.2)$$

where

$$|\log M_N| \le C_1 N^{\frac{1}{2}} \beta \left(C_2 + \sum_{i=1}^k V(y_i) \right)^{\frac{1}{2}}$$

$$|A_N| \le \exp\left(-N\beta \left(g(0) + \sum_{i=1}^k V(y_i) \right) \right),$$
(2.3)

for some constants C_1, C_2 depending only on V, g, d.

The proof is found in Section 5.

We now state the main result of this paper: that the local point process converges to a Poisson point process.

Theorem 1 (Convergence to Poisson point process). Assume that g is weakly interacting, and V is admissible. Assume also that $\operatorname{ent}[\mu_V] < \infty$. Let $x^* \in \mathbf{R}^d$ and define the local point process Ξ by

$$\Xi := \sum_{i=1}^{N} \delta_{N^{\frac{1}{d}}(x_i - x^*)}.$$
(2.4)

Then, for a.e. x^* , if $N^{-1} \ll \beta \ll N^{-\frac{1}{2}}$, Ξ converges to a Poisson point process of intensity $\mu_V(x^*)$ as $N \to \infty$.

The proof is found in Section 6.

Our proof also gives precise a asymptotic for the first marginal of the Gibbs measure, defined as

$$\rho(x) := \int_{\mathbf{R}^{d-1}} \frac{1}{Z_{N,\beta}^{V}} \exp\left(-\beta \mathcal{H}_{N}(x, X_{N-1})\right) \, \mathrm{d}X_{N-1}.$$
(2.5)

The problem of finding an asymptotic for the first marginal was also addressed in [41], motivated by a link to the fractional quantum Hall effect.

Corollary 2.5 (First marginal and confinement). *More specifically, our proof shows the estimate*

$$\rho(x) = M_N'' \mu_\theta(x) + A_N'', \qquad (2.6)$$

for some M''_N, A''_N satisfying

$$\begin{aligned} |\log M_N''| &\leq C_1'' N^{\frac{1}{2}} \beta \left(C_2'' + V(x) \right)^{\frac{1}{2}} \\ |A_N''| &\leq \exp \left(-N \beta C'' \left(g(0) + V(x) \right) \right), \end{aligned}$$
(2.7)

for some constants C'', C''_1, C''_2 depending only on V, g, d.

If V is bounded below by some polynomial, and if $\beta = N^{-s}$ for some $s \in (\frac{1}{2}, 1)$, then equation (2.7) implies a confinement bound, or a bound on the probability that there is one particle outside of a compact set K: Let $K \subset \mathbf{R}^d$ be a compact set such that $V(x) + h^{\mu_V}(x) - c_{\infty}$ (see Lemma 3.1) is bounded below by a positive constant outside of K. Then:

$$\mathbb{P}_{N,\beta}\left(\exp_{N}\left(\mathbf{R}^{d}\setminus K\right)\geq\frac{1}{N}\right)\leq N\int_{\mathbf{R}^{d}\setminus K}\rho(x)\,\mathrm{d}x$$

$$\leq\exp\left(-cN^{\gamma}\right),$$
(2.8)

where $c, \gamma > 0$ depend on V, g, and K.

In the Coulomb case, results from [3] imply that any compact set K that contains a neighborhood of the support of μ_V satisfies that $V(x) + h^{\mu_V}(x) - c_{\infty}$ is bounded below by a positive constant outside of K.

Remark 2.6. Physical considerations lead one to expect that the optimal regime for convergence of the local point process to a Poisson process is $\beta \to 0$ as $N \to \infty$ at any speed. Proving this in a mathematically rigorous way is an interesting direction for future work.

Remark 2.7. Our analysis does not cover Coulomb, Riesz, or 1d-log gases, since their Fourier transform is not integrable. Proving that the local point process of a Coulomb, Riesz, or 1d-log gas converges to a Poisson process in a wider temperature regime than $\beta \simeq \frac{1}{N}$ is work in progress.

3 Preliminaries

Before giving the proof of the main results, we establish some foundational results in interacting particle systems. Some of these results may exist in the literature, but since we were unable to find a source corresponding to our specific case we present arguments for them here. To our knowledge, Lemma 3.6 and Remark 3.7 are new.

During the rest of the paper, we will commit the abuse of notation of not distinguishing between a measure and its density.

Lemma 3.1 (Equilibrium measure). Assume that g is weakly interacting, and V is admissible. Then the functional

$$\mathcal{E}_V(\mu) := \mathcal{E}(\mu) + \int_{\mathbf{R}^d} V \,\mathrm{d}\mu \tag{3.1}$$

has a unique minimizer in the space of probability measures which we call the equilibrium measure and denote μ_V . The equilibrium measure has compact support, denoted Σ , and satisfies the Euler-Lagrange equation

$$h^{\mu_{V}} + \frac{V}{2} - c_{\infty} \ge 0 \text{ in } \mathbf{R}^{d}$$

$$h^{\mu_{V}} + \frac{V}{2} - c_{\infty} = 0 \ \mu_{V} - a.e. \text{ in } \{x \in \mathbf{R}^{d} | \mu_{V}(x) > 0\},$$
(3.2)

where

$$c_{\infty} = \mathcal{E}(\mu_V) + \frac{1}{2} \int_{\mathbf{R}^d} V \,\mathrm{d}\mu.$$
(3.3)

Proof. The analogous argument can be found for the two dimensional log-gas in [44, Chapter 1], for general interactions on compact sets in [6, Chapter 4] and for a general Riesz gas in [47, Chapter 2].

Step 1: Existence and Uniqueness.

First, one notes that \mathcal{E}_V is strictly convex on the space of probability measures $\mathcal{P}(\mathbf{R}^d)$; this follows from the linearity of the map

$$\mu \mapsto \int_{\mathbf{R}^d} V \,\mathrm{d}\mu \tag{3.4}$$

and the positivity of the quadratic map

$$\mu \mapsto \mathcal{E}(\mu), \tag{3.5}$$

where positivity is a result of the positivity of the Fourier transform of g:

$$\iint_{\mathbf{R}^d \times \mathbf{R}^d} g(x-y) \, \mathrm{d}\mu_x d\mu_y = \int_{\mathbf{R}^d} (\mathbf{g} * \mu) \, \mathrm{d}\mu = \langle (\hat{g}\hat{\mu})^{\vee}, \mu \rangle = \langle \hat{g}\hat{\mu}, \overline{\hat{\mu}} \rangle > 0 \tag{3.6}$$

for $\mu \neq 0$ of finite measure. (To be completely rigorous above, one should approximate g by smooth g_n in the L^{∞} norm and send $n \to \infty$.)

Next, we claim that \mathcal{E}_V is lower semicontinuous with compact level sets, i.e. any sequence $\{\mu_n\} \in \mathcal{P}(\mathbf{R}^d)$ with bounded $\{\mathcal{E}_V(\mu_n)\}$ has a weakly convergent subsequence $\mu_{n_k} \rightharpoonup \mu \in \mathcal{P}(\mathbf{R}^d)$ and

$$\liminf_{k \to \infty} \mathcal{E}_V(\mu_{n_k}) \ge \mathcal{E}_V(\mu). \tag{3.7}$$

To prove this claim, note that (3.7) would follow immediately from the weak convergence $\mu_{n_k} \rightharpoonup \mu \in \mathcal{P}(\mathbf{R}^d)$ since V and g are both lower semicontinuous and bounded from below. One demonstrates the weak convergence by showing that the sequence $\{\mu_n\}$ is tight and using Prokhorov's theorem. Tightness follows from the growth of V at infinity coupled with the boundedness of \mathcal{E}_V . In fact, a close inspection of [47, Lemma 2.10] shows we can directly apply [47, Lemma 2.10] as soon as

$$\lim_{|x|,|y| \to +\infty} g(x-y) + \frac{V(x)}{2} + \frac{V(y)}{2} = +\infty$$
(3.8)

and g(x-y) + V(x) + V(y) is bounded below on $\mathbf{R}^d \times \mathbf{R}^d$; both are true for us because of the coercivity of V and the boundedness of g.

Therefore, (3.7) is true and existence of the minimizer μ_V follows immediately from the direct method in the calculus of variations; uniqueness is guaranteed by strict convexity of \mathcal{E}_V .

Step 2: Compact support.

The contradiction argument in [47, Lemma 2.14] goes through exactly to show compact support once one has

$$\lim_{|x|,|y| \to +\infty} g(x-y) + \frac{V(x)}{2} + \frac{V(y)}{2} = +\infty,$$
(3.9)

which we do in our case.

Step 3: Euler-Lagrange equation.

The Euler-Lagrange equation is obtained by taking convex perturbations. We follow the argument in [47, Lemma 2.14], with a few simplifications in our case.

Fix $\nu \in \mathcal{P}(\mathbf{R}^d)$ such that $\mathcal{E}_V(\nu) < +\infty$. If we consider $(1-t)\mu_V + t\nu$ for t > 0, then

$$0 \leq \mathcal{E}_{V}((1-t)\mu_{V}+t\nu) - \mathcal{E}_{V}(\mu_{V})$$

= $(1-t)^{2}\mathcal{E}(\mu_{V}) + t^{2}\mathcal{E}(\nu) + 2t(1-t) \iint_{\mathbf{R}^{d}\times\mathbf{R}^{d}} g(x-y) \,\mathrm{d}\mu_{V}(x) \,\mathrm{d}\nu(y)$
+ $(1-t) \int_{\mathbf{R}^{d}} V \,\mathrm{d}\mu_{V} + t \int_{\mathbf{R}^{d}} V \,\mathrm{d}\nu - \mathcal{E}_{V}(\mu_{V}).$ (3.10)

Expanding and rearranging, we find

$$-2t\mathcal{E}(\mu_V) - t\int_{\mathbf{R}^d} V \,\mathrm{d}\mu_V + 2t \iint_{\mathbf{R}^d \times \mathbf{R}^d} g(x-y) \,\mathrm{d}\mu_V(x) \,\mathrm{d}\nu(y) + t\int_{\mathbf{R}^d} V \,\mathrm{d}\nu + O(t^2) \ge 0 \quad (3.11)$$

or

$$2t \int_{\mathbf{R}^d} \left(h^{\mu_V} + \frac{V}{2} \right) \, \mathrm{d}\nu + O(t^2) \ge 2tc \tag{3.12}$$

with $c_{\infty} := \mathcal{E}(\mu_V) + \int_{\mathbf{R}^d} \frac{V}{2} d\mu_V = \int_{\mathbf{R}^d} \left(h^{\mu_V} + \frac{V}{2} \right) d\mu_V$. Dividing by 2t and taking the limit as $t \downarrow 0$ yields

$$\int_{\mathbf{R}^d} \left(h^{\mu_V} + \frac{V}{2} \right) \, \mathrm{d}\nu \ge c_{\infty}. \tag{3.13}$$

Since Diracs have finite \mathcal{E}_V -energy in our case, we can simplify the rest of [47, Lemma 2.14] and conclude that

$$h^{\mu_V} + \frac{V(x)}{2} \ge c_\infty \tag{3.14}$$

everywhere. Finally, choosing $\nu = \mu_V$ yields

$$c_{\infty} \le \int_{\mathbf{R}^d} \left(h^{\mu_V} + \frac{V}{2} \right) \, \mathrm{d}\mu_V = c_{\infty} \tag{3.15}$$

and since $h^{\mu_V} + \frac{V(x)}{2} \ge c_{\infty}$ everywhere we must have

$$h^{\mu_V} + \frac{V(x)}{2} = c_{\infty} \tag{3.16}$$

almost everywhere on Σ , as required.

Lemma 3.2 (Thermal equilibrium measure). Assume that g is weakly interacting, V is admissible, and let $\theta > 0$. Then the functional

$$\mathcal{E}_{\theta}(\mu) := \mathcal{E}(\mu) + \int_{\mathbf{R}^d} V \,\mathrm{d}\mu + \frac{1}{\theta} \mathrm{ent}[\mu]$$
(3.17)

has a unique minimizer in the space of probability measures which we call thermal equilibrium measure and denote μ_{θ} . The thermal equilibrium measure is everywhere positive, and satisfies the Euler-Lagrange equation

$$\mu_{\theta} = L_{\theta}^{-1} \exp\left(-\theta \left(2h^{\mu_{\theta}} + V\right)\right), \qquad (3.18)$$

where

$$L_{\theta} = \int_{\mathbf{R}^d} \exp\left(-\theta \left(2h^{\mu_{\theta}}(x) + V(x)\right)\right) \,\mathrm{d}x. \tag{3.19}$$

Proof. The existence of the thermal equilibrium measure follows for instance from [3, Lemma 2.1], which only uses that g is bounded from below and that $V \to +\infty$ and satisfies the exponential integrability assumption

$$\int_{|x|\ge 1} e^{-\frac{\theta}{2}V(x)} \,\mathrm{d}x < +\infty.$$
(3.20)

(3.18) then follows immediately from the form of the thermal equilibrium measure given in [24, Proposition A.2], since our kernel g satisfies all of the hypotheses of that proposition.

Once we have existence of the thermal equilibrium measure, $N^2 \mathcal{E}_{\theta}$ gives the leading order asymptotics of the Hamiltonian. To analyze next-order behavior, it is then useful to split off this deterministic term and consider the remainder. This can be accomplished by the following *splitting formula* and its corollary; the version stated below is exactly [2, Lemma 2.1], which only relies on the form of the Hamiltonian \mathcal{H}_N . **Lemma 3.3** (Thermal splitting formula). Let $\theta > 0$, and define

$$\zeta_{\theta} := -\frac{1}{\theta} \log(\mu_{\theta}). \tag{3.21}$$

Then for any point configuration $X_N \in \mathbf{R}^{d \times N}$ the Hamiltonian \mathcal{H}_N can be rewritten (split) as

$$\mathcal{H}_N(X_N) = N^2 \left(\mathcal{E}_\theta(\mu_\theta) + \mathsf{F}_N(X_N, \mu_\theta) + N \sum_{i=1}^N \zeta_\theta(x_i) \right), \tag{3.22}$$

where, given a measure μ ,

$$\mathsf{F}_N(X_N,\mu) := \frac{1}{N^2} \sum_{i \neq j} g(x_i - x_j) + \mathcal{E}(\mu) - \frac{2}{N} \sum_{i=1}^N h^\mu(x_i).$$
(3.23)

A computation then immediately yields the following equivalent definition of the Gibbs measure.

Corollary 3.4. The Gibbs measure may be rewritten as

$$d\mathbb{P}_{N,\beta}(X_N) = \frac{1}{K_{N,\beta}^{\theta}} \exp\left(-N^2\beta \left[\mathsf{F}_N(X_N,\mu_{\theta})\right]\right) \prod_{i=1}^N \mu_{\theta}(x_i) \, \mathrm{d}X_N,\tag{3.24}$$

where

$$K_{N,\beta}^{\theta} = \frac{Z_{N,\beta}^{V}}{\exp\left(N^{2}\beta\mathcal{E}_{\theta}(\mu_{\theta})\right)}.$$
(3.25)

We call $K_{N,\beta}^{\theta}$ the next-order partition function. A useful fact is that its logarithm is always non-negative.

Lemma 3.5. Assume that g is weakly interacting, and V is admissible. Then the next-order partition function is greater than 1, i.e.

$$\log K_{N,\beta}^{\theta} > 0. \tag{3.26}$$

Proof. The proof is as in [32, Proposition 5.10], which considered the analogous question for the Coulomb gas. The result follows immediately from the definition of the Gibbs measure as the probability measure that minimizes the free energy functional

$$\mathbb{Q} \mapsto \int_{\mathbf{R}^{d \times N}} \mathcal{H}_N(X_N) \, \mathrm{d}\mathbb{Q}\left(X_N\right) + \frac{1}{\beta} \mathrm{ent}[\mathbb{Q}] \tag{3.27}$$

over all $\mathbb{Q} \in \mathcal{P}\left(\mathbf{R}^{d \times N}\right)$. Inserting $\mathbb{Q} = \mathbb{P}_{N,\beta}$ yields

$$-\frac{\log Z_{N,\beta}}{\beta} = \inf_{\mu \in \mathcal{P}(\mathbf{R}^{d \times N})} \int_{\mathbf{R}^{d \times N}} \mathcal{H}_N(X_N) \,\mathrm{d}\mu(X_N) + \frac{1}{\beta} \mathrm{ent}[\mu].$$
(3.28)

Inserting instead $\mu_{\theta}^{\otimes N}$ and computing yields

$$-\frac{\log Z_{N,\beta}}{\beta} \le N^2 \mathcal{E}_{\theta}(\mu_{\theta}) - \frac{N}{2} \mathcal{E}(\mu_{\theta}) < N^2 \mathcal{E}_{\theta}(\mu_{\theta})$$
(3.29)

by positivity of g. The splitting formula (3.22) then yields

$$-\frac{\log Z_{N,\beta}}{\beta} = \frac{-\log K_{N,\beta}^{\theta} + \beta N^2 \mathcal{E}(\mu_{\theta})}{\beta} \le N^2 \mathcal{E}_{\theta}(\mu_{\theta}) \implies \log K_{N,\beta}^{\theta} > 0.$$
(3.30)

From this point forward, we present results on the thermal equilibrium measure that are not essentially taken from existing literature.

Lemma 3.6 (Convergence). Assume that g is weakly interacting, and V is admissible. Suppose also that $\operatorname{ent}[\mu_V] < \infty$. Then, as $\theta \to \infty$, μ_{θ} converges to μ_V in the L^1 topology, and $h^{\mu_{\theta}}$ converges to h^{μ_V} in the L^{∞} topology. Furthermore, for a.e. $x \in \mathbf{R}^d$ and any sequence $\delta(\theta)$ such that $\lim_{\theta\to\infty} \delta = 0$,

$$\lim_{\theta \to \infty} \oint_{B(x,\delta)} |\mu_{\theta}(s) - \mu_{V}(x)| \, \mathrm{d}s = 0.$$
(3.31)

Proof. In the case where g is the Coulomb interaction, very precise convergence is established by connections with the classical obstacle problem (cf. [3, Theorem 1]). However, that machinery is not available to us in the general interaction case.

Step 1: We will first prove that $\mu_{\theta} \to \mu_V$ in the topology of weak convergence of probability measures. We will use the language of Γ -convergence (see [9]).

Step 1.1: Γ – lim inf inequality.

The lack of positivity of the entropy presents a difficulty, which we circumvent by a trick as in [3]; namely, we can rewrite the function \mathcal{E}_{θ} as

$$\mathcal{E}_{\theta}(\mu) = \mathcal{E}(\mu) + \left(1 - \frac{\alpha_0}{\theta}\right) \int_{\mathbf{R}^d} V \,\mathrm{d}\mu + \left(\int \frac{\alpha_0}{\theta} V \,\mathrm{d}\mu + \mathrm{ent}[\mu]\right),\tag{3.32}$$

where α_0 is as in Definition 2.1. Consider a sequence $\theta_n \to +\infty$ and suppose that $\mu_n \rightharpoonup \mu$. Then, since g and V are both lower semicontinuous and bounded from below, we have for any $\epsilon > 0$ that

$$\liminf_{n \to \infty} \left(\mathcal{E}(\mu_n) + \int_{\mathbf{R}^d} \left(1 - \frac{\alpha_0}{\theta_n} \right) V \, \mathrm{d}\mu_n \right) \ge \liminf_{n \to \infty} \left(\mathcal{E}(\mu_n) + \int_{\mathbf{R}^d} \left(1 - \epsilon \right) V \, \mathrm{d}\mu_n \right) \qquad (3.33)$$
$$\ge \mathcal{E}(\mu) + \int_{\mathbf{R}^d} (1 - \epsilon) V \, d\mu.$$

Now, if μ_n is not absolutely continuous with respect to Lebesgue measure for all $n \ge N$ for some N, then $\operatorname{ent}[\mu_n]$ is infinite for all $n \ge N$ and so

$$\liminf_{n \to \infty} \left(\int_{\mathbf{R}^d} \frac{\alpha_0}{\theta_n} V \,\mathrm{d}\mu_n + \frac{1}{\theta_n} \mathrm{ent}[\mu_n] \right) = +\infty \ge 0.$$
(3.34)

Otherwise, extracting the subsequence μ_{n_k} that are absolutely continuous to Lebesgue measure yields

$$\liminf_{n \to \infty} \left(\int_{\mathbf{R}^d} \frac{\alpha_0}{\theta_n} V \, \mathrm{d}\mu_n + \frac{1}{\theta_n} \mathrm{ent}[\mu_n] \right) \ge \liminf_{k \to \infty} \int_{\mathbf{R}^d} \left(\frac{\alpha_0 V}{\theta_{n_k}} + \frac{1}{\theta_{n_k}} \log \mu_{n_k} \right) \, d\mu_{n_k}$$
$$\ge \liminf_{k \to \infty} \frac{-1}{\theta_{n_k}} \int_{\mathbf{R}^d} e^{-\alpha_0 V(x) - 1} \, dx = 0$$

where we have used the inequality

$$\frac{c\gamma}{\theta} + \frac{\gamma\log\gamma}{\theta} \ge -\frac{1}{\theta}e^{-c-1},$$

with $c = \alpha_0 V$ and $\gamma = \mu_{n_k}$, which can be seen by minimizing the above as a function of γ . In either case,

$$\liminf_{n \to \infty} \mathcal{E}_{\theta_n}(\mu_n) \ge \mathcal{E}(\mu) + \int_{\mathbf{R}^d} (1 - \epsilon) V \ d\mu.$$

Since $\epsilon > 0$ was arbitrary, we conclude by monotone convergence that $\liminf_{n \to +\infty} \mathcal{E}_{\theta_n}(\mu_n) \geq \mathcal{E}_V(\mu)$.

Step 1.2: Γ – lim sup inequality.

Under the hypothesis that $\operatorname{ent}[\mu_V] < \infty$, it is trivially true that

$$\lim_{\theta \to \infty} \mathcal{E}_{\theta}(\mu_V) = \mathcal{E}_V(\mu_V). \tag{3.35}$$

Step 1.3: Uniform coercivity and conclusion of step 1.

Now we show that $\{\mu_{\theta_n}\}$ is a tight sequence of probability measures. This will follow from the boundedness of the sequence $\{\mathcal{E}_{\theta_n}(\mu_{\theta_n})\}$ in the same way as Lemma 4.1 (following [47, Lemma 2.10]). Without loss of generality, assume $\theta_1 = \inf \theta_n$. First, the boundedness follows from $\frac{1}{\theta_n} \to 0$ and the minimality of μ_{θ_n} :

$$\mathcal{E}_{\theta_n}(\mu_{\theta_n}) \le \mathcal{E}_{\theta_n}(\mu_{\theta_1}) \le \mathcal{E}_{\theta_1}(\mu_{\theta_1}) := C_1 < +\infty$$
(3.36)

for all n. Now, for any $C_2 > 0$, there is a compact set $K \times K$ outside of which

$$g(x-y) + \gamma_n \frac{V(x)}{2} + \gamma_n \frac{V(y)}{2} > C_2$$
(3.37)

with $\gamma_n = \left(1 - \frac{\alpha_0}{\theta_n}\right)$ independently of *n* by our growth assumption on *V* (we can choose this independently of *n* since $\theta_n \to +\infty$). Then, moving some of the potential onto the entropy as above, we have

$$C_{1} \geq \mathcal{E}(\mu_{\theta_{n}}) + \left(1 - \frac{\alpha_{0}}{\theta_{n}}\right) \int_{\mathbf{R}^{d}} V \,\mathrm{d}\mu_{\theta_{n}} + \int_{\mathbf{R}^{d}} \left(\frac{\alpha_{0}}{\theta_{n}}V + \frac{1}{\theta_{n}}\log\mu_{\theta_{n}}\right) \,\mathrm{d}\mu_{\theta_{n}}$$

$$\geq \iint_{\mathbf{R}^{d} \times \mathbf{R}^{d}} \left(g(x - y) + \gamma_{n} \frac{V(x)}{2} + \gamma_{n} \frac{V(y)}{2}\right) \,\mathrm{d}\mu_{\theta_{n}}(x) \,\mathrm{d}\mu_{\theta_{n}}(y) - \frac{1}{\theta_{n}} \int_{\mathbf{R}^{d}} e^{-\alpha_{0}V(x) - 1} \,\mathrm{d}x$$

$$\geq -C_{3} + C_{2}\mu_{\theta_{n}} \otimes \mu_{\theta_{n}}((K \times K)^{c})$$

$$\geq -C_{3} + C_{2}\mu_{\theta_{n}}(K^{c}),$$

where we have used that $g(x-y) + \gamma_n \frac{V(x)}{2} + \gamma_n \frac{V(y)}{2}$ is everywhere bounded below independently of n since $|\gamma_n - 1|$ is small, and $-\frac{1}{\theta_n} \int_{\mathbf{R}^d} e^{-\alpha_0 V(x) - 1} dx$ is bounded below as well since $\theta_n \to \infty$. Since C_2 can be made arbitrarily large, this gives us the tightness of the sequence $\{\mu_{\theta_n}\}$.

We can now conclude as in [9, Theorem 1.21], or as in [47, Theorem 2.2]. Since $\{\mu_{\theta_n}\}$ is a tight sequence, it has a weak limit μ . Furthermore, since μ_V has a density, we have

$$\mathcal{E}_{V}(\mu_{V}) \geq \limsup_{n \to +\infty} \mathcal{E}_{\theta_{n}}(\mu_{V}) \geq \limsup_{n \to +\infty} \mathcal{E}_{\theta_{n}}(\mu_{\theta_{n}}) \geq \mathcal{E}_{V}(\mu).$$
(3.38)

The second inequality follows from the fact that μ_{θ_n} minimizes \mathcal{E}_{θ_n} , and the third inequality is a result of lower semicontinuity. Since the equilibrium measure is unique, we must have $\mu = \mu_V$ and we see that μ_{θ_n} converges weakly to μ_V under the topology of weak convergence. **Step 2:** Now, we prove that $\mu_{\theta} \to \mu_V$ in the L^1 topology.

Note that, by definition of μ_{θ} , we have that, for any $\theta > 0$,

$$\operatorname{ent}[\mu_{\theta}] \le \operatorname{ent}[\mu_{V}] < \infty. \tag{3.39}$$

In particular,

$$\overline{\lim}_{\theta \to \infty} \operatorname{ent}[\mu_{\theta}] \le \operatorname{ent}[\mu_{V}]. \tag{3.40}$$

On the other hand, by l.s.c.,

$$\operatorname{ent}[\mu_V] \le \underline{\lim}_{\theta \to \infty} \operatorname{ent}[\mu_{\theta}], \qquad (3.41)$$

which implies that

$$\lim_{\theta \to \infty} \operatorname{ent}[\mu_{\theta}] = \operatorname{ent}[\mu_{V}]. \tag{3.42}$$

Given $\epsilon > 0$, we define the probability measure μ_V^{ϵ} as

$$\mu_V^{\epsilon} := \frac{\mu_V + \epsilon \exp\left(-\alpha_0 V\right)}{\int_{\mathbf{R}^d} \mu_V + \epsilon \exp\left(-\alpha_0 V\right)},\tag{3.43}$$

where α_0 is as in Definition 2.1. Then, by Pinsker's inequality,

$$\begin{split} \limsup_{\theta \to \infty} \|\mu_{\theta} - \mu_{V}^{\epsilon}\|_{L^{1}} &\leq \limsup_{\theta \to \infty} \sqrt{\int_{\mathbf{R}^{d}} \log\left(\frac{\mu_{\theta}}{\mu_{V}^{\epsilon}}\right) \mu_{\theta}} \\ &= \limsup_{\theta \to \infty} \sqrt{\operatorname{ent}[\mu_{\theta}] - \int_{\mathbf{R}^{d}} \log\left(\mu_{V}^{\epsilon}\right) \mu_{\theta}} \\ &= \sqrt{\operatorname{ent}[\mu_{V}] - \int_{\mathbf{R}^{d}} \log\left(\mu_{V}^{\epsilon}\right) \mu_{V}}. \end{split}$$
(3.44)

Furthermore, by triangle inequality, there holds

$$\limsup_{\theta \to \infty} \|\mu_{\theta} - \mu_{V}\|_{L^{1}} \leq \sqrt{\operatorname{ent}[\mu_{V}]} - \int_{\mathbf{R}^{d}} \log\left(\mu_{V}^{\epsilon}\right) \mu_{V} + \|\mu_{V} - \mu_{V}^{\epsilon}\|_{L^{1}}.$$
 (3.45)

Letting ϵ tend to 0, we may conclude that $\mu_\theta \to \mu_V$ in the L^1 topology.

Note that $||g||_{L^{\infty}} < \infty$ by property 3 of Definition 2.1. Hence, Young's convolution inequality, implies that $h^{\mu_{\theta}} \to h^{\mu_{V}}$ in L^{∞} .

Step 3: Finally, we prove that for a.e. $x \in \mathbf{R}^d$ and any sequence $\delta(\theta)$ such that $\lim_{\theta \to \infty} \delta = 0$,

$$\lim_{\theta \to \infty} \oint_{B(x,\delta)} |\mu_{\theta}(s) - \mu_{V}(x)| \, \mathrm{d}s = 0.$$
(3.46)

To prove this claim, note that, for any $\theta > 0$ and a.e. x, we have

$$\begin{aligned} \oint_{B(x,\delta)} |\mu_{\theta}(s) - \mu_{V}(x)| \, \mathrm{d}s \\ &\leq \int_{B(x,\delta)} |\mu_{\theta}(s) - \mu_{V}(s)| \, \mathrm{d}s + \int_{B(x,\delta)} |\mu_{V}(s) - \mu_{V}(x)| \, \mathrm{d}s \\ &\leq M(\mu_{\theta} - \mu_{V})(x) + \int_{B(x,\delta)} |\mu_{V}(s) - \mu_{V}(x)| \, \mathrm{d}s. \end{aligned}$$
(3.47)

where $M(\cdot)$ denotes the Hardy-Littlewood maximal function. Let $\epsilon>0.$ Then

$$\overline{\lim}_{\theta \to \infty} \left| \left\{ x \in \mathbf{R}^{d} : \int_{B(x,\delta)} |\mu_{\theta}(s) - \mu_{V}(x)| \, \mathrm{d}s > \epsilon \right\} \right| \\
\leq \overline{\lim}_{\theta \to \infty} \left| \left\{ x \in \mathbf{R}^{d} : |M(\mu_{\theta} - \mu_{V})(x)| > \epsilon \right\} \right| \\
+ \overline{\lim}_{\theta \to \infty} \left| \left\{ x \in \mathbf{R}^{d} : \int_{B(x,\delta)} |\mu_{V}(s) - \mu_{V}(x)| \, \mathrm{d}s > \epsilon \right\} \right|.$$
(3.48)

By the Hardy-Littlewood maximal inequality, and step 2, there holds

$$\overline{\lim}_{\theta \to \infty} \left| \left\{ x \in \mathbf{R}^d : |M(\mu_\theta - \mu_V)(x)| > \epsilon \right\} \right| \le \overline{\lim}_{\theta \to \infty} \frac{C_d}{\epsilon} \left\| \mu_\theta - \mu_V \right\|_{L^1} = 0.$$
(3.49)

On the other hand, Lebesgue's Theorem implies that

$$\overline{\lim}_{\theta \to \infty} \left| \left\{ x \in \mathbf{R}^d : \int_{B(x,\delta)} |\mu_V(s) - \mu_V(x)| \, \mathrm{d}s > \epsilon \right\} \right| = 0.$$
(3.50)

Since ϵ is arbitrary, we may conclude.

Remark 3.7 (Qualitative behaviour). The constant L_{θ} defined in (3.19) satisfies that

$$\lim_{\theta \to \infty} -\frac{1}{\theta} \log L_{\theta} = 2c_{\infty}, \qquad (3.51)$$

where c_{∞} is given by (3.3). Furthermore, for all $\theta > 0$ there exists a constant c_{θ} such that, for any $\theta' \geq \theta$ there holds

$$\mu_{\theta'}(x) \le \exp\left(-\theta' \left(h^{\mu_V}(x) + V(x) - 2c_{\theta}\right)\right).$$
(3.52)

Also, $\lim_{\theta\to\infty} c_\theta = c_\infty$.

Proof. Using Lemma 3.6, and equations (3.2) and (3.18) we have that

$$\lim_{\theta \to \infty} -\frac{1}{\theta} \log L_{\theta} = \inf_{x \in \mathbf{R}^d} 2h^{\mu_V}(x) + V(x)$$

= $2c_{\infty}$. (3.53)

The rest of the remark follows from equations (3.18), (3.51) by taking

$$c_{\theta} := \sup_{\theta' \ge \theta} -\frac{1}{2\theta'} \log L_{\theta'} + \|h^{\mu_V} - h^{\mu_{\theta'}}\|_{L^{\infty}}.$$
 (3.54)

4 Proof of Proposition 2.3

We now prove Proposition 2.3, restated here for convenience.

Proposition 4.1 (Concentration bounds for the potential field). Assume that g is weakly interacting, and V is admissible. Let $k \in \mathbf{N}$, $\epsilon > 0$, and $y_1, y_2, ..., y_k \in \mathbf{R}^d$. Then

$$\mathbb{P}_{N,\beta}\left(\left|\sum_{i=1}^{k} h^{\mathrm{emp}_{\mathrm{N}}-\mu_{\theta}}(y_{i})\right| > \epsilon\right) \le \exp\left(-N^{2}\beta\left[\frac{\Lambda\epsilon^{2}}{k^{2}} - \frac{1}{N}g(0)\right]\right),\tag{4.1}$$

for some $\Lambda > 0$ depending only on g.

Proof. Step 1: Starting point.

Using the splitting formula (Lemma 3.3), we have that

$$\mathbb{P}_{N,\beta} \left(\left| \sum_{i=1}^{k} h^{\operatorname{emp}_{N}-\mu_{\theta}}(y_{i}) \right| > \epsilon \right) \\
= \frac{1}{K_{N,\beta}} \int_{\left\{ X_{N}: \left| \sum_{i=1}^{k} h^{\operatorname{emp}_{N}-\mu_{\theta}}(y_{i}) \right| > \epsilon \right\}} \exp\left(-N^{2}\beta \mathsf{F}_{N}(\operatorname{emp}_{N},\mu_{\theta})\right) d\mu_{\theta}^{\otimes N} \\
\leq \frac{1}{K_{N,\beta}} \sup_{\left\{ X_{N}: \left| \sum_{i=1}^{k} h^{\operatorname{emp}_{N}-\mu_{\theta}}(y_{i}) \right| > \epsilon \right\}} \exp\left(-N^{2}\beta \mathsf{F}_{N}(\operatorname{emp}_{N},\mu_{\theta})\right) \\
\leq \frac{1}{K_{N,\beta}} \sup_{\left\{ X_{N}: \left| \sum_{i=1}^{k} h^{\operatorname{emp}_{N}-\mu_{\theta}}(y_{i}) \right| > \epsilon \right\}} \exp\left(-N^{2}\beta \left[\mathcal{E}(\operatorname{emp}_{N}-\mu_{\theta}) - \frac{1}{N}g(0) \right] \right) \\
\leq \frac{1}{K_{N,\beta}} \exp\left(-N^{2}\beta \left[\Phi_{y_{1},\ldots,y_{k}}(\epsilon) - \frac{1}{N}g(0) \right] \right),$$
(4.2)

where

$$\Phi_{y_1,\dots y_k}(\epsilon) := \inf_{\left\{\mu \in \mathcal{M}(\mathbf{R}^d): \left|\sum_{i=1}^k h^\mu(y_i)\right| > \epsilon\right\}} \mathcal{E}(\mu).$$
(4.3)

We will also use the obvious variants of Φ with different values of k.

Step 2: Solving the variational problem.

We claim that for any $y_1, \dots y_k \in \mathbf{R}^d$,

$$\Phi_{y_1,\dots y_k}(\epsilon) \ge \frac{\Lambda \epsilon^2}{k^2},\tag{4.4}$$

for some $\Lambda > 0$.

Substep 2.1: We start by proving that

$$\Lambda := \Phi_0(1) > 0. \tag{4.5}$$

To prove this claim, note that for any $\mu \in \mathcal{M}(\mathbf{R}^d)$,

$$h^{\mu}(0) = \int_{\mathbf{R}^d} \widehat{h^{\mu}}(\xi) \,\mathrm{d}\xi = \int_{\mathbf{R}^d} \widehat{\mu}(\xi) \widehat{g}(\xi) \,\mathrm{d}\xi.$$
(4.6)

On the other hand,

$$\mathcal{E}(\mu) = \int_{\mathbf{R}^d} \left| \widehat{\mu}(\xi) \right|^2 \widehat{g}(\xi) \,\mathrm{d}\xi.$$
(4.7)

By Cauchy-Schwartz, and since \hat{g} is positive a.e., we find

$$\left(\int_{\mathbf{R}^d} \widehat{\mu}(\xi)\widehat{g}(\xi) \,\mathrm{d}\xi\right)^2 \le \left(\int_{\mathbf{R}^d} |\widehat{\mu}(\xi)|^2 \,\widehat{g}(\xi) \,\mathrm{d}\xi\right) \left(\int_{\mathbf{R}^d} \widehat{g}(\xi) \,\mathrm{d}\xi\right),\tag{4.8}$$

and therefore for any $\mu \in \mathcal{M}(\mathbf{R}^d)$ such that $h^{\mu}(0) = 1$, there holds

$$\mathcal{E}(\mu) \ge \left(\int_{\mathbf{R}^d} \widehat{g}(\xi) \,\mathrm{d}\xi\right)^{-1},\tag{4.9}$$

which finishes the substep.

Substep 2.2: We now prove the claim stated at the beginning of step 2.

Note that if $\left|\sum_{i=1}^{k} h^{\mu}(y_i)\right| > \epsilon$ then $|h^{\mu}(y_i)| > \frac{\epsilon}{k}$ for some $i \in \{1, ..., k\}$ by a pigeonhole argument. Clearly, $\Phi_{y_i} = \Phi_0$ by translation invariance and $\epsilon \mapsto \Phi_0(\epsilon)$ is quadratic. It follows

$$\Phi_{y_1,\dots,y_k}(\epsilon) \ge \Phi_0\left(\frac{\epsilon}{k}\right) = \frac{\Phi_0(1)\epsilon^2}{k^2} \ge \frac{\Lambda\epsilon^2}{k^2}.$$
(4.10)

By plugging equation (4.4) into the last line of equation (4.2). we may conclude Proposition 2.3. \Box

5 Proof of Proposition 2.4

In this section, we prove Proposition 2.4, restated here for convenience.

Proposition 5.1 (Asymptotics of the Laplace transform of field fluctuations). Assume that g is weakly interacting, and V is admissible. Let $k \in \mathbb{N}$ and $y_1, y_2, ..., y_k \in \mathbb{R}^d$. Then

$$\mathbb{E}_{N,\beta}\left[\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mathrm{emp}_{N}}(y_{i})\right)\right)\right] = M_{N}\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})\right)\right) + A_{N},\qquad(5.1)$$

where

$$|\log M_N| \le C_1 N^{\frac{1}{2}} \beta \left(C_2 + \sum_{i=1}^k V(y_i) \right)^{\frac{1}{2}}$$

$$|A_N| \le \exp\left(-N\beta \left(g(0) + \sum_{i=1}^k V(y_i) \right) \right),$$
(5.2)

for some constants C_1, C_2 depending only on V, g, d.

Proof. We will only show a complete proof of the lower bound in equation (5.1). The proof of the upper bound is analogous.

Step 1: First simplification.

Using the cake-slicing formula, we have that

$$\mathbb{E}_{N,\beta}\left[\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mathrm{emp}_{\mathrm{N}}}(y_{i})\right)\right)\right] = \int_{0}^{1}\mathbb{P}_{N,\beta}\left(\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mathrm{emp}_{\mathrm{N}}}(y_{i})\right)\right) \ge x\right)\mathsf{d}x.$$
(5.3)

We have used the hypothesis that g > 0 (item 5 of Definition 2.1) to limit the domain of integration to the (0, 1) interval.

We will split the integral at the point $\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})-T\right)\right)$, for T to be determined later. The reason for this is to optimize the application of the concentration bound, Proposition 2.3. Using this splitting of the integral, we have that

$$\mathbb{E}_{N,\beta} \left[\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\exp_{N}}(y_{i})\right)\right) \right] \\= \int_{0}^{\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mu_{\theta}}(y_{i})-T\right)\right)} \mathbb{P}_{N,\beta} \left(\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\exp_{N}}(y_{i})\right)\right) \ge x \right) dx \\+ \int_{\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mu_{\theta}}(y_{i})-T\right)\right)}^{1} \mathbb{P}_{N,\beta} \left(\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\exp_{N}}(y_{i})\right)\right) \ge x \right) dx$$
(5.4)
$$\leq \exp\left(N\beta T\right) \exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mu_{\theta}}(y_{i})\right)\right) \\+ \int_{\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mu_{\theta}}(y_{i})-T\right)\right)}^{1} \mathbb{P}_{N,\beta} \left(\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\exp_{N}}(y_{i})\right)\right) \ge x \right) dx.$$

The rest of the proof will be about estimating the last term in equation (5.4). **Step 2:** Estimating the remaining integral. Using Proposition 2.3, we have that

Using Proposition
$$2.3$$
, we have that

$$\int_{\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})-T\right)\right)}^{1} \mathbb{P}_{N,\beta}\left(\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\exp}(y_{i})\right)\right) \ge x\right) dx$$

$$= \int_{\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})-T\right)\right)}^{1} \mathbb{P}_{N,\beta}\left(\sum_{i=1}^{k}h^{\exp}(y_{i}) \ge \frac{-\log x}{N\beta}\right) dx$$

$$= \int_{\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})-T\right)\right)}^{1} \exp\left(-N^{2}\beta\left[\frac{\Lambda\left(\frac{\log x}{N\beta}-\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})\right)^{2}}{k^{2}}-\frac{1}{N}g(0)\right]\right) dx.$$
(5.5)

Note that, if we define $T := \sqrt{\frac{k^2}{\Lambda N}} \left(2g(0) + \sum_{i=1}^k V(y_i) \right)$, then for $x \in \left(\exp\left(-N\beta\left(\sum_{i=1}^k h^{\mu_\theta}(y_i) - T\right)\right), 1 \right)$, there holds

$$\frac{\Lambda\left(\frac{\log x}{N\beta} - \sum_{i=1}^{k} h^{\mu_{\theta}}(y_i)\right)^2}{k^2} - \frac{1}{N}g(0) \ge \frac{1}{N}\left(g(0) + \sum_{i=1}^{k} V(y_i)\right).$$
(5.6)

Since the integration interval is bounded in measure by 1, we have that

$$\int_{\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\mu_{\theta}}(y_{i})-T\right)\right)}^{1} \mathbb{P}_{N,\beta}\left(\exp\left(-N\beta\left(\sum_{i=1}^{k}h^{\exp}(y_{i})\right)\right) \ge x\right) dx \quad (5.7)$$

$$\leq \exp\left(-N\beta\left(g(0)+\sum_{i=1}^{k}V(y_{i})\right)\right).$$

This concludes the proof of the lower bound of equation (5.1).

Step 3: Upper bound sketch.

In order to prove the upper bound of equation (5.1), we proceed similarly:

$$\mathbb{E}_{N,\beta} \left[\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mathrm{emp}_{N}}(y_{i})\right)\right) \right]$$

$$= \int_{0}^{1} \left[1 - \mathbb{P}_{N,\beta} \left(\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mathrm{emp}_{N}}(y_{i})\right)\right) \le x \right) \right] \mathrm{d}x \qquad (5.8)$$

$$\geq \int_{0}^{\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mu_{\theta}}(y_{i})+T\right)\right)} \left[1 - \mathbb{P}_{N,\beta} \left(\exp\left(-N\beta\left(\sum_{i=1}^{k} h^{\mathrm{emp}_{N}}(y_{i})\right)\right) \le x \right) \right] \mathrm{d}x.$$

The rest of the proof is analogous and hence omitted.

6 Proof of Theorem 1

In this section, we prove the main result of this paper, namely Theorem 1. We will use some background on point processes, which can be reviewed in the Appendix. The main result that we will use is the following lemma:

Lemma 6.1 (Weak convergence and correlation functions). Let \mathfrak{P}_N be a sequence of point processes, and denote their k- point correlation functions by R_k^N . Then \mathfrak{P}_N converges weakly to a point process \mathfrak{P} , with k- point correlation functions denoted by R_k^N if:

- 1. $R_k^N \to R_k$ pointwise almost everywhere as $N \to \infty$.
- 2. For any compact set $\Omega \subset \mathbf{R}^d$ there holds

$$\sup_{N \in \mathbf{N}} \sum_{k=1}^{\infty} \frac{1}{k!} \int_{K} R_k^N \, \mathrm{d}y_1 \dots \, \mathrm{d}y_k < \infty.$$
(6.1)

Equipped with Lemma 6.1, we will prove Theorem 1, restated here for convenience.

Theorem 1 (Convergence to Poisson point process). Assume that g is weakly interacting, and V is admissible. Assume also that $\operatorname{ent}[\mu_V] < \infty$. Let $x^* \in \mathbf{R}^d$ and define the local point process Ξ by

$$\Xi := \sum_{i=1}^{N} \delta_{N^{\frac{1}{d}}(x_i - x^*)}.$$
(6.2)

Then, for a.e. x^* , if $N^{-1} \ll \beta \ll N^{-\frac{1}{2}}$, Ξ converges to a Poisson point process of intensity $\mu_V(x^*)$.

Proof. Step 1: Formula for the correlation functions. Let $Y_N := (y_1, ..., y_N) \in \mathbf{R}^{d \times k}$, $x_i := x^* + N^{-\frac{1}{d}}(y_i - x^*)$, and $X_N = (x_1, ..., x_N)$. Using Proposition 2.4 and Remark 7.4, the *k*-point correlation function can be written as

$$R_k^N(Y_k) = \frac{N!}{(N-k)!} \frac{1}{Z_{N,\beta}} \int_{\mathbf{R}^{d(N-k)}} \exp\left(-\beta \mathcal{H}_N(X_k, x_{k+1}, \dots x_N)\right) \, \mathrm{d}y_{k+1} \dots \, \mathrm{d}y_N$$

$$= \operatorname{err}_N \frac{1}{Z_{N,\beta}} \int_{\mathbf{R}^{d(N-k)}} \exp\left(-\beta \left(\sum_{1 \le i < j \le N} g(x_i - x_j) + N \sum_{i=1}^N V(x_i)\right)\right)\right) \, \mathrm{d}x_{k+1} \dots \, \mathrm{d}x_N$$

$$= \operatorname{err}_N \frac{Z_{N-k,\beta}}{Z_{N,\beta}} \exp\left(-\beta \left(\sum_{1 \le i < j \le k} g(x_i - x_j) + N \sum_{i=1}^k V(x_i)\right)\right)\right)$$

$$\times \mathbb{E}_{N-k,\beta} \left[\exp\left(-(N-k)\beta \left(\sum_{i=1}^k h^{\operatorname{emp}_{N-k}}(x_i)\right)\right)\right]$$

$$= \operatorname{err}_N \frac{Z_{N-k,\beta}}{Z_{N,\beta}} \exp\left(-\beta \left(\sum_{1 \le i < j \le k} g(x_i - x_j) + N \sum_{i=1}^k V(x_i)\right)\right)\right)$$

$$\times \left(M_{N-k} \exp\left(-(N-k)\beta \left(\sum_{i=1}^k h^{\mu_\theta}(x_i)\right)\right) + A_{N-k}\right),$$
(6.3)

where $\operatorname{err}_N := \frac{N!}{(N-k)!} N^{-k}$. Step 2: Estimating the ratio of partition functions.

Integrating equation (6.3), and using that the k-th marginal is a probability measure, we get

$$\frac{Z_{N,\beta}}{Z_{N-k,\beta}} = \operatorname{err}_{N} \int_{\mathbf{R}^{d \times k}} \exp\left(-\beta \left(\sum_{1 \leq i < j \leq k} g(x_{i} - x_{j}) + N \sum_{i=1}^{k} V(x_{i})\right)\right) \times \left(M_{N-k} \exp\left(-(N-k)\beta \left(\sum_{i=1}^{k} h^{\mu_{\theta}}(x_{i})\right)\right) + A_{N-k}\right) dx_{1} \dots dx_{k} \qquad (6.4)$$

$$= M_{N}' \int_{\mathbf{R}^{d \times k}} \exp\left(-N\beta \left(\sum_{i=1}^{k} V(x_{i}) + h^{\mu_{\theta}}(x_{i})\right)\right) dx_{1} \dots dx_{k} + A_{N}' = M_{N}' L_{\theta}^{k} + A_{N}',$$

where

$$\begin{aligned} |\log M'_N| &\leq C'_1 N^{\frac{1}{2}} \beta \\ |A'_N| &\leq \exp\left(-C' N \beta\left(g(0)\right)\right), \end{aligned}$$

$$(6.5)$$

for some new constants C', C'_1, C'_2 depending only on V, g, d.

Recall that, by Remark 3.7, L_{θ} is of order exp $(-N\beta)$. More specifically,

$$\lim_{N \to \infty} \frac{\log L_{\theta}}{N\beta} = -2c_{\infty}.$$
(6.6)

Doing a Taylor expansion of $x \mapsto \frac{1}{x}$, we have that

$$\frac{Z_{N-k,\beta}}{Z_{N,\beta}} = M'_N L_{\theta}^{-k} + A'_N, \tag{6.7}$$

for a new additive error term A'_N satisfying equation (6.5) with a different constant C'.

Step 3: Conclusion.

Subtep 3.1: Pointwise convergence (item 1. of Lemma 6.1). Plugging in equation (6.7) into equation (6.3), and using equation (3.18) we obtain

$$R_k^N(Y_k) = \left(M'_N L_{\theta}^k + A'_N\right) \exp\left(-\beta \left(\sum_{1 \le i < j \le k} g(x_i - x_j) + N \sum_{i=1}^k V(x_i)\right)\right)$$

$$\times \left(M_{N-k} \exp\left(-(N-k)\beta \left(\sum_{i=1}^k h^{\mu_{\theta}}(x_i)\right)\right) + A_{N-k}\right)$$

$$= M''_N L_{\theta}^{-k} \exp\left(-N\beta \left(\sum_{i=1}^k V(x_i) + h^{\mu_{\theta}}(x_i)\right)\right) + A''_N$$

$$= M''_N \prod_{i=1}^k \mu_{\theta}(x_i) + A''_N,$$
(6.8)

where

$$|\log M_N''| \le C_1'' N^{\frac{1}{2}} \beta \left(C_2'' + \sum_{i=1}^k V(x_i) \right)^{\frac{1}{2}} |A_N''| \le \exp\left(-N\beta C'' \left(g(0) + \sum_{i=1}^k V(x_i) \right) \right),$$
(6.9)

for some new constants C'', C''_1, C''_2 . Using Lemma 3.6, we have that $R_k^N(Y_k) \to (\mu_V(x^*))^k$ for a.e. $Y_k \in \mathbf{R}^{d \times k}$. **Substep 3.1:** Summability condition (item 2. of Lemma 6.1). Let $K \subset \mathbf{R}^d$ be a compact set. Then by Lemma 3.6, there holds

$$\sup_{N \in \mathbf{N}} \sum_{k=1}^{\infty} \frac{1}{k!} \int_{K} R_{k}^{N}(Y_{k}) \, \mathrm{d}y_{1} \dots \, \mathrm{d}y_{k} \le C \sum_{k=1}^{\infty} \frac{1}{k!} \left(|K| \mu_{V}(x^{*}) \right)^{k} < \infty.$$
(6.10)

Using Lemma 6.1, we may conclude the proof.

7 **Appendix:** Point processes

In this appendix, we will review the theory on point processes necessary to prove Theorem 1. This appendix is based on [24].

Definition 7.1 (Point process). A point process \mathfrak{P} is a probability measure on the space of locally finite point configurations on \mathbb{R}^d . Alternatively, we may think of a point process \mathfrak{P} as a probability measure on the space of Radon measures consisting of a sum of Dirac deltas.

Definition 7.2 (Laplace functional). We associate to a point process \mathfrak{P} its Laplace functional ψ , defined by

$$\psi(f) := \mathbb{E}_{\mathfrak{P}}\left[\exp\left(-\int_{\mathbf{R}^d} f \,\mathrm{d}X\right)\right],\tag{7.1}$$

for a Borel-measurable $f: \mathbf{R}^d \to [0, \infty)$.

Definition 7.3 (Correlation function). Given a point process, we define the k-point correlation functions $\{R_k\}_{k=1}^{\infty}$ by the condition that

$$\psi(f) := 1 + \sum_{i=1}^{\infty} \frac{1}{k!} \int_{\mathbf{R}^d} \prod_{i=1}^k \left(\exp(-f(x_i)) - 1 \right) R_k(x_1, \dots x_k) \, \mathrm{d}x_1 \dots \, \mathrm{d}x_k.$$
(7.2)

Remark 7.4 (Correlation function of a joint distribution). If \mathfrak{P} is supported on point configurations of N points, and if these N points have a symmetric joint distribution \mathbb{P} , then we can verify that

$$\psi(f) := 1 + \sum_{i=1}^{\infty} {\binom{N}{k}} \int_{\mathbf{R}^d} \prod_{i=1}^k \left(\exp(-f(x_i)) - 1 \right) \mathbb{P}(X_N) \, \mathrm{d}X_N, \tag{7.3}$$

which implies that the correlation functions are given by

$$R_k(x_1, \dots x_k) = \frac{N!}{(N-k)!} \int_{\mathbf{R}^{d \times (N-k)}} \mathbb{P}(x_1, \dots x_k, x_{k+1}, \dots x_N) \, \mathrm{d}x_{k+1} \dots \, \mathrm{d}x_N \tag{7.4}$$

for any $k \leq N$.

Remark 7.5 (Correlation function of a Poisson point process). If \mathfrak{P} is a Poisson process of intensity λ , then its Laplace functional is given by

$$\psi(f) = \exp\left(\int_{\mathbf{R}^d} \left(\exp(-f(x)) - 1\right) \lambda(x) \,\mathrm{d}x\right). \tag{7.5}$$

Furthermore, the correlation functions of \mathfrak{P} are given by

$$R_k(x_1, \dots x_k) = \prod_{i=1}^k \lambda(x_i).$$
(7.6)

Definition 7.6 (Weak convergence). Let \mathfrak{P}_N be a sequence of point processes, and denote their Laplace functional by ψ_N . Then \mathfrak{P}_N converges weakly to a point process \mathfrak{P} with Laplace functional ψ if for any continuous and compactly supported $f: \mathbf{R}^d \to \mathbf{R}^+$ there holds

$$\lim_{N \to \infty} \psi_N(f) = \psi(f). \tag{7.7}$$

Lemma 7.7 (Weak convergence and correlation functions). Let \mathfrak{P}_N be a sequence of point processes, and denote their k- point correlation functions by R_k^N . Then \mathfrak{P}_N converges weakly to a point process \mathfrak{P} , with k- point correlation functions denoted by R_k^N if:

- 1. $R_k^N \to R_k$ pointwise almost everywhere as $N \to \infty$.
- 2. For any compact set $\Omega \subset \mathbf{R}^d$ there holds

$$\sup_{N \in \mathbf{N}} \sum_{k=1}^{\infty} \frac{1}{k!} \int_{K} R_k^N \,\mathrm{d}y_1 \dots \,\mathrm{d}y_k < \infty.$$

$$(7.8)$$

8 Acknowledgements

DPG acknowledges the support of the ERC Starting Grant "Bridging Scales in Random Materials" ERC StG RandSCALES 948819. ET was supported by NSF grant DMS-2303318.

References

- Gernot Akemann and Sung-Soo Byun. The high temperature crossover for general 2d Coulomb gases. *Journal of Statistical Physics*, 175(6):1043–1065, 2019.
- [2] Scott Armstrong and Sylvia Serfaty. Local laws and rigidity for Coulomb gases at any temperature. *The Annals of Probability*, 49(1):46–121, 2021.
- [3] Scott Armstrong and Sylvia Serfaty. Thermal approximation of the equilibrium measure and obstacle problem. In Annales de la Faculté des sciences de Toulouse: Mathématiques, volume 31, pages 1085–1110, 2022.
- [4] Florent Benaych-Georges and Sandrine Péché. Poisson statistics for matrix ensembles at large temperature. Journal of Statistical Physics, 161:633–656, 2015.
- [5] S Borodachov, D Hardin, A Reznikov, and E Saff. Optimal discrete measures for Riesz potentials. *Transactions of the American Mathematical Society*, 370(10):6973–6993, 2018.
- [6] Sergiy Borodachov, Douglas Hardin, and Edward Saff. Discrete Energy on Rectifiable Sets. Springer New York, NY, 2019.
- [7] Jeanne Boursier. Optimal local laws and CLT for the circular Riesz gas. arXiv preprint arXiv:2112.05881, 2021.
- [8] Jeanne Boursier. Decay of correlations and thermodynamic limit for the circular Riesz gas. arXiv preprint arXiv:2209.00396, 2022.
- [9] Andrea Braides. Gamma-Convergence for Beginners. Oxford University Press, 07 2002.
- [10] Djalil Chafaï, Nathael Gozlan, and Pierre-André Zitt. First-order global asymptotics for confined particles with singular pair repulsion. *The Annals of Applied Probability*, 24(6):2371–2413, 2014.
- [11] Djalil Chafaï, Edward B Saff, and Robert S Womersley. On the solution of a Riesz equilibrium problem and integral identities for special functions. *Journal of Mathematical Analysis and Applications*, 515(1):126367, 2022.
- [12] Djalil Chafaï, Edward B Saff, and Robert S Womersley. Threshold condensation to singular support for a Riesz equilibrium problem. Analysis and Mathematical Physics, 13(1):19, 2023.
- [13] David Dereudre, Adrien Hardy, Thomas Leblé, and Mylène Maïda. DLR equations and rigidity for the sine-beta process. *Communications on Pure and Applied Mathematics*, 74(1):172–222, 2021.
- [14] David Dereudre and Thibaut Vasseur. Number-rigidity and β -circular Riesz gas. arXiv preprint arXiv:2104.09408, 2021.

- [15] PD Dragnev, B Fuglede, DP Hardin, EB Saff, and N Zorii. Minimum Riesz energy problems for a condenser with touching plates. *Potential Analysis*, 44:543–577, 2016.
- [16] Peter D Dragnev, Bent Fuglede, Doug P Hardin, Edward B Saff, and Natalia Zorii. Constrained minimum Riesz energy problems for a condenser with intersecting plates. *Journal d'Analyse Mathématique*, 140(1):117–159, 2020.
- [17] David García-Zelada. A large deviation principle for empirical measures on polish spaces: Application to singular Gibbs measures on manifolds. In Annales de l'Institut Henri Poincaré, Probabilités et Statistiques, volume 55, pages 1377–1401. Institut Henri Poincaré, 2019.
- [18] David García-Zelada and David Padilla-Garza. Generalized transport inequalities and concentration bounds for Riesz-type gases. arXiv preprint arXiv:2209.00587, 2022.
- [19] Hans-Otto Georgii. Gibbs measures and phase transitions. In *Gibbs Measures and Phase Transitions*. de Gruyter, 2011.
- [20] Subhroshekhar Ghosh and Yuval Peres. Rigidity and tolerance in point processes: Gaussian zeros and Ginibre eigenvalues. Duke Mathematical Journal, 166(10):1789 1858, 2017.
- [21] Douglas P Hardin, Edward B Saff, and OV Vlasiuk. Generating point configurations via hypersingular Riesz energy with an external field. SIAM Journal on Mathematical Analysis, 49(1):646–673, 2017.
- [22] DP Hardin, A Reznikov, EB Saff, and A Volberg. Local properties of Riesz minimal energy configurations and equilibrium measures. *International Mathematics Research Notices*, 2019(16):5066–5086, 2019.
- [23] Adrien Hardy and Gaultier Lambert. CLT for circular beta-ensembles at high temperature. Journal of Functional Analysis, 280(7):108869, 2021.
- [24] Gaultier Lambert. Poisson statistics for Gibbs measures at high temperature. 57 1 Annales de l'Institut Henri Poicaré Probabilités et Statistiques Vol. 57, No. 1 (February, 2021) 1-602, 57(1):12, 2021.
- [25] Thomas Leblé. Local microscopic behavior for 2d Coulomb gases. Probability Theory and Related Fields, 169:931–976, 2017.
- [26] Thomas Leblé and Sylvia Serfaty. Large deviation principle for empirical fields of log and Riesz gases. *Inventiones mathematicae*, 210:645–757, 2017.
- [27] Thomas Leblé and Sylvia Serfaty. Fluctuations of two dimensional Coulomb gases. Geometric and Functional Analysis, 28:443–508, 2018.
- [28] Fumihiko Nakano and Khanh Duy Trinh. Poisson statistics for beta ensembles on the real line at high temperature. *Journal of Statistical Physics*, 179(2):632–649, 2020.
- [29] Douglas P. Hardin, Edward B. Saff, Brian Z. Simanek, and Yujian Su. Next order energy asymptotics for Riesz potentials on flat tori. *International Mathematics Research Notices*, 2017(12):3529–3556, 2017.

- [30] David Padilla-Garza. Large deviation principle for local empirical measure of Coulomb gases at intermediate temperature regime. *arXiv preprint arXiv:2011.00480*, 2020.
- [31] David Padilla-Garza. Large deviations principle for the tagged empirical field of a general interacting gas. arXiv preprint arXiv:2208.01516, 2022.
- [32] David Padilla-Garza. Concentration inequality around the thermal equilibrium measure of Coulomb gases. *Journal of Functional Analysis*, 284(1):109733, 2023.
- [33] Luke Peilen. Local laws and a mesoscopic CLT for β -ensembles. Communications on Pure and Applied Mathematics, 77(4):2452–2567, 2024.
- [34] Mircea Petrache and Sylvia Serfaty. Next order asymptotics and renormalized energy for Riesz interactions. *Journal of the Institute of Mathematics of Jussieu*, 16(3):501–569, 2017.
- [35] Mircea Petrache and Sylvia Serfaty. Crystallization for Coulomb and Riesz interactions as a consequence of the Cohn-Kumar conjecture. *Proceedings of the American Mathematical Society*, 148(7):3047–3057, 2020.
- [36] Alexander Reznikov, E Saff, and Alexander Volberg. Covering and separation of Chebyshev points for non-integrable Riesz potentials. *Journal of Complexity*, 46:19–44, 2018.
- [37] Grant Rotskoff and Eric Vanden-Eijnden. Parameters as interacting particles: long time convergence and asymptotic error scaling of neural networks. Advances in neural information processing systems, 31, 2018.
- [38] Grant Rotskoff and Eric Vanden-Eijnden. Trainability and accuracy of artificial neural networks: An interacting particle system approach. *Communications on Pure and Applied Mathematics*, 75(9):1889–1935, 2022.
- [39] Grant M Rotskoff and Eric Vanden-Eijnden. Neural networks as interacting particle systems: Asymptotic convexity of the loss landscape and universal scaling of the approximation error. *stat*, 1050:22, 2018.
- [40] Nicolas Rougerie and Sylvia Serfaty. Higher-dimensional Coulomb gases and renormalized energy functionals. Communications on Pure and Applied Mathematics, 69(3):519– 605, 2016.
- [41] Nicolas Rougerie, Sylvia Serfaty, and Jakob Yngvason. Quantum Hall phases and plasma analogy in rotating trapped Bose gases. *Journal of Statistical Physics*, 154:2–50, 2014.
- [42] David Ruelle. A variational formulation of equilibrium statistical mechanics and the Gibbs phase rule. *Communications in Mathematical Physics*, 5:324–329, 1967.
- [43] David Ruelle. Statistical mechanics of a one-dimensional lattice gas. Communications in Mathematical Physics, 9:267–278, 1968.
- [44] Edward Saff and Vilmos Totik. Logarithmic Potentials with External Fields. Springer Berlin, Heidelberg, 1997.

- [45] Etienne Sandier and Sylvia Serfaty. 1d log gases and the renormalized energy: crystallization at vanishing temperature. *Probability Theory and Related Fields*, 162(3):795–846, 2015.
- [46] Etienne Sandier and Sylvia Serfaty. 2d Coulomb gases and the renormalized energy. The Annals of Probability, pages 2026–2083, 2015.
- [47] Sylvia Serfaty. Coulomb gases and Ginzburg-Landau Vortices. European Mathematical Society, 2015.
- [48] Sylvia Serfaty. Gaussian fluctuations and free energy expansion for Coulomb gases at any temperature. In Annales de l'Institut Henri Poincare (B) Probabilites et statistiques, volume 59, pages 1074–1142. Institut Henri Poincaré, 2023.
- [49] Eric Thoma. Non-rigidity properties of the Coulomb gas. arXiv preprint arXiv:2303.11486, 2023.
- [50] Eric Thoma. Overcrowding and separation estimates for the Coulomb gas. Communications on Pure and Applied Mathematics, 2024.
- [51] Yuxiao Wen, Eric Vanden-Eijnden, and Benjamin Peherstorfer. Coupling parameter and particle dynamics for adaptive sampling in neural Galerkin schemes. *Physica D: Nonlinear Phenomena*, page 134129, 2024.

DAVID PADILLA-GARZA Institute of Science and Technology Austria. Email: David.Padilla-Garza@ist.ac.at.

LUKE PEILEN Department of Mathematics, Temple University. Email: luke.peilen@temple.edu.

ERIC THOMA Department of Mathematics, Stanford University. Email: thoma@stanford.edu.