This article is accepted for publishing in the Journal of Fourier and Analysis and Applications

Heyde theorem for locally compact Abelian groups containing no subgroups topologically isomorphic to the 2-dimensional torus

Gennadiy Feldman (ORCID ID https://orcid.org/0000-0001-5163-4079)

Abstract

We prove the following group analogue of the well-known Heyde theorem on a characterization of the Gaussian distribution on the real line. Let X be a second countable locally compact Abelian group containing no subgroups topologically isomorphic to the 2-dimensional torus. Let G be the subgroup of X generated by all elements of X of order 2 and let α be a topological automorphism of the group X such that $\text{Ker}(I + \alpha) = \{0\}$. Let ξ_1 and ξ_2 be independent random variables with values in X and distributions μ_1 and μ_2 with nonvanishing characteristic functions. If the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric, then μ_j are convolutions of Gaussian distributions on X and distributions supported in G. We also prove that this theorem is false if X is the 2-dimensional torus.

Keywords locally compact Abelian group \cdot 2-dimensional torus \cdot topological automorphism \cdot Gaussian distribution \cdot conditional distribution

Mathematics Subject Classification Primary: 43A25 · 43A35 · 60B15 · 62E10

1 Introduction

The following characterization theorem was proved in [15, Theorem 3.1].

Theorem A Let X be a second countable locally compact Abelian group with the connected component of zero of dimension 1. Let G be the subgroup of X generated by all elements of X of order 2 and let α be a topological automorphism of the group X satisfying the condition

$$\operatorname{Ker}(I + \alpha) = \{0\}.\tag{1}$$

Let ξ_1 and ξ_2 be independent random variables with values in X and distributions μ_1 and μ_2 with nonvanishing characteristic functions. If the conditional distribution of the linear form $L_2 = \xi_1 + \alpha\xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric, then μ_j are convolutions of Gaussian distributions on X and distributions supported in G.

Theorem A can be considered as a group analogue for two independent random variables of the well-known theorem of C.C. Heyde ([18], see also [19, \S 13.4.1]), where the Gaussian distribution on the real line is characterized by the symmetry of the conditional distribution of one linear form of independent random variables given another.

Denote by \mathbb{R} the additive group of real numbers and by \mathbb{T} the circle group (the one-dimensional torus), i.e., the multiplicative group of all complex numbers with absolute value 1. The aim of the article is, firstly, to prove that Theorem A is true for a much wider class of locally compact Abelian groups, namely for second countable locally compact Abelian groups containing no subgroups topologically isomorphic to the 2-dimensional torus \mathbb{T}^2 (Theorem 2.1). Secondly, to prove that Theorem A is false

when X is the 2-dimensional torus \mathbb{T}^2 (Theorem 3.1). We emphasize that the proof of Theorem 2.1 given in the article is fundamentally different from the proof of Theorem A, which is based on the proof of the theorem for the group X of the form $X = \mathbb{R} \times D$, where D is a second countable totally disconnected locally compact Abelian group, and uses complex analysis.

Many studies have been devoted to analogues of Heyde's theorem for different classes of locally compact Abelian groups (see e.g. [3–5, 7–13, 20–22], and also [14, Chapter IV], where one can find additional references). In the article we continue this research.

We use in the article standard facts related to abstract harmonic analysis, see e.g. [17]. Let X be a second countable locally compact Abelian group, let $\operatorname{Aut}(X)$ be the group of topological automorphisms of the group X, and let I be the identity automorphism of a group. Denote by $Y = X^*$ the character group of the group X, and by (x, y) the value of a character $y \in Y$ at an element $x \in X$. For a closed subgroup K of the group X, denote by $A(Y, K) = \{y \in Y : (x, y) = 1 \text{ for all } x \in K\}$ its annihilator. The character group of the factor-group X/K is topologically isomorphic to the annihilator A(Y, K). Let $\alpha : X \to X$ be a continuous endomorphism of the group X. The adjoint endomorphism $\tilde{\alpha} : Y \to Y$ is defined as follows: $(\alpha x, y) = (x, \tilde{\alpha} y)$ for all $x \in X$, $y \in Y$. Let n be a natural number. Put $X^{(n)} = \{nx : x \in X\}$. A topological isomorphism of locally compact Abelian groups X_1 and X_2 is denoted as $X_1 \cong X_2$. Denote by \mathbb{Z} the additive group of integers.

Let f(y) be a function on the group Y and let $h \in Y$. Denote by Δ_h the finite difference operator

$$\Delta_h f(y) = f(y+h) - f(y), \quad y \in Y.$$

Denote by $M^1(X)$ the convolution semigroup of probability distributions on the group X. Let $\mu \in M^1(X)$. Denote by

$$\hat{\mu}(y) = \int_X (x, y) d\mu(x), \quad y \in Y,$$

the characteristic function (Fourier transform) of the distribution μ . The characteristic function of a signed measure is defined in the same way. Denote by $\sigma(\mu)$ the support of μ . Define the distribution $\bar{\mu} \in M^1(X)$ by the formula $\bar{\mu}(B) = \mu(-B)$ for any Borel subset B in X. Then $\hat{\mu}(y) = \bar{\mu}(y)$. If F is a Borel subgroup of X, denote by $M^1(F)$ the subsemigroup of $M^1(X)$ consisting of all distributions concentrated on F.

A distribution $\gamma \in M^1(X)$ is called Gaussian ([24, Chapter IV, §6]) if its characteristic function can be represented in the form

$$\hat{\gamma}(y) = (x, y) \exp\{-\varphi(y)\}, \quad y \in Y,$$
(2)

where $x \in X$ and $\varphi(y)$ is a continuous nonnegative function on the group Y satisfying the equation

$$\varphi(u+v) + \varphi(u-v) = 2[\varphi(u) + \varphi(v)], \quad u, v \in Y.$$
(3)

A Gaussian distribution is called symmetric if x = 0 in (2). Denote by $\Gamma(X)$ the set of Gaussian distributions on the group X. Note that in particular, the degenerate distributions are Gaussian. Let $x \in X$. Denote by E_x the degenerate distribution concentrated at the point $x \in X$.

Denote by m_X a Haar measure on the group X. It is well known that m_X is unique up to a positive multiplicative constant. If X is a compact group, then $m_X(X) < \infty$. We suppose that in this case $m_X \in M^1(X)$. Note that, if X is an arbitrary locally compact Abelian group and K is a compact subgroup of X, then the characteristic function $\widehat{m}_K(y)$ is of the form

$$\widehat{m}_{K}(y) = \begin{cases} 1, & \text{if } y \in A(Y, K), \\ 0, & \text{if } y \notin A(Y, K). \end{cases}$$

$$\tag{4}$$

2 Proof of the main theorem

The main result of the article is the following theorem.

Theorem 2.1 Let X be a second countable locally compact Abelian group containing no subgroups topologically isomorphic to the 2-dimensional torus \mathbb{T}^2 . Let G be the subgroup of X generated by all elements of X of order 2 and let α be a topological automorphism of the group X satisfying condition (1). Let ξ_1 and ξ_2 be independent random variables with values in the group X and distributions μ_1 and μ_2 with nonvanishing characteristic functions. If the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric, then $\mu_i \in \Gamma(X) * M^1(G)$, j = 1, 2.

To prove Theorem 2.1 we need some lemmas.

Lemma 2.2 ([21, Lemma 3.8], see also [14, Corollary 9.7]) Let X be a second countable locally compact Abelian group and let α be a topological automorphism of X. Let ξ_1 and ξ_2 be independent random variables with values in the group X. If the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric, then the linear forms $P_1 = (I + \alpha)\xi_1 + 2\alpha\xi_2$ and $P_2 = 2\xi_1 + (I + \alpha)\xi_2$ are independent.

Lemma 2.3 ([6, Lemma 10.1]) Let X be a second countable locally compact Abelian group with character group Y. Let α_j , β_j , j = 1, 2, be continuous endomorphisms of the group X. Let ξ_1 and ξ_2 be independent random variables with values in the group X and distributions μ_1 and μ_2 . The linear forms $L_1 = \alpha_1\xi_1 + \alpha_2\xi_2$ and $L_2 = \beta_1\xi_1 + \beta_2\xi_2$ are independent if and only if the characteristic functions $\hat{\mu}_j(y)$ satisfy the equation

$$\hat{\mu}_1(\widetilde{\alpha}_1 u + \widetilde{\beta}_1 v)\hat{\mu}_2(\widetilde{\alpha}_2 u + \widetilde{\beta}_2 v) = \hat{\mu}_1(\widetilde{\alpha}_1 u)\hat{\mu}_2(\widetilde{\alpha}_2 u)\hat{\mu}_1(\widetilde{\beta}_1 v)\hat{\mu}_2(\widetilde{\beta}_2 v), \quad u, v \in Y.$$
(5)

Lemma 2.4 ([7, Lemma 6], see also [14, Lemma 5.1]) Let Y be a locally compact Abelian group and let A(y) be a continuous function on the group Y satisfying the equation

$$\Delta_{2k}\Delta_h^2 A(y) = 0, \quad y, k, h \in Y,$$

and the conditions A(-y) = A(y), A(0) = 0. Let

$$Y = \bigcup_{\iota} (y_{\iota} + \overline{Y^{(2)}}), \quad y_0 = 0, \tag{6}$$

be a $\overline{Y^{(2)}}$ -coset decomposition of the group Y. Then the function A(y) can be represented in the form

$$A(y) = \varphi(y) + r_{\iota}, \quad y \in y_{\iota} + \overline{Y^{(2)}}, \tag{7}$$

where $\varphi(y)$ is a continuous function on the group Y satisfying equation (3).

Let us recall some definitions. Let X be an Abelian group. We do not assume that X is a topological group. If each element of X has finite order, then we say that X is a torsion group. We say that X is a torsion-free group if each element of X, except zero, has infinite order. The subgroup consisting of all elements of finite order of the group X is called a torsion part of X.

We formulate as a lemma the following corollary of the well-known Baer–Fomin theorem ([16, Theorem 100.1]).

Lemma 2.5 Let Y be an Abelian group and let H be a torsion part of Y. If H is a bounded subgroup, then H is a direct factor of Y.

Using Lemma 2.5 we shall prove the following statement.

Lemma 2.6 Let X be a locally compact Abelian group containing no subgroups topologically isomorphic to the circle group \mathbb{T} . Let K be a closed bounded subgroup of X. Then the factor-group X/K also contains no subgroups topologically isomorphic to the circle group \mathbb{T} .

Proof. The lemma is true if K = X, so we suppose that $K \neq X$. Assume that the factor-group X/K contains a subgroup F such that $F \cong \mathbb{T}$. Let us prove that in this case we arrive at a contradiction. Let $p: X \to X/K$ be the natural homomorphism. Put $S = p^{-1}(F)$. The group S is closed and hence locally compact. Consider the restriction of p to the subgroup S. We have $p: S \to S/K$ and $S/K \cong \mathbb{T}$. Thus, we can prove the lemma assuming that $X/K \cong \mathbb{T}$. By the structure theorem for locally compact Abelian group X is topologically isomorphic to a group of the form $\mathbb{R}^m \times G$, where a locally compact Abelian group G contains a compact open subgroup. Since K is a bounded subgroup, $K \subset G$. It follows from $X/K \cong \mathbb{T}$ that m = 0, i.e., the group X itself contains a compact open subgroup. Denote by Y the character group of the group X.

First suppose that X is a compact group. Then Y is a discrete group. Denote by H the torsion part of the group Y. We have $A(Y, K) \cong (X/K)^*$. Since $X/K \cong \mathbb{T}$, this implies that

$$A(Y,K) \cong \mathbb{Z}.$$
(8)

Inasmuch as K is a bounded group, there is a natural n such that nx = 0 for all $x \in K$. It follows from this that $ny \in A(Y, K)$ for each $y \in Y$, i.e.,

$$Y^{(n)} \subset A(Y, K). \tag{9}$$

In particular,

$$H^{(n)} \subset A(Y, K). \tag{10}$$

In view of (8), (10) implies that

$$H^{(n)} = \{0\}.$$
 (11)

Hence H is a bounded group. By Lemma 2.5, H is a direct factor of Y. We have $Y = H \times L$, where L is a torsion-free group. Taking into account (11), this implies that $Y^{(n)} = L^{(n)}$. For this reason, it follows from (8) and (9) that $L \cong \mathbb{Z}$. This implies that the group X contains a subgroup topologically isomorphic to the circle group \mathbb{T} , contrary to the assumption. Thus, the lemma is proved if the group X is compact.

Consider the general case. Let B be a compact open subgroup of the group X. As far as p is a continuous open epimorphism, p(B) is an open subgroup of the factor-group X/K. Since $X/K \cong \mathbb{T}$, we have p(B) = X/K. Moreover, $p(B) \cong B/(K \cap B)$. Thus, $B/(K \cap B) \cong \mathbb{T}$. As has been shown above, this implies that the group B, and hence the group X, contains a subgroup topologically isomorphic to the circle group \mathbb{T} , contrary to the assumption.

Let \mathbb{R}^{\aleph_0} be the space of all sequences of real numbers considering in the product topology. We will need the definition of the Gaussian distribution in \mathbb{R}^{\aleph_0} . The space \mathbb{R}^{\aleph_0} is one of the simplest examples of a locally convex space. We note that Gaussian distributions in arbitrary locally convex spaces are studied in details in the fundamental monograph [1].

Denote by \mathbb{R}^{\aleph_0*} the space of all finitary sequences of real numbers with the topology of strictly inductive limit of spaces \mathbb{R}^n . Let $t = (t_1, t_2, \ldots, t_n, \ldots) \in \mathbb{R}^{\aleph_0}$ and $s = (s_1, s_2, \ldots, s_n, 0, \ldots) \in \mathbb{R}^{\aleph_0*}$. Set

$$\langle t,s\rangle = \sum_{j=1}^{\infty} t_j s_j.$$

Let μ be a distribution on \mathbb{R}^{\aleph_0} . We define the characteristic function of μ by the formula

$$\hat{\mu}(s) = \int_{\mathbb{R}^{\aleph_0}} \exp\{i\langle t, s \rangle\} d\mu(t), \quad s \in \mathbb{R}^{\aleph_0 *}.$$

Let $A = (\alpha_{ij})_{i,j=1}^{\infty}$ be a symmetric positive semidefinite matrix, i.e., the quadratic form

$$\langle As,s\rangle = \sum_{i,j=1}^\infty \alpha_{ij}s_is_j$$

is nonnegative for all $s = (s_1, s_2, \ldots, s_n, 0, \ldots) \in \mathbb{R}^{\aleph_0 *}$.

A distribution μ on the group \mathbb{R}^{\aleph_0} is called Gaussian if its characteristic function is represented in the form

$$\hat{\mu}(s) = \exp\{i\langle t, s \rangle - \langle As, s \rangle\}, \quad s \in \mathbb{R}^{\aleph_0 *},$$

where $t \in \mathbb{R}^{\aleph_0}$ and $A = (\alpha_{ij})_{i,j=1}^{\infty}$ is a symmetric positive semidefinite matrix.

Lemma 2.7 ([2], see also [6, §3]) Let X be a second countable locally compact Abelian group containing no subgroups topologically isomorphic to the circle group \mathbb{T} . Then there exists a continuous monomorphism $p: E \to X$, where either $E = \mathbb{R}^n$ for some n or $E = \mathbb{R}^{\aleph_0}$, such that if γ is a symmetric Gaussian distribution on X, then $\gamma = p(M)$, where M is a symmetric Gaussian distribution on E.

Lemma 2.8 Let X be a second countable locally compact Abelian group containing no subgroups topologically isomorphic to the circle group \mathbb{T} . Let G be the subgroup of X generated by all elements of X of order 2. Let $\mu \in \Gamma(X) * M^1(G)$ and suppose that the characteristic function of the distribution μ does not vanish. If $\mu = \mu_1 * \mu_2$, where $\mu_j \in M^1(X)$, then $\mu_j \in \Gamma(X) * M^1(G)$, j = 1, 2.

Proof. Since the group X contains no subgroups topologically isomorphic to the circle group \mathbb{T} , let p be a continuous monomorphism $p: E \to X$ which exists by Lemma 2.7. As far as p is a monomorphism, we have $p(E) \cap G = \{0\}$. Hence p can be extended to a continuous monomorphism $\bar{p}: E \times G \to X$ by the formula $\bar{p}(t,g) = p(t) + g$, $t \in E$, $g \in G$, and \bar{p} generates an isomorphism of the semigroups $M^1(E \times G)$ and $M^1(p(E) \times G)$. Let $\mu = \gamma * \omega$, where $\gamma \in \Gamma(X)$, $\omega \in M^1(G)$. We can assume without loss of generality that γ is a symmetric Gaussian distribution. By Lemma 2.7, $\gamma = p(M)$, where $M \in \Gamma(E)$. Hence $\mu = \bar{p}(N)$, where $N = M * \omega$. Obviously, the distribution μ is concentrated on the Borel subgroup $p(E) \times G$ of the group X. Substituting, if it is necessary, the distributions μ_j by their shifts, we can assume that μ_j are also concentrated on the Borel subgroup $p(E) \times G$. Since the semigroups $M^1(E \times G)$ and $M^1(p(E) \times G)$ are isomorphic, we have $\mu_j = \bar{p}(N_j)$, where $N_j \in M^1(E \times G)$ and N_j is a factor of N. By [4, Lemma 5 and Remark 4], we have $N_j = M_j * \omega_j$, where $M_j \in \Gamma(E)$, $\omega_j \in M^1(G)$. Hence

$$\mu_j = \bar{p}(N_j) = \bar{p}(M_j * \omega_j) = \bar{p}(M_j) * \bar{p}(\omega_j) = p(M_j) * \omega_j = \gamma_j * \omega_j,$$

where $\gamma_j = p(M_j)$. Since $\gamma_j \in \Gamma(X)$, the lemma is proved.

Proof of Theorem 2.1 Assume that a locally compact Abelian group X contains no subgroups topologically isomorphic to the 2-dimensional torus \mathbb{T}^2 but contains a subgroup topologically isomorphic to the circle group \mathbb{T} . Obviously, there can be only one such subgroup. Hence it is invariant with respect to each topological automorphism of the group X. Since $\operatorname{Aut}(\mathbb{T}) = \{\pm I\}$, this implies that there is no topological automorphism α of the group X satisfying condition (1). Hence we can prove the theorem

assuming from the beginning that the group X contains no subgroups topologically isomorphic to the circle group \mathbb{T} .

Denote by Y the character group of the group X. By Lemma 2.2, the symmetry of the conditional distribution of the linear form L_2 given L_1 implies that the linear forms $P_1 = (I + \alpha)\xi_1 + 2\alpha\xi_2$ and $P_2 = 2\xi_1 + (I + \alpha)\xi_2$ are independent. By Lemma 2.3, it follows from this that the characteristic functions $\hat{\mu}_i(y)$ satisfy equation (5), which takes the form

$$\hat{\mu}_1((I+\widetilde{\alpha})u+2v)\hat{\mu}_2(2\widetilde{\alpha}u+(I+\widetilde{\alpha})v) = \hat{\mu}_1((I+\widetilde{\alpha})u)\hat{\mu}_2(2\widetilde{\alpha}u)\hat{\mu}_1(2v)\hat{\mu}_2((I+\widetilde{\alpha})v), \quad u,v \in Y.$$
(12)

Put $\nu_j = \mu_j * \bar{\mu}_j$. We have $\hat{\nu}_j(y) = |\hat{\mu}_j(y)|^2 > 0$ for all $y \in Y$, j = 1, 2. The characteristic functions $\hat{\nu}_j(y)$ also satisfy equation (12). Set $\psi_j(y) = -\ln \hat{\nu}_j(y)$, j = 1, 2. It follows from (12) that the functions $\psi_j(y)$ satisfy the equation

$$\psi_1((I+\widetilde{\alpha})u+2v) + \psi_2(2\widetilde{\alpha}u + (I+\widetilde{\alpha})v) = A(u) + B(v), \quad u, v \in Y,$$
(13)

where

$$A(y) = \psi_1((I + \widetilde{\alpha})y) + \psi_2(2\widetilde{\alpha}y), \quad B(y) = \psi_1(2y) + \psi_2((I + \widetilde{\alpha})y), \quad y \in Y.$$

$$\tag{14}$$

Equation (13) has already appeared earlier in the study of Heyde's theorem on various locally compact Abelian groups (see e.g. [10], [11]). For completeness, we give here its solution. We use the finite difference method. Take an arbitrary element h_1 of the group Y. Substitute $u + (I + \tilde{\alpha})h_1$ for u and $v - 2\tilde{\alpha}h_1$ for v in equation (13). Subtracting equation (13) from the obtaining equation we get

$$\Delta_{(I-\widetilde{\alpha})^2h_1}\psi_1((I+\widetilde{\alpha})u+2v) = \Delta_{(I+\widetilde{\alpha})h_1}A(u) + \Delta_{-2\widetilde{\alpha}h_1}B(v), \quad u,v \in Y.$$
(15)

Take an arbitrary element h_2 of the group Y. Substitute $u + 2h_2$ for u and $v - (I + \tilde{\alpha})h_2$ for v in equation (15). Subtracting equation (15) from the resulting equation we obtain

$$\Delta_{2h_2}\Delta_{(I+\tilde{\alpha})h_1}A(u) + \Delta_{-(I+\tilde{\alpha})h_2}\Delta_{-2\tilde{\alpha}h_1}B(v) = 0, \quad u, v \in Y.$$
(16)

Take an arbitrary element h of the group Y. Substituting u + h for u in equation (16) and subtracting equation (16) from the obtaining equation we get

$$\Delta_h \Delta_{2h_2} \Delta_{(I+\widetilde{\alpha})h_1} A(u) = 0, \quad u \in Y.$$
⁽¹⁷⁾

It follows from properties of adjoint homomorphisms that if a topological automorphism α satisfies condition (1), then the subgroup $(I + \tilde{\alpha})(Y)$ is dense in Y. Taking into account that h, h_1, h_2 are arbitrary elements of the group Y, it follows from (17) that the function A(y) satisfies the equation

$$\Delta_{2k}\Delta_h^2 A(y) = 0, \quad y, k, h \in Y.$$
(18)

By Lemma 2.4, (18) implies that the function A(y) is represented in the form (7), where the function $\varphi(y)$, as is easily seen, is nonnegative. Denote by μ the distribution of the random variable $P_1 = (I + \alpha)\xi_1 + 2\alpha\xi_2$. It is obvious that

$$\mu = (I + \alpha)(\mu_1) * (2\alpha)(\mu_2).$$

Put $\nu = \mu * \overline{\mu}$. Then

$$\nu = (I + \alpha)(\mu_1) * (2\alpha)(\mu_2) * (I + \alpha)(\bar{\mu}_1) * (2\alpha)(\bar{\mu}_2).$$
(19)

It follows from (14) and (19) that the characteristic function $\hat{\nu}(y)$ is of the form

$$\hat{\nu}(y) = e^{-A(y)}, \quad y \in Y.$$
(20)

Denote by γ the Gaussian distribution on the group X with the characteristic function

$$\hat{\gamma}(y) = \exp\{-\varphi(y)\}, \quad y \in Y.$$
(21)

Taking into account (6) and (7), define on the group Y the function g(y) by the formula

$$g(y) = \exp\{-r_{\iota}\}, \quad y \in y_{\iota} + \overline{Y^{(2)}}.$$
 (22)

Since $g(y) = \hat{\nu}(y)/\hat{\gamma}(y)$, the function g(y) is continuous. Moreover, (22) implies that the function g(y) is invariant with respect to the subgroup $\overline{Y^{(2)}}$. Check that g(y) is a positive definite function. Note that

$$A(X, \overline{Y^{(2)}}) = G. \tag{23}$$

Consider decomposition (6) and take a finite set of elements y_{i_j} , j = 1, 2, ..., n. Let H be a subgroup of Y generated all cosets $y_{i_j} + \overline{Y^{(2)}}$. Obviously, it suffices to verify that the restriction of the function g(y) to H is a positive definite function. The subgroup H consists of a finite number of cosets $y_i + \overline{Y^{(2)}}$. Put K = A(X, H). Then $H^* \cong X/K$ and obviously, $K \subset G$. Consider the restriction of the function g(y) to H. This restriction is invariant with respect to the subgroup $\overline{Y^{(2)}}$ and hence defines a function on the factor-group $H/\overline{Y^{(2)}}$. We note that $H/\overline{Y^{(2)}}$ is a finite group and all its nonzero elements are of the order 2. It follows from this that any real-valued function on the factor-group $H/\overline{Y^{(2)}}$ is the characteristic function of a signed measure. In particular, the restriction of the function g(y) to the subgroup H is the characteristic function of a signed measure ϖ . Since $(H/\overline{Y^{(2)}})^* \cong A(X/K, \overline{Y^{(2)}})$, we can consider the signed measure ϖ as a signed measure ϖ . Since $(H/\overline{Y^{(2)}})^* \cong A(X/K, \overline{Y^{(2)}})$. It follows from (7) and (20)–(22) that the restriction of the characteristic function $\hat{\nu}(y)$ to H is the characteristic function of the convolution of a Gaussian distribution λ on X/K and the signed measure ϖ on F. We verify that the signed measure ϖ is actually a distribution and this proves that g(y) is a positive definite function.

Since $K \subset G$, by Lemma 2.6, the factor-group X/K contains no subgroups topologically isomorphic to the circle group \mathbb{T} . Then by Lemma 2.7, applying to the group X/K, there exists a continuous monomorphism $p: E \to X/K$, where either $E = \mathbb{R}^n$ for some n or $E = \mathbb{R}^{\aleph_0}$, such that $\lambda = p(M)$, where M is a symmetric Gaussian distribution on E. Hence the Gaussian distribution λ is concentrated on the Borel subgroup p(E) of the group X/K. Note that all nonzero elements of the group F are of the order 2 and hence

$$p(E) \cap F = \{0\}.$$
 (24)

Since the convolution $\lambda * \varpi$ is a distribution, in view of (24), ϖ is also a distribution. Thus, we proved that g(y) is a continuous positive definite function such that g(0) = 0. By the Bochner theorem, there exists a distribution $\omega \in M^1(X)$ such that $\hat{\omega}(y) = g(y), y \in Y$. Since g(y) = 1 for $y \in \overline{Y^{(2)}}$, in view of (23), we have an inclusion

$$\sigma(\omega) \subset A\left(X, \{y \in Y : g(y) = 1\}\right) \subset A(X, Y^{(2)}) = G.$$

It follows from $\hat{\nu}(y) = \hat{\gamma}(y)g(y)$ for all $y \in Y$, that $\nu = \gamma * \omega \in \Gamma(X) * M^1(G)$. By Lemma 2.8, (19) implies that $(I+\alpha)(\mu_1) \in \Gamma(X) * M^1(G)$. Taking into account that $I+\alpha$ is a continuous monomorphism, we get $\mu_1 \in \Gamma(X) * M^1(G)$.

To complete the proof of the theorem, it remains to prove that $\mu_2 \in \Gamma(X) * M^1(G)$. Take an arbitrary element k of the group Y and substitute v + k for u in (16). Subtracting equation (16) from the resulting equation, we obtain

$$\Delta_k \Delta_{-(I+\widetilde{\alpha})h_2} \Delta_{-2\widetilde{\alpha}h_1} B(v) = 0, \quad v \in Y.$$

Consider the distribution of the random variable $P_2 = 2\xi_1 + (I+\alpha)\xi_2$. Arguing in the same way as in the case when we considered the distribution of the random variable P_1 , we prove that $\mu_2 \in \Gamma(X) * M^1(G)$.

Evidently, if the connected component of zero of a second countable locally compact Abelian group X has dimension 1, then X contains no subgroups topologically isomorphic to the 2-dimensional torus \mathbb{T}^2 . For this reason Theorem A follows from Theorem 2.1.

We note that the statement of Theorem 2.1 in the case if a topological automorphism α of the group X satisfies the conditions

$$I \pm \alpha \in \operatorname{Aut}(\mathbf{X}) \tag{25}$$

follows from [7, Theorem 1].

It is obvious that if a locally compact Abelian group X contains no elements of order 2, then X contains no subgroups topologically isomorphic to the 2-dimensional torus \mathbb{T}^2 . Hence Theorem 2.1 implies the following characterization of the Gaussian distribution proved earlier in [10, Theorem 3], see also [14, Theorem 9.9].

Corollary 2.9 Let X be a second countable locally compact Abelian group containing no elements of order 2. Let α be a topological automorphism of the group X satisfying condition (1). Let ξ_1 and ξ_2 be independent random variables with values in the group X and distributions μ_1 and μ_2 with nonvanishing characteristic functions. Then the symmetry of the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ implies that $\mu_j \in \Gamma(X)$, j = 1, 2.

Let X be a second countable locally compact Abelian group, let G be the subgroup of X generated by all elements of X of order 2, and let α be a topological automorphism of the group X. Suppose that $K = \text{Ker}(I + \alpha) \neq \{0\}$. Let ξ_1 and ξ_2 be independent identically distributed random variables with values in the subgroup K and distribution μ . It is obvious that $\alpha x = -x$ for all $x \in K$. It is easy to see that the conditional distribution of the linear form $P_2 = \xi_1 - \xi_2$ given $P_1 = \xi_1 + \xi_2$ is symmetric. Hence if we consider independent random variables ξ_1 and ξ_2 as independent random variables taking values in X, then the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is also symmetric. Since μ is an arbitrary distribution, from what has been said it follows that a necessary condition for a topological automorphism α for Theorem 2.1 to be true is an inclusion

$$\operatorname{Ker}(I+\alpha) \subset G. \tag{26}$$

However, generally speaking, this condition is not sufficient. The corresponding example can be constructed in the case, when the group X is an a-adic solenoid.

Recall the definition of an a-adic solenoid. Let $a = (a_0, a_1, \ldots, a_n, \ldots,)$, where all $a_j \in \mathbb{Z}$ and $a_j > 1$. Let Δ_a be the group of a-adic integers and let B be the subgroup of the group $\mathbb{R} \times \Delta_a$ of the form $B = \{(n, nu)\}_{n=-\infty}^{\infty}$, where $u = (1, 0, \ldots, 0, \ldots)$. The factor-group $\Sigma_a = (\mathbb{R} \times \Delta_a)/B$ is called an a-adic solenoid (see e.g. [17, §10]). The group Σ_a is compact, connected, has dimension 1, and contains no subgroups topologically isomorphic to the circle group \mathbb{T} . Moreover, Σ_a can contain at most one element of order 2. Assume that the group Σ_a contains an element of order 2 and denote by G the subgroup of Σ_a generated by this element. It follows from the results proved in [12], see also [14, Proposition 11.19, Theorem 11.20 and Remark 11.22]) that the following statement holds.

Proposition 2.10 Let Σ_{a} be an *a*-adic solenoid containing an element of order 2. Let α be a topological automorphism of the group Σ_{a} such that

$$\operatorname{Ker}(I + \alpha) = G \tag{27}$$

and hence condition (26) holds. Then there exist independent random variables ξ_1 and ξ_2 with values in the group $\Sigma_{\mathbf{a}}$ and distributions μ_1 and μ_2 with nonvanishing characteristic functions such that the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric, whereas $\mu_j \notin \Gamma(\Sigma_{\mathbf{a}}) * \mathrm{M}^1(G), j = 1, 2.$

An example of an \boldsymbol{a} -adic solenoid $\Sigma_{\boldsymbol{a}}$ and a topological automorphism $\alpha \in \operatorname{Aut}(\Sigma_{\boldsymbol{a}})$ such that (27) holds is $\Sigma_{\boldsymbol{a}}$, where $\boldsymbol{a} = (3, 3, \ldots, 3, \ldots)$, and α is the multiplication by -3.

3 Theorem A is false for the 2-dimensional torus \mathbb{T}^2

We prove in this section that Theorem A is false for the 2-dimensional torus \mathbb{T}^2 . Let $X = \mathbb{T}^2$. Denote by $x = (z, w), z, w \in \mathbb{T}$, elements of the group X. The character group Y of the group X is topologically isomorphic to the group \mathbb{Z}^2 . Denote by $y = (m, n), m, n \in \mathbb{Z}$, elements of the group Y. Every automorphism $\alpha \in \operatorname{Aut}(X)$ is defined by an integer-valued matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$, where |ad - bc| = 1 and α acts on X as follows

$$\alpha(z,w) = (z^a w^c, z^b w^d), \quad (z,w) \in X$$

The adjoint automorphism $\widetilde{\alpha} \in \operatorname{Aut}(Y)$ is of the form

$$\widetilde{\alpha}(m,n) = (am + bn, cm + dn), \quad (m,n) \in Y.$$

We identify α with the matrix $\begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $\tilde{\alpha}$ with the matrix $\begin{pmatrix} a & c \\ b & d \end{pmatrix}$.

It follows from the definition of the Gaussian distribution on a locally compact Abelian group that the characteristic function of a symmetric Gaussian distribution on the group X is of the form

$$\hat{\gamma}(m,n) = \exp\{-\langle A(m,n), (m,n)\rangle\}, \quad (m,n) \in Y,$$

where $\langle \cdot, \cdot \rangle$ is the standard scalar product in \mathbb{R}^2 , $A = (a_{ij})_{i,j=1}^2$ is a symmetric positive semidefinite matrix.

Theorem 3.1 Let $X = \mathbb{T}^2$ and let G be the subgroup of X generated by all elements of X of order 2. Then there exist a topological automorphism α of the group X satisfying condition (1) and independent random variables ξ_1 and ξ_2 with values in X and distributions μ_1 and μ_2 with nonvanishing characteristic functions such that the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric, whereas $\mu_j \notin \Gamma(X) * M^1(G)$, j = 1, 2.

For the proof of Theorem 3.1 we need the following lemmas.

Lemma 3.2 ([23], see also [6, Lemma 11.2]) Consider the 2-dimensional torus \mathbb{T}^2 . Let $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ be a topological automorphism of the group \mathbb{T}^2 . If ad - bc = 1 and a + d < -2, then there exist symmetric positive semidefinite 2×2 matrices A_1 and A_2 such that

$$\det A_1 = \det A_2 > 0 \tag{28}$$

and

$$A_1 + A_2 \widetilde{\alpha} = 0. \tag{29}$$

Lemma 3.3 ([6, Lemma 16.1]) Let X be a second countable locally compact Abelian group with character group Y. Let α be a topological automorphism of X. Let ξ_1 and ξ_2 be independent random variables with values in the group X and distributions μ_1 and μ_2 . The conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric if and only if the characteristic functions $\hat{\mu}_j(y)$ satisfy the equation

$$\hat{\mu}_1(u+v)\hat{\mu}_2(u+\widetilde{\alpha}v) = \hat{\mu}_1(u-v)\hat{\mu}_2(u-\widetilde{\alpha}v), \quad u,v \in Y.$$
(30)

Proof of Theorem 3.1 Consider a topological automorphism $\alpha = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ of the group X such that det $\alpha = 1$ and a + d = -3. It follows from this that det $(I + \alpha) = -1$ and hence $I + \alpha \in \text{Aut}(X)$. For this reason condition (1) holds. By Lemma 3.2, there exist symmetric positive semidefinite 2×2 matrices A_1 and A_2 such that (28) and (29) are valid.

Note that $\det(I - \alpha) = \det(I - \widetilde{\alpha}) = 5$. Hence $I - \widetilde{\alpha} \notin \operatorname{Aut}(X)$ and it follows from this that $H = (I - \widetilde{\alpha})(Y)$ is a proper subgroup of Y. This implies that $K = A(X, H) \neq \{0\}$.

Let us check that $Y^{(2)} \setminus H \neq \emptyset$. Suppose the contrary is true, i.e., $Y^{(2)} \subset H$. Then, on the one hand, for H we have the following possibilities:

$$H = \{(2m, 2n) : m, n \in \mathbb{Z}\}, \quad H = \{(2m, n) : m, n \in \mathbb{Z}\},$$
(31)

$$H = \{ (m, 2n) : m, n \in \mathbb{Z} \}, \quad H = \{ (2m, 2n), (2m - 1, 2n - 1) : m, n \in \mathbb{Z} \}.$$
(32)

On the other hand, $H = \{((1-a)m - bn, -cm + (1-d)n) : m, n \in \mathbb{Z}\}$ and this implies that

$$(-b, 1-d), (1-a, -c) \in H.$$
 (33)

Since a + d = -3, this implies that a and d have different parity. In view of ad - bc = 1, we have either a is even and b, c, d are odd or d is even and a, b, c are odd. Anyway (33) contradicts (31) and (32). Hence $Y^{(2)} \setminus H \neq \emptyset$.

Take $0 < \kappa < 1$ and consider on the group X the distribution

$$\pi_1 = \kappa E_{(1,1)} + (1 - \kappa)m_K$$

and the signed measure

$$\pi_2 = \frac{1}{\kappa} E_{(1,1)} + \frac{\kappa - 1}{\kappa} m_K.$$

Since H = A(Y, K), it follows from (4) that

$$\widehat{m}_{K}(y) = \begin{cases} 1, & \text{if } y \in H, \\ 0, & \text{if } y \notin H, \end{cases}$$

and the characteristic functions $\hat{\pi}_j(y)$ are of the form

$$\hat{\pi}_1(y) = \begin{cases} 1, & \text{if } y \in H, \\ \kappa, & \text{if } y \notin H, \end{cases} \quad \hat{\pi}_2(y) = \begin{cases} 1, & \text{if } y \in H, \\ \frac{1}{\kappa}, & \text{if } y \notin H. \end{cases}$$
(34)

Let us check that the characteristic functions $\hat{\pi}_j(y)$ satisfy equation (30). To see this we verify that for any $u, v \in Y$ both sides of equation (30) are equal to 1. Suppose that there exist $u, v \in Y$ such that $\hat{\pi}_1(u+v)\hat{\pi}_2(u+\tilde{\alpha}v) \neq 1$. In view of (34) then either $u+v \in H, u+\tilde{\alpha}v \notin H$ or $u+v \notin H, u+\tilde{\alpha}v \in H$. In both cases we obtain that $(I-\tilde{\alpha})v \notin H$. This is impossible because $H = (I-\tilde{\alpha})(Y)$. Thus, the left hand-side of equation (30) for any $u, v \in Y$ is equal to 1. Reasoning similarly we check that the right hand-side of equation (30) for any $u, v \in Y$ is also equal 1. Hence the characteristic functions $\hat{\pi}_j(y)$ satisfy equation (30).

It is easy to see that since (29) is valid, the functions

$$h_j(m,n) = \exp\{-\langle A_j(m,n), (m,n) \rangle\}, \quad j = 1, 2,$$

satisfy equation (30). Hence the functions

$$g_j(m,n) = h_j^k(m,n)\hat{\pi}_j(m,n), \quad (m,n) \in Y, \ j = 1,2,$$
(35)

for any natural k also satisfy equation (30).

It follows from (28) that there is $\varepsilon > 0$ such that

$$\langle A_j(m,n),(m,n)\rangle \ge \varepsilon(m^2 + n^2), \quad j = 1, 2.$$
(36)

Since $g_j(0,0) = 1$, inequalities (36) imply that for a big enough k the inequalities

$$\sum_{(m,n)\in Y} g_j(m,n) < 2, \quad j = 1, 2, \tag{37}$$

hold true. Put

$$\rho_j(z,w) = \sum_{(m,n)\in Y} g_j(m,n)\bar{z}^m \bar{w}^n, \quad (z,w)\in X, \ j=1,2.$$

Since $g_j(-y) = g_j(y)$ for all $y \in Y$, it follows from (37) that then $\rho_j(z, w) > 0$ for all $(z, w) \in X$, j = 1, 2. It is also obvious that

$$\int_{X} \rho_j(z, w) dm_X(z, w) = 1, \quad j = 1, 2.$$

Thus, the functions $\rho_j(z, w)$ are densities with respect to m_X of some distributions μ_j on the group X. In so doing, $\hat{\mu}_j(y) = g_j(y)$, j = 1, 2. Let ξ_1 and ξ_2 are independent random variables with values in the group X and distributions μ_1 and μ_2 . Since the characteristic functions $\hat{\mu}_j(y)$ satisfy equation (30), by Lemma 3.3, the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric. Note now that $Y^{(2)} = A(Y, G)$. For this reason if $\mu \in M^1(X)$ and $\mu \in \Gamma(X) * M^1(G)$, then the restriction of the characteristic function $\hat{\mu}(y)$ to the subgroup $Y^{(2)}$ is the characteristic function of a Gaussian distribution. Since $Y^{(2)} \setminus H \neq \emptyset$, it follows from (34) and (35) that $\mu_j \notin \Gamma(X) * M^1(G)$, j = 1, 2.

In view of Theorem 3.1, it is interesting to note that Theorem A holds true for the group \mathbb{T}^2 if we substitute condition (1) for (25) ([4], see also [6, Theorem 16.8]). Namely, the following statement holds.

Theorem B Consider the 2-dimensional torus \mathbb{T}^2 and let G be the subgroup of \mathbb{T}^2 generated by all elements of \mathbb{T}^2 of order 2. Assume that a topological automorphism α of the group \mathbb{T}^2 satisfies conditions (25). Let ξ_1 and ξ_2 be independent random variables with values in the group \mathbb{T}^2 and distributions μ_1 and μ_2 with nonvanishing characteristic functions. If the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ is symmetric, then $\mu_j = \gamma_j * \omega_j$, where $\gamma_j \in \Gamma(\mathbb{T}^2)$, $\omega_j \in \mathrm{M}^1(G)$, j = 1, 2. In so doing the Gaussian distributions γ_j are concentrated on cosets of the same dense one-parameter subgroup in \mathbb{T}^2 . In conclusion, taking into account Theorems 2.1 and 3.1, we formulate the following conjecture.

Conjecture 3.4 Let X be a second countable locally compact Abelian group, let G be the subgroup of X generated by all elements of X of order 2, and let α be a topological automorphism of the group X satisfying condition (1). Let ξ_1 and ξ_2 be independent random variables with values in X and distributions μ_1 and μ_2 with nonvanishing characteristic functions. The symmetry of the conditional distribution of the linear form $L_2 = \xi_1 + \alpha \xi_2$ given $L_1 = \xi_1 + \xi_2$ implies that $\mu_j \in \Gamma(X) * M^1(G)$, j = 1, 2, if and only if the group X contains no subgroups topologically isomorphic to the 2-dimensional torus \mathbb{T}^2 .

The sufficiency in this assertion follows from Theorem 2.1.

Acknowledgements This article was written during my stay at the Department of Mathematics University of Toronto as a Visiting Professor. I am very grateful to Ilia Binder for his invitation and support. I would also like to thank the reviewer for carefully reading the article and useful comments.

References

- Bogachev, V.I.: Gaussian Measures, Mathematical Surveys and Monographs, 62. American Mathematical Society, Providence, RI (1998)
- Feldman, G.M.: Gaussian distributions on locally compact Abelian groups. Theory Probab. Appl. 23, 529–542 (1979). https://doi.org/10.1137/1123062
- Feldman, G.M.: On the Heyde theorem for finite Abelian groups. J. Theoretical Probab. 17, 929–941 (2004). https://doi.org/10.1007/s10959-004-0583-0
- [4] Feldman, G.M.: On a characterization theorem for locally compact abelian groups. Probab. Theory Relat. Fields. 133, 345–357 (2005). https://doi.org/10.1007/s11118-020-09885-x
- [5] Feldman, G.M.: On the Heyde theorem for discrete Abelian groups. Studia Math. 177, 67–79 (2006). https://doi.org/10.4064/sm177-1-5
- [6] Feldman, G.: Functional Equations and Characterization Problems on Locally Compact Abelian Groups, EMS Tracts in Mathematics, 5. European Mathematical Society, Zürich, (2008)
- [7] Feldman, G.M.: The Heyde theorem for locally compact Abelian groups. J. Funct. Anal. 258, 3977–3987 (2010). https://doi.org/10.1016/j.jfa.2010.03.005
- [8] Feldman, G.M.: On a characterization of convolutions of Gaussian and Haar distributions. Math. Nachr. 286, 340–348 (2013). https://doi.org/10.1002/mana.201100320
- Feldman, G.M.: On a characterization theorem for the group of p-adic numbers. Publ. Math. Debrecen. 87, 147–166 (2015). https://doi.org/10.5486/PMD.2015.7100
- [10] Feldman, G.M.: Heyde's characterization theorem on some locally compact Abelian groups. Theory Probab. Appl. 62, 399–412 (2018). https://doi.org/10.1137/S0040585X97T988708
- [11] Feldman, G.M.: On a characterization theorem for probability distributions on discrete Abelian groups. Theory Probab. Appl. 63, 594–612 (2019). https://doi.org/10.1137/S0040585X97T989271

- Feldman, G.M.: On a characterization theorem for connected locally compact Abelian groups. J. Fourier Anal. Appl. 26, no. 1, Paper No. 14, 22 pp., (2020). https://doi.org/10.1007/s00041-019-09721-w
- [13] Feldman, G.M.: A characterization theorem on compact Abelian groups. J. Fourier Anal. Appl. 27, no. 5, Paper No. 86, 23 pp., (2021). https://doi.org/10.1007/s00041-021-09889-0
- [14] Feldman, G.: Characterization of Probability Distributions on Locally Compact Abelian Groups, Mathematical Surveys and Monographs, 273. American Mathematical Society, Providence, RI (2023)
- [15] Feldman, G.M.: Heyde theorem on locally compact Abelian groups with the connected component of zero of dimension 1. Potential Analysis, (2023). https://doi.org/10.1007/s11118-023-10095-4
- [16] Fuchs, L: Infinite Abelian Groups, 2, Pure Applied Math, 36-II. Academic Press, New York (1973)
- [17] Hewitt, E., Ross, K. A.: Abstract Harmonic Analysis, 1: Structure of Topological Groups, Integration Theory, Group Representations, 2nd ed., Grundlehren der Mathematischen Wissenschaften, 115, Springer-Verlag, New York (1994)
- [18] Heyde, C.C.: Characterization of the normal low by the symmetry of a certain conditional distribution. Sankhya. 32, Ser. A. 115–118 (1970). http://www.jstor.org/stable/25049642
- [19] Kagan, A. M., Linnik, Yu. V., Rao C.R.: Characterization Problems in Mathematical Statistics, Wiley Series in Probability and Mathematical Statistics. John Wiley & Sons, New York (1973)
- [20] Myronyuk, M.V.: Heyde's characterization theorem for discrete abelian groups. J. Aust. Math. Soc. 88, 93–102 (2010). https://doi.org/10.1017/S1446788709000378
- [21] Myronyuk, M.V.: The Heyde theorem on a-adic solenoids. Colloquium Mathematicum. 132, 195–210 (2013) https://doi.org/10.4064/cm132-2-3
- [22] Myronyuk, M.V., Feldman, G.M.: On a characterization theorem on finite abelian groups. Siberian Math. J. 46, 315–324 (2005). https://doi.org/10.1007/s11202-005-0033-y
- [23] Myronyuk, M.V., Feldman, G.M.: Independent linear statistics on the two-dimensional torus. Theory Probab. Appl. 52, 78–92 (2008). https://doi.org/10.1137/S0040585X97982785
- [24] Parthasarathy, K.R.: Probability Measures on Metric Spaces. Academic Press, New York (1967)

B. Verkin Institute for Low Temperature Physics and Engineering of the National Academy of Sciences of Ukraine Nauky Ave. 47, 61103 Kharkiv, Ukraine

Department of Mathematics University of Toronto 40 St. George Street Toronto, ON, M5S 2E4 Canada

e-mail: gennadiy_f@yahoo.co.uk