SIMILAR POINT CONFIGURATIONS VIA GROUP ACTIONS

P. BHOWMIK, A. GREENLEAF, A. IOSEVICH, S. MKRTCHYAN, AND F. RAKHMONOV

ABSTRACT. We prove that for $d \geq 2, k \geq 2$, if the Hausdorff dimension of a compact set $E \subset \mathbb{R}^d$ is greater than $\frac{d^2}{2d-1}$, then, for any given r > 0, there exist $(x^1, \ldots, x^{k+1}) \in E^{k+1}, (y^1, \ldots, y^{k+1}) \in E^{k+1}$, a rotation $\theta \in O_d(\mathbb{R})$, and a vector $a \in \mathbb{R}^d$ such that $rx^j = \theta y^j - a$ for $1 \leq j \leq k+1$. Such a result on existence of similar k-simplices in thin sets had previously been established under a more stringent dimensional threshold in Greenleaf, Iosevich and Mkrtchyan [5]. The argument we are use to prove the main result here was previously employed in Bhowmik and Rakhmonov [1] to establish a finite field version. We also show the existence of multi-similarities of arbitrary multiplicity in \mathbb{R}^d , show how to extend these results from similarities to arbitrary proper continuous maps, as well as explore a general group-theoretic formulation of this problem in vector spaces over finite fields.

1. INTRODUCTION

The celebrated Falconer distance conjecture (see e.g. [4], [7]) states that if the Hausdorff dimension $\dim_{\mathcal{H}}(E)$ of a compact set $E \subset \mathbb{R}^d$, $d \geq 2$, is greater than $\frac{d}{2}$, then the Lebesgue measure of the distance set $\Delta(E) := \{|x - y| : x, y \in E\}$ is positive. Currently, the best threshold known for the Falconer distance problem in two dimensions is $\frac{5}{4}$, due to Guth, Iosevich, Ou and Wang ([6]). In higher dimensions, the best exponent known is $\frac{d}{2} + \frac{1}{4} - \frac{1}{8d+4}$, due to Du, Ou, Ren, and Zhang ([2]).

More generally, given a compact set $E \subset \mathbb{R}^d$, $d \geq 2$, let

$$\Delta_k(E) = \{v(x^1, \dots, x^{k+1}) : x^j \in E\} \subset \mathbb{R}^{\binom{k+1}{2}},$$

where $v(x^1, \ldots, x^{k+1})$ is the $\binom{k+1}{2}$ -vector whose entries are $|x^i - x^j|$, $1 \le i < j \le k+1$, listed in the dictionary order.

We can think of $\Delta_k(E)$ as the set of non-congruent (k + 1)-tuples of elements of E (modulo the action of the symmetric group S_{k+1}). More precisely, we say

Date: April 30, 2024.

AG is supported in part by National Science Foundation award DMS-2204943. AI was supported in part by the National Science Foundation under grant no. HDR TRIPODS - 1934962 and by the NSF DMS - 2154232. SM is supported in part by the Simons Foundation Grant no. 523555.

that $(x^1, \ldots, x^{k+1}) \in E^{k+1}$ is congruent to $(y^1, \ldots, y^{k+1}) \in E^{k+1}$ if there exists an orthogonal matrix $\theta \in O_d(\mathbb{R})$ and $a \in \mathbb{R}^d$ such that $y^j = \theta x^j + a$ for $1 \le j \le k+1$. Equivalently, for $k \le d$, one can consider $\Delta_k(E)$ as the set of congruence classes of k-simplices generated by points in E.

One can also consider *similarity* classes of k-simplices. Greenleaf, Iosevich, and Mkrtchyan proved in [5] that if the Hausdorff dimension of E, $\dim_{\mathcal{H}}(E)$, satisfies $\dim_{\mathcal{H}}(E) > s_k := \frac{dk+1}{k+1}$, then

(1.1) for every
$$r > 0$$
, there exist distinct
 $(x^1, \dots, x^{k+1}) \in E^{k+1}, \quad (y^1, \dots, y^{k+1}) \in E^{k+1}$
such that $|y^i - y^j| = r|x^i - x^j|$ for every $1 \le i < j \le k+1$

In other words, given any r > 0, there exist two (k + 1)-tuples of elements of E which are similar via a translation, rotation, and the prescribed scaling factor r. More precisely, the following was proved in [5]: Given E with $\dim_{\mathcal{H}}(E) > s_k$, let $s_k < s < \dim_{\mathcal{H}}(E)$ and μ be a Frostman measure on \mathbb{R}^d , supported on E and of finite *s*-energy, as in Frostman's Lemma (see [7]). From μ one forms the natural configuration measure ν_k on $\Delta_k(E)$, i.e., for any $f \in C_0\left(\mathbb{R}^{\binom{k+1}{2}}\right)$,

$$\int f(\vec{t}) d\nu_k(\vec{t}) = \int \cdots \int f\left(v(x^1, \dots, x^{k+1})\right) d\mu(x^1) d\mu(x^2) \dots d\mu(x^{k+1}).$$

For $0 < r < \infty$, define

(1.2)
$$\Delta_k^r(E) = \left\{ \vec{t} \in \Delta_k(E) : r\vec{t} \in \Delta_k(E) \right\} \subset \Delta_k(E),$$

which records the (k+1)-point configurations in E which also have an r-scaled similar copy in E. Then, in [5] it was shown that if $\dim_{\mathcal{H}}(E) > s_k$, then

(1.3)
$$\nu_k(\Delta_k^r(E)) > 0$$

which implies (and is in fact much stronger than) statement (1.1).

In the first results of the current paper, Theorem 1.1 and Corollary 1.2, we will show that while (1.3) might require the full strength of the machinery used in [5], one can derive the conclusion (1.1) for any k under a much lower, k-independent dimensional threshold, $\dim_{\mathcal{H}}(E) > \frac{d^2}{2d-1}$, and extend it to k > d (for which (k + 1)point sets no longer correspond to k-simplices). In brief, the work in [5] shows, and uses, that if $\dim_{\mathcal{H}}(E) > s_k = \frac{dk+1}{k+1}$ defined above, then the measure ν_k on $\mathbb{R}^{\binom{k+1}{2}}$, supported on $\Delta_k(E)$, is absolutely continuous with respect to Lebesgue measure $d\vec{t}$, with a density in $L^2\left(\mathbb{R}^{\binom{k+1}{2}}\right)$. In contrast the method we employ in this paper merely requires use of the lower threshold, $\dim_{\mathcal{H}}(E) > \frac{d^2}{2d-1}$, above which the configuration measure ν_1 on the distance set $\Delta(E) = \Delta_1(E)$ is known to be in $L^2(\mathbb{R})$. Theorem 1.3 then uses a continuous version of the pigeonhole principle (Proposition 2.3, from [5]) to extend these results to multi-similarities of any multiplicity, while Theorem 1.4 in Section 3 shows how these results can be extended to more general transformations in \mathbb{R}^d .

In addition to the Euclidean setting described so far, we also examine an analogous problem in the context of general transitive group actions on vector spaces over finite fields. See Theorems 1.5 and 1.6, and Corollary 1.7 and their proofs in Section 4. Our goal is to lay the groundwork for future explorations in this broader context, including both finite field and Euclidean problems.

1.1. Results in Euclidean spaces. In the Euclidean setting, our main result is:

Theorem 1.1. Let $E \subset \mathbb{R}^d$, $d \geq 2$, be compact with Hausdorff dimension satisfying $\dim_{\mathcal{H}}(E) > \frac{d^2}{2d-1}$. Then, for any r > 0, there exist $\theta \in O_d(\mathbb{R})$ and $a \in \mathbb{R}^d$ such that $rE \cap (\theta E - a)$ has positive Hausdorff dimension, and is thus uncountably infinite. In particular, for every $k \in \mathbb{N}$, (1.1) holds.

An immediate consequence of Theorem 1.1 is the following.

Corollary 1.2. Let $E \subset \mathbb{R}^d$, $d \geq 2$, be a compact set with $\dim_{\mathcal{H}}(E) > \frac{d^2}{2d-1}$. Then, for any $k \geq 1$ and r > 0, there exist distinct $(x^1, \ldots, x^{k+1}) \in E^{k+1}$ and $(y^1, \ldots, y^{k+1}) \in E^{k+1}$, such that $|x^i - x^j| = r|y^i - y^j|$ for all $1 \leq i < j \leq k+1$. Thus, if $1 \leq k \leq d$, for any r > 0 there exist r-similar k-simplices in E.

As in [5] for the higher threshold s_k , a measure-theoretic pigeonhole principle allows one to leverage the proof of Theorem 1.1 to obtain the existence of *multi-similarities* in E, of arbitrarily large multiplicity, for $\dim_{\mathcal{H}}(E)$ above the new, lower threshold:

Theorem 1.3. Let $E \subset \mathbb{R}^d$, $d \geq 2$, be a compact set with $\dim_{\mathcal{H}}(E) > \frac{d^2}{2d-1}$. Then for all $n \geq 1$ and any compact interval $I \subset (0, \infty)$, there exists $M = M(n, E, I) \in \mathbb{N}$ such that for any distinct $r_1, \ldots, r_M \in I$, there exist distinct r_{i_1}, \ldots, r_{i_n} such that: For every $k \geq 1$, there is a $\vec{t} \in \mathbb{R}^{\binom{k+1}{2}}$ such that

$$\left\{r_{i_1}\vec{t}, r_{i_2}\vec{t}, \ldots, r_{i_n}\vec{t}\right\} \subset \Delta_k(E),$$

i.e., there exists an n-similarity of (k+1)-point sets in E with scaling factors coming from among $\{r_i\}_{i=1}^M$.

See Section 2.2 for the relevant definitions, statements and proof.

We also extend Theorem 1.1 by showing that the dilations in Theorem 1.1 can be replaced by more general transformations:

Theorem 1.4. Suppose $T : \mathbb{R}^d \to \mathbb{R}^d$ is a proper continuous map, and $E \subset \mathbb{R}^d$ is compact. Given $s < \dim_{\mathcal{H}}(E)$, let μ be a Frostman measure μ supported on E and of finite s-energy. Define the pushforward measure μ_T by its action on $f \in C_0(\mathbb{R}^d)$,

$$\int f(x)d\mu_T(x) := \int f(T(x))d\mu(x)$$

If s is such that

$$\int |\widehat{\mu}_T(\xi)|^2 |\xi|^{-s\frac{d-1}{d}} d\xi < \infty$$

then $T(E) \cap (\theta E - a)$ has positive Hausdorff dimension, and is thus uncountably infinite. Thus, for any $k \in \mathbb{N}$, we can find points $(x_1, \ldots, x_{k+1}) \in E^{k+1}$ and $(y_1, \ldots, y_{k+1}) \in E^{k+1}$ such that $T(x_i) = \theta y_i - a$ for all i for some $\theta \in O_d(\mathbb{R})$.

For example, Theorem 1.4 applies if T is a diffeomorphism of \mathbb{R}^d which is the identity outside of a compact set, since such maps preserve Sobolev spaces of all orders.

1.2. Finite field setting and transitive group actions. The idea behind the proof of (1.1) was originally implemented in the finite field setting by Bhowmik and Rakhmonov in [1].

Let \mathbb{F}_q^d be the vector space of dimension $d \geq 2$ over the finite field \mathbb{F}_q with q elements. We define a function $\|\cdot\| : \mathbb{F}_q^d \to \mathbb{F}_q$ by $\|\alpha\| := \alpha_1^2 + \cdots + \alpha_d^2$ for $\alpha = (\alpha_1, \ldots, \alpha_d) \in \mathbb{F}_q^d$. Note that this is not a norm, as it is \mathbb{F}_q -valued (and homogeneous of degree 2), and we do not impose any metric structure on \mathbb{F}_q^d . However, $\|\cdot\|$ does share an important feature of the Euclidean norm: it is invariant under orthogonal transformations.

For $m \geq 2$, let $(\mathbb{F}_q)^m := \{a^m : a \in \mathbb{F}_q\}$; in particular, $(\mathbb{F}_q)^2$ is the set of quadratic residues in \mathbb{F}_q . Then the following holds.

Theorem 1.5. [1, Thm. 1.3] Let $k \ge 1$. Suppose $\emptyset \ne A \subset \{(i, j) : 1 \le i < j \le k+1\}$ and $r \in (\mathbb{F}_q)^2 \setminus \{0\}$. If $E \subset \mathbb{F}_q^d$ with $|E| \ge \sqrt{k+1}q^{\frac{d}{2}}$, then there exist $(x_1, \ldots, x_{k+1}) \in E^{k+1}$ and $(y_1, \ldots, y_{k+1}) \in E^{k+1}$ such that $||y_i - y_j|| = r||x_i - x_j||$ for $(i, j) \in A$, and $x_i \ne x_j, y_i \ne y_j$, for all $1 \le i < j \le k+1$.

As a straightforward corollary, through the variation of the underlying set A, we can establish thresholds for the existence of dilated k-cycles, k-paths, and k-stars

(for $k \geq 3$) with a dilation ratio $r \in (\mathbb{F}_q)^2 \setminus \{0\}$. Notably, Rakhmonov has previously examined the cases of 2-paths, 4-cycles, and k-simplices in [8].

The original idea for a proof of Theorem 1.5 was based on the a group-theoretic approach. (However, in [1], the first and fifth listed authors opted for an alternative, weaker version of this method.) This group-theoretic approach can be generalized to arbitrary transitive group actions, which we state here and prove in Section 4:

Theorem 1.6. Suppose that G is a finite group acting transitively on a set X, and let E and H be subsets of X. Then

(1.4)
$$\max_{g \in G} |H \cap gE| \ge \frac{|H||E|}{|X|}$$

Another application of Theorem 1.6 pertains to $SL_d(\mathbb{F}_q)$ actions:

Corollary 1.7. Suppose $k \ge d$. Let $E \subset \mathbb{F}_q^d$, with $|E| \ge \sqrt{k+1}q^{\frac{d}{2}}$. Then for every $r \in (\mathbb{F}_q)^d \setminus \{0\}$, there exist $(x_1, \ldots, x_{k+1}) \in E^{k+1}$ and $(y_1, \ldots, y_{k+1}) \in E^{k+1}$ such that $\det(x_{i_1}, x_{i_2}, \ldots, x_{i_d}) = r \det(y_{i_1}, y_{i_2}, \ldots, y_{i_d})$

for all d-tuples of elements from $(x_1, \ldots, x_{k+1}) \in E^{k+1}$ and $(y_1, \ldots, y_{k+1}) \in E^{k+1}$, respectively, where det (u_1, u_2, \ldots, u_d) is the determinant of the $d \times d$ matrix where the columns are vectors in \mathbb{F}_q^d .

The paper is organized as follows. In Section 2 we prove Theorem 1.1 in \mathbb{R}^d , and recall in Section 2.2 the material needed to state and prove Theorem 1.3 on multi-similarities. Then, in Section 3 we prove Theorem 1.4, extending the context of Theorem 1.1 from dilations to more general transformations in \mathbb{R}^d . Finally, in Section 4 we prove the group action result Theorem 1.6 and show how it implies Theorem 1.5 and Corollary 1.7 in the finite field setting.

2. Similarities and multi-similarities in \mathbb{R}^d

2.1. **Proof of Theorem 1.1.** Let $E \subset \mathbb{R}^d$ be compact. Then, for any $s < \dim_{\mathcal{H}}(E)$, we can equip E with a Frostman measure μ , a probability measure supported on E, of finite s-energy and satisfying a dimension s ball condition. For $0 < r < \infty$ and $\theta \in O_d(\mathbb{R})$, the group of orthogonal transformations of \mathbb{R}^d , define μ_r and μ^{θ} , also probability measures of dimension s, by their actions on $f \in C_0(\mathbb{R}^d)$,

$$\int f(x)d\mu_r(x) := \int f(rx)d\mu(x), \quad \int f(x)d\mu^{\theta}(x) := \int f(\theta x)d\mu(x),$$

with Fourier transforms $\hat{\mu}_r(\xi) = \hat{\mu}(r\xi)$ and $\hat{\mu}^{\theta}(\xi) = \hat{\mu}(\theta^{-1}\xi)$. Using this notation, consider

(2.1)
$$\int \left(\mu_r * \mu^\theta\right)(x) dx.$$

If the expression in (2.1) converges we are done because, formally, it equals

$$\left(\int d\mu_r(x)\right) \cdot \left(\int d\mu^{\theta}(x)\right) = \int d\mu_r(x) \cdot \int d\mu^{\theta}(x) = 1.$$

However, in order to decouple the integrals, one needs to demonstrate that (2.1) converges, at least for a set of θ of full measure. If this convergence is established, and if we define

$$A_{r,\theta} := \operatorname{supp} \left(\mu_r * \mu^{\theta} \right) \subset \mathbb{R}^d$$

then one has

$$1 = \int \left(\mu_r * \mu^\theta\right)(x) dx = \int \mathbf{1}_{A_{r,\theta}}(x) \cdot \left(\mu_r * \mu^\theta\right)(x) dx.$$

Squaring this, applying Cauchy-Schwarz and then integrating with respect to the normalized Haar measure $d\theta$ on $O_d(\mathbb{R})$, it follows that

$$1 = 1^{2} = \int_{O_{d}(\mathbb{R})} \left(\int_{\mathbb{R}^{d}} \mathbf{1}_{A_{r,\theta}}(x) \cdot \left(\mu_{r} * \mu^{\theta}\right)(x) dx \right)^{2} d\theta$$

$$\leq \int_{O_{d}(\mathbb{R})} \left(\int_{\mathbb{R}^{d}} \mathbf{1}_{A_{r,\theta}}(x)^{2} dx \right) \cdot \left(\int_{\mathbb{R}^{d}} \left(\mu_{r} * \mu^{\theta}\right)(x)^{2} dx \right) d\theta$$

$$= \int \mathcal{L}^{d} \left(A_{r,\theta}\right) \cdot \int \left(\mu_{r} * \mu^{\theta}\right)(x)\right)^{2} dx d\theta$$

$$\leq \left(\sup_{\theta \in O_{d}(\mathbb{R})} \mathcal{L}^{d} \left(A_{r,\theta}\right) \right) \cdot \int \int \left(\mu_{r} * \mu^{\theta}\right)(x)\right)^{2} dx d\theta,$$

where \mathcal{L}^d denotes Lebesgue measure on \mathbb{R}^d . Hence,

(2.2)
$$\sup_{\theta} \mathcal{L}^d(A_{r,\theta}) \ge \frac{1}{\int \int (\mu_r * \mu(\theta \cdot)(x))^2 dx \, d\theta}.$$

To obtain a lower bound for the left hand side of (2.2), we need an upper bound on the denominator $\int \int (\mu_r * \mu(\theta \cdot)(x))^2 dx d\theta$ of the right hand side. To obtain this, we use the following result. **Theorem 2.1.** (Wolff [9] for d = 2; Du and Zhang [3] for $d \ge 3$) Let μ be a measure, supported on a compact set on \mathbb{R}^d and of dimension α for some $\alpha \in (\frac{d}{2}, d)$. Then,

$$\int_{S^{d-1}} |\widehat{\mu}(R\omega)|^2 \, d\omega \le C_{\alpha,\mu} R^{-\alpha \frac{d-1}{d}}$$

Returning to the proof of Theorem 1.1, define $F_{r,\theta}(x) = (\mu_r * \mu^{\theta})(x)$. Then,

$$\int \int F_{r,\theta}^2(x) d\theta \, dx = \int \int |\widehat{\mu}_r(\xi)|^2 |\widehat{\mu}(\theta\xi)|^2 d\theta \, d\xi \lesssim \int |\widehat{\mu}_r(\xi)|^2 |\xi|^{-\alpha \frac{d-1}{d}} d\xi,$$

where for the inequality we have used Theorem 2.1 for any $\frac{d}{2} < \alpha < \dim_{\mathcal{H}}(E)$. Taking $\alpha = s$, the dimension of μ , then if $s > \frac{d^2}{2d-1}$, the denominator of the right side of (2.2) is bounded above if $\dim_{\mathcal{H}}(E) > \frac{d^2}{2d-1}$, and thus the left hand side is bounded below, say by $c_r > 0$. This also shows that in fact the integral in (2.1) converges, thereby justifying the calculations above.

(Note for use in the next subsection that c_r depends on s and the s-energy of μ_r ; thus, as r ranges over any compact interval $I \subset (0, \infty)$, c_r is uniformly bounded away from 0.)

We have shown that, for any r > 0, there exists $\theta \in O_d(\mathbb{R})$ such that

$$\mathcal{L}^d\left(A_{r,\theta}\right) \ge c_r > 0,$$

or equivalently

$$\mathcal{L}^{d}(\{a \in \mathbb{R}^{d} : \mu\{y \in E : (\exists x \in E) (rx = \theta y - a)\} > 0\}) \ge c_{r} > 0$$

Fixing any one such a, it follows that $\{y \in E : (\exists x \in E) rx = \theta y - a\}$ has positive μ measure, thus has positive Hausdorff dimension, and hence is an (uncountably) infinite set. In particular, for any $k \ge 1$, there exist $(x^1, \ldots, x^{k+1}) \in E^{k+1}$, $(y^1, \ldots, y^{k+1}) \in E^{k+1}$, $\theta \in O_d(\mathbb{R})$, and $a \in \mathbb{R}^d$ such that $rx^j = \theta y^j - a$ for $1 \le j \le k+1$. This completes the proof of Theorem 1.1.

2.2. Extension to multi-similarities: Proof of Theorem 1.3. By the parenthetical comment above,

(2.3)
$$(\forall I \subset (0,\infty), I \text{ compact}) \ (\exists c_I > 0) \ (\forall r \in I) \ c_r \ge c_I.$$

Modifying slightly the proof of Theorem 1.1 and using a continuous version of the pigeonhole principle from [5] will allow us to prove the existence of arbitrarily many simultaneous similarities in E. We start with some background.

The set $\Delta_k^r(E)$ defined in (1.2) records the pairs $\{\vec{t}, r\vec{t}\} \subset \Delta_k(E)$ of (k+1)-point sets in E which are similar by an r-scaling and translation. Similarly, one can look

at *n*-tuples of similar (k+1)-point sets. Recall the following definition from [5]; since all line segments are similar to each other, this is only interesting for $k \ge 2$.

Definition 2.2. Let $n \ge 1$. A multi-similarity of multiplicity n of (k + 1)-point sets in E is a set of the form $\{r_1 \vec{t}, r_2 \vec{t}, \ldots, r_n \vec{t}\} \subset \Delta_k(E)$, for some $\vec{t} \in \mathbb{R}^{\binom{k+1}{2}}$ and $r_1, r_2, \ldots, r_n \in (0, \infty)$, pairwise distinct. (If $k \le d$, this is a simultaneous similarity of k-simplices in E under n different scalings.)

In [5], it was shown that the version of our Theorem 1.1 with the higher threshold $\dim_{\mathcal{H}}(E) > s_k$ had as a corollary a multi-similarity version, [5, Thm. 1.7]. This was done by combining a multi-similarity version of (1.3) with a measure-theoretic version of the pigeonhole principle [5, Lem. 2.1]. We can combine this reasoning with a modification of the proof of Theorem 1.1 to obtain Theorem 1.3, i.e., the existence, above the same lower threshold of Theorem 1.1, of multi-similarities of all multiplicities.

We start from the same expression (2.1) as above, but now consider θ as a variable rather than a parameter. Define $D_r = \operatorname{supp}_{x,\theta} (\mu_r * \mu^{\theta}) \subset \mathbb{R}^d \times O_d(\mathbb{R})$. Then, from

$$1 = \int \int_{\mathbb{R}^d \times O_d(\mathbb{R})} \left(\mu_r * \mu^\theta \right)(x) \, dx \, d\theta = \int \int_{\mathbb{R}^d \times O_d(\mathbb{R})} \mathbf{1}_{D_r}(x,\theta) \cdot \left(\mu_r * \mu^\theta \right)(x) \, dx \, d\theta,$$

a variation of the calculation in the previous section yields

$$1 \le \left(\mathcal{L}^d \times d\theta\right) (D_r) \cdot \int \int F_{r,\theta}(x,\theta)^2 \, dx \, d\theta$$

and thus

(2.4) $\left(\mathcal{L}^d \times d\theta\right)(D_r) \ge c_r.$

Now recall the

Proposition 2.3. Measure-theoretic Pigeonhole Principle (5)

Let $\mathcal{X} = (X, \mathcal{M}, \sigma)$ be a measure space with $\sigma(X) < \infty$. For $0 < c < \sigma(X)$, define $\mathcal{M}_c = \{A \in \mathcal{M} : \sigma(A) \ge c\}$. Then, for every $n \ge 1$, there exists an $N = N(\mathcal{X}, c, n)$ such that: for any collection $\{A_1, \ldots, A_N\} \subset \mathcal{M}_c$ of cardinality N, there is a subcollection $\{A_{i_1}, \ldots, A_{i_n}\}$ of cardinality n such that $\sigma(A_{i_1} \cap \cdots \cap A_{i_n}) > 0$ and hence $A_{i_1} \cap \cdots \cap A_{i_n} \neq \emptyset$.

From (2.3), (2.4), we see that $\{D_r\}_{r\in I}$ is a family of measurable subsets of the finite measure space $\overline{B}(0, R) \times O_d(\mathbb{R})$, equipped with $\mathcal{L}^d \times d\theta$, where

$$R = 2(\sup\{r : r \in I\}) \cdot (\sup\{|x| : x \in E\}),$$

with measures bounded below by c_I . Applying Proposition 2.3, there exists an $M \in \mathbb{N}$ such that, for any distinct $\{r_i\}_{i=1}^M \subset I$, there is a subset $\{r_{ij}\}_{j=1}^n$ such that

$$\left(\mathcal{L}^d \times d\theta\right) \left(\bigcap_{j=1}^n D_{r_j}\right) > 0.$$

Thus, there exists a $\theta_0 \in O_d(\mathbb{R})$ such that

$$\mathcal{L}^d\left(\bigcap_{j=1}^n D_{r_j} \cap \{\theta = \theta_0\}\right) > 0.$$

Since $\bigcap_{j=1}^{n} D_{r_j} \cap \{\theta = \theta_0\}$ has positive Lebesgue measure, it is infinite and hence for any $k \in \mathbb{N}$ we can find $\{x^l\}_{l=1}^{k+1}$ and $\{y^l\}_{l=1}^{k+1}$ such that

$$r_{i_j}x^l = \theta_0 y^l - a, \quad \forall 1 \le j \le n, \ 1 \le l \le k+1.$$

This finishes the proof of Theorem 1.3. Note that what was established is considerably stronger than the statement of the theorem.

3. Proof of Theorem 1.4

Since $T : \mathbb{R}^d \to \mathbb{R}^d$ is proper and continuous, the Frostman measure μ on E defines a measure μ_T by

$$\int f(x)d\mu_T(x) := \int f(T(x))d\mu(x), \quad \forall f \in C_0(\mathbb{R}^d).$$

Suppose that

$$\int |\widehat{\mu}_T(\xi)|^2 |\xi|^{-s\frac{d-1}{d}} d\xi < \infty$$

Arguing as in the proof of Theorem 1.1, if we set

$$A_{T,\theta} := \operatorname{supp} \left(\mu_T * \mu^{\theta} \right) \subset \mathbb{R}^d,$$

one sees that there exists a $c_T > 0$ such that

$$\sup_{\theta \in O_d(\mathbb{R})} \mathcal{L}^d \left(A_{T,\theta} \right) \ge c_T,$$

from which it follows that there exists a $\theta_0 \in O_d(\mathbb{R})$ such that

$$\mathcal{L}^{d}\left(\{a \in \mathbb{R}^{d} : \mu\{y \in E : (\exists x \in E) (T(x) = \theta_{0}y - a)\} > 0\}\right) \ge c_{T} > 0.$$

It follows that for all $k \in \mathbb{N}$, there are points $a \in \mathbb{R}^d$ and $x_1, \ldots, x_{k+1}, y_1, \ldots, y_{k+1} \in E$ and such that $T(x_i) = \theta_0 y_i - a$ for all $1 \le i \le k+1$. Remark 3.1. Similarly to how one can go from a single dilation factor in Theorem 1.1 to the multi-similarities in Theorem 1.3, one can boost from the one transformation T in Theorem 1.4 to multiple transformations. Suppose that \mathcal{T} is a family of proper, continuous maps and s is such that there is a uniform bound,

$$\int |\widehat{\mu}_T(\xi)|^2 |\xi|^{-s\frac{d-1}{d}} d\xi \le C, \quad \forall \, 1 \le j \le m, \, T \in \mathcal{T}.$$

Following the proof of Theorem 1.3 in Section 2.2, one can show that for all $n \in \mathbb{N}$, there exists an $M \in \mathbb{N}$ such that for any distinct $\{T_j\}_{j=1}^M \subset \mathcal{T}$, there exists a subset $\{T_{j_l}\}_{l=1}^n$, a $\theta_0 \in O_d(\mathbb{R})$, $a \in \mathbb{R}^d$, and for all $k \in \mathbb{N}$, points $x_1, \ldots, x_{k+1}, y_1, \ldots, y_{k+1} \in E$ such that $T_{j_l}(x_i) = \theta_0 y_i - a$ for all $1 \leq i \leq k+1$, $1 \leq l \leq n$.

4. PROOFS OF THEOREM 1.6, THEOREM 1.5 AND COROLLARY 1.7

We start with the proof of Theorem 1.6, and then show how to apply it to obtain the similarity set results Theorem 1.5 and Corollary 1.7.

Proof of Theorem 1.6. Suppose that G is a finite group acting transitively on a set X, and let H and E be subsets of X. For each $x \in X$, let $\mathcal{O}_x := \{gx : g \in G\}$ be the orbit of x, and $\mathcal{S}_x := \{g \in G : gx = x\}$ be the stabilizer of x. The orbit-stabilizer theorem tells us that $|G| = |\mathcal{O}_x| \cdot |\mathcal{S}_x|$. Since the action of G on X is transitive, then $\mathcal{O}_x = X$, and hence $|\mathcal{S}_x| = |G|/|X|$.

Let $S_{xy} \coloneqq \{g \in G : gx = y\}$. It is not difficult to prove that $|S_{xy}| = |S_x|$. Indeed, consider the map $\phi : S_{xy} \times S_x \to S_{xy}$ defined by $(h,g) \stackrel{\phi}{\mapsto} hg$. We observe that every $u \in S_{xy}$ has exactly $|S_{xy}|$ preimages, i.e., $|\phi^{-1}(\{u\})| = |S_{xy}|$. This immediately implies that $|S_{xy} \times S_x| = |S_{xy}|^2$, and hence $|S_{xy}| = |S_x|$. Therefore, we have shown that $|S_{xy}| = |G|/|X|$ for any $x, y \in X$.

Now, define a set \mathcal{P} by

$$\mathcal{P} = \{ (g, y) \in G \times X : y \in H \cap gE \}.$$

We can compute the cardinality of \mathcal{P} in two ways using a double-counting argument. On one hand, we have:

(4.1)

$$\begin{aligned} |\mathcal{P}| &= \sum_{y \in H} \sum_{\substack{g \in G \\ y \in gE}} 1 = \sum_{y \in H} |\{g \in G : y \in gE\}| \\ &= \sum_{y \in H} \sum_{x \in E} |\{g \in G : y = gx\}| = \sum_{y \in H} \sum_{x \in E} |\mathcal{S}_{xy}| \\ &= \frac{|G||H||E|}{|X|}, \end{aligned}$$

where we have used the fact that $|\mathcal{S}_{xy}| = |G|/|X|$. On the other hand,

(4.2)
$$\begin{aligned} |\mathcal{P}| &= \sum_{g \in G} \sum_{y \in H \cap gE} 1 = \sum_{g \in G} |H \cap gE| \\ &\leq \max_{g \in G} |H \cap gE| \cdot |G|. \end{aligned}$$

Combining (4.1) and (4.2), one finds that

(4.3)
$$\max_{g \in G} |H \cap gE| \ge \frac{|H||E|}{|X|},$$

as claimed in (1.4). This completes the proof of Theorem 1.6.

Proof of Theorem 1.5. Consider $X = \mathbb{F}_q^d$ and $E \subset \mathbb{F}_q^d$. For $r \in (\mathbb{F}_q)^2 \setminus \{0\}$, let \sqrt{r} be any square root of r, and set $H = \sqrt{rE}$. Note that |H| = |E|. Finally, let G be the group of translations of \mathbb{F}_q^d , acting transitively on \mathbb{F}_q^d . Theorem 1.6 implies that

(4.4)
$$\max_{a \in \mathbb{F}_q^d} \left| \left(\sqrt{r} E \right) \cap (E+a) \right| \ge \frac{|E|^2}{q^d}.$$

If $|E| \ge \sqrt{k+1}q^{\frac{d}{2}}$, then $\max_{a \in \mathbb{F}_q^d} |\sqrt{rE} \cap (E+a)| \ge k+1$. Thus, there exists an element $a \in \mathbb{F}_q^d$ such that $|\sqrt{rE} \cap (E+a)| \ge k+1$. Consequently, we can establish the existence of a sequence $\{z_1, \ldots, z_{k+1}\}$ such that $\{z_1, \ldots, z_{k+1}\} \subset \sqrt{rE} \cap (E+a)$. This implies the existence of sequences $\{x_1, \ldots, x_{k+1}\} \subset E$ and $\{y_1, \ldots, y_{k+1}\} \subset E$, such that $z_i = \sqrt{rx_i}$ and $z_i = y_i + a$ for $1 \le i \le k+1$.

In summary, we have demonstrated the existence of (k+1)-tuples $(x_1, \ldots, x_{k+1}) \in E^{k+1}$ and $(y_1, \ldots, y_{k+1}) \in E^{k+1}$ satisfying the following conditions:

- (1) $x_i \neq x_j$ and $y_i \neq y_j$ for $1 \le i < j \le k+1$.
- (2) $y_i + a = \sqrt{r}x_i$ for $i \in \{1, \dots, k+1\}$.

Therefore, for $1 \leq i < j \leq k+1$, we have:

$$||y_i - y_j|| = ||(\sqrt{r}x_i - a) - (\sqrt{r}x_j - a)|| = ||\sqrt{r}x_i - \sqrt{r}x_j||$$

= $(\sqrt{r})^2 ||x_i - x_j|| = r ||x_i - x_j||.$

Since A is a nonempty subset of $\{(i, j) : 1 \le i < j \le k+1\}$, we have shown that there exist two (k+1)-point configurations, $(x_1, \ldots, x_{k+1}) \in E^{k+1}$ and $(y_1, \ldots, y_{k+1}) \in E^{k+1}$, satisfying the following conditions:

(1) $x_i \neq x_j$ and $y_i \neq y_j$ for $1 \le i < j \le k+1$.

(2)
$$||y_i - y_j|| = r ||x_i - x_j||$$
 for $(i, j) \in A$.

This completes the proof of Theorem 1.5.

Proof of Corollary 1.7. Let $G = \operatorname{SL}_d(\mathbb{F}_q)$, $X = \mathbb{F}_q^d$, and suppose that $E \subset \mathbb{F}_q^d$ with $|E| \geq \sqrt{k+1}q^{\frac{d}{2}}$. The action of $\operatorname{SL}_d(\mathbb{F}_q)$ on \mathbb{F}_q^d is transitive for $d \geq 2$ (this is just a special case of the general fact, valid for any field \mathbb{F}). Consequently, Theorem 1.6 is applicable, and we now show that it leads to the conclusion stated in Corollary 1.7.

If we let $r^{\frac{1}{d}}$ be any d^{th} root of $r \in (\mathbb{F}_q)^d \setminus \{0\}$, define $H = r^{\frac{1}{d}}E$. Then Theorem 1.6 yields $\max_{g \in G} |r^{\frac{1}{d}}E \cap gE| \ge k+1$. Let $g \in \text{SL}_d(\mathbb{F}_q)$ be such that $|r^{\frac{1}{d}}E \cap gE| \ge k+1$, so that there is a set

$$\{z_1,\ldots,z_{k+1}\}\subset r^{\frac{1}{d}}E\cap gE.$$

This implies the existence of $\{x_1, \ldots, x_{k+1}\} \subset E$ and $\{y_1, \ldots, y_{k+1}\} \subset E$, such that $z_i = r^{\frac{1}{d}}y_i$ and $z_i = gx_i$ for $1 \leq i \leq k+1$.

Hence,

$$\det(z_{i_1},\ldots,z_{i_d}) = \det(r^{\frac{1}{d}}y_{i_1},\ldots,r^{\frac{1}{d}}y_{i_d}) = r\det(y_{i_1},\ldots,y_{i_d}).$$

On the other hand, since $g \in SL_d$,

$$\det(z_{i_1},\ldots,z_{i_d}) = \det(gx_{i_1},\ldots,gx_{i_d}) = \det(x_{i_1},\ldots,x_{i_d})$$

Comparing the previous two equalities yields the statement of Corollary 1.7.

Remark. Let $X = S^{d-1}$ be a *d*-dimensional sphere in \mathbb{F}_q^d . Then $G = O_d(\mathbb{F}_q)$ acts transitively on S^{d-1} . By Theorem 1.6 we have that if $E \neq H$ and $|E||H| \geq (k+1)q^{d-1}$, then $\max_{g \in O_d} |H \cap gE| \geq k+1$.

References

^[1] P. Bhowmik and F. Rakhmonov, Near optimal thresholds for existence of dilated configurations in \mathbb{F}_a^d , Bulletin of the Australian Mathematical Society. pp. 1-12 (2024) 1, 4, 5

^[2] X. Du, Y. Ou, K. Ren, and R. Zhang, New improvement to Falconer distance set problem in higher dimensions, (arXiv:2309.04501) (2023). 1

- [3] X. Du and R. Zhang, Sharp L² estimate of Schrödinger maximal function in higher dimensions, Ann. of Math. (2) 189 (2019), no. 3, 837-861.
- [4] K. J. Falconer, On the Hausdorff dimensions of distance sets, Mathematika 32 (1986), 206-212.
 1
- [5] A. Greenleaf, A. Iosevich and S. Mkrtchyan, Existence of similar point configurations in thin subsets of ℝ^d, Math. Z. 297 (2021), no. 1-2, 855-865. 1, 2, 3, 7, 8
- [6] L. Guth, A. Iosevich, Y. Ou and H. Wang, On Falconer's distance problem in the plane, Invent. Math. 219 (2020), no. 3, 779-830. 1
- [7] P. Mattila, Geometry of sets and measures in Euclidean spaces, Cambridge Studies in Adv. Math. 44. Cambridge Univ. Press, 1995. 1, 2
- [8] F. Rakhmonov, Distribution of similar configurations in subsets of \mathbb{F}_q^d , Discrete Math. **346**(10) (2023), Paper No. 113571, 21 pages. 5
- T. Wolff, Decay of circular means of Fourier transforms of measures, International Mathematics Research Notices 10 (1999) 547-567.
 - DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY *Email address*: pbhowmik@ur.rochester.edu
 - DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY *Email address*: allan@math.rochester.edu
 - DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY *Email address*: iosevich@math.rochester.edu
 - DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY *Email address*: sevak.mkrtchyan@rochester.edu
 - DEPARTMENT OF MATHEMATICS, UNIVERSITY OF ROCHESTER, ROCHESTER, NY *Email address:* frakhmon@ur.rochester.edu