Moore Determinant of Dual Quaternion Hermitian Matrices

Chunfeng Cui^{*} Liqur

Liqun Qi^{\dagger}

Guangjing Song ‡ and

and Qingwen Wang[§]

May 21, 2024

Abstract

In this paper, we extend the Chen and Moore determinants of quaternion Hermitian matrices to dual quaternion Hermitian matrices. We show the Chen determinant of dual quaternion Hermitian matrices is invariant under addition, switching, multiplication, and unitary operations at the both hand sides. We then show the Chen and Moore determinants of dual quaternion Hermitian matrices are equal to each other, and they are also equal to the products of eigenvalues. The characteristic polynomial of a dual quaternion Hermitian matrix is also studied.

Key words. Moore determinant, dual quaternion, eigenvalue.

MSC subject classifications. 15A18, 15B33, 20G20, 65F40.

1 Introduction

Quaternions are extensions of complex numbers, introduced by the Irish mathematician Hamilton in 1843. A quaternion has three imaginary parts and the quaternion multiplication is noncommutative. Consequently, the determinant

^{*}LMIB of the Ministry of Education, School of Mathematical Sciences, Beihang University, Beijing 100191, China. (chunfengcui@buaa.edu.cn).

[†]Department of Applied Mathematics, The Hong Kong Polytechnic University, Hung Hom, Kowloon, Hong Kong; Department of Mathematics, School of Science, Hangzhou Dianzi University, Hangzhou 310018, China. (maqilq@polyu.edu.hk).

[‡]School of Mathematics and Information Sciences, Weifang University, Weifang 261061, China. (sgjshu@163.com).

[§]Department of Mathematics, Shanghai University, Shanghai 200444, China. (wqw@shu.edu.cn).

of quaternion matrices is much more difficult than their real and complex counterparts since the quaternion numbers. Over the years, many determinants of quaternion matrices with different multiplication orders are proposed, such as the Cayley determinant, the Study determinant, the Moore determinant [14], the determinant and double determinant by Chen [3, 4], the column and row determinants by Kyrchei [7, 8], etc. See [1, 6, 10, 13] for more details.

In 1922, Eliakim Hastings Moore defined the determinant of quaternion Hermitian matrices [14], which is widely known as the Moore determinant. The Moore determinant retains some basic properties of determinants such as Mdet(A) = 0 if and only if A is singular. For more results on the Moore determinant, please refer to [1, 8]. However, Moore determinant is not appliable for arbitrary non-Hermitian quaternion matrices. In 1991, Chen proposed the determinant and double determinant [3, 4]. In 2008 and 2012, Kyrchei proposed the column and row determinants [7, 8].

It was the British mathematician Clifford who introduced dual numbers and dual quaternions in 1873. A **dual quaternion** has two quaternions, which are the standard and dual parts of this dual quaternion, respectively. Recently, eigenvalues of dual quaternion Hermitian matrices were applied to dual unit gain graphs [5] and multi-agent formation control [15]. The latter problem is an important application area in robotics and control [17, 20]. To study more about the eigenvalues of dual quaternion Hermitian matrices, we need to determine the coefficients of the characteristic polynomial of a dual quaternion Hermitian matrix. One possible way to achieve this is to apply the Moore determinant to dual quaternion Hermitian matrices. In this paper, we study the Moore determinant of dual quaternion Hermitian matrices.

In the next section, we review some preliminary knowledge on dual quaternion numbers, the Moore determinant and its extension, the Chen determinant. In Section 3, we show that the Chen determinant of a dual quaternion Hermitian matrix is invariant under the addition, switching, multiplication, and unitary operations at the both hand sides, which is also equal to the product of eigenvalues of that matrix. Then in Section 4, we show that the Moore determinant and the Chen determinant are equal for a dual quaternion Hermitian matrix. Thus, the Moore determinant of a dual quaternion Hermitian matrix is also equal to the product of eigenvalues of that matrix. Finally, we study the characteristic polynomial of a dual quaternion Hermitian matrix in Section 5.

2 Preliminary

2.1 Quaternions and dual quaternions

A quaternion number $q \in \mathbb{H}$ can be written as $q = q_0 + q_1\mathbf{i} + q_2\mathbf{j} + q_3\mathbf{k}$. Here, $\mathbf{i}, \mathbf{j}, \mathbf{k}$ are imaginary units that satisfy $\mathbf{i}^2 = \mathbf{j}^2 = \mathbf{k}^2 = \mathbf{i}\mathbf{j}\mathbf{k} = -1$ and $\mathbf{i}\mathbf{j} = -\mathbf{j}\mathbf{i} = \mathbf{k}$. Thus, the multiplication of quaternion numbers is not commutative. The conjugate of q is $q^* = q_0 - q_1\mathbf{i} - q_2\mathbf{j} - q_3\mathbf{k}$. We have $|q| = \sqrt{q_0^2 + q_1^2 + q_2^2 + q_3^2}$.

A dual quaternion number $q \in \hat{\mathbb{H}}$ can be written as $q = q_s + q_d \epsilon$, where $q_s, q_d \in \mathbb{H}, q_s$ is the standard part of q, q_d is the dual part of q, ϵ is the infinitesimal unit, $\epsilon \neq 0, \epsilon^2 = 0, \epsilon$ is commutative with quaternion numbers. If $q_s \neq 0$, then q is called **appreciable**. The conjugate of q is $q^* = q_s^* + q_d^* \epsilon$. We have

$$|p| := \begin{cases} |p_s| + \frac{(p_s^* p_d + p_d^* p_s)}{2|p_s|} \epsilon, & \text{if } p_s \neq 0, \\ |p_d|\epsilon, & \text{otherwise.} \end{cases}$$
(1)

If both q_s and q_d are real numbers, then $q = q_s + q_d \epsilon$ is called a **dual number**.

Proposition 2.1. Let q_1 and $q_2 \in \hat{\mathbb{H}}$ be two dual quaternion numbers. Then

- (i) $Re(q_1) \leq |q_1|$ and the equality holds if and only if q_1 is a nonnegative dual number.
- (i) $Re(q_1) = Re(q_1^*)$ and $Re(q_1q_2) = Re(q_2q_1)$.

(*ii*)
$$q_1 + q_1^* \in \mathbb{R}$$
.

- (*iii*) $|q_1q_2| = |q_1||q_2|$.
- (*iv*) $|q_1 + q_2| \le |q_1| + |q_2|$.

A dual quaternion vector $\mathbf{x} \in \hat{\mathbb{H}}^n$ can be denoted as $\mathbf{x} = \mathbf{x}_s + \mathbf{x}_d \epsilon$. If $\mathbf{x}_s \neq \mathbf{0}$, we say \mathbf{x} is appreciable. Otherwise, it is infinitesimal. If \mathbf{x} is appreciable, then the 2-norm of \mathbf{x} is defined by $\sqrt{\sum_{i=1}^n |x_i|^2}$. Otherwise, if \mathbf{x} is infinitesimal, then $\|\mathbf{x}\| = \|\mathbf{x}_d\| \epsilon$.

The collection of $m \times n$ dual quaternion matrices is denoted by $\hat{\mathbb{H}}^{m \times n}$. A dual quaternion matrix $A = (a_{ij}) \in \hat{\mathbb{H}}^{m \times n}$ can be denoted as $A = A_s + A_d \epsilon$, where $A_s, A_d \in \mathbb{H}^{m \times n}$. The conjugate transpose of A is $A^* = (a_{ji}^*)$. Let $A \in \hat{\mathbb{H}}^{m \times n}$ and $B \in \hat{\mathbb{H}}^{n \times r}$. Then we have $(AB)^* = B^*A^*$.

Given a square dual quaternion matrix $A \in \hat{\mathbb{H}}^{n \times n}$, it is called invertible (nonsingular) if $AB = BA = I_n$ for some $B \in \hat{\mathbb{H}}^{n \times n}$, where I_n is the $n \times n$ identity matrix. Such B is unique and denoted by A^{-1} . Square dual quaternion matrix A is called Hermitian if $A^* = A$. Then A is Hermitian if and only if both A_s and A_d are quaternion Hermitian matrices. Square dual quaternion matrix A is called unitary if $A^*A = I_n$. Apparently, $A \in \hat{\mathbb{H}}^{n \times n}$ is unitary if and only if its column vectors form an orthonormal basis of $\hat{\mathbb{H}}^n$.

An $n \times n$ dual quaternion Hermitian matrix has exactly n right eigenvalues, which are all dual numbers and also left eigenvalues [16]. We simply call them the eigenvalues of that matrix.

2.2 Quaternion determinants

A **permutation** $\sigma = \{i_1, \ldots, i_n\} \in S_n$ denotes a function σ such that $\sigma(j) = i_j$ for all $j = 1, \ldots, n$. We could also rewrite it as a two-line formulation

$$\begin{pmatrix} 1 & 2 & \cdots & n-1 & n \\ i_1 & i_2 & \cdots & i_{n-1} & i_n \end{pmatrix}$$

A **permutation cycle** $\sigma = (i_1, \ldots, i_k)$ is a subset of a permutation whose elements trade places with one another. Namely, $\sigma(i_j) = i_{j+1}$ for $j = 1, \ldots, k-1$ and $\sigma(i_k) = i_1$. A cycle of length k is called a k-cycle. We could also rewrite the permutation cycle as a two-line formulation

$$\begin{pmatrix} i_1 & i_2 & \cdots & i_{k-1} & i_k \\ i_2 & i_3 & \cdots & i_k & i_1 \end{pmatrix}.$$
 (2)

For instance, the permutation cycle $\sigma = (312)$ represents a permutation $\sigma(3) = 1$, $\sigma(1) = 2$, and $\sigma(2) = 3$. The sign of any k-cycle is $(-1)^{k-1}$. In fact, by implementing k - 1 switches, i.e., i_1 with i_k , i_1 with i_{k-1} , ..., i_1 with i_2 , sequentially, (2) reduces to

$$\begin{pmatrix} i_1 & i_2 & \cdots & i_{k-1} & i_k \\ i_1 & i_2 & \cdots & i_{k-1} & i_k \end{pmatrix},$$

which is corresponding to $\{1, \ldots, k\}$.

The **cyclic decomposition** of a permutation is to decompose the permutation as a product of disjoint cycles,

$$\sigma = (n_{11} \cdots n_{1l_1})(n_{21} \cdots n_{2l_2}) \cdots (n_{r1} \cdots n_{rl_r}).$$
(3)

Every permutation can be written as a product of disjoint cycles. For instance, $\{1, 2, 3\} = (1)(2)(3)$ and $\{2, 1, 3\} = (12)(3)$. The **inverse permutation** σ^{-1} is the inverse of σ such that $\sigma(\sigma^{-1}(i)) = i$ and $\sigma^{-1}(\sigma(i)) = i$ for $i = 1, \ldots, n$. In the cycle form, the inverse permutation just reverses the direction of each cycle. Namely, $\sigma^{-1} = (i_1, i_n, i_{n-1}, \ldots, i_2)$. For more details on the permutation, we refer to [2].

Let $A = (a_{ij})$ be a quaternion Hermitian matrix in $\mathbb{H}^{n \times n}$ and σ be a permutation of $S_n = \{1, \ldots, n\}$. In 1922, Eliakim Hastings Moore defined the **Moore determinant** for quaternion matrices [14] as follows,

$$\operatorname{Mdet}(A) = \sum_{\sigma \in S_n^M} s(\sigma) a_{\sigma}, \tag{4}$$

where the permutation cycle σ is a product of disjoint cycles as in (3) and satisfies

 $n_{i1} < n_{ij}$ for all $j > 1, i = 1, \dots, r$, and $n_{11} > n_{21} > \dots > n_{r1}$, (5)

 $s(\sigma)$ denotes the sign of σ , and

$$a_{\sigma} = (a_{n_{11},n_{12}}a_{n_{12},n_{13}}\cdots a_{n_{1l_1},n_{11}})(a_{n_{21},n_{22}}\cdots a_{n_{2l_2},n_{21}})\cdots(\cdots a_{n_{rl_r},n_{r1}}).$$

Denote S_n^M as the set of permutations satisfying (5). When the matrix is a complex or real matrix, the Moore determinant (4) reduces to the ordinary

determinant. Following [3], we also denote $\sigma_i = (n_{i1} \cdots n_{il_i})$,

$$\langle \sigma_i \rangle = \langle n_{i1} \cdots n_{il_i} n_{i1} \rangle = a_{n_{i1}, n_{i2}} a_{n_{i2}, n_{i3}} \cdots a_{n_{il_i}, n_{i1}}.$$

Thus, $\sigma = \sigma_1 \cdots \sigma_r$ and $\langle \sigma \rangle = \langle \sigma_1 \rangle \cdots \langle \sigma_r \rangle$.

Let $\mathbf{a}_{.j}$ be the *j*th column and \mathbf{a}_{i} . be the *i*th row of a matrix $A \in \mathbb{H}^{n \times n}$ or $A \in \hat{\mathbb{H}}^{n \times n}$. Let $A_{.j}(\mathbf{a})$ be a matrix obtained from replacing the *j*th column of A by the column vector \mathbf{a} , and A_{i} .(b) be a matrix result from replacing the *i*th row of A by the row vector \mathbf{b} . Denote by A^{ij} a submatrix of A obtained by deleting both the *i*th row and the *j*th column. Here, $A_{.j}(\mathbf{a})$ and A_{i} .(b) may not be Hermitian matrices even if A is Hermitian. Such matrices are **almost** Hermitian [6]. We say a matrix is *k*-almost Hermitian if it is self-adjoint except for the *k*-th row or column.

In 1972, Dyson [6] presented an equivalent formulation of the Moore determinant for k-almost Hermitian matrices as follows,

$$\operatorname{Mdet}(A) = a_{kk}\operatorname{Mdet}(A^{kk}) - \sum_{i=1, i \neq k}^{n} a_{ki}\operatorname{Mdet}(A^{kk}_{\cdot i}(a_{\cdot k})).$$
(6)

Here, k is the index in the k-almost Hermitian. If A is Hermitian, then the above value remains the same for all k = 1, ..., n. This formulation is quite useful for the numerical computation.

However, the Moore determinant is only applicable to Hermitian or almost Hermitian matrices. Several researchers have considered determinants of arbitrary quaternion matrices. In 1922, Study [18] proposed determinants of quaternion matrices through their injective algebra homomorphism with complex matrices. Specifically, let $A = A_1 + A_2 \mathbf{j}$. Then

$$\operatorname{Sdet}(A) = \operatorname{det}\left(\begin{bmatrix} A_1 & -\bar{A}_2 \\ A_2 & \bar{A}_1 \end{bmatrix} \right).$$

In 1991, Chen [3, 4] proposed a novel definition of quaternion determinant as follows

$$\operatorname{Cdet}(A) = \sum_{\sigma \in S_n^C} (-1)^{n-r} a_{\sigma}, \tag{7}$$

where the permutation cycle σ is a product of disjoint cycles as in (3) and satisfies

$$n_{i1} > n_{ij}$$
 for all $j > 1, i = 1, \dots, r$, and $n = n_{11} > n_{21} > \dots > n_{r1}$, (8)

and S_n^C denotes the set of permutations satisfying (8). In the following, we call it the **Chen determinant** and denote it by Cdet. In (7), *n* is fixed as the first element in the cycle σ .

In 2008 and also in 2012, Kyrchei [7, 8] generalized the first element in the Chen determinant to any index $i \in \{1, ..., n\}$ and proposed the column and row determinants of arbitrary quaternion matrices as follows

$$\operatorname{Krdet}_{i}(A) = \sum_{\sigma \in S_{n}^{K}} (-1)^{n-r} a_{\sigma}, \qquad (9)$$

where σ is the permutations of $\{1, \ldots, n\}$ satisfying

$$\sigma = (ii_{k_1}i_{k_1+1}\cdots i_{k_1+l_1})(i_{k_2}i_{k_2+1}\cdots i_{k_2+l_2})\cdots (i_{k_r}i_{k_r+1}\cdots i_{k_r+l_r}),$$

$$i_{k_2} < i_{k_3} < \cdots < i_{k_r} \text{ and } i_{k_t} < i_{k_{t+s}} \text{ for all } t = 2, \dots, r, s = 1, \dots, l_t,$$
(10)

and S_n^K denotes the set of permutations satisfying (12). In the following, we call it the **Kyrchei row determinant**.

Similarly, the Kyrchei column determinant is defined by

$$\operatorname{Kcdet}_{j}(A) = \sum_{\sigma \in \bar{S}_{n}^{K}} (-1)^{n-r} a_{\sigma}, \qquad (11)$$

where σ is the permutations of $\{1, \ldots, n\}$ satisfying

$$\sigma = (j_{k_r} j_{k_r+l_r} \cdots j_{k_r+1} j_{k_r}) \cdots (j_{k_2+l_2} \cdots j_{k_2+1} j_{k_2}) (j_{k_1+l_1} \cdots j_{k_1+1} j_{k_1} j),$$
(12)

$$j_{k_2} < j_{k_3} < \cdots < j_{k_r}$$
 and $j_{k_t} < j_{k_t+s}$ for all $t = 2, \ldots, r, s = 1, \ldots, l_t$.

When A is a quaternion Hermitian matrix, all Kyrchei row and column determinants are real numbers and are the same, which also coincide with the Moore determinant. Furthermore, the Kyrchei column and row determinants may derive the formulation of the inverse matrix by the classical adjoint matrix and could be connected to the solution of linear systems. We now review the elementary row and column operations for quaternion matrices. Consider the three elementary row and column operations that can be used to transform a matrix (square or not) into a simple form and can facilitate determinant calculations. In the following definition, we focus on row operations, which are implemented by left multiplying the matrices. Column operations can be defined and used in a similar fashion, while the matrices that implement them act on the right.

Definition 2.2 (Elementary row and column operations). Given a quaternion matrix $A \in \mathbb{H}^{n \times n}$ and $i, j \in \{1, ..., n\}$. Consider the following three elementary transformations.

(i) Switching of two rows. Let $P_{ij} = (a_{kl}) \in \mathbb{R}^{n \times n}$ satisfy

$$a_{kl} = \begin{cases} 1, & \text{if } k = l \neq i \text{ and } k = l \neq j; \\ 1, & \text{if } k = i \text{ and } l = j; \\ 1, & \text{if } k = j \text{ and } l = i; \\ 0, & \text{otherwise.} \end{cases}$$

Then by multiplying P_{ij} on the left, we switch the *i*-th and the *j*-th rows of A. We refer P_{ij} as a switching matrix.

(ii) Multiplication of a row by a scalar. Let $c \in \mathbb{H}$ be any quaternion number and $P_{i;c} = (a_{kl}) \in \mathbb{H}^{n \times n}$ satisfy

$$a_{kl} = \begin{cases} 1, & \text{if } k = l \neq i; \\ c, & \text{if } k = l = i; \\ 0, & \text{otherwise.} \end{cases}$$

Then $P_{i;c}A$ multiplies the *i*-th row of A by c on the left. We refer $P_{i;c}$ as a multiplication matrix.

(iii) Addition of a scalar multiple of one row to another row. Let $c \in \mathbb{H}$ be any quaternion number and $P_{ij;c} = (a_{kl}) \in \mathbb{H}^{n \times n}$ satisfy

$$a_{kl} = \begin{cases} 1, & \text{if } k = l; \\ c, & \text{if } k = j \text{ and } l = i; \\ 0, & \text{otherwise.} \end{cases}$$

Then $P_{ij;c}A$ adds the *j*-th row of A by the *i*-th row of A multiplied with c one the left. We refer $P_{ij;c}$ as an addition matrix.

For more properties and applications of quaternion determinants, people may check [9, 10, 12, 13].

We note that the Moore determinant, the Chen determinant, the Kyrchei determinant, and Definition 2.2 can be extended to dual quaternions without any difficulty. However, the study of properties of dual quaternion determinants may be different. Very recently, Ling and Qi [11] studied the determinant properties of dual complex matrices, and then introduced the concept of quasi-determinant of dual quaternion matrices. Based upon these, they showed the quasi-determinant of a dual quaternion Hermitian matrix is equivalent to the product of the square of the magnitudes of all eigenvalues. In this paper, we focus on the determinants of dual quaternion Hermitian matrix that are equal to the product of eigenvalues, such as the Chen determinant and the Moore determinant.

Next, we present several properties of the Chen determinant for quaternion Hermitian matrices [3, 4].

Proposition 2.3. Let $A = (a_{ij})$ be a quaternion Hermitian matrix in $\mathbb{H}^{n \times n}$, P_{ij} be a switching matrix, $P_{i;\alpha}$ be a multiplication matrix, and $P_{ij;\alpha}$ be an addition matrix. Then the following results hold.

- (i) $\operatorname{Cdet}(P_{ij}^*AP_{ij}) = \operatorname{Cdet}(A);$
- (*ii*) $\operatorname{Cdet}(P_{n;\alpha}A) = \alpha \operatorname{Cdet}(A), \operatorname{Cdet}(AP_{n;\alpha}) = \alpha \operatorname{Cdet}(A), and \operatorname{Cdet}(P_{i;\alpha}^*AP_{i;\alpha}) = \alpha^* \alpha \operatorname{Cdet}(A);$

(*iii*)
$$\operatorname{Cdet}(P_{ij;\alpha}^*AP_{ij;\alpha}) = \operatorname{Cdet}(A);$$

(iv) For any unitary matrix $U \in \mathbb{H}^{n \times n}$, we have $\operatorname{Cdet}(U^*AU) = \operatorname{Cdet}(A)$;

Recall that an $n \times n$ quaternion Hermitian matrix has exactly n right eigenvalues, which are all real numbers and also left eigenvalues [21]. We simply call them eigenvalues of that matrix. Recently, Qi and Luo [16] showed an $n \times n$

dual quaternion Hermitian matrix has exactly n right eigenvalues, which are all dual real numbers and also left eigenvalues. As mentioned early, they are also simply called eigenvalues. In what follows, we show the Chen determinant of dual quaternion Hermitian matrices is equal to the product of these neigenvalues.

3 Properties of the Dual Quaternion Chen Determinant

In this section, we show that the results in Proposition 2.3 can be generalized to dual quaternion Hermitian matrices. We begin with the following lemma.

Lemma 3.1. Let $A = (a_{ij})$ be a dual quaternion Hermitian matrix in $\hat{\mathbb{H}}^{n \times n}$. Suppose $\sigma_0 = (p_1 p_2 \cdots p_s k q_1 \cdots q_t)$ is a cycle factor of the permutation $\sigma \in S_n$. Denote

$$\bar{\sigma}_0 = (p_1 q_t \cdots q_1 k p_s \cdots p_2),$$

$$\sigma_0^+ = (k q_1 \cdots q_t p_1 p_2 \cdots p_s),$$

$$\bar{\sigma}_0^+ = (k p_s \cdots p_2 p_1 q_t \cdots q_1).$$

Then we have

$$\langle \sigma_0 \rangle + \langle \bar{\sigma}_0 \rangle = \langle \sigma_0^+ \rangle + \langle \bar{\sigma}_0^+ \rangle. \tag{13}$$

Proof. By direct computation, we have $\langle \sigma_0 \rangle = a_{p_1 p_2} \cdots a_{p_s k} a_{kq_1} \cdots a_{q_t p_1}$ and $\langle \bar{\sigma}_0 \rangle = a_{p_1 q_t} \cdots a_{q_1 k} a_{kp_s} \cdots a_{p_2 p_1}$. Therefore, we have $\langle \sigma_0 \rangle = \langle \bar{\sigma}_0 \rangle^*$. Similarly, we have $\langle \sigma_0^+ \rangle = \langle \bar{\sigma}_0^+ \rangle^*$. Thus, both the right and left hand sides of (13) are dual numbers.

Furthermore, let $w_1 = a_{p_1p_2} \cdots a_{p_sk}$, $w_2 = a_{kq_1} \cdots a_{q_tp_1}$. By Proposition 2.1, we have $Re(\langle \sigma_0 \rangle) = Re(w_1w_2) = Re(w_2w_1) = Re(\langle \sigma_0^+ \rangle)$. Similarly, there is $Re(\langle \bar{\sigma}_0 \rangle) = Re(\langle \bar{\sigma}_0^+ \rangle)$. Thus, the real part of the right hand side of (13) is equal to that of the left hand side. Combining with the fact that both the right and left hand sides of (13) are dual numbers, we derive (13).

This completes the proof.

Lemma 3.2. Let $A = (a_{ij})$ be a dual quaternion Hermitian matrix in $\hat{\mathbb{H}}^{n \times n}$, and P_{ij} be a switching matrix. Then $\operatorname{Cdet}(P_{ij}^*AP_{ij}) = \operatorname{Cdet}(A)$. Proof. If i, j, n are mutually distinct, then we have $P_{ij} = P_{jn}P_{in}P_{jn}$. Hence, it suffices to show $\operatorname{Cdet}(P_{jn}^*AP_{jn}) = \operatorname{Cdet}(A)$ for $j \neq n$. For any $\sigma = \sigma_1 \cdots \sigma_r \in S_n^C$, denote the apostrophe of σ

$$\sigma' = \sigma|_{n \longleftrightarrow j}$$

as the permutation of S_n derived from interchanging n and j in σ . Since the cycle structures of σ and σ' are the same, they have the same parity. Thus, we have

$$\operatorname{Cdet}(P_{jn}^*AP_{jn}) = \sum_{\sigma \in S_n^C} (-1)^{n-r} a_{\sigma'}.$$

Consider the following two cases.

Case (a). Suppose that n and j are in the same cycle factor. Then there exist nonnegative integers s, t such that $\sigma_1 = (np_1 \cdots p_s jq_1 \cdots q_t)$. Let $\bar{\sigma}_1 = (nq_t \cdots q_1 jp_s \cdots p_1)$, $\delta_1 = (nq_1 \cdots q_t jp_1 \cdots p_s)$, $\bar{\delta}_1 = (np_s \cdots p_1 jq_t \cdots q_1)$ be the first cycle factors of $\bar{\sigma} = \bar{\sigma}_1 \sigma_2 \cdots \sigma_r \in S_n^C$, $\delta = \delta_1 \sigma_2 \cdots \sigma_r \in S_n^C$, $\bar{\delta} = \bar{\delta}_1 \sigma_2 \cdots \sigma_r \in S_n^C$, respectively. Then by Lemma 3.1, we have $\langle \sigma_1 \rangle + \langle \bar{\sigma}_1 \rangle = \langle \delta_1' \rangle + \langle \bar{\delta}_1' \rangle$ and $\langle \delta_1 \rangle + \langle \bar{\delta}_1 \rangle = \langle \sigma_1' \rangle + \langle \bar{\sigma}_1' \rangle$. Here, the apostrophe of a cycle denotes the interchanging of n and j. Thus, we have

$$\langle \sigma \rangle + \langle \bar{\sigma} \rangle + \langle \delta \rangle + \langle \bar{\delta} \rangle = \langle \sigma' \rangle + \langle \bar{\sigma}' \rangle + \langle \delta' \rangle + \langle \bar{\delta}' \rangle.$$

Namely, we have built the correspondence between the permutation terms in S_n^C with the permutation terms of $\text{Cdet}(P_{jn}^*AP_{jn})$.

Case (b). Suppose that n and j are in two distinct cycle factors. Without loss of generality, let

$$\sigma = (np_1 \cdots p_s) \cdots (jq_1 \cdots q_t) \cdots, \ s, t \ge 0,$$

 $j > \max_{i=1}^{s} p_i$, and $j > \max_{i=1}^{t} q_i$. Otherwise, if $j > \max_{i=1}^{s} p_i$ or $j > \max_{i=1}^{t} q_i$ does not hold, we can use the similar technique in Lemma 3.1 to get the corresponding permutations without changing the summation values.

Let

$$\bar{\sigma} = (np_s \cdots p_1) \cdots (jq_1 \cdots q_t) \cdots ,$$
$$\delta = (np_1 \cdots p_s) \cdots (jq_t \cdots q_1) \cdots ,$$

$$\bar{\delta} = (np_s \cdots p_1) \cdots (jq_t \cdots q_1) \cdots,$$

and Σ , $\overline{\Sigma}$, Δ , $\overline{\Delta}$ be the cycles that interchanging $p_1 \cdots p_s$ with $q_1 \cdots q_t$ in σ , $\overline{\sigma}$, δ , $\overline{\delta}$, respectively. Then we have

$$\begin{aligned} \langle \sigma \rangle + \langle \bar{\sigma} \rangle + \langle \delta \rangle + \langle \bar{\delta} \rangle \\ &= [\langle np_1 \cdots p_s n \rangle + \langle np_s \cdots p_1 n \rangle] \cdots [\langle jq_1 \cdots q_t j \rangle + \langle jq_t \cdots q_1 i \rangle] \cdots \\ &= [\langle jq_1 \cdots q_t j \rangle + \langle jq_t \cdots q_1 j \rangle] \cdots [\langle np_1 \cdots p_s n \rangle + \langle np_s \cdots p_1 n \rangle] \cdots \\ &= \langle \Sigma' \rangle + \langle \bar{\Sigma}' \rangle + \langle \Delta' \rangle + \langle \bar{\Delta}' \rangle. \end{aligned}$$

Here, the second equality follows from the fact that the numbers in the square brackets are dual numbers, which are communicative with dual quaternion numbers. Similarly, we have $\langle \Sigma \rangle + \langle \bar{\Sigma} \rangle + \langle \Delta \rangle + \langle \bar{\Delta} \rangle = \langle \sigma' \rangle + \langle \bar{\sigma}' \rangle + \langle \delta' \rangle + \langle \bar{\delta}' \rangle$. Thus, we have built the correspondence between the factors in $\operatorname{Cdet}(P_{ij}^*AP_{ij})$ and $\operatorname{Cdet}(A)$.

Following this scheme, we could show $\operatorname{Cdet}(P_{ij}^*AP_{ij}) = \operatorname{Cdet}(A)$. This completes the proof.

Lemma 3.3. Let $A = (a_{ij})$ be a dual quaternion Hermitian matrix in $\hat{\mathbb{H}}^{n \times n}$, and $P_{n;\alpha}$ be a multiplication matrix. Then $\operatorname{Cdet}(P_{n;\alpha}A) = \alpha \operatorname{Cdet}(A)$, $\operatorname{Cdet}(AP_{n;\alpha}) = \alpha \operatorname{Cdet}(A)$, and $\operatorname{Cdet}(P_{i;\alpha}^*AP_{i;\alpha}) = \alpha^* \alpha \operatorname{Cdet}(A)$ for all $i = 1, \ldots, n$.

Proof. Let $\sigma = \sigma_1 \cdots \sigma_r \in S_n^C$ be a permutation of S_n satisfying (8) and $\sigma_1 = (np_1 \cdots p_s)$. For the *n*-th row multiplication, we have

$$\operatorname{Cdet}(P_{n;\alpha}A) = \sum_{\sigma \in S_n^C} (-1)^{n-r} \alpha a_{np_1} a_{p_1 p_2} \cdots a_{p_s n} \langle \sigma_2 \rangle \cdots \langle \sigma_r \rangle$$
$$= \alpha \left(\sum_{\sigma \in S_n^C} (-1)^{n-r} \langle \sigma_1 \rangle \cdots \langle \sigma_r \rangle \right)$$
$$= \alpha \operatorname{Cdet}(A).$$

In addition, there exists $\bar{\sigma}_1 = (np_s \cdots p_1)$ such that $\bar{\sigma} = \bar{\sigma}_1 \sigma_2 \cdots \sigma_r \in S_n^C$ and

 $\langle \sigma_1 \rangle + \langle \bar{\sigma}_1 \rangle \in \hat{\mathbb{R}}$. For the *n*-th column multiplication, we have

$$Cdet(AP_{n;\alpha}) = \sum_{\sigma \in S_n^C} (-1)^{n-r} a_{np_1} a_{p_1p_2} \cdots a_{p_sn} \alpha \langle \sigma_2 \rangle \cdots \langle \sigma_r \rangle$$

$$= \sum_{\sigma \in S_n^C} (-1)^{n-r} \langle \sigma_1 \rangle \alpha \langle \sigma_2 \rangle \cdots \langle \sigma_r \rangle$$

$$= \frac{1}{2} \sum_{\sigma \in S_n^C} (-1)^{n-r} [\langle \sigma_1 \rangle + \langle \bar{\sigma}_1 \rangle] \alpha \langle \sigma_2 \rangle \cdots \langle \sigma_r \rangle$$

$$= \frac{\alpha}{2} \sum_{\sigma \in S_n^C} (-1)^{n-r} [\langle \sigma_1 \rangle + \langle \bar{\sigma}_1 \rangle] \langle \sigma_2 \rangle \cdots \langle \sigma_r \rangle$$

$$= \alpha Cdet(A).$$

Here, the third equality is true because we have repeated all permutations twice and the fourth equality is true since the number in the square bracket is a dual number.

For the i-th row and column multiplications, we have

$$Cdet(P_{i;\alpha}^*AP_{i;\alpha}) = Cdet(P_{in}^*P_{i;\alpha}^*AP_{i;\alpha}P_{in})$$

=
$$Cdet((P_{i;\alpha}P_{in})^*A(P_{i;\alpha}P_{in}))$$

=
$$Cdet(P_{n;\alpha}^*AP_{n;\alpha})$$

=
$$\alpha^*\alpha Cdet(A).$$

This completes the proof.

Lemma 3.4. Let $A = (a_{ij})$ be a dual quaternion Hermitian matrix in $\hat{\mathbb{H}}^{n \times n}$. Then

$$\operatorname{Cdet}(A_{n \cdot}(a_{k \cdot})) = 0, \quad \forall k = 1, \dots, n-1.$$

Furthermore, we have $\operatorname{Cdet}(A_{n}(\alpha a_{k})) = 0$. In other words, if we replace the nth row of a dual quaternion Hermitian matrix by the k-th row (possibly multiplied by a dual quaternion number α on the left), then the Chen determinant of the result matrix is zero.

Proof. Let s, t be nonnegative integers. We first divide all permutations in S_n^C into three sets depending on the location of k as follows.

Е		-
н		

(i)
$$S_1 = \{\sigma_1(k) : \sigma_1(k) = (np_1 \cdots p_s) \cdots (kq_1 \cdots q_t) \cdots \};$$

(ii) $S_2 = \{\sigma_2(k) : \sigma_2(k) = (np_1 \cdots p_s) \cdots (u_1 \cdots u_l kq_1 \cdots q_t) \cdots \};$
(iii) $S_3 = \{\sigma_3(k) : \sigma_3(k) = (np_1 \cdots p_s kq_1 \cdots q_t) \cdots \}.$

We show this lemma by proving that for any permutation σ in the first two cases, there is a permutation $\bar{\sigma}$ in the same case and permutations σ^+ and $\bar{\sigma}^+$ in the third case such that

$$s(\sigma)\langle\sigma\rangle + s(\bar{\sigma})\langle\bar{\sigma}\rangle = -s(\sigma^+)\langle\sigma^+\rangle - s(\bar{\sigma}^+)\langle\bar{\sigma}^+\rangle.$$
(14)

Case (i). Let $\sigma \in S_1$ and

$$\bar{\sigma} = (np_1 \cdots p_s) \cdots (kq_t \cdots q_1) \cdots,$$
$$\sigma^+ = (nq_1 \cdots q_t kp_1 \cdots p_s) \cdots, \quad \bar{\sigma}^+ = (nq_t \cdots q_1 kp_1 \cdots p_s) \cdots.$$

Then we have

$$\begin{aligned} \langle \sigma \rangle + \langle \bar{\sigma} \rangle &= \langle np_1 \cdots p_s n \rangle \cdots \langle kq_1 \cdots q_t k \rangle \cdots + \langle np_1 \cdots p_s n \rangle \cdots \langle kq_t \cdots q_1 k \rangle \cdots \\ &= \langle kp_1 \cdots p_s n \rangle \cdots \langle kq_1 \cdots q_t k \rangle \cdots + \langle kp_1 \cdots p_s n \rangle \cdots \langle kq_t \cdots q_1 k \rangle \cdots \\ &= \langle kq_1 \cdots q_t kp_1 \cdots p_s n \rangle \cdots + \langle kq_t \cdots q_1 kp_1 \cdots p_s n \rangle \cdots \\ &= \langle nq_1 \cdots q_t kp_1 \cdots p_s n \rangle \cdots + \langle nq_t \cdots q_1 kp_1 \cdots p_s n \rangle \cdots \\ &= \langle \sigma^+ \rangle + \langle \bar{\sigma}^+ \rangle, \end{aligned}$$

where the second and the fourth equalities follow from the fact that the last row of A is replaced by the k-th row, and the third equality follows from the fact that $\langle kq_1 \cdots q_t k \rangle + \langle kq_t \cdots q_1 k \rangle$ is a dual number, which is communicative with dual quaternion numbers and can be moved to the beginning of the products. Furthermore, suppose σ has r distinct cycles. Then $s(\sigma) = s(\bar{\sigma}) = (-1)^{n-r}$ and $s(\sigma^+) = s(\bar{\sigma}^+) = (-1)^{n-r+1}$. Thus, (14) holds.

Case (ii). Let $\sigma \in S_2$ and

$$\bar{\sigma} = (np_1 \cdots p_s) \cdots (u_1 q_t \cdots q_1 k u_l \cdots u_2) \cdots ,$$
$$\sigma^+ = (nq_1 \cdots q_t u_1 u_2 \cdots u_l k p_1 \cdots p_s) \cdots ,$$

and

$$\sigma^+ = (nu_l \cdots u_1 q_t \cdots q_1 k p_1 \cdots p_s) \cdots$$

It follows from Lemma 3.1 that

$$\langle u_1 \cdots u_l k q_1 \cdots q_t u_1 \rangle + \langle u_1 q_t \cdots q_1 k u_l \cdots u_1 \rangle$$

$$= \langle k q_1 \cdots q_t u_1 u_2 \cdots u_l k \rangle + \langle k u_l \cdots u_1 q_t \cdots q_1 k \rangle.$$
(15)

Combining this with the first case derives (14).

The union of all σ^+ and $\bar{\sigma}^+$ defined above is equal to S_3 . In fact, for any permutation $\sigma \in S_3$, if $k \ge p_i$ for $i = 1, \ldots, s$, then it is corresponding to a permutation in the first case. Otherwise, it is corresponding to a permutation in the second case.

Together, we have

$$Cdet(A_{n}.(a_{k}.)) = \sum_{\sigma \in S_{1}} s(\sigma) \langle \sigma \rangle + \sum_{\sigma \in S_{2}} s(\sigma) \langle \sigma \rangle + \sum_{\sigma \in S_{3}} s(\sigma) \langle \sigma \rangle$$
$$= -\sum_{\sigma \in S_{3}} s(\sigma) \langle \sigma \rangle + \sum_{\sigma \in S_{3}} s(\sigma) \langle \sigma \rangle$$
$$= 0.$$

Furthermore, we can verify that $\operatorname{Cdet}(A_{n}(\alpha a_{k})) = \alpha \operatorname{Cdet}(A_{n}(a_{k})) = 0$. This completes the proof.

Lemma 3.5. Let $A = (a_{ij})$ be a dual quaternion Hermitian matrix in $\hat{\mathbb{H}}^{n \times n}$. Then we have $\operatorname{Cdet}(A_{\cdot n}(a_{\cdot k}\alpha)) = 0$ for all $k = 1, \ldots, n$ and $\alpha \in \hat{\mathbb{H}}$.

Proof. Consider the three cases S_1 , S_2 and S_3 in the proof of Lemma 3.4. We now prove that for any permutation $\sigma \in S_1 \cup S_2$, there is a permutation $\bar{\sigma}$ in the same case and permutations $\sigma^+, \bar{\sigma}^+ \in S_3$ such that (14) holds.

Case (i). Let $\sigma \in S_1$ and

$$\bar{\sigma} = (np_1 \cdots p_s) \cdots (kq_t \cdots q_1) \cdots,$$
$$\sigma^+ = (np_1 \cdots p_s kq_1 \cdots q_t) \cdots, \quad \bar{\sigma}^+ = (np_1 \cdots p_s kq_t \cdots q_1) \cdots.$$

Then we have

$$\begin{aligned} \langle \sigma \rangle + \langle \bar{\sigma} \rangle &= \langle np_1 \cdots p_s n \rangle \cdots \langle kq_1 \cdots q_t k \rangle \cdots + \langle np_1 \cdots p_s n \rangle \cdots \langle kq_t \cdots q_1 k \rangle \\ &= \langle np_1 \cdots p_s k \rangle \alpha \cdots \langle kq_1 \cdots q_t k \rangle \cdots + \langle np_1 \cdots p_s k \rangle \alpha \cdots \langle kq_t \cdots q_1 k \rangle \\ &= \langle np_1 \cdots p_s kq_1 \cdots q_t k \rangle \alpha \cdots + \langle np_1 \cdots p_s kq_t \cdots q_1 k \rangle \alpha \cdots \\ &= \langle np_1 \cdots p_s kq_1 \cdots q_t n \rangle \cdots + \langle np_1 \cdots p_s kq_t \cdots q_1 n \rangle \cdots \\ &= \langle \sigma^+ \rangle + \langle \bar{\sigma}^+ \rangle, \end{aligned}$$

where the second and the fourth equalities follow from the fact that the last column of A is replaced by the k-th row multiplied by α , and the third equality follows from the fact that $\langle kq_1 \cdots q_t k \rangle + \langle kq_t \cdots q_1 k \rangle$ is a dual number and is communicative with any dual quaternion numbers. Furthermore, suppose σ has r distinct cycles. Then $s(\sigma) = s(\bar{\sigma}) = (-1)^{n-r}$ and $s(\sigma^+) = s(\bar{\sigma}^+) = (-1)^{n-r+1}$. Thus, (14) holds.

Case (ii). Let $\sigma \in S_2$ and

$$\bar{\sigma} = (np_1 \cdots p_s) \cdots (u_1 q_t \cdots q_1 k u_l \cdots u_2) \cdots ,$$
$$\sigma^+ = (np_1 \cdots p_s k q_1 \cdots q_t u_1 \cdots u_l n) \cdots ,$$

and

 $\sigma^+ = (np_1 \cdots p_s ku_l \cdots u_1 q_t \cdots q_1 n) \cdots$

Combining (15) with the first case derives (14).

The rest of the proof is similar to that of Lemma 3.4 and we omit the details here.

This completes the proof.

Lemma 3.6. Let $A = (a_{ij})$ be a dual quaternion Hermitian matrix in $\hat{\mathbb{H}}^{n \times n}$, and $P_{ij;\alpha}$ be an addition matrix. Then $\operatorname{Cdet}(P_{ij;\alpha}^*AP_{ij;\alpha}) = \operatorname{Cdet}(A)$.

Proof. Without loss of generality, we suppose i < j. Since $P_{ij;\alpha} = P_{jn}P_{in;\alpha}P_{jn}$, we only need to show $\operatorname{Cdet}(P_{in;\alpha}^*AP_{in;\alpha}) = \operatorname{Cdet}(A)$. By direct computation, we have

$$\operatorname{Cdet} \left(P_{in;\alpha}^* A P_{in;\alpha} \right)$$

=
$$\operatorname{Cdet} \left(A \right) + \operatorname{Cdet} \left(A_{n} \left(\alpha^* a_i \right) \right) + \operatorname{Cdet} \left(A_{\cdot n} (a_{\cdot i} \alpha) \right) + \operatorname{Cdet} \left(B(\alpha) \right),$$

where $B(\alpha) = (b_{kl}(\alpha))$ satisfies

$$b_{kl}(\alpha) = \begin{cases} \alpha^* a_{il}, & \text{if } k = n \text{ and } l \neq n; \\ a_{ki}\alpha, & \text{if } l = n \text{ and } k \neq n; \\ \alpha^* a_{ii}\alpha, & \text{if } k = l = n \\ a_{kl}, & \text{otherwise.} \end{cases}$$

In the other words, $B(\alpha) = P_{n;\alpha}^* B(1) P_{n;\alpha}$. Thus, it follows from Lemmas 3.3 and 3.4 that $\operatorname{Cdet}(B(\alpha)) = \alpha^* \alpha \operatorname{Cdet}(B(1)) = 0$. By Lemmas 3.4 and 3.5, we have $\operatorname{Cdet}(A_{n\cdot}(\alpha^*a_{i\cdot})) = \operatorname{Cdet}(A_{\cdot n}(a_{\cdot i}\alpha)) = 0$. Therefore, we have $\operatorname{Cdet}(P_{in;\alpha}^*AP_{in;\alpha}) = \operatorname{Cdet}(A)$. This completes the proof.

Recently, Wang et al. [19] proposed the dual quaternion LU decomposition and the partial pivoting dual quaternion LU decomposition. Specifically, given a dual quaternion matrix $A \in \hat{\mathbb{H}}^{n \times n}$, there is a permutation matrix P, a unit lower triangular dual quaternion matrix $L \in \hat{\mathbb{H}}^{n \times n}$, and an upper triangular dual quaternion matrix $U \in \hat{\mathbb{H}}^{n \times n}$ such that

$$PA = LU$$

Based on this formulation, we have the following lemma.

Lemma 3.7. Suppose $A \in \hat{\mathbb{H}}^{n \times n}$ is a dual quaternion unitary matrix. Then there exists a series of switching matrices and addition matrices P_1, \ldots, P_t such that $P_t \cdots P_1 A = \text{diag}(d_1, \ldots, d_n)$, where $d_1, \ldots, d_n \in \hat{\mathbb{H}}$ and $|d_1 d_2 \cdots d_n| = 1$.

Proof. Let PA = LU be the partial pivoting dual quaternion LU decomposition of A. Since P, A, and L are all invertiable, U is also invertiable. Let $U = U_0D$, where $D = \text{diag}(d_1, \ldots, d_n)$ and $d_i = u_{ii}$ for $i = 1, \ldots, n$. Then we have

$$U_0^{-1}L^{-1}PA = D = \text{diag}(d_1, \dots, d_n).$$

Here, U_0^{-1} and L^{-1} are unit upper and lower triangular dual quaternion matrices, respectively. Thus, they are products of addition matrices.

Furthermore, by the quasi-determinant defined in Ling and Qi [11], we have $\det_q(U_0^{-1}) = 1$, $\det_q(L^{-1}) = 1$, $\det_q(P) = 1$, $\det_q(A^*A) = \det_q(A^*)\det_q(A) = \det_q(A)^2 = 1$, and

$$\det_q(U_0^{-1}L^{-1}PA) = \det_q(U_0^{-1})\det_q(L^{-1})\det_q(P)\det_q(A) = 1.$$

Therefore, $det_q(D) = |d_1 d_2 \cdots d_n|^2 = 1$. This completes the proof.

Based on the above lemmas, we are ready to show our main results.

Theorem 3.8. For any dual quaternion Hermitian matrix $A \in \hat{\mathbb{H}}^{n \times n}$ and any dual quaternion unitary matrix $U \in \hat{\mathbb{H}}^{n \times n}$, we have $\operatorname{Cdet}(U^*AU) = \operatorname{Cdet}(A)$.

Proof. By Lemma 3.7, there are switching matrices and addition matrices P_1, \ldots, P_t such that $U = P_1^{-1} \cdots P_t^{-1} D$, $D = \text{diag}(d_1, \ldots, d_n)$, and $|d_1 \cdots d_n| = 1$. Since $P_{ij}^{-1} = P_{ij}, P_{ij;\alpha}^{-1} = P_{ij;-\alpha}, P_1^{-1}, \ldots, P_t^{-1}$ are also switching or addition matrices. Thus, we have

$$\operatorname{Cdet}(U^*AU) = \operatorname{Cdet}(D^*(P_t^{-1})^* \cdots (P_1^{-1})^* AP_1^{-1} \cdots P_t^{-1}D) = \operatorname{Cdet}(A).$$

Here, the last equality follows directly from Proposition 2.3. This completes the proof. $\hfill \Box$

Theorem 3.9. Given a dual quaternion Hermitian matrix $A \in \hat{\mathbb{H}}^{n \times n}$ and any dual quaternion unitary matrix $U \in \hat{\mathbb{H}}^{n \times n}$, we have

$$\operatorname{Cdet}(A) = \prod_{i=1}^{n} \lambda_i$$

and

$$\operatorname{Cdet}(A^*A) = \operatorname{Cdet}(A)^2 = \prod_{i=1}^n \lambda_i^2,$$

where $\lambda_1, \ldots, \lambda_n$ are eigenvalues of A.

Proof. By [16], there exists a dual quaternion unitary matrix U such that $UAU^* = \Lambda = \operatorname{diag}(\lambda_1, \ldots, \lambda_n)$ and all eigenvalues $\lambda_i, i = 1, \ldots, n$ are dual numbers. Combining this with Theorem 3.8, we have $\operatorname{Cdet}(A) = \operatorname{Cdet}(U^*AU) = \operatorname{Cdet}(\Lambda) = \lambda_n \cdots \lambda_1$. Since dual numbers are communicative, the first conclusion is derived.

The second conclusion follows from $\Lambda^* = \Lambda$, $A^*A = U^*\Lambda^2 U$ and Theorem 3.8. This completes the proof.

4 The Moore Determinant and the Eigenvalues

Theorem 4.1. Let $A = (a_{ij})$ be a dual quaternion Hermitian matrix in $\hat{\mathbb{H}}^{n \times n}$. Then we have

$$Mdet(A) = Cdet(A).$$
(16)

Furthermore, we have

$$\mathrm{Mdet}(A) = \prod_{i=1}^{n} \lambda_i,$$

where $\lambda_1, \ldots, \lambda_n$ are the eigenvalues of A.

Proof. Let $\sigma = \sigma_1 \cdots \sigma_r \in S_n^M$ be a permutation in the Moore determinant. For each $i = 1, \ldots, r$, denote

$$\sigma_i = (p_{i1} \cdots p_{is_i} k_i q_{i1} \cdots q_{it_i}),$$

where p_{i1} and k_i are the minimal and maximal integers in σ_i , respectively. Denote $\bar{\sigma}_i = (p_{i1}q_{it_i}\cdots q_{i1}k_ip_{is_i}\cdots p_{i2}), \ \sigma_i^+ = (k_iq_{i1}\cdots q_{it_i}p_{i1}\cdots p_{is_i}), \ \text{and} \ \bar{\sigma}_i^+ = (k_ip_{is_i}\cdots p_{i1}q_{it_i}\cdots q_{i1}).$ By Lemma 3.1, we have

$$\langle \sigma_i \rangle + \langle \bar{\sigma}_i \rangle = \langle \sigma_i^+ \rangle + \langle \bar{\sigma}_i^+ \rangle \in \hat{\mathbb{R}}.$$

Thus, we have

$$(\langle \sigma_1 \rangle + \langle \bar{\sigma}_1 \rangle) \cdots (\langle \sigma_r \rangle + \langle \bar{\sigma}_r \rangle)$$

= $(\langle \sigma_1^+ \rangle + \langle \bar{\sigma}_1^+ \rangle) \cdots (\langle \sigma_r^+ \rangle + \langle \bar{\sigma}_r^+ \rangle)$
= $(\langle \sigma_{i_1}^+ \rangle + \langle \bar{\sigma}_{i_1}^+ \rangle) \cdots (\langle \sigma_{i_r}^+ \rangle + \langle \bar{\sigma}_{i_r}^+ \rangle)$

Here, i_1, \ldots, i_r are obtained by sorting k_i in the descending order and the second equality holds since the dual numbers are communicative. Let $s_j \in \{\sigma_j, \bar{\sigma}_j\}$ for $j = 1, \ldots, n$. Then $\sigma = s_1 \cdots s_r \in S_n^M$ is a permutation in the Moore determinant. Similarly, let $s_j^+ \in \{\sigma_{i_j}^+, \bar{\sigma}_{i_j}^+\}$ for $j = 1, \ldots, n$. Then $\sigma^+ = s_1^+ \cdots s_r^+ \in S_n^C$ is a permutation in the Chen determinant. Furthermore, those cycle factors are obtained by reversing or switching the original cycle factor. Reversing a cycle factor derives the inverse or conjugate of the corresponding permutation, which does not change the sign. Switching the cycle factors does not change the corresponding permutation. Thus, the signs of all permutations above are the same. By going through all permutations in the Moore determinant, we have Mdet(A) = Cdet(A).

The second conclusion of this theorem holds directly from equation (16) and Theorem 3.9. This completes the proof. $\hfill \Box$

Kyrchei [7, 8] showed that the Kyrchei row and column determinants of a quaternion Hermitian matrix are equal to the Moore determinant. We could also extend this result to dual quaternion Hermitian matrices and show the Kyrchei row and column determinants of a quaternion Hermitian matrix are equal to the Moore determinants, which are also equal to the products of the eigenvalues.

Theorem 4.2. Suppose $A \in \hat{\mathbb{H}}^{n \times n}$ is a dual quaternion Hermitian matrix and $\lambda_1, \ldots, \lambda_n \in \hat{\mathbb{R}}$ are its eigenvalues. Then Mdet(A) = 0 if and only if A_s is singular and A has at least one zero or two infinitesimal eigenvalues; namely, there is $i, j \in \{1, \ldots, n\}$ such that $\lambda_i = 0$ or $\lambda_{i,s} = \lambda_{j,s} = 0$.

Proof. By Theorem 4.1, we have

$$Mdet(A) = \prod_{i=1}^{n} \lambda_i = \prod_{i=1}^{n} \lambda_{is} + \sum_{j=1}^{n} \prod_{i=1, i \neq j}^{n} \lambda_{is} \lambda_{jd} \epsilon.$$

On one hand, if there is $i, j \in \{1, ..., n\}$ such that $\lambda_i = 0$ or $\lambda_{i,s} = \lambda_{j,s} = 0$, we can check that Mdet(A) = 0 directly. On the other hand, if Mdet(A) = 0, then there is $j \in \{1, ..., n\}$ such that $\lambda_{js} = 0$ and $\prod_{i=1, i \neq j}^{n} \lambda_{is} \lambda_{jd} = 0$. Thus, either $\lambda_{jd} = 0$ or there is $i \neq j$ such that $\lambda_{is} = 0$. This completes the proof. \Box

5 The Characteristic Polynomial

Suppose $A \in \hat{\mathbb{H}}^{n \times n}$ is a dual quaternion Hermitian matrix and $\lambda \in \hat{\mathbb{H}}$ is an arbitrary dual number. Define

$$p(\lambda) = \mathrm{Mdet}(\lambda I - A). \tag{17}$$

Assume that $\lambda_1, \ldots, \lambda_n$ are eigenvalues of A. Then (17) is equivalent to

$$p(\lambda) = (\lambda - \lambda_1)(\lambda - \lambda_2) \cdots (\lambda - \lambda_n).$$
(18)

The later definition is also used in Qi and Cui [15]. It was shown in [15] that all eigenvalues of A are also roots of the characteristic polynomial. The reverse does not hold in general. In fact, a dual number λ is a root of $p(\lambda)$ either if λ is an eigenvalue of A or λ_s is a multiple eigenvalue of A_s . However, the reverse holds true when all eigenvalues of A_s are single.

Corollary 5.1. Suppose all eigenvalues of A_s are single. Then a dual number is the root of the characteristic polynomial (17) if and only if it is an eigenvalue of A.

Very recently, Ling and Qi [11] proposed a characteristic polynomial by the quasi-determinant of dual quaternion matrices. In their definition, there is

$$p_q(\lambda) = \det_q(\lambda I - A) = |\lambda - \lambda_1|^2 |\lambda - \lambda_2|^2 \cdots |\lambda - \lambda_n|^2.$$
(19)

We may verify that $p_q(\lambda) = |p(\lambda)|^2$. In other words, the sign of the determinant is lost in the quasi-determinant. Furthermore, it was shown in [11] that all eigenvalues of A are also roots of the characteristic polynomial (19). However, the reverse does not hold, even if all eigenvalues of A_s are single.

Acknowledgment We are grateful to Prof. Zhigang Jia for helpful discussions.

References

- H. Aslaksen, "Quaternionic determinants", The Mathematical Intelligencer 18 (1996) 57-65.
- [2] P. J. Cameron, *Permutation Groups*, Cambridge University Press, Cambridge, 1999.
- [3] L. Chen, "Definition of determinant and Cramer solution over the quaternion field", Acta Mathematica Sinica 7 (1991) 171-180.

- [4] L. Chen, "Inverse matrix and properties of double determinant over quaternion field", Science in China, Series A: 34 (1991) 528-540.
- [5] C. Cui, Y. Lu, L. Qi, and L. Wang, "Spectral properties of dual unit gain graphs", May 2024, arXiv:2402.12988v2.
- [6] F.J. Dyson, "Quaternion determinants", Helvetica Physica Acta 45 (1972) 289-302.
- [7] I.I. Kyrchei, "Cramer's rule for quaternionic systems of linear equations", Journal of Mathematical Sciences 155 (2008) 839-858.
- [8] I.I. Kyrchei, "The theory of the column and row determinants in a quaternion linear algebra", in: A.R. Baswell ed., Advances in Mathematical Research 15, Nova Science Publishers, 2012, pp. 301-358.
- [9] I.I. Kyrchei, E. Treister and V.O. Pelykh, "The determinant of the Laplacian matrix of a quaternion unit gain graph", *Discrete Mathematics* 147 (2024) No. 113955.
- [10] W. Li, Quaternion Matrices, in Chinese, National University of Defence Technology Publisher, Changsha, 2002.
- [11] C. Ling and L. Qi, "Quasi-determinant and right eigenvalues of dual quaternion matrices", April 2024, arXiv:2404.19348.
- [12] Q. Liu, S. Ling and Z. Jia, "Randomized quaternion singular value decomposition for low-rank matrix approximation", SIAM Journal on Scientific Computing, 44 (2022) A870-A900.
- [13] X. Liu, G.J. Song and Y. Zhang, "Determinantal representation of the solutions to systems of generalized Sylvester equations", Advances of Applied Clifford Algebras 30 (2020) No. 12.
- [14] E.H. Moore, "On the determinant of an hermitian matrix of quaternionic elements", Bulletin of the American Mathematical Society 28 (1922) 161-162.

- [15] L. Qi and C. Cui, "Eigenvalues of dual Hermitian matrices with application in formation control", April 2024, arXiv:2403.10308v2.
- [16] L. Qi and Z. Luo, "Eigenvalues and singular values of dual quaternion matrices", *Pacific Journal of Optimization* 19 (2023) 257-272.
- [17] L. Qi, X. Wang and Z. Luo, "Dual quaternion matrices in multi-agent formation control", Communications in Mathematical Sciences 21 (2023) 1865-1874.
- [18] E. Study, "Zur Theorie der linearen Gleichungen", Acta Mathematica 42 (1920) 1-61.
- [19] T. Wang, Y. Li, M. Wei, Y. Xi, and M. Zhang, "Algebraic method for LU decomposition of dual quaternion matrix and its corresponding structure-preserving algorithm", *Numerical Algorithms* https://doi.org/10.1007/s11075-024-01753-8.
- [20] X. Wang, C. Yu and Z. Lin, "A dual quaternion solution to attitude and position control for rigid body coordination", *IEEE Transactions on Robotics* 28 (2012) 1162-1170.
- [21] F. Zhang, "Quaternions and matrices of quaternions", Linear Algebra and its Applications 251 (1997) 21-57.