Kolmogorovian Censorship, Predictive Incompleteness, and the locality loophole in Bell experiments.

Philippe Grangier

Laboratoire Charles Fabry, IOGS, CNRS, Université Paris Saclay, F91127 Palaiseau, France.

In the foundations of quantum mechanics, the Kolmogorovian Censorship (KC) stipulates that quantum probabilities can be identified with classical, Kolmogorovian probabilities when considering a specified measurement context. Then in any given measurement context it is possible to build a Kolmogorovian probability distribution, or equivalently a hidden variable theory; however this distribution must be matched to the chosen context. In a loophole-free Bell test, the remote random choices of measurements (polarizers orientations) have the purpose to prevent that this matching can be obtained from any relativistically causal transmission between the source and the detectors. Then the matching (required to violate Bell's inequalities) may be obtained either by an instantaneous influence at a distance between the source and the detectors (explicit nonlocality), or by assuming that it is pre-established before the actual experiment takes place (super-determinism). If both influence at a distance and super-determinism are not accepted on physical grounds, a third way is still available, called 'predictive incompleteness': it tells that the usual quantum state ψ is incomplete, as long as the measurement context has not been specified. In agreement with the general quantum framework called CSM (Contexts, Systems and Modalities) we argue that predictive incompleteness is the correct quantum way to understand the violation of Bell's inequalities.

I. INTRODUCTION.

In the scientific literature there are still debates to decide whether quantum mechanics (QM) is contextual [1, 2] or noncontextual [3], and whether quantum probabilities are Kolmogorovian [4, 5] or not [6–8]. To some extend, these debates can be considered as matters of definitions, so it is useful to spell out the explicit or implicit assumptions that are are the root of the disagreements. A useful milestone in this debate is the Kolmogorovian Censorship (KC), stating that quantum probabilities can be identified with classical, Kolmogorovian probabilities when the measurement context has been specified, or more generally when a classical probability distribution of measurement contexts has been specified. The KC sounds rather obvious on a physical intuitive basis, and technically it has been demonstrated for a countable number of measurement contexts [9, 10].

II. KOLMOGOROVIAN CENSORSHIP AND BELL EXPERIMENTS.

However there is some controversy on the physical meaning of the KC, as we will explain now. In particular, Szabó et al [4, 5] claim that QM probabilities are Kolmogorovian, not only in a single context, but also in a loophole-free Bell experiment, where a random choice among four different contexts is implemented [13]. Then they conclude that Bell's inequalities are either irrelevant (when considering the four contexts separately), or not violated, when considering the four contexts together: since each one has a 1/4 probability to occur, Bell's S parameter (see Annex) is reduced from $2\sqrt{2}$ down to $\sqrt{2}/2$, that is below 2 as it would be expected classically.

More precisely, it is uncontroversial that the probabilities predicted by QM in a given context are Kolmogorovian. and the problem arises when trying to "put together" probabilities predicted by QM in different contexts. Said otherwise, in any given context it is possible to build a Kolmogorovian probability distribution, or equivalently an hidden variable theory; however this distribution must be adapted to the chosen context.

In a loophole-free Bell test [13], the remote random choices of measurements (polarizers orientations) are designed to forbid that this matching may be done by a relativistically causal transmission between the source and the detectors. Then it may be obtained either by an instantaneous influence between the source and the detectors (explicit nonlocality), or by assuming that it is pre-established before the actual experiment takes place (super-determinism). One has then the following possible options:

1. Szabó et al [4, 5] claim that the only relevant way to speak about QM probabilities in different contexts is to consider a classical probability distribution over these different contexts. This corresponds by construction to a feasible experiment, typically a loophole-free Bell experiment, including the random choice among four different contexts [13]. Then the global probability distribution is Kolmogorovian, and Bell's inequalities (BI) are not violated, because as written above the probabilities in each context are divided by four, as well as the resulting S value. However a drawback of this approach is that the pre-established matching between the source and the measurements is still required, through some kind of "global determinism" [15]. Therefore, though BI are not explicitly violated, the basic physical problem of the origin of the matching is still present.

- 2. Since the averaging over the contexts does not really help, one may calculate the correlation function in each context, which is the very idea of a loophole-free Bell test, and gather them in Bell's S value. However, one may argue that the four correlations functions correspond to four different incompatible experiments, and thus bringing their results together is counterfactual, because they cannot be measured simultaneously; then BI cannot be demonstrated. This is a standard answer to the problem, by simply telling that the question is irrelevant; but without further explanation it sounds more dogmatic than scientific.
- 3. Still an alternative way is to admit that the results in the four contexts can be combined, as it would be the case classically since they apply to the same system; then one gets Bell's inequalities, which are experimentally violated, so one should explain why. We consider that this is a meaningful question, and there are basically three options, spelled out in detail in [14]:
- 3a in the spirit of option 1 above, keep a Kolmogorovian global probability distribution. Then the way to violate Bell's inequalities is by admitting that the system's parameters and the orientations of the polarizers are not independent variables, despite the fact that these orientations are chosen randomly and independently at a large distance. This can be obtained either by admitting superdeterminism (i.e. denying the possibility of independent random choices [15]), or by admitting a non-local influence between the source and the measurements [14]. These two options have recently been shown to be equivalent [16], and they are in our opinion equally undesirable though they are matter of ontological choice, and cannot be proven wrong.
- 3b the third option, known as predictive incompleteness [14, 17], is to recognize that the quantum state by itself is not enough to specify the measured probability distribution, as long as the context has not be specified. Since predictive completeness (also called outcome independence) is a required hypothesis for Bell's theorem, this leads to the conclusion that BI cannot be demonstrated. This conclusion corresponds to the detailed analysis presented in [14], and also the more general framework presented in [18] to "complete" the usual quantum state by specifying the measurement context. These two papers, as well as the arguments above, are consistent within the general quantum framework called CSM (Contexts, Systems and Modalities) [19, 20].

From the above it should be clear that the ontologies underlying either Szabó's position or CSM are quite different. Szabó et al claim that QM probabilities are Kolmogorovian and that QM can ultimately be seen as a (super)deterministic theory. On the other hand, CSM stipulates that QM is fundamentally non-deterministic, due to the conjunction of quantization and contextuality [11, 12], and that quantum probabilities are non-Kolmogorovian, unless restricted to a single context according to the KC. It is true, as claimed by Szabó et al, that the results of a full loophole-free Bell test with random choices of the measurements [13] can be embedded in a Kolmogorovian framework, as far as actual frequencies of observed events are considered.

However, the conflict arises if one considers that this final distribution is obtained from the combination of two Kolmogorovian distributions, one describing the emitted particles, and the other one describing the chosen measurements. If this is done then obtaining the correct final distribution requires either nonlocality or superdeterminism, as written above. This is why the (non-Kolmogorovian) predictive incompleteness of ψ is useful: it leaves enough freedom so that the choice of the measurement can contribute to the determination of the final distribution, avoiding both nonlocality and superdeterminism. It is worth emphasizing again, and spelled out in [14], that predictive incompleteness makes no sense in classical physics, and appears as a specific quantum feature.

III. CONCLUSIONS.

Again, the KC in a single context is easily integrated in the CSM framework, but one should not conclude that quantum probabilities are Kolmogorovian in a classical sense: this would be true in classical physics, because there is only one universal context, but this fails in quantum physics, because there is a continuous infinity of different, incompatible contexts. This point of view is implicit in textbook quantum mechanics, and it is made explicit in the CSM framework, by using operator algebra and infinite tensor products. This framework allows a contextual unification of classical and quantum physics, within a unique macroscopic physical world [22–24].

One may notice that Rédei [10] (see also [25]) considers that the KC is problematic, in particular because it implies that "probabilities are thus not features of quantum systems in and of themselves, they are features that only manifest themselves upon measurement. Philosophers (or physicists) with a robust realist conviction may find unattractive this strongly instrumentalist flavor of interpretation of quantum probability forced upon us by the KC." In the CSM approach, quantum probabilities also get a meaning only upon measurement, due to the predictive incompleteness of ψ , and they are also genuinely non-classical probabilities. Nevertheless, CSM is based on physical realism, as the statement that the purpose of physics is to study entities of the natural world, existing independently from any particular observer's perception, and obeying universal and intelligible rules [18–20]. We claim therefore that even with a "robust realist conviction" one can accept that ψ is predictively incomplete - as the only way to make sense of this conundrum. So here is a message to our philosopher friends: please have a closer look at CSM, and refrain telling too quickly that it is just one more anti-realist Bohrian mantra.

As a final remark, a significant challenge in quantum foundations and reconstructions is to state clearly what are the hypotheses or postulates, and what are their consequences. It should also been made clear whether one looks for a fully deductive reasoning, or a partially inductive one, that is an "Inference to the Best Explanation" [11, 12]. In the CSM point of view a fully deductive approach does not fit, since, quoting Landau [21], 'quantum mechanics (...) contains classical mechanics as a limiting case, yet at the same time it requires this limiting case for its own formulation'. Then what is desired is not a deduction of the Laws of Nature from some postulates in a mathematical sense, but rather a fully consistent construction, including both classical and quantum physics from the beginning, and clearly separating experimentally based evidence from its mathematical description [22–24].

Acknowledgments. The author thanks Mathias Van Den Bossche and Olivier Ezratty for relevant comments on this article, and for many interesting discussions.

Annex. Consider a Bell test with fast random switching of the polarizers orientations [13] between a and a' on one side, b and b' on the other side, where the results A(a), A(a'), B(b), B(b') are denoted ± 1 . The four correlation coefficients like $E(a,b) = \overline{A(a)B(b)}$ are calculated from the number of counts $N_{\pm}(a,b)$, with similar expressions for (a',b), (a,b'), (a',b'), which are all gathered together during the experiment. Then Bell's inequalities tell that $|S| \leq 2$, where S = E(a,b) + E(a',b) + E(a',b') - E(a,b'), in conflict with QM that predicts $S_{max} = 2\sqrt{2}$.

In the references below we quote in italics some relevant sentences extracted from the cited papers. The sentence is meant to represent a crucial thesis in the cited paper, but does not mean that we agree with it.

- [1] C. Budroni, A. Cabello, O. Gühne, M. Kleinmann, J.-A. Larsson, "Kochen-Specker Contextuality", Rev. Mod. Phys. 94, 045007 (2022); https://arxiv.org/abs/2102.13036
- [2] M. Van Den Bossche and P. Grangier, "Revisiting Quantum Contextuality in an Algebraic Framework", J. Phys. Conf. Ser. 2533 (2023) 1, 012008, https://iopscience.iop.org/article/10.1088/1742-6596/2533/1/012008/
- [3] Robert B. Griffiths, "Hilbert Space Quantum Mechanics Is Noncontextual", https://arxiv.org/abs/1201.1510

 It is shown that quantum mechanics is noncontextual if quantum properties are represented by subspaces of the Hilbert space (as proposed by von Neumann) rather than by hidden variables. In particular, a measurement using an appropriately constructed apparatus can be shown to reveal the value of an observable A possessed by the measured system before the measurement took place, whatever other compatible ([B,A]=0) observable B may be measured at the same time.
- [4] László E. Szabó, "Critical reflections on quantum probability theory", in: M. Rédei and M. Stoltzner (eds.), John von Neumann and the Foundations of Quantum Physics, pp 201-219, Kluwer Academic Publishers, 2001.

 My aim is to show in this paper that, beyond its counter intuitiveness, quantum probability theory is inadequate and unnecessary. It is inadequate because there cannot exist events in reality the relative frequencies of which would be equal to quantum probabilities. And it is unnecessary too, because there is no need in quantum mechanics to supersede the Kolmogorov theory of probability; we will see how quantum phenomena can, in general, be accommodated in the classical Kolmogorov theory of probability.
- [5] László E. Szabó, Márton Gömöri, Zalán Gyenis "Questionable and Unquestionable in Quantum Mechanics", https://arxiv.org/abs/2309.01928

 All those investigations remain within the framework of classical Kolmogorovian probability theory, meaning that any physical system (traditionally categorized as classical or quantum) that can be described in operational terms can be described within classical Kolmogorovian probability theory.
- [6] F. H. Holik, "Non-Kolmogorovian Probabilities and Quantum Technologies", Entropy 24(11),1666 (2022) https://www.mdpi.com/1099-4300/24/11/1666
- [7] K. Svozil, "Extending Kolmogorov's Axioms for a Generalized Probability Theory on Collections of Contexts", Entropy 24(9), 1285 (2022) https://www.mdpi.com/1099-4300/24/9/1285
- [8] A nice collection of articles related to the issues discussed here can be found in the book "Quantum, Probability, Logic: The Work and Influence of Itamar Pitowsky", M. Hemmo and O. Shenker editors, Springer Nature Switzerland (2020).
- [9] Gergely Ban and Thomas Durt, "Proof of Kolmogorovian Censorship", Found. Phys. 27:10, 1355 (1997)

 An anonymous referee brought our attention to the fact that (...), although it is impossible to arrange a few noncompatible experiments in a Boolean algebra, compatible algebras always belong to a Boolean algebra. Roughly speaking, when experiments are compatible, their eigenstates can be chosen in such a way that they all belong to the spectral decomposition of a unique observable, and are thus mutually orthogonal. The projectors on arbitrary sets of these eigenstates commute and form a Boolean algebra. According to the referee, the Kolmogorovian censorship expresses, in the formalism of Pitowsky, the fact that "weighted averages (...) of a few noncompatible experiments can be arranged in a single algebra, simply by taking the appropriate mixture on the product space."
- [10] M. Rédei, "Some Historical and Philosophical Aspects of Quantum Probability Theory and its Interpretation". In D. Dieks et al (eds), Probabilities, Laws, and Structures. The Philosophy of Science in a European Perspective, vol 3. Springer, Dordrecht (2012).
 - The idea of Kolmogorovian Censorship hypothesis is that there are in fact no genuinely non-classical probabilities: quantum probabilities are always classical conditional probabilities of outcomes of measurements of quantum observables, where the conditioning events are the events of choosing to set up a measuring device to measure a certain observable.

- [11] A. Auffèves and P. Grangier, "Deriving Born's rule from an Inference to the Best Explanation", Found. Phys. 50, 1781-1793 (2020) [arXiv:1910.13738].
- [12] A. Auffèves and P. Grangier, "Revisiting Born's rule through Uhlhorn's and Gleason's theorems", Entropy 24(2), 199 (2022) https://doi.org/10.3390/e24020199
- [13] A. Aspect, "Closing the Door on Einstein and Bohr's Quantum Debate", Physics 8, 123 (2015) [https://physics.aps.org/articles/v8/123].
- [14] P. Grangier, "Contextual inferences, nonlocality, and the incompleteness of quantum mechanics", Entropy 23 (12), 1660 (2021); https://doi.org/10.3390/e23121660
- [15] T. Durt, "Why God might play dice", Int. J. Theo. Phys. 35:11, 2271 (1996)

 Then our hidden variable would predetermine the result obtained by the freely thinking physicist choosing at random. Such a global determinism is in fact a logically coherent explanation of all the apparently hazardous events occurring in the world and the belief in such a determinism is an old psychological attitude commonly called "fatalism."
- [16] P. Blasiak, E.M. Pothos, J.M. Yearsley, C. Gallus, and E. Borsuk, "Violations of locality and free choice are equivalent resources in Bell experiments", PNAS 118 (17) e2020569118 (2021).
- [17] J. P. Jarrett, "On the physical significance of the locality condition in the Bell arguments", Noûs 18, 569 (1984).
- [18] P. Grangier, "Completing the quantum formalism in a contextually objective framework", Found. Phys. 51, 76 (2021) [arXiv:2003.03121]
- [19] A. Auffèves and P. Grangier, "Contexts, Systems and Modalities: a new ontology for quantum mechanics", Found. Phys. 46, 121 (2016) [arXiv:1409.2120].
- [20] A. Auffèves and P. Grangier, "Extracontextuality and extravalence in quantum mechanics", Phil. Trans. R. Soc. A 376, 20170311 (2018) [arXiv:1801.01398].
- [21] L.D. Landau, E.M. Lifshitz, Quantum Mechanics (1965), available online: https://archive.org/details/ost-physics-landaulifshitz-quantummechanics

 It is clear that, for a system composed only of quantum objects, it would be entirely impossible to construct any logically independent mechanics (...) Thus quantum mechanics occupies a very unusual place among physical theories: it contains classical mechanics as a limiting case, yet at the same time it requires this limiting case for its own formulation.
- [22] M. Van Den Bossche and P. Grangier, "Contextual unification of classical and quantum physics", Found. Phys. 53:45 (2023) [arXiv:2209.01463].
- [23] M. Van Den Bossche and P. Grangier, "Revisiting Quantum Contextuality in an Algebraic Framework", Proceedings of the DICE 2022 Conference, J. Phys. Conf. Ser. 2533 (2023) 1, 012008, https://iopscience.iop.org/article/10.1088/1742-6596/2533/1/012008/
- [24] M. Van Den Bossche and P. Grangier, "Postulating the Unicity of the Macroscopic Physical World", Entropy 25(12), 1600 (2023); https://doi.org/10.3390/e25121600
- [25] M. Gömöri and C. Hoefer, "Classicality and Bell's theorem", Euro. Jnl. Phil. Sci. 13, 45 (2023).