A COMPUTATIONAL APPROACH TO THE HOMOTOPY THEORY OF DG CATEGORIES

DOGANCAN KARABAS AND SANGJIN LEE

ABSTRACT. We give a specific cylinder functor for semifree dg categories. This allows us to construct a homotopy colimit functor explicitly. These two functors are "computable", specifically, the constructed cylinder functor sends a dg category of strictly finite type, i.e., a semifree dg category having finitely many objects and generating morphisms, to a dg category of strictly finite type. The homotopy colimit functor has a similar property. Moreover, using the cylinder functor, we give a cofibration category of semifree dg categories and that of dg categories of strictly finite type, independently from the work of Tabuada [Tab05b]. All the results similarly work for semifree dg algebras. We also describe an application to symplectic topology and provide a toy example.

Contents

1.	Introduction	1
2.	Preliminaries on dg categories	4
3.	Homotopy colimit functor using model structures	7
4.	Cylinder functor for the category of dg categories	11
5.	Homotopy colimit functor on the diagrams of semifree dg categories	17
6.	I-category and cofibration category of semifree dg categories	22
7.	Wrapped Fukaya category of T^*S^n and the reflection functor	32
Re	ferences	40

1. INTRODUCTION

Homotopy theory, originating in algebraic topology, plays a pivotal role in numerous areas of modern mathematics. Specifically, the homotopy theory of differential graded (dg) categories is prominently featured in fields such as algebraic geometry, representation theory, higher categories, and symplectic topology. To explore their homotopy theory up to various weak equivalences, Tabuada [Tab05b, Tab05a] introduced *model structures* for the category of dg categories. These model structures come with auxiliary functors called cofibrations and fibrations, and provide two *functorial factorizations* of functors: the first type factors a functor into a cofibration followed by a weak equivalence, and the second type factors it into a weak equivalence followed by a fibration.

There is another approach to the homotopy theory of dg categories without using a model structure: Starting with cofibrations, instead of considering the entire factorization data, one can focus solely on the functorial factorization of codiagonal maps into a cofibration and a weak equivalence.

Date: May 7, 2024.

This functorial factorization yields a construction of a cylinder functor (Definition 4.1). Once additional axioms (see Definition 6.1) are established, this construction recovers an entire functorial factorization of the first type, representing "half" of a model structure known as a cofibration category in the sense of [Bau89]. Therefore, constructing a cylinder functor allows us to delve into the homotopy theory of dg categories, specifically allowing us to describe the homotopy colimit functor on diagrams of dg categories.

In the current paper, we present a simple construction of a cylinder functor on the category of dg categories, expanding upon our earlier work [KL21] where the cylinder functor was defined solely at the level of objects. Moreover, our cylinder functor leads us to a simple homotopy colimit functor and establishes an *I*-category and a cofibration category of dg categories, offering a computational approach to the homotopy theory of dg categories. Detailed results can be found in Theorems 1-3, which will be discussed following the introduction of our framework. We will also comment on the simplicity of our constructions.

Throughout this paper, we mostly work within the context of the category of semifree dg categories over a commutative ring k, denoted by dgCat_s (see Definition 2.1). We note that every dg category has a semifree resolution (as discussed in [Dri04]), emphasizing the significance of our focus on dgCat_s. Furthermore, we fix weak equivalences as quasi-equivalences, pretriangulated equivalences, or Morita equivalences.

Theorem 1. [Corollary 4.9] Let Cyl: dgCat_s \rightarrow dgCat_s be a functor together with the natural transformations $i_1, i_2: 1_{dgCat_s} \Rightarrow$ Cyl and $p: Cyl \Rightarrow 1_{dgCat_s}$ defined as follows:

- $Cyl(\mathcal{C})$ for any $\mathcal{C} \in dgCat_s$ and the natural transformations are as in Theorem 4.5,
- $\operatorname{Cyl}(F)$ for any morphism (dg functor) $F: \mathcal{A} \to \mathcal{B}$ in dgCat_s is as in Theorem 4.8.

Then, Cyl is a cylinder functor, which means the following conditions are satisfied:

- $p \circ (i_1 \amalg i_2) \colon \mathcal{C} \amalg \mathcal{C} \to \operatorname{Cyl}(\mathcal{C}) \to \mathcal{C}$ is the codiagonal of \mathcal{A} ,
- $i_1 \amalg i_2 \colon \mathcal{C} \amalg \mathcal{C} \to \operatorname{Cyl}(\mathcal{C})$ is a cofibration,
- $p: \operatorname{Cyl}(\mathcal{C}) \to \mathcal{C}$ is a weak equivalence and a fibration.

We note that our cylinder functor differs significantly from the natural cylinder functor induced by Tabuada's model structures. Tabuada employed Quillen's "small object argument" (as detailed in Hovey [Hov07]) to establish a functorial factorization, which results in a cylinder functor as a special case. However, this method (or its refinement by Garner [Gar09]) involves a transfinite construction that is overly complicated from a computational standpoint.

In contrast, our cylinder functor Cyl ensures that the number of generating morphisms of a semifree dg category \mathcal{C} and its image Cyl(\mathcal{C}) are close in size. Specifically, if the former is finite, the latter is also finite, a property that does not hold for Tabuada's natural cylinder. This effectiveness arises from the construction of Cyl, which is specifically tailored for semifree dg categories. Moreover, the computability of our cylinder functor ensures that all our functorial factorizations are computable as can be seen in Theorem 6.9.

Using the cylinder functor described earlier, we can construct the homotopy colimit (pushout) functor for the category of dg categories, where weak equivalences are defined as quasi-equivalences, pretriangulated equivalences, or Morita equivalences:

Theorem 2. [Theorem 5.3] Let J be a category of the form $a \leftarrow c \rightarrow b$, and dgCat^J denote the category of J-diagrams in dgCat. Then, the homotopy colimit functor

hocolim: $\operatorname{Ho}(\operatorname{dgCat}^J) \to \operatorname{Ho}(\operatorname{dgCat})$

can be given such that $\operatorname{hocolim}(\mathcal{C})$ for any $\mathcal{C} \in \operatorname{dgCat}_s$ is defined as in Theorem 5.3(1), and $\operatorname{hocolim}(F)$ for any morphism (dg functor) $F: \mathcal{A} \to \mathcal{B}$ in dgCat_s is defined as in Theorem 5.3(2).

Our homotopy colimit functor is simple in the sense that it produces a semifree dg category of a size close to the total sizes of semifree dg categories in a given diagram. Specifically, the finiteness of the number of generators is preserved under our homotopy colimit functor. In contrast, if we compare this to the Grothendieck construction for homotopy colimit functor (refer to [GPS18]), our method offers several computational advantages. The Grothendieck construction is not explicit; rather, it is expressed as a localization of a non-semifree dg category, which often leads to computational difficulties.

We note that the description in Theorem 2 still holds for the diagrams of the form $\mathcal{A} \leftarrow \mathcal{C} \rightarrow \mathcal{B}$, where \mathcal{A} and \mathcal{B} are not necessarily semifree, if k has flat dimension zero (e.g., if k is a field). See Remark 5.8 for details.

Moreover, the constructions in Theorems 1 and 2 can be adapted for scenarios where the input of the functors is a localization of a semifree dg category. Please refer to Theorems 4.12, 4.13, and 5.7 for more details on these adjustments. The usage of Theorem 5.7 is demonstrated through computations for the case n = 2 in Sections 7.2 and 7.3.

As mentioned earlier, the constructed cylinder functor can define a "half" of a model structure, or more precisely, a cofibration category structure on $dgCat_s$. This structure differs from that induced by [Tab05b] due to its simpler functorial factorization:

Theorem 3.

- (1) (Theorem 6.2) The category of semifree dg categories $dgCat_s$ is an I-category with the structure (cof, I), which is defined as follows:
 - cof : Cofibrations are semifree extensions.
 - I: The functor I is the cylinder functor Cyl: $dgCat_s \rightarrow dgCat_s$ from Theorem 1.
- (2) (Theorem 6.11) The category of semifree dg categories dgCat_s forms a cofibration category, where weak equivalences are quasi-equivalences and cofibrations are semifree extensions. Moreover, every object in dgCat_s is both fibrant and cofibrant. In particular, dgCat_s makes a category of cofibrant objects.

Theorems 1–3 hold in the setting of semifree dg algebras with slight modifications, see Remarks 4.11 and 5.6, and Theorem 6.13.

We note that Theorem 3 is not necessary for constructions such as Theorem 2, as we can use our cylinder functor within Tabuada's model structures to carry out homotopy theory. However, we have established additional axioms for the constructed cylinder functor to explicitly construct a cofibration category of semifree dg categories, which is distinct from the approach in [Tab05b]. In [Tab05b], Quillen's small object argument is employed to construct a model structure and establish the existence of factorizations, but they are not effective or concretely expressible in practice. In contrast, our cylinder functor is explicit and computable, allowing for explicit and computable constructions within the cofibration category dgCat_s. Consequently, we can derive Theorem 2 from Theorem 3 without relying on [Tab05b].

Moreover, we can directly restrict the cofibration category structure given in Theorem 3 to dg categories of strictly finite type, i.e., semifree dg categories with finitely many objects and generating morphisms, as the factorizations preserve finiteness. See Remark 6.14. The same is not true for the model structure of Tabuada [Tab05b], as its given factorizations do not preserve finiteness. We note that dg categories of strictly finite type are, in particular, of finite type. Refer to [Kel07] and [Kon09] for the definition of dg categories of finite type.

DOGANCAN KARABAS AND SANGJIN LEE

One can find a direct application of the above results in symplectic topology.

Symplectic manifolds are associated with a powerful symplectic invariant known as the *Fukaya* category, defined as an A_{∞} -category in [FOOO09a, FOOO09b]. Computing the Fukaya category is typically a notoriously challenging task. However, Ganatra, Pardon, and Shende [GPS20, GPS18] proved that for certain open symplectic manifolds (referred as *Weinstein manifolds*), the (wrapped) Fukaya category can be computed, after a choice of a covering, as a homotopy colimit of the wrapped Fukaya categories of the covering elements.

Furthermore, given the relation of wrapped Fukaya categories to microlocal sheaves by [GPS24], Nadler [Nad17, Nad15] has shown that the wrapped Fukaya category of each covering element can be regarded as a semifree dg category with finitely many objects and generating morphisms (a dg category of strictly finite type) up to pretriangulated (or Morita) equivalence. This result highlights the significance of Theorem 2 in facilitating such computations.

Now, let us consider a symplectomorphism $\phi : W_1 \to W_2$ between two symplectic manifolds. It is known that ϕ induces a functor between Fukaya categories of W_1 and W_2 . If ϕ respects the homotopy colimit diagrams that compute Fukaya categories of W_1 and W_2 , one can obtain a specific description of the induced functor using Theorem 2. For a detailed example and the symplectic topological motivations behind this discussion, please refer to Section 7.

1.1. Acknowledgment. We are grateful to Ezra Getzler for his insightful comments, which inspired the content of Section 6.

The first-named author is supported by World Premier International Research Center Initiative (WPI), MEXT, Japan. The second-named author is supported by a KIAS Individual Grant (MG094401) at Korea Institute for Advanced Study.

2. Preliminaries on DG categories

A differential graded (dg) category is a category enriched over the symmetric monoidal category of complexes over a fixed commutative ring k. It can also be viewed as an A_{∞} -category in which compositions of order greater than 2 are set to vanish. For further details on dg categories, readers may refer to [Kel07], and for a review of A_{∞} -categories, one can consult [Sei08]. We use d for the differential and \circ for compositions of morphisms, and we omit the latter whenever it is convenient. When introducing a dg category, we follow the convention of providing the following five items:

- (i) *Objects:* We list the objects in the category.
- (ii) *Generating morphisms:* We give a set of generating morphisms. They generate all the morphisms as an algebra, not as a module. We will not explicitly mention the existence of identity morphisms, but it should be understood that every object has the identity endomorphism.
- (iii) *Degrees:* For each generating morphism, we specify its degree.
- (iv) Differentials: For each generating morphism, we specify its differential.
- (v) *Relations:* We specify the relations between generating morphisms. This item will be omitted if the generating morphisms freely generate all other morphisms.

Given a dg category \mathcal{C} , we denote by Ob \mathcal{C} (or simply \mathcal{C} when it is clear from the context) the collection of objects in \mathcal{C} , and by Mor \mathcal{C} the collection of morphisms in \mathcal{C} . We use hom^{*}_{\mathcal{C}}(A, B) (or simply hom^{*}(A, B)) to represent the cochain complex of morphisms between the objects A and B of \mathcal{C} .

Next, we introduce a specific class of dg categories and dg functors that will be fundamental throughout the paper. For further details, readers can refer to [KL21].

Definition 2.1.

- (1) A (small) dg category C is called a *semifree dg category* if its morphisms, treated as an algebra, are freely generated by a set of morphisms $\{f_i\}$ (indexed by an ordinal), with the condition that df_i is generated by the set $\{f_j | j < i\}$. In this case, $\{f_i\}$ is called a set of generating morphisms of C.
- (2) A dg functor $F: \mathcal{C} \to \mathcal{D}$ is called a *semifree extension* by a set of objects R and a set of morphisms $S = \{f_i\}$ if it satisfies the following conditions:
 - F is an inclusion.
 - The objects of $F(\mathcal{C})$, along with R, form the objects of \mathcal{D} .
 - The morphisms of \mathcal{D} , treated as an algebra, can be expressed as a free extension of the morphisms of $F(\mathcal{C})$ by $\{f_i\}$ (indexed by an ordinal), with the condition that df_i is generated by the morphisms of $F(\mathcal{C})$ and $\{f_i | j < i\}$.
- (3) A dg category \mathcal{D} is called a *semifree extension* of a dg category \mathcal{C} by a set of objects R and a set of morphisms S if there exists a semifree extension $F: \mathcal{C} \to \mathcal{D}$ as in Definition 2.1(2).

Let dgCat denote the category of small dg categories, where morphisms are dg functors. We aim to invert certain dg functors, referred to as *weak equivalences*, in dgCat. The resulting categories can be studied by introducing *model structures* on dgCat, making dgCat a *model category*. See [DS95], [Hov07], and [Hir03] for a review of model categories. More precisely, upon inverting weak equivalences in dgCat, we obtain the *homotopy category* Ho(dgCat) of the model category dgCat. Refer to Section 3 for further details.

For a given dg category C, we introduce the following notations:

- Tw \mathcal{C} is the dg category of twisted complexes in \mathcal{C} , which is a pretriangulated envelope of \mathcal{C} .
- Perf C is the split-closure (or idempotent completion) of Tw C.

See [Sei08] for more details. With these notations, we can state the following theorem:

Theorem 2.2 ([Tab05b, Tab05a]). The category dgCat admits the following model structures:

- (1) Dwyer-Kan model structure: Weak equivalences are dg functors that are quasi-equivalences, and any dg category is a fibrant object.
- (2) Quasi-equiconic model structure: Weak equivalences are pretriangulated equivalences, which are dg functors $\mathcal{C} \to \mathcal{D}$ that induce a quasi-equivalence $\operatorname{Tw} \mathcal{C} \to \operatorname{Tw} \mathcal{D}$, and fibrant objects are pretriangulated dg categories.
- (3) Morita model structure: Weak equivalences are Morita equivalences, which are dg functors $\mathcal{C} \to \mathcal{D}$ that induce a quasi-equivalence $\operatorname{Perf} \mathcal{C} \to \operatorname{Perf} \mathcal{D}$, and fibrant objects are idempotent complete pretriangulated dg categories.

All three model structures have the same cofibrations, which are retracts of semifree extensions. Consequently, they also share the same cofibrant objects, which are retracts of semifree dg categories.

Remark 2.3. Any quasi-equivalence is a pretriangulated equivalence, and any pretriangulated equivalence is a Morita equivalence.

It is known that any morphism $\mathcal{C} \to \mathcal{D}$ in the homotopy category $Ho(\mathcal{M})$ of a model category \mathcal{M} can be seen as a chain of objects and morphisms in \mathcal{M}

$$\mathcal{C} \xleftarrow{\sim} \mathcal{C}' \to \mathcal{D}' \xleftarrow{\sim} \mathcal{D}$$

for some cofibrant object \mathcal{C}' and fibrant object \mathcal{D}' , and arrows \leftarrow are weak equivalences. Consequently, we can characterize the morphisms in Ho(dgCat) through the following proposition:

Proposition 2.4. For given dg categories C and D, a morphism $C \to D$ in Ho(dgCat) can be characterized as a chain of dg categories and dg functors in the following ways:

- (1) $\mathcal{C} \xleftarrow{\sim} \mathcal{C}' \to \mathcal{D}$, if dgCat is equipped with the Dwyer-Kan model structure.
- (2) $\mathcal{C} \xleftarrow{\sim} \mathcal{C}' \to \operatorname{Tw} \mathcal{D}$, if dgCat is equipped with the quasi-equiconic model structure.
- (3) $\mathcal{C} \xleftarrow{\sim} \mathcal{C}' \to \operatorname{Perf} \mathcal{D}$, if dgCat is equipped with the Morita model structure.

Here, \mathcal{C}' is a cofibrant dg category, and $\stackrel{\sim}{\leftarrow}$ is a weak equivalence in the corresponding model structure.

Remark 2.5. If C is a cofibrant dg category, then we can replace each $C \leftarrow C'$ with C in Proposition 2.4.

Next, we introduce three distinct types of equivalency between dg categories, characterized by becoming isomorphic in the corresponding homotopy category Ho(dgCat):

Definition 2.6. Let C and D be dg categories.

(1) \mathcal{C} and \mathcal{D} are quasi-equivalent if there is a chain of dg categories and dg functors

$$\mathcal{C} \xleftarrow{\sim} \mathcal{C}' \xrightarrow{\sim} \mathcal{D}$$

for some dg category \mathcal{C}' , where each dg functor in the chain is a quasi-equivalence.

- (2) \mathcal{C} and \mathcal{D} are pretriangulated equivalent if $\operatorname{Tw} \mathcal{C}$ and $\operatorname{Tw} \mathcal{D}$ are quasi-equivalent.
- (3) \mathcal{C} and \mathcal{D} are *Morita equivalent* if Perf \mathcal{C} and Perf \mathcal{D} are quasi-equivalent.

As a result, we have two distinct types of generations for a dg category, defined as follows:

Definition 2.7. Let C be a dg category. Let $\{L_i\}$ be a collection of objects in C. We say

- (1) $\{L_i\}$ generates C if the full dg subcategory of C with the objects $\{L_i\}$ is pretriangulated equivalent to C,
- (2) $\{L_i\}$ split-generates C if the full dg subcategory of C with the objects $\{L_i\}$ is Morita equivalent to C.

When C is a dg category, and S is a subset of closed degree zero morphisms in C, there exists a dg category $C[S^{-1}]$, known as the *dg localization* of C at the morphisms in S. This localization is essentially obtained from C by inverting morphisms in S. For a precise definition, one can refer to sources such as [Toë11] or [KL21]. The dg localization is unique up to quasi-equivalence, and its existence is established in [Toë07].

In the case where \mathcal{C} is a semifree dg category, we can explicitly describe $\mathcal{C}[S^{-1}]$:

Proposition 2.8 ([KL21]). When C is a semifree dg category, and $S = \{g_i : A_i \to B_i\}$ is a subset of closed degree zero morphisms in C, the dg localization $C[S^{-1}]$ can be viewed as the semifree extension of C by the morphisms $g'_i, \hat{g}_i, \bar{g}_i$

$$\hat{g}_i
ightarrow A_i \xrightarrow{g_i \\ g_i \\ g_i'} B_i
ightarrow \check{g}_i$$

for each i, with the gradings

$$|g'_i| = 0,$$
 $|\hat{g}_i| = |\check{g}_i| = -1,$ $|\bar{g}_i| = -2,$

and with the differentials

 $dg'_i = 0,$ $d\hat{g}_i = 1_{A_i} - g'_i g_i,$ $d\check{g}_i = 1_{B_i} - g_i g'_i,$ $d\bar{g}_i = g_i \hat{g}_i - \check{g}_i g_i.$

Next, we explore the colimit of diagrams of dg categories:

Proposition 2.9 ([KL21]). Let $G: \mathcal{A} \to \mathcal{C}$ be a dg functor, and $F: \mathcal{A} \to \mathcal{B}$ be a semifree extension by a set of objects R and a set of morphisms S. Then, there exists a pushout (colimit) square

$$\begin{array}{c} \mathcal{B} & \stackrel{G}{\longrightarrow} \mathcal{D} \\ F & & \bar{F} \\ \mathcal{A} & \stackrel{G}{\longrightarrow} \mathcal{C} \end{array}$$

where $\overline{F}: \mathcal{C} \to \mathcal{D}$ is a semifree extension by the set of objects R and a set of morphisms

$$S := \{ f \colon G(A) \to G(B) \, | \, f \colon A \to B \text{ in } S \}$$

with $|\bar{f}| := |f|$ and $d\bar{f} := \bar{G}(df)$, and

$$\bar{G}(A) := \begin{cases} G(A) & \text{if } A \in \operatorname{Ob} \mathcal{A} \\ A & \text{if } A \in R \end{cases}, \quad and \quad \bar{G}(f) := \begin{cases} G(f) & \text{if } f \in \operatorname{Mor} \mathcal{A} \\ \bar{f} & \text{if } f \in S \end{cases}.$$

Remark 2.10. In a more casual sense, the colimit \mathcal{D} in Proposition 2.9 can be thought as $\mathcal{B} \amalg \mathcal{C}$ after the identification of the images of F with the images of G.

Remark 2.11. Given a semifree dg category C with a set of generating morphisms $\{f_i\}$, consider a morphism $f: A \to B$ in $\{f_i\}$ where $A \neq B$. The dg localization $C[f^{-1}]$ can be given by C with the identifications A = B and $f = 1_{A=B}$. This description relies on a description of dg localization through a colimit diagram, as presented in [Toë07], and Proposition 2.9.

Finally, we present two propositions from [Che02] and [EN15], which can be thought as "basis change" and "cancellation of generators" for the morphisms of semifree dg categories, respectively. They are useful when we simplify a given semifree dg category.

Proposition 2.12. Let C be a semifree dg category with a set of generating morphisms $\{f_i\}$ (indexed by an ordinal). Define the morphisms

$$f_i := u_i f_i + g_i$$

where u_i is a unit in the coefficient ring k, and g_i is a morphism in C generated by the set $\{f_j | j < i\}$. Then, the set $\{\tilde{f}_i\}$ also generates the morphisms in C semifreely.

Proposition 2.13. Let C be a semifree dg category, and D be the semifree extension of C by the morphisms $\{a_i, b_i\}$ such that $da_i = b_i$ for all i. Then, C and D are quasi-equivalent.

In the setting of Proposition 2.13, we say \mathcal{D} is obtained from \mathcal{C} by *stabilization*, and \mathcal{C} is obtained from \mathcal{D} by *destabilization*.

3. Homotopy colimit functor using model structures

This section is a review of the homotopy colimit functor and related model-theoretical results. Our main reference is [DS95].

For a given model category \mathcal{M} (more generally, a category \mathcal{M} with weak equivalences), we write Ho(\mathcal{M}) for its *homotopy category*, which is the category

- whose objects are the same as the objects of \mathcal{M} , and
- whose morphisms are generated by the morphisms of \mathcal{M} and the formal inverses of the weak equivalences.

It comes with the *localization functor*

$$l: \mathcal{M} \to \operatorname{Ho}(\mathcal{M})$$

which is the identity on objects, and sending morphisms to themselves.

From now on, let J be the category given by

 $a \leftarrow c \rightarrow b$

where a, b, c are the objects and the arrows are the morphisms.

Definition 3.1. Let \mathcal{M} be a category with weak equivalences, and \mathcal{M}^J be the category of functors $J \to \mathcal{M}$ (*J*-diagrams in \mathcal{M}) whose weak equivalences are the objectwise weak equivalences. The homotopy colimit functor

hocolim:
$$\operatorname{Ho}(\mathcal{M}^J) \to \operatorname{Ho}(\mathcal{M})$$

is defined (up to natural equivalence) as the total left derived functor of the colimit functor

colim: $\mathcal{M}^J \to \mathcal{M}$.

If \mathcal{M} has a model structure, we have a more concrete way to express the homotopy colimit functor. To describe it, we consider an induced model structure on \mathcal{M}^J from the model structure on \mathcal{M} :

Proposition 3.2 ([Hir03], [Dug08]). Let \mathcal{M} be a model category.

(1) \mathcal{M}^J has a model structure, called a Reedy model structure, whose cofibrant objects are the diagrams of the form


$$\mathcal{A} \xleftarrow{\alpha} \mathcal{C} \xrightarrow{\beta} \mathcal{B}$$

where $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are cofibrant objects in \mathcal{M} , and β is a cofibration.

(2) If \mathcal{M}^J is equipped with the Reedy model structure above, then colim: $\mathcal{M}^J \to \mathcal{M}$ preserves cofibrations and acyclic cofibrations.

Before describing the homotopy colimit functor using model structures, we need to define cofibrant resolution functors:

Proposition 3.3 ([DS95]). Let \mathcal{M} be a model category. For any $X \in \mathcal{M}$, there exists a cofibrant object $Q(X) \in \mathcal{M}$ and an acyclic fibration $p_X : Q(X) \to X$ by model category axioms. Then for any morphism $f : X \to Y$, there exists a unique morphism $\tilde{f} : Q(X) \to Q(Y)$ up to right homotopy such that the diagram



commutes.

Definition 3.4. Let \mathcal{M} be a model category.

(1) We define \mathcal{M}_c as the full subcategory of \mathcal{M} consisting of cofibrant objects, and $\pi \mathcal{M}_c$ as the category with the same objects as \mathcal{M}_c whose morphisms are right homotopy classes of morphisms in \mathcal{M}_c .

(2) For any $X \in \mathcal{M}$, there exists a cofibrant object $Q(X) \in \mathcal{M}$ and an acyclic fibration $p_X: Q(X) \to X$ by model category axioms. Then we define a *cofibrant resolution func*tor as

$$Q: \mathcal{M} \to \pi \mathcal{M}_c$$

$$X \mapsto Q(X) \qquad \text{on objects}$$

$$f \mapsto [\tilde{f}] \qquad \text{on morphisms}$$

where given $f: X \to Y$, the morphism \tilde{f} is determined by Proposition 3.3, and $[\tilde{f}]$ is the right homotopy class of \tilde{f} . The well-definedness of this functor follows from Proposition 3.3.

Now, we present the alternative description of the homotopy colimit functor:

Theorem 3.5 ([DS95]). Let \mathcal{M} be a model category, and equip \mathcal{M}^J with the Reedy model structure induced by \mathcal{M} as in Proposition 3.2. The unique lift of the composition

$$\mathcal{M}^J \xrightarrow{Q} \pi(\mathcal{M}^J)_c \xrightarrow{l \circ \operatorname{colim}} \operatorname{Ho}(\mathcal{M})$$


gives the homotopy colimit functor

hocolim:
$$\operatorname{Ho}(\mathcal{M}^J) \to \operatorname{Ho}(\mathcal{M})$$

where Q is a cofibrant resolution functor.

Remark 3.6.

- (1) Although the colimit functor is not well-defined on $\pi(\mathcal{M}^J)_c$, the functor $l \circ \operatorname{colim}$ above is well-defined since it identifies right homotopic maps between cofibrant objects. This follows from the second item in Proposition 3.2. See [DS95] for the details.
- (2) By the lift, we mean that the triangle



commutes. The existence and uniqueness follow from the fact that $l \circ \operatorname{colim} \circ Q$ sends weak equivalences to isomorphisms, which again follows from the second item in Proposition 3.2.

Before presenting a corollary of Theorem 3.5, we describe a functor transforming a given diagram to a more manageable one:

Proposition 3.7.

(1) There exists a functor $T: \mathcal{M}^J \to \mathcal{M}^J$ such that • T sends an object of the form

$$X := (\mathcal{A} \xleftarrow{\alpha} \mathcal{C} \xrightarrow{\beta} \mathcal{B})$$

to the object

$$T(X) := (\mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} \xrightarrow{\nabla_{\mathcal{C}}} \mathcal{C})$$

where $\nabla_{\mathcal{C}}$ is the codiagonal for \mathcal{C} , and

• T sends a morphism of the form

$$\begin{array}{cccc} X & & \mathcal{A} \xleftarrow{\alpha} & \mathcal{C} \xrightarrow{\beta} & \mathcal{B} \\ \downarrow_{F} & := & \downarrow_{F_{\mathcal{A}}} & \downarrow_{F_{\mathcal{C}}} & \downarrow_{F_{\mathcal{B}}} \\ X' & & \mathcal{A}' \xleftarrow{\alpha'} & \mathcal{C}' \xrightarrow{\beta'} & \mathcal{B}' \end{array}$$

to the morphism

$$\begin{array}{cccc} T(X) & & \mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} \xrightarrow{\nabla_{\mathcal{C}}} \mathcal{C} \\ \downarrow_{T(F)} & := & & \downarrow_{F_{\mathcal{A}}\amalg F_{\mathcal{B}}} & \downarrow_{F_{\mathcal{C}}\amalg F_{\mathcal{C}}} & \downarrow_{F_{\mathcal{C}}} \\ T(X') & & \mathcal{A}' \amalg \mathcal{B}' \xleftarrow{\alpha'\amalg \beta'} \mathcal{C}' \amalg \mathcal{C}' \xrightarrow{\nabla_{\mathcal{C}'}} \mathcal{C}' \end{array}$$

(2) The colimit functor satisfies

 $\operatorname{colim} \circ T = \operatorname{colim}.$

(3) T sends weak equivalences to weak equivalences, hence it induces the functor

$$\operatorname{Ho}(T)\colon \operatorname{Ho}(\mathcal{M}^J)\to \operatorname{Ho}(\mathcal{M}^J)$$

satisfying

 $\operatorname{hocolim} \circ \operatorname{Ho}(T) = \operatorname{hocolim}.$

Proof. It is straightforward to check.

Using Proposition 3.7, we have a slight improvement of Theorem 3.5 for constructing the homotopy colimit functor, which only requires us to construct a cofibrant resolution functor Q for the image of the functor T. This will be useful in Section 5.

Corollary 3.8. Let \mathcal{M} be a model category, and equip \mathcal{M}^J with the Reedy model structure induced by \mathcal{M} as in Proposition 3.2. The unique lift of the composition


$$\mathcal{M}^J \xrightarrow{T} \mathcal{M}^J \xrightarrow{Q} \pi(\mathcal{M}^J)_c \xrightarrow{l \circ \operatorname{colim}} \operatorname{Ho}(\mathcal{M})$$

gives the homotopy colimit functor

hocolim:
$$\operatorname{Ho}(\mathcal{M}^J) \to \operatorname{Ho}(\mathcal{M})$$

where T is the functor described in Proposition 3.7, and Q is a cofibrant resolution functor.

Proof. The statement follows from the commutativity of the diagram



by Theorem 3.5 and Proposition 3.7.

4. Cylinder functor for the category of DG categories

Our goal is to give a formula for the homotopy colimit functor hocolim: $\operatorname{Ho}(\mathcal{M}^J) \to \operatorname{Ho}(\mathcal{M})$ for the case $\mathcal{M} = \operatorname{dgCat}$, where dgCat is the model category of small k-linear dg categories for a commutative ring k with Dwyer-Kan, quasi-equiconic, or Morita model structure. Their weak equivalences are quasi-equivalences, pretriangulated equivalences, and Morita equivalences, respectively (see Theorem 2.2). Our formula will work for all these model structures. To express the formula, first we need to describe a cofibrant resolution functor, which will be done in Section 5.

Here, we will define a cylinder functor for the semifree dg categories in dgCat, in other words, we will give a functorial construction of cylinder objects for semifree dg categories. We will use this construction to describe a cofibrant resolution functor in Section 5. Also, in Section 6, the cylinder functor will play a pivotal role in defining an I-category and a cofibration category of semifree dg categories.

We first recall the definition of a cylinder functor:

Definition 4.1. Let \mathcal{M} be a model category. A functor $I: \mathcal{M} \to \mathcal{M}$ is called a *cylinder functor*, if there exists a cofibration

$$i_{\mathcal{C}} \colon \mathcal{C} \amalg \mathcal{C} \to I(\mathcal{C})$$

and an acyclic fibration

 $p_{\mathcal{C}} \colon I(\mathcal{C}) \to \mathcal{C}$

such that $p_{\mathcal{C}} \circ i_{\mathcal{C}} = \nabla_{\mathcal{C}}$ (the codiagonal of \mathcal{C}) for all $\mathcal{C} \in \mathcal{M}$ (in other words, $I(\mathcal{C})$ is a cylinder object for \mathcal{C}), and the diagram

(4.1)

$$\begin{array}{cccc}
\mathcal{C} \amalg \mathcal{C} & \stackrel{F \amalg F}{\longrightarrow} & \mathcal{D} \amalg \mathcal{D} \\
\downarrow_{i_{\mathcal{C}}} & & \downarrow_{i_{\mathcal{D}}} \\
I(\mathcal{C}) & \stackrel{I(F)}{\longrightarrow} & I(\mathcal{D}) \\
\downarrow_{p_{\mathcal{C}}} & & \downarrow_{p_{\mathcal{D}}} \\
\mathcal{C} & \stackrel{F}{\longrightarrow} & \mathcal{D}
\end{array}$$

commutes for any morphism $F: \mathcal{C} \to \mathcal{D}$ in \mathcal{M} .

From now on, we focus on $\mathcal{M} = \text{dgCat}$ with Dwyer-Kan model structure. Everything here still holds if we work with quasi-equiconic or Morita model structure. Before defining a cylinder functor, recall that [KL21] defined a cylinder *object* Cyl(\mathcal{C}) for any semifree dg category \mathcal{C} :

Definition 4.2 ([KL21]). Let C be a semifree dg category, and $i_1, i_2 \colon C \to C \amalg C$ be the inclusions to the first and second copies, respectively. We define Cyl(C) as the semifree extension of $C \amalg C$ by the morphisms comprised of

• the morphisms $t_C, t'_C, \tilde{t}_C, \tilde{t}_C, \bar{t}_C$

$$i_C \stackrel{\overline{i_C}}{\longrightarrow} i_1(C) \xrightarrow{t_C} i_2(C) \xrightarrow{t_C} i_C$$

for each $C \in \mathcal{C}$, with the gradings

$$|t_C| = |t'_C| = 0, \qquad |\hat{t}_C| = |\check{t}_C| = -1, \qquad |\bar{t}_C| = -2,$$

and with the differentials

 $dt_C = dt'_C = 0, \qquad d\hat{t}_C = 1_{i_1(C)} - t'_C \circ t_C, \qquad d\check{t}_C = 1_{i_2(C)} - t_C \circ t'_C, \qquad d\bar{t}_C = t_C \circ \hat{t}_C - \check{t}_C \circ t_C,$ • degree |f| - 1 morphism $t_f : i_1(A) \to i_2(B)$ for each generating morphism $f \in \hom^*_C(A, B),$

• degree |f| - 1 morphism $t_f: i_1(A) \to i_2(B)$ for each generating morphism $f \in \hom_{\mathcal{C}}(A, B)$, with the differential

$$dt_f = (-1)^{|f|} (i_2(f) \circ t_A - t_B \circ i_1(f)) + \text{correction term}$$

where the correction term is 0 if df = 0.

If $df \neq 0$, the correction term is given as follows: If

$$df = c1_A + \sum_{i=1}^m c_i f_{i,n_i} \circ \ldots \circ f_{i,j} \circ \ldots \circ f_{i,1}$$

where $f_{i,j}$ are generating morphisms of C, and $c, c_i \in k$ (c = 0 if $A \neq B$), then

correction term =
$$\sum_{i=1}^{m} c_i \sum_{j=1}^{n_i} (-1)^{|f_{i,j-1}| + \dots + |f_{i,1}|} i_2(f_{i,n_i}) \dots i_2(f_{i,j+1}) \circ t_{f_{i,j}} \circ i_1(f_{i,j-1}) \dots i_1(f_{i,1}).$$

Remark 4.3. The semifree dg category $Cyl(\mathcal{C})$, associated with a given semifree dg category \mathcal{C} , is well-defined up to isomorphism. In other words, it does not depend on the choice of generating morphisms of \mathcal{C} . This can be verified directly or by considering Remark 4.10.

Remark 4.4. By the description of the dg localization in Proposition 2.8, $\operatorname{Cyl}(\mathcal{C})$ is the dg localization $\operatorname{Cyl}_0(\mathcal{C})[S^{-1}]$ (up to quasi-equivalence), where $\operatorname{Cyl}_0(\mathcal{C})$ is the semifree extension of $\mathcal{C} \amalg \mathcal{C}$ by the morphisms comprised of

- closed degree zero morphism $t_C: i_1(C) \to i_2(C)$ for each $C \in \mathcal{C}$,
- degree |f| 1 morphism $t_f: i_1(A) \to i_2(B)$ for each generating morphism $f \in \hom^*_{\mathcal{C}}(A, B)$, with the differential given in Definition 4.2,

and $S = \{t_C \mid C \in \mathcal{C}\}.$

Theorem 4.5 ([KL21]). Let C be a semifree dg category. Cyl(C) defined in Definition 4.2 is a cylinder object for C. That is, the semifree extension

$$i_{\mathcal{C}} := i_1 \amalg i_2 \colon \mathcal{C} \amalg \mathcal{C} \to \operatorname{Cyl}(\mathcal{C})$$

is a cofibration, and the functor

$$p_{\mathcal{C}} \colon \operatorname{Cyl}(\mathcal{C}) \to \mathcal{C}$$

$$i_1(C), i_2(C) \mapsto C, \quad t_C, t'_C \mapsto 1_C, \quad \hat{t}_C, \bar{t}_C, \bar{t}_C \mapsto 0 \quad \text{for each object } C \in \mathcal{C}$$

$$i_1(f), i_2(f) \mapsto f, \quad t_f \mapsto 0 \quad \text{for each generating morphism } f \text{ in } \mathcal{C}$$

is an acyclic fibration, and they satisfy $p_{\mathcal{C}} \circ i_{\mathcal{C}} = \nabla_{\mathcal{C}}$.

So, Theorem 4.5 builds the cylinder functor on objects of dgCat, but not on morphisms. The functoriality of the cylinder object construction is not discussed in [KL21]. Here, we will make the construction functorial. This is the main goal of this section. First, we need some definitions and properties.

Given a semifree dg category \mathcal{C} , Definition 4.2 defines $\operatorname{Cyl}(\mathcal{C})$ and for each generating morphism f of \mathcal{C} , it specifies a generating morphism t_f of $\operatorname{Cyl}(\mathcal{C})$. Here, we extend this and define a morphism t_θ in $\operatorname{Cyl}(\mathcal{C})$ for each morphism θ in \mathcal{C} , which will be useful later:

Definition 4.6. Let \mathcal{C} be a semifree dg category. Let $\theta \in \hom^{\ast}_{\mathcal{C}}(A, B)$ be given by

$$\theta = c1_A + \sum_{i=1}^m c_i f_{i,n_i} \circ \ldots \circ f_{i,j} \circ \ldots \circ f_{i,1}$$

for some generating morphisms $f_{i,j}$ of C and $c, c_i \in k$ (c = 0 if $A \neq B$). We define degree $|\theta| - 1$ morphism $t_{\theta} \in \hom^*_{Cvl(\mathcal{C})}(i_1(A), i_2(B))$ by

$$t_{\theta} := \sum_{i=1}^{m} c_i \sum_{j=1}^{n_i} (-1)^{|f_{i,j-1}| + \dots + |f_{i,1}|} i_2(f_{i,n_i}) \circ \dots \circ i_2(f_{i,j+1}) \circ t_{f_{i,j}} \circ i_1(f_{i,j-1}) \circ \dots \circ i_1(f_{i,1})$$

Proposition 4.7. Let C be a semifree dg category, and let $\theta \in \hom^*_{\mathcal{C}}(A, B)$. We have the identity

(4.2)
$$dt_{\theta} = (-1)^{|\theta|} (i_2(\theta) \circ t_A - t_B \circ i_1(\theta)) + t_{d\theta}.$$

Moreover, if

(4.3)
$$\theta = c1_A + \sum_{i=1}^m c_i \theta_{i,n_i} \circ \ldots \circ \theta_{i,j} \circ \ldots \circ \theta_{i,1}$$

for some morphisms $\theta_{i,j}$ in C (not necessarily generating morphisms) and $c, c_i \in k$ (c = 0 if $A \neq B$), we have the identity

$$(4.4) t_{\theta} = \sum_{i=1}^{m} c_i \sum_{j=1}^{n_i} (-1)^{|\theta_{i,j-1}| + \dots + |\theta_{i,1}|} i_2(\theta_{i,n_i}) \circ \dots \circ i_2(\theta_{i,j+1}) \circ t_{\theta_{i,j}} \circ i_1(\theta_{i,j-1}) \circ \dots \circ i_1(\theta_{i,1}).$$

Proof. First, we remark that (4.2) holds by definition when θ is a generating morphism. Now, assume θ is as in Definition 4.6. Then the assignment $\theta \mapsto t_{\theta}$ is linear. Hence, it is enough to prove (4.2) for $\theta = 1_A$ and $\theta = f_n \circ \ldots \circ f_1$ for any generating morphisms f_i of C. This is straightforward to check using the remark in the beginning of the proof.

Now, assume θ is given as in (4.3). By the linearity of the assignment $\theta \mapsto t_{\theta}$, we only need to prove (4.4) for $\theta = \theta_n \circ \ldots \circ \theta_1$ for any given morphisms θ_i . Furthermore, again by the linearity, we can assume that each θ_i is given as a product of generating morphisms f_j of C. Then, it is straightforward to check that (4.4) holds.

We are now ready to state one of the main results of this section:

Theorem 4.8. Let $F: \mathcal{C} \to \mathcal{D}$ be a dg functor between semifree dg categories.

(1) There is a dg functor $\operatorname{Cyl}(F)$: $\operatorname{Cyl}(\mathcal{C}) \to \operatorname{Cyl}(\mathcal{D})$ that is an extension of the dg functor $F \amalg F : \mathcal{C} \amalg \mathcal{C} \to \mathcal{D} \amalg \mathcal{D}$

$$\begin{split} i.e. \\ \mathrm{Cyl}(F)\colon \mathrm{Cyl}(\mathcal{C}) &\to \mathrm{Cyl}(\mathcal{D}) \\ i_1(C), i_2(C) &\mapsto i_1(F(C)), i_2(F(C)) \\ i_1(\theta), i_2(\theta) &\mapsto i_1(F(\theta)), i_2(F(\theta)) \\ by \ additionally \ specifying \\ t_C, t'_C, \hat{t}_C, \check{t}_C, \check{t}_C &\mapsto t_{F(C)}, t'_{F(C)}, \hat{t}_{F(C)}, \bar{t}_{F(C)}, \bar{t}_{F(C)} \\ t_f &\mapsto t_{F(f)} \\ \end{split}$$

where $t_{F(f)}$ is defined according to Definition 4.6.

(2) The dg functor Cyl(F) satisfies

(4.5)
$$\operatorname{Cyl}(F)(t_{\theta}) = t_{F(\theta)}$$

for any morphism θ in C.

(3) The diagram

(4.6)
$$\begin{array}{ccc} \mathcal{C} \amalg \mathcal{C} & \xrightarrow{F \amalg F} \mathcal{D} \amalg \mathcal{D} \\ & \downarrow^{i_{\mathcal{C}}} & & \downarrow^{i_{\mathcal{D}}} \\ \operatorname{Cyl}(\mathcal{C}) & \xrightarrow{\operatorname{Cyl}(F)} \operatorname{Cyl}(\mathcal{D}) \\ & \downarrow^{p_{\mathcal{C}}} & & \downarrow^{p_{\mathcal{D}}} \\ & \mathcal{C} & \xrightarrow{F} & \mathcal{D} \end{array}$$

commutes.

Proof. We need to show that $\operatorname{Cyl}(F)$ is indeed a dg functor (it is already a functor by definition). The only nontrivial part is showing that $d(\operatorname{Cyl}(F)(t_f)) = \operatorname{Cyl}(F)(dt_f)$ for any generating morphism f in \mathcal{C} . To verify this, we need to prove the identity

$$\operatorname{Cyl}(F)(t_{\theta}) = t_{F(\theta)}$$

for any morphism θ in C. Assume without loss of generality that θ is given by

$$\theta = f_n \circ \ldots \circ f_1$$

for some generating morphisms f_j of \mathcal{C} . Then, since $\operatorname{Cyl}(F)$ is a functor, we have

$$Cyl(F)(t_{\theta}) = Cyl(F) \left(\sum_{j=1}^{n} (-1)^{|f_{j-1}| + \dots + |f_1|} i_2(f_n) \circ \dots \circ i_2(f_{j+1}) \circ t_{f_j} \circ i_1(f_{j-1}) \circ \dots \circ i_1(f_1) \right)$$

=
$$\sum_{j=1}^{n} (-1)^{|f_{j-1}| + \dots + |f_1|} i_2(F(f_n)) \circ \dots \circ i_2(F(f_{j+1})) \circ t_{F(f_j)} \circ i_1(F(f_{j-1})) \circ \dots \circ i_1(F(f_1))$$

=
$$t_{F(\theta)}$$

by the identity (4.4) since $F(\theta) = F(f_n) \circ \ldots \circ F(f_1)$.

Using this and the identity (4.2), for any generating morphism $f: A \to B$ in \mathcal{C} , we see that

$$\begin{aligned} d(\operatorname{Cyl}(F)(t_f)) &= dt_{F(f)} \\ &= (-1)^{|F(f)|} (i_2(F(f)) \circ t_{F(A)} - t_{F(B)} \circ i_1(F(f))) + t_{d(F(f))} \\ &= (-1)^{|f|} (i_2(F(f)) \circ t_{F(A)} - t_{F(B)} \circ i_1(F(f))) + t_{F(df)} \\ &= \operatorname{Cyl}(F)((-1)^{|f|} (i_2(f) \circ t_A - t_B \circ i_1(f)) + t_{df}) \\ &= \operatorname{Cyl}(F)(dt_f) \end{aligned}$$

which shows that Cyl(F) is a dg functor.

Finally, the commutation of the diagram (4.6) is obvious from the construction.

Theorem 4.8 suffices for the purpose of defining the homotopy colimit functor in Section 5. However, we proceed to establish a cylinder functor for semifree dg categories in Corollary 4.9. This construction plays a key role in defining an *I*-category (a category with a natural cylinder functor), thus giving rise to a cofibration category of semifree dg categories in Section 6.

Corollary 4.9. The assignment

$$\begin{array}{l} \text{Cyl: } \mathrm{dgCat}_s \to \mathrm{dgCat}_s \\ \mathcal{C} \mapsto \mathrm{Cyl}(\mathcal{C}) \\ F \mapsto \mathrm{Cyl}(F) \end{array}$$

is a cylinder functor for $dgCat_s$, where $dgCat_s$ is the full subcategory of the model category of dg categories dgCat (with Dwyer-Kan, quasi-equiconic, or Morita model structure) consisting of semifree dg categories, and Cyl(C) is as defined in Definition 4.2, and Cyl(F) is as defined in Theorem 4.8.

Proof. Clearly, $\operatorname{Cyl}(1_{\mathcal{C}}) = 1_{\operatorname{Cyl}(\mathcal{C})}$ where $1_{\mathcal{C}} : \mathcal{C} \to \mathcal{C}$ is the identity functor on \mathcal{C} . Also, the equality $\operatorname{Cyl}(G \circ F) = \operatorname{Cyl}(G) \circ \operatorname{Cyl}(F)$ directly follows from the definition of $\operatorname{Cyl}(F)$ and the identity (4.5), hence Cyl is a functor. Then, the commutative diagram (4.6) shows that Cyl is indeed a cylinder functor in the sense of Definition 4.1.

Remark 4.10. To establish the well-definedness of the functor Cyl up to natural isomorphism, let us denote the application of Cyl to a semifree dg category \mathcal{C} with a predetermined set of generating morphisms $\{f_i\}$ by Cyl $(\mathcal{C}, \{f_i\})$. By applying Cyl to the identity functor $1_{\mathcal{C}}: \mathcal{C} \to \mathcal{C}$, we get an isomorphism

$$\operatorname{Cyl}(1_{\mathcal{C}})\colon \operatorname{Cyl}(\mathcal{C}, \{f_i\}) \xrightarrow{\sim} \operatorname{Cyl}(\mathcal{C}, \{g_i\})$$

for any set of generating morphisms $\{g_i\}$ of \mathcal{C} , which follows from the fact that Cyl is a functor. Hence, $\text{Cyl}(\mathcal{C})$ is well-defined up to isomorphism. Lastly, if there is a functor $F: \mathcal{C} \to \mathcal{D}$ between semifree dg categories, it induces a commutative square

$$\begin{array}{c} \operatorname{Cyl}(\mathcal{C}, \{f_i\}) \xrightarrow{\operatorname{Cyl}(1_{\mathcal{C}})} \operatorname{Cyl}(\mathcal{C}, \{g_i\}) \\ & \downarrow^{\operatorname{Cyl}(F)} & \downarrow^{\operatorname{Cyl}(F)} \\ \operatorname{Cyl}(\mathcal{D}, \{f'_i\}) \xrightarrow{\operatorname{Cyl}(1_{\mathcal{D}})} \operatorname{Cyl}(\mathcal{D}, \{g'_i\}) \end{array}$$

Here, $\{f'_i\}$ and $\{g'_i\}$ are two arbitrary sets of generating morphisms of \mathcal{D} . Hence, Cyl is well-defined up to natural isomorphism.

Remark 4.11. The cylinder object and functor can be also defined for the model category of dg algebras. In that case, for a given semifree dg algebra \mathcal{C} , $\mathcal{C} \amalg \mathcal{C}$ is the semifree dg algebra (with the same unique object as \mathcal{C} by definition) whose generating morphisms are doubled, and $i_1, i_2: \mathcal{C} \to \mathcal{C} \amalg \mathcal{C}$ are the obvious inclusions. Cyl(\mathcal{C}) is defined as the semifree extension of $\mathcal{C} \amalg \mathcal{C}$ by the morphisms t_f , where f is a generating morphism of \mathcal{C} (no $t_C, t'_C, \hat{t}_C, \bar{t}_C, \bar{t}_C$ are used for $C \in \mathcal{C}$). All the formulas still hold by setting

$$t_C = 1$$
, $t'_C = 1$, $\hat{t}_C = 0$, $\check{t}_C = 0$, $\bar{t}_C = 0$.

Given a morphism $F: \mathcal{C} \to \mathcal{D}$ between semifree dg algebras, Cyl(F) is just defined by

$$\operatorname{Cyl}(F)(t_f) = t_{F(f)}$$

for every generating morphism f in C. Hence, this defines a cylinder functor for the model category of dg algebras as in Corollary 4.9. The proofs are the similar in the case of semifree dg algebras.

Finally, we want to discuss the cylinder object for the dg localization $C[S^{-1}]$, where C is a semifree dg category and S is a subset of closed degree zero morphisms in C. We note that since $C[S^{-1}]$ can be also seen as a semifree dg category, one can apply Theorem 4.5 and 4.8 to $C[S^{-1}]$. But we would like to discuss a simpler cylinder object for $C[S^{-1}]$ for later convenience.

By the description of dg localization in Proposition 2.8, we know that $\mathcal{C}[S^{-1}]$ can be express as a semifree extension of \mathcal{C} by the morphisms

$$g' \colon B \to A, \quad \hat{g} \colon A \to A, \quad \check{g} \colon B \to B, \quad \bar{g} \colon A \to B$$

for every $g: A \to B$ in S, whose gradings and differentials are given as in Proposition 2.8. Hence, the generating morphisms of $\mathcal{C}[S^{-1}]$ are the generating morphisms of \mathcal{C} and the morphisms $g', \hat{g}, \check{g}, \check{g}$ for each morphism g in S.

Recall that $i_1, i_2: \mathcal{C} \to \mathcal{C} \amalg \mathcal{C} \hookrightarrow \operatorname{Cyl}(\mathcal{C})$ are the inclusions to the first and second copies, respectively. Then, it is easy to see that $\operatorname{Cyl}(\mathcal{C}[S^{-1}])$ is the semifree extension of $\operatorname{Cyl}(\mathcal{C})[(i_1(S) \sqcup i_2(S))^{-1}]$ by the morphisms

 $t_{g'} \colon i_1(B) \to i_2(A), \quad t_{\hat{g}} \colon i_1(A) \to i_2(A), \quad t_{\check{g}} \colon i_1(B) \to i_2(B), \quad t_{\bar{g}} \colon i_1(A) \to i_2(B)$

for every $g: A \to B$ in S.

Note that $\operatorname{Cyl}(\mathcal{C})[(i_1(S) \sqcup i_2(S))^{-1}]$ is the semifree extension of $\mathcal{C}[S^{-1}] \amalg \mathcal{C}[S^{-1}]$ by the morphisms

- $t_C, t'_C, \tilde{t}_C, \tilde{t}_C, \bar{t}_C$ for each $C \in \mathcal{C}$, and
- t_f for each generating morphism f in C.

In [KL21], it is shown that one can choose a cylinder object for $\mathcal{C}[S^{-1}]$ that is simpler than $\operatorname{Cyl}(\mathcal{C}[S^{-1}])$ in the sense that it does not need the generating morphisms $t_{g'}, t_{\hat{g}}, t_{\tilde{g}}, t_{\tilde{g}}$:

Theorem 4.12 ([KL21]). Let C be a semifree dg category, and S be a subset of closed degree zero morphisms in C. Then $\operatorname{Cyl}(C)[(i_1(S) \sqcup i_2(S))^{-1}]$ is a cylinder object for $C[S^{-1}]$. That is, the semifree extension

$$\mathcal{C}[S^{-1}] := i_1 \amalg i_2 \colon \mathcal{C}[S^{-1}] \amalg \mathcal{C}[S^{-1}] \to \operatorname{Cyl}(\mathcal{C})[(i_1(S) \sqcup i_2(S))^{-1}]$$

is a cofibration, and the functor

$$\begin{array}{ll} p_{\mathcal{C}[S^{-1}]} \colon \mathrm{Cyl}(\mathcal{C})[(i_1(S) \sqcup i_2(S))^{-1}] \to \mathcal{C}[S^{-1}] \\ i_1(C), i_2(C) \mapsto C, \quad t_C, t'_C \mapsto 1_C, \quad \hat{t}_C, \check{t}_C, \bar{t}_C \mapsto 0 & \text{for each object } C \in \mathcal{C} \\ t_f \mapsto 0 & \text{for each generating morphism } f \text{ in } \mathcal{C} \\ i_1(\theta), i_2(\theta) \mapsto \theta & \text{for each morphism } \theta \text{ in } \mathcal{C}[S^{-1}] \end{array}$$

is an acyclic fibration, and they satisfy $p_{\mathcal{C}[S^{-1}]} \circ i_{\mathcal{C}[S^{-1}]} = \nabla_{\mathcal{C}[S^{-1}]}$.

Let \mathcal{C} and \mathcal{D} be a semifree dg categories, and $S_{\mathcal{C}}$ and $S_{\mathcal{D}}$ be subsets of closed degree zero morphisms in \mathcal{C} and \mathcal{D} , respectively. For a given functor $F: \mathcal{C}[S_{\mathcal{C}}^{-1}] \to \mathcal{D}[S_{\mathcal{D}}^{-1}]$, Theorem 4.8 constructs the dg functor

$$\operatorname{Cyl}(F)\colon \operatorname{Cyl}(\mathcal{C}[S_{\mathcal{C}}^{-1}]) \to \operatorname{Cyl}(\mathcal{D}[S_{\mathcal{D}}^{-1}]).$$

However, in such cases, i.e., when a dg category is given as a dg localization, we want to work with its simpler cylinder object given by Theorem 4.12. Hence, we conclude this section with a construction of a dg functor between the simpler cylinder objects:

Theorem 4.13. Let \mathcal{C} and \mathcal{D} be a semifree dg categories, and $S_{\mathcal{C}}$ and $S_{\mathcal{D}}$ be subsets of closed degree zero morphisms in \mathcal{C} and \mathcal{D} , respectively. Let $F : \mathcal{C}[S_{\mathcal{C}}^{-1}] \to \mathcal{D}[S_{\mathcal{D}}^{-1}]$ be a dg functor.

(1) There is a dg functor

$$\operatorname{Cyl}(F) \colon \operatorname{Cyl}(\mathcal{C})[(i_1(S_{\mathcal{C}}) \sqcup i_2(S_{\mathcal{C}}))^{-1}] \to \operatorname{Cyl}(\mathcal{D})[(i_1(S_{\mathcal{D}}) \sqcup i_2(S_{\mathcal{D}}))^{-1}]$$

between cylinder objects of $C[S_{\mathcal{C}}^{-1}]$ and $\mathcal{D}[S_{\mathcal{D}}^{-1}]$ given by Theorem 4.12, which is an extension of the dg functor

$$F \amalg F : \mathcal{C}[S_{\mathcal{C}}^{-1}] \amalg \mathcal{C}[S_{\mathcal{C}}^{-1}] \to \mathcal{D}[S_{\mathcal{D}}^{-1}] \amalg \mathcal{D}[S_{\mathcal{D}}^{-1}]$$

by additionally specifying

$$\begin{array}{ll}t_C, t'_C, \hat{t}_C, \check{t}_C, \bar{t}_C \mapsto t_{F(C)}, t'_{F(C)}, \hat{t}_{F(C)}, \check{t}_{F(C)}, \bar{t}_{F(C)} & respectively, \ for \ each \ object \ C \in \mathcal{C}\\ t_f \mapsto t_{F(f)} & for \ each \ generating \ morphism \ f \ in \ \mathcal{C}\end{array}$$

where $t_{F(f)}$ is defined according to Definition 4.6, and for any $g: A \to B$ in $S_{\mathcal{D}}$, we define the morphisms

$$(4.7) \quad t_{g'} := -i_2(g') \circ t_g \circ i_1(g') - i_2(\hat{g}) \circ t_A \circ i_1(g') + i_2(g') \circ t_B \circ i_1(\check{g}) + i_2(\hat{g} \circ g' - g' \circ \check{g}) \circ t_B \\ t_{\hat{g}} := i_2(g') \circ t_g \circ i_1(\hat{g}) + i_2(\hat{g}) \circ t_A \circ i_1(\hat{g}) + i_2(g') \circ t_B \circ i_1(\bar{g}) - i_2(\hat{g} \circ \hat{g} + g' \circ \bar{g}) \circ t_A \\ - i_2(\hat{g} \circ g' - g' \circ \check{g}) \circ t_g \\ t_{\check{g}} := -i_2(\check{g}) \circ t_g \circ i_1(g') + i_2(\check{g}) \circ t_B \circ i_1(\check{g}) + i_2(\bar{g}) \circ t_A \circ i_1(g') - i_2(\check{g} \circ \check{g} + \bar{g} \circ g') \circ t_B \\ t_{\bar{g}} := -i_2(\check{g}) \circ t_g \circ i_1(\hat{g}) - i_2(\check{g}) \circ t_B \circ i_1(\bar{g}) + i_2(\bar{g}) \circ t_A \circ i_1(\hat{g}) + i_2(\check{g} \circ \bar{g} - \bar{g} \circ \hat{g}) \circ t_A \\ - i_2(\check{g} \circ \check{g} + \bar{g} \circ g') \circ t_B \\ t_{\bar{g}} := -i_2(\check{g}) \circ t_g \circ i_1(\hat{g}) - i_2(\check{g}) \circ t_B \circ i_1(\bar{g}) + i_2(\bar{g}) \circ t_A \circ i_1(\hat{g}) + i_2(\check{g} \circ \bar{g} - \bar{g} \circ \hat{g}) \circ t_A \\ - i_2(\check{g} \circ \check{g} + \bar{g} \circ g') \circ t_g \\ \end{array}$$

in $\operatorname{Cyl}(\mathcal{D})[(i_1(S_{\mathcal{D}}) \sqcup i_2(S_{\mathcal{D}}))^{-1}].$ (2) The dg functor $\operatorname{Cyl}(F)$ satisfies

$$\operatorname{Cyl}(F)(t_{\theta}) = t_{F(\theta)}$$

for any morphism θ in $\mathcal{C}[S_{\mathcal{C}}^{-1}]$.

(3) The diagram

$$\begin{split} \mathcal{C}[S_{\mathcal{C}}^{-1}] \amalg \mathcal{C}[S_{\mathcal{C}}^{-1}] & \xrightarrow{F \amalg F} \mathcal{D}[S_{\mathcal{D}}^{-1}] \amalg \mathcal{D}[S_{\mathcal{D}}^{-1}] \\ & \downarrow^{i_{\mathcal{C}}[S_{\mathcal{C}}^{-1}]} & \downarrow^{i_{\mathcal{D}}[S_{\mathcal{D}}^{-1}]} \\ \operatorname{Cyl}(\mathcal{C})[(i_{1}(S_{\mathcal{C}}) \sqcup i_{2}(S_{\mathcal{C}}))^{-1}] \xrightarrow{\operatorname{Cyl}(F)} \operatorname{Cyl}(\mathcal{D})[(i_{1}(S_{\mathcal{D}}) \sqcup i_{2}(S_{\mathcal{D}}))^{-1}] \\ & \downarrow^{p_{\mathcal{C}}[S_{\mathcal{C}}^{-1}]} & \downarrow^{p_{\mathcal{D}}[S_{\mathcal{D}}^{-1}]} \\ \mathcal{C}[S_{\mathcal{C}}^{-1}] \xrightarrow{F} \mathcal{D}[S_{\mathcal{D}}^{-1}] \end{split}$$

commutes.

Proof. First, it is straightforward (although tedious) to check that (4.2) holds for $t_{g'}, t_{\hat{g}}, t_{\tilde{g}}, t_{\tilde{g}}$ for every morphism g in $S_{\mathcal{D}}$ (and $S_{\mathcal{C}}$). Then, one can similarly prove Proposition 4.7 for the morphisms in $\mathcal{D}[S_{\mathcal{D}}^{-1}]$ (and $\mathcal{C}[S_{\mathcal{C}}^{-1}]$). Hence, the first two items of Theorem 4.13 can be proven similar to Theorem 4.8. The last item, the commutation of the diagram, is straightforward to check.

5. Homotopy colimit functor on the diagrams of semifree DG categories

Our goal in this section is to construct the homotopy colimit functor $Ho(dgCat^J) \rightarrow Ho(dgCat)$, where dgCat is the model category of dg categories with Dwyer-Kan, quasi-equiconic, or Morita model structure (see Theorem 2.2), and J is a category given as follows:

$$a \leftarrow c \rightarrow b$$

Note that we equip $dgCat^J$ with the Reedy model structure as in Proposition 3.2.

To achieve our goal, we will explicitly construct the homotopy colimit functor on $(\text{dgCat}_s)^J$ in Theorem 5.3 and Theorem 5.7, where dgCat_s is the full subcategory of dgCat consisting of semifree dg categories. The category $(\text{dgCat}_s)^J$ can be seen as a subcategory of $\text{Ho}(\text{dgCat}^J)$, and there is a way to lift the homotopy colimit functor on $(\text{dgCat}_s)^J$ to $\text{Ho}(\text{dgCat}^J)$ via the commuting diagram

$$(\mathrm{dgCat}_{s})^{J} \xrightarrow{j} \mathrm{dgCat}^{J} \xrightarrow{T} \mathrm{dgCat}^{J} \xrightarrow{Q} \pi(\mathrm{dgCat}^{J})_{c}$$

$$\downarrow l \qquad \qquad \qquad \downarrow l \circ \operatorname{colim}$$

$$\mathrm{Ho}(\mathrm{dgCat}^{J}) \xrightarrow{\mathrm{hocolim}} \mathrm{Ho}(\mathrm{dgCat})$$

where j is the inclusion functor, T is the functor given in Proposition 3.7, and Q is a cofibrant resolution functor, which we will give in Lemma 5.1. The reason is as follows: hocolim (up to natural equivalence) is the unique lift of the composition $l \circ \operatorname{colim} \circ Q \circ T$ by Corollary 3.8, and each object of dgCat^J is weakly equivalent to an object of (dgCat_s)^J.

Therefore, we only need to describe a cofibrant resolution functor Q on the image of the functor $T \circ j$:

Lemma 5.1. There is a cofibrant resolution functor $Q: (\mathrm{dgCat})^J \to \pi (\mathrm{dgCat}^J)_c$ satisfying the following:

(1) Q sends an object of the form

$$X := (\mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} \xrightarrow{\nabla_{\mathcal{C}}} \mathcal{C})$$

where $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are semifree dg categories and α, β are dg functors, to the object

$$Q(X) := (\mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} \xrightarrow{i_{\mathcal{C}}} \operatorname{Cyl}(\mathcal{C}))$$

where $\operatorname{Cyl}(\mathcal{C})$ is defined as in Definition 4.2 (along with the functors $i_{\mathcal{C}}$ and $p_{\mathcal{C}}$ given in Theorem 4.5).

(2) Q sends a morphism of the form

$$\begin{array}{cccc} X & & \mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} & \stackrel{\nabla_{\mathcal{C}}}{\longrightarrow} \mathcal{C} \\ \downarrow_{F} & := & & \downarrow_{F_{\mathcal{A}}\amalg F_{\mathcal{B}}} & \downarrow_{F_{\mathcal{C}}\amalg F_{\mathcal{C}}} & \downarrow_{F_{\mathcal{C}}} \\ X' & & \mathcal{A}' \amalg \mathcal{B}' \xleftarrow{\alpha'\amalg\beta'} \mathcal{C}' \amalg \mathcal{C}' & \stackrel{\nabla_{\mathcal{C}'}}{\longrightarrow} \mathcal{C}' \end{array}$$

where $F_{\mathcal{A}}, F_{\mathcal{B}}, F_{\mathcal{C}}$ are dg functors, to the morphism

$$Q(X) \qquad \qquad \mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} \xrightarrow{i_{\mathcal{C}}} \operatorname{Cyl}(\mathcal{C}) \\ \downarrow_{Q(F)} \qquad := \qquad \qquad \downarrow_{F_{\mathcal{A}}\amalg F_{\mathcal{B}}} \qquad \downarrow_{F_{\mathcal{C}}\amalg F_{\mathcal{C}}} \qquad \downarrow_{\operatorname{Cyl}(F_{\mathcal{C}})} \\ Q(X') \qquad \qquad \mathcal{A}' \amalg \mathcal{B}' \xleftarrow{\alpha' \amalg \beta'} \mathcal{C}' \amalg \mathcal{C}' \xrightarrow{i_{\mathcal{C}'}} \operatorname{Cyl}(\mathcal{C}') \end{cases}$$

where $\operatorname{Cyl}(F_{\mathcal{C}})$ is defined as in Theorem 4.8.

Proof. For every object X given in the lemma, Q(X) is a cofibrant object by Proposition 3.2 and Theorem 4.5. Moreover, consider the morphism $p_X: Q(X) \to X$ given by

$$Q(X) \qquad \qquad \mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} \xrightarrow{i_{\mathcal{C}}} \operatorname{Cyl}(\mathcal{C}) \\ \downarrow_{p_{X}} \qquad := \qquad \qquad \downarrow_{1_{\mathcal{A}}\amalg 1_{\mathcal{B}}} \qquad \qquad \downarrow_{1_{\mathcal{C}}\amalg 1_{\mathcal{C}}} \qquad \downarrow_{p_{\mathcal{C}}} \\ \mathcal{A} \amalg \mathcal{B} \xleftarrow{\alpha \amalg \beta} \mathcal{C} \amalg \mathcal{C} \xrightarrow{\nabla_{\mathcal{C}}} \mathcal{C}$$

which is indeed a morphism and an acyclic (Reedy) fibration (see [Hir03]) since $p_{\mathcal{C}} \circ i_{\mathcal{C}} = \nabla_{\mathcal{C}}$ and $p_{\mathcal{C}}$ is an acyclic fibration by Theorem 4.5. Then, the diagram

$$Q(X) \xrightarrow{Q(F)} Q(X')$$
$$\downarrow^{p_X} \qquad \qquad \downarrow^{p_{X'}}$$
$$X \xrightarrow{F} \qquad X'$$

commutes for every morphism $F: X \to X'$ given in the lemma since the diagram (4.6) commutes by Theorem 4.8. Note that Q(F) is indeed a morphisms also because the diagram (4.6) commutes. Therefore, by Proposition 3.3 and Definition 3.4, there is a cofibration functor Q with the properties given in the lemma.

Remark 5.2. Assume that in Lemma 5.1, we replace \mathcal{C} and \mathcal{C}' by $\mathcal{C}[S_{\mathcal{C}}^{-1}]$ and $\mathcal{C}'[S_{\mathcal{C}'}^{-1}]$ for some subsets of degree zero closed morphisms $S_{\mathcal{C}}$ and $S_{\mathcal{C}'}$ in \mathcal{C} and \mathcal{C}' , respectively. Then, instead of using $\operatorname{Cyl}(F_{\mathcal{C}[S_{\mathcal{C}}^{-1}]})$: $\operatorname{Cyl}(\mathcal{C}[S_{\mathcal{C}}^{-1}]) \to \operatorname{Cyl}(\mathcal{C}'[S_{\mathcal{C}'}^{-1}])$ when defining Q, we can use the dg functor

$$\operatorname{Cyl}(\mathcal{C})[(i_1(S_{\mathcal{C}}) \sqcup i_2(S_{\mathcal{C}}))^{-1}] \to \operatorname{Cyl}(\mathcal{C}')[(i_1(S_{\mathcal{C}'}) \sqcup i_2(S_{\mathcal{C}'}))^{-1}]$$

given in Theorem 4.13, which gives a simpler cofibrant resolution functor Q. The proof will be the same as the proof of Lemma 5.1 after replacing Theorem 4.8 with Theorem 4.13 in the proof.

Finally, we can state the formula for the homotopy colimit functor on the diagrams of semifree dg categories:

Theorem 5.3. The homotopy colimit functor hocolim: $Ho(dgCat^J) \rightarrow Ho(dgCat)$ (up to natural equivalence) satisfies the following:

(1) hocolim sends an object of the form

$$X := (\mathcal{A} \xleftarrow{\alpha} \mathcal{C} \xrightarrow{\beta} \mathcal{B})$$

where $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are semifree dg categories and α, β are dg functors, to the object (semifree dg category)

hocolim(X)

which is the semifree extension of $\mathcal{A}\amalg\mathcal{B}$ by

- closed degree zero morphism $t_C \colon \alpha(C) \to \beta(C)$ for each object $C \in \mathcal{C}$,
- morphisms $t'_C, \hat{t}_C, \check{t}_C, \check{t}_C$ for each object $C \in \mathcal{C}$ as in Definition 4.2 (after replacing i_1 with α and i_2 with β),
- degree |f| 1 morphism $t_f: \alpha(A) \to \beta(B)$ for each generating morphism $f: A \to B$ in \mathcal{C} whose differential is given as in Definition 4.2 (after replacing i_1 with α and i_2 with β).

(2) hocolim sends a morphism of the form

$$\begin{array}{cccc} X & & \mathcal{A} \xleftarrow{\alpha} & \mathcal{C} \xrightarrow{\beta} & \mathcal{B} \\ \downarrow_{F} & := & \downarrow_{F_{\mathcal{A}}} & \downarrow_{F_{\mathcal{C}}} & \downarrow_{F_{\mathcal{B}}} \\ X' & & \mathcal{A}' \xleftarrow{\alpha'} & \mathcal{C}' \xrightarrow{\beta'} & \mathcal{B}' \end{array}$$

where $F_{\mathcal{A}}, F_{\mathcal{B}}, F_{\mathcal{C}}$ are dg functors, to the morphism (dg functor)

$\operatorname{hocolim}(F) \colon \operatorname{hocolim}(X) \to \operatorname{hocolim}(X')$	
$A\mapsto F_{\mathcal{A}}(A)$	for any object $A \in \mathcal{A}$
$B \mapsto F_{\mathcal{B}}(B)$	for any object $B \in \mathcal{B}$
$a \mapsto F_{\mathcal{A}}(a)$	for any morphism a in \mathcal{A}
$b \mapsto F_{\mathcal{B}}(b)$	for any morphism b in $\mathcal B$
$t_C, t'_C, \hat{t}_C, \check{t}_C, \bar{t}_C \mapsto t_{F_{\mathcal{C}}(C)}, t'_{F_{\mathcal{C}}(C)}, \hat{t}_{F_{\mathcal{C}}(C)}, \check{t}_{F_{\mathcal{C}}(C)}, \bar{t}_{F_{\mathcal{C}}(C)}, \bar{t}_{F_{\mathcal{C}$	respectively, for any object $C \in \mathcal{C}$
$t_f \mapsto t_{F_{\mathcal{C}}(f)}$	for any generating morphism f in C ,

where for any generating morphism $f: A \to B$ in C, the degree |f| - 1 morphism $t_{F_{\mathcal{C}}(f)}$ is defined as

$$t_{F_{\mathcal{C}}(f)} := \sum_{i=1}^{m} c_i \sum_{j=1}^{n_i} (-1)^{|f_{i,j-1}| + \dots + |f_{i,1}|} \beta'(f_{i,n_i}) \circ \dots \circ \beta'(f_{i,j+1}) \circ t_{f_{i,j}} \circ \alpha'(f_{i,j-1}) \circ \dots \circ \alpha'(f_{i,1})$$

if $F_{\mathcal{C}}(f)$ is given by

$$F_{\mathcal{C}}(f) = c1_{F_{\mathcal{C}}(A)} + \sum_{i=1}^{m} c_i f_{i,n_i} \circ \ldots \circ f_{i,j} \circ \ldots \circ f_{i,1} \in \hom_{\mathcal{C}'}^*(F_{\mathcal{C}}(A), F_{\mathcal{C}}(B))$$

for some generating morphisms $f_{i,j}$ of \mathcal{C}' and $c, c_i \in k$ $(c = 0 \text{ if } F_{\mathcal{C}}(A) \neq F_{\mathcal{C}}(B)).$

Proof. First, we note that the description of hocolim(X) is given in [KL21]. We can also see it here as follows: By Corollary 3.8, we have

$$\operatorname{hocolim}(X) = l \circ \operatorname{colim} \circ Q \circ T(X)$$

where T is a functor as in Proposition 3.7, Q is the cofibration functor given in Lemma 5.1. Using the description of colim given in Proposition 2.9, it is straightforward to check that hocolim(X) is as described in the first item.

For the second item (which does not appear in [KL21]), by Corollary 3.8 again, we have

$$\operatorname{hocolim}(F) = l \circ \operatorname{colim} \circ Q \circ T(F).$$

Then, considering that $\operatorname{hocolim}(X)$ is the semifree extension of $\mathcal{A} \amalg \mathcal{B}$ by the morphims given in the theorem, we see that $\operatorname{hocolim}(F)$: $\operatorname{hocolim}(X) \to \operatorname{hocolim}(X')$ acts like $F_{\mathcal{A}}$ on \mathcal{A} , $F_{\mathcal{B}}$ on \mathcal{B} , and $\operatorname{Cyl}(F_{\mathcal{C}})$ (which is given in Theorem 4.8) on the added morphisms. \Box

Remark 5.4. In Theorem 5.3, we could have expressed $\operatorname{hocolim}(X)$ as the semifree dg category obtained by first taking the semifree extension of $\mathcal{A} \amalg \mathcal{B}$ by the morphisms t_C and t_f for each $C \in \mathcal{C}$ and for each generating morphism f in \mathcal{C} , and then taking the dg localization of the resulting category at the morphisms $\{t_C \mid C \in \mathcal{C}\}$ as in Remark 4.4. Hence, up to natural equivalence, the images of the morphisms $t'_C, \hat{t}_C, \tilde{t}_C, \tilde{t}_C$ under the homotopy colimit functor are uniquely determined for every $C \in \mathcal{C}$.

Remark 5.5. Theorem 5.3 implies that for any morphism θ in C, we have

 $\operatorname{hocolim}(F)(t_{\theta}) = t_{F_{\mathcal{C}}(\theta)}.$

This follows from the identity (4.5).

Remark 5.6. Similar to Remark 4.11, we can modify Theorem 5.3 to describe the homotopy colimit functor $Ho(dgAlg^J) \rightarrow Ho(dgAlg)$ for the model category of dg algebras dgAlg. All the formulas still hold by setting

$$t_C = 1, \quad t'_C = 1, \quad \hat{t}_C = 0, \quad \check{t}_C = 0, \quad \bar{t}_C = 0.$$

Considering Theorem 4.13 and Remark 5.2, we have the following version of Theorem 5.3 when C is given as a dg localization:

Theorem 5.7. The homotopy colimit functor hocolim: $Ho(dgCat^J) \rightarrow Ho(dgCat)$ (up to natural equivalence) satisfies the following:

(1) hocolim sends an object of the form

$$X := (\mathcal{A} \stackrel{\alpha}{\leftarrow} \mathcal{C}[S_{\mathcal{C}}^{-1}] \stackrel{\beta}{\to} \mathcal{B})$$

where $\mathcal{A}, \mathcal{B}, \mathcal{C}$ are semifree dg categories, $S_{\mathcal{C}}$ is a subset of degree zero morphisms in \mathcal{C} , and α, β are dg functors, to the object (semifree dg category)

$$\operatorname{hocolim}(X) = \operatorname{hocolim}(\mathcal{A} \xleftarrow{\alpha} \mathcal{C} \xrightarrow{\beta} \mathcal{B})$$

which is given by Theorem 5.3.

(2) hocolim sends a morphism of the form

$$\begin{array}{cccc} X & & \mathcal{A} \xleftarrow{\alpha} \mathcal{C}[S_{\mathcal{C}}^{-1}] \xrightarrow{\beta} \mathcal{B} \\ \downarrow_{F} & := & \downarrow_{F_{\mathcal{A}}} & \downarrow_{F_{\mathcal{C}}} & \downarrow_{F_{\mathcal{B}}} \\ X' & & \mathcal{A}' \xleftarrow{\alpha'} \mathcal{C}'[S_{\mathcal{C}'}^{-1}] \xrightarrow{\beta'} \mathcal{B}' \end{array}$$

where for any generating morphism $f: A \to B$ in C, the degree |f| - 1 morphism $t_{F_{C}(f)}$ is defined as

$$t_{F_{\mathcal{C}}(f)} := \sum_{i=1}^{m} c_i \sum_{j=1}^{n_i} (-1)^{|f_{i,j-1}| + \ldots + |f_{i,1}|} \beta'(f_{i,n_i}) \circ \ldots \circ \beta'(f_{i,j+1}) \circ t_{f_{i,j}} \circ \alpha'(f_{i,j-1}) \circ \ldots \circ \alpha'(f_{i,1})$$

if $F_{\mathcal{C}}(f)$ is given by

$$F_{\mathcal{C}}(f) = c1_{F_{\mathcal{C}}(A)} + \sum_{i=1}^{m} c_i f_{i,n_i} \circ \ldots \circ f_{i,j} \circ \ldots \circ f_{i,1} \in \hom_{\mathcal{C}'[S_{\mathcal{C}'}^{-1}]}^{*}(F_{\mathcal{C}}(A), F_{\mathcal{C}}(B))$$

for some generating morphisms $f_{i,j}$ of $\mathcal{C}'[S_{\mathcal{C}'}^{-1}]$ and $c, c_i \in k$ $(c = 0 \text{ if } F_{\mathcal{C}}(A) \neq F_{\mathcal{C}}(B))$. The generating morphisms of $\mathcal{C}'[S_{\mathcal{C}'}^{-1}]$ are given by the generating morphisms of \mathcal{C}' and the morphisms $g', \hat{g}, \check{g}, \bar{g}$ for every morphism g in $S_{\mathcal{C}'}$ as in Proposition 2.8. The morphisms

 $t_{g'}, t_{\hat{g}}, t_{\check{g}}, t_{\bar{g}}$

are given in terms of t_g as in (4.7) and t_g is given as in Definition 4.6 (after replacing i_1 with α' and i_2 with β').

Proof. The first item appears in [KL21]. We can also see it here as in the proof of Theorem 5.3 after replacing the cofibrant resolution functor Q with the one given in Remark 5.2.

The second item (which does not appear in [KL21]) can be again shown as in the proof of Theorem 5.3 after replacing the cofibrant resolution functor Q with the one given in Remark 5.2. In particular, we replace the functor Cyl(F) given in Theorem 4.8 to define Q with the one given in Theorem 4.13.

Remark 5.8. According to [Hol14], if the coefficient ring k has flat dimension 0 (e.g., k is a field), then the model category dgCat with weak equivalences being quasi-equivalences, pretriangulated equivalences, or Morita equivalences is left proper. This implies that the formulas in Theorems 5.3 and 5.7 still hold if we choose arbitrary dg categories $\mathcal{A}, \mathcal{A}', \mathcal{B}, \mathcal{B}'$ instead of semifree ones, given that k has flat dimension 0.

In Section 7, we will demonstrate a simple application of Theorem 5.3 and Theorem 5.7 to symplectic geometry.

6. I-CATEGORY AND COFIBRATION CATEGORY OF SEMIFREE DG CATEGORIES

The purpose of this section is to elaborate on the construction of the cylinder functor introduced in Section 4 to approach the homotopy theory of dg categories up to quasi-equivalence. This approach is independent of the celebrated result by Tabuada (Theorem 2.2(1)). The advantage of our approach is that the simplicity of our cylinder functor makes many constructions explicit and computable. It is worth noting that readers can skip to Section 7 without losing continuity in the paper.

We will define an *I*-category structure on the category of semifree dg categories in Theorem 6.2 by introducing cofibrations and a natural cylinder functor. This structure enables us to establish a cofibration category of semifree dg categories in Theorem 6.11, following the framework outlined in [Bau89]. Conceptually, this category corresponds to half of a model category. Alternatively, it can be regarded as a category of cofibrant objects first introduced by Brown in [Bro73] and further studied, for example, in [KP97].

These frameworks enable the construction of various homotopical operations, such as homotopy colimits (Remark 6.12), without relying on the existence of a model structure. Thus, it provides an alternative pathway to achieve the most of the results outlined in the preceding sections without relying on [Tab05b]. We will also present an explicit functorial factorization of dg functors into a cofibration and a quasi-equivalence in Theorem 6.9.

We recall that k is a commutative ring, and dgCat_s represents the category of (small) k-linear semifree dg categories, with morphisms defined as dg functors. The definitions of semifree dg categories and semifree extensions can be revisited from Definition 2.1. Note that every dg category is quasi-equivalent to a semifree one, thus justifying our focus on dgCat_s.

We now introduce the definition of an *I*-category as outlined in [Bau89]. Then, we will show how the category of semifree dg categories can be made an *I*-category:

Definition 6.1. An *I*-category is a category \mathcal{M} with an initial object \emptyset and equipped with a structure (cof, *I*). The term cof represents a class of morphisms in \mathcal{M} identified as cofibrations, and *I* is a functor from \mathcal{M} to itself. The structure satisfies the following set of axioms:

(1) Cylinder Axiom: $I: \mathcal{M} \to \mathcal{M}$ is a functor together with natural transformations

$$i_1, i_2 : \operatorname{id}_{\mathcal{M}} \Rightarrow I, \quad p : I \Rightarrow \operatorname{id}_{\mathcal{M}},$$


such that for all $\mathcal{A} \in \mathcal{M}$ and $r \in \{1, 2\}$, $p \circ i_r \colon \mathcal{A} \to I(\mathcal{A}) \to \mathcal{A}$ is the identity of \mathcal{A} .

(2) Pushout Axiom: For a cofibration $F: \mathcal{A} \to \mathcal{B}$ and a morphism $G: \mathcal{A} \to \mathcal{C}$ in \mathcal{M} , there exists a pushout in \mathcal{M}

$$\begin{array}{c} \mathcal{B} & \stackrel{\bar{G}}{\longrightarrow} \mathcal{B} \cup_{\mathcal{A}} \mathcal{C} \\ F & & \bar{F} \\ \mathcal{A} & \stackrel{G}{\longrightarrow} \mathcal{C} \end{array}$$


where \overline{F} is also a cofibration. Moreover, $I(\emptyset) = \emptyset$, and I sends the pushout diagram into a pushout diagram, i.e., $I(\mathcal{B} \cup_{\mathcal{A}} \mathcal{C}) = I(\mathcal{B}) \cup_{I(\mathcal{A})} I(\mathcal{C})$.

(3) Cofibration Axiom: Each isomorphism in \mathcal{M} is a cofibration, and for each $\mathcal{A} \in \mathcal{M}$, the morphism $\emptyset \to \mathcal{A}$ is a cofibration. The composition of cofibrations is a cofibration. Moreover, any cofibration $F: \mathcal{A} \to \mathcal{B}$ in \mathcal{M} has the following homotopy extension property: For each $r \in \{1, 2\}$, and for each commutative diagram in \mathcal{M}



there is a morphism $E: I(\mathcal{B}) \to \mathcal{C}$ with $E \circ i_r = G$ and $E \circ I(F) = H$.

(4) Relative Cylinder Axiom: For a cofibration $F: \mathcal{A} \to \mathcal{B}$ in \mathcal{M} , the morphism G defined by the pushout diagram



is a cofibration.

(5) Interchange Axiom: For all $\mathcal{A} \in \mathcal{M}$, there exists a morphism $T: I(I(\mathcal{A})) \to I(I(\mathcal{A}))$ such that $T \circ i_r = I(i_r)$ and $T \circ I(i_r) = i_r$ for any $r \in \{1, 2\}$.

Theorem 6.2. The category of semifree dg categories $dgCat_s$ is an *I*-category with the structure (cof, I), which is defined as follows:

- cof : Cofibrations are semifree extensions.
- I: The functor I is the cylinder functor Cyl: $dgCat_s \rightarrow dgCat_s$ introduced in Corollary 4.9.

Proof. First, note that the empty category \emptyset is an initial object of dgCat_s. We introduce the following conventions and review some definitions:

- Given a semifree dg category \mathcal{A} , we fix a set of generating morphisms $\{a_i \colon X_i \to Y_i\}$ (indexed by an ordinal). Then, the semifree dg category $Cyl(\mathcal{A})$ is given as follows:
 - (i) Objects: A^1, A^2 for each $A \in \mathcal{A}$.
 - (ii) Generating morphisms: For each object $A \in \mathcal{A}$,

$$t_A \colon A^1 \to A^2, \quad t'_A \colon A^2 \to A^1, \quad \hat{t}_A \colon A^1 \to A^1, \quad \check{t}_A \colon A^2 \to A^2, \quad \bar{t}_A \colon A^1 \to A^2,$$

and for each generating morphism $a_i \colon X_i \to Y_i$ of \mathcal{A} ,

$$a_i^1 \colon X_i^1 \to Y_i^1, \quad a_i^2 \colon X_i^2 \to Y_i^2, \quad t_{a_i} \colon X_i^1 \to Y_i^2.$$

(iii) Degrees:

$$|t_A| = |t'_A| = 0, \quad |\hat{t}_A| = |\check{t}_A| = -1, \quad |\bar{t}_A| = -2, \quad |a_i^1| = |a_i^2| = |a_i|, \quad |t_{a_i}| = |a_i| - 1.$$

(iv) Differentials:

$$dt_A = dt'_A = 0, \quad d\hat{t}_A = 1_{A^1} - t'_A \circ t_A, \quad d\check{t}_A = 1_{A^2} - t_A \circ t'_A, \quad d\bar{t}_A = t_A \circ \hat{t}_A - \check{t}_A \circ t_A, \\ da_i^1 = (da_i)^1, \quad da_i^2 = (da_i)^2, \quad dt_{a_i} = (-1)^{|a_i|} (a_i^2 \circ t_{X_i} - t_{Y_i} \circ a_i^1) + t_{da_i},$$

where t_{da_i} is defined according to Definition 4.6, and for each $r \in \{1, 2\}$, $(da_i)^r$ is defined via the dg functor

$$i_r \colon \mathcal{A} \to \operatorname{Cyl}(\mathcal{A})$$

 $A \mapsto A^r, \quad a_i \mapsto a_i^r$

by setting $a^r := i_r(a)$ for each morphism a of \mathcal{A} .

• Given a semifree extension $F: \mathcal{A} \to \mathcal{B}$ by a set of objects R_b and a set of morphisms $S_b = \{b_i: U_i \to V_i\}$, we select a set of generating morphisms of \mathcal{B} as

 $\{F(a_i) \mid a_i \text{ is a generating morphism of } \mathcal{A}\} \cup S_b.$

It is important to note that the functor Cyl may initially depend on the choice of generating morphisms, but it ultimately does not (up to natural isomorphism), as discussed in Remark 4.10.

• Given a dg functor $G: \mathcal{A} \to \mathcal{C}$ between semifree dg categories, the dg functor Cyl(G) is defined as follows:

$$Cyl(G): Cyl(\mathcal{A}) \to Cyl(\mathcal{C})$$
$$A^r \mapsto G(A)^r, \quad a_i^r \mapsto G(a_i)^r, \quad t_{a_i} \mapsto t_{G(a_i)},$$
$$t_A \mapsto t_{G(A)}, \quad t'_A \mapsto t'_{G(A)}, \quad \hat{t}_A \mapsto \hat{t}_{G(A)}, \quad \check{t}_A \mapsto \check{t}_{G(A)}, \quad \bar{t}_A \mapsto \bar{t}_{G(A)},$$

where $r \in \{1, 2\}$.

With the notations established, we proceed to verify the axioms outlined in Definition 6.1 for $\mathcal{M} = \text{dgCat}_s$ with cofibrations defined as semifree extensions and I = Cyl:

- (1) Cylinder Axiom: The functor I = Cyl satisfies the conditions of Definition 4.1 by Corollary 4.9, thus fulfilling the Cylinder Axiom. Note that Definition 4.1 provides more than what is necessary for the axiom.
- (2) Pushout Axiom: The existence of the pushout in Definition 6.1(2) is given by Proposition 2.9. Also, we have $\operatorname{Cyl}(\emptyset) = \emptyset$ by definition. Moreover, if $F \colon \mathcal{A} \to \mathcal{B}$ is a semifree extension by a set of objects R_b and a set of morphisms S_b , then $\operatorname{Cyl}(F) \colon \operatorname{Cyl}(\mathcal{A}) \to \operatorname{Cyl}(\mathcal{B})$ is a semifree extension by the set of objects $\{B^1, B^2 \mid B \in R_b\}$ and the set of morphisms

$$\{t_B, t'_B, \hat{t}_B, \check{t}_B, \bar{t}_B | B \in R_b\} \cup \{b_i^1, b_i^2, t_{b_i} | b_i \in S_b\}.$$

Then, given a pushout square

$$\begin{array}{c} \mathcal{B} \xrightarrow{\overline{G}} \mathcal{D} \\ F \uparrow & \bar{F} \uparrow \\ \mathcal{A} \xrightarrow{\overline{G}} \mathcal{C} \end{array}$$

as in Proposition 2.9, applying the functor Cyl yields a commutative diagram

$$\begin{array}{c} \operatorname{Cyl}(\mathcal{B}) \xrightarrow{\operatorname{Cyl}(G)} \operatorname{Cyl}(\mathcal{D}) \\ \\ \operatorname{Cyl}(F) & \xrightarrow{\operatorname{Cyl}(F)} \\ \\ \operatorname{Cyl}(\mathcal{A}) \xrightarrow{\operatorname{Cyl}(G)} \operatorname{Cyl}(\mathcal{C}) \end{array}$$

where $\operatorname{Cyl}(\overline{F})$ is a semifree extension, as \overline{F} is. Also, it is straightforward to verify that $\operatorname{Cyl}(\overline{F}) = \overline{\operatorname{Cyl}(F)}$ and $\operatorname{Cyl}(\overline{G}) = \overline{\operatorname{Cyl}(G)}$. Thus, the above diagram forms a pushout square.

(3) Cofibration Axiom: Clearly, each isomorphism in dgCat_s is a semifree extension, and for each $\mathcal{A} \in \text{dgCat}_s$, $\emptyset \to \mathcal{A}$ is a semifree extension. Moreover, the composition of semifree extensions is a semifree extension.

Regarding the homotopy extension property, let us first set r = 1 without loss of generality. Consider the commutative diagram described in Definition 6.1(3), where I = Cyl. Then, we define the dg functor $E: \text{Cyl}(\mathcal{B}) \to \mathcal{C}$ as follows: Since $\text{Cyl}(F): \text{Cyl}(\mathcal{A}) \to \text{Cyl}(\mathcal{B})$ is a semifree extension, we can view $\text{Cyl}(\mathcal{A})$ as a full dg subcategory of $\text{Cyl}(\mathcal{B})$. We define

$$E|_{\operatorname{Cvl}(\mathcal{A})} := H.$$

Moreover, we set

 $E(B^1) = E(B^2) := G(B), \quad E(t_B) = E(t'_B) := 1_{G(B)}, \quad E(\hat{t}_B) = E(\check{t}_B) = E(\bar{t}_B) := 0$ for each $B \in R_b$, and

$$E(b_i^1) := G(b_i)$$

for each $b_i \in S_b$. We will define $E(b_i^2)$ and $E(t_{b_i})$ by transfinite induction: Let us give an ordering on the set of morphisms $S_b = \{b_i : U_i \to V_i\}$ (indexed by an ordinal) such that each db_i in \mathcal{B} is generated by the morphisms of \mathcal{A} and $\{b_j | j < i\}$. Assume that E is defined on the morphisms generated by the morphisms of $Cyl(\mathcal{A})$ and

$$\{t_B, t'_B, \hat{t}_B, \check{t}_B, \bar{t}_B \mid B \in R_b\} \cup \{b_i^1 \mid b_i \in S_b\} \cup \{b_j^2, t_{b_j} \mid j < i\}.$$

Then, we define

$$E(b_i^2) := E(t_{V_i} \circ b_i^1 \circ t'_{U_i} - (-1)^{|b_i|} t_{db_i} \circ t'_{U_i} - (-1)^{|b_i|} (db_i)^2 \circ \check{t}_{U_i}),$$

$$E(t_{b_i}) := E(-t_{V_i} \circ b_i^1 \circ \hat{t}_{U_i} + (-1)^{|b_i|} t_{db_i} \circ \hat{t}_{U_i} - (-1)^{|b_i|} (db_i)^2 \circ \bar{t}_{U_i}).$$

It is straightforward to check that E is a dg functor, as $dE(b_i^2) = E(db_i^2) = E((db_i)^2)$ and $dE(t_{b_i}) = E(dt_{b_i}) = E((-1)^{|b_i|}(b_i^2 \circ t_{U_i} - t_{V_i} \circ b_i^1) + t_{db_i})$, and it satisfies $E \circ i_1 = G$ and $E \circ \text{Cyl}(F) = H$.

(4) Relative Cylinder Axiom: Consider the diagram described in Definition 6.1(4), where I =Cyl. The dg category $\mathcal{B} \cup_{\mathcal{A}} \text{Cyl}(\mathcal{A}) \cup_{\mathcal{A}} \mathcal{B}$ is a semifree extension of $\text{Cyl}(\mathcal{A})$ by the set of objects $\{B^1, B^2 | B \in R_b\}$ and the set of morphisms $\{b_i^1, b_i^2 | b_i \in S_b\}$. Then, the dg functor $G: \mathcal{B} \cup_{\mathcal{A}} \text{Cyl}(\mathcal{A}) \cup_{\mathcal{A}} \mathcal{B} \to \text{Cyl}(\mathcal{B})$ is a semifree extension by the morphisms

$$\{t_B, t'_B, \hat{t}_B, \check{t}_B, \bar{t}_B \mid B \in R_b\} \cup \{t_{b_i} \mid b_i \in S_b\},$$

Hence, G is a cofibration.

(5) Interchange Axiom: First, note that for $s \in \{1, 2\}$, we have the dg functor

$$i_{s} \colon \operatorname{Cyl}(\mathcal{A}) \to \operatorname{Cyl}(\operatorname{Cyl}(\mathcal{A}))$$

$$A^{r} \mapsto A^{rs} \qquad \text{if } A^{r} \in \operatorname{Ob}(\operatorname{Cyl}(\mathcal{A})) \text{ and } r \in \{1, 2\},$$

$$\theta \mapsto \theta^{s} \qquad \text{if } \theta \in \operatorname{Mor}(\operatorname{Cyl}(\mathcal{A})),$$

and the commutative diagram

$$\begin{array}{c} \mathcal{A} \xrightarrow{i_r} & \operatorname{Cyl}(\mathcal{A}) \\ \downarrow^{i_s} & \downarrow^{i_s} \\ \operatorname{Cyl}(\mathcal{A}) \xrightarrow{\operatorname{Cyl}(i_r)} & \operatorname{Cyl}(\operatorname{Cyl}(\mathcal{A})) \end{array}$$

for any $r, s \in \{1, 2\}$ by Cylinder Axiom. Then, the semifree dg category $Cyl(Cyl(\mathcal{A}))$ can be given as follows:

- (i) Objects: $A^{rs} := i_s(i_r(A)) = \operatorname{Cyl}(i_r)(i_s(A))$ for each $A \in \mathcal{A}$ and $r, s \in \{1, 2\}$.
- (ii) Generating morphisms: For each object $A \in \mathcal{A}$ and $r \in \{1, 2\}$,

$$(t_A^*)^r := i_r(t_A^*), \quad t_{A^r}^* := \operatorname{Cyl}(i_r)(t_A^*),$$

$$t_{t_A} \colon A^{11} \to A^{22}, \quad t_{t'_A} \colon A^{21} \to A^{12}, \quad t_{\hat{t}_A} \colon A^{11} \to A^{12}, \quad t_{\tilde{t}_A} \colon A^{21} \to A^{22}, \quad t_{\bar{t}_A} \colon A^{11} \to A^{22},$$

where $t^* \in \{t, t', \hat{t}, \check{t}, \check{t}\}$, and for each generating morphism $a_i \colon X_i \to Y_i$ of \mathcal{A} and $r, s \in \{1, 2\}$,

$$\begin{aligned} a_i^{rs} &:= i_s(i_r(a_i)) = \operatorname{Cyl}(i_r)(i_s(a_i)), \quad (t_{a_i})^r := i_r(t_{a_i}), \quad t_{a_i^r} := \operatorname{Cyl}(i_r)(t_{a_i}), \quad t_{t_{a_i}} : X_i^{11} \to Y_i^{22}. \\ (\text{iii) } Degrees: \ |(t_A^*)^r| = |t_{A^r}^*| = |t_A^*|, \ |t_{t_A^*}| = |t_A^*| - 1, \ |(t_{a_i}^*)^r| = |t_{a_i^r}^*| = |a_i| - 1, \ |t_{t_{a_i}}| = |a_i| - 2. \\ (\text{iv) } Differentials: \end{aligned}$$

$$\begin{split} d(t_A^*)^r &= i_r(dt_A^*), \quad d(t_{A^r}^*) = \operatorname{Cyl}(i_r)(dt_A^*), \\ dt_{t_A} &= (t_A)^2 \circ t_{A^1} - t_{A^2} \circ (t_A)^1, \quad dt_{t'_A} &= (t'_A)^2 \circ t_{A^2} - t_{A^1} \circ (t'_A)^1, \\ dt_{\hat{t}_A} &= -(\hat{t}_A)^2 \circ t_{A^1} + t_{A^1} \circ (\hat{t}_A)^1 - (t'_A)^2 \circ t_{t_A} - t_{t'_A} \circ (t_A)^1, \\ dt_{\tilde{t}_A} &= -(\check{t}_A)^2 \circ t_{A^2} + t_{A^2} \circ (\check{t}_A)^1 - (t_A)^2 \circ t_{t'_A} - t_{t_A} \circ (t'_A)^1, \\ dt_{\bar{t}_A} &= (\bar{t}_A)^2 \circ t_{A^1} - t_{A^2} \circ (\bar{t}_A)^1 + (t_A)^2 \circ t_{\hat{t}_A} - t_{t_A} \circ (\hat{t}_A)^1 - (\check{t}_A)^2 \circ t_{t_A} - t_{\tilde{t}_A} \circ (t_A)^1, \\ dt_{\bar{t}_A} &= (\bar{t}_A)^2 \circ t_{A^1} - t_{A^2} \circ (\bar{t}_A)^1 + (t_A)^2 \circ t_{\hat{t}_A} - t_{t_A} \circ (\hat{t}_A)^1 - (\check{t}_A)^2 \circ t_{t_A} - t_{\tilde{t}_A} \circ (t_A)^1, \\ da_i^{rs} &= i_s(i_r(da_i)) = \operatorname{Cyl}(i_r)(i_s(da_i)), \quad d(t_{a_i})^r &= i_r(dt_{a_i}), \quad dt_{a_i^r} = \operatorname{Cyl}(i_r)(dt_{a_i}), \\ dt_{t_{a_i}} &= (-1)^{|a_i|}(-(t_{a_i})^2 \circ t_{X_i^1}^1 + t_{Y_i^2}^2 \circ (t_{a_i})^1 + a_i^{22} \circ t_{t_{X_i}} + t_{a_i^2}^2 \circ (t_{X_i})^1 - (t_{Y_i})^2 \circ t_{a_i^1}) - t_{t_{Y_i}} \circ a_i^{11} + t_{t_{da_i}}. \end{split}$$

Given this expression of $Cyl(Cyl(\mathcal{A}))$, we define a functor

$$T: \operatorname{Cyl}(\operatorname{Cyl}(\mathcal{A})) \to \operatorname{Cyl}(\operatorname{Cyl}(\mathcal{A})) \\ A^{rs} \mapsto A^{sr}, \quad (t_A^*)^r \mapsto t_{A^r}^*, \quad t_{A^r}^* \mapsto (t_A^*)^r \\ t_{t_A} \mapsto -t_{t_A}, \quad t_{t'_A} \mapsto t'_{A^2} \circ t_{t_A} \circ t'_{A^1} - \hat{t}_{A^2} \circ (t_A)^1 \circ t'_{A^1} + t'_{A^2} \circ (t_A)^2 \circ \check{t}_{A^1}, \\ t_{\hat{t}_A} \mapsto -t'_{A^2} \circ t_{t_A} \circ \hat{t}_{A^1} + \hat{t}_{A^2} \circ (t_A)^1 \circ \hat{t}_{A^1} + t'_{A^2} \circ (t_A)^2 \circ \bar{t}_{A^1}, \\ t_{\check{t}_A} \mapsto \check{t}_{A^2} \circ t_{t_A} \circ t'_{A^1} + \check{t}_{A^2} \circ (t_A)^2 \circ \check{t}_{A^1} + \bar{t}_{A^2} \circ (t_A)^1 \circ t'_{A^1}, \\ t_{\bar{t}_A} \mapsto \check{t}_{A^2} \circ t_{t_A} \circ \hat{t}_{A^1} + \bar{t}_{A^2} \circ (t_A)^1 \circ \hat{t}_{A^1} - \check{t}_{A^2} \circ (t_A)^2 \circ \bar{t}_{A^1}, \\ a_i^{rs} \mapsto a_i^{sr}, \quad (t_{a_i})^r \mapsto t_{a_i^r}, \quad t_{a_i^r} \mapsto (t_{a_i})^r, \quad t_{t_{a_i}} \mapsto -t_{t_{a_i}}, \end{cases}$$

where $r, s \in \{1, 2\}$. The functor T clearly satisfies $T \circ i_r = \text{Cyl}(i_r)$ and $T \circ \text{Cyl}(i_r) = i_r$ for any $r \in \{1, 2\}$. Confirming that T is a dg functor involves a straightforward computation, with the exception of showing $dT(t_{t_{a_i}}) = T(dt_{t_{a_i}})$. Upon observing that $T(t_{t_{d_a_i}}) = -t_{t_{d_a_i}}$, this also becomes trivial.

Remark 6.3. The dg functor T introduced in the proof of Interchange Axiom acts as an involution as expected, except for the morphisms $t_{t'_A}, t_{\tilde{t}_A}, t_{\tilde{t}_A}, t_{\tilde{t}_A}$. These morphisms lack counterparts such as $t'_{t_A}, \hat{t}_{t_A}, \tilde{t}_{t_A}, \tilde{t}_{t_A}, \tilde{t}_{t_A}$. However, they are superfluous in Cyl(Cyl(\mathcal{A})) (compare with Theorem 4.12). More explicitly, $t_{t'_A}, t_{\tilde{t}_A}, t_{\tilde{t}_A}, t_{\tilde{t}_A}$ are homotopic to $-T(t_{t'_A}), -T(t_{\tilde{t}_A}), -T(t_{\tilde{t}_A}), -T(t_{\tilde{t}_A})$, respectively. Therefore, T acts like an involution, but up to homotopy.

In order to show that $dgCat_s$ is a cofibration category using Theorem 6.2, we first need to establish an analogue of Whitehead theorem for semifree dg categories. Before presenting the theorem, we need to introduce some definitions and lemmas:

Definition 6.4. Let \mathcal{M} be an *I*-category with a structure (cof, *I*).

- (1) Two morphisms $F_1, F_2: \mathcal{A} \to \mathcal{B}$ in \mathcal{M} are *homotopic*, which is denoted by $F_1 \approx F_2$, if there exists a morphism $H: I(\mathcal{A}) \to \mathcal{B}$ in \mathcal{M} such that $H \circ i_1 = F_1$ and $H \circ i_2 = F_2$.
- (2) A morphism $F: \mathcal{A} \to \mathcal{B}$ in \mathcal{M} is a homotopy equivalence if there exists a morphism $G: \mathcal{B} \to \mathcal{A}$ in \mathcal{M} such that $G \circ F \approx 1_{\mathcal{A}}$ and $F \circ G \approx 1_{\mathcal{B}}$.

Lemma 6.5. Let \mathcal{A} be a dg category. If $A, C \in \mathcal{A}$ are isomorphic in $H^0(\mathcal{A})$, then there exist morphisms

$$\begin{aligned} r: A \to C, \quad r': C \to A, \quad \hat{r}: A \to A, \quad \check{r}: C \to C, \quad \bar{r}: A \to C \\ with \ the \ gradings \ |r| = |r'| = 0, \ |\hat{r}| = |\check{r}| = -1, \ |\bar{r}| = -2, \ and \ the \ differentials \\ dr = dr' = 0, \quad d\hat{r} = 1_A - r' \circ r, \quad \check{r} = 1_C - r \circ r', \quad d\bar{r} = r \circ \hat{r} - \check{r} \circ r. \end{aligned}$$

Proof. Assume $A, C \in \mathcal{A}$ are isomorphic in $H^0(\mathcal{A})$. Then, there exist morphisms

 $s: A \to C, \quad s': C \to A, \quad \hat{s}: A \to A, \quad \check{s}: C \to C$

with the gradings |s| = |s'| = 0, $|\hat{s}| = |\check{s}| = -1$, and the differentials

 $ds = ds' = 0, \quad d\hat{s} = 1_A - s' \circ s, \quad \check{s} = 1_C - s \circ s'.$

We get the desired morphisms by defining

 $r := s, \quad r' := s' \circ s \circ s', \quad \hat{r} := \hat{s} + s' \circ \check{s} \circ s, \quad \check{r} := \check{s} + s \circ \hat{s} \circ s', \quad \bar{r} := \check{s} \circ s \circ \hat{s} - s \circ \hat{s} \circ \hat{s} - \check{s} \circ \check{s} \circ s.$

Lemma 6.6. Let \mathcal{A} and \mathcal{B} be dg categories, and $F: \mathcal{A} \to \mathcal{B}$ be a dg functor such that $H^*(F)$ is full and faithful. If da = 0 and db = F(a) for some $a \in Mor(\mathcal{A})$ and $b \in Mor(\mathcal{B})$, then there exist $c \in Mor(\mathcal{A})$ and $e \in Mor(\mathcal{B})$ such that dc = a and de = b - F(c).

Proof. Since $H^*(F)$ is faithful, da = 0 and db = F(a), there exists $c_1 \in Mor(\mathcal{A})$ such that $dc_1 = a$. Then, $d(b - F(c_1)) = 0$, and since $H^*(F)$ is full, there exist $c_2 \in Mor(\mathcal{A})$ and $e \in Mor(\mathcal{B})$ such that $dc_2 = 0$ and $de = b - F(c_1) - F(c_2)$. Setting $c := c_1 + c_2$ concludes the proof.

We are now prepared to present an analogue of Whitehead's theorem for semifree dg categories:

Theorem 6.7. Consider dgCat_s as an *I*-category with the structure (cof, I = Cyl) as in Theorem 6.2. Let $F: \mathcal{A} \to \mathcal{B}$ be a morphism in dgCat_s, i.e., a dg functor between semifree dg categories. Then, F is a homotopy equivalence if and only if F is a quasi-equivalence.

Proof. First, let us assume that $F: \mathcal{A} \to \mathcal{B}$ is a homotopy equivalence. Then, there exists dg functors $G: \mathcal{B} \to \mathcal{A}, H: \operatorname{Cyl}(\mathcal{A}) \to \mathcal{A}$ and $K: \operatorname{Cyl}(\mathcal{B}) \to \mathcal{B}$ such that $H \circ i_1 = G \circ F, H \circ i_2 = 1_{\mathcal{A}}, K \circ i_1 = F \circ G$ and $K \circ i_2 = 1_{\mathcal{B}}$.

- $H^0(F): H^0(\mathcal{A}) \to H^0(\mathcal{B})$ is essentially surjective: Let $B \in \mathcal{B}$. Since K is a dg functor, we have the morphisms $K(t_B): F(G(B)) \to B$ and $K(t'_B): B \to F(G(B))$ with $dK(t_B) = dK(t'_B) = 0, dK(\hat{t}_B) = 1_{F(G(B))} - K(t'_B) \circ K(t_B)$ and $dK(\check{t}_B) = 1_B - K(t_B) \circ K(t'_B)$. Hence, $H^0(F)(G(B)) = F(G(B))$ is isomorphic to B in $H^0(\mathcal{B})$.
- $H^*(F): H^*(\mathcal{A}) \to H^*(\mathcal{B})$ is faithful: Let $a: A \to C$ be a morphism in \mathcal{A} with da = 0 such that there exists a morphism $h: F(A) \to F(C)$ with dh = F(a). By Proposition 4.7 and applying H, we have $dH(t_a) = (-1)^{|a|} (a \circ H(t_A) H(t_C) \circ G(F(a)))$. Consequently, we get

$$d((-1)^{|a|}a \circ H(\check{t}_A) + (-1)^{|a|}H(t_a \circ t'_A) + H(t_C) \circ G(h) \circ H(t'_A)) = a.$$

• $H^*(F): H^*(\mathcal{A}) \to H^*(\mathcal{B})$ is full: Let $A, C \in \mathcal{A}$ and $b: F(A) \to F(C)$ be a morphism with db = 0. Define $y := K(t_{F(C)}) \circ F(H(t'_C)) \circ b \circ F(H(t_A)) \circ K(t'_{F(A)}): F(A) \to F(C)$, which satisfies dy = 0. By Proposition 4.7 and applying K, we have $dK(t_y) = (-1)^{|y|}(y \circ K(t_{F(A)}) - K(t_{F(C)}) \circ F(G(y)))$. Consequently, we get $dh = F(H(t'_C)) \circ b \circ F(H(t_A)) - F(G(y))$ where

$$\begin{split} h &:= (-1)^{|y|} K(t'_{F(C)} \circ t_y) - K(\hat{t}_{F(C)}) \circ F(G(y)) + K(\hat{t}_{F(C)}) \circ F(H(t'_C)) \circ b \circ F(H(t_A)) \circ K(t'_{F(A)} \circ t_{F(A)}) \\ &+ (-1)^{|b|} F(H(t'_C)) \circ b \circ F(H(t_A)) \circ K(\hat{t}_{F(A)}). \end{split}$$

Hence, we have $dk = b - F(H(t_C) \circ G(y) \circ H(t'_A))$ where

$$k := F(H(t_C)) \circ h \circ F(H(t'_A)) + F(H(\check{t}_C)) \circ b \circ F(H(t_A \circ t'_A)) + (-1)^{|b|} b \circ F(H(\check{t}_A))$$

These items prove that F is a quasi-equivalence.

Conversely, assume that $F: \mathcal{A} \to \mathcal{B}$ is a quasi-equivalence. We first define a dg functor $G: \mathcal{B} \to \mathcal{A}$ as follows: Assume $\{b_i: B_i \to D_i\}$ (indexed by an ordinal) is a set of generating morphisms of \mathcal{B} .

• Let $B \in \mathcal{B}$. Since $H^0(F)$ is essentially surjective, there exists $C \in \mathcal{A}$ such that F(C) is isomorphic to B in $H^0(\mathcal{B})$. We set G(B) := C. Then, by Lemma 6.5, there exist morphisms

$$r_B \colon F(G(B)) \to B, r'_B \colon B \to F(G(B)), \, \hat{r}_B \colon F(G(B)) \to F(G(B)), \, \check{r}_B \colon B \to B, \, \bar{r}_B \colon F(G(B)) \to B,$$

where their gradings and differentials are given as in Lemma 6.5.

• Let $b_i: B_i \to D_i$ be a generating morphism such that $db_i = 0$. Then, $r'_{D_i} \circ b_i \circ r_{B_i}: F(G(B_i)) \to F(G(D_i))$ is a morphism with degree $|b_i|$ and zero differential. Since $H^*(F)$ is full, there exist morphisms $c_i: G(B_i) \to G(D_i)$ and $s_{b_i}: F(G(B_i)) \to F(G(D_i))$ such that $|c_i| = |b_i|$, $|s_{b_i}| = |b_i| - 1$, $dc_i = 0$ and $ds_{b_i} = r'_{D_i} \circ b_i \circ r_{B_i} - F(c_i)$. We set $G(b_i) := c_i$. Then, there exists a morphism $r_{b_i}: F(G(B_i)) \to D_i$ such that $|r_{b_i}| = |b_i| - 1$ and

$$dr_{b_i} = (-1)^{|b_i|} (b_i \circ r_{B_i} - r_{D_i} \circ F(G(b_i))).$$

• We proceed with transfinite induction: Assume G is defined on morphisms generated by $\{b_j: B_j \to D_j \mid j < i\}$ as a dg functor, and assume that for j < i, there exist morphisms $r_{b_j}: F(G(B_j)) \to D_j$ such that $|r_{b_j}| = |b_j| - 1$ and

$$dr_{b_j} = (-1)^{|b_j|} (b_j \circ r_{B_j} - r_{D_j} \circ F(G(b_j))) + r_{db_j},$$

where r_b for any morphism b generated by $\{b_j | j < i\}$ is defined as in Definition 4.6 after replacing t with r, i_1 with $F \circ G$, and i_2 with $1_{\mathcal{B}}$. Also, Proposition 4.7 applies to the morphisms r_b after such replacement. We want to define $G(b_i)$ and r_{b_i} .

We have $d(r'_{D_i} \circ b_i \circ r_{B_i} + \hat{r}_{D_i} \circ F(G(db_i)) + (-1)^{|b_i|} r'_{D_i} \circ r_{db_i}) = F(G(db_i))$. Since $H^*(F)$ is full and faithful, by Lemma 6.6, there exist morphisms c_i and s_{b_i} such that $|c_i| = |b_i|$, $|s_{b_i}| = |b_i| - 1$, $dc_i = G(db_i)$ and $ds_{b_i} = r'_{D_i} \circ b_i \circ r_{B_i} + \hat{r}_{D_i} \circ F(G(db_i)) + (-1)^{|b_i|} r'_{D_i} \circ r_{db_i} - F(c_i)$. We set $G(b_i) := c_i$. We also define

$$r_{b_i} := (-1)^{|b_i|} r_{D_i} \circ s_{b_i} + (-1)^{|b_i|} \check{r}_{D_i} \circ b_i \circ r_{B_i} + \check{r}_{D_i} \circ r_{db_i} - (-1)^{|b_i|} \bar{r}_{D_i} \circ F(G(db_i)),$$

which satisfies $|r_{b_i}| = |b_i| - 1$ and $dr_{b_i} = (-1)^{|b_i|} (b_i \circ r_{B_i} - r_{D_i} \circ F(G(b_i))) + r_{db_i}$. Note that we can also define r_b for any morphisms $b: B \to D$ in \mathcal{B} in this case as in Definition 4.6 (after replacing t with r, i_1 with $F \circ G$, and i_2 with $1_{\mathcal{B}}$), which satisfies $|r_b| = |b| - 1$ and $dr_b = (-1)^{|b|} (b \circ r_B - r_D \circ F(G(b))) + r_{db}$ by Proposition 4.7.

Having defined the dg functor $G: \mathcal{B} \to \mathcal{A}$ and morphisms $r_b: F(G(B)) \to D$, it is straightforward to check that $F \circ G \approx 1_{\mathcal{B}}$ by constructing a dg functor $K: \operatorname{Cyl}(\mathcal{B}) \to \mathcal{B}$ satisfying $K(t_B^*) = r_B^*$ for each $B \in \mathcal{B}$ and $t^* \in \{t, t', \hat{t}, \check{t}, \check{t}\}$, and $K(t_{b_i}) = r_{b_i}$ for each generating morphism b_i of \mathcal{B} . Finally, we want to show that $G \circ F \approx 1_{\mathcal{A}}$. For that, we need to construct a dg functor $H: \operatorname{Cyl}(\mathcal{A}) \to \mathcal{A}$ such that $H \circ i_1 = G \circ F$ and $H \circ i_2 = 1_{\mathcal{A}}$. Hence, we only need to define $H(t_A^*)$ for any $A \in \mathcal{A}$ satisfying $|H(t_A^*)| = |t_A^*|$ and $dH(t_A^*) = H(dt_A^*)$, and $H(t_{a_i})$ for any generating morphism a_i of \mathcal{A} satisfying $|H(t_{a_i})| = |t_{a_i}|$ and $dH(t_{a_i}) = H(dt_{a_i})$:

• Let $A \in \mathcal{A}$. Since $H^*(F)$ is full, there exist morphisms

 $u_A \colon G(F(A)) \to A, \quad v_A \colon F(G(F(A))) \to F(A), \quad u'_A \colon A \to G(F(A)), \quad v'_A \colon F(A) \to F(G(F(A)))$

such that $dv_A = r_{F(A)} - F(u_A)$ and $dv'_A = r'_{F(A)} - F(u'_A)$. Next, since $H^*(F)$ is full and faithful and $d(\hat{r}_{F(A)} + r'_{F(A)} \circ v_A + v'_A \circ F(u_A)) = F(1_{G(F(A))} - u'_A \circ u_A)$, by Lemma 6.6, there exist morphisms \hat{u}_A and \hat{v}_A such that $d\hat{u}_A = 1_{G(F(A))} - u'_A \circ u_A$ and $d\hat{v}_A =$ $\hat{r}_{F(A)} + r'_{F(A)} \circ v_A + v'_A \circ F(u_A) - F(\hat{u}_A)$. Similarly, we can define \check{u}_A and \check{v}_A satisfying $d\check{u}_A = 1_A - u_A \circ u'_A$ and $d\check{v}_A = \check{r}_{F(A)} + v_A \circ r'_{F(A)} + F(u_A) \circ v'_A - F(\check{u}_A)$. Finally, by Lemma 6.6 again, we can show the existence of morphisms \bar{u}_A and \bar{v}_A satisfying $d\bar{u}_A = u_A \circ \hat{u}_A - \check{u}_A \circ u_A$ and $d\bar{v}_A = \bar{r}_{F(A)} + \check{v}_A \circ F(u_A) - F(u_A) \circ \hat{v}_A - \check{r}_{F(A)} \circ v_A - v_A \circ \hat{r}_{F(A)} - v_A \circ r'_{F(A)} \circ v_A - F(\bar{u}_A)$. Then, we set

$$H(t_A) := u_A, \quad H(t'_A) := u'_A, \quad H(\hat{t}_A) := \hat{u}_A, \quad H(\check{t}_A) := \check{u}_A, \quad H(\bar{t}_A) := \bar{u}_A.$$

• Let $a_i: A_i \to C_i$ be a generating morphism of \mathcal{A} satisfying $da_i = 0$. First, observe that $d(r_{F(a_i)} - F(a_i) \circ v_{A_i} + (-1)^{|a_i|} v_{C_i} \circ F(G(F(a_i)))) = F((-1)^{|a_i|} (a_i \circ u_{A_i} - u_{C_i} \circ G(F(a_i))))$. Hence, by Lemma 6.6, there exist u_{a_i} and v_{a_i} satisfying

$$du_{a_i} = (-1)^{|a_i|} (a_i \circ u_{A_i} - u_{C_i} \circ G(F(a_i))) = H(dt_{a_i})$$

and $dv_{a_i} = r_{F(a_i)} - F(a_i) \circ v_{A_i} + (-1)^{|a_i|} v_{C_i} \circ F(G(F(a_i))) - F(u_{a_i})$. We set $H(t_{a_i}) := u_{a_i}$. • We proceed with transfinite induction: Let $a_i : A_i \to C_i$ be a generating morphism of \mathcal{A} . Assume $H(t_{a_j}) := u_{a_j}$ and v_{a_j} are defined for generating morphisms $\{a_j : A_j \to C_j \mid j < i\}$

with
$$du_{a_j} = (-1)^{|a_j|} (a_j \circ u_{A_j} - u_{C_j} \circ G(F(a_j))) + u_{da_j}$$
 and

$$dv_{a_j} = r_{F(a_j)} - F(a_j) \circ v_{A_j} + (-1)^{|a_j|} v_{C_j} \circ F(G(F(a_j))) - v_{da_j} - F(u_{a_j}).$$

Here, we set $u_a := H(t_a)$ for any morphism *a* generated by $\{a_j \mid j < i\}$, and for any morphism $a = a_{j_n} \circ \ldots \circ a_{j_1}$ with $a_{j_l} \in \{a_j \mid j < i\}$, we define

$$v_a := \sum_{l=1}^n (-1)^{|a|-|a_l|} F(a_{j_n} \circ \ldots \circ a_{j_{l+1}}) \circ v_{a_{j_l}} \circ F(G(F(a_{j_{l-1}} \circ \ldots \circ a_{j_1}))).$$

The definition extends linearly to define v_a for any morphism a generated by $\{a_j | j < i\}$. Then, a version of Proposition 4.7 holds for the morphisms v_a . In particular, we have

$$dv_a = r_{F(a)} - F(a) \circ v_A + (-1)^{|a|} v_C \circ F(G(F(a))) - v_{da} - F(u_a)$$

for any morphism $a: A \to C$ generated by $\{a_j | j < i\}$. We want to show that there exist morphisms u_{a_i} and v_{a_i} satisfying $du_{a_i} = (-1)^{|a_i|} (a_i \circ u_{A_i} - u_{C_i} \circ G(F(a_i))) + u_{da_i}$ and

$$dv_{a_i} = r_{F(a_i)} - F(a_i) \circ v_{A_i} + (-1)^{|a_i|} v_{C_i} \circ F(G(F(a_i))) - v_{da_i} - F(u_{a_i}).$$

We observe that

$$d(r_{F(a_i)} - F(a_i) \circ v_{A_i} + (-1)^{|a_i|} v_{C_i} \circ F(G(F(a_i))) - v_{da_i}) = F((-1)^{|a_i|} (a_i \circ u_{A_i} - u_{C_i} \circ G(F(a_i))) + u_{da_i}).$$

Hence, by Lemma 6.6, the desired morphisms u_{a_i} and v_{a_i} exist. We set $H(t_{a_i}) := u_{a_i}$.

These items define the dg functor $H: \operatorname{Cyl}(\mathcal{A}) \to \mathcal{A}$ satisfying $H \circ i_1 = G \circ F$ and $H \circ i_2 = 1_{\mathcal{A}}$, which implies $G \circ F \approx 1_{\mathcal{A}}$. Hence, F is a homotopy equivalence.

At this stage, we can provide a computable functorial factorizations of dg functors as a composition of a cofibration and a quasi-equivalence. We will achieve this using mapping cylinders of dg functors, which we will define below:

Definition 6.8. Let \mathcal{A} be a semifree dg category and $F: \mathcal{A} \to \mathcal{B}$ be a dg functor.

(1) We define the mapping cylinder M_F of F as the pushout

$$M_F := \operatorname{colim}(\operatorname{Cyl}(\mathcal{A}) \xleftarrow{^{i_1}} \mathcal{A} \xrightarrow{F} \mathcal{B}).$$

Explicitly, M_F is given as the semifree extension of $\mathcal{B} \amalg \mathcal{A}$ by the morphisms

 $t_A \colon F(A) \to A, \quad t'_A \colon A \to F(A), \quad \hat{t}_A \colon F(A) \to F(A), \quad \check{t}_A \colon A \to A, \quad \bar{t}_A \colon F(A) \to A$ for each $A \in \mathcal{A}$, and the morphism

$$t_{a_i} \colon F(X_i) \to Y_i$$

for each generating morphism $a_i \colon X_i \to Y_i$ of \mathcal{A} , with the gradings

$$|t_A| = |t'_A| = 0, \quad |\hat{t}_A| = |\check{t}_A| = -1, \quad |\bar{t}_A| = -2, \quad |t_{a_i}| = |a_i| - 1$$

and differentials

$$dt_A = dt'_A = 0, \quad d\hat{t}_A = 1_{F(A)} - t'_A \circ t_A, \quad d\check{t}_A = 1_A - t_A \circ t'_A, \quad d\bar{t}_A = t_A \circ \hat{t}_A - \check{t}_A \circ t_A, \\ dt_{a_i} = (-1)^{|a_i|} (a_i \circ t_{X_i} - t_{Y_i} \circ F(a_i)) + t_{da_i},$$

where for any morphism a in \mathcal{A} , t_a is defined as in Definition 4.6 (after replacing i_1 with F and i_2 with $1_{\mathcal{A}}$).

(2) For any commutative diagram

$$\begin{array}{ccc} \mathcal{A} & \stackrel{F}{\longrightarrow} & \mathcal{B} \\ \downarrow^{\alpha} & & \downarrow^{\beta} \\ \mathcal{A}' & \stackrel{F'}{\longrightarrow} & \mathcal{B}' \end{array}$$

in dgCat_s, we define a dg functor $M(\alpha, \beta) \colon M_F \to M_{F'}$ as the image of the morphism of the diagrams

$$\begin{array}{ccc} \operatorname{Cyl}(\mathcal{A}) & \xleftarrow{i_1} & \mathcal{A} & \xrightarrow{F} & \mathcal{B} \\ & & & \downarrow^{\operatorname{Cyl}(\alpha)} & & \downarrow^{\alpha} & & \downarrow^{\beta} \\ \operatorname{Cyl}(\mathcal{A}') & \xleftarrow{i_1} & \mathcal{A}' & \xrightarrow{F'} & \mathcal{B}' \end{array}$$

under the colimit functor. Explicitly, $M(\alpha, \beta)$ is given by

$$M(\alpha,\beta)|_{\mathcal{B}\amalg\mathcal{A}} = \beta \amalg \alpha, \quad M(\alpha,\beta)(t_A^*) = t_{\alpha(A)}^*, \quad M(\alpha,\beta)(t_{a_i}) = t_{\alpha(a_i)},$$

where $t^* \in \{t, t', \hat{t}, \check{t}, \bar{t}\}$.

Theorem 6.9. Let $F: \mathcal{A} \to \mathcal{B}$ be a dg functor between semifree dg categories, M_F be the mapping cylinder of F, and $M(\alpha, \beta): M_F \to M_{F'}$ be a dg functor as in Definition 6.8.

(1) There exists a factorization of F by dg functors

$$F: \mathcal{A} \xrightarrow{\jmath} M_F \xrightarrow{q} \mathcal{B},$$

where j is defined as the composition $\mathcal{A} \hookrightarrow \mathcal{B} \amalg \mathcal{A} \hookrightarrow M_F$, and q is defined as

$$q|_{\mathcal{B}\amalg\mathcal{A}} := 1_{\mathcal{B}}\amalg F, \quad q(t_A) = q(t'_A) := 1_{F(A)}, \quad q(\hat{t}_A) = q(\check{t}_A) = q(\bar{t}_A) = q(t_{a_i}) := 0$$

Moreover, j is a cofibration and q is a quasi-equivalence.

(2) The factorization above is functorial. In other words, for every commutative square

$$\begin{array}{c} \mathcal{A} \xrightarrow{F} \mathcal{B} \\ \downarrow^{\alpha} \qquad \qquad \downarrow^{\beta} \\ \mathcal{A}' \xrightarrow{F'} \mathcal{B}' \end{array}$$

in $dgCat_s$, the diagram

$$\begin{array}{cccc} \mathcal{A} & \stackrel{j}{\longrightarrow} & M_{F} & \stackrel{q}{\longrightarrow} & \mathcal{B} \\ & & & & \downarrow^{M(\alpha,\beta)} & & \downarrow^{\beta} \\ \mathcal{A}' & \stackrel{j'}{\longrightarrow} & M_{F'} & \stackrel{q'}{\longrightarrow} & \mathcal{B}' \end{array}$$

commutes, $M(1_{\mathcal{A}}, 1_{\mathcal{A}}) = 1_{M_F}$, and $M(\alpha', \beta') \circ M(\alpha, \beta) = M(\alpha' \circ \alpha, \beta' \circ \beta)$.

Proof. Since dgCat_s is an *I*-category with I = Cyl by Theorem 6.2, [Bau89, Lemma 3.12] shows that the first item holds with the distinction that q is a homotopy equivalence. Then, Theorem 6.7 implies that q is a quasi-equivalence. As for the second item, it is straightforward to check.

Remark 6.10. It is easy to check that the dg functor q above is a fibration in the Dwyer-Kan model category of dg categories given in Theorem 2.2. See e.g. [KL21] for the definition of fibrations. Hence, Theorem 6.9 gives an explicit factorization of F into a cofibration and an acyclic fibration in the Dwyer-Kan model category. This factorization also holds in quasi-equiconic and Morita model categories of dg categories given in Theorem 2.2, as they are left Bousfield localization of the Dwyer-Kan model category.

We are ready to present one of the main results of this section:

Theorem 6.11. The category of semifree dg categories $dgCat_s$ forms a cofibration category, where weak equivalences are quasi-equivalences and cofibrations are semifree extensions. Moreover, every object in $dgCat_s$ is both fibrant and cofibrant. In particular, $dgCat_s$ makes a category of cofibrant objects.

Proof of Theorem 6.11. According to [Bau89], every *I*-category is a cofibration category with the same cofibrations, and its weak equivalences are homotopy equivalences. Additionally, each of its objects is both fibrant and cofibrant. Since homotopy equivalences between semifree dg categories are exactly quasi-equivalences by Theorem 6.7, the result follows. Finally, cofibrant objects in a cofibration category makes a category with cofibrant objects, see e.g. [KP97] for the definition. \Box

Remark 6.12. Since Theorem 6.11 realizes $dgCat_s$ as a cofibration category, we can consider homotopical constructions such as homotopy pushouts in $dgCat_s$, see [Bau89]. Specifically, we can define the homotopy pushout as

 $\operatorname{hocolim}(\mathcal{A} \xleftarrow{\alpha} \mathcal{C} \xrightarrow{\beta} \mathcal{B}) := \operatorname{colim}(M_{\alpha} \xleftarrow{j} \mathcal{C} \xrightarrow{\beta} \mathcal{B})$

up to quasi-equivalence using Theorem 6.9(1). As a consequence, we can directly obtain Theorem 5.3 for dgCat_s using Theorem 6.9(1) and Theorem 6.9(2).

We conclude this section by stating the main theorems for semifree dg algebras and dg categories/algebras of strictly finite type. Their proofs follow a similar approach to those for semifree dg categories.

Theorem 6.13.

- (1) The category of semifree dg algebras $dgAlg_s$ is an I-category with the structure (cof, I), which is defined as follows:
 - cof : Cofibrations are semifree extensions.
 - I: The functor I is the cylinder functor Cyl: $dgAlg_s \rightarrow dgAlg_s$ described in Remark 4.11.

(2) The category of semifree dg algebras dgAlg_s forms a cofibration category, where weak equivalences are quasi-equivalences and cofibrations are semifree extensions. Moreover, every object in dgAlg_s is both fibrant and cofibrant. In particular, dgAlg_s makes a category of cofibrant objects.

Remark 6.14. Let $dgCat_f$ be the category of dg categories of strictly finite type, i.e., semifree dg categories with finitely many objects and generating morphisms. It is easy to check that Theorem 6.2 and 6.11 hold if we replace $dgCat_s$ with $dgCat_f$. The same is true if we consider the category $dgAlg_f$ of dg algebras of strictly finite type, i.e., semifree dg algebras with finitely many generating morphisms: Theorem 6.13 still holds if we replace $dgAlg_s$ with $dgAlg_f$.

7. Wrapped Fukaya category of T^*S^n and the reflection functor

In this section, we give a simple application of Theorem 5.3 and 5.7 within the context of symplectic geometry.

Consider two symplectic manifolds W_1 and W_2 , and a symplectic map $f : W_1 \to W_2$, which is a smooth map respecting their symplectic structures. It is well-known that f induces a functor between their Fukaya categories. It is a natural question to ask how to construct the induced functor from f.

For a Weinstein manifold, i.e., an open symplectic manifold satisfying certain conditions, [GPS18] proved that the corresponding wrapped Fukaya category can be computed, after a choice of a covering, as a homotopy colimit of the wrapped Fukaya categories of the covering elements. Furthermore, if f respects the coverings, applying Theorems 5.3 or 5.7 allows us to describe the functor induced by f. To illustrate this, we provide an example of constructing an induced functor using Theorems 5.3 and 5.7. We consider the symplectic manifold T^*S^n , where S^n is the *n*-dimensional sphere.

Before going further, let us discuss motivations behind considering these induced functors.

The first motivation stems from a conjecture written in [Kon09], which questions whether the group of autofunctors of the Fukaya category coincides with the (stabilized) group of symplectic automorphisms. If this conjecture holds true, the induced functor would encapsulate the same information as the original symplectic automorphism. Thus, studying induced functors becomes instrumental in exploring symplectic automorphisms.

The second motivation is the homotopy colimit computation of the wrapped Fukaya category, established in [GPS18]. This computation involves a homotopy colimit diagram comprising categories and functors between them. For instance, in Theorems 5.3 and 5.7, the homotopy colimit diagram includes functors denoted as α and β . Therefore, to fully describe a homotopy colimit diagram and subsequently compute the wrapped Fukaya category via a homotopy colimit, it is essential to specify these corresponding functors.

In Section 7.1, we recall some facts about the wrapped Fukaya category of Weinstein manifolds. In Section 7.2, we calculate the wrapped Fukaya category $\mathcal{W}(T^*S^n)$ of the cotangent bundle T^*S^n . In Section 7.3, we present the dg functor $R_n: \mathcal{W}(T^*S^n) \to \mathcal{W}(T^*S^n)$ that is induced from the reflection map $r_n: T^*S^n \to T^*S^n$ along an axis.

7.1. **Preliminaries on wrapped Fukaya categories.** In this subsection, we will review some key aspects of a powerful invariant known as the "wrapped Fukaya category" $\mathcal{W}(W)$ of a Weinstein manifold/sector W. A Weinstein manifold is an exact symplectic manifold whose Liouville vector field is both complete and gradient-like for an exhausting Morse function. A Weinstein sector is a Weinstein manifold with boundary, where its Liouville vector field satisfies certain conditions near the boundary. For a more detailed exposition, readers can consult [CE12] and [GPS20].

32

The wrapped Fukaya category $\mathcal{W}(W)$ of W is an A_{∞} -category whose objects are certain exact Lagrangians in W with cylindrical ends, equipped with additional data. Morphisms in this category are generated by the intersections of Lagrangians after perturbing them through a process known as "wrapping". A_{∞} -operations arise from counting pseudoholomorphic polygons bounded by Lagrangians (edges) and their intersections (corners). For a rigorous definition, refer to [GPS20] and [Sei08].

The subsequent subsections will focus on the case where W is a cotangent bundle, specifically, $W = T^*S^n$ for some n.

Remark 7.1.

- (1) Any A_{∞} -category C can be regarded as a dg category up to quasi-equivalence by replacing C with its image under the A_{∞} -Yoneda embedding. Hence, wrapped Fukaya categories can be regarded as dg categories up to quasi-equivalence.
- (2) The wrapped Fukaya category $\mathcal{W}(W)$ can be given \mathbb{Z} -grading if $2c_1(W) = 0 \in H^2(W; \mathbb{Z})$. Also, the definition of $\mathcal{W}(W)$ depends on the classes $\eta \in H^1(W; \mathbb{Z})$ (grading structure) and $b \in H^2(W; \mathbb{Z}/2)$ (background class), which are used to give gradings on the Lagrangian intersections and orientations of moduli spaces of pseudoholomorphic disks, respectively. See [Sei08] for more details, or [BCJ⁺22] for a quick overview.

The *skeleton* of a Weinstein manifold/sector W can be defined as the union of the stable manifolds of the zeroes of the Liouville vector field of W. Then, we have the following theorem:

Theorem 7.2 ([CRGG17, GPS18]). Let W be a Weinstein manifold (or sector) of dimension 2n. Then W(W) is generated by Lagrangian cocores, which are Lagrangian disks dual to n dimensional strata of the skeleton of W.


Remark 7.3. When $W = T^*M$ for a smooth manifold M, its skeleton is the zero section M, and Lagrangian cocores correspond to cotangent fibers. Then, Theorem 7.2 implies that the wrapped Fukaya category $\mathcal{W}(T^*M)$ is generated by cotangent fibers of T^*M , which is originally due to [Abo11]. Moreover, if $\mathcal{W}(T^*M)$ is equipped with the standard grading structure and the background class as described in [NZ09], [Abo12] (or [GPS18, Example 1.36]) shows that

$$\hom^*(L_p, L_p) \simeq C_{-*}(\Omega_p(M))$$

where L_p is the cotangent fiber at $p \in M$, and $C_{-*}(\Omega_p(M))$ is the normalized cubical chains on the based (Moore) loop space $\Omega_p(M)$ of M at p. The product structure on $C_{-*}(\Omega_p(M))$ is induced by the concatenation of Moore loops, which is strictly associative.

Given an inclusion of Weinstein sectors $F \hookrightarrow W$, there is an induced A_{∞} -functor $\mathcal{W}(F) \to \mathcal{W}(W)$ as described in [GPS20]. Then, we state the following theorem, which will serve as our primary tool for computing wrapped Fukaya categories:

Theorem 7.4 ([GPS18]). Let W be a Weinstein manifold (or sector). Suppose $W = W_1 \cup W_2$ for some Weinstein sectors W_1 and W_2 , and $W_1 \cap W_2$ is a hypersurface in W whose neighborhood can be written as $F \times T^*[0,1]$, where F is Weinstein (up to deformation). Then there is a pretriangulated equivalence



where the arrows are induced by the inclusion of Weinstein sectors $F \hookrightarrow W_i$ for i = 1, 2.

7.2. Wrapped Fukaya category $\mathcal{W}(T^*S^n)$ of T^*S^n . First, note that by Remark 7.1(2), the wrapped Fukaya category $\mathcal{W}(T^*S^n)$ can be given \mathbb{Z} -grading since $2c_1(T^*S^n) = 0 \in H^2(T^*S^n;\mathbb{Z})$ for any $n \geq 1$. Also by Remark 7.1(2), the definition of $\mathcal{W}(T^*S^n)$ depends on the grading structure $\eta \in H^1(T^*S^n;\mathbb{Z})$ and the background class $b \in H^2(T^*S^n;\mathbb{Z}/2)$. They are uniquely determined for T^*S^n for $n \geq 3$. For T^*S^1 and T^*S^2 , we choose the standard grading structure and the background class as in [NZ09]. We will comment on the nonstandard choices in Remark 7.6.

Proposition 7.5. Let $n \ge 1$. The wrapped Fukaya category of T^*S^n is given, up to pretriangulated equivalence, by

$$\mathcal{W}(T^*S^n) \simeq \begin{cases} \mathcal{C}_1[z^{-1}] & \text{if } n = 1\\ \mathcal{C}_n & \text{if } n \ge 2 \end{cases}$$

where C_n is the semifree dg category given as follows:

- (i) Objects: L (representing a cotangent fiber of T^*S^n).
- (ii) Generating morphisms: $z \in \hom^*(L, L)$.
- (*iii*) Degrees: |z| = 1 n.
- (*iv*) Differentials: dz = 0.

Remark 7.6.

- (1) For T^*S^1 , there are \mathbb{Z} -many grading structures to define $\mathcal{W}(T^*S^1)$. To capture the nonstandard ones, one needs to let |z| = m for $m \in \mathbb{Z}$ instead of |z| = 0.
- (2) For T^*S^2 , there is another background class to define $\mathcal{W}(T^*S^2)$. To capture the nonstandard one, one needs to replace $dz = 1_L 1_L = 0$ with $dz = 1_L + 1_L = 2 \cdot 1_L$.

Proof of Proposition 7.5. The cases n = 1 and n = 2 are proved in e.g. [Kar18] using microlocal sheaves, but we will prove it independently for any $n \ge 1$. By Theorem 7.4, we have the pretriangulated equivalence

(7.1)

$$\mathcal{W}(T^*S^n) \simeq \operatorname{hocolim}(\mathcal{W}(T^*D^n) \leftarrow \mathcal{W}(T^*S^{n-1}) \to \mathcal{W}(T^*D^n))$$

$$\simeq \operatorname{hocolim}(\operatorname{Tw}(\mathcal{A}_1(1)) \leftarrow \mathcal{W}(T^*S^{n-1}) \to \operatorname{Tw}(\mathcal{A}_1(2)))$$

where we used the pretriangulated equivalence $\mathcal{W}(T^*D^n) \simeq \mathcal{A}_1(i)$ for any *i*, where $\mathcal{A}_1(i)$ is the semifree dg category with a single object K_i (representing a cotangent fiber) and no generating morphisms. Note that we use *i* just to distinguish multiple \mathcal{A}_1 's.

Case n = 1: We have the pretriangulated equivalence $\mathcal{W}(T^*S^0) \simeq \mathcal{A}_1(3) \amalg \mathcal{A}_1(4)$ because S^0 is a disjoint union of two points, hence (7.1) becomes

$$\mathcal{W}(T^*S^1) \simeq \operatorname{hocolim}(\operatorname{Tw}(\mathcal{A}_1(1)) \leftarrow \operatorname{Tw}(\mathcal{A}_1(3) \amalg \mathcal{A}_1(4)) \to \operatorname{Tw}(\mathcal{A}_1(2)))$$

$$\simeq \operatorname{hocolim}(\mathcal{A}_1(1) \leftarrow \mathcal{A}_1(3) \amalg \mathcal{A}_1(4) \to \mathcal{A}_1(2)).$$

The second pretriangulated equivalence makes sense since the functors $\operatorname{Tw}(\mathcal{A}_1(3) \amalg \mathcal{A}_1(4)) \to \operatorname{Tw}(\mathcal{A}_1(i))$ for i = 1, 2 are induced by the dg functors

$$\mathcal{A}_1(3) \amalg \mathcal{A}_1(4) \to \mathcal{A}_1(i)$$
$$K_3, K_4 \mapsto K_i.$$

Then [KL21] (or Theorem 5.3(1)) gives the pretriangulated equivalence

$$\mathcal{W}(T^*S^1) \simeq \mathcal{C}'_1[\{t_{K_3}, t_{K_4}\}^{-1}]$$

where C'_1 is the semifree dg category given as follows:

- (i) Objects: K_1, K_2 .
- (ii) Generating morphisms: $t_{K_3}, t_{K_4} \in \hom^*(K_1, K_2)$.

- (iii) Degrees: $|t_{K_3}| = |t_{K_4}| = 0.$
- (iv) Differentials: $dt_{K_3} = dt_{K_4} = 0$.

Since K_1 and K_2 are homotopy equivalent in $\mathcal{C}'_1[t_{K_4}^{-1}]$, we have the quasi-equivalence

 $\begin{aligned} \mathcal{C}_1'[t_{K_4}^{-1}] \xrightarrow{\sim} \mathcal{C}_1 \\ K_1, K_2 \mapsto L, \quad t_{K_3} \mapsto z, \quad t_{K_4} \mapsto 1_L \end{aligned}$

hence we have the quasi-equivalence

$$C'_1[\{t_{K_3}, t_{K_4}\}^{-1}] \simeq C_1[z^{-1}]$$

which then gives the pretriangulated equivalence

$$\mathcal{W}(T^*S^1) \simeq \mathcal{C}_1[z^{-1}].$$

Case n = 2: By the previous case (by renaming z by x for notational purposes), (7.1) becomes

$$\mathcal{W}(T^*S^2) \simeq \operatorname{hocolim}(\operatorname{Tw}(\mathcal{A}_1(1)) \leftarrow \operatorname{Tw}(\mathcal{C}_1[x^{-1}]) \to \operatorname{Tw}(\mathcal{A}_1(2)))$$
$$\simeq \operatorname{hocolim}(\mathcal{A}_1(1) \leftarrow \mathcal{C}_1[x^{-1}] \to \mathcal{A}_1(2)).$$

The second pretriangulated equivalence makes sense since the functors $\operatorname{Tw}(\mathcal{C}_1[x^{-1}]) \to \operatorname{Tw}(\mathcal{A}_1(i))$ for i = 1, 2 are induced by the dg functors

$$\mathcal{C}_1[x^{-1}] \to \mathcal{A}_1(i)$$
$$L \mapsto K_i, \quad x \mapsto 1_{K_i}$$

Then [KL21] (or Theorem 5.7(1)) gives the pretriangulated equivalence

$$\mathcal{W}(T^*S^2) \simeq \mathcal{C}_2'[t_L^{-1}]$$

where C'_2 is the semifree dg category given as follows:

- (i) Objects: K_1, K_2 .
- (ii) Generating morphisms: $t_L, t_x \in \hom^*(K_1, K_2)$.
- (iii) Degrees: $|t_L| = 0$, $|t_x| = -1$.
- (iv) Differentials: $dt_L = 0$, $dt_x = t_L t_L = 0$.

Since K_1 and K_2 are homotopy equivalent in $\mathcal{C}'_2[t_L^{-1}]$, we have the quasi-equivalence

$$\begin{aligned} \mathcal{C}_2'[t_L^{-1}] \xrightarrow{\sim} \mathcal{C}_2 \\ K_1, K_2 \mapsto L, \quad t_L \mapsto 1_L, \quad t_x \mapsto z \end{aligned}$$

which then gives the pretriangulated equivalence

$$\mathcal{W}(T^*S^2) \simeq \mathcal{C}_2.$$

Case $n \ge 3$: Assume the proposition holds for n-1, i.e. we have the pretriangulated equivalence $\mathcal{W}(T^*S^{n-1}) \simeq \mathcal{C}_{n-1}$ (rename z in \mathcal{C}_{n-1} by x for notational purposes). Then (7.1) becomes

$$\mathcal{W}(T^*S^n) \simeq \operatorname{hocolim}(\operatorname{Tw}(\mathcal{A}_1(1)) \leftarrow \operatorname{Tw}(\mathcal{C}_{n-1}) \to \operatorname{Tw}(\mathcal{A}_1(2)))$$
$$\simeq \operatorname{hocolim}(\mathcal{A}_1(1) \leftarrow \mathcal{C}_{n-1} \to \mathcal{A}_1(2)).$$

The second pretriangulated equivalence makes sense since the functors $\operatorname{Tw}(\mathcal{C}_{n-1}) \to \operatorname{Tw}(\mathcal{A}_1(i))$ for i = 1, 2 are induced by the dg functors

$$\begin{aligned} \mathcal{C}_{n-1} &\to \mathcal{A}_1(i) \\ L &\mapsto K_i, \quad x \mapsto 0. \end{aligned}$$

Then [KL21] (or Theorem 5.3(1)) gives the pretriangulated equivalence

$$\mathcal{W}(T^*S^n) \simeq \mathcal{C}'_n[t_L^{-1}]$$

where \mathcal{C}'_n is the semifree dg category given as follows:

- (i) Objects: K_1, K_2 .
- (ii) Generating morphisms: $t_L, t_x \in \hom^*(K_1, K_2)$.
- (iii) Degrees: $|t_L| = 0$, $|t_x| = 1 n$.
- (iv) Differentials: $dt_L = 0$, $dt_x = 0$.

Since K_1 and K_2 are homotopy equivalent in $\mathcal{C}'_n[t_L^{-1}]$, we have the quasi-equivalence

$$\mathcal{C}'_n[t_L^{-1}] \xrightarrow{\sim} \mathcal{C}_n$$

$$K_1, K_2 \mapsto L, \quad t_L \mapsto 1_L, \quad t_x \mapsto z$$

which then gives the pretriangulated equivalence

$$\mathcal{W}(T^*S^n) \simeq \mathcal{C}_n.$$

7.3. The reflection functor on $\mathcal{W}(T^*S^n)$. There is a reflection map

$$r_n \colon T^* \mathbb{R}^{n+1} \to T^* \mathbb{R}^{n+1}$$
$$(x_1, x_2, \dots, x_{n+1}, y_1, y_2, \dots, y_{n+1}) \mapsto (-x_1, x_2, \dots, x_{n+1}, -y_1, y_2, \dots, y_{n+1})$$

where x_i are the base coordinates and y_i are fiber coordinates. By considering its restriction to T^*S^n via $\sum_{i=1}^n x_i^2 = 1$ and $\sum_{i=1}^n x_i y_i = 0$, we get an exact symplectomorphism

 $r_n: T^*S^n \to T^*S^n.$

Since r_n is an exact symplectomorphism, it induces an A_{∞} -quasi-equivalence

$$R_n: \mathcal{W}(T^*S^n) \to \mathcal{W}(T^*S^n).$$

For convenience, we will use the term "the reflection map/functor" to denote the exact symplectomorphism r_n and its induced functor R_n . Our goal in the subsection is to understand the reflection functor R_n explicitly using the pretriangulated equivalences $\mathcal{W}(T^*S^1) \simeq \mathcal{C}_1[z^{-1}]$ and $\mathcal{W}(T^*S^n) \simeq \mathcal{C}_n$ for $n \geq 2$ given by Proposition 7.5. We will use Theorem 5.3 and 5.7 to achieve this. The result is given by the following proposition:

Proposition 7.7. The reflection map $r_n: T^*S^n \to T^*S^n$ induces a dg functor (up to A_{∞} -natural equivalence)

$$R_n \colon \mathcal{W}(T^*S^n) \to \mathcal{W}(T^*S^n)$$

which is given as follows:

(1) If n = 1, then

$$R_1 \colon \operatorname{Tw}(\mathcal{C}_1[z^{-1}]) \to \operatorname{Tw}(\mathcal{C}_1[z^{-1}])$$
$$L \mapsto L$$
$$z \mapsto z'$$

where C_1 is the semifree dg category defined in Proposition 7.5, and z' is the inverse of z up to homotopy.

(2) If $n \geq 2$, then

$$R_n \colon \operatorname{Tw}(\mathcal{C}_n) \to \operatorname{Tw}(\mathcal{C}_n)$$
$$L \mapsto L$$
$$z \mapsto -z$$

where C_n is the semifree dg category defined in Proposition 7.5.

Proof. The reflection map $r_n: T^*S^n \to T^*S^n$ sends a cotangent fiber to a cotangent fiber, hence $R_n(L) = L[m]$ for some $m \in \mathbb{Z}$. Since $r_n \circ r_n = 1$, we have $R_n \circ R_n \simeq 1$, and consequently, we must have m = 0. Hence, $R_n(L) = L$.

Since L generates $\mathcal{W}(T^*S^n)$, R_n is determined by R_n : hom^{*}(L, L) \rightarrow hom^{*}(L, L), i.e., we need to understand the A_{∞} -functors

$$R_1: \mathcal{C}_1[z^{-1}] \to \mathcal{C}_1[z^{-1}]$$
 and $R_n: \mathcal{C}_n \to \mathcal{C}_n$ for $n \ge 2$

by Proposition 7.5. In fact, we can regard R_n as a dg functor, since C_n is a semifree dg category (in particular, cofibrant).

To determine R_n , first divide S^n with the hyperplane $\{x_{n+1} = 0\}$ into two *n*-dimensional closed disks D^n whose intersection is S^{n-1} . This induces the gluing diagram of Weinstein sectors for T^*S^n

$$T^*D^n \leftarrow T^*S^{n-1} \to T^*D^n$$

The reflection map $r_n: T^*S^n \to T^*S^n$ can be restricted to the gluing diagram such that the maps $(r_n)_{T^*D^n}: T^*D^n \to T^*D^n$ and $(r_n)_{T^*S^{n-1}} = r_{n-1}: T^*S^{n-1} \to T^*S^{n-1}$ are reflection maps, and the following diagram commutes:

$$T^*D^n \longleftarrow T^*S^{n-1} \longrightarrow T^*D^n$$

$$\downarrow^{(r_n)_{T^*D^n}} \qquad \downarrow^{r_{n-1}} \qquad \downarrow^{(r_n)_{T^*D^n}}$$

$$T^*D^n \longleftarrow S^{n-1} \longrightarrow T^*D^n$$

The induced functor by $(r_n)_{T^*D^n}$ is clearly identity on $\mathcal{W}(T^*D^n)$. Hence, the crucial data is the induced functor $R_{n-1}: \mathcal{W}(T^*S^{n-1}) \to \mathcal{W}(T^*S^{n-1})$, or equivalently, the dg functors

- $R_0: \mathcal{A}_1(3) \amalg \mathcal{A}_1(4) \to \mathcal{A}_1(3) \amalg \mathcal{A}_1(4)$ when n = 1, where $\mathcal{A}_1(i)$ is the semifree dg category with a single object K_i with no generating morphisms,
- $R_1: \mathcal{C}_1[z^{-1}] \to \mathcal{C}_1[z^{-1}]$ when n = 2,
- $R_{n-1}: \mathcal{C}_{n-1} \to \mathcal{C}_{n-1}$ when $n \ge 3$.

There is the induced morphism between diagrams

(7.2)
$$\begin{array}{c} \mathcal{A}_{1}(1) \longleftarrow \mathcal{C}_{n-1} \longrightarrow \mathcal{A}_{1}(2) \\ \downarrow^{1} \qquad \qquad \downarrow^{R_{n-1}} \qquad \qquad \downarrow^{1} \\ \mathcal{A}_{1}(1) \longleftarrow \mathcal{C}_{n-1} \longrightarrow \mathcal{A}_{1}(2) \end{array}$$

for $n \geq 3$ (replace C_{n-1} and R_{n-1} with $C_1[z^{-1}]$ and R_1 when n = 2, and with $\mathcal{A}_1(3) \amalg \mathcal{A}_1(4)$ and R_0 when n = 1), and the homotopy colimit functor sends it to $R_n \colon C_n \to C_n$ when $n \geq 2$ $(R_1 \colon C_1[z^{-1}] \to C_1[z^{-1}]$ when n = 1). So, using the homotopy colimit functor described in Theorem 5.3 and Theorem 5.7, we can determine R_n inductively. **Case** n = 1: There is the pretriangulated equivalence $\mathcal{W}(T^*S^0) \simeq \mathcal{A}_1(3) \amalg \mathcal{A}_1(4)$. Then, it is easy to see that the reflection map $r_0: T^*S^0 \to T^*S^0$ induces the dg functor

$$\begin{aligned} R_0 \colon \mathcal{A}_1(3) \amalg \mathcal{A}_1(4) &\to \mathcal{A}_1(3) \amalg \mathcal{A}_1(4) \\ K_3 &\mapsto K_4, \quad K_4 \mapsto K_3. \end{aligned}$$

We note that K_3 may be actually mapped to $K_4[m]$ for some $m \in \mathbb{Z}$, and K_4 to $K_3[-m]$, however, we can rename $K_4[m]$ as K_4 in that case. By the proof of Proposition 7.5, we have the quasi-equivalence

$$\operatorname{hocolim}(\mathcal{A}_1(1) \leftarrow \mathcal{A}_1(3) \amalg \mathcal{A}_1(4) \to \mathcal{A}_1(2)) \simeq \mathcal{C}'_1[\{t_{K_3}, t_{K_4}\}^{-1}]$$

where C'_1 is defined in the proof. Then, by applying the homotopy colimit functor to the morphism of diagrams in (7.2) for n = 1, Theorem 5.3 gives us

Finally, we need to interpret this functor on $C_1[z^{-1}]$ using the quasi-equivalence given in the proof of Proposition 7.5:

$$\begin{aligned} \mathcal{C}_1'[\{t_{K_3}, t_{K_4}\}^{-1}] &\xrightarrow{\sim} \mathcal{C}_1[z^{-1}] \\ K_1 \mapsto L, \quad K_2 \mapsto L, \quad t_{K_3} \mapsto z, \quad t_{K_4} \mapsto 1_L \\ \text{(implies } t'_{K_3} \mapsto z' \text{and } t'_{K_4} \mapsto 1_L) \end{aligned}$$

It has a quasi-inverse

$$\mathcal{C}_1[z^{-1}] \xrightarrow{\sim} \mathcal{C}'_1[\{t_{K_3}, t_{K_4}\}^{-1}]$$
$$L \mapsto K_1, \quad z \mapsto t'_{K_4} t_{K_3}.$$

This gives us the dg functor

$$R_1: \mathcal{C}_1[z^{-1}] \xrightarrow{\sim} \mathcal{C}_1'[\{t_{K_3}, t_{K_4}\}^{-1}] \xrightarrow{R_1} \mathcal{C}_1'[\{t_{K_3}, t_{K_4}\}^{-1}] \xrightarrow{\sim} \mathcal{C}_1[z^{-1}]$$

$$L \mapsto K_1 \mapsto K_1 \mapsto L$$

$$z \mapsto t'_{K_4} t_{K_3} \mapsto t'_{K_3} t_{K_4} \mapsto z'$$

as desired.

Case n = 2: We know that $\mathcal{W}(T^*S^1) \simeq \mathcal{C}_1[x^{-1}]$ (note that we replaced z with x for notational purposes), and from the case n = 1, we have the functor $R_1: \mathcal{C}_1[x^{-1}] \to \mathcal{C}_1[x^{-1}]$ such that $R_1(L) = L$ and $R_1(x) = x'$. By the proof of Proposition 7.5, we have the quasi-equivalence

$$\operatorname{hocolim}(\mathcal{A}_1(1) \xleftarrow{\alpha} \mathcal{C}_1[x^{-1}] \xrightarrow{\beta} \mathcal{A}_1(2)) \simeq \mathcal{C}'_2[t_L^{-1}]$$

where C'_2 is defined in the proof. Then, by applying the homotopy colimit functor to the morphism of diagrams in (7.2) for n = 2, Theorem 5.7 gives us

$$\begin{aligned} R_2 \colon \mathcal{C}'_2[t_L^{-1}] &\to \mathcal{C}'_2[t_L^{-1}] \\ K_1 &\mapsto K_1, \quad K_2 \mapsto K_2, \quad t_L \mapsto t_{R_1(L)} = t_L, \quad t_x \mapsto t_{R_1(x)} = t_{x'}, \\ t'_L, \hat{t}_L, \check{t}_L \mapsto t'_L, \hat{t}_L, \check{t}_L, \check{t}_L & \text{respectively,} \end{aligned}$$

where $t_{x'}$ is determined by the formula in (4.7) (after replacing i_1 with α , i_2 with β). It is given by

$$t_{x'} = -\beta(x') \circ t_x \circ \alpha(x') - \beta(\hat{x}) \circ t_L \circ \alpha(x') + \beta(x') \circ t_L \circ \alpha(\check{x}) + \beta(\hat{x} \circ x' - x' \circ \check{x}) \circ t_L$$

= $-1_{K_2} \circ t_x \circ 1_{K_1} - 0 \circ t_L \circ 1_{K_1} + 1_{K_2} \circ t_L \circ 0 + 0 \circ t_L$
= $-t_x$,

since $\alpha(x) = \alpha(x') = 1_{K_1}$, $\beta(x) = \beta(x') = 1_{K_2}$, and $\alpha(\hat{x}) = \alpha(\check{x}) = \beta(\check{x}) = \beta(\check{x}) = 0$. Finally, we need to interpret this functor on \mathcal{C}_2 using the quasi-equivalence given in the proof of Proposition 7.5:

$$\begin{array}{c} \mathcal{C}_2'[t_L^{-1}] \xrightarrow{\sim} \mathcal{C}_2 \\ K_1 \mapsto L, \quad K_2 \mapsto L, \quad t_L \mapsto 1_L, \quad t_x \mapsto z \\ \text{(implies } t_L' \mapsto 1_L) \end{array}$$

It has a quasi-inverse

$$\mathcal{C}_2 \xrightarrow{\sim} \mathcal{C}'_2[t_L^{-1}] \\ L \mapsto K_1, \quad z \mapsto t'_L t_x$$

This gives us the dg functor

$$R_2 \colon \mathcal{C}_2 \xrightarrow{\sim} \mathcal{C}'_2[t_L^{-1}] \xrightarrow{R_2} \mathcal{C}'_2[t_L^{-1}] \xrightarrow{\sim} \mathcal{C}_2$$
$$L \mapsto K_1 \mapsto K_1 \mapsto L$$
$$z \mapsto t'_L t_x \mapsto -t'_L t_x \mapsto -z$$

as desired.

Case $n \geq 3$: We know that $\mathcal{W}(T^*S^{n-1}) \simeq \mathcal{C}_{n-1}$ (note that we replaced z in \mathcal{C}_{n-1} with x for notational purposes), and by the induction hypothesis, we have the functor $R_{n-1}: \mathcal{C}_{n-1} \to \mathcal{C}_{n-1}$ such that $R_{n-1}(L) = L$ and $R_{n-1}(x) = -x$. By the proof of Proposition 7.5, we have the quasi-equivalence

$$\operatorname{hocolim}(\mathcal{A}_1(1) \xleftarrow{\alpha} \mathcal{C}_{n-1} \xrightarrow{\beta} \mathcal{A}_1(2)) \simeq \mathcal{C}'_n[t_L^{-1}]$$

where C'_n is defined in the proof. Then, by applying the homotopy colimit functor to the morphism of diagrams in (7.2) for $n \ge 3$, Theorem 5.3 gives us

$$\begin{aligned} R_n \colon \mathcal{C}'_n[t_L^{-1}] &\to \mathcal{C}'_n[t_L^{-1}] \\ K_1 \mapsto K_1, \quad K_2 \mapsto K_2, \quad t_L \mapsto t_{R_{n-1}(L)} = t_L, \quad t_x \mapsto t_{R_{n-1}(x)} = t_{-x} = -t_x, \\ t'_L, \hat{t}_L, \check{t}_L, \check{t}_L \mapsto t'_L, \hat{t}_L, \check{t}_L, \check{t}_L & \text{respectively,} \end{aligned}$$

where $t_{-x} = -t_x$ is determined by the formula in Definition 4.6. Finally, we need to interpret this functor on C_n using the quasi-equivalence given in the proof of Proposition 7.5:

$$\begin{aligned} & \mathcal{C}'_n[t_L^{-1}] \xrightarrow{\sim} \mathcal{C}_n \\ & K_1 \mapsto L, \quad K_2 \mapsto L, \quad t_L \mapsto 1_L, \quad t_x \mapsto z \\ & \text{(implies } t'_L \mapsto 1_L) \end{aligned}$$

It has a quasi-inverse

$$\begin{array}{c} \mathcal{C}_n \xrightarrow{\sim} \mathcal{C}'_n[t_L^{-1}] \\ L \mapsto K_1, \quad z \mapsto t'_L t_x \end{array}$$

This gives us the dg functor

$$R_n \colon \mathcal{C}_n \xrightarrow{\sim} \mathcal{C}'_n[t_L^{-1}] \xrightarrow{R_n} \mathcal{C}'_n[t_L^{-1}] \xrightarrow{\sim} \mathcal{C}_n$$
$$L \mapsto K_1 \mapsto K_1 \mapsto L$$
$$z \mapsto t'_L t_x \mapsto -t'_L t_x \mapsto -z$$

as desired. This concludes the proof.

References

- [Abo11] Mohammed Abouzaid, A cotangent fibre generates the Fukaya category, Advances in Mathematics **228** (2011), no. 2, 894–939.
- [Abo12] _____, On the wrapped Fukaya category and based loops, J. Symplectic Geom. 10 (2012), no. 1, 27–79. MR 2904032
- [Bau89] Hans J Baues, Algebraic homotopy, vol. 15, Cambridge university press, 1989.
- [BCJ⁺22] Hanwool Bae, Dongwook Choa, Wonbo Jeong, Dogancan Karabas, and Sangjin Lee, *Categorical entropies* on symplectic manifolds, arXiv preprint arXiv:2203.12205 (2022).
- [Bro73] Kenneth S Brown, Abstract homotopy theory and generalized sheaf cohomology, Transactions of the American Mathematical Society **186** (1973), 419–458.
- [CE12] Kai Cieliebak and Yakov Eliashberg, From Stein to Weinstein and back: Symplectic geometry of affine complex manifolds, vol. 59, American Mathematical Soc., 2012.
- [Che02] Yuri Chekanov, Differential algebra of legendrian links, Invent. Math 150 (2002), 441–483.
- [CRGG17] Baptiste Chantraine, Georgios Dimitroglou Rizell, Paolo Ghiggini, and Roman Golovko, Geometric generation of the wrapped Fukaya category of Weinstein manifolds and sectors, arXiv preprint arXiv:1712.09126 (2017).
- [Dri04] Vladimir Drinfeld, DG quotients of DG categories, Journal of Algebra 272 (2004), no. 2, 643–691.
- [DS95] William G Dwyer and Jan Spalinski, *Homotopy theories and model categories*, Handbook of algebraic topology **73** (1995), no. 126, 21.
- [Dug08] Daniel Dugger, A primer on homotopy colimits, preprint (2008).
- [EN15] Tobias Ekholm and Lenhard Ng, Legendrian contact homology in the boundary of a subcritical Weinstein 4-manifold, Journal of Differential Geometry **101** (2015), no. 1, 67–157.
- [FOOO09a] Kenji Fukaya, Yong-Geun Oh, Hiroshi Ohta, and Kaoru Ono, Lagrangian intersection Floer theory: anomaly and obstruction. Part I, AMS/IP Studies in Advanced Mathematics, vol. 46.1, American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2009. MR 2553465
- [FOO009b] _____, Lagrangian intersection Floer theory: anomaly and obstruction. Part II, AMS/IP Studies in Advanced Mathematics, vol. 46.2, American Mathematical Society, Providence, RI; International Press, Somerville, MA, 2009. MR 2548482
- [Gar09] Richard Garner, Understanding the small object argument, Applied categorical structures **17** (2009), no. 3, 247–285.
- [GPS18] Sheel Ganatra, John Pardon, and Vivek Shende, Sectorial descent for wrapped Fukaya categories, arXiv preprint arXiv:1809.03427 (2018).
- [GPS20] _____, Covariantly functorial wrapped Floer theory on Liouville sectors, Publ. Math. Inst. Hautes Études Sci. 131 (2020), 73–200. MR 4106794
- [GPS24] _____, Microlocal Morse theory of wrapped Fukaya categories, Annals of Mathematics **199** (2024), no. 3, 943–1042.
- [Hir03] Philip S Hirschhorn, Model categories and their localizations, no. 99, American Mathematical Soc., 2003.
- [Hol14] Julian V. S. Holstein, Properness and simplicial resolutions in the model category dgCat, Homology, Homotopy and Applications 16 (2014), no. 2, 263–273.
- [Hov07] Mark Hovey, *Model categories*, no. 63, American Mathematical Soc., 2007.
- [Kar18] Dogancan Karabas, *Microlocal sheaves on pinwheels*, arXiv preprint arXiv:1810.09021 (2018).
- [Kel07] Bernhard Keller, On differential graded categories, Proceedings of the International Congress of Mathematicians Madrid, August 22–30, 2006, 2007, pp. 151–190.
- [KL21] Dogancan Karabas and Sangjin Lee, Homotopy colimits of semifree dg categories and Fukaya categories of cotangent bundles of lens spaces, arXiv preprint arXiv:2109.03411 (2021).
- [Kon09] Maxim Kontsevich, Symplectic geometry of homological algebra.
- [KP97] K Heiner Kamps and Timothy Porter, *Abstract homotopy and simple homotopy theory*, World Scientific, 1997.

- [Nad15] David Nadler, Non-characteristic expansions of Legendrian singularities, arXiv preprint arXiv:1507.01513 (2015).
- [Nad17] _____, Arboreal singularities, Geometry & Topology **21** (2017), no. 2, 1231–1274.
- [NZ09] David Nadler and Eric Zaslow, Constructible sheaves and the Fukaya category, Journal of the American Mathematical Society **22** (2009), no. 1, 233–286.
- [Sei08] Paul Seidel, Fukaya categories and Picard-Lefschetz theory, vol. 10, European Mathematical Society, 2008.
- [Tab05a] Gonçalo Tabuada, Invariants additifs de dg-catégories, International Mathematics Research Notices **2005** (2005), no. 53, 3309–3339.
- [Tab05b] _____, Une structure de catégorie de modeles de Quillen sur la catégorie des dg-catégories, Comptes Rendus Mathematique **340** (2005), no. 1, 15–19.
- [Toë07] Bertrand Toën, The homotopy theory of dg-categories and derived Morita theory, Inventiones mathematicae **167** (2007), no. 3, 615–667.
- [Toë11] _____, Lectures on dg-categories, Topics in algebraic and topological K-theory, Springer, 2011, pp. 243–302.