# On de Bruijn Rings and Families of Almost Perfect Maps

Peer Stelldinger<sup>1</sup>

<sup>1</sup> Department of Computer Science, HAW Hamburg, Germany <sup>2</sup> Hamburg University of Technology, Germany peer.stelldinger@haw-hamburg.de

Abstract. De Bruijn tori, also called perfect maps, are two-dimensional periodic arrays of letters drawn from a given finite alphabet, such that each possible pattern of a given shape (m, n) appears exactly once within one period of the torus. It is still unknown if de Bruijn tori of some certain size exist, like e.g. square shaped de Bruijn Tori with odd  $m = n \in \{3, 5, 7\}$  and an even alphabet size k. However, in certain applications like positional coding, sub-perfect maps are sufficient, i.e. one does not need every possible (m, n)-pattern to appear, as long as a sufficient large number of such patterns is captured and every pattern occurs at most once. We show, that given any m = n and a square alphabet size  $k^2$ , one can efficiently construct a sub-perfect map which is almost perfect, i.e. of almost maximal size. We do this by introducing de Bruijn rings, i.e. sub-perfect maps of minimal height, and providing an efficient construction method for them. We extend our results to non-square torus shapes and arbitrary non-prime alphabet sizes.

## 1 Introduction

De Bruijn tori, also called perfect maps, are two-dimensional cyclic matrices with their entries being drawn from some alphabet, such that every pattern of a given rectangular shape occurs exactly once within one period.

One of the most prominent use cases of de Bruijn tori is in the spatial coding context, e.g. for robot localization based on some optical ground pattern [17,18].

In such a use case, it is not required that every pattern occurs exactly once. It is sufficient, that every pattern occurs at most once. Such maps are called subperfect maps [11]. For optical robot localization, a spatial code based on some (m, n)-pattern of letters from an alphabet of size k should have the following properties:

- 1. every (m, n)-pattern occurs at most once. I.e. a local pattern uniquely determines the position.
- 2. (Almost) every possible (m, n)-pattern occurs somewhere. I.e. a large area can be covered.
- 3. m, n, k are small numbers. I.e. the pattern to be captured locally is of a manageable size.

- 2 P. Stelldinger
- 4. The code can efficiently be generated for different m, n, k. I.e. it can be streamlined for the specific application.
- 5. The decoding should be efficient. I.e. the amount of data to be stored for decoding the position given some local pattern should be much smaller than the whole code and the time complexity should be as low as possible.

The first two properties point to sub-perfect maps. Nevertheless, even in spatial coding, it is advantageous when the majority if not all patterns occur in the code. However, it is still unknown if certain types of de Bruijn tori exist, e.g. no square shaped de Bruijn torus with pattern shape (n, n) is known for  $n \in \{3, 5, 7\}$  and even alphabet size [8]. For such cases, it is of interest to be able to generate sub-perfect maps which are not far away from perfect maps in that they contain almost every possible local pattern.

For these applications, we provide a new efficient construction method for sub-perfect maps including square shaped maps for any square pattern shape (n, n) (thus including  $n \in \{3, 5, 7\}$ ). Moreover, we prove, that these maps are almost perfect, i.e. for increasing m, n, the percentage of not occurring patterns tends to zero.

## 2 Basic Definitions and Related Work for the 1D Case

Let an *alphabet* be a totally ordered, nonempty finite set  $\Sigma$  with its elements being called *letters*. A *word* is a finite sequence of letters  $[a_1, a_2, \ldots, a_m]$  with  $a_i \in \Sigma, 1 \leq i \leq m$ . A word is a *Lyndon word*, if it is strictly less than any of the rotations  $[a_i, \ldots, a_m, a_1, \ldots, a_{i-1}]$   $(2 \leq i < m)$  with respect to the induced lexicographic order. With lexmin(w) being the lexicographically smallest rotation of a word w, we call a word w aperiodic, if lexmin(w) is a Lyndon word, otherwise w is *periodic*. The equivalence classes of words w modulo cyclic rotation are called *necklaces* with lexmin(w) being a natural choice of representatives of these classes. Then the representatives of aperiodic necklaces are the Lyndon words.

For  $|\Sigma| = k$ , there are  $k^m$  words of length m in total. The number of Lyndon words is given by the necklace polynomial M(k,m) [13] (also known as Witt's formula [10]), as defined by

$$\mathbf{M}(k,m) = \frac{1}{m} \sum_{d|m} \mu\left(\frac{m}{d}\right) k^d$$

where  $\mu$  is the classic Möbius function [12]

$$\mu(n) = \begin{cases} 1, & \text{if } n = 1\\ (-1)^i, & \text{if } n \text{ is the product of } i \text{ different primes}\\ 0, & \text{otherwise.} \end{cases}$$

For m prime this simplifies to  $M(k,m) = \frac{1}{m}(k^m - k)$ . Note, that  $m \cdot M(k,m)$ , i.e. the number of aperiodic words of a given length m, quickly tends to  $k^m$  for

increasing m, or in other words, the majority of words of size m are aperiodic and fall into equivalence classes of size m under rotation [16]. As Riedel [16] only describes the asymptotic behaviour, we give an explicit (although not very strict) lower bound:

**Lemma 1.** Let  $k, m \geq 2$  be natural numbers. Then M(k,m) is smaller than  $\frac{1}{m}k^m$  and larger than  $\frac{1}{m}(k^m - k^{\lfloor m/2 \rfloor + 1})$ .

*Proof.* It is obvious that  $M(k, m) < \frac{1}{m}k^m$  since there are  $k^m$  words of length m, each aperiodic necklace is an equivalence class of m words and there are periodic necklaces for any  $m \ge 2$ . For the lower bound we have:

$$\mathbf{M}(k,m) = \frac{1}{m} \sum_{d|m} \mu\left(\frac{m}{d}\right) k^d \tag{1}$$

$$=\frac{1}{m}\left(k^{m}+\sum_{d\mid m,\ d\neq m}\mu\left(\frac{m}{d}\right)k^{d}\right)$$
(2)

$$\geq \frac{1}{m} \left( k^m - \sum_{d|m, \ d \neq m} k^d \right) \tag{3}$$

$$\geq \frac{1}{m} \left( k^m - \sum_{i=1}^{\lfloor m/2 \rfloor} k^i \right) \tag{4}$$

$$=\frac{1}{m}\left(k^m - \frac{k\left(k^{\lfloor m/2 \rfloor} - 1\right)}{k-1}\right) \tag{5}$$

$$> \frac{1}{m} \left( k^m - \frac{k \left( k^{\lfloor m/2 \rfloor} \right)}{k-1} \right) \tag{6}$$

$$> \frac{1}{m} \left( k^m - k^{\lfloor m/2 \rfloor + 1} \right) \tag{7}$$

Since the probability of a random word of length m given an alphabet of size  $k \ge 2$  being periodic is given by  $1 - mM(k, m)/k^m$ , it follows corollary 1:

**Corollary 1.** Given an alphabet  $\Sigma$  with  $\Sigma = |k| \ge 2$ , the probability of a random word of length m being periodic does not exceed  $k^{1-\lceil m/2 \rceil}$ .

A de Bruijn sequence of type  $(M; m)_k$  is a cyclic sequence of letters from an alphabet  $\Sigma$  of size k in which every possible word of length m occurs exactly once. Such a sequence always exist and is of length  $M = k^m$  [3].

A well-known method to generate de Bruijn sequences is the construction of an Eulerian cycle or of a Hamiltonian cycle in a de Bruijn digraph [9]: Given an alphabet  $\Sigma$  with  $|\Sigma| = k$ , the *m*-dimensional de Bruijn digraph is defined by the vertex set  $V = \Sigma^m$  of words of length *m* and the edge set

 $E = \{(v, w) \mid v = [v_1, v_2, \ldots, v_m] \in V; w = [v_2, \ldots, v_m, v_{m+1}] \in V\}$ . Any such graph is Hamiltonian and each Hamiltonian cycle defines a de Bruijn sequence of type  $(k^m; m)_k$ , given by  $[a_1, a_2, a_3, \ldots, a_{k^m}]$  where each sub-sequence  $[a_i, a_{i+1}, \ldots, a_{i+m}]$  of the cyclic sequence is the *i*th vertex of the Hamiltonian cycle  $(1 \leq i \leq k^m)$ . Moreover, the (m-1)-dimensional de Bruijn digraph is Eulerian and each Euler cycle defines a de Bruijn sequence of type  $(k^m; m)_k$ , given by  $[a_1, a_2, a_3, \ldots, a_{k^m}]$  where each length-(m-1)-subword  $[a_i, a_{i+1}, \ldots, a_{i+m-1}]$  of the cyclic sequence is the *i*th vertex of the Eulerian cycle.

#### 3 Basic Definitions and Related Work for the 2D Case

Given an alphabet  $\Sigma$ , a pattern of shape (m, n) is a finite array of letters

$$P = \begin{bmatrix} p_{1,1} \cdots p_{1,n} \\ \vdots & \ddots & \vdots \\ p_{m,1} \cdots p_{m,n} \end{bmatrix}$$

with  $p_{i,j} \in \Sigma$ ,  $1 \le i \le m$  and  $1 \le j \le n$ . A pattern is a row-Lyndon pattern, if the word  $[[p_{1,1}, \ldots, p_{1,n}], \ldots, [p_{m,1}, \ldots, p_{m,n}]]$  is a Lyndon word given the alphabet  $\Sigma^n$ . Further, we define lexmin(P) as the rotation of rows

$$\begin{bmatrix} p_{i,1} & \cdots & p_{i,n} \\ \vdots & \ddots & \vdots \\ p_{m,1} & \cdots & p_{m,n} \\ p_{1,1} & \cdots & p_{1,n} \\ \vdots & \ddots & \vdots \\ p_{i-1,1} & \cdots & p_{i-1,n}, \end{bmatrix}$$

such that the corresponding word  $[[p_{i,1}, \ldots, p_{i,n}], \ldots, [p_{i-1,1}, \ldots, p_{i-1,n}]]$  is a lexicographically smallest one. We call a pattern *row-aperiodic* if lexmin(w) is a row-Lyndon pattern, otherwise it is *row-periodic*.

A cyclic two-dimensional array of shape (M, N) of letters from an alphabet  $\Sigma$  of size  $|\Sigma| = k$ , is called a *sub-perfect map of type*  $(M, N; m, n)_k$ , if every (m, n)-pattern  $p \in \Sigma^{m,n}$  occurs at most once. It is called a *perfect map* or a *de Bruijn torus of type*  $(M, N; m, n)_k$  if every (m, n)-pattern occurs exactly once.

It is well known [15,8], that (omitting the trivial cases M = m = 1 and/or N = n = 1) every de Bruijn torus of type  $(M, N; m, n)_k$  must fulfill

$$M > m, N > n \text{ and } MN = k^{mn}.$$
(8)

For k = 2 it has been shown constructively, that these necessary conditions are also sufficient [15]. For k > 2, the question is not that simple. However, there are a lot of methods known for the construction of de Bruijn tori.

e.g. Cook proves that for every m, n and k (except n = 2 if k even), there exists a de Bruijn torus of type  $(k^m, k^{mn-m}; m, n)_k$  [2].

As shown by Hurlbert and Isaak [8], for k even and  $n \notin \{3, 5, 7, 9\}$ , for k = 2and for k odd there exists a de Bruijn torus of type  $(k^{n^2/2}, k^{n^2/2}; n, n)_k$  if and only if n is even or k is a perfect square. They conjecture that there are also such square shaped de Bruijn tori for k even and  $n \in \{3, 5, 7, 9\}$  iff k is a perfect square, but this is still not proven.

In the following we show on how to construct square shaped sub-perfect maps for these and other cases. We do this by introducing de Bruijn rings.

# 4 De Bruijn Rings

When replacing the necessary condition M > m in equation 8 by M = m, one gets a sub-perfect map of minimal height. We denote such thin but long sub-perfect maps as de Bruijn rings. The reason for the name *ring* is as follows:

When thinking of the geometrical shape of a torus one usually has a torus of revolution in mind, which can be described as the two dimensional surface in 3d space which one gets when rotating a circle of minor radius r around a coplanar axis of distance R > r from the center of the circle (with R being called the major radius). Although 'torus' is just the latin word for 'ring', r is often assumed to be small compared to R when one has a physical ring in mind (while for greater r one would more prefer to call this a 'donut'). We therefore define a de Bruijn ring to be a sub-perfect map of minimal minor and maximal major radius:

**Definition 1 (De Bruijn ring).** A sub-perfect map of type  $(m, N; m, n)_k$  is called an  $(m, n)_k$ -de Bruijn ring if there does not exist any sub-perfect map of type  $(m, N'; m, n)_k$  with N' > N.

For m = 1, a  $(m, n)_k$ -de Bruijn ring simplifies to a one-dimensional de Bruijn sequence of type  $(k^n; n)_k$ . Therefore, we assume for the rest of the paper, that m > 1. For similar reasons we also omit the trivial cases n = 1 and k = 1. As an example, a  $(2, 2)_2$ -de Bruijn ring is given by

$$\begin{bmatrix} 0 & 0 & 1 & 1 & 0 & 0 \\ 0 & 1 & 0 & 1 & 1 & 1 \end{bmatrix}$$

This is a sub-perfect map which contains every possible binary (2,2) pattern except of

$\begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix},$	$\begin{bmatrix} 0 \ 1 \\ 0 \ 1 \end{bmatrix}$	,	$egin{array}{c} 1 \ 0 \\ 1 \ 0 \end{array}$	and	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \end{bmatrix}$	,
---	--	---	---	-----	--	---

since each of these patterns would occur twice within a cyclic map of height two. It is obvious, that a de Bruijn Ring of type  $(m, N; m, n)_k$  can only contain row-aperiodic (m, n)-patterns, since row-periodic patterns would occur more than once in the cyclic map. e.g. for m = 3, n = 2 and k = 2, there are  $2^6 = 64$  patterns in total, whereas the following 4 patterns are the only row-periodic

ones and thus can not occur in a  $(3,2)_2$ -de Bruijn Ring:

$\begin{bmatrix} 0 & 0 \end{bmatrix}$	[0 1]		$[1 \ 0]$		[1 1]
00 ,	01	,	$1 \ 0$	,	11
0 0	$\begin{bmatrix} 0 & 1 \end{bmatrix}$		$1 \ 0$		11

In case of m = 4, n = 2 and k = 2, the following 16 out of the possible 256 patterns are row-periodic:

$\begin{bmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \end{bmatrix}$	,	$\begin{bmatrix} 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \\ 0 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 0 & 0 \\ 0 & 1 \\ 0 & 0 \\ 0 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 0 & 1 \\ 0 & 0 \\ 0 & 1 \\ 0 & 0 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 1 & 0 \\ 1 & 0 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 1 \\ 1 & 1 \\ 1 & 1 \\ 1 & 1 \\ 1 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 0 \\ 1 & 1 \\ 1 & 0 \\ 1 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 1 \\ 1 & 0 \\ 1 & 1 \\ 1 & 0 \end{bmatrix}$	,
$\begin{bmatrix} 0 & 0 \\ 1 & 0 \\ 0 & 0 \\ 1 & 0 \end{bmatrix}$	,	$\begin{bmatrix} 0 & 1 \\ 1 & 1 \\ 0 & 1 \\ 1 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 0 & 0 \\ 1 & 1 \\ 0 & 0 \\ 1 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 0 & 1 \\ 1 & 0 \\ 0 & 1 \\ 1 & 0 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 0 \\ 0 & 0 \\ 1 & 0 \\ 0 & 0 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 1 \\ 0 & 1 \\ 1 & 1 \\ 0 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 0 \\ 0 & 1 \\ 1 & 0 \\ 0 & 1 \end{bmatrix}$	,	$\begin{bmatrix} 1 & 1 \\ 0 & 0 \\ 1 & 1 \\ 0 & 0 \end{bmatrix}$	

In general, given some alphabet size k, there are  $k^{mn}$  patterns of shape (m, n), of which  $m \cdot M(k^n, m)$  are row-aperiodic (each being one of the m vertical rotations of a row-Lyndon pattern) and  $k^{mn} - m \cdot M(k^n, m)$  are row-periodic. The ratio of row-aperiodic patterns  $ap(m, n, k) := (m \cdot M(k^n, m))/k^{mn}$  quickly grows towards 1 with increasing m, n and/or k, as

$$1 \ge ap(m,n,k) > \frac{k^{mn} - k^{(\lfloor m/2 \rfloor + 1)n}}{k^{mn}},$$

and the limit of this fraction is 1 for any of  $m \to \infty$ ,  $n \to \infty$  or  $k \to \infty$ . Table 1 illustrates the ratio of row-aperiodic patterns for  $2 \le m, n \le 6$  and  $2 \le k \le 5$ .

Further examples are given in the following for de Bruijn rings of different sizes:

$$(2,2)_3: \begin{bmatrix} 000121111110200021010112201000102021\\ 011010201211212200200221221202221222 \end{bmatrix}$$
(10)

Each of these  $(m, n)_k$ -de Bruijn rings contains every row-aperiodic  $(m, n)_k$ pattern, such that it is obvious, that these sub-perfect maps are of maximal
width. The following theorem 1 states, that this is always the case, i.e. there
always exists a sub-perfect map of type  $(m, M(k^n, m); m, n)_k$ .

m	n	k = 2	k = 3	k = 4	k = 5
2	2	$\frac{12}{16}$	$\frac{72}{81}$	$\frac{240}{256}$	$\frac{600}{625}$
2	3	$\frac{56}{64}$	$\frac{702}{729}$	$\frac{4032}{4096}$	$\frac{15500}{15625}$
2	4	$\frac{240}{256}$	6480 6561	<u>65280</u> 65536	<u>390000</u> <u>390625</u>
2	5	$\frac{992}{1024}$	58806 59049	$\frac{1047552}{1048576}$	<u>9762500</u> 9765625
2	6	$\frac{4032}{4096}$	$\frac{530712}{531441}$	$\frac{16773120}{16777216}$	$\frac{244125000}{244140625}$
3	2	$\frac{60}{64}$	$\frac{720}{729}$	$\frac{4080}{4096}$	$\frac{15600}{15625}$
3	3	$\frac{504}{512}$	$\frac{19656}{19683}$	$\frac{262080}{262144}$	$\frac{1953000}{1953125}$
3	4	$\frac{4080}{4096}$	$\frac{531360}{531441}$	$\frac{16776960}{16777216}$	$\frac{244140000}{244140625}$
3	5	$\frac{32736}{32768}$	$\frac{14348664}{14348907}$	$\frac{1073740800}{1073741824}$	$\frac{30517575000}{30517578125}$
3	6	$\frac{262080}{262144}$	$\frac{387419760}{387420489}$	$\frac{68719472640}{68719476736}$	$\frac{3814697250000}{3814697265625}$
4	2	$\frac{240}{256}$	$\frac{6480}{6561}$	<u>65280</u> 65536	$\frac{390000}{390625}$
4	3	$\frac{4032}{4096}$	$\frac{530712}{531441}$	$\frac{16773120}{16777216}$	$\frac{244125000}{244140625}$
4	4	<u>65280</u> 65536	$\frac{43040160}{43046721}$	<u>4294901760</u> 4294967296	<u>152587500000</u> 152587890625
4	5	$\frac{1047552}{1048576}$	<u>3486725352</u> 3486784401	<u>1099510579200</u> 1099511627776	<u>95367421875000</u> 95367431640625
4	6	$\frac{16773120}{16777216}$	282429005040 282429536481	$\frac{281474959933440}{281474976710656}$	$\frac{59604644531250000}{59604644775390624}$
5	2	$\frac{1020}{1024}$	$\frac{59040}{59049}$	$\frac{1048560}{1048576}$	$\frac{9765600}{9765625}$
5	3	$\frac{32760}{32768}$	$\frac{14348880}{14348907}$	$\frac{1073741760}{1073741824}$	<u>30517578000</u> 30517578125
5	4	$\frac{1048560}{1048576}$	3486784320 3486784401	$\frac{1099511627520}{1099511627776}$	95367431640000 95367431640625
5	5	<u>33554400</u> 33554432	847288609200 847288609443	<u>1125899906841600</u> 1125899906842624	298023223876950016 298023223876953152
5	6	$\frac{1073741760}{1073741824}$	$\frac{205891132093920}{205891132094649}$	$\frac{1152921504606846976}{1152921504606846976}$	$\frac{931322574615478534144}{931322574615478534144}$
6	2	<u>4020</u> 4006	<u>530640</u> 521441	<u>16772880</u> 16772916	244124400
6	3	<u>261576</u> <u>260144</u>	<u>387400104</u> 297420420	<u>68719210560</u> <u>69710476726</u>	<u>3814695297000</u> 3914695265695
6	4	<u>16772880</u>	<u>282428998560</u> 282428998560	<u>281474959868160</u> 201474959868160	3814697265625 59604644530860000
6	5	$\frac{16777216}{1073708064}$	282429536481 205891117686936	281474976710656 1152921503532057600	59604644775390624 931322574584951078912
6	6	$     1073741824 \\     \underline{68719210560} \\     \underline{6871920} \\     \underline{6871920} \\     \underline{68719} \\      \underline{68719} \\      \underline{68719} \\      \underline{68719} \\      \underline{68719} \\      \underline{68719} \\      \underline{68719} \\       \underline{68719} \\       \underline{68719} \\       \underline{68719} \\       \underline{68719} \\       \underline{68719} \\       \underline{68719} \\        \underline{68719} \\        \underline{68719} \\        \underline{68719} \\        \underline{68719} \\        \underline{68719} \\        \underline{68719} \\        \underline{68719} \\        \underline{68719} \\                                    $	205891132094649 150094634909047936	$\frac{1152921504606846976}{4722366482800908959744}$	931322574615478534144 14551915228363036345499648
	ĭ	68719476736	150094635296999136	4722366482869645213696	14551915228366852423942144

**Table 1.** Ratio of row-aperiodic to all patterns of shape (m, n) and alphabet size k.

**Theorem 1.** For any alphabet size k and any pattern size (m, n), there exists an  $(m, M(k^n, m); m, n)_k$  sub-perfect map. This map is an  $(m, n)_k$ -de Bruijn ring.

It is obvious, that only the  $m \cdot M(k^n, m)$  row-aperiodic patterns of size (m, n) can occur in a map of height m. Thus, a sub-perfect map of height m has a width of at most  $M(k^n, m)$ . In order to prove that such a map exists and that it can efficiently be constructed, we introduce  $(m, n)_k$ -ring graphs, which generalize the construction of de Bruijn sequences using Euler cycles to two dimensions. In contrast to the one-dimensional case, these ring graphs are not necessarily Hamiltonian.

**Definition 2 (Ring graph).** A  $(m,n)_k$ -ring graph is a labeled multidigraph G = (V, E) with the vertices  $V = \{lexmin(S) \mid S \in \Sigma^{m,n-1}\}$  and E being the set of labeled edges  $(P_1, P_2)$  with label R, such that there exist patterns  $L, R \in \Sigma^{m,1}, C \in \Sigma^{m,n-2}$ , such that

- 1.  $P_1 = [L, C] \in V, P_2 = lexmin([C, R]) \in V,$
- 2. lexmin([L, C, R]) is a row-Lyndon pattern of shape (m, n) and
- 3. there is no lexicographically smaller R' < R with  $R' \in \Sigma^{m,1}$  and lexmin([L, C, R']) = lexmin([L, C, R]).

We set one separate edge (each with a different label) for every  $R \in \Sigma^{m,1}$  between  $P_1$  and  $P_2$  if more than one are applicable.

The nodes are representatives of the vertical rotation groups of patterns of shape (m, n-1). I.e. each node is the representative of a set of vertical rotations:

$\begin{bmatrix} p_{1,1} \cdots p_{1,n-1} \\ \vdots & \vdots \end{bmatrix}$	$\begin{array}{c}p_{2,1} \cdots p_{2,n-1}\\\vdots & \vdots\\\end{array}$		$p_{m,1} \cdots p_{m,n-1}$ $p_{1,1} \cdots p_{1,n-1}$
$\begin{vmatrix} \vdots & \ddots & \vdots \\ n & 1 & \cdots & n & 1 \end{vmatrix}$	$p_{m,1}\cdots p_{m,n-1}$	····	÷ •. ÷
[Pm,1  Pm,n-1]	$p_{1,1} \cdots p_{1,n-1}$		$p_{m-1,1}\cdots p_{m-1,n-1}$

The number of elements in each vertical rotation group is a divisor of m and depends on the vertical periodicity of the group elements.

Moreover, each edge stands for the vertical rotation group of a unique pattern of shape (m, n) and one gets a group representative by concatenating the outgoing vertex label with the edge label.

While any vertex P is equal to lexmin(P), this is not necessarily true for the edge labels. A vertex will have an outgoing edge for any  $R \in \Sigma^{m,1}$ , which completes the vertex label to a unique row-aperiodic pattern. E.g. in case of m = 2, n = 3 and k = 2, there are two edges from the vertex  $\begin{bmatrix} 10\\11 \end{bmatrix}$  to the vertex  $\begin{bmatrix} 10\\11 \end{bmatrix}$ , with one edge being labeled as  $\begin{bmatrix} 0\\11 \end{bmatrix}$  and the other as  $\begin{bmatrix} 1\\0 \end{bmatrix}$ . These edges represent the row-Lyndon patterns  $\begin{bmatrix} 010\\111 \end{bmatrix}$  and  $\begin{bmatrix} 011\\110 \end{bmatrix}$ . However, due to the third condition, there is only one edge from the vertex  $\begin{bmatrix} 00\\00 \end{bmatrix}$  to the vertex  $\begin{bmatrix} 00\\01 \end{bmatrix}$  with label  $\begin{bmatrix} 0\\1 \end{bmatrix}$ , which represents the row-Lyndon pattern  $\begin{bmatrix} 000\\001 \end{bmatrix}$ . Figures 1 to 5 show the ring graphs which correspond to the de Bruijn rings given above.

The following lemmata describe some interesting properties of ring graphs, which we will use for proving theorem 1:

**Lemma 2.** Any  $(m, n)_k$ -ring graph has exactly  $M(k^n, m)$  edges.

*Proof.* By definition of the edges of a ring graph, each row-Lyndon pattern P of shape (m, n) corresponds to exactly one edge of label R from a vertex V to some other vertex by P = lexmin([V, R]). This implies the lemma.

#### **Lemma 3.** Any $(m, n)_k$ -ring graph is weakly connected.

Proof. Let  $0, 1 \in \Sigma$  be two different characters with 0 < 1. Now let V be an arbitrary vertex of the ring graph. Since V = lexmin(V) and  $R := [0, ..., 0, 1]^T$  is row-Lyndon, the concatenation [V, R] is row-Lyndon as well and thus V has an outgoing edge being labelled with R. This edge directs to a vertex V', of which the number of rightmost columns being equal to R is increased by one compared to V itself if not all columns of V are already equal to R. In that last case, the edge goes from V to V itself. It follows that there is a directed path from every vertex to the vertex [R, R, ..., R].

#### **Lemma 4.** Any $(m, n)_k$ -ring graph is Eulerian.

Proof. Let [L, C] be a vertex of a ring graph with  $L \in \Sigma^{m,1}$  and  $C \in \Sigma^{m,n-2}$ . Further, let  $R \in \Sigma^{m,1}$ . Then lexmin([L, C, R]) is a row-Lyndon pattern iff lexmin([R, L, C]) is a low-Lyndon pattern, since the row-periodicity does not depend on the order of the columns. With lexmin([L, C, R]) being row-Lyndon, let  $L' \in \Sigma^{m,1}, C' \in \Sigma^{m,n-2}$  and  $R' \in \Sigma^{m,1}$  be uniquely defined by lexmin([R, L, C]) = lexmin([L', C', R']), [L', C'] = lexmin([L', C']), and R' being the lexicographically smallest such choice if <math>[L', C'] is row-periodic. It follows that there is an edge of label R from [L, C] to lexmin([C, R]) iff there is an edge of label R' from [L', C', R'] = [L, C]. This defines a one-to-one mapping of ingoing and outgoing edges of the arbitrarily chosen vertex [L, C]. Thus, each vertex of the ring graph has its in-degree equal to its out-degree, and due to the ring graph being weakly connected, each ring graph is Eulerian.

Now we prove theorem 1 by constructing  $(m, n)_k$ -de Bruijn rings based on Eulerian cycles of  $(m, n)_k$ -ring graphs:

Proof (of Theorem 1). Let  $C = (V_0, R_1, V_1, R_2, V_2, \ldots, R_{M(k^n,m)}, V_{M(k^n,m)} = V_0)$ be an Eulerian cycle of an  $(m, n)_k$ -ring graph which starts and ends in the vertex  $V_0 := 0^{m,n-1}$  with  $0 \in \Sigma$ . Such a cycle can be constructed in linear time by using Hierholzers algorithm [7]. We now build a sub-perfect map using algorithm 1.

In every round of the for loop, algorithm 1 adds a new column to Ring. For i = 1, Ring is set equal to  $[V_0, R_1]$  with  $R_1$  being an edge from  $V_0$  to  $V_1$ .  $[V_0, R_1]$  is a row-Lyndon pattern which ends with  $V_1$ . Now suppose, after i - 1 rounds of the loop, Ring ends with  $V_{i-1}$  for some  $2 \le i \le M(k^n, m) - (n-1)$ . Then there exists an edge  $R_i$  from  $V_{i-1}$  to  $V_i$  and  $lexmin([V_{i-1}, R_i])$  is row-Lyndon. Moreover, the last n-1 columns of Ring are equal to  $V_i$  up to vertical rotation. Thus, vertically rotating Ring accordingly lets it end with  $V_i$ . It follows by induction, that each n-column subset of Ring is a vertical rotation of a unique row-Lyndon pattern. After  $M(k^n, m) - (n-1)$  loops, Ring is of length  $M(k^n, m)$ 

Algorithm 1 Generation of a de Bruijn ring.

**Require:** An Eulerian cycle  $\overline{C} = (V_0, R_1, V_1, R_2, V_2, \dots, R_{M(k^n, m)}, V_{M(k^n, m)})$  of an  $(m, n)_k$ -ring graph using the alphabet  $\Sigma$ . **Require:**  $V_0 = 0^{m, n-1}$  for some  $0 \in \Sigma$ .

 $Ring \leftarrow V_0$ for i = 1 to  $M(k^n, m) - (n - 1)$  do  $Ring \leftarrow [Ring, R_i]$ Rotate Ring vertically, such that it ends with  $V_i$ end for return Ring

and thus contains every row-aperiodic  $(m, n)_k$  pattern. Moreover it is a cyclic map, since the remaining n-1 edges in C lead to  $V_{\rm M}(k^n, m) = V_0$  and thus are labelled with  $0^m$ . Thus, *Ring* is an  $(m, {\rm M}(k^n, m); m, n)_k$  sub-perfect map. Since there cannot be a bigger sub-perfect map of height m, it is an  $(m, n)_k$ -de Bruijn ring.



Fig. 1. The  $(2, 2)_2$ -ring graph.

Note, that the generation of de Bruijn rings is both time- and space-efficient, as the construction of ring graphs, the construction of one of their Euler cycles using the Hierholzer algorithm and algorithm 1 all have linear time and space complexity with respect to the number of aperiodic patterns.

In the following, we will use de Bruijn rings for constructing sub-perfect maps of more general sizes.

## 5 From Rings to larger Sub-Perfect Maps

Two sub-perfect maps with patterns of the same shape(m, n) but not necessarily different alphabet sizes  $k_1, k_2$  can be easily combined to a larger sub-perfect map with patterns of the same shape but larger alphabet size  $k_1k_2$ :

**Lemma 5.** Let  $A_1$  be a sub-perfect map of type  $(M_1, N_1; m, n)_{k_1}$  and  $A_2$  be a sub-perfect map of type  $(M_2, N_2; m, n)_{k_2}$ . Then there exists a sub-perfect map A of type  $(\operatorname{lcm}(M_1, M_2), \operatorname{lcm}(N_1, N_2); m, n)_{k_1k_2}$ .





**Fig. 3.** The  $(2, 2)_3$ -ring graph. If no label is given at an edge, the label is equal to the target node label. A bidirectional edge represents two edges in opposing directions (with not necessarily equal labels).



Fig. 4. The  $(2,3)_2$ -ring graph. If no label is given at an edge, the label follows uniquely from the rules given in definition 2.



**Fig. 5.** The  $(4, 2)_2$ -ring graph. For better readability, the edge labels (which are rotations of the vertex label the edge is pointing to) are omitted and only the number of labels per edge and direction is given.

*Proof.* Let  $\Sigma_1$  and  $\Sigma_2$  be the alphabets of  $A_1$  and  $A_2$ , respectively. We now simply lay the cyclic maps onto each other by combining  $\Sigma_1$  and  $\Sigma_2$  to the product alphabet  $\Sigma_1 \times \Sigma_2$ , which has  $k_1 k_2$  elements. This new map has a vertical period length of  $(\operatorname{lcm}(M_1, M_2))$  and a horizontal period length of  $\operatorname{lcm}(N_1, N_2)$ .

Note, that if both  $A_1$  and  $A_2$  are perfect maps, the combined map A is a perfect map as well, iff  $M_1$  and  $M_2$  are coprime and  $N_1$  and  $N_2$  are coprime.

Lemma 5 can easily be applied to de Bruijn rings. E.g. when rotating one of two de Bruijn rings by 90°, it follows corollary 2.

**Corollary 2.** Given two alphabets  $\Sigma_1$  and  $\Sigma_2$  with  $|\Sigma_1| = k_1$  and  $|\Sigma_2| = k_2$ , there exists a sub-perfect map of type  $(M, N; m, n)_{k_1k_2}$  with

 $M = \text{lcm}(m, M(k_2^m, n)) \text{ and } N = \text{lcm}(n, M(k_1^n, m)).$ 

*Proof.* According to theorem 1, there exists an  $(m, n)_{k_1}$ -de Bruijn ring  $A_1$  and an  $(n, m)_{k_2}$ -de Bruijn ring  $A_2$ . By rotating  $A_2$  by 90°, we get a sub-perfect map  $A'_2$  of type  $(M(k_2^m, n), n; m, n)_k$ . By stacking  $A_1$  and  $A'_2$  onto each other and combining the alphabets to  $\Sigma = \Sigma_1 \times \Sigma_2$ , we get a sub-perfect map of type  $(\operatorname{lcm}(m, M(k_2^m, n)), \operatorname{lcm}(n, M(k_1^n, m)); m, n)_{k_1k_2}$ .

For example, using the  $(2, 2)_2$ -de Bruijn ring given above for both  $A_1$  and  $A_2$ , and when interpreting the tuples of  $\Sigma = \Sigma_1 \times \Sigma_2$  as two-digit binary numbers and writing them as decimals, one gets the following sub-perfect map of type  $(6, 6; 2, 2)_4$ :

[0]	0	2	2	0	0
1	2	1	2	3	2
0	1	2	3	0	1
1	3	1	3	3	3
1	0	3	2	1	0
1	2	1	2	3	2

However, since the with 6 and the height 2 are not coprime, the resulting map is far away from containing almost all patterns. In fact, it contains only 36 of 256 possible (2, 2)-patterns of a 4-letter alphabet. A simple trick is to remove the fifth column of the de Bruijn ring and use this sub-perfect map as  $A_1$  and  $A_2$ . Now, the width and height are coprime and the resulting combined sub-perfect map has size (10, 10):

$\begin{bmatrix} 0 & 0 & 2 & 2 & 0 & 0 & 0 & 2 & 2 & 0 \end{bmatrix}$
$1\ 2\ 1\ 2\ 3\ 0\ 3\ 0\ 3\ 2$
0123010321
$1\ 3\ 1\ 3\ 3\ 1\ 3\ 1\ 3\ 3$
$1\ 0\ 3\ 2\ 1\ 0\ 1\ 2\ 3\ 0$
0202202022
$1\ 0\ 3\ 2\ 1\ 0\ 1\ 2\ 3\ 0$
0303212123
1133111331
$\begin{bmatrix} 1 \ 2 \ 1 \ 2 \ 3 \ 0 \ 3 \ 0 \ 3 \ 2 \end{bmatrix}$

In this case, the numbered of covered patterns increases to almost three times the original number, but is still less than half the number of all possible patterns. However, as we show, the trick of removing columns to ensure coprime dimensions of  $A_1$  and  $A_2$  can be used in general, and in case of larger rings, the percentage of covered patterns quickly grows near 100% with increasing m, n. E.g. when using the already given  $(3, 2)^2$ -de Bruijn ring as  $A_1$  and the  $(2, 3)_2$ -de Bruijn ring as  $A_2$ , one gets a sub-perfect map of type  $(84, 20; 3, 2)_4$  without using the trick and a sub-perfect map of type  $(84, 38; 3, 2)_2$  by removing the second column from  $A_1$ . This map covers 3192 of 4096 patterns.

In general, if m and  $M(k_2^m, n)$  are not coprime or if n and  $M(k_1^n, m)$  are not coprime, then the resulting sub-perfect map contains only a fraction of all possible (m, n)-patterns over  $\Sigma$  (at most half of them).

However, we are interested in families of almost perfect maps, where the maps contain almost every pattern.

**Definition 3 (Almost perfect map).** Let k > 2 and  $\mathcal{M}$  be a set of subperfect maps for an alphabet of size k, such that for any tuple  $m', n' \ge 1$  there exists a sup-perfect map of type  $(M, N; m, n)_k$  in  $\mathcal{M}$  with m > m' and n > n'. A sequence  $(A_i)_{i \in \mathbb{N}}$  of sub-perfect maps for an alphabet of size k is called an increasing sequence if for any elements  $A_i$  of type  $(M_i, N_i; m_i, n_i)_k$  and  $A_j$  of type  $(M_j, N_j; m_j, n_j)_k$  with j > i it follows  $m_j > m_i$  and  $n_j > n_i$ . We call a set of sub-perfect maps  $\mathcal{M}$  a family of almost perfect maps, if for any increasing sequence  $(A_i)_{i \in \mathbb{N}}$  with  $A_i$  being of type  $(M_i, N_i; m_i, n_i)_k$  it follows

$$\lim_{i \to \infty} \frac{M_i N_i}{k^{m_i n_i}} = 1.$$

We call  $\mathcal{M}$  a family of perfect maps, if  $M_i N_i = k^{m_i n_i}$  for any element of type  $(M_i, N_i; m_i, n_i)_k$ .

Theorem 1 directly implies that  $(m, n)_k$ -de Bruijn rings are a family of almost perfect maps for any k, while the construction method given in lemma 5 does not give a family of almost perfect maps since m and  $M(k_2^m, n)$ , as well as n and  $M(k_1^n, m)$  are not necessarily coprime. However, the trick being used above can be used for arbitrary de Bruijn rings, such that stacking them leads to families of almost perfect maps:

**Theorem 2.** Given two alphabets  $\Sigma_1$  and  $\Sigma_2$  with  $|\Sigma_1| = k_1 \ge 2$  and  $|\Sigma_2| = k_2 \ge 2$ , and given a pattern shape (m, n) with  $|m - n| \le 2$ , there exists a subperfect map of type  $(M, N; m, n)_{k_1k_2}$  with

$$M = m \cdot (\mathbf{M}(k_2^m, n) - (|m' - 1| \cdot [m' < m - 1]))$$

$$\geq m \cdot (\mathbf{M}(k_2^m, n) - \max(1, m - 3))$$
(12)

and

$$N = n \cdot (\mathcal{M}(k_1^n, m) - (|n' - 1| \cdot [n' < n - 1]))$$

$$\geq n \cdot (\mathcal{M}(k_1^n, m) - \max(1, n - 3)),$$
(13)

where

$$m' := \operatorname{M}(k_2^m, n) \mod m$$

and

 $n' := \mathcal{M}(k_1^n, m) \bmod n.$ 

*Proof.* By definition of m' it follows that  $M(k_2^m, n) - m'$  is a multiple of m. Thus,  $M(k_2^m, n) - m' \pm 1$  and m are coprime. Now let

$$m'' := |m' - 1| \cdot [m' < m - 1]$$

Then m' and m'' either differ by one, or m' = m-1 and m'' = 0. Thus  $M(k_2^m, n) - m''$  and m are coprime. Since  $0 \le m'' \le \max(1, m-3)$ , it follows

$$M := \operatorname{lcm}(m, \operatorname{M}(k_2^m, n) - m'')$$
(14)

$$= m \cdot (\mathbf{M}(k_2^m, n) - m'') \tag{15}$$

$$\geq m \cdot (M(k_2^m, n) - \max(1, m - 3)).$$

Analogously it follows

$$N := \operatorname{lcm}(n, \operatorname{M}(k_1^n, m) - n'')$$
(16)

$$= n \cdot (\mathbf{M}(k_1^n, m) - n'') \tag{17}$$

$$\geq n \cdot (\mathcal{M}(k_1^n, m) - \max(1, n-3))$$

with

$$n'' := |(\mathbf{M}(k_1^n, m) \mod n) - 1| \cdot [n' < n - 1].$$

It remains to be shown, that in case of n'' > 0 and/or m'' > 0 there exist sub-perfect maps of types

$$(m, \mathcal{M}(k_1^n, m) - m''; m, n)_{k_1}$$
 and  
 $(n, \mathcal{M}(k_2^m, n) - n''; n, m)_{k_2}$ .

Given some alphabet  $\Sigma_1$ ,  $|\Sigma_1| = k_1$ , let A be an  $(m, n)_{k_1}$ -de Bruijn ring, i.e. a sub-perfect map of type  $(m, M(k_1^n, m); m, n)_{k_1}$ . Now let  $0, 1 \in \Sigma_1$ . Then each of the (m, n)-patterns with the first i rows  $(1 \leq i < m)$  being equal to 0 and the last m - i rows being equal to 1 is contained in A. Removing one column of one such pattern from A removes this pattern and its vertical rotations from A but still results in a sub-perfect map. Since there are m - 1 such patterns, we can construct a sub-perfect map of type  $(m, M(k_1^n, m) - i; m, n)_{k_1}$  for any  $1 \leq i < m$  which includes  $i := n'' \leq \max(1, n - 3) \leq m - 1$ .

Analogously, it follows the existence of a sub-perfect map of type  $(n, \mathcal{M}(k_2^m, n) - j; n, m)_{k_2}$  for any  $1 \leq j < n$  which includes  $j := m'' \leq \max(1, m - 3) \leq n - 1$ .

**Theorem 3.** The set of sub-perfect maps given in theorem 2 for  $k = k_1k_2$  with  $k_1, k_2 \ge 2$  is a family of almost perfect maps.

*Proof.* Let  $(A_i)_{i \in \mathbb{N}}$  be an increasing sequence of such sub-perfect maps of type  $(M_i, N_i; m_i, n_i)_k$ . Then for  $k = k_1 k_2$  it follows

$$\frac{M_i N_i}{k^{m_i n_i}} \ge \frac{m_i \left(\mathcal{M}(k_2^{m_i}, n_i) - m_i + 3\right) \cdot n_i \left(\mathcal{M}(k_1^{n_i}, m_i) - n_i + 3\right)}{(k_1 k_2)^{m_i n_i}} \tag{18}$$

$$=\frac{\left(m_{i}\mathrm{M}(k_{2}^{m_{i}},n_{i})-m_{i}^{2}+3m_{i}\right)\cdot\left(n_{i}\mathrm{M}(k_{1}^{n_{i}},m_{i})-n_{i}^{2}+3n_{i}\right)}{k_{2}^{m_{i}n_{i}}k_{1,i}^{m_{i}n_{i}}}$$
(19)

$$> \frac{k_2^{m_i n_i} - k_2^{(\lfloor m_i/2 \rfloor + 1)n_i} - m_i^2 + 3m_i}{k_2^{m_i n_i}} \cdot \frac{k_1^{n_i m_i} - k_1^{(\lfloor n_i/2 \rfloor + 1)m_i} - n_i^2 + 3n_i}{k_1^{m_i n_i}}$$
(20)

For  $k_1, k_2 \geq 2$  and  $m_i, n_i \to \infty$ , the last term reduces to  $\frac{k_2^{m_i n_i}}{k_2^{m_i n_i}} \cdot \frac{k_1^{m_i n_i}}{k_1^{m_i n_i}}$ , thus  $\lim_{i\to\infty} \frac{M_i N_i}{k_1^{m_i n_i}} = 1$ .

In case of m = n and  $k_1 = k_2$ , i.e. for square shaped patterns of perfect square alphabet sizes we get square shaped maps:

**Corollary 3.** Given an alphabet  $\Sigma$  with  $|\Sigma| = k \ge 2$ , there exists for any  $n \ge 2$  a sub-perfect map of type  $(N, N; n, n)_{k^2}$  with

$$N = n \cdot (\mathcal{M}(k^n, n) - (|n'-1| \cdot [n' < n-1]))$$
(21)

$$\geq n \cdot (\mathbf{M}(k^n, n) - \max(1, n-3))$$

$$> k^{n^2} - k^{(\lfloor n/2 \rfloor + 1)n} - \max(n, n^2 - 3n),$$
(22)

with

$$n' = \mathcal{M}(k^n, n) \mod n$$

and this set of sub-perfect maps is a family of almost perfect maps.

**Theorem 4.** Given any alphabet  $\Sigma$  with  $|\Sigma| = k \ge 2$  and any prime p, there exists a sub-perfect map of type  $(N, N; p, p)_{k^2}$  with  $N = p \cdot (M(k^p, p) - 1) = k^{p^2} - k^p - p$ .

*Proof.* Since p is prime, it follows  $M(k^p, p) = \frac{k^{p^2} - k^p}{p}$ . If the nominator  $k^{p^2} - k^p$  is a multiple of  $p^2$ , it follows, that  $M(k^p, p)$  is a multiple of p and thus  $M(k^p, p) - 1$  is coprime to p. Then, according to corollary 3 there exists a sub-perfect map of type  $(N, N; p, p)_{k^2}$  with  $N = p \cdot (M(k^p, p) - 1) = k^{p^2} - k^p - p$ . Thus, we only need to show that  $k^{p^2} \equiv k^p \pmod{p^2}$ , or equivalently the p-adic valuation  $\nu_p(k^{p^2} - k^p)$  is at least 2.

Depending on p, the lifting-the-exponent (LTE) lemma [14] implies the following:

-  $p \mid k$ : In this case obviously  $p^2 \mid k^p$  and  $p^2 \mid k^{p^2}$ , which implies  $0 \equiv k^{p^2} \equiv k^p \pmod{p^2}$ .

17

- -p k, p = 2: Due to the LTE lemma it follows  $ν_2(k^{p^2} k^p) = ν_2(k^2 k) + ν_2(k^2 + k) + ν_2(2) 1 = ν_2(k^2 k) + ν_2(k^2 + k)$ , which is at least 2 since  $k^2 \pm k$  is always even.
- $-p \nmid k, p \text{ odd: This implies } p \nmid k^p$ . In this case, the LTE lemma gives  $\nu_p(k^{p^2} k^p) = \nu_p(k^p k) + \nu_p(p) = \nu_p(k^p k) + 1$ , which is at least 2 due to Fermat's little theorem.

As an example, theorem 4 says, that for p = 3, there is a sub-perfect map type (501, 501; 3, 3)<sub>4</sub>, which covers more than 95.7% of all possible (3, 3) patterns of an alphabet of size  $2^2 = 4$ . Such a map requires just to build a  $(3,3)_2$ -de Bruijn ring, which is a matrix of size (3, 168) and which can efficiently be constructed.

Analogously, a  $(5,5)_2$ -de Bruijn ring, which is of size (5,6710880), can be used to construct a sub-perfect map of type  $(33554395, 33554395; 5,5)_4$ , which covers more than 99.9997% of all (5,5) patterns of an alphabet of size 4, and a  $(7,7)_2$ -de Bruijn ring of size (7,80421421917312) gives a sub-perfect map of type  $(562949953421177, 562949953421177; 7,7)_4$  covering more than 99.99999999995% of all patterns. Table 2 gives some further examples.

Theorems 2 and 4 allow to generate sub-perfect maps which are perfectly suitable for For optical localization tasks, since:

- 1. every local (m, n)-pattern occurs at most once.
- 2. Almost every possible (m, n)-pattern is contained in the map.
- 3. The code can efficiently be generated for arbitrary  $m, n, k \ge 2$  with k being not prime (such that it can be decomposed into some  $k_1$  and  $k_2$ ).
- 4. The decoding is quite efficient, as one only needs to store a  $(m, n)_k$ -de Bruijn ring, which reduces the number of entries to the square root of the whole pattern.

In the spacial coding context, the decoding process should be efficient, i.e. given some local pattern it should easily be possible to derive its position in the whole map. For sub-perfect maps being constructed according to theorem 2 and its specializations, this can be done as follows: The local pattern of alphabet size  $k = k_1k_2$  is first separated into the two patterns of alphabet sizes  $k_1$  and  $k_2$ , then by using a lookup table of size of the modified de Bruijn rings used, one gets the position in these modified rings, which can directly be combined to the position in the combined map. Such a decoding algorithm has time complexity O(mn) while the space complexity is linear with regard to the size of the de Bruijn rings and thus of order  $O(\sqrt{MN})$ .

If space complexity needs to be further restricted at the cost of time complexity, one could generate the de Bruijn rings implicitly for every lookup (with choosing a fixed selection scheme for each next edge in the Hierholzer algorithm) which results in time complexity  $O(\sqrt{MN})$  and space complexity O(mn) when the ring graphs are not stored permanently.

## 6 Conclusion and Future Work

We have introduced de Bruijn rings, i.e. sub-perfect maps of minimal height, which can be constructed by generating Eulerian cycles in a graph, which gener-

k	n	$N/ ilde{N}$	$N^2/\tilde{N}^2$ in %
2	2	$\frac{10}{16}$	39.06250000000000%
2	3	$\frac{501}{512}$	95.749282836914063%
2	4	$\tfrac{65276}{65536}$	99.208116903901100%
2	5	$\frac{33554395}{33554432}$	99.999779462935919%
2	6	$\tfrac{68719210554}{68719476736}$	99.999225309920504%
3	2	$\frac{70}{81}$	74.683737235177560%
3	3	$\frac{19653}{19683}$	99.695400725179510%
3	4	$\frac{43040156}{43046721}$	99.969500583377069%
3	5	$\frac{847288609195}{847288609443}$	99.999999941460331%
3	6	$rac{150094634909047936}{150094635296999136}$	99.999999483057906%
4	2	$\frac{238}{256}$	86.431884765625000%
4	3	$\frac{262077}{262144}$	99.948889588995371%
4	4	$\tfrac{4294901756}{4294967296}$	99.996948079208892%
4	5	$\tfrac{1125899906841595}{1125899906842624}$	99.999999999817220%
4	6	$\frac{4722366482800908959744}{4722366482869645213696}$	99.999999997088906%
5	2	$\frac{598}{625}$	91.546623999999994%
5	3	$\frac{1952997}{1953125}$	99.986893229496729%
5	4	$\tfrac{152587499996}{152587890625}$	99.999487995412494%
5	5	$\frac{298023223876950016}{298023223876953152}$	99.999999999997897%
5	6	$\frac{14551915228363036345499648}{14551915228366852423942144}$	99.999999999947562%

**Table 2.** Size comparison of square shaped sub-perfect maps according to theorem 2 and (not guaranteed to exist) de Bruijn tori of the same type. When N is the side length of a sub-perfect map,  $\tilde{N}$  denotes the side length of the corresponding de Bruijn torus. The second column gives the ratio of covered to all  $(n, n)_{k^2}$ -patterns.

alizes a construction method of de Bruijn sequences to two dimensions. The time complexity of this construction is linear in the number of contained patterns and a de Bruijn ring contains every pattern which does not only consists of constant columns.

We further introduced the concept of a family of almost perfect maps and show, that de Bruijn rings are almost perfect in that sense, which practically means, that they contain almost all possible patterns. Moreover, we showed, how de Bruijn rings can be used to construct families of almost perfect maps of other rectangular shapes, including the cases for which perfect maps are not yet known. This construction allows a decoding scheme which is sublinear (with regard to the total numper of patterns) in time and space complexity. As a side product, we derive an explicit lower bound for the Mobius function.

The findings are are especially of practical interest in the spacial coding context, since here a map does not need to be perfect, but it should be efficiently constructible and decodible. The findings can easily be generalized to higher dimensions.

#### References

- Bailey, D.F.: Some Binomial Coefficient Congruences. Applied Mathematic Letters, 4(4), pp. 1–5 (1991).
- Cook, J.C.: Toroidal tilings from de Bruijn-Good cyclic sequences. Discrete Mathematics, 70, pp. 209-210 (1988).
- De Bruijn, N. G.: A combinatorial problem. Proceedings of the Section of Sciences of the Koninklijke Nederlandse Akademie van Wetenschappen te Amsterdam,49(7), pp. 758–764 (1946).
- Flye Sainte-Marie, C.: Solution to question nr. 48. l'Intermediaire des Mathematiciens, pp. 107–110 (1894).
- Fredricksen, H.: The lexicographically least de Bruijn cycle. Journal of Combinatorial Theory, 9(1), pp. 1–5 (1970). doi:10.1016/S0021-9800(70)80050-3
- Fredricksen, H., Maiorana, J.: Necklaces of Beads in k colors and k-ary de Bruijn sequences. Discrete Mathematics, 23(3), pp. 207—210 (1978). doi:10.1016/ 0012-365X(78)90002-X
- Hierholzer, C.: Über die Möglichkeit, einen Linienzug ohne Wiederholung und ohne Unterbrechung zu umfahren", Mathematische Annalen, 6 (1), pp. 30–32 (1873)
- Hurlbert, G., Isaak, G.: On the de Bruijn Torus Problem. Journal of Combinatorial Theory, Series A (64), pp. 50–62 (1993).
- 9. Klein, A.: Stream Ciphers. Springer, p. 59 (2013).
- Lothaire, M.: Combinatorics on words. Encyclopedia of Mathematics and Its Applications, 17, pp. 79–84. (1997).
- Mitchell, C.J., Paterson, K.G.: Decoding Perfect Maps. Designs, Codes and Cryptography 4, pp. 11-30 (1994).
- Möbius, A.F.: Über eine besondere Art von Umkehrung der Reihen. Journal für die reine und angewandte Mathematik, 9, pp. 105–123 (1832).
- Moreau, C.: Sur les permutations circulaires distinctes. Nouvelles Annales de Mathématiques, Série 2, 11, pp. 309 – 314 (1872).
- 14. Parvadi, A.H.: Lifting The Exponent Lemma (LTE), version 6 (04/2011). https://s3.amazonaws.com/aops-cdn.artofproblemsolving.com/resources/ articles/lifting-the-exponent.pdf (retrieved 02/2024).

- 20 P. Stelldinger
- Paterson, K.G.: Perfect Maps. IEEE Transactions on Information Theory, 40(3), pp. 743–753 (1994).
- Riedel, M.: Pólya's enumeration theorem and the symbolic method. Technical Report, University of Stuttgart (2021). pnp.mathematik.uni-stuttgart.de/iadm/ Riedel/papers/collier.pdf
- 17. Scheinerman, E.R.: Determining Planar Location Via Complement-Free de Bruijn Sequences Using Discrete Optical Sensors. IEEE Transactions on Robotics and Automation, 17(6), pp. 883–889 (2001).
- Schüsselbauer, D., Schmid, A:, Wimmer, R.: Dothraki: Tracking Tangibles Atop Tabletops Through De-Bruijn Tori. 15th Conference on Tangible, Embedded, and Embodies Interaction (2021).