

TROPICAL VECTOR BUNDLES

BIVAS KHAN AND DIANE MACLAGAN

ABSTRACT. The goal of this paper is to introduce a construction of a vector bundle on a tropical variety. When the base is a tropical toric variety these tropicalize toric vector bundles, and are described by the data of a valuated matroid and some flats in the lattice of flats of the underlying matroid. The fibers are tropical linear spaces. We define global sections for tropical toric vector bundles, stability, and Jordan-Hölder and Harder-Narasimhan filtrations. Many of these require additional modularity assumptions on the defining matroid.

1. INTRODUCTION

The goal of this paper is to introduce a notion of vector bundles on tropical varieties.

The theory of divisors, or equivalently line bundles, on tropical curves/metric graphs has been intensively studied since the beginning of tropical geometry. This has been particularly successful in Brill-Noether theory [CDPR12, JR21] and other aspects of the geometry of the moduli space M_g [FJP00].

The theory of higher rank vector bundles, by contrast, is much less well developed. After early work by Allermann [All12], the story was restarted recently in the context of metric graphs by Gross, Ulirsch, and Zakharov [GUZ22]. In [JMT24] Jun, Mincheva, and Tolliver defined and studied vector bundles on tropical toric varieties (in a slightly more general framework). However their construction has the drawback that all vector bundles are direct sums of line bundles. Similarly, all vector bundles on a simply connected tropical variety are trivial in Allermann's theory.

In this paper we give a definition of a vector bundle on a tropical toric variety that does not have these limitations, and define the tropicalization of an equivariant vector bundle on a toric variety. This definition then extends to vector bundles on subschemes of tropical toric varieties, in the sense of [MR18], and to the tropicalization of some vector bundles on subschemes of toric varieties.

Our approach is to first tropicalize presentations of toric vector bundles on a toric variety X_Σ . By [Cox95, Per03, Per04] an equivariant sheaf on a toric variety can be described by an $\mathbb{Z}^{|\Sigma(1)|}$ -graded module over the Cox ring of X_Σ , which is a polynomial ring with variables indexed by the rays of Σ . An equivariant surjection

$$(1.1) \quad \bigoplus_{\mathbf{w} \in \mathcal{G}} \mathcal{O}(D_{\mathbf{w}}) \rightarrow \mathcal{E}$$

from a direct sum of line bundles onto a toric vector bundle \mathcal{E} gives rise to a presentation of this module (see Section 4). This presentation can then be tropicalized in the sense of [MR18, MR22]. The surjection of line bundles induces the structure of a matroid on the index set \mathcal{G} , in the sense of DiRocco, Jabbusch, and Smith [DJS]. It also gives a linear embedding of the fibers into $\mathbb{A}^{|\mathcal{G}|}$, which tropicalizes to a tropical linear space in $\overline{\mathbb{R}}^{|\mathcal{G}|}$. See §3 and §5.4 respectively for more details.

Our definition of a tropical toric vector bundle on a tropical toric variety takes these properties as part of the definition, building on the Klyachko classification of toric vector bundles [Kly89]. We write $\text{trop}(X_\Sigma)$ for the tropical toric variety [Kaj08, Pay08] with fan Σ .

Definition 1.1. Let $\text{trop}(X_\Sigma)$ be a tropical toric variety. A *tropical toric reflexive sheaf* $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ on $\text{trop}(X_\Sigma)$ is given by a simple valuated matroid \mathcal{M} of rank r on a ground set \mathcal{G} , and for each ray ρ_i of Σ a collection $\{E^i(j) : j \in \mathbb{Z}\}$ of flats of the underlying matroid $\underline{\mathcal{M}}$ with the property that $E^i(j) \leq E^i(j')$ if $j > j'$, $E^i(j) = \emptyset$ for $j \gg 0$, and $E^i(j) = \mathcal{G}$ for $j \ll 0$.

The reflexive sheaf \mathcal{E} is a *tropical toric vector bundle* if in addition for every maximal cone σ of Σ there is a multiset $\mathbf{u}(\sigma) \subseteq M$, and a basis $B_\sigma = \{\mathbf{w}_{\mathbf{u}} : \mathbf{u} \in \mathbf{u}(\sigma)\}$ of $\underline{\mathcal{M}}$ such that for every ray $\rho_i \in \sigma$

and $j \in \mathbb{Z}$ we have

$$(1.2) \quad E^i(j) = \bigvee_{\mathbf{u} \cdot \mathbf{v}_i \geq j} \mathbf{w}_{\mathbf{u}},$$

is the smallest flat containing all the $\mathbf{w}_{\mathbf{u}}$ with $\mathbf{u} \cdot \mathbf{v}_i \geq j$.

Klyachko [Kly89] showed that the category of toric vector bundles with equivariant morphisms is equivalent to a category of compatible filtrations of a vector space. Our definition is an abstraction of Klyachko's classification. The poset of all intersections of filtered pieces, ordered by inclusions, embeds into the lattice of flats of a matroid in the sense of [DJS]. See §3 for more details. In our setting the [DJS] matroid is the underlying matroid of \mathcal{M} . The compatibility relation (1.2) is the abstraction of the usual Klyachko compatibility relation.

A surjection of the form (1.1) induces a realizable valuated matroid \mathcal{M} , and thus a tropical toric vector bundle (see Definition 5.8). Tropical toric vector bundles that arise in this fashion are called *realizable*. This tropicalization depends on the choice of surjection. This is analogous to the way in which the tropicalization of a projective variety depends on its projective embedding, including the choice of coordinates on \mathbb{P}^n . However some tropical invariants are constant for any, or for many, choices of surjection.

While the tropicalization of toric vector bundles is the main motivation for our construction, not all tropical toric vector bundles are realizable. A tropical toric vector bundle can fail to be realizable in two different ways: if the valuated matroid \mathcal{M} is not realizable, or if the choice of flats $E^i(j)$ cannot come from a filtration of subspaces. See Examples 5.15, 5.29, and 6.5(2) for more on this phenomenon.

A coherent sheaf on a toric variety can be described by a module over the Cox ring of the toric variety. We describe this explicitly for toric vector bundles.

Proposition 1.2. *Let S be the Cox ring of a smooth toric variety X_{Σ} , and let \mathcal{E} be a rank- r toric vector bundle on X_{Σ} with an equivariant surjection of the form (1.1). Let G be an $r \times m$ matrix whose columns are a realization of the associated valuated matroid \mathcal{M} with ground set \mathcal{G} . Then \mathcal{E} is the sheafification \tilde{P} of the $\mathbb{Z}^{|\Sigma(1)|}$ -graded module P given by*

$$P = \bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}})/R,$$

where the $\mathbf{d}_{\mathbf{w}} \in \mathbb{Z}^{|\Sigma(1)|}$ encode the subspaces $E^i(j)$, and R is a submodule of $\bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}})$ obtained by homogenizing $\ker(G)$.

A precise statement is given in Proposition 4.3. This motivates the following definition in the tropical setting.

Definition 1.3. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric vector bundle on a tropical toric variety $\text{trop}(X_{\Sigma})$, and let $S = \overline{\mathbb{R}}[x_1, \dots, x_s]$ be the Cox semiring of $\text{trop}(X_{\Sigma})$. Then \mathcal{E} can be described by the semimodule

$$(1.3) \quad P = \bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}})/\mathcal{B}(R),$$

where $\mathcal{B}()$ is the Giansiracusa bend congruence, and R is a homogenization of the equations for the tropical linear space of \mathcal{M} .

A consequence of this is a description of the total space of a tropical toric vector bundle as a tropical subvariety of $\text{trop}(X_{\Sigma}) \times \overline{\mathbb{R}}^m$. The fiber over a point in $\text{trop}(X_{\Sigma})$ is a tropical linear space in $\overline{\mathbb{R}}^m$. When the point is the tropicalization of the identity of the torus of X_{Σ} the corresponding tropical linear space is the one defined by the valuated matroid \mathcal{M} . See § 5.4 for more details.

Tropical toric vector bundles have the following properties that are familiar from toric vector bundles.

Theorem 1.4. (1) *There is a definition of $h^0(\mathcal{E})$ for a tropical toric vector bundle. When $\mathcal{E} = \text{trop}(\mathcal{F})$ and the matroid for \mathcal{F} is chosen carefully, we have $h^0(\mathcal{E}) = h^0(\mathcal{F})$ (Definition 6.13 and Lemma 6.15).*

- (2) *With extra modularity and minimality assumptions on \mathcal{M} , tropical toric vector bundles on $\text{trop}(\mathbb{P}^1)$ split as a direct sum of line bundles (Proposition 6.4).*
- (3) *There is a notion of stability for tropical toric vector bundles (see Section 7). When $\mathcal{E} = \text{trop}(\mathcal{F})$ stability for \mathcal{F} implies stability for \mathcal{E} . If the matroid for \mathcal{F} is chosen carefully, \mathcal{E} is stable if and only if \mathcal{F} is.*
- (4) *With extra modularity assumptions on \mathcal{M} , Jordan-Hölder and Harder-Narasimhan filtrations exist (Theorems 7.23 and 7.25).*

Part 1 of Theorem 1.4 is inspired by and generalizes [DJS]. Part 3 generalizes [Dev22] and [DDK20, DDK21]. The classical proof that every vector bundle on \mathbb{P}^1 splits as a direct sum of line bundles already relies in a crucial way in some foundational results from lattice theory, and Part 2 of the theorem can be regarded as abstracting back to those results.

Modularity is a property of lattices that abstracts the linear algebra fact that $\dim(V + W) + \dim(V \cap W) = \dim(V) + \dim(W)$ for two subspaces V, W of a finite-dimensional vector space. Lattices of flats of matroids are always *semimodular* (the analogous relation holds with \leq instead of $=$), but are much less likely to be modular, even when realizable. While we do not require complete modularity in Theorem 1.4, some form is necessary, as Part 2 is false without the modularity assumption (see Example 6.5). More work is required to completely understand this condition for realizable tropical toric vector bundles.

The work of Jun, Mincheva, and Tolliver [JMT24] also tropicalizes toric vector bundles. However in their setting all tropical vector bundles on $\text{trop}(X_\Sigma)$ are a direct sum of line bundles, which is far from the case in non-tropical algebraic geometry. The issue here is one would expect transition functions for a rank- r tropical vector bundle to live in a tropicalization of $\text{GL}(r)$ over a valued field K . However the naive form of tropical $\text{GL}(r)$ consists of tropical $r \times r$ matrices with only one entry not equal to ∞ in each row and column, so is isomorphic to $S_r \rtimes \mathbb{R}^r$. This is the tropicalization of those elements of $\text{GL}(r)$ that preserve the torus $(K^*)^r$ when acting on \mathbb{A}^r . A toric vector bundle is determined by a choice of basis for the fiber E of \mathcal{E} over the identity of the torus for each cone $\sigma \in \Sigma$, and such a matrix only permutes the elements of this basis. Thus we would have the same basis for every cone of Σ , and so the bundle \mathcal{E} would decompose as a direct sum of line bundles.

By contrast, in the present work we use the tropicalization of $\text{GL}(m)$ for transition maps, where $m = |\mathcal{G}|$ is the size of the ground set of the valuated matroid \mathcal{M} . This allows the construction of indecomposable tropical toric vector bundles. One way to think of the matroid \mathcal{M} in the realizable case is that we only require finitely many matrices in $\text{GL}(r)$ to construct the transition functions of a vector bundle, and a realization of \mathcal{M} can be thought of as a map from K^m to K^r that has all of the columns of these matrices among the images of the standard basis vectors.

We also extend the definition of tropical vector bundles to more general bases. Embedded tropicalization of a variety X in the sense of [MS15] proceeds by first embedding X into a toric variety. In an analogous fashion we *define* a tropical vector bundle on a tropical subscheme of a tropical toric variety to be the restriction of a tropical toric variety on the ambient toric variety. Vector bundles on X that can be tropicalized in this fashion are those that are the pullback of a toric vector bundle \mathcal{E} on a toric variety X_Σ under *some* embedding of X into X_Σ . Note that every line bundle on X has this form; for example, when \mathcal{L} is a very ample line bundle corresponding to an embedding $\phi: X \rightarrow \mathbb{P}^m$, we have $\mathcal{L} = \phi^*(\mathcal{O}_{\mathbb{P}^m}(1))$. As with the tropicalization of X , the freedom in the choice of embedding is crucial to get interesting answers. It would be interesting to understand which vector bundles can be constructed in this fashion. This is covered in Section 8.

The tropicalization of vector bundles on curves was considered already by Gross, Ulirsch, and Zakharov [GUZ22], in the context of tropicalization taking curves to metric graphs. Their vector bundles are constructed as pushforwards of tropical line bundles on a cover of the metric graph, and the vector bundles on a classical curve that can be tropicalized are those that can be realized as a pushforward in this sense. It will also be important to understand the interaction between this approach and the one of this paper, and in particular which vector bundles on curves can be tropicalized in each sense.

As this paper was nearing completion we became aware of the similar project of Kiumars Kaveh and Chris Manon on *Toric matroid bundles* [KM24]. Their toric matroid bundles are the same as our tropical

toric vector bundles in the case that the valuated matroid structure is trivial. Both works have the same definition of first Chern class and global sections, and highlight the importance of modularity.

The structure of this paper is as follows. In Section 2 we recall the necessary background on tropical geometry and lattice theory needed in the paper. In Section 3 we explain the connection between toric vector bundles and matroids. Section 4 describes the Cox module of a toric vector bundle. The tropical material begins in Section 5, with the definition of a tropical toric reflexive sheaf and a tropical toric vector bundle, and a description of the total space of a tropical toric vector bundle. Further properties of tropical toric vector bundles are explained in Section 6: decomposition into direct sums, tensor product with line bundles, and parliaments of polytopes. Stability of tropical toric vector bundles, and the Jordan-Hölder and Harder-Narasimhan filtrations, are covered in Section 7. Finally, Section 8 extends these definition to tropical vector bundles on more general tropical schemes.

Acknowledgements: We thank Michel Brion for pointing out Remark 8.9(2), and Lucie Devey for conversations about [Dev22]. Thanks are also due to Chris Manon for explaining his earlier work with Kiumars Kaveh, and to both Kaveh and Manon for explaining their forthcoming work. MacLagan also thanks Milena Hering and Greg Smith for many useful conversations over the years about toric vector bundles, and Georg Loho for useful conversations about polymatroids. Both authors were partially supported by EPSRC grant EP/R02300X/1. Khan was also supported by postdoctoral fellowships from the National Board for Higher Mathematics and a grant from the Infosys Foundation, and MacLagan was also supported by ESPRC Fellowship EP/X02752X/1.

Notation: We follow standard notation for toric geometry as in [Ful93] or [CLS11], so $N \cong \mathbb{Z}^n$ always denotes a rank- n lattice, and $M = \text{Hom}(N, \mathbb{Z})$ is the dual lattice. We write $s = |\Sigma(1)|$ for the number of rays of the fan Σ .

We use \mathcal{M} to denote a valuated matroid, and $\underline{\mathcal{M}}$ to denote the corresponding underlying matroid of \mathcal{M} . The ground set \mathcal{G} of \mathcal{M} has size m . Note the typographical distinction between the lattice $M \cong \mathbb{Z}^n$ and the valuated matroid \mathcal{M} .

The symbol \sqcup denotes disjoint union of sets.

2. BACKGROUND ON TROPICAL AND LATTICE TECHNIQUES

We recall here three key concepts from tropical geometry: tropical toric varieties, valuated matroids, and the Giansiracusa bend congruence. We also recall the basics of lattice theory that we will need.

We follow standard notation for toric varieties from [Ful93] or [CLS11]. We write n for the dimension of the toric variety X_Σ , $s = |\Sigma(1)|$ for the number of rays, $\mathbf{v}_1, \dots, \mathbf{v}_s$ for the first lattice points on the rays ρ_1, \dots, ρ_s of the fan Σ , and D_i for the torus-invariant divisor corresponding to the i th ray of Σ . The lattice of X_Σ is $N \cong \mathbb{Z}^n$, and $M = \text{Hom}(N, \mathbb{Z})$ is the dual lattice, where the homomorphism is as abelian groups. We denote by $N_{\mathbb{R}}$ the vector space $N \otimes \mathbb{R}$. We assume unless otherwise noted that the fan Σ is complete.

We denote the tropical semiring by $\overline{\mathbb{R}} = (\mathbb{R} \cup \{\infty\}, \oplus, \odot)$, where $\oplus = \min$, and $\odot = +$. Let K be a field equipped with a valuation $\text{val} : K \rightarrow \overline{\mathbb{R}}$. For a vector $\mathbf{v} \in \overline{\mathbb{R}}^n$ we denote by $\text{supp}(\mathbf{v})$ the set $\{i : v_i \neq \infty\}$.

2.1. Tropical toric varieties. Recall that for a pointed rational cone $\sigma \subseteq N_{\mathbb{R}}$ the affine toric variety U_σ is $\text{Spec}(K[\sigma^\vee \cap M])$, with K -rational points $\text{Hom}(K[\sigma^\vee \cap M], K) = \text{Hom}(\sigma^\vee \cap M, K)$, where the first homomorphism is as K -algebras, and the second is as semigroup homomorphisms from $(\sigma^\vee \cap M, +)$ to (K, \times) . The *tropical affine toric variety* $\text{trop}(U_\sigma)$ is

$$\text{Hom}(\sigma^\vee \cap M, \overline{\mathbb{R}}),$$

where the homomorphism is as semigroups from $(\sigma^\vee \cap M, +)$ to $(\overline{\mathbb{R}}, \odot)$. The topology on $\text{trop}(U_\sigma)$ comes from viewing it as a subspace of $\overline{\mathbb{R}}^{\sigma^\vee \cap M}$. Given a rational polyhedral fan Σ , the *tropical toric variety* $\text{trop}(X_\Sigma)$ is made by gluing together the sets $\text{trop}(U_\sigma)$ along faces as in the standard case: if τ is a face of σ , then σ^\vee is a sub-semigroup of τ^\vee , so $\text{Hom}(\tau^\vee \cap M, \overline{\mathbb{R}})$ is a subset of $\text{Hom}(\sigma^\vee \cap M, \overline{\mathbb{R}})$. The coordinate ring of $\text{trop}(U_\sigma)$ is the semiring $\overline{\mathbb{R}}[\sigma^\vee \cap M]$.

The Cox semiring of a tropical toric variety is

$$\text{Cox}(\text{trop}(X_\Sigma)) = \overline{\mathbb{R}}[x_1, \dots, x_s].$$

This is graded by the class group of X_Σ :

$$(2.1) \quad A^1(\Sigma) = \text{coker}(M \xrightarrow{\phi} \mathbb{Z}^s).$$

Here $(\phi(\mathbf{u}))_i = \mathbf{u} \cdot \mathbf{v}_i$, and the grading is $\deg(x_i) = \mathbf{e}_i + \text{im } \phi$.

For $\sigma \in \Sigma$ write $\mathbf{x}^\sigma = \prod_{\rho_i \notin \sigma} x_i$. The irrelevant ideal B_Σ is $\langle \mathbf{x}^\sigma : \sigma \in \Sigma \rangle \subseteq \overline{\mathbb{R}}[x_1, \dots, x_s]$. The tropical Cox construction of $\text{trop}(X_\Sigma)$ is

$$(\overline{\mathbb{R}}^s \setminus V(B_\Sigma))/H,$$

where $H = \text{im}(\phi \otimes \mathbb{R}) \subseteq \mathbb{R}^s$ is the tropical dual of the class group. For more details on tropical toric varieties, see [Kaj08], [Pay09], or [MS15, §6.2].

2.2. Valuated matroids. Valuated matroids are a generalization of matroids introduced by Dress and Wenzel [DW92] that have proved particularly useful in tropical geometry. We assume familiarity with the basics of matroid theory as in [Oxl92].

A valuated matroid \mathcal{M} of rank r on a finite set \mathcal{G} is given by a basis valuation function

$$\nu: \binom{\mathcal{G}}{r} \rightarrow \overline{\mathbb{R}}$$

satisfying that $\nu(B) \neq \infty$ for some $B \in \binom{\mathcal{G}}{r}$ and for all $B_1, B_2 \in \binom{\mathcal{G}}{r}$ and $i \in B_1 \setminus B_2$ there is $j \in B_2 \setminus B_1$ with

$$\nu(B_1) \circ \nu(B_2) \geq \nu(B_1 \setminus \{i\} \cup \{j\}) \circ \nu(B_2 \setminus \{j\} \cup \{i\}).$$

Any matroid can be given a valuated matroid structure by setting $\nu(B) = 0$ if B is a basis of the matroid, and $\nu(B) = \infty$ otherwise. We call this the trivial valuated matroid structure on the matroid. Conversely, every valuated matroid \mathcal{M} has an underlying matroid $\underline{\mathcal{M}}$ with bases $\{B \in \binom{\mathcal{G}}{r} : \nu(B) \neq \infty\}$. Two different functions ν_1, ν_2 define the same valuated matroid if there is a constant $\lambda \neq \infty$ such that $\nu_1(B) = \lambda \circ \nu_2(B)$. The matroids of ν_1 and ν_2 are *projectively equivalent* if there is $\mathbf{b} \in \mathbb{R}^{|\mathcal{G}|}$ with $\nu_1(B) = \nu_2(B) + \sum_{i \in B} b_i$ for all subsets B .

The *circuits* of a valuated matroid are the elements of $\overline{\mathbb{R}}^{|\mathcal{G}|}$ of the form

$$(2.2) \quad (C_{B,i})_j = \begin{cases} \lambda \circ \nu(B \cup \{i\} \setminus \{j\}) & j \in B \cup \{i\} \\ \infty & \text{otherwise} \end{cases}$$

for $\lambda \in \mathbb{R}$, B a basis of $\underline{\mathcal{M}}$, and $i \in \mathcal{G} \setminus B$. The circuit $C_{B,i}$ with $\lambda = -\nu(B)$ is the *fundamental circuit* of i over B . Note that we may have $C_{B,i} = C_{B',i'}$ if $B \neq B'$. A *vector* of \mathcal{M} is a tropical sum of circuits.

The set of circuits can be characterised by the circuit axioms [MT01]. These say that a collection \mathcal{C} of elements of $\overline{\mathbb{R}}^{|\mathcal{G}|}$ is the set of circuits of a valuated matroid if:

- (1) $(\infty, \dots, \infty) \notin \mathcal{C}$,
- (2) if $\mathbf{c} \in \mathcal{C}$, then $\lambda \circ \mathbf{c} \in \mathcal{C}$ for all $\lambda \in \mathbb{R}$,
- (3) we do not have $\text{supp}(\mathbf{c}) \subsetneq \text{supp}(\mathbf{c}')$ for $\mathbf{c}, \mathbf{c}' \in \mathcal{C}$, and
- (4) whenever $\mathbf{c}, \mathbf{c}' \in \mathcal{C}$ with $c_i = c'_i < \infty$ and $c_j < c'_j$ there is $\mathbf{c}'' \in \mathcal{C}$ with $\mathbf{c}'' \geq \mathbf{c} \oplus \mathbf{c}'$, $c''_i = \infty$, and $c''_j = c_j$.

An r -dimensional subspace of K^n gives rise to a valuated matroid on the ground set $\mathcal{G} = \{1, \dots, n\}$ by fixing an $r \times n$ matrix A whose rows form a basis for the subspace, and setting

$$\nu(B) = \text{val}(\det(A_B)),$$

where A_B denotes the submatrix of A with columns indexed by B . Different choices of the matrix A may scale ν by a constant. A collection of n vectors \mathcal{G} in K^r also induces a rank- r valuated matroid on \mathcal{G} in the same way by forming the matrix A with columns the vectors in \mathcal{G} . Valuated matroids coming from a matrix A in these ways are called *realizable*.

Deletion and contraction are fundamental operations for matroids, and these have versions for valuated matroids which we will need. Let $\mathcal{G}' \subseteq \mathcal{G}$ be a subset of the ground set \mathcal{G} of a valuated matroid \mathcal{M} of rank r . The *restriction* $\mathcal{M}|_{\mathcal{G}'}$, or deletion of $\mathcal{G} \setminus \mathcal{G}'$, is a valuated matroid on the ground set \mathcal{G}' with basis valuation function defined as follows. Let $r' = \text{rk}_{\underline{\mathcal{M}}}(\mathcal{G}')$. Fix $E \subseteq \mathcal{G} \setminus \mathcal{G}'$ with $\text{rk}(\mathcal{G}' \cup E) = \text{rk}(\mathcal{M})$. We define a basis valuation function $\nu': \binom{\mathcal{G}'}{r'} \rightarrow \overline{\mathbb{R}}$ by

$$\nu'(B) = \nu(B \cup E).$$

See [DW92, Proposition 1.2] for a proof that this defines a valuated matroid on \mathcal{G}' . While it depends on the choice of set E , all such choices define the same valuated matroid. The underlying matroid of $\mathcal{M}|_{\mathcal{G}'}$ is the restriction $\underline{\mathcal{M}}|_{\mathcal{G}'}$ of the underlying matroid of \mathcal{M} .

The *contraction* \mathcal{M}/\mathcal{G}' is the valuated matroid on $\mathcal{G} \setminus \mathcal{G}'$ with basis valuation function defined as follows. Let r'' be the rank of the contraction $\underline{\mathcal{M}}/\mathcal{G}'$ of the underlying matroid $\underline{\mathcal{M}}$. Fix $E \subseteq \mathcal{G}'$ with $|E| = \text{rk}(E) = r - r''$. We define a basis valuation function $\nu'': \binom{\mathcal{G} \setminus \mathcal{G}'}{r''} \rightarrow \overline{\mathbb{R}}$ by

$$\nu''(B) = \nu(B \cup E).$$

See [DW92, Proposition 1.3] for a proof that this defines a valuated matroid on $\mathcal{G} \setminus \mathcal{G}'$. While it depends on the choice of set E , all such choices define the same valuated matroid. The underlying matroid of \mathcal{M}/\mathcal{G}' is the contraction $\underline{\mathcal{M}}/\mathcal{G}'$ of the underlying matroid of \mathcal{M} .

2.3. The bend congruence. When working over semirings instead of rings, the notion of taking the quotient of a semimodule by a submodule is replaced by taking the quotient by a *congruence*. This is an equivalence relation \sim on the semimodule that is compatible with addition and multiplication by the semiring: $a \sim b, c \sim d$ implies $a + c \sim b + d$ and $\lambda a \sim \lambda b$.

In [GG16] Jeff and Noah Giansiracusa introduce a particular congruence on submodules of R^n when R is an idempotent semiring (such as $\overline{\mathbb{R}}$ or $\overline{\mathbb{R}}[x_1, \dots, x_s]$) that is particularly useful in tropical geometry. We now recall this in the context of the tropical semiring.

Definition 2.1. Let P be an $\overline{\mathbb{R}}$ -subsemimodule of $\overline{\mathbb{R}}^m$. We write $p \in P$ as $p = \sum_{i=1}^m p_i \mathbf{e}_i$ for $p_i \in \overline{\mathbb{R}}$, and set $p^{\hat{i}} = \sum_{j \neq i} p_j \mathbf{e}_j$. The *bend congruence* of P is the congruence $\mathcal{B}(P)$ on $\overline{\mathbb{R}}^m$ generated by

$$\{p \sim p^{\hat{i}} : p \in P, 1 \leq i \leq m\}.$$

This concept generalizes to ideals in polynomial semirings, or more generally monoid semirings. For a (tropical) polynomial $f = \sum c_{\mathbf{u}} \mathbf{x}^{\mathbf{u}} \in \overline{\mathbb{R}}[x_1, \dots, x_s]$ we denote by $f_{\hat{\mathbf{v}}}$ the polynomial

$$f_{\hat{\mathbf{v}}} = \sum_{\mathbf{u} \neq \mathbf{v}} c_{\mathbf{u}} \mathbf{x}^{\mathbf{u}}.$$

Definition 2.2. Fix a pointed rational polyhedral cone, and let I be an ideal in $\overline{\mathbb{R}}[\sigma^{\vee} \cap M]$. The *bend congruence* $\mathcal{B}(I)$ of I is the congruence on $\overline{\mathbb{R}}[\sigma^{\vee} \cap M]$ generated by

$$\{f \sim f_{\hat{\mathbf{v}}} : f \in I\}.$$

2.4. Lattices and modularity. We recall the notion and some basic properties of modular flats in geometric lattice from [Zie91, Chapter 3; Sta71]. Modular flats play an important role in our theory. Recall that a lattice is a poset where every pair of elements has a unique least upper bound (join \vee), and a unique greatest lower bound (meet \wedge). Geometric lattices are atomic lattices (every element is the join of rank one elements) that are semimodular ($\text{rk}(x \wedge y) + \text{rk}(x \vee y) \leq \text{rk}(x) + \text{rk}(y)$) for all x, y in the lattice). These are precisely the lattices of flats of matroids [Sta12, §3.3]. Hence, we also call elements of L flats.

A lattice is modular if

$$(2.3) \quad \text{rk}(x \vee y) + \text{rk}(x \wedge y) = \text{rk}(x) + \text{rk}(y)$$

for all x, y in the lattice. Lattices of flats of matroids are always semimodular, but not always modular.

Definition 2.3. We say that a pair (x, y) in a semimodular lattice is modular if the equality (2.3) holds [Bir67, §IV.3]. An element x is modular if (x, y) is a modular pair for *all* elements $y \in L$.

Proposition 2.4. [Zie91, Definition 2.1] *Let L be a geometric lattice. The following conditions are equivalent:*

- (1) *A flat $F \in L$ is modular.*
- (2) *For $G \leq H$ in L , $G \vee (F \wedge H) = (G \vee F) \wedge H$.*
- (3) *For $G \in L$ and $H \leq F$, $H \vee (G \wedge F) = (H \vee G) \wedge F$.*

Proposition 2.5. (1) [Zie91, Proposition 2.2(2)] *Let L be a geometric lattice and $F \in L$ be a modular flat. If G and H are flats in L with $G \leq H$, then*

$$G \vee (F \wedge H) = (G \vee F) \wedge H$$

is a modular flat of $[G, H]$.

- (2) [Zie91, Proposition 2.3] *Let \mathcal{M} be a matroid on a ground set \mathcal{G} . Then*
 - \emptyset, \mathcal{G} and all atoms are modular flats.
 - *The meet of modular flats is always modular.*

Remark 2.6. Let F be a modular flat in the lattice of flats of a matroid \mathcal{M} and F' be any other flat.

- (1) Then taking $G = \emptyset$, and $H = F'$ in Proposition 2.5(1), we see $F' \wedge F$ is a modular flat of $[\emptyset, F']$.
- (2) Taking $G = F'$, and $H = \mathcal{G}$ in Proposition 2.5(1), we see $E^i(j) \vee F$ is a modular flat of $[F', \mathcal{G}]$.

3. MATROIDS OF TORIC VECTOR BUNDLES

In this section we discuss matroids associated to toric vector bundles in the spirit of [DJS].

3.1. The polymatroid of a toric vector bundle. Let X_Σ be an n -dimensional toric variety. A toric vector bundle \mathcal{E} of rank r on a toric variety X_Σ is described by the following data, due to Klyachko ([Kly89]): For each ray of the fan we associate a descending filtration $E^i(j) \supseteq E^{i+1}(j)$ of a vector space $E \cong K^r$, with $E^i(j) = E$ for $j \ll 0$ and $E^i(j) = \{\mathbf{0}\}$ for $j \gg 0$. These filtrations need to be *compatible* in the sense that for every cone $\sigma \in \Sigma$, we can write $E = \bigoplus_{\mathbf{u} \in M} E_{\mathbf{u}}^\sigma$ so that $E^i(j) = \sum_{\mathbf{u} \cdot \mathbf{v}_i \geq j} E_{\mathbf{u}}^\sigma$ when $i \in \sigma$.

When X_Σ is smooth, so σ is generated by part of a basis for N , this is equivalent to requiring that there is a basis for E such that each $E^i(j)$ with $\rho_i \in \sigma$ is spanned by part of the basis (cf. [Kly89, Remark 2.2.2], [KD19, Page 76-77]).

If the filtrations do not satisfy these compatibility requirements we get a toric reflexive sheaf, instead of a toric vector bundle; see [Per03, §4.5].

The subspaces $E^i(j)$ induce a *polymatroid* as follows. Let \mathcal{H} be the set of all intersections $\cap_i E^i(j_i)$ of the $E^i(j)$ not equal to $\{\mathbf{0}\}$ or E . We define the rank function $\text{rk}: 2^{\mathcal{H}} \rightarrow \mathbb{N}$ by

$$(3.1) \quad \text{rk}(A) = \dim \left(\sum_{V \in A} V \right).$$

This satisfies the polymatroid axioms:

- (1) $\text{rk}(\emptyset) = 0$,
- (2) $\text{rk}(A) \leq \text{rk}(B)$ when $A \subseteq B$, and
- (3) $\text{rk}(A \cap B) + \text{rk}(A \cup B) \leq \text{rk}(A) + \text{rk}(B)$ for all $A, B \subseteq \mathcal{H}$.

Definition 3.1. The *polymatroid* of a toric vector bundle \mathcal{E} on a toric variety X_Σ is the polymatroid on the set \mathcal{H} of all intersections of the $E^i(j)$ not equal to $\{\mathbf{0}\}$ or E with rank function given by (3.1).

A flat of this polymatroid is a set $A \subseteq \mathcal{H}$ that is maximal among sets of its rank. We can identify each intersection of $E^i(j) \in \mathcal{H}$ with the smallest flat containing it. The poset of all flats under inclusion is a lattice, with meet given by intersection. This poset can be identified with the lattice $L(\mathcal{E})$ whose elements are all spans of intersections of subsets of the $E^i(j)$, ordered by inclusion.

Note that for each $1 \leq i \leq s$ the $E^i(j)$ for all $j \in \mathbb{N}$ form a chain in this lattice.




FIGURE 1




FIGURE 2

Example 3.2. Write $\rho_0 = \text{cone}(-1, -1)$, $\rho_1 = \text{cone}(1, 0)$, and $\rho_2 = \text{cone}(0, 1)$ for the rays of the fan of \mathbb{P}^2 as a toric variety. Let \mathcal{E} be the toric vector bundle on \mathbb{P}^2 given by the following filtrations on $E = K^3$ with basis $\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3$:

$$E^0(j) = \begin{cases} E & j \leq 0 \\ \text{span}(\mathbf{e}_1, \mathbf{e}_2) & j = 1 \\ 0 & j \geq 2 \end{cases}, \quad E^1(j) = \begin{cases} E & j \leq 0 \\ \text{span}(\mathbf{e}_1 + \mathbf{e}_3) & j = 1 \\ 0 & j \geq 2 \end{cases}, \quad E^2(j) = \begin{cases} E & j \leq 0 \\ \text{span}(\mathbf{e}_3) & j = 1 \\ 0 & j \geq 2 \end{cases}.$$

We have $\mathcal{H} = \{a_1 = \text{span}(\mathbf{e}_1, \mathbf{e}_2), a_2 = \text{span}(\mathbf{e}_1 + \mathbf{e}_3), a_3 = \text{span}(\mathbf{e}_3)\}$. The function $\text{rk}: 2^{\mathcal{H}} \rightarrow \mathbb{N}$ is:

$$\begin{aligned} \text{rk}(\{a_1, a_2, a_3\}) &= \text{rk}(\{a_1, a_2\}) = \text{rk}(\{a_1, a_3\}) = 3 \\ \text{rk}(\{a_1\}) &= \text{rk}(\{a_2, a_3\}) = 2 \\ \text{rk}(\{a_2\}) &= \text{rk}(\{a_3\}) = 1. \end{aligned}$$

The sets $\{a_1, a_2, a_3\}, \{a_2, a_3\}, \{a_1\}, \{a_2\}, \{a_3\}$ are flats. Figure 1 shows the poset of intersections of the $E^i(j)$, and the lattice of flats is shown in Figure 2. \diamond

3.2. Matroids of a toric vector bundle. In [DJS] the authors associate a representable matroid $\underline{\mathcal{M}}(\mathcal{E})$ to a toric vector bundle \mathcal{E} . The key property of this matroid is that all intersections of the subspaces $E^i(j)$ occur as flats of this matroid.

Let $L'(\mathcal{E})$ be the poset of all intersections $\cap_i E^i(j_i)$, ordered by inclusion. This is isomorphic to a subposet of the lattice of flats of the polymatroid of \mathcal{E} of Definition 3.1. It is well known that every polymatroid rank function is induced from that of a matroid. This is a special case of the fact that every lattice embeds into a geometric lattice. We now recall this, in a version that is relevant for this paper.

A function $\phi: L \rightarrow L'$ is an embedding of lattices if ϕ is an injection, $\phi(a \vee b) = \phi(a) \vee \phi(b)$, and $\phi(a \wedge b) = \phi(a) \wedge \phi(b)$. Note that this implies that $a < b$ if and only if $\phi(a) < \phi(b)$, as $a < b$ if and only if $a \vee b = b$. The original observation that lattices can be embedded into geometric lattices goes back to Dilworth [CD73, Chapter 14]. See Remark 3.4 for more references.

Recall that given two matroids $\underline{\mathcal{M}}_1, \underline{\mathcal{M}}_2$ on a ground set \mathcal{G} we say that $\underline{\mathcal{M}}_1 < \underline{\mathcal{M}}_2$ in the weak order if every basis of $\underline{\mathcal{M}}_1$ is a basis of $\underline{\mathcal{M}}_2$.

Theorem 3.3. *Let L be a lattice with $\hat{0}$ and $\hat{1}$ and a function $r: L \rightarrow \mathbb{N}$ that satisfies*

- (1) (normalization) $r(\hat{0}) = 0$;
- (2) (increasing) If $x < y$ then $r(x) < r(y)$;
- (3) (semimodularity) For any $x, y \in L$, we have $r(x \wedge y) + r(x \vee y) \leq r(x) + r(y)$.

Then there is a matroid $\underline{\mathcal{M}}$ for which L embeds into the lattice of flats $\mathcal{L}(\underline{\mathcal{M}})$ of $\underline{\mathcal{M}}$ as a lattice, with $\text{rk}(x) = r(x)$ for all $x \in L$.

There is a unique choice of such matroid $\underline{\mathcal{M}}$ for which

- (1) the ground set of $\underline{\mathcal{M}}$ has minimal size, and

(2) for any other matroid \underline{M}' whose lattice of flats contains L as a meet-semilattice with the same ground set, we have $\underline{M}' < \underline{M}$ in the weak order.

This matroid \underline{M} has no loops or parallel elements.

Proof. The main theorem of [Sim77] shows that there is a matroid \underline{M} for which L is isomorphic to the lattice of *cyclic flats* of \underline{M} . These are flats (also called “fully dependent”) that are unions of circuits. In addition, $\text{rk}_{\underline{M}}(x) = r(x)$ for all $x \in L$. This shows the existence claim of the theorem.

We write \mathcal{G} for the ground set of \underline{M} and $\phi(x)$ for the flat of \underline{M} corresponding to $x \in L$. In the proof of [Sim77, Theorem 1], Step C shows that the matroid \underline{M} constructed has rank function given by $\text{rk}(S) = \min_{x \in L} (r(x) + |S \setminus \phi(x)|)$ for all $S \subseteq \mathcal{G}$. If $B \subseteq \mathcal{G}$ satisfies

$$(3.2) \quad |B \cap \phi(x)| \leq r(x) \text{ for all } x \in L,$$

then $\text{rk}(B) = r(y) + |B \setminus \phi(y)|$ for some $y \in L$, and thus $\text{rk}(B) = r(y) + |B| - |B \cap \phi(y)| \geq r(y) + |B| - r(y) = |B|$, so all such B are independent in Sim’s matroid \underline{M} .

To show that there is a minimal choice for \underline{M} , let \underline{M} be any matroid satisfying the conditions of the theorem. We now construct a subset \mathcal{G}' of the ground set \mathcal{G} of \underline{M} for which L is still isomorphic to a sublattice of the lattice of flats of $\underline{M}|_{\mathcal{G}'}$. Let \mathcal{J} be the poset of join irreducibles (elements that are not the join of other subsets of L) in L . For $j \in \mathcal{J}$, set $s_j = \bigvee_{j' < j, j' \in \mathcal{J}} j'$. The set \mathcal{G}' we construct will have size $\sum_{j \in \mathcal{J}} r(j) - r(s_j)$. Since this will show that every choice of matroid \underline{M} satisfying the conditions of the theorem has a subset \mathcal{G}' of this size, we can conclude that this is the minimal possible size.

The construction is by induction on \mathcal{J} . For each minimal element j of \mathcal{J} , choose a maximal independent subset of $\phi(j)$, and set \mathcal{G}' initially to be the union of all such subsets. For $j \in \mathcal{J}$ for which all smaller $j' \in \mathcal{J}$ have been considered, choose a complement $W_j \in \mathcal{L}(\underline{M})$ to $\phi(j)$ with respect to $\phi(s_j)$. This has the property that $W_j \vee \phi(s_j) = \phi(j)$, while $W_j \wedge \phi(s_j) = \hat{0}$; it exists by [Whi86, Proposition 3.4.4]. Add a maximal independent subset \mathcal{G}'_j of W_j to \mathcal{G}' . We claim that when this procedure is finished, with all $j \in \mathcal{J}$ considered, L is a sublattice of the lattice of flats of $\underline{M}|_{\mathcal{G}'}$. Indeed, for $x \in L$ set $\psi(x) = \phi(x) \cap \mathcal{G}'$. Then $\psi(x)$ is a flat of $\underline{M}|_{\mathcal{G}'}$, since if there was a circuit in $\underline{M}|_{\mathcal{G}'}$ with exactly one element not in $\psi(x)$, this would be a circuit with exactly one element not in the flat $\phi(x)$ of \underline{M} , which is a contradiction. In addition, if $y < x$ then $\psi(y) \subsetneq \psi(x)$. Note also that this construction removes any loops or parallel elements.

Finally, we note that $\text{rk}(\psi(x)) = r(x)$ for all $x \in L$. The proof is by induction on L . When $x \in L$ is minimal, it is a minimal element of the poset \mathcal{J} as well, so the rank is correct by construction. Suppose now that the claim is true for all $y < x$. If $x \in \mathcal{J}$, then by induction we have $\text{rk}(\psi(s_x)) = r(s_x)$. During the construction of \mathcal{G}' we add a set \mathcal{G}'_x of size $r(x) - r(s_x)$ to \mathcal{G}' . This has the property that the closure in \underline{M} of $\psi(s_x) \cup \mathcal{G}'_x$, which is also the closure of $\phi(s_x) \cup \mathcal{G}'_x$, is $\phi(s_x) \vee W_x = \phi(x)$, which has rank $r(x)$, so the union of a maximal independent subset of $\psi(s_x)$ and \mathcal{G}'_x is independent, and thus $\text{rk}(\psi(x)) = r(x)$. If $x \notin \mathcal{J}$, write $x = j_1 \vee \cdots \vee j_\ell$ for $j_i \in \mathcal{J}$ less than x . By induction $\text{rk}(\psi(j_i)) = r(j_i)$, and thus the closure in \underline{M} of $\psi(j_i)$ is $\phi(j_i)$ for $1 \leq i \leq \ell$. Thus the closure in \underline{M} of $\bigcup_{i=1}^\ell \psi(j_i)$ equals the closure of $\bigcup_{i=1}^\ell \phi(j_i)$, which is $\phi(j_1) \vee \cdots \vee \phi(j_\ell) = \phi(j_1 \vee \cdots \vee j_\ell) = \phi(x)$. This shows that $\text{rk}(\bigcup_{i=1}^\ell \psi(j_i)) = \text{rk}(\phi(x)) = r(x)$. As $\text{rk}(\bigcup_{i=1}^\ell \psi(j_i)) \leq \text{rk}(\psi(x)) \leq \text{rk}(\phi(x))$, we conclude that $\text{rk}(\psi(x)) = r(x)$ as required.

Note that when this minimalization procedure is applied to Sim’s matroid \underline{M} , a set $B \subseteq \mathcal{G}'$ is still independent whenever the condition (3.2) holds. For any matroid satisfying the conditions of the theorem with ground set \mathcal{G} , and $x \in L$, if a set $B \subseteq \mathcal{G}$ is independent then we must have $|B \cap \phi(x)| \leq r(x)$. In particular, if $\mathcal{G} = \mathcal{G}'$, then every basis of the matroid is a basis of the minimalization of \underline{M} , and thus this minimalization is the largest element in the weak order of matroids of minimal ground set satisfying the conditions of the theorem. \square

Remark 3.4. The first proof that all lattices can be embedded into lattices of flats of matroids is attributed to earlier unpublished work of Dilworth in [CD73, Chapter 14]. Sims’ proof in [Sim77] follows the general outline given in [CD73], which is also present in the minimalization part of our proof of Theorem 3.3. A different short proof is given in [BdM08]. The fact that there is a unique choice of matroid with minimal ground set that is largest in the weak order is essentially shown by Nguyen in

[Whi86, Chapter 10], in the setting that the lattice L is atomic (pointed), and our proof mimics the one given there. The case that the lattice is the lattice of flats of a polymatroid was observed by Helgason [Hel74] (who calls polymatroids “hypermatroids”). Expositions are given in [Lov83] and [Oxl92, §12.1]. See also [CHLSW22] for a tropical connection. Note that the matroid constructed in these latter cases is rarely the minimal one of Theorem 3.3. See Remark 3.7 for the realizable case.

Theorem 3.3 allows us to associate a matroid to a toric vector bundle.

Definition 3.5. Let \mathcal{E} be a toric vector bundle on a toric variety X_Σ , with associated filtrations $E^i(j)$. Let $L'(\mathcal{E})$ be the set of all intersections $\cap_i E^i(j_i)$, ordered by inclusion, and $L(\mathcal{E})$ be the poset consisting of all spans of subsets of $L'(\mathcal{E})$. Note that E is the maximal element $\hat{1}$ of this poset, and the empty span $\{0\}$ is the minimal element $\hat{0}$. The poset $L(\mathcal{E})$ is a join-semilattice by construction (as the poset of all subspaces of E is lattice with join equal to span), and thus is a lattice by [Sta12, Proposition 3.3.1]. Define $r: L(\mathcal{E}) \rightarrow \mathbb{N}$ by $r(x) = \dim(x)$. As join in $L(\mathcal{E})$ is span (join) of the two subspaces, and the meet is contained in the intersection (which may not be in $L(\mathcal{E})$ if the subspaces are not in $L'(\mathcal{E})$) the function r is semimodular. We say that $\underline{\mathcal{M}}$ is a matroid for \mathcal{E} if $L(\mathcal{E})$ is isomorphic to a join-sublattice of the lattice of flats of \mathcal{E} . Such a matroid always exists by applying Theorem 3.3 to $L(\mathcal{E})$. We say that $\underline{\mathcal{M}}$ is a minimal matroid of \mathcal{E} if the ground set of $\underline{\mathcal{M}}$ has the minimal possible size for such a matroid.

Example 3.6. Let \mathcal{E} be the toric vector bundle on \mathbb{P}^2 of Example 3.2. The lattice $L'(\mathcal{E})$ of Definition 3.5 is shown in Figure 1. The lattice $L(\mathcal{E})$ is shown in Figure 2.

We give three examples of matroids for \mathcal{E} .

- (1) The unique matroid $\underline{\mathcal{M}}_1$ constructed by Theorem 3.3 for $L(\mathcal{E})$ with function r given by dimension has ground set $\mathcal{G} = \{e_0^1, e_0^2, e_1^1, e_2^2\}$. The bases of $\underline{\mathcal{M}}_1$ are all subsets of \mathcal{G} of size 3, so $\underline{\mathcal{M}}_1$ is the uniform matroid $U(3, 4)$. When constructing a realization of $\underline{\mathcal{M}}_1$, we choose e_1^1 to be a multiple of $\mathbf{e}_1 + \mathbf{e}_3$, and e_2^2 to be a multiple of \mathbf{e}_3 . Since $\{e_0^i, e_1^1, e_2^2\}$ is independent for $i = 1, 2$, we must then choose e_0^1, e_0^2 outside $\text{span}(e_1^1, e_2^2)$, so not a multiple of \mathbf{e}_1 . One such choice is $e_0^1 = \mathbf{e}_1 + \mathbf{e}_2$, and $e_0^2 = \mathbf{e}_2$, so

$$\mathcal{G} = \{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}.$$

- (2) Another choice is the matroid $\underline{\mathcal{M}}_2$ with ground set $\mathcal{G} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$. We have $\underline{\mathcal{M}}_2 \neq \underline{\mathcal{M}}_1$, as there is a circuit $\{\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$ of size three in $\underline{\mathcal{M}}_2$, so it is not the uniform matroid, which is the largest rank-3 matroid on 4 elements in the weak order.
- (3) Let $\underline{\mathcal{M}}_3$ be the matroid with ground set $\mathcal{G} = \{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3\}$. Then $L(\mathcal{E})$ is identified with the flats $\{\text{span}(\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2), \text{span}(\mathbf{e}_1 + \mathbf{e}_3), \text{span}(\mathbf{e}_3), \{0\}, E\}$.

◇

Remark 3.7. When the lattice L is a join-sublattice of the lattice of all subspaces of E , meaning that the join in L is the join in the subspace lattice, but the meet in the subspace lattice might not lie in L , then the matroid of Theorem 3.3 can be chosen to be realizable. This is shown in [Zie92, Theorem 4.9], and also by [DJS, Algorithm 3.2].

Note that for each ray ρ_i of Σ , each filtered piece $E^i(j)$ is an element of the lattice of flats $\mathcal{L}(\underline{\mathcal{M}})$ of any matroid for \mathcal{E} , and the set of all $E^i(j)$ as j decreases is a chain in $\mathcal{L}(\underline{\mathcal{M}})$.

Remark 3.8. The matroid associated to a toric vector bundle was first introduced in [DJS]. Theorem 3.3 differs from that presentation as follows. Firstly, [DJS] only requires an embedding of the meet-semilattice $L'(\mathcal{E})$; however their embedding actually embeds $L(\mathcal{E})$, and respects joins. Adding extra elements to the ground set of the matroid in our sense does not change the join, but might change the meet. Secondly, note that for the toric vector bundle of Example 3.6, the minimal possible size of the ground set is 4 for an embedding of either $L(\mathcal{E})$ as a join-semilattice, or $L'(\mathcal{E})$ as a meet-semilattice. However both $\underline{\mathcal{M}}_1$ and $\underline{\mathcal{M}}_2$ have one circuit: $\{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$ for $\underline{\mathcal{M}}_1$, and $\{\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$ for $\underline{\mathcal{M}}_2$, so the claim in [DJS, Proposition 3.1] that there is a unique matroid with the number of elements in the ground set being minimal, and the number of circuits minimal does not hold.

The issue here is [DJS, Algorithm 3.2] does not construct a unique matroid. Given $L(\mathcal{E})$, we initialize the G from that algorithm to contain multiples of $\mathbf{e}_1 + \mathbf{e}_3$ and \mathbf{e}_3 . The algorithm then considers

$V = \text{span}(\mathbf{e}_1, \mathbf{e}_2)$. The set G' of elements of G lying in V is empty, so $\text{span}(G') = \{0\}$. The algorithm then tells us to append to G a basis for a complementary subspace to $\{0\} = \text{span}(G')$ in V , which is a basis for V . However the choice of basis at this stage affects the matroid. The matroids $\underline{\mathcal{M}}_1$ and $\underline{\mathcal{M}}_2$ are the results of two different choices for a basis for V .

Other than the fact that references to “the matroid” should be replaced by “a matroid”, we are not aware that this issue substantively affects any results in [DJS]. In particular, as we note in Remark 6.16, the results about parliaments of polytopes (which are no longer unique) encoding global sections still hold.

As in Theorem 3.3 we can reimpose uniqueness on the matroid by asking that it be maximal in the weak order on matroids. However this may not always be geometrically desirable. For example, the toric vector bundle of Example 3.6 is a direct sum of a rank-2 bundle and a line bundle. This can be seen by noting that every subspace in the filtration is a direct sum of a subspace in $\text{span}(\mathbf{e}_1, \mathbf{e}_3)$ and one in $\text{span}(\mathbf{e}_2)$. However the matroid $\underline{\mathcal{M}}_1$, which is the largest in the weak order, is not a direct sum of matroids of lower ranks, while the matroid $\underline{\mathcal{M}}_2$, which is smaller in the weak order, is a direct sum. See Section 6.1 for more on direct sums in the context of this paper. A theme of the approach taken here is that the correct choice of matroid depends on the property desired. See Remark 7.15 for more on this in the context of stability of tropical toric vector bundles.

4. TORIC VECTOR BUNDLES AS MODULES OVER THE COX RING

In this section we give a description of a toric vector bundle as a module over the Cox ring of the base toric variety. Let \mathcal{E} be a rank- r toric vector bundle on toric variety X_Σ , and let $\underline{\mathcal{M}}(\mathcal{E})$ be a matroid for \mathcal{E} in the sense of Definition 3.5, which has a realization $\mathcal{G} \subset K^r$. For each $\mathbf{w} \in \mathcal{G}$, set $\mathbf{d}_\mathbf{w} \in \mathbb{Z}^{|\Sigma(1)|}$ to be the vector with

$$(4.1) \quad (\mathbf{d}_\mathbf{w})_i = \max(j : \mathbf{w} \in E^i(j)).$$

Note that the vectors $\mathbf{d}_\mathbf{w}$, together with the realization \mathcal{G} , actually determine the filtrations $E^i(j)$.

Lemma 4.1. *Let \mathcal{E} be a toric vector bundle on X_Σ given by filtrations $(E, \{E^i(j)\})$. Then*

$$E^i(j) = \text{span}(\mathbf{w} \in \mathcal{G} : (\mathbf{d}_\mathbf{w})_i \geq j).$$

Proof. By definition, if $(\mathbf{d}_\mathbf{w})_i \geq j$, we have $\mathbf{w} \in E^i(j)$, so the right-hand side is contained in $E^i(j)$. For the converse, suppose $\mathbf{w} \in \mathcal{G}$ lies in $E^i(j)$. This implies that $(\mathbf{d}_\mathbf{w})_i \geq j$, so \mathbf{w} is in the right-hand side. As $E^i(j)$ is a flat in the atomic lattice of flats of the matroid $\underline{\mathcal{M}}(\mathcal{E})$, it is the span of those $\mathbf{w} \in \mathcal{G}$ it contains, so it follows that $E^i(j)$ is contained in the right-hand side. \square

Remark 4.2. In [KM22], Kaveh and Manon give an equivalent description of toric vector bundles in terms of piecewise linear functions, and introduce the *diagram* of a toric vector bundle. This is the $|\Sigma(1)| \times |\mathcal{G}|$ matrix with (i, \mathbf{w}) th entry $(\mathbf{d}_\mathbf{w})_i$.

Write G for the $r \times m$ matrix with columns the vectors in \mathcal{G} , and \mathbf{w}_i for the i th column of G . For a collection of vectors $A = \{\mathbf{a}_1, \dots, \mathbf{a}_l\} \subset \mathbb{Z}^{|\Sigma(1)|}$, we write $\min A$ for the vector with $(\min A)_k = \min_j ((\mathbf{a}_j)_k)$. We write V for the $|\Sigma(1)| \times n$ matrix with rows the first lattice points on the rays of Σ .

We assume for now that X_Σ is smooth, so all Weil divisors are Cartier. Recall that Cox [Cox95] showed that in this setting there is a correspondence between coherent sheaves on X_Σ and class-group-graded modules over a polynomial ring, now called the Cox ring, with one variable for each ray of the fan Σ . This generalizes the analogous correspondence for projective space. When the sheaf is torus equivariant, the module has a finer $\mathbb{Z}^{|\Sigma(1)|}$ -grading coming from the surjection $\mathbb{Z}^{|\Sigma(1)|} \rightarrow A^1(\Sigma)$ of (2.1) (see [Per03, Section 4.8]).

We write x_1, \dots, x_s for the variables of the Cox ring of X_Σ , and t_1, \dots, t_n for the coordinates on the torus of X_Σ .

Proposition 4.3. *Let $S = K[x_1, \dots, x_s]$ be the Cox ring of a smooth toric variety X_Σ , let \mathcal{E} be a toric vector bundle on X_Σ , and let $\mathcal{G} \subseteq K^r$ be the ground set of a matroid for \mathcal{E} in the sense of Definition 3.5.*

For $\mathbf{c} \in \ker(G)$, and $1 \leq i \leq m$, set $\mathbf{u}_i = \mathbf{d}_{\mathbf{w}_i} - \min(\mathbf{d}_{\mathbf{w}_i} : c_l \neq 0)$. Then \mathcal{E} is the sheafification \tilde{P} of the $\mathbb{Z}^{|\Sigma(1)|}$ -graded module P given by

$$P = \bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}})/R,$$

where R is the submodule of $\bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}})$ generated by

$$\left\{ \sum_{i=1}^m c_i \mathbf{x}^{\mathbf{u}_i} \mathbf{e}_i : (c_1, \dots, c_m) \in \ker(G) \right\}.$$

Proof. Let $D_{\mathbf{w}}$ be the torus-invariant divisor $\sum (\mathbf{d}_{\mathbf{w}})_i D_i$ on X_{Σ} .

By [DJS, Remark 3.6] there is a surjection

$$\eta: \mathcal{F} := \bigoplus_{\mathbf{w} \in \mathcal{G}} \mathcal{O}(D_{\mathbf{w}}) \rightarrow \mathcal{E}.$$

This follows, via Klyachko's equivalence of categories, from the surjection $\eta_e: K^{|\mathcal{G}|} \rightarrow E$ given by $\mathbf{e}_{\mathbf{w}} \mapsto \mathbf{w}$.

Let $Q = \bigoplus_{\alpha \in A^1(X_{\Sigma})} H^0(X_{\Sigma}, \mathcal{E} \otimes_{\mathcal{O}_{X_{\Sigma}}} \mathcal{O}_{X_{\Sigma}}(\alpha))$. The proof of [Cox95, Theorem 3.2] shows that the sheafification \tilde{Q} of Q equals \mathcal{E} . The map η induces a linear transformation

$$\bigoplus_{\mathbf{w} \in \mathcal{G}} H^0(X_{\Sigma}, \mathcal{O}(D_{\mathbf{w}})) \rightarrow H^0(X_{\Sigma}, \mathcal{E}),$$

and thus, after taking the sum over all twists by $\mathcal{O}(\alpha)$, we get a homomorphism of S -modules

$$\eta': P' := \bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}}) \rightarrow Q.$$

We begin by showing that for fixed $\sigma \in \Sigma$ we have $(R_{\mathbf{x}^{\sigma}})_{\mathbf{0}} \subseteq (P'_{\mathbf{x}^{\sigma}})_{\mathbf{0}} \cong \mathcal{F}(U_{\sigma})$ equal to $\ker(\eta(U_{\sigma}))$.

Recall that we have $(S_{\mathbf{x}^{\sigma}})_{\mathbf{0}} \cong K[U_{\sigma}]$, via the map that takes $\mathbf{x}^{\mathbf{v}}$ to $\mathbf{t}^{\mathbf{u}} \in K[U_{\sigma}]$, where $\mathbf{u} \in M$ is the unique element with $V\mathbf{u} = \mathbf{v}$. For each $\mathbf{w} \in \mathcal{G}$, choose $\mathbf{x}^{\mathbf{v}_{\mathbf{w}}} \in S_{\mathbf{x}^{\sigma}}$ with $\deg(\mathbf{x}^{\mathbf{v}_{\mathbf{w}}}) = [D_{\mathbf{w}}] \in A^1(X_{\Sigma})$ and $(\mathbf{v}_{\mathbf{w}})_i = 0$ whenever $\mathbf{v}_i \in \sigma$. This is possible since X_{Σ} is smooth. Then $(S(\mathbf{d}_{\mathbf{w}})_{\mathbf{x}^{\sigma}})_{\mathbf{0}} \cong \mathbf{x}^{\mathbf{v}_{\mathbf{w}}}(S_{\mathbf{x}^{\sigma}})_{\mathbf{0}}$. Since $\deg(\mathbf{x}^{\mathbf{v}_{\mathbf{w}}} - \mathbf{d}_{\mathbf{w}}) = \mathbf{0}$, there is $\mathbf{u}_{\mathbf{w}} \in M$ with $V\mathbf{u}_{\mathbf{w}} = \mathbf{v}_{\mathbf{w}} - \mathbf{d}_{\mathbf{w}}$. Thus as an M -graded $K[U_{\sigma}]$ module, $(S(\mathbf{d}_{\mathbf{w}})_{\mathbf{x}^{\sigma}})_{\mathbf{0}}$ is isomorphic to $\mathbf{t}^{\mathbf{u}_{\mathbf{w}}} K[U_{\sigma}]$ [Ful93, p62].

Thus

$$(4.2) \quad \mathcal{F}(U_{\sigma}) = (P'_{\mathbf{x}^{\sigma}})_{\mathbf{0}} \cong \bigoplus_{\mathbf{w} \in \mathcal{G}} \mathbf{t}^{\mathbf{u}_{\mathbf{w}}} K[U_{\sigma}].$$

We also have that \mathcal{E} is a direct sum of line bundles on U_{σ} , so $\mathcal{E}(U_{\sigma}) \cong \bigoplus_i \mathbf{t}^{\mathbf{u}'_i} K[U_{\sigma}]$ as an M -graded module. We write $\sum_{\mathbf{w} \in \mathcal{G}} \mathbf{t}^{\mathbf{u}_{\mathbf{w}}} g_{\mathbf{w}} \mathbf{f}_{\mathbf{w}}$, with $g_{\mathbf{w}} \in K[U_{\sigma}]$, for an element of $\mathcal{F}(U_{\sigma})$, and $\sum_i \mathbf{t}^{\mathbf{u}'_i} g_i \mathbf{e}'_i$ for an element of $\mathcal{E}(U_{\sigma})$, where the $\mathbf{f}_{\mathbf{w}}$ and \mathbf{e}'_i are basis vectors. The morphism $\eta(U_{\sigma})$ is given by an $r \times |\mathcal{G}|$ matrix A with entries in $K[U_{\sigma}]$, with $\eta(U_{\sigma})(\mathbf{t}^{\mathbf{u}_{\mathbf{w}}} \mathbf{f}_{\mathbf{w}}) = \sum_i A_{i\mathbf{w}} \mathbf{t}^{\mathbf{u}'_i} \mathbf{e}'_i$. Since η is a morphism of toric vector bundles, $\eta(U_{\sigma})$ is equivariant, so each entry of A has the form $a\mathbf{t}^{\mathbf{u}}$ for some $a \in K$, $\mathbf{u} \in \sigma^{\vee} \cap M$. The morphism η_e is $\eta \otimes (K[T]/\langle t_i - 1 : 1 \leq i \leq n \rangle)_{\langle t_i - 1 : 1 \leq i \leq n \rangle}$, so $A_{i\mathbf{w}} = w_i \mathbf{t}^{\mathbf{u}}$ for some \mathbf{u} . Since $\eta(U_{\sigma})$ is a degree-zero homomorphism, we have $\mathbf{u} = \mathbf{u}_{\mathbf{w}} - \mathbf{u}'_i$. The submodule $\ker(\eta(U_{\sigma}))$ of $\mathcal{F}(U_{\sigma})$ is thus generated by

$$\left\{ \sum_{\mathbf{w} \in \mathcal{G}} a_{\mathbf{w}} \mathbf{t}^{\mathbf{u}_{\mathbf{w}}} \mathbf{f}_{\mathbf{w}} : (a_{\mathbf{w}}) \in \ker(G) \right\}.$$

The compatibility condition for \mathcal{E} implies that there is a basis B_{σ} for $\underline{\mathcal{M}}(\mathcal{E})$ with each $E^i(j)$ with $\rho_i \in \sigma$ spanned by a collection of elements of B_{σ} . Fix $k \notin B_{\sigma}$. Let $\mathbf{c}^k \in \ker(G)$ be the vector whose tropicalization is the fundamental circuit of k over B_{σ} , so $c_{\mathbf{w}_k}^k = 1$, $c_{\mathbf{w}}^k = 0$ for $\mathbf{w} \notin B_{\sigma} \cup \{k\}$, and

$$\mathbf{w}_k = - \sum_{\mathbf{w} \in B_{\sigma}} c_{\mathbf{w}}^k \mathbf{w}.$$

The elements

$$(4.3) \quad \left\{ \sum_{\mathbf{w}} c_{\mathbf{w}}^k \mathbf{t}^{\mathbf{u}_{\mathbf{w}}} \mathbf{f}_{\mathbf{w}} : \mathbf{c}^k \text{ is the fundamental circuit of } k \text{ over } B_{\sigma} \text{ for all } k \notin B_{\sigma} \right\}$$

generate $\ker(\eta(U_{\sigma}))$.

We now check that these elements also generate $(R_{\mathbf{x}^{\hat{\sigma}}})_{\mathbf{0}}$. For fixed $i \in \sigma$, $j \in \mathbb{Z}$, and $k \notin B_\sigma$, if $\mathbf{w} \notin E^i(j)$ for some $\mathbf{w} \in \mathcal{B}_\sigma$ with $c_{\mathbf{w}}^k \neq 0$, then since $\{\mathbf{w} \in B_\sigma\} \subseteq K^r$ is linearly independent and contains a basis for $E^i(j)$, we have $\mathbf{w}_k \notin E^i(j)$. This implies that $(\mathbf{d}_{\mathbf{w}_k})_i \leq (\mathbf{d}_{\mathbf{w}})_i$. However if all $\mathbf{w} \in \mathcal{B}_\sigma$ with $c_{\mathbf{w}}^k \neq 0$ lie in $E^i(j)$, then we also have $\mathbf{w}_k \in E^i(j)$. Thus $(\mathbf{d}_{\mathbf{w}_k})_i = \min((\mathbf{d}_{\mathbf{w}})_i : \mathbf{w} \in B_\sigma, c_{\mathbf{w}}^k \neq 0)$, so $(\mathbf{u}_k)_i = 0$.

This means that $(\mathbf{u}_k)_i = 0$ if $\rho_i \in \sigma$. Thus $\mathbf{x}^{-\mathbf{u}_k} \in S_{\mathbf{x}^{\hat{\sigma}}}$, and so $\mathbf{e}_k + \sum_{j \in B_\sigma} c_j \mathbf{x}^{\mathbf{u}_j - \mathbf{u}_k} \mathbf{e}_j \in P'_{\mathbf{x}^{\hat{\sigma}}}$. We claim that these elements, as k varies over the complement of B_σ , generate $\tilde{R}_{\mathbf{x}^{\hat{\sigma}}}$. If not, after subtracting multiples of these elements from a generator, we would have an element $\sum_{j \in B_\sigma} c'_j \mathbf{x}^{\mathbf{v}_j} \mathbf{e}_j \in R_{\mathbf{x}^{\hat{\sigma}}}$. Since the coefficients of elements of R lie in $\ker(G)$, this would contradict B_σ being a basis.

Recall that $\mathbf{v}_{\mathbf{w}_k}$ has $\deg(\mathbf{x}^{\mathbf{v}_{\mathbf{w}_k}}) = [D_{\mathbf{w}_k}]$ and $(\mathbf{v}_{\mathbf{w}_k})_i = 0$ when $\rho_i \in \sigma$, so $\mathbf{x}^{\mathbf{v}_{\mathbf{w}_k}}$ is a unit in $S_{\mathbf{x}^{\hat{\sigma}}}$. Thus the elements $\mathbf{x}^{\mathbf{v}_{\mathbf{w}_k}}(\mathbf{e}_{\mathbf{w}_k} + \sum_{\mathbf{w} \in B_\sigma} c_{\mathbf{w}_k} \mathbf{x}^{\mathbf{u}_{\mathbf{w}} - \mathbf{u}_k} \mathbf{e}_{\mathbf{w}})$ for $k \in B_\sigma$ have degree $\mathbf{0} \in A^1(X_\Sigma)$, and also generate $R_{\mathbf{x}^{\hat{\sigma}}}$, so generate $(R_{\mathbf{x}^{\hat{\sigma}}})_{\mathbf{0}}$. Under the isomorphism (4.2), these are taken to the elements (4.3). This shows that $(R_{\mathbf{x}^{\hat{\sigma}}})_{\mathbf{0}} \subseteq (P'_{\mathbf{x}^{\hat{\sigma}}})_{\mathbf{0}} \cong \mathcal{F}(U_\sigma)$ is equal to $\ker(\eta(U_\sigma))$ as claimed.

We thus have $\tilde{R} = \ker(\eta)$, so

$$0 \rightarrow \tilde{R} \rightarrow \tilde{P}' \cong \mathcal{F} \xrightarrow{\eta} \mathcal{E} \rightarrow 0$$

is exact. Since the sheafification map is exact [Cox95, Proposition 3.1], it follows that

$$\tilde{P} = \widetilde{P'/R} \cong \mathcal{E}$$

as required. \square

Example 4.4. Consider the tangent bundle \mathcal{T} on \mathbb{P}^2 . We use the notation of [DJS, Example 3.8]. The matrix G equals

$$G = \begin{pmatrix} 1 & 0 & -1 \\ 0 & 1 & -1 \end{pmatrix}.$$

The degrees $\mathbf{d}_{\mathbf{e}_i}$ are $(1, 0, 0)$, $(0, 1, 0)$, and $(0, 0, 1)$. The Cox ring of \mathbb{P}^2 is $S = K[x, y, z]$. We have \mathcal{T} equal to \tilde{P} , where P is the S -module

$$P = (S(1, 0, 0) \oplus S(0, 1, 0) \oplus S(0, 0, 1)) / \langle (x\mathbf{e}_1 + y\mathbf{e}_2 + z\mathbf{e}_3) \rangle.$$

Note that on the chart $x \neq 0$ this becomes

$$(4.4) \quad (P_x)_{\mathbf{0}} = xK[y/x, z/x] \oplus yK[y/x, z/x] \oplus zK[y/x, z/x] / \langle (\mathbf{e}_1 + y/x\mathbf{e}_2 + z/x\mathbf{e}_3) \rangle \\ = xK[y/x, z/x] \oplus yK[y/x, z/x].$$

This is free of rank 2 as expected. \diamond

Remark 4.5. A similar construction occurs in [GM23a, GM23b, GM23c], where George and Manon construct the Cox rings of some families of projectivized toric vector bundles. For those families, the Cox module described in Proposition 4.3 is a graded submodule of the associated Cox ring.

5. TROPICAL TORIC VECTOR BUNDLES

In this section we make the main definition of the paper: in Definition 5.4 we define a *tropical toric vector bundle* on a tropical toric variety.

5.1. Tropical toric reflexive sheaves. We first define a tropical toric reflexive sheaf on $\text{trop}(X_\Sigma)$.

Definition 5.1. A tropical toric reflexive sheaf \mathcal{E} on $\text{trop}(X_\Sigma)$ is given by the data of a simple (no loops or parallel elements) valuated matroid \mathcal{M} of rank r on a ground set \mathcal{G} , together with for each ray ρ_i of Σ a collection $\{E^i(j) : j \in \mathbb{Z}\}$ of flats of the underlying matroid $\underline{\mathcal{M}}$ with the property that $E^i(j) \leq E^i(j')$ if $j > j'$, $E^i(j) = \emptyset$ for $j \gg 0$, and $E^i(j) = \mathcal{G}$ for $j \ll 0$.

Moreover, we call a tropical toric reflexive sheaf *tropically minimal* if the lattice of flats of $\underline{\mathcal{M}}$ does not contain a proper subposet containing the $E^i(j)$ that is the lattice of flats of a matroid.

We denote the tropical toric reflexive sheaf \mathcal{E} by $(\mathcal{M}, \mathcal{G}, \{E^i(j)\})$.

For $\mathbf{w} \in \mathcal{G}$, set $\mathbf{d}_{\mathbf{w}} \in \mathbb{Z}^{|\Sigma(1)|}$ to be the vector with $(\mathbf{d}_{\mathbf{w}})_i = \max(j : \mathbf{w} \in E^i(j))$. This is directly analogous to (4.1), however here each $E^i(j)$ is a subset of the set \mathcal{G} , and $\mathbf{w} \in \mathcal{G}$.

Let $S = \text{Cox}(\text{trop}(X_{\Sigma})) = \overline{\mathbb{R}}[x_1, \dots, x_s]$. Fix $m = |\mathcal{G}|$, and let R be the S -subsemimodule of $\bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}})$ generated by

$$\left\{ \sum_{k=1}^m c_i \mathbf{x}^{\mathbf{u}_k} \mathbf{e}_k : (c_1, \dots, c_m) \text{ is a vector of the valuated matroid } \mathcal{M} \text{ and } \mathbf{u}_k = \mathbf{d}_{\mathbf{w}_k} - \min(\mathbf{d}_{\mathbf{w}_l} : c_l \neq \infty) \right\}.$$

The tropical toric reflexive sheaf on $\text{trop}(X_{\Sigma})$ given by this data is described by the $\mathbb{Z}^{|\Sigma(1)|}$ -graded S -semimodule

$$(5.1) \quad P = \bigoplus_{\mathbf{w} \in \mathcal{G}} S(\mathbf{d}_{\mathbf{w}}) / \mathcal{B}(R),$$

where $\mathcal{B}()$ is the Giansiracusa bend congruence; see Section 2.3. The semimodule P is the tropicalization of the module P of Proposition 4.3.

Remark 5.2. Note that we can recover the filtrations $E^i(j)$ from $\{\mathbf{d}_{\mathbf{w}} : \mathbf{w} \in \mathcal{G}\}$ as in Lemma 4.1. Indeed, for $\rho_i \in \Sigma(1)$ and $j \in \mathbb{Z}$ set

$$E_j^i = \bigvee_{(\mathbf{d}_{\mathbf{w}})_i \geq j} \mathbf{w},$$

where \vee is join in the lattice $\mathcal{L}(\mathcal{M})$ of flats of \mathcal{M} . We will show that $E_j^i = E^i(j)$. If $(\mathbf{d}_{\mathbf{w}})_i \geq j$ for some $\mathbf{w} \in \mathcal{G}$, then $\mathbf{w} \in E^i(j)$ by the definition of $\mathbf{d}_{\mathbf{w}}$, so $E_j^i \leq E^i(j)$. Conversely, if $\mathbf{w} \in \mathcal{G}$ is contained in $E^i(j)$ for some i, j , then $(\mathbf{d}_{\mathbf{w}})_i \geq j$ by the definition of $\mathbf{d}_{\mathbf{w}}$, so $\mathbf{w} \in E_j^i$. This shows that $E^i(j) \leq E_j^i$, and thus $E^i(j) = E_j^i$.

Example 5.3. (1) A rank-1 tropical reflexive sheaf is given by the data $(\mathcal{M}, \mathcal{G}, \{E^i(j)\})$, where $\text{rk}(\mathcal{M}) = 1$. The only information in the filtrations $\{E^i(j)\}$ is the smallest integer a_i for which $E^i(a_i) \neq \emptyset$, so we may identify such a sheaf with the vector $\mathbf{a} = (a_1, \dots, a_s) \in \mathbb{Z}^s$. This is the tropical version of $\mathcal{O}(\sum_{i=1}^s a_i D_i)$. In view of [CLS11, Theorem 8.2.3], we can define tropical canonical sheaf $\text{trop}(\omega_{X_{\Sigma}})$ as the rank-1 sheaf where all a_i equal -1 .

(2) (Tropical tangent sheaf). Let X_{Σ} be a complete toric variety. We define the tropical tangent reflexive sheaf $\text{trop}(\mathcal{T})$ on $\text{trop}(X_{\Sigma})$ as follows. Let $\mathcal{G} = \{\mathbf{v}_1, \dots, \mathbf{v}_s\} \subseteq N \otimes_{\mathbb{Z}} K$. Consider the simple representable matroid on the ground set \mathcal{G} (i.e., we identify \mathbf{v}_i with \mathbf{v}_j if they are scalar multiples of each other). Let \mathcal{M} be the trivial valuated matroid structure on this matroid. Consider the following filtration of flats

$$\mathcal{T}^i(j) = \begin{cases} \mathcal{G} & i \leq 0 \\ \{\mathbf{v}_i\} & j = 1 \\ \emptyset & j > 1 \end{cases}.$$

Then $\text{trop}(\mathcal{T}) = (\mathcal{M}, \mathcal{G}, \{\mathcal{T}^i(j)\})$ is a tropical reflexive sheaf on $\text{trop}(X_{\Sigma})$. See Example 5.9 (1) for an interpretation of this as a tropicalization of the tangent sheaf.

(3) (Uniform matroid as tangent bundle on \mathbb{P}^n). Consider the uniform matroid $U_{n, n+1}$ on the ground set $\mathcal{G} = \{\mathbf{w}_0, \dots, \mathbf{w}_n\}$, where any proper subset of \mathcal{G} is independent, and let \mathcal{M} be the trivial valuated matroid structure on this uniform matroid. Let $X_{\Sigma} = \mathbb{P}^n$ be projective space. Let $\mathbf{e}_1, \dots, \mathbf{e}_n$ denote the standard basis of \mathbb{Z}^n . The fan Σ consists of rays $\mathbf{v}_1 = \mathbf{e}_1, \dots, \mathbf{v}_n = \mathbf{e}_n$ and $\mathbf{v}_0 = -\mathbf{e}_1 - \dots - \mathbf{e}_n$ and the maximal cones generated by proper subsets of $\{\mathbf{v}_0, \dots, \mathbf{v}_n\}$. Consider the family of filtrations

$$E^{\mathbf{v}_i}(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{w}_i\} & j = 1 \text{ for } i = 0, \dots, n \\ \emptyset & j > 1, \end{cases}$$

This defines a tropical reflexive sheaf on $\text{trop}(\mathbb{P}^n)$, which is the tangent sheaf by Example 5.3 (2).




FIGURE 3




FIGURE 4

- (4) Consider the singular toric surface X_Σ , where the rays of the fan are given by $\mathbf{v}_1 = (1, 0)$, $\mathbf{v}_2 = (1, 2)$ and $\mathbf{v}_3 = (-1, 1)$. The maximal cones of Σ are given by

$$\sigma_1 = \text{cone}(\mathbf{v}_1, \mathbf{v}_2), \sigma_2 = \text{cone}(\mathbf{v}_2, \mathbf{v}_3) \text{ and } \sigma_3 = \text{cone}(\mathbf{v}_1, \mathbf{v}_3).$$

Let $\mathbf{e}_1, \mathbf{e}_2$ denote the standard basis of $E = K^2$. Consider the representable matroid $\underline{\mathcal{M}}$ of rank 2 on the ground set

$$\mathcal{G} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_2\}.$$

Consider the following family of filtrations

$$E^1(j) = \begin{cases} \mathcal{G} & j \leq 1 \\ \{\mathbf{e}_1\} & j = 2 \\ \emptyset & j > 2 \end{cases}, \quad E^2(j) = \begin{cases} \mathcal{G} & j \leq 1 \\ \{\mathbf{e}_2\} & j = 2 \\ \emptyset & j > 2 \end{cases}, \quad E^3(j) = \begin{cases} \mathcal{G} & j \leq 1 \\ \{\mathbf{e}_1 + \mathbf{e}_2\} & j = 2 \\ \emptyset & j > 2 \end{cases}.$$

- (5) Let $X_\Sigma = \mathbb{P}^1$. Let \mathcal{M} be the valuated matroid (with the trivial valuation) corresponding to the five vectors in K^3 : $\mathcal{G} = \{\mathbf{w}_1 = (1, 0, 0), \mathbf{w}_2 = (1, 1, 0), \mathbf{w}_3 = (0, 1, 0), \mathbf{w}_4 = (0, 0, 1), \mathbf{w}_5 = (0, 1, 1)\}$. This has the lattice of flats shown in Figure 3.

The fan of \mathbb{P}^1 has two rays: ρ_0 and ρ_1 . Consider the filtrations

$$E^0(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3\} & j = 1 \\ \{\mathbf{w}_1\} & j = 2 \\ \emptyset & j > 2 \end{cases}, \quad E^1(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{w}_3, \mathbf{w}_4, \mathbf{w}_5\} & j = 1 \\ \{\mathbf{w}_3\} & j = 2 \\ \emptyset & j > 2 \end{cases}.$$

Note that \mathcal{E} is not tropically minimal, as the subposet of Figure 4 is the lattice of flats of the uniform matroid $U(3, 3)$. Replacing \mathcal{M} by the (trivially valuated) valuated matroid corresponding to the vectors $\{\mathbf{w}_1 = (1, 0, 0), \mathbf{w}_3 = (0, 0, 1), \mathbf{w}_4 = (0, 0, 1)\}$ also gives a tropical reflexive sheaf.

◇

5.2. Tropical toric vector bundles. We now generalize Definition 5.1 to define tropical toric vector bundles.

Definition 5.4. A rank- r tropical toric vector bundle on $\text{trop}(X_\Sigma)$ is a rank- r tropical toric reflexive sheaf $(\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ that satisfies the following compatibility condition:

For any $\sigma \in \Sigma$ there is a multiset $\mathbf{u}(\sigma) \subseteq M$, and a basis $B_\sigma = \{\mathbf{w}_\mathbf{u} : \mathbf{u} \in \mathbf{u}(\sigma)\}$ of $\underline{\mathcal{M}}$ such that for every ray $\rho_i \in \sigma$ and $j \in \mathbb{Z}$ we have

$$(5.2) \quad E^i(j) = \bigvee_{\mathbf{u} \in B_\sigma, \mathbf{u} \cdot \mathbf{v}_i \geq j} \mathbf{w}_\mathbf{u}.$$

When the toric variety X_Σ is smooth, we can simplify the compatibility condition (5.2), as in the case of usual toric vector bundles. This should be compared with [Kly89, Remark 2.2.2].

Lemma 5.5. *If X_Σ is a smooth toric variety then condition (5.2) is equivalent to the condition that*

$$(5.3) \quad \text{for every } \sigma \in \Sigma \text{ there is a basis } B_\sigma \text{ of } \underline{\mathcal{M}} \text{ such that each } E^i(j) \text{ for } i \in \sigma \text{ is the join of elements of } B_\sigma.$$

Proof. Suppose that X_Σ is smooth, and the data $\mathcal{E} = \{\mathcal{M}, \mathcal{G}, E^i(j)\}$ defines a tropical toric reflexive sheaf. We first suppose that the tropical reflexive sheaf satisfies (5.2). Condition 5.3 follows immediately, as it is an a priori weaker condition than (5.2), and the same basis B_σ works.

Now suppose that the tropical reflexive sheaf \mathcal{E} satisfies Condition 5.3. Fix $\sigma \in \Sigma$, and a basis B_σ of $\underline{\mathcal{M}}$ such that Condition 5.3 holds. For $\mathbf{w} \in B_\sigma$, consider $(\mathbf{d}_\mathbf{w})_i = \max(j : \mathbf{w} \in E^i(j))$. We then have $\mathbf{w} \in B_\sigma \cap E^i(j)$ if and only if $j \leq (\mathbf{d}_\mathbf{w})_i$. Since σ is a unimodular cone, for each $\mathbf{w} \in B_\sigma$ there is $\mathbf{u}_\mathbf{w} \in M$ with $\mathbf{u}_\mathbf{w} \cdot \mathbf{v}_i = (\mathbf{d}_\mathbf{w})_i$ for all i with $\rho_i \in \sigma$. By construction $\mathbf{w} \in B_\sigma \cap E^i(j)$ if and only if $\mathbf{u}_\mathbf{w} \cdot \mathbf{v}_i \geq j$. Since $E^i(j)$ is the join of elements of B_σ , and B_σ is a basis and thus independent, we have that $E^i(j)$ is the join of all elements of B_σ that it contains. Thus the multiset $\mathbf{u}(\sigma) := \{\mathbf{u}_\mathbf{w} : \mathbf{w} \in B_\sigma\} \subseteq M$, and the basis B_σ satisfies (5.2). \square

As an application of the above lemma we determine which of the tropical reflexive sheaves in Example 5.3 are in fact vector bundles.

Example 5.6. (1) When X_Σ is smooth, then using Lemma 5.5, we see that the tropical reflexive sheaves in Example 5.3 (2) and (3) are examples of tropical vector bundles. In particular, the tropical tangent sheaf on $\text{trop}(\mathbb{P}^n)$ in Example 5.3 (3) is a tropical toric vector bundle. More generally, observe that any choice of filtrations of flats the uniform matroid $U_{n,n+1}$ will give rise to a tropical vector bundle on $\text{trop}(\mathbb{P}^n)$ as the compatibility condition will always hold using Lemma 5.5.

- (2) The filtrations in Example 5.3 (4) do not satisfy the compatibility condition for the cone σ_1 . If they did, there would be weights $\mathbf{u}_{\mathbf{w}_1} = (a_1, b_1)$ and $\mathbf{u}_{\mathbf{w}_2} = (a_2, b_2)$ in \mathbb{Z}^2 . From the filtration $E^{\mathbf{v}_1}(j)$, we have

$$a_1 = 2 \text{ and } a_2 = 1.$$

Similarly, from the filtration $E^{\mathbf{v}_2}(j)$, we have

$$a_1 + 2b_1 = 1 \text{ and } a_2 + 2b_2 = 2.$$

Note that these equations do not have an integer solution. Hence, this is an example of tropical reflexive sheaf that is not a vector bundle.

- (3) Consider the representable matroid $\underline{\mathcal{M}}$ of rank-2 on the ground set

$$\mathcal{G} = \{\mathbf{w}_1 = \mathbf{e}_1, \mathbf{w}_2 = \mathbf{e}_2, \mathbf{w}_3 = \mathbf{e}_1 + \mathbf{e}_2\}.$$

We define a tropical toric reflexive sheaf on the tropical toric variety $\text{trop}(\mathbb{A}^3) = \overline{\mathbb{R}}^3$, whose fan has three rays that span a single cone. Consider the filtrations on these rays:

$$E^1(j) = \begin{cases} \mathcal{G} & j \leq 1 \\ \{\mathbf{w}_1\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^2(j) = \begin{cases} \mathcal{G} & j \leq 1 \\ \{\mathbf{w}_2\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^3(j) = \begin{cases} \mathcal{G} & j \leq 1 \\ \{\mathbf{w}_3\} & j = 2 \\ \emptyset & j > 2 \end{cases}.$$

These filtrations define a tropical toric reflexive sheaf, but not a tropical toric vector bundle. If they defined a tropical toric vector bundle, there would be a basis for $\underline{\mathcal{M}}$ for which $E^1(2) = \{\mathbf{w}_1\}$, $E^2(2) = \{\mathbf{w}_2\}$, and $E^3(2) = \{\mathbf{w}_3\}$ are all joins of elements of this basis. But this would mean that the three elements $\{\mathbf{w}_1, \mathbf{w}_2, \mathbf{w}_3\}$ are part of a basis for the rank-2 matroid $\underline{\mathcal{M}}$.

- (4) The tropical reflexive sheaf in Example 5.3 (5) is a tropical toric vector bundle as Lemma 5.5 is satisfied with the basis $\{\mathbf{w}_1, \mathbf{w}_3, \mathbf{w}_4\}$ for both cones. \diamond

Remark 5.7. For toric vector bundles, Klyachko [Kly89, Remark 2.2.2] shows that Condition 5.3 is also equivalent to the fact that the subspaces $E^i(j)$ generate a *distributive lattice*. Recall that a lattice L is *distributive* if $x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$ for all $x, y, z \in L$, or equivalently $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z)$ for all $x, y, z \in L$. This is no longer equivalent in this setting.

To see this, let \mathcal{M} be the Vamos matroid with the trivial valuated matroid structure. This is the rank-4 matroid with ground set $\mathcal{G} = \{1, \dots, 8\}$, and all subsets of size at most 4 independent except

$$\{\{1, 2, 3, 4\}, \{1, 4, 5, 6\}, \{1, 4, 7, 8\}, \{2, 3, 5, 6\}, \{2, 3, 7, 8\}\}.$$




FIGURE 5




FIGURE 6

Let \mathcal{E} be the tropical toric reflexive sheaf on $\text{trop}(\mathbb{A}^2)$ with

$$E^1(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{1, 2\} & j = 1, \\ \emptyset & j > 1 \end{cases}, \quad E^2(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{3, 4\} & j = 1, \\ \emptyset & j > 1 \end{cases}.$$

The sublattice of $\mathcal{L}(\underline{\mathcal{M}})$ generated by $\mathcal{L}(\mathcal{E})$ is shown in Figure 5.

Note that this lattice is distributive. However there is no basis B for $\underline{\mathcal{M}}$ with all $E^i(j)$ a join of elements of B , so \mathcal{E} is not a tropical toric vector bundle.

This contrasts with the case of usual toric vector bundles, where every toric reflexive sheaf on a surface is a toric vector bundle (see [Kly89, Example 2.3 (4)]). In Corollary 5.19 we recover a tropical version of that result by placing an additional assumption on the reflexive sheaf.

A toric vector bundle on X_Σ gives rise to a tropical toric vector bundle on $\text{trop}(X_\Sigma)$ as follows.

Definition 5.8. Let X_Σ be a toric variety and let \mathcal{E} be a toric vector bundle of rank r on X_Σ . Let $\underline{\mathcal{M}}$ be a realizable matroid for \mathcal{E} in the sense of Definition 3.5, with realization $\mathcal{G} \subseteq K^r$. The tropicalization $\text{trop}(\mathcal{E})$ of the pair $(\mathcal{E}, \mathcal{G})$ is the tropical toric vector bundle corresponding to the valuated matroid $\underline{\mathcal{M}}$ induced by \mathcal{G} . This has underlying matroid $\underline{\mathcal{M}}$. The subspaces $E^i(j)$ of the filtration defining \mathcal{E} correspond to flats of $\underline{\mathcal{M}}$ by construction, and the compatibility condition (5.2) is satisfied for $\text{trop}(\mathcal{E})$ because it is satisfied for \mathcal{E} . Tropical toric vector bundles of the form $\text{trop}(\mathcal{E})$ are called realizable.

The module (5.1) is the tropicalization of the Cox module for \mathcal{E} as given in Proposition 4.3.

Example 5.9. (1) The tropical tangent sheaf in Example 5.3 (2) is the tropicalization of the tangent sheaf on the toric variety X_Σ . See [Kly89, Example 2.3 (5)] for the filtration of the tangent sheaf.
 (2) Consider the representable matroid $\underline{\mathcal{M}}$ of rank 3 on the ground set

$$\mathcal{G} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2 - \mathbf{e}_3\} \subseteq K^3.$$

The lattice of flats of $\underline{\mathcal{M}}$ is given in Figure 6.

Consider the following filtrations of flats.

$$E^1(j) = \begin{cases} \mathcal{G} & j \leq -1 \\ \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_1 - \mathbf{e}_2\} & j = 0 \\ \{\mathbf{e}_1\} & 0 < j \leq 4 \\ \emptyset & j > 4 \end{cases}, \quad E^2(j) = \begin{cases} \mathcal{G} & j \leq -2 \\ \{\mathbf{e}_2, \mathbf{e}_3, \mathbf{e}_2 - \mathbf{e}_3\} & -2 < j \leq 0 \\ \{\mathbf{e}_3\} & 0 < j \leq 3 \\ \emptyset & j > 3 \end{cases}, \quad E^0(j) = \begin{cases} \mathcal{G} & j \leq -1 \\ \{\mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2 - \mathbf{e}_3\} & -1 < j \leq 2 \\ \{\mathbf{e}_2 - \mathbf{e}_3\} & j = 3 \\ \emptyset & j > 3 \end{cases}.$$

Then $(\underline{\mathcal{M}}, \mathcal{G}, \{E^i(j)\})$ defines a tropical toric vector bundle on $\text{trop}(\mathbb{P}^2)$. This tropical sheaf is obtained by tropicalizing the toric vector bundle in [DJS, Example 4.2].

◇

Remark 5.10. Note that the tropicalization of a toric vector bundle depends not only on the choice of matroid $\underline{\mathcal{M}}$ for \mathcal{E} , but also on the choice of realization of that matroid, as that affects the valuated matroid.





FIGURE 7. Fano Matroid

FIGURE 8. Fan for X_Σ

Example 5.11. Example 3.6 gives three different choices of matroid for a rank-3 toric vector bundle \mathcal{E} on \mathbb{P}^2 . This gives rise to three different choices of tropicalization of \mathcal{E} . We emphasize that the choice of matroid is part of the data of the tropical vector bundle. \diamond

Remark 5.12. Given a choice of a realization \mathcal{G} for a matroid for a rank- r toric vector bundle \mathcal{E} on a toric variety X_Σ , following [DJS, Remark 3.6] we can construct a surjection

$$(5.4) \quad \bigoplus_{\mathbf{w} \in \mathcal{G}} \mathcal{O}(D_{\mathbf{w}}) \rightarrow \mathcal{E}$$

as follows. Recall that $\bigoplus_{\mathbf{w} \in \mathcal{G}} \mathcal{O}(D_{\mathbf{w}})$ is a toric vector bundle, with corresponding filtration on $K^{|\mathcal{G}|}$ the direct sum of the filtrations corresponding to each $\mathcal{O}(D_{\mathbf{w}})$ as given in Example 5.3 (1); the vector \mathbf{a} there records the coefficients of torus invariant divisors in $D_{\mathbf{w}}$. Consider the linear map $K^{|\mathcal{G}|} \rightarrow K^r$ by $\mathbf{e}_i \mapsto \mathbf{w}_i$. The definition of $D_{\mathbf{w}}$ implies that this map respects the filtrations on $K^{|\mathcal{G}|}$ and K^r , so by Klyachko's equivalence of categories induces a morphism $\bigoplus_{\mathbf{w} \in \mathcal{G}} \mathcal{O}(D_{\mathbf{w}}) \rightarrow \mathcal{E}$ as required. Since each $E^i(j)$ is a flat of the matroid, the linear map is surjective on the filtration, so the morphism is surjective. The tropicalization of \mathcal{E} can thus be thought of as tropicalizing the presentation 5.4.

Remark 5.13. The semimodule corresponding to $\text{trop}(\mathcal{E})$ is

$$(5.5) \quad \bigoplus_{i=1}^m S(\mathbf{d}_{\mathbf{w}_i}) / \mathcal{B}(\text{trop}(R)),$$

where $\mathcal{B}()$ denotes the bend locus of [GG16], R is as in Proposition 4.3, and $\text{trop}(R)$ is the submodule obtained by taking $\sum_{k=1}^m c_k \mathbf{x}^{\mathbf{u}_k}$ to $\bigoplus_{k=1}^m \text{val}(c_k) \circ \mathbf{x}^{\mathbf{u}_k} \in S^m$.

Example 5.14. For the tropicalization of the tangent bundle to \mathbb{P}^2 , as in Example 5.3 (2), the semimodule defining $\text{trop}(\mathcal{T})$ over $\text{Cox}(\text{trop}(\mathbb{P}^2)) = \overline{\mathbb{R}}[x, y, z]$ is

$$(\overline{\mathbb{R}}[x, y, z](1, 0, 0) \oplus \overline{\mathbb{R}}[x, y, z](0, 1, 0) \oplus \overline{\mathbb{R}}[x, y, z](0, 1, 0)) / \mathcal{B}(\langle x\mathbf{e}_1 + y\mathbf{e}_2 + z\mathbf{e}_3 \rangle)$$

over $\text{Cox}(\text{trop}(\mathbb{P}^2)) = \overline{\mathbb{R}}[x, y, z]$. Note that on the affine chart $x \neq 0$, this becomes

$$x\overline{\mathbb{R}}[y/x, z/x] \oplus y\overline{\mathbb{R}}[y/x, z/x] \oplus z\overline{\mathbb{R}}[y/x, z/x] / \mathcal{B}(\mathbf{e}_1 + y/x\mathbf{e}_2 + z/x\mathbf{e}_3).$$

Note that this tropicalization, and presentation of the Cox module, comes from tropicalizing the presentation for $\text{trop}(\mathcal{T})$ coming from the Euler sequence as discussed in Remark 5.12. \diamond

However, there are tropical toric vector bundles that are not the tropicalization of a toric vector bundle on some toric variety, as the following example shows.

Example 5.15. Let \mathcal{M} be the trivial valuated matroid structure on the Fano matroid on $\mathcal{G} = \{1, \dots, 7\}$. The Fano matroid is the rank-3 matroid whose dependent sets of size three are the lines in Figure 7.

Let X_Σ be the smooth toric surface whose rays are the columns of the matrix

$$\begin{pmatrix} 1 & 1 & 0 & -1 & -1 & -1 & 0 \\ 0 & 1 & 1 & 1 & 0 & -1 & -1 \end{pmatrix}.$$

The corresponding fan is shown on the right of Figure 8.

We define a tropical toric vector bundle by setting

$$E^1(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{1, 2, 3\} & j = 1 \\ \{1\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^2(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{2, 4, 6\} & j = 1 \\ \{2\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^3(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{3, 4, 5\} & j = 1 \\ \{3\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^4(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{1, 4, 7\} & j = 1 \\ \{4\} & j = 2 \\ \emptyset & j > 2 \end{cases},$$

$$E^5(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{2, 5, 7\} & j = 1 \\ \{5\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^6(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{1, 5, 6\} & j = 1 \\ \{6\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^7(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{3, 6, 7\} & j = 1 \\ \{7\} & j = 2 \\ \emptyset & j > 2 \end{cases}.$$

Numbering the rays of the fan Σ from 1 to 7, using that X_Σ is smooth, we can check the compatibility condition (5.3) as follows:

For $E^1(j)$ and $E^2(j)$ the basis is given by $B = \{1, 2, 4\}$

For $E^2(j)$ and $E^3(j)$ the basis is given by $B = \{2, 3, 4\}$

For $E^3(j)$ and $E^4(j)$ the basis is given by $B = \{1, 4, 5\}$

For $E^4(j)$ and $E^5(j)$ the basis is given by $B = \{4, 5, 7\}$

For $E^5(j)$ and $E^6(j)$ the basis is given by $B = \{2, 5, 6\}$

For $E^6(j)$ and $E^7(j)$ the basis is given by $B = \{1, 6, 7\}$

For $E^7(j)$ and $E^1(j)$ the basis is given by $B = \{1, 3, 7\}$.

We claim that \mathcal{E} is not realizable over any field of characteristic not equal to 2. This is immediate from the fact that \mathcal{M} is not realizable over any field of characteristic not equal to 2, since the Fano matroid is the only rank-3 matroid on 7 elements containing the given flats of rank one and two in our choice of $E^i(j)$, so this failure of realizability cannot be fixed by choosing another matroid containing $\mathcal{L}(\mathcal{E})$. \diamond

5.3. Partially modular tropical toric vector bundles. We now introduce a subclass of tropical toric vector bundles that behave more like the tropicalizations of toric vector bundles.

Definition 5.16. A tropical toric reflexive sheaf or tropical toric vector bundle is *partially modular* if the following condition holds for all $1 \leq i, j \leq s$, and $k, l \in \mathbb{Z}$:

$$(5.6) \quad \begin{aligned} & \text{rk}((E^i(k-1) \wedge E^j(l)) \vee (E^i(k) \wedge E^j(l-1))) \\ &= \text{rk}(E^i(k-1) \wedge E^j(l)) + \text{rk}(E^i(k) \wedge E^j(l-1)) - \text{rk}(E^i(k) \wedge E^j(l)). \end{aligned}$$

Condition (5.6) requires that $(E^i(k-1) \wedge E^j(l), E^i(k) \wedge E^j(l-1))$ is a modular pair for all $1 \leq i, j \leq s$ and $k, l \in \mathbb{Z}$; see §2.4.

The following lemma is then a consequence of the fact that the lattice of all subspaces of a vector space is modular.

Lemma 5.17. *Realizable tropical toric vector bundles are partially modular.*

Proof. For any matroid constructed from a toric vector bundle the meet of two flats of the form $E^i(k)$ is their intersection, and the join is their sum. The rank of a flat is the dimension of the corresponding subspace. We then have

$$\begin{aligned} & \dim((E^i(k-1) \cap E^j(l)) + (E^i(k) \cap E^j(l-1))) \\ &= \dim(E^i(k-1) \cap E^j(l)) + \dim(E^i(k) \cap E^j(l-1)) - \dim((E^i(k-1) \cap E^j(l)) \cap (E^i(k) \cap E^j(l-1))) \\ &= \dim(E^i(k-1) \cap E^j(l)) + \dim(E^i(k) \cap E^j(l-1)) - \dim(E^i(k) \cap E^j(l)). \end{aligned}$$

□

Lemma 5.18. *Let \mathcal{E} be a partially modular tropical toric reflexive sheaf on a tropical toric variety. Fix $1 \leq i < j \leq s$. Then there is a basis B of the matroid $\underline{\mathcal{M}}(\mathcal{E})$ of \mathcal{E} with each $E^i(k)$ or $E^j(l)$ for $k, l \in \mathbb{Z}$ the join of a subset of B .*

Proof. Write $a_0 \leq \cdots \leq a_q$ for the distinct $E^i(k)$, where $a_0 = \emptyset$, and $a_q = \mathcal{G}$, and $b_0 \leq \cdots \leq b_p$ for the distinct $E^j(l)$, where again $b_0 = \emptyset$ and $b_p = \mathcal{G}$. For $1 \leq i \leq q$ and $1 \leq j \leq p$ we have

$$(a_{i-1} \wedge b_j) \vee (a_i \wedge b_{j-1}) \leq a_i \wedge b_j.$$

If the inequality is strict, then since lattice of flats of matroids are relatively complemented (see [Whi86, Proposition 3.4.4]) there is $w_{ij} \in \mathcal{L}(\underline{\mathcal{M}})$ with

$$w_{ij} \wedge ((a_{i-1} \wedge b_j) \vee (a_i \wedge b_{j-1})) = \emptyset, \text{ and } w_{ij} \vee ((a_{i-1} \wedge b_j) \vee (a_i \wedge b_{j-1})) = a_i \wedge b_j,$$

and $\text{rk}(w_{ij}) = \text{rk}(a_i \wedge b_j) - \text{rk}((a_{i-1} \wedge b_j) \vee (a_i \wedge b_{j-1}))$. If the inequality is not strict, we set $w_{ij} = \emptyset$.

We now show that for any $1 \leq i \leq q$ and $1 \leq j \leq p$ we have

$$(5.7) \quad a_i \wedge b_j = \vee_{k \leq i, l \leq j} w_{kl}, \text{ and } \text{rk}(a_i \wedge b_j) = \sum_{k \leq i, l \leq j} \text{rk}(w_{kl}).$$

The proof is by induction on $i + j$. The base case is when $i = j = 1$, when $w_{11} = a_1 \wedge b_1$, so the claim is true. Suppose the claim is true for smaller $i + j$. We then have

$$\begin{aligned} a_i \wedge b_j &= w_{ij} \vee ((a_{i-1} \wedge b_j) \vee (a_i \wedge b_{j-1})) \\ &= w_{ij} \vee (\vee_{k \leq i-1, l \leq j} w_{kl}) \vee (\vee_{k \leq i, l \leq j-1} w_{kl}) \\ &= \vee_{k \leq i, l \leq j} w_{kl}. \end{aligned}$$

In addition,

$$\begin{aligned} \text{rk}(a_i \wedge b_j) &= \text{rk}((a_{i-1} \wedge b_j) \vee (a_i \wedge b_{j-1})) + \text{rk}(w_{ij}) \\ &= \text{rk}(a_{i-1} \wedge b_j) + \text{rk}(a_i \wedge b_{j-1}) - \text{rk}(a_{i-1} \wedge b_{j-1}) + \text{rk}(w_{ij}) \\ &= \sum_{k \leq i-1, l \leq j} \text{rk}(w_{kl}) + \sum_{k \leq i, l \leq j-1} \text{rk}(w_{kl}) - \sum_{k \leq i-1, l \leq j-1} \text{rk}(w_{kl}) + \text{rk}(w_{ij}) \\ &= \sum_{k \leq i, l \leq j} \text{rk}(w_{kl}). \end{aligned}$$

The case $i = q, j = p$ of (5.7) states that \mathcal{G} is the join of all the w_{kl} , and the sum of their ranks is the rank of $\underline{\mathcal{M}}(\mathcal{E})$. For each $1 \leq i \leq q$ and $1 \leq j \leq p$, choose a maximal independent subset of \mathcal{G} whose join is w_{ij} , and let B be the union of all such subsets. Then $|B| = \sum \text{rk}(w_{ij}) = \text{rk } \underline{\mathcal{M}}(\mathcal{E})$, and the join of the elements of B is \mathcal{G} , so $\text{rk}(B) = \text{rk } \underline{\mathcal{M}}(\mathcal{E})$. Thus B is a basis for $\underline{\mathcal{M}}(\mathcal{E})$. Since each a_i equals $a_i \wedge b_p$, and $b_j = a_q \wedge b_j$, the lemma follows. \square

One consequence of Lemma 5.18 is the following tropical version of [Kly89, Example 2.3(4)].

Corollary 5.19. *A partially modular tropical toric reflexive sheaf \mathcal{E} on the tropicalization of a smooth toric surface $\text{trop}(X_\Sigma)$ is a tropical toric vector bundle.*

Proof. From Lemma 5.18 we know that for all cones $\sigma \in \Sigma$ there is a basis B_σ of $\underline{\mathcal{M}}(\mathcal{E})$ with every $E^i(j)$ for all $\rho_i \in \sigma$ and $j \in \mathbb{Z}$ the join of a subset of B_σ . Then, Lemma 5.5 implies that condition (5.2) holds, so \mathcal{E} is a tropical toric vector bundle. \square

See Example 6.5 for examples and non-examples of partially modular tropical reflexive sheaves.

5.4. The total space of a tropical toric vector bundle. The total space of a rank- r tropical toric vector bundle on $\text{trop}(X_\Sigma)$ is a subset of $\text{trop}(X_\Sigma) \times \overline{\mathbb{R}}^{|\mathcal{G}|}$, which we now define.

Let $\mathbf{z} \in \overline{\mathbb{R}}^s$ be the Cox coordinates of a point in $\text{trop}(X_\Sigma)$. The fiber over \mathbf{z} of a tropical toric vector bundle $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ on $\text{trop}(X_\Sigma)$ is the tropical linear space in $\overline{\mathbb{R}}^{|\mathcal{G}|}$ constructed as follows. Set $m = |\mathcal{G}|$. Recall from Proposition 4.3 that for a vector $\mathbf{c} = (c_1, \dots, c_m)$ and $1 \leq i \leq m$ we set $\mathbf{u}_i = \mathbf{d}_{\mathbf{w}_i} - \min(\mathbf{d}_{\mathbf{w}_i} : c_l \neq \infty)$. We define $\mathbf{c}(\mathbf{z}) \in \overline{\mathbb{R}}^m$ to be

$$\mathbf{c}(\mathbf{z}) := \sum_{l=1}^m c_l \mathbf{z}^{\mathbf{u}_l} \mathbf{e}_l,$$

where all operations are tropical, so $c_l \mathbf{z}^{\mathbf{u}_l}$ is $c_l + \mathbf{z} \cdot \mathbf{u}_l$ in usual arithmetic. Note that $\mathbf{z}^{\mathbf{u}_k} = \infty$ if $z_k = \infty$ for some k with $(\mathbf{u}_k)_k > 0$.

Lemma 5.20. *For any point $\mathbf{z} \in \text{trop}(X_\Sigma)$ the set $\mathcal{C}_{\mathbf{z}}$ of elements of minimal nonempty support from $\{\mathbf{c}(\mathbf{z}) : \mathbf{c} \text{ is a circuit of } \mathcal{M}\}$ is the collection of circuits of a valuated matroid $\mathcal{M}_{\mathbf{z}}$.*

Proof. We show that the valuated circuit axioms hold for $\mathcal{C}_{\mathbf{z}}$ (see §2.2). We have $(\infty, \dots, \infty) \notin \mathcal{C}_{\mathbf{z}}$, and the elements of $\mathcal{C}_{\mathbf{z}}$ have minimal support by construction. The fact that $\mu \circ \mathbf{c}(\mathbf{z}) \in \mathcal{C}_{\mathbf{z}}$ for $\mu \in \mathbb{R}$ follows from the analogous axiom for \mathcal{M} , as $\mu \circ \mathbf{c}(\mathbf{z}) = (\mu \circ \mathbf{c})(\mathbf{z})$. Suppose now that $\mathbf{c}_1(\mathbf{z}), \mathbf{c}_2(\mathbf{z}) \in \mathcal{C}_{\mathbf{z}}$ with $\mathbf{c}_1(\mathbf{z})_i = \mathbf{c}_2(\mathbf{z})_i < \infty$ and $\mathbf{c}_1(\mathbf{z})_j < \mathbf{c}_2(\mathbf{z})_j$.

Write \mathbf{u}_i^α for the vector \mathbf{u}_i corresponding to \mathbf{c}_α for $\alpha = 1, 2$. Since $\mathbf{c}_\alpha(\mathbf{z})_i < \infty$ for $\alpha = 1, 2$, we have we have $(\mathbf{u}_i^\alpha)_k = 0$ whenever $z_k = \infty$, so $(\mathbf{d}_{\mathbf{w}_i})_k = \min((\mathbf{d}_{\mathbf{w}_i})_k : c_{\alpha l} \neq \infty)$ whenever $z_k = \infty$. Thus $\min((\mathbf{d}_{\mathbf{w}_i}) : c_{1l} \neq \infty)_k = \min((\mathbf{d}_{\mathbf{w}_i}) : c_{2l} \neq \infty)_k$ for all such k . For any $\mathbf{c} \in \overline{\mathbb{R}}^m$, set

$$\lambda_{\mathbf{c}} = \ominus_{k:z_k \neq \infty} \min((\mathbf{d}_{\mathbf{w}_i} : c_{1l} \neq \infty)_k).$$

Since $\mathbf{c}_1(\mathbf{z})_i = c_{1i} \mathbf{z}^{\mathbf{u}_i^1} = \mathbf{c}_2(\mathbf{z})_i = c_{2i} \mathbf{z}^{\mathbf{u}_i^2} < \infty$, we have $\lambda_{\mathbf{c}_\alpha} \circ \mathbf{z}^{\mathbf{u}_i^\alpha} = \mathbf{z}^{\mathbf{d}_{\mathbf{w}_i}}$ for $\alpha = 1, 2$, and thus $c_{1i} \circ \lambda_{\mathbf{c}_2} = c_{2i} \circ \lambda_{\mathbf{c}_1}$. Similarly, either $c_{1j} \circ \lambda_{\mathbf{c}_2} < c_{2j} \circ \lambda_{\mathbf{c}_1}$ or $\mathbf{c}_2(\mathbf{z})_j = \infty$.

By the circuit elimination axiom for \mathcal{M} applied to the circuits $\lambda_{\mathbf{c}_2} \circ \mathbf{c}_1$ and $\lambda_{\mathbf{c}_1} \circ \mathbf{c}_2$, there is a circuit \mathbf{c}_3 of \mathcal{M} with $c_{3i} = \infty$, $\mathbf{c}_3_j = \lambda_{\mathbf{c}_2} \circ c_{1j}$, and $\mathbf{c}_3 \geq \lambda_{\mathbf{c}_2} \circ \mathbf{c}_1 \oplus \lambda_{\mathbf{c}_1} \circ \mathbf{c}_2$ coordinatewise. We may assume that \mathbf{c}_3 is chosen to have $\text{supp}(\mathbf{c}_3(\mathbf{z}))$ minimal among all such choices for \mathbf{c}_3 . Set $\mathbf{c}_4 = (\lambda_{\mathbf{c}_3} - \lambda_{\mathbf{c}_1} - \lambda_{\mathbf{c}_2}) \circ \mathbf{c}_3$. We now show that, if its support is minimal, $\mathbf{c}_4(\mathbf{z})$ is the required circuit.

We have $c_{4i} = \infty$, and since $\text{supp}(\mathbf{c}_4) = \text{supp}(\mathbf{c}_3)$ we have $\lambda_{\mathbf{c}_4} = \lambda_{\mathbf{c}_3}$. In addition, for any $1 \leq t \leq m$,

$$\begin{aligned} c_{4t} &= \lambda_{\mathbf{c}_3} - \lambda_{\mathbf{c}_1} - \lambda_{\mathbf{c}_2} + c_{3t} \\ (5.8) \quad &\geq \lambda_{\mathbf{c}_3} - \lambda_{\mathbf{c}_1} - \lambda_{\mathbf{c}_2} + \min(\lambda_{\mathbf{c}_2} + c_{1t}, \lambda_{\mathbf{c}_1} + c_{2t}) \\ &= \lambda_{\mathbf{c}_4} + \min(c_{1t} - \lambda_{\mathbf{c}_1}, c_{2t} - \lambda_{\mathbf{c}_2}), \end{aligned}$$

with equality if $t = j$. Thus $c_{4t} - \lambda_{\mathbf{c}_4} \geq \min(c_{1t} - \lambda_{\mathbf{c}_1}, c_{2t} - \lambda_{\mathbf{c}_2})$, with equality if $t = j$.

We now show that $\mathbf{c}_4(\mathbf{z}) \geq \mathbf{c}_1(\mathbf{z}) \oplus \mathbf{c}_2(\mathbf{z})$. Consider $t \neq i$ with $\min(\mathbf{c}_1(\mathbf{z})_t, \mathbf{c}_2(\mathbf{z})_t) < \infty$. Write $\mathbf{c}_\alpha(\mathbf{z})_t$ for this minimum, where $\alpha \in \{1, 2\}$. Since $\mathbf{c}_\alpha(\mathbf{z})_t < \infty$, it follows that

$$(5.9) \quad (\mathbf{d}_{\mathbf{w}_t})_k = \min((\mathbf{d}_{\mathbf{w}_l})_k : c_{\alpha l} \neq \infty) = (\mathbf{d}_{\mathbf{w}_i})_k$$

for all k with $z_k = \infty$. As $(\mathbf{d}_{\mathbf{w}_i})_k$ also equals $\min((\mathbf{d}_{\mathbf{w}_l})_k : c_{\beta l} \neq \infty)$ for $\beta = \{1, 2\} \setminus \{\alpha\}$ and such k , adding $\ominus_{k:z_k \neq \infty} z_k^{(\mathbf{d}_{\mathbf{w}_t})_k}$ to both sides of (5.8) yields

$$(5.10) \quad c_{4t} + \ominus_{k:z_k \neq \infty} z_k^{(\mathbf{d}_{\mathbf{w}_t})_k} - \lambda_{\mathbf{c}_4} \geq \min(\mathbf{c}_1(\mathbf{z})_t, \mathbf{c}_2(\mathbf{z})_t)$$

for t where the right-hand side is not equal to ∞ . We have equality when $t = j$. When $c_{4t} < \infty$, since $t \in \text{supp}(\mathbf{c}_4) \subsetneq \text{supp}(\mathbf{c}_1) \cup \text{supp}(\mathbf{c}_2)$, the equality (5.9) implies that $\min(\mathbf{d}_{\mathbf{w}_l} : c_{4l} \neq \infty)_k = (\mathbf{d}_{\mathbf{w}_t})_k$ for all k with $z_k = \infty$. Thus the left-hand side of (5.10) equals $\mathbf{c}_4(\mathbf{z})$. This means that

$$\mathbf{c}_4(\mathbf{z})_t \geq \min(\mathbf{c}_1(\mathbf{z})_t, \mathbf{c}_2(\mathbf{z})_t)$$

with equality when $t = j$, whenever the right-hand side is not equal to ∞ .

If the right-hand side of (5.10) is equal to ∞ , so $\mathbf{c}_1(\mathbf{z})_t = \mathbf{c}_2(\mathbf{z})_t = \infty$, then either $c_{1t} = c_{2t} = \infty$, so $c_{4t} = \infty$ and thus $\mathbf{c}_4(\mathbf{z})_t = \infty$, or $(\mathbf{d}_{\mathbf{w}_t})_k > \min((\mathbf{d}_{\mathbf{w}_l})_k : c_{1l} \neq \infty) = \min((\mathbf{d}_{\mathbf{w}_l})_k : c_{2l} \neq \infty) = (\mathbf{d}_{\mathbf{w}_i})_k = (\mathbf{d}_{\mathbf{w}_j})_k$

for some k with $z_k = \infty$. However $\mathbf{c}_4(\mathbf{z})_j < \infty$, so we must also have $\min((\mathbf{d}_{\mathbf{w}_l})_k : c_{4l} \neq \infty) = (\mathbf{d}_{\mathbf{w}_j})_k$, and so $(\mathbf{d}_{\mathbf{w}_t})_k > \min((\mathbf{d}_{\mathbf{w}_l})_k : c_{4l} \neq \infty)$. So we conclude that if $\mathbf{c}_1(\mathbf{z})_t = \mathbf{c}_2(\mathbf{z})_t = \infty$ then $\mathbf{c}_4(\mathbf{z})_t = \infty$, and so

$$\mathbf{c}_4(\mathbf{z}) \geq \mathbf{c}_1(\mathbf{z}) \oplus \mathbf{c}_2(\mathbf{z}).$$

If $\mathbf{c}_4(\mathbf{z}) \in \mathcal{C}_{\mathbf{z}}$ then we are done. Otherwise there would be a circuit \mathbf{c}_5 of \mathcal{M} with $\text{supp}(\mathbf{c}_5(\mathbf{z}))$ minimal and $\text{supp}(\mathbf{c}_5(\mathbf{z})) \subsetneq \text{supp}(\mathbf{c}_4(\mathbf{z}))$. After scaling we may assume that $\mathbf{c}_5(\mathbf{z}) \geq \mathbf{c}_4(\mathbf{z})$, and $\mathbf{c}_5(\mathbf{z})_l = \mathbf{c}_4(\mathbf{z})_l$ for some l . If we cannot choose $l = j$, then repeating the above procedure with \mathbf{c}_4 and \mathbf{c}_5 playing the roles of \mathbf{c}_1 and \mathbf{c}_2 , with i replaced by l would yield a circuit \mathbf{c}_6 with $\text{supp}(\mathbf{c}_6(\mathbf{z})) \subsetneq \text{supp}(\mathbf{c}_4(\mathbf{z}))$, $\mathbf{c}_6(\mathbf{z})_i = \infty$, and $\mathbf{c}_6(\mathbf{z})_j = \mathbf{c}_1(\mathbf{z})_j$, which contradicts the minimality assumption on the choice of \mathbf{c}_3 . We thus conclude that $\mathbf{c}_5(\mathbf{z})_j$ is the desired elimination. \square

Remark 5.21. When $\mathbf{c}(\mathbf{z}) \notin \mathcal{C}_{\mathbf{z}}$ for a circuit \mathbf{c} of \mathcal{M} , it is a tropical sum of elements of $\mathcal{C}_{\mathbf{z}}$, so a vector of $\mathcal{M}_{\mathbf{z}}$. This follows from repeated applications of the technique of the last paragraph of the proof of Lemma 5.20 to get circuits \mathbf{c}_j with $\mathbf{c}_j(\mathbf{z}) \geq \mathbf{c}(\mathbf{z})$ and $\mathbf{c}_j(\mathbf{z})_j = \mathbf{c}(\mathbf{z})_j$ for all $j \in \text{supp}(\mathbf{c}(\mathbf{z}))$. The vector $\mathbf{c}(\mathbf{z})$ is then the tropical sum of all such $\mathbf{c}_j(\mathbf{z})$.

In light of Lemma 5.20, we make the following definition.

Definition 5.22. Let \mathcal{E} be a tropical toric vector bundle on $\text{trop}(X_{\Sigma})$. The fiber of \mathcal{E} over a point $\mathbf{z} \in \overline{\mathbb{R}}^s$ giving the Cox coordinates of a point in $\text{trop}(X_{\Sigma})$ is the *tropical linear space* in $\overline{\mathbb{R}}^{|\mathcal{G}|}$ of the valuated matroid $\mathcal{M}_{\mathbf{z}}$. This is the set of all $\mathbf{a} \in \overline{\mathbb{R}}^{|\mathcal{G}|}$ tropically orthogonal to the circuits $\mathbf{c}(\mathbf{z})$ of $\mathcal{M}_{\mathbf{z}}$:

$$\{\mathbf{a} \in \overline{\mathbb{R}}^{|\mathcal{G}|} : \text{either } \bigoplus_{\mathbf{w} \in \mathcal{G}} \mathbf{c}(\mathbf{z})_{\mathbf{w}} a_{\mathbf{w}} = \infty, \text{ or the minimum is achieved at least twice}\}.$$

Let

$$\mathcal{C} = \{c_1 \mathbf{z}^{\mathbf{u}_1} y_1 \oplus \cdots \oplus c_m \mathbf{z}^{\mathbf{u}_m} y_m : \mathbf{c} \text{ is a circuit of } \mathcal{M}\} \subseteq \overline{\mathbb{R}}[z_1, \dots, z_s, y_1, \dots, y_m].$$

The total space of \mathcal{E} is the quotient of

$$V(\mathcal{C}) \setminus V(B_{\Sigma}) \subseteq \overline{\mathbb{R}}^s \times \overline{\mathbb{R}}^m$$

by the tropical dual of the class group as in §2.1. This is a subset of $\text{trop}(X_{\Sigma}) \times \overline{\mathbb{R}}^m$. The fiber over a point $\mathbf{z} \in \overline{\mathbb{R}}^s$ under the projection to the first coordinates is the fiber defined above.

Remark 5.23. The fact that the fiber over a point $\mathbf{z} \in \overline{\mathbb{R}}^s$ under the projection to the first coordinates is the tropical linear space of $\mathcal{M}_{\mathbf{z}}$ follows from the fact that the specializations of the elements of \mathcal{C} at $\mathbf{z} \in \overline{\mathbb{R}}^s$ are the elements $\mathbf{c}(\mathbf{z})$ for circuits \mathbf{c} of \mathcal{M} . The tropical prevariety defined by the collection $\mathcal{C}_{\mathbf{z}}$ is the tropical linear space of $\mathcal{M}_{\mathbf{z}}$. When $\mathbf{c}(\mathbf{z}) \notin \mathcal{C}_{\mathbf{z}}$ for some circuit \mathbf{c} , it is the tropical sum of circuits of $\mathcal{M}_{\mathbf{z}}$, so the tropical hyperplane corresponding to $\mathbf{c}(\mathbf{z})$ contains the fiber. Thus the tropical prevariety defined by all the $\mathbf{c}(\mathbf{z})$ is the fiber over \mathbf{z} .

Remark 5.24. When \mathcal{E} is realizable, so $\mathcal{E} = \text{trop}(\mathcal{F})$ for a toric vector bundle \mathcal{F} , the fiber of \mathcal{E} at a point of $\text{trop}(X_{\Sigma})$ with Cox coordinates \mathbf{z} is the tropicalization of the fiber of \mathcal{F} at any point of X_{Σ} with Cox coordinates \mathbf{x} with $\text{val}(\mathbf{x}) = \mathbf{z}$. To see this, note that when \mathbf{x} corresponds to a point in U_{σ} for $\sigma \in \Sigma$, the fiber over \mathbf{x} is

$$(P_{\mathbf{x}^{\sigma}})_{\mathbf{0}} \otimes_{(S_{\mathbf{x}^{\sigma}})_{\mathbf{0}}} ((S/\mathfrak{m}(\mathbf{x}))_{\mathbf{x}^{\sigma}})_{\mathbf{0}},$$

where P is the Cox-module of \mathcal{F} , as in Proposition 4.3, S is the Cox ring of X_{Σ} , and $\mathfrak{m}(\mathbf{x})$ is the ideal of all polynomials in S , homogeneous with respect to the grading on S , that vanish at \mathbf{x} . Given the description of P in Proposition 4.3, this is isomorphic to

$$K^m / R_{\mathbf{x}},$$

where $R_{\mathbf{x}}$ is the submodule generated by

$$\left\{ \sum_{i=1}^m c_i \mathbf{x}^{\mathbf{u}_i} \mathbf{e}_i : (c_1, \dots, c_m) \in \ker(G) \right\}.$$

The valuated matroid associated to the images of standard basis vectors of K^m in the r -dimensional vector space $K^m/R_{\mathbf{x}}$ is $\mathcal{M}_{\mathbf{z}}$. When \mathbf{x} corresponds to the identity of the torus of X_{Σ} , $\mathcal{M}_{\mathbf{z}}$ is the matroid \mathcal{M} associated to \mathcal{E} .

Remark 5.25. When \mathcal{E} is a realizable tropical toric vector bundle, there is a subset of the tropicalization of the fiber over the identity of the torus that does not depend on the choice of tropicalization. Consider, for example, the case that the valuated matroid \mathcal{M} has a trivial valuated matroid structure (so the valuation of every basis is 0). Then the tropical linear space of \mathcal{M} is the cone over the order complex of the lattice of flats of $\underline{\mathcal{M}}$ by [AK06]. As the poset $L'(\mathcal{E})$ of all intersections of the $E^i(j)$ is a subposet of this lattice of flats, its order complex is a subcomplex of the order complex of $\mathcal{L}(\underline{\mathcal{M}})$, so the cone over the order complex of $L'(\mathcal{E})$ is a subfan of the fiber. We call this the *intrinsic subcomplex*. As noted in the following examples, this often has smaller dimension than the actual fiber. This can be regarded as the polymatroid fiber (but should not be confused with the Bergman fan of the polymatroid in the sense of [CHLSW22]).

Example 5.26. The trivial rank- r tropical toric vector bundle with valuated matroid \mathcal{M} on a tropical toric variety $\text{trop}(X_{\Sigma})$ is given by the filtration with $E^i(j) = \mathcal{G}$ for $j \leq 0$, and $E^i(j) = \emptyset$ for $j > 0$ for $1 \leq i \leq s$. In this case we have $\mathbf{d}_{\mathbf{w}} = \mathbf{0} \in \mathbb{Z}^s$ for all $\mathbf{w} \in \mathcal{G}$, so $\mathbf{c}(\mathbf{z}) = \mathbf{c}$ for all $\mathbf{z} \in \overline{\mathbb{R}}^s$. This means that $\mathcal{M}_{\mathbf{z}} = \mathcal{M}$ for all \mathbf{z} , and the total space of \mathcal{E} is $\text{trop}(X_{\Sigma}) \times \Gamma$, where Γ is the tropical linear space of \mathcal{M} . In this case the intrinsic subcomplex is the origin $\mathbf{0} \in \overline{\mathbb{R}}^m$. \diamond

Example 5.27. When \mathcal{E} is the tropical tangent sheaf on $\text{trop}(\mathbb{P}^2)$, as in Example 5.3 (2), the valuated matroid \mathcal{M} has one valuated circuit up to scaling: $(0, 0, 0)$. The underlying matroid is $U(2, 3)$. At a point $[z_0 : z_1 : z_2]$ of $\text{trop}(\mathbb{P}^2)$ with $z_0, z_1, z_2 \neq \infty$ the unique (up to scaling) circuit is (z_0, z_1, z_2) , so $\mathcal{M}_{[z_0 : z_1 : z_2]}$ is a projectively equivalent valuated matroid. The fiber in $\overline{\mathbb{R}}^3$ has three cones: $(-z_0, -z_1, -z_2) + \text{cone}(\mathbf{e}_i) + \text{span}((1, 1, 1))$ for $i = 0, 1, 2$. The quotient of the fiber in $\text{trop}(\mathbb{P}^2)$ is the standard tropical line with vertex $[-z_0 : -z_1 : -z_2]$. All three cones lie in the intrinsic subcomplex.

When $z_0 = \infty$, but $z_1, z_2 \neq \infty$ we get the valuated matroid with unique (up to scaling) circuit (∞, z_1, z_2) . This has underlying matroid $U(1, 1) \oplus U(1, 2)$. The fiber in $\overline{\mathbb{R}}^3$ is the hyperplane $z_1 + y_1 = z_2 + y_2$.

When $z_0 = z_1 = \infty$, the valuated matroid has the unique (up to scaling) circuit (∞, ∞, z_2) , so the underlying matroid is $U(2, 2) \oplus U(0, 1)$. The fiber in $\overline{\mathbb{R}}^3$ is the hyperplane $y_2 = \infty$.

Recall that $\text{trop}(\mathbb{P}^2)$ is $(\overline{\mathbb{R}}^3 \setminus \{(\infty, \infty, \infty)\}) / \text{span}((1, 1, 1))$. Thus the total space of the tropical tangent sheaf is

$$(V(z_0 y_0 \oplus z_1 y_1 \oplus z_2 y_2) \setminus \{(\infty, \infty, \infty, a_1, a_2, a_3) : a_i \in \overline{\mathbb{R}}\}) / \text{span}((1, 1, 1, 0, 0, 0)) \subseteq \text{trop}(\mathbb{P}^2) \times \overline{\mathbb{R}}^3.$$

\diamond

Example 5.28. As noted in Example 5.11, the three different matroids given in Example 3.6 for that toric vector bundle \mathcal{E} on \mathbb{P}^2 give three different tropicalizations. The total spaces are as follows.

- (1) For the matroid $\underline{\mathcal{M}}_1$ with the choice of basis $\mathcal{G} = \{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$, and the corresponding trivial valuated matroid structure, we have a unique circuit $\mathbf{c} = (0, 0, 0, 0)$, and $\mathbf{d}_{\mathbf{e}_1 + \mathbf{e}_2} = (1, 0, 0)$, $\mathbf{d}_{\mathbf{e}_2} = (1, 0, 0)$, $\mathbf{d}_{\mathbf{e}_1 + \mathbf{e}_3} = (0, 1, 0)$, and $\mathbf{d}_{\mathbf{e}_3} = (0, 0, 1)$. Thus $\mathbf{c}(\mathbf{z}) = (z_0, z_0, z_1, z_2)$, so the total space is $V(z_0 y_1 \oplus z_0 y_2 \oplus z_1 y_3 \oplus z_2 y_4) \subseteq \text{trop}(\mathbb{P}^2) \times \overline{\mathbb{R}}^4$. The fiber over $z_0 = \infty$ is the classical hyperplane $z_1 + y_3 = z_2 + y_4$, while the fiber over $z_1 = \infty$ is $V(z_0 y_1 \oplus z_0 y_2 \oplus z_2 y_4)$, which is the tropical cone over a tropical line times \mathbb{R} . The intrinsic subcomplex consists of the three cones $\text{cone}((1, 1, 0, 0)) + \text{span}(1, 1, 1, 1)$, $\text{cone}((0, 0, 1, 0) + \text{span}((1, 1, 1, 1)))$, and $\text{cone}((0, 0, 0, 1)) + \text{span}((1, 1, 1, 1))$.
- (2) For the matroid $\underline{\mathcal{M}}_2$ with the choice of basis $\mathcal{G} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$, and the corresponding trivial valuated matroid structure, we have a unique circuit $\mathbf{c} = (0, \infty, 0, 0)$, and $\mathbf{d}_{\mathbf{e}_1} = (1, 0, 0)$. Thus $\mathbf{c}(\mathbf{z}) = (z_0, \infty, z_1, z_2)$, so the total space is $V(z_0 y_1 \oplus z_1 y_3 \oplus z_2 y_4) \subseteq \text{trop}(\mathbb{P}^2) \times \overline{\mathbb{R}}^3$. The fiber over $z_0 = \infty$ is again the classical hyperplane $z_1 + y_3 = z_2 + y_4$, but the fiber over $z_1 = \infty$ is the classical hyperplane $z_0 + y_1 = z_2 + y_4$. The intrinsic subcomplex is as in the previous part.

- (3) For the matroid $\underline{\mathcal{M}}_2$ with the choice of basis $\mathcal{G} = \{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3\}$, and the corresponding trivial valuated matroid structure, we have 9 circuits. We have $\mathbf{d}_{\mathbf{e}_1 - \mathbf{e}_2} = (1, 0, 0)$, and $\mathbf{d}_{\mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3} = (0, 0, 0)$. The total space is thus

$$\begin{aligned} V(y_1 \oplus y_2 \oplus y_3, z_0 y_3 \oplus z_1 y_4 \oplus y_6, z_0 y_1 \oplus z_2 y_5 \oplus y_6, z_0 y_1 \oplus z_0 y_2 \oplus z_1 y_4 \oplus z_2 y_5, \\ z_0 y_1 \oplus z_0 y_2 \oplus z_1 y_4 \oplus y_6, z_0 y_1 \oplus z_0 y_3 \oplus z_1 y_4 \oplus z_2 y_5, z_0 y_2 \oplus z_0 y_3 \oplus z_1 y_4 \oplus z_2 y_5, \\ z_0 y_2 \oplus z_0 y_3 \oplus z_2 y_5 \oplus y_6, z_0 y_2 \oplus z_1 y_4 \oplus z_2 y_5 \oplus y_6) \subseteq \text{trop}(\mathbb{P}^2) \times \overline{\mathbb{R}}^6. \end{aligned}$$

The intrinsic subcomplex consists of the three cones $\text{cone}((1, 1, 1, 0, 0, 0)) + \text{span}(1, 1, 1, 1, 1, 1)$, $\text{cone}((0, 0, 0, 1, 0, 0)) + \text{span}((1, 1, 1, 1, 1, 1))$, and $\text{cone}((0, 0, 0, 0, 1, 0)) + \text{span}((1, 1, 1, 1, 1, 1))$ in $\overline{\mathbb{R}}^6$. The fiber over $z_0 = \infty$ is

$$V(y_1 \oplus y_2 \oplus y_3, z_1 y_4 \oplus y_6, z_2 y_5 \oplus y_6, z_1 y_4 \oplus z_2 y_5),$$

which is the tropical cone over a tropical line times \mathbb{R} , and the fiber over $z_1 = \infty$ is

$$\begin{aligned} V(y_1 \oplus y_2 \oplus y_3, z_0 y_3 \oplus y_6, z_0 y_1 \oplus z_2 y_5 \oplus y_6, z_0 y_1 \oplus z_0 y_2 \oplus z_2 y_5, \\ z_0 y_1 \oplus z_0 y_2 \oplus y_6, z_0 y_1 \oplus z_0 y_3 \oplus z_2 y_5, z_0 y_2 \oplus z_0 y_3 \oplus z_2 y_5, \\ z_0 y_2 \oplus z_0 y_3 \oplus z_2 y_5 \oplus y_6, z_0 y_2 \oplus z_1 y_4 \oplus z_2 y_5 \oplus y_6) \end{aligned}$$

◇

Example 5.29. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^1(j)\})$ be the tropical toric vector bundle on $\text{trop}(\mathbb{A}^1)$ with \mathcal{M} the trivial valuated matroid structure on the Vamos matroid as in Remark 5.7, and $E^1(j) = \mathcal{G}$ for $j \leq 0$, and $E^1(j) = \emptyset$ for $j > 0$. Then the fiber over a point in tropical torus is the tropical cone in $\overline{\mathbb{R}}^8$ over Bergman fan of the Vamos matroid. ◇

6. PROPERTIES OF TROPICAL TORIC VECTOR BUNDLES

In this section we extend some standard properties of vector bundles to tropical toric vector bundles: decompositions into direct sums, tensoring by line bundles, and global sections.

6.1. Decomposition into direct sums. We now give a definition for a tropical toric vector bundle to decompose as a direct sum of other tropical toric vector bundles.

We first recall the notion of the direct sums of valuated matroids; see, for example [HLSV22]. If \mathcal{M}_1 is a valuated matroid of rank r_1 on a ground set \mathcal{G}_1 with basis valuation function ν_1 , and \mathcal{M}_2 is a valuated matroid of rank r_2 on a ground set \mathcal{G}_2 with basis valuation function ν_2 , then $\mathcal{M}_1 \oplus \mathcal{M}_2$ is a valuated matroid on the ground set $\mathcal{G}_1 \sqcup \mathcal{G}_2$ with basis valuation function $\nu(I_1 \cup I_2) = \nu_1(I_1) + \nu_2(I_2)$ for all sets $I_1 \subseteq \mathcal{G}_1$ of size r_1 and $I_2 \subseteq \mathcal{G}_2$ of size r_2 . The lattice of flats $\mathcal{L}(\mathcal{M}_1 \oplus \mathcal{M}_2)$ is isomorphic to $\mathcal{L}(\mathcal{M}_1) \times \mathcal{L}(\mathcal{M}_2)$.

This lets us define the direct sum of tropical toric vector bundles.

Definition 6.1. Let $\mathcal{E}_1 = (\mathcal{M}_1, \mathcal{G}_1, \{E_{\mathcal{M}_1}^i(j)\})$ and $\mathcal{E}_2 = (\mathcal{M}_2, \mathcal{G}_2, \{E_{\mathcal{M}_2}^i(j)\})$ be tropical toric vector bundles on $\text{trop}(X_\Sigma)$. The tropical toric vector bundle $\mathcal{E}_1 \oplus \mathcal{E}_2$ is given by the valuated matroid $\mathcal{M}_1 \oplus \mathcal{M}_2$ on the ground set $\mathcal{G}_1 \sqcup \mathcal{G}_2$ with $E^i(j) = E_{\mathcal{M}_1}^i(j) \sqcup E_{\mathcal{M}_2}^i(j)$. Compatibility for $\mathcal{E}_1 \oplus \mathcal{E}_2$ follows by setting $B_\sigma = B_{\sigma, \mathcal{M}_1} \sqcup B_{\sigma, \mathcal{M}_2}$, and $\mathbf{u}(\sigma) = \mathbf{u}(\sigma)_{\mathcal{M}_1} \cup \mathbf{u}(\sigma)_{\mathcal{M}_2}$.

A tropical toric vector bundle \mathcal{E} is *decomposable* if it is the direct sum of two tropical toric vector bundles of smaller rank. A tropical toric vector bundle is *indecomposable* if it is not decomposable.

Recall that a matroid is *connected* if it is not a nontrivial direct sum of two matroids of lower ranks. The following lemma shows that the Krull-Schmidt theorem (see [Per03, Section 4.3]) holds for tropical toric vector bundles.

Lemma 6.2. *A tropical toric vector bundle $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ is indecomposable if and only if the underlying matroid $\underline{\mathcal{M}}$ is connected. Every tropical toric vector bundle can be written uniquely as the direct sum of indecomposable toric vector bundles.*

Proof. The first claim follows from the fact that a valuated matroid \mathcal{M} is a direct sum of two valuated matroids of smaller ranks if and only if the underlying matroid $\underline{\mathcal{M}}$ is not connected. If the valuated matroid decomposes, then each flat in the lattice of flats of the underlying matroid is a union of flats of the two constituent matroids, which lets us define the summand tropical toric vector bundles. The second claim then follows from the fact that matroids have well-defined connected components. \square

Remark 6.3. If \mathcal{E} is an indecomposable toric vector bundle, then $\text{trop}(\mathcal{E})$ is also indecomposable. To see this, the key observation is that when a realizable matroid is decomposable, it is decomposable as a direct sum of *realizable* matroids. Thus if $\text{trop}(\mathcal{E})$ is decomposable, the realizable matroid $\underline{\mathcal{M}}(\mathcal{E})$ is a direct sum of matroids of smaller ranks, which are themselves realizable. This means that the vector space E decomposes as a direct sum of subspaces spanned by the realizations of each of the summand matroids. This induces a decomposition of \mathcal{E} as a direct sum of toric vector bundles.

Note, however, that the tropicalization of a decomposable vector bundle might not be decomposable; this depends on the choice of matroid for the bundle \mathcal{E} . For example, consider the toric vector bundle \mathcal{E} of Example 3.2. This is decomposable: we can write $K^3 = \text{span}(\mathbf{e}_1, \mathbf{e}_3) \oplus \text{span}(\mathbf{e}_2)$. This is reflected in the matroid \mathcal{M}_2 from that example, which is the direct sum $U_{2,3} \oplus U_{1,1}$. The matroids \mathcal{M}_1 and \mathcal{M}_3 do not decompose as direct sums, however. Given a decomposable toric vector bundle $\mathcal{E} \cong \mathcal{E}_1 \oplus \mathcal{E}_2$ we can always *choose* a valuated matroid \mathcal{M} for \mathcal{E} for which $\text{trop}(\mathcal{E})$ is decomposable.

A tropical toric vector bundle \mathcal{E} *splits as a direct sum of line bundles* if there are $\mathcal{E}_1, \dots, \mathcal{E}_r$ all of rank one with $\mathcal{E} = \mathcal{E}_1 \oplus \dots \oplus \mathcal{E}_r$. If \mathcal{E} is tropically minimal we then have $\underline{\mathcal{M}}(\mathcal{E})$ equal to the uniform matroid $U_{r,r}$, as a tropically minimal tropical vector bundle of rank one has matroid $U_{1,1}$.

We now see the use of the partially modular condition of Definition 5.16 for decomposing into direct sums.

Proposition 6.4 (Tropical version of Grothendieck's theorem). (*cf.* [Kly89, Example 2.3 (3)], [GUZ22, Example 3.3]) *Let \mathcal{E} be a partially modular tropical toric vector bundle on $\text{trop}(\mathbb{P}^1)$ that is tropically minimal. Then \mathcal{E} splits as a direct sum of line bundles.*

Proof. Since the fan of \mathbb{P}^1 has only two rays, we have two flags of flats $E^0(j)$ and $E^1(k)$ for $j, k \in \mathbb{Z}$. By Lemma 5.18 there is a basis B of the lattice of flats $\mathcal{L}(\underline{\mathcal{M}})$ for which each $E^0(j)$ and $E^1(k)$ is the join of a subset of B . Since \mathcal{E} is tropically minimal, the lattice of flats must equal the lattice of subsets of B , so the underlying matroid is the uniform matroid $U_{r,r}$, where r is the rank of \mathcal{E} . This is the direct sum $\oplus_{i=1}^r U_{1,1}$, so we conclude that \mathcal{E} splits as the direct sum of line bundles. \square

Example 6.5. (1) Consider the tropically minimal tropical toric vector bundle \mathcal{E} on $\text{trop}(\mathbb{P}^1)$ of Example 5.3 (5). The lattice of flats is shown in Figure 4. The underlying matroid decomposes as $U_{3,3} = \{1\} \oplus \{3\} \oplus \{4\}$. Consider the filtrations

$$L_1^1(j) = \begin{cases} \{1\} & j \leq 2 \\ \emptyset & j > 2 \end{cases}, L_2^1(j) = \begin{cases} \{3\} & j \leq 1 \\ \emptyset & j > 1 \end{cases}, L_3^1(j) = \begin{cases} \{4\} & j \leq 0 \\ \emptyset & j > 0 \end{cases},$$

$$L_1^2(j) = \begin{cases} \{3\} & j \leq 2 \\ \emptyset & j > 2 \end{cases}, L_2^2(j) = \begin{cases} \{4\} & j \leq 1 \\ \emptyset & j > 1 \end{cases}, L_3^2(j) = \begin{cases} \{1\} & j \leq 0 \\ \emptyset & j > 0 \end{cases}.$$

These give rise to tropical toric line bundles \mathcal{L}_1 , \mathcal{L}_2 and \mathcal{L}_3 respectively. Note that $\mathcal{E} = \mathcal{L}_1 \oplus \mathcal{L}_2 \oplus \mathcal{L}_3$.

The tropical toric vector bundle \mathcal{E} is partially modular; in fact, the entire lattice of flats in Figure 4 is a modular lattice.

(2) Consider the rank-3 realizable matroid $\underline{\mathcal{M}}$ with ground set $\mathcal{G} = \{\mathbf{w}_1 = (1, 0, 0), \mathbf{w}_2 = (0, 1, 0), \mathbf{w}_3 = (0, 0, 1), \mathbf{w}_4 = (1, 1, 1)\}$. Let \mathcal{E} be the tropical toric vector bundle on $\text{trop}(\mathbb{P}^1)$ given by

$$E^0(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{w}_1, \mathbf{w}_2\} & j = 1 \\ \emptyset & j > 1 \end{cases}, E^1(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{w}_3, \mathbf{w}_4\} & j = 1 \\ \emptyset & j > 1 \end{cases}.$$

Then \mathcal{E} is not decomposable. It is also not partially modular, as

$$\begin{aligned} 3 &= \text{rk}((E^0(0) \wedge E^1(1)) \vee (E^0(1) \wedge E^1(0))) \\ &\leq \text{rk}(E^0(0) \wedge E^1(1)) + \text{rk}(E^0(1) \wedge E^1(0)) - \text{rk}(E^0(1) \wedge E^1(1)) \\ &= 4 \end{aligned}$$

Note that while $\underline{\mathcal{M}}$ is realizable, \mathcal{E} is not realizable. This is because in the realizable case we impose the condition that all intersections of $E^i(j)$ are flats of the matroid, which is not the case here, as $\text{span}(E^0(1)) \cap \text{span}(E^1(1)) = \text{span}((1, 1, 0))$, and a multiple of $(1, 1, 0)$ is not in the ground set of $\underline{\mathcal{M}}$. This shows that a condition like partial modularity is necessary for Proposition 6.4.

(3) Consider $\mathcal{O}(aD_0 + bD_1)$ on \mathbb{P}^1 as a toric vector bundle. This has

$$E^0(j) = \begin{cases} K & j \leq a \\ 0 & j > a \end{cases}, E^1(j) = \begin{cases} K & j \leq b \\ 0 & j > b \end{cases}.$$

for $i = 0, 1$. Let $\mathcal{E} = \mathcal{O}(D_0 + 2D_1) \oplus \mathcal{O}(2D_0 + D_1)$, and choose the matroid $\mathcal{G} = \{(1, 0), (0, 1), (1, 1)\}$ for \mathcal{E} . Then $\text{trop}(\mathcal{E})$ is not decomposable, so not a direct sum of line bundles. In this case $\text{trop}(\mathcal{E})$ is partially modular, but not tropically minimal; the subposet given by the matroid of $\{(1, 0), (0, 1)\}$ contains the $E^i(j)$, so this shows the necessity of the tropically minimal condition for Proposition 6.4.

◇

Remark 6.6. Note that Example 6.5 (2) shows that a tropical vector bundle whose underlying matroid is realizable need not be a realizable tropical vector bundle.

6.2. Tensor product with line bundles. We now define the tensor product of a tropical toric vector bundle by a tropical line bundle.

Recall that for a toric vector bundle \mathcal{E} on a toric variety X_Σ with associated filtrations $(E, \{E^i(j)\})$, the tensor product of \mathcal{E} by a torus-invariant line bundle is obtained by shifting the filtration. Specifically, consider the line bundle $\mathcal{O}_X(D)$, for some torus-invariant Cartier divisor $D = \sum_{i \in \Sigma(1)} a_i D_i$ with the associated filtrations $(L, \{L^i(j)\})$. Then the filtrations associated to the tensor product $\mathcal{E} \otimes \mathcal{O}_X(D)$ are given by $(E, \{E^i(j - a_i)\})$ (see [Gon11, Example II.9.]). Motivated by this we define the tensor product by a line bundle as follows.

Definition 6.7. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric vector bundle on tropical toric variety $\text{trop}(X_\Sigma)$. A tropical line bundle \mathcal{L} is determined by a sequence of integers a_1, \dots, a_s (see Example 5.3 (1)). We define the tensor product $\mathcal{E} \otimes \mathcal{L}$ by the data $(\mathcal{M}, \mathcal{G}, \{E^i(j - a_i)\})$: for the ray ρ_i the j th piece of the filtration for $\mathcal{E} \otimes \mathcal{L}$ is the $(j - a_i)$ th piece $E^i(j - a_i)$ of the filtration for \mathcal{E} .

We now define the notion of isomorphism of tropical toric vector bundle motivated by the following result on toric vector bundles.

Proposition 6.8. [Kly89, Corollary 1.2.4] *Let \mathcal{E} and \mathcal{F} be indecomposable toric vector bundles on a complete toric variety X_Σ . If \mathcal{E} and \mathcal{F} are isomorphic as ordinary bundles, then \mathcal{E} is equivariantly isomorphic to $\mathcal{F} \otimes \text{div}(\chi^m)$ for some character $m \in M$.*

Note that in view of Lemma 6.2 it suffices to define isomorphism of tropical toric vector bundle for indecomposable bundles.

Definition 6.9. Two indecomposable tropical toric vector bundles $\mathcal{E}_1 = (\mathcal{M}_1, \mathcal{G}_1, \{E_1^i(j)\})$ and $\mathcal{E}_2 = (\mathcal{M}_2, \mathcal{G}_2, \{E_2^i(j)\})$ on a tropical toric variety $\text{trop}(X_\Sigma)$ are said to be isomorphic if the valuated matroids $(\mathcal{M}_1, \mathcal{G}_1)$ and $(\mathcal{M}_2, \mathcal{G}_2)$ are isomorphic and under the identification of the lattice of flats $\mathcal{L}(\underline{\mathcal{M}}_1)$ and $\mathcal{L}(\underline{\mathcal{M}}_2)$, we have

$$E_1^i(j) = E_2^i(j + \mathbf{u} \cdot \mathbf{v}_i)$$

for some $\mathbf{u} \in M$ and for all i and j .

Definition 6.10. The Picard group of a tropical toric variety $\text{trop}(X_\Sigma)$, denoted by $\text{Pic}(\text{trop}(X_\Sigma))$, is the set of isomorphism classes of tropical toric line bundles under tensor product. Any tropical line bundle on a tropical toric variety $\text{trop}(X_\Sigma)$ is realizable (see Definition 5.8) as any rank 1 matroid is realizable. Hence, we have $\text{Pic}(X_\Sigma) = \text{Pic}(\text{trop}(X_\Sigma))$, as expected from [JMT19].

Example 6.11. Consider the toric variety \mathbb{P}^2 . Any tropical toric line bundle \mathcal{L} is determined by a vector $(a_0, a_1, a_2) \in \mathbb{Z}^3$. Let $\mathbf{u} = (-a_1, -a_2) \in M = \mathbb{Z}^2$. Then \mathcal{L} is isomorphic to the vector $(a_0 + a_1 + a_2, 0, 0)$ under the isomorphism defined in Definition 6.9. Hence, the group $\text{Pic}(\text{trop}(\mathbb{P}^2))$ is isomorphic to \mathbb{Z} . \diamond

Remark 6.12. There is not a simple definition of the tensor product of tropical toric vector bundles. This is because there is not a tensor product of matroids in general. In [LV81] Las Vergnas showed that, given two matroids $\mathcal{M}_1, \mathcal{M}_2$, there is in general no definition of a matroid with ground set the product $\mathcal{G}_1 \times \mathcal{G}_2$ of the two ground sets, rank the product of the ranks, the product of two flats a flat, and the restriction of the matroid to sets of the form $\{x\} \times \mathcal{M}_2$ or $\mathcal{M}_1 \times \{y\}$ is a flat. These are all natural requirements for the tensor product of matroids, and thus for the tensor product of tropical toric vector bundles, so any definition of “the” or “a” tensor product needs to relax one of them.

When \mathcal{E} and \mathcal{F} are toric vector bundles on a toric variety X_Σ , so define realizable tropical vector bundles, the choices of realizations of matroids for \mathcal{E} and \mathcal{F} give rise to a matroid for $\text{trop}(\mathcal{E} \otimes \mathcal{F})$. However this matroid may depend on the choice of realization of the two matroids. For example, suppose that the matroid of both \mathcal{E} and \mathcal{F} is the uniform matroid $U(2, 4)$, with realizations

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & a \end{pmatrix} \text{ and } \begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 1 & 1 & b \end{pmatrix}.$$

We write $\mathbf{v}_1, \dots, \mathbf{v}_4$ for the columns of the first matrix, and $\mathbf{v}'_1, \dots, \mathbf{v}'_4$ for the columns of the second matrix. The tensor product of these two vector configurations defines a matroid on ground set $\{\mathbf{v}_i \otimes \mathbf{v}'_j : 1 \leq i, j \leq 4\}$ of size 16. The subconfiguration $\{\mathbf{v}_1 \otimes \mathbf{v}'_1, \mathbf{v}_2 \otimes \mathbf{v}'_2, \mathbf{v}_3 \otimes \mathbf{v}'_3, \mathbf{v}_4 \otimes \mathbf{v}'_4\}$ has matrix

$$\begin{pmatrix} 1 & 0 & 1 & 1 \\ 0 & 0 & 1 & b \\ 0 & 0 & 1 & a \\ 0 & 1 & 1 & ab \end{pmatrix},$$

which has determinant $a - b$. Thus this subset is a basis for the matroid of $\text{trop}(\mathcal{E} \otimes \mathcal{F})$ if and only if $a \neq b$, so the tropicalization of the tensor product depends on the choice of realization.

In this case there is a generic choice for the matroid, which is the matroid of the tensor product when none of $\{a - b, ab - 1, ab - a - b, ab - a + 1, ab - b + 1, a + b - 1\}$ are zero. However this is a consequence of the realization space of the matroid $U(2, 4)$ being irreducible. When the realization spaces of the matroids for \mathcal{E} and \mathcal{F} are reducible there may be no choice of generic matroid for the tensor product.

6.3. Parliaments of polytopes. In [DJS] Di Rocco, Jabbusch, and Smith associate a collection of polytopes to a toric vector bundle that encode the global sections of the bundle. We now give a tropical formulation of this, which lets us *define* $h^0(\mathcal{E})$ for a tropical toric vector bundle. For the rest of this section we assume that X_Σ is smooth and complete.

Recall that for a tropical toric vector bundle \mathcal{E} with associated filtrations $\{E^i(j)\}$ and \mathbf{w} in the ground set of $\underline{\mathcal{M}}(\mathcal{E})$ we have $\mathbf{d}_{\mathbf{w}} \in \mathbb{Z}^{|\Sigma(1)|}$ given by $(\mathbf{d}_{\mathbf{w}})_i = \max\{j : \mathbf{w} \in E^i(j)\}$.

Definition 6.13. Let \mathcal{E} be a tropical toric vector bundle. For each \mathbf{w} in the ground set \mathcal{G} of $\underline{\mathcal{M}}(\mathcal{E})$ we set

$$P_{\mathbf{w}} = \{\mathbf{u} \in M_{\mathbb{R}} : \mathbf{u} \cdot \mathbf{v}_i \leq (\mathbf{d}_{\mathbf{w}})_i \text{ for all } 1 \leq i \leq s\}.$$


We call $\{P_{\mathbf{w}} : \mathbf{w} \in \mathcal{G}\}$ the *parliament of polytopes* associated to \mathcal{E} . For $\mathbf{u} \in M$, we define

$$(6.1) \quad h^0(\mathcal{E})_{\mathbf{u}} = \text{rk} \left(\bigvee_{\mathbf{w} \in P_{\mathbf{w}}} \mathbf{w} \right),$$

and

$$h^0(\mathcal{E}) = \sum_{\mathbf{u} \in M} h^0(\mathcal{E})_{\mathbf{u}}.$$

FIGURE 9. The parliament of polytopes for Example 6.14(1)



Note that the assumption that Σ is complete implies that each $P_{\mathbf{w}}$ is bounded, so a polytope. Thus $h^0(\mathcal{E})_{\mathbf{u}}$ is zero for all but finitely many \mathbf{u} , and so the last sum is finite.

Example 6.14. We consider the tropicalization of the toric vector bundle \mathcal{E} of Example 3.6.

- (1) For the matroid with ground set $\{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$ in Example 3.6 (1), we have

$$\mathbf{d}_{\mathbf{e}_1+\mathbf{e}_2} = \mathbf{d}_{\mathbf{e}_2} = (1, 0, 0), \mathbf{d}_{\mathbf{e}_1+\mathbf{e}_3} = (0, 1, 0) \text{ and } \mathbf{d}_{\mathbf{e}_3} = (0, 0, 1).$$

The parliament of polytopes is

$$\begin{aligned} P_{\mathbf{e}_1+\mathbf{e}_2} &= P_{\mathbf{e}_2} = \{(u_1, u_2) \in M_{\mathbb{R}} : -u_1 - u_2 \leq 1, u_1 \leq 0, u_2 \leq 0\}, \\ P_{\mathbf{e}_1+\mathbf{e}_3} &= \{(u_1, u_2) \in M_{\mathbb{R}} : -u_1 - u_2 \leq 0, u_1 \leq 1, u_2 \leq 0\} \text{ and} \\ P_{\mathbf{e}_3} &= \{(u_1, u_2) \in M_{\mathbb{R}} : -u_1 - u_2 \leq 0, u_1 \leq 0, u_2 \leq 1\}. \end{aligned}$$

These are shown in Figure 9.

Note that

$$\begin{aligned} h^0(\text{trop}(\mathcal{E})) &= h^0(\text{trop}(\mathcal{E}))_{(0,0)} + h^0(\text{trop}(\mathcal{E}))_{(1,0)} + h^0(\text{trop}(\mathcal{E}))_{(1,-1)} + h^0(\text{trop}(\mathcal{E}))_{(0,1)} \\ &\quad + h^0(\text{trop}(\mathcal{E}))_{(-1,1)} + h^0(\text{trop}(\mathcal{E}))_{(-1,0)} + h^0(\text{trop}(\mathcal{E}))_{(0,-1)} \\ &= 3 + 1 + 1 + 1 + 1 + 2 + 2 = 11. \end{aligned}$$

- (2) For the matroid with ground set $\{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$ in Example 3.6 (2), we have $\mathbf{d}_{\mathbf{e}_1} = (1, 0, 0)$, so $P_{\mathbf{e}_1} = P_{\mathbf{e}_2}$, $P_{\mathbf{e}_1+\mathbf{e}_3}$ and $P_{\mathbf{e}_3}$ are as above. Thus $h^0(\text{trop}(\mathcal{E}))$ is again 11.
- (3) For the matroid with ground set $\{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_1 - \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3, \mathbf{e}_1 + \mathbf{e}_2 + \mathbf{e}_3\}$ in Example 3.6 (3), $P_{\mathbf{e}_1-\mathbf{e}_2} = P_{\mathbf{e}_2} = P_{\mathbf{e}_1+\mathbf{e}_2}$. We have $\mathbf{d}_{\mathbf{e}_1+\mathbf{e}_2+\mathbf{e}_3} = (0, 0, 0)$, so

$$P_{\mathbf{e}_1+\mathbf{e}_2+\mathbf{e}_3} = \{(u_1, u_2) \in M_{\mathbb{R}} : -u_1 - u_2 \leq 0, u_1 \leq 0, u_2 \leq 0\} = \{\mathbf{0}\}.$$

We then get

$$\begin{aligned} h^0(\text{trop}(\mathcal{E})) &= h^0(\text{trop}(\mathcal{E}))_{(0,0)} + h^0(\text{trop}(\mathcal{E}))_{(1,0)} + h^0(\text{trop}(\mathcal{E}))_{(1,-1)} + h^0(\text{trop}(\mathcal{E}))_{(0,1)} \\ &\quad + h^0(\text{trop}(\mathcal{E}))_{(-1,1)} + h^0(\text{trop}(\mathcal{E}))_{(-1,0)} + h^0(\text{trop}(\mathcal{E}))_{(0,-1)} \\ &= 3 + 1 + 1 + 1 + 1 + 2 + 2 = 11. \end{aligned}$$

Note that in this example dimension of the space of global sections does not depend on the choice of the matroid. The following lemma shows that this is true in general. \diamond

Lemma 6.15. *When $\mathcal{E} = \text{trop}(\mathcal{F})$ for a toric vector bundle \mathcal{F} on X_{Σ} we have*

$$h^0(\mathcal{E})_{\mathbf{u}} = h^0(\mathcal{F})_{\mathbf{u}} \text{ for } \mathbf{u} \in M \text{ and thus } h^0(\mathcal{E}) = h^0(\mathcal{F}).$$

Proof. The proof of Proposition 1.1 of [DJS] gives an isomorphism $H^0(X, \mathcal{E})_{\mathbf{u}} \cong \bigcap_{i=1}^s E^i(\mathbf{u} \cdot \mathbf{v}_i)$, essentially following [Kly89, Corollary 4.13]. Since a spanning set of $\bigcap_{i=1}^s E^i(\mathbf{u} \cdot \mathbf{v}_i)$ belongs to the lattice of flats of the underlying matroid of $\mathcal{E} = \text{trop}(\mathcal{F})$ (see Definition 3.5), the lemma follows from the proof of [DJS, Proposition 1.1]. \square

Remark 6.16. A version of the formula (6.1) is found in the work of Kaveh and Manon connecting toric vector bundles with piecewise linear valuations; see [KM23, Proposition 3.12]. Note that one corollary of Lemma 6.15 is that the dimension of $h^0(\mathcal{E})$ for a toric vector bundle as given in [DJS, Proposition 1.1] does not depend on the choice of matroid of \mathcal{E} .

7. STABILITY

In this section we give a definition of (semi)stability for a tropical toric reflexive sheaf, and construct Jordan-Hölder and Harder-Narasimhan filtrations. The concept of modularity from lattice theory plays an essential role here; see §2.4. Throughout this section we assume that the fan Σ defines a *projective* toric variety. For convenience we also assume that Σ is simplicial, though this is only needed for defining the polarization (7.3).

7.1. Definition of stability. Recall that a reflexive sheaf \mathcal{E} on a nonsingular projective n -dimensional variety with distinguished very ample divisor H is *stable* if the slope

$$\mu(\mathcal{F}) = \frac{c_1(\mathcal{F}) \cdot H^{n-1}}{\text{rk}(\mathcal{F})}$$

of a torsion-free subsheaf \mathcal{F} of \mathcal{E} with $0 < \text{rk}(\mathcal{F}) < \text{rk}(\mathcal{E})$ and first Chern class $c_1(\mathcal{F})$ satisfies

$$\mu(\mathcal{F}) < \mu(\mathcal{E}).$$

The sheaf \mathcal{E} is *semistable* if the condition holds with $<$ replaced by \leq .

Note that for a toric variety it suffices to consider only equivariant reflexive subsheaves of \mathcal{E} (see [DDK20, Remark 2.3.2]) to check (semi)stability. We now formulate a tropical version of this concept. This first requires the definition of a reflexive subsheaf of a tropical reflexive sheaf. We are motivated by the following proposition which establishes a correspondence between equivariant subsheaves and subspaces.

Proposition 7.1 ([DDK21, Corollary 0.0.2]). *Let \mathcal{E} be an equivariant reflexive sheaf on a toric variety X_{Σ} with associated filtrations $(E, \{E^i(j)\})$. Then equivariant saturated subsheaves of \mathcal{E} are in one-to-one correspondence with induced subfiltrations $(F, \{F^i(j)\})$ of $(E, \{E^i(j)\})$, where F is a subspace of E and $F^i(j) = E^i(j) \cap F$.*

See also [HNS22, Proposition 2.3].

Definition 7.2. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical reflexive sheaf on a tropical toric variety $\text{trop}(X_{\Sigma})$ as in Definition 5.1. We associate to a flat $F \in \mathcal{L}(\underline{\mathcal{M}})$ a tropical reflexive sheaf as follows. Let $\mathcal{M}|_F$ be the restriction of the valuated matroid \mathcal{M} (see §2.2). Note that the lattice of flats $\mathcal{L}(\underline{\mathcal{M}}|_F)$ is the interval $[\emptyset, F]$ in $\mathcal{L}(\underline{\mathcal{M}})$. For any ray $\rho_i \in \Sigma(1)$ and $j \in \mathbb{Z}$, set

$$F^i(j) = F \wedge E^i(j) \in \mathcal{L}(\underline{\mathcal{M}}).$$

The data $\mathcal{E}_F := (\mathcal{M}|_F, F, \{F^i(j)\})$ defines a tropical toric reflexive sheaf, which we call a reflexive subsheaf, or just a subsheaf, of \mathcal{E} .

Remark 7.3. In [GUZ22, Proposition 2.5], the authors showed that every subbundle \mathcal{F} of a vector bundle \mathcal{E} on a metric graph Γ is split: there exists another subbundle \mathcal{H} of \mathcal{E} such that $\mathcal{E} \cong \mathcal{F} \oplus \mathcal{H}$. This phenomenon does not occur for tropical toric vector bundles. Consider the Hirzebruch surface $X = \mathbb{F}_1$, whose fan has rays spanned by $\mathbf{v}_1 = (1, 0)$, $\mathbf{v}_2 = (0, 1)$, $\mathbf{v}_3 = (-1, 1)$ and $\mathbf{v}_4 = (0, -1)$. Consider the tangent bundle $\mathcal{E} = \mathcal{T}_X$ with filtrations given by $(E, \{E^i(j)\})$, where

$$E^i(j) = \begin{cases} E = \text{span}(\mathbf{v}_1, \mathbf{v}_2) & j \leq 0 \\ \text{span}(\mathbf{v}_i) & j = 1 \\ 0 & j > 1 \end{cases} .$$

Then \mathcal{E} has as realizable matroid the matroid $\underline{\mathcal{M}}(\mathcal{E})$ with ground set $\mathcal{G} = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. Note that we do not include \mathbf{v}_4 as $\text{span}(\mathbf{v}_4) = \text{span}(\mathbf{v}_2)$. Consider the flat $F = \text{span}(\mathbf{v}_2)$. Then, by [KD19, Proposition 4.1.1] F corresponds to a subbundle \mathcal{F} with associated filtrations $(F, \{F \cap E^i(j)\})$. If there is a subsheaf \mathcal{H} of \mathcal{E} such that $\mathcal{E} \cong \mathcal{F} \oplus \mathcal{H}$, we must have $H^1(1) = \text{span}(\mathbf{v}_1)$. From the direct sum we also need $H^3(1) = \text{span}(\mathbf{v}_3)$, which is not possible as \mathcal{H} has rank one. In [Dev22, Section 3.2], the author also constructed a similar example. Now tropicalizing, we get the tropical toric vector bundle $\text{trop}(\mathcal{E}) = (\underline{\mathcal{M}}(\mathcal{E}), \mathcal{E}, \{E^i(j)\})$ on $\text{trop}(X_\Sigma)$. The flat F corresponds to a tropical toric subsheaf $\mathcal{E}_F = \text{trop}(\mathcal{F})$, which is not split as the above argument suggests.

Recall that the first Chern class of a toric vector bundle \mathcal{E} associated with Klyachko data $(E, \{E^i(j)\})$ has the following combinatorial description (see [Kly89, Remark 3.2.4]):

$$(7.1) \quad c_1(\mathcal{E}) = \sum_{\mathbf{v}_i \in \Sigma(1), j \in \mathbb{Z}} j \dim \left(\frac{E^i(j)}{E^i(j+1)} \right) D_i,$$

We thus *define* the first Chern class of a tropical toric reflexive sheaf as follows.

Definition 7.4. Let \mathcal{E} be a tropical reflexive sheaf on a tropical toric variety $\text{trop}(X_\Sigma)$ given by the data $(\mathcal{M}, \mathcal{G}, \{E^i(j)\})$. Recall that we have the exact sequence (2.1)

$$0 \longrightarrow M \longrightarrow \mathbb{Z}^s \xrightarrow{\phi} A^1(\Sigma) \longrightarrow 0.$$

Let $\mathbf{c} \in \mathbb{Z}^s$ be the vector with i th component

$$(7.2) \quad c_i = \sum_{j \in \mathbb{Z}} j \left(\text{rk}(E^i(j)) - \text{rk}(E^i(j+1)) \right).$$

We define the *first Chern class* $c_1(\mathcal{E})$ as the image of \mathbf{c} in $A^1(\Sigma)$ under the map ϕ . We now define the degree of a tropical reflexive sheaf \mathcal{E} . We fix a polarization H , i.e., an ample line bundle on the toric variety X_Σ . This gives rise to a vector

$$(7.3) \quad \mathbf{h} := (h_1, \dots, h_s) \in \mathbb{N}^s, \text{ where } h_i = D_i \cdot H^{n-1}.$$

We define the degree as follows:

$$\text{deg}(\mathcal{E}) := \sum_{\rho_i \in \Sigma(1), j \in \mathbb{Z}} j \left(\text{rk}(E^i(j)) - \text{rk}(E^i(j+1)) \right) h_i \in \mathbb{N}.$$

We fix such a vector \mathbf{h} throughout the rest of the section, and all definitions implicitly depend on the choice of \mathbf{h} . We define the slope as the quotient

$$\mu(\mathcal{E}) := \frac{\text{deg}(\mathcal{E})}{\text{rank}(\mathcal{E})}.$$

Example 7.5. When \mathcal{E} is a rank-1 toric vector bundle, given by $\mathbf{a} \in \mathbb{Z}^s$ as in Example 5.3 (1), we have $c_1(\mathcal{E})$ equal to the image of \mathbf{a} in $A^1(\Sigma)$. \diamond

Remark 7.6. Let \mathcal{E} be a toric vector bundle on a toric variety X_Σ . Consider the tropicalization $\text{trop}(\mathcal{E})$ defined in Definition 5.8. Then $c_1(\mathcal{E}) = c_1(\text{trop}(\mathcal{E}))$, as the rank of the flat $E^i(j)$ in the tropicalization equals the dimension of the subspace that it corresponds to, so (7.1) agrees with (7.2). Since the definition of degree in both cases only depends on a class in $A^1(X_\Sigma) \cong A^1(\Sigma)$, we also have $\text{deg}(\mathcal{E}) = \text{deg}(\text{trop}(\mathcal{E}))$, and so $\mu(\mathcal{E}) = \mu(\text{trop}(\mathcal{E}))$. Note that this does not depend on the choice of matroid for \mathcal{E} , and thus on the precise tropicalization.

Definition 7.7. Let \mathcal{E} be a tropical reflexive sheaf on a tropical toric variety $\text{trop}(X_\Sigma)$ given by the data $(\mathcal{M}, \mathcal{G}, \{E^i(j)\})$. We say that \mathcal{E} is *stable* if

$$\mu(\mathcal{E}_F) < \mu(\mathcal{E})$$

for all proper nonempty flats F of the lattice of flats of the underlying matroid $\underline{\mathcal{M}}$. The sheaf \mathcal{E} is *semistable* if the condition holds with $<$ replaced by \leq .

Example 7.8. Let \mathcal{E} be a rank-1 tropical toric reflexive sheaf on a tropical toric variety $\text{trop}(X_\Sigma)$ given by the data $(\mathcal{M}, \mathcal{G}, \{E^i(j)\})$. Since the underlying matroid $\underline{\mathcal{M}}$ does not contain any proper flats, \mathcal{E} is vacuously stable. \diamond

Example 7.9. Let $X_\Sigma = \mathbb{P}^n$ be projective space, and consider the tropical tangent bundle $\text{trop}(\mathcal{T})$ defined in Example 5.3(3). The only j which contributes to the first Chern class computation is $j = 1$, for which $j(\text{rk}(E^i(j)) - \text{rk}(E^i(j+1))) = 1$, so the first Chern class $c_1(\text{trop}(\mathcal{T}))$ is the image of $(1, \dots, 1)$ in $A^1(\mathbb{P}^n) \cong \mathbb{Z}$, which is $1 \in \mathbb{Z}$. For the polarization $\mathbf{h} = (1, \dots, 1)$, corresponding to $\mathcal{O}(1)$ on \mathbb{P}^n , the degree is then $n + 1$, so the slope of $\text{trop}(\mathcal{T})$ is given by

$$\mu(\text{trop}(\mathcal{T})) = \frac{n+1}{n}.$$

Let F be a proper flat of rank $l < n$. Then $|F| = l$ as any proper subset of the ground set of the matroid of $\text{trop}(\mathcal{T})$ is linearly independent. Hence, the slope of the subsheaf corresponding to F is $\mu(\mathcal{T}_F) = 1 < \mu(\text{trop}(\mathcal{T}))$. Thus, $\text{trop}(\mathcal{T})$ is stable. Note that this is also true for the tangent bundle on (nontropical) projective space (see [OSS11, Theorem 1.3.2]). \diamond

Example 7.10. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric vector bundle on a tropical toric variety $\text{trop}(X_\Sigma)$ and let \mathcal{L} be a tropical line bundle, determined by $\mathbf{a} \in \mathbb{Z}^s$ as in Example 5.3(1). We have

$$\begin{aligned} \deg(\mathcal{E} \otimes \mathcal{L}) &= \sum_{\rho_i \in \Sigma, j \in \mathbb{Z}} j \left(\text{rk}(E^i(j - a_i)) - \text{rk}(E^i(j - a_i + 1)) \right) h_i \\ &= \sum_{\rho_i \in \Sigma} \left(\sum_{j \in \mathbb{Z}} (j - a_i) \left(\text{rk}(E^i(j - a_i)) - \text{rk}(E^i(j - a_i + 1)) \right) h_i \right. \\ &\quad \left. + a_i \sum_{j \in \mathbb{Z}} \left(\text{rk}(E^i(j - a_i)) - \text{rk}(E^i(j - a_i + 1)) \right) h_i \right) \\ &= \deg(\mathcal{E}) + \sum_{\rho_i \in \Sigma} a_i \sum_{j \in \mathbb{Z}} \left(\text{rk}(E^i(j - a_i)) - \text{rk}(E^i(j - a_i + 1)) \right) h_i \\ &= \deg(\mathcal{E}) + \sum_{\rho_i \in \Sigma} a_i \text{rk}(\mathcal{G}) h_i \\ &= \deg(\mathcal{E}) + \text{rk}(\mathcal{G}) \deg(\mathcal{L}). \end{aligned}$$

Thus we have $\mu(\mathcal{E} \otimes \mathcal{L}) = \mu(\mathcal{E}) + \mu(\mathcal{L})$. Hence, a tropical toric vector bundle \mathcal{E} is (semi)stable if and only if $\mathcal{E} \otimes \mathcal{L}$ is (semi)stable for any tropical line bundle \mathcal{L} . \diamond

Proposition 7.11. *Stable tropical toric vector bundles are indecomposable.*

Proof. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a stable tropical toric vector bundle of rank r . Suppose that \mathcal{E} is decomposable. Then $\mathcal{M} = \mathcal{M}_1 \oplus \mathcal{M}_2$, for valuated matroids \mathcal{M}_1 and \mathcal{M}_2 of ranks r_1, r_2 respectively on ground sets \mathcal{G}_1 and \mathcal{G}_2 with $\mathcal{G} = \mathcal{G}_1 \sqcup \mathcal{G}_2$. Note that their lattice of flats are related as follows:

$$\mathcal{L}(\mathcal{M}) = \mathcal{L}(\mathcal{M}_1) \times \mathcal{L}(\mathcal{M}_2).$$

Then, $E^i(j) = E_1^i(j) \times E_2^i(j)$, where $E_l^i(j) \in \mathcal{L}(\mathcal{M}_l)$ for $l = 1, 2$. So we can consider the tropical reflexive sheaves $\mathcal{E}_l = (\mathcal{M}_l, \mathcal{G}_l, \{E_l^i(j)\})$ for $l = 1, 2$. Note that we have

$$\text{rk}(E^i(j)) = \text{rk}(E_1^i(j)) + \text{rk}(E_2^i(j)).$$


Hence, from the definition of the first Chern class (Definition 7.4), we see that

$$c_1(\mathcal{E}) = c_1(\mathcal{E}_1) + c_1(\mathcal{E}_2).$$

This implies that

$$\deg(\mathcal{E}) = \deg(\mathcal{E}_1) + \deg(\mathcal{E}_2).$$

FIGURE 10



Without loss of generality we may assume that $\mu(\mathcal{E}_1) \leq \mu(\mathcal{E}_2)$. Then

$$\begin{aligned} \mu(\mathcal{E}) &= \frac{\deg(\mathcal{E}_1) + \deg(\mathcal{E}_2)}{r_1 + r_2} \\ &= \frac{r_1}{r} \frac{\deg(\mathcal{E}_1)}{r_1} + \frac{r_2}{r} \frac{\deg(\mathcal{E}_2)}{r_2} \\ &\leq \left(\frac{r_1}{r} + \frac{r_2}{r} \right) \mu(\mathcal{E}_2) = \mu(\mathcal{E}_2). \end{aligned}$$

Since $\underline{\mathcal{M}}_2 = \underline{\mathcal{M}}|_F$, where F is the flat $\emptyset \times \mathcal{G}_2 \in \mathcal{L}(\underline{\mathcal{M}})$, this contradicts the stability of \mathcal{E} . Thus \mathcal{E} must be indecomposable. \square

Remark 7.12. Note that any matroid $\underline{\mathcal{M}}$ can be written as a direct sum $\underline{\mathcal{M}} \cong \underline{\mathcal{M}}' \oplus (U_{0,1})^{\oplus l} \oplus (U_{1,1})^{\oplus c}$, where $\underline{\mathcal{M}}'$ has neither loops nor coloops (see [Kat16, Proposition 5.3]). Proposition 7.11 shows that a necessary condition for a tropical toric vector bundle to be stable is that the underlying matroid should, in addition to being loopfree as required in Definition 5.1, also be coloopfree.

Example 7.13. (Behavior of (semi)stability under tropicalization).

- (1) Example 7.9 shows that tangent bundle on \mathbb{P}^n remains stable under tropicalization.
- (2) Let \mathcal{E} be the toric vector bundle on \mathbb{P}^2 considered in Example 3.6. The slope is given by

$$\mu(\mathcal{E}) = \frac{2 + 1 + 1}{3} = 1 + \frac{1}{3}.$$

Consider the subspace $F = \text{span}(\mathbf{e}_1, \mathbf{e}_3) \subseteq E$. The corresponding subsheaf is given by the following filtrations:

$$F^0(j) = \begin{cases} F & j \leq 0 \\ \text{span}(\mathbf{e}_1) & j = 1 \\ 0 & j > 1 \end{cases}, E^1(j) = \begin{cases} F & j \leq 0 \\ \text{span}(\mathbf{e}_1 + \mathbf{e}_3) & j = 1 \\ 0 & j > 1 \end{cases}, E^2(j) = \begin{cases} F & j \leq 0 \\ \text{span}(\mathbf{e}_3) & j = 1 \\ 0 & j > 1 \end{cases}.$$

The slope is given by

$$\mu(\mathcal{F}) = \frac{1 + 1 + 1}{2} = 1 + \frac{1}{2}.$$

Thus \mathcal{E} is not semistable.

Now we consider the tropicalization from Example 3.6(1) and show that it is stable. Let \mathcal{M} be the rank-3 matroid on the ground set $\mathcal{G} = \{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$. The lattice of flats is given in Figure 10. The tropicalization is the tropical toric vector bundle $\text{trop}(\mathcal{E}) = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$, where the filtrations of the flats are given as follows:

$$E^0(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{e}_1 + \mathbf{e}_2, \mathbf{e}_2\} & j = 1 \\ \emptyset & j > 1 \end{cases}, E^1(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{e}_1 + \mathbf{e}_3\} & j = 1 \\ \emptyset & j > 1 \end{cases}, E^2(j) = \begin{cases} \mathcal{G} & j \leq 0 \\ \{\mathbf{e}_3\} & j = 1 \\ \emptyset & j > 1 \end{cases}.$$

The slope is given by

$$\mu(\text{trop}(\mathcal{E})) = 1 + \frac{1}{3}.$$

All the rank-1 and rank-2 flats have slope 1. Hence, $\text{trop}(\mathcal{E})$ is stable.

The issue here is that the destabilizing subsheaf corresponding to the subspace $F = \text{span}(\mathbf{e}_1, \mathbf{e}_3)$ does not tropicalize to a subsheaf of $\text{trop}(\mathcal{E})$. We have $F \cap \mathcal{G} = \{\mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$, which is a flat of $\mathcal{L}(\underline{\mathcal{M}})$, but the restriction of \mathcal{M} to F has lattice of flats $[\emptyset, F]$, which just has two elements of rank 1. However the subsheaf of \mathcal{E} determined by F has three one-dimensional flats: in addition to $\text{span}(\mathbf{e}_1 + \mathbf{e}_3)$ and $\text{span}(\mathbf{e}_3)$ in $[\emptyset, F]$, there is also $\text{span}(\mathbf{e}_1) = E^0(2) \cap F$. This is a consequence of the lattice of flats not being modular: we have $E^0(2) \wedge (E^1(3) \vee E^2(1)) = \emptyset$, and $E^0(2) \vee (E^1(3) \vee E^2(1)) = \mathcal{G}$, so the sum of these ranks is $0 + 3 = 3$, which is less than $\text{rk}(E^0(2)) + \text{rk}(E^1(3) \vee E^2(1)) = 2 + 2 = 4$.

- (3) If we choose the tropicalization of the toric vector bundle \mathcal{E} in Example 7.13(2) to be given by the matroid of Example 3.6(2), then we see $\text{trop}(\mathcal{E})$ is not semistable as well. Note that here modularity assumption is satisfied by Proposition 2.5(2). ◇

Lemma 7.14. *Let \mathcal{E} be a stable, or semistable, toric vector bundle. Then $\text{trop}(\mathcal{E})$ is stable, respectively semistable.*

Proof. Write $\mathcal{F} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ for $\text{trop}(\mathcal{E})$, where the ground set \mathcal{G} is a subset of the vector space E . If \mathcal{E} is stable then for every subsheaf \mathcal{E}_F induced by a subspace $F \subseteq E$ in the sense of Proposition 7.1 or [HNS22, Proposition 2.3] we have $\mu(\mathcal{E}_F) < \mu(\mathcal{E})$. In particular this is true for the subspaces spanned by subsets of \mathcal{G} , which correspond to flats of $\underline{\mathcal{M}}$.

By construction $c_1(\mathcal{F}) = c_1(\mathcal{E})$, so $\mu(\mathcal{F}) = \mu(\mathcal{E})$. For a flat of $\underline{\mathcal{M}}$ we write F for the subspace and $\tilde{F} \subseteq \mathcal{G}$ for the set; in particular we also use this notation to make a distinction between $E^i(j) \subseteq E$ and $\tilde{E}^i(j) \subseteq \mathcal{G}$. To show that \mathcal{F} is stable it suffices to show that $\mu(\mathcal{F}_{\tilde{F}}) \leq \mu(\text{trop}(\mathcal{E}_F))$, as that will show that slope of every subsheaf induced by a flat of $\underline{\mathcal{M}}$ is less than $\mu(\mathcal{F})$. Since the rank of the flat \tilde{F} equals the dimension of the subspace $F = \text{span}(\tilde{F})$, and $h_i \geq 0$ for all $1 \leq i \leq s$, to show this it suffices to show that $c_1(\mathcal{F}_{\tilde{F}})_i \leq c_1(\mathcal{E}_F)_i$ for all i . This follows, using [HNS22, Lemma 2.5], from the observation that $\text{rk}(\tilde{E}^i(j) \wedge \tilde{F}) \leq \dim(E^i(j) \cap F)$. □

Remark 7.15. The converse of Lemma 7.14 is not true: if \mathcal{E} is unstable, then $\text{trop}(\mathcal{E})$ might not be unstable. One example is given by Example 7.13(2), where the toric vector bundle \mathcal{E} is not stable, but the first tropicalization is.

There are two ways in which the tropicalization of an unstable toric vector bundle can fail to be unstable. The first, which is what occurs in Example 7.13(2), is that the subspace of E corresponding to a destabilizing subsheaf may not be a flat of $\underline{\mathcal{M}}$. The second is that the subspace may be a flat of $\underline{\mathcal{M}}$, but the inequality $\text{rk}(\tilde{E}^i(j) \wedge \tilde{F}) \leq \dim(E^i(j) \cap F)$ may be strict in a way that means that the slope of the $\text{trop}(\mathcal{E})_{\tilde{F}}$ is less than the slope of \mathcal{E}_F . This does not happen when $\tilde{E}^i(j)$ and \tilde{F} are a modular pair.

However, if \mathcal{E} is toric vector bundle that is not (semi)stable, it is always possible to choose a matroid for \mathcal{E} in such a way that the tropicalization $\text{trop}(\mathcal{E})$ is not (semi)stable. This is inspired by [Dev22, Theorem 2.18], though see Remark 7.16 for some differences. If \mathcal{E} is not stable, then by [BDGP21, Theorem 2.1] there is an equivariant destabilizing subsheaf \mathcal{F} . Such a sheaf is given by filtrations $\{F_{\mathbf{u}}^\sigma\}_{\sigma \in \Sigma, \mathbf{u} \in M}$ [Per03, Chapter 4]. By [Koo11, Corollary 3.18] the first Chern class of \mathcal{F} only depends on the $F_{\mathbf{u}}^{\rho_i}$ and the formula is the same as for the reflexive sheaf with filtrations $F^i(j) := F_{\mathbf{u}}^{\rho_i}$ when $j = \mathbf{u} \cdot \mathbf{v}_i$. Here we use that $F_{\mathbf{u}}^{\rho_i} = F_{\mathbf{u}'}^{\rho_i}$ if $\mathbf{u} \cdot \mathbf{v}_i = \mathbf{u}' \cdot \mathbf{v}_i$. Let \mathcal{F}' be this toric reflexive sheaf. By construction we have $\mu(\mathcal{F}) = \mu(\mathcal{F}')$, so \mathcal{F}' is also destabilizing. After passing to its saturation if necessary, we have $F^i(j) = E^i(j) \cap F$ for some subspace F of E . Given any matroid for \mathcal{E} , we can augment \mathcal{G} by adding a basis for each $F \cap E^i(j)$ to \mathcal{G} . This ensures that $\text{trop}(\mathcal{F}) = \text{trop}(\mathcal{E})_{\tilde{F}}$, where \tilde{F} is the flat of the matroid $\underline{\mathcal{M}}$ for \mathcal{E} corresponding to F . This means that $\mu(\text{trop}(\mathcal{E})_{\tilde{F}}) = \mu(\mathcal{F})$, so the flat \tilde{F} is destabilizing for $\text{trop}(\mathcal{E})$.

Remark 7.16. The first tropicalization in Example 7.13(2) shows that the whether a rank-3 toric vector bundle is stable can not always be detected from its tropicalization. This contrasts with [Dev22, Theorem 2.23], where Devey shows that (semi)stability of a toric vector bundle \mathcal{E} of rank at most 3 can be determined by considering flats of *any* choice of the associated matroid $\underline{\mathcal{M}}$. The distinction is that in [Dev22] the flat is identified with the subspace: then content of [Dev22, Theorem 2.23] is that if there is a destabilizing sheaf, it is \mathcal{E}_F for a subspace $F \subseteq E$ corresponding to a flat of the chosen matroid. However the stability of $\text{trop}(\mathcal{E})$ depends on $\text{trop}(\mathcal{E})_{\bar{F}}$, which may not equal $\text{trop}(\mathcal{E}_F)$.

7.2. Jordan-Hölder and Harder-Narasimhan filtrations. In this section we construct Jordan-Hölder and Harder-Narasimhan filtrations for tropical toric reflexive sheaves. This allows us to reduce the study of tropical toric vector bundles to (semi)stable ones. In usual algebraic geometry these justify the study of moduli spaces of stable vector bundles, and we hope that the same will be true in the tropical setting.

We defined the notion of a subsheaf of a tropical reflexive sheaf in Definition 7.2. We now formalise the notion of the quotient of a tropical reflexive sheaf. This uses the notion of the *quotient* of valuated matroids; see §2.2.

Definition 7.17. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric reflexive sheaf. A flat $F \in \mathcal{L}(\underline{\mathcal{M}})$ corresponds to a subsheaf $\mathcal{E}_F := (\mathcal{M}|_F, F, \{F^i(j)\})$ (see Definition 7.2). Consider the contraction of the valuated matroid \mathcal{M} by the flat F , denoted by \mathcal{M}/F , as recalled in §2.2. This is a matroid on the ground set $\mathcal{G} \setminus F$ with lattice of flats $\mathcal{L}(\underline{\mathcal{M}}/F)$ isomorphic to the interval $[F, \mathcal{G}]$ in $\mathcal{L}(\underline{\mathcal{M}})$. Note that $(\mathcal{M}/F, \mathcal{G} \setminus F, \{E^i(j)\})$ is a tropical toric reflexive sheaf, where

$$E^i(j) = (E^i(j) \vee F).$$

Here we are using the isomorphism $\mathcal{L}(\underline{\mathcal{M}}/F) \cong [F, \mathcal{G}]$ to regard $E^i(j)$ as a subset of $\mathcal{G} \setminus F$. We call this the quotient of \mathcal{E} by the subsheaf \mathcal{E}_F and denote it by \mathcal{E}/F . Note that

$$(7.4) \quad \begin{aligned} \text{rk}(\mathcal{M}/F) &= \text{rk}(\underline{\mathcal{M}}) - \text{rk}(F), \text{ and} \\ \text{rk}_{\mathcal{M}/F}((E^i(j) \vee F)) &= \text{rk}_{\underline{\mathcal{M}}}(E^i(j) \vee F) - \text{rk}_{\underline{\mathcal{M}}}(F) \end{aligned}$$

(see [Whi86, Proposition 7.4.3]). Here we have used the subscripts \mathcal{M} and \mathcal{M}/F to denote the corresponding rank functions. We usually omit the subscript for the rank function of \mathcal{M} . Let $F' \in \mathcal{L}(\underline{\mathcal{M}})$ be another flat containing the flat F . We have $F' \in \mathcal{L}(\underline{\mathcal{M}}|_{F'})$ and we call the contraction

$$\left((\mathcal{M}|_{F'})/F, F' \setminus F, \{((E^i(j) \wedge F') \vee F)\} \right),$$

the quotient of $\mathcal{E}_{F'}$ by \mathcal{E}_F . This is denoted by $\mathcal{E}_{F'/F}$. Note that $(\underline{\mathcal{M}}|_{F'})/F$ has rank $\text{rk}(F') - \text{rk}(F)$ and

$$(7.5) \quad \text{rk}_{\mathcal{M}_{F'/F}}(((E^i(j) \wedge F') \vee F)) = \text{rk}((E^i(j) \wedge F') \vee F) - \text{rk}(F).$$

Remark 7.18 (Comparison with realizable tropical vector bundles). The definition of matroid quotient is consistent with the realizable case. Let \mathcal{E} be a toric vector bundle with associated filtrations $(E, \{E^i(j)\})$ and let \mathcal{F} be a subsheaf corresponding to the subspace $F \subseteq E$. Then the filtrations associated to the quotient \mathcal{E}/\mathcal{F} are given by $\left(\frac{E}{F}, \left\{ \frac{E^i(j)+F}{F} \right\} \right)$.

Suppose $\underline{\mathcal{M}}$ is a realizable matroid for \mathcal{E} , with ground set $\mathcal{G} \subseteq E$, and F is the span of a subset of \mathcal{G} , so corresponds to a flat of $\underline{\mathcal{M}}$, which we will also call F . Then $E^i(j) + F$ corresponds to the flat $E^i(j) \vee F$. Quotienting E by F corresponds to contracting $\underline{\mathcal{M}}$ by F . The quotient $\underline{\mathcal{M}}/F$ has lattice of flats isomorphic to the interval $[F, \mathcal{G}]$ in $\mathcal{L}(\underline{\mathcal{M}})$. Thus $\{E^i(j) \vee F\}$ is the induced filtration on the quotient.

Remark 7.19. Using (7.4), we have

$$\begin{aligned} \deg(\mathcal{E}/F) &= \sum_{\rho_i \in \Sigma(1), j \in \mathbb{Z}} j \left(\text{rk}_{\mathcal{M}/F}(E^i(j)) - \text{rk}_{\mathcal{M}/F}(E^i(j+1)) \right) h_i \\ &= \sum_{\rho_i \in \Sigma(1), j \in \mathbb{Z}} j \left(\text{rk}(E^i(j) \vee F) - \text{rk}(E^i(j+1) \vee F) \right) h_i \end{aligned}$$

Note that the lattice of flats of $\underline{\mathcal{M}}' := (\underline{\mathcal{M}}|_{F'})/F$ is isomorphic to the interval $[F, F']$ in $\mathcal{L}(\underline{\mathcal{M}})$. Using (7.5), we have

$$\deg(\mathcal{E}_{F'/F}) = \sum_{\rho_i \in \Sigma(1), j \in \mathbb{Z}} j \left(\text{rk}((E^i(j) \wedge F') \vee F) - \text{rk}((E^i(j+1) \wedge F') \vee F) \right) h_i.$$

Lemma 7.20. *Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric reflexive sheaf of rank r . Let $F \in \mathcal{L}(\underline{\mathcal{M}})$ be a flat of rank r_1 such $(F, E^i(j))$ is a modular pair for all $1 \leq i \leq s$ and all $j \in \mathbb{Z}$, so the following condition holds*

$$(7.6) \quad \text{rk}(F) + \text{rk}(E^i(j)) = \text{rk}(E^i(j) \vee F) + \text{rk}(E^i(j) \wedge F)$$

for all $1 \leq i \leq s$ and $j \in \mathbb{Z}$. Let $r_2 = r - r_1$. Then we have

$$\mu(\mathcal{E}) = \frac{r_1}{r} \mu(\mathcal{E}_F) + \frac{r_2}{r} \mu(\mathcal{E}/F).$$

Proof. First we show that $\deg(\mathcal{E}) = \deg(\mathcal{E}_F) + \deg(\mathcal{E}/F)$. Consider the condition (7.6) for $j+1$,

$$(7.7) \quad \text{rk}(F) + \text{rk}(E^i(j+1)) = \text{rk}(E^i(j+1) \vee F) + \text{rk}(E^i(j+1) \wedge F).$$

From the equations (7.6) and (7.7), we get

$$\begin{aligned} \text{rk}(E^i(j)) - \text{rk}(E^i(j+1)) &= \left(\text{rk}(E^i(j) \vee F) - \text{rk}(E^i(j+1) \vee F) \right) \\ &\quad + \left(\text{rk}(E^i(j) \wedge F) - \text{rk}(E^i(j+1) \wedge F) \right). \end{aligned}$$

Thus, using Remark 7.19, we have $\deg(\mathcal{E}) = \deg(\mathcal{E}_F) + \deg(\mathcal{E}/F)$. Now consider

$$\begin{aligned} \mu(\mathcal{E}) &= \frac{\deg(\mathcal{E})}{r} = \frac{\deg(\mathcal{E}_F) + \deg(\mathcal{E}/F)}{r} \\ &= \frac{r_1}{r} \frac{\deg(\mathcal{E}_F)}{r_1} + \frac{r_2}{r} \frac{\deg(\mathcal{E}/F)}{r_2} \\ &= \frac{r_1}{r} \mu(\mathcal{E}_F) + \frac{r_2}{r} \mu(\mathcal{E}/F). \end{aligned}$$

□

The following immediate corollary, called the see-saw property, is extremely crucial:

Corollary 7.21. *Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric reflexive sheaf. Let $F \in \mathcal{L}(\underline{\mathcal{M}})$ be a flat satisfying condition (7.6). Then*

$$\begin{aligned} \mu(\mathcal{E}_F) < \mu(\mathcal{E}) &\iff \mu(\mathcal{E}) < \mu(\mathcal{E}/F), \text{ and} \\ \mu(\mathcal{E}_F) > \mu(\mathcal{E}) &\iff \mu(\mathcal{E}) > \mu(\mathcal{E}/F). \end{aligned}$$

Lemma 7.22. *Let \mathcal{M} be a valuated matroid of rank r on the ground set \mathcal{G} with basis valuation function ν . Let $F_1 \in \mathcal{L}(\underline{\mathcal{M}})$ be a flat of rank r_1 . Let $F_2 \in \mathcal{L}(\underline{\mathcal{M}})$ be another flat of rank r_2 with $F_1 \subsetneq F_2$. Then we have $(\mathcal{M}/F_1)/F_2 = \mathcal{M}/F_2$. Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric reflexive sheaf, then we have $(\mathcal{E}/F_1)/F_2 = \mathcal{E}/F_2$. Moreover, if $E^i(j)$ are modular flats we have $\mathcal{E}'_{F_2} = \mathcal{E}_{F_2/F_1}$, where $\mathcal{E}' = \mathcal{E}/F_1$.*

Proof. Recall from §2.2 that the contraction $\mathcal{M}' := \mathcal{M}/F_1$ is a valuated matroid of rank $r - r_1$ on the ground set $\mathcal{G} \setminus F_1$ with the basis valuation function given as follows:

$$\nu_{\mathcal{M}'} : \binom{\mathcal{G} \setminus F_1}{r - r_1} \longrightarrow \overline{\mathbb{R}}, \quad B \longmapsto \nu(B \cup A),$$

where $A \subseteq F_1$ is such that $|A| = \text{rk}_{\mathcal{M}}(A) = r_1$ and $B \subseteq \mathcal{G} \setminus F_1$. Note that $F_2 \in \mathcal{L}(\underline{\mathcal{M}}/F_1) = [F_1, \mathcal{G}]$ and $\text{rk}_{\mathcal{M}'}(F_2) = r_2 - r_1$. Then the contraction \mathcal{M}'/F_2 is a valuated matroid on the ground set $(\mathcal{G} \setminus F_1) \setminus F_2 = \mathcal{G} \setminus F_2$ of rank $r - r_1 - (r_2 - r_1) = r - r_2$. We can choose an independent subset $A' \subseteq \mathcal{G}$ of rank r_2 in \mathcal{M} satisfying $A \subsetneq A' \subseteq F_2$. Let $A'' := A' \setminus F_1 \subseteq F_2 \setminus F_1$. Then, we have

$\text{rk}_{\mathcal{M}'}(A'') = \text{rk}(A') - r_1 = r_2 - r_1$. Also, $|A''| = |A'| - |F_1| \leq r_2 - r_1$. Hence, A'' must be independent. Then the basis valuation function for \mathcal{M}'/F_2 is given as follows: for any $B \subseteq \mathcal{G} \setminus F_2$

$$\nu_{\mathcal{M}'/F_2}(B) = \nu_{\mathcal{M}'}(B \cup A'') = \nu(B \cup A'' \cup A) = \nu(B \cup A').$$

This shows that $\nu_{\mathcal{M}'/F_2}$ coincides with the basis valuation function for the matroid \mathcal{M}/F_2 . Hence, we have $(\mathcal{M}/F_1)/F_2 = \mathcal{M}/F_2$ as valuated matroids. Recall that the quotient \mathcal{E}/F_2 is given by $(\mathcal{M}/F_2, \mathcal{G} \setminus F_2, \{E^i(j) \vee F_2\})$. Note that $\mathcal{L}(\underline{\mathcal{M}}/F_2)$ is the interval $[F_2, \mathcal{G}]$ in $\mathcal{L}(\underline{\mathcal{M}})$ and we have

$$\text{rk}_{\underline{\mathcal{M}}/F_2}(E^i(j) \vee F_2) = \text{rk}(E^i(j) \vee F_2) - \text{rk}(F_2).$$

The tropical toric reflexive sheaf $(\mathcal{E}/F_1)/F_2$ is given by

$$\left(\mathcal{M}/F_2, \mathcal{G} \setminus F_2, \{E^i(j) \vee F_2\} \right),$$

as $F_1 \subsetneq F_2$. This shows that the tropical toric reflexive sheaves $(\mathcal{E}/F_1)/F_2$ and \mathcal{E}/F_2 are same.

To see the last sentence of the lemma, note that the tropical toric reflexive sheaves \mathcal{E}_{F_2/F_1} and \mathcal{E}'_{F_2} are given by the data

$$\begin{aligned} & \left((\mathcal{M}|_{F_2})/F_1, F_2 \setminus F_1, \{(E^i(j) \wedge F_2) \vee F_1\} \right) \text{ and} \\ & \left((\mathcal{M}/F_1)|_{F_2}, F_2 \setminus F_1, \{(E^i(j) \vee F_1) \wedge F_2\} \right), \end{aligned}$$

respectively. Since $E^i(j)$ are modular flats, we have

$$(E^i(j) \wedge F_2) \vee F_1 = (E^i(j) \vee F_1) \wedge F_2.$$

Hence, it suffices to check that both the matroids $(\mathcal{M}|_{F_2})/F_1$ and $(\mathcal{M}/F_1)|_{F_2}$ have the same basis valuation function. Choose

$$(7.8) \quad A \subseteq \mathcal{G} \setminus F_2 \text{ such that } \text{rk}(A \cup F_2) = r.$$

Then the basis valuation function for $\mathcal{M}|_{F_2}$ is given by

$$\nu_{F_2} : \begin{pmatrix} F_2 \\ r_2 \end{pmatrix} \longrightarrow \overline{\mathbb{R}}, B_2 \longmapsto \nu(B_2 \cup A),$$

for all $B_2 \subseteq F_2$. Note that $\mathcal{L}(\underline{\mathcal{M}}|_{F_2}) = [\emptyset, F_2]$ and $F_1 \in \mathcal{L}(\underline{\mathcal{M}}|_{F_2})$ is a flat of rank r_1 . Choose $A' \subseteq F_1$ independent with $\text{rk}(A') = r_1$. Then the contraction $(\mathcal{M}|_{F_2})/F_1$ is a valuated matroid on the ground set $F_2 \setminus F_1$ with the basis valuation function given by

$$(7.9) \quad \nu'' : \begin{pmatrix} F_2 \setminus F_1 \\ r_2 - r_1 \end{pmatrix} \longrightarrow \overline{\mathbb{R}}, B \longmapsto \nu_{F_2}(B \cup A') = \nu(B \cup A' \cup A).$$

The contraction $\mathcal{M}' := \mathcal{M}/F_1$ is a valuated matroid of rank $r - r_1$ on the ground set $\mathcal{G} \setminus F_1$ with the basis valuation function given as follows:

$$\nu_{\mathcal{M}'} : \begin{pmatrix} \mathcal{G} \setminus F_1 \\ r - r_1 \end{pmatrix} \longrightarrow \overline{\mathbb{R}}, B_1 \longmapsto \nu(B_1 \cup A'),$$

where $B_1 \subseteq F_1$. Note that using (7.8), we have

$$\text{rk}_{\mathcal{M}'}(A \cup F_2) = \text{rk}(A \cup F_2) - r_1 = r - r_1 = \text{rk}(\mathcal{M}').$$

Now the restriction $\mathcal{M}'|_{F_2}$ is a valuated matroid on the ground set $F_2 \setminus F_1$ with basis valuation function given by

$$(7.10) \quad \nu_{\mathcal{M}'|_{F_2}} : \begin{pmatrix} F_2 \setminus F_1 \\ r_2 - r_1 \end{pmatrix} \longrightarrow \overline{\mathbb{R}}, \quad \nu_{\mathcal{M}'|_{F_2}}(B) = \nu_{\mathcal{M}'}(B \cup A) = \nu(B \cup A \cup A'),$$

where $B \subseteq F_2 \setminus F_1$. Thus, from (7.9) and (7.10) we get the valuated matroids $(\mathcal{M}|_{F_2})/F_1$ and $(\mathcal{M}/F_1)|_{F_2}$ are same. \square

Theorem 7.23 (Jordan-Hölder filtration). *Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a semistable tropical toric vector bundle on tropical toric variety $\text{trop}(X_\Sigma)$. Assume that the $E^i(j)$ are modular flats of $\mathcal{L}(\underline{\mathcal{M}})$. Then, there is an increasing filtration by flats of the underlying matroid $\underline{\mathcal{M}}$*

$$\emptyset = F_0 \subsetneq F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_k = \mathcal{G}$$

for which the quotients $gr_i := \mathcal{E}_{F_i/F_{i-1}}$ are stable with slope $\mu(\mathcal{E})$ for $1 \leq i \leq k$.

Proof. If \mathcal{E} is stable, we set $F_1 = \mathcal{G}$ and we are done. Suppose that \mathcal{E} is not stable. Since \mathcal{E} is semistable there is a flat $G_1 \in \mathcal{L}(\underline{\mathcal{M}})$ with $\mu(\mathcal{E}) = \mu(\mathcal{E}_{G_1})$. If \mathcal{E}_{G_1} is not stable we can choose a flat $G_2 \in \mathcal{L}(\underline{\mathcal{M}})$ with $G_2 \subsetneq G_1$ and $\mu(\mathcal{E}_{G_2}) = \mu(\mathcal{E}_{G_1})$. Since the ranks of these flats are decreasing, this cannot continue forever, so we must have a nonempty flat $F_1 \in \mathcal{L}(\underline{\mathcal{M}})$ with \mathcal{E}_{F_1} stable and $\mu(\mathcal{E}) = \mu(\mathcal{E}_{F_1})$. Now we show that the quotient \mathcal{E}/F_1 is semistable with slope μ . That the slope of \mathcal{E}/F_1 is μ follows from Lemma 7.20. Suppose \mathcal{E}/F_1 is not semistable. Then using Corollary 7.21, we have a flat $G \in [F_1, \mathcal{G}]$ such that $\mu(\mathcal{E}/F_1) > \mu((\mathcal{E}/F_1)/G)$. Thus we have $\mu(\mathcal{E}) > \mu(\mathcal{E}/G)$ using Lemma 7.22. Corollary 7.21 then contradicts that \mathcal{E} is semistable, so we conclude that \mathcal{E}/F_1 is semistable. We denote by \mathcal{E}' the quotient \mathcal{E}/F_1 and repeat the same procedure. Let $F_2 \in \mathcal{L}(\underline{\mathcal{M}}/F_1)$ be a flat such that \mathcal{E}'_{F_2} is stable with slope μ . Then we have $\mathcal{E}'_{F_2} = \mathcal{E}_{F_2/F_1}$ by Lemma 7.22. Hence, we get $\emptyset \subsetneq F_1 \subsetneq F_2$ with \mathcal{E}_{F_1} and \mathcal{E}_{F_2/F_1} are both stable with same slope μ . Since $\mathcal{E}' = \mathcal{E}/F_1$ is semistable and \mathcal{E}'_{F_2} is stable, we have \mathcal{E}'/F_2 is again semistable, using Remark 2.6(2). Note that $\mathcal{E}'/F_2 = \mathcal{E}/F_2$ by Lemma 7.22. Hence, we can repeat the procedure together with Remark 2.6 to get the desired filtration. \square

Lemma 7.24. *Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric reflexive sheaf on tropical toric variety $\text{trop}(X_\Sigma)$. Assume that $E^i(j)$ are modular flats of the lattice of flats of $\underline{\mathcal{M}}$. Let $F, G \in \mathcal{L}(\underline{\mathcal{M}})$ be flats such that \mathcal{E}/F is semistable. Assume further that one of the flats F or G is also modular. Then, we have*

$$\mu(\mathcal{E}_{G/G \wedge F}) \leq \mu(\mathcal{E}/F).$$

Proof. Recall that the tropical toric reflexive sheaf $\mathcal{E}_{G/G \wedge F}$ is given by

$$\left((\mathcal{M}|_G)/(G \wedge F), G \setminus (G \wedge F), \{(E^i(j) \wedge G) \vee (G \wedge F)\} \right).$$

Note that using the modularity of the flat $E^i(j)$ (see Proposition 2.4(2)), we have

$$(7.11) \quad E^i(j) := (G \wedge F) \vee (E^i(j) \wedge G) = ((G \wedge F) \vee E^i(j)) \wedge G.$$

Let $A = F \vee G \in [F, \mathcal{G}] = \mathcal{L}(\underline{\mathcal{M}}/F)$. The tropical reflexive subsheaf of \mathcal{E}/F corresponding to A is given by

$$\left((\mathcal{M}/F)|_A, A \setminus F, \{(E^i(j) \vee F) \wedge A\} \right).$$

We have $E^i(j) \leq (E^i(j) \vee F) \wedge A$ and under the additional assumption of modularity, we get

$$\begin{aligned} \text{rk}((\mathcal{M}|_G)/(G \wedge F)) &= \text{rk}(G) - \text{rk}(G \wedge F) \\ &= \text{rk}(F \vee G) - \text{rk}(F) = \text{rk}((\mathcal{M}/F)|_A). \end{aligned}$$

Hence, using [HNS22, Lemma 2.5] we have $\mu(\mathcal{E}_{G/G \wedge F}) \leq \mu((\mathcal{E}/F)_A)$. Finally, by semistability of \mathcal{E}/F we get $\mu((\mathcal{E}/F)_A) \leq \mu(\mathcal{E}/F)$. Hence, the lemma follows. \square

Theorem 7.25 (Harder-Narasimhan filtration). *Let $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$ be a tropical toric reflexive sheaf on tropical toric variety $\text{trop}(X_\Sigma)$. Assume that the $E^i(j)$ are modular flats of the lattice of flats of $\underline{\mathcal{M}}$. Then there is an increasing filtration by flats of the underlying matroid $\underline{\mathcal{M}}$*

$$\emptyset = F_0 \subsetneq F_1 \subsetneq F_2 \subsetneq \dots \subsetneq F_k = \mathcal{G}$$

where the quotients $gr_i := \mathcal{E}_{F_i/F_{i-1}}$, $i = 1, \dots, k$ satisfies the following conditions:

- (i) the quotient gr_i is semistable, and
- (ii) $\mu(gr_i) > \mu(gr_{i+1})$ for $i = 1, \dots, k-1$.

This filtration is unique under the additional assumption that the flats F_1, \dots, F_k are also modular.

Proof. If \mathcal{E} is semistable, then we set $k = 1$ and $F_1 = \mathcal{G}$. Now suppose that \mathcal{E} is not semistable. Since there are only finitely many flats in the lattice of flats of $\underline{\mathcal{M}}$, we may choose a flat $F_1 \in \mathcal{L}(\underline{\mathcal{M}})$ of maximal slope and maximal rank amongst flats of maximal slope. Here by slope of a flat F_1 , we mean slope of \mathcal{E}_{F_1} . Then \mathcal{E}_{F_1} is semistable by the choice of the flat F_1 . Now consider the quotient $\mathcal{E}' := \mathcal{E}/F_1$ on the ground set $\mathcal{G} \setminus F_1$. As before choose a flat $F_2 \in \mathcal{L}(\underline{\mathcal{M}}/F_1)$ of maximum slope and maximum rank amongst flats of maximal slope of $\underline{\mathcal{M}}/F_1$. Note that, as before \mathcal{E}'_{F_2} is semistable by the choice of F_2 . Using Lemma 7.22, we get $\mathcal{E}'_{F_2} = \mathcal{E}_{F_2/F_1}$. Next we show that

$$\mu(\mathcal{E}_{F_1}) > \mu(\mathcal{E}_{F_2/F_1}).$$

Since $\text{rk}(F_2) > \text{rk}(F_1)$, by the choice of F_1 , we have $\mu(\mathcal{E}_{F_2}) < \mu(\mathcal{E}_{F_1})$. By Remark 2.6(1), $E^i(j) \vee F$ is a modular flat of \mathcal{M}/F , so by Corollary 7.21 and we get $\mu(\mathcal{E}_{F_2}) > \mu(\mathcal{E}_{F_2/F_1})$. Thus we get $\mu(\mathcal{E}_{F_1}) > \mu(\mathcal{E}_{F_2/F_1})$. To continue consider \mathcal{E}'/F_2 . Note that $\mathcal{E}'/F_2 = \mathcal{E}/F_2$ by Lemma 7.22. Hence, we can repeat the procedure to obtain the desired filtration.

To see the uniqueness, note that if $(F_i)_{i=1,\dots,k}$ is a filtration which satisfies the desired properties and if F' is another flat in $\mathcal{L}(\underline{\mathcal{M}})$ then this filtration induces a filtration $(F'_i = F_i \wedge F')_{i=1,\dots,k}$ of F' . Let $\text{gr}'_i = \mathcal{E}_{F'_i/F'_{i-1}}$ be the corresponding quotients. If they are all non-zero, by taking $\mathcal{E} = \mathcal{E}_{F_i}$, $F = F_{i-1}$ and $G = F' \wedge F_i$ in Lemma 7.24 together with Remark 2.6(1), we have

$$(7.12) \quad \mu(\text{gr}'_i) \leq \mu(\text{gr}_i) \leq \mu(\mathcal{E}_{F_1})$$

Here we have used the additional assumption that the flats F_i are also modular. Now applying Lemma 7.20 together with Remark 2.6(1), we get

$$\begin{aligned} \mu(\mathcal{E}_{F'_2}) &= \frac{r_1}{r'} \mu(\mathcal{E}_{F'_1}) + \frac{r_2}{r'} \mu(\mathcal{E}_{F'_2/F'_1}) \\ &\leq \frac{r_1}{r'} \mu(\mathcal{E}_{F_1}) + \frac{r_2}{r'} \mu(\mathcal{E}_{F_1}) \quad (\text{using (7.12) and semistability of } \mathcal{E}_{F_1}) \\ &= \mu(\mathcal{E}_{F_1}), \end{aligned}$$

here $r_1 = \text{rk}(F'_1)$, $r' = \text{rk}(F'_2)$ and $r_2 = r' - r_1$. Inducting on k , we get

$$\mu(\mathcal{E}_{F'}) = \mu(\mathcal{E}_{F'_k}) \leq \mu(\mathcal{E}_{F_1}).$$

Moreover, if the equality holds, it forces all gr'_i to be zero for all $i \geq 2$, in other words $F' \subseteq F_1$. Now if $(G_i)_{i=1,\dots,k}$ is another filtration with these properties then the slopes of \mathcal{E}_{F_1} and \mathcal{E}_{G_1} , we have $F_1 = G_1$. Now we continue applying the same argument to \mathcal{E}/F_1 . Note that we can apply the same argument in view of Remark 2.6. Hence, we conclude the filtration is unique. \square

Example 7.26. (1) Consider the tropical tangent sheaf $\text{trop}(\mathcal{T})$ on $\text{trop}(X_\Sigma)$ from Example 7.9, which is given by the representable matroid $\underline{\mathcal{M}}$ on the ground set $\mathcal{G} = \{\mathbf{v}_1, \dots, \mathbf{v}_s\}$. Note that by Proposition 2.5(2) \emptyset , \mathcal{G} and atoms in a lattice are always modular flats.


Hence using Theorem 7.25 a Harder-Narasimhan filtration of $\text{trop}(\mathcal{T})$ corresponds to a particular increasing sequence of flats of $\underline{\mathcal{M}}$. This can be thought of as a tropical analogue of [Pan15, Corollary 2.1.7]. which shows that a Harder-Narasimhan filtration of the tangent bundle on a toric variety corresponds to an increasing filtration of subspaces of $N_{\mathbb{R}}$.

- (2) Using Proposition 2.5(2), we get that any rank-2 tropical toric reflexive sheaf admits a Harder-Narasimhan filtration as described in Theorem 7.25.
- (3) Let $\mathcal{E} = (\underline{\mathcal{M}}, \mathcal{G}, \{E^i(j)\})$ be a tropical reflexive sheaf on tropical toric variety $\text{trop}(X_\Sigma)$. Suppose that there is a modular flat F of maximal slope such that the quotient is \mathcal{E}/F is semistable and $\mu(\mathcal{E}) < \mu(\mathcal{E}_F)$. Then $\emptyset \subsetneq F \subsetneq \mathcal{G}$ is a Harder-Narasimhan filtration. To see this it suffices to check

$$\mu(\mathcal{E}_F) > \mu(\mathcal{E}/F).$$

This follows from Corollary 7.21. Note that this case does not require the $E^i(j)$ to be modular flats. \diamond

FIGURE 11



Example 7.27. Consider the representable matroid \mathcal{M} on the ground set $\mathcal{G} = \{\mathbf{e}_1, \mathbf{e}_2, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$ introduced in Example 3.6 (2). The lattice of flats is given in Figure 11. Consider the tropical toric vector bundle $\mathcal{E} = (\mathcal{M}, \mathcal{G}, \{E^i(j)\})$, where the filtrations of the flats are given as follows:

$$E^0(j) = \begin{cases} \mathcal{G} & i \leq 1 \\ \{\mathbf{e}_1, \mathbf{e}_2\} & j = 2 \\ \emptyset & j > 2 \end{cases}, E^1(j) = \begin{cases} \mathcal{G} & i \leq 2 \\ \{\mathbf{e}_1 + \mathbf{e}_3\} & j = 3 \\ \emptyset & j > 3 \end{cases}, E^2(j) = \begin{cases} \mathcal{G} & i \leq 0 \\ \{\mathbf{e}_3\} & j = 1 \\ \emptyset & j > 1 \end{cases}.$$

The slope is given by

$$\mu(\mathcal{E}) = \frac{(1+4) + (4+3) + 1}{3} = \frac{13}{3}.$$

Note that $F = \{\mathbf{e}_1, \mathbf{e}_1 + \mathbf{e}_3, \mathbf{e}_3\}$ is the unique flat with maximum slope and maximum rank. Also, the flat F is modular and the quotient \mathcal{E}/F is rank 1 so stable. Hence, by Example 7.26 (3) we get that $\emptyset \subsetneq F \subsetneq \mathcal{G}$ is the Harder-Narasimhan filtration for \mathcal{E} . \diamond

Example 7.28. Consider the tropical toric sheaf $(\mathcal{E}, \mathcal{G}, \{E^i(j)\})$ from Remark 5.7. Note that here the $E^i(j)$ are not all modular flats. The slope of the bundle is given by

$$\mu(\mathcal{E}) = \frac{2+2}{4} = 1.$$

Note that the rank-1 flats are singleton subsets of \mathcal{G} . For the flats $\{1\}, \{2\}, \{3\}$ and $\{4\}$ the slope of the corresponding tropical toric subsheaf is 1. All other rank-1 flats have slope 0. The rank-2 flats are subsets of \mathcal{G} with cardinality 2 as there are no circuits with cardinality 3. All of them have slope less than or equal to 1.

The rank-3 flats are $\{1, 2, 3, 4\}, \{1, 4, 5, 6\}, \{1, 4, 7, 8\}, \{2, 3, 5, 6\}$ and $\{2, 3, 7, 8\}$ and all subsets of cardinality 3 of \mathcal{G} not contained in any of the previous 4-element flats. Consider the flat $F = \{1, 2, 3, 4\}$ and the corresponding filtrations are given by

$$F^0(j) = \begin{cases} F & j \leq 0 \\ \{1, 2\} & j = 1 \\ \emptyset & j > 1 \end{cases}, F^1(j) = \begin{cases} F & j \leq 0 \\ \{3, 4\} & j = 1 \\ \emptyset & j > 1 \end{cases}.$$

The slope is given by

$$\mu(\mathcal{E}_F) = \frac{2+2}{3} = 1 + \frac{1}{3} > \mu(\mathcal{E}).$$

Hence, the bundle \mathcal{E} is not semistable. Note that F is the unique flat with maximum slope and maximum rank. Note that F is not a modular flat. Consider the quotient $\mathcal{E}/F = (\mathcal{M}/F, \{5, 6, 7, 8\}, \{E^i(j) \vee F\})$, where

$$E^0(j) \vee F = \begin{cases} \mathcal{G} & j \leq 0 \\ \{1, 2, 3, 4\} & j > 0 \end{cases}, E^1(j) \vee F = \begin{cases} \mathcal{G} & j \leq 0 \\ \{1, 2, 3, 4\} & j > 0 \end{cases}.$$

Thus the slope $\mu(\mathcal{E}/F) = 0 < \mu(\mathcal{E}_F)$. Since the quotient \mathcal{E}/F is rank 1, it is stable. Hence, the Harder-Narasimhan filtration is given by $\emptyset \subsetneq F \subsetneq \mathcal{G}$. This Harder-Narasimhan filtration is unique, even

though $E^1(1)$ and $E^2(1)$ are not modular flats. Since there are only finitely many possibilities for a filtration, this can be checked directly. \diamond

8. TROPICAL VECTOR BUNDLES ON OTHER VARIETIES

In this section we extend the definition of tropical vector bundles from bundles on tropical toric varieties to more general bases. A subscheme X of a tropical toric variety $\text{trop}(X_\Sigma)$ is given by a tropical ideal in the sense of [MR18]. Given a tropical toric vector bundle on $\text{trop}(X_\Sigma)$ we can restrict it to X . Algebraically, this means replacing the Cox semiring S in (5.5) by the quotient of S by the bend locus of the ideal for X .

We first recall the definition of a subscheme of a tropical toric variety.

Definition 8.1. [MR18, §4] Let Σ be a rational polyhedral fan in $N_{\mathbb{R}}$. Let σ be a cone of Σ . An ideal $J \subseteq \overline{\mathbb{R}}[\sigma^\vee \cap M]$ is a *tropical ideal* if for every finite set E of monomials in $\overline{\mathbb{R}}[\sigma^\vee \cap M]$ the set of polynomials with support in E is the collection of vectors of a valuated matroid with ground set E . An ideal $I \subseteq \text{Cox}(\text{trop}(X_\Sigma)) = \overline{\mathbb{R}}[x_1, \dots, x_s]$ that is homogeneous with respect to the $A^1(\Sigma)$ -grading is a *locally tropical ideal* if for every $\sigma \in \Sigma$ the ideal $(I \text{Cox}(\text{trop}(X_\Sigma))_{\mathbf{x}^\sigma})_{\mathbf{0}} \subseteq \overline{\mathbb{R}}[\sigma^\vee \cap M]$ is a tropical ideal.

A subscheme X of a tropical toric variety X_Σ is determined by a locally tropical ideal $I \subseteq \text{Cox}(\text{trop}(X_\Sigma))$ as follows. On the local chart $\text{trop}(U_\sigma)$ the subscheme $X \cap \text{trop}(U_\sigma)$ has coordinate semiring

$$\overline{\mathbb{R}}[\sigma^\vee \cap M] / \mathcal{B}(I \text{Cox}(\text{trop}(X_\Sigma))_{\mathbf{x}^\sigma})_{\mathbf{0}},$$

where $\mathcal{B}()$ is the Giansiracusa bend congruence of §2.3.

Example 8.2. Let $X \subseteq \text{trop}(\mathbb{P}^4)$ be the tropicalization of the embedding of \mathbb{P}^1 into \mathbb{P}^4 given by $\phi: [x : y] \mapsto [x : y : x - y : x - 2y : x - 5y]$. Then X is given by the tropical ideal $\text{trop}(J)$ for $J = \langle u - x + y, v - x + 2y, w - x + 5y \rangle \subseteq K[x, y, u, v, w]$. On the chart $x \neq \infty$ the coordinate ring of X is

$$\overline{\mathbb{R}}[y/x, z/x, u/x, v/x, w/x] / \mathcal{R},$$

where \mathcal{R} is the bend congruence of the ideal $\text{trop}(\langle u/x - 1 + y/x, v/x - 1 + 2y/x, w/x - 1 + 5y/x \rangle)$. This contains relations such as

$$u/x \oplus 0 \oplus y/x \sim u/x \oplus 0 \sim u/x \oplus y/x \sim 0 \oplus y/x.$$

\diamond

We then define tropical vector bundles on subschemes of tropical toric varieties to be the restriction of tropical vector bundles on the ambient toric variety.

Definition 8.3. Let X be a tropical subscheme of a tropical toric variety $\text{trop}(X_\Sigma)$. A tropical toric vector bundle on X is given by the data of a tropical toric vector bundle on the ambient variety $\text{trop}(X_\Sigma)$.

Example 8.4. Let $Y = V(x^3 + y^3 + z^3) \subseteq \mathbb{P}^2$, and let $X = \text{trop}(Y) \subseteq \text{trop}(\mathbb{P}^2)$. Let \mathcal{E} be the tropical tangent bundle on \mathbb{P}^2 . The restriction of \mathcal{E} to X is described by the \mathbb{Z} -graded semimodule

$$(\overline{\mathbb{R}}[x, y, z] / \mathcal{X})^3 / R,$$

where \mathcal{X} is the bend congruence $\{x^3 \oplus y^3 \oplus z^3 \sim x^3 \oplus y^3, \dots\}$ of X , and R is the congruence generated by $\{x\mathbf{e}_1 \oplus y\mathbf{e}_2 \sim x\mathbf{e}_1 \oplus z\mathbf{e}_3 \sim y\mathbf{e}_2 \oplus z\mathbf{e}_3\}$. \diamond

We can tropicalize vector bundles on projective varieties as follows.

Definition 8.5. Let \mathcal{F} be a vector bundle on a projective variety Y . Suppose

$$(8.1) \quad \begin{aligned} &\text{there is an embedding } \phi: Y \rightarrow X_\Sigma \text{ of } Y \text{ into a toric variety } X_\Sigma, \\ &\text{and a toric vector bundle } \mathcal{E} \text{ on } X_\Sigma \text{ for which } \mathcal{F} = \phi^*(\mathcal{E}). \end{aligned}$$

Then a tropicalization of \mathcal{F} is given by the tropicalization of the subscheme $Y \subseteq X_\Sigma$, and the data of a tropicalization of \mathcal{E} in the sense of Definition 5.8.

Condition 8.1 does not hold for all embeddings of Y into a toric variety, as the following example shows.

Example 8.6. Let \mathcal{F} be the coherent sheaf on \mathbb{P}^1 given by $\mathcal{F} = \tilde{P}$ for the module

$$P = K[x, y]^3 / \langle (x - y)\mathbf{e}_1 + (x - 2y)\mathbf{e}_2 + (x - 5y)\mathbf{e}_3 \rangle.$$

over $K[x, y]$. Note that \mathcal{F} is locally free, as $\mathbb{P}^1 = D(x - y) \cup D(x - 2y) \cup D(x - 5y)$ and on each of these open sets \mathcal{F} is free. For example,

$$\begin{aligned} \mathcal{F}(D(x - y)) &= (P_{x-y})_{\mathbf{0}} \\ &= K[x/(x - y), y/(x - y)]^3 / \langle \mathbf{e}_1 + (x - 2y)/(x - y)\mathbf{e}_2 + (x - 5y)/(x - y)\mathbf{e}_3 \rangle \\ &\cong K[x/(x - y), y/(x - y)]\mathbf{e}_2 \oplus K[x/(x - y), y/(x - y)]\mathbf{e}_3. \end{aligned}$$

Thus \mathcal{F} defines a rank-2 vector bundle on \mathbb{P}^1 . However \mathcal{F} is not presented as an equivariant vector bundle, so the embedding $\phi: \mathbb{P}^1 \rightarrow \mathbb{P}^1$ does not satisfy 8.1. \diamond

Question 8.7. Given a vector bundle \mathcal{E} on a projective variety Y , does there exist an embedding $i: Y \rightarrow X_{\Sigma}$ of Y into a toric variety X_{Σ} with the property that $\mathcal{E} = i^*\mathcal{F}$ for a toric vector bundle \mathcal{F} on X_{Σ} ?

The vector bundles for which the answer to Question 8.7 is yes are those that we can currently tropicalize.

It is important to note that Question 8.7 does not fix the ambient toric variety. Given one embedding of Y into a toric variety, we may want to reembed Y into a larger toric variety to achieve success. We illustrate this with Example 8.6.

Example 8.8. Consider the example of Example 8.6. We embed \mathbb{P}^1 into \mathbb{P}^4 by $\phi: [x : y] \mapsto [x : y : x - y : x - 2y : x - 5y]$. We then have \mathbb{P}^1 is the subscheme defined by the ideal $J = \langle u - x + y, v - x + 2y, w - x + 5y \rangle \subseteq S = K[x, y, u, v, w]$. We have

$$P \cong (S/J)^3 / \langle u\mathbf{e}_1 + v\mathbf{e}_2 + w\mathbf{e}_3 \rangle.$$

Let X_{Σ} be \mathbb{P}^4 with all torus invariant planes removed; this has fan the 1-skeleton of the fan of \mathbb{P}^4 , and contains $\phi(\mathbb{P}^1)$. The lift

$$\tilde{P} = S^3 / \langle u\mathbf{e}_1 + v\mathbf{e}_2 + w\mathbf{e}_3 \rangle$$

defines a coherent sheaf \mathcal{F} on \mathbb{P}^4 , and thus on X_{Σ} . Note that it is not a vector bundle on \mathbb{P}^4 , but is on X_{Σ} , and \mathcal{E} is the restriction of \mathcal{F} to $\phi(\mathbb{P}^1) \cong \mathbb{P}^1$. \diamond

Remark 8.9. (1) The answer to Question 8.7 is yes almost by definition when \mathcal{E} is a very ample line bundle: in that case $\mathcal{E} = \phi^*(\mathcal{O}(1))$ for the embedding $\phi: Y \rightarrow \mathbb{P}^{h^0(\mathcal{F})-1}$ given by \mathcal{E} . This extends easily to all line bundles.

(2) However the answer to Question 8.7 is not uniformly positive, at least if we assume that the toric variety X_{Σ} is smooth. The Chow ring of a smooth toric variety is generated by the classes of divisors, so in particular all Chern classes of a toric vector bundle \mathcal{F} are polynomials in classes of divisors on X_{Σ} . Since Chern classes of vector bundles pullback, any vector bundle for which the answer to Question 8.7 is yes must have its Chern classes in the subring of the Chow ring of Y generated by divisors. As this is not the case for most tautological bundles of Grassmannians (see, for example, [EH16, Theorem 5.26]), we see that the answer to Question 8.7 is not always yes. We thank Michel Brion for pointing this out to us.

REFERENCES

- [All12] Lars Allermann, *Chern classes of tropical vector bundles*, Ark. Mat. **50** (2012), no. 2, 237–258.
- [AK06] Federico Ardila and Caroline J. Klivans, *The Bergman complex of a matroid and phylogenetic trees*, J. Combin. Theory Ser. B **96** (2006), no. 1, 38–49.
- [Bir67] Garrett Birkhoff, *Lattice theory*, 3rd ed., American Mathematical Society Colloquium Publications, vol. Vol. XXV, American Mathematical Society, Providence, RI, 1967.

- [BDGP21] Indranil Biswas, Arijit Dey, Ozhan Genc, and Mainak Poddar, *On stability of tangent bundle of toric varieties*, Proc. Indian Acad. Sci. Math. Sci. **131** (2021), no. 2, Paper No. 36, 21.
- [BdM08] Joseph E. Bonin and Anna de Mier, *The lattice of cyclic flats of a matroid*, Ann. Comb. **12** (2008), no. 2, 155–170.
- [CDPR12] Filip Cools, Jan Draisma, Sam Payne, and Elina Robeva, *A tropical proof of the Brill-Noether theorem*, Adv. Math. **230** (2012), no. 2, 759–776.
- [Cox95] David A. Cox, *The homogeneous coordinate ring of a toric variety*, J. Algebraic Geom. **4** (1995), no. 1, 17–50.
- [CLS11] David A. Cox, John B. Little, and Henry K. Schenck, *Toric varieties*, Graduate Studies in Mathematics, vol. 124, American Mathematical Society, Providence, RI, 2011.
- [CD73] Peter Crawley and Robert P. Dilworth, *Algebraic Theory of Lattices*, the University of Michigan, Prentice-Hall, 1973.
- [CHLSW22] Colin Crowley, June Huh, Matt Larson, Connor Simpson, and Botong Wang, *The Bergman fan of a polymatroid*, 2022. arXiv:2207.08764.
- [DDK20] Jyoti Dasgupta, Arijit Dey, and Bivas Khan, *Stability of equivariant vector bundles over toric varieties*, Doc. Math. **25** (2020), 1787–1833.
- [DDK21] ———, *Erratum for “Stability of equivariant vector bundles over toric varieties”*, Doc. Math. **26** (2021), 1271–1274.
- [Dev22] Lucie Devey, *A combinatorial description of stability for toric vector bundles*, 2022. arXiv:2212.11020.
- [DJS] Sandra Di Rocco, Kelly Jabbusch, and Gregory G. Smith, *Toric vector bundles and parliaments of polytopes*, Trans. Amer. Math. Soc. **370** (2018), no. 11, 7715–7741.
- [DW92] Andreas W. M. Dress and Walter Wenzel, *Valuated matroids*, Adv. Math. **93** (1992), no. 2, 214–250.
- [EH16] David Eisenbud and Joe Harris, *3264 and all that—a second course in algebraic geometry*, Cambridge University Press, Cambridge, 2016.
- [FJP00] Gavril Farkas, David Jensen, and Sam Payne, *The Kodaira dimensions of $\overline{\mathcal{M}}_{22}$ and $\overline{\mathcal{M}}_{23}$* , 2000. arXiv:2005.00622.
- [Ful93] William Fulton, *Introduction to toric varieties*, Annals of Mathematics Studies, vol. 131, Princeton University Press, Princeton, NJ, 1993. The William H. Roever Lectures in Geometry.
- [GM23a] Courtney George and Christopher Manon, *Cox rings of projectivized toric vector bundles and toric flag bundles*, J. Pure Appl. Algebra **227** (2023), no. 11, Paper No. 107437, 25.
- [GM23b] Courtney George and Christopher Manon, *Positivity properties of divisors on Toric Vector Bundles*, 2023. arXiv:2308.09014.
- [GM23c] Courtney George and Christopher Manon, *Algebra and Geometry of Irreducible toric vector bundles of rank n on \mathbb{P}^n* , 2023. arXiv:2308.09017.
- [GG16] Jeffrey Giansiracusa and Noah Giansiracusa, *Equations of tropical varieties*, Duke Math. J. **165** (2016), no. 18, 3379–3433.
- [Gon11] Jose L. Gonzalez, *Toric projective bundles*, ProQuest LLC, Ann Arbor, MI, 2011. Thesis (Ph.D.)—University of Michigan.
- [GUZ22] Andreas Gross, Martin Ulirsch, and Dmitry Zakharov, *Principal bundles on metric graphs: the GL_n case*, Adv. Math. **411** (2022), Paper No. 108775, 45.
- [Hel74] Thorkell Helgason, *Aspects of the theory of hypermatroids*, Hypergraph Seminar (Proc. First Working Sem., Ohio State Univ., Columbus, Ohio, 1972; dedicated to Arnold Ross), Lecture Notes in Math., vol. Vol. 411, Springer, Berlin-New York, 1974, pp. 191–213.
- [HNS22] Milena Hering, Benjamin Nill, and Hendrik Süß, *Stability of tangent bundles on smooth toric Picard-rank-2 varieties and surfaces*, Facets of algebraic geometry. Vol. II, London Math. Soc. Lecture Note Ser., vol. 473, Cambridge Univ. Press, Cambridge, 2022, pp. 1–25.
- [HLSV22] Edin Husić, Georg Loho, Ben Smith, and László A. Végh, *On complete classes of valuated matroids*, Proceedings of the 2022 Annual ACM-SIAM Symposium on Discrete Algorithms (SODA), [Society for Industrial and Applied Mathematics (SIAM)], Philadelphia, PA, 2022, pp. 945–962.
- [JR21] David Jensen and Dhruv Ranganathan, *Brill-Noether theory for curves of a fixed gonality*, Forum Math. Pi **9** (2021), Paper No. e1, 33.
- [JMT19] Jaiung Jun, Kalina Mincheva, and Jeffrey Tolliver, *Picard groups for tropical toric schemes*, Manuscripta Math. **160** (2019), no. 3–4, 339–357.
- [JMT24] ———, *Vector bundles on tropical schemes*, J. Algebra **637** (2024), 1–46.
- [Kaj08] Takeshi Kajiwara, *Tropical toric geometry*, Toric topology, Contemp. Math., vol. 460, Amer. Math. Soc., Providence, RI, 2008, pp. 197–207.
- [Kat16] Eric Katz, *Matroid theory for algebraic geometers*, Nonarchimedean and tropical geometry, Simons Symp., Springer, [Cham], 2016, pp. 435–517.
- [KM22] Kiumars Kaveh and Christopher Manon, *Toric flat families, valuations, and applications to projectivized toric vector bundles*, 2019. arXiv:1907.00543.
- [KM23] ———, *Toric vector bundles, valuations and tropical geometry*, 2023. arXiv:2304.11211.
- [KM24] ———, *Toric Matroid Bundles*, 2024. Preprint.

- [KD19] Bivas Khan and Jyoti Dasgupta, *Toric vector bundles on Bott tower*, Bull. Sci. Math. **155** (2019), 74–91.
- [Kly89] Aleksander A. Klyachko, *Equivariant bundles over toric varieties*, Izv. Akad. Nauk SSSR Ser. Mat. **53** (1989), no. 5, 1001–1039, 1135.
- [Koo11] Martijn Kool, *Fixed point loci of moduli spaces of sheaves on toric varieties*, Adv. Math. **227** (2011), no. 4, 1700–1755.
- [LV81] Michel Las Vergnas, *On products of matroids*, Discrete Math. **36** (1981), no. 1, 49–55.
- [Lov83] László Lovász, *Submodular functions and convexity*, Mathematical programming: the state of the art (Bonn, 1982), Springer, Berlin, 1983, pp. 235–257.
- [MS15] Diane Maclagan and Bernd Sturmfels, *Introduction to tropical geometry*, Graduate Studies in Mathematics, vol. 161, American Mathematical Society, Providence, RI, 2015.
- [MR18] Diane Maclagan and Felipe Rincón, *Tropical ideals*, Compos. Math. **154** (2018), no. 3, 640–670.
- [MR22] ———, *Varieties of tropical ideals are balanced*, Adv. Math. **410** (2022), Paper No. 108713, 44.
- [MT01] Kazuo Murota and Akihisa Tamura, *On circuit valuation of matroids*, Adv. in Appl. Math. **26** (2001), no. 3, 192–225. MR1818743
- [OSS11] Christian Okonek, Michael Schneider, and Heinz Spindler, *Vector bundles on complex projective spaces*, Modern Birkhäuser Classics, Birkhäuser/Springer Basel AG, Basel, 2011. Corrected reprint of the 1988 edition; With an appendix by S. I. Gelfand.
- [Oxl92] James G. Oxley, *Matroid theory*, Oxford Science Publications, The Clarendon Press, Oxford University Press, New York, 1992.
- [Pan15] Thiam-Sun Pang, *The Harder-Narasimhan filtrations and rational contractions*, Ph.D. Thesis, Albert-Ludwigs-Universität Freiburg im Breisgau, 2015.
- [Pay08] Sam Payne, *Moduli of toric vector bundles*, Compos. Math. **144** (2008), no. 5, 1199–1213.
- [Pay09] ———, *Analytification is the limit of all tropicalizations*, Math. Res. Lett. **16** (2009), no. 3, 543–556.
- [Per03] Markus Perling, *Resolutions and Moduli for Equivariant Sheaves over Toric Varieties*, Ph.D. thesis, University of Kaiserslautern, 2003.
- [Per04] ———, *Graded rings and equivariant sheaves on toric varieties*, Math. Nachr. **263/264** (2004), 181–197.
- [Sim77] Julie A. Sims, *An extension of Dilworth’s theorem*, J. London Math. Soc. (2) **16** (1977), no. 3, 393–396.
- [Sta71] Richard P. Stanley, *Modular elements of geometric lattices*, Algebra Universalis **1** (1971/72), 214–217, DOI 10.1007/BF02944981.
- [Sta12] ———, *Enumerative combinatorics. Volume 1*, Second, Cambridge Studies in Advanced Mathematics, vol. 49, Cambridge University Press, Cambridge, 2012.
- [Whi86] Neil White, *Theory of Matroids (Encyclopedia of Mathematics and its Applications)*., Cambridge: Cambridge University Press., 1986.
- [Zie91] Günter M. Ziegler, *Binary supersolvable matroids and modular constructions*, Proc. Amer. Math. Soc. **113** (1991), no. 3, 817–829.
- [Zie92] ———, *Combinatorial Models for Subspace Arrangements*, 1992.

DEPARTMENT OF MATHEMATICS, CHENNAI MATHEMATICAL INSTITUTE, CHENNAI, INDIA
 Email address: bivaskhan10@gmail.com

MATHEMATICS INSTITUTE, UNIVERSITY OF WARWICK, COVENTRY, CV4 7AL, UNITED KINGDOM
 Email address: D.Maclagan@warwick.ac.uk