### EXAMPLES OF INCLUSIONS OF UNITAL C\*-ALGEBRAS OF INDEX-FINITE TYPE WITH THE ROKHLIN PROPERTY

HYUN HO LEE, HIROYUKI OSAKA, AND TAMOTSU TERUYA

ABSTRACT. In this note, we provide several examples of inclusions of unital  $C^*$ algebras of index finite-type with the Rokhlin property lacking in literature and more importantly we construct the example which does not belong to the fixed point algebra model which means that for the inclusion  $P \subset A$  there is neither finite group action  $G \curvearrowright A$  such that  $P = A^G$  nor finite dimensional  $C^*$ - Hopf algebra action  $H \curvearrowright A$  such that  $P = A^H$ .

#### 1. INTRODUCTION

In [11], H. Osaka, K. Kodaka, T. Teruya introduced the Rokhlin property for an inclusion of unital  $C^*$ -algebras and its dual notion so called the approximate representability following Izumi's pioneering work on the Rokhlin property of a finite group action [5,6]. Along the developments in the finite group action case, the authors investigated several aspects of this new notion for inclusion of unital  $C^*$ -algebras of index-finite type. In particular, they showed that if there is a finite group G, a group action  $\alpha: G \to A$  on a unital C<sup>\*</sup>-algebra A with the Rokhlin property, then the canonical expectation  $E_{\alpha}$  from A to  $A^{\alpha}$  the fixed point algebra has the Rokhlin property, so thereby they provided the examples for the inclusion of unital  $C^*$ -algebras of index-finite type with the Rokhlin property. To get examples outside group actions, they broaden the scope from finite group actions to finite dimensional  $C^*$ -Hopf algebra actions and extend the Rokhlin property accordingly. In this way, they observe that if the finite dimensional  $C^*$ -Hopf algebra H which acts on a simple unital  $C^*$ algebra A is not commutative, the fixed point algebra  $A^H$ -model, i.e.  $A^H \subset A$ , does not come from a finite group action. Although it is not completely out of interest, it is unsatisfactory in the sense that they introduced a new category which is much larger than the category of finite groups and introduce the new Rokhlin property for such  $C^*$ -Hopf algebra actions extending the Rokhlin property for the group action. Nontheless, such examples still belongs to the fixed point algebra models.

Though the Rokhlin property for an inclusion of  $C^*$ -algebras of index-finite type was extended to the case of  $\sigma$ -unital  $C^*$ -algebra [10] or to the case of allowing the tracial approximation in the sense H. Lin [8,9], the examples beyond the fixed point algebra models have been unknown until recently. This has been a disadvantage for attracting more attention from the  $C^*$ -dynamics research group. In search of such examples, we consider several examples of inclusions of unital  $C^*$ -algebras with the

<sup>2010</sup> Mathematics Subject Classification. Primary:46L55. Secondary:47C15, 46L35.

Key words and phrases. Inclusions of unital  $C^*$ -algebras, Watatani index, Rokhlin property.

The second author is partially supported by the JSPS grant for Scientific Research No.20K03644.

Rokhlin property whose constructions do not rely on the explicit group action on the original algebra; without the prescribed group and its action we define the conditional expectation E from A and its image P together and show that E has the Rokhlin property. It is interesting that all our examples with the integer index are associated with (intrinsic) group actions so that the fixed point algebra models are recovered. This phenomenon was not observed in the past and we think that it is worthwhile to record them here. Our main work is the following; based on Izumi's example of an inclusion of unital  $C^*$ -algebras with a non-integer Jones index we construct an inclusion of unital  $C^*$ -algebras with the Rokhlin property (see Proposition 4.5 and Corollary 4.7) and such an example does not belong to the fixed point algebra models which come from either finite group actions or finite dimensional  $C^*$ -Hopf algebra actions since it is irreducible and has a non-integer index value. Moreover, due to the duality between the Rokhlin property and the approximate representability [11, Proposition 3.4] it follows that there exists an inclusion of unital  $C^*$ -algebras of index-finite type with the approximate representability, which does not come from the group action. Thus we resolve an issue which has been curious among experts.

# 2. Examples of inclusions of unital $C^*$ -algebras of index-finite type with the Rokhlin property

Since we only deal with unital  $C^*$ -algebras, we introduce the notion of index for an inclusion  $P \subset A$  in the sense of Watatani [13].

**Definition 2.1** (Y. Watatani). Let  $P \subset A$  be an inclusion of unital  $C^*$ -algebras with a conditional expectation  $E: A \to P$ , which is a completely positive contraction such that E(b) = b, E(xb) = E(x)b and E(cx) = cE(x) for all  $x \in A$  and  $b, c \in P$  or shortly projection map of norm one [1, II.6.10.1]. A quasi-basis for E is a finite set of order pairs  $\{(u_i, v_i)\}_{i=1}^n$  such that for every  $a \in A$ 

$$a = \sum_{i=1}^{n} u_i E(v_i a) = \sum_{i=1}^{n} E(au_i)v_i.$$

When  $\{(u_i, v_i)\}_{i=1}^n$  is a quasi-basis for E, Index E is defined by Index  $E = \sum_{i=1}^{n} u_i v_i$  which is independent of the choice of the quasi-basis. In this case, it is said that the inclusion  $P \subset A$  is of index-finite type [13, Definition 1.2.2].

For a  $C^*$ -algebra, recall that  $A_{\infty} = l^{\infty}(\mathbb{N}, A)/c_0(A)$  where  $c_0(A)$  is the ideal of sequences whose norm limit is 0. We embed A into  $A_{\infty}$  by the equivalence classes of constant sequences and call  $A_{\infty} \cap A'$  the central sequence algebra of A.

**Definition 2.2.** [11, Definition 3.1] Let  $P \subset A$  be an inclusion of unital  $C^*$ -algebras of index-finite type. Then we say that E has the Rokhlin property if there exist a nonzero projection e in  $A_{\infty} \cap A'$ , the central sequence algebra of A, satisfying the following two conditions;

(2.1) 
$$(\operatorname{Index} E)E^{\infty}(e) = 1_{A_{\infty}},$$

and

(2.2) the map 
$$A \ni x \mapsto xe \in A_{\infty}$$
 is injective.

We call e the Rokhlin projection.

Let us begin with an elementary example but which is heuristically useful to understand the Rokhlin property.

**Example 2.3.** Let  $X_1, X_2, \ldots, X_n$  be compact Hausdorff spaces such that there are homeomorphisms from  $X_i$  to  $X_j$  which is denoted by  $\sigma_i^j$ , for instance  $X_i = [i - i]$ 1, i - 2/3 and  $\sigma_i^j(t) = t + (j - i)$  where  $1 \leq i, j \leq n$  or more generally when there are pull-back and push-forward maps which are not precisely homeomorphisms. Note that  $\sigma_i^k \circ \sigma_i^j = \sigma_i^k$ . Now let  $X = \bigcup_{i=1}^n X_i$  and consider A = C(X) and for  $f \in A$  we write  $f = (f_1, f_2, \ldots, f_n)$  where each  $f_i$  is the element of  $C(X_i)$  and set  $P = \{(f_1, f_2, \dots, f_n) \in A \mid f_i(\sigma_i^i(t)) = f_j(t) \quad \forall i, j\}.$  Then we define a conditional expectation from A to P as follows; given  $f = (f_1, \ldots, f_n) \in A$ ,

(2.3) 
$$E(f)(t) = \frac{\sum_{k} f_k(\sigma_i^k(t))}{n} \quad \text{for} \quad t \in X_i.$$

Indeed, if we write  $E(f) = (g_1, \ldots, g_n)$ , then

$$g_i(\sigma_j^i(t)) = \frac{\sum_k f_k((\sigma_i^k \circ \sigma_j^i)(t))}{n}$$
$$= \frac{\sum_k f_k(\sigma_j^k(t))}{n}$$
$$= g_j(t) \quad \text{for} \quad t \in X_j,$$
so it is well-defined. Then define  $e_j = \begin{cases} 1 & \text{if } x \in X_j \\ 0 & \text{otherwise} \end{cases}$ , or  $e_j = (0, 0, \dots, 1, 0, \dots, 0).$ Then  $E(e_j) = \frac{1}{n}(1, 1, \dots, 1)$  or  $E(e_j) = \frac{1}{n}1_A$  for all  $j$ .  $E$  is of index finite-type and

has the Rokhlin property.

 $\mathbf{SO}$ 

Let  $\{(u_i = ne_i, v_i = e_i) \mid n = 1, \dots, n\}$ , then it is a quasi-basis for E. Let  $f = (f_1, \ldots, f_n)$ . Then

$$\sum_{i=1}^{n} u_i E(v_i f) = \sum_{i=1}^{n} ne_i \left( \frac{f_i \circ \sigma_i^1}{n}, \dots, \frac{f_i \circ \sigma_n^i}{n} \right)$$
$$= \sum_{i=1}^{n} (0, \dots, 0, f_i, 0, \dots, 0) = (f_1, \dots, f_n) = f.$$

Note that Index  $E = \sum_{i=1}^{n} u_i v_i = n I_A$  and let e be an element of  $A_{\omega}$  defined by the following periodic sequence  $(e_1, e_2, \ldots, e_n, e_1, e_2, \ldots, e_n, \ldots)$ . Since A is commutative, automatically  $e \in A_{\infty} \cap A'$ . Moreover,

$$E^{\infty}(e) = [(E(e_1), E(e_2), \dots, E(e_n), E(e_1), \dots)] = \frac{1}{n} I_{A_{\infty}}.$$

Thus,  $(\operatorname{Index} E)E^{\infty}(e) = I_{A_{\infty}}$ . Finally, we show that the map  $A \ni f \mapsto fe \in A_{\infty}$  is injective. For  $f = (f_1, \ldots, f_n)$ ,  $f = [(\mathbf{f}_1, \mathbf{f}_2, \ldots, \mathbf{f}_n, \mathbf{f}_1, \ldots)]$  where  $\mathbf{f}_j$  is the function such that  $f_j$  on  $X_j$  and vanishes elsewhere. Write fe as  $[(\mathbf{f}_k)]$  where  $\mathbf{f}_{nq+l} = \mathbf{f}_l$  for all  $q, l \in \mathbf{N}$ . Let fe = 0, then for any  $\epsilon > 0$ , there is a natural number N such that

 $\|\mathbf{f}_k\| < \epsilon$  for all  $k \ge N$ . Write N = nq + r for some natural numbers  $q \ge 0$  and r < n. Then we have

$$\|\mathbf{f}_{n(q+1)+j}\| < \epsilon$$

for  $j = 1, 2, \ldots, n$ . Consequently,

$$\|\mathbf{f}_j\| < \epsilon$$

for j = 1, 2, ..., n which implies  $||f|| < \epsilon$  in A. Thus f = 0 in A. So we are done.

If we forget homeomorphisms between equivalent spaces, or regard all spaces as same, thereby no actions are needed, then we have an example of inclusions of unital  $C^*$ -algebras of index finite-type with the Rokhlin property irrelavant of group actions.

**Example 2.4.** Let A be a unital C<sup>\*</sup>-algebra and for any  $n \in \mathbb{N}$  with  $n \geq 2$  consider  $B = A^n = A \oplus A \oplus \cdots \oplus A$ , the *n*-direct sum of A's. Let  $D = \{(x, x, \dots, x) \mid x \in A\}$  be a unital sub-C<sup>\*</sup>-algebra of B. We define a conditional expectation  $E : B \to D$  by

(2.4) 
$$E(x_1, x_2, \cdots, x_n) = \left(\frac{\sum_{i=1}^n x_i}{n}, \cdots, \frac{\sum_{i=1}^n x_i}{n}\right)$$

Let  $e_1 = (1, 0, \ldots, 0), e_2 = (0, 1, 0, \ldots, 0), \ldots, e_n = (0, \ldots, 0, 1)$  are the canonical central projections for  $A^n$ , then define  $u_i = ne_i$  and  $v_i = e_i$  for  $i \le i \le n$ . We claim that  $\{(u_i, v_i) \mid i = 1, \ldots, n\}$  is a quasi-basis for E as follows; for any  $x = (x_i) \in B$ ,

$$\sum_{i=1}^{n} u_i E(v_i x) = \sum_{i=1}^{n} n e_i \left(\frac{x_i}{n}, \frac{x_i}{n}, \dots, \frac{x_i}{n}\right)$$
$$= \sum_{i=1}^{n} n \left(0, \dots, 0, \frac{x_i}{n}, 0, \dots, 0\right) = (x_1, \dots, x_n) = x.$$

Similarly,  $\sum_{i=1}^{n} E(xu_i)v_i = x$ . Note that Index  $E = \sum_{i=1}^{n} u_i v_i = n$ .

We claim that E has the Rokhlin property: Note that  $E(e_i) = \frac{1}{n} \mathbb{1}_B$  for every  $i = 1, \ldots, n$ . We define a Rokhlin projection e by a periodic infinite sequence  $(e_1, e_2, \ldots, e_n, e_1, e_2, \ldots, e_n, e_1, \ldots)$ . Since  $e_i$ 's are central projections, it follows that  $e \in B_{\infty} \cap B'$  and  $E^{\infty}(e) = [(E_i(e_j)_{j=1}^{\infty}] = \frac{1}{n} \mathbb{1}_{B_{\infty}}$ , thus  $(\text{Index } E)E^{\infty}(e) = \mathbb{1}_{B_{\infty}}$ . Finally, we check that the map  $B \ni x \mapsto xe \in B_{\infty}$  is injective. Let  $x = (x_1, x_2, \ldots, x_n)$  and  $\mathbf{x}_i = (0, \ldots, 0, x_i, 0, \ldots, 0) = xe_i$  for each i, then xe is also represented by a periodic sequence  $(\mathbf{x}_1, \mathbf{x}_2, \cdots, \mathbf{x}_n, \mathbf{x}_1, \ldots)$ . Now suppose that xe = 0, then for any  $\epsilon > 0$ , there is a natural number N such that  $||\mathbf{x}_k|| < \epsilon$  for all  $k \ge N$ . Write N = nq + r for some natural numbers  $q \ge 0$  and r < n. Then we have

 $\|\mathbf{x}_{n(q+1)+j}\| < \epsilon$ 

for  $j = 1, 2, \ldots, n$ . Consequently,

$$\|\mathbf{x}_{j}\| < \epsilon$$

for j = 1, 2, ..., n which implies  $||x|| < \epsilon$  in B. Thus x = 0 in B.

**Remark 2.5.** If we consider the mod *n*- shift action or the circular action on  $B = A^n$  in the above, then we have a  $\mathbb{Z}/n\mathbb{Z}$ -action  $\alpha$  on B and it is easily checked that  $D = B^{\alpha}$  and  $E = E_{\alpha}$ .

The examples that we have seen so far are canonical in some sense, now we show that there is a nontrivial example from the hyperfinite  $II_1$  factor appeared in Vaughan Jones' celebrated work [7].

**Example 2.6.** Consider  $e_1, e_2, \ldots$  be a sequence of Jones' projections in a hyperfinite factor M of type  $II_1$  which satisfies the following relations;

(2.5) 
$$\begin{aligned} e_i e_{i\pm 1} e_i &= \tau e_i \\ e_i e_j &= e_j e_i \quad \text{for} \quad |i-j| \ge 2. \end{aligned}$$

and  $M = \{e_1, e_2, \dots\}''$ , where  $\tau^{-1}$  is a fixed constant in  $\{4\cos^2\frac{\pi}{n} \mid n = 3, 4, \dots\}$ . In this case, when  $\mathcal{B}$  is the  $C^*$ -algebra generated by  $1, e_1, e_2, \dots$ , and  $\mathcal{A}$  is the sub- $C^*$ -algebra generated by  $1, e_2, e_3, \dots$ , there exists a conditional expectation  $E : \mathcal{B} \to \mathcal{A}$  such that

(2.6) 
$$E(xe_1y) = \tau xy \text{ for } x, y \in \mathcal{A}.$$

Now we consider the  $C^*$ -algebra generated by  $1, e_1, e_3, e_4, \ldots$ , denoted by B, and the  $C^*$ -algebra generated by  $1, e_3, e_4, \ldots$ , denoted by A. Then  $B \subset \mathcal{B}$  and  $A \subset \mathcal{A}$  and E(B) = A. Thus if we restrict E on B, then we obtain the conditional expectation from B onto A and still denote it by E.

Let  $u_1 = \frac{e_1}{\tau}, v_1 = e_1, u_2 = \frac{1 - e_1}{1 - \tau}, v_2 = 1 - e_1$ . Then  $\{(u_1, v_1), (u_2, v_2)\}$  is a quasibasis for E as follows; for any  $x \in A$ 

$$u_1 E(v_1 x) + u_2 E(v_2 x) = \frac{e_1}{\tau} E(e_1 x) + \frac{1 - e_1}{1 - \tau} E((1 - e_1) x)$$
$$= \frac{e_1}{\tau} \tau x + \frac{1 - e_1}{1 - \tau} (1 - \tau) x \quad \text{by (3.1)}$$
$$= e_1 x + (1 - e_1) x = x.$$

Similarly,  $E(xu_1)v_1 + E(xu_2)v_2 = x$ . Thus Index  $E = \frac{e_1}{\tau} + \frac{1-e_1}{1-\tau} = \frac{1}{\tau}$  if  $\tau = 1-\tau$ or  $\tau = 1/2$  which is indeed the case  $\tau^{-1} = 4\cos^2\frac{\pi}{n}$  and n = 4. One more benefit of such a choice is that  $E(e_1) = E(1-e_1) = \tau$ .

Now we claim that E has the Rokhlin property; we take e in  $B_{\infty}$  which is represented by a periodic sequence  $(e_1, 1 - e_1, e_1, 1 - e_1, \ldots)$ . Then from (2.5)  $e \in B_{\infty} \cap B'$ . It also follows that  $E^{\infty}(e) = \tau I_{B_{\infty}}$ . Moreover, as we have seen in the above,  $(\operatorname{Index} E)E^{\infty}(e) = 1_{B_{\infty}}$ . By the same argument in Example 2.4, the map  $B \ni x \mapsto xe$  is injective. So we are done.

**Remark 2.7.** If we define an automorphism  $\alpha$  on B by sending  $e_1$  to  $1 - e_1$  and other elements fixed, then we can easily see that  $\alpha$  is of order 2 and  $E = E_{\alpha}$  in this case.

## 3. Examples of inclusions of unital $C^*$ -algebras of index-finite type without the Rokhlin property

In this section, we give elementary examples of inclusions of unital  $C^*$ -algebras of index-finite type which do not have the Rokhlin property. Of oourse, if there are no projections in the algebra, then we may not expect that the Rokhlin projection exists. However, even though there are abundant projections, it could be impossible to find a projection in  $A_{\infty} \cap A'$  satisfying the conditions of Definition 2.2 or the proper

projection in A satisfying Lemma 3.1.

In many cases, the following equivalent condition is much more practical to check whether a conditional expectation E possesses the Rokhlin property.

**Lemma 3.1.** Let A be a separable simple  $C^*$ -algebra and  $P \subset A$  be an inclusion of unital  $C^*$ -algebras and  $E : A \to P$  be a conditional expectation of index-finite type. A conditional expectation E has the Rokhlin property if for every  $\epsilon > 0$  and a finite subset  $\mathcal{F} \subset A$  there exists a projection (positive contraction) e such that

- (1)  $\|ea ae\| < \epsilon$  for every  $a \in \mathcal{F}$
- (2)  $||1 (\text{Index } E)E(e)|| < \epsilon.$

The following may be known to experts in this field and we may refer [13, Section 2.4] about the conditional expectation determined by the trace between matrix algebras. However, we present the proof directly.

**Proposition 3.2.** Let B be a unital C<sup>\*</sup>-algebra and  $A = M_n(B)$  where n = k + l ( $k \ge 1, l \ge 1$ ), then let  $P = M_k(B) \oplus M_l(B) \subset A$ . We define a linear projection E from A to P by

(3.1) 
$$E\left(\begin{pmatrix} S & T \\ V & W \end{pmatrix}\right) = \begin{pmatrix} S & 0 \\ 0 & W \end{pmatrix}$$

Then it is a conditional expectation of index-finite type. In addition, Index E = 2.

*Proof.* It is not difficult to check that for any  $b \in P$  and  $a \in A$ 

$$E(ba) = bE(a)$$
$$E(ab) = E(a)b.$$

Now we set  $u_1 = I_k, v_1 = I_k, u_2 = I_l, v_2 = I_l$  where  $I_k$  is the identity matrix in  $M_k(B), I_l$  is the identity matrix in  $M_l(B)$ . Moreover, fix  $i_0$  such that  $1 \le i_0 \le k$  then we take  $u_{j-k+2} = e_{ji_0}, v_{j-k+2} = e_{i_0j}$  for  $k+1 \le j \le n$ . Similarly, fix  $i_1$  such that  $k+1 \le i_1 \le l$  then we take  $u_{l+2+j} = e_{ji_1}, v_{l+2+j} = e_{i_1j}$  for  $1 \le j \le k$ . Then we claim that  $\{(u_i, v_i) \mid 1 \le i \le n+2\}$  is a quasi-basis for E; let  $a = \begin{pmatrix} S & T \\ V & W \end{pmatrix}$ . Note that

when  $e_{ij}$  acts on the matrix a by the left multiplication, it moves the j-th row to i-th row. Therefore,  $E(e_{ij}a)$  leaves the i-th row in part, more precisely if  $k + 1 \leq j \leq n$  and  $1 \leq i \leq k \; a_{j1}, a_{j2}, \ldots, a_{jk}$  are remained in the i-th row after taking  $E(e_{ij}a)$ . Then  $e_{ji}E(e_{ij}a)$  returns the i-row to the j-row so that the off-diagonal part V is recovered. Similarly, if  $k + 1 \leq i \leq n$  and  $1 \leq j \leq k$  then  $e_{ji}E(e_{ij}a)$  recovers T. Also, we easily check that  $u_1E(v_1a)$  and  $u_2E(v_2a)$  captures S and W respectively. Thus we conclude that  $\sum_{i=1}^{n+2} u_i E(v_i a) = a$ .

On the other hand, when  $e_{ji}$  acts on the matrix a by the right multiplication, it moves the *j*-column to the *i*-th column. Therefore  $E(ae_{ji})$  leaves the *i*-th column in part, more precisely if  $k + 1 \leq j \leq n$  and  $1 \leq i \leq k a_{1j}, a_{2j}, \ldots, a_{kj}$  are remained in the *i*-th column after taking  $E(ae_{ji})$ . Then  $E(ae_{ji})e_{ij}$  returns the *i*-column to the *j*-column so that the off-diagonal part T is recovered. Similarly, if  $k + 1 \leq j \leq n$  and  $1 \leq i \leq k$  then  $E(ae_{ij})e_{ji}$  recovers V. Also, we easily check that  $E(au_1)v_1$  and  $E(au_2)v_2$  captures S and W respectively. Thus we conclude that  $\sum_{i=1}^{n+2} E(au_i)v_i = a$ . This shows that  $\{(u_i, v_i) \mid 1 \le i \le n+2\}$  is a quasi-basis for E. Finally, Index  $E = \sum_{i=1}^{n+2} u_i v_i = I_k + I_l + \sum_{j=k+1}^n e_{ji_0} e_{i_0j} + \sum_{j=1}^k e_{ji_1} e_{i_1j} = I_k + I_l + I_k = 2I_n$ .

**Remark 3.3.** In fact, if we define E from  $A = M_n(B)$ , where n = k + l + m such that k > l > m and k - l > m, to  $P = M_k(B) \oplus M_l(B) \oplus M_m(B)$  by taking diagonal blocks as in (3.1), we can show that it is of index-finite type and Index E = 3. Moreover, there is no projection e such that  $(\text{Index } E)E(e) = 1_{M_3(B)}$ .

There is a family of inclusions of  $C^*$ -algebras of index-finite type without the Rokhlin property.

**Corollary 3.4.** Let  $A = M_n(\mathbb{C})$  where  $n = k + l \ (k \ge 2, l \ge 2)$  and k = l, then let  $P = M_k(\mathbb{C}) \oplus M_l(\mathbb{C}) \subset A$ . Let E be a conditional expectation from A to P given by (3.1) as in Proposition 3.2. Then E does not have the Rokhlin property.

*Proof.* Suppose that E has the Rokhlin property. Then for  $0 < \epsilon < \frac{1}{1024}$  and a single set  $\left\{ \begin{pmatrix} 1_k & 0\\ 0 & 0 \end{pmatrix} \right\}$  there must be a projection *e* such that

$$\|1_A - (\operatorname{Index} E)E(e)\| < \epsilon$$

(2)

$$\begin{pmatrix} 1_k & 0\\ 0 & 0 \end{pmatrix} e - e \begin{pmatrix} 1_k & 0\\ 0 & 0 \end{pmatrix} \| < \epsilon.$$

From (1), if we write  $e = \begin{pmatrix} D & B \\ B & D \end{pmatrix}$ , then  $\|D - \frac{I_k}{2}\| < \epsilon$ . Therefore if we onsider  $e' = \begin{pmatrix} \frac{1_k}{2} & B\\ B & \frac{1_k}{2} \end{pmatrix}$ , then  $||e - e'|| < \epsilon$ . It follows that  $||e'^2 - e'|| < 4\epsilon$ . This implies that ε.

(3.2) 
$$||B^2 - \frac{I_k}{4}|| < 4\epsilon$$

But from (2) we also have

$$\left\| \begin{pmatrix} 1_k & 0\\ 0 & 0 \end{pmatrix} \begin{pmatrix} D & B\\ B & D \end{pmatrix} - \begin{pmatrix} D & B\\ B & D \end{pmatrix} \begin{pmatrix} 1_k & 0\\ 0 & 0 \end{pmatrix} \right\| < \epsilon$$

This implies that

(3.3) 
$$||B|| < \epsilon \quad \text{or} \quad ||B||^2 < \epsilon^2.$$

But then

$$\frac{1}{4} - 4\epsilon < \|\frac{1_k}{4}\| - \|B^2 - \frac{I_k}{4}\| \le \|B\|^2 < \epsilon^2$$

which is a contradiction. So we prove that E does not have the Rokhlin property.  $\Box$ 

In the setting of Proposition 3.2, the case k = 1, l = 2 or k = 2, l = 1 the analysis is much easier. Even we can show that E does not satisfy the tracial Rokhlin property which means that for any finite set  $F \subset A$ ,  $\epsilon > 0$  there exists a projection e such that  $||ea - ae|| < \epsilon$  for all  $a \in F$  and  $tr(1 - (Index E)E(e)) < \epsilon$  where tr is the normalized trace since we deal with the finite simple  $C^*$ -algebra with a unique tracial state.

**Example 3.5.** In the setting of Corollary 3.4, let  $B = \mathbb{C}$ , n = 3 and k = 2, l = 1; we set  $\mathcal{F}$  to be the set of matrix units  $\{e_{ij} \mid 1 \leq i, j \leq 3\}$  and  $\epsilon < \frac{1}{12}$ . If E has the tracial Rokhlin property, then a self-adjoint element  $e = \begin{pmatrix} \lambda_1 & a & b \\ a & \lambda_2 & c \\ b & c & \lambda_3 \end{pmatrix}$  must satisfy that

(3.4)

$$e^{2} = 2, \text{ or } \begin{pmatrix} \lambda_{1}^{2} + a^{2} + b^{2} & a(\lambda_{1} + \lambda_{2}) + bc & ac + b(\lambda_{1} + \lambda_{3}) \\ a(\lambda_{1} + \lambda_{2}) + cb & \lambda_{2}^{2} + a^{2} + c^{2} & ab + c(\lambda_{2} + \lambda_{3}) \\ b(\lambda_{1} + \lambda_{3}) + ac & ba + c(\lambda_{2} + \lambda_{3}) & \lambda_{3}^{2} + b^{2} + c^{2} \end{pmatrix} = \begin{pmatrix} \lambda_{1} & a & b \\ a & \lambda_{2} & c \\ b & c & \lambda_{3} \end{pmatrix},$$

(3.5) 
$$||ee_{ij} - e_{ij}e|| < 1/4 \text{ for } i, j = 1, 2, 3,$$

and

(3.6) 
$$\tau \left( \begin{pmatrix} 1 - 2\lambda_1 & -2a & 0\\ -2a & 1 - 2\lambda_2 & 0\\ 0 & 0 & 1 - 2\lambda_3 \end{pmatrix} \right) < \frac{1}{12}.$$

From (3.4), note that  $b^2 + c^2 = \lambda_3 - \lambda_3^2$ . From (3.5),  $||e_{13}e - ee_{13}|| < 1/4$  implies that  $b^2 < \frac{1}{16}$  and  $c^2 < \frac{1}{16}$ .

Since 1 - (Index E) E(e) is positive,  $1 - 2\lambda_1 + 1 - 2\lambda_2 \ge 0$ . Thus from (3.6)

$$\frac{1}{3}(1 - 2\lambda_1 + 1 - 2\lambda_2 + 1 - 2\lambda_3) < \frac{1}{12}$$
$$(1 - 2\lambda_1 + 1 - 2\lambda_2 + 1 - 2\lambda_3) < \frac{1}{4}$$
$$0 \le 1 - 2\lambda_3 < \frac{1}{4}.$$

Consequently,

$$(3.7)\qquad \qquad \frac{3}{8} < \lambda_3 \le \frac{1}{2}$$

Then  $\lambda_3^2 - \lambda_3 > \frac{15}{64}$ . But  $\lambda_3 - \lambda_3^2 = b^2 + c^2 < \frac{1}{8}$  by (3.4) and (3.5). This is a contradiction. Hence *E* does not have the tracial Rokhlin property in this case.

### 4. Examples of inclusions of unital $C^*$ -algebras index-finite type with the Rokhlin property which does not come from the finite group action

Finally, in this section we construct an inclusions of unital  $C^*$ -algebras of indexfinite type with the Rokhlin property which does not come from the finite group action. To do so, we combine two ideas; the first is to find the inclusion of unital  $C^*$ -algebras with a non-integer index and the second is to use the infinite tensor product to find a central sequence. For the first we borrow Izumi's constructions and summarize them in the below. **Theorem 4.1.** Let A be the Cuntz algebra  $O_2$  whose generators are  $S_1, S_2$  and  $\rho$ :  $O_2 \rightarrow O_2$  the endomorphism defined by

$$\rho(S_1) = \frac{1}{d}S_1 + \frac{1}{\sqrt{d}}S_2S_2$$
  
$$\rho(S_2) = \left(\frac{1}{\sqrt{d}}S_1 - \frac{1}{d}S_2S_2\right)S_2^* + S_2S_1S_1^*$$

where d is the real number satisfies  $1 = \frac{1}{d} + \frac{1}{d^2}$ . Then the conditional expectation  $E_{\rho}: O_2 \to \text{Image } \rho$  defined by  $E_{\rho}(x) = \rho(S_1^*\rho(x)S_1)$  has  $\text{Index}E_{\rho} = d^2$  and its quasibasis is  $\{dS_1^*, dS_1\}$ . Moreover,  $e = S_1S_1^*$  satisfies that  $\text{Index}(E_{\rho})E_{\rho}(e) = 1$ .

Proof. See [4, Example 3.2].

We need Elliott's theorem  $O_2 \otimes O_2 \cong O_2$ , which we utilize to construct an inductive system for  $\bigotimes_{i=1}^{\infty} O_2$ . The following is well-known but the detail of the proof is important to us.

**Proposition 4.2.** Let  $\phi: O_2 \to O_2 \otimes O_2$  be the injective \*-homomorphism given by the second factor embedding or  $\phi(x) = 1 \otimes x$ . Then there is a sequence of unitaries  $\{v_n\}_{n=1}^{\infty}$  in  $O_2 \otimes O_2$  such that

$$\lim_{n \to \infty} \|v_n \phi(a) - \phi(a)v_n\| = 0, \quad \lim_{n \to \infty} \operatorname{dist}(v_n^* b v_n, \phi(O_2)) = 0$$

for all  $a \in O_2$  and all  $b \in O_2 \otimes O_2$ . Consequently, there is an isomorphism  $\psi : O_2 \to O_2 \otimes O_2$  which is approximately unitarily equivalent to  $\phi$  denoted by  $\phi \approx_u \psi$ .

Proof.  $v_n$ 's in the above essentially come from an asymptotically central sequence of endomorphisms,  $\{\rho_n\}_{n=1}^{\infty}$  (see [12, Lemma 5.2.3]). In fact, for an arbitrary  $\delta > 0$  there is a unitary w in  $O_2 \otimes O_2$  such that  $||w(1 \otimes S_j)w^* - S_j \otimes 1|| < \delta$  and  $||w(1 \otimes S_j^*)w^* - S_j^* \otimes 1|| < \delta$  for j = 1, 2 by the uniqueness result for  $O_2$  or the approximately inner half flip property of  $O_2$ . So we take  $w_n = (\mathrm{id} \otimes \rho_n)(w)$ . Then

$$\|w_n(1 \otimes S_j) - (1 \otimes S_j)w_n\| \to 0 \quad \text{as} \quad n \to \infty$$
$$\|w_n(1 \otimes S_j^*) - (1 \otimes S_j^*)w_n\| \to 0 \quad \text{as} \quad n \to \infty$$

for j = 1, 2. Also,

$$dist(w_n^*(S_j \otimes 1)w_n, 1 \otimes O_2) \le \|w_n^*(S_j \otimes 1)w_n - 1 \otimes \rho_n(S_j)\|$$
$$\le \|(id \otimes \rho_n)(w^*(S_j \otimes 1)w) - (id \otimes \rho_n)(1 \otimes S_j)\|$$
$$< \delta$$

for j = 1, 2. Then the conclusion follows form [12, Proposition 2.3.5]. We note that  $\psi$  is the limit of the maps of the form  $w_{n_1}w_{n_2}\cdots w_{n_k}\phi(\cdot)w_{n_k}^*\cdots w_{n_2}^*w_{n_1}^*$  for a suitable choice of  $\{w_{n_k}\}$  which is a subsequence of  $\{w_n\}$ .

**Corollary 4.3.** For each  $n \geq 2$  we inductively let  $\phi_n : \bigotimes_{i=1}^n O_2 \to O_2 \otimes (\bigotimes_{i=1}^n O_2)$  be the second factor embedding as in Proposition 4.2. Viewing  $\bigotimes_{i=1}^n O_2 \cong O_2$  via  $\psi_{n-1}$ , we can apply Proposition 4.2 to the injective map  $\phi_n \circ \psi_{n-1} : O_2 \to \bigotimes_{i=1}^{n+1} O_2$ . Therefore there is an isomorphism  $\psi_n : O_2 \to \bigotimes_{i=1}^{n+1} O_2$  which is approximately unitarily equivalent to  $\phi_n \circ \psi_{n-1}$ .

**Remark 4.4.** In the above inductive process, in each *n*-th step it is identical to use the approximately inner half flip property of  $O_2$  and an asymptotically central sequence of endomorphisms  $\rho_n$ 's while replacing  $S_j$  and  $S_j^*$  by  $\psi_{n-1}(S_j)$  and  $\psi_{n-1}(S_j^*)$  respectively.

We put  $\psi_1 = \psi$  corresponding to  $\phi_1 = \phi$  in Proposition 4.2. If we inductively replace  $\psi_n$  by Ad  $W_n \circ \psi_n$  by suitable unitary  $W_n$  in  $\bigotimes_{i=1}^{n+1}O_2$ , we have the following diagram for the inductive limit of a system  $(\bigotimes_{i=1}^n O_2, \phi_n : \bigotimes_{i=1}^n O_2 \to \bigotimes_{i=1}^{n+1} O_2)$  which is approximate intertwining in the sense of Elliott (here we still abuse  $\psi_n$  for Ad  $W_n \circ \psi_n$ without confusion);



**Proposition 4.5.** Consider  $E_{\rho}(x) = \rho(S_1^*\rho(x)S_1)$  where  $\rho$  is the endomorphism of  $O_2$  given in Theorem 4.1. Then we define  $E_{n+1} : \bigotimes_{i=1}^{n+1} O_2 \to \psi_n(\operatorname{Image} \rho)$  by  $E_{n+1} = \psi_n \circ E_{\rho} \circ \psi_n^{-1}$  for each  $n \in \mathbb{N}$  and  $E : \bigotimes_{i=1}^{\infty} O_2 \to O_2$  by  $\lim_{n \to \infty} E_n$ . It follows that Index  $E = \operatorname{Index} E_{\rho}$  and E has the Rokhlin property.

Proof. Note that for each  $n E_{n+1}$  is the conditional expectation such that Index  $E_{n+1} =$ Index  $E_{\rho}$ . Since we can easily check that  $E_{n+1} \circ \phi_n = \phi_n \circ E_n$  approximately, E is well-defined as the inductive limit of  $E_n$  (see [2, pp181 2.3]). It remains to show that E has the Rokhlin property: for given a finite set  $F \subset \bigotimes_{i=1}^{\infty} O_2$  and  $\epsilon > 0$ , there is a  $M \in \mathbb{N}$  such that  $F \subset \bigotimes_{i=1}^{M} O_2$ . Then we consider a projection M times

 $\widehat{e} = \underbrace{1 \otimes \cdots \otimes 1}_{k \in \mathbb{N}} \otimes \psi_N(e) \in \bigotimes_{i=1}^{N+M+1} O_2 \text{ for some } N \text{ where } e = S_1 S_1^*, \text{ then } \widehat{e} \text{ commutes with elements in } F.$  Note that  $\widehat{e} = (\phi_{N+M} \circ \phi_{N+M-1} \circ \cdots \circ \phi_{N+1})(\psi_N(e)).$ Thus if we choose N large enough,  $\widehat{e}$  is close to  $\psi_{N+M}(e)$  up to  $\epsilon$  by the approximate intertwining of the diagram (4.1). If necessary, we can choose N such that  $\|(\operatorname{Index} E)E(\psi_{N+M}(e)) - (\operatorname{Index} E)E_{N+M+1}(\psi_{N+M}(e))\| < \epsilon.$ Then

$$\begin{aligned} \|1 - (\operatorname{Index} E)E(\widehat{e})\| &\leq \|1 - (\operatorname{Index} E)E_{N+M+1}(\psi_{N+M}(e))\| \\ &+ \|(\operatorname{Index} E)E(\psi_{N+M}(e)) - (\operatorname{Index} E)E_{N+M+1}(\psi_{N+M}(e))\| \\ &+ \|(\operatorname{Index} E)E(\psi_{N+M}(e)) - (\operatorname{Index} E)E(\widehat{e})\| \\ &< 0 + \epsilon + (\operatorname{Index} E)\epsilon = (1 + \operatorname{Index} E)\epsilon. \end{aligned}$$

Since  $\bigotimes_{i=1}^{\infty} O_2 \cong O_2$  is simple, we are done by Lemma 3.1.

We need to make sure that the above construction of  $P = \text{Image } E \subset \bigotimes_{i=1}^{\infty} O_2$  with the conditional expectation E as in Proposition 4.5 is irreducible. To show it, we begin with a simple observation.

**Lemma 4.6.** Let  $P \subset A$  an inclusion of simple  $C^*$  algebras and  $E : A \to P$  a conditional expectation from A onto P with Index  $E < \infty$ . If Index E is less than 4, then  $P \subset A$  is irreducible, i.e.,  $P' \cap A \simeq \mathbb{C}$ .

*Proof.* Since Index E is a element of  $A' \cap A \simeq \mathbb{C}$ , we have

Index 
$$E = 4\cos^2 \frac{\pi}{n}, \quad n = 3, 4, 5, \cdots$$

by [13, Theorem 2.3.6]. Therefore the principal graph of  $P \subset A$  is in  $A_n, D_n, E_n$  by [3, Theorem 1.4.3]. In any cases, there is only one edge connected to \* and hence  $P' \cap A \simeq \mathbb{C}$ , i.e.,  $P \subset A$  is irreducible.

**Corollary 4.7.** Let  $P \subset \bigotimes_{i=1}^{\infty} O_2$  be an inclusion given in Proposition 4.5. Then,  $P' \cap \bigotimes_{i=1}^{\infty} O_2 = \mathbb{C}$ . In other words, this inclusion is irreducible.

*Proof.* Let  $E : \bigotimes_{i=1}^{\infty} O_2 \to P$ . Then, since  $\operatorname{Index} E < 4$ , from Lemma 4.6 we know that  $P' \cap \bigotimes_{i=1}^{\infty} O_2 = \mathbb{C}$ .

**Remark 4.8.** Using the irreducibility of Image  $\rho \subset O_2$  in Theorem 4.1 we can deduce the irreducibility of  $P \subset \bigotimes_{i=1}^{\infty} O_2$  directly; note that  $P = \psi(O_2)$  where  $\psi$  is the isomorphism obtained in the diagram (4.1). Let  $y \in P' \cap \bigotimes_{i=1}^{\infty} O_2$ . Then, there is a  $x \in O_2$  such that  $y = \psi(x)$ . Since  $y\psi(\rho(a)) = \psi(\rho(a))y$  for any  $a \in O_2$ ,  $\psi(x)\psi(\rho(a)) = \psi(\rho(a))\psi(x)$ , that is,  $x\rho(a) = \rho(a)x$  for any  $a \in O_2$ . Since  $\rho(O_2)' \cap$  $O_2 = \mathbb{C}$  [4, Theorem 5.6(1)],  $x \in \mathbb{C}$ . Hence  $y = \psi(x) \in \mathbb{C}$ . That is,  $P' \cap \bigotimes_{i=1}^{\infty} O_2 = \mathbb{C}$ .

#### References

- [1] B. Blackadar, *Operator algebras*, Encyclopaedia of Mathematical Scinences, Vol. 122, Springer.
- [2] G. Elliott On the classification of C\*-algebras of real rank zero, J. Reine. Angew. Math., 443(1993) 179 - 219.
- [3] F. Goodman, P. de la Harpe, and V. F. R. Jones, Coxeter graphs and towers of algebras, MSRI publications 14, Springer, 1989.
- [4] M. Izumi, Subalgebras of Infinie C\*-Algebras with Finite Watatani Indices I. Cuntz Algebras, Comm. Math. Phys. 155(1993), 157 – 182.
- [5] M. Izumi, Finite group actions on C\*-algebras with the Rokhlin property I, Duke Math. J. 122(2004), no.2, 233 – 280.
- [6] M. Izumi, Finite group actions on C<sup>\*</sup>-algebras with the Rokhlin property II, Adv. Math. 184(2004), no.1, 119 – 160.
- [7] V. Jones, Index for subfactors, Invent. Math. 72(1983), 1 25.
- [8] H. Osaka and T. Teruya, The Jiang-Su absorption for inclusions of C<sup>\*</sup>-algebras, Canad. J. Math. 70 (2018), no.2, 400–425.
- H. Osaka and T. Teruya, Erratum: The Jiang-Su absorption for inclusions of C<sup>\*</sup>-algebras, Canad. J. Math. 73(2021) no. 1, 293–295, DOI:https://doi.org/10.4153/S0008414X20000310.
- [10] H. Osaka and T. Teruya, The Rokhlin property for inclusion of C<sup>\*</sup>-algebras, Rocky Mount. J. Math. 50 (2020), no. 5. 1785 –1792.
- [11] H. Osaka, K. Kodaka, T. Teruya, The Rokhlin property for inclusion of C<sup>\*</sup>-algebras with a finite Watatani index, Contem. Math. 503(2009), 177–195.
- [12] M. Rørdam, Classification of nuclear simple C\*-aglebras, Encyclopedia of Mathematical Sciences 126, Springer-Verlag, Berlin, Heidelberg 2002
- [13] Y. Watatani, Index for C<sup>\*</sup>-subalgebras, Mem. Amer. Math. Soc. **424**(1990), AMS.

Department of Mathematics, University of Ulsan, Ulsan, 44610 South Korea & School of Mathematics, Korea institute of Advance Study, Seoul, 02455 South Korea

 $Email \ address: \verb+hadamard@ulsan.ac.kr$ 

Department of Mathematical Sciences, Ritsumeikan University, Kusatsu, Shiaga, Japan

Email address: osaka@se.ritsumei.ac.jp

FACULTY OF EDUCATION, GUNMA UNIVERSITY, MAEBASHI CITY, GUNMA, JAPAN *Email address*: teruya@gunma-u.ac.jp