MEET-U Project I: The key drivers of the preference for dynamic dark energy

Zhiqi Huang, Jianqi Liu, Jianfeng Mo, Yan Su, Junchao Wang, Yanhong Yao, Guangyao Yu, and Zhengxin Zhu School of Physics and Astronomy, Sun Yat-sen University, 2 Daxue Road, Zhuhai, 519082, CHINA*

Zhuoyang Li

Department of Astronomy, Tsinghua University, Beijing 100084, China

Zhenjie Liu

Department of Astronomy, Shanghai Jiaotong University, Shanghai, 200240, CHINA

Haitao Miao

National Astronomical Observatories, Chinese Academy of Sciences, Beijing 100101, People's Republic of China

Hui Tong

School of Physics and Astronomy, Monash University, Vic 3800, Australia and

OzGrav: The ARC Centre of Excellence for Gravitational Wave Discovery, Clayton VIC 3800, Australia

(Dated: May 8, 2024)

Joint analysis of the baryon acoustic oscillations (BAO) measurement by the Dark Energy Spectroscopic Instrument (DESI) first data release, Type Ia supernovae (SNe) of the Dark Energy Survey Year 5 (DES5YR) release and cosmic microwave background (CMB) data favors a quintom-like dynamic dark energy model over the standard Lambda cold dark matter (Λ CDM) model at 3.9 σ level (Adame et al. 2024). We demonstrate that the preference for dynamic dark energy does not rely on the detailed modeling of CMB physics and remains at 3.2 σ level when the full CMB likelihood is replaced by a CMB acoustic-oscillation angle (θ_{\star}) prior and a baryon abundance ($\Omega_b h^2$) prior. By comparing the data with over 10⁴ Λ CDM-based simulations, we find that both the SNe and BAO data contribute significantly to the preference for dynamic dark energy. The preference for dynamic dark energy is unlikely (probability $\lesssim 0.02$) due to unknown systematics in DES5YR SNe and statistical fluctuations in DESI BAO, or vice versa.

I. INTRODUCTION

The past quarter century has witnessed great success of the standard Lambda cold dark matter (Λ CDM) model, where Λ stands for the cosmological constant that acts as dark energy driving the late-time cosmic acceleration. The six-parameter Λ CDM model has been confronted with, and passed the tests of a broad range of cosmological measurements that contain billions of bits of information [1–8]. The cosmic microwave background (CMB) temperature and polarization anisotropies measured by Planck space mission [4] and ground-based telescopes [3, 8] have determined the Λ CDM parameters to percentage-level accuracy, ushering in the era of precision cosmology.

As the accuracy of observations continues to improve, however, some data that do not conform to Λ CDM have emerged [9–12]. While most studies on cosmological discordance focus on checking self consistency of Λ CDM without involving an alternative model, some observational data provide hints of dynamic dark energy model that is beyond Λ CDM [13–17]. The most recent revival of dynamic dark energy comes from the measurement of baryon acoustic oscillations (BAO) by the Dark Energy Spectroscopic Instrument (DESI) first data release [17]. The DESI BAO data, when combined with CMB and type Ia supernovae data from Dark Energy Survey Year 5 (DES5YR) release [18], favors a dynamic dark energy model with the equation of state (EOS) of dark energy parameterized as $w = w_0 + w_a(1-a)$, where *a* is the scale factor [19, 20]. The joint analysis of DESI BAO + DES5YR SNe + CMB gives $w_0 = -0.727 \pm 0.067$ and $w_a = -1.05^{+0.31}_{-0.27}$, rejecting the Λ CDM model ($w_0 = -1$ and $w_a = 0$) at 3.9 σ confidence level [17]. The significance level, however, drops to 3.5σ and 2.5σ when DES5YR SNe is replaced by the Union3 [21] and Pantheon+ supernova data [22], respectively.

While a Luminous Red Galaxy (LRG) data point at effective redshift $z_{\rm eff} = 0.51$ has been identified to be responsible for the preference for dynamic dark energy [17, 23], it remains unclear why the statistical significance is sensitive to the choice of supernova data. It has also been shown that the significance level has subtle dependence on the prior range of dark energy parameters and whether Atacama Cosmology Telescope (ACT) data release 6 CMB lensing data is included [24]. To address these confusions, the present work aims to identify which data set is the key driver of the preference for dynamic dark energy, and to investigate the possibility that the $\gtrsim 3\sigma$ rejection of Λ CDM model is due to some unknown systematics in DESI BAO or DES5YR SNe.

Throughout the paper we work with natural units c =

^{*} huangzhq25@mail.sysu.edu.cn

 $\hbar = k_B = 1$ and a spatially flat Friedmann-Robertson-Walker (FRW) background metric. The Hubble constant H_0 is written as 100h km s⁻¹Mpc⁻¹, where the dimensionless parameter h is the reduced Hubble constant. The abundance parameters Ω_c , Ω_b , $\Omega_{\rm DE}$, Ω_γ , $\Omega_{\nu i}$ (i = 1, 2, 3) are defined as the present fractional background density of cold dark matter, ordinary matter, dark energy, radiation, and the *i*-th neutrino, respectively. The matter abundance is defined as $\Omega_m = \Omega_b + \Omega_c$.

II. METHOD

For spatially flat $w_0 w_a$ CDM cosmology, the Hubble expansion rate H(z) > 0 is determined by the first Friedmann equation

$$\left[\frac{H(z)}{H_0}\right]^2 = \Omega_m (1+z)^3 + \Omega_{\rm DE} e^{3\left[(1+w_0+w_a)\ln(1+z)-w_a\frac{z}{1+z}\right]} \\ + \left[\Omega_\gamma + \sum_{i=1}^3 \Omega_{\nu,i} \frac{I_\rho\left(\frac{m_{\nu,i}}{(1+z)T_{\rm CNB}}\right)}{I_\rho\left(\frac{m_{\nu,i}}{T_{\rm CNB}}\right)}\right] (1+z)^4,$$
(1)

where z is the cosmological redshift; $m_{\nu,i}$ is the neutrino mass of the *i*-th specie; $T_{\rm CNB} = T_{\rm CMB} \left(\frac{4}{11}\right)^{1/3} \approx 1.95 \,\rm K$ is the effective temperature for neutrino momentum distribution. The neutrino density integral is

$$I_{\rho}(\lambda) \equiv \frac{1}{2\pi^2} \int_0^\infty \frac{x^2 \sqrt{x^2 + \lambda^2}}{e^x + 1} dx.$$
 (2)

As in the standard scenarios ≤ 0.1 eV neutrino masses have almost no impact on the determination of dark energy parameters [4, 25], we fix three neutrino species with masses 0.05eV, 0.009eV, 0.001eV for simplicity. The abundance parameter $\Omega_{\nu,i}$ is related to the mass parameter m_i via

$$\Omega_{\nu,i}h^2 = 1.981 \times 10^{-5} I_\rho \left(\frac{m_{\nu,i}}{T_{\rm CNB}}\right),\tag{3}$$

where a standard CMB temperature $T_{\rm CMB} = 2.726 \,\rm K$ effective neutrino number $N_{\rm eff} = 3.044$ have been used. The comoving angular diameter distance is given by

$$D_M(z) = \int_0^z \frac{dz'}{H(z')}.$$
 (4)

The sound horizon at the baryon drag epoch, r_d , is approximated by [26]

$$\frac{r_d}{147.05\,\mathrm{Mpc}} = \left(\frac{\Omega_m h^2}{0.1432}\right)^{-0.23} \left(\frac{N_{\mathrm{eff}}}{3.04}\right)^{-0.1} \left(\frac{\Omega_b h^2}{0.02236}\right)^{-0.13}$$
(5)

The BAO data measures D_M/r_d , $1/(Hr_d)$ and $D_V \equiv (zD_M^2/H)^{1/3}$. We apply multi-variable Gaussian likelihood for the data and covariance matrix listed in Table 1 of DESI 2024 paper [17], but with a better decimal precision obtained via private communication.

The distance modulus of SNe is

$$\mu(z) = 5 \log_{10} \frac{(1+z_{\rm hel})D_M(z_{\rm CMB})}{\rm Mpc} + 25, \qquad (6)$$

where z_{hel} is the heliocentric redshift and z_{CMB} is the CMB dipole-causing peculiar redshift removed. A supernova data set with observed distance moduli μ_i^{obs} $(i = 1, 2, \ldots, N)$ and their covariance matrix C has a chi-square function (defined as -2 times the logarithm of likelihood)

$$\chi^{2} = \sum_{i,j} \left(\mu_{i}^{\text{obs}} - \mu_{i}^{\text{th}} - \Delta \right) \left(C^{-1} \right)_{ij} \left(\mu_{j}^{\text{obs}} - \mu_{j}^{\text{th}} - \Delta \right),$$
(7)

where μ^{th} is the theoretical distance modulus and

$$\Delta = \frac{\sum_{i,j} \left(\mu_i^{\text{obs}} - \mu_i^{\text{th}} \right) \left(C^{-1} \right)_{ij}}{\sum_{i,j} \left(C^{-1} \right)_{ij}}$$
(8)

can be considered as the average difference between observation and theory. Because we are marginalizing over the absolute magnitude of SNe, the average difference Δ enters Eq. (7) in such a way that the likelihood is invariant under a global translation of μ^{th} . Eq. (8) can be used to define a local Δ_{bin} for a subset of SN data in a redshift bin. The difference $\Delta_{\text{bin}} - \Delta$, which we dub "binned distance modulus", measures how the data in the redshift bin deviates from theoretical prediction.

The list of z_{hel} , z_{CMB} , μ^{obs} and the covariance matrix C for DES5YR SNe data are all publicly available at https://github.com/des-science/DES-SN5YR/.

Due to strong degeneracy between parameters, a single probe of SNe, BAO or CMB cannot effectively constrain the $w_0 w_a$ CDM model. Joint analysis with CMB + BAO + SNe breaks the degeneracy and gives a prior-insensitive constraint [17, 24]. In this way, the constraint on dark energy parameters involves CMB physics and relies on many details of the cosmological model. The robustness of dark-energy constraint can be vastly enhanced if we treat CMB as a BAO data point at recombination (redshift $z \sim 1090$) and only use the constraint on the angular extension of the sound horizon on the last scattering surface, i.e., the θ_{\star} parameter. The CMB constraint on θ_{\star} is only sensitive to the spatial curvature and the earlyuniverse physics, and is almost model-independent if we work with the standard scenario of Big Bang Nucleosynthesis (BBN) in a spatially flat universe. Based on the Planck constraint [4], we adopt

$$\theta_{\star} = 0.0104092 \pm 0.0000032. \tag{9}$$

To use the CMB and BAO standard ruler informa- \cdot tion, it is necessary to determine the sound horizon at the recombination epoch, which depends on the baryon abundance $\Omega_b h^2$. We quote a BBN constraint [27]

$$\Omega_b h^2 = 0.02196 \pm 0.00063. \tag{10}$$

Hereafter we denote the θ_{\star} and $\Omega_b h^2$ constraints simply as " θ_{\star} " when brevity is needed.

TABLE I. cosmological parameters

parameter	definition	prior range
Ω_m	matter abundance	[0.05, 0.8]
w_0	dark energy EOS at $\boldsymbol{z}=\boldsymbol{0}$	[-3, 1]
w_a	dark energy EOS slope	[-3, 3]
h	reduced Hubble constant	[0.5, 0.9]
$\Omega_b h^2$	baryon density	[0.015, 0.03]



FIG. 1. Marginalized 68.3% and 95.4% confidence constraints on w_0 and w_a .

With the DESI BAO and DES5YR SNe likelihoods and constraints on θ_{\star} and $\Omega_b h^2$, we run Markov Chain Monte Carlo (MCMC) simulations to measure the cosmological parameters that are listed in Table I.

III. RESULTS

As Figure 1 shows, both $BAO+\theta_{\star}$ and $SNe+\theta_{\star}$ are consistent with ΛCDM , but their combination, namely $BAO+SNe+\theta_{\star}$ gives $w_0 = -0.756^{+0.089}_{-0.087}$ and $w_a =$ $-0.904^{+0.489}_{-0.494}$, excluding ΛCDM model at 3.2σ significance level. The BAO+SNe+ θ_{\star} constraint is similar to the BAO+SNe+CMB result presented in Ref. [17], but is more robust in the sense that it only depends on the standard BBN picture and the directly observable acoustic angular scale in the CMB maps.

While nothing precludes using the particular combination of DESI BAO and DES5YR SNe and confront it with the particular w_0w_a CDM model, given that there is yet no independent confirmation from other cosmological probes, it is reasonable to speculate unknown systematic



FIG. 2. Histogram of significance level of favoring $w_0 w_a \text{CDM}$ over ΛCDM , for DESI BAO + SN mock + θ_{\star} (upper panel) and DES5YR SNe + BAO mock + θ_{\star} (lower panel), respectively. The dotted vertical line (3.2 σ) is the significance level from the real data (DESI BAO + DES5YR SNe + θ_{\star}).

errors in either the DESI BAO or DES5YR SNe [28–31]. When Λ CDM is taken as a pre-assumption, such speculation is in fact testable. If the preference for dynamic dark energy is driven by, for instance, some unknown systematics in DESI BAO, we can replace DES5YR SNe with random Λ CDM mocks at the same redshifts and with the same error matrix. There should be some chance that DESI BAO + SN mock + θ_{\star} favors w_0w_a CDM over Λ CDM at a significance level equal to or greater than the real-data case. Similarly if some unknown systematics in DES5YR SNe is at play, there should be some chance that DES5YR SNe + BAO mock + θ_{\star} favors w_0w_a CDM over Λ CDM at a higher significance level.

We produce 5329 SN mocks and 5263 BAO mocks based on DESI/DES5YR error matrices and Planck bestfit Λ CDM cosmology ($\Omega_m = 0.3149, \Omega_b h^2 = 0.02238$ and h = 0.6732). For each SN mock, we compute the significance level, in terms of number of sigmas, of DESI BAO + SN mock + θ_{\star} favoring $w_0 w_a$ CDM over Λ CDM. Likewisely DES5YR SNe + BAO mock + θ_{\star} constraint is calculated for each BAO mock. Figure 2 shows the histogram of the results. When either DESI BAO or DES5YR SNe is replaced by Λ CDM mocks, the probability of favoring $w_0 w_a \text{CDM}$ over ΛCDM at $\geq 3.2\sigma$ confidence level is small (0.00919 for DESI BAO + SN mock) $+ \theta_{\star}$ and 0.02109 for DES5YR SNe + BAO mock + θ_{\star} , respectively). This result indicates that the preference for dynamic dark energy is unlikely due to unknown systematics in one data set and statistical fluctuations in another.

Since both DESI BAO and DES5YR SNe are likely to be the key drivers of the preference for dynamic dark energy, it is instructive to compare the two data set in a same observable space and see if there is any signal in common. If we take r_dh as a free parameter and



FIG. 3. Binned distance moduli and $\ln H$. The Planck best-fit Λ CDM and a global difference is subtracted. The green solid line is the prediction of the DES5YR SNe + DESI BAO + θ_{\star} best-fit w_0w_a CDM model, where $w_0 = -0.76$, $w_a = -0.88$, and $\Omega_m = 0.314$.

marginalize over it, the BAO measurements of D_M/r_d can be translated to constraints on distance moduli in the same form of SNe data. The upper panel of Figure 3 shows the binned distance moduli of SNe and BAO, with the Planck best-fit Λ CDM prediction and a global difference subtracted. We do not find similar patterns in the distance-modulus chart for DES5YR SNe and DESI BAO. While the DES5YR SNe distance-modulus data clearly follow the prediction of DES5YR SNe + DESI BAO + θ_{\star} best-fit w_0w_a CDM model, the DESI BAO D_M/r_d data do not. What pushes an overall fit to w_0w_a CDM in the DESI BAO data is the anomalously large H at effective redshift $z_{\rm eff} = 0.51$ [17, 23], as shown in the lower panel of Figure 3.

IV. DISCUSSION AND CONCLUSIONS

In this work, we demonstrate that the recently claimed preference for dynamic dark energy in DESI+DES5YR+CMB does not depend on the physical details of CMB. Replacing the full CMB information with a simple and model-independent θ_{\star} constraint leads to similar results. The simplified likelihood only involves the background geometry of the universe and can therefore be evaluated with little computational resources. We then run over 10⁴ ACDM-based simulations and compare them to the real data. We find that the preference for dynamic dark energy can hardly be interpreted as unknown systematics in DESI BAO and statistical fluctuations in DES5YR SNe, or vice versa.

Although both DESI BAO and DES5YR SNe have a trend of favoring w_0w_a CDM over Λ CDM, the signals are from different observable spaces. The BAO D_M/r_d data do not have a trend to follow the pattern of the binned distance moduli of DES5YR SNe. We also check that a different SN data set Pantheon+ does not follow this pattern, neither. It is therefore yet too early to claim that the data consistently point to the dynamic dark energy model.

V. ACKNOWLEDGEMENTS

This work is supported by the National Natural Science Foundation of China (NSFC) under Grant No. 12073088, National key R&D Program of China (Grant No. 2020YFC2201600), and National SKA Program of China No. 2020SKA0110402. The computing resources are supported by Weide Liang, Mingcheng Zhu, Ren-Peng Zhou, Zhenyang Huang, Penghui Dai and Huan Zhou via the MEET-U ("Make Everyone Explore The Universe") outreach website (http://zhiqihuang.top/ MEETU), where we make the simulation source code publicly available and allow astronomy amateurs to contribute their computing resources. We are grateful to Kyle Dawson for kindly sharing the data used in the DESI 2024 release.

- A. G. Riess, A. V. Filippenko, P. Challis, *et al.*, Astronomical Journal **116**, 1009 (1998), arXiv:astroph/9805201 [astro-ph].
- [2] D. J. Eisenstein, I. Zehavi, D. W. Hogg, *et al.*, Astrophys. J. **633**, 560 (2005), arXiv:astro-ph/0501171 [astro-ph].
- [3] S. Raghunathan, S. Patil, E. Baxter, B. A. Benson, et al., Phys. Rev. Lett. **123**, 181301 (2019), arXiv:1907.08605 [astro-ph.CO].
- [4] N. Aghanim, Y. Akrami, M. Ashdown, *et al.*, Astronomy & Astrophysics **641**, A6 (2020), arXiv:1807.06209 [astroph.CO].
- [5] S. Alam, M. Aubert, S. Avila, et al., Phys. Rev. D 103, 083533 (2021), arXiv:2007.08991 [astro-ph.CO].

- [6] J. Hou, A. G. Sánchez, A. J. Ross, et al., MNRAS 500, 1201 (2021), arXiv:2007.08998 [astro-ph.CO].
- [7] T. M. C. Abbott, M. Aguena, A. Alarcon, *et al.*, Phys. Rev. D **105**, 023520 (2022), arXiv:2105.13549 [astroph.CO].
- [8] M. S. Madhavacheril, F. J. Qu, B. D. Sherwin, *et al.*, Astrophys. J. **962**, 113 (2024), arXiv:2304.05203 [astroph.CO].
- [9] A. G. Riess, W. Yuan, L. M. Macri, *et al.*, Astrophysical Journal Letters **934**, L7 (2022), arXiv:2112.04510 [astroph.CO].
- [10] M. Asgari, C.-A. Lin, B. Joachimi, et al., Astronomy & Astrophysics 645, A104 (2021), arXiv:2007.15633 [astro-

ph.CO].

- [11] L. Perivolaropoulos and F. Skara, New Astronomy Reviews 95, 101659 (2022), arXiv:2105.05208 [astroph.CO].
- [12] J.-C. Wang, Z.-Q. Huang, L. Huang, and J. Liu, Research in Astronomy and Astrophysics 24, 045001 (2024), arXiv:2311.02866 [astro-ph.CO].
- [13] G.-B. Zhao, M. Raveri, L. Pogosian, et al., Nature Astronomy 1, 627 (2017), arXiv:1701.08165 [astro-ph.CO].
- [14] X.-D. Li, C. G. Sabiu, C. Park, Y. Wang, G.-b. Zhao, H. Park, A. Shafieloo, J. Kim, and S. E. Hong, Astrophys. J. 856, 88 (2018), arXiv:1803.01851 [astro-ph.CO].
- [15] J. Solà Peracaula, J. de Cruz Pérez, and A. Gómez-Valent, EPL (Europhysics Letters) 121, 39001 (2018).
- [16] D. Wang, Phys. Rev. D 106, 063515 (2022), arXiv:2207.07164 [astro-ph.CO].
- [17] A. G. Adame, J. Aguilar, S. Ahlen, et al., arXiv e-prints, arXiv:2404.03002 (2024), arXiv:2404.03002 [astro-ph.CO].
- [18] T. M. C. Abbott, M. Acevedo, M. Aguena, *et al.*, arXiv e-prints , arXiv:2401.02929 (2024), arXiv:2401.02929 [astro-ph.CO].
- [19] M. Chevallier and D. Polarski, Int. J. Mod. Phys. D 10, 213 (2001), arXiv:gr-qc/0009008.
- [20] E. V. Linder, Phys. Rev. Lett. 90, 091301 (2003), arXiv:astro-ph/0208512.

- [21] D. Rubin, G. Aldering, M. Betoule, *et al.*, arXiv e-prints , arXiv:2311.12098 (2023), arXiv:2311.12098 [astro-ph.CO].
- [22] D. Brout, D. Scolnic, B. Popovic, et al., Astrophys. J.
 938, 110 (2022), arXiv:2202.04077 [astro-ph.CO].
- [23] E. Ó. Colgáin, M. G. Dainotti, S. Capozziello, S. Pourojaghi, M. M. Sheikh-Jabbari, and D. Stojkovic, arXiv e-prints, arXiv:2404.08633 (2024), arXiv:2404.08633 [astro-ph.CO].
- [24] D. Wang, arXiv e-prints , arXiv:2404.13833 (2024), arXiv:2404.13833 [astro-ph.CO].
- [25] J.-Q. Liu, Z.-Q. Huang, and Y. Su, Research in Astronomy and Astrophysics 24, 045002 (2024), arXiv:2311.09703 [astro-ph.CO].
- [26] S. Brieden, H. Gil-Marín, and L. Verde, JCAP 2023, 023 (2023), arXiv:2212.04522 [astro-ph.CO].
- [27] N. Schöneberg, arXiv e-prints, arXiv:2401.15054 (2024), arXiv:2401.15054 [astro-ph.CO].
- [28] Z. Ding, H.-J. Seo, Z. Vlah, Y. Feng, M. Schmittfull, and F. Beutler, MNRAS **479**, 1021 (2018), arXiv:1708.01297 [astro-ph.CO].
- [29] S.-F. Chen, C. Howlett, M. White, *et al.*, arXiv e-prints , arXiv:2402.14070 (2024), arXiv:2402.14070 [astro-ph.CO].
- [30] J. Wang, Z. Huang, and L. Huang, Science China Physics, Mechanics, and Astronomy 66, 129511 (2023), arXiv:2303.15267 [astro-ph.CO].
- [31] Y. Yao, Y. Wang, and H. Miao, Science China Physics, Mechanics, and Astronomy 67, 219533 (2024).