

HARPER'S BEYOND SQUARE-ROOT CONJECTURE

VICTOR Y. WANG AND MAX WENQIANG XU

ABSTRACT. We explain how the (shifted) Ratios Conjecture for $L(s, \chi)$ would extend a randomization argument of Harper from a conductor-limited range to an unlimited range of “beyond square-root cancellation” for character twists of the Liouville function. As a corollary, the Liouville function would have nontrivial cancellation in arithmetic progressions of modulus just exceeding the well-known square-root barrier. Morally, the paper passes from random matrices to random multiplicative functions.

1. INTRODUCTION

Studying character sums is a classical and central topic in number theory. Recently, in [18], Harper established a rather striking new phenomenon that the typical character sums have better than square-root cancellation, which can be viewed as a successful de-randomized version of his earlier celebrated result in the random setting [17]. Precisely, he [18, Theorem 1.1] showed that if $1 \leq x \leq r$ and $\min(x, r/x) \rightarrow +\infty$, then

$$\mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} \chi(n) \right| = o(\sqrt{x}),$$

where \mathbb{E}_χ denotes $\frac{1}{r-1} \sum_\chi$, and the summation is over all Dirichlet characters of a given modulus r , where r is a prime for convenience. Moreover, if $c \in \{\mu, \lambda\}$ is either the *Möbius function* μ , or the closely related *Liouville function* λ , he also established that

$$\mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} c(n) \chi(n) \right| = o(\sqrt{x}) \quad (1.1)$$

for the same range of x [18, Theorem 3]. As noted in [18, paragraph after Theorem 3], the same phenomenon could also be proven for the continuous character family $\chi = n^{it}$. That is, if $1 \leq x \leq T$ and $\min(x, T/x) \rightarrow +\infty$, then

$$\frac{1}{T} \int_0^T \left| \sum_{1 \leq n \leq x} c(n) n^{it} \right| dt = o(\sqrt{x}).$$

The restrictions $x \leq r$ and $x \leq T$ in the above results naturally appeared in Harper's proof, where he needed such a restriction to perform a clever perfect orthogonality trick. However, he believed that the same sort of result should also hold for a wider range of x , even though perfect orthogonality itself no longer holds. This is Conjecture 1.1:

2020 *Mathematics Subject Classification.* Primary 11L40; Secondary 11K65, 11M06, 11M50, 11N37.

Key words and phrases. Classical L -functions, random matrices, random multiplicative functions.

Conjecture 1.1 (Harper). *Fix $c \in \{\mu, \lambda\}$. Let A be a fixed positive constant. Then*

$$\mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} c(n) \chi(n) \right| = o(\sqrt{x}) \quad (1.2)$$

as $x \rightarrow +\infty$, provided $1 \leq x \leq r^A$. Similarly, if $1 \leq x \leq T^A$ and $x \rightarrow +\infty$, then

$$\frac{1}{T} \int_0^T \left| \sum_{1 \leq n \leq x} c(n) n^{it} \right| dt = o(\sqrt{x}). \quad (1.3)$$

Moreover, in both (1.2) and (1.3), the quantitative upper bound $O\left(\frac{\sqrt{x}}{(\log \log x)^{1/4}}\right)$ holds.

This conjecture is due to Harper [18, (1.2)], though strictly speaking he only records the conjecture in the Möbius case $c = \mu$. Given c , we refer to (1.2) and (1.3) as the *discrete* and *continuous* cases, respectively. We concentrate on $c = \lambda$ and discrete χ , which allows for the most elegant treatment out of all four possible cases. We expect that with more technical work, the ideas involved would extend to the other three cases, and to other families such as quadratic Dirichlet characters (when $c = \mu$).

Theorem 1.2. *Let $c = \lambda$. In the discrete case (1.2), Conjecture 1.1 is true under GRH and the Ratios Conjecture [9, (5.6)] for the Dirichlet L -functions $L(s, \chi)$.*

We will prove this in a stronger form, in Theorem 3.1. Conjecture 2.1 is the precise form of the Ratios Conjecture that we use. See [8–10] for details on the general Moments and Ratios Conjectures, and their rich history based on random matrices.

It is natural to hope that GRH alone might suffice, by adapting techniques of [16, 30, 32] from moments to ratios. See [5, 6, 11], and references within, for progress in this direction. It would be interesting to pursue this further. Let us also mention that over function fields $\mathbb{F}_q(t)$, there is now enough progress on the Ratios Conjecture that one can likely prove an unconditional version of Theorem 1.2 in one or more families for $q \gg_A 1$. See [2, 26, 29, 33], and references within, for some relevant developments on ratios.

We believe Theorem 1.2 is interesting and surprising in its own right, but there is also a significant application at $A \approx 2$. Harper [18] pointed out that by Perron’s formula, one can use the continuous case (1.3) of Conjecture 1.1 for $c = \mu$ to establish nontrivial cancellation of $\mu(n)$ in short intervals breaking the “square-root barrier”. Precisely, this means

$$\sum_{x \leq n \leq x+y} \mu(n) = o(y) \quad (1.4)$$

for all x and y as long as $y \gg \sqrt{x}/W(x)$ for some $W(x) \rightarrow +\infty$ as $x \rightarrow +\infty$. It is known that the Riemann Hypothesis (RH) implies (1.4) if $y \gg x^{1/2+\varepsilon}$; in fact, $y \gg x^{1/2} e^{(\log x)^{1/2+\varepsilon}}$ suffices, by [31, Theorem 1] of Soundararajan. The best unconditional result [22], due to Matomäki and Teräväinen, gives an exponent around 0.55. We also remark that a key feature here is that we require (1.4) to hold for “all” short intervals with length beyond certain threshold. If we only require “almost all”, then celebrated work of Matomäki–Radziwiłł [21] shows that it is sufficient to only require $y \rightarrow +\infty$.

The situation for arithmetic progressions $a \bmod r$ is similar under the Generalized Riemann Hypothesis (GRH), by Ye [35], but more complicated unconditionally, due to uniformity issues near the 1-line. To our knowledge, there is no unconditional analog of (1.4) available when $r = x^{0.01}$, but results are available for $r = x^{o(1)}$ (see [1] for instance). This is loosely related to the limited range $r \ll (\log x)^A$ in the Siegel–Walfisz Theorem. However, we note that if one allows to take an average over r , then the Bombieri–Vinogradov large sieve inequality is as good as what the GRH gives. We refer the reader to some recent developments by Maynard [23–25] on various special cases where better results can be obtained when averaging r and references therein. Other related results beyond the square-root range are available in works such as [12–14].

For simplicity and clarity, we conditionally break the “square-root barrier” in the arithmetic-progression analog of (1.4) for $c = \lambda$, as a corollary of Theorem 1.2.

Corollary 1.3. *Under the same assumptions as in Theorem 1.2, we have as $x \rightarrow +\infty$*

$$\sum_{\substack{1 \leq n \leq x \\ n \equiv a \bmod r}} \lambda(n) = o\left(\frac{x}{r}\right)$$

for $1 \leq a \leq r \leq W(x)\sqrt{x}$, for some function $W(x) \rightarrow +\infty$ as $x \rightarrow +\infty$.

Proof. We first prove this assuming $r \geq x^{0.49}$. By complete multiplicativity of λ , it suffices to treat the case $\gcd(a, r) = 1$. In this case, $\mathbf{1}_{n \equiv a \bmod r} = \mathbb{E}_\chi \bar{\chi}(a) \chi(n)$. Therefore,

$$\sum_{\substack{1 \leq n \leq x \\ n \equiv a \bmod r}} \lambda(n) = \mathbb{E}_\chi \bar{\chi}(a) \sum_{1 \leq n \leq x} \lambda(n) \chi(n) \ll \mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} \lambda(n) \chi(n) \right| \ll \frac{\sqrt{x}}{W(x)^{1+\varepsilon}},$$

say, by Theorem 1.2 with $A = 2.05$. This is $O\left(\frac{x}{rW(x)^\varepsilon}\right) = o\left(\frac{x}{r}\right)$, since $r \leq W(x)\sqrt{x}$.

(In fact, by Theorem 3.1, we can take $W(x) = (\log \log x)^b$ for any fixed $b < 1/4$.)

On the other hand, the range $r \leq x^{0.49}$ can be handled by GRH. \square

This application is less interesting in $\mathbb{F}_q[t]$, where stronger techniques are available due to Sawin [29]. We also note that the Ratios Conjecture can be directly used to estimate averages like $\mathbb{E}_\chi \sum_{1 \leq n \leq x} \lambda(n) \chi(n)$, but not to estimate weighted averages like $\mathbb{E}_\chi \bar{\chi}(a) \sum_{1 \leq n \leq x} \lambda(n) \chi(n)$ in general, except for special values of a like $a = 1$.

Remark 1.4. The analog of Corollary 1.3 for the short-interval case (1.4) is likely susceptible to similar methods. Conditionally on RH and the Ratios Conjecture for the continuous family of L -functions $\zeta(s + it_0)$ indexed by $t_0 \in \mathbb{R}$, one should be able to prove (1.4) for $y \gg \sqrt{x}/(\log \log x)^{1/4-\epsilon}$. It would be interesting to work out the details.

To understand how universal the phenomena above are, it would be desirable to prove a random matrix analog of Conjecture 1.1, in the $N \times N$ unitary groups $U(N)$. In this setting, partial sums $\sum_{1 \leq n \leq x} c(n) \chi(n)$ roughly correspond to characters $\text{tr}(A, \text{Sym}^k \mathbb{C}^N)$, i.e. Schur polynomials, for $k \geq 0$, associated to symmetric powers $\text{Sym}^k \mathbb{C}^N$ of the standard representation \mathbb{C}^N . Let ν_N be the Haar probability measure on $U(N)$.

Conjecture 1.5. Fix $B > 0$. If $0 \leq k \leq BN$ and $k \rightarrow +\infty$, then

$$\int_{U(N)} |\mathrm{tr}(A, \mathrm{Sym}^k \mathbb{C}^N)| d\nu_N(A) = o(1).$$

Here k is analogous to $\log x$ in Conjecture 1.1, and N is analogous to $\log r$. Note that the range of B is unrestricted, unlike in [28], which studies $\bigwedge^k \mathbb{C}^N$ instead of $\mathrm{Sym}^k \mathbb{C}^N$. The connection to Möbius, or $1/L(s, \chi)$, is roughly given by the identity

$$\frac{1}{\det(1 - e^{1/2-s} A)} = \sum_{k \geq 0} \mathrm{tr}(A, \mathrm{Sym}^k \mathbb{C}^N) e^{(1/2-s)k}.$$

Also, note that $\int_{U(N)} |\mathrm{tr}(A, \mathrm{Sym}^k \mathbb{C}^N)|^2 d\nu_N(A) = 1$, since $\mathrm{Sym}^k \mathbb{C}^N$ is irreducible. So the $o(1)$ in Conjecture 1.5 is the analog of $o(\sqrt{x})$ in Conjecture 1.1.

We leave this interesting challenge, Conjecture 1.5, open for now. A similar conjecture may hold in the symplectic case $A \in USp(2N)$, or in the orthogonal case $A \in O(N, \mathbb{R})$ after replacing $\mathrm{Sym}^k \mathbb{C}^N$ with the kernel of the contraction map $\mathrm{Sym}^k \mathbb{C}^N \rightarrow \mathrm{Sym}^{k-2} \mathbb{C}^N$.

Strategy. We sketch the proof ideas. Our proof is based on Harper's strategy in [18]. Roughly speaking, Harper made a remarkable randomization argument to show that when summing over the given family of Dirichlet characters, the typical behavior of the character sums $\sum_n \chi(n)$ is very close to the random sums of Steinhaus Random multiplicative functions (RMF) $\sum_n f(n)$. The latter is well studied in Harper's earlier paper [17] and in particular, a conjecture of Helson [19] is proved:

$$\mathbb{E} \left| \sum_{1 \leq n \leq x} f(n) \right| = o(\sqrt{x}),$$

which shows the “better than square-root” cancellation phenomenon holds.¹ Once the link between the character sums and partial sums of RMF is built, the rest of the proof in [18] closely follows the work [17] in the RMF setting.

A key obstruction in Harper's strategy above, that prevented him from establishing Conjecture 1.1 for larger x is that, in order to show that the character sums typically behave like the random sums of RMF, Harper crucially used the perfect orthogonality of the character sums. To make use of this, there is a natural restriction on the ranges of the parameters, e.g. the relation between the moduli r and the length of the sum x . Our innovation is to use a more complex-analytic approach, namely by assuming the Ratios Conjecture to avoid using the orthogonality and thus we can extend the range of the key parameters to establish Conjecture 1.1 conditionally.

However, certain auxiliary estimates and parameters in our work are more delicate than those in Harper's paper [18]. Some of the changes required are listed before Proposition 2.8 below. A key ingredient is to replace *perfect* orthogonality with *approximate* orthogonality on average. This is achieved in a new even-moment estimate, (2.9), which has a genuine error term, unlike in Harper's work. To establish this estimate, we must

¹See [7, 15, 28, 34] for further developments about this phenomenon and its universality.

carefully distinguish between $c(n)\chi(n)$ and $\chi(n)$.² We remark that a main different feature of the two cases is that, in the unweighted case $\sum_{1 \leq n \leq x} \chi(n)$, one needs $r/x \rightarrow +\infty$ (see [18, Theorems 1 and 2]) in order to get extra cancellation due to the “Fourier flip”; while in the weighted case $\sum_{1 \leq n \leq x} c(n)\chi(n)$, we do not expect such restriction required. The Ratios Conjecture ultimately provides a source of randomness for the weighted case but not present for the unweighted case.

Our work can be interpreted as saying that a *random matrix model* for $\lambda(n)\chi(n)$, namely the Ratios Conjecture, justifies the *Steinhaus model* for $\lambda(n)\chi(n)$, for low-moment upper-bound statistics of the partial sums $\sum_{1 \leq n \leq x} \lambda(n)\chi(n)$.

Notation. Our notation $\ll, \gg, \asymp, O(\cdot), o(\cdot), \lfloor \cdot \rfloor$ is standard. We let $P(n)$ denote the maximum prime divisor of n . We let $\mathbf{1}_E$ denote 1 if an event E holds, and 0 otherwise. The only other convention that deserves comment is the *Steinhaus average* $\mathbb{E}_f A(f)$, or $\mathbb{E}A(f)$ for short, which we always write with the lowercase letter f . Here f denotes a completely multiplicative function.

Usually, $A(f)$ depends only on $f(p)$ for finitely many primes p . In this case, $\mathbb{E}_f A(f)$ denotes the expected value of $A(f)$ when each $f(p)$ is drawn uniformly, independently, from $\{z \in \mathbb{C} : |z| = 1\}$. This is the only case used in Harper’s work [18].

However, in § 2, we will allow $A(f)$ to be a Dirichlet series of the form

$$A(f, \mathbf{z}, \mathbf{s}) = \sum_{(\mathbf{m}, \mathbf{n}) \in \mathbb{N}^k \times \mathbb{N}^l} \frac{a_{\mathbf{m}, \mathbf{n}}(f)}{m_1^{z_1} \cdots m_k^{z_k} n_1^{s_1} \cdots n_l^{s_l}},$$

where $a_{\mathbf{m}, \mathbf{n}}(f)$ depends only on (\mathbf{m}, \mathbf{n}) and on $f(p)$ for $p \mid m_1 \cdots m_k n_1 \cdots n_l$. Here $k, l \geq 0$, and $(\mathbf{z}, \mathbf{s}) \in \mathbb{C}^k \times \mathbb{C}^l$. Then we let

$$\mathbb{E}_f A(f, \mathbf{z}, \mathbf{s}) := \sum_{(\mathbf{m}, \mathbf{n}) \in \mathbb{N}^k \times \mathbb{N}^l} \frac{\mathbb{E}_f a_{\mathbf{m}, \mathbf{n}}(f)}{m_1^{z_1} \cdots m_k^{z_k} n_1^{s_1} \cdots n_l^{s_l}}.$$

We note that the series $\mathbb{E}_f A(f, \mathbf{z}, \mathbf{s})$ may converge absolutely in a strictly larger domain than the series $A(f, \mathbf{z}, \mathbf{s})$ does for individual f .

Acknowledgements. We thank Paul Bourgade and Kannan Soundararajan for discussions on random matrices and probability, Alexandra Florea for helpful comments on the Ratios Conjecture, and Joni Teräväinen for providing several references. We are also grateful to Alexandra Florea, Adam Harper, and Joni Teräväinen for helpful comments on earlier drafts. The first author is supported by the European Union’s Horizon 2020 research and innovation program under the Marie Skłodowska-Curie Grant Agreement No. 101034413. The second author is partially supported by the Cuthbert C. Hurd Graduate Fellowship in the Mathematical Sciences, Stanford.

²Indeed, (2.9) would be false if we had $c = 1$ instead of $c \in \{\mu, \lambda\}$.

2. CHARACTER TWISTS BEHAVE LIKE RANDOM MODEL

In this section, we establish that certain mean values of twists behave like a random model (Proposition 2.8). Instead of using perfect orthogonality, we use the Ratios Conjecture to break through the barrier for the length of the twisted sum. In order for the Ratios Conjecture to make sense, we assume GRH for $L(s, \chi)$ throughout this section. Let χ_0 be the principal character modulo r , and define $\mathbb{E}_{\chi \neq \chi_0} := \frac{1}{r-2} \sum_{\chi \neq \chi_0}$. It will be sometimes convenient to exclude χ_0 from statements, and sometimes convenient to include it, so both $\mathbb{E}_{\chi \neq \chi_0}$ and $\mathbb{E}_\chi = \frac{1}{r-1} \sum_\chi$ will appear in our work below.

For every completely multiplicative function $f(n)$, we formally let

$$L(s, f) := \sum_{n \geq 1} f(n) n^{-s} = \prod_p (1 - f(p) p^{-s})^{-1},$$

$$\sum_{n \geq 1} \mu_f(n) n^{-s} := 1/L(s, f) = \prod_p (1 - f(p) p^{-s}).$$

Since f is completely multiplicative, we have $\mu_f(n) = \mu(n)f(n)$. By definition, $\mathbb{E}_f \frac{L(z_1, f)L(z_2, \bar{f})}{L(s_1, f)L(s_2, \bar{f})}$ is the meromorphic function obtained by formally expanding

$$\frac{L(z_1, f)L(z_2, \bar{f})}{L(s_1, f)L(s_2, \bar{f})} = \sum_{m_1, m_2, n_1, n_2 \geq 1} \frac{f(m_1)\bar{f}(m_2)\mu_f(n_1)\bar{\mu}_f(n_2)}{m_1^{z_1} m_2^{z_2} n_1^{s_1} n_2^{s_2}}$$

as a 4-variable Dirichlet series, and applying the average \mathbb{E}_f to each coefficient. In terms of the meromorphic function G_ζ defined in [9, (5.10)], we have

$$\mathbb{E}_f \frac{L(z_1, f)L(z_2, \bar{f})}{L(s_1, f)L(s_2, \bar{f})} = G_\zeta(z_1 - \tfrac{1}{2}; z_2 - \tfrac{1}{2}; s_1 - \tfrac{1}{2}; s_2 - \tfrac{1}{2}). \quad (2.1)$$

For the reader's convenience, we record the definition of G_ζ here:

$$G_\zeta(\alpha; \beta; \gamma; \delta) := \sum_{\substack{m_1, m_2, n_1, n_2 \geq 1 \\ m_1 n_1 = m_2 n_2}} \frac{\mu(n_1)\mu(n_2)}{m_1^{1/2+\alpha} m_2^{1/2+\beta} n_1^{1/2+\gamma} n_2^{1/2+\delta}},$$

where the variables m_1, m_2, n_1, n_2 in our notation correspond respectively to m, n, h, j in the notation of [9, (5.10)]. The series G_ζ converges absolutely on the region

$$\operatorname{Re}(\alpha), \operatorname{Re}(\beta), \operatorname{Re}(\gamma), \operatorname{Re}(\delta) > 0,$$

for instance, by a short calculation with the divisor bound. However, G_ζ is meromorphic on a larger region, as is discussed in [9, 10] in great depth.

A special case of the Ratios Conjecture [9, (5.6)] for $L(s, \chi)$ is the following:³

³To avoid smoothing issues, which are orthogonal to the main point of the paper, we put $O_\varepsilon((1+T)^\varepsilon)$ in the error term. Morally, $O((1+T)^{O(1)})$ should be enough to say something interesting, but might require smoothing the sums over $n \leq x$ in Conjecture 1.1. Cf. the situation of [3, Theorem 1].

Conjecture 2.1. *Let $\operatorname{Re}(z_1) = \operatorname{Re}(z_2) = \frac{1}{2}$ and $\operatorname{Re}(s_1) = \operatorname{Re}(s_2) = \frac{1}{2} + \varepsilon$. Let $T := \max(|\operatorname{Im}(z_1)|, |\operatorname{Im}(z_2)|, |\operatorname{Im}(s_1)|, |\operatorname{Im}(s_2)|)$. Then for some absolute constant $\omega \in (0, \frac{1}{2}]$ (independent of r and ε), we have for all sufficiently small $\varepsilon > 0$*

$$\mathbb{E}_{\chi \neq \chi_0} \frac{L(z_1, \chi)L(z_2, \bar{\chi})}{L(s_1, \chi)L(s_2, \bar{\chi})} = \mathbf{MT} + \frac{O_\varepsilon((1+T)^\varepsilon)}{r^\omega},$$

where

$$\mathbf{MT} := \mathbb{E}_f \frac{L(z_1, f)L(z_2, \bar{f})}{L(s_1, f)L(s_2, \bar{f})} + \frac{H}{r^{z_1+z_2-1}} \mathbb{E}_f \frac{L(1-z_2, f)L(1-z_1, \bar{f})}{L(s_1, f)L(s_2, \bar{f})},$$

$$\text{where } H := \frac{(2\pi)^{z_1+z_2}}{\pi^2} \Gamma(1-z_1)\Gamma(1-z_2) \frac{\sum_{a \in \{0,1\}} \sin\left(\frac{\pi}{2}(1-z_1+a)\right) \sin\left(\frac{\pi}{2}(1-z_2+a)\right)}{2}.$$

Derivation via the Ratios Recipe. In addition to the general recipe [9, (5.6)], see [8, § 4.3] for some details on the family $L(s, \chi)$. Write $\chi(-1) = (-1)^a$ with $a \in \{0, 1\}$. Let $\tau(\chi) := \sum_{1 \leq x \leq r} \chi(x) e^{2\pi i x/r}$. It is known that

$$L(s, \chi) = w_\chi X_\chi(s) L(1-s, \bar{\chi}),$$

where $w_\chi := \frac{\tau(\chi)}{i^a r^{1/2}}$ and $X_\chi(s) := 2^s \pi^{s-1} r^{1/2-s} \sin\left(\frac{\pi}{2}(s+a)\right) \Gamma(1-s)$. Moreover, by standard properties of Gauss sums, we have $w_{\bar{\chi}} = \overline{w_\chi} = w_\chi^{-1}$. On multiplying out

$$\frac{\left(L(z_1, \chi) + w_\chi X_\chi(z_1) L(1-z_1, \bar{\chi})\right) \left(L(z_2, \bar{\chi}) + w_{\bar{\chi}} X_{\bar{\chi}}(z_2) L(1-z_2, \chi)\right)}{L(s_1, \chi) L(s_2, \bar{\chi})}$$

and formally averaging over χ , as described in [9, (5.6)], with cancellation over root numbers as in [8, § 4.3], we get the desired Ratios Conjecture. \square

Conjecture 2.1 is an asymptotic for $T \leq r^{0.99\omega/\varepsilon}$, for any $\varepsilon > 0$, whereas for Theorem 1.2 it would actually suffice to have an asymptotic for $T \leq r^{A/2+0.01}$.⁴ However, we do not wish to optimize the T -aspect at all in this paper. Similarly, it could be interesting to weaken the saving r^ω required in the asymptotic, along the lines of [4, Conjecture 3.6 (R2)] for instance. However, our assumption that ω is independent of r and ε is very convenient.

The quantity $H = H(z_1, z_2)$ is independent of r , and can be bounded as follows.

Lemma 2.2. *The function H is holomorphic for $\operatorname{Re}(z_1), \operatorname{Re}(z_2) < 1$. Moreover, if $\operatorname{Re}(z_1), \operatorname{Re}(z_2) \in \{\frac{1}{2} - \delta, \frac{1}{2} + \delta\}$ with $\delta \in [0, \frac{1}{2})$, then $H \ll_\delta (1 + |\operatorname{Im}(z_1)|)^\delta (1 + |\operatorname{Im}(z_2)|)^\delta$.*

Proof. The first part is clear since Γ is holomorphic away from $\mathbb{Z} \setminus \mathbb{N}$. The second formula follows from Stirling's bound $\Gamma(s) \ll_\delta (1 + |\operatorname{Im}(s)|)^{\operatorname{Re}(s)-1/2} e^{-\frac{\pi}{2}|\operatorname{Im}(s)|}$, a consequence of [20, (5.113)], on $\operatorname{Re}(s) \in \{\frac{1}{2} - \delta, \frac{1}{2} + \delta\}$. Note that $\sin\left(\frac{\pi}{2}(s+a)\right) \ll_\delta e^{\frac{\pi}{2}|\operatorname{Im}(s)|}$ for $a \in \{0, 1\}$, and the exponential growth here cancels out the exponential decay factor for $\Gamma(s)$. \square

The usual Ratios Conjecture implies a *twisted* Ratios Conjecture:

⁴We actually only really apply Conjecture 2.1 with $\varepsilon \asymp \omega/A$.

Conjecture 2.3. *Let $\operatorname{Re}(s_1) = \operatorname{Re}(s_2) = \frac{1}{2} + \varepsilon$ and $1 \leq m_1, m_2 \leq r^h$. Let $T := \max(|\operatorname{Im}(s_1)|, |\operatorname{Im}(s_2)|)$. If $\varepsilon, h > 0$ are sufficiently small, then for some absolute constant $\omega' \in (0, \frac{1}{2}]$ (independent of r, ε, h), we have*

$$\mathbb{E}_{\chi \neq \chi_0} \frac{\chi(m_1) \overline{\chi}(m_2)}{L(s_1, \chi) L(s_2, \overline{\chi})} = \mathbb{E}_f \frac{f(m_1) \overline{f}(m_2)}{L(s_1, f) L(s_2, \overline{f})} + O_{\varepsilon, h}((1+T)^\varepsilon r^{-\omega'}).$$

Proof assuming the Ratios Conjecture 2.1. We first discuss the convergence regions for various quantities that appear in the proof, as preparation for subsequent contour-shifting arguments. By (2.1), we have

$$\mathbb{E}_f \frac{L(1-z_2, f) L(1-z_1, \overline{f})}{L(s_1, f) L(s_2, \overline{f})} = G_\zeta(\tfrac{1}{2} - z_2; \tfrac{1}{2} - z_1; s_1 - \tfrac{1}{2}; s_2 - \tfrac{1}{2}). \quad (2.2)$$

However, in the notation of [9, (5.12)–(5.14)], we have $G_\zeta(\alpha; \beta; \gamma; \delta) = A_\zeta(\alpha; \beta; \gamma; \delta) Y_U$, where A_ζ is an explicit Euler product known to be absolutely convergent on the region $|\operatorname{Re}(\alpha)|, \dots, |\operatorname{Re}(\delta)| < \frac{1}{4}$ (see [10, Remark 2.3]), and where

$$Y_U := \frac{\zeta(1+\alpha+\beta)\zeta(1+\gamma+\delta)}{\zeta(1+\alpha+\delta)\zeta(1+\beta+\gamma)}. \quad (2.3)$$

Since $\operatorname{Re}(s_1), \operatorname{Re}(s_2) > \frac{1}{2}$, it follows that the main term of Conjecture 2.1,

$$\mathbf{MT} = G_\zeta(z_1 - \tfrac{1}{2}; z_2 - \tfrac{1}{2}; s_1 - \tfrac{1}{2}; s_2 - \tfrac{1}{2}) + \frac{H}{r^{z_1+z_2-1}} G_\zeta(\tfrac{1}{2} - z_2; \tfrac{1}{2} - z_1; s_1 - \tfrac{1}{2}; s_2 - \tfrac{1}{2}),$$

is holomorphic for $\operatorname{Re}(z_1), \operatorname{Re}(z_2) \in (\frac{1}{4}, \frac{3}{4})$, and that $\mathbb{E}_f \frac{L(z_1, f) L(z_2, \overline{f})}{L(s_1, f) L(s_2, \overline{f})}$ itself is holomorphic for $\operatorname{Re}(z_1), \operatorname{Re}(z_2) > \frac{1}{2}$. We note that in \mathbf{MT} , there is a now-familiar cancellation of poles, observed by [8] in the case of moments and by [9, 10] in the case of ratios.

We now do a contour integral to extract $m_1^{-z_1} m_2^{-z_2}$ terms. Let $g_0(x)$ be a bump function supported on $(-\frac{1}{2}, \frac{1}{2})$, with $g_0(0) = 1$, and let $g_m(x) := g_0(x - m)$. The Mellin transform $g_m^\vee(z) := \int_0^\infty g_m(x) x^{z-1} dx$ satisfies the bound $g_m^\vee(z) \ll_B m^{\operatorname{Re}(z)} \min(1, m/|\operatorname{Im}(z)|)^B$, by repeated integration by parts in x if $m/|\operatorname{Im}(z)| > 1$. By Mellin inversion,

$$\iint_{2-i\infty}^{2+i\infty} \frac{\prod_{j=1}^2 g_{m_j}^\vee(z_j)}{(2\pi i)^2} \mathbb{E}_f \frac{L(z_1, f) L(z_2, \overline{f})}{L(s_1, f) L(s_2, \overline{f})} dz_1 dz_2 = \mathbb{E}_f \frac{f(m_1) \overline{f}(m_2)}{L(s_1, f) L(s_2, \overline{f})}, \quad (2.4)$$

since both sides equal the $m_1^{-z_1} m_2^{-z_2}$ coefficient of $\mathbb{E}_f \frac{L(z_1, f) L(z_2, \overline{f})}{L(s_1, f) L(s_2, \overline{f})}$. Indeed, (2.4) is an identity of holomorphic functions on $\operatorname{Re}(s_1), \operatorname{Re}(s_2) > \frac{1}{2}$, and can thus be checked for $\operatorname{Re}(s_1), \operatorname{Re}(s_2) \geq 2$, say, where all convergence issues become trivial.

The deterministic side requires more care, due to the shape of \mathbf{MT} . By Mellin inversion,

$$\mathbb{E}_{\chi \neq \chi_0} \frac{\chi(m_1) \overline{\chi}(m_2)}{L(s_1, \chi) L(s_2, \overline{\chi})} = \mathbb{E}_{\chi \neq \chi_0} \iint_{2-i\infty}^{2+i\infty} \frac{\prod_{j=1}^2 g_{m_j}^\vee(z_j)}{(2\pi i)^2} \frac{L(z_1, \chi) L(z_2, \overline{\chi})}{L(s_1, \chi) L(s_2, \overline{\chi})} dz_1 dz_2;$$

this is absolutely convergent, by GRH, so by Fubini we may move \mathbb{E}_χ inside the integral. Now, we shift the integral first to $\operatorname{Re}(z_1) = \operatorname{Re}(z_2) = \frac{1}{2}$, to apply Conjecture 2.1, then

shift the main term MT to $\operatorname{Re}(z_1) = \operatorname{Re}(z_2) = \frac{1}{2} + \frac{1}{50}$, say (this is possible by the first paragraph of the proof). This leads to the estimate

$$\begin{aligned} & \mathbb{E}_{\chi \neq \chi_0} \frac{\chi(m_1)\overline{\chi}(m_2)}{L(s_1, \chi)L(s_2, \overline{\chi})} - \iint_{2-i\infty}^{2+i\infty} \frac{\prod_{j=1}^2 g_{m_j}^\vee(z_j)}{(2\pi i)^2} \mathbb{E}_f \frac{L(z_1, f)L(z_2, \overline{f})}{L(s_1, f)L(s_2, \overline{f})} dz_1 dz_2 \\ & \ll_{B, \varepsilon} \iint_{\frac{1}{2} + \frac{1}{50} - i\infty}^{\frac{1}{2} + \frac{1}{50} + i\infty} \frac{(1+T)^\varepsilon \prod_{j=1}^2 (1 + |\operatorname{Im}(z_j)|)^{\frac{1}{50} + \varepsilon} m_j^{\operatorname{Re}(z_j)} \min(1, \frac{m_j}{|\operatorname{Im}(z_j)|})^B |dz_j|}{\min(r^\omega, r^{\operatorname{Re}(z_1) + \operatorname{Re}(z_2) - 1})} \end{aligned} \quad (2.5)$$

by (2.2) and (2.3), because for $\operatorname{Re}(z_1) = \operatorname{Re}(z_2) = \frac{1}{2} + \frac{1}{50}$, we have (by Lemma 2.2)

$$H \ll (1 + |\operatorname{Im}(z_1)|)^{\frac{1}{50}} (1 + |\operatorname{Im}(z_2)|)^{\frac{1}{50}}$$

and (by using RH near the 1-line to bound the ζ and $1/\zeta$ factors in Y_U)

$$Y_U(\tfrac{1}{2} - z_2; \tfrac{1}{2} - z_1; s_1 - \tfrac{1}{2}; s_2 - \tfrac{1}{2}) \ll_\varepsilon (1 + |\operatorname{Im}(z_1)|)^\varepsilon (1 + |\operatorname{Im}(z_2)|)^\varepsilon (1 + T)^\varepsilon.$$

We note that the right-hand side of (2.5) takes into account both the error term $(1 + T)^\varepsilon / r^\omega$ from Conjecture 2.1, and the second (or “dual”) term in MT.

Taking $B = 2$, and integrating over z_j (the dominant contribution coming from $|\operatorname{Im}(z_j)| \asymp m_j$), we find that the right-hand side of the inequality (2.5) is

$$\ll \frac{(1+T)^\varepsilon}{\min(r^\omega, r^{\frac{1}{50} + \frac{1}{50}})} \prod_{1 \leq j \leq 2} m_j^{1 + \frac{1}{50} + \varepsilon + \frac{1}{2} + \frac{1}{50}} \leq \frac{(1+T)^\varepsilon r^{4\hbar}}{\min(r^\omega, r^{\frac{1}{25}})} \leq \frac{(1+T)^\varepsilon}{r^{\omega'}},$$

for suitable $\omega' > 0$, provided ε and \hbar are sufficiently small. Conjecture 2.3 follows, upon plugging (2.4) into (2.5). \square

Next, we pass to a version for λ . Assume Conjecture 2.3 for the rest of the section.

Proposition 2.4. *Assume Conjecture 2.3. Let $\operatorname{Re}(s_1) = \operatorname{Re}(s_2) = \frac{1}{2} + \varepsilon$ and $1 \leq m_1, m_2 \leq r^\hbar$. Let $T := \max(|\operatorname{Im}(s_1)|, |\operatorname{Im}(s_2)|)$ and $L^\flat(s, \psi) := L(s, \psi)/L(2s, \psi^2)$.⁵ If $\varepsilon, \hbar > 0$ are sufficiently small, then for some absolute constant $\eta \in (0, \omega']$ (independent of r, ε, \hbar), we have*

$$\mathbb{E}_{\chi \neq \chi_0} \frac{\chi(m_1)\overline{\chi}(m_2)}{L^\flat(s_1, \chi)L^\flat(s_2, \overline{\chi})} = \mathbb{E}_f \frac{f(m_1)\overline{f}(m_2)}{L^\flat(s_1, f)L^\flat(s_2, \overline{f})} + O_{\varepsilon, \hbar}((1+T)^\varepsilon r^{-\eta}).$$

Proof. Let η be small in terms of ω' . Let ε and \hbar be small in terms of η . Each χ^2 is non-principal, since r is assumed to be prime and we may assume $r > 2$. Let $M \geq 1$ be a parameter. Fix $\kappa > 0$ small in terms of ε . By GRH, $\sum_{d \leq N} \chi^2(d) d^{-2it_j} \ll (1 + |t_j|)^\kappa r^\kappa \cdot N^{1/2 + \kappa}$ for all $N \geq 1$, since the vertically shifted L -function $L(s + 2t_j, \chi^2)$ has analytic conductor $\ll (1 + |t_j|)r$ in the sense of [20, Chapter 5]. By partial summation over d , we conclude that $L(2s_j, \chi^2) - \sum_{d \leq M} \frac{\chi^2(d)}{d^{2s_j}} = \sum_{d > M} \frac{\chi^2(d)}{d^{2s_j}} \ll \frac{(1+T)^\kappa r^\kappa}{M^{\operatorname{Re}(2s_j)}} \cdot M^{1/2 + \kappa}$ for $j \in \{1, 2\}$.

⁵This definition ensures that $1/L^\flat(s, \psi) = \sum_{n \geq 1} \lambda(n)\psi(n)$ for any completely multiplicative ψ .

Also, $\sum_{d \leq M} \frac{\chi^2(d)}{d^{2s}} \ll_\varepsilon 1$ trivially. It follows that, under GRH,

$$L(2s_1, \chi^2)L(2s_2, \chi^2) - \sum_{d_1, d_2 \leq M} \frac{\chi(d_1^2)\bar{\chi}(d_2^2)}{d_1^{2s_1}d_2^{2s_2}} \ll 1 \cdot \frac{(1+T)^\kappa r^\kappa}{M^{1/2}} + \left(\frac{(1+T)^\kappa r^\kappa}{M^{1/2}} \right)^2.$$

Multiplying both sides by $\chi(m_1)\bar{\chi}(m_2)L(s_1, \chi)^{-1}L(s_2, \chi)^{-1}$ and using the GRH bound $1/L(s, \chi) \ll (1+T)^\kappa r^\kappa$, we get

$$\mathbb{E}_{\chi \neq \chi_0} \frac{\chi(m_1)\bar{\chi}(m_2)}{L^b(s_1, \chi)L^b(s_2, \bar{\chi})} - \sum_{d_1, d_2 \leq M} \mathbb{E}_{\chi \neq \chi_0} \frac{\chi(d_1^2 m_1)\bar{\chi}(d_2^2 m_2)}{d_1^{2s_1}d_2^{2s_2}L(s_1, \chi)L(s_2, \bar{\chi})} \ll \frac{(1+T)^{4\kappa} r^{4\kappa}}{M^{1/2}}.$$

If $M \leq r^{3\eta}$, then on plugging in Conjecture 2.3 (with $6\eta + \hbar$ in place of \hbar) and summing over $d_1, d_2 \leq M$, we get

$$\begin{aligned} & \mathbb{E}_{\chi \neq \chi_0} \frac{\chi(m_1)\bar{\chi}(m_2)}{L^b(s_1, \chi)L^b(s_2, \bar{\chi})} - \sum_{d_1, d_2 \leq M} \mathbb{E}_f \frac{f(d_1^2 m_1)\bar{f}(d_2^2 m_2)}{d_1^{2s_1}d_2^{2s_2}L(s_1, f)L(s_2, \bar{f})} \\ & \ll \frac{(1+T)^{4\kappa} r^{4\kappa}}{M^{1/2}} + \frac{(1+T)^\varepsilon}{r^{\omega'}} \ll \frac{(1+T)^\varepsilon r^\varepsilon}{M^{1/2}}. \end{aligned} \quad (2.6)$$

For any prime $r' \geq r$, subtracting (2.6) for r and r' , with $M := r^{3\eta}$, gives

$$(\mathbb{E}_{\chi \neq \chi_0}^{[r]} - \mathbb{E}_{\chi \neq \chi_0}^{[r']}) \frac{\chi(m_1)\bar{\chi}(m_2)}{L^b(s_1, \chi)L^b(s_2, \bar{\chi})} \ll \frac{(1+T)^\varepsilon (r')^\varepsilon}{r^{3\eta/2}},$$

where the superscripts $[r]$ and $[r']$ indicate that χ has modulus r and r' , respectively. Iterating this over an infinite sequence of primes r_0, r_1, r_2, \dots with $r_0 = r$ and $r_{i+1} \in [2r_i, 4r_i]$, as is possible by Bertrand's postulate, we find by telescoping that the sequence of quantities $\mathbb{E}_{\chi \neq \chi_0}^{[r_i]} \frac{\chi(m_1)\bar{\chi}(m_2)}{L^b(s_1, \chi)L^b(s_2, \bar{\chi})}$ forms a Cauchy sequence, and moreover

$$(\mathbb{E}_{\chi \neq \chi_0}^{[r]} - \lim_{i \rightarrow \infty} \mathbb{E}_{\chi \neq \chi_0}^{[r_i]}) \frac{\chi(m_1)\bar{\chi}(m_2)}{L^b(s_1, \chi)L^b(s_2, \bar{\chi})} \ll \frac{1}{1 - 2^{\varepsilon - 3\eta/2}} \frac{(1+T)^\varepsilon (4r)^\varepsilon}{r^{3\eta/2}}. \quad (2.7)$$

In particular, $\lim_{i \rightarrow \infty} \mathbb{E}_{\chi \neq \chi_0}^{[r_i]} \frac{\chi(m_1)\bar{\chi}(m_2)}{L^b(s_1, \chi)L^b(s_2, \bar{\chi})}$ exists, so taking $r \rightarrow \infty$ in (2.6) with $M := r^{3\eta}$ gives

$$\begin{aligned} \lim_{r \rightarrow \infty} \mathbb{E}_{\chi \neq \chi_0}^{[r]} \frac{\chi(m_1)\bar{\chi}(m_2)}{L^b(s_1, \chi)L^b(s_2, \bar{\chi})} &= \sum_{d_1, d_2 \geq 1} \mathbb{E}_f \frac{f(d_1^2 m_1)\bar{f}(d_2^2 m_2)}{d_1^{2s_1}d_2^{2s_2}L(s_1, f)L(s_2, \bar{f})} \\ &= \sum_{\substack{d_1, d_2, n_1, n_2 \geq 1 \\ d_1^2 m_1 n_1 = d_2^2 m_2 n_2}} \frac{\mu(n_1)\mu(n_2)}{d_1^{2s_1}d_2^{2s_2}n_1^{s_1}n_2^{s_2}} \\ &= \mathbb{E}_f \frac{f(m_1)\bar{f}(m_2)}{L^b(s_1, f)L^b(s_2, \bar{f})}, \end{aligned} \quad (2.8)$$

since the series over $d_1, d_2, n_1, n_2 \geq 1$ in (2.8) is absolutely convergent.

Plugging (2.8) into (2.7) yields Proposition 2.4. \square

We now prove an unrestricted version of [18, Lemma 1]. As we remarked in the introduction, one may think of (2.9) as an approximate orthogonality relation on average.

Lemma 2.5 (Approximate orthogonality and even-moment estimate). *Let $c \in \{\mu, \lambda\}$. Let \mathcal{P} be a finite nonempty set of primes. Let $\mathcal{Q} := \mathcal{P} \cup \{p^2 : p \in \mathcal{P}\}$ and $U := \max(\mathcal{Q})$. Let $Q(f) := \sum_{q \in \mathcal{Q}} q^{-1/2} a(q) f(q)$ where $a(q)$ are any complex numbers. Assume $U^k \leq r^\varepsilon$ and $1 \leq x \leq r^A$, where $A > 0$ is fixed.⁶ Assume Conjecture 2.3. Let η be as in the statement of Proposition 2.4. Then*

$$\mathbb{E}_\chi \chi(m_1) \overline{\chi}(m_2) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 - \mathbb{E}_f f(m_1) \overline{f}(m_2) \left| \sum_{n \leq x} c(n) f(n) \right|^2 \ll_A \frac{x}{r^{\eta/2}}, \quad (2.9)$$

uniformly for $1 \leq m_1, m_2 \leq U^k$, provided $\varepsilon \ll_A 1$.⁷ Moreover,

$$\mathbb{E}_f |Q(f)|^{2k} \left| \sum_{n \leq x} c(n) f(n) \right|^2 \ll \sum_{n \leq x} \tilde{d}(n) |c(n)|^2 \cdot (k!) \left(2 \sum_{q \in \mathcal{Q}} q^{-1} v_q |a(q)|^2 \right)^k, \quad (2.10)$$

where $\tilde{d}(n) := \sum_{d|n} \mathbf{1}_{p|n \Rightarrow p \in \mathcal{P}}$, and $v_q := \mathbf{1}_{q \in \mathcal{P}} + 6 \cdot \mathbf{1}_{q \notin \mathcal{P}}$.

Proof. If $r > xU^k$, then (2.9) holds with no error, by perfect orthogonality. Therefore, by taking r large enough, we see that (2.10) follows immediately from [18, Lemma 1]. It remains to prove (2.9) in general, for $U^k \leq r^\varepsilon$ and $1 \leq x \leq r^A$.

For convenience, let $L^c(s, \psi) := L(s, \psi) \mathbf{1}_{c=\mu} + L^b(s, \psi) \mathbf{1}_{c=\lambda}$. By Perron's formula in the form of [27, Theorem 5.2 and Corollary 5.3], we have

$$\sum_{n \leq x} c(n) \chi(n) - \int_{1+\varepsilon-iT_0}^{1+\varepsilon+iT_0} \frac{1}{L^c(s, \chi)} \frac{y^s}{s} \frac{ds}{2\pi i} \ll \frac{y \log y}{T_0} + \frac{(4+y)^{1+\varepsilon}}{T_0} \ll \frac{y^{1+\varepsilon}}{T_0},$$

where $y := \lfloor x \rfloor + 0.5 \geq 1.5$. But by contour shifting and GRH, we have

$$\left(\int_{1+\varepsilon-iT_0}^{1+\varepsilon+iT_0} - \int_{\frac{1}{2}+\varepsilon-iT_0}^{\frac{1}{2}+\varepsilon+iT_0} \right) \frac{\mathbf{1}_{\chi \neq \chi_0}}{L^c(s, \chi)} \frac{y^s}{s} \frac{ds}{2\pi i} \ll (rT_0)^\varepsilon \frac{y^{1+\varepsilon}}{T_0}.$$

Since $|\sum_{n \leq x} c(n) \chi(n) \mathbf{1}_{\chi \neq \chi_0}| \ll_\varepsilon r^\varepsilon x^{0.5+\varepsilon}$ by GRH, we thus have, for each $j \in \{0, 1\}$,

$$(-1)^j \sum_{n \leq x} c(n) \chi(n) \mathbf{1}_{\chi \neq \chi_0} + \int_{\frac{1}{2}+\varepsilon-iT_0}^{\frac{1}{2}+\varepsilon+iT_0} \frac{\mathbf{1}_{\chi \neq \chi_0}}{L^c(s, \chi)} \frac{y^s}{s} \frac{ds}{2\pi i} \ll r^\varepsilon x^{0.5+\varepsilon} \mathbf{1}_{j=0} + (rT_0)^\varepsilon \frac{y^{1+\varepsilon}}{T_0}.$$

Let $T_0 := x^{(0.5+\eta)/(1-\varepsilon)}$. Since $||z|^2 - |w|^2| \leq |z^2 - w^2| = |z - w||z + w|$, it follows that

$$\left| \sum_{n \leq x} c(n) \chi(n) \mathbf{1}_{\chi \neq \chi_0} \right|^2 - \left| \int_{\frac{1}{2}+\varepsilon-iT_0}^{\frac{1}{2}+\varepsilon+iT_0} \frac{\mathbf{1}_{\chi \neq \chi_0}}{L^c(s, \chi)} \frac{y^s}{s} \frac{ds}{2\pi i} \right|^2 \ll (r^2 T_0)^\varepsilon \frac{x^{1.5+2\varepsilon}}{T_0}.$$

⁶We do not assume $xU^k < r$ like Harper does. On the other hand, our definition of $Q(f)$ agrees with Harper's definition when $f = \chi$.

⁷Of course, we could write $1 \leq m_1, m_2 \leq r^\varepsilon$ instead, and thus eliminate the role of U^k in the statement. However, we prefer to keep it, in order to highlight the connection to [18, Lemma 1].

Whenever $1 \leq m_1, m_2 \leq U^k$, we thus have

$$\begin{aligned} & \mathbb{E}_{\chi \neq \chi_0} \chi(m_1) \bar{\chi}(m_2) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 - \iint_{\frac{1}{2} + \varepsilon - iT_0}^{\frac{1}{2} + \varepsilon + iT_0} \mathbb{E}_f \frac{f(m_1) \bar{f}(m_2) y^{s_1} \bar{y}^{\bar{s}_2} ds_1 d\bar{s}_2}{L^c(s_1, f) L^c(\bar{s}_2, \bar{f}) s_1 \bar{s}_2 |2\pi i|^2} \\ & \ll (r^2 T_0)^\varepsilon \frac{x^{1.5+2\varepsilon}}{T_0} + \iint_{\frac{1}{2} + \varepsilon - iT_0}^{\frac{1}{2} + \varepsilon + iT_0} \frac{(1+T_0)^\varepsilon}{r^\eta} y^{\operatorname{Re}(s_1+s_2)} \frac{|ds_1 ds_2|}{|s_1 s_2|}, \end{aligned} \quad (2.11)$$

by Conjecture 2.3 if $c = \mu$, or by Proposition 2.4 if $c = \lambda$. In each case, we take $\hbar := \varepsilon$.

The right-hand side of (2.11) is $\ll r^{2\varepsilon} \frac{x^{1.5+2\varepsilon}}{T_0^{1-\varepsilon}} + \frac{(1+T_0)^\varepsilon}{r^\eta} x^{1+2\varepsilon} (\log T_0)^2$. Let r' be the smallest prime exceeding $\max(xU^k, r)$. By perfect orthogonality,

$$\mathbb{E}_{\chi \bmod r'} \chi(m_1) \bar{\chi}(m_2) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 = \mathbb{E}_f f(m_1) \bar{f}(m_2) \left| \sum_{n \leq x} c(n) f(n) \right|^2.$$

On subtracting (2.11) by its value for r' , we conclude that

$$\begin{aligned} & \frac{r-1}{r-2} \mathbb{E}_\chi \chi(m_1) \bar{\chi}(m_2) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 - \frac{r'-1}{r'-2} \mathbb{E}_f f(m_1) \bar{f}(m_2) \left| \sum_{n \leq x} c(n) f(n) \right|^2 \\ & \ll \frac{r^{2\varepsilon} x^{1+4\varepsilon}}{x^\eta} + \frac{x^{1+5\varepsilon}}{r^\eta} + \frac{|\sum_{n \leq x, \gcd(n,r)=1} c(n)|^2}{r-2} + \frac{|\sum_{n \leq x, \gcd(n,r')=1} c(n)|^2}{r'-2}, \end{aligned} \quad (2.12)$$

where the last two terms account for the missing $\chi = \chi_0$ contributions. By RH, we have $|\sum_{n \leq x, \gcd(n,r)=1} c(n)| \ll r^\varepsilon x^{1/2+\varepsilon}$. Moreover, $\frac{r'-1}{r'-2} - \frac{r-1}{r-2} \ll \frac{1}{r}$ and

$$\mathbb{E}_f f(m_1) \bar{f}(m_2) \left| \sum_{n \leq x} c(n) f(n) \right|^2 \ll \mathbb{E}_f \left| \sum_{n \leq x} c(n) f(n) \right|^2 \leq x,$$

by orthogonality. Therefore, after multiplying (2.12) by $\frac{r-2}{r-1}$, we get

$$\mathbb{E}_\chi \chi(m_1) \bar{\chi}(m_2) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 - \mathbb{E}_f f(m_1) \bar{f}(m_2) \left| \sum_{n \leq x} c(n) f(n) \right|^2 \ll \frac{r^{2\varepsilon} x^{1+4\varepsilon}}{x^\eta} + \frac{x^{1+5\varepsilon}}{r^\eta},$$

since $x/r \leq x^{1+2\varepsilon}/r^{1-2\varepsilon} \leq x^{1+5\varepsilon}/r^\eta$ and $x^{1+2\varepsilon}/(r')^{1-2\varepsilon} \leq x^{1+2\varepsilon}/r^{1-2\varepsilon}$.

We have already justified (2.9) for $r > xU^k$ at the beginning of the proof. Now assume $r \leq xU^k$. Then $x \geq r/U^k \geq r^{1-\varepsilon}$, so the last display is $\ll x/r^{\eta/2}$, provided $\varepsilon \ll_A 1$. So (2.9) holds. \square

For the next lemma, let $\Psi(x, y) := \#\{n \leq x : p \mid n \Rightarrow p \leq y\}$.

Lemma 2.6 (Smooth number bound). *Fix $\theta \in (0, \frac{1}{2})$ and $\varepsilon > 0$. If $\log Q \ll (\log x)^\theta$, then $\Psi(x, Q) \leq x/e^{(\log x)^{1-\theta-\varepsilon}}$ for all large enough x .*

Proof. Let x be large. Increasing Q if necessary, we may assume $\log Q \asymp (\log x)^\theta$. We have $Q = (\log x)^a$, where $a = (\log Q)/(\log \log x)$ satisfies $1 \leq a \leq (\log x)^{1/2}/(2 \log \log x)$. By Corollary 7.9 of [27], we have

$$\Psi(x, Q) \leq x^{1-1/a} x^{((\log a) + O(1))/(a \log \log x)} \leq x^{1-1/a} x^{(\theta+\varepsilon)/a} = x/e^{(1-\theta-\varepsilon)(\log x)/a},$$

because $(\log a) + O(1) \leq (\theta + \varepsilon)(\log \log x)$. Since $(\log x)/a \gg (\log x)^{1-\theta}$ as $x \rightarrow \infty$, it follows that

$$\Psi(x, Q) \leq x/e^{(\log x)^{1-\theta-\varepsilon}},$$

as desired. \square

We are now prepared to give a variant, Proposition 2.8, of [18, Proposition 1], with the following main differences:

- (1) Most importantly, we allow $x \leq r^A$ for any fixed $A > 0$. The cost is that we only allow *special* sequences of coefficients $c(n)$. More precisely, we either take $c(n) = \mu(n)$ for all n , or take $c(n) = \lambda(n)$ for all n .
- (2) We introduce a restriction $P^{400(Y/\delta)^2 \log(N \log P)} \leq r^\varepsilon$, in order to be able to apply the Ratios Conjecture. We will actually assume the conditions (2.14) and (2.15), which are nonetheless satisfied for the key choice of parameters in (2.13).
- (3) The deterministic side is supported on all moduli $n \leq x$, while the random side is supported on $\{n \leq x : P(n) > Q\}$ for some parameter $Q \leq e^{(\log x)^{1/3}}$. The perfect character orthogonality used in [18, § 3.2] (to pass to $P(n) > Q$) is no longer available for large x , so it seems more natural for us to pass to $P(n) > Q$ on the random side, not the deterministic side.

The cost of this asymmetry is that we can no longer easily⁸ take $Q = x^{1/\log \log x} = e^{(\log x)/\log \log x}$ like Harper does in [18, § 3.2]. Moreover, in our § 3.4 below, we will need to take X a bit smaller than Harper's choice of $e^{(\log x)^{1/2}}$ in [18, § 3.4].

- (4) We get a worse error term, with $x/(N \log P)^{10Y}$ in place of $x/(N \log P)^{Y/\delta^2}$. Essentially, the effect is that an error term of the form $x \left(\frac{2N+2}{(N \log P)^{1/\delta^2}} \right)^Y$ in [18, § 3.3] becomes $x \left(\frac{2N+2}{(N \log P)^{10}} \right)^Y$ in § 3.3 of our work below.

At this point it may help to note that we will eventually apply the proposition with the same parameters as Harper eventually does:

$$\log P \asymp (\log x)^{1/6}, \quad Y \asymp (\log P)^{1.02}, \quad N \asymp \log \log P, \quad \delta \asymp (\log P)^{-1.3}. \quad (2.13)$$

However, we will state it in slightly greater generality for the reader's convenience.

We begin by recalling the approximation result in [18].

Lemma 2.7 (Harper [18, Approximation Result 1]). *Let $N \in \mathbb{N}$ be large, and $\delta > 0$ be small. There exist functions $g : \mathbb{R} \rightarrow \mathbb{R}$ (depending on δ) and $g_{N+1} : \mathbb{R} \rightarrow \mathbb{R}$ (depending on δ and N) such that, if we define $g_j(x) = g(x - j)$ for all integers $|j| \leq N$, we have the following properties:*

- (i) $\sum_{|j| \leq N} g_j(x) + g_{N+1}(x) = 1$ for all $x \in \mathbb{R}$;
- (ii) $g(x) \geq 0$ for all $x \in \mathbb{R}$, and $g(x) \leq \delta$ whenever $|x| > 1$;
- (iii) $g_{N+1}(x) \geq 0$ for all $x \in \mathbb{R}$, and $g_{N+1}(x) \leq \delta$ whenever $|x| \leq N$;

⁸Harper notes that $\Psi(x, x^{1/\log \log x}) \ll x/(\log x)^{c \log \log \log x}$, but this bound does not fit into the error term of our proposition, in the key setting (2.13).

(iv) for all $\ell \in \mathbb{N}$ and all $x \in \mathbb{R}$, we have $|\frac{d^\ell}{dx^\ell} g(x)| \leq \frac{1}{\pi(\ell+1)} \left(\frac{2\pi}{\delta}\right)^{\ell+1}$.

Proposition 2.8. Assume Conjecture 2.3. Let $x, N, \delta^{-1}, P \geq 1$ be large real numbers, with $N \in \mathbb{Z}$. Let $c \in \{\mu, \lambda\}$ and $0 \leq Q \leq e^{(\log x)^{1/3}}$. Let $g_j: \mathbb{R} \rightarrow \mathbb{R}$, for $j \in [-N, N+1]$, be functions as in Lemma 2.7, with associated parameters N and δ . Let $Y \in \mathbb{N}$. Fix $A > 0$, suppose $x \leq r^A$, and assume, for ε as in Lemma 2.5, that

$$\max(P, \delta^{-1}, N)^{400(Y/\delta)^2 \log(N \log P)} \leq r^\varepsilon, \quad (2.14)$$

$$20Y \log(N \log P) \leq (\log x)^{1/2}. \quad (2.15)$$

Let $j(1), \dots, j(Y) \in [-N, N+1]$ be indices. Let

$$G_{f,j}(\mathbf{a}) := \prod_{1 \leq i \leq Y} g_{j(i)} \left(\operatorname{Re} \sum_{p \leq P} \frac{a_i(p)f(p)}{p^{1/2}} + \frac{a_i(p^2)f(p^2)}{p} \right),$$

where $a_i(p), a_i(p^2) \in \{z \in \mathbb{C} : |z| \leq 1\}$ for all i and p . Then

$$\mathbb{E}_\chi G_{\chi,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 = \mathbb{E}_f G_{f,j}(\mathbf{a}) \left| \sum_{\substack{n \leq x: \\ P(n) > Q}} c(n) f(n) \right|^2 + O\left(\frac{x}{(N \log P)^{10Y}}\right). \quad (2.16)$$

Proof. Before proceeding, we note that $0 \leq g_j(y) \leq 1$ for all $j \in [-N, N+1]$ and $y \in \mathbb{R}$, by Lemma 2.7 properties (i)–(iii). In particular, $0 \leq G_{f,j}(\mathbf{a}) \leq 1$. Also, by properties (iv) and (i) in Lemma 2.7, we have for all $l \geq 0$

$$\frac{|g_j^{(l)}(0)|}{l!} \leq \mathbf{1}_{l=0} + \frac{(2N+1)(2\pi/\delta)^{l+1}}{(l+1)!} \mathbf{1}_{l \geq 1}. \quad (2.17)$$

For the main proof, we first reduce to the case $Q = 0$, i.e. we bound the “missing” contribution from $P(n) \leq Q$ on the right-hand side of (2.16). Squaring both sides of

$$\sum_{\substack{n \leq x: \\ P(n) > Q}} c(n) f(n) = \sum_{n \leq x} c(n) f(n) - \sum_{\substack{n \leq x: \\ P(n) \leq Q}} c(n) f(n),$$

then using the bound $|G_{f,j}(\mathbf{a})| \leq 1$ and the triangle inequality, we find that

$$\mathbb{E}_f G_{f,j}(\mathbf{a}) \left| \sum_{\substack{n \leq x: \\ P(n) > Q}} c(n) f(n) \right|^2 - \mathbb{E}_f G_{f,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) f(n) \right|^2 \ll S_{Q,Q} + S_{Q,x},$$

where

$$S_{Q,R} := \mathbb{E}_f \left| \sum_{\substack{n \leq x: \\ P(n) \leq Q}} c(n) f(n) \right| \left| \sum_{\substack{n \leq x: \\ P(n) \leq R}} c(n) f(n) \right|.$$

By orthogonality over f , we have $S_{R,R} \leq \Psi(x, R)$. By the Cauchy–Schwarz inequality, $S_{Q,x} \leq S_{Q,Q}^{1/2} S_{x,x}^{1/2} \leq \Psi(x, Q)^{1/2} x^{1/2}$. By Lemma 2.6 with $\theta = 1/3$, we have $\Psi(x, Q) \ll x/e^{(\log x)^{1/2}}$, say. So

$$S_{Q,Q} + S_{Q,x} \leq \Psi(x, Q) + \Psi(x, Q)^{1/2} x^{1/2} \ll x/e^{\frac{1}{2}(\log x)^{1/2}}.$$

This fits into the error term of (2.16), by (2.15).

Therefore, it remains to prove

$$\mathbb{E}_\chi G_{\chi,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 = \mathbb{E}_f G_{f,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) f(n) \right|^2 + O\left(\frac{x}{(N \log P)^{10Y}}\right), \quad (2.18)$$

i.e. the $Q = 0$ case. The estimate (2.18) will be proven using Lemma 2.5. The main new difficulty is to account for the error term in (2.9); the contribution from (2.10) will essentially match the error terms Harper already accounted for. We begin by using Lemma 2.7 as Harper does in [18, proof of Proposition 1]. Let

$$S := 100Y \lfloor (1/\delta)^2 \log(N \log P) \rfloor \geq 100. \quad (2.19)$$

For all $y \in \mathbb{R}$, we then have $g_j(y) = \tilde{g}_j(y) + r_j(y)$, where $\tilde{g}_j(y) := \sum_{0 \leq l \leq 2S-1} g_j^{(l)}(0) \frac{y^l}{l!}$ and

$$|r_j(y)| \leq \alpha \frac{N |2\pi y/\delta|^{2S}}{\delta (2S+1)!},$$

by [18, first paragraph of the proof of Proposition 1], for some absolute constant $\alpha > 0$ that we name for later convenience. The bound (2.17) implies

$$\sum_{0 \leq l \leq 2S-1} \frac{|g_j^{(l)}(0) y^l|}{l!} \leq 1 + \sum_{0 \leq l \leq 2S-1} \frac{(2N+1)(2\pi) |2\pi y/\delta|^l}{\delta (l+1)!} \ll \frac{N \max(4S, |2\pi y/\delta|)^{2S}}{\delta (2S)!},$$

where in the final step we note that $\frac{\max(4S, |2\pi y/\delta|)^l}{(l+1)!} \leq \frac{1}{2} \frac{\max(4S, |2\pi y/\delta|)^{l+1}}{(l+2)!}$ for $0 \leq l \leq 2S-2$, and that $1 \ll \frac{N}{\delta}$ since N, δ^{-1} are large. Let

$$\tilde{G}_{f,j}(\mathbf{a}) := \prod_{1 \leq i \leq Y} \tilde{g}_{j(i)} \left(\operatorname{Re} \sum_{p \leq P} \frac{a_i(p) f(p)}{p^{1/2}} + \frac{a_i(p^2) f(p^2)}{p} \right).$$

Now write $\operatorname{Re} z = \frac{1}{2}(z + \bar{z})$, and expand \tilde{G} trivially. Note that (2.14) and (2.19) imply $P^{4SY} \leq r^\varepsilon$. By (2.9) with $U = P^{4SY} \leq r^\varepsilon$ and $k = 1$, we get

$$\mathbb{E}_\chi \tilde{G}_{\chi,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 - \mathbb{E}_f \tilde{G}_{f,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) f(n) \right|^2 \ll O(1)^Y R_0(Y), \quad (2.20)$$

where for any real $t \geq 0$ we define, for later convenience,

$$R_0(t) := \frac{x}{r^{\eta/2}} \left(2 + \frac{N \max(4S, |2\pi \sum_{p \leq P} (p^{-1/2} + p^{-1})/\delta|)^{2S}}{\delta (2S)!} \right)^t. \quad (2.21)$$

Next, the bound $|\tilde{g}_j(y)| \leq |g_j(y)| + |r_j(y)| \leq 1 + \alpha \frac{N |2\pi y/\delta|^{2S}}{\delta (2S+1)!}$ observed by Harper gives

$$\mathbb{E}_\chi |\tilde{G}_{\chi,j}(\mathbf{a}) - G_{\chi,j}(\mathbf{a})| \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 \leq \sum_{1 \leq i \leq Y} \mathbb{E}_\chi H_1(\chi, i) \quad (2.22)$$

by the triangle inequality, where for any function f we let $H_1(f, i)$ denote the quantity

$$\left| \sum_{n \leq x} c(n) f(n) \right|^2 \prod_{1 \leq l \leq i} \left(\mathbf{1}_{l < i} + \frac{\alpha N |2\pi \sum_{p \leq P} (p^{-1/2} a_l(p) f(p) + p^{-1} a_l(p^2) f(p^2))/\delta|^{2S}}{\delta (2S+1)!} \right).$$

Similarly,

$$\mathbb{E}_f |\tilde{G}_{f,j}(\mathbf{a}) - G_{f,j}(\mathbf{a})| \left| \sum_{n \leq x} c(n) f(n) \right|^2 \leq \sum_{1 \leq i \leq Y} \mathbb{E}_f H_1(f, i). \quad (2.23)$$

However, if we trivially expand H_1 , after replacing $\mathbf{1}_{l < i}$ with its upper bound 1 even for $l = i$, then an application of (2.9) with $U = P^{4Si} \leq P^{4SY} \leq r^\varepsilon$ and $k = 1$ shows that

$$\mathbb{E}_\chi H_1(\chi, i) - \mathbb{E}_f H_1(f, i) \ll O(1)^i R_0(i),$$

whence

$$\sum_{1 \leq i \leq Y} \mathbb{E}_\chi H_1(\chi, i) - \sum_{1 \leq i \leq Y} \mathbb{E}_f H_1(f, i) \ll \sum_{1 \leq i \leq Y} O(1)^i R_0(i) \ll O(1)^Y R_0(Y), \quad (2.24)$$

since $R_0(i+1)/R_0(i) \geq 2$. Next, we bound $\mathbb{E}_f H_1(f, i)$ as Harper does, using the fact that $\mathbf{1}_{l < i} = 0$ for $l = i$. That is, we first expand the product $\prod_{1 \leq l \leq i}$ in $H_1(f, i)$ as a sum of 2^{i-1} terms, then apply Hölder's inequality (as Harper does implicitly) in the form

$$\begin{aligned} & \mathbb{E}_f \left| \sum_{n \leq x} c(n) f(n) \right|^2 \prod_{l \in J} \left| \sum_{p \leq P} (p^{-1/2} a_l(p) f(p) + p^{-1} a_l(p^2) f(p^2)) \right|^{2S} \\ & \leq \prod_{l \in J} \left(\mathbb{E}_f \left| \sum_{n \leq x} c(n) f(n) \right|^2 \left| \sum_{p \leq P} (p^{-1/2} a_l(p) f(p) + p^{-1} a_l(p^2) f(p^2)) \right|^{2S|J|} \right)^{1/|J|} \end{aligned}$$

for various sets $J \subseteq \{1, \dots, i\}$ with $i \in J$, and finally use (2.10), to obtain the bound

$$\sum_{1 \leq i \leq Y} \mathbb{E}_f H_1(f, i) \ll O(1)^Y H_2, \quad (2.25)$$

where

$$H_2 := \sum_{n \leq x} \tilde{d}(n) |c(n)|^2 \sum_{1 \leq j \leq i \leq Y} \binom{i-1}{j-1} (jS)! \left(\frac{N |2\pi/\delta|^{2S} (2 \sum_{p \leq P} (p^{-1} + p^{-2}))^S}{(2S+1)!} \right)^j.$$

The factor $\binom{i-1}{j-1}$ represents the number of sets $J \subseteq \{1, \dots, i\}$ with $i \in J$ and $|J| = j$.

Combining (2.20), (2.22), and (2.23), via the triangle inequality, we get

$$\begin{aligned} & \mathbb{E}_\chi G_{\chi,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) \chi(n) \right|^2 - \mathbb{E}_f G_{f,j}(\mathbf{a}) \left| \sum_{n \leq x} c(n) f(n) \right|^2 \\ & \ll O(1)^Y R_0(Y) + \sum_{1 \leq i \leq Y} \mathbb{E}_\chi H_1(\chi, i) + \sum_{1 \leq i \leq Y} \mathbb{E}_f H_1(f, i) \ll O(1)^Y (R_0(Y) + H_2), \end{aligned} \quad (2.26)$$

where in the final step we bound the H_1 contributions using (2.24) and (2.25). To estimate the terms $R_0(Y)$ and H_2 in (2.26), we will repeatedly use the well-known bounds $n! \geq (n/e)^n$ and $n! \ll n^{1/2} (n/e)^n$ for integers $n \geq 1$. By (2.21),

$$R_0(Y) \leq \frac{x}{r^{\eta/2}} \left(2 + \frac{N (4S)^{2S} + O(P^{1/2}/\delta)^{2S}}{(2S/e)^{2S}} \right)^Y \leq \frac{O(N/\delta)^Y x}{r^{\eta/2}} ((2e)^{2S} + O(P^{1/2}/\delta)^{2S})^Y,$$

since $N \gg 1 \gg \delta$. By (2.14) we have $\max(P, \delta^{-1}, N)^{4SY} \leq r^\varepsilon$, so this means

$$O(1)^Y R_0(Y) \leq \frac{x}{r^{\eta/2}} (PN\delta^{-1})^{4SY} \leq \frac{r^{3\varepsilon} x}{r^{\eta/2}} \leq \frac{x}{r^{5\varepsilon}} \leq \frac{x}{(NP)^{10SY}} \leq \frac{x}{(N \log P)^{10Y}},$$

because $(NP)^{2SY} \leq r^\varepsilon$ and $S \geq 1$.

We now estimate H_2 as Harper does, after summing over $i \in [j, Y]$ using the hockey-stick identity $\sum_{j \leq i \leq Y} \binom{i-1}{j-1} = \binom{Y}{j}$. Noting that $\sum_{n \leq x} \tilde{d}(n) \leq x \prod_{p \leq P} (1 - p^{-1})^{-1} \ll x \log P$ and $\sum_{p \leq P} p^{-1} = \log \log P + O(1)$, we get

$$\begin{aligned} H_2 &\ll x \log P \sum_{1 \leq j \leq Y} \binom{Y}{j} (jS)^{1/2} (jS/e)^{jS} \left(\frac{N}{\delta} \frac{|2\pi/\delta|^{2S} (2.1 \log \log P)^S}{(2S/e)^{2S}} \right)^j \\ &\ll x \log P \sum_{1 \leq j \leq Y} \binom{Y}{j} (jS)^{1/2} \frac{(2.1\pi^2 e j)^{jS}}{(S\delta^2)^{jS}} \left(\frac{N}{\delta} (\log \log P)^S \right)^j. \end{aligned}$$

Since $j \leq Y$, and $S\delta^2 \geq 99Y \log(N \log P)$ by (2.19), we get

$$\begin{aligned} O(1)^Y H_2 &\ll O(1)^Y (x \log P) 2^Y (YS)^{1/2} \max_{1 \leq j \leq Y} \left(\frac{N}{\delta} \frac{(\log \log P)^S}{(1.7 \log(N \log P))^S} \right)^j \\ &\ll O(1)^Y (x \log P) S^{1/2} \left(\frac{N/\delta}{1.7^S} + \left(\frac{N/\delta}{1.7^S} \right)^Y \right). \end{aligned}$$

By Lemma 2.9 below, $N/\delta, O(1)^Y, S, \log P \leq 1.1^S$, so the last line is $\ll x(1.1^4/1.7)^S \leq x/1.1^S \leq x/(N \log P)^{Y/\delta^2}$, because $99 \log 1.1 \geq 1$. This suffices, because $\delta^{-2} \geq 10$. \square

Lemma 2.9. *Fix $A > 0$. Suppose $P, N, \delta^{-1} \geq 1$ are large. Let $Y \geq 1$. Let $S = 100Y \lfloor (1/\delta)^2 \log(N \log P) \rfloor$. Then $N/\delta, O(1)^Y, S, \log P \leq 1.1^S$.*

Proof. This is clear logarithmically:

$$\log N + \log(1/\delta), Y \log O(1), \log S, \log \log P \leq S \log 1.1,$$

because P, N, δ^{-1}, S are large. \square

While [18, Proposition 1] would not suffice for our purposes below, because of its restrictions on x , we note that [18, Proposition 2] will still apply unconditionally, without change, because its statement does not involve x at all.

3. PROOF OF THEOREM 1.2

3.1. Overview. In this section, we prove Theorem 1.2, in the following stronger form:

Theorem 3.1. *Each of the following implies the next:*

- (1) GRH for $L(s, \chi)$, and the Ratios Conjecture 2.1 for $L\bar{L}/L\bar{L}$, hold.
- (2) GRH for $L(s, \chi)$, and the twisted Ratios Conjecture 2.3 for $\chi\bar{\chi}/L\bar{L}$, hold.
- (3) Lemma 2.5 holds.
- (4) Proposition 2.8 holds.

(5) Fix a constant $A > 0$. Let $10 \leq x \leq r^A$ and $q \in [0, 1]$. Then

$$\mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} \lambda(n) \chi(n) \right|^{2q} \ll \left(\frac{x}{1 + (1 - q)(\log \log x)^{1/2}} \right)^q.$$

The implications (1) \Rightarrow (2), (2) \Rightarrow (3), (3) \Rightarrow (4) have already been proven in the previous section. Therefore, it only remains to prove the implication (4) \Rightarrow (5). Let $c(n) := \lambda(n)$. The key tool is Proposition 2.8, which helps us to pass the study of the deterministic side to the random side.

By using Hölder's inequality, it is sufficient to establish that for all $q \in [2/3, 1]$,

$$\mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} c(n) \chi(n) \right|^{2q} \ll \left(\frac{x}{1 + (1 - q)(\log \log x)^{1/2}} \right)^q.$$

3.2. The conditioning argument. We first do the “condition” argument. More precisely, we use the idea from [18], using a partition function to split the character sum into several pieces, based on values taken by certain character sum over primes. The proof is almost identical to § 3.2 in [18]. The only difference is that we do not restrict ourselves to the integers with an extra condition $P(n) > Q$ here.

We define

$$S_k(\chi, c) := \operatorname{Re} \sum_{p \leq P} \left(\frac{\chi(p)c(p)}{p^{1/2+ik/\log^{1.01} P}} + \frac{\chi(p)^2 c(p)^2}{p^{1+2ik/\log^{1.01} P}} \right)$$

for all $|k| \leq M := 2 \log^{1.02} P \in \mathbb{Z}$. Let \mathbf{j} be a $2M+1$ -vector and W be any function. We next apply the approximation result to rewrite the sum $\mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} c(n) \chi(n) \right|^{2q}$ as

$$\begin{aligned} & \mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} c(n) \chi(n) \right|^{2q} \prod_{i=-M}^{i=M} \sum_{j=-N}^{N+1} g_j(S_i(\chi, c)) \\ &= \sum_{-N \leq j(-M), \dots, j(M) \leq N+1} \mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} c(n) \chi(n) \right|^{2q} \prod_{i=-M}^M g_{j(i)}(S_i(\chi, c)) \\ &= \sum_{-N \leq j(-M), \dots, j(M) \leq N+1} \sigma(\mathbf{j}) \mathbb{E}_\chi^{\mathbf{j}} \left| \sum_{1 \leq n \leq x} \chi(n) c(n) \right|^{2q}, \end{aligned}$$

where we use the notation

$$\sigma(\mathbf{j}) := \mathbb{E}_\chi \prod_{i=-M}^M g_{j(i)}(S_i(\chi, c)), \quad \mathbb{E}_\chi^{\mathbf{j}} W := \frac{\mathbf{1}_{\sigma(\mathbf{j}) \neq 0}}{\sigma(\mathbf{j})} \mathbb{E}_\chi W \prod_{i=-M}^M g_{j(i)}(S_i(\chi, c)).$$

(Note that $g_{j(i)} \geq 0$, so if $\sigma(\mathbf{j}) = 0$, then $\mathbb{E}_\chi W \prod_{i=-M}^M g_{j(i)}(S_i(\chi, c)) = 0$ for all W .) We next apply the Hölder inequality to $\mathbb{E}_\chi^{\mathbf{j}}$ and conclude that

$$\mathbb{E}_\chi \left| \sum_{1 \leq n \leq x} \chi(n) c(n) \right|^{2q} \leq \sum_{-N \leq j(-M), \dots, j(M) \leq N+1} \sigma(\mathbf{j}) \left(\mathbb{E}_\chi^{\mathbf{j}} \left| \sum_{1 \leq n \leq x} \chi(n) c(n) \right|^2 \right)^q. \quad (3.1)$$

3.3. Passing to the random side. We next use Proposition 2.8 to pass to the random case. Let $Y := 2M + 1$ and $Q = Q(x) := e^{(\log x)^{1/3}}$, and assume P, N, δ satisfy (2.13). Then the requirements in (2.14) and (2.15) hold, so we may apply Proposition 2.8 to get

$$\begin{aligned} & \mathbb{E}_{\chi}^j \left| \sum_{1 \leq n \leq x} \chi(n) c(n) \right|^2 \\ &= \frac{\mathbf{1}_{\sigma(j) \neq 0}}{\sigma(j)} \left(\mathbb{E} \prod_{i=-M}^M g_{j(i)}(S_i(f, 1)) \left| \sum_{\substack{n \leq x \\ P(n) > Q}} f(n) \right|^2 + O\left(\frac{x}{(N \log P)^{10Y}}\right) \right). \end{aligned} \quad (3.2)$$

Note that we have replaced $f(n)c(n)$ on the right with $f(n)$ (in S_i and in the sum over n), which is possible because $c = \lambda$ is completely multiplicative and f is Steinhaus.

We note that our Q is smaller than Harper's $x^{1/\log \log x}$, and that our error term is worse, with $10Y$ instead of Y/δ^2 . Thus, we will focus our exposition on aspects where we lose something compared to Harper, to clarify why we still win overall.

We next notice that $\sum_j \sigma(j) = 1$ and apply the Hölder inequality to the $\sigma(j)$. This lets us conclude that the contribution from the “big Oh” term in (3.2) to (3.1) is at most

$$\left(\sum_{-N \leq j(-M), \dots, j(M) \leq N+1} \sigma(j) \frac{\mathbf{1}_{\sigma(j) \neq 0}}{\sigma(j)} \frac{x}{(N \log P)^{10Y}} \right)^q \leq \left(\frac{x(2N+2)^Y}{(N \log P)^{10Y}} \right)^q \ll \left(\frac{x}{\log P} \right)^q,$$

which is $\ll \left(\frac{x}{1+(1-q)(\log \log P)^{1/2}} \right)^q$. Thus, we can ignore the “big Oh” term in (3.2).

We next define

$$\sigma^{\text{random}}(j) := \mathbb{E} \prod_{i=-M}^M g_{j(i)}(S_i(f, 1))$$

for all $(2M+1)$ -vectors j where f is a Steinhaus random multiplicative function. We then use [18, Proposition 2] to get that $\sigma(j)^{1-q} \ll \sigma^{\text{random}}(j)^{1-q} + \left(\frac{1}{(N \log P)^{(2M+1)(1/\delta)^2}} \right)^{1-q}$. This implies that

$$\sum_{-N \leq j(-M), \dots, j(M) \leq N+1} \sigma(j) \left(\frac{\mathbf{1}_{\sigma(j) \neq 0}}{\sigma(j)} \mathbb{E} \prod_{i=-M}^M g_{j(i)}(S_i(f, 1)) \left| \sum_{\substack{n \leq x \\ P(n) > Q}} f(n) \right|^2 \right)^q \ll T_1 + T_2,$$

where

$$\begin{aligned} T_1 &:= \sum_j \sigma^{\text{random}}(j) \left(\frac{\mathbf{1}_{\sigma^{\text{random}}(j) \neq 0}}{\sigma^{\text{random}}(j)} \mathbb{E} \prod_{i=-M}^M g_{j(i)}(S_i(f, 1)) \left| \sum_{\substack{n \leq x \\ P(n) > Q}} f(n) \right|^2 \right)^q, \\ T_2 &:= \left(\frac{1}{(N \log P)^{(2M+1)(1/\delta)^2}} \right)^{1-q} \sum_j \left(\mathbb{E} \prod_{i=-M}^M g_{j(i)}(S_i(f, 1)) \left| \sum_{\substack{n \leq x \\ P(n) > Q}} f(n) \right|^2 \right)^q, \end{aligned}$$

where $\sum_{\mathbf{j}} := \sum_{-N \leq j(-M), \dots, j(M) \leq N+1}$. To deal with T_2 , we apply Hölder's inequality to the sum over \mathbf{j} and use the fact that g_j form a partition of unity. This gives us

$$\begin{aligned} T_2 &\ll \left(\frac{1}{(N \log P)^{(2M+1)(1/\delta)^2}} \right)^{1-q} ((2N+2)^{2M+1})^{1-q} \left(\sum_{\mathbf{j}} \mathbb{E} \prod_{i=-M}^M g_{j(i)}(S_i(f, 1)) \left| \sum_{\substack{n \leq x \\ P(n) > Q}} f(n) \right|^2 \right)^q \\ &= \left(\left(\frac{2N+2}{(N \log P)^{(1/\delta)^2}} \right)^{2M+1} \right)^{1-q} \left(\mathbb{E} \left| \sum_{\substack{n \leq x \\ P(n) > Q}} f(n) \right|^2 \right)^q \leq (\log P)^{-(1-q)} x^q, \end{aligned}$$

which is again $\ll \left(\frac{x}{1+(1-q)(\log \log P)^{1/2}} \right)^q$. In summary, we can ignore the contribution from T_2 . If we define the notation $\mathbb{E}^{\mathbf{j}, \text{rand}} W := \frac{\mathbf{1}_{\sigma^{\text{random}}(\mathbf{j}) \neq 0}}{\sigma^{\text{random}}(\mathbf{j})} \mathbb{E} W \prod_{i=-M}^M g_{j(i)}(S_i(f, 1))$ for all random variables W , then by (3.1) and (3.2), it suffices to show that

$$T_1 = \sum_{\mathbf{j}} \sigma^{\text{random}}(\mathbf{j}) \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \left| \sum_{\substack{n \leq x \\ P(n) > Q}} f(n) \right|^2 \right)^q \ll \left(\frac{x}{1+(1-q)(\log \log P)^{1/2}} \right)^q, \quad (3.3)$$

in order to complete the proof of Theorem 3.1. The point is that from now on we only need to focus on T_1 , which is purely about random multiplicative functions.

3.4. Passing to Euler products. Recall that $Q(x) = e^{(\log x)^{1/3}}$. Let $X = e^{(\log x)^{1/100}}$. In the corresponding section [18, § 3.4], Harper uses a standard sieve bound [27, Theorem 3.6] several times in the course of the argument. Since our parameters are slightly different than Harper's, it is worth spelling these out more explicitly.

Lemma 3.2 (Applied sieve bounds). *We have*

$$\sum_{\substack{Q < m \leq x \\ p|m \Rightarrow p > P}} \frac{X}{m} \int_m^{(1+1/X)m} \left(\frac{x}{m} - \frac{x}{t} + 1 \right) dt \ll \frac{x}{\log P}.$$

Also, for $t \geq Q$, we have

$$\sum_{\substack{t/(1+1/X) < m \leq t \\ p|m \Rightarrow p > P}} \frac{X}{m} \ll (\log P)^{-1}.$$

Proof. In the first display, we have $\frac{x}{m} - \frac{x}{t} + 1 = \frac{(t-m)x}{mt} + 1 \ll \frac{x}{mX} + 1$. Therefore, we only need to show that

$$\sum_{\substack{Q < m \leq x \\ p|m \Rightarrow p > P}} \left(1 + \frac{x}{mX} \right) \ll \frac{x}{\log P}.$$

Notice that sum involving x/Xm is crudely bounded by $\ll x \log x/X$ which is more than acceptable by our choice of X and P . For the other term, we just use the upper bound

sieve [27, Theorem 3.6], noting here $P \ll \sqrt{x - Q}$ by (2.13), to conclude that this is at most $\ll x \prod_{p \leq P} (1 - p^{-1}) \ll x / \log P$.

We now turn to the second display. We have the bound

$$\sum_{\substack{t/(1+1/X) < m \leq t \\ p|m \Rightarrow p > P}} \frac{X}{m} \ll \frac{X}{t} \sum_{\substack{t/(1+1/X) < m \leq t \\ p|m \Rightarrow p > P}} 1.$$

By using the upper bound sieve again, noting that $P \ll \sqrt{t - t/(1 + 1/X)}$ by (2.13), the inner sum is at most $\ll \frac{t}{X} \prod_{p \leq P} (1 - p^{-1}) \ll \frac{t}{X \log P}$ and the conclusion follows. \square

We now return to our goal, (3.3). Expanding the square in T_1 and noticing that $g_{j(i)}(S_i(f, 1))$ only depends on $f(p)$ for $p \leq P$, we get

$$T_1 = \sum_j \sigma^{\text{random}}(\mathbf{j}) \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \sum_{\substack{m \leq x \\ P(m) > Q \\ p|m \Rightarrow p > P}} \left| \sum_{\substack{n \leq x/m \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 \right)^q. \quad (3.4)$$

Note that we have used $P < Q$ above. The next step is to replace the discrete sum with a smooth version. Replacing the $P(m) > Q$ by the weaker condition $m > Q$, we find that the above bracket term, $(\mathbb{E}^{\mathbf{j}, \text{rand}} \sum \dots)^q$, in (3.4) is at most

$$\begin{aligned} &\ll \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \sum_{\substack{Q < m \leq x \\ p|m \Rightarrow p > P}} \frac{X}{m} \int_m^{m(1+1/X)} \left| \sum_{\substack{n \leq x/t \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 dt \right)^q \\ &+ \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \sum_{\substack{Q < m \leq x \\ p|m \Rightarrow p > P}} \frac{X}{m} \int_m^{m(1+1/X)} \left| \sum_{\substack{x/t \leq n \leq x/m \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 dt \right)^q. \end{aligned} \quad (3.5)$$

We first estimate the contribution of the second term in (3.5) to (3.4), which is at most, via Hölder's inequality (using $\sum_j \sigma^{\text{random}}(\mathbf{j}) = 1$),

$$\begin{aligned} &\ll \left(\sum_j \sigma^{\text{random}}(\mathbf{j}) \mathbb{E}^{\mathbf{j}, \text{rand}} \sum_{\substack{Q < m \leq x \\ p|m \Rightarrow p > P}} \frac{X}{m} \int_m^{m(1+1/X)} \left| \sum_{\substack{x/t \leq n \leq x/m \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 dt \right)^q \\ &= \left(\mathbb{E} \sum_{\substack{Q < m \leq x \\ p|m \Rightarrow p > P}} \frac{X}{m} \int_m^{m(1+1/X)} \left| \sum_{\substack{x/t \leq n \leq x/m \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 dt \right)^q, \end{aligned}$$

by the definition of $\mathbb{E}^{\mathbf{j}, \text{rand}}$. Next we use the orthogonality to expand the square and the above is $\ll (\frac{x}{\log P})^q$ by applying Lemma 3.2. This is $\ll (\frac{x}{1+(1-q)(\log \log P)^{1/2}})^q$.

We next estimate the contribution from the first term in (3.5) to (3.4). This is at most

$$\begin{aligned}
& \ll \sum_j \sigma^{\text{random}}(\mathbf{j}) \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \int_Q^x \left| \sum_{\substack{n \leq x/t \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 \sum_{\substack{t/(1+1/X) < m \leq x \\ p|m \Rightarrow p > P}} \frac{X}{m} dt \right)^q \\
& \ll \sum_j \sigma^{\text{random}}(\mathbf{j}) \left(\frac{1}{\log P} \right)^q \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \int_Q^x \left| \sum_{\substack{n \leq x/t \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 dt \right)^q \\
& = \sum_j \sigma^{\text{random}}(\mathbf{j}) \left(\frac{x}{\log P} \right)^q \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \int_1^{x/Q} \left| \sum_{\substack{n \leq z \\ n \text{ is } P\text{-smooth}}} f(n) \right|^2 \frac{dz}{z^2} \right)^q,
\end{aligned}$$

where in the second line we applied Lemma 3.2 and the in last line we used $z := x/t$. Then we can apply [18, Harmonic Analysis Result 1] to bound the last line by

$$\ll \left(\frac{x}{\log P} \right)^q \sum_j \sigma^{\text{random}}(\mathbf{j}) \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \int_{\mathbb{R}} \frac{|F_P^{\text{rand}}(1/2 + it)|^2}{|1/2 + it|^2} dt \right)^q, \quad (3.6)$$

where $F_P^{\text{rand}}(s) := \prod_{p \leq P} (1 - \frac{f(p)}{p^s})^{-1}$. Note that $F_P^{\text{rand}}(s)$ is a truncated version of $L(s, f)$.

The quantity in (3.6) is identical to the quantity in [18, (3.4)], because our choice of P is identical to Harper's. Therefore, by [18, (3.5) and the paragraph before it], we have

$$\sum_j \sigma^{\text{random}}(\mathbf{j}) \left(\mathbb{E}^{\mathbf{j}, \text{rand}} \int_{\mathbb{R}} \frac{|F_P^{\text{rand}}(1/2 + it)|^2}{|1/2 + it|^2} dt \right)^q \ll \left(\frac{\log P}{1 + (1 - q)(\log \log P)^{1/2}} \right)^q.$$

Plugging this into (3.6), and recalling our bounds for T_1 based on (3.4) and (3.5), we conclude that (3.3) holds. This completes the proof of Theorem 3.1.

REFERENCES

- [1] A. Balog, A. Granville, and K. Soundararajan. Multiplicative functions in arithmetic progressions. *Ann. Math. Qué.*, 37(1):3–30, 2013. [3](#)
- [2] J. Bergström, A. Diaconu, D. Petersen, and C. Westerland. Hyperelliptic curves, the scanning map, and moments of families of quadratic L -functions. [arXiv:2302.07664](#), 2023. [2](#)
- [3] S. Bettin and J. B. Conrey. Averages of long Dirichlet polynomials. *Riv. Math. Univ. Parma (N.S.)*, 12(1):1–27, 2021. [6](#)
- [4] T. Browning, J. Glas, and V. Y. Wang. Sums of three cubes over a function field. [arXiv:2402.07146](#), 2024. [7](#)
- [5] H. M. Bui and A. Florea. Negative moments of the Riemann zeta-function. *J. Reine Angew. Math.*, 806:247–288, 2024. [2](#)
- [6] H. M. Bui, A. Florea, and J. P. Keating. The Ratios Conjecture and upper bounds for negative moments of L -functions over function fields. *Trans. Amer. Math. Soc.*, 376(6):4453–4510, 2023. [2](#)
- [7] R. Caich. Random multiplicative functions and typical size of character in short intervals. [arXiv:2402.06426](#), 2024. [4](#)
- [8] J. B. Conrey, D. W. Farmer, J. P. Keating, M. O. Rubinstein, and N. C. Snaith. Integral moments of L -functions. *Proc. London Math. Soc. (3)*, 91(1):33–104, 2005. [2](#), [7](#), [8](#)

- [9] J. B. Conrey, D. W. Farmer, and M. R. Zirnbauer. Autocorrelation of ratios of L -functions. *Commun. Number Theory Phys.*, 2(3):593–636, 2008. [2](#), [6](#), [7](#), [8](#)
- [10] J. B. Conrey and N. C. Snaith. Applications of the L -functions ratios conjectures. *Proc. Lond. Math. Soc.* (3), 94(3):594–646, 2007. [2](#), [6](#), [8](#)
- [11] A. Florea. Negative moments of L -functions with small shifts over function fields. *Int. Math. Res. Not. IMRN*, 2024(3):2298–2337, 2024. [2](#)
- [12] É. Fouvry and M. Radziwiłł. Level of distribution of unbalanced convolutions. *Ann. Sci. Éc. Norm. Supér.* (4), 55(2):537–568, 2022. [3](#)
- [13] A. Granville and X. Shao. Bombieri-Vinogradov for multiplicative functions, and beyond the $x^{1/2}$ -barrier. *Adv. Math.*, 350:304–358, 2019. [3](#)
- [14] B. Green. A note on multiplicative functions on progressions to large moduli. *Proc. Roy. Soc. Edinburgh Sect. A*, 148(1):63–77, 2018. [3](#)
- [15] H. Gu and Z. Zhang. Universality and phase transitions in low moments of secular coefficients of critical holomorphic multiplicative chaos. [arXiv:2401.05681](#), 2024. [4](#)
- [16] A. J. Harper. Sharp conditional bounds for moments of the Riemann zeta function. [arXiv:1305.4618](#), 2013. [2](#)
- [17] A. J. Harper. Moments of random multiplicative functions, I: Low moments, better than squareroot cancellation, and critical multiplicative chaos. *Forum Math. Pi*, 8:e1, 95, 2020. [1](#), [4](#)
- [18] A. J. Harper. The typical size of character and zeta sums is $o(\sqrt{x})$. [arXiv:2301.04390](#), 2023. [1](#), [2](#), [4](#), [5](#), [10](#), [11](#), [13](#), [15](#), [17](#), [18](#), [19](#), [20](#), [22](#)
- [19] H. Helson. Hankel forms. *Studia Math.*, 198(1):79–84, 2010. [4](#)
- [20] H. Iwaniec and E. Kowalski. *Analytic number theory*, volume 53 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, 2004. [7](#), [9](#)
- [21] K. Matomäki and M. Radziwiłł. Multiplicative functions in short intervals. *Ann. of Math.* (2), 183(3):1015–1056, 2016. [2](#)
- [22] K. Matomäki and J. Teräväinen. On the Möbius function in all short intervals. *J. Eur. Math. Soc. (JEMS)*, 25(4):1207–1225, 2023. [2](#)
- [23] J. Maynard. Primes in arithmetic progressions to large moduli II: Well-factorable estimates. *Mem. AMS*; [arXiv:2006.07088](#), 2020. [3](#)
- [24] J. Maynard. Primes in arithmetic progressions to large moduli III: Uniform residue classes. *Mem. AMS*; [arXiv:2006.08250](#), 2020. [3](#)
- [25] J. Maynard. Primes in arithmetic progressions to large moduli I: Fixed residue classes. *Mem. AMS*; [arXiv:2006.06572](#), 2021. [3](#)
- [26] J. Miller, P. Patzt, D. Petersen, and O. Randal-Williams. Uniform twisted homological stability. [arXiv:2402.00354](#), 2024. [2](#)
- [27] H. L. Montgomery and R. C. Vaughan. *Multiplicative number theory. I. Classical theory*, volume 97 of *Cambridge Studies in Advanced Mathematics*. Cambridge University Press, 2007. [11](#), [12](#), [20](#), [21](#)
- [28] J. Najnudel, E. Paquette, and N. Simm. Secular coefficients and the holomorphic multiplicative chaos. *Ann. Probab.*, 51(4):1193–1248, 2023. [4](#)
- [29] W. Sawin. Square-root cancellation for sums of factorization functions over squarefree progressions in $\mathbb{F}_q[t]$. *Acta Math*; [arXiv:2102.09730](#), 2022. [2](#), [3](#)
- [30] K. Soundararajan. Moments of the Riemann zeta function. *Ann. of Math.* (2), 170(2):981–993, 2009. [2](#)
- [31] K. Soundararajan. Partial sums of the Möbius function. *J. Reine Angew. Math.*, 631:141–152, 2009. [2](#)
- [32] B. Szabó. High moments of theta functions and character sums. *Mathematika*, 70(2):Paper No. e12242, 37, 2024. [2](#)
- [33] V. Y. Wang. Notes on zeta ratio stabilization. [arXiv:2402.01214](#), 2024. [2](#)

- [34] M. W. Xu. Better than square-root cancellation for random multiplicative functions. *Trans. Amer. Math. Soc. Ser. B*, 11:482–507, 2024. [4](#)
- [35] L. Ye. Bounding sums of the Möbius function over arithmetic progressions. [arXiv:1406.7326](#), 2014. [3](#)

COURANT INSTITUTE, 251 MERCER STREET, NEW YORK 10012, USA

IST AUSTRIA, AM CAMPUS 1, 3400 KLOSTERNEUBURG, AUSTRIA

Email address: `vywang@alum.mit.edu`

DEPARTMENT OF MATHEMATICS, STANFORD UNIVERSITY, STANFORD, CA, USA

Email address: `maxxu@stanford.edu`