

# THE TAIL DISTRIBUTION OF THE PARTITION FUNCTION FOR DIRECTED POLYMER IN THE WEAK DISORDER PHASE

STEFAN JUNK AND HUBERT LACOIN

**ABSTRACT.** We investigate the upper tail distribution of the partition function of the directed polymer in a random environment on  $\mathbb{Z}^d$  in the weak disorder phase. We show that the distribution of the infinite volume partition function  $W_\infty^\beta$  displays a power-law decay, with an exponent  $p^*(\beta) \in [1 + \frac{2}{d}, \infty)$ . We also prove that the distribution of the suprema of the point-to-point and point-to-line partition functions display the same behavior. On the way to these results, we prove a technical estimate of independent interest: the  $L^p$ -norm of the partition function at the time when it overshoots a high value  $A$  is comparable to  $A$ . We use this estimate to extend the validity of many recent results that were proved under the assumption that the environment is upper bounded.

2010 *Mathematics Subject Classification:* 60K35, 60K37, 82B26, 82B27, 82B44.

*Keywords:* Disordered models, Directed polymers, Weak disorder.

## 1. INTRODUCTION

The directed polymer in random environment (or DPRE) on  $\mathbb{Z}^d$  is a model in statistical mechanics involving a random walk (or polymer) interacting with a disordered medium. It was introduced [HH85], for  $d = 1$ , as a simplified model to describe the interfaces of the planar Ising model with random coupling constants at low temperature, and was generalized to higher dimension soon afterwards [IS88].

The DPRE and variations of the model have received much attention from the mathematical community for a wide variety of reasons – we refer to [Zyg24] for a recent survey. In this introduction, we discuss mainly the model in spatial dimension  $d \geq 3$  as this is the object of the present work. In that setup, the DPRE undergoes a phase transition from a high-temperature, weak disorder phase to a low-temperature, strong disorder phase, each phase exhibiting radically different behavior. In the weak disorder phase, on large scales, the polymer trajectory is not affected by the disorder and displays the same behavior as a simple random walk. In particular, it converges to a standard Brownian motion under a diffusive scaling (see [CY06] and references therein).

On the other hand, in the strong disorder phase, it is conjectured that there exists a corridor where the environment is particularly favorable and around which the trajectories localize with high probability. It is further predicted that this corridor is *superdiffusive* in the sense that its transversal fluctuation is much larger than  $\sqrt{N}$ . While the rigorous understanding of this *pathwise* localization phenomenon is still rudimentary and usually requires assumptions beyond strong disorder (see e.g. [CC13, Bat21] for progresses in that direction), a more precise picture has emerged concerning localization for the *end point* of the polymer (see for instance [CH02, CSY03, BC20]).

The weak and strong disorder regimes are defined in terms of the asymptotic behavior of the partition function  $W_n^\beta$  (defined in Equation (2.3) below). Weak disorder holds if the sequence  $(W_n^\beta)_{n \geq 1}$  is uniformly integrable and converge to a non-trivial limit  $W_\infty^\beta$ , while strong disorder

holds if  $W_n^\beta$  converges to zero. There exists a critical value of  $\beta_c$  which separates the weak and strong disorder phases. The understanding of the phase transition from weak to strong disorder has been significantly improved in [JL24] by proving (under technical assumption on the distribution of  $\omega$ ) that whenever  $W_n^\beta$  decays to zero, it does so exponentially fast, and that consequently, weak disorder holds at  $\beta_c$ . However, most questions concerning this transition remain widely open. Let us mention two of them:

- (A) How regular is the free energy curve around  $\beta_c$ ? The free energy, defined as  $F(\beta) := \lim_{n \rightarrow \infty} n^{-1} \log W_n^\beta$ , is equal to zero for  $\beta \leq \beta_c$  and is negative for  $\beta > \beta_c$ , hence it is not analytic at  $\beta_c$ , but currently not much is known beyond this.
- (B) What features, if any, distinguish the behavior of the system at  $\beta_c$  from the interior of the weak disorder phase?

In the present work, we study the probability distribution of the limit  $W_\infty^\beta$  in the weak disorder phase. We prove that  $P(W_\infty^\beta > u)$  decays, up to a multiplicative constant, like  $u^{-p^*(\beta)}$  as  $u \rightarrow \infty$  for an exponent  $p^*(\beta) \in [1 + \frac{2}{d}, \infty)$ . In the process we also obtain comparable bounds for  $\sup_{n \geq 1} W_n^\beta$  and for the supremum over all point-to-point partition functions.

Many properties of the function  $\beta \mapsto p^*(\beta)$  have been proved in earlier works [Jun22a, Jun23a, Jun23b, JL24] (with an appropriate definition for  $p^*(\beta)$  see Equation (2.5)) sometimes with additional technical assumptions concerning the distribution on the environment  $\omega$ . More precisely in many instance  $\omega$  is assumed to be upper bounded. In this paper, we provide alternative proofs for some of these statements which do not require any assumption besides exponential integrability of  $\omega$ .

The key estimates that allows us to prove our results with greater generality is Proposition 2.3 which controls the amount of overshoot of  $W_n^\beta$  at time when it crosses a large value  $A$ . This technical result is likely to find further application in the study of directed polymer, and could also be generalized and helpful in the study of other disordered models.

## 2. MODEL AND RESULTS

**2.1. Definition and previous results.** Let  $X = (X_k)_{k \geq 0}$  be the nearest neighbor simple random walk on  $\mathbb{Z}^d$  starting from the origin and  $P$  its law. We have  $P(X_0 = 0) = 1$  and the increments  $(X_{k+1} - X_k)_{k \geq 1}$  are independent and identically distributed (i.i.d.) with distribution

$$P(X_1 = x) = \frac{\mathbf{1}_{\{|x|=1\}}}{2d}$$

where  $|\cdot|$  denotes the  $\ell_1$  distance on  $\mathbb{Z}^d$ . Given a collection  $\omega = (\omega_{k,x})_{k \geq 1, x \in \mathbb{Z}^d}$  of real-valued weights (the environment), a parameter  $\beta \geq 0$  (the inverse temperature) and  $n \geq 1$  (the polymer length), we define the polymer measure  $P_n^{\beta, \omega}$  as a modification of the distribution  $P$  which favors trajectories that visit sites where  $\omega$  is large. More precisely, to each path  $\pi: \mathbb{N} \rightarrow \mathbb{Z}^d$  we associate an energy  $H_n(\omega, \pi) := \sum_{i=1}^n \omega_{i, \pi(i)}$ , and we set

$$P_n^{\beta, \omega}(dX) := \frac{1}{Z_n^\beta} e^{\beta H_n(\omega, X)} P(dX), \quad \text{where} \quad Z_n^\beta := E \left[ e^{\beta H_n(\omega, X)} \right]. \quad (2.1)$$

The quantity  $Z_n^\beta$  is referred to as the partition function of the model (note that  $Z_n^\beta$  depends on  $\omega$ ). In what follows, we assume that the environment  $\omega$  is given by a fixed realization of an i.i.d. random field on  $\mathbb{N} \times \mathbb{Z}^d$  with law  $\mathbb{P}$ . We assume that  $\omega$  has finite exponential moments of all order, that is

$$\forall \beta \in \mathbb{R}, \quad \lambda(\beta) := \log \mathbb{E}[e^{\beta \omega_{1,0}}] < \infty. \quad (2.2)$$

By Fubini, it is easy to check that  $\mathbb{E}[Z_n^\beta] = e^{n\lambda(\beta)}$ , and it is thus natural to define the normalized partition function  $W_n^\beta$  by setting

$$W_n^\beta = \frac{Z_n^\beta}{\mathbb{E}[Z_n^\beta]} = E \left[ e^{\beta H_n(\omega, X) - n\lambda(\beta)} \right]. \quad (2.3)$$

The normalized partition function  $W_n^\beta$  encodes essential information on the typical behavior of  $X$  under  $P_n^{\beta, \omega}$  and thus has been a central object of attention in the study of this model. An important observation made in [Bol89] is that  $W_n^\beta$  is a martingale with respect to the filtration  $\mathcal{F}_n := \sigma(\omega_{k,x} : k \leq n)$ , and hence converges almost surely to a limit  $W_\infty^\beta \in [0, \infty)$ . Furthermore we have  $\mathbb{P}(W_\infty^\beta > 0) \in \{0, 1\}$  since the event is measurable w.r.t. the tail  $\sigma$ -algebra. We say that *weak disorder* holds if  $W_\infty^\beta > 0$  while *strong disorder* holds if  $W_\infty^\beta = 0$ .

Another observation from [Bol89] is that  $(W_n^\beta)_{n \in \mathbb{N}}$  is bounded in  $L^2$  when  $d \geq 3$  and  $\beta$  is sufficiently small. Indeed setting  $\beta_2 := \sup\{\beta \geq 0 : \sup_{n \geq 0} \mathbb{E}[(W_n^\beta)^2] < \infty\}$ , an explicit computation yields that  $\beta_2$  either satisfies the following identity

$$e^{\lambda(2\beta_2) - 2\lambda(\beta_2)} = \frac{1}{P^{\otimes 2}(\exists n \geq 1, X_n^{(1)} = X_n^{(2)})}. \quad (2.4)$$

or is equal to infinity when (2.4) has no solution. In particular,  $\beta_2 > 0$  when  $d \geq 3$  and weak disorder hold for  $\beta \in (0, \beta_2)$ . On the other hand,  $\beta_c = 0$  when  $d = 1, 2$ , and it was shown in that case [CH02, CSY03] that strong disorder holds for all  $\beta > 0$ . As mentioned earlier, this work is concerned with the weak disorder phase and we thus restrict ourselves to  $d \geq 3$ . The following result indicates that the influence of the disorder is monotone in  $\beta$ :

**Theorem A** ([CY06, JL24]). *Assume  $d \geq 3$ . There exists  $\beta_c \in (0, \infty]$  such that weak disorder holds for  $\beta < \beta_c$  and strong disorder holds for  $\beta > \beta_c$ . If  $\beta_c < \infty$  and  $\omega$  is bounded from above, i.e.,  $\text{ess sup } \omega_{1,0} < \infty$ , then weak disorder also holds at  $\beta = \beta_c$ .*

It is furthermore known that  $\beta_c > \beta_2$  (see [BS10] for a proof of this statement when  $d \geq 4$ , for dimension 3 we refer to [BT10] and [JL24, Theorem B]) i.e. there exists an interval of  $\beta$  such that weak disorder holds but  $(W_n^\beta)_{n \in \mathbb{N}}$  is not bounded in  $L^2$ . The integrability of  $W_\infty^\beta$  in the weak disorder phase, and in particular in the interval  $[\beta_2, \beta_c]$ , has been an object of interest. In order to quantify it, the following integrability exponent has been introduced in [Jun22a]

$$p^*(\beta) := \inf \left\{ p > 0 : \sup_{n \geq 0} \mathbb{E}[(W_n^\beta)^p] = \infty \right\}. \quad (2.5)$$

We always have  $p^*(\beta) \geq 1$  and as a consequence of [CY06, Lemma 3.3],  $\beta \mapsto p^*(\beta)$  is nonincreasing. Clearly we have  $p^*(\beta) = 1$  in strong disorder and  $p^*(\beta) \geq 2$  for  $\beta < \beta_2$ , but a priori there is no guarantee that  $p^*(\beta) > 1$  in the remainder of the weak disorder. The following result summarizes what is known about the integrability of  $W_\infty^\beta$ .

**Theorem B** ([CY06, Jun22b]). *Weak disorder is equivalent to  $\mathbb{E}[\sup_n W_n^\beta] < \infty$ , and in particular  $W_n^\beta$  converges to  $W_\infty^\beta$  in  $L^1$  in weak disorder. If furthermore  $\text{ess sup } \omega_{1,0} < \infty$ , then weak disorder implies that  $p^*(\beta) > 1$ .*

**2.2. Power tail asymptotics for the partition function.** Our first result establishes that, up to a multiplicative constant, the tail distribution  $P(W_\infty^\beta > u)$  decays like a power of  $u$ . The corresponding exponent depends  $\beta$  and is equal to  $p^*(\beta)$ . We obtain asymptotics of the same

order for the suprema of partition functions

$$W_{\infty}^{\beta,*} = \sup_{n \geq 0} W_n^{\beta,*} \quad \text{and} \quad \widehat{W}_{\infty}^{\beta,*} = \sup_{(n,x) \in \mathbb{N} \times \mathbb{Z}^d} \widehat{W}_n^{\beta}(x), \quad (2.6)$$

where

$$\widehat{W}_n^{\beta}(x) := E \left[ e^{\beta H_n(\omega, X) - n\lambda(\beta)} \mathbf{1}_{\{X_n = x\}} \right]. \quad (2.7)$$

Given two positive functions  $f$  and  $g$  defined on  $\mathbb{R}_+$  we say that  $f$  and  $g$  are comparable at infinity and write  $f(u) \asymp g(u)$  if there exist  $C > 1$  and  $u_0 > 0$  such such that

$$\forall u \geq u_0, \quad \frac{g(u)}{C} \leq f(u) \leq Cg(u).$$

**Theorem 2.1.** *When weak disorder holds, we have the following:*

$$\mathbb{P} \left( W_{\infty}^{\beta,*} > u \right) \asymp \mathbb{P} \left( \widehat{W}_{\infty}^{\beta,*} > u \right) \asymp \mathbb{P} \left( W_{\infty}^{\beta} > u \right) \asymp u^{-p^*(\beta)}. \quad (2.8)$$

**Remark 2.2.** *Save for a couple special cases, the value of  $p^*(\beta)$  is not known in general. A first special case is the  $L^2$ -threshold  $\beta_2$  defined in Equation (2.4): we have  $p^*(\beta_2) = 2$ . A second one is the critical threshold  $\beta_c$ . In [JL24, Corollary 2.2] it is stated that under the assumption that  $\omega$  is upper bounded we have  $p^*(\beta_c) = 1 + \frac{2}{d}$ . Hence in that case Theorem 2.1 implies that*

$$\mathbb{P} \left( W_{\infty}^{\beta_c} \geq u \right) \asymp u^{-\frac{d+2}{d}}. \quad (2.9)$$

*We believe that such a precise information may be helpful in attempts to address Questions (A) and (B) raised in the introduction.*

**2.3. Overshoot considerations.** We present next a technical result which not only plays a key role in our proof but has many other potential applications.

In recent years, many significant results concerning directed polymers have been obtained under some restriction concerning the distribution on  $\omega$ . Either assuming that  $\omega$  is bounded from above [Jun22b, Jun22a, Jun23a, JL24], or some regularity on the tail distribution of  $\omega$  [FJ23]. One of the reasons (and in many occurrences the main one) for imposing these restriction is to have a control on the value of the partition function as it overshoots a given threshold  $A > 1$ . To illustrate better what we mean, let us define  $\tau_A := \inf\{n \geq 1 : W_n^{\beta} \geq A\}$  (with the convention  $\inf \emptyset = \infty$ ). If one assume that  $\omega$  is bounded from above then we have

$$W_{\tau_A} \leq LA \quad \text{with} \quad L := e^{\beta_{\text{ess sup}}(\omega_{1,0}) - \lambda(\beta)}. \quad (2.10)$$

The information that  $W_{\tau_A}$  is comparable to  $A$  turns out to have many practical applications. Of course (2.10) is false if the environment is not bounded from above. Using only the assumption (2.2), we obtain a result in the same spirit with a control of the  $L^p$  norm of  $W_{\tau_A}$ , for arbitrary  $p \in [1, \infty)$ , instead of the supremum norm.

**Proposition 2.3.** *For any  $p \geq 1$  there exists a constant  $C_p > 0$  (depending also on  $\beta$  and on  $\mathbb{P}$ ) such that for every  $A > 1$  we have*

$$\mathbb{E} \left[ (W_{\tau_A}^{\beta})^p \mid \tau_A < \infty \right] < C_p A^p. \quad (2.11)$$

**Remark 2.4.** *In the course of the proof of Proposition 2.3, we will see that the constant  $C_p$  can be taken of the form*

$$C_p = \left( 2^p + C e^{\frac{\lambda(2\beta p)}{2} - p\lambda(\beta)} \right). \quad (2.12)$$

*where  $C$  depends on  $\beta$  and on  $\mathbb{P}$  but not on  $p$ . In particular,  $C_p$  can be taken continuous and increasing in  $p$ .*

With the help of Proposition 2.3, we prove the required lower bound for  $\mathbb{P}(W_\infty^{\beta,*} \geq u)$  in (2.8).

**Corollary 2.5.** *There exists a constant  $c > 0$  (depending on  $\beta$  and on  $\mathbb{P}$ ) such that for every  $u \geq 1$  we have*

$$\mathbb{P}(\tau_A < \infty) = \mathbb{P}(W_\infty^{\beta,*} \geq A) \geq cA^{-p^*(\beta)}. \quad (2.13)$$

Note that we do not require that weak disorder holds and that (2.13) is also valid in the strong disorder regime, in which case  $p^*(\beta) = 1$ .

**Remark 2.6.** *The bound (2.13) has been proved under the assumption that  $p^*(\beta) \leq 2$  and that the environment is upper bounded as [Jun22a, Theorem B].*

**2.4. First consequences of the main results.** The main results have a couple of rather direct consequences. First, we obtain a lower bound for  $p^*(\beta)$  in the weak disorder phase.

**Corollary 2.7.** *Assume that weak disorder holds at  $\beta$ .*

- (i) *It holds that  $p^*(\beta) > 1$ .*
- (ii) *It holds that  $p^*(\beta) \geq 1 + \frac{2}{d}$ .*

Of course, claim (ii) implies claim (i) and there is no need to state it separately. We included it here because of the simplicity of its proof, which is immediate from (2.8). By [JL24, Corollary 2.2], the bound from part (ii) is sharp in the case of a bounded environment.

**Remark 2.8.** *Item (i) has been proved as [Jun22b, Theorem 1.1(ii)] under the assumption that the environment is upper-bounded and in [FJ23, Theorem 1.1] under a weaker assumption on the regularity of the tail of  $\omega$  at infinity. Item (ii) was proved assuming an upper-bounded environment as [Jun22a, Corollary 1.3]. The proof of this lower-bound was accomplished indirectly via the study of a fluctuation exponent associated with the partition function. This approach extends to general environment, see the comment after Corollary 2.13, but we present an alternative, direct proof, which partially relies on ideas developed in [JL24].*

Next, we introduce the critical threshold for exponential growth of moments,

$$q^*(\beta) := \inf \left\{ p \geq 0 : \lim_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{E}[(W_n^\beta)^p] > 0 \right\}. \quad (2.14)$$

The existence of the above limit follows from the subadditivity/superadditivity of the sequence  $\log \mathbb{E}[(W_n^\beta)^p]$  (depending on whether  $p \leq 1$  or  $p \geq 1$ ). The inequality  $p^*(\beta) \leq q^*(\beta)$  is trivial.

**Corollary 2.9.** *The following hold:*

- (i) *If weak disorder holds at  $\beta$  then  $\lim_{\beta' \uparrow \beta} p^*(\beta') = p^*(\beta)$ .  
In particular,  $\beta \mapsto p^*(\beta)$  is left-continuous in  $(0, \beta_c)$ .*
- (ii) *If weak disorder holds at  $\beta$  then  $p^*(\beta) = q^*(\beta)$ .*

**Remark 2.10.** *We expect that the identity  $p^*(\beta) = q^*(\beta)$  holds in the strong disorder regime (recall that in that case  $p^*(\beta) = 1$ ) but at the moment cannot prove this with full generality. Using exponential concentration of  $\log W_n$  around its mean [LW09, Section 6] it is possible to show that  $q^*(\beta) = 1$  whenever  $\limsup_{n \rightarrow \infty} \frac{1}{n} \log W_n^\beta < 0$  a regime known as very strong disorder. The equivalence of strong disorder and very strong disorder has been established [JL24, Theorem 2.1] under the assumption that the environment is upper-bounded.*

**Remark 2.11.** *Item (i) has been proved as [Jun23a, Theorem 1.2, item (ii)] under the additional assumption that  $p^*(\beta) \leq 2$  and that the environment is upper bounded. Item (ii) has been proved under the same assumption as [Jun22a, Theorem 1.5]. The validity of part (ii) for general environments has been mentioned as an open problem in [Zyg24, Question 6].*

Part (ii) of the previous corollary shows that  $\mathbb{E}[(W_n^\beta)^p]$  grows exponentially fast for any  $p > p^*(\beta)$ , and by definition  $\mathbb{E}[(W_n^\beta)^p]$  is bounded when  $p < p^*(\beta)$ . When  $p = p^*(\beta)$ , Theorem 2.1 immediately implies that  $\lim_{n \rightarrow \infty} \mathbb{E}[(W_n^\beta)^{p^*(\beta)}] = \infty$  but identifying the growth of  $\mathbb{E}[(W_n^\beta)^{p^*(\beta)}]$  turns out to be a challenging task. As a consequence of Theorem 2.1, we can prove that this growth is at most linear in  $n$ .

**Corollary 2.12.** *For any  $\beta$ , there exists  $C > 0$  such that*

$$\mathbb{E} \left[ (W_n^\beta)^{p^*(\beta)} \right] \leq Cn. \quad (2.15)$$

We conjecture in fact that

$$\mathbb{E} \left[ (W_n^\beta)^{p^*(\beta)} \right] = n^{\kappa(\beta) + o(1)}$$

where  $\kappa(\beta) \in (0, 1]$  when  $\beta \in (0, \beta_c)$  and  $\kappa(\beta_c) = 0$ . To motivate this, let us explicitly compute  $\mathbb{E}[(W_n^\beta)^{p^*(\beta)}]$  in the specific case of  $\beta = \beta_2$  (recall (2.4)). The second moment of the partition function  $\mathbb{E}[(W_n^{\beta_2})^2]$  coincides with the partition function of a homogeneous pinning model and thus can be explicitly computed explicitly:  $\mathbb{E}[(W_n^{\beta_2})^2]$  grows exponentially in  $n$  for  $\beta > \beta_2$  and when  $\beta = \beta_2$  we have

$$\mathbb{E}[(W_n^{\beta_2})^2] \sim \begin{cases} C_3 n^{1/2} & \text{if } d = 3, \\ C_4 \frac{n}{\log n} & \text{if } d = 4, \\ C_d n & \text{if } d \geq 5. \end{cases} \quad (2.16)$$

For a proof of (2.16) we refer to [Gia07, Theorem 2.2], applied to the renewal function

$$K(n) := P^{\otimes 2} \left( n = \inf \{ i \geq 1 : X_i^{(1)} = X_i^{(2)} \} \right)$$

The above information, combined with the fact that weak disorder holds at  $\beta_2$  and Corollary 2.9, implies that  $p^*(\beta_2) = q^*(\beta_2) = 2$ . The asymptotic (2.16) illustrates in particular that (2.15) can be sharp in some situations (namely  $\beta = \beta_2$  and  $d \geq 5$ ). The identification of the growth rate in the general case, and particularly for  $\beta = \beta_c$ , remains a challenging open problem that we believe to be quite important for understanding the phase transition of the model, in particular for Questions (A) and (B) mentioned in the introduction.

**2.5. The fluctuation field.** To introduce the next corollary, let us digress a bit on the connection between the DPPE and the Stochastic Heat Equation with multiplicative noise (SHE). This connection can be seen for instance by writing the recursion equation which is satisfied by the point-to-point partition function (or some variant, for instance, considering shifted partition function, see Equation (2.19) below). The resulting equation is a discrete analogue of the SHE.

In dimension 1, the solution of the SHE can be obtained as a scaling limit of the point to point partition function of the directed polymer by considering diffusive scaling and taking  $\beta$  proportional to  $n^{-1/4}$  where  $n$  length of the polymer (see [AKQ14]).

When  $d = 2$ , the SHE, as is, is ill defined and the directed polymer model has been used as instrument to define a two dimensional version of the SHE via scaling limit (in that case  $\beta$  has to be proportional to  $(\log n)^{-1/2}$  see [CSZ23] as well as references therein).

In dimension  $d \geq 3$ , when weak disorder holds, the system homogenizes and, at first order, the disorder disappears under diffusive scaling (the scaling limit is simply the heat equation without noise). In that case, a natural question to investigate is that of the amplitude and distribution of the random fluctuations around this deterministic limit. It has been done in [MSZ16, GRZ18, CN21, CNN22, LZ22, Jun22a].

To state the result, let us introduce the translation operator  $\theta_{m,y}$ . For  $m \geq 0$  and  $y \in \mathbb{Z}^d$  the environment  $\theta_{m,y} \omega$  is defined by setting

$$(\theta_{m,y} \omega)_{n,x} := \omega_{n+m,x+y}. \quad (2.17)$$

We also let  $\theta_{m,y}$  act on functions of  $\omega$  by setting  $\theta_{m,y} f(\omega) := f(\theta_{m,y} \omega)$ . We are interested in investigating the scaling limit of the field  $(Y_\beta(n, x))_{x \in \mathbb{Z}^d}$  defined by

$$Y_\beta(n, x) := \theta_{0,x} W_n^\beta \quad (2.18)$$

Note that, by time reversal, for a fixed  $n$ ,  $Y_\beta(\cdot, x)$  has the same distribution as  $\tilde{Y}_\beta(n, \cdot)$  defined by

$$\begin{aligned} \tilde{Y}_\beta(0, x) &= 1, & \forall x \in \mathbb{Z}^d, \\ \tilde{Y}_\beta(n+1, x) &= e^{\beta \omega_{n+1,x} - \lambda(\beta)} D \tilde{Y}_\beta(n, x). \end{aligned} \quad (2.19)$$

where the operator  $D$  is the transition matrix of the simple random walk on  $\mathbb{Z}^d$ , that is to say

$$Df(x) = \frac{1}{2d} \sum_{y \in \mathbb{Z}^d} f(y) \mathbf{1}_{|x-y|_1=1}. \quad (2.20)$$

Thus (2.19) corresponds to a discrete analogue of the stochastic heat equation with multiplicative noise. In the weak disorder phase, it has been established (see [Jun22a, Theorem C(i)] and [CNN22, Theorem 2.1] for an continuum analogue) that homogenization occurs in the weak disorder phase in the sense that for any  $f \in C_c(\mathbb{R}^d)$  (continuous and compactly supported functions  $f: \mathbb{R}^d \rightarrow \mathbb{R}$ ) we have, in  $L^1$ ,

$$\lim_{n \rightarrow \infty} n^{-d/2} \sum_{x \in \mathbb{Z}^d} f(x/\sqrt{n}) Y_\beta(n, x) = \int_{\mathbb{R}^d} f(x) dx. \quad (2.21)$$

We define the fluctuation field  $\mathcal{X}_n^\beta$  around this limit by setting for  $f \in C_c(\mathbb{R}^d)$

$$\mathcal{X}_n^\beta(f) := n^{-d/2} \sum_{x \in \mathbb{Z}^d} f(x/\sqrt{n}) (Y_\beta(n, x) - 1).$$

In the case where  $\beta < \beta_2$  (that is to say  $p^*(\beta) > 2$ ), the exact scaling limit of  $\mathcal{X}_n^\beta$  has been identified. More precisely it has been proved that the process  $(n^{\frac{d-2}{4}} \mathcal{X}_n^\beta(f))_{f \in C_c(\mathbb{R}^d)}$  converges to a random Gaussian field whose covariance can be expressed in terms of the  $d$ -dimensional heat kernel (see [LZ22, Theorem 1.1]). A consequence of this is that when  $p^*(\beta) > 2$  we have

$$\lim_{\varepsilon \rightarrow 0} \lim_{n \rightarrow \infty} \mathbb{P} \left( \varepsilon n^{-\frac{d-2}{4}} \leq |\mathcal{X}_n^\beta(f)| \leq \frac{1}{\varepsilon} n^{-\frac{d-2}{4}} \right) = 1. \quad (2.22)$$

When  $p^*(\beta) < 2$ , since the field  $Y_\beta(n, x)$  is not  $L^2$ -integrable, one may expect larger fluctuations for the field. Indeed, in [Jun22a, Theorem 1.1], the correct fluctuation exponent has been identified in this regime, showing that  $|\mathcal{X}_n^\beta(f)|$  is of order  $n^{-\xi(\beta)+o(1)}$  where

$$\xi(\beta) := \frac{d}{2} - \frac{2+d}{2(p^*(\beta) \wedge 2)}.$$

However, for technical reasons, the result was proved under the assumption that the environment  $\omega$  is upper bounded. The only reason for this technical limitation is that the identity  $p^*(\beta) = q^*(\beta)$  was proved in [Jun22a] under this assumption. Hence, using Corollary 2.9, we can extend the validity of the result. We record this as our last corollary.

**Corollary 2.13.** *When weak disorder holds we have, for any  $\varepsilon > 0$  and  $f \neq 0$ ,*

$$\lim_{n \rightarrow \infty} \mathbb{P}\left(n^{-\xi(\beta)-\varepsilon} \leq |\mathcal{X}_n^\beta(f)| \leq n^{-\xi(\beta)+\varepsilon}\right) = 1, \quad (2.23)$$

Note that (2.23) only identifies the correct fluctuation exponent but it leaves open the question of identifying the exact order of magnitude of  $\mathcal{X}_n^\beta(f)$  (as in (2.22)) and that of identifying the scaling limit of  $\mathcal{X}_n^\beta(f)$  as in [LZ22].

**Remark 2.14.** *Note that the combination of (2.21) and (2.23) implies that we have necessarily  $\xi(\beta) \geq 0$ . This yields an alternative proof of Corollary 2.7(ii).*

*Proof of Corollary 2.13.* When  $p^*(\beta) > 2$  there is nothing to prove since (2.23) follows directly from (2.22). When  $p^*(\beta) \leq 2$ , we can use [Jun22a, Theorem 1.4] which states – under the assumption that  $p^*(\beta) > 1$ , which holds by Corollary 2.7(i) – that

$$\begin{aligned} \lim_{n \rightarrow \infty} \mathbb{P}\left(|\mathcal{X}_n^\beta(f)| \leq n^{-\xi^*(\beta)+\varepsilon}\right) &= 1, \\ \lim_{n \rightarrow \infty} \mathbb{P}\left(|\mathcal{X}_n^\beta(f)| \geq n^{-\xi(\beta)-\varepsilon}\right) &= 1, \end{aligned} \quad (2.24)$$

where  $\xi^*(\beta) := \frac{d}{2} - \frac{d+2}{2q^*(\beta)}$  (recall the definition (2.14)). Since by Corollary 2.9(i) we have  $q^*(\beta) = p^*(\beta)$  in weak disorder, this allows to conclude.  $\square$

**2.6. Organization of the paper.** In Section 3, we prove Proposition 2.3 and Corollary 2.5. In Section 4, we prove Theorem 2.1 and Corollaries 2.5 and 2.7(i). The remaining corollaries are proved in Section 5.

**2.7. Notation.** Throughout the paper, we make use of the notation  $\llbracket a, b \rrbracket := [a, b] \cap \mathbb{Z}$ . We let  $\mu_n^\beta$  denote the endpoint distribution associated with the polymer measure  $P_{\omega, n}^\beta$  that is to say

$$\mu_n^\beta(x) = P_n^{\beta, \omega}(X_n = x). \quad (2.25)$$

### 3. PROOF OF THE RESULTS OF SECTION 2.3

**3.1. Probability of doubling the mean of a linear combination of i.i.d. random variables.** We start this section introducing a key technical lemma used in the proof. Let us introduce first some context. Recalling the definition 2.20, we have

$$\frac{W_n^\beta}{W_{n-1}^\beta} = \sum_{x \in \mathbb{Z}^d} D\mu_{n-1}^\beta(x) e^{\beta\omega_{x, n} - \lambda(\beta)}. \quad (3.1)$$

The coefficients  $D\widehat{W}_{n-1}^\beta(x)$  are  $\mathcal{F}_{n-1}$  measurable while the variables  $e^{\beta\omega_{x, n} - \lambda(\beta)}$  are i.i.d. and independent of  $\mathcal{F}_{n-1}$ . The following result can be used to estimate the conditional probability of “doubling” the partition function at step  $n$ .

**Lemma 3.1.** *Let  $(Y_i)_{i \geq 1}$  be a sequence of i.i.d. nonnegative random variables with moments of all orders and such that  $\mathbb{E}[Y_1] = 1$ . Then for any  $q \geq 1$  there exists  $C_q$  (depending on the distribution of  $Y_1$ ) such that for any sequence  $(\alpha_i)_{i \geq 1}$  of nonnegative real numbers satisfying  $\sum_{i \geq 1} \alpha_i \leq 1$*

$$\mathbb{P}\left(\sum_{i \geq 1} \alpha_i Y_i \geq 2\right) \leq C_q \alpha_{\max}^q, \quad (3.2)$$

where  $\alpha_{\max} = \max_{i \geq 1} \alpha_i$ .



*Proof.* We want to derive the inequality from a Chernov type bound. Since a priori the variables  $Y_i$  are not exponentially integrable, we first need to apply a truncation. We observe that

$$\mathbb{P} \left( \sum_{i \geq 1} \alpha_i Y_i \geq 2 \right) \leq \mathbb{P} \left( \max_{i \geq 1} \alpha_i Y_i > \sqrt{\alpha_{\max}} \right) + \mathbb{P} \left( \sum_{i \geq 1} (\alpha_i Y_i) \wedge \sqrt{\alpha_{\max}} \geq 2 \right). \quad (3.3)$$

We can bound the first term in the r.h.s. as follows

$$\mathbb{P} \left( \max_{i \geq 1} \alpha_i Y_i > \sqrt{\alpha_{\max}} \right) \leq \sum_{i \geq 1} \mathbb{P} (\alpha_i Y_i > \sqrt{\alpha_{\max}}) \leq \mathbb{E}[Y_1^{2(q+1)}] \sum_{i \geq 1} \frac{\alpha_i^{2q+2}}{\alpha_{\max}^{q+1}} \leq \mathbb{E}[Y_1^{2q+2}] \alpha_{\max}^q,$$

where in the last inequality simply uses that  $\alpha_i^{2q+2} \leq \alpha_{\max}^{2q+1} \alpha_i$ . Now to bound the second term in the r.h.s. of (3.3), we observe that for any  $\lambda > 0$  we have

$$\mathbb{P} \left( \sum_{i \geq 1} (\alpha_i Y_i) \wedge \sqrt{\alpha_{\max}} \geq 2 \right) \leq \mathbb{E} \left[ e^{\lambda [\sum_{i \geq 1} ((\alpha_i Y_i) \wedge \sqrt{\alpha_{\max}}) - 2]} \right]. \quad (3.4)$$

Using the inequality  $e^u \leq 1 + u + u^2$  valid for  $u \leq 1$  we obtain that for all  $\lambda \leq \alpha_{\max}^{-1/2}$ ,

$$\mathbb{E} \left[ e^{\lambda ((\alpha_i Y_i) \wedge \sqrt{\alpha_{\max}})} \right] \leq \mathbb{E} [1 + \lambda \alpha_i Y_i + \lambda^2 \alpha_i^2 Y_i^2] \leq e^{\lambda \alpha_i + \lambda^2 \alpha_i^2 \mathbb{E}[Y_i^2]}. \quad (3.5)$$

This implies that

$$\mathbb{E} \left[ e^{\lambda [\sum_{i \geq 1} ((\alpha_i Y_i) \wedge \sqrt{\alpha_{\max}}) - 2]} \right] \leq e^{\lambda [\sum_{i \geq 1} \alpha_i - 2] + \lambda^2 \mathbb{E}[Y_1^2] \sum_{i \geq 1} (\alpha_i)^2}. \quad (3.6)$$

Using  $\sum_{i \geq 1} \alpha_i \leq 1$  and  $\sum_{i \geq 1} \alpha_i^2 \leq \alpha_{\max}$  and taking  $\lambda = \alpha_{\max}^{-1/2}$ , we obtain

$$\mathbb{P} \left( \sum_{i \geq 1} (\alpha_i Y_i) \wedge \sqrt{\alpha_{\max}} \geq 2 \right) \leq e^{\mathbb{E}[Y_1^2] - \alpha_{\max}^{-1/2}} \leq e^{\mathbb{E}[Y_1^2] [2q]! \alpha_{\max}^{-q}}, \quad (3.7)$$

which concludes the proof.  $\square$

**3.2. Proof of Proposition 2.3.** Since is trivial if  $\omega$  is bounded from above, we may assume that  $\mathbb{P}(\omega_{1,0} > t) > 0$  for all  $t > 0$ . We set  $\tau := \tau_A$  for notational simplicity. We want to discard first the contribution of “small overshoot” above the value  $A$  using the following decomposition

$$\begin{aligned} \mathbb{E} \left[ (W_\tau^\beta)^p \mathbf{1}_{\{\tau < \infty\}} \right] &= \mathbb{E} \left[ (W_\tau^\beta)^p (\mathbf{1}_{\{\tau < \infty ; W_\tau \in [A, 2A]\}} + \mathbf{1}_{\{\tau < \infty ; W_\tau \geq 2A\}}) \right] \\ &\leq (2A)^p \mathbb{P}(\tau < \infty) + \mathbb{E} \left[ (W_\tau^\beta)^p \mathbf{1}_{\{\tau < \infty ; W_\tau \geq 2A\}} \right]. \end{aligned} \quad (3.8)$$

Then we estimate the second term decomposing according to the value of  $\tau$ . We have

$$\begin{aligned} \mathbb{E} \left[ (W_\tau^\beta)^p \mathbf{1}_{\{\tau < \infty ; W_\tau \geq 2A\}} \right] &= \sum_{n \geq 1} \mathbb{E} \left[ (W_n^\beta)^p \mathbf{1}_{\{\tau \geq n ; W_n \geq 2A\}} \right] \\ &= \sum_{n \geq 1} \mathbb{E} \left[ \mathbb{E} \left[ (W_n^\beta)^p \mathbf{1}_{\{W_n \geq 2A\}} \mid \mathcal{F}_{n-1} \right] \mathbf{1}_{\{\tau \geq n\}} \right]. \end{aligned} \quad (3.9)$$

Next we apply Cauchy-Schwarz and obtain

$$\mathbb{E} \left[ (W_n^\beta)^p \mathbf{1}_{\{W_n \geq 2A\}} \mid \mathcal{F}_{n-1} \right] \leq \sqrt{\mathbb{E} \left[ (W_n^\beta)^{2p} \mid \mathcal{F}_{n-1} \right] \mathbb{P}(W_n \geq 2A \mid \mathcal{F}_{n-1})} \quad (3.10)$$

Now setting  $\xi_x := e^{\beta\omega_{n,x}-\lambda(\beta)}$  and recalling (3.1) we obtain that (using Jensen's inequality for the probability  $D\widehat{W}_{n-1}^\beta(x)/W_{n-1}^\beta$ ) on the event  $\tau \geq n$

$$\begin{aligned} \mathbb{E} \left[ (W_n^\beta)^{2p} \mid \mathcal{F}_{n-1} \right] &= (W_{n-1}^\beta)^{2p} \mathbb{E} \left[ \left( \sum_{x \in \mathbb{Z}^d} \frac{D\widehat{W}_{n-1}^\beta(x)}{W_{n-1}^\beta} \xi_x \right)^{2p} \mid \mathcal{F}_{n-1} \right] \\ &\leq e^{\lambda(2p\beta)-2p\lambda(\beta)} A^{2p}. \end{aligned} \quad (3.11)$$

Next we use Lemma 3.1 with coefficients  $\alpha_x := A^{-1}D\widehat{W}_{n-1}^\beta(x)$  and

$$q := -4 \log_2 \kappa \quad \text{with} \quad \kappa := \mathbb{P}(e^{\beta\omega_{0,1}-\lambda(\beta)} \geq 4d) \wedge \frac{1}{2}, \quad (3.12)$$

( $\kappa > 0$  since we assumed that  $\omega$  was unbounded from above). We obtain that for a constant  $C$  that depends only on  $\beta$  and of the distribution of  $\omega$ , on the event  $\tau \geq n$  (which guarantees that  $W_{n-1}^\beta \leq A$  and hence that the coefficient's sum is smaller than one) we have

$$\mathbb{P}(W_n \geq 2A \mid \mathcal{F}_{n-1}) \leq C \left( \max_{x \in \mathbb{Z}^d} A^{-1} D\widehat{W}_{n-1}^\beta(x) \right)^q. \quad (3.13)$$

Combining (3.9), (3.10) and (3.11) we obtain that

$$\mathbb{E} \left[ (W_n^\beta)^p \mathbf{1}_{\{\tau < \infty ; W_n \geq 2A\}} \right] \leq C e^{\frac{\lambda(2p\beta)}{2} - p\lambda(\beta)} A^p \sum_{n \geq 1} \mathbb{E} \left[ \mathbf{1}_{\{\tau \geq n\}} \left( \max_{x \in \mathbb{Z}^d} \frac{D\widehat{W}_{n-1}^\beta(x)}{A} \right)^{q/2} \right]. \quad (3.14)$$

To conclude the proof we are going to show that – with the above choice of  $q$  – we have (for a different  $C > 0$ )

$$\sum_{n \geq 1} \mathbb{E} \left[ \mathbf{1}_{\{\tau \geq n\}} \left( \max_{x \in \mathbb{Z}^d} D\widehat{W}_{n-1}^\beta(x) \right)^{q/2} \right] \leq C \mathbb{P}(\tau < \infty). \quad (3.15)$$

Combining (3.8), (3.14) and (3.15) we obtain that for a constant  $C$  that depends on the distribution of  $\omega$  and on  $\beta$  but not on  $p$  we have

$$\mathbb{E} \left[ (W_\tau^\beta)^p \mathbf{1}_{\{\tau < \infty\}} \right] \leq \left( 2^p + C e^{\frac{\lambda(2p\beta)}{2} - p\lambda(\beta)} \right) \mathbb{P}(\tau < \infty). \quad (3.16)$$

Let us now prove (3.15). Recalling the definition of  $\kappa$  in (3.12), we observe that for any  $k \geq 1$ ,

$$\mathbb{P} \left[ W_{n+k-1} \geq \left( 2^{k+1} d \right) \max_{x \in \mathbb{Z}^d} D\widehat{W}_{n-1}^\beta(x) \mid \mathcal{F}_{n-1} \right] \geq \kappa^k. \quad (3.17)$$

Indeed, assuming that  $\max_{x \in \mathbb{Z}^d} D\widehat{W}_{n-1}^\beta(x)$  is attained at  $x_0$ , the event on the right-hand side is satisfied if

$$\forall i \in \llbracket 1, k \rrbracket, \quad e^{\beta\omega_{n+i-1, x_0 + (i-1)\mathbf{e}_1} - \lambda(\beta)} \geq 4d.$$

Of course (3.17) remains valid if  $k$  is replaced by any  $(\mathcal{F}_{n-1})$ -measurable random quantity. With this in mind we set

$$\Theta_n := \left\lceil \log_2 \left( \frac{A}{d \max_{x \in \mathbb{Z}^d} D\widehat{W}_{n-1}^\beta(x)} \right) \right\rceil \geq 1.$$

We obtain that on the event  $\{\tau \geq n\}$  we have

$$\kappa^{\Theta_n} \leq \mathbb{P}[W_{n+\Theta_n-1} \geq A \mid \mathcal{F}_{n-1}] \leq \mathbb{P}[\tau \in \llbracket n, n + \Theta_n - 1 \rrbracket \mid \mathcal{F}_n]. \quad (3.18)$$

Hence we have

$$\begin{aligned} \mathbb{E} \left[ \mathbf{1}_{\{\tau \geq n\}} \left( \max_{x \in \mathbb{Z}^d} A^{-1} D \widehat{W}_{n-1}^\beta(x) \right)^{q/2} \right] &\leq (2d)^{q/2} \mathbb{E} \left[ \mathbf{1}_{\{\tau \geq n\}} 2^{-\frac{\Theta_n q}{2}} \kappa^{-\Theta_n + \Theta_n} \right] \\ &\leq (2d)^{q/2} \mathbb{E} \left[ \mathbf{1}_{\{\tau \in \llbracket n, n + \Theta_n - 1 \rrbracket\}} (2^{q/2} \kappa)^{-\Theta_n} \right]. \end{aligned} \quad (3.19)$$

Taking the sum over  $n$  we obtain

$$\sum_{n \geq 1} \mathbb{E} \left[ \mathbf{1}_{\{\tau \geq n\}} \left( \max_{x \in \mathbb{Z}^d} A^{-1} D \widehat{W}_{n-1}^\beta(x) \right)^{q/2} \right] \leq (2d)^{q/2} \sum_{n \geq 1} \mathbb{E} \left[ \Theta_n (2^{q/2} \kappa)^{-\Theta_n} \right]. \quad (3.20)$$

and with our choice for  $q$  (3.12) the r.h.s. is finite. This concludes the proof of (3.15).  $\square$

**3.3. Proof of Corollary 2.5.** Recalling (2.17) and (2.25) we have for any  $k \leq n$

$$W_n^\beta / W_k^\beta := \sum_{x \in \mathbb{Z}^d} \mu_k^\beta(x) \theta_{k,x}(W_{n-k}^\beta). \quad (3.21)$$

Hence for  $p \geq 1$ , Jensen's inequality applied twice yields

$$\mathbb{E} \left[ \left( W_n^\beta / W_k^\beta \right)^p \mid \mathcal{F}_k \right] \leq \mathbb{E}[(W_{n-k}^\beta)^p] \leq \mathbb{E}[(W_n^\beta)^p]. \quad (3.22)$$

Hence for any  $A > 1$ ,  $p > 1$  and  $n \in \mathbb{N}$ , we have

$$\begin{aligned} \mathbb{E}[(W_n^\beta)^p] &= \mathbb{E} \left[ (W_n^\beta)^p \mathbf{1}_{\{\tau_A > n\}} \right] + \sum_{k=1}^n \mathbb{E}[(W_n^\beta)^p \mathbf{1}_{\{\tau_A = k\}}] \\ &\leq A^p + \sum_{k \leq n} \mathbb{E} \left[ (W_k^\beta)^p \mathbf{1}_{\{\tau_A = k\}} \mathbb{E} \left[ \left( W_n^\beta / W_k^\beta \right)^p \mid \mathcal{F}_k \right] \right] \\ &\leq A^p + \mathbb{E}[(W_n^\beta)^p] \sum_{k \leq n} \mathbb{E}[(W_k^\beta)^p \mathbf{1}_{\{\tau_A = k\}}] \\ &\leq A^p + \mathbb{E}[(W_n^\beta)^p] \mathbb{E}[(W_{\tau_A}^\beta)^p \mathbf{1}_{\{\tau_A < \infty\}}] \\ &\leq A^p + C_p A^p \mathbb{P}(\tau_A < \infty) \mathbb{E}[(W_n^\beta)^p], \end{aligned} \quad (3.23)$$

where we have used (3.22) in the first inequality and (2.11) in the last one. Then, since  $\mathbb{E}[(W_n^\beta)^p]$  is unbounded for  $p > p^*(\beta)$ , (3.23) implies that

$$C_p A^p \mathbb{P}(\tau_A < \infty) \geq 1. \quad (3.24)$$

Taking the limit as  $p \downarrow p^*(\beta)$  (recall (2.12)) yields the desired result with  $c = (C_{p^*(\beta)})^{-1}$ .  $\square$

**3.4. Proof of Corollary 2.7(i).** We simply observe that

$$\mathbb{E} \left[ \sup_{n \geq 1} W_n^\beta \right] = 1 + \int_1^\infty \mathbb{P}[\tau_A < \infty] dA.$$

Since by [Jun22b, Theorem 1.1(i)],  $\mathbb{E}[\sup_{n \geq 1} W_n^\beta] < \infty$  in the weak disorder phase, in view of (2.13) we must necessarily have  $p^*(\beta) > 1$ .  $\square$

## 4. PROVING THEOREM 2.1

**4.1. Organizing the proof of Theorem 2.1.** Note that (2.8) displays a total of 6 comparisons of the form  $f(u) \asymp g(u)$ , corresponding to 12 inequalities. Of course some of these inequalities are redundant and in the end, it is sufficient to prove 6 “independent” inequalities. Before we begin the formal proof, let us briefly expose which inequalities we prove. First we observe that

$$\widehat{W}_\infty^{\beta,*} \leq W_\infty^{\beta,*} \quad \text{and} \quad W_\infty^\beta \leq W_\infty^{\beta,*}, \quad (4.1)$$

and hence two inequalities are immediate. A third inequality, that is (2.13) in Corollary 2.5, has been proved in the previous section. This leaves us with three remaining inequalities to prove. The first one is proved in Section 4.2, using the supermultiplicativity of the point-to-point partition functions.

**Lemma 4.1.** *In the weak disorder regime, we have for any  $u > 1$ ,*

$$\mathbb{P} \left( \widehat{W}_\infty^{\beta,*} > u \right) \leq u^{-p^*(\beta)}. \quad (4.2)$$

The next lemma does not exactly match an inequality in (2.8) (because of the factor 4 in the r.h.s.) but combined with the others it is sufficient to prove the theorem. It is proved in Section 4.3.

**Lemma 4.2.** *There exists a constant  $c$  such that for every  $u > 1$*

$$\mathbb{P} \left[ W_\infty^\beta \geq u \right] \geq c \mathbb{P} [W_\infty^{\beta,*} \geq 4u]. \quad (4.3)$$

The most technical part of the proof is the comparison of the tail of  $\widehat{W}_\infty^{\beta,*}$  (the maximum over point-to-point partition functions) with that of  $W_\infty^{\beta,*}$  (the maximum over point-to-plane partition functions). We achieve this by proving a localization result for the endpoint at time  $\tau_u$ , which is interesting in its own right. The outcome of our proof (presented in Sections 4.4 and 4.5) is the following comparison which combined with the five previous inequalities allows to complete the proof of Theorem 2.1.

**Proposition 4.3.** *There exist constants  $C > 1$ ,  $\delta > 0$  and a closed set  $\mathcal{U} \subset [1, \infty)$  satisfying*

$$\forall v \geq 1, \quad [v, Cv] \cap \mathcal{U} \neq \emptyset, \quad (4.4)$$

*(the set  $\{\log u : u \in \mathcal{U}\}$  has no gap larger than  $\log C$ ) such that the following holds*

$$\forall u \in \mathcal{U}, \quad \mathbb{P} \left[ \max_{x \in \mathbb{Z}^d} \mu_{\tau_u}(x) \geq \delta \mid \tau_u < \infty \right] \geq \delta. \quad (4.5)$$

**Remark 4.4.** *Observe that the above proposition is an intermediate result on the path of obtaining Theorem 2.1. Once Theorem 2.1 is proven, one obtains that for  $\delta > 0$  sufficiently small*

$$\forall u \geq 1, \quad \mathbb{P} \left[ \max_{x \in \mathbb{Z}^d} \mu_{\tau_u}(x) \geq \delta \mid \tau_u < \infty \right] \geq \delta, \quad (4.6)$$

*with no need for a restriction to a specific set  $\mathcal{U}$ .*

We end up this subsection by showing that indeed the combination of all the element exposed above yields our main result.

*Proof of Theorem 2.1.* Let us first show that  $\mathbb{P} \left[ W_\infty^{\beta,*} \geq u \right]$  and  $\mathbb{P} \left[ W_\infty^\beta \geq u \right]$  are comparable to  $u^{-p^*}$ . As two of the four required inequalities are provided by (4.2) and (2.13), it only remains

to show that for some constant  $\kappa > 0$  we have

$$\begin{aligned} \forall u \geq 1, \quad \mathbb{P} \left[ W_{\infty}^{\beta,*} \geq u \right] &\leq \frac{1}{\kappa} u^{-p^*(\beta)}, \\ \forall u \geq 1, \quad \mathbb{P} \left[ \widehat{W}_{\infty}^{\beta,*} \geq u \right] &\geq \kappa u^{-p^*(\beta)}. \end{aligned} \quad (4.7)$$

From Proposition 4.3, we can find  $C, \delta$  and  $\mathcal{U}$  such that (4.4) and (4.5) hold. Let us define  $u'$  and  $u''$  to be the nearest points to  $u$  in  $\mathcal{U}$

$$u' := \max\{v \in \mathcal{U} : v \leq u\} \text{ and } u'' := \min\{v \in \mathcal{U} : \delta v \geq u\} \quad (4.8)$$

Note that from (4.4) we have  $u/u' \in [1, C]$  and  $u''/u \in [1/\delta, C/\delta]$ . Both  $\mathbb{P} \left[ W_{\infty}^{\beta,*} \geq u \right]$  and  $\mathbb{P} \left[ \widehat{W}_{\infty}^{\beta,*} \geq u \right]$  are decreasing in  $u$ . Thus, at the cost of changing the value of the constant  $\kappa$ , it is sufficient to show that the first line in (4.7) is valid for  $u'$  and that the second line is valid for  $\delta u''$ . From (4.5) we have for any  $v \in \mathcal{U}$

$$\mathbb{P} \left( \widehat{W}_{\infty}^{\beta,*} \geq \delta v \right) \geq \mathbb{P} \left( \max_{x \in \mathbb{Z}^d} \widehat{W}_{\tau_v}^{\beta}(x) \geq \delta v ; \tau_v < \infty \right) \geq \delta \mathbb{P}(\tau_v < \infty). \quad (4.9)$$

Combining this with (2.13) (and writing  $p^*$  for  $p^*(\beta)$  for better readability) we obtain that

$$\mathbb{P} \left( \widehat{W}_{\infty}^{\beta,*} \geq u \right) \geq \mathbb{P} \left( \widehat{W}_{\infty}^{\beta,*} \geq \delta u'' \right) \geq \delta c (u'')^{-p^*} \geq c C^{-p^*} \delta^{1+p^*} u^{-p^*}. \quad (4.10)$$

On the other hand we have also from (4.9), using this time (4.2),

$$\mathbb{P} \left[ W_{\infty}^{\beta,*} \geq u \right] \leq \mathbb{P}(\tau_{u'} < \infty) \leq \delta^{-1} \mathbb{P} \left( \widehat{W}_{\infty}^{\beta,*} \geq \delta u' \right) \leq \delta^{-1-p^*} (u')^{-p^*} \leq \delta^{-1-p^*} u^{-p^*}, \quad (4.11)$$

finishing the proof of (4.7).

To conclude, let us prove that  $\mathbb{P} \left[ W_{\infty}^{\beta} \geq u \right] \asymp u^{-p^*}$ . The upper bound is a direct consequence of the first line in (4.7) since  $W_{\infty}^{\beta} \leq W_{\infty}^{\beta,*}$ . The lower bound it is obtained by combining Lemma 4.2 with (2.13).  $\square$

**4.2. Proof of Lemma 4.1.** We can focus on the case  $p^*(\beta) > 1$  since when  $p^*(\beta) = 1$  the result is an immediate consequence of Markov's inequality. The result follows from the observation that setting  $\theta(u) := \mathbb{P} \left( \sup_{n \geq 0, x \in \mathbb{Z}^d} \widehat{W}_n^{\beta}(x) > u \right)$  we have

$$\forall u, v > 1, \quad \theta(uv) \geq \theta(u)\theta(v) \quad (4.12)$$

Now, for any  $u > 1$ ,  $k \geq 1$  and  $p \in (1, p^*(\beta))$  we have

$$(\theta(u)u^p)^k \leq \theta(u^k)u^{kp} \leq \mathbb{E} \left[ \left( \widehat{W}_{\infty}^{\beta,*} \right)^p \right] \leq \mathbb{E} \left[ \left( W_{\infty}^{\beta,*} \right)^p \right] < \infty. \quad (4.13)$$

By taking  $k \rightarrow \infty$ , this implies that  $\theta(u) \leq u^{-p}$ , and we obtain the result by taking the limit  $p \uparrow p^*(\beta)$ . Now let us justify (4.12). Set  $B_u := \{\sup_{n \geq 0, x \in \mathbb{Z}^d} \widehat{W}_n^{\beta}(x) > u\}$ , so that  $\theta(u) = \mathbb{P}(B_u)$ . We define

$$(\sigma, Z) := \min \left\{ (n, x) \in \mathbb{N} \times \mathbb{Z}^d : \widehat{W}_n(x) > u \right\}$$

where min refers to the minimal element of the set for the lexicographical order on  $\mathbb{N} \times \mathbb{Z}^d$ . By convention  $(\sigma, Z) = (\infty, 0)$  on the event  $B_u^c$ . Recalling the definition (2.17) we have, for any  $n, m \geq 0$  and  $x, z \in \mathbb{Z}^d$ ,

$$\widehat{W}_{n+m}(x+z) \geq \mathbb{E} \left[ e^{\beta H_{n+m}(X) - (n+m)\lambda(\beta)} \mathbf{1}_{\{X_m=z, X_{n+m}=x+z\}} \right] = \widehat{W}_m(z) \theta_{m,z} \widehat{W}_n(x). \quad (4.14)$$

Hence  $\widehat{W}_{n+m}(x+z) \geq uv$  if  $\widehat{W}_m(z) \geq u$  and  $\theta_{m,z}\widehat{W}_n(x) \geq v$ . Decomposing over the possible values of  $(\sigma, Z)$  we have

$$\bigcup_{\substack{m \geq 1, z \in \mathbb{Z}^d \\ n \geq 1, x \in \mathbb{Z}^d}} \{(\sigma, Z) = (m, z)\} \cap \theta_{m,z}B_v \subset B_{uv}. \quad (4.15)$$

Now the events  $(\sigma, Z) = (m, z)$  and  $\theta_{m,z}B_v$  rely on disjoint regions of the i.i.d. environment and are hence independent. Combining this observation with translation invariance and the fact that  $\{(\sigma, Z) = (m, z)\}_{m \geq 1, z \in \mathbb{Z}^d}$  is a partition of the event  $B_u$ , we obtain that

$$\begin{aligned} \mathbb{P}(B_{uv}) &\geq \sum_{m \geq 1, z \in \mathbb{Z}^d} \mathbb{P}(\{(\sigma, Z) = (m, z)\} \cap \theta_{m,z}B_v) \\ &= \sum_{m \geq 1, z \in \mathbb{Z}^d} \mathbb{P}((\sigma, Z) = (m, z)) \mathbb{P}(B_v) = \mathbb{P}(B_u) \mathbb{P}(B_v). \end{aligned} \quad \square$$

**4.3. Proof of Lemma 4.2.** Set  $A = 4u$ . On the event  $\tau_A < \infty$  we have

$$W_\infty^\beta = W_{\tau_A}^\beta \sum_{x \in \mathbb{Z}^d} D\mu_{\tau_A-1}(x) \theta_{\tau_A, x} W_\infty^\beta \geq A \sum_{x \in \mathbb{Z}^d} D\mu_{\tau_A-1}(x) \theta_{\tau_A, x} (W_\infty^\beta). \quad (4.16)$$

Let us set  $Z^A(x) := (\theta_{\tau_A, x} W_\infty^\beta)$ . Note that  $(Z_x^A)_{x \in \mathbb{Z}}$  is independent of  $\mathcal{F}_{\tau_A}$  and distributed like  $(Z(x))_{x \in \mathbb{Z}^d} := (\theta_{0, x} W_\infty^\beta)_{x \in \mathbb{Z}^d}$ . For this reason we have for any  $u \geq 0$

$$\mathbb{P}(W_\infty^\beta \geq u) \geq \mathbb{P}[\tau_A < \infty] \inf_{\alpha \in \mathcal{P}(\mathbb{Z}^d)} \mathbb{P}\left(\sum_{x \in \mathbb{Z}^d} \alpha(x) Z(x) \geq 1/4\right) \quad (4.17)$$

where the infimum is taken over all probability measures  $\alpha$  on  $\mathbb{Z}^d$ . To conclude we just need to show that this infimum is positive. Now if one sets  $\mathcal{W}_\alpha := \sum_{x \in \mathbb{Z}^d} \alpha(x) Z(x)$ , it is immediate to check that the collection of variables  $(\mathcal{W}_\alpha)_{\alpha \in \mathcal{P}_{\mathbb{Z}^d}}$  is uniformly integrable. Indeed if we let  $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$  be a convex function such that  $\mathbb{E}[\varphi(W_\infty^\beta)] < \infty$  and  $\lim_{x \rightarrow \infty} \varphi(x)/x = \infty$  (such a  $\varphi$  exists since  $W_\infty^\beta \in L^1$ ), Jensen's inequality yields that for any  $\alpha \in \mathcal{P}(\mathbb{Z}^d)$

$$\mathbb{E}[\varphi(\mathcal{W}_\alpha)] \leq \sum_{x \in \mathbb{Z}^d} \alpha(x) \mathbb{E}[\varphi(Z(x))] = \mathbb{E}[\varphi(W_\infty^\beta)], \quad (4.18)$$

implying the desired uniform integrability. Since  $\mathbb{E}[\mathcal{W}_\alpha] = 1$ , using uniform integrability there exists  $M > 0$  such that for every  $\alpha \in \mathcal{P}(\mathbb{Z}^d)$

$$\begin{aligned} \mathbb{E}[\mathcal{W}_\alpha \mathbf{1}_{\{\mathcal{W}_\alpha \in [1/4, M]\}}] &= 1 - \mathbb{E}[\mathcal{W}_\alpha \mathbf{1}_{\{\mathcal{W}_\alpha < 1/4\}}] - \mathbb{E}[\mathcal{W}_\alpha \mathbf{1}_{\{\mathcal{W}_\alpha > M\}}] \\ &\geq 3/4 - \mathbb{E}[\mathcal{W}_\alpha \mathbf{1}_{\{\mathcal{W}_\alpha > M\}}] \geq 1/2, \end{aligned} \quad (4.19)$$

and thus we have

$$\mathbb{P}(\mathcal{W}_\alpha \geq 1/4) \geq \frac{1}{M} \mathbb{E}[\mathcal{W}_\alpha \mathbf{1}_{\{\mathcal{W}_\alpha \in [1/4, M]\}}] \geq \frac{1}{2M}.$$

□

**4.4. Proof of Proposition 4.3.** We start by fixing two parameters

$$q > p^*(\beta) \quad \text{and} \quad D := 3C_{2q}2^{2q} \quad (4.20)$$

where  $C_{2q} \geq 1$  comes from Proposition 2.3. The reason for this peculiar choice for  $D$  will become apparent in the course of the proof of Proposition 4.5 below. We start with the geometric sequence/set  $\mathcal{U}_0 := \{D^k : k \geq 0\}$ . Note that  $\mathcal{U}_0$  clearly satisfies the density requirement (4.4). We are going to show (4.5) is valid for a “large” subset of  $\mathcal{U}_0$  such that (4.4) remains true. Given  $q > p^*$ , we set

$$\mathcal{V} := \{u > 1 : \mathbb{P}(\tau_{Du} < \infty) \geq D^{-q}\mathbb{P}(\tau_u < \infty)\} \quad \text{and} \quad \mathcal{U}_1 := \mathcal{U}_0 \cap \mathcal{V}. \quad (4.21)$$

Note that  $\mathcal{U}_1$  is infinite. Indeed if this was not the case, then there would exist some  $k_0$  such that the sequence  $(D^{kq}\mathbb{P}(\tau_{D^k} < \infty))_{k \geq k_0}$  is non-increasing (and hence bounded). Since  $q > p^*$ , this would be a contradiction to (2.13). The next step is then to prove that  $u \in \mathcal{V}$  satisfies the desired localization property (4.5). This is the most important step, its proof is postponed to the next subsection.

**Proposition 4.5.** *There exist  $\delta > 0$  and  $D > 2$  such that for every  $u \in \mathcal{V}$*

$$\mathbb{P}\left(\max_{x \in \mathbb{Z}^d} \mu_{\tau_u}(x) \geq \delta \mid \tau_u < \infty\right) \geq \delta. \quad (4.22)$$

**Remark 4.6.** *Note that a posteriori, Theorem 2.1 implies that  $\mathcal{V} = [1, \infty)$  if  $D$  is chosen sufficiently large. Hence Proposition 4.5 implies that (4.6) holds.*

Finally to conclude we need to show that there are no big gaps in  $\mathcal{U}_1$ . This is the role of the following lemma.

**Lemma 4.7.** *There exists an integer  $k_0 \geq 1$  such that*

$$\forall u \in \mathcal{U}_1, \quad \{uD^i : i \in \llbracket 1, k_0 \rrbracket\} \cap \mathcal{U}_1 \neq \emptyset \quad (4.23)$$

*In particular  $\mathcal{U}_1$  satisfies (4.4) with  $C = D^{k_0}$ .*

Proposition 4.3 follows immediately from the combination of Proposition 4.5 and Lemma 4.7.

*Proof of Lemma 4.7.* For  $k_0$  to be chosen later, let us assume that we can find  $u \in \mathcal{U}_1$  such that (4.23) does not hold. Then we have

$$\mathbb{P}(\tau_{uD^{k_0}} < \infty) \leq D^{-q(k_0-1)}\mathbb{P}(\tau_{uD} < \infty) \leq D^{-q(k_0-1)}\mathbb{P}(\tau_u < \infty). \quad (4.24)$$

On the other hand, since  $u \in \mathcal{U}_1$ , Proposition 4.5 implies that (4.5) holds. Recalling the computation (4.9) this implies that

$$\mathbb{P}(\tau_u < \infty) \leq \delta^{-1}\mathbb{P}\left(\widehat{W}_\infty^{\beta,*} \geq \delta u\right) \leq \delta^{-p^*-1}u^{-p^*}, \quad (4.25)$$

where the last bound is simply (4.2). We conclude that

$$\mathbb{P}(\tau_{uD^{k_0}} < \infty) \leq D^{-q(k_0-1)}\delta^{-p^*-1}u^{-p^*}. \quad (4.26)$$

Now recalling that  $q > p^*$ , the bound contradicts (2.13) if  $k_0$  is large enough that

$$D^{-q(k_0-1)}\delta^{-p^*-1} \leq cD^{-p^*k_0},$$

where  $c$  is the constant in (2.13). □

**4.5. Proof of Proposition 4.5.** The idea of the proof is to show that if the probability  $\mu_{\tau_u}$  is too spread-out then the conditional probability  $\mathbb{P}(\tau_{Du} < \infty \mid \tau_u < \infty)$  is small, which in turns implies that  $u \notin \mathcal{V}$ . Note that on the event  $\tau_{Du} < \infty$ , either  $W_{\tau_u}$  is much larger than  $u$  or  $(W_{n+\tau_u}/W_{\tau_u})_{n \geq 1}$  overshoots 2,

$$\mathbb{P}(\tau_{Du} < \infty \mid \tau_u < \infty) \leq \mathbb{P}\left(W_{\tau_u} \geq \frac{Du}{2} \mid \tau_u < \infty\right) + \mathbb{P}\left(\sup_{n \geq 0} \frac{W_{n+\tau_u}}{W_{\tau_u}} \geq 2 \mid \tau_u < \infty\right). \quad (4.27)$$

The first term can be controlled by Proposition 2.3. We have

$$\mathbb{P}\left(W_{\tau_u} \geq \frac{Du}{2} \mid \tau_u < \infty\right) \leq \left(\frac{Du}{2}\right)^{-2q} \mathbb{E}[|W_{\tau_u}|^{2q} \mid \tau_u < \infty] \leq C_{2q} 2^{2q} D^{-2q} \leq \frac{D^{-q}}{3}, \quad (4.28)$$

where the last inequality is valid thanks to our choice for  $D$  (4.20). We are going to show that if (4.22) is violated for a value of  $\delta$  to be determined in the course of the proof, then we would also have

$$\mathbb{P}\left(\sup_{n \geq 0} \frac{W_{n+\tau_u}}{W_{\tau_u}} \geq 2 \mid \tau_u < \infty\right) \leq \frac{D^{-q}}{3}. \quad (4.29)$$

Combined with (4.27) and (4.28), this implies that  $u \notin \mathcal{V}$  and hence a contradiction. We first note that

$$\frac{W_{n+\tau_u}}{W_{\tau_u}} = \sum_{x \in \mathbb{Z}^d} \mu_{\tau_u}(x) \theta_{\tau_u, x} W_n^\beta. \quad (4.30)$$

Now by Markov's property  $(\theta_{\tau_u, x} W_n^\beta)_{n \geq 0, x \in \mathbb{Z}^d}$  is independent from  $\mathcal{F}_{\tau_u}$ . For this reason we define  $\Gamma: \mathcal{P}(\alpha) \rightarrow \mathbb{R}_+$  by

$$\Gamma(\alpha) := \mathbb{P}\left(\sup_{n \geq 0} \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0, x} W_n^\beta \geq 2\right), \quad (4.31)$$

so that, on the event  $\tau_u < \infty$ , we have

$$\mathbb{P}\left[\sup_{n \geq 0} \frac{W_{n+\tau_u}}{W_{\tau_u}} \geq 2 \mid \mathcal{F}_{\tau_u}\right] = \Gamma(\mu_{\tau_u}). \quad (4.32)$$

In particular, we have

$$\mathbb{P}\left(\sup_{n \geq 0} \frac{W_{n+\tau_u}}{W_{\tau_u}} \geq 2 \mid \tau_u < \infty\right) = \mathbb{E}[\Gamma(\mu_{\tau_u}) \mid \tau_u < \infty]. \quad (4.33)$$

Now the idea is that if the probability  $\alpha$  is very spread out, good mixing properties of the field  $(\theta_{0, x} W_n^\beta)_{x \in \mathbb{Z}^d}$  (here we use slightly more than ergodicity) imply that  $\sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0, x} W_n^\beta$  is concentrated around its mean which is equal to one and thus that  $\Gamma(\alpha)$  is small. This is stated in the following lemma, whose proof is postponed to the end of this section.

**Lemma 4.8.** *For any  $\varepsilon > 0$ , there exists  $\eta = \eta(\varepsilon)$  such that for any  $\alpha \in \mathcal{P}(\mathbb{Z}^d)$*

$$\max_{x \in \mathbb{Z}^d} \alpha(x) \leq \eta \quad \Rightarrow \quad \Gamma(\alpha) \leq \varepsilon. \quad (4.34)$$

To conclude, let us fix  $\delta := \min(\eta(D^{-2q}), D^{-2q})$  – where  $\eta$  is the function given by the above lemma – and assume that (4.22) is violated for this value of  $\delta$ . Since  $\Gamma(\mu_{\tau_u}) \leq 1$ , the combination of (4.33) and (4.34) yields

$$\mathbb{E}[\Gamma(\mu_{\tau_u}) \mid \tau_u < \infty] \leq D^{-2q} + \mathbb{P}\left(\max_{x \in \mathbb{Z}^d} \mu_{\tau_u}(x) > \delta \mid \tau_u < \infty\right) \leq D^{-2q} + \delta \leq 2D^{-2q}, \quad (4.35)$$

which, with our choice for  $D$  (4.20), concludes the proof of (4.29) and hence of the proposition.  $\square$



*Proof of Lemma 4.8.* We prove that there exists  $\eta$  such that if  $\max_{x \in \mathbb{Z}^d} \alpha(x) \leq \eta$  then

$$\mathbb{E} \left[ \sup_{n \geq 0} \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_n^\beta - 1 \right] \leq \varepsilon \quad (4.36)$$

and then conclude using Markov's inequality (note that the supremum is non-negative). The first step is to reduce the range over which the sup is taken. For any  $M$ , we have

$$\begin{aligned} \mathbb{E} \left[ \sup_{n \geq 0} \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_n^\beta \right] &\leq \mathbb{E} \left[ \sup_{n \in \llbracket 0, M \rrbracket} \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_n^\beta \right] \\ &\quad + \sum_{x \in \mathbb{Z}^d} \alpha(x) \mathbb{E} \left[ \sup_{i \geq 0} \theta_{0,x} W_{M+i}^\beta - \theta_{0,x} W_M^\beta \right] \end{aligned} \quad (4.37)$$

Now since  $\alpha$  is a probability, by translation invariance we have

$$\sum_{x \in \mathbb{Z}^d} \alpha(x) \mathbb{E} \left[ \sup_{i \geq 1} \theta_{0,x} W_{M+i}^\beta - \theta_{0,x} W_M^\beta \right] = \mathbb{E} \left[ \sup_{i \geq 0} W_{M+i}^\beta - W_M^\beta \right]. \quad (4.38)$$

Note that we have

$$\lim_{M \rightarrow \infty} \left( \sup_{i \geq 0} W_{M+i}^\beta - W_M^\beta \right) = 0 \quad \text{and} \quad \sup_{i \geq 0} \left| W_{M+i}^\beta - W_M^\beta \right| \leq W_\infty^{\beta,*} \quad (4.39)$$

Since by [Jun22b, Theorem 2.1(i)],  $W_\infty^{\beta,*} \in L^1$ , we can use dominated convergence to obtain that for  $M$  sufficiently large we have

$$\mathbb{E} \left[ \sup_{i \geq 0} W_{M+i}^\beta - W_M^\beta \right] \leq \varepsilon/2 \quad (4.40)$$

and we are left with estimating the first term in (4.37). First observe that (by Cauchy-Schwarz)

$$\mathbb{E} \left[ \sup_{n \in \llbracket 0, M \rrbracket} \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_n^\beta - 1 \right] \leq \mathbb{E} \left[ \sup_{n \in \llbracket 0, M \rrbracket} \left( \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_n^\beta - 1 \right)^2 \right]^{1/2} \quad (4.41)$$

Then we use Doob's  $L^2$ -inequality (for the martingale  $\sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_n^\beta - 1$ ) and obtain that

$$\mathbb{E} \left[ \sup_{n \in \llbracket 0, M \rrbracket} \left( \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_n^\beta - 1 \right)^2 \right] \leq 4 \mathbb{E} \left[ \left( \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_M^\beta - 1 \right)^2 \right] \quad (4.42)$$

Now we can conclude by observing that

$$\begin{aligned} \mathbb{E} \left[ \left( \sum_{x \in \mathbb{Z}^d} \alpha(x) \theta_{0,x} W_M^\beta - 1 \right)^2 \right] &\leq \sum_{\substack{x, x' \in \mathbb{Z}^d \\ |x-x'| \leq 2M}} \alpha(x) \alpha(x') \mathbb{E} \left[ \left( W_M^\beta - 1 \right)^2 \right] \\ &\leq (4M+1)^d e^{M(\lambda(2\beta)-2\lambda(\beta))} \eta, \end{aligned} \quad (4.43)$$

which is bounded by  $\frac{\varepsilon}{2}$  if  $\eta$  is sufficiently small.  $\square$

## 5. PROOF OF THE COROLLARIES FROM SECTION 2.4

**5.1. Preparation for the proof of Corollary 2.7(ii).** We are going to prove that  $p^*(\beta) < 1 + \frac{2}{d}$  implies strong disorder. To do so we borrow a couple of elements of proof from [JL24]. This includes using a characterization of strong disorder via the size-biased measure, as well as an estimate of the tail distribution of the supremum of the partition function  $W_n^\beta$  take over a finitely many values of  $n$ .

The size-biased measure  $\tilde{\mathbb{P}}_n$  defined by  $d\tilde{\mathbb{P}}_n := W_n^\beta d\mathbb{P}$ . When weak disorder holds,  $\tilde{\mathbb{P}}_n$  converges when  $n \rightarrow \infty$  to a measure  $\tilde{\mathbb{P}}_\infty$  which is absolutely continuous w.r.t.  $\mathbb{P}$ . On the other hand, when strong disorder holds,  $\tilde{\mathbb{P}}_n$  and  $\mathbb{P}$  become increasingly singular as  $n$  grows. Hence to prove that strong disorder holds, it is sufficient to find an event  $A_n$  which has large probability under  $\mathbb{P}$  and small probability under  $\tilde{\mathbb{P}}_n$ . The following lemma, although not strictly necessary, helps to understand this dichotomy. Its proof can be found in [JL24, Lemma 3.2].

**Lemma 5.1.** *For any event  $A$  we have*

$$\mathbb{P} \left[ (W_n^\beta)^{1/2} \right] \leq \sqrt{\mathbb{P}(A^c)} + \sqrt{\tilde{\mathbb{P}}_n(A)} \quad (5.1)$$

Our goal is thus to identify a sequence of events  $A_n$  such that

$$\lim_{n \rightarrow \infty} \mathbb{P}(A_n^c) = \lim_{n \rightarrow \infty} \tilde{\mathbb{P}}_n(A_n) = 0. \quad (5.2)$$

Before introducing the event  $A_n$ , we state a key technical result that will be essential to obtain an estimate on  $\tilde{\mathbb{P}}_n(A_n)$ .

**Lemma 5.2.** *Assume that weak disorder holds at  $\beta$ . Given  $\varepsilon > 0$ , there exist constants  $C_\varepsilon > 0$  and  $u_0(\varepsilon) > 0$  such that for every  $u \geq u_0(\varepsilon)$*

$$\exists m \in \llbracket 0, C_\varepsilon \log u \rrbracket, \quad \mathbb{P} \left( W_m^\beta \geq u \right) \geq u^{-p^*(\beta) - \varepsilon}. \quad (5.3)$$

The above result is the analogue of [JL24, Proposition 4.2] which states that (5.3) is valid when strong disorder holds (in which case  $p^*(\beta) = 1$ ) under the additional assumption that the environment is upper-bounded. The proof of the present lemma is almost identical, except that we use Theorem 2.1 (more specifically, the asymptotic behavior of  $\widehat{W}_n^{\beta,*}$ ) in place of [JL24, Theorem 4.4] as an input. We therefore only sketch the argument.

*Proof.* First, by an inductive argument similar to the one used to prove (4.12) (we refer to the proof of [JL24, Proposition 6.1] for details), we obtain, for any  $k, T \in \mathbb{N}$  and  $u > 1$ ,

$$\mathbb{P} \left( \max_{n \in \llbracket 1, kT \rrbracket, x \in \mathbb{Z}^d} \widehat{W}_n^\beta(x) > v^k \right) \geq \mathbb{P} \left( \max_{n \in \llbracket 1, T \rrbracket, x \in \mathbb{Z}^d} \widehat{W}_n^\beta(x) \geq v \right)^k. \quad (5.4)$$

Then we fix  $v_1$  large enough that  $v_1^{-\varepsilon/2} \leq \frac{c}{2}$ , where  $c$  is such that  $\mathbb{P} \left( \widehat{W}_\infty^{\beta,*} > u \right) \geq cu^{-p^*}$  holds for all  $u > 1$  (such a constant exists due to Theorem 2.1). By choosing  $T$  large enough, we have

$$\mathbb{P} \left( \max_{n \in \llbracket 1, T \rrbracket, x \in \mathbb{Z}^d} \widehat{W}_n(x) \geq v_1 \right) \geq \frac{c}{2} v_1^{-p^*}. \quad (5.5)$$

Now given  $u$  we set  $k = k_u := \lfloor \log u / \log v_1 \rfloor$ . By a union bound and (5.4)–(5.5), we obtain

$$\begin{aligned} \max_{n \in \llbracket 1, kT \rrbracket} \mathbb{P} \left( W_n^\beta > u \right) &\geq \frac{1}{kT} \mathbb{P} \left( \max_{n \in \llbracket 1, kT \rrbracket} W_n^\beta > u \right) \\ &\geq \frac{1}{kT} \mathbb{P} \left( \max_{n \in \llbracket kT \rrbracket, x \in \mathbb{Z}^d} \widehat{W}_n^\beta(x) > v_1^k \right) \geq \left( \frac{c}{2} v_1^{-p^*} \right)^k \geq v_1^{-k(p^* + \varepsilon/2)} \geq \left( \frac{u}{v_1} \right)^{-(p^* + \varepsilon/2)} \end{aligned} \quad (5.6)$$

It is then easy to see that the claim holds with  $u_0 = (v_1)^{\frac{2p^*}{\varepsilon}+1}$  and  $C_\varepsilon := \frac{T}{\log v_1}$ .  $\square$

The last ingredient we need is a well-known construction which allows for a simple representation of the size-biased measure, which is sometimes called the *spine construction*, see for instance [Bir04, Lemma 1]. We define  $(\widehat{\omega}_i)_{i \geq 1}$  as a sequence of i.i.d. random variable with marginal distribution given by

$$\widehat{\mathbb{P}}[\widehat{\omega}_1 \in \cdot] = \mathbb{E} \left[ e^{\beta \omega_{1,0} - \lambda(\beta)} \mathbf{1}_{\{\omega_{1,0} \in \cdot\}} \right], \quad (5.7)$$

and  $X$  a simple random walk ( $\widehat{\mathbb{P}}$  and  $P$  denote the respective distributions). Given  $\omega$ ,  $\widehat{\omega}$  and  $X$ , all sampled independently, we define a new environment  $\widetilde{\omega} = \widetilde{\omega}(X, \omega, \widehat{\omega})$  by

$$\widetilde{\omega}_{i,x} := \begin{cases} \omega_{i,x} & \text{if } x \neq X_i, \\ \widehat{\omega}_i & \text{if } x = X_i. \end{cases} \quad (5.8)$$

In words,  $\widetilde{\omega}$  is obtained by tilting the distribution of the environment on the graph of  $(i, X_i)_{i=1}^\infty$ . The distribution of  $\widetilde{\omega}$  under  $P \otimes \mathbb{P} \otimes \widehat{\mathbb{P}}$  corresponds to that of  $\omega$  under the size-biased measure.

**Lemma 5.3.** *It holds that*

$$\widehat{\mathbb{P}}_n[(\omega_{i,x})_{i \in [1,n], x \in \mathbb{Z}^d} \in \cdot] = P \otimes \mathbb{P} \otimes \widehat{\mathbb{P}}[(\widetilde{\omega}_{i,x})_{i \in [1,n], x \in \mathbb{Z}^d} \in \cdot]. \quad (5.9)$$

We refer to [JL24, Lemma 3.3] for a proof of the above using the same notation.

**5.2. Proof of Corollary 2.7(ii).** Assuming that  $p^*(\beta) < 1 + \frac{2}{d}$ , we fix three parameters

$$\zeta \in \left( p^*(\beta), 1 + \frac{2}{d} \right), \quad \alpha \in \left( \frac{2+d}{2\zeta}, \frac{1}{\zeta-1} \right) \quad \text{and} \quad \xi \in \left( \frac{1}{2}, \frac{1-\alpha\zeta}{d} \right). \quad (5.10)$$

Using Lemma 5.1, for  $n \geq n_0$  sufficiently large and some  $C > 0$  we let  $m$  be such that

$$m \leq C \log n \quad \text{and} \quad \mathbb{P} \left( W_m^\beta \geq n^\alpha \right) \geq n^{-\alpha\zeta}. \quad (5.11)$$

Now we define the event  $A_n$  as follows

$$A_n := \left\{ \forall (x, r) \in [0, n-m] \times [-n^\xi, n^\xi]^d, \quad \theta_{x,r} W_m^\beta < n^{-\alpha} \right\}. \quad (5.12)$$

Our aim is to prove that both  $\mathbb{P}(A_n^c)$  and  $\widehat{\mathbb{P}}_n(A_n)$  tend to zero (cf. (5.2)). Since  $((W_n^\beta)^{1/2})_{n \in \mathbb{N}}$  is uniformly integrable,  $\lim_{n \rightarrow \infty} \mathbb{E}[(W_n^\beta)^{1/2}] = 0$  implies strong disorder, as desired. Using a union bound and translation invariance we have (using Theorem 2.1)

$$\mathbb{P}(A_n^c) \leq n^{d\xi+1} \mathbb{P} \left( W_m^\beta \geq n^\alpha \right) \leq C n^{d\xi+1-\alpha\zeta}. \quad (5.13)$$

With our choice of parameters (5.10), the exponent in the r.h.s. is negative and hence  $\mathbb{P}(A_n^c)$  tends to zero. The remainder of the proof is to show that the same holds for  $\widehat{\mathbb{P}}_n(A_n)$ .

The key idea is to show that, in some sense that, under the size biased measure  $\widehat{\mathbb{P}}_n$ , we can extract  $n^{1-o(1)}$  variables amongst the collection  $(\theta_{r,x} W_m^\beta)_{x \in \mathbb{Z}^d, r \in [0, m-m]}$ , whose distribution is i.i.d. and given by  $\widehat{\mathbb{P}}_m(\widetilde{W}_m \in \cdot)$ . This is where the spine representation from Lemma 5.3 is useful. Let  $\theta_{x,r} \widetilde{W}_m^\beta$  denote the partition function built with environment  $\widetilde{\omega}(X, \omega, \widehat{\omega})$  that is,

$$\theta_{x,r} \widetilde{W}_m^\beta := E' \left[ e^{\sum_{i=1}^m \beta \widetilde{\omega}_{r+m, x+X'_i} - m\lambda(\beta)} \right] \quad (5.14)$$

where  $X'$  is a simple random walk with the same law as  $X$  ( $X$  appears in the definition  $\tilde{\omega}$  so it is not available as a variable of integration). From Lemma 5.3 we have

$$\tilde{\mathbb{P}}_n(A_n) = \hat{\mathbb{P}} \otimes \mathbb{P} \otimes P(\tilde{\omega} \in A_n).$$

Now we observe that if  $\tilde{\omega}$  satisfies  $A_n$  then either  $\theta_{r,x} \tilde{W}_m^\beta$  has to be small when  $(x, r)$  runs along the graph of  $X$ , or the graph of  $X$  must leave the box  $\llbracket 1, n \rrbracket \times \llbracket -n^\xi, n^\xi \rrbracket^d$ , i.e.,

$$\{\tilde{\omega} \in A_n\} \subset \left\{ \forall r \in \llbracket 0, n-m \rrbracket, \theta_{r,X_r} \tilde{W}_m^\beta < n^{-\alpha} \right\} \cup \left\{ \exists i \in \llbracket 1, n \rrbracket, X_i \notin \llbracket -n^\xi, n^\xi \rrbracket^d \right\}. \quad (5.15)$$

Setting  $j_n := \lfloor n/m \rfloor$  and restricting the range of  $r$  to multiples of  $m$  we obtain that

$$\begin{aligned} \hat{\mathbb{P}} \otimes \mathbb{P} \otimes P(\tilde{\omega} \in A_n^c) &\leq P\left(\exists i \in \llbracket 1, n \rrbracket, X_i \notin \llbracket -n^\xi, n^\xi \rrbracket^d\right) \\ &\quad + \hat{\mathbb{P}} \otimes \mathbb{P} \otimes P\left(\forall i \in \llbracket 0, j_n - 1 \rrbracket, \theta_{im, X_{im}} \tilde{W}_m^\beta < n^{-\alpha}\right) \end{aligned} \quad (5.16)$$

Since  $\xi > 1/2$ , the probability of the first term goes to zero and we can conclude by showing that

$$\lim_{n \rightarrow \infty} \hat{\mathbb{P}} \otimes \mathbb{P} \otimes P\left(\forall i \in \llbracket 0, j_n - 1 \rrbracket, \theta_{im, X_{im}} \tilde{W}_m^\beta < n^{-\alpha}\right) = 0. \quad (5.17)$$

By construction (using the Markov property for the random walk  $X$ ) the sequence of environment

$$\left( (\theta_{im, X_{im}} \tilde{\omega}_{n,x})_{(n,x) \in \llbracket 1, m \rrbracket \times \mathbb{Z}^d} \right)_{i \geq 0}$$

is independent and identically distributed. In particular, the sequence of variables  $(\theta_{im, X_{im}} \tilde{W}_m^\beta)_{i \geq 1}$  is i.i.d. and from Lemma 5.3 the marginal distribution is given by

$$P \otimes \mathbb{P} \otimes \hat{\mathbb{P}}\left(\tilde{W}_m^\beta \in \cdot\right) = \tilde{\mathbb{P}}_m\left(W_m^\beta \in \cdot\right). \quad (5.18)$$

Observing that, from (5.11) we have

$$\tilde{\mathbb{P}}_m(W_m \geq n^{-\alpha}) = \mathbb{E}\left[W_m \mathbf{1}_{\{W_m \geq n^{-\alpha}\}}\right] \geq n^{-\alpha(\zeta-1)},$$

we obtain that for any  $j \geq 1$  we have

$$\hat{\mathbb{P}} \otimes \mathbb{P} \otimes P\left(\forall i \in \llbracket 0, j-1 \rrbracket, \theta_{im, X_{im}} \tilde{W}_m^\beta < n^{-\alpha}\right) = \tilde{\mathbb{P}}_m(W_m < n^{-\alpha})^j \leq e^{-jn^{-\alpha(\zeta-1)}}. \quad (5.19)$$

We apply the above formula for  $j_n$  and we can conclude using the fact that  $\alpha(\zeta-1) < 1$  (due to our choice of (5.10)), hence  $\lim_{n \rightarrow \infty} j_n n^{-\alpha(\zeta-1)} = \infty$ .  $\square$

**5.3. Proof of Corollary 2.9.** For part (i), since  $p^*$  is decreasing, it is enough to show that for any  $\varepsilon > 0$  there exists  $\beta' < \beta$  such that  $p^*(\beta') \leq p^*(\beta) + \varepsilon$ , that is

$$\sup_n \mathbb{E}[(W_n^{\beta'})^{p^*(\beta)+\varepsilon}] = \infty. \quad (5.20)$$

For this purpose, let  $u_0$  be large enough that  $u_0^\varepsilon \geq \frac{8}{c_1}$ , where  $c_1$  is such that  $\mathbb{P}(\widehat{W}_n^{\beta,*} \geq u) \geq c_1 u^{-p^*}$  holds for all  $u > 1$ . We then choose  $T_0$  large enough so that

$$\mathbb{P}\left(\sup_{x \in \mathbb{Z}^d, t \leq T_0} \widehat{W}_t^\beta(x) > u_0\right) \geq \frac{1}{2} \mathbb{P}\left(\sup_{x \in \mathbb{Z}^d, t \in \mathbb{N}} \widehat{W}_t^\beta(x) > u_0\right) \geq \frac{c_1}{2} u_0^{-p^*}.$$

The probability on the left-hand side being continuous in  $\beta$ , we can find  $\beta' < \beta$  such that

$$\mathbb{P}\left(\sup_{x \in \mathbb{Z}^d, t \leq T_0} \widehat{W}_t^{\beta'}(x) > u_0\right) \geq \frac{c_1}{4} u_0^{-p^*}. \quad (5.21)$$

Using (5.4), we have

$$\mathbb{P}\left(W_{kT_0}^{\beta'} \geq u_0^k\right) \geq \mathbb{P}\left(\sup_{x \in \mathbb{Z}^d} \widehat{W}_{kT_0}^{\beta'}(x) \geq u_0^k\right) \geq \mathbb{P}\left(\sup_{x \in \mathbb{Z}^d, t \leq T_0} \widehat{W}_t^{\beta'}(x) > u_0\right)^k \quad (5.22)$$

and thus combining (5.21) and (5.22) we obtain

$$\mathbb{E}\left[(W_{kT_0}^{\beta'})^{p^*+\varepsilon}\right] \geq (u_0)^{(p^*+\varepsilon)k} \mathbb{P}\left(W_{kT_0}^{\beta'} \geq u_0^k\right) \geq \left(\frac{c_1}{4} u_0^\varepsilon\right)^k \geq 2^k \quad (5.23)$$

where the last inequality comes from our choice for  $u_0$ . This proves (5.20). For part (ii) we repeat the above argument with  $\beta'$  equal replaced by  $\beta$ , which gives

$$\limsup_{n \rightarrow \infty} \frac{1}{n} \log \mathbb{E}[(W_n^\beta)^{p^*+\varepsilon}] \geq \lim_{k \rightarrow \infty} \frac{1}{kT_0} \log \left[\left(\frac{c_1}{2} u_0^\varepsilon\right)^k\right] \geq \frac{\log 4}{T_0} > 0.$$

□

**5.4. Proof of Corollary 2.12.** We observe that

$$\mathbb{E}\left[(W_n^\beta)^{p^*}\right] = \int_0^\infty p^* u^{p^*-1} \mathbb{P}\left(W_n^\beta \geq u\right) du \leq 1 + \int_1^\infty p^* u^{p^*-1} \mathbb{P}\left(W_n^\beta \geq u\right) du. \quad (5.24)$$

Now Theorem 2.1 implies that, for any  $u \geq 1$  and some constant  $C > 0$ ,

$$u^{p^*-1} \mathbb{P}\left(W_n^\beta \geq u\right) \leq C u^{-1}. \quad (5.25)$$

We are going to show that there exists a constant  $K > 0$  such that for every  $n$  we have

$$\forall u \geq \exp(Kn), \quad u^{p^*-1} \mathbb{P}\left(W_n^\beta \geq u\right) \leq u^{-2}. \quad (5.26)$$

Combining (5.24), (5.25) and (5.26) we obtain that

$$\mathbb{E}\left[(W_n^\beta)^{p^*}\right] \leq 1 + C p^* \int_1^{e^{Kn}} u^{-1} du + p^* \int_{e^{Kn}}^\infty u^{-2} du \leq 1 + p^* + C K p^* n. \quad (5.27)$$

Finally, to prove (5.26), we simply observe that

$$\mathbb{P}\left(W_n^\beta \geq u\right) \leq u^{-p} \mathbb{E}\left[(W_n^\beta)^p\right] \leq u^{-p} e^{n(\lambda(\beta p) - p\lambda(\beta))}. \quad (5.28)$$

and conclude by applying the above to  $p := p^* + 2$ , which implies (5.26) for  $K = \lambda(\beta p) - p\lambda(\beta)$ . □

#### ACKNOWLEDGEMENTS

This work was partially realized as H.L. was visiting the Research Institute for Mathematical Sciences, an International Joint Usage/Research Center located in Kyoto University and he acknowledges kind hospitality and support. This research was supported by JSPS Grant-in-Aid for Scientific Research 23K12984 and 21K03286. H.L. also acknowledges the support of a productivity grand from CNQq and of a CNE grant from FAPERj.

## REFERENCES

- [AKQ14] Tom Alberts, Konstantin Khanin, and Jeremy Quastel. The intermediate disorder regime for directed polymers in dimension  $1 + 1$ . *Ann. Probab.*, 42(3):1212–1256, 2014.
- [Bat21] Erik Bates. Full-path localization of directed polymers. *Electron. J. Probab.*, 26:Paper No. 74, 24, 2021.
- [BC20] Erik Bates and Sourav Chatterjee. The endpoint distribution of directed polymers. *The Annals of Probability*, 48(2):817 – 871, 2020.
- [Bir04] Matthias Birkner. A condition for weak disorder for directed polymers in random environment. *Electron. Comm. Probab.*, 9:22–25, 2004.
- [Bol89] Erwin Bolthausen. A note on the diffusion of directed polymers in a random environment. *Comm. Math. Phys.*, 123(4):529–534, 1989.
- [BS10] Matthias Birkner and Rongfeng Sun. Annealed vs quenched critical points for a random walk pinning model. *Ann. Inst. H. Poincaré Probab. Statist.*, 46(2):414 – 441, 2010.
- [BT10] Quentin Berger and Fabio Toninelli. On the Critical Point of the Random Walk Pinning Model in Dimension  $d=3$ . *Electron. J. Probab.*, 15(none):654 – 683, 2010.
- [CC13] Francis Comets and Michael Cranston. Overlaps and pathwise localization in the Anderson polymer model. *Stochastic Process. Appl.*, 123(6):2446–2471, 2013.
- [CH02] Philippe Carmona and Yueyun Hu. On the partition function of a directed polymer in a Gaussian random environment. *Probab. Theory Related Fields*, 124(3):431–457, 2002.
- [CN21] Clément Cosco and Shuta Nakajima. Gaussian fluctuations for the directed polymer partition function in dimension  $d \geq 3$  and in the whole  $L^2$ -region. *Ann. Inst. Henri Poincaré Probab. Stat.*, 57(2):872–889, 2021.
- [CNN22] Clément Cosco, Shuta Nakajima, and Makoto Nakashima. Law of large numbers and fluctuations in the sub-critical and  $L^2$  regions for SHE and KPZ equation in dimension  $d \geq 3$ . *Stochastic Process. Appl.*, 151:127–173, 2022.
- [CSY03] Francis Comets, Tokuzo Shiga, and Nobuo Yoshida. Directed polymers in a random environment: path localization and strong disorder. *Bernoulli*, 9(4):705–723, 2003.
- [CSZ23] Francesco Caravenna, Rongfeng Sun, and Nikos Zygouras. The critical 2d stochastic heat flow. *Invent. Math.*, 233(1):325–460, 2023.
- [CY06] Francis Comets and Nobuo Yoshida. Directed polymers in random environment are diffusive at weak disorder. *Ann. Probab.*, 34(5):1746–1770, 2006.
- [FJ23] Ryoki Fukushima and Stefan Junk. Moment characterization of the weak disorder phase for directed polymers in a class of unbounded environments. *Electron. Commun. Probab.*, 28:Paper No. 41, 9, 2023.
- [Gia07] Giambattista Giacomin. *Random polymer models*. Imperial College Press, London, 2007.
- [GRZ18] Yu Gu, Lenya Ryzhik, and Ofer Zeitouni. The Edwards-Wilkinson limit of the random heat equation in dimensions three and higher. *Comm. Math. Phys.*, 363(2):351–388, 2018.
- [HH85] David A. Huse and Christopher L. Henley. Pinning and roughening of domain walls in ising systems due to random impurities. *Phys. Rev. Lett.*, 54:2708–2711, Jun 1985.
- [IS88] John Z. Imbrie and Thomas Spencer. Diffusion of directed polymers in a random environment. *J. Statist. Phys.*, 52(3-4):609–626, 1988.
- [JL24] Stefan Junk and Hubert Lacoïn. Strong disorder and very strong disorder are equivalent for directed polymers. *arXiv e-prints*, page arXiv:2402.02562, February 2024.
- [Jun22a] Stefan Junk. Fluctuations of partition functions of directed polymers in weak disorder beyond the  $L^2$ -phase. *arXiv e-prints*, page arXiv:2202.02907, February 2022.
- [Jun22b] Stefan Junk. New Characterization of the Weak Disorder Phase of Directed Polymers in Bounded Random Environments. *Comm. Math. Phys.*, 389(2):1087–1097, 2022.
- [Jun23a] Stefan Junk. Local limit theorem for directed polymers beyond the  $L^2$ -phase. *arXiv e-prints*, page arXiv:2307.05097, July 2023.
- [Jun23b] Stefan Junk. Stability of weak disorder phase for directed polymer with applications to limit theorems. *ALEA Lat. Am. J. Probab. Math. Stat.*, 20:861–883, 2023.
- [LW09] Quansheng Liu and Frédérique Watbled. Exponential inequalities for martingales and asymptotic properties of the free energy of directed polymers in a random environment. *Stochastic Process. Appl.*, 119(10):3101–3132, 2009.
- [LZ22] Dimitris Lygkonis and Nikos Zygouras. Edwards-Wilkinson fluctuations for the directed polymer in the full  $L^2$ -regime for dimensions  $d \geq 3$ . *Ann. Inst. Henri Poincaré Probab. Stat.*, 58(1):65–104, 2022.
- [MSZ16] Chiranjib Mukherjee, Alexander Shamov, and Ofer Zeitouni. Weak and strong disorder for the stochastic heat equation and continuous directed polymers in  $d \geq 3$ . *Electron. Commun. Probab.*, 21:Paper No. 61, 12, 2016.

[Zyg24] Nikos Zygouras. Directed polymers in a random environment: a review of the phase transitions. *arXiv e-prints*, page arXiv:2401.01757, January 2024.

GAKUSHUIN UNIVERSITY, 1-5-1 MEJIRO, TOSHIMA-KU, TOKYO 171-8588 JAPAN

*Email address:* `sjunk@math.gakushuin.ac.jp`

IMPA, ESTRADA DONA CASTORINA 110, RIO DE JANEIRO RJ-22460-320- BRASIL

*Email address:* `lacoim@impa.br`