Evaluation of integrals for the emptiness formation probability in the square-ice model

F. Colomo and A. G. Pronko

ABSTRACT. We study the emptiness formation probability (EFP) in the sixvertex model with domain wall boundary conditions. We present a conjecture according to which at the ice point, i.e., when all the Boltzmann weights are equal, the known multiple integral representation (MIR) for the EFP can be given as a finite-size matrix determinant of Fredholm type. Our conjecture is based on the explicit evaluation of the MIR for particular values of geometric parameters and on two kinds of identities for the boundary correlation function. The obtained representation can be further written as the Fredholm determinant of some linear integral operator. We show that as the geometric parameters of the EFP are tuned to the vicinity of the arctic curve arising in the scaling limit, the conjectured determinant turns into the GUE Tracy–Widom distribution.

Contents

1. Introduction	1
2. MIRs for EFP	3
2.1. The state sum	3
2.2. Boundary one-point function	4
2.3. Two types of identities	6
2.4. EFP	8
3. Evaluation of the integrals	9
3.1. Deformation of the contours	9
3.2. Determinant structures	10
3.3. Particular cases of $s = 1, \ldots, 4$	11
3.4. Case of generic s	15
4. EFP as a Fredholm determinant	16
4.1. Integral form of the matrices	17
4.2. Fredholm determinants	18
4.3. Scaling limit	19
4.4. Tracy–Widom distribution	21
Acknowledgments	24
Appendix A. Asymptotic expansion of $h_{r+1}(z)$ as $r \to \infty$	24
References	25

1. Introduction

A useful tool in study of the six-vertex model with domain wall boundary conditions is the 'emptiness formation probability' (EFP), a non-local correlation function describing the probability of obtaining a region of the lattice with all vertices in the same state. It can be seen as a cumulative distribution function with respect to the geometric parameters describing the size of the 'frozen' region. The name 'EFP' originated in the context of Heisenberg spin chains (equivalent to the six-vertex model on a torus), where a similar correlation function deserved a lot of attention, see, e.g., [1–7].

In [8], we have derived various representations for the EFP of the six-vertex model with domain wall boundary conditions, in terms of determinants, orthogonal polynomials, and multiple contour integrals. The multiple integral representation (MIR) has turned out to be most useful, as it allowed us to derive the arctic curve of the model [9, 10]. The proposed approach found later a simplified formulation with the tangent method [11]. As a further development toward better understanding of the structures arising in non-local correlation functions, an alternative derivation of the known MIR based on certain antisymmetrization relation have been provided [12]. Recently, it was been shown that this antisymmetrization relation can be used to obtain a new, alternative, MIR for the EFP [13]. Furthermore, it appears that the equivalence of the two MIRs implies for the boundary one-point function entering these MIRs the existence of nontrivial relations, which can be viewed as some sort of 'sum rule' identities.

In the present paper, to elaborate further on these results, we focus on a very interesting special case, known as the 'ice point', where all the six-vertex model Boltzmann weights are equal to each other. We find, that besides the sum rule type identities, there exist one more type of relations for the one-point boundary correlation function, that follows from the fact that at the ice point it can be given in terms the Gauss hypergeometric function. Our main result is an explicit, although conjectural expression for the EFP, as a finite-size matrix determinant of the Fredholm type. Our conjecture is based on the direct evaluation of the contour integrals in the first MIR, and the above mentioned two types of identities. Next we investigate the Fredholm determinant of the corresponding integral operator. We show that, as the system size becomes large, and the geometric parameters of the EFP are tuned to the vicinity of the arctic curve, and appropriately scaled, the kernel of this integral operator turns into the so-called Airy kernel.

It is well known that the Fredholm determinant of the Airy kernel determines the celebrated Tracy–Widom distribution describing fluctuations of the largest eigenvalue of the Gaussian unitary ensemble (GUE) [14, 15]. The Tracy–Widom distribution has been observed to arise in a variety of models and it is commonly believed to describe the crossover between weakly- and strongly-coupled phases in various probabilistic problems, see, e.g., [16] and reference therein. In particular, it has been shown to arise in the discrete random matrix model with Hahn polynomials type measure [17] and in the closely related domino tilings of the Aztec diamond [18, 19]. The quantity studied there can be viewed as the EFP of the six-vertex model with domain wall boundary conditions and with Boltzmann weights restricted by the free-fermion condition [20]. Appearance of the Tracy–Widom distribution in the free-fermion six-vertex model context can also be derived by means of a correspondence with non-intersecting paths [21] and from the theory of Painlevé equations [22].

An important open problem actively addressed recently is how far this law remains valid away from the free-fermionic case. Examples where it has been shown to hold are the asymmetric simple exclusion process [23–25] and the six-vertex model with Boltzmann weights restricted by the stochastic condition [26]. As for the general six-vertex model, numerical simulations suggest a positive answer as well [27,28]. A significant analytical achievement is the proof that fluctuations for the maximum of the top path in alternating-sign matrices, or, in other words, in the six-vertex model at its ice point with domain wall boundary condition, are governed by the Tracy–Widom distribution for the Gaussian orthogonal ensemble (GOE) [29] (see Theorem 2.4 therein). As conjectured in that paper, this would suggest the validity of the GUE Tracy–Widom distribution for the EFP (Conjecture 2.5 therein). The present paper is strongly inspired by this observation. Our main result here leads exactly to the same statement, i.e., that the behaviour of the EFP of the ice model with domain wall boundary conditions, in the vicinity of the arctic curve, is described by the GUE Tracy–Widom distribution.

The paper is organized as follows. In the next section we collect all necessary ingredients for calculations from the previous studies and explain the origin of the two types of identities for the boundary one-point function. In section 3, we explain how we compute integrals and use these identities to simplify the expressions, and provide the main result that the EFP is given by a Fredholm-type determinant of a finite-size matrix, see Conjecture 3.1. In section 4, we rewrite the conjectured result in the form of the Fredholm determinant of a linear integral operator, study it in the limit of large system size, and show that when geometric parameters are tuned to the vicinity of the arctic curve, and suitably rescaled, the kernel of the integral operator turns into the Airy kernel, and thus the EFP into the GUE Tracy–Widom distribution.

2. MIRs for EFP

In this section we collect all the necessary input information about the model: configurations of the model, the one-point boundary correlation function, two types of identities, and MIRs for the EFP.

2.1. The state sum. Configurations of the six-vertex model are usually depicted in terms of arrows aligned along edges of a square lattice (or a four-valent graph). Allowed configurations are subject to the 'ice rule': the number of incoming and outgoing arrows at each lattice vertex is equal. Sometimes it is more convenient to use a description of states in terms of occupation number variables, namely, an edge is considered 'empty' if it carries an up or right arrow, and 'occupied' if it carries a left or down arrow. Graphically, it corresponds to drawing a solid lines on the lattice. The ice rule guaranties that the lines flow from the top and right to the down and left. The six typical vertices and their Boltzmann weights in the arrow-reversal symmetric model are shown in Fig. 1.

We consider the six-vertex model with homogeneous (vertex position independent) Boltzmann weights. The three weight functions a, b, and c denoting the Boltzmann weights in the standard notation (see, e.g. [30], for a detailed description) can be parameterized (modulo overall normalization) by two parameters

$$\Delta = \frac{a^2 + b^2 - c^2}{2ab}, \qquad t = \frac{b}{a}.$$

The so-called "ice point" corresponds to all weights equal, a = b = c, that is $\Delta = 1/2$ and t = 1. Our main results will be obtained for this case, though some considerations appear to be valid for arbitrary Δ and t.

Domain wall boundary conditions [31–33] can be imposed for the six-vertex model defined on a square domain of the square lattice, consisting of N horizontal



FIGURE 1. The six vertices of the six-vertex model in terms of lines and their Boltzmann weights.



FIGURE 2. Domain wall boundary conditions (N = 7). Configurations of the model are obtained by filling edges by thin or thick lines.

and N vertical lines (the ' $N \times N$ lattice'). With these conditions, each of the four boundaries has all its edges carrying the same states and these states are opposite to each other on opposite boundaries.

The partition function (the sum over states or 'state sum') is defined, as usual, as the sum over all configurations consistent with the imposed boundary conditions, each configuration being assigned its Boltzmann weight, given by the product of the weights of all vertices. We will consider the partition function, normalized by the factor $a^{N(N-1)}c^N$. We denote this quantity by S_N . The chosen normalization ensures that $S_N = S_N(t, \Delta)$ is a polynomial in t and Δ . The first few values are:

$$S_{1} = 1,$$

$$S_{2} = 1 + t^{2},$$

$$S_{3} = (1 + t^{2})^{3} - 2\Delta t^{3},$$

$$S_{4} = (1 + t^{2})^{6} - 8\Delta (1 + 2t^{2} + 2t^{4} + t^{6}) t^{3} + 4\Delta^{2} (1 + t^{4}) t^{4}.$$

At the ice point, t = 1 and $\Delta = 1/2$, S_N is equal to the number of $N \times N$ alternating-sign matrices, $S_1 = 1$, $S_2 = 2$, $S_3 = 7$, $S_4 = 42$, etc. In general,

$$S_N(1,1/2) = A_N, \qquad A_N = \prod_{j=0}^{N-1} \frac{(3j-1)!}{(2N-j)!}.$$

On the connection with the alternating-sign matrices, see [34–38].

2.2. Boundary one-point function. A simple but important correlation function in the model is the so-called boundary one-point function, originally introduced in [39]. It is usually denoted by $H_N^{(r)}$, and gives the probability that the *r*th vertical edge (e.g., from the right) in the first horizontal row of the $N \times N$ lattice contains an altered (e.g., empty vs occupied) state, see Fig. 3. Note that



FIGURE 3. Boundary one-point correlation function (N = 7, r = 5).

because of the boundary conditions there is exactly one such a state in this row (in general, the nth row from the boundary contains n altered states).

It is customary to deal with the related generating function of the boundary one-point correlation function,

$$h_N(z) = \sum_{r=1}^N H_N^{(r)} z^{r-1}, \qquad h_N(1) = 1.$$
 (2.1)

The function $h_N(z)$ admits an interpretation as the Izergin-Korepin partition function [32, 33] with just one inhomogeneity parameter [8, 40].

To write down a general formula for $h_N(z)$, we introduce a sequence of functions $\phi_j(t), j = 0, 1, 2, \ldots$, defining them in a recursive way as follows [41]:

$$\phi_{j+1}(t) = t\partial_t(t - 2\Delta + t^{-1})\phi_j(t), \qquad \phi_0 \equiv 1.$$

Then,

$$h_N(z) = (N-1)! \left((t-2\Delta+t^{-1})\frac{z}{z-1} \right)^{N-1} \\ \times \frac{\det \left[\begin{cases} \phi_{i+j-2}(t) & j=1,\dots,N-1 \\ \phi_{i-1}(tz) & j=N \end{cases} \right]_{i,j=1,\dots,N}}{\det[\phi_{i+j-2}(t)]_{i,j=1,\dots,N}}.$$

The first few functions $h_N(z)$ are:

$$\begin{split} h_1(z) &= 1, \\ h_2(z) &= \frac{1+t^2 z}{S_2}, \\ h_3(z) &= \frac{S_2 + 2 \left(1+t^2 - \Delta t\right) t^2 z + S_2 t^4 z^2}{S_3}, \\ h_4(z) &= \frac{1}{S_4} \Big\{ S_3 + \Big[3 \left(1+t^2\right)^3 t^2 - 2\Delta \left(3+6t^2+2t^4\right) t^3 + 4\Delta^2 t^4 \Big] z \\ &+ \Big(3 \left(1+t^2\right)^3 t^2 - 2\Delta \left(2+6t^2+3t^4\right) t^3 + 4\Delta^2 t^4 \Big) t^4 z^2 + S_3 t^6 z^3 \Big\}. \end{split}$$

Here, S_i 's are the state sums given above.

In some special cases the function $h_N(z)$ can be written explicitly for generic N. For example, for $\Delta = 0$, known as the free-fermion point of the model, $h_N(z) = [(1 + t^2 z)/(1 + t^2)]^{N-1}$.

At the ice point, t = 1 and $\Delta = 1/2$, it reads [9]

$$h_N(z) = {}_2F_1\left(\begin{array}{c} -N+1, \ N\\ 2N \end{array} \middle| 1-z\right).$$
(2.2)

or

$$h_N(z) = \frac{(N)_{N-1}}{(2N)_{N-1}} {}_2F_1 \left(\begin{array}{c} -N+1, N\\ -2N+2 \end{array} \middle| z \right).$$
(2.3)

It can also be written explicitly at the 'dual ice point', t = 1 and $\Delta = -1/2$, though the expressions are rather bulky, see [42] for details.

The coefficients $H_N^{(r)}$ in (2.1) in the ice-point case $(t = 1, \Delta = 1/2)$ can also be represented as

$$H_N^{(r)} = \frac{A_{N,r}}{A_N}.$$

Here, A_N is number of alternating-sign matrices of size $N \times N$, and $A_{N,r}$ is their refined enumeration, i.e., the number of such matrices with the sole '1' entry in the first row being at the *r*th position [37]. In a more explicit form,

$$H_N^{(r)} = \binom{N+r-2}{N-1} \binom{2N-1-r}{N-1} / \binom{3N-2}{N-1},$$

that can be seen as another form of writing (2.2) or (2.3).

2.3. Two types of identities. To shorten the formulas a bit, we will skip the argument of functions whenever it is 0, e.g., writing h_N for $h_N(0)$, h'_N for $h'_N(0)$, etc.

In the first type of identities the derivatives of $h_N(z)$ at the point z = 1 are expressed in terms of the derivatives of $h_N(z)$ at the point z = 0. The first three relations read:

$$\begin{aligned} h'_{N-1}(1) &= \frac{1}{1 - 2\Delta t + t^2} \left\{ \frac{h'_N}{h_N} - t^2 \right\}, \\ h''_{N-2}(1) &= \frac{1}{(1 - 2\Delta t + t^2)^2} \left\{ -\frac{h''_N}{h_N} + 2\frac{h'_{N-1}h'_N}{h_{N-1}h_N} - 2\left(1 - 2\Delta t + 2t^2\right)\frac{h'_{N-1}}{h_{N-1}} \right. \\ &\quad + 2\frac{h'_N}{h_N} - 2t^2 + 2t^4 \right\}, \\ h'''_{N-3}(1) &= \frac{1}{(1 - 2\Delta t + t^2)^2} \left\{ \frac{h'''_N}{h_N} - 3\frac{h'_{N-2}h'_N}{h_{N-2}h_N} - 3\frac{h''_{N-1}h'_N}{h_{N-1}h_N} \right. \\ &\quad + 3\left(2 + 3t^2 - 4t\Delta\right)\frac{h''_{N-1}}{h_{N-1}} - 6\frac{h''_N}{h_N} + 6\frac{h'_{N-2}h'_{N-1}h'_N}{h_{N-2}h_{N-1}h_N} \\ &\quad - 6\left(2 + 3t^2 - 4t\Delta\right)\frac{h'_{N-2}h'_{N-1}}{h_{N-2}h_{N-1}} + 6\frac{h'_{N-2}h'_N}{h_{N-2}h_N} + 6\frac{h'_{N-1}h'_N}{h_{N-1}h_N} \\ &\quad + 6\left(1 + 2t^2 + 3t^4 - 4t\Delta - 6t^3\Delta + 4t^2\Delta^2\right)\frac{h'_{N-2}}{h_{N-2}} \\ &\quad - 6\left(2 + 3t^2 - 4t\Delta\right)\frac{h'_{N-1}}{h_{N-1}} + 6\frac{h'_N}{h_N} + 18t^4 - 6t^6 - 12t^3\Delta \right\}. \end{aligned}$$

In view of (2.1), these identities express sums over the set of functions $H_N^{(r)}$, $r = 1, \ldots, N$, in terms of first few of them, i.e., they are sum rule identities. As explained in the next section, these identities follows from the existence of two different MIRs for the EFP.

Another sort of identities involves only higher derivatives at z = 0. It has to be stressed, however, that we are able at the moment to establish them only for the special case of the ice point, t = 1 and $\Delta = 1/2$. In this case one can use the explicit expression (2.3).

For example, from (2.3) one can easily find for the first derivative of $h_N(z; 1, 1/2)$ at z = 0:

$$\frac{h'_N}{h_N} = \frac{N}{2}.$$

Hence, we have

$$\frac{h'_N}{h_N} - \frac{h'_{N-1}}{h_{N-1}} - \frac{1}{2} = 0.$$
(2.5)

Clearly, relation (2.5) shows that the identities in (2.4) at the ice point can be somewhat simplified, e.g., by expressing h'_{N-1}/h_{N-1} , h'_{N-2}/h_{N-2} , etc, in terms of h'_N/h_N .

For the second derivative we get

$$\frac{h_N''}{h_N} = \frac{(N-2)N(N+1)}{2(2N-3)}.$$

Taking into account that

$$\frac{h_{N-1}}{h_N} = \frac{3(3N-2)(3N-4)}{4(2N-1)(2N-3)}$$

one can find the following identity:

$$\frac{h_N''}{h_N} - \frac{h_{N-1}''}{h_{N-1}} - \frac{h_N'}{h_N} - 2\frac{h_{N-2}}{h_{N-1}} + \frac{7}{2} = 0.$$
(2.6)

Note that the coefficients are all independent of N.

For the third derivative we have

$$\frac{h_N''}{h_N} = \frac{(N-3)N(N+1)(N+2)}{4(2N-3)}.$$

In the similar manner, we get

$$\frac{h_{N}^{\prime\prime\prime}}{h_{N}} - \frac{h_{N-1}^{\prime\prime\prime}}{h_{N-1}} - \frac{3}{2} \left(\frac{h_{N}^{\prime}}{h_{N}}\right)^{2} - \frac{21}{2} \left(\frac{h_{N-2}}{h_{N-1}} - \frac{7}{4}\right) = 0.$$

Clearly, this game can be continued.

Note that the first type of identities make it possible to express $h_{N-s}^{(s)}(1)$ in terms of the quantities $h_{N-k}^{(j)}$, with j = 0, ..., s and k = 0, ..., s - 1, such that $0 \leq j + k \leq s$. In turn, the second type of identities make it possible to express these quantities in terms of $h_{N-s+1}, ..., h_N, h'_N, ..., h_N^{(s)}$, thus reducing the number of formally independent objects from s(s+3)/2 to 2s.



FIGURE 4. Emptiness formation probability (N = 7, r = 4, s = 2).

2.4. EFP. The EFP, denoted as $F_N^{(r,s)}$, can be defined as the probability of obtaining a domain of size $s \times (N - r)$ attached to a corner of the $N \times N$ lattice, with all its horizontal and vertical edges having the same states as those at the boundary. In other words, all vertices belonging to this $s \times (N - r)$ domain are all *a*- or *b*-weight vertices; this domain can be regarded as an 'empty' region. The vertices belonging to this domain can be removed, giving rise to so-called L-shaped domain. The partition function of the model on this domain is exactly the EFP, modulo the factor $a^{s(N-r)}$ or $b^{s(N-r)}$ times the partition function of the model on the original $N \times N$ lattice.

A simple property of the EFP is that it vanishes identically whenever s > r. This corresponds to a situation where the frozen rectangular domain entering the definition of the EFP extends beyond the diagonal of the $N \times N$ lattice. In terms of the model in the L-shaped domain, it is clear that whenever s > r, the number of allowed configurations vanishes.

Here, for definiteness we stick at the conventions used in Refs. [8, 10, 13], and define EFP such that the frozen domain is located at the top left corner of the $N \times N$ lattice, with all vertices of this domain to be *a*-weight vertices, see Fig. 4. In [8], the following MIR have been derived:

$$F_N^{(r,s)} = (-1)^s \oint_{C_0} \cdots \oint_{C_0} \prod_{j=1}^s \frac{[(t^2 - 2\Delta t)z_j + 1]^{s-j}}{z_j^r (z_j - 1)^{s-j+1}} \\ \times \prod_{1 \le j < k \le s} \frac{z_j - z_k}{t^2 z_j z_k - 2\Delta t z_j + 1} h_{N,s}(z_1, \dots, z_s) \frac{\mathrm{d}^s z}{(2\pi \mathrm{i})^s}.$$
(2.7)

Here, C_0 denotes a small simple anticlockwise oriented contour around the point z = 0. The functions $h_{N,s}(z_1, \ldots, z_s)$, $s = 1, \ldots, N$, are symmetric polynomials of the degree N - 1 in each of their variables and defined in terms of s single-variable functions $h_{N-s+1}(z), \ldots, h_N(z)$ by the formula

$$h_{N,s}(z_1, \dots, z_s) = \frac{\det\left[(z_j - 1)^{s-i} z_j^{i-1} h_{N-i+1}(z_j)\right]_{i,j=1,\dots,s}}{\prod_{1 \le i < j \le s} (z_i - z_j)}.$$
 (2.8)

If, say, $z_s = 1$, then

$$h_{N,s}(z_1,\ldots,z_{s-1},1) = h_{N,s-1}(z_1,\ldots,z_{s-1}).$$

If $z_s = 0$, then

$$h_{N,s}(z_1,\ldots,z_{s-1},0) = h_N(0)h_{N-1,s-1}(z_1,\ldots,z_{s-1})$$

i.e., the degree in all the remaining variables decreases by one.

In [13], it was proven that for generic values of t and Δ , together with MIR (2.7), EFP admits also the following MIR:

$$F_N^{(r,s)} = \frac{h_N(0)\cdots h_{N-s+1}(0)t^{(r-s)(r-s-1)}}{(r-s)!h_1(0)\cdots h_r(0)} \oint_{C_1} \cdots \oint_{C_1} \prod_{j=1}^{r-s} \frac{1}{z_j - 1} \\ \times \prod_{\substack{j,k=1\\j\neq k}}^{r-s} \frac{z_j - z_k}{t^2 z_j z_k - 2\Delta t z_j + 1} h_{N-s,r-s} (z_1, \dots, z_{r-s}) \\ \times h_{r,r-s} \left(\frac{t^2 z_1 - 2\Delta t + 1}{t^2 (z_1 - 1)}, \dots, \frac{t^2 z_{r-s} - 2\Delta t + 1}{t^2 (z_{r-s} - 1)}\right) \frac{\mathrm{d}^{r-s} z}{(2\pi \mathrm{i})^{r-s}}, \quad (2.9)$$

where C_1 is a (small) contour encircling the point z = 1. In comparison with (2.7), the number of integrations in (2.9) is r - s, that is the lattice distance of the point (r, s) from the antidiagonal, rather than s, the lattice distance from the top boundary.

As far as MIRs (2.7) and (2.9) describe the same quantity and both are valid for all allowed values of the geometric parameters, they imply some relations for the function $h_N(z)$. These relations are exactly the first type identities, (2.4). Indeed, setting r = s + 1, one gets for the second MIR just single integration around the pole at the point z = 1, while the first MIR contains *s* integrations around the poles at the points $z_1, \ldots, z_s = 0$. Considering the cases $s = 1, 2, 3, \ldots$ one arrives at the identities listed in (2.4).

3. Evaluation of the integrals

In this section we address the problem of evaluation of the MIR (2.7) for s = 1, 2, 3, 4 and assuming that r = N-s, in the case of the ice point $(t = 1, \Delta = 1/2)$. In principle, this calculation can be done for generic values of the Boltzmann weights, but at the present moment we able to formulate a conjecture only for this special case.

3.1. Deformation of the contours. We start with making a change of the variables $z_j \mapsto z_j^{-1}$, $j = 1, \ldots, s$ in (2.7), that gives

$$F_N^{(r,s)} = \oint_{C_\infty} \cdots \oint_{C_\infty} J_N^{(r,s)}(z_1,\ldots,z_s) \,\mathrm{d}^s z,$$

where C_{∞} denotes a very large contour, anticlockwise oriented, and

$$J_N^{(r,s)}(z_1,\ldots,z_s) = (2\pi i)^{-s} \prod_{j=1}^s \frac{[t^2 - 2\Delta t + z_j]^{s-j}}{z_j^{N-r}(z_j-1)^{s-j+1}} \prod_{1 \le j < k \le s} \frac{z_j - z_k}{t^2 - 2\Delta t z_k + z_j z_k} \times \tilde{h}_{N,s}(z_1,\ldots,z_s).$$
(3.1)

Here,

$$\tilde{h}_{N,s}(z_1,\ldots,z_s) \equiv (z_1\cdots z_s)^{N-1}h_{N,s}(z_1^{-1},\ldots,z_s^{-1}).$$

These functions can also be written as determinants, similarly to (2.8),

$$\tilde{h}_{N,s}(z_1,\ldots,z_s) = \frac{\det\left[(z_j-1)^{s-i}z_j^{i-1}h_{N-i+1}(z_j)\right]_{i,j=1,\ldots,s}}{\prod_{1 \le i < j \le s}(z_i-z_j)}.$$

The function $\tilde{h}_N(z)$ is defined by

$$\tilde{h}_N(z) \equiv z^{N-1} h_N(z^{-1})$$

and it can also be related to $h_N(z)$ by mapping $t \mapsto t^{-1}$,

$$h_N(z;t) = h_N(z;t^{-1}).$$

Hence, in the symmetric case t = 1 the tildes over the functions can be lifted.

We continue transforming the expression by deforming the integration contours, shrinking them down to encircle the sole poles at z = 0 and z = 1. An important property of the integrand (3.1) is that in the deformation of the contours all terms deriving from the evaluation of the residues at mutual poles (those due to the double product in the denominator) vanish [13]. As a result, we have for the EFP

$$F_N^{(r,s)} = \oint_{C_1 \cup C_0} \cdots \oint_{C_1 \cup C_0} J_N^{(r,s)}(z_1, \dots, z_s) \,\mathrm{d}^s z.$$
(3.2)

This formula implies that the EFP is essentially the sum of s + 1 terms:

$$F_N^{(r,s)} = \sum_{k=0}^s I_k,$$

where I_k is the sum of integrals with k variables integrated over C_0 and with the remaining s - k variables integrated over C_1 . The following two simple lemmas show that $I_0 = 1$ and establish the value of I_s in the case of our interest.

LEMMA 3.1. For arbitrary values of all parameters, a cumulative residue at z = 1 in (3.1) is equal to one,

$$\operatorname{res}_{z_1=1} \dots \operatorname{res}_{z_s=1} J_N^{(r,s)}(z_1, \dots, z_s) = 1.$$

PROOF. To prove this statement, evaluate the residues in the shown order. At each step the pole in the corresponding variable at the point z = 1 is simple, and the result follows due to $h_{N,s}(1, \ldots, 1) = 1$.

LEMMA 3.2. At the ice point, $\Delta = 1/2, t = 1$, and for N - r = s the cumulative residue in (3.1) at the point z = 0 equals:

$$\operatorname{res}_{z_1=0} \dots \operatorname{res}_{z_s=0} J_N^{(N-s,s)}(z_1,\dots,z_s) = (-1)^s h_N \cdots h_{N-s+1},$$

where $h_N \equiv h_N(0)$, etc.

PROOF. The proof is similar to that of the previous lemma.

3.2. Determinant structures. Our strategy in computing the EFP is based on the observation (see also [24, 43]) that there exists an $s \times s$ matrix A such that

$$\sum_{k=0}^{s} \lambda^{k} I_{k} = \det_{s} (I - \lambda A), \qquad (3.3)$$

where I is the $s \times s$ identity matrix and λ is a formal parameter. In the examples below matrix A is such that by eliminating its last row and column, the reduction

 $s \mapsto s-1$, $N \mapsto N-1$ is made. As we shall see, this reduction is a key property of the expressions for I_k 's computed from (3.2). We have also found that the matrix A can be given explicitly in a factorized form

$$A = D L U, \tag{3.4}$$

where D, L, and U is a diagonal, lower-triangle, and upper-triangle matrix, respectively.

Representation (3.3) is not unique and one can also to try to find a similar formula

$$\sum_{k=0}^{s} \lambda^k I_k = \det_s (I - \lambda V).$$

where the matrix V has entries independent of s. We find such a matrix and it is in fact related to A. Matrix V can be given as a product of lower-triangle, diagonal, and upper-triangle matrices:

$$V = \widetilde{L} \, \widetilde{D} \, \widetilde{U}. \tag{3.5}$$

They appear to be related to L, D, and U by

$$\widetilde{L} = \Omega L^{\mathsf{T}} \Omega, \qquad \widetilde{D} = \Omega D \Omega, \qquad \widetilde{U} = \Omega U^{\mathsf{T}} \Omega,$$
(3.6)

for s even, and

$$\widetilde{L} = \Omega U \Omega, \qquad \widetilde{D} = \Omega D \Omega, \qquad \widetilde{U} = \Omega L \Omega,$$
(3.7)

for s odd. Here T denotes matrix transposition, and Ω is the $s \times s$ 'unit anti-diagonal' matrix,

$$\Omega_{ij} = \delta_{i+j,s+1}, \qquad \Omega^{\mathsf{T}} = \Omega, \qquad \Omega^2 = I.$$

The sum-rule relations (2.4) are essentially used in obtaining expressions for quantities I_k in terms of the values of derivatives of the function $h_N(z)$ at z = 0. To find the matrix A, starting from the s = 3 case, we need to use also relations between these values. These are the second type of identities; by reducing the number of formally independent quantities from s(s + 1)/2 to 2s - 1, they play a crucial role in the construction of our main conjecture about entries of the matrices A and V.

To proceed with particular cases, we need some more notation. To shorten formulas below, we denote

$$b_i \equiv h_{N-i}, \qquad i = 0, 1, 2, \dots, s-1,$$

and (we limit ourselves below in writing explicitly terms with three derivatives)

$$\kappa'_{i} = \frac{h'_{N-i}}{h_{N-i}}, \quad \kappa''_{i} = \frac{h''_{N-i}}{h_{N-i}}, \quad \kappa'''_{i} = \frac{h''_{N-i}}{h_{N-i}}.$$

We recall that $h_N \equiv h_N(0), h'_N \equiv h'_N(0)$, etc.

3.3. Particular cases of s = 1, ..., 4. We now turn to evaluation of the integrals in the case of s = 1, ..., 4. These expressions appear to be very instructive in our conjecture of the result for a generic s, considered next.

3.3.1. Case s = 1. We start with the simplest but useful case of s = 1. Evaluating the integral for r = N - 1 and using that $\tilde{h}_N(1) = 1$, we get for the EFP (for generic Δ and t) the expression

$$F_N^{(N-1,1)} = 1 - \tilde{h}_N$$

At t = 1 we have $\tilde{h}_N = h_N \equiv b_0$ and so

$$I_0 = 1, \qquad I_1 = -b_0.$$
 (3.8)

Hence,

$$A = V = b_0.$$
 (3.9)

3.3.2. Case s = 2. This is very instructive example and so we will be able to explain main ideas of further calculations. Evaluation of the integrals in (3.2) in this case, using that $\tilde{h}_N(1) = 1$, yields

$$\begin{split} I_0 &= 1, \\ I_1 &= -\frac{t^4 - 2\Delta t^3 + 4\Delta^2 t^2 - 1}{t^4} \,\tilde{h}_N - \frac{2\Delta}{t} \,\tilde{h}'_N - \frac{t^2 - 2\Delta t + 1}{t^2} \,\tilde{h}_{N-1} \\ &- \left(\frac{2\Delta t - 1}{t^2} \,\tilde{h}_N + \tilde{h}'_N\right) \frac{t^2 - 2\Delta t + 1}{t^2} \,\tilde{h}'_{N-1}(1), \\ I_2 &= \frac{t^2 - 4\Delta t + 4\Delta^2 + 1}{t^2} \tilde{h}_N \tilde{h}_{N-1}. \end{split}$$

Now, setting t = 1 and $\Delta = 1/2$ (note that we must keep generic t or Δ , or both, before evaluation of integrals, in order to avoid erroneous contributions from the term $t^2 - 2\Delta tz_2 + z_1z_2$ when computing I_1), and using the first identity from (2.4), we get

$$I_0 = 1, \qquad I_1 = -b_1 - b_0 (\kappa'_0)^2, \qquad I_2 = b_0 b_1.$$
 (3.10)

Note that if we put here $b_0 = 0$, then we get the s = 1 result (3.8) in which b_0 is replaced by b_1 , or $N \mapsto N - 1$. This means that if an 2×2 matrix A exists such that (3.3) holds, then it would be desirable that its top-left entry is b_1 . In turn, this choice fixes the bottom-right entry from the relation tr $A = -I_1$ to be $b_0(\kappa'_0)^2$. The off-diagonal entries must satisfy det $A = I_2$, that modulo diagonal similarity transformation (and up to matrix transposition) leads us to

$$A = \begin{pmatrix} b_1 & b_1(\kappa'_0 - 1) \\ b_0(\kappa'_0 + 1) & b_0(\kappa'_0)^2 \end{pmatrix}.$$

Clearly, this matrix possesses the DLU-factorization (3.4), where

$$D = \begin{pmatrix} b_1 & 0 \\ 0 & b_0 \end{pmatrix}, \quad L = \begin{pmatrix} 1 & 0 \\ \kappa'_0 + 1 & 1 \end{pmatrix}, \quad U = \begin{pmatrix} 1 & \kappa'_0 - 1 \\ 0 & 1 \end{pmatrix}.$$

Another way to write the result of integration in a determinant form is with a matrix V such that s determines only its size, but not entries. For such a matrix then, first from (3.9) follows $V_{11} = b_0$, next from tr $V = -I_1$ follows $V_{22} = b_0 \left((\kappa'_0)^2 - 1 \right) + b_1$, and finally from det $V = I_2$ follows $V_{12}V_{21} = b_0^2 \left((\kappa'_0)^2 - 1 \right)$.

Modulo diagonal transformation and matrix transposition,

$$V = \begin{pmatrix} b_0 & b_0(\kappa'_0 - 1) \\ b_0(\kappa'_0 + 1) & b_0((\kappa'_0)^2 - 1) + b_1 \end{pmatrix}.$$

This matrix admits LDU-factorization (3.5), where

$$\widetilde{L} = \begin{pmatrix} 1 & 0 \\ \kappa'_0 + 1 & 1 \end{pmatrix}, \quad \widetilde{D} = \begin{pmatrix} b_0 & 0 \\ 0 & b_1 \end{pmatrix}, \quad \widetilde{U} = \begin{pmatrix} 1 & \kappa'_0 - 1 \\ 0 & 1 \end{pmatrix}.$$

These matrices are related to L, D, and U by (3.6).

3.3.3. Case s = 3. In this case, after evaluating the integrals in MIR (3.2), using $\tilde{h}_N(1) = 1$, applying the first two identities from (2.4), and putting t = 1, $\Delta = 1/2$, we get

$$I_{0} = 1,$$

$$I_{1} = -b_{2} - b_{1} (\kappa_{1}')^{2} - b_{0} \left[\left(\frac{\kappa_{0}''}{2} - \kappa_{0}' \right)^{2} + 2\kappa_{0}' - 1 \right],$$

$$I_{2} = b_{1}b_{2} + b_{0}b_{2} (\kappa_{0}')^{2} + b_{0}b_{1} \left[1 - \kappa_{0}'(1 + \kappa_{1}') + \frac{\kappa_{0}''}{2} \right]^{2},$$

$$I_{3} = -b_{0}b_{1}b_{2}.$$
(3.11)

If we put $b_0 = 0$ in (3.11) then we get (3.10) in which $b_0, b_1, \kappa'_0 \mapsto b_1, b_2, \kappa'_1$, respectively, that is, $N \mapsto N - 1$.

Thus, we can construct A by choosing its top-left 2×2 block as the matrix A from the case s = 2, with $b_0, b_1, \kappa'_0 \mapsto b_1, b_2, \kappa'_1$, that also fixes A_{33} entry from tr $A = -I_1$. We write A = DLU, where $D = \text{diag}(b_2, b_1, b_0)$, and for the matrices L and U we take

$$L = \begin{pmatrix} 1 & 0 & 0 \\ \kappa'_1 + 1 & 1 & 0 \\ \frac{1}{2}\kappa''_0 - 2\kappa'_0 + 1 & \kappa'_0 - 1 & 1 \end{pmatrix}, \qquad U = \begin{pmatrix} 1 & \kappa'_1 - 1 & \frac{1}{2}\kappa''_0 - 1 \\ 0 & 1 & \kappa'_0 + 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

Clearly, it can easily seen that this choice reproduces values I_0 , I_1 and I_3 , but not I_2 , with the difference of some complicated factor times $\kappa'_1 - \kappa'_0 + \frac{1}{2}$. But it vanishes due to identity (2.5)!

The similar construction in the case of the matrix V, whose entries are required to be independent of s, leads us to (3.5), where $\tilde{D} = \text{diag}(b_0, b_1, b_2)$ and

$$\widetilde{L} = \begin{pmatrix} 1 & 0 & 0 \\ \kappa'_0 + 1 & 1 & 0 \\ \frac{1}{2}\kappa''_0 - 1 & \kappa'_1 - 1 & 1 \end{pmatrix}, \qquad \widetilde{U} = \begin{pmatrix} 1 & \kappa'_0 - 1 & \frac{1}{2}\kappa''_0 - 2\kappa'_0 + 1 \\ 0 & 1 & \kappa'_1 + 1 \\ 0 & 0 & 1 \end{pmatrix}.$$

These matrices are related to D, L, U by (3.7).

3.3.4. Case s = 4. In this case, we obtain

$$\begin{split} I_{0} &= 1, \\ I_{1} &= -b_{3} - b_{2} \left(\kappa_{2}'\right)^{2} + b_{1} \left[1 - 2\kappa_{1}' - \left(\kappa_{1}' - \frac{\kappa_{1}''}{2}\right)^{2} \right] \\ &+ b_{0} \left[2 - 6\kappa_{0}' + 3 \left(\kappa_{0}'\right)^{2} + (1 - \kappa_{0}') \kappa_{0}'' - \left(\kappa_{0}' - \kappa_{0}'' + \frac{\kappa_{0}''}{6}\right)^{2} \right], \\ I_{2} &= b_{2}b_{3} + b_{1}b_{3} \left(\kappa_{1}'\right)^{2} + b_{0}b_{3} \left[-1 + 2\kappa_{0}' + \left(\kappa_{0}' - \frac{\kappa_{0}''}{2}\right)^{2} \right] \\ &+ b_{1}b_{2} \left[1 - \kappa_{1}' \left(1 + \kappa_{2}'\right) + \frac{\kappa_{1}''}{2} \right]^{2} \\ &+ b_{0}b_{2} \left\{ - \left(5 + \kappa_{2}'\right) \left(1 + \kappa_{2}'\right) + 2\kappa_{0}' \left(1 + \kappa_{2}'\right)^{2} \\ &+ \left[1 + \left(2 + \kappa_{2}'\right) \left(-\kappa_{0}' + \frac{\kappa_{0}''}{2} \right) - \frac{\kappa_{0}'''}{6} \right]^{2} \right\} \\ &+ b_{0}b_{1} \left\{ 4 - 2\kappa_{0}' + \frac{\kappa_{0}''}{3} + \left(1 - \kappa_{0}' + \frac{\kappa_{0}''}{2} \right)^{2} \\ &+ 2\kappa_{1}' \left[2 + \left(2 - \kappa_{0}' + \frac{\kappa_{0}''}{2} \right) \left(1 - \kappa_{0}' + \frac{\kappa_{0}''}{2} - \frac{\kappa_{0}'''}{6} \right) \right] \\ &+ \kappa_{1}'' \left[\left(\kappa_{0}' - \frac{\kappa_{0}''}{2} \right)^{2} - 3\kappa_{0}' + \frac{3\kappa_{0}''}{2} - \frac{\kappa_{0}'''}{6} \right)^{2} \\ &+ \left(\kappa_{1}')^{2} \left(1 - \kappa_{0}' + \frac{\kappa_{0}''}{2} - \frac{\kappa_{0}'''}{6} \right)^{2} \right] \\ &- \kappa_{1}'\kappa_{1}'' \left[2 + \left(\kappa_{0}' - \frac{\kappa_{0}''}{2} \right) \left(1 - \kappa_{0}' + \frac{\kappa_{0}''}{2} - \frac{\kappa_{0}'''}{6} \right) \right] \right\}, \\ I_{3} &= -b_{1}b_{2}b_{3} - b_{0}b_{2}b_{3} \left(\kappa_{0}' \right)^{2} - b_{0}b_{1}b_{2} \left\{ 1 + \kappa_{0}'' + \kappa_{2}' \left(1 + \frac{\kappa_{0}''}{2} \right) - \kappa_{0}' \left[\left(1 + \kappa_{1}' \right) \left(1 + \kappa_{2}' \right) - \frac{\kappa_{1}'''}{6} \right]^{2} \right\}, \end{aligned}$$

Repeating the procedure from the s = 2 and s = 3 cases, we write matrix A in the form (3.4), with

,

$$D = \operatorname{diag}(b_3, b_2, b_1, b_0),$$

$$\begin{split} L &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ \kappa_2' + 1 & 1 & 0 & 0 \\ \frac{1}{2}\kappa_1'' - 2\kappa_1' + 1 & \kappa_1' - 1 & 1 & 0 \\ \frac{1}{6}\kappa_0''' - \frac{1}{2}\kappa_0'' - \kappa_0' + 1 & \frac{1}{2}\kappa_0'' - 1 & \kappa_0' + 1 & 1 \end{pmatrix}, \\ U &= \begin{pmatrix} 1 & \kappa_2' - 1 & \frac{1}{2}\kappa_1'' - 1 & \frac{1}{6}\kappa_0''' - \frac{3}{2}\kappa_0'' + 3\kappa_0' - 1 \\ 0 & 1 & \kappa_1' + 1 & \frac{1}{2}\kappa_0'' - 2\kappa_0' + 1 \\ 0 & 0 & 1 & \kappa_0' - 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \end{split}$$

This choice indeed works, but one have to use the identity (2.5) to express now κ'_1 and κ'_2 , for the latter twice, to get them both expressed in terms of κ'_0 . Furthermore, the identity (2.6) is also needed, to express κ''_1 in terms of κ''_0 , b_2 and b_1 , which are already involved.

As for the matrix V we get (3.5), where

$$\begin{split} \widetilde{L} &= \begin{pmatrix} 1 & 0 & 0 & 0 \\ \kappa_0' + 1 & 1 & 0 & 0 \\ \frac{1}{2}\kappa_0'' - 1 & \kappa_1' - 1 & 1 & 0 \\ \frac{1}{6}\kappa_0''' - \frac{1}{2}\kappa_0'' - \kappa_0' + 1 & \frac{1}{2}\kappa_1'' - 2\kappa_1' + 1 & \kappa_2' + 1 & 1 \end{pmatrix}, \\ \widetilde{D} &= \operatorname{diag}(b_0, b_1, b_2, b_3), \\ \widetilde{U} &= \begin{pmatrix} 1 & \kappa_0' - 1 & \frac{1}{2}\kappa_0'' - 2\kappa_0' + 1 & \frac{1}{6}\kappa_0''' - \frac{3}{2}\kappa_0'' + 3\kappa_0' - 1 \\ 0 & 1 & \kappa_1' + 1 & \frac{1}{2}\kappa_1'' - 1 \\ 0 & 0 & 1 & \kappa_2' - 1 \\ 0 & 0 & 0 & 1 \end{pmatrix}. \end{split}$$

These matrices are related to L, D, and U by (3.6).

3.4. Case of generic s. To formulate a conjecture about entries of the matrices L and U (while this is not difficult for D), we recall a formula for generalized Laguerre polynomials (see, e.g., [44])

$$L_{n}^{(\alpha)}(x) = \sum_{k=0}^{n} \binom{n+\alpha}{n-k} \frac{(-x)^{k}}{k!},$$
(3.12)

The special case $L_n(x) \equiv L_n^{(0)}(x)$ is so familiar

$$L_0(x) = 1,$$

$$L_1(x) = -x + 1,$$

$$L_2(x) = \frac{x^2}{2} - 2x + 1,$$

$$L_3(x) = -\frac{x^3}{6} + \frac{3x^2}{2} - 3x + 1,$$

that it is easy to recognize them in the basement of half of entries. The second sequence appearing there, x + 1, $\frac{1}{2}x^2 - 1$, $\frac{1}{6}x^3 - \frac{x^2}{2} - x + 1$, seem at the first glance not to fall into the Laguerre case. However, a simple observation shows that this also the case:

$$L_1(x) - 2L_0(x) = -x - 1,$$

$$L_2(x) - 2L_1(x) = \frac{x^2}{2} - 1,$$

$$L_3(x) - 2L_2(x) = -\frac{x^3}{6} + \frac{x^2}{2} + x - 1.$$

To proceed, we recall that the generalized Laguerre polynomials are subject, among many others, to the relation

$$L_n^{(\alpha-1)}(x) = L_n^{(\alpha)}(x) - L_{n-1}^{(\alpha)}(x).$$

We thus meet in our case two linear combinations of the polynomials $L_n^{(-1)}(x)$ and $L_{n-1}^{(0)}(x)$, namely, they are $L_n^{(-1)}(x) \pm L_{n-1}^{(0)}(x)$. Our main result here is the following.

(0) 0

CONJECTURE 3.1. For t = 1 and $\Delta = 1/2$, and for r = N - s, the EFP can be given as the determinant of the $s \times s$ matrix I - A, where A = DLU and

$$D_{ij} = h_{r+i}(0) \,\delta_{ij},$$

$$L_{ij} = \frac{(-1)^{i-j}}{h_{r+i}(0)} \left[L_{i-j}^{(-1)}(\partial_z) + (-1)^{i-1} L_{i-j-1}^{(0)}(\partial_z) \right] h_{r+i}(z) \Big|_{z=0},$$

$$U_{ij} = \frac{(-1)^{i-j}}{h_{r+j}(0)} \left[L_{j-i}^{(-1)}(\partial_z) + (-1)^j L_{j-i-1}^{(0)}(\partial_z) \right] h_{r+j}(z) \Big|_{z=0}.$$
(3.13)

Equivalently, the EFP is given by the determinant of the $s \times s$ matrix I - V, where V = LDU, and

$$\widetilde{L}_{ij} = \frac{(-1)^{i-j}}{h_{N-i+1}(0)} \left[L_{i-j}^{(-1)}(\partial_z) + (-1)^j L_{i-j-1}^{(0)}(\partial_z) \right] h_{N-i+1}(z) \Big|_{z=0},
\widetilde{D}_{ij} = h_{N-i+1}(0) \,\delta_{ij},
\widetilde{U}_{ij} = \frac{(-1)^{i-j}}{h_{N-j+1}(0)} \left[L_{j-i}^{(-1)}(\partial_z) + (-1)^{i-1} L_{j-i-1}^{(0)}(\partial_z) \right] h_{N-j+1}(z) \Big|_{z=0}.$$
(3.14)

Function $h_N(z)$ is the Gauss hypergeometric function given in (2.2) or (2.3).

Note that the entries of the matrix A are independent of s, if one takes r = N - sas an independent geometric parameter.

To verify our conjecture, we have also performed exact evaluation (with the help of symbolic manipulation software) of the MIR (2.7) in the case of s = 5 for $N \leq 13$. The results are given in terms of ratios of integers and we find that the conjectural determinant representation for the EFP reproduces them exactly.

4. EFP as a Fredholm determinant

In this section our aim is to proceed with the determinant formulas for the EFP and obtain some other representations, specifically, in the form of Fredholm determinants of linear integral operators. We also show that in the scaling limit $(r, s \to \infty$ with their ratio fixed) some expressions simplify, and, moreover, at the critical value of s/r corresponding to the arctic curve, there exists local scaling where the Fredholm determinant turns into the celebrated formula in terms of the Airy kernel for the Tracy–Widom distribution.

4.1. Integral form of the matrices. We start with the explicit formula for the Laguerre polynomials, see (3.12). It can be easily seen that for a trial function f(z), regular at the point z = 0, the following holds:

$$\oint_{C_0} \frac{(1-z)^{n+\alpha}}{z^{n+1}} f(z) \frac{\mathrm{d}z}{2\pi \mathrm{i}} = (-1)^n L_n^{(\alpha)}(\partial_z) f(z) \big|_{z=0}.$$
(4.1)

We are interested in the special cases $\alpha = 0$ and $\alpha = -1$.

Reading formula (4.1) in the reverse order, for the entries of the matrices L and U given in (3.13) we can write

$$L_{ij} = \frac{1}{h_{r+i}(0)} \oint_{C_0} \left(\frac{(1-z)^{i-j-1}}{z^{i-j+1}} + (-1)^i \frac{(1-z)^{i-j-1}}{z^{i-j}} \right) h_{r+i}(z) \frac{\mathrm{d}z}{2\pi \mathrm{i}},$$
$$U_{ij} = \frac{1}{h_{r+j}(0)} \oint_{C_0} \left(\frac{(1-z)^{j-i-1}}{z^{j-i+1}} + (-1)^{j+1} \frac{(1-z)^{j-i-1}}{z^{j-i}} \right) h_{r+j}(z) \frac{\mathrm{d}z}{2\pi \mathrm{i}}$$

Let us consider now the matrix A = DLU, where $D_{ij} = \delta_{ij}h_{r+j}(0)$, see (3.13). Slightly simplifying expressions, we get

$$A_{ij} = \sum_{l=1}^{\min(i,j)} \frac{1}{h_{r+j}(0)} \oint_{C_0} \frac{(1-z)^{i-l-1}}{z^{i-l+1}} \left(1 + (-1)^i z\right) h_{r+i}(z) \frac{\mathrm{d}z}{2\pi \mathrm{i}} \\ \times \oint_{C_0} \frac{(1-w)^{j-l-1}}{w^{j-l+1}} \left(1 + (-1)^{j+1} w\right) h_{r+j}(w) \frac{\mathrm{d}w}{2\pi \mathrm{i}}.$$
(4.2)

Let us now transform expression (4.2). First, we note that the sum over l in (4.2) can be extended to infinity without altering the result, since the actual value of the upper limit is controlled by the integrals with the respect to z and w. Next, we evaluate this sum, that yields

$$\sum_{l=1}^{\infty} \left(\frac{zw}{(1-z)(1-w)}\right)^l = \frac{zw}{1-z-w}$$

Finally, introducing the functions

$$e_i^L(z) = \frac{(1-z)^{i-1}}{z^i} \left(1 + (-1)^i z\right) h_{r+i}(z),$$

$$e_j^U(w) = \frac{(1-w)^{j-1}}{h_{r+j}(0)w^j} \left(1 + (-1)^{j+1}w\right) h_{r+j}(w),$$
(4.3)

for entries of the matrix A we arrive at the formula

$$A_{ij} = \oint_{C_0} \oint_{C_0} \frac{e_i^L(z)e_j^U(w)}{1 - z - w} \frac{\mathrm{d}z\mathrm{d}w}{(2\pi\mathrm{i})^2}.$$
(4.4)

We will use this formula as the main input for what follows.

As for the matrix V, see (3.14), one can easily obtain a formula similar to (4.4).

4.2. Fredholm determinants. Let \hat{K}_{Γ} denote a linear integral operator with the kernel K(z, w) acting on functions on the contour Γ ,

$$(\hat{K}_{\Gamma}f)(z) = \int_{\Gamma} K(z,w)f(w) \,\mathrm{d}w.$$

The Fredholm determinant of this operator is usually defined as

$$\det\left(1-\hat{K}_{\Gamma}\right)=1+\sum_{n=1}^{\infty}\frac{(-1)^n}{n!}\int_{\Gamma}\cdots\int_{\Gamma}\det_{1\leqslant i,j\leqslant n}\left[K(w_i,w_j)\right]\,\mathrm{d}w_1\cdots\mathrm{d}w_n.$$

Another way is to use the identity

t

$$\det \left(1 - \hat{K}_{\Gamma}\right) = \exp \left\{ \operatorname{tr} \log \left(1 - \hat{K}_{\Gamma}\right) \right\}$$

and defining the function $\log(1 - \hat{K}_{\Gamma})$ by its power series expansion in powers of \hat{K}_{Γ} , one can also write

$$\det\left(1-\hat{K}_{\Gamma}\right) = \exp\left\{-\sum_{n=1}^{\infty}\frac{1}{n}\operatorname{tr}\left(\hat{K}_{\Gamma}\right)^{n}\right\},\,$$

where

$$\mathbf{r}\left(\hat{K}_{\Gamma}\right)^{n} = \int_{\Gamma} \cdots \int_{\Gamma} K(w_{1}, w_{2}) \cdots K(w_{n}, w_{1}) \,\mathrm{d}w_{1} \cdots \mathrm{d}w_{n}$$

One can say that the Fredholm determinants of an $s \times s$ matrix A and linear integral operator \hat{K}_{Γ} are equal to each other,

$$\det_s(I-A) = \det\left(1 - \hat{K}_{\Gamma}\right),\tag{4.5}$$

where the dependence on s is somehow encoded into \hat{K}_{Γ} , when

$$\operatorname{tr}_{s} A^{n} = \operatorname{tr} \left(\hat{K}_{\Gamma} \right)^{n}, \qquad n = 1, 2, \dots$$

Here, the subscript s recalls that the matrix A is taken to be $s \times s$.

Let us now consider our matrix A given by (4.4). We can immediately claim that we have the identity (4.5), where $\Gamma = C_0$ and the kernel is given by

$$K(z_1, z_2) = \sum_{j=1}^{s} \frac{1}{(2\pi i)^2} \oint_{C_0} \frac{e_j^U(w) e_j^L(z_2)}{1 - z_1 - w} \, \mathrm{d}w, \qquad z_1, z_2 \in C_0.$$
(4.6)

Another choice could be with the functions e_j^U and e_j^L being exchanged. Note that the dependence on s is encoded into the kernel in the sum over j.

To make our considerations below slightly simpler, we find it useful at this stage to introduce an integral operator acting on functions on the real half-axis, $[0, \infty)$. Since z and w lie on the contour C_0 , they can be chosen such that $|z|, |w| \ll 1$, and we can use the formula

$$\frac{1}{1-z-w} = \int_0^\infty e^{(z+w-1)t} \,\mathrm{d}t, \qquad \operatorname{Re}\left(z+w\right) < 1.$$
(4.7)

We can rewrite entries of the matrix A in the form

$$A_{ij} = \int_0^\infty \mathrm{e}^{-t} E_i^L(t) E_j^U(t) \,\mathrm{d}t$$

where we have introduced the functions

$$E_{j}^{L,U}(t) = \oint_{C_{0}} e^{wt} e_{j}^{L,U}(w) \frac{\mathrm{d}w}{2\pi \mathrm{i}}.$$
18

As a result, the EFP can be given as the Fredholm determinant

$$F_N^{(N-s,s)} = \det\left(1 - \hat{K}_{[0,\infty)}^E\right),$$

where the kernel is

$$K^{E}(t_{1}, t_{2}) = e^{-\frac{1}{2}(t_{1}+t_{2})} \sum_{j=1}^{s} E_{j}^{L}(t_{1})E_{j}^{U}(t_{2}).$$

More explicitly,

$$K^{E}(t_{1}, t_{2}) = \oint_{C_{0}} \oint_{C_{0}} e^{\left(z - \frac{1}{2}\right)t_{1} + \left(w - \frac{1}{2}\right)t_{2}} \sum_{j=1}^{s} e_{j}^{L}(z) e_{j}^{U}(w) \frac{\mathrm{d}z\mathrm{d}w}{(2\pi\mathrm{i})^{2}}.$$
 (4.8)

Clearly, the kernel $K^E(t_1, t_2)$ can get a simplified form when the saddle-point analysis is applied to the contour integrals. This is what we address below.

4.3. Scaling limit. Let us now consider the situation where the size of the system, N, is large. To have an interesting picture, r and s have to be taken large as well. We recall that we consider here only the 'symmetric' case, where r = N - s.

An important ingredient for subsequent considerations is the following asymptotic result for the function $h_N(z)$.

Proposition 4.1. As $r \to \infty$,

$$h_{r+1}(z) = [\rho(z)]^r C(z) \left(1 + O(r^{-1})\right), \qquad (4.9)$$

where

$$\rho(z) = 4 \frac{(1-2z)(2-z)(1+z) + 2(1-z+z^2)^{3/2}}{27(1-z)^2}$$
(4.10)

and

$$C(z) = \frac{2z}{\sqrt{2(1-z+z^2)^2 - (2-z)(1-2z)(1+z)\sqrt{1-z+z^2}}}.$$
 (4.11)

We give a proof of this result in Appendix.

Consider now the sum over j in (4.8) or in (4.6). From (4.3) we have

$$\sum_{j=1}^{s} e_j^L(z) e_j^U(w) = \sum_{j=1}^{s} [1 - zw + (-1)^j (z - w)] \times \frac{[(1 - z)(1 - w)]^{j-1}}{(zw)^j} \frac{h_{r+j}(z)h_{r+j}(w)}{h_{r+j}(0)}.$$

For r large, we can use (4.9). Denoting

$$\psi(z,w) = \frac{\rho(z)\rho(w)}{\rho(0)},$$
¹⁹

we get

$$\begin{split} \sum_{j=1}^{s} e_{j}^{L}(z) e_{j}^{U}(w) &\sim \frac{C(z)C(w) \left[(1-z)(1-w)\right]^{s-1} \left[\psi(z,w)\right]^{r+s-1}}{C(0)(zw)^{s}} \\ &\times \left\{ (1-zw) \frac{1 - \left(\frac{zw}{(1-z)(1-w)\psi(z,w)}\right)^{s}}{1 - \frac{zw}{(1-z)(1-w)\psi(z,w)}} \right. \\ &+ (-1)^{s}(w-z) \frac{1 - \left(-\frac{zw}{(1-z)(1-w)\psi(z,w)}\right)^{s}}{1 + \frac{zw}{(1-z)(1-w)\psi(z,w)}} \right\}. \quad (4.12) \end{split}$$

By symbol ~ we denote that the two expressions are equal as $s, r \to \infty$ up to terms which vanish in the limit; more exactly, $f \sim g$ means that $f/g \to 1$.

The expression (4.12) can be simplified by noting that z and w are integrated around the origin in (4.8) and in (4.6). Since $\psi(z, w)$ is regular at the points z = 0and w = 0, from (4.8) we get

$$K^{E}(t_{1}, t_{2}) \sim \oint_{C_{0}} \oint_{C_{0}} e^{\left(z - \frac{1}{2}\right)t_{1} + \left(w - \frac{1}{2}\right)t_{2}} \\ \times \frac{C(z)C(w)\left[(1 - z)(1 - w)\right]^{s-1}\left[\psi(z, w)\right]^{r+s-1}}{C(0)(zw)^{s}} \\ \times \left\{\frac{1 - zw}{1 - \frac{zw}{(1 - z)(1 - w)\psi(z, w)}} + \frac{(-1)^{s}(w - z)}{1 + \frac{zw}{(1 - z)(1 - w)\psi(z, w)}}\right\} \frac{\mathrm{d}z\mathrm{d}w}{(2\pi\mathrm{i})^{2}}.$$
 (4.13)

When r and s are large, the integrals over z and w can be approximated by the saddle-point method. Moreover, both integrals in (4.13) contain as the main factors in their integrands the same function,

$$\left(\frac{1-w}{w}\right)^s \left[\frac{\rho(w)}{\sqrt{\rho(0)}}\right]^N =: \exp\{Ng(w)\},\tag{4.14}$$

where we have used that r + s = N and included the factor $\sqrt{\rho(0)} = 4/3\sqrt{3}$ for a later convenience (it simplifies a constant term in g(w)).

Let us find the saddle points of the function g(w). Denoting

$$y = \frac{s}{N},$$

 $y \in (0, 1/2]$, we have

$$g(w) = y \log \frac{1-w}{w} + \log \frac{(1-2w)(2-w)(1+w) + 2(1-w+w^2)^{3/2}}{3\sqrt{3}(1-w)^2}.$$

We get

$$g'(w) = \frac{y}{w(w-1)} - \frac{1 - \sqrt{1 - w + w^2}}{w(w-1)},$$

and hence the saddle point equation g'(w) = 0 possesses two solutions $w = w_{\pm}$, where

$$w_{\pm} = \frac{1 \pm \sqrt{1 - 8y + 4y^2}}{20}.$$

The two solutions collide if $1 - 8y + 4y^2 = 0$, and choosing the root of this equation that lies in the interval (0, 1/2], we find that $w_+ = w_-$ when $y = y_c$, where

$$y_{\rm c} = 1 - \frac{\sqrt{3}}{2}.\tag{4.15}$$

The critical value y_c separating two asymptotic regimes has a very clear meaning in terms of the so-called arctic curve. This curve describes spatial separation between the ordered and disordered phases of the model in the thermodynamic limit. In the language of dimer models, it is the frozen boundary of the limit shape. For the six-vertex model with domain wall boundary conditions (with a generic choice of weights corresponding to $\Delta < 1$), it is a curve inscribed into the unit square, consisting of four portions joining the four contact points (see [10] for more details). All the four portions are related to each other by simple symmetry transformations. They are described in general by a non-algebraic equation given in a parametric form. For the ice point ($\Delta = 1/2$ and t = 1) this equation appear to be algebraic, and, moreover, quadratic [9, 45],

$$4x(1-x) + 4y(1-y) + 4xy = 1, \qquad x, y \in [0, 1/2]. \tag{4.16}$$

The arctic curve in the ice-point case is constructed by taking the arc between the points (x, y) = (0, 1/2) and (x, y) = (1/2, 0), given by (4.16), and applying to it the maps $(x, y) \mapsto (x, 1-y), (x, y) \mapsto (1-x, y)$, and $(x, y) \mapsto (1-x, 1-y)$. This yields all four portions of the arctic curve.

In the context of the EFP, we need only the arc (4.16). Recalling that in the scaling limit the geometric parameters of the EFP scales as (N - r)/N =: x and s/N =: y, we find that the case r = N - s which we consider here, means simply that x = y. Setting x = y in (4.16) we immediately get the equation $4y^2 - 8y + 1 = 0$, i.e., we get $y = y_c$ where y_c is given by (4.15). Thus the coincidence of the two roots w_{\pm} corresponds to the situation where the frozen region of the EFP touches the arctic curve. The case $y \in (0, y_c)$ corresponds to the frozen region of the EFP fully lying in the ordered region, outside the arctic curve, while the case $y \in (y_c, 1/2]$ corresponds to the frozen region of the EFP partially overlapping with the disordered region, inside the arctic curve.

The two regimes $y \in (0, y_c)$ and $y \in (y_c, 1/2]$ correspond to two different behaviours of the EFP. The values $y \in (0, y_c)$ correspond to w_{\pm} real. In this case one finds an exponentially decaying asymptotic behaviour of the integrals. Among the two points, only the closest to origin, w_- , is relevant in the saddle-point approximation. The values $y \in (y_c, 1/2]$ correspond to w_{\pm} complex, $w_- = w_{\pm}^*$. In this case both saddle points are relevant and the leading term in the asymptotic expansion of the integrals is oscillatory, giving non-vanishing contribution in the limit of large N (and s). Analogous behaviours occur in the context of dimer models [46].

4.4. Tracy–Widom distribution. Let us now focus on the vicinity of the point $y = y_c$. We introduce a new temporary geometric parameter η ,

$$y = y_{\rm c} - \eta$$

As N is large, η is assumed to be small, below we find that $\eta \propto N^{-2/3}$.

Let us consider the behaviour of the function g(w) in the vicinity of the double saddle point $w_+ = w_- = 1/2$. Setting $w = 1/2 + \lambda$, and expanding in the Taylor



FIGURE 5. Deformation of the integration contour in the *w*-complex plane; shaded area is a vicinity of the saddle point at w = 1/2.

series in λ , we find

$$g(w)\Big|_{w=\frac{1}{2}+\lambda} = 4\eta\lambda - \frac{4}{3\sqrt{3}}\lambda^3 + O(\lambda^4).$$

Absence of a constant term here is due to the factor $\sqrt{\rho(0)}$ in (4.14).

Hence, setting

$$\tilde{\lambda} = q\lambda, \qquad q = \frac{2^{2/3}}{3^{1/6}} N^{1/3},$$
(4.17)

and

$$\sigma = \frac{4N}{q}\eta = 2^{4/3}3^{1/6}N^{2/3}\eta,$$

we get

$$Ng(w)\Big|_{w=\frac{1}{2}+\lambda} = \sigma \tilde{\lambda} - \frac{1}{3}\tilde{\lambda}^3 + O(N^{-1/3}),$$
 (4.18)

where we write the *O*-term assuming that σ and $\tilde{\lambda}$ are both of O(1) as $N \to \infty$. This means that in the saddle-point approximation we are considering here $\eta \propto N^{-2/3}$ and $\lambda \propto N^{-1/3}$. Moreover, expression (4.18) fixes the location of the contour of integration along the steepest descent directions.

We recall that the initial contour of integration is around the point w = 0. One can deform the contour such that it arrives at the point w = 1/2 from direction $-2\pi/3$ and departs from it in the direction $2\pi/3$, see Fig. 5. The main contribution to the integral comes from that part of the deformed contour lying in the vicinity of the double saddle point w = 1/2. Let us denote it by γ and let the vicinity of the saddle point be a disk of some radius a (shown in light grey in the picture). Then,

$$\gamma = (e^{-2i\pi/3}a, 0) \cup (0, e^{2i\pi/3}a).$$

From (4.17) it follows that for the scaled variable $\tilde{\lambda}$ the contour of integration is γ , but scaled by the factor of q. Let us denote this contour by $\tilde{\gamma}$. Since $q \to \infty$ as $N \to \infty$, the end-points of $\tilde{\gamma}$ tend to infinity, and therefore, as it can be shown that up to exponentially small corrections to the resulting integral,

$$\tilde{\gamma} = (e^{-2i\pi/3}\infty, 0) \cup (0, e^{2i\pi/3}\infty).$$
 (4.19)

Similar considerations hold for the integration variable z. We set $z = 1/2 + \mu$, and introduce the scaled variable $\tilde{\mu}$ related to μ by $\mu = q\tilde{\mu}$, see (4.17). The contour of integration for $\tilde{\mu}$ is $\tilde{\gamma}$.

Let us now inspect the expression in the braces in (4.13). Since λ and μ scale as $N^{-1/3}$, we have

$$\frac{zw}{(1-z)(1-w)\psi(z,w)}\Big|_{w=\frac{1}{2}+\lambda,z=\frac{1}{2}+\mu} = 1 + 2\sqrt{3}(\lambda+\mu) + O(N^{-2/3}).$$

The first term in the braces in (4.13) behaves as $1/(\lambda + \mu) = O(N^{1/3})$, and the second term is just of $O(N^{-1/3})$ thus contributing only to sub-leading terms. As for the first term, taking also into account the prefactor in the integrand for which we can use

$$C(1/2) = \frac{2\sqrt{2}}{3}, \qquad C(0) = \frac{4}{3\sqrt{3}}, \qquad \psi(1/2, 1/2) = 1,$$

we get

$$\frac{C(w)C(z)}{C(0)(1-z)(1-w)\psi(z,w)}\frac{1-wz}{1-\frac{zw}{(1-z)(1-w)\psi(z,w)}}\bigg|_{w=\frac{1}{2}+\lambda,z=\frac{1}{2}+\mu} = -\frac{1}{\lambda+\mu}+O(1).$$

As an intermediate result, we thus have in the leading order

$$K^{E}(t_{1},t_{2}) = -\int_{\gamma} \int_{\gamma} \frac{\mathrm{e}^{\mu t_{1} + \lambda t_{2} + \sigma(\tilde{\lambda} + \tilde{\mu}) - (\tilde{\lambda}^{3} + \tilde{\mu}^{3})/3}}{\lambda + \mu} \frac{\mathrm{d}\lambda \mathrm{d}\mu}{(2\pi\mathrm{i})^{2}}.$$

To turn to the scaled integration variables λ and $\tilde{\mu}$, we change the variables of the kernel, $t_1 \mapsto qt_1, t_2 \mapsto qt_2$ and introduce the new kernel

$$\widetilde{K}^{E}(t_{1}, t_{2}) = qK^{E}(qt_{1}, qt_{2}), \qquad q > 0, \qquad t_{1}, t_{2} \in [0, \infty).$$

Note that the Fredholm determinants for the kernels $\tilde{K}^{E}(t_1, t_2)$ and $K^{E}(t_1, t_2)$ are equal to each other, because this holds for the traces of arbitrary positive integer powers of the corresponding integral operators. We have

$$\widetilde{K}^{E}(t_{1},t_{2}) = -\int_{\widetilde{\gamma}} \int_{\widetilde{\gamma}} \frac{\mathrm{e}^{\widetilde{\mu}t_{1}+\widetilde{\lambda}t_{2}+\sigma(\widetilde{\lambda}+\widetilde{\mu})-(\widetilde{\lambda}^{3}+\widetilde{\mu}^{3})/3}}{\widetilde{\lambda}+\widetilde{\mu}} \frac{\mathrm{d}\widetilde{\lambda}\mathrm{d}\widetilde{\mu}}{(2\pi\mathrm{i})^{2}}$$

Summing up, we may write

$$\lim_{N \to \infty} \left(F_N^{(N-s,s)} \Big|_{s=N\left(1-\frac{\sqrt{3}}{2}\right) - \frac{N^{1/3}}{2^{4/3}3^{1/6}}\sigma} \right) = \det\left(1 - \hat{\widetilde{K}}^E_{[0,\infty)}\right).$$

From this point we shall work with these Fredholm determinant and kernel.

We transform the expression for $\tilde{K}^E(t_1, t_2)$ by noting that the contour $\tilde{\gamma}$, see (4.19), can be slightly deformed such that it passes near the origin ($\tilde{\lambda} = 0$, i.e., $w = \frac{1}{2}$) from the left, so that it can be made to lie entirely in the left half-plane. Note that this is in agreement also with the condition in (4.7). Then, we can use

$$\int_{\sigma}^{\infty} \mathrm{e}^{(\tilde{\lambda}+\tilde{\mu})t} \, \mathrm{d}t = -\frac{\mathrm{e}^{(\tilde{\lambda}+\tilde{\mu})\sigma}}{\tilde{\lambda}+\tilde{\mu}}, \qquad \mathrm{Re}\left(\tilde{\lambda}+\tilde{\mu}\right) < 0,$$
²³

that is valid since $\operatorname{Re} \tilde{\lambda} < 0$ and $\operatorname{Re} \tilde{\mu} < 0$ along the contour $\tilde{\gamma}$. We also use the fact that

$$\int_{\tilde{\gamma}} e^{t\tilde{\lambda} - \tilde{\lambda}^3/3} \frac{\mathrm{d}\tilde{\lambda}}{2\pi\mathrm{i}} = \mathrm{Ai}(t),$$

where Ai(t) is the Airy function, that follows from the well-known contour integral definition of the Airy function in which the change $\tilde{\lambda} \mapsto -\tilde{\lambda}$ has been made. As a result, for the kernel $\tilde{K}^E(t_1, t_2)$ we obtain the expression

$$\widetilde{K}^E(t_1, t_2) = \int_{\sigma}^{\infty} \operatorname{Ai}(t_1 + t) \operatorname{Ai}(t_2 + t) dt.$$

As a final step, we note that

$$\det\left(1-\hat{\tilde{K}}^{E}_{[0,\infty)}\right) = \det\left(1-\hat{K}^{\mathrm{Ai}}_{[\sigma,\infty)}\right) =: \mathcal{F}_{2}(\sigma),$$

where the kernel $K^{Ai}(t_1, t_2)$ is given by

$$K^{\rm Ai}(t_1, t_2) = \int_0^\infty \operatorname{Ai}(t_1 + t) \operatorname{Ai}(t_2 + t) \,\mathrm{d}t, \qquad (4.20)$$

and the function $\mathcal{F}_2(\sigma)$ is the celebrated GUE Tracy–Widom distribution (for the Gaussian unitary ensemble, $\beta = 2$). The kernel (4.20) is also known in the form

$$K^{\text{Ai}}(t_1, t_2) = \frac{\text{Ai}(t_1) \text{Ai}'(t_2) - \text{Ai}'(t_1) \text{Ai}(t_2)}{t_1 - t_2}$$

The proof of equivalence of the two expressions for $K^{Ai}(t_1, t_2)$ can be found, e.g., in [15].

Summarizing, we conclude that representation (3.13) for the EFP in the sixvertex model with domain wall boundary condition at the ice point ($\Delta = 1/2$ and t = 1) implies that

$$\lim_{N \to \infty} \left(F_N^{(N-s,s)} \Big|_{s=N\left(1-\frac{\sqrt{3}}{2}\right)-\frac{N^{1/3}}{2^{4/3}3^{1/6}}\sigma} \right) = \mathcal{F}_2(\sigma).$$

This result is in a full agreement with the one proposed in [29] (see Conjecture 2.5 therein) and with the numerical simulations in [27,28].

Acknowledgments

We are indebted to N. M. Bogoliubov, L. Cantini, S. Chhita, H. Spohn, and J. Viti, for stimulating discussions. AGP is grateful to INFN, Sezione di Firenze, for hospitality and partial support during his stay in Florence, Italy, where a part of this work was done. AGP also acknowledges partial support from the Theoretical Physics and Mathematics Advancement Foundation «BASIS».

Appendix A. Asymptotic expansion of $h_{r+1}(z)$ as $r \to \infty$

We have the following expression [9]:

$$h_{r+1}(z) = \frac{\Gamma(2r+2)}{\left[\Gamma(r+1)\right]^2} \int_0^1 \left\{ t(1-t) \left[1 - (1-z)t\right] \right\}^r \, \mathrm{d}t.$$
(A.1)

Consider first the prefactor in this representation. From the Stirling formula

$$\log \Gamma(z+a) = \left(z+a - \frac{1}{2}\right) \log z - z + \log \sqrt{2\pi} + O(1/z)$$
24

we have, as $r \to \infty$,

$$\frac{\Gamma(2r+2)}{\left[\Gamma(r+1)\right]^2} = 4^r \sqrt{\frac{4r}{\pi}} \left(1 + O(r^{-1})\right).$$

Consider now the integral in (A.1). The standard saddle-point analysis yields

$$\int_0^1 \left\{ t(1-t) \left[1 - (1-z)t \right] \right\}^r \, \mathrm{d}t = \mathrm{e}^{rf(t_0)} \sqrt{\frac{2\pi}{|f''(t_0)|r|}} \left(1 + O(r^{-1}) \right),$$

where we have denoted

$$f(t) = \log(t(1-t)[1-(1-z)t]),$$

and $t_0 = t_0(z)$ is the saddle point, $f'(t_0) = 0$, given by

$$t_0 = \frac{2 - z - \sqrt{1 - z + z^2}}{3(1 - z)}.$$
 (A.2)

Note that t_0 lies on the interval of integration in (A.1) for broad values of z, including the interval [0, 1]. We also have

$$f''(t_0) = -2\frac{2(1-z+z^2)^2 - (2-z)(1-2z)(1+z)\sqrt{1-z+z^2}}{z^2}.$$
 (A.3)

It is easy to see that $f''(t_0)$ is negative, and so the original contour goes along the steepest descent directions. Note that there exists another solution of the equation f'(t) = 0, but it is not relevant (both for analytical and topological reasons).

Collecting things together, we obtain

$$h_{r+1}(z) = [\rho(z)]^r C(z) \left(1 + O(r^{-1})\right),$$

where

$$\rho(z) = 4e^{f(t_0)} = 4t_0(1-t_0)\left[1 - (1-z)t_0\right]$$

and

$$C(z) = \sqrt{\frac{8}{|f''(t_0)|}}.$$

Substitution of (A.2) yields the expression (4.10) for the function $\rho(z)$, and (A.3) leads to (4.11) for C(z).

References

- V. E. Korepin, N. M. Bogoliubov, and A. G. Izergin, Quantum inverse scattering method and correlation functions, Cambridge University Press, Cambridge, 1993.
- [2] H. E. Boos and V. E. Korepin, Evaluation of integrals representing correlations in the XXX Heisenberg spin chain, In: MathPhys Odyssey 2001: Integrable Models and Beyond In Honor of Barry M. McCoy (M. Kashiwara and T. Miwa, eds.), Progress in Mathematical Physics, vol. 23, Birkhäuser Boston, MA, 2002, pp. 65–108, doi:10.1007/978-1-4612-0087-1_4, arXiv: hep-th/0105144.
- [3] H. E. Boos, V. E. Korepin, Y. Nishiyama, and M. Shiroishi, *Quantum correlations and number theory*, J. Phys. A **35** (2002), 4443–4451, doi:10.1088/0305-4470/35/20/305, arXiv: cond-mat/0202346.
- [4] V.E. Korepin, S. Lukyanov, Y. Nishiyama, and M. Shiroishi, Asymptotic behavior of the emptiness formation probability in the critical phase of XXZ spin chain, Phys. Lett. A 312 (2003), 21–26, doi:10.1016/S0375-9601(03)00616-9, arXiv:cond-mat/0210140.
- [5] H. E. Boos, V. E. Korepin, and F. A. Smirnov, Emptiness formation probability and quantum Knizhnik-Zamolodchikov equation, Nucl. Phys. B 658 (2003), 417–439, doi: 10.1016/S0550-3213(03)00153-6, arXiv:hep-th/0209246.

- [6] F. Göhmann, A. Klümper, and A. Seel, Emptiness formation probability at finite temperature for the isotropic Heisenberg chain, Physica B 359-361 (2005), 807-809, doi:10.1016/j.physb. 2005.01.234, arXiv:cond-mat/0406611.
- [7] N. Kitanine, K. K. Kozlowski, J. M. Maillet, N. A. Slavnov, and V. Terras, Algebraic Bethe ansatz approach to the asymptotic behavior of correlation functions, J. Stat. Mech.: Theor. Exp. 2009 (2009), P04003, doi:10.1088/1742-5468/2009/04/P04003, arXiv:0808.0227.
- [8] F. Colomo and A. G. Pronko, Emptiness formation probability in the domain-wall six-vertex model, Nucl. Phys. B 798 (2008), 340-362, doi:10.1016/j.nuclphysb.2007.12.016, arXiv: 0712.1524.
- F. Colomo and A. G. Pronko, The limit shape of large alternating-sign matrices, SIAM J. Discrete Math. 24 (2010), 1558–1571, doi:10.1137/080730639, arXiv:0803.2697.
- [10] F. Colomo and A. G. Pronko, The arctic curve of the domain-wall six-vertex model, J. Stat. Phys. 138 (2010), 662–700, doi:10.1007/s10955-009-9902-2, arXiv:0907.1264.
- [11] F. Colomo and A. Sportiello, Arctic curves of the six-vertex model on generic domains: the tangent method, J. Stat. Phys. 164 (2016), 1488–1523, doi:10.1007/s10955-016-1590-0, arXiv:1605.01388.
- [12] L. Cantini, F. Colomo, and A. G. Pronko, Integral formulas and antisymmetrization relations for the six-vertex model, Ann. Henry Poincaré 21 (2020), 865–884, doi:10.1007/ s00023-019-00856-6, arXiv:1906.07636.
- [13] F. Colomo, G. Di Giulio, and A. G. Pronko, Six-vertex model on a finite lattice: Integral representations for nonlocal correlation functions, Nucl. Phys. B 972 (2021), 115535 (42 pp.), doi:10.1016/j.nuclphysb.2021.115535, arXiv:2107.13358.
- [14] C. A. Tracy and H. Widom, Level spacing distributions and the Airy kernel, Comm. Math. Phys. 159 (1994), 151–174, doi:10.1007/BF02100489, arXiv:hep-th/9211141.
- [15] L. M. Mehta, Random matrices, 3rd ed., Pure and Applied Mathematics, vol. 142, Elsevier and Academic Press, Amsterdam, 2004.
- [16] P. Deift, Universality for mathematical and physical systems, In: Proceedings of the International Congress of Mathematicians, Madrid, 2006 (M. Sanz-Solé, J. Soria, J. L. Varona, and J. Verdera, eds.), European Mathematical Society, Zurich, 2007, pp. 125–152, doi: 10.4171/022-1/7, arXiv:math-ph/0603038.
- [17] K. Johansson, Shape fluctuations and random matrices, Comm. Math. Phys. 209 (2000), 437–476, doi:10.1007/s002200050027, arXiv:math/9903134.
- [18] K. Johansson, Non-intersecting paths, random tilings and random matrices, Probab. Theory Relat. Fields 123 (2002), 225-280, doi:10.1007/s004400100187, arXiv:math.PR/0011250.
- [19] K. Johansson, The arctic circle boundary and the Airy process, Ann. Probab. 33 (2005), 1–30, doi:10.1214/009117904000000937, arXiv:math.PR/0306216.
- [20] F. Colomo and A. G. Pronko, Third-order phase transition in random tilings, Phys. Rev. E 88 (2013), 042125, doi:10.1103/PhysRevE.88.042125, arXiv:1306.6207.
- [21] P. L. Ferrari and H. Spohn, Domino tilings and the six-vertex model at its free fermion point, J. Phys. A **39** (2006), 10297–10306, doi:10.1088/0305-4470/39/33/003, arXiv:cond-mat/ 0605406.
- [22] A. V. Kitaev and A. G. Pronko, Emptiness formation probability of the six-vertex model and the sixth Painlevé equation, Comm. Math. Phys. 345 (2016), 305–354, doi:10.1007/ s00220-016-2636-5, arXiv:1505.00032.
- [23] C. A. Tracy and H. Widom, Integral formulas for the asymmetric simple exclusion process, Comm. Math. Phys. 279 (2008), 815–844, doi:10.1007/s00220-008-0443-3, arXiv:0704.2633.
- [24] C. A. Tracy and H. Widom, A Fredholm determinant representation in ASEP, J. Stat. Phys. 132 (2008), 291–300, doi:10.1007/s10955-008-9562-7, arXiv:0804.1379.
- [25] C. A. Tracy and H. Widom, Asymptotics in ASEP with step initial condition, Comm. Math. Phys. 290 (2009), 129–154, doi:10.1007/s00220-009-0761-0, arXiv:0807.1713.
- [26] A. Borodin, I. Corwin, and V. Gorin, Stochastic six-vertex model, Duke Math. J. 165 (2016), 563–624, doi:10.1215/00127094-3166843, arXiv:1407.6729.
- [27] I. Lyberg, V. Korepin, and J. Viti, Fluctuation of the phase boundary in the six-vertex model with domain wall boundary conditions: a Monte Carlo study, J. Phys. A: Math. Theor. 56 (2023), 495002, doi:10.1088/1751-8121/ad0a43, arXiv:2303.14669.
- [28] M. Prähofer and H. Spohn, Domain wall fluctuations of the six-vertex model at the ice point, J. Phys. A: Math. Theor. 57 (2023), 025001, doi:10.1088/1751-8121/ad13b4, arXiv:2305.09502.

- [29] A. Ayyer, S. Chhita, and K. Johansson, GOE fluctuations for the maximum of the top path in alternating sign matrices, Duke Math. J. 172 (2023), 1961–2104, doi:10.1215/ 00127094-2022-0075, arXiv:2109.02422.
- [30] R. J. Baxter, Exactly solved models in statistical mechanics, Academic Press, San Diego, CA, 1982.
- [31] V. E. Korepin, Calculations of norms of Bethe wave functions, Commun. Math. Phys. 86 (1982), 391–418, doi:10.1007/BF01212176.
- [32] A. G. Izergin, Partition function of the six-vertex model in a finite volume, Sov. Phys. Dokl. 32 (1987), 878–879.
- [33] A. G. Izergin, D. A. Coker, and V. E. Korepin, Determinant formula for the six-vertex model, J. Phys. A: Math. Gen. 25 (1992), 4315–4334, doi:10.1088/0305-4470/25/16/010.
- [34] N. Elkies, G. Kuperberg, M. Larsen, and J. Propp, Alternating-sign matrices and domino tilings (Part 1), J. Algebraic Combin. 1 (1992), 111–132, doi:10.1023/A:1022420103267, arXiv: math/9201305.
- [35] N. Elkies, G. Kuperberg, M. Larsen, and J. Propp, Alternating-sign matrices and domino tilings (Part 2), J. Algebraic Combin. 1 (1992), 219–234, doi:10.1023/A:1022483817303, arXiv: math/9201305.
- [36] G. Kuperberg, Another proof of the alternative-sign matrix conjecture, Int. Math. Res. Not. 1996 (1996), 139–150, doi:10.1155/S1073792896000128, arXiv:math/9712207.
- [37] D. Zeilberger, Proof of the refined alternating sign matrix conjecture, New York J. Math. 2 (1996), 59–68, arXiv:math/9606224.
- [38] D. M. Bressoud, Proofs and confirmations: The story of the alternating sign matrix conjecture, Cambridge University Press, Cambridge, 1999.
- [39] N. M. Bogoliubov, A. G. Pronko, and M. B. Zvonarev, Boundary correlation functions of the six-vertex model, J. Phys. A: Math. Gen. 35 (2002), 5525–5541, doi:10.1088/0305-4470/35/ 27/301, arXiv:math-ph/0203025.
- [40] Yu. G. Stroganov, Izergin-Korepin determinant at a third root of unity, Theor. Math. Phys. 146 (2006), 53–62, doi:10.1007/s11232-006-0006-8, arXiv:math-ph/0204042.
- [41] A. G. Pronko and G. P. Pronko, Off-shell Bethe states and the six-vertex model, J. Math. Sci. 242 (2019), 742–752, doi:10.1007/s10958-019-04511-7.
- [42] F. Colomo and A. G. Pronko, On the refined 3-enumeration of alternating sign matrices, Adv. in Appl. Math. 34 (2005), 798–811, doi:10.1016/j.aam.2004.09.007, arXiv:math-ph/0404045.
- [43] A. Saenz, C. A. Tracy, and H. Widom, Domain walls in the Heisenberg-Ising spin-¹/₂ chain, In: Toeplitz Operators and Random Matrices: In Memory of Harold Widom (E. Basor, A. Böttcher, T. Ehrhardt, and C. A. Tracy, eds.), Operator Theory: Advances and Applications, vol. 289, Birkhäuser, Cham, 2022, pp. 9–47, doi:10.1007/978-3-031-13851-5_2, arXiv:2202. 07695.
- [44] I. S. Gradshteyn and I. M. Ryzhik, Table of integrals, series, and products, 8th ed., Academic Press, 2015.
- [45] A. Aggarwal, Arctic boundaries of the ice model on three-bundle domains, Invent. Math. 220 (2019), 611–671, doi:10.1007/s00222-019-00938-6, arXiv:1812.03847.
- [46] R. Kenyon, A. Okounkov, and S. Sheffield, *Dimers and amoebae*, Ann. of Math. 163 (2006), 1019–1056, doi:10.4007/annals.2006.163.1019, arXiv:math-ph/0311005.

INFN, SEZIONE DI FIRENZE VIA G. SANSONE 1, I-50019 SESTO FIORENTINO (FI), ITALY *Email address*: colomo@fi.infn.it

STEKLOV MATHEMATICAL INSTITUTE, FONTANKA 27, 191023 SAINT PETERSBURG, RUSSIA *Email address*: a.g.pronko@gmail.com