A RANDOM WALK APPROACH TO BROADCASTING ON RANDOM RECURSIVE TREES

A PREPRINT

Ernst Althaus Institute of Computer Science Johannes Gutenberg-University Mainz, Germany Lisa Hartung Institute of Mathematics Johannes Gutenberg-University Mainz, Germany Rebecca Steiner Institute of Computer Science Johannes Gutenberg-University Mainz, Germany

May 7, 2024

ABSTRACT

In the broadcasting problem on trees, a $\{0, 1\}$ -message originating in an unknown node is passed along the tree with a certain error probability q. The goal is to estimate the original message without knowing the order in which the nodes were informed. A variation of the problem is considering this broadcasting process on a randomly growing tree, which Addario-Berry et al. [2] have investigated for uniform and linear preferential attachment recursive trees. We extend their studies of the majority estimator to the entire group of very simple increasing trees as well as shape exchangeable trees using the connection to inhomogeneous random walks and other stochastic processes with memory effects such as Pólya Urns.

1 Introduction

Incrementally growing random trees and networks are important building blocks in understanding the formation of networks and their structural properties. In this article we study the broadcasting process for two classes of growing random trees, namely very simple increasing trees, see for example [15], and shape exchangeable trees [10]. The broadcasting process on a growing tree can informally be described as follows: Consider a sequence of trees $(T_n)_{n \in \mathbb{N}}$ where T_{n+1} is obtained by adding one vertex (and edge) to the tree T_n . At the beginning, the tree consists of only the root vertex which receives one of two available colors, say red and blue. Each timestep, one new vertex attaches itself to the tree and receives the color of its parent with probability 1 - q and the opposite color with probability q, independent of the other vertices and their colors. The key question we study in this paper is the influence of the root color on the global appearance of the tree. Specifically, we investigate the relation between the color majority and the root color when growing the tree to infinite size. This can be understood as both a reconstruction problem and as a question of local choices influencing global behavior.

The two classes of random trees we consider are defined as follows. Let $(T_n)_{n \in \mathbb{N}}$ be a sequence of growing trees as above, with T_n containing n vertices. Then the vertices of T_n can naturally be labeled by $\{1, \ldots, n\}$ according to their arrival time, making each T_n a random recursive tree [14] of size n.

Definition 1 (Very Simple Increasing Tree). A very simple increasing (v.s.i.) tree is a random recursive tree $(T_n)_{n \in \mathbb{N}}$ that can be grown iteratively with the following attachment probability distribution for each new vertex:

$$\forall v \in \{1, \dots, n\}: \mathbb{P}(n+1 \sim v \mid T_n) = \frac{\alpha \mathrm{deg}_n^+(v) + 1}{\alpha(n-1) + n},\tag{1}$$

where

$$\alpha \in \left\{ \frac{-1}{d} \mid d \in \mathbb{N}_{>1} \right\} \cup [0, \infty).$$
⁽²⁾

Definition 2 (Shape Exchangeable Tree). A shape exchangeable (s.e.) tree is a random recursive tree $(T_n)_{n \in \mathbb{N}}$ that can be grown iteratively with the following attachment probability distribution for each new vertex:

$$\forall v \in \{1, \dots, n\}: \mathbb{P}(n+1 \sim v \mid T_n) = \frac{\alpha \deg_n(v) + 1}{2\alpha(n-1) + n},\tag{3}$$

where

$$\alpha \in \left\{ \frac{-1}{d} \mid d \in \mathbb{N}_{>1} \right\} \cup [0, \infty).$$
(4)

The difference between these two models is found in their treatment of the root vertex - the root only has outgoing edges, while all other vertices have one ingoing edge. This causes differing attachment probabilities: Consider for example a tree T_2 consisting of two connected vertices. In the shape exchangeable case, these two will be indistinguishable while in very simple increasing trees, vertex 1 will have a larger attachment probability than vertex 2. Thus the root is harder to distinguish from the other vertices in shape exchangeable trees, making the estimation problem more interesting.

Definition 3 (Broadcasting Process). The broadcasting process $(\mathcal{T}_n)_{n \in \mathbb{N}}$ with $\mathcal{T}_n = T_n \times \{-1, 1\}^n$ is a combination of a growing tree process T_n and a coloring $\{-1, 1\}^n$. \mathcal{T}_{n+1} is obtained from \mathcal{T}_n as follows: At time n + 1, the new vertex n + 1 will first choose its parent p_{n+1} according to the attachment distribution given by the tree process. It will then inherit its parent's color $B_{p_{n+1}}$ with probability 1 - q and flip to the other color with probability q independently of all other vertices and their colors. The parameter q is called the bit-flipping probability and one realisation of this process is called a broadcasting tree.

Additionally, vertex 1 has no parent and is therefore assigned a randomly chosen color at time 1. By symmetry, we may simply call this color "red". As mentioned above, the problem that we want to study can be viewed as the reconstruction of the color at the root vertex [2]. Here, we observe the unrooted, unlabeled and entirely colored tree of size N. In particular, we do not have any information about which vertex is the root. The estimator we consider is the *majority estimator*:

Definition 4. The majority estimator $b_{maj}(N,q)$ is defined on a given broadcasting tree of size N with bit-flipping probability q as follows:

$$b_{\text{maj}}(N,q) \coloneqq \begin{cases} \operatorname{sgn}\left(\sum_{u \in T_N} B_u\right) & \text{if not } 0, \\ \operatorname{Rad}\left(\frac{1}{2}\right) & \text{otherwise.} \end{cases}$$
(5)

This estimator either outputs the color majority in a broadcasting tree of size N or, if there is a tie, it makes a random guess. Its error probability is given by

$$\mathbb{P}(b_{\text{maj}}(N,q) \neq B_1) \eqqcolon R_{\text{maj}}(N,q).$$
(6)

We are interested in analysing the limiting behavior of the error probability in relation to q, that is,

$$R_{\text{maj}}(q) \coloneqq \limsup_{N \to \infty} R_{\text{maj}}(N, q).$$
⁽⁷⁾

Related results As we consider the root-bit reconstruction problem, it is naturally linked to root-finding algorithms. For very simple increasing trees it was shown in [7, 25] that there is a set of constant size of vertices that contains the root with high probability and a more general inference problem has been studied for both very simple increasing and shape exchangeable trees in [10]. Root reconstruction is also linked to the question of how a given finite seed graph influences the shape and structure of the resulting tree or graph. We refer to [8, 9, 11, 28], where this problem is studied mainly for random recursive trees. For general networks this may evolve into studying *hubs* or the position of a central vertex, see [3, 4, 24].

The broadcasting process and root color reconstruction have also been investigated on a wide range of random tree [1, 2, 13, 18, 19] and random graph models [6, 30]. For some statistical hardness results for the reconstruction of the root color from the leaf bits, we refer to [16, 21, 26, 35]. Similar problems can be investigated on the stochastic block model [1, 34] and in models arising from statistical physics, such as the Ising model [5, 18]. Finally, the question of the color majority is connected to the study of Pólya Urns [17, 23, 29] and inhomogeneous random walks [22, 31, 32], especially to (reinforced) Elephant Random Walks [12, 27]: Here, the one-dimensional walker remembers a randomly chosen point in the past before each step. With probability 1 - q, it repeats this past step and with probability q it moves in the opposite direction. The relation to the broadcasting process is quite natural: The remembered point in the past corresponds to the chosen parent of the new vertex and the direction the walker moves in is the change in the color difference. To encompass the non-uniform setting, it remains to weigh the past timepoints accordingly. This view of the color majority process gives a phase transition depending on the relation between the model parameter α and the bit-flipping probability q. If q is too large, the process is *diffusive*, while it is *superdiffusive* for small values of q [27]. So far, not much is known about the behavior in the superdiffusive case, but recent work has shed some light on this [20].

To our knowledge, the color majority estimator as we defined it was first investigated on random trees in [2] for a subgroup of the very simple increasing tree model. We aim to complete the picture given by the authors and to provide

a more model-agnostic approach to the problem. Even though our bound on the error probability is not as tight, our approach directly covers both very simple increasing and shape exchangeable trees and we are confident that the random walk perspective we present can be applied to other models as well. For further references, we also refer to their paper.

Outline of the article In Section 2 we present our modelling approaches for the majority estimator and our results on its performance in relation to the bit-flipping probability q, which we then prove in Sections 3 and 4.

Acknowledgement. We thank Stephan Wagner for useful discussions on the broadcasting problem and on [15]. RS thanks Cécile Mailler for useful discussions on elephant random walks. This research is supported by the internal research funding (Stufe I) at Johannes Gutenberg-University, the TOP-ML project and the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) through Project-ID 233630050 - TRR 146.

2 **Results and Preliminaries**

2.1 Main Results

Theorem 5. *Let*

$$f(\alpha) = \begin{cases} \frac{\alpha+1}{4} & \text{for very simple inc. trees} \\ \frac{2\alpha+1}{4(\alpha+1)} & \text{for shape ex. trees.} \end{cases}$$
(8)

For shape exchangeable and very simple increasing trees, it holds that for $q \ge f(\alpha)$,

$$R_{\rm maj}(q) = \frac{1}{2}.\tag{9}$$

Theorem 6. For shape exchangeable and very simple increasing trees, it holds that for each allowed α there exists $c_{\alpha} > 0$ such that

$$R_{\rm maj}(q) \le c_{\alpha} \sqrt{q}.\tag{10}$$

To prove these results, we analyze the majority estimator from two different viewpoints.

2.2 Color Majority as an Inhomogeneous Random Walk

Calculating the color majority of a broadcasting tree does not require any information about the tree structure, only the vertex colors. Therefore, we may consider the process describing the evolution of the *color difference*: We denote by $\#_{red}(n)$ and $\#_{blue}(n)$ the number of red, respectively blue, vertices at time n. Again, by symmetry, we may simply call the color of the root "red". Let

$$\Delta_1(n) \coloneqq \#_{\text{red}}(n) - \#_{\text{blue}}(n) \quad \text{for all } n \le N,$$
(11)

be the color difference at time n, with $\Delta_1(1) = 1$.

Lemma 7.

$$R_{\text{maj}}(q) \le \limsup_{N \to \infty} \mathbb{P}(\Delta_1(N) \le 0).$$
(12)

Proof. We observe that

$$\Delta_1(N) < 0 \implies b_{\text{maj}}(N,q) \neq B_1 \quad \text{and} \quad \Delta_1(N) = 0 \implies b_{\text{maj}}(N,q) = \text{Rad}(\frac{1}{2}), \quad (13)$$

which implies

$$R_{\text{maj}}(N,q) = \mathbb{P}(\Delta_1(N) < 0) + \frac{1}{2}\mathbb{P}(\Delta_1(N) = 0) \le \mathbb{P}(\Delta_1(N) \le 0)$$
(14)
laim follows.

for all $N \in \mathbb{N}$ and the claim follows.

If the underlying tree process T(n) uses uniform attachment ($\alpha = 0$) over the existing vertices, the number of red and blue vertices present at time n - or rather the difference between them - is enough to fully describe the evolution of Δ_1 . In the non-uniform case, the color difference alone does not contain enough information as the attachment distribution depends on the amount of children the vertices have (see also [2]). Therefore, we define a process for the *weight difference*, which we call Δ_2 . In very simple increasing trees, it is given by

$$\Delta_2(n) \coloneqq \#_{\text{children of red vertices}}(n) - \#_{\text{children of blue vertices}}(n) \tag{15}$$

and in shape exchangeable trees by

$$\Delta_2(n) \coloneqq \#_{\text{edges of red vertices}}(n) - \#_{\text{edges of blue vertices}}(n).$$
(16)

 Δ_1 and Δ_2 together with the model parameter α are sufficient to fully describe the attachment distribution on a given broadcasting tree \mathcal{T}_n of size n.

Definition 8. Let

$$Z_{\alpha}(n) \coloneqq \begin{cases} \alpha(1-\frac{1}{n})+1 & \text{for simple inc. trees} \\ 2\alpha(1-\frac{1}{n})+1 & \text{for shape ex. trees.} \end{cases}$$
(17)

Lemma 9.

$$\mathbb{P}(n+1 \sim \text{red vertex} \mid \mathcal{T}_n) = \frac{1}{2} \left(1 + \frac{\Delta_1(n) + \alpha \Delta_2(n)}{Z_\alpha(n)n} \right).$$
(18)

Proof. For very simple increasing trees the attachment distribution is given by:

$$\mathbb{P}(n+1 \sim \text{red vertex} \mid \mathcal{T}_n) = \frac{\sum_{v \text{ red}} (\alpha \deg_n^+(v) + 1)}{\sum_{u \in \mathcal{T}_n} (\alpha \deg_n^+(v) + 1)}$$
$$= \frac{\sum_{v \text{ red}} (\alpha \deg_n^+(v) + 1)}{\alpha(n-1) + n}$$
$$= \frac{1}{2} \left(1 + \frac{\Delta_1(n) + \alpha \Delta_2(n)}{(\alpha(1-\frac{1}{n}) + 1)n} \right) =: p_{vs}(\alpha, n).$$
(19)

Similarly, for shape exchangeable trees,

$$\mathbb{P}(n+1 \sim \text{red vertex} \mid \mathcal{T}_n) = \frac{1}{2} \left(1 + \frac{\Delta_1(n) + \alpha \Delta_2(n)}{(2\alpha(1-\frac{1}{n})+1)n} \right) =: p_{\text{se}}(\alpha, n).$$
(20)

Adding an independent Ber(q)-distributed coin flip lets us fully describe the color difference process Δ_1 . Specifically, $\Delta_1(n)$ and $\Delta_2(n)$ together form a two-dimensional time-inhomogeneous Markov random walk

$$\Delta(n) \coloneqq \begin{pmatrix} \Delta_1(n) \\ \Delta_2(n) \end{pmatrix},\tag{21}$$

with

$$\Delta_1(1) = 1 \qquad \Delta_2(1) = 0 \qquad \text{and} \qquad \Delta(n+1) = \Delta(n) + D(n), \tag{22}$$

where the distribution of D(n) distinguishes the models from another. Let $\mathcal{F}(n)$ be the natural filtration of $\Delta(n)$. Then, in very simple increasing trees D(n) is distributed as

$$\mathbb{P}\left(D(n) = \begin{pmatrix} D_1(n) \\ 1 \end{pmatrix} \middle| \mathcal{F}(n)\right) = p_{vs}(\alpha, n) \cdot \begin{cases} 1-q & \text{if } D_1(n) = 1\\ q & \text{if } D_1(n) = -1 \end{cases}$$

$$\mathbb{P}\left(D(n) = \begin{pmatrix} D_1(n) \\ -1 \end{pmatrix} \middle| \mathcal{F}(n)\right) = (1-p_{vs}(\alpha, n)) \cdot \begin{cases} 1-q & \text{if } D_1(n) = 1\\ q & \text{if } D_1(n) = -1 \end{cases}$$
(23)

and in shape exchangeable trees as

$$\mathbb{P}\left(D(n) = \begin{pmatrix} 1\\2 \end{pmatrix} \middle| \mathcal{F}(n)\right) = p_{se}(\alpha, n) \cdot (1-q)$$

$$\mathbb{P}\left(D(n) = \begin{pmatrix} -1\\0 \end{pmatrix} \middle| \mathcal{F}(n)\right) = p_{se}(\alpha, n) \cdot q$$

$$\mathbb{P}\left(D(n) = \begin{pmatrix} 1\\0 \end{pmatrix} \middle| \mathcal{F}(n)\right) = (1 - p_{se}(\alpha, n)) \cdot q$$

$$\mathbb{P}\left(D(n) = \begin{pmatrix} -1\\-2 \end{pmatrix} \middle| \mathcal{F}(n)\right) = (1 - p_{se}(\alpha, n)) \cdot (1-q).$$
(24)

2.3 Color Majority as a Pólya Urn with randomized replacement

One can also imagine the above process as coming from a *Pólya Urn* with randomized replacement: At each timestep, we draw a ball of one color from the urn and add a new ball of the same color with probability 1 - q or one of the opposite color with probability q. To formalize this, we use the notation from [23] which is also used by e. g. [2, 13]. Generally, an *m*-type Pólya Urn process is given by

$$X(n) = \left((X_{i,n})_{i=1}^m \right)_{n \in \mathbb{N}},$$

where $X_{i,n}$ is the random variable describing the amount of balls of type *i* in the urn at time *n*. The evolution of the Pólya process is given by the *replacement vectors* ξ_j - if a ball of type *j* is drawn at time *n*, then

$$X_{i,n+1} = X_{i,n} + \xi_{j,i}.$$

Again, we need to represent both the amount of vertices of each color and the respective attachment weights to have the full picture. In the Pólya Urn model we achieve this by associating two types to each color, a *weight type* and a *count type*. The weight types $\{r_w, b_w\}$ should fulfill that the total amount of balls of one type is proportional to the attachment probability of the represented color, while the count types $\{r_c, b_c\}$ count the red, respectively blue, vertices. As [13], we set the activities of these types to $a_{r_w} = a_{b_w} = 1$, $a_{r_c} = a_{b_c} = 0$ and number them $r_w = 1$, $b_w = 2$, $r_c = 3$, $b_c = 4$. As is well known, the *(expected) replacement matrix A* given by

$$A \coloneqq \left(a_j \mathbb{E}[\xi_{j,i}]\right)_{i,j}$$

is quite important for the analysis of the Pólya Urn process. Note that we put the expected replacement vectors in the columns of the matrix as in [23]. For very simple increasing trees, this associated Pólya Urn has the following expected replacement matrix [2, 13]:

$$A_{\rm vs} = \begin{pmatrix} \alpha + 1 - q & q & 0 & 0 \\ q & \alpha + 1 - q & 0 & 0 \\ 1 - q & q & 0 & 0 \\ q & 1 - q & 0 & 0 \end{pmatrix}$$
(25)

and initial vector X(0) = (1, 0, 1, 0). For shape exchangeable trees, we follow the same modelling idea, but with a slightly different replacement rule: After time 1, each new vertex starts with attachment weight $\alpha + 1$ instead of 1, which changes the expected replacement matrix to

$$A_{\rm se} = \begin{pmatrix} \alpha + (1-q)(\alpha+1) & q(\alpha+1) & 0 & 0\\ q(\alpha+1) & \alpha + (1-q)(\alpha+1) & 0 & 0\\ 1-q & q & 0 & 0\\ q & 1-q & 0 & 0. \end{pmatrix}$$
(26)

The initial vector X(0) = (1, 0, 1, 0) is the same since the root has degree 0 at time 1. Then, for both tree models,

$$\Delta(n) = \begin{pmatrix} X_{3,n} - X_{4,n} \\ (X_{1,n} - X_{3,n}) - (X_{2,n} - X_{4,n}) \end{pmatrix}.$$
(27)

Theorem 5 is proven in Section 3. The proof of Theorem 6 is given in Section 4.

3 Proving Theorem **5**

To apply the convergence results from [23, Thms. 3.22-3.24] (see also [13, Thm. 3.1]), we need to check that the expected replacement matrices fulfill the necessary assumptions [13, (A1)-(A8)]:

Lemma 10. The Pólya Urns described in Subsection 2.3 with expected replacement matrices given in Eqs. (25) and (26) exhibit the following convergence behavior: Let $\lambda_1 > \lambda_2$ be the first two eigenvalues. Then,

$$\frac{X(N) - N\lambda_1 v_1}{\sqrt{N\ln(N)}} \xrightarrow{d} \mathcal{N}(0, \Sigma_I)$$
(28)

2. and if
$$\lambda_1 > 2\lambda_2$$
:

$$\frac{X(N) - N\lambda_1 v_1}{\sqrt{N}} \xrightarrow{d} \mathcal{N}(0, \Sigma_{II})$$
⁽²⁹⁾

with $\Sigma_{I,II}$ as defined in [23] or [13, Section 3].

Proof. If $q \neq \frac{\alpha+1}{2}$, A_{vs} is diagonalizable with eigenvalues $\Lambda = \{\alpha + 1, \alpha + 1 - 2q, 0, 0\}$. If $q = \frac{\alpha+1}{2}$, A_{vs} has eigenvalues $\{\alpha + 1, 0, 0\}$. Similarly, A_{se} is diagonalizable for $q \neq \frac{2\alpha+1}{2(\alpha+1)}$ with eigenvalues $\Lambda = \{2\alpha + 1, 2\alpha + 1 - 2q - 2\alpha q, 0, 0\}$. If $q = \frac{2\alpha+1}{2(\alpha+1)}$, then $\Lambda = \{\frac{2\alpha^2+3\alpha+1}{\alpha+1}, 0, 0\}$. One easily checks that the remaining conditions hold for both matrices (see also [13]).

As we can see in Lemma 10, the ratio between λ_1 and λ_2 is essential in determining the convergence behavior. Note that

$$\lambda_1 \ge 2\lambda_2 \iff q \ge f(\alpha) = \begin{cases} \frac{\alpha+1}{4} & \text{for very simple inc. trees} \\ \frac{2\alpha+1}{4(\alpha+1)} & \text{for shape ex. trees.} \end{cases}$$
(30)

Additionally, we remark that whether the matrices are diagonalizable or not, the first right eigenvector, v_1 , always fulfills $v_{1,3} = v_{1,4} = 1$. With this, Theorem 5 follows directly.

Proof of Theorem 5. For $q \ge f(\alpha)$ let

$$g(N) = \begin{cases} \sqrt{N} & \text{if } \lambda_1 = 2\lambda_2\\ \sqrt{N\ln(N)} & \text{if } \lambda_1 > 2\lambda_2 \end{cases}$$
(31)

and define

$$\widetilde{X}_{3,N} := \frac{X_{3,N} - N\lambda_1}{g(N)}, \quad \widetilde{X}_{4,N} := \frac{X_{4,N} - N\lambda_1}{g(N)}, \tag{32}$$

Then $(X(N)-N\lambda_1v_1)/g(N)$ converges jointly to a normal distribution, implying (since $v_{1,3} = 1 = v_{1,4}$ as mentioned above)

$$\begin{pmatrix} \widetilde{X}_{3,N} \\ \widetilde{X}_{4,N} \end{pmatrix} \xrightarrow{\mathbf{d}} \begin{pmatrix} \widetilde{X}_3 \\ \widetilde{X}_4 \end{pmatrix} \sim \mathcal{N}(0, \Sigma')$$
(33)

where calculating the covariance matrices $\Sigma_{I,II}$ gives

$$\Sigma' = \sigma(\alpha, q) \begin{pmatrix} 1 & -1 \\ -1 & 1 \end{pmatrix}$$
(34)

for both tree models. With this covariance structure $\widetilde{X}_3 - \widetilde{X}_4$ is also normal-distributed with mean 0. Analogous to Lemma 7, it holds that $\liminf_{N\to\infty} R_{\text{maj}}(N,q) \ge \liminf_{N\to\infty} \mathbb{P}(\Delta(N) < 0)$, giving [2]

$$\liminf_{N \to \infty} R_{\text{maj}}(N, q) \ge \liminf_{N \to \infty} \mathbb{P}(\Delta(N) < 0)$$
$$= \liminf_{N \to \infty} \mathbb{P}\left(\widetilde{X}_{3,N} - \widetilde{X}_{4,N} < 0\right)$$
$$= \mathbb{P}\left(\widetilde{X}_3 - \widetilde{X}_4 \le 0\right) = \frac{1}{2}$$
(35)

and similarly

$$\limsup_{N \to \infty} R_{\text{maj}}(N, q) = R_{\text{maj}}(q) \le \limsup_{n \to \infty} \mathbb{P}(\Delta(N) \le 0) = \frac{1}{2}.$$
(36)

Therefore, $R_{\text{maj}}(q) = \frac{1}{2}$ for $q \ge f(\alpha)$ follows.

4 **Proving Theorem 6**

The proof of Theorem 6 uses the random walk model presented in Subsection 2.2. We recall Lemma 7 and investigate the event $\{\Delta_1(N) > 0\}$.

Definition 11. We define the following stopping times for any A > B > 1:

$$\tau_{\text{good}}(A) \coloneqq \inf \left\{ n > 0 \mid \Delta_1(n) + \alpha \Delta_2(n) > A Z_\alpha(n) \sqrt{n} \right\}$$

$$\tau_{\text{bad}}(B) \coloneqq \inf \left\{ n > \tau_{\text{good}} \mid \Delta_1(n) + \alpha \Delta_2(n) \le B Z_\alpha(n) \sqrt{n} \right\}.$$
(37)

With this notation, there exists $N_0 > 0$ such that for all $N > N_0$

$$\mathbb{P}(\Delta_1(N) > 0) \ge \underbrace{\mathbb{P}(\Delta_1(N) > 0 \mid \tau_{\text{good}}(A) \le N, \tau_{\text{bad}}(B) > N)}_{(1)} \underbrace{\mathbb{P}(\tau_{\text{good}}(A) \le N, \tau_{\text{bad}}(B) > N)}_{(2)}.$$
(38)

Lemma 12. For all $q \in [0, 1]$ and each allowed value of α , there exists $c_{\alpha,1} > 0$ such that

$$\liminf_{N \to \infty} (1) \ge 1 - c_{\alpha, 1} \sqrt{q}.$$
(39)



Figure 1. Illustrating the two events in Eq. (38). In (a), the combined process $\Delta_1(n) + \alpha \Delta_2(n)$ passes the $\tau_{\text{good}}(A)$ boundary before the timehorizon N and does not drop below the $\tau_{\text{bad}}(B)$ -boundary again. In (b), we have the same behavior as in (a) and additionally see that the isolated process $\Delta_1(n)$ is above zero at time N.

Lemma 13. For all $q \in [0, 1]$ and each allowed value of α , there exists $c_{\alpha,2} > 0$ such that

$$\liminf_{N \to \infty} \left(2 \right) \ge 1 - c_{\alpha,2} \sqrt{q}. \tag{40}$$

Together, Lemmata 12 and 13 imply our theorem:

Proof of Theorem 6.

$$\liminf_{N \to \infty} \mathbb{P}(\Delta_1(N) > 0) \ge \liminf_{N \to \infty} \left((1 \cdot (2)) \right)$$
$$\ge (1 - c_{\alpha,1}\sqrt{q})(1 - c_{\alpha,2}\sqrt{q})$$
$$\ge 1 - c_{\alpha}\sqrt{q}. \tag{41}$$

With $1 - R_{\text{maj}}(q) \ge \liminf_{N \to \infty} \mathbb{P}(\Delta_1(N) > 0)$, the claim follows.

In the following subsections, we first prove Lemma 13 and then Lemma 12. For this, we set the parameters A and B to fixed values.

Definition 14. For $\tilde{c}_{\alpha} > 0$ set

$$B \coloneqq \frac{1}{Z_{\alpha}(n)} \sqrt{\left(3M_2 + \tilde{c}_{\alpha}\right) / \frac{\alpha(1 + \frac{1}{n}) + \frac{1}{2}}{Z_{\alpha}(n)}} \vee 1.$$

$$\tag{42}$$

and for $\gamma \in (0, \frac{1}{2})$ set

$$A \coloneqq \frac{1}{q^{\frac{1}{2} - \gamma}}.\tag{43}$$

Remark 15. With

$$\lim_{N \to \infty} Z_{\alpha}(N) \eqqcolon Z_{\alpha} = \begin{cases} \alpha + 1 & \text{for simple inc. trees} \\ 2\alpha + 1 & \text{for shape ex. trees.} \end{cases},$$
(44)

it holds that

$$\lim_{N \to \infty} B = \frac{1}{Z_{\alpha}} \sqrt{\left(3M_2 + \tilde{c}_{\alpha}\right) / \frac{\alpha + \frac{1}{2}}{Z_{\alpha}}} \vee 1.$$
(45)

4.1 Proving Lemma 13

To prove Lemma 13, we adopt a line of argumentation presented by Menshikov and Volkov [32] and consider the auxilliary process

$$Y(n) \coloneqq \frac{n}{(\Delta_1(n) + \alpha \Delta_2(n))^2},\tag{46}$$

which is adapted to $\mathcal{F}(n)$, the filtration generated by $\Delta(n)$.

Definition 16. Let

$$M_2 \coloneqq \max_{\omega} \left(D_1(n) + \alpha D_2(n) \right)^2(\omega). \tag{47}$$

Lemma 17. For any $\gamma \in (0, \frac{1}{2})$ there exists a threshold q_0 such that for all $q < q_0$: A > B > 1 and the stopped process $Y(\tau_{good}(A) \lor n \land \tau_{bad}(B))$ is a nonnegative supermartingale on $\mathcal{F}(n)$.

Proof. As we only consider $\tau_{\text{bad}}(B) \ge n \ge \tau_{\text{good}}(A)$, $\Delta_1(n) + \alpha \Delta_2(n)$ is nonzero and Y(n) is well-defined. If $n \land \tau_{\text{bad}}(B) = \tau_{\text{bad}}(B)$,

$$Y((n+1) \wedge \tau_{bad}(B)) - Y(n \wedge \tau_{bad}(B)) = Y(\tau_{bad}(B)) - Y(\tau_{bad}(B)) = 0$$
(48)

and for $n \wedge \tau_{\text{bad}}(B) = n$,

$$Y((n+1) \wedge \tau_{\text{bad}}(B)) - Y(n \wedge \tau_{\text{bad}}(B)) = Y(n+1) - Y(n)$$

$$= \frac{n+1}{(\Delta_1(n+1) + \alpha \Delta_2(n+1))^2} - \frac{n}{(\Delta_1(n) + \alpha \Delta_2(n))^2}$$

$$= \frac{n+1}{(\Delta_1(n) + D_1(n) + \alpha \Delta_2(n) + \alpha D_2(n))^2} - \frac{n}{(\Delta_1(n) + \alpha \Delta_2(n))^2}$$

$$= \frac{1}{(\Delta_1(n) + \alpha \Delta_2(n))^2} \left(\frac{n+1}{(1 + \frac{D_1(n) + \alpha D_2(n)}{\Delta_1(n) + \alpha \Delta_2(n)})^2} - n \right)$$

$$= \frac{n+1}{(\Delta_1(n) + \alpha \Delta_2(n))^2} \left(\frac{1}{(1 + \frac{D_1(n) + D_2(n)}{\Delta_1(n) + \alpha \Delta_2(n)})^2} - \frac{n}{n+1} \right). \quad (49)$$

Taking the conditional expectation with respect to $\mathcal{F}(n)$, we have

$$\mathbb{E}[Y(n+1) - Y(n) \mid \mathcal{F}(n)] = \frac{n+1}{(\Delta_1(n) + \alpha \Delta_2(n))^2} \mathbb{E}\left[\frac{1}{(1 + \frac{D_1(n) + D_2(n)}{\Delta_1(n) + \alpha \Delta_2(n)})^2} - \frac{n}{n+1} \mid \mathcal{F}(n)\right].$$
 (50)

Note that the first factor is always positive and we can therefore focus solely on the second factor. Set

$$f(x) \coloneqq \frac{1}{(1+x)^2} - \frac{n}{n+1} \text{ for } x > -1.$$
(51)

By a second order Taylor expansion around zero,

$$f(x) = 1 - \frac{n}{n+1} - 2x + 3x^2 + R_2 f(x;0),$$
(52)

where $R_2 f(x; 0)$ is the remainder term, which we bound uniformly in x using its Lagrangian form. If x > 0, there exists $\xi \in [0, x]$ such that

$$R_2 f(x;0) = -24 \underbrace{(1+\xi)^{-5}}_{\geq 0} \underbrace{x^3}_{\geq 0} \le 0.$$
(53)

If x < 0, there exists $\xi \in [x, 0] \subset (-1, 0]$ such that

$$R_2 f(x;0) = -24(1+\xi)^{-5} x^3.$$
(54)

Setting $x = \frac{D_1(n) + \alpha D_2(n)}{\Delta_1(n) + \alpha \Delta_2(n)}$ for $\tau_{\text{good}}(A) \le n \le \tau_{\text{bad}}(B)$, there exists $\tilde{c}_{\alpha} > 0$ such that (54) is bounded from above by

$$-24(1+\xi)^{-5} \left(\frac{-M_2}{\Delta_1(n) + \alpha \Delta_2(n)}\right)^3 = \tilde{c}_\alpha \left(\frac{1}{\Delta_1(n) + \alpha \Delta_2(n)}\right)^3$$
$$\leq \tilde{c}_\alpha \left(\frac{1}{\Delta_1(n) + \alpha \Delta_2(n)}\right)^2, \tag{55}$$

where the last inequality follows because $\tau_{\text{good}}(A) \leq n \leq \tau_{\text{bad}}(B)$ and therefore $\Delta_1(n) + \alpha \Delta_2(n) > 0$ almost surely. And so, uniformly over all n between $\tau_{\text{good}}(A)$ and $\tau_{\text{bad}}(B)$ and all realisations of $\Delta(n)$ (a.s.),

$$f\left(\frac{D_1(n) + \alpha D_2(n)}{\Delta_1(n) + \alpha \Delta_2(n)}\right)$$

$$\leq 1 - \frac{n}{n+1} - 2\frac{D_1(n) + \alpha D_2(n)}{\Delta_1(n) + \alpha \Delta_2(n)} + 3\left(\frac{D_1(n) + \alpha D_2(n)}{\Delta_1(n) + \alpha \Delta_2(n)}\right)^2 + \tilde{c}_{\alpha}\left(\frac{1}{\Delta_1(n) + \alpha \Delta_2(n)}\right)^2.$$
(56)

Hence, the conditional expectation on the righthandside of (50) is bounded from above by

$$\mathbb{E}\left[\frac{1}{n+1} - 2\frac{D_{1}(n) + \alpha D_{2}(n)}{\Delta_{1}(n) + \alpha \Delta_{2}(n)} + 3\left(\frac{D_{1}(n) + \alpha D_{2}(n)}{\Delta_{1}(n) + \alpha \Delta_{2}(n)}\right)^{2} + \tilde{c}_{\alpha}\left(\frac{1}{\Delta_{1}(n) + \alpha \Delta_{2}(n)}\right)^{2} \middle| \mathcal{F}(n)\right] \\
= \frac{1}{n+1} - \frac{2\mathbb{E}[D_{1}(n) + \alpha D_{2}(n) | \mathcal{F}(n)]}{\Delta_{1}(n) + \alpha \Delta_{2}(n)} + \frac{3\mathbb{E}[(D_{1}(n) + \alpha D_{2}(n))^{2} | \mathcal{F}(n)]}{(\Delta_{1}(n) + \alpha \Delta_{2}(n))^{2}} + \tilde{c}_{\alpha}\left(\frac{1}{\Delta_{1}(n) + \alpha \Delta_{2}(n)}\right)^{2}.$$
(57)

The first moment is

$$\mathbb{E}[D_1(n) + \alpha D_2(n) \mid \mathcal{F}(n)] = \frac{\rho}{Z_\alpha(n)} \frac{1}{n} (\Delta_1(n) + \alpha \Delta_2(n)),$$
(58)

with ρ defined as

$$\rho \coloneqq \begin{cases} \alpha + 1 - 2q & \text{for very simple increasing trees} \\ 2\alpha + 1 - 2q(\alpha + 1) & \text{for shape exchangeable trees} \end{cases}$$
(59)

and the second moment is bounded from above by M_2 . Therefore, Eq. (57) is bounded from above by

$$\frac{1}{n} - 2\frac{\rho}{Z_{\alpha}(n)}\frac{1}{n} + \frac{3M_2 + \tilde{c}_{\alpha}}{(\Delta_1(n) + \alpha\Delta_2(n))^2} = \frac{1}{n} \left(1 - 2\frac{\rho}{Z_{\alpha}(n)} + (3M_2 + \tilde{c}_{\alpha})Y(n)\right),\tag{60}$$

which gives

$$Y(n) \le -\left(1 - 2\frac{\rho}{Z_{\alpha}(n)}\right) / (3M_2 + \tilde{c}_{\alpha}) \implies \mathbb{E}[Y(n+1) - Y(n) \mid \mathcal{F}(n)] \le 0.$$
(61)

It holds that

$$-\left(1-2\frac{\rho}{Z_{\alpha}(n)}\right) > 0 \iff q < \begin{cases} \frac{\alpha(1+\frac{1}{n})+1}{4} & \text{for v.s.i. trees} \\ \frac{2\alpha(1+\frac{1}{n})+1}{4(\alpha+1)} & \text{for s.e. trees,} \end{cases}$$
(62)

meaning that the numerator is always positive for small enough values of q in both groups of models - this is important since Y(n) is a positive process. Additionally, because B only depends on α and A increases when q decreases, A > B > 1 for small enough values of q. Take q small enough to fulfill both these constraints. For n between $\tau_{good}(A)$ and $\tau_{bad}(B)$ it holds that

$$\Delta_1(n) + \alpha \Delta_2(n) > BZ_\alpha(n)\sqrt{n},\tag{63}$$

implying

$$Y(n) \leq \frac{n}{\left(BZ_{\alpha}(n)\sqrt{n}\right)^{2}}$$

$$= \frac{1}{\left(BZ_{\alpha}(n)\right)^{2}}$$

$$\leq \frac{\alpha(1+\frac{1}{n})+\frac{1}{2}}{Z_{\alpha}(n)} / (3M_{2}+\tilde{c}_{\alpha}).$$
(64)

Further, for $q < \frac{1}{8}$ it holds in very simple increasing trees,

$$\frac{\alpha(1+\frac{1}{n})+\frac{1}{2}}{Z_{\alpha}(n)} \leq \frac{\alpha(1+\frac{1}{n})+1-4q}{Z_{\alpha}(n)}$$
$$= -\left(1-2\frac{\alpha+1-2q}{Z_{\alpha}(n)}\right)$$
$$= -\left(1-2\frac{\rho}{Z_{\alpha}(n)}\right)$$
(65)

and in shape exchangeable trees, for $q < \frac{2\alpha(1+\frac{1}{n})+1}{8(\alpha+1)},$

$$\frac{\alpha(1+\frac{1}{n})+\frac{1}{2}}{Z_{\alpha}(n)} \leq \frac{2\alpha(1+\frac{1}{n}-2q)+1-4q}{Z_{\alpha}(n)} \\ = -\left(1-2\frac{2\alpha+1-2q(\alpha+1)}{Z_{\alpha}(n)}\right) \\ = -\left(1-2\frac{\rho}{Z_{\alpha}(n)}\right).$$
(66)

Therefore there exists a q_0 such that for all $q < q_0$, A > B > 1 and $Y(\tau_{good}(A) \lor n \land \tau_{bad}(B))$ is a supermartingale. This finishes the proof of Lemma 17.

Proof of Lemma 13. To prove the lower bound for

$$(2) \coloneqq \mathbb{P}(\tau_{\text{good}}(A) \le N, \tau_{\text{bad}}(B) > N),$$

we write

$$\mathbb{P}(\tau_{\text{good}}(A) \le N, \tau_{\text{bad}}(B) > N) = 1 - \mathbb{P}(\tau_{\text{good}}(A) > N \text{ or } \tau_{\text{bad}}(B) \le N)$$
$$= 1 - \mathbb{P}(\tau_{\text{good}}(A) > N) - \mathbb{P}(\tau_{\text{bad}}(B) \le N).$$
(67)

For $\mathbb{P}(\tau_{\text{bad}}(B) \leq N)$, we use $Y(\tau_{\text{good}}(A) \vee n \wedge \tau_{\text{bad}}(B))$, where we omit the parameters A and B in the notation whereever they are not relevant. Remember that we consider Y(n) as a process adapted to $\mathcal{F}(n)$, the natural filtration of $\Delta(n)$. Note first that

$$Y(\tau_{\text{good}}) \le \frac{\tau_{\text{good}}}{(AZ_{\alpha}(\tau_{\text{good}})\sqrt{\tau_{\text{good}}})^2} = \frac{1}{A^2 Z_{\alpha}(\tau_{\text{good}})^2} \le \frac{1}{A^2},\tag{68}$$

$$Y(\tau_{\text{bad}}) \ge \frac{\tau_{\text{bad}}}{(BZ_{\alpha}(\tau_{\text{bad}})\sqrt{\tau_{\text{bad}}})^2} = \frac{1}{B^2 Z_{\alpha}(\tau_{\text{bad}})^2} \ge \begin{cases} \frac{1}{B^2(\alpha+2)^2} & \text{for simple inc. trees} \\ \frac{1}{B^2(2\alpha+2)^2} & \text{for shape ex. trees} \end{cases}$$
(69)

and note additionally that

$$\mathbb{P}(\tau_{\text{bad}} \le N) = \mathbb{E}[\mathbb{1}_{N \land \tau_{\text{bad}} = \tau_{\text{bad}}}] = \mathbb{E}\left[\mathbb{E}[\mathbb{1}_{N \land \tau_{\text{bad}} = \tau_{\text{bad}}} | \mathcal{F}(\tau_{\text{good}})]\right].$$
(70)

On the event $\tau_{\text{good}} \leq N$, we get by a variant of the optimal stopping theorem [33, Theorem 28, Chapter V] $\mathbb{E}[Y(N \wedge \tau_{\text{bad}}) \mid \mathcal{F}(\tau_{\text{good}})] \leq Y(\tau_{\text{good}}). \tag{71}$

Further, since $Y(\tau_{\text{good}} \lor n \land \tau_{\text{bad}})$ is always positive,

$$\mathbb{E}[Y(N \wedge \tau_{\text{bad}}) | \mathcal{F}(\tau_{\text{good}})] = \mathbb{E}[Y(N) \mathbb{1}_{N \wedge \tau_{\text{bad}} = N} + Y(\tau_{\text{bad}}) \mathbb{1}_{N \wedge \tau_{\text{bad}} = \tau_{\text{bad}}} | \mathcal{F}(\tau_{\text{good}})] \\ \geq \mathbb{E}[Y(\tau_{\text{bad}}) \mathbb{1}_{N \wedge \tau_{\text{bad}} = \tau_{\text{bad}}} | \mathcal{F}(\tau_{\text{good}})] \\ \geq \begin{cases} \frac{1}{B^2(\alpha+2)^2} \mathbb{E}[\mathbb{1}_{N \wedge \tau_{\text{bad}} = \tau_{\text{bad}}} | \mathcal{F}(\tau_{\text{good}})] & \text{for simple inc. trees} \\ \frac{1}{B^2(2\alpha+2)^2} \mathbb{E}[\mathbb{1}_{N \wedge \tau_{\text{bad}} = \tau_{\text{bad}}} | \mathcal{F}(\tau_{\text{good}})] & \text{for shape ex. trees.} \end{cases}$$
(72)

Therefore,

$$\frac{1}{A^2} \ge \mathbb{E}[Y(\tau_{\text{good}})] \ge \mathbb{E}\left[\mathbb{E}\left[\mathbbm{1}_{N \land \tau_{\text{bad}} = \tau_{\text{bad}}} \mid \mathcal{F}(\tau_{\text{good}})\right]\right] \cdot \begin{cases} \frac{1}{B^2(\alpha+2)^2} & \text{for simple inc. trees} \\ \frac{1}{B^2(2\alpha+2)^2} & \text{for shape ex. trees} \end{cases}$$

$$\iff \frac{1}{A^2} \ge \mathbb{P}(\tau_{\text{bad}} \le N) \cdot \begin{cases} \frac{1}{B^2(\alpha+2)^2} & \text{for simple inc. trees} \\ \frac{1}{B^2(2\alpha+2)^2} & \text{for shape ex. trees,} \end{cases}$$
(73)

which by definition of A and B implies

$$\mathbb{P}(\tau_{\text{bad}} \le N) \le q^{1-2\gamma} \cdot \begin{cases} B^2(\alpha+2)^2 & \text{for simple inc. trees} \\ B^2(2\alpha+2)^2 & \text{for shape ex. trees.} \end{cases}$$
(74)

To bound $\mathbb{P}(\tau_{\text{good}}(A) \leq N)$, we consider the first point at which $\Delta_1(n) + \alpha \Delta_2(n)$ may reach this boundary. For it to happen at time n_0 , $\Delta_1(n_0) = n_0$, $\Delta_2(n_0) = n_0 - 1$ must hold, implying $\Delta_1(n_0) + \alpha \Delta_2(n_0) = Z_\alpha(n)n_0$. For both model groups,

$$Z_{\alpha}(n_0)n_0 > AZ_{\alpha}(n_0)\sqrt{n_0} \iff \sqrt{n_0} > A, \tag{75}$$

which is fulfilled for $n_0 > A^2$. Note that, since $\Delta_1(1) = 1$ and, by definition of D(n), $\Delta_2(2) = 1$ a.s., it holds that

$$\mathbb{P}(\Delta_1(n_0) + \alpha \Delta_2(n_0) = Z_\alpha(n_0)n_0) = (1-q)^{n_0-1}.$$
(76)

Therefore, the probability that the process will not reach the $\tau_{good}(A)$ -boundary before time N is bounded from above by

$$\mathbb{P}(\tau_{\text{good}}(A) > N) \le 1 - (1-q)^{\lceil A^2 \rceil - 1} \le (A^2 + 1)q - q = q^{2\gamma}.$$
(77)

Plugging Eqs. (74) and (77) into Eq. (67) gives

$$(2) \ge \begin{cases} 1 - q^{2\gamma} + q - B^2 (\alpha + 2)^2 q^{1-2\gamma} & \text{for very simple inc. trees} \\ 1 - q^{2\gamma} + q - B^2 (2\alpha + 2)^2 q^{1-2\gamma} & \text{for shape ex. trees.} \end{cases}$$
(78)

With Remark 15,

$$\liminf_{N \to \infty} \left(2 \right) \ge 1 - c_{\alpha,2} \sqrt{q} \tag{79}$$

follows.

4.2 Proving Lemma 12

Recall

$$(1) \coloneqq \mathbb{P}(\Delta_1(N) > 0 \mid \tau_{\text{good}} \le N, \tau_{\text{bad}} > N).$$
(80)

Proof of Lemma 12 for $\alpha = 0$ *.*

For $\alpha = 0$, it holds that $\tau_{\text{good}}(A) = \inf\{n > 0 \mid \Delta_1(n) > A\sqrt{n}\}$ and $\tau_{\text{bad}}(B) = \inf\{n > \tau_{\text{good}} \mid \Delta_1(n) \le B\sqrt{n}\}$ (since $Z_{\alpha=0}(n) \equiv 1$) and therefore

$$\{\tau_{\text{good}} \le N, \tau_{\text{bad}} > N\} \subset \{\Delta_1(N) > 0\},\tag{81}$$

which implies $1 \ge 1 - c_{1,\alpha=0}\sqrt{q}$ for any $c_{1,\alpha=0} > 0$.

For $\alpha \neq 0$, Eq. (81) does not hold. The core idea of the following argument is that with high enough probability, Δ_1 and Δ_2 will not stray too far from each other.

Definition 18. Let $[N] := \{1, 2, ..., N\}$. We define the following random variables representing the decisions made by $\Delta(n)$ up to time N.

- (r, r)([N]), (r, b)([N]), (b, r)([N]), (b, b)([N]): The first entry in the tuple represents the color of the attached-to vertex and the second entry the color of the attaching vertex. Each random variable takes on values in $\mathcal{P}(\{2, ..., N\})$ such that, e. g., iff $k \in (b, r)[N]$), a new red vertex attached to an existing blue vertex at the transition from time k - 1 to k.
- $(r, _)([N]), (_, r)([N]), (b, _)([N]), (_, b)([N])$: These random variables are defined as (disjoint) unions of the random variables defined above. For example, $(r, _)([N]) = (r, r)([N]) \cup (r, b)([N])$.

Additionally, we define the event

$$\mathcal{A} \coloneqq \left\{ \#_{(r, \cdot)([N])} \in \left[\frac{\#_{(\cdot, r)([N])} - Nq - a\sqrt{N}}{1 - 2q}, \frac{\#_{(\cdot, r)([N])} - Nq + a\sqrt{N}}{1 - 2q} \right] \right\},\tag{82}$$

where $a = \frac{BZ_{\alpha}(N)}{4|\alpha|} > 0$ and we write $\#_A$ to denote the cardinality of a set A. Finally, we introduce the shorthand notation

$$\mathcal{C} = \{\tau_{\text{good}}(A) \le N, \tau_{\text{bad}}(B) > N\}.$$
(83)

With these definitions in place, we are ready to prove Lemma 12 for $\alpha \neq 0$ as well.

Proof of Lemma 12 for $\alpha \neq 0$. We show

$$\liminf_{N \to \infty} \mathbb{P}(\mathcal{A} \mid \mathcal{C}) \ge 1 - c_{\alpha, 1} \sqrt{q} \quad \text{and} \quad \mathbb{P}(\Delta_1(N) > 0 \mid \mathcal{A} \cap \mathcal{C}) = 1,$$
(84)

which implies the claim via

$$1 = \mathbb{P}(\Delta_1(N) > 0 \mid \mathcal{C}) \ge \mathbb{P}(\Delta_1(N) > 0 \mid \mathcal{A} \cap \mathcal{C})\mathbb{P}(\mathcal{A} \mid \mathcal{C}).$$
(85)

It holds that

$$\Delta_{1}(N) = \#_{(.,r)([N])} - \#_{(.,b)([N])}$$

= $\#_{(.,r)([N])} - (N - \#_{(.,r)([N])})$
= $2\#_{(.,r)([N])} - N.$ (86)

Further, in the very simple increasing tree case,

$$\Delta_{2}(N) = \#_{(r,-)([N])} - \#_{(b,-)([N])}$$

= $\#_{(r,-)([N])} - (N - \#_{(r,-)([N])})$
= $2\#_{(r,-)([N])} - N.$ (87)

In shape exchangeable trees we get

$$\Delta_{2}(N) = 2\#_{(r,r)([N])} + 0\#_{(r,b)([N])} + 0\#_{(b,r)([N])} - 2\#_{(b,b)([N])}$$

= $2\#_{(r,r)([N])} + 0\#_{(r,b)([N])} + 0\#_{(b,r)([N])} - 2(N - \#_{(r,b)([N])} - \#_{(b,r)([N])} - \#_{(r,r)([N])})$
= $2\#_{(r,\cdot)([N])} + 2\#_{(-,r)([N])} - 2N.$ (88)

To lower-bound $\mathbb{P}(\mathcal{A} \mid \mathcal{C})$, note that $\mathbb{P}(\mathcal{A} \mid \mathcal{C}) \geq \mathbb{P}(\mathcal{A}) - \mathbb{P}(\mathcal{C}^c)$ and that given the color of the drawn vertex, the color of the new vertex is simply an independent coin flip. Let \mathbb{P}_i be the probability measure conditioned on $\#_{(,,r)([N])} = i$:

$$\mathbb{E}_i \left[\#_{(,r)([N])} \right] = (1-q)i + q(N-i) = (1-2q)i + Nq$$
(89)

and

$$\operatorname{Var}_{i}\left[\#_{(,r)([N])}\right] = (1-q)qi + (1-q)q(N-i) \le qi + q(N-i) = qN.$$
(90)

By definition of \mathcal{A} ,

$$\mathbb{P}_{i}(\mathcal{A}^{c}) = \mathbb{P}_{i}\Big(\big|\#_{(,r)([N])} - \mathbb{E}_{i}\big[\#_{(,r)([N])}\big]\big| > a\sqrt{N}\Big).$$

$$\tag{91}$$

With this, it holds since a > 0

$$\mathbb{P}_{i}\Big(\big|\#_{(-,r)([N])} - \mathbb{E}_{i}\big[\#_{(-,r)([N])}\big]\big| > a\sqrt{N}\Big) \le \frac{qN}{a^{2}N} = \frac{1}{a^{2}}q$$
(92)

and thereby

$$\mathbb{P}_{i}(\mathcal{A}) \ge 1 - \left(\frac{4|\alpha|}{BZ_{\alpha}(N)}\right)^{2} q \tag{93}$$

uniformly in i (since the righthandside does not depend on q), bringing us together with Remark 15 to

$$\liminf_{N \to \infty} \mathbb{P}(\mathcal{A} \mid \mathcal{C}) \ge 1 - c_{\alpha, 1} \sqrt{q}.$$
(94)

It remains to prove

$$\mathbb{P}(\Delta_1(N) > 0 \mid \mathcal{A} \cap \mathcal{C}) = 1.$$
(95)

Note that $\{\Delta_1(N) > 0\} = \{\#_{(,r)([N])} > \frac{N}{2}\}$ and let $\omega \in \mathcal{A} \cap \mathcal{C}$. By Definition 18 we have

$$\frac{\#_{(r,.)([N])}(\omega) - Nq - a\sqrt{N}}{1 - 2q} \le \#_{(r,.)([N])}(\omega) \le \frac{\#_{(r,.)([N])}(\omega) - Nq + a\sqrt{N}}{1 - 2q}$$
(96)

and

$$\Delta_1(N)(\omega) + \alpha \Delta_2(N)(\omega) > BZ_\alpha(N)\sqrt{N}$$
(97)

(we now omit (ω) for the sake of readability). Eqs. (86) and (97) together with Eq. (87) give for very simple increasing trees

$$0 < \Delta_1(N) + \alpha \Delta_2(N) - BZ_{\alpha}(N)\sqrt{N}$$

= $2(\#_{(-,r)([N])} + \alpha \#_{(r,-)([N])}) - (1+\alpha)N - BZ_{\alpha}(N)\sqrt{N}.$ (98)
For (88) to get

Replace Eq. (87) with Eq. (88) to get

$$0 < \Delta_1(N) + \alpha \Delta_2(N) - BZ_{\alpha}(N)\sqrt{N} = (2 + 2\alpha) \#_{(r,r)([N])} + 2\alpha \#_{(r,r)([N])} - (1 + 2\alpha)N - BZ_{\alpha}(N)\sqrt{N}$$
(99)

for shape exchangeable trees. If
$$\alpha > 0$$
, we apply the upper interval bound from Eq. (96), which together with Eq. (98) gives for very simple increasing trees:

$$0 < 2(\#_{(.,r)([N])} + \alpha \#_{(r,.)([N])}) - (1+\alpha)N - BZ_{\alpha}(N)\sqrt{N}$$

$$\leq 2\left(\#_{(.,r)([N])} + \alpha \frac{\#_{(.,r)([N])} - Nq + a\sqrt{N}}{1 - 2q}\right) - (1+\alpha)N - BZ_{\alpha}(N)\sqrt{N}$$

$$= 2\#_{(.,r)([N])}\left(1 + \alpha \frac{1}{1 - 2q}\right) - \left(1 + \alpha + 2\alpha \frac{q}{1 - 2q}\right)N - \left(BZ_{\alpha}(N) - \frac{2\alpha a}{1 - 2q}\right)\sqrt{N}.$$
(100)

and for shape exchangeable trees, we continue from Eq. (99):

$$0 < (2+2\alpha)\#_{(,,r)([N])} + 2\alpha\#_{(r,.)([N])} - (1+2\alpha)N - BZ_{\alpha}(N)\sqrt{N}$$

$$\leq (2+2\alpha)\#_{(,,r)([N])} + 2\alpha\left(\frac{\#_{(,,r)([N])} - Nq + a\sqrt{N}}{1-2q}\right) - (1+2\alpha)N - BZ_{\alpha}(N)\sqrt{N}$$

$$= 2\#_{(,,r)([N])}\left(1 + \alpha + \frac{\alpha}{1-2q}\right) - \left(1 + 2\alpha + \frac{2\alpha q}{1-2q}\right)N - \left(BZ_{\alpha}(N) - \frac{2\alpha a}{1-2q}\right)\sqrt{N}.$$
(101)

It holds that

$$a = \frac{BZ_{\alpha}(N)}{4\alpha} \stackrel{q < \frac{1}{2}}{\Longrightarrow} a < (1 - 2q) \frac{BZ_{\alpha}(N)}{2\alpha}$$
$$\stackrel{\alpha > 0}{\Longleftrightarrow} BZ_{\alpha}(N) > 2\alpha \frac{a}{1 - 2q}$$
$$\iff BZ_{\alpha}(N) - 2\alpha \frac{a}{1 - 2q} > 0.$$
(102)

In the very simple increasing tree case we continue from Eq. (100)

$$0 < 2\#_{(,,r)([N])} \left(1 + \alpha \frac{1}{1 - 2q}\right) - \left(1 + \alpha + 2\alpha \frac{q}{1 - 2q}\right)N - \left(BZ_{\alpha}(N) + 2\alpha \frac{a}{1 - 2q}\right)\sqrt{N}$$

$$< 2\#_{(,,r)([N])} \left(1 + \alpha \frac{1}{1 - 2q}\right) - \left(1 + \alpha + 2\alpha \frac{q}{1 - 2q}\right)N$$

$$= \left(1 + \frac{\alpha}{1 - 2q}\right)(2\#_{(,,r)([N])} - N)$$
(103)

and in shape exchangeable trees from Eq. (101)

$$0 < \left(2 + 2\alpha + \frac{2\alpha}{1 - 2q}\right) \#_{(.,r)([N])} - \left(1 + 2\alpha + \frac{2\alpha q}{1 - 2q}\right) N - \left(BZ_{\alpha}(N) - \frac{2\alpha a}{1 - 2q}\right) \sqrt{N} < \left(2 + 2\alpha + \frac{2\alpha}{1 - 2q}\right) \#_{(.,r)([N])} - \left(1 + 2\alpha + \frac{2\alpha q}{1 - 2q}\right) N = \left(2 + \frac{2\alpha - 2\alpha q}{1 - 2q}\right) (2\#_{(.,r)([N])} - N).$$
(104)

Both Eqs. (103) and (104) imply $\#_{(.,r)([N])} > \frac{N}{2}$ for $q < \frac{1}{2}$. For negative values of α , we use the lower interval bound from Eq. (96) which only changes the \sqrt{N} -term, giving us for very simple increasing trees

$$0 < 2(\#_{(,,r)([N])} + \alpha \#_{(r,.)([N])}) - (1+\alpha)N - BZ_{\alpha}(N)\sqrt{N}$$

$$\leq 2\#_{(,,r)([N])} \left(1 + \alpha \frac{1}{1-2q}\right) - \left(1 + \alpha + 2\alpha \frac{q}{1-2q}\right)N - \left(BZ_{\alpha}(N) + 2\alpha \frac{a}{1-2q}\right)\sqrt{N}$$
(105)

and for shape exchangeable trees

$$0 < (2+2\alpha) \#_{(,,r)([N])} + 2\alpha \#_{(r,-)([N])} - (1+2\alpha)N - BZ_{\alpha}(N)\sqrt{N}$$

$$\leq \left(2+2\alpha + \frac{2\alpha}{1-2q}\right) \#_{(,,r)([N])} - \left(1+2\alpha + \frac{2\alpha q}{1-2q}\right)N - \left(BZ_{\alpha}(N) + \frac{2\alpha a}{1-2q}\right)\sqrt{N}.$$
(106)

Analogous to above, the \sqrt{N} -term is positive by our choice of a, resulting in the following equations for very simple increasing trees

$$0 < 2(\#_{(.,r)([N])} + \alpha \#_{(r,.)([N])}) - (1+\alpha)N - BZ_{\alpha}(N)\sqrt{N} < 2\#_{(.,r)([N])} \left(1 + \alpha \frac{1}{1-2q}\right) - \left(1 + \alpha + 2\alpha \frac{q}{1-2q}\right)N$$
(107)

and shape exchangeable trees

$$0 < (2+2\alpha) \#_{(,,r)([N])} + 2\alpha \#_{(r,-)([N])} - (1+2\alpha)N - BZ_{\alpha}(N)\sqrt{N} < \left(2+2\alpha + \frac{2\alpha}{1-2q}\right) \#_{(,,r)([N])} - \left(1+2\alpha + \frac{2\alpha q}{1-2q}\right)N.$$
(108)

We see that these correspond to the $\alpha > 0$ case (cf. Eqs. (100) and (101) ff.), which finishes the proof.

References

- E. Abbe, E. Cornacchia, Y. Gu, and Y. Polyanskiy. "Stochastic block model entropy and broadcasting on trees with survey". In: *Conference on Learning Theory*. Ed. by M. Belkin and S. Kpotufe. Vol. 134. Proceedings of Machine Learning Research. PMLR, 2021, pp. 1–25.
- [2] L. Addario-Berry, L. Devroye, G. Lugosi, and V. Velona. "Broadcasting on random recursive trees". In: *The Annals of Applied Probability* 32.1 (2022), pp. 497–528.
- [3] S. Banerjee and S. Bhamidi. "Persistence of hubs in growing random networks". In: *Probab. Theory Related Fields* 180.3-4 (2021), pp. 891–953.
- [4] S. Banerjee and S. Bhamidi. "Root finding algorithms and persistence of Jordan centrality in growing random trees". In: *Ann. Appl. Probab.* 32.3 (2022), pp. 2180–2210.
- [5] P. M. Bleher, J. Ruiz, and V. A. Zagrebnov. "On the purity of the limiting Gibbs state for the Ising model on the Bethe lattice". In: *Journal of Statistical Physics* 79 (1995), pp. 473–482.
- [6] S. Briend, L. Devroye, and G. Lugosi. "Broadcasting in random recursive dags". In: *arXiv preprint* arXiv:2306.01727 (2023).
- [7] S. Bubeck, L. Devroye, and G. Lugosi. "Finding Adam in random growing trees". In: *Random Structures & Algorithms* 50.2 (2017), pp. 158–172.
- [8] S. Bubeck, R. Eldan, E. Mossel, and M. Z. Rácz. "From trees to seeds: on the inference of the seed from large trees in the uniform attachment model". In: *Bernoulli* 23.4A (2017), pp. 2887–2916.
- [9] S. Bubeck, E. Mossel, and M. Z. Rácz. "On the influence of the seed graph in the preferential attachment model". In: *IEEE Trans. Network Sci. Eng.* 2.1 (2015), pp. 30–39.
- [10] H. Crane and M. Xu. "Inference on the history of a randomly growing tree". In: J. R. Stat. Soc. Ser. B. Stat. Methodol. 83.4 (2021), pp. 639–668.
- [11] N. Curien, T. Duquesne, I. Kortchemski, and I. Manolescu. "Scaling limits and influence of the seed graph in preferential attachment trees". In: J. Éc. polytech. Math. 2 (2015), pp. 1–34.
- [12] N. Curien and L. Laulin. "Recurrence of the plane Elephant random walk". In: (2023). arXiv: 2310.10314 [math.PR].
- [13] C. Desmarais, C. Holmgren, and S. Wagner. "Broadcasting-induced colorings of preferential attachment trees". In: *Random Structures & Algorithms* (2023). eprint: https://onlinelibrary.wiley.com/doi/pdf/10. 1002/rsa.21142.
- [14] M. Drmota. Random trees: an interplay between combinatorics and probability. Springer Science & Business Media, 2009.
- [15] K. Durant and S. Wagner. "On the centroid of increasing trees". In: Discrete Mathematics & Theoretical Computer Science 21.4 (2019), Paper No. 8, 29.
- [16] C. Efthymiou. *Reconstruction/Non-reconstruction Thresholds for Colourings of General Galton-Watson Trees*. 2014.

- [17] F. Eggenberger and G. Pólya. "Über die Statistik verketteter Vorgänge". In: ZAMM-Journal of Applied Mathematics and Mechanics/Zeitschrift für Angewandte Mathematik und Mechanik 3.4 (1923), pp. 279–289.
- [18] W. Evans, C. Kenyon, Y. Peres, and L. J. Schulman. "Broadcasting on trees and the Ising model". In: Annals of Applied Probability (2000), pp. 410–433.
- [19] Y. Gu, H. Roozbehani, and Y. Polyanskiy. "Broadcasting on trees near criticality". In: 2020 IEEE International Symposium on Information Theory (ISIT). IEEE. 2020, pp. 1504–1509.
- [20] H. Guérin, L. Laulin, and K. Raschel. A fixed-point equation approach for the superdiffusive elephant random walk. 2024. arXiv: 2308.14630 [math.PR].
- [21] H. Huang and E. Mossel. Low Degree Hardness for Broadcasting on Trees. 2024. arXiv: 2402.13359 [math.PR].
- [22] T. E. Huillet. "On Bagchi–Pal urn models and related Pólya–Friedman ones". In: *Journal of Statistical Mechanics: Theory and Experiment* 2017.9 (2017), p. 093211.
- [23] S. Janson. "Functional limit theorems for multitype branching processes and generalized Pólya urns". In: *Stochastic Processes and their Applications* 110.2 (2004), pp. 177–245.
- [24] V. Jog and P.-L. Loh. "Analysis of centrality in sublinear preferential attachment trees via the Crump-Mode-Jagers branching process". In: *IEEE Trans. Network Sci. Eng.* 4.1 (2017), pp. 1–12.
- [25] J. Khim and P.-L. Loh. "Confidence Sets for the Source of a Diffusion in Regular Trees". In: *IEEE Transactions* on Network Science and Engineering 4.1 (Jan. 2017), pp. 27–40.
- [26] F. Koehler and E. Mossel. "Reconstruction on Trees and Low-Degree Polynomials". In: *Advances in Neural Information Processing Systems*. Ed. by A. H. Oh, A. Agarwal, D. Belgrave, and K. Cho. 2022.
- [27] L. Laulin. "About the elephant random walk". PhD thesis. Université de Bordeaux, 2022.
- [28] G. Lugosi and A. S. Pereira. "Finding the seed of uniform attachment trees". In: *Electronic Journal of Probability* 24.none (2019), pp. 1–15.
- [29] H. Mahmoud. Pólya urn models. CRC press, 2008.
- [30] A. Makur, E. Mossel, and Y. Polyanskiy. "Broadcasting on random directed acyclic graphs". In: *IEEE Trans. Inform. Theory* 66.2 (2020), pp. 780–812.
- [31] M. Menshikov, S. Popov, and A. Wade. *Non-homogeneous random walks: Lyapunov function methods for near-critical stochastic systems*. Vol. 209. Cambridge University Press, 2016.
- [32] M. Menshikov and S. Volkov. "Urn-related random walk with drift $\rho x^{\alpha}/t^{\beta}$ ". In: *Electron. J. Probab.* 13 (2008), no. 31, 944–960.
- [33] P. Meyer. *Probability and Potentials*. Blaisdell book in pure and applied mathematics. Blaisdell Publishing Company, 1966.
- [34] E. Mossel, J. Neeman, and A. Sly. "Belief propagation, robust reconstruction and optimal recovery of block models". In: *Proceedings of The 27th Conference on Learning Theory, COLT 2014, Barcelona, Spain, June* 13-15, 2014. Ed. by M. Balcan, V. Feldman, and C. Szepesvári. Vol. 35. JMLR Workshop and Conference Proceedings. JMLR.org, 2014, pp. 356–370.
- [35] E. Mossel, A. Sly, and Y. Sohn. "Exact Phase Transitions for Stochastic Block Models and Reconstruction on Trees". In: *Proceedings of the 55th Annual ACM Symposium on Theory of Computing*. STOC 2023. Orlando, FL, USA: Association for Computing Machinery, 2023, pp. 96–102.