RÉNYI DIVERGENCE GUARANTEES FOR HASHING WITH LINEAR CODES

MADHURA PATHEGAMA AND ALEXANDER BARG

ABSTRACT. We consider the problem of distilling uniform random bits from an unknown source with a given p-entropy using linear hashing. As our main result, we estimate the expected p-divergence from the uniform distribution over the ensemble of random linear codes for all integer $p \geq 2$. The proof relies on analyzing how additive noise, determined by a random element of the code from the ensemble, acts on the source distribution. This action leads to the transformation of the source distribution into an approximately uniform one, a process commonly referred to as distribution smoothing. We also show that hashing with Reed-Muller matrices reaches intrinsic randomness of memoryless Bernoulli sources in the l_p sense for all integer $p \geq 2$.

1. Introduction

Uniform random bit-strings are an essential resource in both computer science and cryptography. In computer science, numerous algorithms rely on randomization to efficiently solve problems. These algorithms encompass a wide range of applications, including primality testing [28, 33], finding minimum-cut in a graph [24] and random shuffling [15]. Moreover, uniform random bits are indispensable in many cryptographic applications. Examples include universal hash functions [8], pseudo-random generators [19], randomized encryption schemes[36], and randomized signature algorithms [45]. In essence, uniform random bits provide the necessary unpredictability and confidentiality required for robust cryptographic systems. They safeguard sensitive information and thwart attacks, making them an essential resource.

One way of distilling a uniform distribution is to use a weak random source, i.e., a source with low entropy, and convert its randomness to a uniform distribution. For a weak random source with a known distribution, the amount of uniform bits that can be distilled from the source is called the intrinsic randomness of the source [44] and it is possible to construct a deterministic function that transforms almost all the entropy of the source to uniform q-ary symbols. In many cryptographic problems, however, the distribution of the source remains unknown. Usually we assume that we know some quantitative measure of randomness, for instance, the entropy of the source. A (random) function that is capable of distilling uniform bits from an unknown source Z with a given (Rényi) entropy $H_p(Z)$ is called a randomness extractor [30]. If the inherent randomness of this function is nearly independent of the output, such a function is called a strong extractor in the literature. Note that almost always, it is impossible to convert all the randomness of a weak source into a perfect uniform distribution; what we usually obtain is an approximation. Proximity to uniformity can be measured in several ways. For instance, in many cryptographic applications, the distance between probability distributions is often measured using the total variation distance or KL divergence.

A classic result randomness extraction, known as the leftover hash lemma or LHL [23, 43], implies that universal hash functions are capable of converting almost all min-entropy of the source to a bit string whose distribution is close to uniform in the sense of total variation distance. The assumption of this lemma can be relaxed from the min-entropy to p-Rényi entropy in [6] and (for p = 2) in [3].

In some instances, uniformity guarantees of the total variation distance or KL divergence are insufficient for the applications [7, 25, 14, 41]. Motivated by these shortcomings, the authors of [21] obtained asymptotic

1

The authors are with the Department of ECE and Institute for Systems Research, University of Maryland, College Park, MD 20817. Emails: {madhura,abarg}@umd.edu. Research supported by NSF grants CCF-2110113 (NSF-BSF) and CCF-2104489.

p-Rényi divergence-based ($p \in [0,2]$) uniformity guarantees for universal hash functions. They further extended the analysis to the case p > 2 in [42].

In this paper, we focus on distilling an approximately uniform distribution from a weak source with the help of a special class of hash functions, namely random linear codes $\{\mathscr{C}\}$, relying on the p-norms, $p = 2, 3, \ldots, \infty$ to quantify the distance to uniformity. In contrast to [21, 42], we do not make any assumptions about the source, and our bounds work both for the finite and asymptotic settings. Relying on properties of random linear codes, we obtain better bounds compared to [42] for p > 2.

An informal statement of the described results amounts to the following. LHL states that, for a source Z and a family of universal hash functions with range of m bits, $d_{\text{TV}}(P_{f(Z),P_{U_m}}) \leq \epsilon$ if $m \leq H_{\infty}(Z) - \log(1/\epsilon)$, and [3] proves a similar result, replacing $H_{\infty}(Z)$ with $H_2(Z)$ and d_{TV} with the KL divergence. We extend these claims to all integer orders of Rényi divergence $p \geq 2$ with linear families of hash functions.

An early study of linear codes as hash functions was performed by Alon *et al.* [1]. In particular, within certain parameter regimes, [1] derived bounds for the expected *maximum bucket size*, i.e., the largest set of source sequences having the same syndrome. This result effectively compares the distribution of hash values to the uniform distribution in l_{∞} sense. Extending this work, Dhar and Dvir [13] showed that for almost all the linear hash functions, the size of the largest hash bucket is close to the expected size. Both works assume that the source sequence is sampled uniformly from a subset of the universe (in our case, \mathbb{F}_q^n), limiting the scope of the results. In contrast, we allow any distribution of the source, without assuming anything beyond knowing its p-Rényi entropy $H_p(Z)$.

As our main result, we estimate the expected l_p -distance (equivalently, the p-Rényi divergence) of the hashed source to the uniform distribution on \mathbb{F}_q^m . The assumption of knowing $H_p(Z)$ is weaker than the conditions in [1] and [13]. In terms of the bounds on the distance to uniformity, previous results covered only the cases $p=1,\infty$; here we address all the intermediate integer values. Extending our bounds for finite values of p to $p=\infty$, we also estimate the size of the largest hash bucket that complements the existing results.

The main tool we use to prove our results is often called called smoothing of distributions. We say a distribution P on \mathbb{F}_q^n is (ϵ, p) -smoothable with respect to \mathscr{C} if $\|q^nP*P_{\mathscr{C}}\|_p-1\leq \epsilon$, where $P_{\mathscr{C}}$ is the uniform distribution over the code \mathscr{C} . This metric is easily seen to measure closeness to uniformity; as explained formally in Sec. 2.2, it is also equivalent to p-Rényi divergence. Smoothing of distributions [11, 27] has been also studied in information theory under the names of channel resolvability [17], or noisy functions [38, 39]. It has applications in information-theoretic security [20, 4, 31], coding theory [22, 34, 31], converse coding theorems of information theory [2, 32], strong coordination [10, 4], secret key generation [9, 26], and worst-to-average case reductions in cryptography [27, 5, 12].

In this work, we use smoothing as a way to distill a uniform distribution from a weak (low-entropy) source. We start with an ensemble of random linear codes $\{\mathscr{C}\}$. Consider two random vectors X and Z, where X is distributed uniformly over a code \mathscr{C} sampled from the ensemble and Z is distributed according to P. If P is smoothable with respect to \mathscr{C} , then X+Z is approximately uniformly distributed. Pursuing strong extraction, we also require that the (nearly) uniform vector X+Z be independent of the code \mathscr{C} . We show that by projecting the output random vector onto the syndrome space, we can eliminate the randomness borrowed from the choice of the codeword. We also show that the resulting random vector is almost independent of the choice of the code itself.

In addition to general sources, we consider the binary case, whereby the random source generates a sequence of independent (biased) Bernoulli trials. Simulating an approximate uniform distribution using such a source was studied in [44, 17]. The authors of [46] considered this problem under Rényi divergence proximity measures and obtained the largest rate of uniform bits that can be produced in such instances, called the intrinsic randomness. Their results rely on a computationally involved rearrangement of the distributions. Taking the view of efficient distillation of uniformity, we show that almost all randomness

(Rényi entropy) the Bernoulli source can be converted into a uniform distribution using a low-complexity deterministic mechanism, given by the parity-check matrices of Reed-Muller (RM) codes. In other words, we show that RM codes achieve p-Rényi resolvability rates of binary Bernoulli sources for integer $p \ge 2$.

2. Preliminaries

We begin with setting up the notation for the rest of the paper. Let $q \geq 2$ be a prime power and let \mathbb{F}_q^n be the n-dimensional vector space over the field \mathbb{F}_q , equipped with the Hamming metric. A linear subspace of \mathbb{F}_q^n is called a linear code. Below we use the letter \mathscr{C} to refer to various codes considered in the paper, including random codes. Unless specified otherwise, the codes are always assumed to be linear.

Probability distributions in this paper are supported on linear spaces over the finite field \mathbb{F}_q . For a random variable/vector Z, P_Z denotes the probability mass function (pmf) of Z. Sometimes we use Z to refer to random vectors from \mathbb{F}_q^n , using the same notation P_Z for their distributions. For a given probability distribution P, we write $Z \sim P$ to express the fact that $P_Z = P$. If Z is distributed uniformly over a subset $S \subset \mathbb{F}_q^n$, with some abuse of notation we write $Z \sim S$. The uniform random variable on \mathbb{F}_q^n is denoted by U_n , and P_{U_n} refers to its distribution. For a code $\mathscr{C} \in \mathbb{F}_q^n$, $P_{\mathscr{C}}$ denotes the uniform distribution on the code \mathscr{C} and $X_{\mathscr{C}}$ is a random codeword of \mathscr{C} .

2.1. **Measures of randomness.** Let X be a random variable. Our measure of choice for quantifying the amount of randomness in X is the Rényi entropy H_p of order p. For $p \in (1, \infty)$, it is defined as follows

(1)
$$H_p(X) = \frac{1}{1-p} \log_q \left(\sum_x P_X(x)^p \right).$$

Taking the limits, we also find

$$H_1(X) = -\sum_{x} P_X(x) \log_q P_X(x)$$

$$H_{\infty}(X) = \min_{x} (-\log_q P_X(x)),$$

so for p=1, the Rényi entropy is the familiar Shannon entropy. If this is the case, below we omit the subscript. The entropy H_{∞} is often called min-entropy in the literature.

We note that $H_p(X)$ is a decreasing function of p, while the function $\frac{p-1}{p}H_p(X)$ increases with p. These relations allow us to bound Rényi entropies of different orders through each other, leading to general bounds for uniformity of smoothed sources.

Extending the notion of Rényi entropy to a pair of distributions, one also defines the *Rényi divergence*. For two discrete distributions $P \ll Q$ on the same probability space \mathfrak{X} and for $p \in (1, \infty)$, let

$$D_p(P||Q) = \frac{1}{p-1} \log_q \sum_x P(x)^p Q(x)^{-(p-1)}.$$

As above, we can take the limits to obtain

$$D_1(P||Q) = \sum_{x} P(x) \log_q \frac{P(x)}{Q(x)}$$

$$D_{\infty}(P||Q) = \max_{x} \log_{q} \frac{P(x)}{Q(x)}.$$

If p=1, below we omit the subscript and simply write $D(\cdot||\cdot)$ to refer to the KL divergence.

2.2. Proximity of distributions and measures of uniformity. Since we are interested in quantifying the closeness of distributions to P_{U_n} , we start with introducing some measures of proximity. Before defining them, recall the expression for the p-norm (p-th moment) of a function $f: \mathbb{F}_q^n \to \mathbb{R}$:

(2)
$$||f||_p = \begin{cases} \left(\frac{1}{q^n} \sum_{x \in \mathbb{F}_q^n} |f(x)|^p\right)^{1/p} & \text{for } p \in (0, \infty) \\ \max_{x \in \mathbb{F}_q^n} |f(x)| & \text{for } p = \infty. \end{cases}$$

Under the uniform distribution U_n on \mathbb{F}_q^n , f becomes a random variable, and $||f||_p = (\mathbb{E}_{P_{U_n}}|f|^p)^{\frac{1}{p}}$, 0 . If <math>f = P is a pmf on \mathbb{F}_q^n , then

(3)
$$||P||_p^p = q^{-(p-1)H_p(P)-n}$$

$$||P||_1 = q^{-n}.$$

The case of p=1 leads to a distance between distributions used in many applications. Define the *total* variation distance as

$$d_{\text{TV}}(P,Q) = \max_{A \in \mathbb{F}_q^n} |P(A) - Q(A)|.$$

Its relation to the 1-norm becomes apparent once we write it in a different form:

$$d_{\text{TV}}(P,Q) = \frac{q^n}{2} ||P - Q||_1.$$

There are more than a few options to measure the distance from a given distribution P to the uniform distribution. We will use the following three metrics. The first one, which we also call D_p -smoothness, is the p-divergence, $D_p(P||P_{U_n}), p \in [0, \infty]$. It is easy to see that $D_p(P||P_{U_n}) = n - H_p(P)$, so it is is an increasing function of p.

Another measure of uniformity is the l_p distance $\|P - P_{U_n}\|_p$. For computational convenience, we remove the dependence on n, normalizing by the expectation $\mathbb{E}P = \|P\|_1$, so we will use $q^n \|P - P_{U_n}\|_p = \|q^n P - 1\|_p$.

To introduce our last version of closeness to uniformity, observe that

$$||q^n P||_p = \frac{||P||_p}{||P||_1} \ge 1$$
 for $p \in (1, \infty]$

with equality iff $P = P_{U_n}$. Thus, the better the distribution P approximates the uniform distribution, the closer is $||q^nP||_p$ to 1. Therefore, $\Delta_p^{(n)}(P) := ||q^nP||_p - 1$, $p \neq 1$ can be considered as another measure of uniformity. We call it the l_p -smoothness of the distribution P.

We say that the uniformity measures $m_1(P)$ and $m_2(P)$ are equivalent if $m_1(P) \leq \epsilon$ implies that $m_2(P) = O_p(\epsilon)$ and the same holds upon interchanging the roles of m_1 and m_2 . In the next proposition we show that the measures introduced above are equivalent for $p \in (1, \infty)$. In one part of the proof we rely on Claskson's inequalities [40, p. 388], which state that for functions f and g on \mathbb{F}_q^n ,

where p' = p/(p-1).

Proposition 2.1. Let P be a distribution on \mathbb{F}_q^n and let $p \in (1, \infty)$. Then

(7a)
$$\Delta_p^{(n)}(P) \le \epsilon \implies D_p(P \| P_{U_n}) \le \frac{p}{p-1} \log_q(1+\epsilon)$$

(7b)
$$D_p(P||P_{U_n}) \le \epsilon \implies \Delta_p^{(n)}(P) \le q^{\frac{p-1}{p}\epsilon} - 1$$

and

(8a)
$$\Delta_p^{(n)}(P) \le \epsilon \implies ||q^n P - 1||_p \le \phi_p(\epsilon)$$

(8b)
$$||q^n P - 1||_p \le \epsilon \Rightarrow \Delta_p^{(n)}(P) \le \epsilon,$$

where

$$\phi_p(\epsilon) = \begin{cases} 2\left(\left(\frac{(1+\epsilon)^p + 1}{2}\right)^{p'/p} - 1\right)^{1/p'} & 1$$

Proof. To prove relations (7a)-(7b) all we have to do is to write the divergence in a slightly different form:

(9)
$$D_{p}(P||P_{U_{n}}) = \begin{cases} \frac{p}{p-1} \log_{q} ||q^{n}P||_{p} & \text{for } p \in (0,1) \cap (1,\infty) \\ \lim_{p' \to 1} \frac{p'}{p'-1} \log_{q} ||q^{n}P||_{p'} & \text{for } p = 1 \\ \log_{q} ||q^{n}P||_{\infty} & \text{for } p = \infty. \end{cases}$$

Let us show relations (8a)-(8b). Let $p \in (1,2)$, p' = p/(p-1), then from (5) we have

$$(10) 1 + \left\| \frac{q^n P - 1}{2} \right\|_p^{p'} \le \left\| \frac{q^n P + 1}{2} \right\|_p^{p'} + \left\| \frac{q^n P - 1}{2} \right\|_p^{p'} \le \left(\frac{1}{2} (\|q^n P\|_p^p + 1) \right)^{p'/p},$$

or

$$||q^n P - 1||_p \le 2\left(\left(\frac{||q^n P||_p^p + 1}{2}\right)^{p'/p} - 1\right)^{1/p'}.$$

For $p \in [2, \infty)$, using (6), we obtain

(11)
$$1 + \left\| \frac{q^n P - 1}{2} \right\|_p^p \le \left\| \frac{q^n P + 1}{2} \right\|_p^p + \left\| \frac{q^n P - 1}{2} \right\|_p^p \le \frac{1}{2} (\|q^n P\|_p^p + 1).$$

This yields

(12)
$$||q^n P - 1||_p \le 2 \left(\frac{||q^n P||_p^p - 1}{2} \right)^{1/p}.$$

Finally, for $p \in [1, \infty)$, by Minkowski's inequality we have

$$||q^n P||_n - 1 < ||q^n P - 1||_n$$
. \square

This equivalence allows us to choose the most convenient metric, and below we formulate our results based on the l_n -smoothness.

Observe that for p > 1, the D_p - or l_p -smoothness can be treated as a stronger measure of uniformity compared to the total variation distance because convergence in l_p norm is stronger than convergence in l_1 norm. Therefore, our work can be interpreted as characterizing sufficient conditions for extracting uniformity in a strong sense compared to the TV distance.

2.3. **Universal hash functions.** In this section we recall the known results for extracting uniformity based on universal hash functions.

Below Z is a random vector on \mathbb{F}_q^n with some underlying probability distribution P_Z .

Definition 2.1. A set $\mathscr{F}_{n,m}$ of functions from $\mathbb{F}_2^n \to \mathbb{F}_2^m$ is said to form a universal hash family (UHF) if for all $u, v \in \mathbb{F}_2^n$ with $u \neq v$, $\Pr_{f \sim \mathscr{F}}(f(u) = f(v)) \leq \frac{1}{2m}$.

The randomness extraction property of hash functions relies on the following classic result [23], [43, p.122].

Theorem 2.2. (Leftover Hash Lemma) Let $\epsilon > 0$ and let n, m, t be positive integers. Let $\mathscr{F}_{n,m}$ be a UHF and $f \sim \mathscr{F}_{n,m}$. If $m \leq H_{\infty}(Z) - 2\log(1/\epsilon)$, then,

(13)
$$d_{\text{TV}}(P_{f(Z),f}, P_{U_m} \times P_f) \le \epsilon/2,$$

Note that the condition in (13) can be also written as follows:

$$E_{f \sim \mathscr{F}} \| q^m P_{f(Z)} - 1 \|_1 \le \epsilon.$$

We note another well-known result [3] that established the randomness extracting property of hash functions in a different sense. Rephrased to match our notation, it has the following form.

Theorem 2.3 ([3]). With the notation of Theorem 2.2, if $m \le H_2(Z) - \log(1/\epsilon)$, then

$$E_{f \sim \mathscr{F}}[D(P_{f(Z)} || P_{U_m})] \le \frac{\epsilon}{\ln 2}.$$

The uniformity measures in this theorem and in the leftover hash lemma are, respectively, the KL divergence and the total variation distance, and they are essentially equivalent because of Pinsker's inequalities. At the same time, Theorem 2.3 relies on a somewhat weaker measure of randomness, namely, the 2-Rényi entropy, while yielding essentially the same uniformity claim, so it forms a slightly stronger claim than Theorem 2.2.

Pursuing the line of thought expounded in the introduction, we aim to characterize uniformity in a more stringent way by moving from the TV distance to p-norms with p > 1.

3. Smoothing-based randomness extraction

In this section, we show that random linear codes over \mathbb{F}_q are capable of extracting randomness measured by Rényi entropy. Below the parameters of the code are written as $[n,k]_q$, where n is the length k is the dimension. For positive integers $m \leq n$, denote by \mathfrak{C} the set of all $[n,n-m]_q$ linear codes and for a code $\mathscr{C} \in \mathfrak{C}$ denote by H its parity-check matrix (for brevity, we suppress the dependence on the code from the notation). We assume that H is of dimensions $m \times n$ and note that $\mathrm{rank}(H) = m$. If \mathscr{C} is a random code and Z is a random vector, then P_{HZ} denotes the induced distribution on \mathbb{F}_q^m .

The following theorem forms the main result of our work.

Theorem 3.1. Let $\epsilon > 0$ and let $p \geq 2$ be an integer. If Z is a random vector from \mathbb{F}_q^n with $Z \sim P_Z$ and $m \leq H_p(Z) - p - \log_q(1/\epsilon)$, then

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\Delta_n^{(m)}(P_{\mathsf{H}Z})] \leq \epsilon.$$

The following corollary provides a similar result for the other two types of uniformity measures defined above.

Corollary 3.2. With the assumptions of Theorem 3.1, we have

(14)
$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[D_p(P_{\mathsf{H}Z} || P_{U_m})] \le \frac{p\epsilon}{(p-1)\ln q}$$

(15)
$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \| q^m P_{\mathsf{H}Z} - 1 \|_p \le 2^{1 - 1/p} ((1 + \epsilon)^p - 1)^{1/p}.$$

Proof. Using the inequality $\ln x \le x - 1$ in the first relation in (9), we find that

$$\frac{p-1}{p}(\ln q)D_p(P_{\mathsf{H}Z}||P_{U_m}) \le ||q^m P_{\mathsf{H}Z}||_p - 1,$$

which proves (14). The second inequality is proved as follows. From (11),

$$\left[2\left\|\frac{q^{m}P_{\mathsf{H}Z}-1}{2}\right\|_{p}^{p}+1\right]^{1/p} \leq \|q^{m}P_{\mathsf{H}Z}\|_{p}$$

From the assumptions, we have

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \Big[2 \Big\| \frac{q^m P_{\mathsf{H}Z} - 1}{2} \Big\|_p^p + 1 \Big]^{1/p} \leq 1 + \epsilon.$$

For brevity, let $X := 2^{1/p} \left\| \frac{q^m P_{\mathsf{H}Z} - 1}{2} \right\|_p$, then the above inequality takes the form

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[X^p + 1]^{1/p} \le 1 + \epsilon.$$

Applying Jensen's inequality we obtain

$$([\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} X]^p + 1)^{1/p} \le \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} [X^p + 1]^{1/p} \le 1 + \epsilon.$$

This implies the relation,

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} X \le ((1+\epsilon)^p - 1)^{1/p},$$

which proves (15).

Remark 1. The condition $\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z} - 1\|_p] \leq \epsilon$ not only guarantees that the distribution $P_{\mathsf{H}Z}$ is close to uniform on average, but also says that $P_{\mathsf{H}Z}$ is almost independent of the choice $\mathscr{C} \sim \mathfrak{C}$. Indeed, since $P_{\mathsf{H}Z}$ is close to the uniform distribution, it does not depend on \mathscr{C} for almost all the codes. As mentioned in the introduction, the uniformity of hash functions is typically measured in total variation distance. As a result, the approximate independence is also measured in the same metric. Since $\|q^m P_{\mathsf{H}Z} - 1\|_p$ is increasing in p, it is a stronger measure of independence compared to the total variation distance.

The main ingredient of the proof of Theorem 3.1 is the following result on smoothing a source using random linear codes.

Theorem 3.3. Let Z be a random vector in \mathbb{F}_q^n . Let \mathfrak{C} be the set of all $[n,k]_q$ linear codes. Then for all natural $p \geq 2$,

(16)
$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^n P_{X_{\mathscr{C}} + Z}\|_p^p] \leq \sum_{d=0}^p \binom{p}{d} q^{(p-d)(d+n-k-H_p(Z))},$$

where $X_{\mathscr{C}}$ is a uniform random codeword of \mathscr{C} .

Remark 2. This result is independent of its applications to hashing, and it can be strengthened if random linear codes are replaced with the ensemble of all binary codes of the same cardinality. More precisely, if \mathfrak{C}' is the set of all $(n, q^k)_q$ codes, then

(17)
$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}'}[\|q^n P_{X_{\mathscr{C}} + Z}\|_p^p] \le \sum_{d=0}^p \begin{Bmatrix} p \\ d \end{Bmatrix} q^{(p-d)(n-k-H_p(Z))},$$

where $\binom{p}{d}$ is the Stirling number of second kind. From [35, Theorem 3], we have $\binom{p}{d} \leq \binom{p}{d} d^{p-d} \leq \binom{p}{d} q^{d(p-d)}$, which shows that (17) is a better bound than (16). Since this result does not have direct implication for hashing, we will omit the proof.

Below we will say that an unordered tuple of vectors (v_1, \ldots, v_p) is a [p, d]-tuple if their linear span has dimension d. If $[v_1, \ldots, v_p]$ is the $n \times p$ matrix whose columns are the vectors v_1, \ldots, v_p , then this means that $\operatorname{rank}[v_1, \ldots, v_p] = d$. To shorten the notation, below we will write " $(v_1^p \text{ is } [p, d])$ ".

The proof of Theorem 3.3 is based on the following lemmas.

Lemma 3.4. Let \mathfrak{C} be the set of $[n,k]_q$ linear codes in \mathbb{F}_q^n . Let $p \geq 2$ be an integer. For any function $f:(\mathbb{F}_q^n)^p \to [0,\infty)$,

(18)
$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \left[\sum_{z_1, \dots, z_p \in \mathscr{C}} f(z_1, \dots, z_p) \right] \leq \sum_{d=0}^{\min\{k, p\}} q^{d(k-n)} \sum_{(v_1^p \text{ is } [p, d])} f(v_1, \dots, v_p).$$

Lemma 3.5. Let $1 \leq d < p$. Then for any function $f : \mathbb{F}_q^n \to [0, \infty)$,

$$\sum_{\substack{(v_1^p \text{ is } [p,d])}} \prod_{1 \le j \le p} f(v_j) \le \binom{p}{d} \sum_{m=0}^{p-d} \binom{p-d}{m} (q^d-1)^{p-d-m} q^{nd} f(0)^m \|f\|_1^{d-1} \|f\|_{p-d-m+1}^{p-d-m+1}.$$

Proofs of Lemmas 3.4 and 3.5 are deferred to Appendix A.

Remark 3. Lemma 3.4 is equivalent to the following fact. For a fixed $u_1, \ldots, u_p \in \mathbb{F}_q^n$,

(19)
$$\Pr_{\mathscr{L}_{a,\sigma}}\{(u_1,\ldots,u_p)\in\mathscr{C}\}\leq q^{-\operatorname{rank}[u_1,\ldots,u_p](n-k)}.$$

This version of Lemma 3.4 has been used in the literature to prove that some ensembles of linear codes achieve list decoding capacity [29]. In particular, it is known that some random ensembles such as LDPC codes [29] and randomly punctured low-bias codes [16] approximately satisfy (19). The aforementioned papers used this fact to show that these code ensembles also achieve list decoding capacity. Since the only property of random codes that we use in the proof of Theorem 3.1 is the one mentioned in Lemma 3.4, in our main result we can replace the ensemble of random linear codes with smaller ensembles such as LDPC or low-bias codes.

Next, assuming validity of Lemmas 3.4 and 3.5, let us give a proof of Theorem 3.3.

Proof. Recall that P_Z refers to the distribution of the vector Z sampled from \mathbb{F}_q^n . Below we abbreviate $\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}$ to $\mathbb{E}_{\mathscr{C}}$. We have

$$\mathbb{E}_{\mathscr{C}}[\|q^{n}P_{X_{\mathscr{C}}+Z}\|_{p}^{p}] = \mathbb{E}_{\mathscr{C}}[\|q^{n}P_{X_{\mathscr{C}}+Z}\|_{p}^{p}] = \mathbb{E}_{\mathscr{C}}\left[\frac{1}{q^{n}}\sum_{x\in\mathbb{F}_{q}^{n}}(q^{n}P_{Z}*P_{\mathscr{C}}(x))^{p}\right]$$

$$= \mathbb{E}_{\mathscr{C}}\left[q^{n(p-1)}\sum_{x\in\mathbb{F}_{q}^{n}}\sum_{z_{1},...,z_{p}\in\mathbb{F}_{q}^{n}}\prod_{j=1}^{p}P_{Z}(x-z_{j})P_{\mathscr{C}}(z_{j})\right]$$

$$= \sum_{x\in\mathbb{F}_{q}^{n}}\frac{q^{n(p-1)}}{q^{kp}}\mathbb{E}_{\mathscr{C}}\left[\sum_{z_{1},...,z_{p}\in\mathscr{C}}\prod_{l=1}^{p}P_{Z}(x-z_{l})\right]$$

$$\leq \sum_{d=0}^{p}q^{(p-d)(n-k)-n}\sum_{x\in\mathbb{F}_{q}^{n}}\sum_{(v_{1}^{p}\text{ is }[p,d])}\prod_{l=1}^{p}P_{Z}(x-v_{l}),$$

$$(20)$$

where (20) follows by Lemma 3.4. Let us introduce the notation

(21)
$$g(d) := \frac{1}{q^n} \sum_{x \in \mathbb{F}_q^n} \sum_{(v_1^p \text{ is } [p,d])} \prod_{l=1}^p P_Z(x - v_l).$$

Then the bound (20) can be written as

(22)
$$\mathbb{E}_{\mathscr{C}}[\|q^n P_{X_{\mathscr{C}}+Z}\|_p^p] \le \sum_{d=0}^p q^{(p-d)(n-k)} g(d).$$

The theorem will be proved if we show that

(23)
$$g(d) \le \binom{p}{d} q^{(p-d)(d-H_p(P_Z))}.$$

For d = 0, p this is immediate from the definition:

(24)
$$g(0) = \frac{1}{q^n} \sum_{x \in \mathbb{F}_q^n} P_Z(x)^p = q^{-(p-1)H_p(P_Z) - n} \le q^{-pH_p(P_Z)}$$

(25)
$$g(p) \le \frac{1}{q^n} \sum_{x \in \mathbb{F}_q^n} \prod_{l=1}^p \sum_{v \in \mathbb{F}_q^n} P_Z(x-v) = 1 = q^{(p-p)H_p(P_Z)}.$$

Now let us consider the case 0 < d < p. To shorten the writing, for the remainder of the proof we put $\tau = p - d - m - 1$. Using Lemma 3.5 and (4), we have

$$g(d) = \frac{1}{q^{n}} \sum_{x \in \mathbb{F}_{q}^{n}} \sum_{(v_{1}^{p} \text{ is } [p,d])} \prod_{l=1}^{p} P_{Z}(x - v_{l})$$

$$\leq \frac{1}{q^{n}} \sum_{x \in \mathbb{F}_{q}^{n}} \binom{p}{d} \sum_{m=0}^{p-d} \binom{p-d}{m} (q^{d} - 1)^{p-d-m} q^{nd} P_{Z}(x)^{m} \|P_{Z}\|_{1}^{d-1} \|P_{Z}\|_{\tau}^{\tau}$$

$$= \binom{p}{d} \sum_{m=0}^{p-d} \binom{p-d}{m} (q^{d} - 1)^{p-d-m} \sum_{x} P_{Z}(x)^{m} \|P_{Z}\|_{\tau}^{\tau}.$$
(26)

First let m = 0. Using (3),

$$\sum_{x} P_Z(x)^m \|P_Z\|_{\tau}^{\tau} = q^n \|P_Z\|_{p-d-1}^{p-d-1} = q^{-(p-d)H_p(P_Z)}.$$

Further for $m \ge 1$, again using (3) and the inequalities $m < p, \tau < p$, we obtain

$$\sum_{\tau} P_Z(x)^m \|P_Z\|_{\tau}^{\tau} = q^{-(m-1)H_m(P_Z) - (\tau+1)H_{\tau}(P_Z) - n} \le q^{-(p-d)H_p(P_Z)}.$$

Using these results in (26), we obtain

$$g(d) \le \binom{p}{d} q^{-(p-d)H_p(P_Z)} \sum_{m=0}^{p-d} \binom{p-d}{m} (q^d-1)^{p-d-m},$$

which gives the right-hand side of (23). This concludes the proof of Theorem 3.3.

To prove Theorem 3.1, we need an additional lemma that establishes a connection between smoothing and linear hashing.

Lemma 3.6. Let \mathscr{C} be an $[n,k]_q$ linear code, and let H be its parity check matrix. Let $X_{\mathscr{C}}$ be a uniform random codeword of \mathscr{C} and $Z \sim P_Z$ be a random vector in \mathbb{F}_q^n . Then for $p \in (0,\infty]$

$$||q^{n-k}P_{\mathsf{H}Z}||_p = ||q^nP_{X_{\mathscr{C}}+Z}||_p.$$

Proof. Define $\mathcal{S} := \{ Hy : y \in \mathbb{F}_q^n \}$ to be the set of syndromes corresponding to code \mathscr{C} . Observe that

$$P_{\mathsf{H}(X_\mathscr{C}+Z)}(u) = \sum_{y: \mathsf{H}y=u} P_{X_\mathscr{C}+Z}(y) = |\mathscr{C}| P_{X_\mathscr{C}+Z}(y_u),$$

where y_u is a representative of the coset defined by u. The last equality is due to the fact that

$$P_{X_{\mathscr{C}}+Z}(y) = P_{\mathscr{C}} * P_Z(y) = P_{\mathscr{C}} * P_Z(y+c) = P_{X_{\mathscr{C}}+Z}(y+c)$$

for any $c \in \mathscr{C}$. Hence, for $p \in (0, \infty)$,

$$\|q^{n-k}P_{\mathsf{H}(X_{\mathscr{C}}+Z)}\|_p^p = q^{(n-k)(p-1)} \sum_{u \in \mathbb{F}_q^{n-k}} P_{\mathsf{H}(X_{\mathscr{C}}+Z)}(u)^p$$

$$= q^{(n-k)(p-1)} \sum_{y \in \mathbb{F}_q^n} \frac{(|\mathscr{C}|P_{X_{\mathscr{C}}+Z}(y))^p}{|\mathscr{C}|}$$

$$= \|q^n P_{X_{\mathscr{C}}+Z}\|_p^p.$$

Finally, since $H(X_{\mathscr{C}} + Z) = HZ$, we have proven the desired statement for $p \in (0, \infty)$. The case $p = \infty$ is obtained by taking a continuous limit.

Now Theorem 3.1 follows from Theorem 3.3 and Lemma 3.6 as follows:

Proof. (of Theorem 3.1) By Lemma 3.6

$$\begin{split} \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z}\|_p^p] &= \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^n P_{X_{\mathscr{C}} + Z}\|_p^p] \\ &\leq \sum_{d=0}^p \binom{p}{d} q^{d(p-d)} q^{(p-d)(m-H_p(Z))} \\ &\leq \left(1 + q^{m-H_p(Z) + p}\right)^p. \end{split} \tag{Theorem 3.3}$$

Therefore, we have

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z}\|_p] - 1 \le q^{m - H_p(Z) + p}.$$

Choosing, m such that $m \leq H_p(Z) - p - \log_q(1/\epsilon)$ we obtain $\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \|q^m P_{\mathsf{H}Z}\|_p - 1 \leq \epsilon$ proving the desired result. \square

Remark 4. (Symbol loss) If it were possible to distill all the randomness of $H_p(Z)$ into uniform symbols, this would yield $H_p(Z)$ random symbols. However Theorem 3.1 (equivalently Corollary 3.2) states that we can achieve the prescribed uniformity if we distill $m \leq H_p(Z) - p - \log_q(1/\epsilon)$ symbols. This is at least $p + \log_q(1/\epsilon)$ symbols away from the ideal target.

More precisely, we say that hashing achieves (ϵ, p) -uniformity for a source Z with a loss of $s = H_p(Z) - m$ symbols if

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\Delta_n^{(m)}(P_{\mathsf{H}Z})] \le \epsilon.$$

In these terms, Theorem 3.1 states that (ϵ, p) -uniformity is attained with an at most $p + \log_q(1/\epsilon)$ symbol loss. Since this is a constant, this loss is negligible in the setting of large t.

However, when $H_p(Z)$ is not very large, this symbol loss may be significant. We show that it can be reduced for the case of collision entropy, i.e., p = 2, saving more than two symbols.

Proposition 3.7. With the assumptions of Theorem 3.1, linear hashing achieves $(\epsilon, 2)$ -uniformity with a $\log_q \frac{1}{(1+\epsilon)^2-1}$ loss.

Proof. Defining $(P_Z \circ P_Z)(x) = \sum_{y \in \mathbb{F}_q^n} P_Z(y) P_Z(x+y)$, we proceed as follows:

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^{m}P_{\mathsf{H}Z}\|_{2}^{2}] = \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^{n}P_{X_{\mathscr{C}}+Z}\|_{2}^{2}]$$

$$= \frac{1}{q^{n}} \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \sum_{y \in \mathbb{F}_{q}^{n}} \sum_{x_{1} \in \mathbb{F}_{q}^{n}} q^{n}P_{Z}(y - x_{1})P_{\mathscr{C}}(x_{1}) \sum_{x_{2} \in \mathbb{F}_{q}^{n}} q^{n}P_{Z}(y - x_{2})P_{\mathscr{C}}(x_{2})$$

$$= \frac{1}{q^{n}} \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \sum_{x_{1} \in \mathscr{C}} \sum_{x_{2} \in \mathscr{C}} \sum_{y \in \mathbb{F}_{q}^{n}} \frac{q^{n}P_{Z}(y - x_{1})}{|\mathscr{C}|} \frac{q^{n}P_{Z}(y - x_{2})}{|\mathscr{C}|}$$

$$= q^{n-2k} \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \sum_{x_{1} \in \mathscr{C}} \sum_{x_{2} \in \mathscr{C}} (P_{Z} \circ P_{Z})(x_{1} - x_{2}) \quad (k := n - m)$$

$$= q^{n-k} \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \sum_{z \in \mathscr{C}} (P_{Z} \circ P_{Z})(z)$$

$$\leq q^{n-k} \left((P_{Z} \circ P_{Z})(0) + q^{k-n} \sum_{z \in \mathbb{F}_{q}^{n}} (P_{Z} \circ P_{Z})(z) \right)$$

$$= q^{n-k} \left(\sum_{z \in \mathbb{F}_{q}^{n}} P_{Z}(z)^{2} + q^{k-n} \right)$$

$$= 1 + q^{m-H_{2}(Z)},$$

$$(28)$$

where the inequality follows from Lemmma 3.4. Therefore if $m \leq H_2(Z) - \log_q \frac{1}{(1+\epsilon)^2-1}$, then

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\Delta_p^{(m)}(P_{\mathsf{H}Z})] \le \epsilon. \quad \Box$$

Similar improvements can be obtained for the first few integer values of $p \ge 3$. At the same time, the proof of Prop. 3.8 suggests that for large p it is desirable to have a loss which is a sublinear function of p. We do not know if our results can be tightened so as to attain this loss scaling.

A natural question that arises in our line of work is whether our results on D_p - (or l_p -) smoothness improve the conclusions of classical theorems such as Theorems 2.2 or 2.3. The closeness to uniformity in these theorems is measured by the total variation distance and KL divergence (Shannon entropy), respectively. Note that these quantities are bounded by the l_p distance and D_p -smoothness due to the monotonicity with respect to p. Specifically, we have

$$d_{\mathsf{TV}}(P_{\mathsf{H}(Z),\mathscr{C}} \| P_{U_m} \times P_{\mathscr{C}}) = 2\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \| q^m P_{\mathsf{H}(Z)} - 1 \|_1 \le 2\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} \| q^m P_{\mathsf{H}(Z)} - 1 \|_p$$
$$D(P_{\mathsf{H}(Z)} \| P_{U_m} \times P_{\mathscr{C}}) = \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} [D(P_{\mathsf{H}(Z)} \| P_{U_m})] \le \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}} [D_p(P_{\mathsf{H}(Z)} \| P_{U_m})].$$

However, these inequalities do not enable us to improve upon the claim of Theorem 2.3. For instance, using 2-entropy and D_2 -smoothness in Theorem 3.1 (or in (28)), and using the second inequality essentially reproduces the claim of this theorem, which is already known for any UHF. At the same time, this theorem does not imply our results for p > 2 because the only assumption used in its proof is that of general UHFs. To establish our results we additionally employ p-balancedness (defined in the appendix, see Def. A.1), which is not readily available under the general approach.

3.1. Expected size of the largest hash bucket. Given a source Z, the hashing function maps n-vectors generated by it to the syndromes of a randomly chosen [n,n-m] linear code $\mathscr C$. If $\mathscr C$ is fixed, then the expectation with respect to P_Z of the largest number of source sequences mapped to the same syndrome s is $P_{\mathsf{H}Z}(s)$. Therefore the expected maximum bucket size is $\|P_{\mathsf{H}Z}\|_{\infty}$. Taking a random code $\mathscr C$ results in the expected maximum hash bucket having the size $\mathbb E_{\mathscr C \sim \mathfrak C}\|P_{\mathsf{H}Z}\|_{\infty}$. The average size of the hash bucket is q^{-m} , and so we study the quotient of these two quantities, given by $\mathbb E_{\mathscr C \sim \mathfrak C}[\|q^m P_{\mathsf{H}Z}\|_{\infty}]$. Our main result is

a new bound on this quantity when $m \leq H_{\infty}(Z) - n\epsilon$. Note that our main result, Theorem 3.1, does not cover the case $p = \infty$. We will circumvent this obstacle by extending the bounds for finite p to infinity.

Given a source Z, one would expect to distill $H_\infty(Z)$ approximately uniform q-ary symbols. Intuitively, if we aim at distilling fewer than $H_\infty(Z)$ random symbols, we can make the resulting vector to be 'more' uniform. On the other hand, attempting to distill more than $H_\infty(Z)$ symbols forces the resulting distribution to deviate from the uniformity. In the existing literature, this intuitive reasoning has been rigorously quantified for different regimes of H(Z) and m [1, 13]. In terms of $\|q^m P_{\mathsf{H}Z}\|_\infty$, the bound of [1] for the expected size of the largest hash bucket has the form

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z}\|_{\infty}] = O(H_{\infty}(Z) \log H_{\infty}(Z)).$$

Here $m=H_{\infty}(Z)$ and the sequences of the source Z are drawn uniformly from a subset of the universe (e.g., \mathbb{F}_q^n).

For m slightly smaller than the source entropy, namely $m + \log m = H_{\infty}(Z)$, [1] has the bound

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|2^m P_{\mathsf{H}Z}\|_{\infty}] = O(\log H_{\infty}(Z)).$$

We consider the case of the gap between m and the min-entropy proportional to n, showing that then then the expected size is bounded by an absolute constant.

Proposition 3.8. Let $\epsilon > 0$ and let n be a positive integer. Let $Z \sim P_Z$ be a random vector in \mathbb{F}_q^n . Choose m such that $m \leq H_{\infty}(Z) - n\epsilon$, and define \mathfrak{C} the set of $[n, n-m]_q$ linear codes. Then

(29)
$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z}\|_{\infty}] = O(1).$$

Proof. From the fact $H_p(Z) \leq \frac{p}{p-1} H_{\infty}(Z)$, we have

(30)
$$pD_{\infty}(P_Z||P_{U_n}) - n \le (p-1)D_p(P_Z||P_{U_n}).$$

Further,

$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^{m}P_{\mathsf{H}Z}\|_{\infty}] = \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^{n}P_{X_{\mathscr{C}}+Z}\|_{\infty}] \qquad \text{(from Lemma 3.6)}$$

$$= \mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[q^{D_{\infty}(P_{X_{\mathscr{C}}+Z}\|P_{U_{n}})}]$$

$$\leq q^{\frac{n}{p}}\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[q^{\frac{p-1}{p}D_{p}(P_{X_{\mathscr{C}}+Z}\|P_{U_{n}})}] \quad \text{(by (30))}$$

$$= q^{\frac{n}{p}}\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^{n}P_{X_{\mathscr{C}}+Z}\|_{p}]$$

$$\leq q^{\frac{n}{p}}(1+q^{m-H_{p}(Z)+p}) \quad \text{(by (27))}$$

$$\leq q^{\frac{n}{p}}(1+q^{m-H_{\infty}(Z)+p}).$$

Now, choose m such that $m = H_{\infty}(Z) - n\epsilon$. Then by choosing $p = \epsilon n/2$, we obtain

(31)
$$\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z}\|_{\infty}] \le q^{2/\epsilon} (1 + q^{-\epsilon n/2}),$$

proving the desired result.

The recent work [13] established a high-probability estimate for the maximum size of a hash bucket. In this work, too, the random vector Z was assumed to be uniform over a subset. In particular, for the case $m \leq H_{\infty}(Z) - n\epsilon$ their results imply that

(32)
$$\Pr_{\mathscr{C} \sim \mathfrak{C}}(\|q^m P_{\mathsf{H}Z} - 1\|_{\infty} \ge \epsilon) \le 2^{-\zeta n}$$

for some constant $\zeta = \zeta(\epsilon)$. We note that neither of the estimates (29) and (32) implies the other one, for the following reasons. The probabilistic estimate in (32) states that the fraction of codes that result in an approximately uniform P_{HZ} is close to one. This discounts the outliers that that yield conditional distributions P_{HZ} very different from the uniform one. On the other hand, the quantity $\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{HZ}\|_{\infty}]$

takes all these outliers into account, so (32) does not imply (29). In the other direction, if we had a stronger estimate of the form $\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z} - 1\|_{\infty}] = o(1)$, Markov's inequality would yield a result comparable to (32). As it stands, the estimate in (29) is not strong enough to imply (32).

An interesting open question is whether it is possible to improve (29) to $\mathbb{E}_{\mathscr{C} \sim \mathfrak{C}}[\|q^m P_{\mathsf{H}Z}\|_{\infty}] = 1 + o(1)$. If true, it would give strong guarantees for both uniformity and independence (see Remark 1).

Remark 5. In the language of smoothing, $\|q^n P_{X_{\mathscr{C}}+Z}\|_{\infty} - 1$ is the l_{∞} -smoothness of $P_{X_{\mathscr{C}}+Z}$. In [31], we proved that smoothing of P_Z with respect to a random code \mathscr{C} of dimension $k \geq n - H_{\infty}(P_Z) + \epsilon n$ yields $\mathbb{E}_{\mathscr{C}}[\|q^n P_{X_{\mathscr{C}}+Z}\|_{\infty}] = 1 + o_n(1)$. We do not see a way of obtaining a similar estimate for linear codes, leaving this as an open problem.

4. RANDOM BITS FROM BERNOULLI SOURCES WITH RM CODES

In addition to the problem of smoothing an arbitrary source, one can also consider the same question for a source with a known distribution. For a fixed source, the amount of randomness that can be converted to (approximate) uniformity is sometimes called the intrinsic randomness [44]. The authors of [46] proposed a way of approximately simulating a given memoryless distribution with a known distribution, including the uniform one, using another memoryless source and a carefully chosen mapping, and measuring proximity by the Rényi divergence. We will state their result for Bernoulli sources, starting with the definition of intrinsic randomness of a Bernoulli random variable.

Definition 4.1. Let Z be a Bernoulli(δ) random variable. The p-Rényi intrinsic randomness of Z (in more detail, of a memoryless source given by Z) is defined as

(33)
$$I_p(Z) = \sup\{R : \lim_{n \to \infty} \sup_{(k_n) : \frac{k_n}{n} \to R} \sup_{f_n \in F(n, k_n)} D_p(P_{f_n(Z_n)} || P_{U_{k_n}}) = 0\},$$

where F(n,m) is the set of all functions from \mathbb{F}_2^n to \mathbb{F}_2^m and Z_n is a vector formed of n independent copies of Z.

Paper [46] established the following result, stated here for a particular case of Bernoulli sources.

Proposition 4.1. [46, Theorem 10] Let Z be a Bernoulli(δ) random variable. Then

(34)
$$I_p(Z) = \begin{cases} h_p(\delta) & \text{for } p \in \{0\} \cup [1, \infty] \\ h(\delta) & \text{for } p \in (0, 1), \end{cases}$$

where $h_p(\delta) = \frac{1}{1-p} \log_2(\delta^p + (1-\delta)^p)$ is the two-point Rényi entropy and $h = h_1$.

Remark 6. Because of the equivalence stated in Proposition 2.1, if we replace D_P with l_p distance or l_p -smoothness in Definition 4.1, we still obtain $I_p(Z) = h_p(\delta)$ for $p \in (1, \infty)$.

The construction of the uniform uniform distribution in [46] involves rearranging the masses of the Bernoulli distribution into $2^{n(1-R)}$ bins having approximately equal probabilities, which is a computationally involved procedure. In this section, we show that RM codes are capable of extracting randomness from Bernoulli sources. The following theorem is a consequence of [31, Theorem 6].

Theorem 4.2. Let $R \in (0,1)$ and let \mathcal{C}_n be a sequence of RM codes whose rate R_n approaches R. Let H_n be the parity check matrix of \mathcal{C}_n and let Z_n be a binary vector formed of independent Bernoulli(δ) random bits. If $R > 1 - h_p(\delta)$, then

$$\lim_{n \to \infty} D_p(P_{\mathsf{H}_n Z_n} || P_{U_{n(1-R_n)}}) = 0, \quad p \in \{2, \dots, \infty\}$$

If p = 1 and $R > (1 - 2\delta)^2$, then

$$\lim_{n \to \infty} D(P_{\mathsf{H}_n Z_n} || P_{U_{n(1-R_n)}}) = 0.$$

Proof. From Lemma 3.6, we know that for all $p \in (0, \infty]$,

$$||2^{n(1-R_n)}P_{\mathsf{H}_nZ_n}||_p = ||2^nP_{X_{\mathscr{C}_n}+Z_n}||_p.$$

Equivalence of the l_p -smoothness and D_p -smoothness (9) implies that

$$D_p(P_{\mathsf{H}_n Z_n} || P_{U_{n(1-R_n)}}) = D_p(P_{X_{\mathscr{C}_n} + Z_n} || P_{U_n}).$$

Therefore, it suffices to show that for $p \in \{2, ..., \infty\}$ and $R > 1 - h_p(\delta)$,

$$\lim_{n \to \infty} D_p(P_{X_{\mathscr{C}_n} + Z_n} || P_{U_n}) = 0,$$

and that for $R > (1 - 2\delta)^2$,

(36)
$$\lim_{n \to \infty} D(P_{X_{\mathscr{C}_n} + Z_n} || P_{U_n}) = 0,$$

In [31, Theorem 6] we proved exactly this result for a general class of codes that includes RM codes. \Box

Setting $\hat{R}_n = 1 - R_n$ in Theorem 4.2, we observe that for $\hat{R} := 1 - R > h_{\alpha}(\delta)$ and for $p \in \{2, 3, \dots, \infty\}$,

$$\lim_{n\to\infty} D_p(P_{\mathsf{H}_n Z_n} || P_{U_{n\hat{R}_n}}) = 0,$$

i.e., the intrinsic randomness is achievable by setting $f_n(Z_n) = H_n Z_n$. To summarize, we observe that RM codes yield a computationally efficient alternative for distilling a string of (nearly) uniform bits from a Bernoulli source at rate close the intrinsic randomness $I_n(Z)$ of the source.

5. CONCLUDING REMARKS

The obtained results suggest the following open questions, already mentioned in the main text. It is of interest to attempt better bounds for the symbol loss than those mentioned in Remark 4, which may result in better bounds for l_{∞} -smoothness. Another question is to derive similar results for ensembles of linear codes smaller than the ensemble of all linear codes of fixed dimension considered here. Finally, it would be interesting to show that RM codes achieve intrinsic randomness of Bernoulli sources $I_p(Z)$ for p=1. Per the results of [31], this is equivalent to showing that nested sequences of RM codes achieve secrecy capacity of the binary wiretap channel in the strong sense.

APPENDIX A. OMITTED PROOFS

A.1. **Proof of Lemma 3.4.** Let us begin with the following definition that allows using certain symmetry properties of random linear codes.

Definition A.1. Let p be a positive integer and let $\mathfrak C$ be a family of $[n,k]_q$ linear codes in $\mathbb F_q^n$. We call $\mathfrak C$ a p-balanced code family if for all $d \in [0,p]$, any [p,d]-tuple (v_1,\ldots,v_p) appears in the same number of codes from $\mathfrak C$.

Observe that the set of all $[n, k]_q$ linear codes forms a p-balanced collection for all integer $p \ge 1$. For p = 1 this definition appears routinely in proofs of the GV bound.

Let $T_d(n,p) = T_d^{(q)}(n,p)$ be the number of [p,d]-tuples in \mathbb{F}_q^n . If \mathscr{C} is a k-dimensional linear code in \mathbb{F}_q^n , the number of [p,d]-tuples formed by codewords in \mathscr{C} is given by $T_d^{(q)}(k,p)$. With these definitions, we proceed to the following lemma.

Lemma A.1. Let $\mathfrak C$ be an p-balanced collection of codes in $\mathbb F_q^n$. For any function $f:\mathbb F_q^{n\times p}\to\mathbb R$,

(37)
$$\frac{1}{|\mathfrak{C}|} \sum_{\mathscr{C} \in \mathfrak{C}} \sum_{z_1, \dots, z_p \in \mathscr{C}} f(z_1, \dots, z_p) = \sum_{d=0}^p \frac{T_d(k, p)}{T_d(n, p)} \sum_{(v_1^p \text{ is } [p, d])} f(v_1, \dots, v_p).$$

Proof. Suppose that each d-dimensional tuple (v_1, \ldots, v_p) appears S_d times in the family \mathfrak{C} for some number $S_d \geq 1$. Then we find that

(38)
$$\sum_{\mathscr{C} \in \mathfrak{C}} \sum_{z_1, \dots, z_p \in \mathscr{C}} f(z_1, \dots, z_p) = \sum_{d=0}^p S_d \sum_{(v_1, \dots, v_p) \text{ is } [p, d]} f(v_1, \dots, v_p).$$

Letting $f(v_1, \ldots, v_p) = \mathbb{1}\{\operatorname{rank}[v_1, \ldots, v_p] = d\}$, we obtain

$$|\mathfrak{C}|T_d(k,p) = S_dT_d(n,p).$$

Using this in (38) completes the proof.

Now let us estimate $T_d(n,p)/T_d(k,p)$. Observe that $T_d^{(q)}(n,p)$ can be interpreted as the number of $n \times p$ matrices (over \mathbb{F}_q) whose rank is exactly d.

It is well known that for $d \in [0, \min\{n, p\}]$

(39)
$$T_d(n,p) = \prod_{j=0}^{d-1} \frac{(q^n - q^j)(q^p - q^j)}{q^d - q^j}.$$

With this, we have

$$\frac{T_d(k,p)}{T_d(n,p)} = \frac{\prod_{j=0}^{d-1} (q^k - q^j)}{\prod_{j=0}^{d-1} (q^n - q^j)}$$
$$= \prod_{j=0}^{d-1} \frac{q^{k-j} - 1}{q^{n-j} - 1}$$
$$\leq q^{d(k-n)}.$$

Noticing that $T_d(k, p) = 0$, for $d > \min\{k, p\}$, we conclude the proof.

A.2. **Proof of Lemma 3.5.** As the first step of proving Lemma 3.5, we will establish an auxiliary estimate. In the proof we will need the classical *rearrangement inequality* [18, Sec. 10.2]: given two sequences of nonnegative numbers (a_1, \ldots, a_n) and (b_1, \ldots, b_n) , we would like to permute them to maximize the sum $\sum_i a_{\tau(i)} b_{\sigma(i)}$. The inequality states that the maximum is attained when both sequences are arranged in a monotone (nondecreasing or nonincreasing) order. This statement extends to multiple sequences, see e.g., [37].

Lemma A.2. Let $d \in \{1, ..., p-1\}$ be fixed and let $v_j \in \mathbb{F}_q^n, j \in [d]$. Choose arbitrary p-d vectors $x_i \in \mathbb{F}_q^d \setminus \{0\}, i \in [d+1, p]$ and construct vectors $v_{d+1}, ..., v_p$ as linear combinations of the vectors $v_i, i \in [d]$ as follows:

(40)
$$v_i = \sum_{j=1}^d x_{i,j} v_j, \quad i = d+1, \dots, p.$$

Then for a function $f: \mathbb{F}_q^n \to [0, \infty)$

$$\frac{1}{q^{nd}} \sum_{v_1, \dots, v_d \in \mathbb{F}_q^n} f(v_1) f(v_2) \dots f(v_p) \le ||f||_1^{d-1} ||f||_{p-d+1}^{p-d+1},$$

Proof. Let us partition [p] into d classes as follows:

$$B_d = \{i \in [d+1, p] : x_i(d) \neq 0\} \cup \{d\}$$

$$B_j = \{i \in [d+1, p] : x_i(j) \neq 0, i \notin \cap_{l=j+1}^d B_l\} \cup \{j\}, \quad j \in [d-1].$$

In words, class B_j contains j and the indices of the vectors $v_i, i \in [p]$ whose expansion (40) includes v_j , but not v_{j+1}, \ldots, v_d . Note that some of the vectors $v_i, i \geq d+1$ may be simply copies of one of the vectors $v_i, i \in [d]$. Therefore,

$$\sum_{v_1, \dots, v_d \in \mathbb{F}_q^n} f(v_1) f(v_2) \dots f(v_p) = \prod_{i=1}^d \sum_{v_i \in \mathbb{F}_q^n} \prod_{l \in B_i} f(v_l).$$

Observe that when $v_i, i \in [d]$ runs through all the elements in \mathbb{F}_q^n , while keeping the other vectors $v_j, j \in [i-1]$ fixed, the vector $v_l, l \in B_i$ also runs through all the elements in \mathbb{F}_q^n . i.e., $f(v_i)$ and $f(v_l), l \in B_i$ as function evaluations of v_i are permutations of each other. Applying the rearrangement inequality , we find that

$$\sum_{v_i \in \mathbb{F}_q^n} \prod_{l \in B_i} f(v_l) \le \sum_{v_i \in \mathbb{F}_q^n} f(v_i)^{|B_i|}.$$

Denoting $b_i := B_i$, we obtain

$$\frac{1}{q^{nd}} \sum_{v_1, \dots, v_d \in \mathbb{F}_a^n} f(v_1) f(v_2) \dots f(v_p) \le \prod_{i=1}^d ||f||_{b_i}^{b_i}.$$

Now all that is left is to show that $\prod_{i=1}^d \|f\|_{b_i}^{b_i} \leq \|f\|_1^{d-1} \|f\|_{p-d+1}^{p-d+1}$. This is immediate from Lyapunov's inequality¹, which implies that

$$\|f\|_{b_i}^{b_i} \leq \|f\|_1^{\frac{p-d+1-b_i}{p-d}} \|f\|_{p-d+1}^{\frac{(p-d+1)(b_i-1)}{p-d}}.$$

Therefore,

$$\prod_{l=1}^{d} \|f\|_{b_{i}}^{b_{i}} \leq \|f\|_{1}^{\frac{(p-d+1)d-p}{p-d}} \|f\|_{p-d+1}^{\frac{(p-d+1)(p-d)}{p-d}} = \|f\|_{1}^{d-1} \|f\|_{p-d+1}^{p-d+1},$$

concluding the proof.

Now let us prove the bound in Lemma 3.5, copied here for readers' convenience: for $1 \le d \le p-1$

(41)
$$\sum_{(v_1^p \text{ is } [p,d])} f(v_1) \dots f(v_p)$$

$$\leq \binom{p}{d} \sum_{m=0}^{p-d} \binom{p-d}{m} (q^d-1)^{p-d-m} q^{nd} f(0)^m ||f||_1^{d-1} ||f||_{p-d-m+1}^{p-d-m+1}.$$

 $^{^{1} \}text{The Lyapunov inequality [18, Thm.17] states that if } 0 < r < s < t \text{, then } \|f\|_{s}^{s} \leq \|f\|_{r}^{r\frac{t-s}{t-r}} \|f\|_{t}^{t\frac{s-r}{t-r}}$

Proof. (of (41)) Any [p,d]-tuple can be written as a permutation of p vectors, of which the first d are linearly independent and the remaining ones are their linear combinations. Let $\mathcal{V}_d := \{v_1, \dots, v_d\}$ and let $\langle \mathcal{V}_d \rangle$ be its linear span. We have

$$\sum_{(v_1^p \text{ is } [p,d])} f(v_1) \dots f(v_p) \\
\leq \binom{p}{d} \sum_{(v_1^p \text{ is } [p,d])} f(v_1) \dots f(v_d) \sum_{v_{d+1}, \dots, v_p \in \langle \mathcal{V}_d \rangle} f(v_{d+1}) \dots f(v_p) \\
\leq \binom{p}{d} \sum_{v_1, \dots, v_d \in \mathbb{F}_q^n} f(v_1) \dots f(v_d) \sum_{v_{d+1}, \dots, v_p \in \langle \mathcal{V}_d \rangle} f(v_{d+1}) \dots f(v_p).$$

Note that $(v_i)_{d+1}^n$ are linear combinations of $(v_i)_i^d$. Let $G = [v_1, \dots, v_d]$. Then for $j \in [d+1, n]$, we can write $v_j = Gx_j$ for some $x_j \in \mathbb{F}_q^d$. This implies

$$\sum_{v_{d+1},\dots,v_p\in\langle\mathcal{V}_d\rangle} f(v_{d+1})\dots f(v_p) = \sum_{x_{d+1},\dots,x_p\in\mathbb{F}_q^d} f(Gx_{d+1})\dots f(Gv_p).$$

Therefore,

$$\begin{split} \sum_{v_1, \dots, v_d} f(v_1) \dots f(v_d) & \sum_{v_{d+1}, \dots, v_p \in \langle \mathcal{Y}_d \rangle} f(v_{d+1}) \dots f(v_p) \\ &= \sum_{v_1, \dots, v_d} f(v_1) \dots f(v_d) \sum_{x_{d+1}, \dots, x_p \in \mathbb{F}_q^d} f(Gx_{d+1}) \dots f(Gx_p) \\ &= \sum_{x_{d+1}, \dots, x_p \in \mathbb{F}_q^d} \sum_{v_1, \dots, v_d} f(v_1) \dots f(v_d) f(Gx_{d+1}) \dots f(Gx_p). \end{split}$$

Now consider the inner sum $\sum_{v_1,\dots,v_d} f(v_1)\dots f(v_d) f(Gx_{d+1})\dots f(Gx_p)$, for a fixed choice of x_{d+1},\dots,x_p , where m variables in $\{x_{d+1},\dots,x_p\}$ are set to 0. Without loss of generality, we may assume that $x_i=0$ for $i\in[d+1,d+m]$. In this case,

$$\sum_{v_1,\dots,v_d} f(v_1) \dots f(v_d) f(Gx_{d+1}) \dots f(Gx_p)$$

$$= f(0)^m \sum_{v_1,\dots,v_d} f(v_1) \dots f(v_d) f(Gx_{d+m+1}) \dots f(Gx_p)$$

$$\leq q^{nd} f(0)^m ||f||_1^{d-m-1} ||f||_{p-d+1}^{p-d+1} \quad \text{(from Lemma A.2)}.$$

Hence,

$$\sum_{v_1,\dots,v_d} f(v_1) \dots f(v_d) \sum_{\substack{v_{d+1},\dots,v_p \\ \in span(v_1,\dots,v_d)}} f(v_{d+1}) \dots f(v_p)$$

$$= \sum_{x_{d+1},\dots,x_p \in \mathbb{F}_q^d} \sum_{v_1,\dots,v_d} f(v_1) \dots f(v_d) f(Gx_{d+1}) \dots f(Gx_p)$$

$$\leq \sum_{m=0}^{p-d} \binom{p-d}{m} (q^d-1)^{p-d-m} q^{nd} f(0)^m ||f||_1^{d-1} ||f||_{p-d-m+1}^{p-d-m+1},$$

completing the proof.

REFERENCES

- [1] N. Alon, M. Dietzfelbinger, P. B. Miltersen, E. Petrank, and G. Tardos. Linear hash functions. *Journal of the ACM (JACM)*, 46(5):667–683, 1999.
- [2] S. Arimoto. On the converse to the coding theorem for discrete memoryless channels. *IEEE Transactions on Information Theory*, 19(3):357–359, 1973.
- [3] C. H. Bennett, G. Brassard, C. Crépeau, and U. M. Maurer. Generalized privacy amplification. *IEEE Transactions on Information Theory*, 41(6):1915–1923, 1995.
- [4] M. R. Bloch and J. N. Laneman. Strong secrecy from channel resolvability. *IEEE Transactions on Information Theory*, 59(12):8077–8098, 2013.
- [5] Z. Brakerski, V. Lyubashevsky, V. Vaikuntanathan, and D. Wichs. Worst-case hardness for LPN and cryptographic hashing via code smoothing. In *Annual International Conference on the Theory and Applications of Cryptographic Techniques*, pages 619–635. Springer, 2019.
- [6] C. Cachin. Entropy measures and unconditional security in cryptography. PhD thesis, ETH Zurich, 1997.
- [7] R. Canetti, S. Halevi, and M. Steiner. Mitigating dictionary attacks on password-protected local storage. In *Advances in Cryptology-CRYPTO 2006: 26th Annual International Cryptology Conference, Santa Barbara, California, USA, August 20-24, 2006. Proceedings 26*, pages 160–179. Springer, 2006.
- [8] J. L. Carter and M. N. Wegman. Universal classes of hash functions. In *Proceedings of the ninth annual ACM symposium on Theory of computing*, pages 106–112, 1977.
- [9] R. A. Chou, M. R. Bloch, and E. Abbe. Polar coding for secret-key generation. *IEEE Transactions on Information Theory*, 61(11):6213–6237, 2015.
- [10] T. M. Cover and H. H. Permuter. Capacity of coordinated actions. In 2007 IEEE International Symposium on Information Theory, pages 2701–2705. IEEE, 2007.
- [11] T. Debris-Alazard, L. Ducas, N. Resch, and J.-P. Tillich. Smoothing codes and lattices: Systematic study and new bounds. *IEEE Transactions on Information Theory*, 69(9):6006–6027, 2023.
- [12] T. Debris-Alazard and N. Resch. Worst and average case hardness of decoding via smoothing bounds. *Cryptology ePrint Archive*, 2022.
- [13] M. Dhar and Z. Dvir. Linear hashing with l_{∞} guarantees and two-sided Kakeya bounds. In 2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS), pages 419–428. IEEE, 2022.
- [14] Y. Dodis, S. J. Ong, M. Prabhakaran, and A. Sahai. On the (im)possibility of cryptography with imperfect randomness. In 45th Annual IEEE Symposium on Foundations of Computer Science, pages 196–205. IEEE, 2004.
- [15] R. A. Fisher, F. Yates, et al. *Statistical tables for biological, agricultural and medical research*. Edinburgh: Oliver and Boyd, 1963
- [16] V. Guruswami and J. Mosheiff. Punctured low-bias codes behave like random linear codes. In 2022 IEEE 63rd Annual Symposium on Foundations of Computer Science (FOCS), pages 36–45. IEEE, 2022.
- [17] T. S. Han and S. Verdú. Approximation theory of output statistics. IEEE Transactions on Information Theory, 39(3):752–772, 1993.
- [18] G. H. Hardy, J. E. Littlewood, and G. Pólya. *Inequalities*. Cambridge University Press, 1952.
- [19] J. Håstad, R. Impagliazzo, L. A. Levin, and M. Luby. A pseudorandom generator from any one-way function. SIAM Journal on Computing, 28(4):1364–1396, 1999.
- [20] M. Hayashi. General nonasymptotic and asymptotic formulas in channel resolvability and identification capacity and their application to the wiretap channel. *IEEE Transactions on Information Theory*, 52(4):1562–1575, 2006.
- [21] M. Hayashi and V. Y. Tan. Equivocations, exponents, and second-order coding rates under various Rényi information measures. *IEEE Transactions on Information Theory*, 63(2):975–1005, 2016.
- [22] J. Hązła, A. Samorodnitsky, and O. Sberlo. On codes decoding a constant fraction of errors on the BSC. In *Proceedings of the 53rd Annual ACM SIGACT Symposium on Theory of Computing*, pages 1479–1488, 2021.
- [23] R. Impagliazzo, L. A. Levin, and M. Luby. Pseudo-random generation from one-way functions. In *Proceedings of the Twenty-First Annual ACM Symposium on Theory of Computing*, pages 12–24, 1989.
- [24] D. R. Karger. Global min-cuts in RNC, and other ramifications of a simple min-cut algorithm. In *Soda*, volume 93, pages 21–30. Citeseer, 1993.
- [25] I. Kaslasi, R. D. Rothblum, and P. N. Vasudevanr. Public-coin statistical zero-knowledge batch verification against malicious verifiers. In Annual International Conference on the Theory and Applications of Cryptographic Techniques, pages 219–246. Springer, 2021.
- [26] L. Luzzi, C. Ling, and M. R. Bloch. Optimal rate-limited secret key generation from Gaussian sources using lattices. *IEEE Transactions on Information Theory*, 69(8):4944–4960, 2023.

- [27] D. Micciancio and O. Regev. Worst-case to average-case reductions based on Gaussian measures. SIAM Journal on Computing, 37(1):267–302, 2007.
- [28] G. L. Miller. Riemann's hypothesis and tests for primality. In *Proceedings of the Seventh Annual ACM Symposium on Theory of Computing*, pages 234–239, 1975.
- [29] J. Mosheiff, N. Resch, N. Ron-Zewi, S. Silas, and M. Wootters. LDPC codes achieve list decoding capacity. In 2020 IEEE 61st Annual Symposium on Foundations of Computer Science (FOCS), pages 458–469. IEEE, 2020.
- [30] N. Nisan. Extracting randomness: how and why. a survey. Proceedings of Computational Complexity (Formerly Structure in Complexity Theory), pages 44–58, 1996.
- [31] M. Pathegama and A. Barg. Smoothing of binary codes, uniform distributions, and applications. *Entropy*, 25(11):1515, 2023.
- [32] Y. Polyanskiy and S. Verdú. Arimoto channel coding converse and Rényi divergence. In 2010 48th Annual Allerton Conference on Communication, Control, and Computing, pages 1327–1333. IEEE, 2010.
- [33] M. O. Rabin. Probabilistic algorithm for testing primality. Journal of Nnumber Theory, 12(1):128–138, 1980.
- [34] A. Rao and O. Sprumont. A criterion for decoding on the binary symmetric channel. *Advances in Mathematics of Communications*, pages 0–0, 2024.
- [35] B. C. Rennie and A. J. Dobson. On Stirling numbers of the second kind. *Journal of Combinatorial Theory*, 7(2):116–121, 1969.
- [36] R. L. Rivest and A. T. Sherman. Randomized encryption techniques. In Advances in Cryptology: Proceedings of Crypto 82, pages 145–163. Springer, 1983.
- [37] H. D. Ruderman. Two new inequalities. The American Mathematical Monthly, 59(1):29–32, 1952.
- [38] A. Samorodnitsky. On the entropy of a noisy function. IEEE Transactions on Information Theory, 62(10):5446–5464, 2016.
- [39] A. Samorodnitsky. An upper bound on ℓ_q norms of noisy functions. *IEEE Transactions on Information Theory*, 66(2):742–748, 2019.
- [40] B. Simon. Real analysis: A comprehensive course in analysis, Part 1. American Mathematical Society, 2015.
- [41] M. Skórski. Shannon entropy versus Rényi entropy from a cryptographic viewpoint. In *IMA International Conference on Cryptography and Coding*, pages 257–274. Springer, 2015.
- [42] V. Y. Tan and M. Hayashi. Analysis of remaining uncertainties and exponents under various conditional Rényi entropies. *IEEE Transactions on Information Theory*, 64(5):3734–3755, 2018.
- [43] H. Tyagi and S. Watanabe. Information-theoretic Cryptography. Cambridge University Press, 2023.
- [44] S. Vembu and S. Verdú. Generating random bits from an arbitrary source: Fundamental limits. IEEE Transactions on Information Theory, 41(5):1322–1332, 1995.
- [45] B. Waters. Dual system encryption: Realizing fully secure IBE and HIBE under simple assumptions. In *Annual International Cryptology Conference*, pages 619–636. Springer, 2009.
- [46] L. Yu and V. Y. Tan. Simulation of random variables under Rényi divergence measures of all orders. *IEEE Transactions on Information Theory*, 65(6):3349–3383, 2019.