EKEDAHL-OORT STRATA AND THE SUPERSINGULAR LOCUS IN THE GU(q - 2, 2) SHIMURA VARIETY

EMERALD ANNE, DEEWANG BHAMIDIPATI, MARIA FOX, HEIDI GOODSON, STEVEN GROEN, AND SANDRA NAIR

ABSTRACT. This paper concerns the characteristic-p fibers of GU(q - 2, 2) Shimura varieties, which classify abelian varieties with additional structure. These Shimura varieties admit two stratifications of interest: the Ekedahl-Oort stratification, based on the isomorphism class of the p-torsion subgroup scheme, and the Newton stratification, based on the isogeny class of the p-divisible group. It is natural to ask which Ekedahl-Oort strata intersect the unique closed Newton stratum, called the *supersingular locus*. In this paper, we present several novel techniques that give information about the interaction between the two stratifications for general signature (q - 2, 2), and as an application, we completely answer this question for the signature (3, 2).

1. INTRODUCTION

This paper studies the characteristic-*p* fibers of Shimura varieties of PEL type. Shimura varieties of PEL type are moduli spaces of abelian varieties with additional structure, and this moduli interpretation gives rise to two stratifications of the characteristic-*p* fiber: the Newton stratification and the Ekedahl-Oort stratification. The Ekedahl-Oort stratification is based on the isomorphism class of the *p*-torsion subgroup scheme of the parameterized abelian varieties, while the Newton stratification is based on the isogeny class of their *p*-divisible group. Though each stratification is compelling in its own right, it is also interesting to study the interactions between the two stratifications. For instance, the two stratifications coincide on the modular curve, both distinguishing between *ordinary* and *supersingular* elliptic curves. However, when studying abelian varieties of increasing dimension, the stratifications diverge further away from one another.

The interaction between the Ekedahl-Oort and Newton stratifications on general Shimura varieties of PEL type has been widely studied (see, for example, the presentation of results in [VW13]). More precise statements can be made when one restricts attention to a smaller class of Shimura varieties. In this paper, we study the interaction between these stratifications for *unitary Shimura varieties* $\mathcal{M}(q - a, a)$. These are moduli spaces of abelian varieties of dimension q with an action of an imaginary quadratic field K that meets the "signature (q - a, a)" condition. In [Woo16], Wooding studies several aspects of the Ekedahl-Oort and Newton stratifications of $\mathcal{M}(q - a, a)$. However, the following question is still open in general.

Question 1.1. Which Ekedahl-Oort strata of $\mathcal{M}(q - a, a)$ intersect the supersingular locus?

The answer to this question is known for some signatures. For example, in signature (q, 0) the Shimura variety is zero-dimensional, consisting of a single Ekedahl-Oort stratum. By a classical theorem of Deuring, this stratum coincides with the supersingular locus if and only if p does not split in K. The signature (q - 1, 1) case is already much more complex, but it is established in [VW11] which of the Ekedahl-Oort strata intersect the supersingular locus. Following from the results of [HP14], it is known which Ekedahl-Oort strata of $\mathcal{M}(2, 2)$ intersect the supersingular locus. It is known ([GH15, Theorem A]) that these are the only signatures for which the supersingular locus is a union of Ekedahl-Oort strata, and so answering Question 1.1 should ostensibly be significantly more complicated for Shimura varieties of other signatures.

This paper focuses on unitary Shimura varieties of signature (q - 2, 2) at a prime inert in the relevant quadratic imaginary field. The geometry of the supersingular locus of $\mathcal{M}(q - 2, 2)$ is described in [FHI23], but little is known about which Ekedahl-Oort strata intersect the supersingular locus. In the tradition of [VW11], such information may be useful when studying more subtle aspects of the supersingular locus. We

present several techniques that reveal information about the interaction between the Ekedahl-Oort stratification and the Newton stratification. As an application of these general methods, we give a complete answer to Question 1.1 when the signature is (3,2).

Theorem A (Theorem 6.7). Four of the Ekedahl-Oort strata of $\mathcal{M}(3,2)$ do not intersect the supersingular locus. The remaining six Ekedahl-Oort strata of $\mathcal{M}(3,2)$ do intersect the supersingular locus. Furthermore, of these six, two strata are completely contained in the supersingular locus and one intersects but is not contained in the supersingular locus.

The precise statement of Theorem 6.7 identifies each relevant Ekedahl-Oort stratum in terms of the corresponding *p*-torsion group scheme, using the notation that we develop in Section 3.

The structure of our paper is as follows. We first recall some relevant notation and background information in Section 2 that will be used throughout the paper. In Section 3, we study the basic properties of the Ekedahl-Oort stratification. Using results of Moonen (see [Moo01, Theorem 6.7]), we index the Ekedahl-Oort strata of $\mathcal{M}(q - 2, 2)$ by certain Weyl group cosets and, via Dieudonné theory, concretely describe the *p*-torsion subgroup of the corresponding Ekedahl-Oort stratum. In Section 3.4, we study the topological closure relations among the strata and prove that a number of topological closure relations hold (see Theorems 3.9 and 3.15).

In Sections 4 and 5, we obtain information about the Ekedahl-Oort stratification of $\mathcal{M}(q-2,2)$ by leveraging its relation to Shimura varieties whose Ekedahl-Oort stratifications are better understood. First, in Section 4, we relate $\mathcal{M}(q-2,2)$ to simpler unitary Shimura varieties via the natural product maps

$$\mathcal{M}(a,b) \times \mathcal{M}(c,d) \to \mathcal{M}(q-2,2),$$

induced by taking products of the parameterized abelian varieties. In Theorems 4.10 and 4.16, we explicitly describe the effect of these product maps on the Ekedahl-Oort stratifications.

In Section 5, we relate $\mathcal{M}(q-2,2)$ to the Siegel modular variety via a forgetful map

$$\mathcal{M}(q-2,2) \to \mathcal{A}_q,$$

which "forgets" the unitary structure of $\mathcal{M}(q-2,2)$.

More precisely, in Theorem 5.2, we concretely describe the effect of this map on the Ekedahl-Oort stratifications.

By taking advantage of prior results on the interaction between the Ekedahl-Oort and Newton stratifications of these better-understood Shimura varieties, the techniques in Sections 4 and 5 both yield information regarding the interaction between the Ekedahl-Oort stratification and the supersingular locus in signature (q - 2, 2).

In Section 6, we apply the amassed tools to the unitary Shimura variety of signature (3, 2) to prove Theorem A. For all but two of the Ekedahl-Oort strata of $\mathcal{M}(3, 2)$, the results in Sections 4 and 5 can be used to detect whether or not these strata intersect the supersingular locus. We prove that the remaining two strata intersect the supersingular locus by explicitly constructing a point in the intersection. Our classification result for signature (3, 2) is summarized in Table 3.

Acknowledgements. This project started at the Rethinking Number Theory workshop in June 2022. We want to thank the organizers for the workshop, and are grateful for the supportive, collaborative research environment this workshop provided. The workshop was supported by the Number Theory Foundation, the American Institute of Mathematics, and the University of Wisconsin-Eau Claire. Additionally, we would like to thank Rachel Pries, Damiano Testa, and Martin Weissman for useful conversations. We would also like to thank Satyam Patel for programming assistance and Mohamed Tawfik for his early contributions to the project.

H.G. was supported by NSF grant DMS-2201085 and a PSC-CUNY Award, jointly funded by The Professional Staff Congress and The City University of New York. M.F. was supported in part by NSF MSPRF Grant 2103150.

2. BACKGROUND

2.1. **Unitary Shimura varieties.** Unitary Shimura varieties are moduli spaces of abelian varieties equipped with extra structure, including an action of a quadratic imaginary field. To define an integral model of such a Shimura variety, we fix a prime p > 2, a positive integer q, non-negative integers a and b such that a + b = q, and a quadratic imaginary field K. We further assume that the prime p is inert in K, and so we identify $\mathcal{O}_K/(p)$ as \mathbb{F}_{p^2} throughout.

Definition 2.1. We use the **PEL datum** $(K, \mathcal{O}_K \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}, *, V, (\cdot, \cdot), \Lambda, \mathbf{G}, h)$ of Kottwitz [Kot92], defined as follows:

- *K* is the quadratic imaginary field introduced above, with * being the nontrivial automorphism of *K* over Q.
- *V* is a *K*-vector space of dimension *q*, equipped with a perfect alternating \mathbb{Q} -bilinear pairing (\cdot, \cdot) : $V \times V \to \mathbb{Q}$ such that $(xv, w) = (v, x^*w)$ for all $x \in K$ and $v, w \in V$.
- **G** is the algebraic group of *K*-linear symplectic similitudes of $(V, (\cdot, \cdot))$. We assume that **G**_R is isomorphic to the real algebraic group GU(a, b).
- Λ is an $\mathcal{O}_K \otimes_\mathbb{Z} \mathbb{Z}_{(p)}$ -invariant lattice of $V \otimes_\mathbb{Q} \mathbb{Q}_p$ such that the alternating form induced by (\cdot, \cdot) is a perfect \mathbb{Z}_p -form.
- $h : \operatorname{Res}_{\mathbb{C}/\mathbb{R}}(\mathbb{G}_{m,\mathbb{C}}) \to \mathbb{G}_{\mathbb{R}}$ is the homomorphism of real algebraic groups that maps $z \in \mathbb{C}^{\times}$ to $\operatorname{diag}(z^{a}, \overline{z}^{b})$.

Let *L* be the reflex field associated to the PEL-datum $(K, \mathcal{O}_K \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}, *, V, (\cdot, \cdot), \Lambda, \mathbf{G}, h)$; if a = b then $L = \mathbb{Q}$, and L = K otherwise. Let \mathbb{A}_f^p denote the ring of finite adeles with a trivial component at p. Fix a compact open subgroup $C^p \subset \mathbf{G}(\mathbb{A}_f^p)$. For C^p small enough, the construction of Kottwitz [Kot92] attaches to this PEL datum a smooth, quasi-projective scheme $\mathbf{M}(a, b)_{C^p}$ over $\operatorname{Spec}(\mathcal{O}_{L,(p)})$ with the following moduli interpretation.

Let *S* be an $\mathcal{O}_{L,(p)}$ -scheme. Then the set $\mathbf{M}(a, b)_{C^p}(S)$ parameterizes isomorphism classes of tuples (A, ι, λ, ξ) , where:

- *A* is an abelian variety over *S* of dimension *q*.
- $\iota : \mathcal{O}_K \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)} \to \operatorname{End}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ is a nonzero homomorphism of $\mathbb{Z}_{(p)}$ -algebras such that the Rosati involution on $\operatorname{End}(A) \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ induces the involution * on $\mathcal{O}_K \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$.
- λ is a one-dimensional Q-subspace of Hom $(A, A^{\vee}) \otimes_{\mathbb{Z}} \mathbb{Q}$ that contains a *p*-principal $\mathcal{O}_K \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ -linear polarization.
- $\xi : H_1(A, \mathbb{A}_f^p) \to V \otimes_{\mathbb{Q}} \mathbb{A}_f^p \mod C^p$ is a C^p -level structure.

We also require that (A, ι) meets Kottwitz's determinant condition of signature (a, b). Two tuples (A, ι, λ, ξ) and $(A', \iota', \lambda', \xi')$ are isomorphic if there exists a prime-to-*p* isogeny from *A* to *A'*, commuting with the action of $\mathcal{O}_K \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$, mapping ξ to ξ' and λ to λ' .

The integral model $\mathbf{M}(a, b)_{C^p}$ has relative dimension of ab. The main object of study for this paper is the characteristic p unitary Shimura variety, denoted by $\mathcal{M}(a, b)$, which is the fiber at p of $\mathbf{M}(a, b)_{C^p}$. In particular, $\mathcal{M}(a, b)$ is defined over the residue field \mathbb{F} of L at p and is of dimension ab. Since $\mathbf{M}(a, b)_{C^p} \cong \mathbf{M}(b, a)_{C^p}$, we assume without loss of generality that $0 \le b \le a$.

Frequently, important properties of $\mathcal{M}(a, b)$ can be understood in terms of its geometric points. For this reason, we fix an algebraic closure \Bbbk of \mathbb{F} .

2.2. Ekedahl-Oort and Newton Stratifications. In this section, we recall the definitions of the Ekedahl-Oort and Newton stratifications of $\mathcal{M}(a, b)$. For more details, see [VW13].

The Ekedahl-Oort stratification is based on the isomorphism class of the *p*-torsion group scheme of the parameterized abelian varieties. Two field-valued points (A, ι, λ, ξ) and $(A', \iota', \lambda', \xi')$ of $\mathcal{M}(a, b)$ are in the same **Ekedahl-Oort stratum** if and only if the *p*-torsion group schemes equipped with induced action and polarization, $(A[p], \iota, \lambda)$ and $(A'[p], \iota', \lambda')$, are isomorphic over k. The Ekedahl-Oort strata are locally closed, and the closure of each Ekedahl-Oort stratum is a union of Ekedahl-Oort strata.

The Newton stratification is based on the isogeny class of the *p*-divisible group of the parameterized abelian varieties. Two field-valued points (A, ι, λ, ξ) and $(A', \iota', \lambda', \xi')$ of $\mathcal{M}(a, b)$ are in the same **Newton stratum** if and only if the *p*-divisible groups equipped with induced action and polarization, $(A[p^{\infty}], \iota, \lambda)$ and $(A'[p^{\infty}], \iota', \lambda')$, are isogenous (in a way that respects the actions and polarizations) over \Bbbk .

The Newton strata are locally closed, and the closure of each Newton stratum is a union of Newton strata. The unique closed Newton stratum of $\mathcal{M}(a, b)$ is the supersingular locus, which we denote as $\mathcal{M}(a, b)^{ss}$. In particular, a point (A, ι, λ, ξ) of $\mathcal{M}(a, b)(\Bbbk)$ is contained in the supersingular locus if and only if A is a supersingular abelian variety.

2.3. **Weyl Group Cosets.** Results of [Moo01] relate the study of the Ekedahl-Oort strata to cosets in a certain Weyl group. This section introduces the relevant Weyl group cosets and their minimal-length coset representatives.

The Weyl group that is relevant for the study of $\mathcal{M}(a, b)$ is $W = \mathfrak{S}_q$, the symmetric group on q elements. We consider W as a Coxeter group with a set of **simple reflections**

$$S = \{s_1, \ldots, s_{q-1}\}$$
, where $s_i = (i, i+1)$.

The **length** of $w \in W$, denoted $\ell(w)$, is the length of a shortest expression for w as a product of simple reflections. It is proved in [BB06, Proposition 1.5.2] that the length of an element w in \mathfrak{S}_q can be computed as the number of inversions, i.e., the cardinality of the set

(2.3.1) $Inv(w) := \{(i, j) \mid i < j \text{ and } w(i) > w(j)\}.$

In particular, *W* has a unique element w_0 of maximal length, where $w_0(k) = q + 1 - k$.

For $J \subset S$, let W_J denote the subgroup of W generated by J. Note that W_J is a parabolic subgroup of W, and it follows from Proposition 2.4.4 of [BB06] that every coset of $W_J \setminus W$ contains a unique minimal-length coset representative. Let JW be the collection of such minimal-length coset representatives for $W_J \setminus W$.

For the subset $J_{(a,b)} = \{s_1, ..., s_{q-1}\} \setminus \{s_b\}$ of *S*, we let $W_{(a,b)} := W_{J_{(a,b)}}$, and $\mathbf{W}(a,b) := J_{(a,b)}W$. The following theorem is paraphrased from Theorem 6.7 of [Moo01]:

Theorem 2.2 (Moonen). *There is a bijection of sets:*

{*Ekedahl-Oort Strata of* $\mathcal{M}(a, b)$ } \longleftrightarrow **W**(a, b).

We now recall some properties of Moonen's construction. (See also [Woo16].) Let *G* be the group of $\mathcal{O}_K \otimes_{\mathbb{Z}} \mathbb{Z}_{(p)}$ -linear symplectic similitudes of Λ . Then *G* is a group scheme over \mathbb{Z}_p , and we let \overline{G} be its special fiber. Moonen [Moo01] gives an explicit identification of the Ekedahl-Oort strata with $W_X \setminus W_{\overline{G}}$, for a certain subgroup W_X depending on the signature. Concretely, $W_X \setminus W_{\overline{G}}$ can be described as

$$\{(w_1, w_2) \in \mathbf{W}(a, b) \times \mathbf{W}(b, a) \mid w_2 = w_0 w_1 w_0\}$$

As the map $\pi \mapsto w_0 \pi w_0$ gives an isomorphism between W(a, b) and W(b, a), one has $W_X \setminus W_{\overline{G}} \cong W(a, b)$ which agrees with our statement of Theorem 2.2.

For any $w \in \mathbf{W}(a, b)$, let $\mathcal{M}(a, b)_w$ denote the corresponding Ekedahl-Oort stratum and $(G_w, \iota_w, \lambda_w)$ denote the corresponding *p*-torsion group scheme. One observes that since G_w is a *p*-torsion group scheme, the

action of \mathcal{O}_K on G_w , via ι_w , factors through $\mathcal{O}_K/(p)$. Since p is assumed to be inert, $\mathcal{O}_K/(p) = \mathbb{F}_{p^2}$, and we abuse notation and refer to this induced action of \mathbb{F}_{p^2} as ι_w as well. Moonen uses (contravariant) Dieudonné theory to describe $(G_w, \iota_w, \lambda_w)$. For each $w \in \mathbf{W}(a, b)$, he constructs the **standard object** (N_w, F, V) . This is the Dieudonné module of G_w , consisting of a vector space N_w of dimension 2q over \Bbbk , F a Frob \Bbbk -semilinear operator on N_w , and V a Frob \Bbbk^{-1} -semilinear operator on N_w , described explicitly on a basis. The action ι_w is recorded by a splitting $N_w = N_{w,1} \oplus N_{w,2}$. By Theorem 6.7 of [Moo01], λ_w is uniquely determined by (G_w, ι_w) , so it is unnecessary to record the corresponding polarization of N_w .

We now outline the inverse of the assignment $\omega \mapsto N_{\omega}$. Given a Dieudonné module (N, F, V) with an \mathbb{F}_{v^2} -action, one constructs a *final filtration*

$$0 \subset W_1 \subset W_2 \subset \cdots \subset W_{2q} = N$$

that is stable under *F* and *V*⁻¹ and has the property dim_k(W_i) = *i*.

The action of \mathbb{F}_{p^2} on N induces a decomposition $N = N_1 \oplus N_2$. Intersecting the filtration W_{\bullet} of N with the subspace N_i (for i = 1, 2) gives:

$$0 \subset C_{i,1} \subset C_{i,2} \subset \cdots \subset C_{i,q} = N_i.$$

From here we define the functions

$$\eta_i(j) = \dim(C_{i,j} \cap N[F]).$$

We focus on η_1 , as η_2 is determined by η_1 via the Rosati involution condition. Since

$$\eta_1(q) = \dim(N_1[F]) = b,$$

there are *b* indices where η_1 jumps (meaning $\eta_1(j) = \eta_1(j-1) + 1$). We denote these integers by $1 \le j_1 \le \ldots \le j_b \le q$ and the remaining integers by $1 \le i_1 \le \ldots i_a \le q$. The permutation $\omega_N \in \mathbf{W}(a, b)$ is

(2.3.2)
$$\omega_N(j_l) = l \text{ and } \omega_N(i_m) = b + m.$$

By construction, *N* is the standard object corresponding to the permutation ω_N .

In Section 3, we give more explicit descriptions in signature (q - 2, 2) of the minimal length coset representatives of $W_I \setminus W$ and the standard objects.

3. STRUCTURE OF EKEDAHL-OORT STRATA FOR SIGNATURE (q - 2, 2)

The goal of this section is to index the Ekedahl-Oort strata of $\mathcal{M}(q-2,2)$ and study the topological closure relations between them. First, we explicitly describe the elements of the set W(q-2,2). Following that, we recall the Bruhat partial ordering, and its generalization due to He [He07] that corresponds to topological closure relations between the Ekedahl-Oort strata. Finally, we present some results on the closure relations between the elements of W(q-2,2).

3.1. Index Set. In this section, we explicitly describe the set W(q - 2, 2).

For $1 \leq u < v \leq q$ we define the following element of *W*

(3.1.1)
$$\gamma_{u,v} := (2, 3, \dots, v)(1, 2, \dots, u).$$

We note that $\gamma_{u,v}$ fixes every element in the set $\{j \mid v < j \leq q\}$ and

(3.1.2)
$$\gamma_{u,v}(j) = \begin{cases} j+2 & \text{if } 1 \leq j \leq u-1, \\ 1 & \text{if } j = u, \\ j+1 & \text{if } u+1 \leq j \leq v-1, \\ 2 & \text{if } j = v. \end{cases}$$

Lemma 3.1. $W(q-2,2) = \{\gamma_{u,v} \mid 1 \le u < v \le q\}$ with $\ell(\gamma_{u,v}) = u + v - 3$.

Proof. Recall that the length satisfies $\ell(\gamma_{u,v}) = \#\text{Inv}(\gamma_{u,v})$ where $\text{Inv}(\gamma_{u,v})$ is the set of inversions, as defined in Equation (2.3.1). It is apparent from the description of $\gamma_{u,v}$ as a function in Equation (3.1.2) that (understood appropriately for u = 1)

$$Inv(\gamma_{u,v}) = \{(j, u), (j, v), (u+k, v) \mid 1 \le j \le u-1, 1 \le k \le v-u-1\}.$$

Thus, $\ell(\gamma_{u,v}) = u + v - 3$. Further, we note that for $\sigma \in W_{(q-2,2)}$ we obtain $\text{Inv}(\gamma_{u,v}) \subseteq \text{Inv}(\sigma\gamma_{u,v})$ and thus $\ell(\gamma_{u,v})$ is minimal in the coset $W_{(q-2,2)}\gamma_{u,v}$.

Lastly, to prove that any two distinct $\gamma_{u_1,v_1}, \gamma_{u_2,v_2}$ lie in distinct cosets one can verify that $\gamma_{u_2,v_2}\gamma_{u_1,v_1}^{-1}$ is not an element of $W_{(q-2,2)}$.

Remark 3.2. Each $\gamma_{u,v} \in W(q-2,2)$ corresponds to an Ekedahl-Oort stratum labelled $\mathcal{M}(q-2,2)_{\gamma_{u,v}}$. By [Moo01], the dimension of $\mathcal{M}(q-2,2)_{\gamma_{u,v}}$ is equal to $\ell(\gamma_{u,v})$. Let n_d be the number of Ekedahl-Oort strata there are of dimension $0 \leq d \leq 2(q-2)$, i.e.,

$$n_d = \# \{ \gamma_{u,v} \mid \ell(\gamma_{u,v}) = d \}.$$

Since $\ell(\gamma_{u,v}) = u + v - 3$, we have that

$$n_d = \begin{cases} \lfloor d/2 \rfloor + 1 & \text{if } d \leq q-2, \\ \lfloor d/2 \rfloor + 1 - (d - (q-2)) & \text{if } d > q-2. \end{cases}$$

3.2. **Standard Objects.** In this section, we describe the standard objects of the Ekedahl-Oort strata of $\mathcal{M}(q - 2, 2)$. These standard objects are mod-*p* Dieudonné modules, and by Dieudonné theory each determines a corresponding *p*-torsion group scheme.

Lemma 3.3. Let $(N_{\gamma_{u,v}}, F, V)$ be the standard object corresponding to $\gamma_{u,v}$. Then,

$$N_{\gamma_{u,v}} = \operatorname{Span}_{\Bbbk} \{e_{i,j}\},$$

where $1 \leq i \leq 2$ and $1 \leq j \leq q$. Further, F is the $\operatorname{Frob}_{\Bbbk}$ -semilinear operator on $N_{\gamma_{u,v}}$ and V is the $\operatorname{Frob}_{\Bbbk}^{-1}$ -semilinear operator on $N_{\gamma_{u,v}}$ extended from the following action on the basis:

$$F(e_{1,j}) = \begin{cases} 0 & \text{if } j = u, j = v \\ e_{2,j} & \text{if } 1 \leq j \leq u - 1 \\ e_{2,j-1} & \text{if } u < j \leq v - 1 \\ e_{2,j-2} & \text{if } j > v \end{cases} \qquad V(e_{1,j}) = \begin{cases} 0 & \text{if } j = 1, 2 \\ e_{2,j-2} & \text{if } 2 < j \leq q - v + 2 \\ e_{2,j-1} & \text{if } q - v + 2 < j \leq q - u + 1 \\ e_{2,j} & \text{if } j > q - u + 1 \end{cases}$$
$$F(e_{2,j}) = \begin{cases} e_{1,1} & \text{if } j = q - v + 1 \\ e_{1,2} & \text{if } j = q - u + 1 \\ 0 & \text{otherwise} \end{cases} \qquad V(e_{2,j}) = \begin{cases} e_{1,u} & \text{if } j = q - 1 \\ e_{1,v} & \text{if } j = q \\ 0 & \text{otherwise.} \end{cases}$$

Proof. We use the explicit description of $\gamma_{u,v}$ to unwind the description of the corresponding standard object given in [Moo01, 4.9], under the assumption that the prime *p* is inert in the quadratic imaginary field *K*.

3.3. **Bruhat Order.** In this section we use the index set defined in Section 3.1 to give an explicit description of the Bruhat order on W(q - 2, 2). We first recall the definition of the Bruhat order on a Weyl group.

Definition 3.4 (The Bruhat order). Consider the Coxeter system (*W*, *S*). For *w* and *w'* in *W*, we say $w' \leq w$, if $\ell(w') \leq \ell(w)$ and there exists a sequence $w' = v_0, v_1, \ldots, v_m = w$ such that

- $\ell(v_{i-1}) \leq \ell(v_i)$; and
- $v_{i-1}^{-1}v_i$ is a reflection.

By [BB06, Proposition 2.5.1], the Bruhat order on W induces a partial order on W(q - 2, 2), also called the Bruhat order and denoted \leq . Using the notation in Section 3.1, we can completely characterize the Bruhat relations for the Weyl group cosets W(q - 2, 2) associated to $\mathcal{M}(q - 2, 2)$.

Proposition 3.5. Let γ_{u_1,v_1} and γ_{u_2,v_2} be elements of $\mathbf{W}(q-2,2)$. Then $\gamma_{u_1,v_1} \leq \gamma_{u_2,v_2}$ with respect to the Bruhat order if and only if $u_1 \leq u_2$ and $v_1 \leq v_2$.

Proof. For a permutation $x \in \mathfrak{S}_n$, define

 $x[i,j] := \#\{a \leq i \mid x(a) \geq j\}.$

Then $x \leq y$ with respect to the Bruhat order if and only if $x[i, j] \leq y[i, j]$ for all $i, j \leq n$ (see Theorem 2.1.5 of [BB06]). Using the definition of $\gamma_{u,v}$ and its inverse we see that

$$\gamma_{u,v}^{-1}[i,j] = \begin{cases} i-j+1 & \text{if } j < i \\ 1 & \text{if } j \leqslant u \text{ and } i = 1 \\ 2 & \text{if } j \leqslant u \text{ and } 1 < i \leqslant j \\ 0 & \text{if } u < j \leqslant v \text{ and } i = 1 \\ 1 & \text{if } u < j \leqslant v \text{ and } i = 1 \\ 1 & \text{if } u < j \leqslant v \text{ and } 1 < i \leqslant j \\ 0 & \text{if } j > v \text{ and } 1 < i \leqslant j \\ 1 & \text{if } j > v \text{ and } i < j \end{cases}$$

Comparing the values $\gamma_{u_1,v_1}^{-1}[i,j]$ and $\gamma_{u_2,v_2}^{-1}[i,j]$, we see that $\gamma_{u_1,v_1}^{-1} \leq \gamma_{u_2,v_2}^{-1}$ if and only if $u_1 \leq u_2$ and $v_1 \leq v_2$. Since $\gamma_{u_1,v_1}^{-1} \leq \gamma_{u_2,v_2}^{-1}$ is equivalent to $\gamma_{u_1,v_1} \leq \gamma_{u_2,v_2}$, the statement follows.

3.4. The Closure Order.

3.4.1. *Definition and properties.* There are two partial orders on W(q - 2, 2) that we are concerned with in this paper: the Bruhat order defined above, and a generalization of this due to He [He07] that we refer to as the Closure order. In this section we define the Closure order and discuss how both orders relate to Ekedahl-Oort strata. As in Section 3.3, we focus on W(q - 2, 2) but the constructions here can all be re-framed for more general W(a, b). We refer to [He07, PWZ11, VW13, Wed05, Woo16] for more information.

Recall that the Frobenius automorphism *F* of the algebraic group \overline{G} induces a $\psi \in Aut(W, S)$ of the Coxeter system: for any element $w \in W$, $\psi(w) = w_0 w w_0$, where w_0 is the unique maximal element in *W*.

As in Section 2.3, $W_{(q-2,2)}$ denotes the subgroup of the Weyl group W generated by $J_{(a,b)} = \{s_1, ..., s_{q-1}\} \setminus \{s_b\}$. For ease of notation, let $J := J_{(q-2,2)}$. The unique maximal element in $W_{(q-2,2)}$, with respect to the Bruhat order, is denoted by $w_{0,J}$. Let $W^{\psi(J)}$ be the set of elements $w \in W$ that have minimal length in their coset $wW_{\psi(J)}$, and let $J' := {}^{w_0}\psi(J)$ denote the set $w_0\psi(J)w_0$. In the notation of Section 2.3, $J'W^{\psi(J)}$ denotes the collection of minimal-length coset representatives for $W_{I'} \setminus W/\psi(J)$.

The definition of the Closure order involves the element *x* of minimal length in the double coset $W_{J'}w_0W_{\psi(J)}$. One can show that this element satisfies

$$x = w_{0,I'}w_0 = w_0w_{0,\psi(I)}$$

and so *x* is the unique maximal length element amongst the minimal length coset representatives $J'W^{\psi(J)}$.

We now define the **Closure order** corresponding to ψ on **W**(q – 2, 2).

Definition 3.6 (The Closure order \preccurlyeq). Let w, w' be elements of W(q - 2, 2). Then $w \preccurlyeq w'$ if there exists an $h \in W_{(q-2,2)}$ such that

$$hwx\psi(h^{-1})x^{-1} \leqslant w',$$

where \leq denotes the Bruhat order¹.

Using the fact that $\psi(h) = w_0 h w_0$, we have

$$x\psi(h^{-1})x^{-1} = (w_0w_{0,\psi(J)})(w_0h^{-1}w_0)(w_{0,\psi(J)}w_0) = w_{0,J}h^{-1}w_{0,J},$$

and so Inequality (3.4.1) can be rewritten as

$$(3.4.2) hww_{0,I}h^{-1}w_{0,I} \leqslant w'.$$

Note that the Bruhat order is the special case of the Closure order where Inequality (3.4.1) is satisfied for *h* equal to the identity in $W_{(q-2,2)}$.

In Section 3.4 we give some partial results on the Closure order on W(q - 2, 2). One difficulty that arises in giving a complete description of the Closure order on W(q - 2, 2) is that the expression $hww_{0,J}h^{-1}w_{0,J}$ is not a coset representative in W(q - 2, 2). In other words, the Bruhat relation that we are checking in Inequality (3.4.1) is in the Weyl group W and not in W(q - 2, 2), and so we are not able to apply our results on the Bruhat order in Section 3.3 to the Closure order.

The Closure order is of interest because it captures topological closure relations among the Ekedahl-Oort strata. Let $\overline{\mathcal{M}(q-2,2)}_{\gamma_{u',v'}}$ denote the topological closure of the stratum $\mathcal{M}(q-2,2)_{\gamma_{u',v'}}$. By [VW13, Theorem 1.2],

$$\overline{\mathscr{M}(q-2,2)}_{\gamma_{u',v'}} = \bigcup_{\gamma_{u,v} \preccurlyeq \gamma_{u',v'}} \mathscr{M}(q-2,2)_{\gamma_{u,v}}.$$

Thus, to understand closure relations among Ekedahl-Oort strata, it suffices to analyze the Closure order relations in W(q - 2, 2).

3.4.2. *Closure relations.* In this section, we prove several results that give a partial classification of the Closure order for signature (q - 2, 2). We begin with a lemma that we will use in order to prove the two sets of closure relations in Theorems 3.9 and 3.15.

Lemma 3.7. Let s_k with k > 2 denote the simple reflection (k, k+1). For $1 \le u < v \le q$, the following hold:

- $\gamma_{u,v+1}s_v = \gamma_{u,v}$ and $s_v\gamma_{u,v+1}^{-1} = \gamma_{u,v}^{-1}$,
- $\gamma_{u+1,v}s_u = \gamma_{u,v}$ and $s_u\gamma_{u+1,v}^{-1} = \gamma_{u,v}^{-1}$.

Furthermore, $s_{q-(j-1)}s_{q-(j-2)}\cdots s_{q-(j-i)}\gamma_{j+1,q+1-j+i}^{-1} = \gamma_{j+1,q+1-j}^{-1}$ for any $4 \le j \le \frac{q}{2}$ and $1 \le i \le j-3$.

Proof. We can prove these results using the non-disjoint cycle definition of $\gamma_{u,v}$:

 $\gamma_{u,v} = (2, 3, \dots, v - 1, v)(1, 2, \dots, u - 1, u).$

For example, to prove the first equality, note that since v > u, the reflection s_v commutes with (1, 2, ..., u - 1, u). Hence,

$$\begin{aligned} \gamma_{u,v+1}s_v &= (2,3,\ldots,v,v+1)(1,2,\ldots,u-1,u)(v,v+1) \\ &= (2,3,\ldots,v,v+1)(v,v+1)(1,2,\ldots,u-1,u) \\ &= (2,3,\ldots,v)(1,2,\ldots,u-1,u) \end{aligned}$$

The other relations in the bullet-points are proved in a similar manner, and the last result follows from applying these repeatedly. \Box

¹In the notation of [He07], Inequality (3.4.1) actually implies $wx \preccurlyeq w'x$. In our work, as in [VW13], we drop the *x* from the notation and simply write $w \preccurlyeq w'$.

We define an action on elements $w \in \mathbf{W}(q-2,2)$ by elements $h \in W_{(q-2,2)}$ by

$$h \bullet w := h w w_{0,l} h^{-1} w_{0,l}$$

which is the expression that appears on the left side of Inequality 3.4.2. In other words, $\gamma_{u_1,v_1} \preccurlyeq \gamma_{u_2,v_2}$ if there is an $h \in W_{(q-2,2)}$ such that the Bruhat relation $h \bullet \gamma_{u_1,v_1} \leqslant \gamma_{u_2,v_2}$ holds. In our work, we express h as a product of simple reflections, and the following lemma will be used throughout.

Lemma 3.8. For any simple reflection $s_k = (k, k+1)$ with k > 2,

$$w_{0,J}s_kw_{0,J} = s_{q+2-k}$$

Proof. Since $w_{0,I}$, defined as $w_{0,I}(x) = q + 3 - x$ for all $3 \le x \le q$, is of order two and k > 2, we have

$$w_{0,I}s_kw_{0,I} = (w_{0,I}(k), w_{0,I}(k+1)) = s_{q+2-k}.$$

The following theorem gives the first set of closure relations.

Theorem 3.9. Let $q \ge 5$. Then the following relations hold

- (1) $\gamma_{j+1,q+1-j} \preccurlyeq \gamma_{j,q+3-j}$ for $3 \leqslant j < q/2$,
- (2) $\gamma_{q-j,j+1} \preccurlyeq \gamma_{q-j+2,j}$ for $q/2 + 1 < j \leqslant q 1$.

Remark 3.10. For both relations, the difference in lengths between the coset representatives being compared is 1. We restrict to $q \ge 5$ since the specific $\gamma_{u,v}$ in the statement of the theorem are not defined for $q \le 4$. For smaller values of q, there are no closure relations other than those coming from Bruhat relations.

In many computations, it is easier to work with the inverses $\gamma_{u,v}^{-1}$ of the coset representatives. To prove the above result, Theorem 3.9, we will demonstrate that

$$s_j \bullet \gamma_{j+1,q+1-j}^{-1} \leqslant \gamma_{j,q+3-j}^{-1} \quad \text{and} \quad s_j \bullet \gamma_{q-j,j+1}^{-1} \leqslant \gamma_{q+2-j,j+1}^{-1}$$

with respect to the Bruhat order. There are two conditions to check for each of the relations in Theorem 3.9, and we split these into separate lemmata. We first compute the action by s_j on $\gamma_{j+1,q+1-j}^{-1}$ and $\gamma_{q-j,j+1}^{-1}$.

Lemma 3.11. Let $3 \leq j < q/2$, and let τ_i be defined by

Then $s_j \bullet \gamma_{j+1,q+1-j}^{-1} = \tau_j$.

Proof. Applying Lemma 3.7 and Lemma 3.8 yields

$$s_j \bullet \gamma_{j+1,q+1-j}^{-1} = s_j \gamma_{j+1,q+1-j}^{-1} s_{q+2-j} = \gamma_{j,q+1-j}^{-1} s_{q+2-j}.$$

Recall that $\gamma_{u,v} = (2, 3, ..., v)(1, 2, ..., u)$. Thus, the expression above can be written as

$$(1,2,\ldots,j)^{-1}(2,3,\ldots,q+1-j)^{-1}s_{q+2-j}.$$

We now determine the image of each *k* under the above permutation, focusing first on small values of *k*. Moving right to left through the permutations, we see that 1 is mapped as: $1 \mapsto j$. The element 2 is mapped as: $2 \mapsto q + 1 - j$. For all other $k \leq j + 1$, we have $k \mapsto k - 1 \mapsto k - 2$ via $(1, 2, ..., j)^{-1}(2, 3, ..., q + 1 - j)^{-1}$.

For $j + 2 \le k \le q + 1 - j$, we have $k \mapsto k - 1$ via $(2, 3, ..., q + 1 - j)^{-1}$. Since s_{q+2-j} is disjoint from the rest of the permutations, q + 2 - j and q + 3 - j are swapped. Finally, all k > q + 3 - j are fixed.

Lemma 3.12. Let $q/2 + 1 < j \leq q - 1$, and let τ_j be defined by

(3.4.5)
$$\tau_{j}(k) = \begin{cases} q-j & \text{for } k = 1, \\ j & \text{for } k = 2, \\ k-2 & \text{for } 3 \leq k \leq j+1, \\ q+2-j & \text{for } k = q+2-j, \\ q+1-j & \text{for } k = q+3-j, \\ k-1 & \text{for } q+4-j \leq k \leq j, \\ k & \text{for } k > q+3-j. \end{cases}$$

Then $s_j \bullet \gamma_{q-j,j+1}^{-1} = \tau_j$.

Proof. Applying Lemma 3.7 and Lemma 3.8 yields

$$s_j \bullet \gamma_{q-j,j+1}^{-1} = s_j \gamma_{q-j,j+1}^{-1} s_{q+2-j} = \gamma_{q-j,j}^{-1} s_{q+2-j},$$

which can be written as

$$(1, 2, \ldots, q-j)^{-1}(2, 3, \ldots, j)^{-1}s_{q+2-j}.$$

The result then follows using techniques similar to those used in the proof of Lemma 3.11.

We now prove the following result for the lengths of the τ_i 's.

Lemma 3.13. If
$$3 \le j < q/2$$
, then $\ell(s_j \bullet \gamma_{j+1,q+1-j}^{-1}) = \ell(\gamma_{j+1,q+1-j}^{-1})$.
If $q/2 + 1 < j \le q - 1$, then $\ell(s_j \bullet \gamma_{q-i,j+1}^{-1}) = \ell(\gamma_{q-i,j+1}^{-1})$.

Proof. Recall that one may compute the length of a permutation as the cardinality of its set of inversions (Equation 2.3.1). We obtain the result by a straightforward computation of the relevant sets of inversions. \Box

The following is the final ingredient needed in order to prove Theorem 3.9.

Lemma 3.14. If $3 \le j < q/2$ then $(s_j \bullet \gamma_{j+1,q+1-j}^{-1})^{-1}\gamma_{j,q+3-j}^{-1} = (2, q+2-j)$. If $q/2 + 1 < j \le q-1$ then $(s_j \bullet \gamma_{q-j,j+1}^{-1})^{-1}\gamma_{q+2-j,j}^{-1} = (1, q+2-j)$.

Proof. The proofs of these equalities use the above lemmata and simple permutation composition. For the first equality, we use the definition of τ_i in Equation (3.4.4) to write

$$(s_j \bullet \gamma_{j+1,q+1-j}^{-1})^{-1} \gamma_{j,q+3-j}^{-1} = \tau_j^{-1} (1, 2, \dots, j)^{-1} (2, 3, \dots, q+3-j)^{-1}$$

We verify that this simplifies to (2, q + 2 - j) by composition of permutations. For example, $\gamma_{j,q+3-j}^{-1}$ maps $2 \mapsto q + 3 - j$, which then maps to q + 2 - j via τ_j^{-1} . Similarly, $\gamma_{j,q+3-j}^{-1}$ maps $q + 2 - j \mapsto q + 1 - j$, which is then mapped to 2 via τ_j^{-1} . All other values are fixed by the composition.

For the second equality, we use the definition of τ_i in Equation (3.4.5) to write

$$(s_j \bullet \gamma_{q-j,j+1}^{-1})^{-1} \gamma_{q+2-j,j}^{-1} = \tau_j^{-1} (1, 2, \dots, q+2-j)^{-1} (2, 3, \dots, j)^{-1}.$$

We can use the same method as above to prove the desired result. For example, we see that $1 \mapsto q + 2 - j$ via $\gamma_{q+2-j,j}^{-1}$, which is then fixed by τ_j^{-1} . We also have that $\gamma_{q+2-j,j}^{-1}$ maps $q + 2 - j \mapsto q + 1 - j \mapsto q - j$, which is then mapped to 1 via τ_i^{-1} . All other values are fixed by the composition.

These results combine to give a proof of Theorem 3.9.

Proof of Theorem 3.9. The above lemmata demonstrate that $s_j \bullet \gamma_{j+1,q+1-j}^{-1} \leq \gamma_{j,q+3-j}^{-1}$ and $s_j \bullet \gamma_{q-j,j+1}^{-1} \leq \gamma_{q+2-j,j}^{-1}$, for the appropriately chosen values of *j*, with respect to the Bruhat order, which proves the two closure relation statements.

The following theorem gives the second set of closure relations.

Theorem 3.15. *Let* $q \ge 7$ *. Then the following relations hold*

(1) Let
$$4 \le j < q/2$$
 and $1 \le i \le j-3$. Then $\gamma_{j+1,q+1-j+i} \le \gamma_{j,q+3-j+i}$.
(2) Let $q/2 + 2 < j \le q-1$ and $1 \le i < j-q/2 - 1$. Then $\gamma_{q-j,j+1-i} \le \gamma_{q+2-j,j-i}$.

Remark 3.16. As in Theorem 3.9, the difference in lengths between the coset representatives being compared is 1. We again restrict to the values of q, this time to $q \ge 7$, since the specific $\gamma_{u,v}$ in the statement of the theorem are not defined for smaller q. Note that letting i = 0 yields the two relations given in Theorem 3.9, however we need to act by different permutations in order to realize the relations.

As we did for Theorem 3.9, we will prove the result, Theorem 3.15, by working with the inverses of the coset representatives and by proving the corresponding Bruhat relations. There are two conditions to check for each of the relations in Theorem 3.15, and we split these into separate lemmata. We first compute the action by certain permutations $h_{i,j}$ (defined below) on $\gamma_{j+1,q+1-j+i}^{-1}$ and $\gamma_{q-j-i,j+1-i}^{-1}$.

Lemma 3.17. Let $4 \leq j < q/2$ and $1 \leq i \leq j-3$. Define $h_{i,j}$ to be the permutation

(3.4.6) $h_{i,j} = s_j s_{j-1} \cdots s_{j-i} \cdot s_{q-(j-1)} s_{q-(j-2)} \cdots s_{q-(j-i)}$

and let $\tau_{i,i}$ be defined by

Then $h_{i,j} \bullet \gamma_{j+1,q+1-j+i}^{-1} = \tau_{i,j}$.

Proof. The proof of this result relies on computing permutation compositions. We provide some simplifications of the expression to make the computations easier. First, use Lemma 3.7 to write

$$h_{i,j} \bullet \gamma_{j+1,q+1-j+i}^{-1} = s_j s_{j-1} \cdots s_{j-i} \gamma_{j+1,q+1-j}^{-1} (w_{0,j} h_{i,j} w_{0,j})^{-1},$$

where $(w_{0,j} h_{i,j} w_{0,j})^{-1} = s_{j+2-i} \cdots s_j s_{j+1} s_{q+2-(j-i)} \cdots s_{q+2-(j-1)} s_{q+2-j}.$
Recall that $\gamma_{u,v} = (2, 3, ..., v)(1, 2, ..., u)$. Thus, the expression above can be written as

$$s_j s_{j-1} \cdots s_{j-i} (1, 2, \dots, j+1)^{-1} (2, 3, \dots, q+1-j)^{-1} (w_{0,j} h_{i,j} w_{0,j})^{-1}$$

We now determine the image of each k under the above permutation, focusing first on small values of k. First note that the permutation $(w_{0,j}h_{i,j}w_{0,j})^{-1}$ fixes all values less than j + 2 - i. Moving right to left through the permutations, we see that 1 is mapped as: $1 \mapsto j + 1 \mapsto j$. The element 2 is mapped as: $2 \mapsto q + 1 - j$. For all other $k \leq j + 2 - i$, we have $k \mapsto k - 1 \mapsto k - 2$ via $(1, 2, \dots, j + 1)^{-1}(2, 3, \dots, q + 1 - j)^{-1}$. Since k - 2 < j - i, it is fixed by $s_j s_{j-1} \cdots s_{j-i}$.

Now suppose $j - i + 2 \le k < j + 2$. The permutation $(w_{0,J}h_{i,j}w_{0,J})^{-1}$ maps k to k + 1. This is then mapped as $k + 1 \mapsto k \mapsto k - 1$ by $\gamma_{j+1,q+1-j}^{-1}$. Moving right to left, we see that k - 1 is mapped to k - 2 by $s_{j-1} \cdots s_{j-i}$.

As a special case, note that the permutation $(w_{0,J}h_{i,j}w_{0,J})^{-1}$ maps j + 2 as follows: $j + 2 \mapsto j + 1 \mapsto \cdots \mapsto j + 2 - i$. This is then mapped as $j + 2 - i \mapsto j + 1 - i \mapsto j - i$ by $\gamma_{j+1,q+1-j}^{-1}$. The permutation $s_{j-1} \cdots s_{j-i}$ maps j - i as follows: $j - i \mapsto j - i + 1 \mapsto \cdots \mapsto j \mapsto j + 1$.

We now consider larger values of *k*. For j + 2 < k < q + 2 - j, we see that *k* is again fixed by $(w_{0,J}h_{i,j}w_{0,J})^{-1}$. Moving right to left from here, we see that we have $k \mapsto k - 1$ by $\gamma_{i+1,a+1-i}^{-1}$.

As another special case, we may also see that $(w_{0,J}h_{i,j}w_{0,J})^{-1}$ maps q + 2 - j as follows: $q + 2 - j \mapsto q + 3 - j \mapsto q + 4 - j \dots \mapsto q + 2 - (j - i) + 1 = q + 3 - j + i$. Moving right to left, we see that q + 3 - j + i is fixed by the remaining permutations.

For $q + 3 - j \le k \le q + 3 - (j - i)$, the permutation $(w_{0,J}h_{i,j}w_{0,J})^{-1}$ maps k to k - 1. This is then fixed by $\gamma_{i+1,q+1-i}^{-1}$. Lastly, it is easy to check that all k > q + 3 - (j - i) are fixed by the permutation.

Lemma 3.18. Let $q/2 + 2 < j \le q - 1$ and $1 \le i < j - q/2 - 1$. Define $h_{i,j}$ to be the permutation (3.4.8) $h_{i,j} = s_j s_{j-1} \cdots s_{j-i} \cdot s_{q+1-(j-1)} s_{q+1-(j-2)} \cdots s_{q+1-(j-i)}$,

and let $\tau_{i,j}$ be defined by

(3.4.9)
$$\tau_{i,j}(k) = \begin{cases} q-j & \text{for } k = 1, \\ j-i & \text{for } k = 2, \\ k-2 & \text{for } 3 \leq k \leq q+1-j, \\ q+2-j+i & \text{for } k = q+2-j, \\ q+1-j+i & \text{for } k = q+3-j, \\ k-1 & \text{for } q+4-j \leq k \leq j-i, \\ k & \text{for } k > j-i. \end{cases}$$

Then $h_{i,j} \bullet \gamma_{q-j,j+1-i}^{-1} = \tau_{i,j}$.

Proof. We omit the proof of the result since it is similar to the proof Lemma 3.17.

Lemma 3.17 and Lemma 3.18 lead to the following corollary.

Corollary 3.19. *If* $4 \le j < q/2$ *and* $1 \le i \le j - 3$ *, then*

$$(h_{i,j} \bullet \gamma_{j+1,q+1-j+i}^{-1})^{-1} \gamma_{j,q+3-j+i}^{-1} = (2, q+2-j)$$

for $h_{i,j}$ *defined in Equation* (3.4.6). *If* $q/2 + 2 < j \le q - 1$ *and* $1 \le i < j - q/2 - 1$ *then*

$$(h_{i,j} \bullet \gamma_{q-j,j+1-i}^{-1})^{-1} \gamma_{q+2-j,j-i}^{-1} = (1, q+2-j)$$

for $h_{i,i}$ defined in Equation (3.4.8).

We have the following result for the lengths of the elements $\tau_{i,j}$.

Lemma 3.20. If $4 \le j < q/2$ and $1 \le i \le j-3$, then $\ell(h_{i,j} \bullet \gamma_{j+1,q+1-j+i}^{-1}) = \ell(\gamma_{j+1,q+1-j+i}^{-1})$ for $h_{i,j}$ defined in Equation (3.4.6). If $q/2 + 2 < j \le q-1$ and $1 \le i < j-q/2 - 1$ then

$$\ell(h_{i,j} \bullet \gamma_{q-j,j+1-i}^{-1}) = \ell(\gamma_{q-j,j+1-i}^{-1})$$

for $h_{i,j}$ defined in Equation (3.4.8).

Proof. Recall that one may compute the length of a permutation as the cardinality of its set of inversions (see Equation (2.3.1)). We obtain the result by a straightforward computation of the relevant sets of inversions.

These results combine to give a proof of Theorem 3.15.

Proof of Theorem 3.15. The above lemmata demonstrate that, for the appropriate values of *i* and *j* and for the appropriate $h_{i,j}$ in Equations (3.4.6) and (3.4.8), $h_{i,j} \bullet \gamma_{j+1,q+1-j+i}^{-1} \leq \gamma_{j,q+3-j+i}^{-1}$ and $h_{i,j} \bullet \gamma_{q-j,j+1-i}^{-1} \leq \gamma_{q+2-i,j-i'}^{-1}$, which proves the two closure relation statements.

3.4.3. *Example:* q = 11. We demonstrate the above theorems with the following example, where q = 11. Each node (u, v) of the diagram in Figure 1 represents a coset representative $\gamma_{u,v}$. We draw an arrow from (u_2, v_2) to (u_1, v_1) if $\gamma_{u_1, v_1} \preccurlyeq \gamma_{u_2, v_2}$. The vertical position of a node is determined by the length of the coset representative, descending from $\ell = 18$ to $\ell = 0$. Note that in the middle of the diagram, there is an "equator" where the organization of nodes shifts for ease of reading.

Observe that there are three types of relations shown, as follows:

- Bruhat relations are represented by black, straight arrows, and represent relations of the following form: $\gamma_{u,v} \preccurlyeq \gamma_{u+1,v}$ and $\gamma_{u,v} \preccurlyeq \gamma_{u,v+1}$. These are the relations given in Proposition 3.5.
- The orange squiggle arrows indicate closure relations of the form γ_{u+1,v-2} ≤ γ_{u,v}. These are above the equator, and point down and to the right.
- The purple squiggle arrows indicate closure relations of the form $\gamma_{u-2,v+1} \preccurlyeq \gamma_{u,v}$. These are below the equator, and point down and to the left.

These last two relations are given in Theorems 3.9 and 3.15.

3.4.4. *Concluding remarks on the closure order*. Based on experiments done in Sage[The23], we believe that the relations in Theorems 3.9 and 3.15 are the only non-Bruhat closure relations between coset representatives whose lengths differ by 1, and that all other closure relations comes from a chain of such relations combined with Bruhat relations. We summarize this in the following conjecture.

Conjecture 3.21. Let γ_{u_1,v_1} and γ_{u_2,v_2} be elements of $\mathbf{W}(q-2,2)$ satisfying $\gamma_{u_1,v_1} \preccurlyeq \gamma_{u_2,v_2}$. Then either we have $\gamma_{u_1,v_1} \leqslant \gamma_{u_2,v_2}$ with respect to the Bruhat order, or γ_{u_1,v_1} and γ_{u_2,v_2} satisfy the closure relations in Theorem 3.9 or 3.15, or γ_{u_1,v_1} and γ_{u_2,v_2} are related by a chain of such relations.

We have checked these claims in Sage for $q \leq 11$, but it was difficult to fully verify these statements for larger values of q. Up to q = 20 we confirmed that we do not get further relations from acting by a single simple transposition s_k . Checking the action for all possible $h \in W_{(q-2,2)}$, which has size 2!(q - 2)!, is computationally challenging. For example, even for q = 13 there are over 79 million possible h. Nevertheless, our numerical experiments and our extensive work with the action in Equation (3.4.3) lead us to believe that Conjecture 3.21 is true and that further work in this area would be fruitful.

Remark 3.22. There are $2\lceil q/2 \rceil - 5$ numbers *j* that satisfy the conditions of Theorem 3.9. Moreover, there are $(\lceil q/2 \rceil - 3)^2$ pairs (i, j) that satisfy the conditions of Theorem 3.15. Combined, this gives $(\lceil q/2 \rceil - 2)^2$ closure relations that are not Bruhat relations. Conjecture 3.21 predicts that this is the total number of closure relations that are not Bruhat relations.



FIGURE 1. Closure relations for q = 11.

4. PRODUCT MAPS: RELATIONS TO OTHER UNITARY SHIMURA VARIETIES

In this section, we begin our study of the interaction between Ekedahl-Oort strata and the supersingular locus by considering products of abelian varieties. Note that when an abelian variety *A* decomposes as a product of abelian varieties, the *p*-torsion group scheme of *A* also decomposes as a product. Because of this, the natural "product map" from a pair of unitary Shimura varieties to our Shimura variety of interest, $\mathcal{M}(q - 2, 2)$, will also induce a map in terms of the Ekedahl-Oort strata of these Shimura varieties. In Theorems 4.10 and 4.16, we explicitly describe these induced product maps on the index sets $\mathbf{W}(a, b)$ for the Ekedahl-Oort strata that must intersect the supersingular locus of $\mathcal{M}(q - 2, 2)$, using the product maps and the observation that when two abelian varieties are supersingular, their product is also supersingular.

4.1. **Background.** For any $m_1, m_2, n_1, n_2 \in \mathbb{N}$, there is a natural *product map*

$$\Phi: \mathscr{M}(m_1, n_1) \times \mathscr{M}(m_2, n_2) \to \mathscr{M}(m_1 + m_2, n_1 + n_2)$$
$$((A_1, \lambda_1, \iota_1, \xi_1), (A_2, \lambda_2, \iota_2, \xi_2)) \mapsto (A_1 \times A_2, \lambda_1 \times \lambda_2, \iota_1 \times \iota_2, \xi_1 \times \xi_2)$$

where $\lambda_1 \times \lambda_2$, $\iota_1 \times \iota_2$, and $\xi_1 \times \xi_2$ are (respectively) the natural product polarization, action, and level structure on the abelian variety $A_1 \times A_2$.

Consider an abelian variety *A* which decomposes as a product $A \cong A_1 \times A_2$, and note that the *p*-torsion group scheme satisfies $A[p] \cong A_1[p] \times A_2[p]$. Replacing A_1 by an abelian variety B_1 such that $A_1[p] \cong B_1[p]$ does not affect the Ekedahl-Oort stratum of *A*.

As a result of this, given Ekedahl-Oort strata $\mathcal{M}(m_1, n_1)_{\omega_1}$ and $\mathcal{M}(m_2, n_2)_{\omega_2}$, there is a unique stratum $\mathcal{M}(m_1 + m_2, n_1 + n_2)_{\omega}$ such that:

$$\Phi(\mathscr{M}(m_1,n_1)_{\omega_1}\times\mathscr{M}(m_2,n_2)_{\omega_2})\subseteq\mathscr{M}(m_1+m_2,n_1+n_2)_{\omega}.$$

In particular, the product map Φ induces a map φ on the index sets for the Ekedahl-Oort strata:

$$\varphi: \mathbf{W}(m_1, n_1) \times \mathbf{W}(m_2, n_2) \rightarrow \mathbf{W}(m_1 + m_2, n_1 + n_2),$$

where $\varphi(\omega_1, \omega_2)$ is the index for the unique Ekedahl-Oort stratum containing $\Phi(\mathcal{M}(m_1, n_1)_{\omega_1} \times \mathcal{M}(m_2, n_2)_{\omega_2})$.

The goal of this section is to explicitly describe the product map on the level of Weyl group cosets under the condition $(m_1 + m_2, n_1 + n_2) = (q - 2, 2)$. This is done by first constructing *standard objects* of the strata $\mathcal{M}(m_1, n_1)_{\omega_1}$ and $\mathcal{M}(m_2, n_2)_{\omega_2}$ and then computing the permutation corresponding to the sum of these standard objects.

Without loss of generality, we assume $n_2 \leq n_1$, so that either $n_1 = 1$ or $n_1 = 2$.

The former case is treated in Section 4.2 and the latter in Section 4.3.

4.2. The 1×1 Multiplication Map.

4.2.1. *General approach*. In this section we study the product map

$$\Phi: \mathscr{M}(m,1) \times \mathscr{M}(n,1) \to \mathscr{M}(m+n,2),$$

under the condition m + n = q - 2. We compute the induced map on Ekedahl-Oort strata

$$\varphi: \mathbf{W}(m,1) \times \mathbf{W}(n,1) \to \mathbf{W}(m+n,2).$$

By [Moo01, Theorem 6.7], the Ekedahl-Oort strata of $\mathcal{M}(m, 1)$ are in correspondence with cosets in the quotient $(\mathfrak{S}_1 \mathfrak{S}_m) \setminus \mathfrak{S}_{m+1}$. Each coset has minimal-length representative given by a cyclic permutation

$$\delta_a := (1, 2, \ldots, a+1)$$

for $0 \leq a \leq m$. We therefore have

$$\mathbf{W}(m,1) = \{\delta_a \mid 0 \leqslant a \leqslant m\}.$$

Note that the length of δ_a is *a*, and so the corresponding stratum $\mathcal{M}(m, 1)_{\delta_a}$ has dimension *a*.

In [VW11, Theorem F], the interaction between the Ekedahl-Oort stratification and the Newton stratification of $\mathcal{M}(m, 1)$ is completely described. If $a \leq m/2$, then we have containment:

$$\mathcal{M}(m,1)_{\delta_a} \subseteq \mathcal{M}(m,1)^{ss}.$$

On the other hand, if a > m/2, then we have disjointedness:

$$\mathcal{M}(m,1)_{\delta_a} \cap \mathcal{M}(m,1)^{ss} = \emptyset.$$

In particular, from here on, given a Dieudonné module *M* arising as the standard object for an Ekedahl-Oort stratum $\mathcal{M}(m, 1)_{\delta_a}$, we say that *M* is **supersingular** if $a \leq m/2$.

Fix $0 \le a \le m$ and $0 \le b \le n$. We aim to describe $\varphi(\delta_a, \delta_b) \in \mathbf{W}(m + n, 2)$. Recall the bijection between $\mathbf{W}(a, b)$ and Ekedahl-Oort strata of $\mathcal{M}(a, b)$ outlined below Theorem 2.2. Let the Dieudonné module M be the standard object in the Ekedahl-Oort stratum $\mathcal{M}(m, 1)_{\delta_a}$. Recall that the \mathbb{F}_{p^2} -action induces a decomposition $M = M_1 \oplus M_2$. We fix a final filtration W_{\bullet} of M, which induces filtrations $C_{i,\bullet} = W_{\bullet} \cap M_i$ of M_i for i = 1, 2. In our specific case, we have (a, b) = (m, 1) and $\dim(M_1[F]) = 1$, so the function

$$\eta_1^M(j) = \dim(C_{1,j} \cap M[F])$$

jumps, i.e., increases by 1, at exactly one index. This index is $j_1 = a + 1$ by the bijection in Theorem 2.2.

Similarly, we let $\delta_b \in \mathbf{W}(n, 1)$ represent an Ekedahl-Oort stratum in $\mathcal{M}(n, 1)$, with standard object *N*. Let $D_{i,\bullet}$ be filtrations of N_i for i = 1, 2 coming from a final filtration of *N*. Let $L := M \oplus N$ and let $E_{i,\bullet}$ be

the filtrations of L_i coming from a final filtration of L. Our goal is to determine the Weyl group coset corresponding to the Dieudonné module L via Theorem 2.2. We do this by computing the function

$$\eta_1^L(j) = \dim(E_{1,j} \cap L[F]).$$

As noted in Section 2.3, it is sufficient to compute η_1 in our work, and so this determines the permutation $\gamma_{u,v} = \varphi(\delta_a, \delta_b)$ representing the Ekedahl-Oort stratum of *L*. The signature of *L* is (m + n, 2), and so we have

$$\eta_1^L(q) = \dim(L_1[F]) = 2$$

Since the function η_1^L is clearly non-decreasing, it suffices to find the two integers where η_1^L jumps. Then this function η_1^L corresponds to a Weyl group coset $\gamma_{u,v}$. It can be seen from the description of the standard object in Lemma 3.3 that *u* and *v* are precisely the two integers where η_1^L jumps.

Depending on whether *M* and *N* are supersingular or not, there are slight differences in our method of determining the two places where η_1^L jumps. These cases are treated separately in Propositions 4.7, 4.8, and 4.9 (and summarized in Theorem 4.10). In Section 4.2.2, the necessary parts of the canonical filtration of *L* are constructed. In Section 4.2.3, this information is used to determine the jumps of η_1^L , which in turn yields the resulting Weyl group coset representative $\gamma_{u,v}$ of *L*, such that

$$\Phi\left(\mathscr{M}(m,1)_{\delta_a}\times\mathscr{M}(n,1)_{\delta_b}\right)\subseteq\mathscr{M}(m+n,2)_{\gamma_{u,v}}.$$

4.2.2. *The canonical filtration*. Using the standard objects of *M* and *N*, we form the canonical filtration of *L*, which is crucial in computing the Ekedahl-Oort stratum. Let *M* be the standard object of the Ekedahl-Oort stratum $\mathcal{M}(m, 1)_{\delta_a}$. [Moo01, 4.9] provides the action of *F* and *V* on a basis $\{e_{i,j} \mid 1 \leq i \leq 2, 1 \leq j \leq q\}$ of *M*.

$$F(e_{1,j}) = \begin{cases} e_{2,j} & \text{if } 1 \leq j \leq a \\ 0 & \text{if } j = a + 1 \\ e_{2,j-1} & \text{if } j > a + 1 \end{cases} \qquad V(e_{1,j}) = \begin{cases} 0 & \text{if } j = 1 \\ e_{2,j-1} & \text{if } 1 < j \leq m + 1 - a \\ e_{2,j} & \text{if } j > m + 1 - a \end{cases}$$
$$F(e_{2,j}) = \begin{cases} 0 & \text{if } j \neq m + 1 - a \\ e_{1,1} & \text{if } j = m + 1 - a \end{cases} \qquad V(e_{2,j}) = \begin{cases} 0 & \text{if } j \neq m + 1 - a \\ e_{1,a+1} & \text{if } j = m + 1. \end{cases}$$

The following lemma describes the action of F and V^{-1} on the subspaces $\langle e_{i,l} | 1 \leq l \leq j \rangle$. Since applying F or V^{-1} always gives another such subspace, we pick a final filtration W_{\bullet} of M such that the filtrations $C_{i,\bullet} = W_{\bullet} \cap M_i$ are given by

$$C_{i,j} = \langle e_{i,l} \mid 1 \leq l \leq j \rangle.$$

Lemma 4.1. *F* and V^{-1} have the following actions on the spaces $C_{i,j}$:

$$\begin{split} F(C_{1,j}) &= \begin{cases} C_{2,j} & \text{if } j \leq a \\ C_{2,j-1} & \text{if } j > a \end{cases} \qquad \qquad V^{-1}(C_{1,j}) \cap M_2 = \begin{cases} C_{2,m} & \text{if } j \leq a \\ C_{2,m} & \text{if } j > a \end{cases} \\ F(C_{2,j}) &= \begin{cases} C_{1,0} & \text{if } j < m+1-a \\ C_{1,1} & \text{if } j \geq m+1-a \end{cases} \qquad \qquad V^{-1}(C_{2,j}) \cap M_1 = \begin{cases} C_{1,j+1} & \text{if } j < m+1-a \\ C_{1,j} & \text{if } j \geq m+1-a. \end{cases} \end{split}$$

Proof. This is a direct application of the description, given above, of the standard object *M*.

The same result applies to the standard object *N* of the Ekedahl-Oort stratum $\mathcal{M}(n, 1)_{\delta_b}$, upon replacing *a* by *b* and *m* by *n*.

For the remainder of this section, let $s_1 := \min\{a+1, m+1-a\}$ and $s_2 := \max\{a, m+1-a\}$.

Lemma 4.2. *For* $1 \leq j \leq q$ *,*

$$V^{-1}(F(C_{1,j})) \cap M_1 = \begin{cases} C_{1,j+1} & j < s_1 \\ C_{1,j} & j = s_1 \\ C_{1,j-1} & j > s_2 \\ C_{1,j} & j = s_2 \end{cases}$$

Proof. Suppose $j < s_1 = \min\{a + 1, m + 1 - a\}$. Using Lemma 4.1, we obtain $F(C_{1,j}) = C_{2,j}$ and $V^{-1}(C_{2,j}) \cap M_1 = C_{1,j+1}$ as desired.

We check the second assertion for two cases of s_1 . In the case $s_1 = a + 1 \le m + 1 - a$ we get $F(C_{1,a+1}) = C_{2,a}$ and $V^{-1}(C_{2,a}) \cap M_1 = C_{1,a+1}$. In the case $s_1 = m + 1 - a < a + 1$, we get $F(C_{1,m+1-a}) = C_{2,m+1-a}$ and $V^{-1}(C_{2,m+1-a}) \cap M_1 = C_{1,m+1-a}$.

We now treat the analogue with decreasing index: assume $j > s_2 = \max\{a, m + 1 - a\}$. In this case, we have $F(C_{1,j}) = C_{2,j-1}$ and, subsequently, $V^{-1}(C_{2,j-1}) \cap M_1 = C_{1,j-1}$.

Finally, we prove the last assertion for both values of s_2 . In the case $s_2 = a \ge m + 1 - a$, we get $F(C_{1,a}) = C_{2,a}$ and $V^{-1}(C_{2,a}) \cap M_1 = C_{1,a}$. In the case $s_2 = m + 1 - a > a$ we obtain $F(C_{1,m+1-a}) = C_{2,m-a}$ and $V^{-1}(C_{2,m-a}) \cap M_1 = C_{1,m+1-a}$, as desired.

We now prove a corollary that allows us to 'move up' from $C_{1,0}$ to C_{1,s_1} , step by step.

Corollary 4.3. *For* $c \in \mathbb{Z}_{\geq 0}$ *, we have*

$$(V^{-1}F)^{c}(0) \cap M_{1} = C_{1,\min\{c,s_{1}\}}.$$

Proof. Apply Lemma 4.2 for $0 \le j \le s_1$ repeatedly, so that the result follows by induction starting at j = 0.

Similarly we can 'move down' from $C_{1,m+1}$ to C_{1,s_2} , as the following corollary records.

Corollary 4.4. *For* $c \in \mathbb{Z}_{\geq 0}$ *, we have*

$$(V^{-1}F)^{c}(M) \cap M_{1} = C_{1,\max\{m+1-c,s_{2}\}}$$

Proof. Apply Lemma 4.2 for $s_2 \le l \le m + 1$ repeatedly, starting at l = m + 1 and proceeding by decreasing induction.

In the rest of this section, we use these corollaries to construct the crucial parts of the filtration $E_{1,\bullet}$ of $L_1 = M_1 \oplus N_1$, induced by intersecting a final filtration of *L* with L_1 . Let s_1 and s_2 be defined as above and let $u_1 := \min\{b+1, n+1-b\}$ and $u_2 := \max\{b, n+1-b\}$.

Lemma 4.5. *For* $c \in \mathbb{Z}_{\geq 0}$ *, we have*

$$(V^{-1}F)^{c}(0) \cap L_{1} = C_{1,\min\{c,s_{1}\}} \oplus D_{1,\min\{c,u_{1}\}} = E_{1,\min\{c,s_{1}\}+\min\{c,u_{1}\}}$$
$$(V^{-1}F)^{c}(L) \cap L_{1} = C_{1,\max\{m+1-c,s_{2}\}} \oplus D_{1,\max\{n+1-c,u_{2}\}} = E_{1,\max\{m+1-c,s_{2}\}+\max\{n+1-c,u_{2}\}}.$$

Proof. This is a straightforward application of Corollary 4.3 and Corollary 4.4. For the first equality, note

$$(V^{-1}F)^{c}(0) \cap L_{1} = \left((V^{-1}F)^{c}(0) \cap M_{1} \right) \oplus \left((V^{-1}F)^{c}(0) \cap N_{1} \right)$$

Both summands are computed in Corollary 4.3. Similarly, for the second equality observe

$$(V^{-1}F)^{c}(L) \cap L_{1} = \left((V^{-1}F)^{c}(M) \cap M_{1} \right) \oplus \left((V^{-1}F)^{c}(N) \cap N_{1} \right).$$

Both summands are computed in Corollary 4.4.

In some cases, Lemma 4.5 determines every value of $\eta_1^L(j) = \dim(E_{1,j} \cap L[F])$, providing all the information we need. In general, we need to construct more subspaces of L_1 . For that, we turn to the following lemma.

Lemma 4.6. Suppose one of the following conditions holds:

- (i) M and N are both supersingular and a > b.
- (ii) *M* and *N* are both not supersingular and m a < n b.
- (iii) M is not supersingular and N is supersingular.

Then we have that

$$E_{1,1} = C_{1,1} \oplus D_{1,0}$$
 and $E_{1,q-1} = C_{1,m} \oplus D_{1,n+1}$.

Proof. We claim that, in all three cases, it suffices to construct a subspace of the form $C_{2,l} \oplus D_{2,n-b}$, for some l > m - a, in the filtration $E_{2,\bullet}$. We then show that such a subspace can be constructed in all three cases.

Assume that $C_{2,l} \oplus D_{2,n-b}$ has been constructed with l > m - a. Then Lemma 4.1 yields

$$F(C_{2,l} \oplus D_{2,n-b}) = C_{1,1} \oplus D_{1,0} = E_{1,1},$$

proving the claim. For the second statement, we first construct $E_{2,q-1}$ using symplectic complements. Note that M and N are each equipped with a symplectic pairing coming from the polarization of the abelian varieties, and so L is equipped with the product of these two pairings. By the Rosati involution condition, we have $M_1^{\perp} = M_1$. Furthermore, we may assume that the final filtration W_{\bullet} of M is stable under taking symplectic complements. Therefore, for any $1 \leq j \leq q$ we have $C_{1,j}^{\perp} = M_1 \oplus C_{2,q-j}$ and likewise for N. This implies

$$E_{1,1}^{\perp} \cap L_2 = (C_{1,1}^{\perp} \cap M_2) \oplus (D_{1,0}^{\perp} \cap N_2) = C_{2,m} \oplus D_{2,n+1} = E_{2,q-1}.$$

All three cases imply a > 0, and so applying Lemma 4.1 yields

$$V^{-1}(E_{2,q-1}) = C_{1,m} \oplus D_{1,n+1} = E_{1,q-1},$$

which finishes the proof of the second statement.

We now construct $C_{2,l} \oplus D_{2,n-b}$ in each individual case. In case (i), Lemma 4.5 allows us to move down to

$$(V^{-1}F)^b(L) \cap L_1 = C_{1,m+1-b} \oplus D_{1,n+1-b} = E_{1,q-2b}.$$

Here we have used m + 1 - b > m + 1 - a, since a > b. Furthermore, $m + 1 - a \ge a$ since M is supersingular, and $n + 1 - b \ge b$ since N is supersingular. Together these observations imply $m + 1 - b \ge s_2$ and $n + 1 - b \ge u_2$. We then construct

$$F(C_{1,m+1-b} \oplus D_{1,n+1-b}) = C_{2,m-b} \oplus D_{2,n-b}$$

Since m - b > m - a, we are done.

In case (ii), Lemma 4.5 allows us to move up to

$$(V^{-1}F)^{n-b}(0) \cap L_1 = C_{1,m+1-a} \oplus D_{1,n-b} = E_{1,m+n+1-a-b}$$

where we use: $s_1 = m + 1 - a$ and m - a < n - b. Applying *F*, using $m + 1 - a \leq a$ and $n - b \leq b$, yields

$$F(C_{1,m+1-a} \oplus D_{1,n-b}) = C_{2,m+1-a} \oplus D_{2,n-b}$$

This subspace satisfies our requirements, as m + 1 - a > m - a.

Finally, in case (iii), Lemma 4.5 allows us to move down to

$$C_{1,s_2} \oplus D_{1,u_2} = C_{1,a} \oplus D_{1,n+1-b}$$

Applying F yields

$$F(C_{1,a} \oplus D_{1,n+1-b}) = C_{2,a} \oplus D_{2,n-b}$$
18

which satisfies our requirement since *M* is not supersingular, implying a > m - a.

After constructing $E_{1,1}$ and $E_{1,q-1}$ if necessary, under the conditions of Lemma 4.6, we use the proof of Lemma 4.5 to construct

$$(4.2.1) (V^{-1}F)^{c}(E_{1,1}) \cap L_{1} = C_{1,\min\{1+c,s_{1}\}} \oplus D_{1,\min\{c,u_{1}\}} = E_{1,\min\{1+c,s_{1}\}} + \min\{c,u_{1}\},$$

$$(4.2.2) (V^{-1}F)^{c}(E_{1,q-1}) \cap L_{1} = C_{1,\max\{m-c,s_{2}\}} \oplus D_{1,\max\{n+1-c,u_{2}\}} = E_{1,\max\{m-c,s_{2}\}+\max\{n+1-c,u_{2}\}}.$$

The information that these lemmata provide about the canonical filtration of *L* is sufficient to compute its Ekedahl-Oort stratum.

4.2.3. *Ekedahl-Oort strata*. We now have the tools to completely describe the product map on the level of Weyl group cosets

$$\varphi: \mathbf{W}(m,1) \times \mathbf{W}(n,1) \to \mathbf{W}(m+n,2)$$

In particular, we compute $\varphi(\delta_a, \delta_b)$, where $\mathcal{M}(m, 1)_{\delta_a}$ is the unique Ekedahl-Oort stratum of dimension a and likewise for δ_b . Let M be a standard object for the stratum δ_a and let N be a standard object for the stratum δ_b . All possibilities for M and N (up to exchanging M and N) are treated in this section. Recall that M is supersingular if and only if $a \leq m/2$ or, equivalently, $s_1 = a + 1$. N is supersingular if and only if $b \leq n/2$ or, equivalently, $u_1 = b + 1$.

We first treat the case when *M* and *N* are both supersingular in full detail. Exchanging *M* and *N* if necessary, we assume $a \ge b$.

Proposition 4.7. Assume M and N are both supersingular, with $a \ge b$. Then the Ekedahl-Oort stratum of L is represented by the Weyl group coset of $\gamma_{2b+1,2b+2}$ if a = b and $\gamma_{2b+2,a+b+2}$ if a > b.

Proof. We compute the two places j_1 and j_2 where η_1^L jumps. Using Lemma 4.5 we 'move up' to construct the following parts of the canonical filtration:

$$\begin{split} (V^{-1}F)^b(0) \cap L_1 &= C_{1,b} \oplus D_{1,b} = E_{1,2b}, \\ (V^{-1}F)^{b+1}(0) \cap L_1 &= C_{1,b+1} \oplus D_{1,b+1} = E_{1,2b+2}, \\ (V^{-1}F)^a(0) \cap L_1 &= C_{1,a} \oplus D_{1,b+1} = E_{1,a+b+1} & \text{ in the case } a > b, \\ (V^{-1}F)^{a+1}(0) \cap L_1 &= C_{1,a+1} \oplus D_{1,b+1} = E_{1,a+b+2}. \end{split}$$

We use the explicit description $M_1[F] = \langle e_{1,a+1} \rangle$ (see Lemma 4.1), and similarly for $N_1[F]$. Thus, we compute the following values of η_1^L :

$$\begin{split} \eta_1^L(2b) &= 0, \\ \eta_1^L(2b+2) &= \begin{cases} 1 & \text{if } a > b, \\ 2 & \text{if } a = b, \end{cases} \\ \eta_1^L(a+b+1) &= 1, \\ \eta_1^L(a+b+2) &= 2. \end{split}$$

Hence η_1^L jumps at $j_2 = a + b + 2$. In the case a = b, it follows that η_1^L must also jump at $j_1 = 2b + 1$, since $\eta_1^L(2b) = 0$.

When a > b, it is not yet determined whether the first jump occurs at 2b + 1 or 2b + 2 and more work is required. In this case, Lemma 4.6 and Equation (4.2.1) imply $E_{1,1} = C_{1,1} \oplus D_{1,0}$, and so

$$(V^{-1}F)^{b}(E_{1,1}) = C_{1,\min\{b+1,a+1,m+1-a\}} \oplus D_{1,\min\{b,a+1,m+1-a\}}$$
$$= C_{1,b+1} \oplus D_{1,b} = E_{1,2b+1}.$$

Finally, using the explicit description of $L_1[F]$, we infer $\eta_1^L(2b+1) = 0$ and hence η_1^L jumps at $j_1 = 2b + 2$.

Finally, we prove the statement about the Weyl group coset representative $\gamma_{u,v}$. Recall that $\omega = \gamma_{u,v}$ is determined by the jumps of η_L^1 via Equation (2.3.2). Comparing this with Equation (3.1.1) defining $\gamma_{u,v}$ yields $u = j_1$ and $v = j_2$, completing the proof.

We now treat the case when *M* and *N* are both not supersingular, which is essentially the mirror image of the preceding case. Exchanging *M* and *N* if necessary, assume that $m - a \leq n - b$.

Proposition 4.8. Assume M and N are both not supersingular, with $m - a \leq n - b$. Then the Ekedahl-Oort stratum of L is represented by the Weyl group coset of $\gamma_{a+b+1,2a+n-m+2}$.

Proof. Using the 'moving down' of Lemma 4.5, we compute

$$(V^{-1}F)^{m-a}(L) \cap L_1 = C_{1,a+1} \oplus D_{1,n+1-m+a} = E_{1,2a+n-m+2},$$

$$(V^{-1}F)^{m-a+1}(L) \cap L_1 = C_{1,a} \oplus D_{1,n-m+a} = E_{1,2a+n-m},$$

$$(V^{-1}F)^{n-b}(L) \cap L_1 = C_{1,a} \oplus D_{1,b+1} = E_{1,a+b+1}$$
 in the case $m - a < n - b$,

$$(V^{-1}F)^{n-b+1}(L) \cap L_1 = C_{1,a} \oplus D_{1,b} = E_{1,a+b}.$$

This yields the values

$$\begin{split} \eta_1^L(a+b) &= 0, \\ \eta_1^L(a+b+1) &= 1, \\ \eta_1^L(2a+n-m) &= 1, \\ \eta_1^L(2a+n-m+2) &= 2. \end{split}$$

This implies that η_1^L jumps at $j_1 = a + b + 1$. In the case m - a = n - b, the second jump is at $j_2 = 2a + n - m + 2 = a + b + 2$.

When m - a < n - b, Lemma 4.6 and Equation (4.2.2) imply $E_{1,q-1} = C_{1,m} \oplus D_{1,n+1}$, and so

$$(V^{-1}F)^{m-a}(E_{1,q-1}) = C_{1,a} \oplus D_{1,n+1-m+a} = E_{1,2a+n-m+1}.$$

Using the explicit description of $L_1[F]$, it follows that $\eta_1^L(2a + n - m + 1) = 1$. Hence in that case we also have $j_2 = 2a + n - m + 2$.

The statement about Weyl group coset representatives follows from the formulae $u = j_1$ and $v = j_2$.

Finally, we treat the case when exactly one of the summands *M* and *N* is supersingular. Exchanging *M* and *N* if necessary, we assume that *M* is not supersingular and *N* is supersingular.

Proposition 4.9. Assume *M* is not supersingular and *N* is supersingular. Then the Ekedahl-Oort stratum of *L* is represented by the Weyl group coset of $\gamma_{b+2+\min\{b,m-a\},a+2+\max\{n-m+a,n-b\}}$.

Proof. Using Lemma 4.5, Lemma 4.6, Equation (4.2.1) and Equation (4.2.2), we compute

$$(V^{-1}F)^{b}(E_{1,1}) \cap L_{1} = C_{1,\min\{b+1,m+1-a\}} \oplus D_{1,b} = E_{1,b+1+\min\{b,m-a\}},$$

$$(V^{-1}F)^{b+1}(0) \cap L_{1} = C_{1,\min\{b+1,m+1-a\}} \oplus D_{1,b+1} = E_{1,b+2+\min\{b,m-a\}},$$

$$(V^{-1}F)^{m-a}(E_{1,q-1}) \cap L_{1} = C_{1,a} \oplus D_{1,a+1+\max\{n-m+a,n-b\}} = E_{1,a+1+\max\{n-m+a,n-b\}},$$

$$(V^{-1}F)^{m-a}(L) \cap L_{1} = C_{1,a+1} \oplus D_{1,\max\{n+1-m+a,n+1-b\}} = E_{1,a+2+\max\{n-m+a,n-b\}}.$$

Using the explicit description of $L_1[F]$, we infer that the jumps of η_1^L are

$$a = j_1 = b + 2 + \min\{b, m - a\}, \text{ and }$$

$$v = j_2 = a + 2 + \max\{n - m + a, n - b\}.$$

Proposition 4.7, Proposition 4.8 and Proposition 4.9 are summarized in Theorem 4.10.

Theorem 4.10. *The Ekedahl-Oort stratum of* $L = M \oplus N$ *is given as follows.*

- (A) If M and N are both supersingular, with $a \ge b$, then the Ekedahl-Oort stratum of L is represented by the Weyl group coset of $\gamma_{2b+1,2b+2}$ if a = b and $\gamma_{2b+2,a+b+2}$ if a > b.
- (B) If M and N are both not supersingular, with $m a \le n b$, then the Ekedahl-Oort stratum of L is represented by the Weyl group coset of $\gamma_{a+b+1,2a+n-m+2}$.
- (C) If M is not supersingular and N is supersingular. Then the Ekedahl-Oort stratum of L is represented by the Weyl group coset of $\gamma_{b+2+\min\{b,m-a\},a+2+\max\{n-m+a,n-b\}}$.

Theorem 4.10 has the following implication on the supersingular locus $\mathcal{M}(q-2,2)^{ss}$.

Corollary 4.11. *Let m*, *n*, *a*, *b be integers satisfying:*

$$m+n=q-2$$
, $a \leq m/2$, $b \leq n/2$, and $a \geq b$.

Then the Ekedahl-Oort stratum

$$\gamma_{u,v} = \begin{cases} \gamma_{2b+1,2b+2} & \text{if } a = b\\ \gamma_{2b+2,a+b+2} & \text{if } a > b \end{cases}$$

intersects $\mathcal{M}(q-2,2)^{ss}$.

Proof. A point (A, λ, ι, ξ) in the intersection $\mathcal{M}(q-2, 2)_{\gamma_{u,v}} \cap \mathcal{M}(q-2, 2)^{ss}$ is constructed as the product

 $(A, \lambda, \iota, \xi) = (A_1 \times A_2, \lambda_1 \times \lambda_2, \iota_1 \times \iota_2, \xi_1 \times \xi_2),$

where $(A_1, \lambda_1, \iota_1, \xi_1) \in \mathcal{M}(m, 1)_{\delta_a}$ and $(A_2, \lambda_2, \iota_2, \xi_2) \in \mathcal{M}(n, 1)_{\delta_b}$. It follows from Theorem 4.10 (A) that (A, λ, ι, ξ) lies in the Ekedahl-Oort stratum $\mathcal{M}(q - 2, 2)_{\gamma_{u,v}}$. Moreover, $A = A_1 \times A_2$ is supersingular, as A_1 and A_2 are supersingular by the conditions $a \leq m/2$ and $b \leq n/2$ respectively.

4.3. The 2×0 Multiplication Map.

4.3.1. General approach. We now study the product map

$$\Phi: \mathscr{M}(m,2) \times \mathscr{M}(n,0) \to \mathscr{M}(q-2,2),$$

where m + n = q - 2. Let $\gamma_{u,v}$ represent an Ekedahl-Oort stratum of $\mathcal{M}(m, 2)$, with standard object M. It is well known that the Shimura variety $\mathcal{M}(n, 0)$ consists only of the superspecial Ekedahl-Oort stratum $\mathcal{M}(n, 0)_{id} = \mathcal{M}(n, 0)$ which is characterized by the property $F^2 = 0$. The standard object of that stratum is $N^n = \bigoplus_{l=1}^n N$, where N is the mod-p Dieudonné module arising from a supersingular elliptic curve that has an action of \mathcal{O}_K with signature (1, 0). Hence, $\mathcal{M}(n, 0) = \mathcal{M}(n, 0)^{ss}$. Through a study of the sum $L := M \oplus N^n$, we compute the permutation $\gamma_{y,s} := \varphi(\gamma_{u,v}, id)$, such that

$$\Phi(\mathscr{M}(m,2)_{\gamma_{u,v}},\mathscr{M}(n,0)_{\mathrm{id}})\subseteq \mathscr{M}(q-2,2)_{\gamma_{u,s}}$$

As a result, we obtain information about whether $\mathcal{M}(q-2,2)_{\gamma_{y,s}}$ intersects the supersingular locus. If $\mathcal{M}(m,2)_{\gamma_{u,v}}$ intersects the supersingular locus, then so does $\mathcal{M}(q-2,2)_{\gamma_{y,s}}$. Conversely, if $\mathcal{M}(m,2)_{\gamma_{u,v}}$ is not contained in the supersingular locus, then neither is $\mathcal{M}(q-2,2)_{\gamma_{u,v}}$. While we do not always know how $\mathcal{M}(m,2)_{\gamma_{u,v}}$ interacts with the supersingular locus, our methods allow us to draw conclusions in some cases (see, for example, Table 3).

Our method for computing $\varphi(\gamma_{u,v}, id)$ is essentially the same as the method used in Section 4.2. We describe the standard object *M* corresponding to $\mathcal{M}(m, 2)_{\gamma_{u,v}}$ and then compute the permutation corresponding to $L = M \oplus N^n$. In this context, set $q_M := m + 2$ to be the dimension of *M*. As before, let $C_{i,\bullet}$ be filtrations of M_i coming from a final filtration of *M*. In the same way, define the filtrations $D_{i,\bullet}$ of N_i and $E_{i,\bullet}$ of L_i . As before, define $\eta_1^L(j) := \dim(E_{1,j} \cap L[F])$, and in this setting $L_1[F] = M_1[F] = \langle e_{1,u}, e_{1,v} \rangle$. It suffices to pinpoint the two places where η_1^L increases, as this determines the permutation $\gamma_{y,s}$ representing the Ekedahl-Oort stratum of *L* by Equation (2.3.2).

4.3.2. *The canonical filtration.* We construct parts of the canonical filtration of *L* by analysing the action of *F* and V^{-1} on both *M* and *N*. First, *F* and *V* act on *N* as follows:

(4.3.1)
$$F(N_1) = V(N_1) = N_2$$
 and $F(N_2) = V(N_2) = 0$.

The following lemma gives an analogous description for the action of *F* and V^{-1} on *M*.

Lemma 4.12. *F* and V^{-1} have the following actions on the spaces $C_{i,i}$:

$$F(C_{1,j}) = \begin{cases} C_{2,j} & \text{if } j < u \\ C_{2,j-1} & \text{if } u \leq j < v \\ C_{2,j-2} & \text{if } j \geq v \end{cases} \qquad V^{-1}(C_{1,j}) \cap M_2 = \begin{cases} C_{2,q_M-2} & \text{if } j < u \\ C_{2,q_M-1} & \text{if } u \leq j < v \\ C_{2,q_M} & \text{if } j \geq v \end{cases}$$
$$F(C_{2,j}) = \begin{cases} C_{1,0} & \text{if } j \leq q_M - v \\ C_{1,1} & \text{if } q_M - v < j \leq q_M - u \\ C_{1,2} & \text{if } j > q_M - u \end{cases} \qquad V^{-1}(C_{2,j}) \cap M_1 = \begin{cases} C_{1,j+2} & \text{if } j \leq q_M - v \\ C_{1,j+1} & \text{if } q_M - v < j \leq q_M - u \\ C_{1,j} & \text{if } j > q_M - u. \end{cases}$$

Proof. The result follows from applying Lemma 3.3.

A variant of 'moving up' can be used to construct parts of the canonical filtration of *L*. To record this, define the multiset $S_{up} = \{u, v, q_M - v + 2, q_M - u + 1\}$ and let $r_2 := \min(S_{up} \setminus \{\min S_{up}\})$, i.e., the second element from the multiset when the elements are listed in increasing order.

Lemma 4.13. For a sufficiently large integer *c*, we have that $(V^{-1}F)^c(0) \cap L_1 = C_{1,r_2} \oplus D_{1,0}^n$.

Proof. We compute both summands of

(4.3.2)
$$(V^{-1}F)^{c}(0) \cap L_{1} = ((V^{-1}F)^{c}(C_{1,0}) \cap M_{1}) \oplus ((V^{-1}F)^{c}(D_{1,0}) \cap N_{1})^{n}.$$

First note that $F(D_{1,0}) = D_{2,0}$ and $V^{-1}(D_{2,0}) \cap N_1 = D_{1,0}$ by Equation (4.3.1). Hence

$$V^{-1}(F(D_{1,0})) \cap N_1 = D_{1,0}$$

which implies the second summand of Equation (4.3.2). We use Lemma 4.12 to compute the first summand.

First, we assume $j < r_2$ and show

(4.3.3)

$$V^{-1}(F(C_{1,j})) \cap M_1 \supseteq C_{1,j+1}.$$

We begin by computing

$$F(C_{1,j}) = \begin{cases} C_{2,j} & \text{if } j < u \\ C_{2,j-1} & \text{if } u \leq j < v. \end{cases}$$

In the case j < u, one verifies

$$V^{-1}(F(C_{1,j})) \cap M_1 = V^{-1}(C_{2,j}) \cap M_1 = \begin{cases} C_{1,j+2} & \text{if } j \leq q_M - v, \\ C_{1,j+1} & \text{if } j > q_M - v, \end{cases}$$

so that Equation (4.3.3) is satisfied. In the case $j \ge u$, it follows from the definition of r_2 that $j \le q_M - v + 2$, and therefore

$$V^{-1}(F(C_{1,j})) \cap M_1 = V^{-1}(C_{2,j-1}) \cap M_1 = C_{1,j+1}$$

by Lemma 4.12 and, again, Equation (4.3.3) is satisfied. Thus applying $V^{-1}F$ increases the index in $C_{1,\bullet}$ for $j < r_2$. To prove it does not increase further, we show

(4.3.4)
$$V^{-1}(F(C_{1,r_2})) \cap M_1 = C_{1,r_2}.$$

First, Lemma 4.12 yields

$$F(C_{1,r_2}) = \begin{cases} C_{2,r_2} & \text{if } r_2 < u, \\ C_{2,r_2-1} & \text{if } u \leq r_2 < v, \\ C_{2,r_2-2} & \text{if } r_2 = v. \end{cases}$$

In the first case we have $r_2 < u < v$ and it follows that $r_2 = q_M - u + 1$. Hence

$$V^{-1}(F(C_{1,r_2})) \cap M_1 = V^{-1}(C_{2,r_2}) \cap M_1 = C_{1,r_2}.$$

In the second case we have $u \leq r_2 < v$ and it follows that $q_M - v + 2 \leq r_2 \leq q_M - u + 1$. Therefore

$$V^{-1}(F(C_{1,r_2})) \cap M_1 = V^{-1}(C_{2,r_2-1}) \cap M_1 = C_{1,r_2}.$$

Finally, in the third case we have $r_2 = v$ and it follows that $r_2 \leq q_M - v + 2$. We obtain

$$W^{-1}(F(C_{1,r_2})) \cap M_1 = V^{-1}(C_{2,r_2-2}) \cap M_1 = C_{1,r_2}.$$

We conclude that applying $V^{-1}F$ allows us to 'move up' along $C_{1,\bullet}$, precisely until we reach C_{1,r_2} . This proves the lemma.

Analogously, we record the effect of 'moving down'. Define the multiset $S_{\text{down}} := \{u - 1, v - 1, q_M - v + 2, q_M - u + 1\}$ and let $r_3 := \max(S_{\text{down}} \setminus \{\max S_{\text{down}}\})$, i.e., the second element from the multiset when the elements are listed in increasing order.

Lemma 4.14. For a sufficiently large integer *c*, we have $(V^{-1}F)^{c}(L) \cap L_{1} = C_{1,r_{3}} \oplus D_{1,1}^{n}$.

Proof. The proof is similar to the proof of Lemma 4.13. We compute both summands of

(4.3.5)
$$(V^{-1}F)^{c}(L) \cap L_{1} = ((V^{-1}F)^{c}(C_{1,q_{M}}) \cap M_{1}) \oplus ((V^{-1}F)^{c}(D_{1,1}) \cap N_{1})^{n}.$$

First, using Equation (4.3.1) we obtain $V^{-1}(F(D_{1,1})) \cap N_1 = D_{1,1}$ for the second summand of Equation (4.3.5). We now compute the first summand using Lemma 4.12. Assume $l > r_3$, so that l exceeds at least three of the elements of S_{down} . We show

(4.3.6) $V^{-1}(F(C_{1,l})) \cap M_1 \subseteq C_{1,l-1}.$

First, applying *F* yields

$$F(C_{1,l}) = \begin{cases} C_{2,l-1} & \text{if } l < v \\ C_{2,l-2} & \text{if } l \ge v \end{cases}$$

In the first case, it follows that $l > q_M - u + 1$ and hence

$$W^{-1}(F(C_{1,l})) \cap M_1 = V^{-1}(C_{2,l-1}) \cap M_1 = C_{1,l-1}.$$

In the second case, we use $l > q_M - v + 2$ to obtain

$$W^{-1}(F(C_{1,l})) \cap M_1 = V^{-1}(C_{2,l-2}) \cap M_1 \subseteq C_{1,l-1}$$

In both cases Equation (4.3.6) is satisfied. Thus applying $V^{-1}F$ decreases the index in $C_{1,\bullet}$ for $l > r_3$. We show it cannot decrease further, meaning

(4.3.7)
$$V^{-1}(F(C_{1,r_3})) \cap M_1 = C_{1,r_3}.$$

We begin by applying *F*:

$$F(C_{1,r_3}) = \begin{cases} C_{2,r_3} & \text{if } r_3 = u - 1 \\ C_{2,r_3 - 1} & \text{if } u \leqslant r_3 < v \\ C_{2,r_3 - 2} & \text{if } r_3 \geqslant v. \end{cases}$$

In the case $r_3 = u - 1$, it follows that $r_3 \ge q_M - u + 1$ and therefore

$$V^{-1}(F(C_{1,r_3})) \cap M_1 = V^{-1}(C_{2,r_3}) \cap M_1 = C_{1,r_3},$$

as desired. In the second case we have $u \leq r_3 < v$ and it follows that $q_M - v + 2 \leq r_3 \leq q_M - u + 1$. Hence

$$V^{-1}(F(C_{1,r_3})) \cap M_1 = V^{-1}(C_{2,r_3-1}) \cap M_1 = C_{1,r_3}.$$

Finally, in the third case we have $r_3 > v$. It follows that $r_3 = q_M - v + 2$ and thus

$$V^{-1}(F(C_{1,r_3})) \cap M_1 = V^{-1}(C_{2,r_3-2}) \cap M_1 = C_{1,r_3}.$$

In each case, Equation (4.3.7) holds, showing that we cannot 'move down' further down than C_{1,r_3} . This finishes the proof of the lemma.

In order to describe the Ekedahl-Oort stratum of L, one more lemma is needed.

Lemma 4.15. We have $C_{1,r_2} \cap M[F] = C_{1,r_3} \cap M[F]$.

Proof. Since $r_2 \leq r_3$ and, hence, $C_{1,r_2} \subseteq C_{1,r_3}$ the inclusion " \subseteq " is immediate, so we focus on the inclusion " \supseteq ". Recall that $M_1[F] = \langle e_{1,u}, e_{1,v} \rangle$. Given the definitions of S_{up} and S_{down} one can check that it is not possible to have $r_2 < u < r_3$ or $r_2 < v < r_3$. Therefore there is no element of M[F] gained when moving from C_{1,r_2} to C_{1,r_3} . In other words, we have $C_{1,r_3} \cap M[F] \subseteq C_{1,r_2}$, which proves the lemma.

4.3.3. *Ekedahl-Oort strata*. Recall the setting. *M* is the standard object of $\mathcal{M}(m, 2)_{\gamma_{u,v}}$, and $C_{i,\bullet}$ are filtrations of M_i coming from a final filtration on *M*. *N* is the standard object of the unique Ekedahl-Oort stratum of $\mathcal{M}(1,0)$, with the filtration $D_{i,\bullet}$ of N_i . We form $L := M \oplus N^n$, with the filtrations $E_{1,\bullet}$ of L_i . We define

$$\eta_1^M(j) = \dim(C_{1,j} \cap M[F]) \eta_1^L(j) = \dim(E_{1,j} \cap L[F]).$$

Recall that $r_2 = \min(S_{up \setminus \min S_{up}})$, the second smallest element of the multiset $S_{up} := \{u, v, q_M - v + 2, q_M - u + 1\}$. We now completely describe the Ekedahl-Oort stratum of *L*.

Theorem 4.16. The Ekedahl-Oort stratum of $L := M \oplus N^n$ in $\mathcal{M}(m + n, 2)$ is represented by the permutation $\gamma_{y,s}$, with

$$y = \begin{cases} u & \text{if } u \leqslant r_2 \\ u+n & \text{if } u > r_2 \end{cases} \qquad s = \begin{cases} v & \text{if } v \leqslant r_2 \\ v+n & \text{if } v > r_2. \end{cases}$$

Proof. To prove the result, we compute the function η_1^L . For $j \leq r_2$, Lemma 4.13 implies

$$E_{1,j} = C_{1,j} \oplus D_{1,0}^n$$

Hence, $\eta_1^L(j) = \eta_1^M(j)$, since $D_{1,0} \cap N[F] = 0$ by Equation (4.3.1). For $j \ge r_3 + n$, Lemma 4.14 implies

$$E_{1,j}=C_{1,j-n}\oplus D_{1,1}^n$$

Hence, $\eta_1^L(j) = \eta_1^M(j-n)$, since $D_{1,1} \cap N[F] = 0$ by Equation (4.3.1). Finally, Lemma 4.15 implies that η_1^L stays constant between r_2 and r_3 . In summary:

(4.3.8)
$$\eta_1^L(j) = \begin{cases} \eta_1^M(j) & \text{if } j \leq r_2, \\ \eta_1^M(r_2) & \text{if } r_2 < j < r_3 + n, \\ \eta_1^M(j-n) & \text{if } j \geq r_3 + n. \end{cases}$$

Note that η_1^M jumps at u and v. If $u \leq r_2$, then we are in case 1 of Equation (4.3.8) and η_1^L jumps at j = u. If $u > r_2$ then, as noted in the proof of Lemma 4.15, u must also be greater than r_3 . Hence, we are in case 3 of Equation (4.3.8) and η_1^L jumps at j = u + n. Similarly, η_1^L jumps at either v or v + n if $v \leq r_2$ or $v > r_2$, respectively. This results in the statement of the theorem.

This theorem has the following implication for supersingular loci.

Corollary 4.17. Let m + n = q - 2 and $\mathcal{M}(m, 2)_{\gamma_{u,v}} \cap \mathcal{M}(m, 2)^{ss} \neq \emptyset$. Then we have

$$\mathcal{M}(q-2,2)_{\gamma_{\mu,s}}\cap \mathcal{M}(q-2,2)^{ss}\neq \emptyset,$$

where y and s are computed from u and v as in Theorem 4.16.

Proof. This follows immediately from Theorem 4.16, the fact $\mathcal{M}(n, 0) = \mathcal{M}(n, 0)^{ss}$, and the observation that the product of two supersingular abelian varieties is supersingular.

Note that Corollary 4.17 allows one to gain information about some Ekedahl-Oort strata that intersect $\mathcal{M}(q-2,2)$, using information about which Ekedahl-Oort strata intersect $\mathcal{M}(m,2)^{ss}$, for *m* smaller than q-2. Fortunately, there is a complete classification of Ekedahl-Oort strata that intersect the supersingular locus of $\mathcal{M}(m,2)$ for very small *m*. We first recall results for m = 0 and 1.

The Shimura variety $\mathcal{M}(0,2)$ is isomorphic to $\mathcal{M}(2,0)$. As mentioned in Subsection 4.3.1, $\mathcal{M}(0,2)$ has a unique Ekedahl-Oort stratum. This must coincide with the supersingular locus, and so $\mathcal{M}(0,2) = \mathcal{M}(0,2)^{ss}$.

The Shimura variety $\mathcal{M}(2,1)$ is isomorphic to $\mathcal{M}(1,2)$, and the interaction between the Ekedahl-Oort strata and the supersingular locus in this case is summarized in Subsection 4.2.1.

Proposition 4.18. *The supersingular locus of* $\mathcal{M}(2,2)$ *is exactly the union of those Ekedahl-Oort strata of dimension two or less:*

$$\mathscr{M}(2,2)^{ss} = \mathscr{M}(2,2)_{\gamma_{1,2}} \sqcup \mathscr{M}(2,2)_{\gamma_{1,3}} \sqcup \mathscr{M}(2,2)_{\gamma_{1,4}} \sqcup \mathscr{M}(2,2)_{\gamma_{2,3}}$$

Proof. By [GH15], the Shimura variety $\mathcal{M}(2,2)$ is of Coxeter type. In particular, the supersingular locus is a union of Ekedahl-Oort strata since every Ekedahl-Oort stratum intersecting the supersingular locus is entirely contained in the supersingular locus.

There are six Ekedahl-Oort strata of $\mathcal{M}(2,2)$, including exactly two strata of dimension two: those indexed by $\gamma_{2,3}$ and $\gamma_{1,4}$. It is known by results of [HP14] that the supersingular locus $\mathcal{M}(2,2)^{ss}$ is two-dimensional, and so at least one of $\mathcal{M}(2,2)_{\gamma_{1,4}}$ or $\mathcal{M}(2,2)_{\gamma_{2,3}}$ must be contained in the supersingular locus. In what follows, we observe that, in fact, both are.

There is a natural automorphism of $\mathcal{M}(2,2)(\Bbbk)$, taking (A, ι, λ, ξ) to $(A, \overline{\iota}, \lambda, \xi)$, where $\overline{\iota}$ is the action ι composed with the nontrivial automorphism of K. This action stabilizes the supersingular locus, but has the effect of conjugating the action on the *p*-torsion group schemes defining the Ekedahl-Oort strata. On the level of the standard objects $N_{\gamma_{u,v}}$ corresponding to $\gamma_{u,v}$, this action interchanges the roles of the basis vectors $e_{1,j}$ and $e_{2,j}$ for each $1 \leq j \leq q$.

By applying Lemma 3.3 to compute the standard objects $N_{1,4}$ and $N_{2,3}$, one can see immediately that interchanging the roles of $e_{1,j}$ and $e_{2,j}$ defines an isomorphism between $N_{1,4}$ and $N_{2,3}$. Therefore, both $\mathcal{M}(2,2)_{\gamma_{1,4}}$ and $\mathcal{M}(2,2)_{\gamma_{2,3}}$ are contained in the supersingular locus.

Since $\mathcal{M}(2,2)_{\gamma_{1,4}}$ and $\mathcal{M}(2,2)_{\gamma_{2,3}}$ are contained in the supersingular locus and the supersingular locus is closed, the closure of these strata are also contained in the supersingular locus. By Proposition 3.5,

$$\mathscr{M}(2,2)_{\gamma_{1,4}} \bigsqcup \mathscr{M}(2,2)_{\gamma_{2,3}} = \mathscr{M}(2,2)_{\gamma_{1,2}} \sqcup \mathscr{M}(2,2)_{\gamma_{1,3}} \sqcup \mathscr{M}(2,2)_{\gamma_{1,4}} \sqcup \mathscr{M}(2,2)_{\gamma_{2,3}}$$

i.e., exactly those strata of dimension less than or equal to two.

Finally, the strata indexed by $\gamma_{2,4}$ and $\gamma_{3,4}$ have dimension greater than two, and so they cannot be contained in the supersingular locus.

5. THE FORGETFUL MAP: RELATION TO SIEGEL MODULAR VARIETY

In the previous section we obtained information about the Shimura variety $\mathcal{M}(q-2,2)$ via product maps that we developed. The results of that section, particularly Corollary 4.11 and Corollary 4.17, allow us to draw conclusions about the interaction between Ekedahl-Oort strata and the supersingular locus in some cases. In this section, we further our study of this interaction by relating our Shimura variety $\mathcal{M}(q-2,2)$ to the Siegel modular variety \mathcal{A}_q .

5.1. **Background.** By "forgetting" the unitary structure of the abelian varieties we are considering, we obtain a point on the Siegel modular variety \mathcal{A}_q . In other words, there is a **forgetful map**

The map maintains the polarisation λ and the level structure ξ of the abelian variety A, but drops the unitary structure $\iota : \mathcal{O}_K \to \text{End}(A)$ of signature (q - 2, 2). This forgetful map induces a map on Ekedahl-Oort strata

$$\psi: \mathbf{W}(q-2,2) \to \mathbf{W}_q,$$

where \mathbf{W}_q is the set of minimal length Weyl group coset representatives indexing the Ekedahl-Oort strata of \mathscr{A}_q . By [Moo01, 3.6], the set \mathbf{W}_q consists of the permutations $\omega \in \mathfrak{S}_{2q}$ satisfying $\omega^{-1}(1) < \omega^{-1}(2) < \cdots < \omega^{-1}(q)$ and $\omega(i) + \omega(2q + 1 - i) = 2q + 1$. This section is devoted to computing $\psi(\gamma_{u,v})$ for $\gamma_{u,v} \in \mathbf{W}(q-2,2)$ with the goal of obtaining information about $\mathscr{M}(q-2,2)$ from various results about \mathscr{A}_q .

We now outline how, given $\gamma_{u,v} \in \mathbf{W}(q-2,2)$, the permutation $\psi(\gamma_{u,v}) \in \mathbf{W}_q$ is computed. Using the bijection in Theorem 2.2 and Lemma 3.3, we construct the standard object *M* corresponding to the Ekedahl-Oort stratum $\mathscr{M}(q-2,2)_{\gamma_{u,v}}$. Upon forgetting the \mathbb{F}_{p^2} -action of *M*, we observe that *M* is the Dieudonné module of an Ekedahl-Oort stratum of \mathscr{A}_q . By [Moo01, 3.6], this Ekedahl-Oort stratum corresponds to an element of \mathbf{W}_q . This correspondence works as follows. Let W_{\bullet} be a final filtration of *M* and define the non-decreasing function

$$\eta(j) := \dim(W_j \cap M[F])$$

for $0 \le j \le 2q$. Note that $\eta(2q) = \dim(M[F]) = q$. Let $1 \le j_1 < \cdots < j_q \le 2q$ be the indices where η jumps, meaning $\eta(j_l) = \eta(j_l - 1) + 1$, and let $1 \le i_1 < \cdots < i_q \le 2q$ be the remaining indices. We define $\omega_{u,v} \in \mathfrak{S}_{2q}$ by $\omega_{u,v}(j_l) = l$ and $\omega_{u,v}(i_m) = m + q$. By construction, $\omega_{u,v}$ is an element of \mathbf{W}_q and $\psi(\gamma_{u,v}) = \omega_{u,v}$.

The duality coming from the symplectic pairing on *M* implies that $\eta(j) = \eta(j-1)$ holds if and only if $\eta(2q+1-j) = \eta(2q-j) + 1$ holds. By induction, it follows that $\eta(j) + q = \eta(2q-j) + j$. The identity $i_m = 2q + 1 - j_{q-m}$ then follows from induction, implying that

(5.1.1)
$$\omega_{u,v}(i) + \omega_{u,v}(2q+1-i) = 2q+1.$$

The preceding paragraphs show that the function $\eta(j) = \dim(W_j \cap M[F])$ determines the permutation $\omega_{u,v} = \psi(\gamma_{u,v})$, and so we focus on determining all the values of η . Lemma 3.3 yields the description

$$M[F] = \langle e_{1,u}, e_{1,v} \rangle \oplus \langle e_{2,j} \mid j \notin \{q+1-u, q+1-v\} \rangle.$$

Recall the filtrations $C_{i,\bullet} = W_{\bullet} \cap M_i$ and note that any W_j can be written as $W_j = C_{1,l_1} \oplus C_{2,l_2}$ for some l_1 and l_2 satisfying $l_1 + l_2 = j$. In Proposition 5.1, this decomposition is made explicit, and as a result a final filtration of M is given in terms of $C_{1,\bullet}$ and $C_{2,\bullet}$. The shape of this final filtration depends on u and v. In Theorem 5.2, this final filtration is used to determine the permutation $\omega_{u,v} \in \mathbf{W}_q$.

There are three possibilities for the Ekedahl-Oort stratum of $(\mathscr{A}_q)_{\omega_{u,v}}$. The first possibility is that $(\mathscr{A}_q)_{\omega_{u,v}}$ is contained in \mathscr{A}_q^{ss} . In that case, it follows that $\mathscr{M}(q-2,2)_{\gamma_{u,v}}$ is contained in $\mathscr{M}(q-2,2)^{ss}$. Corollary 5.4 records the Ekedahl-Oort strata of $\mathscr{M}(q-2,2)$ that occur in in this way. Another possibility is that $(\mathscr{A}_q)_{\omega_{u,v}}$ is disjoint from \mathscr{A}_q^{ss} . In that case, it follows that $\mathscr{M}(q-2,2)_{\gamma_{u,v}}$ is disjoint from $\mathscr{M}(q-2,2)^{ss}$. Proposition 5.6 and Corollary 5.12 record the Ekedahl-Oort strata of $\mathscr{M}(q-2,2)$ with this property. Finally, it is possible that the Ekedahl-Oort stratum $(\mathscr{A}_q)_{\omega_{u,v}}$ intersects \mathscr{A}_q^{ss} , but is not contained in it. In this case, we do not obtain information about the Ekedahl-Oort stratum $\mathscr{M}(a,b)_{\gamma_{u,v}}$, due to the highly non-surjective nature of the forgetful map. It could, for instance, be that there exist supersingular abelian varieties in $(\mathcal{A}_q)_{\omega_{u,v}}$, but that none of them admit a compatible unitary structure of signature (q - 2, 2). For this reason, our work in this section focuses on the cases when the Ekedahl-Oort $(\mathcal{A}_q)_{\omega_{u,v}}$ is either completely contained in \mathcal{A}_q^{ss} or completely disjoint from it.

5.2. A Final Filtration of the Standard Object. In order to compute $\eta(j) = \dim(W_j \cap M[F])$, we construct a final filtration W_{\bullet} of M in terms of $C_{1,\bullet}$ and $C_{2,\bullet}$. Essentially this is done by applying Lemma 3.3 (or its consequence Lemma 4.12) repeatedly. We could give a canonical filtration instead, but giving a final filtration makes the exposition shorter and later computations easier. It is sufficient to construct W_j , where $1 \leq j \leq q$, as the remaining parts of the filtration can be constructed by taking symplectic complements. As seen in the proof of Lemma 4.6, we have

$$W_{j}^{\perp} = (C_{1,l_{1}} \oplus C_{2,l_{2}})^{\perp} = C_{1,l_{1}}^{\perp} \cap C_{2,l_{2}}^{\perp} = (M_{1} \oplus C_{2,q-l_{1}}) \cap (C_{1,q-l_{2}} \oplus M_{2}) = C_{1,q-l_{2}} \oplus C_{1,q-l_{1}} = W_{2q-j}.$$

The following proposition gives a final filtration W_{\bullet} of M that is dependent on u and v. When there is only one way to fill up a gap between two subspaces, for instance between $C_{1,1} \oplus C_{2,0}$ and $C_{1,1} \oplus C_{2,q-2}$, then the subspaces in between (of the form $C_{1,1} \oplus C_{2,l}$ for 1 < l < q - 2) are omitted from the notation. In each case, Lemma 3.3 yields

(5.2.1)
$$W_{q} = F(M) = \langle e_{1,1}, e_{1,2} \rangle \oplus \langle e_{2,j} \mid j \leq q - 2 \rangle = C_{1,2} \oplus C_{2,q-2},$$
$$M[F] = M[F]_{1} \oplus M[F]_{2} = \langle e_{1,u}, e_{1,v} \rangle \oplus \langle e_{2,j} \mid j \notin \{q + 1 - v, q + 1 - u\} \rangle.$$

Proposition 5.1. Let *M* be the standard object of $\mathcal{M}(q-2,2)_{\gamma_{u,v}}$. In the case u = 1, the first half of a final filtration of *M* is

$$0 \subset C_{1,1} \oplus C_{2,0} \subset \cdots \subset C_{1,1} \oplus C_{2,q-2} \subset W_q.$$

In the case u = 2, the first half of a final filtration is

$$0 \subset C_{1,0} \oplus C_{2,1} \subset C_{1,1} \oplus C_{2,1} \subset \cdots \subset C_{1,1} \oplus C_{2,s_1} \subset C_{1,2} \oplus C_{2,s_1} \subset \cdots \subset W_q$$

In the case u > 2 and v < q - 1, the first half of a final filtration is

$$0 \subset \cdots \subset C_{1,0} \oplus C_{2,2} \subset \cdots \subset C_{1,2} \oplus C_{2,2} \subset \cdots \subset W_q.$$

In the case u > 2 and $v \ge q - 1$, the first half of a final filtration is

$$0 \subset C_{1,0} \oplus C_{2,1} \subset C_{1,1} \oplus C_{2,1} \subset C_{1,1} \oplus C_{2,2} \subset C_{1,2} \oplus C_{2,2} \subset \cdots \subset W_q.$$

Proof. We prove the proposition by constructing the necessary parts of the canonical filtration. The standard object from Lemma 3.3 is used for determining the effect on *F* or V^{-1} on a subspace. We split the proof up into three cases: u = 1, u = 2 and u > 2.

First, assume u = 1. In that case, we construct

$$F(W_q) = \begin{cases} C_{1,0} \oplus C_{2,0} = W_0 & \text{if } v = 2, \\ C_{1,1} \oplus C_{2,1} = W_2 & \text{if } v > 2. \end{cases}$$

In the case (u, v) = (1, 2), we have $W_q = M[F]$ and therefore $\eta(q) = q$, implying that η has to increase by 1 at each index between 0 and q. It is not possible to construct a subspace between 0 and W_q using F and V^{-1} , so we can extend the canonical filtration to a final filtration in any way we like; it will not influence the function η . In the case v > 2, we construct

$$F(W_2) = \begin{cases} C_{1,0} \oplus C_{2,0} = W_0 & \text{if } v < q, \\ C_{1,1} \oplus C_{2,0} = W_1 & \text{if } v = q. \end{cases}$$

Here we have used that $F(C_{1,1}) = 0$, since u = 1, and $F(C_{2,1}) = 0$ if and only if v < q. In the first case, the canonical filtration cannot produce W_1 . This is not required, as $\eta(2) = 2$ implies $\eta(1) = 1$. In both cases, we can form a final filtration containing W_1 .

Next, we fill in the canonical filtration between W_2 and W_q . This amounts to discovering when $C_{1,1}$ changes to $C_{1,2}$ in the filtration. We construct

$$V^{-1}(W_q) = C_{1,q-1} \oplus C_{2,q-1} = W_{2q-2} = W_2^{\perp}$$

We then apply *F* to obtain

$$F(W_{2q-2}) = \begin{cases} C_{1,1} \oplus C_{2,q-2} = W_{q-1} & \text{if } v = q, \\ C_{1,1} \oplus C_{2,q-3} = W_{q-2} & \text{if } v < q. \end{cases}$$

Thus far, we have constructed W_0 , W_1 , W_2 , W_{q-1} , and W_q . This forces all the intermediate $W_i = C_{1,1} \oplus C_{2,i-1}$ due to dimension reasons, as we showed that all have the $C_{1,1}$ term in the first half. In the case v = q, we are done: the first half of the canonical filtration is

$$(5.2.2) 0 \subset C_{1,1} \oplus C_{2,0} \subset C_{1,1} \oplus C_{2,1} \subset \cdots \subset C_{1,1} \oplus C_{2,q-2} \subset C_{1,2} \oplus C_{2,q-2} = W_q.$$

In the case v = 3, we compute

$$\eta(q-2) = \dim \left(W_{q-2} \cap M[F] \right) = \eta_1(1) + \eta_2(q-3) = 1 + (q-3) = q-2.$$

Together with $\eta(q) = q - 2$, this implies $\eta(q - 1) = q - 2$, meaning that W_{q-1} does not need to be constructed. The final filtration in Equation (5.2.2) is still valid in this case.

In the case 3 < v < q, one more step is needed. We construct

$$V^{-1}(W_2) = V^{-1}(C_{1,1} \oplus C_{2,1}) = C_{1,3} \oplus C_{2,q-1} = W_{q+2},$$

$$V^{-1}(W_{q+2}) = V^{-1}(C_{1,3} \oplus C_{2,q-1}) = C_{1,q} \oplus C_{2,q-1} = W_{2q-1},$$

$$F(W_{2q-1}) = F(C_{1,q} \oplus C_{2,q-1}) = C_{1,1} \oplus C_{2,q-2} = W_{q-1}.$$

We conclude that the final filtration given in Equation (5.2.2) always works in the case u = 1.

We now treat the case u = 2, which is the most challenging. We again construct

$$F(W_q) = C_{1,1} \oplus C_{2,1} = W_2.$$

In the case v = q, we compute $\eta(2) = 0$. Combining this with $\eta(q) = q - 2$ fixes the behavior of η everywhere; it has to jump everywhere between $\eta(2) = 0$ and $\eta(q) = q - 2$. Therefore we are finished with this case and assume 2 < v < q from here on. Applying *F* to W_2 yields

$$F(W_2) = C_{1,0} \oplus C_{2,1} = W_1$$

This implies that in the case u = 2 the final filtration must be of the form

$$0 \subset C_{1,0} \oplus C_{2,1} \subset C_{1,1} \oplus C_{2,1} \subset C_{1,0} \oplus C_{2,2} \subset C_{1,1} \oplus C_{2,2} \subset \cdots$$

$$\cdots \subset C_{1,1} \oplus C_{2,m} \subset C_{1,2} \oplus C_{2,m} \oplus \cdots \subset C_{1,2} \oplus C_{2,q-2} = W_q$$

for some (not necessarily unique) integer *m*. The remaining goal is to find such an integer *m*. We do this by constructing $C_{1,1} \oplus C_{2,j}$ for increasing *j* (by 'moving up') and constructing $C_{1,2} \oplus C_{2,j}$ for decreasing *j* (by 'moving down').

We first outline the moving down procedure. Define $s_2 := \max\{v - 2, q - v\}$. Starting from $W_q = C_{1,2} \oplus C_{2,q-2}$, we construct $W_{j+2} = C_{1,2} \oplus C_{2,j}$ for decreasing *j*, by applying FV^{-1} , until we reach $C_{1,2} \oplus C_{2,s_2}$. The induction step goes as follows. Assume $j > s_2$. Since j > q - v, we have

$$V^{-1}(W_{j+2}) = V^{-1}(C_{1,2} \oplus C_{2,j}) = C_{1,j+1} \oplus C_{2,q-1} = W_{q+j}$$

As additionally $j + 1 \ge v$ holds, we have

$$F(C_{1,j+1} \oplus C_{2,q-1}) = C_{1,2} \oplus C_{2,j-1} = W_{j+1}$$

In this way $C_{1,2} \oplus C_{2,j}$ is constructed for decreasing *j*. We now show that this process halts when $j = s_2$ is reached. If j = q - v, then we have

$$V^{-1}(C_{1,2} \oplus C_{2,j}) = C_{1,j+2} \oplus C_{2,q-1} = W_{q+j+1}.$$

and we cannot move down further. Similarly, if j = v - 2, then we obtain

$$F(V^{-1}(C_{1,2} \oplus C_{2,j})) = F(C_{1,j+1} \oplus C_{2,q-1}) = C_{1,2} \oplus C_{2,j}.$$

We conclude that we can move down to $W_{2+s_2} = C_{1,2} \oplus C_{2,s_2}$ and no further.

The moving up procedure works analogously. Define $s_1 := \min\{v - 2, q + 1 - v\}$. We begin with $W_2 = C_{1,1} \oplus C_{2,1}$ and construct $W_{j+1} = C_{1,1} \oplus C_{2,j}$ for increasing j, by applying FV^{-1} , until we reach $C_{1,1} \oplus C_{2,s_1}$. Now, for the induction step we assume $j < s_1$. Since $j \leq q - v$, we have

$$V^{-1}(W_{j+1}) = V^{-1}(C_{1,1} \oplus C_{2,j}) = C_{1,j+2} \oplus C_{2,q-2} = W_{q+j}.$$

Then, as j + 2 < v holds, we obtain

$$F(W_{q+j}) = F(C_{1,j+2} \oplus C_{2,q-2}) = C_{1,1} \oplus C_{2,j+1} = W_{j+2}$$

Thus $C_{1,1} \oplus C_{2,j}$ is constructed for increasing *j*. We now show that this procedure stops when $j = s_1$ is reached. In the case j = q + 1 - v, we have

$$V^{-1}(W_{j+1}) = V^{-1}(C_{1,1} \oplus C_{2,j}) = C_{1,j+1} \oplus C_{2,q-2} = W_{q+j-1}.$$

On the other hand, in the case j = v - 2, we have

$$F(W_{q+j}) = F(C_{1,j+2} \oplus C_{2,q-2}) = C_{1,1} \oplus C_{2,j}.$$

Finally, it is left to prove that this construction suffices, in the sense that it is not necessary to construct more parts of the filtration in order to compute the function η . In general there may be subspaces between W_{1+s_1} and W_{2+s_2} . We now show that this does not influence the behavior of η .

In the case $v - 2 \ge q + 1 - v$, we have $W_{1+s_1} = W_{q+2-v} = C_{1,1} \oplus C_{2,q+1-v}$. Since $e_{2,q+1-v} \notin M[F]$, we obtain that $\eta(q+2-v) = q - v$. Then η must keep increasing till $\eta(q) = q - 2$, so the canonical filtration does not develop further.

On the other hand, in the case v - 2 < q + 1 - v we have $W_{2+s_2} = W_{q+2-v} = C_{1,2} \oplus C_{2,q-v}$. We compute and obtain $\eta(q+2-v) = q+1-v$. This implies that η must keep increasing between $\eta(2) = 1$ and $\eta(q+2-v) = q+1-v$. Hence the canonical filtration does not develop further. In both cases the first half of a final filtration of *M* is given by

$$0 \subset C_{1,0} \oplus C_{2,1} \subset C_{1,1} \oplus C_{2,1} \subset \cdots \subset C_{1,1} \oplus C_{2,s_1} \subset C_{1,2} \oplus C_{2,s_1} \subset \cdots \subset W_q$$

The last case to treat is u > 2. Here we obtain

$$F(W_q) = C_{1,2} \oplus C_{2,2} = W_4,$$

which already fixes the final filtration between W_4 and W_q : everything must be of the form $C_{1,2} \oplus C_{2,j}$ for $2 \le j \le q-2$. The only task at hand is to compute $\eta(1)$, $\eta(2)$ and $\eta(3)$. In the case u = q-1, we have $F(W_4) = W_4$ and therefore $\eta(4) = 0$, which implies that $\eta(1) = \eta(2) = \eta(3) = 0$. We now assume u < q-1 from here on and compute

$$F(W_4) = \begin{cases} C_{1,0} \oplus C_{2,2} = W_2 & \text{if } 2 \leq q - v, \\ C_{1,1} \oplus C_{2,2} = W_3 & \text{if } 2 > q - v. \end{cases}$$

First, we treat the case $2 \le q - v$. From the expression for W_2 and the fact that a final filtration is a filtration, it follows what W_1 and W_3 must be. This gives the final filtration

$$0 \subset \cdots \subset C_{1,0} \oplus C_{2,2} \subset \cdots \subset C_{1,2} \oplus C_{2,2} \subset \cdots \subset W_q.$$

We finally treat the case 2 > q - v or equivalently $v \ge q - 1$. Applying *F* again gives

$$F(W_3) = C_{1,1} \oplus C_{2,1} = W_2, \text{ and}$$

$$F(W_2) = \begin{cases} C_{1,0} \oplus C_{2,1} = W_1 & \text{if } v = q - 1, \\ C_{1,1} \oplus C_{2,1} = W_2 & \text{if } v = q. \end{cases}$$

When v = q, we have $\eta(2) = 0$ and therefore $\eta(1) = 0$. On the other hand, in the case v = q - 1 we have constructed W_1 , W_2 and W_3 , which clearly suffices. In both cases, there is a final filtration

$$0 \subset C_{1,0} \oplus C_{2,1} \subset C_{1,1} \oplus C_{2,1} \subset C_{1,1} \oplus C_{2,2} \subset C_{1,2} \oplus C_{2,2} \subset \cdots \subset W_q. \quad \Box$$

5.3. Weyl group cosets. We now compute the permutation $\omega_{u,v} = \psi(\gamma_{u,v}) \in \mathbf{W}_q \subset \mathfrak{S}_{2q}$ using the results of Proposition 5.1. The resulting permutation represents the Ekedahl-Oort stratum of *M* in \mathscr{A}_q .

In the following theorem, only the action of $\omega_{u,v}$ on the integers $1 \leq i \leq q$ is given, as the remaining information can be retrieved using Equation (5.1.1).

Theorem 5.2. In the case (u, v) = (1, 2), we have $\omega_{u,v} = id$. For various other choices of u, v, the following tables describe the corresponding $\omega_{u,v}$:

	u = 1 and $v > 2$		
$\omega_{u,v}(i)$	$\begin{cases} i & i \\ q+1 & i \\ i-1 & i \\ q+2 & i \end{cases}$	$\begin{aligned} fi &< q+2-v\\ fi &= q+2-v\\ fq+2-v &< i < q\\ fi &= q. \end{aligned}$	

	u = 2 and $v = q$	$u = 2 and 1 < q + 1 - v \leq v - 2$	u = 2 and $q + 1 - v > v - 2$
$\omega_{u,v}(i)$	$\begin{cases} q+i & \text{if } i \leq 2\\ i-2 & \text{if } 2 < i \leq q. \end{cases}$	$\begin{cases} 1 & \text{if } i = 1 \\ q+1 & \text{if } i = 2 \\ i-1 & \text{if } 2 < i < q+2-v \\ q+2 & \text{if } i = q+2-v \\ i-2 & \text{if } q+2-v < i \leqslant q \end{cases}$	$\begin{cases} 1 & if \ i = 1 \\ q+1 & if \ i = 2 \\ i-1 & if \ 2 < i < q+3-v \\ q+2 & if \ i = q+3-v \\ i-2 & if \ q+3-v < i \leqslant q \end{cases}$

	2 < u < q - 1 and $v = q - 1$	2 < u < q - 1 and $v = q$	2 < u < q - 1 and $v < q - 1$	
$\omega_{u,v}(i)$	$\begin{cases} 1 & \text{if } i = 1 \\ q + i - 1 & \text{if } 1 < i \leq 4 \\ i - 3 & \text{if } 4 < i < q + 3 - u \\ q + 4 & \text{if } i = q + 3 - u \\ i - 4 & \text{if } q + 3 - u < i \leq q. \end{cases}$	$\begin{cases} q+i & if \ 1 \leqslant i \leqslant 2\\ i-2 & if \ i=3\\ q+3 & if \ i=4\\ i-3 & if \ 4 < i \leqslant q+3-u\\ q+4 & if \ i=q+3-u\\ i-4 & if \ q+3-u < i \leqslant q. \end{cases}$	$\begin{cases} i & \text{if } 1 \leqslant i \leqslant 2\\ q+i-2 & \text{if } 3 \leqslant i \leqslant 4\\ i-2 & \text{if } 4 < i \leqslant q+3-v\\ q+3 & \text{if } i=q+3-v\\ i-3 & \text{if } q+3-v < i < q+3-u\\ q+4 & \text{if } i=q+3-u\\ i-4 & \text{if } q+3-u < i \leqslant q. \end{cases}$	

	u = q - 1		
(i)	$\int q + i if \ 1 \leqslant i \leqslant 4$		
$\omega_{u,v}(\iota)$	$i-4$ if $4 < i \leq q$.		

Proof. In the case (u, v) = (1, 2), we have $\eta(q) = q$. Therefore the function η jumps at every integer $1 \le j \le q$. Hence we obtain $j_l = l$ and $i_m = m + q$. We conclude $\omega_{u,v} = id$.

In the case u = 1 and v > 2, we follow the final filtration given in Proposition 5.1. We compute

$$\eta(j) = \begin{cases} 1 & \text{if } j = 1, \\ 1 + \eta_2(j-1) & \text{if } 1 < j < q, \\ 1 + \eta_2(j-2) & \text{if } j = q \end{cases}$$

using the information we have on η_1 and η_2 . We see that η jumps everywhere except at the indices where $e_{1,2}$ and $e_{2,q+1-v}$ are added. The computation above shows where this happens: $i_1 = q + 2 - v$ and $i_2 = q$. This is the information needed to compute $\omega_{u,v}$:

$$\omega_{u,v}(i) = \begin{cases} i & \text{if } i < q+2-v, \\ q+1 & \text{if } i = q+2-v, \\ i-1 & \text{if } q+2-v < i < q, \\ q+2 & \text{if } i = q. \end{cases}$$

as asserted.

In the case u = 2, set $s_1 = \min\{v - 2, q + 1 - v\}$. Then by Proposition 5.1 it follows that

$$\eta(j) = \begin{cases} \eta_2(j) & \text{if } j = 1, \\ \eta_2(j-1) & \text{if } 1 < j \le s_1 + 1, \\ 1 + \eta_2(j-2) & \text{if } s_1 + 1 < j \le q. \end{cases}$$

The integers i_1 and i_2 correspond to the indices where $e_{1,1}$ and $e_{2,q+1-v}$ are added. In the case v = q, this implies $i_1 = 1$ and $i_2 = 2$. Otherwise, we have $i_1 = 2$. In the case $1 < q + 1 - v \leq v - 2$, we have $i_2 = q + 2 - v$. In the case q + 1 - v > v - 2, we have $i_2 = q + 3 - v$.

In the case 2 < u < q - 1 and v < q - 1, the final filtration in Proposition 5.1 implies

$$\eta(j) = \begin{cases} \eta_2(j) & \text{if } j \leq 2, \\ \eta_2(j-1) & \text{if } j = 3, \\ \eta_2(j-2) & \text{if } 3 < j \leq q. \end{cases}$$

Thus η jumps everywhere except at the indices $\{3, 4, q + 3 - v, q + 3 - u\}$, resulting in the permutation $\omega_{u,v}$ from the theorem.

Finally, in the case u > 2 and $v \ge q - 1$, Proposition 5.1 provides

$$\eta(j) = \begin{cases} \eta_2(j) & \text{if } j = 1, \\ \eta_2(j-1) & \text{if } j \in \{2,3\}, \\ \eta_2(j-2) & \text{if } 3 < j \leq q. \end{cases}$$

Now there are four indices where η does not jump: these are the indices when $e_{1,1}$, $e_{1,2}$, $e_{2,q+1-u}$ and $e_{2,q+1-v}$ are added. In each case, 2 and 4 are among these indices. In the case u = q - 1, the remaining indices are 1 and 3. In the case u < q - 1 and v = q, the remaining indices are 1 and q + 3 - u. Finally, in the case u < q - 1 and v = q - 1, the remaining indices are 3 and q + 3 - u.

5.4. Ekedahl-Oort strata contained in the supersingular locus. Using the main result of [Hoe09], we pinpoint exactly which $\omega_{u,v}$ represent an Ekedahl-Oort stratum of \mathscr{A}_q that is contained in \mathscr{A}_q^{ss} .

Proposition 5.3. The Ekedahl-Oort stratum $(\mathcal{A}_q)_{\omega_{u,v}}$ is contained in \mathcal{A}_q^{ss} if and only if u = 1 and $v < \lfloor q/2 \rfloor + 2$.

Proof. By [Hoe09, Theorem 1.2], the containment holds if and only if $\omega_{u,v}(i) = i$ for each $1 \le i \le \lceil q/2 \rceil$. For u = 1, by Theorem 5.2 this holds exactly when $\lceil q/2 \rceil < q + 2 - v$, i.e., when $v < \lfloor q/2 \rfloor + 2$.

We now show that containment is not possible for u > 1. In the case u = 2, we have $\omega_{u,v}(2) > q$, so the containment can only hold if $q \leq 2$. That, however, contradicts the assumption u = 2.

In the case u > 2, we also have $\omega_{u,v}(2) > q$, except when 2 < u < q - 1 and v < q - 1. In this case we have $\omega_{u,v}(3) > q$ and hence $\lceil q/2 \rceil < 3$. This implies $q \leq 4$, which contradicts $2 < u < v < q - 1 \leq 3$.

Corollary 5.4. Assume u = 1 and $v < \lfloor q/2 \rfloor + 2$. Then the Ekedahl-Oort stratum $\mathcal{M}(q-2,2)_{\gamma_{u,v}}$ is contained in $\mathcal{M}(q-2,2)^{ss}$.

Proof. Proposition 5.3 shows that $\omega_{u,v}$ is completely contained in \mathscr{A}_q^{ss} under these assumptions. Therefore any 4-tuple (A, λ, ι, ξ) in $\mathscr{M}(q - 2, 2)_{\gamma_{u,v}}$ has the property that (A, λ, ι) is supersingular. This implies that the 4-tuple is supersingular.

5.5. **Ekedahl-Oort strata on which** *F* and *V* are not nilpotent. If A[p] has a non-trivial subgroup scheme on which *F* or *V* act bijectively, then the slopes 0 and 1 occur in the Newton polygon of *A*. Thus *F* and *V* act nilpotently on supersingular abelian varieties (equivalently, their so-called *p*-*rank* is zero). By duality, *F* is nilpotent if and only if *V* is nilpotent. The following lemma shows how this is measured by the permutation $\omega_{u,v}$.

Lemma 5.5. The action of F is nilpotent on M if and only if $\omega_{u,v}(1) = 1$.

Proof. Assume the action of *F* is nilpotent on *M*. Then we must have $F(W_1) = 0$, because otherwise *F* acts bijectively on W_1 . Hence

$$\eta(1) = \dim(W_1 \cap M[F]) = 1$$

and therefore $j_1 = 1$ and $\omega_{u,v}(1) = 1$.

On the other hand, assume that *F* is not nilpotent on *M*. This implies that $F^n(M) \neq 0$ for every *n*. Since applying *F* gives a subspace in the canonical filtration, we infer that there is an l > 0 such that $F(W_l) = W_l$. Thus $\eta(l) = 0$ and, therefore, $i_1 = 1$ and $\omega_{u,v}(1) = q + 1$.

This allows us show that several Ekedahl-Oort strata cannot intersect the supersingular locus.

Proposition 5.6. Assume u > 1. Then the Ekedahl-Oort stratum $\mathcal{M}(q-2,2)_{\gamma_{u,q}}$ does not intersect $\mathcal{M}(q-2,2)^{ss}$.

Proof. By Lemma 5.5, it suffices to check that $\omega_{u,q}(1) > 1$. Appealing to Theorem 5.2, we conclude that this happens exactly when u > 1 and v = q. Since $(\mathscr{A}_q)_{\omega_{u,q}}$ does not intersect \mathscr{A}_q^{ss} , it follows that $\mathscr{M}(q-2,2)_{\gamma_{u,q}}$ does not intersect $\mathscr{M}(q-2,2)^{ss}$.

5.6. **Minimal Ekedahl-Oort strata of non-supersingular Newton strata.** We now describe another method that allows us to show Ekedahl-Oort strata are disjoint from the supersingular locus. This method is based on *minimal* Ekedahl-Oort strata, which are completely contained in a non-supersingular Newton stratum.

Definition 5.7. An Ekedahl-Oort stratum *S* of a Shimura variety of PEL type \mathcal{M} is *minimal* if $A[p^{\infty}] \cong B[p^{\infty}]$ holds for any $A, B \in S(\Bbbk)$.

It follows immediately from this definition that the Ekedahl-Oort stratum *S* is completely contained in one Newton stratum. Unfortunately, it is not known what the minimal Ekedahl-Oort strata of $\mathcal{M}(q-2,2)$ look like, or whether each Newton stratum of $\mathcal{M}(q-2,2)$ contains a (unique) minimal Ekedahl-Oort stratum, since the group GU(q-2,2) is not split for q > 4 (see [VW13]).

Luckily things are different for \mathcal{A}_q . Oort proved in [Oor05a, Oor05b] that each Newton stratum of \mathcal{A}_q contains a unique minimal Ekedahl-Oort stratum. Given a Newton polygon, results of [dJO00, 5.3] give an explicit description of the *p*-divisible group and the *p*-torsion group scheme of the minimal Ekedahl-Oort stratum contained in that Newton stratum. Hence, the minimal Ekedahl-Oort stratum of a non-supersingular Newton stratum does not intersect \mathcal{A}_q^{ss} , and we can use Theorem 5.2 to determine Ekedahl-Oort strata of $\mathcal{M}(q-2,2)$ that do not intersect $\mathcal{M}(q-2,2)^{ss}$.

Definition 5.8. Let *m* and *n* be non-negative integers. Define the mod-*p* Dieudonné module $M_{m,n} := \text{span}_{\Bbbk} \{e_0, \ldots, e_{m+n-1}\}$ with following action of *F* and *V*:

(5.6.1)
$$F(e_i) = \begin{cases} e_{i+n} & \text{if } i \leq m-1, \\ 0 & \text{else,} \end{cases}$$

(5.6.2)
$$V(e_i) = \begin{cases} e_{i+m} & \text{if } i \leq n-1, \\ 0 & \text{else.} \end{cases}$$

Let α be a Newton stratum of \mathscr{A}_q with slopes $\frac{n_1}{m_1+n_1}, \ldots, \frac{n_r}{m_r+n_r}$, where each slope may occur with multiplicity greater than or equal to 1. Oort shows in [Oor05a] that the unique minimal Ekedahl-Oort stratum in α has the Dieudonné module

$$(5.6.3) M_{\alpha} := \bigoplus_{l=1}^{\prime} M_{m_l, n_l}$$

The Dieudonné module M_{α} corresponds to a permutation $\omega_{\alpha} \in \mathbf{W}_{q}$, which we determine in the results below.

Lemma 5.9. $M_{m,n}$ has a final filtration given by

$$W_j^{m,n} := \operatorname{span}_{\Bbbk} \{ e_{m+n-j}, \dots, e_{m+n-1} \}$$

(5.6.4)
$$F(W_j^{m,n}) = W_{\max\{0,j-n\}'}^{m,n}$$

(5.6.5)
$$V^{-1}(W_j^{m,n}) = W_{\min\{m+n,j+m\}}^{m,n},$$

(5.6.6)
$$M_{m,n}[F] = W_n^{m,n}.$$

Proof. Equation (5.6.4) follows from Equation (5.6.1); all indices are shifted by *n* when *F* is applied, except if the resulting index exceeds m + n - 1, in which case *F* acts as 0. Similarly, Equation (5.6.5) is proved using Equation (5.6.2). Since the filtration $W_{\bullet}^{m,n}$ is stable under *F* and V^{-1} and $\dim_{\mathbb{K}}(W_j) = j$, it is a final filtration of $M_{m,n}$. Finally, Equation (5.6.6) also follows directly from Equation (5.6.1).

Given integers *m*, *n* and a word *w* in the letters *F* and V^{-1} , define the integer w(m, n) recursively as follows:

- If *w* is the empty word, then w(m, n) = m + n,
- $(Fw)(m,n) = \max\{0, w(m,n) n\},\$
- $(V^{-1}w)(m,n) = \min\{m+n, w(m,n)+m\}.$

Corollary 5.10. We have $w(M_{m,n}) = W_{w(m,n)}^{m,n}$.

Proof. Apply Lemma 5.9 repeatedly.

Recall the definition of the mod-*p* Dieudonné module M_{α} from Equation (5.6.3). Combining this with Corollary 5.10 gives, for any word *w* in the alphabet {*F*, *V*⁻¹}, the formula

$$w(M_{\alpha}) = w\left(\bigoplus_{l=1}^{r} M_{m_l,n_l}\right) = \bigoplus_{l=1}^{r} w(M_{m_l,n_l}) = \bigoplus_{l=1}^{r} W_{w(m_l,n_l)}^{m_l,n_l}$$

We now introduce the function $\eta_{\alpha}(j) = \dim (M_{\alpha} \cap M[F])$. We have the following restrictions on η_{α} :

$$\eta_{\alpha}\left(\sum_{l=1}^{r} w(m_{l}, n_{l})\right) = \dim_{\mathbb{k}}\left(\left(\bigoplus_{l=1}^{r} W_{w(m_{l}, n_{l})}^{m_{l}, n_{l}}\right) \cap M_{\alpha}[F]\right)$$
$$= \dim_{\mathbb{k}}\left(\bigoplus_{l=1}^{r} W_{w(m_{l}, n_{l})}^{m_{l}, n_{l}} \cap W_{n_{l}}^{m_{l}, n_{l}}\right)$$
33

	1
	L
	L

$$= \sum_{l=1}^{r} \min\{n_l, w(m_l, n_l)\}.$$

By letting *w* range over sufficiently many words, we obtain enough restrictions to determine the nondecreasing function $\eta_{\alpha} : \{1, ..., 2q\} \rightarrow \{1, ..., q\}$ uniquely. This is because η_{α} is determined by the canonical filtration of M_{α} . Finally, this function η_{α} gives rise to a permutation $\omega_{\alpha} \in \mathbf{W}_q$ via the steps given in [Moo01, 3.6].

Proposition 5.11. Let n_1, \ldots, n_r be non-negative integers satisfying the following conditions:

- (*i*) $\sum_{l=1}^{r} n_l = q$,
- (*ii*) $GCD(n_l, n_{r+1-l}) = 1$ for every $l \in \{1, ..., r\}$,
- (iii) there exists $l \in \{1, ..., r\}$ such that $n_l + n_{r+1-l} \neq 2$.

Define the mod-p Dieudonné module $M_{\alpha} := \bigoplus M_{n_{r+1-l},n_l}$ and let $\omega_{\alpha} \in \mathfrak{S}_{2q}$ be as above. Then the Ekedahl-Oort stratum $\mathscr{A}_{\omega_{\alpha}}$ does not intersect \mathscr{A}_{q}^{ss} .

Proof. By conditions (i) and (ii), there exists a symmetric Newton polygon from (0,0) to (2q,q) with slopes $\frac{n_l}{n_l+n_{r+1-l}}$ (possibly with multiplicity greater than 1). By condition (iii), this Newton polygon has a slope that is not 1/2. By construction, M_{α} and ω_{α} correspond to a minimal Ekedahl-Oort stratum contained in a Newton stratum that is not supersingular. Thus it does not intersect \mathscr{A}_q^{ss} .

Corollary 5.12. Suppose $\omega_{u,v} = \omega_{\alpha}$ under the conditions of Proposition 5.11. Then the Ekedahl-Oort stratum of $\mathcal{M}(q-2,2)_{\gamma_{u,v}}$ does not intersect $\mathcal{M}(q-2,2)^{ss}$.

Proof. This follows directly from Proposition 5.11, as (forgetting) the unitary structure does not affect whether an abelian variety is supersingular or not. \Box

6. Ekedahl-Oort Strata of $\mathcal{M}(3,2)$ and the Supersingular Locus

Recall our motivating question.

Question 1.1. Which Ekedahl-Oort strata of $\mathcal{M}(q - a, a)$ intersect the supersingular locus?

As discussed in the introduction, the lowest-dimensional unitary Shimura variety without a complete answer to the above question is $\mathcal{M}(3, 2)$. This Shimura variety is also the lowest-dimensional unitary Shimura variety that is not of "Coxeter type" (see [GH15]). Because $\mathcal{M}(3, 2)$ is not of Coxeter type, by [GH15] it is known that there must exist some Ekedahl-Oort stratum that intersects the supersingular locus, but is not entirely contained in the supersingular locus. In this sense, the interaction between the Ekedahl-Oort strata and the supersingular locus is significantly more complicated for $\mathcal{M}(3, 2)$ than for previously studied unitary Shimura varieties.

While the techniques explored in the previous sections (including closure properties of the Ekedahl-Oort strata, the product maps, and the forgetful map) can be used to study the intersection of Ekedahl-Oort strata with the supersingular locus for unitary Shimura varieties of signature (q - 2, 2) for any $q \ge 2$, in this section we consider $\mathcal{M}(3, 2)$, i.e. q = 5, as a concrete example.

Of the ten Ekedahl-Oort strata of $\mathcal{M}(3, 2)$, all but two can be studied with the methods developed so far; the remaining strata are those indexed by $\gamma_{1,4}$ and $\gamma_{3,4}$. We will first show that these strata do in fact intersect the supersingular locus, by directly constructing a point in the intersection. Then, we will combine this approach with the previous techniques to give a complete answer to Question 1.1 for the Shimura variety $\mathcal{M}(3, 2)$, in Theorem 6.7.

6.1. **Explicit Construction of Points.** In this section we explicitly construct k-points in the intersections $\mathcal{M}(3,2)_{\gamma_{1,4}} \cap \mathcal{M}(3,2)^{ss}$ and $\mathcal{M}(3,2)_{\gamma_{3,4}} \cap \mathcal{M}(3,2)^{ss}$. To do this, we will first use *p*-adic Dieudonné theory to construct the *p*-divisible groups of these points; then we will use the *p*-divisible groups and Rapoport-Zink uniformization to construct points of $\mathcal{M}(3,2)$.

Let $\mathbb{Z}_p = W(\mathbb{k})$ be the ring of Witt vectors of \mathbb{k} , denote by $\operatorname{Frob} = W(\operatorname{Frob}_{\mathbb{k}})$ the lift of $\operatorname{Frob}_{\mathbb{k}}$ to \mathbb{Z}_p , and let $\mathbb{Q}_p = \mathbb{Z}_p[\frac{1}{p}]$. Note that \mathbb{Q}_p is isomorphic to the completion of the maximal unramified extension of \mathbb{Q}_p , with ring of integers isomorphic to \mathbb{Z}_p , and that the residue field of \mathbb{Z}_p is \mathbb{k} . Let φ_1, φ_2 be the two embeddings of \mathcal{O}_K into \mathbb{Q}_p .

Definition 6.1. For any scheme *S* over \Bbbk , a *unitary p-divisible group of signature* (3,2) over *S* is a triple (X, ι_X, λ_X) , where:

- *X* is a *p*-divisible group over *S*.
- $\iota_X : \mathcal{O}_K \otimes_\mathbb{Z} \mathbb{Z}_p \to \text{End}(X)$ is an action satisfying the signature (3,2) condition, for all $a \in \mathcal{O}_K$:

charpol($\iota(a) \mid \operatorname{Lie}(X)$) = $(T - \varphi_1(a))^3 (T - \varphi_2(a))^2 \in \mathbb{Z}_p[T].$

• $\lambda_X : X \to X^{\vee}$ is a *p*-principal polarization, meeting the following \mathcal{O}_K -linearity condition, for all $a \in \mathcal{O}_K$:

$$\lambda_X \circ \iota_X(a) = \iota_X(\overline{a})^{\vee} \circ \lambda_X.$$

Over an algebraically closed field, we may study unitary *p*-divisible groups linear-algebraically:

Definition 6.2. A unitary *p*-adic Dieudonné module of signature (3,2) over \Bbbk is a tuple $(M, M = M_1 \oplus M_2, F, V, \langle \cdot, \cdot \rangle)$, where:

- *M* is a free \mathbb{Z}_p -module of rank 10,
- $M = M_1 \oplus M_2$ is a decomposition into rank-5 summands,
- $F : M \to M$ is a Frob-semilinear operator, $V : M \to M$ is a Frob⁻¹-semilinear operator, with $F \circ V = V \circ F = p$,
- $\langle \cdot, \cdot \rangle$ is a perfect alternating \mathbb{Z}_p -bilinear pairing on M such that $\langle Fx, y \rangle = \langle x, Vy \rangle^{\text{Frob}}$, for all $x, y \in M$,
- $\dim_{\Bbbk}(M_1/FM_2) = 3$ and $\dim_{\Bbbk}(M_2/FM_1) = 2$,

and further *F* and *V* are homogeneous of degree 1 with respect to the decomposition $M = M_1 \oplus M_2$, and M_1 and M_2 are each totally isotropic with respect to $\langle \cdot, \cdot \rangle$.

Using contravariant *p*-adic Dieudonné theory, there is an equivalence of categories between unitary *p*-divisible groups of signature (3,2) over k and unitary *p*-adic Dieudonné module of signature (3,2) over k. When no confusion is possible, we will drop the extra structure and abbreviate unitary *p*-divisible groups simply as *X* and unitary *p*-adic Dieudonné modules simply as *M*.

Lemma 6.3. Let $G_{\gamma_{1,4}}$ be the *p*-torsion group scheme occurring as *p*-torsion subgroup for points in the Ekedahl-Oort stratum $\mathcal{M}(3,2)_{\gamma_{1,4}}$. There exists a supersingular unitary *p*-divisible group $X_{1,4}$ of signature (3,2) over \Bbbk such that $X_{1,4}[p] \cong G_{\gamma_{1,4}}$, respecting action and polarization.

Proof. We aim to construct a supersingular unitary *p*-divisible group $X_{1,4}$ of signature (3,2) such that $X_{1,4}[p] \cong G_{\gamma_{1,4}}$. Using contravariant *p*-adic Dieudonné theory, it suffices to construct a unitary *p*-adic Dieudonné module *M* of signature (3,2) such that all of the slopes of the isocrystal $M[\frac{1}{p}]$ are equal to $\frac{1}{2}$ and such that M/pM is isomorphic to the mod-*p* Dieudonné module of $G_{\gamma_{1,4}}$.

Let *M* be the free \mathbb{Z}_p -module with basis $\{e_i, f_i\}_1^5$. Let M_1 be the submodule spanned by $\{e_i\}_1^5$ and let M_2 be the submodule spanned by $\{f_i\}_1^5$. Define *F* (resp. *V*) as the Frob-semilinear (resp. Frob⁻¹-semilinear) operator defined on the basis of *M* as in Table 1 below.

$F(e_1) = f_5$	$V(e_1) = pf_2$	$F(f_1) = -pe_5$	$V(f_1) = e_2$
$F(e_2) = pf_1$	$V(e_2) = pf_3$	$F(f_2) = e_1$	$V(f_2) = e_3$
$F(e_3) = pf_2$	$V(e_3) = f_4$	$F(f_3) = e_2$	$V(f_3) = pe_4$
$F(e_4) = f_3$	$V(e_4) = f_5$	$F(f_4) = pe_3$	$V(f_4) = pe_5$
$F(e_5) = f_4$	$V(e_5) = -f_1$	$F(f_5) = pe_4$	$V(f_5) = pe_1$

TABLE 1. *F* and *V* on *M* for $\gamma_{1,4}$

We define an alternating pairing $\langle \cdot, \cdot \rangle$ on M by the condition that $\langle e_i, f_j \rangle = (-1)^{i-1} \delta_{ij}$ and claim that $(M, M = M_1 \oplus M_2, F, V, \langle \cdot, \cdot \rangle)$ is a unitary *p*-adic Dieudonné module module of signature (3, 2). Indeed:

- $M = M_1 \oplus M_2$, and M_1 and M_2 both have rank 5, so M has rank 10.
- The operators *F* and *V* are homogeneous of degree 1 with respect to the decomposition $M = M_1 \oplus M_2$. Let \mathbf{A}_F and \mathbf{A}_V be the matrices given by the action of *F* and *V*, respectively, on the chosen basis. Since \mathbf{A}_F and \mathbf{A}_V have integer entries, to check that $F \circ V = V \circ F = p$, it suffices to verify that $\mathbf{A}_F \mathbf{A}_V = \mathbf{A}_V \mathbf{A}_F = p \mathrm{Id}$, which is true by construction.
- The condition that ⟨e_i, f_j⟩ = (−1)^{i−1}δ_{ij} extends uniquely to a perfect alternating Ž_p-bilinear pairing on *M*, and under this pairing M₁ and M₂ are each totally isotropic.

Let **B** be the matrix of this alternating form, and note that **B** has integer entries. To check that $\langle Fx, y \rangle = \langle x, Vy \rangle^{\text{Frob}}$, it suffices to verify that $\mathbf{A}_F^T \mathbf{B} = \mathbf{B} \mathbf{A}_V$, which can be verified using the description of *F* and *V* in Table 1 and the definition of $\langle \cdot, \cdot \rangle$.

• From the definition of *F*, we have that $M_1/FM_2 \cong \text{Span}_{\Bbbk}\{e_3, e_4, e_5\}$ and $M_1/FM_2 \cong \text{Span}_{\Bbbk}\{f_1, f_2\}$. In particular, $\dim_{\Bbbk}(M_1/FM_2) = 3$ and $\dim_{\Bbbk}(M_2/FM_1) = 2$.

Thus, by contravariant *p*-adic Dieudonné theory, *M* defines a unitary *p*-divisible group $X_{1,4}$ of signature (3, 2).

We use Lemma 6.12 in [Zin84] to compute the slopes of the isocrystal $M[\frac{1}{p}]$. Note that for any positive integer *m*, we find:

$$V^{10m}(M) = p^{5m}M,$$

and so $\frac{1}{10m} \max\{k \in \mathbb{Z} : V^{10m} M \subset p^k M\} = \frac{1}{2}$. Therefore,

$$\lim_{n\to\infty}\frac{1}{n}\max\{k\in\mathbb{Z}:V^nM\subset p^kM\}=\frac{1}{2},$$

and by [Zin84] all slopes of $M[\frac{1}{p}]$ are equal to $\frac{1}{2}$. Accordingly, the *p*-divisible group $X_{1,4}$ is supersingular.

Define *N* as $M/pM \cong \text{Span}_{\Bbbk} \{e_i, f_i\}_{1}^{5}$, with splitting $N = N_1 \oplus N_2$ and *F* and *V* operators induced from those on *M*. We will follow the procedure of Theorem 2.2 to compute the permutation $\omega \in \mathbf{W}(3,2)$ associated to *N*. Following from the definition of *F* and *V*, the Dieudonné module *N* has final filtration:

$$0 \subset \langle e_2 \rangle \subset \langle e_2, f_5 \rangle \subset \langle e_2, f_3, f_5 \rangle \subset \langle e_2, f_3, f_4, f_5 \rangle \subset \langle e_1, e_2, f_3, f_4, f_5 \rangle \subset \langle e_1, e_2, f_1, f_3, f_4, f_5 \rangle \subset \langle e_1, e_2, e_4, f_1, f_3, f_4, f_5 \rangle \subset \langle e_1, e_2, e_3, e_4, f_1, f_3, f_4, f_5 \rangle \subset \langle e_1, e_2, e_3, e_4, e_5, f_1, f_3, f_4, f_5 \rangle \subset N$$

Intersecting with N_1 gives the filtration $C_{1,\bullet}$:

$$0 \subset \langle e_2 \rangle \subset \langle e_1, e_2 \rangle \subset \langle e_1, e_2, e_4 \rangle \subset \langle e_1, e_2, e_3, e_4 \rangle \subset N_1$$

If we define the function $\eta_1(j) = \dim(C_{1,j} \cap N[F])$, then:

$$\eta_1(1) = \eta_1(2) = \eta_1(3) = 1,$$

 $\eta_1(4) = \eta_1(5) = 2.$

The permutation ω corresponding to η is (2,3,4). As $\gamma_{1,4}$ is also equal to (2,3,4), the mod-p Dieudonné module of $G_{\gamma_{1,4}}$ is isomorphic to N. Therefore, by contravariant Dieudonné theory, $X_{1,4}[p] \cong G_{\gamma_{1,4}}$.

Lemma 6.4. Let $G_{\gamma_{3,4}}$ be the p-torsion group scheme occurring as the p-torsion subgroup for points in the Ekedahl-Oort stratum $\mathcal{M}(3,2)_{\gamma_{3,4}}$. There exists a supersingular unitary p-divisible group $X_{3,4}$ of signature (3,2) over \Bbbk such that $X_{3,4}[p] \cong G_{\gamma_{3,4}}$, respecting action and polarization.

Proof. Similar to the proof of Lemma 6.3 let M be the free \mathbb{Z}_p -module with basis $\{e_i, f_i\}_1^5$. Let M_1 be the submodule spanned by $\{e_i\}_1^5$ and let M_2 be the submodule spanned by $\{f_i\}_1^5$. Define F (resp. V) as the Frob-semilinear (resp. Frob⁻¹-semilinear) operator defined on the basis of M as in Table 2 below, and let $\langle \cdot, \cdot \rangle$ be the alternating form on M defined by the condition that $\langle e_i, f_i \rangle = (-1)^{i-1} \delta_{ij}$.

$F(e_1) = f_5$	$V(e_1) = pf_2$	$F(f_1) = -pe_5$	$V(f_1) = e_2$
$F(e_2) = pf_1$	$V(e_2) = f_3$	$F(f_2) = e_1$	$V(f_2) = pe_3$
$F(e_3) = f_2$	$V(e_3) = f_4$	$F(f_3) = pe_2$	$V(f_3) = pe_4$
$F(e_4) = f_3$	$V(e_4) = pf_5$	$F(f_4) = pe_3$	$V(f_4) = e_5$
$F(e_5) = pf_4$	$V(e_5) = -f_1$	$F(f_5) = e_4$	$V(f_5) = pe_1$

TABLE 2. *F* and *V* on *M* for $\gamma_{3,4}$

By exactly analogous computations, M defines a supersingular p-divisible group $X_{3,4}$.

Define *N* as $M/pM \cong \text{Span}_{\Bbbk} \{e_i, f_i\}_{1}^{5}$, with splitting $N = N_1 \oplus N_2$ and *F* and *V* operators induced from those on *M*. The Dieudonné module *N* has final filtration:

$$0 \subset \langle f_3 \rangle \subset \langle e_4, f_3 \rangle \subset \langle e_4, f_3, f_5 \rangle \subset \langle e_1, e_4, f_3, f_5 \rangle \subset \langle e_1, e_4, f_2, f_3, f_5 \rangle \subset \langle e_1, e_2, e_4, f_2, f_3, f_5 \rangle \subset \langle e_1, e_2, e_4, f_1, f_2, f_3, f_5 \rangle \subset \langle e_1, e_2, e_4, e_5, f_1, f_2, f_3, f_5 \rangle \subset \langle e_1, e_2, e_4, e_5, f_1, f_2, f_3, f_5 \rangle \subset N.$$

Intersecting with N_1 gives the filtration $C_{1,\bullet}$:

$$0 \subset \langle e_4 \rangle \subset \langle e_1, e_4 \rangle \subset \langle e_1, e_2, e_4 \rangle \subset \langle e_1, e_2, e_4, e_5 \rangle \subset N_1.$$

The function $\eta_1(j) = \dim(C_{1,j} \cap N[F])$ is then given by

$$\begin{split} \eta_1(1) &= \eta_1(2) = 0, \\ \eta_1(3) &= 1, \\ \eta_1(4) &= \eta_1(5) = 2. \end{split}$$

The permutation ω corresponding to η is (13)(24). As $\gamma_{3,4}$ is also equal to (13)(24), $X_{3,4}[p] \cong G_{\gamma_{3,4}}$.

Using Lemmas 6.3 and 6.4, we will show that the Ekedahl-Oort strata $\mathcal{M}(3,2)_{\gamma_{1,4}}$ and $\mathcal{M}(3,2)_{\gamma_{3,4}}$ intersect the supersingular locus. Recall that the supersingular locus $\mathcal{M}(3,2)^{ss}$ is uniformized by a formal scheme called a Rapoport-Zink space. As a framing object, let $(\mathbb{X}, \iota_{\mathbb{X}}, \lambda_{\mathbb{X}})$ be a fixed supersingular unitary *p*-divisible group of signature (3, 2) over \Bbbk .

Definition 6.5. For any scheme *S* over \Bbbk , denote by $\mathcal{N}(3,2)(S)$ the set of isomorphism classes of tuples $(X, \iota_X, \lambda_X, \rho_X)$, where:

- (X, ι_X, λ_X) is a unitary *p*-divisible group of signature (3, 2) over *S*,
- $\rho_X : X \to X$ is an \mathcal{O}_K -linear quasi-isogeny identifying λ_X and λ_X up to scaling in \mathbb{Q}_p^{\times} .

By [RZ96], the functor defined above is represented by a formal scheme over k which is locally formally of finite type; we will also denote the underlying reduced scheme of this representing object (a "signature (3, 2) unitary Rapoport-Zink space") as $\mathcal{N}(3, 2)$.

Proposition 6.6. The Ekedahl-Oort strata $\mathcal{M}(3,2)_{\gamma_{1,4}}$ and $\mathcal{M}(3,2)_{\gamma_{3,4}}$ have non-empty intersection with the supersingular locus. Moreover, $\mathcal{M}(3,2)_{\gamma_{3,4}}$ intersects the supersingular locus without being contained in the supersingular locus.

Proof. By the uniformizaton theorem of Rapoport and Zink [RZ96], there exist groups $\{\Gamma_j\}_{j=1}^n$ (arising as subgroups of the \mathbb{Q}_p -points of the algebraic group defining the automorphisms of \mathbb{X} , and depending on the level structure implicit in the definition of $\mathcal{M}(3,2)$) such that there is an isomorphism of schemes over \Bbbk :

$$\bigsqcup_{j=1}^{n} \mathcal{N}(3,2) / \Gamma_{j} \cong \mathcal{M}(3,2)^{ss}.$$

In particular, there is a surjection of k-points:

$$\bigsqcup_{j=1}^{n} \mathscr{N}(3,2)(\Bbbk) \twoheadrightarrow \mathscr{M}(3,2)^{ss}(\Bbbk).$$

Let $G_{\gamma_{1,4}}$ be the *p*-torsion group scheme (with extra structure) occurring as *p*-torsion subgroup for points in the Ekedahl-Oort stratum $\mathcal{M}(3,2)_{\gamma_{1,4}}$. By Lemma 6.3, there exists a supersingular unitary *p*-divisible group $X_{1,4}$ of signature (3,2) over k such that $X_{1,4}[p] \cong G_{\gamma_{1,4}}$.

By Lemma 6.1 of [VW11], there exists a quasi-isogeny $\rho_{X_{1,4}} : X_{1,4} \to \mathbb{X}$ that is \mathcal{O}_K -linear and identifies the polarizations, up to \mathbb{Q}_p^{\times} -scaling. That is, $(X_{1,4}, \iota_{X_{1,4}}, \lambda_{X_{1,4}}, \rho_{X_{1,4}})$ defines a k-point of $\mathcal{N}(3,2)$. Let $(A_{1,4}, \iota_{A_{1,4}}, \lambda_{A_{1,4}}, \xi_{A_{1,4}})$ be the image of $(X_{1,4}, \iota_{X_{1,4}}, \lambda_{X_{1,4}}, \rho_{X_{1,4}})$ in $\mathcal{M}(3,2)^{ss}(\mathbb{k})$.

Since $A_{1,4}[p] \cong X_{1,4}[p] \cong G_{\gamma_{1,4}}$, as the *p*-torsion group schemes equipped with extra structure, the abelian variety $(A_{1,4}, \iota_{A_{1,4}}, \lambda_{A_{1,4}}, \xi_{A_{1,4}})$ lies in the Ekedahl-Oort stratum indexed by $\gamma_{1,4}$. That is, $(A_{1,4}, \iota_{A_{1,4}}, \lambda_{A_{1,4}}, \xi_{A_{1,4}})$ is an explicit point in the intersection $\mathcal{M}(3, 2)_{\gamma_{1,4}} \cap \mathcal{M}(3, 2)^{ss}$.

Similarly, the *p*-divisible group $X_{3,4}$ of Lemma 6.4 gives rise to an explicit point in the intersection $\mathcal{M}(3,2)_{\gamma_{3,4}} \cap \mathcal{M}(3,2)^{ss}$.

Finally, recall that dim $(\gamma_{u,v}) = u + v - 3$ and dim $(\mathcal{M}(a, b)^{ss}) = \lfloor \frac{ab}{2} \rfloor$, so dim $(\gamma_{3,4}) = 4 > 3 = \lfloor \frac{3 \times 2}{2} \rfloor = \dim(\mathcal{M}(3, 2)^{ss})$. Thus, $\mathcal{M}(3, 2)_{\gamma_{3,4}} \not\subseteq \mathcal{M}(3, 2)^{ss}$.

6.2. **Results for** $\mathcal{M}(3,2)$ **.**

Theorem 6.7. The Ekedahl-Oort strata of $\mathcal{M}(3,2)$ that intersect the supersingular locus are those indexed by

*γ*1,2,*γ*1,3,*γ*1,4,*γ*2,3,*γ*1,5,*γ*3,4.

Furthermore, the stratum indexed by $\gamma_{3,4}$ intersects but is not contained in the supersingular locus.

The remaining Ekedahl-Oort strata of $\mathcal{M}(3,2)$ *, indexed by*

72,4, 72,5, 73,5, 74,5

do not intersect the supersingular locus.

Proof. We observe the following:

• The stratum $\mathcal{M}(3,2)_{\gamma_{1,2}}$ is entirely contained in the supersingular locus by Corollary 5.4, because u = 1 and $v = 2 < \lfloor 5/2 \rfloor + 2$. This follows from properties of the forgetful map $\Psi : \mathcal{M}(3,2) \to \mathcal{A}_5$. By Theorem 5.2,

$$\Psi(\mathscr{M}(3,2)_{\gamma_{1,2}}) \subseteq (\mathscr{A}_5)_{\mathrm{id}},$$

and as $(\mathscr{A}_5)_{id} \subseteq \mathscr{A}_5^{ss}$, we have $\mathscr{M}(3,2)_{\gamma_{1,2}} \subseteq \mathscr{M}(3,2)^{ss}$ as well.

• Similarly, $\mathcal{M}(3,2)_{\gamma_{1,3}}$ is entirely contained in the supersingular locus by Corollary 5.4, because u = 1 and $v = 3 < \lfloor 5/2 \rfloor + 2$. In this case,

$$\Psi(\mathscr{M}(3,2)_{\gamma_{1,2}}) \subseteq (\mathscr{A}_5)_{(46)(57)},$$

which is also contained in \mathscr{A}_{5}^{ss} , and so $\mathscr{M}(3,2)_{\gamma_{1,3}} \subseteq \mathscr{M}(3,2)^{ss}$.

- The stratum *M*(3,2)_{γ1,4} has a non-empty intersection with the supersingular locus because we explicitly constructed a k-point of the intersection in Proposition 6.6.
- The stratum $\mathcal{M}(3,2)_{\gamma_{2,3}}$ has a non-empty intersection with the supersingular locus by applying either Corollary 4.11 or Corollary 4.17. For Corollary 4.11, we may take m = 2, n = 1, a = 1, and b = 0. This is because if

$$\Phi: \mathscr{M}(2,1) \times \mathscr{M}(1,1) \to \mathscr{M}(3,2)$$

is the natural product map, then

$$\Phi(\mathscr{M}(2,1)_{\delta_1} \times \mathscr{M}(1,1)_{\delta_0}) \subseteq \mathscr{M}(3,2)_{\gamma_{2,3}}.$$

Since both $\mathcal{M}(2,1)_{\delta_1}$ and $\mathcal{M}(1,1)_{\delta_0}$ intersect the supersingular locus, $\mathcal{M}(3,2)_{\gamma_{2,3}}$ does as well. (Corollary 4.17 is a reflection of a similar fact using the product map on $\mathcal{M}(2,2) \times \mathcal{M}(1,0)$.)

• The stratum $\mathcal{M}(3,2)_{\gamma_{1,5}}$ has a non-empty intersection with the supersingular locus by applying Corollary 4.17. If

$$\Phi: \mathscr{M}(2,2) \times \mathscr{M}(1,0) \to \mathscr{M}(3,2)$$

is the natural product map, then

$$\Phi(\mathscr{M}(2,2)_{\gamma_{1,4}} \times \mathscr{M}(1,0)_{\delta_0}) \subseteq \mathscr{M}(3,2)_{\gamma_{1,5}}$$

Since (by Proposition 4.18) both $\mathcal{M}(2,2)_{\gamma_{1,4}}$ and $\mathcal{M}(1,0)_{\delta_0}$ intersect the supersingular locus, $\mathcal{M}(3,2)_{\gamma_{1,5}}$ does as well.

- The stratum $\mathcal{M}(3,2)_{\gamma_{2,4}}$ does not intersect the supersingular locus by the minimality result of Corollary 5.12. Letting $(n_1, n_2, n_3) = (1, 1, 3)$, we can compute the associated minimal stratum ω_{α} using Proposition 5.11, with the slope sequence $\alpha = [1/4, 1/2, 3/4]$. We can then use Theorem 5.2 to verify that $\psi(\gamma_{2,4}) = \omega_{\alpha}$. Thus $(\mathcal{A}_q)_{\psi(\gamma_{2,4})}$ is completely contained in the non-supersingular Newton stratum \mathcal{A}_q^{α} and therefore $\mathcal{M}(3, 2)_{\gamma_{2,4}}$ does not intersect $\mathcal{M}(3, 2)^{ss}$.
- The stratum $\mathcal{M}(3,2)_{\gamma_{2,5}}$ does not intersect the supersingular locus by Proposition 5.6, because in this case u = 2 > 1 and v = q. Recall that Proposition 5.6 is reflecting the fact that *F* does not act nilpotently on the Dieudonné module corresponding to $\gamma_{2,5}$, and so there cannot exist a supersingular abelian variety *A* with *p*-torsion group scheme indexed by $\gamma_{2,5}$.

One could also apply Corollary 5.12 to see that $\mathcal{M}(3,2)_{\gamma_{2,5}}$ does not intersect the supersingular locus. In this case,

$$\Psi(\mathscr{M}(3,2)_{\gamma_{2,5}}) \subseteq (\mathscr{A}_5)_w,$$

The stratum M(3,2)_{γ3,4} intersects the supersingular locus, because we explicitly constructed a k-point of the intersection in Proposition 6.6. This Ekedahl-Oort stratum is not completely contained in the supersingular locus, because dim(M(3,2)_{γ3,4}) = 4, while dim(M(3,2)^{ss}) = 3.

- Similar to $\mathcal{M}(3,2)_{\gamma_{2,5}}$, the stratum $\mathcal{M}(3,2)_{\gamma_{3,5}}$ does not intersect the supersingular locus by Proposition 5.6, reflecting the fact that *F* does not act nilpotently on the Dieudonné module corresponding to $\gamma_{3,5}$.
- The stratum $\mathcal{M}(3,2)_{\gamma_{4,5}}$ does not intersect the supersingular locus by Proposition 5.6, reflecting the fact that *F* does not act nilpotently on the that Dieudonné module corresponding to $\gamma_{4,5}$. Alternately, one can see that this stratum does not intersect the supersingular locus by applying Corollary 5.12. In this case,

$$\Psi(\mathscr{M}(3,2)_{\gamma_{4,5}})\subseteq (\mathscr{A}_5)_w,$$

where $w = (1 \ 6 \ 10 \ 5)(2 \ 7)(3 \ 8)(4 \ 9)$, and $(\mathscr{A}_5)_w$ does not intersect the supersingular locus of \mathscr{A}_5 , because it is a minimal Ekedahl-Oort stratum for the slope sequence $\left[\frac{0}{1}, \frac{0}{1}, \frac{0}{1}, \frac{1}{2}, \frac{1}{1}, \frac{1}{1}, \frac{1}{1}, \frac{1}{1}\right]$.

We summarize the results of Theorem 6.7 in the following table.

Stratum	Dim.	Intersection w/ ss locus	Method(s)	Reference(s)
Ŷ1,2	0	containment	Forgetful	Corollary 5.4
Ŷ1,3	1	containment	Forgetful	Corollary 5.4
$\gamma_{1,4}$	2	non-empty	Explicit	Proposition 6.6
Y2,3	2	non-empty	Product _{1,1} , Product _{2,0}	Corollary 4.11, Corollary 4.17
$\gamma_{1,5}$	3	non-empty	Product _{2,0}	Corollary 4.17
$\gamma_{2,4}$	3	empty	Minimality	Corollary 5.12
Y2,5	4	empty	Nilpotence, Minimality	Proposition 5.6, Corollary 5.12
$\gamma_{3,4}$	4	proper, non-empty	Explicit	Proposition 6.6
Y3,5	5	empty	Nilpotence	Proposition 5.6
$\gamma_{4,5}$	6	empty	Nilpotence, Minimality	Proposition 5.6, Corollary 5.12

TABLE 3. Classification of Ekedahl-Oort strata of $\mathcal{M}(3,2)$

REFERENCES

- [BB06] A. Bjorner and F. Brenti, *Combinatorics of Coxeter Groups*, Graduate Texts in Mathematics, Springer Berlin Heidelberg, 2006.
- [dJO00] Aise Johan de Jong and Frans Oort, *Purity of the stratification by newton polygons*, Journal of the American Mathematical Society **13** (2000), no. 1, 209–241.
- [FHI23] Maria Fox, Benjamin Howard, and Naoki Imai, *Rapoport-zink spaces of type gu*(2,*n*-2), 2023.
- [GH15] Ulrich Görtz and Xuhua He, *Basic loci of Coxeter type in Shimura varieties*, Camb. J. Math. **3** (2015), no. 3, 323–353. MR 3393024
- [He07] Xuhua He, *The G-stable pieces of the wonderful compactification*, Trans. Amer. Math. Soc. **359** (2007), no. 7, 3005–3024. MR 2299444
- [Hoe09] Maarten Hoeve, Ekedahl-oort strata in the supersingular locus, Journal of the London Mathematical Society 81 (2009), no. 1, 129–141.
- [HP14] Benjamin Howard and Georgios Pappas, *On the supersingular locus of the GU*(2, 2) *shimura variety*, Algebra Number Theory **8** (2014), no. 7, 1659–1699.

- [Kot92] Robert E. Kottwitz, Points on some Shimura varieties over finite fields, J. Amer. Math. Soc. 5 (1992), no. 2, 373–444. MR 1124982
- [Moo01] Ben Moonen, Group Schemes with Additional Structures and Weyl Group Cosets, pp. 255–298, Birkhäuser Basel, Basel, 2001.
- [Oor05a] Frans Oort, Minimal p-divisible groups, Annals of Mathematics 161 (2005), 1021–1036.
- [Oor05b] _____, Simple p-kernels of p-divisible groups, Advances in Mathematics 198 (2005), 275–310.
- [PWZ11] Richard Pink, Torsten Wedhorn, and Paul Ziegler, *Algebraic zip data*, Doc. Math. **16** (2011), 253–300. MR 2804513
- [RZ96] M. Rapoport and Th. Zink, Period spaces for p-divisible groups, Annals of Mathematics Studies, vol. 141, Princeton University Press, Princeton, NJ, 1996. MR 1393439
- [The23] The Sage Developers, Sagemath, the Sage Mathematics Software System (Version 9.8), 2023, https://www.sagemath.org.
- [VW11] Inken Vollaard and Torsten Wedhorn, *The supersingular locus of the Shimura variety of* GU(1, n 1) *II*, Inventiones Mathematicae **184** (2011), 591–627.
- [VW13] Eva Viehmann and Torsten Wedhorn, Ekedahl-Oort and Newton strata for Shimura varieties of PEL type, Math. Ann. 356 (2013), no. 4, 1493–1550. MR 3072810
- [Wed05] Torsten Wedhorn, Specialization of F-zips, 2005.
- [Woo16] Amy Wai Ling Jane Wooding, *The Ekedahl-Oort stratification of unitary Shimura varieties*, Ph.D. thesis, McGill University, 2016.
- [Zin84] Thomas Zink, Cartiertheorie kommutativer former gruppen, Teubner-Texte zur Mathematik, 1984.

Department of Mathematics and Computer Science, Washington College, Chestertown, Maryland 21620, USA Mar

Email address: estacy2@washcoll.edu

MATHEMATICS DEPARTMENT, UNIVERSITY OF CALIFORNIA, SANTA CRUZ, CA 95064, USA

Email address: dbhamidi@ucsc.edu

DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, STILLWATER, OK 74078, USA

Email address: maria.fox@okstate.edu

DEPARTMENT OF MATHEMATICS, BROOKLYN COLLEGE, CITY UNIVERSITY OF NEW YORK, BROOKLYN, NY 11210 USA

Email address: heidi.goodson@brooklyn.cuny.edu

DEPARTMENT OF MATHEMATICS, LEHIGH UNIVERSITY, BETHLEHEM, PA 18015, USA

Email address: stg323@lehigh.edu

DEPARTMENT OF MATHEMATICS, COLORADO STATE UNIVERSITY, FORT COLLINS, CO 80523, USA

Email address: sandra.nair@colostate.edu