Limiting distributions of the Spherical model and Spin O(N) model: Appearance of GFF

Juhan Aru, Aleksandra Korzhenkova

Abstract

We revisit the relation between the spherical model of Berlin-Kac and the spin O(N) model in the limit $N \to \infty$ and explain how they are related via the discrete Gaussian free field (GFF).

More precisely, using probabilistic limit theorems and concentration of measure we first prove that the infinite volume limit of the spherical model on a d-dimensional torus is a massive GFF in the high-temperature regime, a standard GFF at the critical temperature, and a standard GFF plus a Rademacher random constant in the low-temperature regime. The proof in the case of the critical temperature appears to be new and requires a fine understanding of the zero-average Green's function on the torus.

For the spin O(N) model, we study the model in the double limit of the spin-dimensionality and the torus size. We take the limit as first the spin-dimension N goes to infinity, and then the size of the torus, and obtain that the different spin coordinates become i.i.d. fields, whose distribution in the high-temperature regime is a massive GFF, a standard GFF at the critical temperature, and a standard GFF plus a Gaussian random constant in the low-temperature regime.

In particular, this means that although the limiting free energies per site of the two models agree at all temperatures, their actual finite-dimensional laws still differ in terms of their zero modes in the low-temperature regime.

1 Introduction

1.1 Spherical model

Let \mathbb{T}_n^d be a d-dimensional discrete torus with side-length n. To each vertex $x \in \Lambda$, associate a continuous "spin" $\theta_x \in \mathbb{R}$ in such a way that the whole configuration $\boldsymbol{\theta} = (\theta_x)_{x \in \mathbb{T}_n^d}$ lives on the $(n^d - 1)$ -dimensional spin of radius $\sqrt{n^d}$, i.e., $\|\boldsymbol{\theta}\|_2^2 = \sum_x \theta_x^2 = n^d$. We call $\boldsymbol{\theta}$ a configuration of the *spherical model* on \mathbb{T}_n^d at the inverse temperature $\beta \geq 0$ if

$$\boldsymbol{\theta} \sim \nu_{\mathbb{T}_n^d, \beta}(\mathrm{d}\boldsymbol{\theta}) = \frac{1}{Z_{\mathbb{T}_n^d, \beta}} \exp\left(\frac{\beta}{2} \sum_{x \sim y} \theta_x \theta_y\right) \mathrm{Unif}_{\sqrt{n^d} \mathbb{S}^{n^d - 1}}(\mathrm{d}\boldsymbol{\theta}),$$

where $x \sim y$ stands for the neighbouring vertices. This model was first introduced by Berlin and Kac [BK52] in 1952 as a simplification of the Ising model that still exhibits a phase transition in $d \geq 3$, but whose free energy can be analytically studied in the infinite volume limit. In the original work Berlin and Kac used variations of the steepest-descent method to calculate the limiting free energy, to observe the phase transition in the case of three-dimensional lattice, and to argue that the distributional limit of the model below the critical temperature is non-Gaussian. In a short note Molchanov and Sudarev [SM75] stated that this method further would give various results about the limiting distribution of the spherical model, but the proofs never appeared. In this article, we rigorously prove a subset of these results using probabilistic tools instead (cf. Theorem 1.1), which we believe also help clarify the conceptual picture.

An alternative approach that also implicitly leads in [BD87] to the results of our theorem under unverified technical assumptions is the so-called mean-spherical approach, where instead of conditioning on the spin configuration, one adds a mass term $(s\sum_x \theta_x^2)$ to the Hamiltonian in such a way that on average $\|\boldsymbol{\theta}\|_2^2$ equals exactly n^d . This method was proposed by Lewis and Wannier [LW52] shortly after the appearance of the original paper [BK52]; they observed that computations of various quantities of the model, such as correlations or the free energy, are significantly simpler in the mean-spherical model. The method works perfectly well in the high-temperature regime, however, as it was quickly realized [LW53, Lax55], it can not be directly applied in the setting of zero (magnetic) field below the critical temperature, as some discrepancies with the original spherical model arise. Still, in the presence of an external magnetic field $h \neq 0$, it was shown [YW04] that the thermodynamic limits of the spherical model and the mean-spherical model agree and further, provided that first the thermodynamic limit is taken and then $h \downarrow 0$, the case of zero magnetic field of the spherical model can be recovered. Since then, this approach has been used in several papers [KT77, Shc88, BD87], but none of them seem to contain a fully rigorous proof for the critical and low-temperature regimes in the case of zero magnetic field.

Finally, the low-temperature regime of the spherical model can be seen as a toy model for observing the condensation phenomena. In this wider context, in [Luk20] the author studies, as an example, a complexified version of the spherical model below the critical temperature. The author derives convergence of local correlation functions, which combined with tightness, in turn, implies (local) convergence in law in the aforementioned regime. While this proof also works for the usual spherical model, it is not sufficient to show convergence at the critical temperature.

Let us now state our main result about the limiting distribution of the spherical model and explain the reasoning leading to the proof.

Theorem 1.1 (Infinite volume limit of the spherical model). Let $d \geq 2$. The spherical model on \mathbb{T}_n^d at inverse temperature $\beta > 0$, $[\boldsymbol{\theta}]_n = (\theta_x)_{x \in \Lambda}$, converges in law uniformly over compact subsets of \mathbb{Z}^d as $n \to \infty$ to:

- 1. $\beta < \beta_c$: a massive Gaussian free field (GFF) on \mathbb{Z}^d scaled by $1/\sqrt{\beta}$ with the mass $m^2 > 0$ depending on β and d in a specific way;
- 2. $\beta = \beta_c$: a GFF on \mathbb{Z}^d scaled by $1/\sqrt{\beta_c}$;
- 3. $\beta > \beta_c$: a GFF on \mathbb{Z}^d scaled by $1/\sqrt{\beta}$ plus an independent constant random drift $\sqrt{\frac{\beta \beta_c}{\beta}} X$ with X being a Rademacher random variable variable.

Furthermore, all local correlations of spins converge.

We start by observing that for any n, since $\sum_{x} \theta_{x}^{2} = n^{d}$

$$\nu_{\mathbb{T}_n^d,\beta}(\mathrm{d}\boldsymbol{\theta}) \propto \exp\bigg(-\frac{\beta}{2}\langle\boldsymbol{\theta},(-\Delta+m^2)\boldsymbol{\theta}\rangle\bigg)\mathrm{Unif}_{\sqrt{n^d}\mathbb{S}^{n^d-1}}(\mathrm{d}\boldsymbol{\theta})$$

for any $m^2 > 0$. Hence, the law of $\sqrt{\beta}[\theta]_n$ is related to a massive GFF φ (for an arbitrary mass $m^2 > 0$) through conditioning on its norm to be equal to $\sqrt{\beta n^d}$, i.e., $\|\varphi\|_2^2 = \beta n^d$ (see Section 2.3). Now the idea is somewhat similar to the one of Lewis and Wannier [LW52], but from a probabilistic point of view: namely, we choose mass m_n^2 in such a way that $\mathbb{E}[\|\varphi\|_2^2] = \beta n^d$ (or equivalently, $G_{\mathbb{T}_n^d, m_n^2}(0, 0) = \beta$ for the massive Green's function $G_{\mathbb{T}_n^d, m_n^2}$ on \mathbb{T}_n^d), which is natural due to concentration of Gaussian measure.

For large n and fixed m>0, $G_{\mathbb{T}_n^d,m^2}$ is roughly comparable to $G_{\mathbb{Z}^d,m^2}$, and thus, the question of existence of a critical point boils down to the understanding of whether there is an $m^2\geq 0$ such that $G_{\mathbb{Z}^d,m^2}=\beta$ or not. Observe that this explains the difference between the 2D and higher dimensional cases. Indeed, since in d=2, $G_{\mathbb{Z}^d,0}(0,0)$ is infinite, by choosing the mass small enough, one could get an arbitrarily large value of the variance $G_{\mathbb{Z}^d,m^2}(0,0)$; whereas in $d\geq 3$, there is a maximal possible value $G_{\mathbb{Z}^d,0}(0,0)<\infty$.

More precisely, we observe that in the high-temperature regime $\beta < \beta_c := G_{\mathbb{Z}^d,0}(0,0)$, the aforementioned sequence of masses $(m_n^2)_n$ converges to a positive number $m^2 > 0$ in such a way that

$$G_{\mathbb{T}_n^d, m_n^2}(0, 0) \to G_{\mathbb{Z}^d, m^2}(0, 0) = \beta.$$

In this case, the concentration of measure for $\|\varphi\|_2^2$ is sufficient to make the conditioning disappear in the limit, and we obtain a purely Gaussian limit. Somewhat refined versions of classical limiting theorems help us handle this phase rather directly.

The low-temperature phase $\beta > \beta_c$ is already trickier. One can check that

$$G_{\mathbb{T}_n^d,m_n^2}(x,y) \sim G_{\mathbb{T}_n^d}^{0\text{-avg.}}(x,y) + \beta - \beta_c \overset{n \to \infty}{\sim} G_{\mathbb{Z}^d,0}(x,y) + \beta - \beta_c,$$

where $G_{\mathbb{T}_n^d}^{0\text{-avg.}}$ is the correlation function of the zero-average GFF on torus. This heuristically lets us restate the problem in terms of the zero-average GFF γ plus an independent zero mode, a constant (in space) Gaussian drift $Z\mathbf{1}_{T_n^d}$, conditioned on the norm of the sum to be $\sqrt{\beta n^d}$. As the drift is constant, almost surely $\frac{1}{n^d}\|\gamma + Z\mathbf{1}\|_2^2 = \frac{1}{n^d}\|\gamma\|_2^2 + |Z|^2$. Due to high concentration of $\frac{1}{n^d}\|\gamma\|_2^2$ around β_c , the conditioning on the norm of the sum forces |Z| to become constant in the limit; and by symmetry we obtain a Rademacher random variable for the zero mode. Making this precise is slightly more subtle, but after finding the right angle, the proof follows from combining basic concentration of measure results with relatively direct density bounds.

Finally, in the critical case, we observe that $G_{\mathbb{T}_n^d}^{0\text{-avg.}}(x,y) \sim G_{\mathbb{Z}^d,0}(x,y)$, however, it is necessary to refine this relation further. In this direction we prove Proposition 2.14. First, it improves the error estimates on the zero-average Green's function on the torus in $d \geq 3$ – we show that $G_{\mathbb{T}_n^d}^{0\text{-avg.}}(x,y) = G_{\mathbb{Z}^d,0}(x,y) + O(n^{2-d})$. Second, in the case of d=3, as we further need to determine the sign of the error on the diagonal, we prove that $G_{\mathbb{T}_n^d}^{0\text{-avg.}}(x,x) < G_{\mathbb{Z}^d,0}(x,x)$ uniformly in large n. Obtaining the sign was surprisingly tricky and is related to what is called Madelung constant for electrostatic potential in certain salts – a quantity of interest for chemical physics introduced in the beginning of the 20th century [Mad19]. See the subsection 'Further questions and wider context' for a discussion on this connection.

To deduce convergence of local spin correlations from convergence in law (Corollary 3.2), we prove the existence of moments of single spherical spins (see Lemma 3.4), inspired by the analogous result about complexified spherical spins in [Luk20].

1.2 Spin O(N) model

The spin O(N) model (on \mathbb{Z}^d), introduced by Stanley in 1968 [Sta68a], is a fundamental model in statistical mechanics: for N=1, one recovers the Ising model; for N=2, the XY model; and for N=3, the classical Heisenberg model. Here we consider the infinite spin-dimensionality limit, $N\to\infty$, and its relation to the spherical model.

A connection between the spin O(N) model in the $N \to \infty$ limit and the spherical model was first suggested by Stanley [Sta68b] and later proved for any fixed non-critical temperature

by Kac and Thompson [KT71], see also [Shc88] and [GP93] that together also cover the case of the critical temperature. More precisely, combining the steepest-descent method with an approach similar to the aforementioned mean-spherical one, they showed the equality of the free energies per site of the two models: if $\mathbf{S} \in \prod_{x \in \mathbb{T}_n^d} (\sqrt{N} \mathbb{S}^{N-1})$ is a configuration of the $spin\ O(N)\ model$ on \mathbb{T}_n^d at inverse temperature $\beta \geq 0$, i.e.,

$$\mathbf{S} \sim \frac{1}{Z_{\mathbb{T}_n^d, N, \beta}} \exp \left(\frac{\beta}{2} \sum_{i \sim j} \mathbf{S}_i \cdot \mathbf{S}_j \right) \prod_{x \in \mathbb{T}_n^d} \mathrm{Unif}_{\sqrt{N} \mathbb{S}^{N-1}} (\mathrm{d} \mathbf{S}_x),$$

then (for $\beta \neq \beta_c$ in [KT71]),

$$\lim_{n \to \infty} \frac{1}{n^d N} \log Z_{\mathbb{T}_n^d, N, \beta} = \lim_{n \to \infty} \frac{1}{n^d} \log Z_{\mathbb{T}_n^d, \beta};$$

for any ordering of the limits $N, n \to \infty$. In this article we try to look at this connection more closely, and observe that it actually passes by the Gaussian free field. In the following theorem, we describe the local limit of the spin O(N) model when we first take the infinite spin-dimensionality limit $N \to \infty$ and then the thermodynamic limit $\mathbb{T}_n^d \to \mathbb{Z}^d$.

Theorem 1.2 (Infinite spin-dimensionality thermodynamic limit of the spin O(N) model). Any projection on the finitely many coordinates of the spin O(N) model on \mathbb{T}_n^d at inverse temperature $\beta > 0$, $\overline{\mathbf{S}}^M \coloneqq (S_x^i)_{x \in \Lambda}^{i=1,\dots,M}$ converges in law as $N \to \infty$ to that of an M-vectorial massive GFF scaled by $1/\sqrt{\beta}$ with the mass m_n^2 depending on β, d and n in a specific way. The consequent local (uniform on compacts of \mathbb{Z}^d) infinite volume distributional limit $(n \to \infty)$ has the law of:

- 1. $\beta < \beta_c$: an M-vectorial massive GFF on \mathbb{Z}^d scaled by $1/\sqrt{\beta}$ with the mass m^2 depending on β and d in a specific way;
- 2. $\beta = \beta_c$: an M-vectorial GFF on \mathbb{Z}^d scaled by $1/\sqrt{\beta}$;
- 3. $\beta > \beta_c$: an M-vectorial GFF on \mathbb{Z}^d scaled by $1/\sqrt{\beta}$ plus an independent constant random drift $\sqrt{\frac{\beta-\beta_c}{\beta}}\overline{Z}^M$ with \overline{Z}^M being an M-dimensional standard normal vector.

As a corollary we see, maybe a little surprisingly, that at least in the order of limits we take, although the free energies per site of the two models agree at all temperatures, their limiting laws are, in fact, not equal in the low-temperature regime. We expect this to be true in whatever order we take the limits. The a posteriori explanation of this appearing discrepancy in the low-temperature regime is not particularly hard: the information about a single degree of freedom, for instance, the observed difference in law of the drift part of the limiting fields, just disappears when one calculates the limiting free energy per site.

Corollary 1.3. In the high-temperature regime and at the critical temperature $\beta_c = G_{\mathbb{Z}^d,0}(0,0)$, the local distributional limit of the spherical model and the corresponding limiting law (obtained by first taking $N \to \infty$ and then the volume of the torus to infinity) of a single spin-coordinate process of the spin O(N) model agree; while in the low-temperature regime they differ in the law of the global zero mode.

The proof of the theorem consists of two steps: first we show that as the spin-dimensionality N tends to infinity, the law converges locally (in spin-dimensionality) to a vector-valued massive GFF with mass m_n^2 chosen in such a way that $G_{\mathbb{T}_n^d,m_n^2}(0,0)=\beta$; and then take the infinite volume limit of the Gaussian measure of m_n^2 -massive GFF on torus. The first step

follows our proof of the high-temperature regime of the spherical model, the proof of the latter step is quite simple and requires only a few well-known results about Green's functions on discrete lattices (torus and \mathbb{Z}^d).

We believe that the theorem should also hold in the opposite order of limits. In fact in the high-temperature regime this would follow directly from the work of A. Kupiainen [Kup80]; and we are hopeful that by combining some of his ideas with our approach, we would be able to extend this interchange of limits to all temperatures.

1.3 Further questions and wider context

To complete the probabilistic picture of the spherical model, which at the very least offers a statistical physics model that can be fully studied in all dimensions and phases, one would need to investigate it across various domains with different boundary conditions and in the presence of a magnetic field; ideally, extending the study to more general graphs as well. It is not particularly hard to predict the outcomes of adding boundary conditions or a constant magnetic field – and, indeed, our proofs can adapt to these changes; however, more general (also weighted) graphs might present a challenge.

As mentioned before, an interesting aspect in dealing with the critical case of the spherical model was the study of zero-average Green's function on the torus and determining whether it is smaller than the lattice Green's function. It is a mathematically delicate question that is related to the so-called Madelung constant in chemical physics and mathematical chemistry [Mad19, BGM⁺13]. Let us explain this connection here: one way to try to obtain the sign of the difference between the two mentioned Green's functions would be by taking an appropriate scaling limit and understanding the zero-average continuum Green's function $G_{\mathbb{T}^d}(0,x)$ for $x \in \partial[-1/2,1/2]^d$ viewed as points on the torus \mathbb{T}^d of side-length one. These quantities give us forms of the Madelung constant – for example, d(2), c(2) and b(2) in [BGM⁺13, (1.3.29) and Table 1.4] correspond precisely to $(2\pi)^2 G_{\mathbb{T}^3}(0,x)$ with x=(1/2,1/2,1/2), x=(0,1/2,1/2)and x = (0, 0, 1/2), respectively. Determining/approximating the Madelung constants is a well-known problem, and although there are explicit expressions using, for instance, Fourier series, the series expansions are not absolutely convergent and might diverge when one takes some very natural truncation! In our case, we find a way to prove the needed inequality on the lattice directly, but several interesting questions of potential analysis pose themselves in this direction. For example, it would be interesting to better understand the negativity region of the zero-average Green's function in all dimensions $d \geq 3$.

Yet another interesting aspect about the spherical model is that it can be viewed as a simple model for observing the condensation phenomena [Luk20]. The links between the ideal Bose gas and the spherical model through their critical behaviour [GB68] and appearance of Bose-Einstein condensation [GP93] were noted in the late sixties and early nineties, respectively, and have recently regained some attention [CSZ19, Luk20]. It might be interesting to revisit this connection, as well as to look at some other models forming a condensate, from the perspective of probabilistic limit theorems and concentration of measure, as used in the current paper.

Let us now turn to questions and topics related to the spin O(N) model. To round out the picture of the limiting distribution of the spin O(N) model, the primary objective would be finding a way of interchanging the order of taking infinite spin-dimensionality and infinite volume limits at all temperatures. We are currently working on this and expect it to be within reach.

In a broader context, an intriguing question is whether our approach could provide new

insight into Polyakov's conjecture [Pol75] on the exponential decay of correlation functions in 2D spin O(N) models with $N \geq 3$. It is hard to see how our approaches and techniques could help to solve the conjecture, however, one could possibly hope to extract useful quantitative bounds on critical temperatures in the 2D spin O(N) model as $N \to \infty$ (cf. [Kup80]).

Lastly, our results on the spin O(N) model (especially, Proposition 4.3) could be considered in the context of the propagation of chaos [Szn91]. The most famous example of this phenomena is the Poincaré lemma, which states that the marginals of the spherical measure in high dimensions become independent Gaussians. In our setting, we observe that throughout all the temperature regimes, the spin O(N) measures are chaotic: a sequence of symmetric probability measures on the product spaces Ω^N is called chaotic if, in the limit $N \to \infty$, any projection to finitely many coordinates becomes the product measure. One could certainly obtain similar results for spin models whose Hamiltonians additionally contain non-quadratic interaction terms, e.g., quartic terms; however, it is not yet clear, whether any interesting new phenomena could be found in these generalisations.

1.4 Outline

Section 2 defines precise setting of the spherical and spin O(N) models, provides a necessary background on various versions of GFF and explains their connection to the two models. It also contains some new results on the zero-average Green's function. In Section 3 we deal more closely with the spherical model, in particular, we prove Theorem 1.1 and convergence of correlations of any finite number of its spins. The object of interest in Section 4 is the spin O(N) model, especially Theorem 1.2, which is proven there. To keep this article concise we moved some of the proofs less relevant to the topic to Appendix A.

1.5 Acknowledgements

We would like to thank A. Prévost for interesting discussions. Both authors are supported by Eccellenza grant 194648 of the Swiss National Science Foundation and are members of NCCR Swissmap.

2 Setup and preliminaries

In this section we introduce notation that will be used in the rest of the article, as well as a few versions of Gaussian free fields that are closely related to our models of interest, namely spherical and spin O(N). We further recall some of the properties of these GFFs and their corresponding Green's functions (correlation structure) that later will be relevant for our proofs.

The only new contribution of this section is Proposition 2.14, which proves some finer bounds on the zero-average Green's function on the discrete torus and compares it to the Green's function on \mathbb{Z}^d .

2.1 Notation, definitions and setup of our two models

Let $d \geq 2$. We consider a discrete d-dimensional domain Λ to either be a discrete torus $\mathbb{T}_n^d = (\mathbb{Z}/n\mathbb{Z})^d$ for some $n \in \mathbb{N}$ or \mathbb{Z}^d , both endowed with the usual nearest-neighbour graph structure. Let $x, y \in \Lambda$, we write d(x, y) for the graph distance and |x - y| for the Euclidean distance between the points. $x \sim y$ denotes two neighbouring vertices of Λ . Given $A \subset \Lambda$, $A^c = \Lambda \setminus A$ stands for the complement of A in Λ , ∂A for the outer boundary of A in Λ , i.e.,

 $\partial A = \{y \in A^c : \exists x \in A \text{ s.t. } y \sim x\}, \ \overline{A} := A \cup \partial A, \ |A| \text{ for the cardinality of } A, \text{ and } \mathbf{1}_A \text{ for the vector } (1)_{x \in A}. \text{ If } \Lambda = \mathbb{T}_n^d \text{ and } \pi_n^d : \mathbb{Z}^d \to \Lambda \text{ is the canonical projection, for each } x \in \Lambda, \text{ we set } \hat{x} \text{ to be the unique element of } (\pi_n^d)^{-1}(\{x\}) \cap [-n/2, n/2)^d. \text{ Analogously, for } A \subset \mathbb{T}_n^d, \text{ we define } \hat{A} = \{\hat{x} \in [-n/2, n/2)^d \cap \mathbb{Z}^d : x \in A\} \subset \mathbb{Z}^d. \text{ We further call } A \text{ properly contained in } \mathbb{T}_n^d \text{ if } \partial \hat{A} \subset [-n/2, n/2)^d \cap \mathbb{Z}^d.$

For the convenience of the reader we recall the definitions of the spherical and spin O(N) models in the setting of our interest.

Definition 2.1 (Spherical model). Let $\beta > 0, n \in \mathbb{N}$ and $\Lambda = \mathbb{T}_n^d$. We call $\boldsymbol{\theta} := (\theta_x)_{x \in \Lambda} \in \sqrt{n^d} \mathbb{S}^{n^d-1}$ a configuration of the spherical model on Λ at inverse temperature β if $\boldsymbol{\theta} \sim \nu_{\Lambda,\beta}$ with

$$\nu_{\Lambda,\beta}(\mathrm{d}\boldsymbol{\theta}) = \frac{1}{Z_{\Lambda,\beta}} \exp\left(\frac{\beta}{2} \sum_{x \sim y} \theta_x \theta_y\right) \lambda_{\sqrt{n^d} \, \mathbb{S}^{n^d - 1}}(\mathrm{d}\boldsymbol{\theta}). \tag{2.1}$$

Here, $\lambda_{\sqrt{n^d} \mathbb{S}^{n^d-1}}$ denotes the uniform measure on the sphere $\sqrt{n^d} \mathbb{S}^{n^d-1} \subset \mathbb{R}^{n^d}$ of radius $\sqrt{n^d}$, $\sum_{x \sim y} runs$ over all neighbouring points $x, y \in \Lambda$ and $Z_{\Lambda,\beta}$ is the normalizing constant, i.e.,

$$Z_{\Lambda,\beta} = \int_{\sqrt{n^d} \, \mathbb{S}^{n^d - 1}} \exp\left(\frac{\beta}{2} \sum_{x \sim y} \theta_x \theta_y\right) \lambda_{\sqrt{n^d} \, \mathbb{S}^{n^d - 1}} (\mathrm{d}\boldsymbol{\theta}).$$

Definition 2.2 (Spin O(N) **model).** Let $\beta > 0, n, N \in \mathbb{N}$ and $\Lambda = \mathbb{T}_n^d$. We call $\mathbf{S} := (S_x)_{x \in \Lambda} \in (\sqrt{N} \mathbb{S}^{N-1})^{n^d}$ a configuration of the spin O(N) model on Λ at inverse temperature β if $\mathbf{S} \sim \mu_{\Lambda,N,\beta}$ with

$$\mu_{\Lambda,N,\beta}(\mathrm{d}\mathbf{S}) = \frac{1}{Z_{\Lambda,N,\beta}} \exp\left(\frac{\beta}{2} \sum_{x \sim y} S_x \cdot S_y\right) \prod_{x \in \Lambda} \lambda_{\sqrt{N} \, \mathbb{S}^{N-1}}(\mathrm{d}S_x). \tag{2.2}$$

Here, $\lambda_{\sqrt{N} \mathbb{S}^{N-1}}$ denotes the uniform measure on the sphere $\sqrt{N} \mathbb{S}^{N-1} \subset \mathbb{R}^N$ of radius \sqrt{N} , $\sum_{x \sim y}$ runs over all neighbouring points $x, y \in \Lambda$, $S_x \cdot S_y$ stands for the inner product in \mathbb{R}^N and $Z_{\Lambda,N,\beta}$ is the normalizing constant, i.e.,

$$Z_{\Lambda,N,\beta} = \int \cdots \int_{\Pi_{x \in \Lambda}} \exp\left(\frac{\beta}{2} \sum_{x \sim y} S_x \cdot S_y\right) \prod_{x \in \Lambda} \lambda_{\sqrt{N} \, \mathbb{S}^{N-1}} (\mathrm{d}S_x).$$

In the sequel we will always refer to a configuration of the spherical model as $\boldsymbol{\theta}$ and of the spin O(N) model as \mathbf{S} .

2.2 Massive GFF, its vectorial descendant and zero-average GFF

In the present subsection we review a few different GFFs, which as we will see later are closely related to our models of interest both in the finite setting and in the distributional limit.

Definition 2.3 (Massive GFF on torus). Let $n \in \mathbb{N}$ and $\Lambda = \mathbb{T}_n^d$. Let U be a subset of Λ , possibly $U = \emptyset$, $m^2 > 0$ if $U = \emptyset$ and $m^2 \geq 0$ otherwise. We call a real-valued function on Λ , denoted by $\varphi^U := (\varphi_x^U)_{x \in \Lambda} \in \mathbb{R}^{n^d}$, an m^2 -massive GFF on Λ with Dirichlet boundary

condition on U if it is distributed according to

$$\mathbb{P}_{U^c, m^2}(\mathrm{d}\boldsymbol{\varphi}) \propto \exp\left(-\frac{1}{4} \sum_{x \sim y} (\varphi_x - \varphi_y)^2 - \frac{m^2}{2} \sum_{x \in \Lambda} \varphi_x^2\right) \prod_{x \in U} \delta_0(\mathrm{d}\varphi_x) \prod_{x \in U^c} \mathrm{d}\varphi_x, \tag{2.3}$$

where $\sum_{x\sim y}$ runs over all neighbouring points $x,y\in\Lambda$. If $U=\emptyset$, we write φ instead of φ^{\emptyset} and call it an m^2 -massive GFF. If U is such that $\hat{U}=\partial_{\mathrm{inner}}\hat{\Lambda}:=\{x\in\hat{\Lambda}:\exists y\in\mathbb{Z}^d\setminus\hat{\Lambda}:y\sim x\}$, we call it an m^2 -massive Dirichlet GFF on $\hat{\Lambda}$.

Definition 2.4 (N-vectorial massive GFF on torus). Let $n, N \in \mathbb{N}$ and $\Lambda = \mathbb{T}_n^d$. Let U be a subset of Λ , possibly $U = \emptyset$, $m^2 > 0$ if $U = \emptyset$ and $m^2 \geq 0$ otherwise. We call an \mathbb{R}^N -valued function on Λ , denoted by $\Phi^U := (\Phi_x^U)_{x \in \Lambda} \in (\mathbb{R}^N)^{n^d}$, an N-vectorial m^2 -massive GFF on Λ with Dirichlet boundary condition on U if it is distributed according to

$$\mathbb{P}_{U^c,N,m^2}(\mathrm{d}\boldsymbol{\Phi}) \propto \exp\left(-\frac{1}{4}\sum_{x\sim y}\|\Phi_x - \Phi_y\|^2 - \frac{m^2}{2}\sum_{x\in\Lambda}\|\Phi_x\|^2\right) \prod_{x\in U} \delta_{\mathbf{0}}(\mathrm{d}^N\Phi_x) \prod_{x\in U^c} \mathrm{d}^N\Phi_x,$$

where $\sum_{x\sim y}$ runs over all neighbouring points $x,y\in\Lambda$, d^N denotes the Lebesgue measure on \mathbb{R}^N and $\|\cdot\|$ is the Euclidean norm on \mathbb{R}^N . If $U=\emptyset$, we write Φ instead of Φ^\emptyset and call it an N-vectorial m^2 -massive GFF. If U is such that $\hat{U}=\partial_{\mathrm{inner}}\hat{\Lambda}:=\{x\in\hat{\Lambda}:\exists y\in\mathbb{Z}^d\setminus\hat{\Lambda}:y\sim x\}$, we call it an N-vectorial m^2 -massive Dirichlet GFF on $\hat{\Lambda}$.

Remark 1. Note that we could have equivalently defined an N-vectorial m^2 -massive GFF with Dirichlet boundary condition on U as a vector with N i.i.d. coordinate processes, each distributed as an m^2 -massive GFF with Dirichlet boundary condition on U.

Taking into account the above remark, it suffices to understand the covariance structure of an m^2 -massive GFF with Dirichlet boundary condition on U. Note that we can rewrite the expression in the exponent of (2.3) as follows

$$-\frac{1}{4}\sum_{x\sim y}(\varphi_x-\varphi_y)^2 - \frac{m^2}{2}\sum_{x\in\Lambda}\varphi_x^2 = -\frac{1}{2}\langle \varphi', (-\Delta'_{U^c}+m^2)\varphi'\rangle, \tag{2.4}$$

where $\langle \cdot, \cdot \rangle$ stands for the inner product on $\mathbb{R}^{n^d-|U|}$, $\varphi' := (\varphi_x)_{x \in U^c}$ and Δ'_{U^c} is defined as

$$(\Delta'_{U^c}f)(x) \coloneqq \sum_{y \in U^c: y \sim x} f(y) - 2df(x) \text{ for any } x \in U^c, f: U^c \to \mathbb{R}.$$

When considered as an operator on the set \mathcal{F}_U of functions $f: \Lambda \to \mathbb{R}$ satisfying $f|_U \equiv 0$, it is denoted by Δ_{U^c} and called discrete Laplacian with Dirichlet boundary condition on U. Note that if we view $(-\Delta'_{U^c} + m^2)$ as an $\mathbb{R}^{(n^d - |U|) \times (n^d - |U|)}$ -matrix, (2.4) proves that it is positive definite, hence, invertible. This implies that the normalizing constant in (2.3) is $\sqrt{\det G'_{U^c,m^2}}$, where $G'_{U^c,m^2} \coloneqq (-\Delta'_{U^c} + m^2)^{-1} \in \mathbb{R}^{(n^d - |U|) \times (n^d - |U|)}$. Its extension $(G_{U^c,m^2}(x,y))_{x,y \in \Lambda}$ by zeros to an $\mathbb{R}^{n^d \times n^d}$ -matrix, i.e., $G_{U^c,m^2}(x,y) = G'_{U^c,m^2}(x,y) \mathbb{1}_{x,y \in U^c}$ for all $x,y \in \Lambda$, is the so-called massive Green's function on Λ with Dirichlet boundary condition on U. It is a well-known fact (see for instance [Rod17, Section 1]) that it is related to random walks in the

following way: for all $x, y \in \Lambda$,

$$G_{U^{c},m^{2}}(x,y) = \frac{1}{2d}G_{\alpha,U}(x,y) := \frac{1-\alpha}{2d} \sum_{k\geq 0} \mathbb{P}_{\alpha,U}^{x}[X_{k} = y]$$

$$= \frac{1-\alpha}{2d} \sum_{k\geq 0} (1-\alpha)^{k} \mathbb{P}_{0}^{x}[X_{k} = y, k < H_{U}],$$
(2.5)

where $\mathbb{P}^x_{\alpha,U}$ denotes the canonical law of a random walk **X** on the graph $\Lambda \cup \{\dagger\}$ (obtained from Λ viewed as a graph by adding an edge connecting each vertex of Λ with \dagger) starting at x with transition probabilities

$$p_{x,y} = \frac{1}{2d} (1 - \alpha) \mathbb{1}_{\{x \sim y, x \notin U\}}, \ p_{x,\dagger} = \alpha \mathbb{1}_{\{x \notin U\}}, \ p_{z,z} = 1, \text{ for all } x, y \in \Lambda, z \in U \cup \{\dagger\},$$

where $\alpha = \frac{m^2}{2d+m^2}$, $G_{\alpha,U}$ is the so-called Green's function of the described walk, $H_U := \inf\{n \ge 0 : X_n \in U\}$ the first hitting time of U, and \mathbb{P}_0^x is the law of a simple random walk on Λ started at x. $G_{\alpha,U}$ is also well-defined if $\Lambda = \mathbb{Z}^d$.

Remark 2. Observe that the assumption $m^2 > 0$ is mandatory if we work on \mathbb{T}_n^d with $U = \emptyset$ since simple random walk on torus in any dimension is recurrent. This also explains why if $U \neq \emptyset$, we might consider $m^2 = 0$. If we start directly with $\Lambda = \mathbb{Z}^d$ using (2.5) as the definition, then even if $U = \emptyset$, for $d \geq 3$, we as well might assume $m^2 = 0$ since a simple random walk on \mathbb{Z}^d for $d \geq 3$ is transient.

Therefore, we can also define a (massive) GFF on \mathbb{Z}^d .

Definition 2.5 ((N-vectorial) massive GFF on \mathbb{Z}^d). Let U be a subset of \mathbb{Z}^d , possibly $U=\emptyset$. Let $m^2>0$ if d=2 and $U=\emptyset$ or $m^2\geq 0$ otherwise. An m^2 -massive GFF on \mathbb{Z}^d with Dirichlet boundary condition on U is a centered Gaussian process indexed by \mathbb{Z}^d , denoted by $\psi^U:=(\psi_x)_{x\in\mathbb{Z}^d}$, with covariance structure $(G_{U^c,m^2}(x,y))_{x,y\in\mathbb{Z}^d}$. If $U=\emptyset$, we write ψ instead of ψ^\emptyset and call it an m^2 -massive GFF on \mathbb{Z}^d . If $m^2=0$, we call this process simply a GFF on \mathbb{Z}^d (with Dirichlet boundary condition on U). Let N>0 and let $(\psi_i^U)_{i=1}^N$ be N i.i.d. m^2 -massive GFFs (with Dirichlet boundary condition at U). The vector-valued Gaussian process $\Psi^U=(\Psi^U_x)_{x\in\mathbb{Z}^d}\coloneqq (\psi^U_i)_{i=1}^N$ is called an N-vectorial m^2 -massive GFF (with Dirichlet boundary condition on U).

It is now an easy observation (from (2.5)) that for any fixed finite $U \subset \mathbb{Z}^d$ (considered for each n as a subset of \mathbb{T}_n^d) and a fixed $m^2 > 0$, as n tends to infinity, an m^2 -massive GFF φ^U on \mathbb{T}_n^d with Dirichlet boundary condition on U converges in law uniformly on compact subsets of \mathbb{Z}^d towards an m^2 -massive GFF ψ^U on \mathbb{Z}^d with Dirichlet boundary condition on U. Moreover, we can extend the argument to a sequence of m_n^2 -massive GFFs $([\varphi^U]_n)_n$, each defined on \mathbb{T}_n^d , under the assumption that $m_n^2 \to m^2$ as $n \to \infty$. Indeed, this follows from the second line of (2.5) since it guarantees existence of C > 0 uniform in n such that $|G_{\mathbb{T}_n^d \setminus U, m_n^2} - G_{\mathbb{T}_n^d \setminus U, m^2}| \le C|\alpha - \alpha_n|G_{\mathbb{T}_n^d \setminus U, m^2}$. The latter in turn converges to zero uniformly over any compact subset of \mathbb{Z}^d as n tends to infinity.

Let us end the discussion of a massive GFF in this subsection by stating one more of its properties that will be frequently used in the sequel. Its proof may be found in [Rod17, Lemma 1].

Proposition 2.6 (Weak domain Markov property of massive GFF). Let Λ be either \mathbb{T}_n^d or \mathbb{Z}^d and U be a subset of Λ . In the former case or if d=2 and $U=\emptyset$, take $m^2>0$,

otherwise $m^2 \geq 0$. Let φ^U ($\varphi^\emptyset = \varphi$) be an m^2 -massive GFF on Λ with Dirichlet boundary condition on U. For a finite subset K of Λ , define $\tilde{\varphi}^{U,K} = (\tilde{\varphi}^{U,K}_x)_{x \in \Lambda}$ ($\tilde{\varphi}^{\emptyset,K} = \tilde{\varphi}^K$) by

$$\varphi_x^U = \tilde{\varphi}_x^{U,K} + h_x^{\varphi_X^U} \quad for \ all \ x \in \Lambda,$$

where $\mathbf{h}^{\boldsymbol{\varphi}_K^U} = (h_x^{\boldsymbol{\varphi}_K^U})_{x \in \Lambda}$ is the $\sigma(\varphi_x^U : x \in K)$ -measurable map given by

$$h_x^{\varphi_K^U} = \sum_{y \in K} \mathbb{P}_{\alpha, U}^x [H_K < \infty, X_{H_K} = y] \, \varphi_y^U \quad \text{for all } x \in \Lambda,$$

with $\alpha = \frac{m^2}{2d+m^2}$. Then under \mathbb{P}_{U^c,m^2} – the canonical law of φ^U – the field $\tilde{\varphi}^{U,K}$ is independent of $\sigma(\varphi_x^U: x \in K)$ and has the law of an m^2 -massive GFF on Λ with Dirichlet boundary condition on $U \cup K$.

Clearly, we have a fully analogous statement for a vector-valued massive GFF in view of Remark 1.

Remark 3. Note that the function h^{φ_K} is the so-called m^2 -massive harmonic extension of φ_K on K to Λ , i.e., h^{φ_K} solves

$$\begin{cases} (-\Delta_{\Lambda} + m^2)h(x) = 0 & \text{for } x \in K^c; \\ h(y) = \varphi_y & \text{for } y \in K. \end{cases}$$

This is an easy consequence of the Markov property of a random walk.

We conclude the present subsection by introducing one further version of a GFF on torus with $m^2=0$. As we noticed in Remark 2, it is impossible to define the corresponding measure living on the space of full $(n^d$ -)dimensionality, but realizable on the space of lower dimension. Before we assumed for example that $\varphi_x=0$ for some $x\in\mathbb{T}_n^d$. Another canonical way to do so is to require $\sum_x \varphi_x=0$. This leads to the following definition.

Definition 2.7 (Zero-average GFF on torus). Let $n \in \mathbb{N}, d \geq 3$ and $\Lambda = \mathbb{T}_n^d$. We call a real-valued function on Λ , denoted by $\gamma := (\gamma_x)_{x \in \Lambda} \in \mathbb{R}^{n^d}$, a zero-average GFF on Λ if it is distributed according to

$$\mathbb{P}_{\Lambda,0\text{-}avg.}(\mathrm{d}\boldsymbol{\gamma}) \propto \exp\left(-\frac{1}{4}\sum_{x\sim y}(\gamma_x - \gamma_y)^2\right)\delta_0\left(\frac{1}{\sqrt{n^d}}\sum_x \gamma_x\right)\prod_{x\in\Lambda}\mathrm{d}\gamma_x,\tag{2.6}$$

where $\sum_{x\sim y}$ runs over all neighbouring points $x,y\in\Lambda$.

Let $0 = \eta_1 < \eta_2 \le \ldots \le \eta_{n^d}$ be the eigenvalues of $-\Delta_{\Lambda}$ and u^1, \ldots, u^{n^d} be the corresponding orthogonal eigenvectors. Note that $u^1 = \frac{1}{\sqrt{n^d}}(1, \ldots, 1)^T$ and set $Q := (u^1, \ldots, u^{n^d})$. Then,

$$e^{-\frac{1}{4}\sum_{x\sim y}(\gamma_x-\gamma_y)^2}\delta_0\left(\frac{1}{\sqrt{n^d}}\sum_{x}\gamma_x\right)\mathrm{d}\gamma = e^{-\frac{1}{2}\langle Q^T\gamma,\mathrm{diag}\left(0,(\eta_k)_{k=2}^{n^d}\right)Q^T\gamma\rangle}\delta_0\left((Q^T\gamma)_1\right)\mathrm{d}\gamma.$$

This implies that the covariance matrix of $Q^T \gamma$ is $\operatorname{diag}(0, (1/\eta_k)_{k=2}^{n^d})$, and hence, the one of γ is $Q\operatorname{diag}(0, (1/\eta_k)_{k=2}^{n^d})Q^T =: \left(G_{\Lambda}^{0-\operatorname{avg.}}(x,y)\right)_{x,y\in\Lambda}$. The latter matrix is the so-called zero-average Green's function. We recollect and prove a few of its properties in Subsection 2.4.

2.3 Massive GFF, its vectorial descendant and zero-average GFF: relation to the spherical and spin O(N) models

Let $d \geq 2$, n and $N \in \mathbb{N}$. Within this subsection we assume that $\Lambda = \mathbb{T}_n^d$ is the d-dimensional discrete torus.

Proposition 2.8 (Relation between massive GFF and spherical model). Let $m^2 > 0$ and φ be an m^2 -massive GFF on Λ , $\beta > 0$. Then the law of $\varphi/\sqrt{\beta}$ conditionally on $\|\varphi/\sqrt{\beta}\|_2 = \sqrt{n^d}$ is that of the spherical model on Λ at inverse temperature β . Here $\|\cdot\|_2$ stands for the norm associated with the inner product $\langle \cdot, \cdot \rangle$ on \mathbb{R}^{n^d} .

Proof. Using polar transformation of coordinates we can easily compute the densities of $(X/\|X\|_2, \|X\|_2)$ and $\|X\|_2$ for $X = \varphi/\sqrt{\beta}$:

$$f_{(\frac{X}{\|X\|_{2}},\|X\|_{2})}(\boldsymbol{\sigma},t) = t^{n^{d}-1} f_{X}(t\boldsymbol{\sigma}), \quad \text{for } \boldsymbol{\sigma} \in \mathbb{S}^{n^{d}-1}, t > 0;$$

$$f_{\|X\|_{2}}(t) = \frac{\mathrm{d}}{\mathrm{d}t} \int_{0}^{t} \int_{\partial B^{n^{d}}(0,s)} f_{X}(\boldsymbol{\sigma}_{s}) \mathrm{d}\boldsymbol{\sigma}_{s} \mathrm{d}s$$

$$= \int_{\partial B^{n^{d}}(0,t)} f_{X}(\boldsymbol{\sigma}_{t}) \mathrm{d}\boldsymbol{\sigma}_{t}, \quad \text{for } t > 0,$$

where $d\sigma_t$ is the Hausdorff measure on $t \mathbb{S}^{n^d-1} = \partial B^{n^d}(0,t)$. Thus, for any $\gamma \in \sqrt{n^d} \mathbb{S}^{n^d-1}$,

$$f_{\frac{\varphi}{\sqrt{\beta}}} \Big|_{\|\frac{\varphi}{\sqrt{\beta}}\|_{2} = \sqrt{n^{d}}}(\gamma) d\gamma = \frac{f_{\frac{\varphi}{\sqrt{\beta}}}(\gamma)}{\int_{\partial B^{n^{d}}(0,\sqrt{n^{d}})} f_{\frac{\varphi}{\sqrt{\beta}}}(\sigma) d\sigma} d\gamma$$

$$= \frac{\exp\left(-\frac{\beta}{2} \langle \gamma, (-\Delta_{\Lambda} + m^{2})\gamma \rangle\right) \lambda_{\sqrt{n^{d}} \mathbb{S}^{n^{d}-1}}(d\gamma)}{\int \exp\left(-\frac{\beta}{2} \langle \sigma, (-\Delta_{\Lambda} + m^{2})\sigma \rangle\right) \lambda_{\sqrt{n^{d}} \mathbb{S}^{n^{d}-1}}(d\sigma)}$$

$$= \frac{\exp\left(\frac{\beta}{2} \sum_{x \sim y} \gamma_{x} \gamma_{y}\right) \lambda_{\sqrt{n^{d}} \mathbb{S}^{n^{d}-1}}(d\gamma)}{\int \exp\left(\frac{\beta}{2} \sum_{x \sim y} \sigma_{x} \sigma_{y}\right) \lambda_{\sqrt{n^{d}} \mathbb{S}^{n^{d}-1}}(d\sigma)} = \nu_{\Lambda,\beta}(d\gamma),$$

where we used that $\langle \boldsymbol{\gamma}, \boldsymbol{\gamma} \rangle = \langle \boldsymbol{\sigma}, \boldsymbol{\sigma} \rangle = n^d$.

We also have a similar statement for the spin O(N) model, which was implicitly used already in [Kup80] and proven in a slightly different setting in [AGS22, Proposition 2.3].

Proposition 2.9 (Relation between N-vectorial massive GFF and spin O(N) model). Let $m^2 > 0$ and Φ be an N-vectorial m^2 -massive GFF on Λ , $\beta > 0$. Then the law of $\Phi/\sqrt{\beta}$ conditionally on $\|\Phi_x/\sqrt{\beta}\| = \sqrt{N}$ for all $x \in \Lambda$ is that of the spin O(N) model on Λ at inverse temperature β . Here $\|\cdot\|$ stands for the Euclidean norm on \mathbb{R}^N .

We omit the proof of this result as it is fully analogous to the one for the spherical model and also can be easily adjusted from [AGS22, Proposition 2.3].

Proposition 2.10 (Relation between zero-average GFF and spherical model). Let $d \geq 3$ and γ be a zero-average GFF on $\Lambda = \mathbb{T}_n^d$, $\beta > 0$. Then the distribution of the spherical model at the inverse temperature β is equal to the conditional law of $(\gamma + c\mathbf{1}_{\Lambda})/\sqrt{\beta}$ given $\|\gamma + c\mathbf{1}_{\Lambda}\|_2/\sqrt{\beta} = \sqrt{n^d}$, where c is an "independent" Lebesgue constant, i.e., the law of (γ, c)

is
$$\mathcal{N}(0, G_{\Lambda}^{0\text{-}avg.}) \times \mathrm{d}c$$
.

Proof. Notice that equivalently we can work with orthonormal transformations of the corresponding laws. Let Q be the orthogonal matrix diagonalizing $G_{\Lambda}^{0\text{-avg.}}$. It is an easy observation that the "law" of $\gamma_{c,Q} := Q^T(\gamma + c\mathbf{1}_{\Lambda})$ is given by $\frac{1}{\sqrt{n^d}} \mathrm{d}c \times \times_{k=2}^{n^d} \mathbb{N}(0, \eta_k^{-1})$, where $(\eta_k)_{k\geq 2}$ are the non-zero eigenvalues of $-\Delta_{\Lambda}$ ordered non-decreasingly. Restricted to a finite ball and normalized accordingly, it is a probability measure that we denote by ρ . For our purposes we can consider, e.g., $B^{n^d}(0, 2\sqrt{\beta n^d})$. Then, fully analogously to the proof of Proposition 2.8, for any Borel-measurable $A \subset \mathbb{R}^{n^d}$,

$$\rho[A \mid \|\boldsymbol{\gamma}_{c,Q}\|_{2} = \sqrt{\beta n^{d}}] = \frac{\int \mathbb{1}_{A}(\boldsymbol{x}) \prod_{k=2}^{n^{d}} \exp\left(-\frac{\beta}{2} \eta_{k} x_{k}^{2}\right) \lambda_{\sqrt{n^{d}} \, \mathbb{S}^{n^{d}-1}}(\mathrm{d}\boldsymbol{x})}{\int \prod_{k=2}^{n^{d}} \exp\left(-\frac{\beta}{2} \eta_{k} y_{k}^{2}\right) \lambda_{\sqrt{n^{d}} \, \mathbb{S}^{n^{d}-1}}(\mathrm{d}\boldsymbol{y})}$$

$$= \frac{\int \mathbb{1}_{QA}(\boldsymbol{\sigma}) \exp\left(\frac{\beta}{2} \sum_{x \sim y} \sigma_{x} \sigma_{y}\right) \lambda_{\sqrt{n^{d}} \, \mathbb{S}^{n^{d}-1}}(\mathrm{d}\boldsymbol{\sigma})}{\int \exp\left(\frac{\beta}{2} \sum_{x \sim y} \tilde{\sigma}_{x} \tilde{\sigma}_{y}\right) \lambda_{\sqrt{n^{d}} \, \mathbb{S}^{n^{d}-1}}(\mathrm{d}\tilde{\boldsymbol{\sigma}})} = \nu_{\Lambda,\beta}[QA]$$

as desired. \Box

2.4 Preliminary estimates on Green's functions

This subsection mainly recalls but also proves some further results about Green's functions and their eigenvalues. We will start off with the massive Green's function and then turn to the zero-average Green's function.

2.4.1 Massive Green's function

We start with some canonical results whose proofs can be found, e.g., in [Rod17, Section 1] and [FV17, Proposition 8.30].

Lemma 2.11 (Properties of massive Green's function). Let $n \in \mathbb{N}$, $\Lambda = \mathbb{T}_n^d$ or \mathbb{Z}^d and U be a subset of Λ , possibly $U = \emptyset$. Let $m^2 \geq 0$ if $U \neq \emptyset$ and $m^2 > 0$ otherwise.

1. For any $K \subset U^c$ finite,

$$G_{U^c,m^2}(x,y) = G_{(U \cup K)^c,m^2} + \mathbb{E}_{\alpha,U}^x [\mathbb{1}_{\{H_K < \infty\}} G_{U^c,m^2}(X_{H_K},y)]. \tag{2.7}$$

2. If $m^2 > 0$, there exist two constants c, C > 0 depending only on m^2 and d such that

$$G_{U^c,m^2}(x,y) \le Ce^{-c|x-y|}, \text{ for all } U \subset \Lambda, x,y \in \Lambda.$$
 (2.8)

We will also often find it helpful to calculate with the basis of the eigenfunctions of the Green's function.

Lemma 2.12 (Eigenvalues of massive Green's function). Let $n \in \mathbb{N}$, $m^2 > 0$, $\Lambda = \mathbb{T}_n^d$, U be a finite subset of \mathbb{Z}^d viewed as a subset of Λ for each n. Set $u_n := |U^c|$.

1. $-\Delta_{\Lambda}$ has one $\eta_0 = 0$ and $(n^d - 1)$ positive eigenvalues given by

$$\eta_w = 2\sum_{i=1}^d \left(1 - \cos\left(2\pi \frac{w_i}{n}\right)\right) \text{ for } w = (w_i)_{i=1}^d \in \hat{\Lambda} \setminus \{0\} + \left(\left\lfloor \frac{n}{2}\right\rfloor, \dots, \left\lfloor \frac{n}{2}\right\rfloor\right).$$

The corresponding eigenvectors $(q^w)_{w \in [0,n)^d \cap \mathbb{Z}^d}$ building an orthonormal system are given by

$$q_x^w = \frac{1}{\sqrt{n^d}} \prod_{i=1}^d \left(\cos \left(2\pi \frac{x_i w_i}{n} \right) + \sin \left(2\pi \frac{x_i w_i}{n} \right) \right) \text{ for all } x \in \hat{\Lambda} + \left(\left\lfloor \frac{n}{2} \right\rfloor, \dots, \left\lfloor \frac{n}{2} \right\rfloor \right).$$

- 2. Let $(\eta_k)_{k\geq 2}$ be the positive eigenvalues of $-\Delta_{\Lambda}$ ordered non-decreasingly. Then, $\eta_k = \Theta(k^{2/d}/n^2)$.
- 3. The eigenvalues of G_{Λ,m^2} are given by $(1/(m^2 + \eta_x))_{x \in [0,n)^d \cap \mathbb{Z}^d}$.
- 4. Let $\mu_1 \geq \mu_2 \geq \ldots \geq \mu_{u_n}$ be the eigenvalues of $G_{U^c,m^2}|_{U^c \times U^c}$. There exists $L = L(U,m^2,d) > 0$ such that

$$\mu_i \in [1/(m^2 + 4d), 1/m^2] \quad \text{for all } |U| + 1 \le i \le u_n - |U|;
\mu_i \in [1/(m^2 + 4d), 1/m^2 + L] \quad \text{for all } i.$$
(2.9)

Proof. The eigenvalues of $-\Delta_{\Lambda}$ can be found for instance in [AF02, Example 5.17] or easily derived from [BK52, Appendix A]; the eigenvectors are also classical.

Using explicit formulae for $(\eta_x)_{x\in[0,n/2]^d\cap\mathbb{Z}^d}$ (any remaining x gives an eigenvalue coinciding with one indexed by some point in $[0,n/2]^d\cap\mathbb{Z}^d$), we obtain $\eta_x=\Theta(|x|^2/n^2)$. This together with symmetries of the torus and properties of cos-function implies that $\eta_k=\Theta(k^{2/d}/n^2)$. The third point follows immediately since G_{Λ,m^2} is the inverse of $-\Delta_{\Lambda}+m^2$.

As for the last property, recall that by (2.7), since U is finite,

$$G_{\Lambda,m^2}(x,y) = G_{U^c,m^2}(x,y) + \underbrace{\mathbb{E}_{\alpha}^x [\mathbb{1}_{\{H_U < \infty\}} G_{\Lambda,m^2}(X_{H_U}, y)]}_{=:M(x,y)}$$
(2.10)

for $x,y\in U^c$. Note that both $G_{\Lambda,m^2}|_{U^c\times U^c}$ and $G_{U^c,m^2}|_{U^c\times U^c}$ as principal submatrices of G_{Λ,m^2} and G'_{U^c,m^2} , respectively, are positive definite, and thus, symmetric. The latter is then also true for $M\in\mathbb{R}^{u_n\times u_n}$. Let $(\tilde{\lambda}_i)_{i\leq u_n}$ and $(\alpha_i)_{i\leq u_n}$ ordered non-increasingly denote the eigenvalues of $G_{\Lambda,m^2}|_{U^c\times U^c}$ and M, respectively. It is clear that the rank of M is at most |U|, and hence, there are at most as many non-zero eigenvalues. Furthermore, the minimal and maximal eigenvalues, $\alpha_{u_n}\leq 0$, $\alpha_1\geq 0$, satisfy

$$\max(\alpha_{1}, -\alpha_{u_{n}}) \leq \max_{f: \|f\|_{2}=1} |f^{T}Mf| \leq \max_{f: \|f\|_{2}=1} \sum_{z \in U} \sum_{x, y \in V} |f_{x}f_{y}| \mathbb{P}_{\alpha}^{x} [H_{U} < \infty] G_{\Lambda, m^{2}}(z, y)$$
$$\leq (2d + m^{2}) \left(\sum_{z \in U} \sum_{y \in V} G_{\Lambda, m^{2}}(z, y) \right)^{2} \leq L$$

for a constant $L(U, m^2, d) > 0$ uniform in n. In the second line we first used the weak Markov property of the random walk to conclude that

$$\mathbb{P}_{\alpha}^{x}[H_{U} < \infty] = \sum_{w \in U} \mathbb{P}_{\alpha}^{w}[\tilde{H}_{U} = \infty] \underbrace{\frac{2d}{1 - \alpha}}_{=2d + m^{2}} G_{\Lambda, m^{2}}(x, w) \leq (2d + m^{2}) \sum_{w \in U} G_{\Lambda, m^{2}}(x, w);$$

and then (2.8) to get the last inequality. From Weyl's inequality [HJ12, Theorem 4.3.1] and the fact that $\alpha_k = 0$ for all $u_n - |U| \ge k > |U|$, we conclude that $\mu_i \in [\tilde{\lambda}_{i+|U|}, \tilde{\lambda}_{i-|U|}]$ for all $|U| + 1 \le i \le u_n - |U|$ and $\mu_i \in [\tilde{\lambda}_i - \alpha_1, \tilde{\lambda}_i - \alpha_{u_n}]$ for all $i \le u_n$. Furthermore, by [HJ12, Theorem 4.3.28], $\lambda_{n^d-u_n+i} \le \tilde{\lambda}_i \le \lambda_i$ for all $i \le u_n$, where $(\lambda_j = (\eta_j + m^2)^{-1})_{j=1}^{n^d}$ are the eigenvalues of G_{Λ,m^2} ordered non-increasingly. Altogether, since $\eta_k \in [0,4d]$ for all k,

$$\mu_i \in [1/(m^2 + 4d), 1/m^2]$$
 for all $|U| + 1 \le i \le u_n - |U|$.

Moreover, for the remaining values of i, $\mu_i \leq \lambda_i - \alpha_{|U^c|} \leq 1/m^2 + L$. Since $G_{U^c,m^2}|_{U^c \times U^c} = (-\Delta'_{U^c} + m^2)^{-1}|_{U^c \times U^c}$, $\mu_{u_n} = \lambda_{\max}^{-1}(-\Delta'_{U^c} + m^2) = (m^2 + \lambda_{\max}(-\Delta'_{U^c}))^{-1} \geq (m^2 + 4d)^{-1}$. Indeed, by setting $f_x := 0$ on U,

$$\lambda_{\max}(-\Delta'_{U^c}) = \max_{\|f\|_2 = 1} \langle f, -\Delta'_{U^c} f \rangle \le 2d + 2 \max_{\|f\|_2 = 1} \Big| \sum_{e \in E(U^c \cup U)} f_{e-} f_{e+} \Big| \le 4d$$

since $\left|\sum_{e \in E(U^c \cup U)} f_{e-} f_{e+}\right| \leq \left(\sum_{e \in E(U^c \cup U)} f_{e-}^2\right)^{1/2} \left(\sum_{e \in E(U^c \cup U)} f_{e+}^2\right)^{1/2} = d\|f\|_2^2$. The desired result follows.

Remark 4. Note that equivalently the eigenvalues and the corresponding eigenfunctions as well as their domains can be indexed by $\hat{\Lambda}$ instead of $[0,n)^d \cap \mathbb{Z}^d$ due to the invariance of cosine and sine functions under translation of the argument by $2\pi\mathbb{Z}$. More precisely, for $w, x \in [0,n)^d \cap \mathbb{Z}^d$, $\eta_w = \eta_{\tilde{w}}$ and $q_x^w = q_{\tilde{x}}^{\tilde{w}}$, where $\tilde{y}_i = y_i$ if $y_i < n/2$ and $y_i - n$ otherwise for y = x, w.

2.4.2 Zero-average Green's function

We now turn to the zero-average Green's function, whose detailed understanding is a bit trickier, and start with the recollection of some known results stemming from [Abä17]:

Proposition 2.13 (Properties of zero-average Green's function). Let $d \geq 3, n \in \mathbb{N}$, $\Lambda = \mathbb{T}_n^d$. G_{Λ}^{0-avg} . satisfies the following properties:

- 1. For any $y \in \Lambda$, $\sum_{x \in \Lambda} G_{\Lambda}^{0-avg.}(x,y) = 0$.
- 2. Let $\overline{\mathbf{X}} = (\overline{X}_t)_{t \geq 0}$ be the continuous-time simple random walk on Λ viewed as a graph. We denote its law when started at x by \mathbb{P}^x_{Λ} . Then, for all $x, y \in \Lambda$,

$$G_{\Lambda}^{0\text{-}avg.}(x,y) = \frac{1}{2d} \int_0^{\infty} \left(\mathbb{P}_{\Lambda}^x [\overline{X}_t = y] - \frac{1}{n^d} \right) dt.$$

3. For any $U \subsetneq \Lambda$, $x, y \in \Lambda$,

$$G_{\Lambda}^{0-avg.}(x,y) = G_{U^c,0}(x,y) + \mathbb{E}_0^x [G_{\Lambda}^{0-avg.}(X_{H_U},y)] - \frac{1}{2dn^d} \mathbb{E}_0^x [H_U], \tag{2.11}$$

where $H_U := \inf\{n \geq 0 : X_n \in U\}$ and \mathbb{P}_0^x is the canonical law of a simple random walk on Λ started at x.

- 4. $G_{\Lambda}^{0\text{-}avg.}$ converges to $G_{\mathbb{Z}^d} := G_{\mathbb{Z}^d,0}$ uniformly over compact subsets of \mathbb{Z}^d viewed as properly contained in $\Lambda = \mathbb{T}_n^d$ for all n sufficiently large.
- 5. There exists C > 0 such that for all $x, y \in \Lambda, n \in \mathbb{N}$,

$$|G_{\Lambda}^{0-avg.}(x,y)| \le C(\log n)^{3d/2} d(x,y)^{2-d}.$$
 (2.12)

In most of our paper, the estimate (2.12) would have been sufficient, however, not for the critical regime in dimensions 3 and 4. Therefore, we need to improve on this bound.

Proposition 2.14 (Improved estimates on zero-average Green's function). Let $d \ge 3, n \in \mathbb{N}, \ \Lambda = \mathbb{T}_n^d$. $G_{\Lambda}^{0\text{-}avg.}$ satisfies the following properties:

1. There exists C = C(d) > 0 such that for all n sufficiently large and $y \in \hat{\Lambda}$,

$$|G_{\Lambda}^{0-avg.}(0,y) - G_{\mathbb{Z}^d}(0,y)| \le Cn^{2-d}.$$
 (2.13)

In particular, for $|y| = \Theta(n)$, $G_{\Lambda}^{0\text{-}avg.}(0,y) = \mathfrak{O}(n^{2-d})$.

2. For d=3 and any $x \in \mathbb{Z}^d$,

$$G_{\Lambda}^{0\text{-}avg.}(x,x) - G_{\mathbb{Z}^d}(x,x) = \Theta(n^{2-d}) < 0$$
 uniformly for large n. (2.14)

Proof. The proof of the estimate (2.13) is rather technical and is given in Appendix A.1; it mainly follows the argument verifying (2.12) in [Abä17, Proposition 1.5], but with greater precision.

(2.14) for d=3 can be obtained as follows. Let us fix x=0 and recall that $\hat{\Lambda} \subset [-n/2, n/2)^d \cap \mathbb{Z}^d$ is a canonical projection of Λ onto \mathbb{Z}^d . Let $\hat{U}=(\partial[-\lfloor n/2\rfloor, \lfloor n/2\rfloor]^d) \cap \mathbb{Z}^d$ and U be the corresponding preimage under the canonical projection. Then, since the random walk on torus killed upon entering U and the one on \mathbb{Z}^d (started in the interior of U) killed on \hat{U} have the same law, by (2.11) and 2.7:

$$G_{\Lambda}^{0\text{-avg.}}(0,0) = G_{U^{c},0}(0,0) + \mathbb{E}_{0}^{0}[G_{\Lambda}^{0\text{-avg.}}(X_{H_{U}},0)] - \frac{1}{2dn^{d}}\mathbb{E}_{0}^{0}[H_{U}]$$

$$= G_{\mathbb{Z}^{d}}(0,0) - \mathbb{E}_{0}^{0,\mathbb{Z}^{d}}[G_{\mathbb{Z}^{d}}(X_{H_{U}},0)\mathbb{1}_{\{H_{U}<\infty\}}] + \mathbb{E}_{0}^{0}[G_{\Lambda}^{0\text{-avg.}}(X_{H_{U}},0)] - \frac{1}{2dn^{d}}\mathbb{E}_{0}^{0}[H_{U}]$$

$$= G_{\mathbb{Z}^{d}}(0,0) - \mathbb{E}_{0}^{0,\mathbb{Z}^{d}}[G_{\mathbb{Z}^{d}}(X_{H_{U}},0)] + \mathbb{E}_{0}^{0}[G_{\Lambda}^{0\text{-avg.}}(X_{H_{U}},0)] - \frac{1}{2dn^{d}}\mathbb{E}_{0}^{0,\mathbb{Z}^{d}}[H_{U}]$$

(cf. [DH18, (15)]). By a standard argument (see e.g. [Law12, (1.21)]) since $|S_n|^2 - n$ is a martingale under $\mathbb{P}_0^{0,\mathbb{Z}^d}$, $\mathbb{E}_0^{0,\mathbb{Z}^d}[H_U] = \Theta(n^2)$. By [Law12, Theorem 1.5.4] we further know that $G_{\mathbb{Z}^d}(0,x) \sim C(d)|x|^{2-d}$ with $C(d) = \frac{1}{2d}\frac{d}{2}\Gamma(\frac{d}{2}-1)\pi^{-d/2}$ for $d \geq 3$. In A.2 (cf. (A.4)) we further show that $G_{\Lambda}^{0-\text{avg.}}(0,y)$ for any $y \in \hat{U}$ is bounded from above by $\left(\frac{1-0.04\pi+4\pi\log(3/2)+\pi e^{-1}}{(2\pi)^2}+c(\varepsilon)\right)n^{2-d}$ with $c(\varepsilon)>0$ that can be made arbitrary small for all sufficiently large n (as it in turn would allow the choice of small $\varepsilon>0$). With the above this implies that

$$G_{\Lambda}^{0-\text{avg.}}(0,0) - G_{\mathbb{Z}^{3}}(0,0) \leq -\inf_{r \in \left[\lfloor \frac{n}{2} \rfloor, \frac{\sqrt{3}n}{2} \right]} \left(\frac{C(3)}{r} + \frac{1}{6n^{3}} r^{2} \right)$$

$$+ \frac{1}{n} \left(\frac{1 - 0.04\pi + 4\pi \log(3/2) + \pi e^{-1}}{(2\pi)^{2}} + c(\varepsilon) \right)$$

$$= \frac{1}{n} \left(-\frac{(3C(3))^{2/3}}{2} + \frac{1 - 0.04\pi + 4\pi \log(3/2) + \pi e^{-1}}{(2\pi)^{2}} + c(\varepsilon) \right)$$

$$= \frac{1}{n} \left(-\frac{3^{2/3}}{2(4\pi)^{2/3}} + \frac{1 - 0.04\pi + 4\pi \log(3/2) + \pi e^{-1}}{(2\pi)^{2}} + c(\varepsilon) \right)$$

$$\leq -\frac{1}{100n}.$$

The infimum is attained at $r = (3C(3))^{1/3}n = \left(\frac{3}{4\pi}\right)^{1/3}n \in \left[\frac{n}{2}, \frac{\sqrt{3}n}{2}\right].$

We conclude this section with the following result on the eigenvalues of submatrices of $G_{\Lambda}^{0\text{-}\mathrm{avg.}}$.

Lemma 2.15 (Eigenvalues of zero-average Green's function). Let d, n and Λ be as in the previous proposition. Let U be a finite non-empty subset of \mathbb{Z}^d viewed as a subset of Λ for each n. Set $u_n := |U^c|$.

- 1. $G_{\Lambda}^{0-avg.}$ is positive semi-definite with eigenvalues given by 0 and $(1/\eta_k)_{k=2,...,n^d}$ with the latter as in Lemma 2.12.
- 2. Let $(\mu_k)_{k \leq u_n}$ ordered non-increasingly denote the eigenvalues of $G_{\Lambda}^{0\text{-}avg.}|_{U^c \times U^c} G_{\Lambda}^{0\text{-}avg.}|_{U^c \times U}G_{\Lambda}^{0\text{-}avg.}|_{U^2}G_{\Lambda}^{0\text{-}avg.}|_{U \times U^c}$. For all $k \leq u_n |U|$,

$$\mu_k \in \left[\frac{1}{\eta_{n^d - u_n + k + |U| + 1}}, \frac{1}{\eta_{k+1}}\right] \quad \left(=\left[0, \frac{1}{\eta_{k+1}}\right] \text{ for } k = u_n - |U|\right).$$
 (2.15)

Proof. The first item follows directly from the discussion in Subsection 2.2. Let $(\alpha_k)_{k \leq u_n}$ and $(\lambda_k)_{k \leq u_n}$ ordered non-increasingly denote the eigenvalues of $G_{\Lambda}^{0\text{-avg.}}|_{U^c \times U}G_{\Lambda}^{0\text{-avg.}}|_{U^2}G_{\Lambda}^{0\text{-avg.}}|_{U \times U^c}$ and $G_{\Lambda}^{0\text{-avg.}}|_{U^c \times U^c}$, respectively. Since the matrix $G_{\Lambda}^{0\text{-avg.}}|_{U^c \times U}G_{\Lambda}^{0\text{-avg.}}|_{U^2}G_{\Lambda}^{0\text{-avg.}}|_{U \times U^c}$ is positive semi-definite and has rank at most |U|, $\alpha_k = 0$ for all k > |U|. By Weyl's inequality [HJ12, Theorem 4.3.1], $\mu_k \in [\lambda_{k+|U|}, \lambda_k]$ for all $k \leq u_n - |U|$ and $\mu_k \in [\lambda_k - \alpha_1, \lambda_k]$ for all $k \leq u_n$. Furthermore, by [HJ12, Theorem 4.3.28], since $(1/\eta_k)_{k \geq 2}$ and 0 are the eigenvalues of $G_{\Lambda}^{0\text{-avg.}}$, $1/\eta_{n^d-u_n+k+1} \leq \lambda_k \leq 1/\eta_{k+1}$ for all $1 \leq k \leq u_n - 1$ and $0 \leq \lambda_{u_n} \leq 1/\eta_{u_n+1}$. (2.15) follows.

3 The infinite volume limit of the Spherical model

The main objective of this section is to understand the law of the infinite volume limit of the spherical model. To be exact, we prove the following result, which is a preciser version of Theorem 1.1.

Theorem 3.1 (Infinite volume limit of spherical model). Let $\Lambda_n = \mathbb{T}_n^d$ and $\boldsymbol{\theta} = (\theta_x)_{x \in \Lambda_n} \in \sqrt{n^d} \mathbb{S}^{n^d-1}$ be a configuration of the spherical model at inverse temperature $\beta > 0$, i.e., $\boldsymbol{\theta} \sim \nu_{\Lambda_n,\beta}$. Then for any finite box $U \subset \mathbb{Z}^d$ considered for each n sufficiently large as a subset of $\hat{\Lambda}_n$, $[\boldsymbol{\theta}_U]_n := [(\theta_x)_{x \in U}]_n$ converges in law as $n \to \infty$ to:

1. $\beta < \beta_c$: an m^2 -massive GFF on \mathbb{Z}^d restricted to U scaled by $1/\sqrt{\beta}$ with the mass m^2 depending on β and d in such a way that

$$G_{\mathbb{Z}^d,m^2}(x,x) = \beta \text{ for all } x \in \mathbb{Z}^d;$$

- 2. $\beta = \beta_c$: a GFF on \mathbb{Z}^d restricted to U scaled by $1/\sqrt{\beta}$;
- 3. $\beta > \beta_c$: a GFF on \mathbb{Z}^d restricted to U scaled by $1/\sqrt{\beta}$ plus an independent constant (in space) random drift of the form $\sqrt{\frac{\beta-\beta_c}{\beta}}X\mathbf{1}_U$ with X being a Rademacher random variable, i.e., $\mathbb{P}[X=1] = \mathbb{P}[X=-1] = \frac{1}{2}$.

The proof consists of three parts corresponding to each of the regimes, which are discussed separately in the following three subsections.

Based on this result we further establish convergence of local covariance functions of the spherical model. More precisely, in Section 3.4 we prove the following.

Corollary 3.2. Let Λ_n , $\boldsymbol{\theta} = (\theta_x)_{x \in \Lambda_n}$ and $U \subset \mathbb{Z}^d$ be as above, $(i_x)_{x \in U} \in \mathbb{N}_0^{|U|}$ be an arbitrary but fixed vector of non-negative integers. Then the following holds:

$$\nu_{\Lambda_n,\beta} \bigg[\prod_{x \in U} \theta_x^{i_x} \bigg] \xrightarrow{n \to \infty} \mathbb{E} \bigg[\prod_{x \in U} \alpha_x^{i_x} \bigg],$$

where $\sqrt{\beta}\alpha$ in correspondence with Theorem 3.1 is either an m^2 -massive GFF on \mathbb{Z}^d if $\beta < \beta_c$, a GFF on \mathbb{Z}^d if $\beta = \beta_c$ or a GFF on \mathbb{Z}^d plus an independent drift $\sqrt{\beta - \beta_c} X \mathbf{1}_{\mathbb{Z}^d}$ with a Rademacher random variable X if $\beta > \beta_c$.

3.1 The high-temperature regime

This subsection presents a proof of $\beta < \beta_c$ case of Theorem 3.1. We start by recalling that Proposition 2.8 gives us,

$$\operatorname{Law}(\boldsymbol{\theta}) = \operatorname{Law}\left(\boldsymbol{\varphi}/\sqrt{\beta} \mid \|\boldsymbol{\varphi}\|_2 = \sqrt{\beta n^d}\right)$$

for any m_n^2 -massive GFF φ with $m_n^2 > 0$. Thus, for any finite box $U \subset \mathbb{Z}^d$ viewed for each n as a subset of Λ_n , we have the following equality of Lebesgue densities on $\mathbb{R}^{|U|}$:

$$f_{\boldsymbol{\theta}_{U}}(\gamma) = f_{\frac{\boldsymbol{\varphi}_{U}}{\sqrt{\beta}} | \|\boldsymbol{\varphi}\|_{2} = \sqrt{\beta n^{d}}}(\gamma) = f_{\frac{\boldsymbol{\varphi}_{U}}{\sqrt{\beta}}}(\gamma) \underbrace{\frac{f_{\|(\tilde{\boldsymbol{\varphi}}^{U} + h^{\sqrt{\beta}\gamma})\|_{2}^{2}}(\beta n^{d})}{f_{\|\boldsymbol{\varphi}\|_{2}^{2}}(\beta n^{d})}}_{J_{r}}, \tag{3.1}$$

where $\gamma \coloneqq (\gamma_x)_{x \in U} \in \mathbb{R}^{|U|}$, $\|\cdot\|_2$ interchangeably denotes Euclidean norms on $\mathbb{R}^{|\Lambda_n|}$ and $\mathbb{R}^{|U|}$. In the second equality we applied Proposition 2.6 using its notation with K = U. Since φ_U is a centered Gaussian vector with covariance $(G_{\Lambda_n,m_n^2}(x,y))_{x,y \in U}$, our discussion right after Definition 2.5 yields its convergence in law towards a centered Gaussian vector with covariance $(G_{\mathbb{Z}^d,m^2}(x,y))_{x,y \in U}$, which in turn is an m^2 -massive GFF ψ on \mathbb{Z}^d restricted to U, as long as $m_n^2 > 0$ converges to some $m^2 > 0$. As we are working with Gaussians, it also implies pointwise convergence of the corresponding densities. In particular, $f_{\frac{\varphi_U}{\sqrt{\beta}}}(\gamma) \to f_{\frac{\psi_U}{\sqrt{\beta}}}(\gamma)$. We should now choose $m_n^2 > 0$ so that the ratio in (3.1) tends to 1. Since $\mathbb{E}[\|\varphi\|_2^2] = G_{\Lambda_n,m_n^2}(0,0)n^d$, a natural guess to get rid of conditioning on the norm of φ in the limit would be to pick $m_n^2, m^2 > 0$ such that $\lim_n G_{\Lambda_n,m_n^2}(0,0) = G_{\mathbb{Z}^d,m^2}(0,0) = \beta$. This is indeed possible since $\beta < \beta_c$ and the critical inverse temperature $\beta_c \in (0,\infty]$ in terms of Green's functions corresponds to $G_{\mathbb{Z}^d}(0,0)$. In particular, $\beta_c = \infty$ if d = 2, $\beta_c \in (0,\infty)$ for $d \geq 3$. We can further pick the sequence $(m_n^2)_n$ such that $G_{\Lambda_n,m_n^2}(0,0) = \beta$ for all n.

Let us conclude the proof of this step by showing that with the choice of $(m_n^2)_n$ as above, the ratio in (3.1) converges to one. Let $Q \in \mathbb{R}^{U^c \times U^c}$, $P \in \mathbb{R}^{\Lambda_n \times \Lambda_n}$ be the orthonormal matrices diagonalizing $G_{U^c, m_n^2}|_{U^c \times U^c}$ and G_{Λ_n, m_n^2} , respectively. Then,

$$Q^{T}(G_{U^{c},m_{n}^{2}}|_{U^{c}\times U^{c}})Q = \operatorname{diag}\left((\mu_{i})_{i=1,\dots,|U^{c}|}\right),$$
$$P^{T}G_{\Lambda_{n},m_{n}^{2}}P = \operatorname{diag}\left(\lambda_{i} := (\eta_{i} + m_{n}^{2})^{-1}\right)_{i=1}^{n^{d}}$$

where $\mu_1 \geq \ldots \geq \mu_{|U^c|} > 0$ are the eigenvalues of $G_{U^c, m_n^2}|_{U^c \times U^c}$, $0 = \eta_1 < \eta_2 \leq \ldots \eta_{n^d}$ those of $-\Delta_{\Lambda_n}$. We set $u_n \coloneqq |U^c|$, $\tilde{h} \coloneqq Q^T h_{U^c}^{\sqrt{\beta}\gamma}$, $Z_{i,u_n} \coloneqq (\tilde{h}_i + \sqrt{\mu_i} Y_{i,u_n})^2 - \mu_i - \tilde{h}_i^2$ for a triangle array $(Y_{i,u_n})_{i\leq u_n,u_n\in\mathbb{N}}$ of i.i.d. standard normal random variables. Then,

$$J_n(u) = \frac{f_{\|\tilde{h} + \mathcal{N}(0, \operatorname{diag}(\mu_j)_{j=1}^{u_n})\|_2^2}(\beta n^d - \beta \|\gamma\|_2^2)}{f_{\|\mathcal{N}(0, \operatorname{diag}(\lambda_j)_{j=1}^{n^d})\|_2^2}(\beta n^d)}$$

$$= \frac{f_{\sum_{i=1}^{u_n} Z_{i, u_n}}(\beta n^d - \sum_{i=1}^{u_n} \mu_i - \|\tilde{h}\|_2^2 - \beta \|\gamma\|_2^2)}{f_{\sum_{i=1}^{n^d} \lambda_i(Y_{i, n^d}^2 - 1)}(0)}.$$

Here we used that $\beta n^d = \text{Trace}(G_{\Lambda_n, m_n^2}) = \sum_{i=1}^{n^d} \lambda_i$. Recall also that by (2.7),

$$\begin{split} \sum_{i=1}^{u_n} \mu_i + \|\tilde{h}\|_2^2 &= \operatorname{Trace}\left(G_{U^c, m_n^2}|_{U^c \times U^c}\right) + \|h_{U^c}^{\sqrt{\beta}\gamma}\|_2^2 \\ &= \operatorname{Trace}\left(G_{\Lambda_n, m_n^2}|_{U^c \times U^c}\right) - \sum_{x \in U^c} \mathbb{E}_{\alpha_n}^x [\mathbb{1}_{\{H_U < \infty\}} G_{\Lambda_n, m_n^2}(X_{H_U}, x)] + \|h_{U^c}^{\sqrt{\beta}\gamma}\|_2^2 \\ &= (n^d - |U|)\beta - \sum_{y \in U} \sum_{x \in U^c} \mathbb{P}_{\alpha_n}^x [H_U < \infty, X_{H_U} = y] G_{\Lambda_n, m_n^2}(y, x) + \|h_{U^c}^{\sqrt{\beta}\gamma}\|_2^2 \\ &=: n^d \beta - R_{U,n}. \end{split}$$

Note that $0 \le |R_{U,n}| \le \beta |U| + \sum_{y \in U} \sum_{x \in U^c} Ce^{-c|x-y|} + ||h_{U^c}^{\sqrt{\beta}\gamma}||_2^2 \le M$ for an absolute constant M > 0. Indeed, by Jensen's inequality

$$\|h_{U^c}^{\sqrt{\beta}\gamma}\|_2^2 \leq \|\gamma\|_2^2 \sum_{x \in U^c} \mathbb{P}_{\alpha_n}^x [H_U < \infty]^2,$$

where $\tilde{H}_U := \{n \geq 1 : X_n \in U\}$. The latter probability is bounded by $Ce^{-c \operatorname{dist}(x,U)}$ for some c, C > 0 depending only on m_n^2, d (this follows from the weak Markov property of the random walk $\mathbf{X} \sim \mathbb{P}_{\alpha_n}^x$ and (2.8)):

$$\begin{split} \mathbb{P}^x_{\alpha_n}[H_U < \infty] &= \sum_{k \geq 0} \sum_{y \in U} \mathbb{P}^x_{\alpha_n}[X_k = y, \tilde{H}_U \circ \tau_k = \infty] \\ &\leq \sum_{y \in U} \mathbb{P}^y_{\alpha_n}[\tilde{H}_U = \infty] \underbrace{\sum_{k \geq 0} \mathbb{P}^x_{\alpha_n}[X_k = y]}_{= \frac{2d}{1 - \alpha_n} G_{\Lambda_n, m_n^2}(x, y)} \\ &\stackrel{\text{(2.8)}}{\leq} \frac{2d}{1 - \alpha_n} \sum_{y \in U} C(m^2, d) e^{-c(m^2, d)|x - y|} \\ &\leq \tilde{C}|U| \exp\left(-c \min_{y \in U} |x - y|\right). \end{split}$$

Let us introduce the variances of the two sums $\sum_{i=1}^{u_n} Z_{i,u_n}$ and $\sum_{i=1}^{n^d} \lambda_i(Y_{i,n^d}^2 - 1)$, denoted s_n^2 and σ_n^2 , respectively. Using the bounds obtained in Lemma 2.12, we can deduce that

$$\sigma_n^2 = 2\sum_{k=1}^{n^d} \lambda_k^2 = \Theta(n^d); \qquad s_n^2 = 2\sum_{k=1}^{u_n} \mu_k^2 + 4\sum_{j=1}^{u_n} \mu_j \tilde{h}_j^2 = \sigma_n^2 (1 - o(1)).$$

Then.

$$J_n(u) = (1 + o(1)) \frac{s_n f_{\sum_{i=1}^{u_n} Z_{i,u_n}} \left(s_n \frac{R_{U,n} - \beta ||\gamma||_2^2}{s_n} \right)}{\sigma_n f_{\sum_{i=1}^{n^d} \lambda_i (Y_{i,n^d}^2 - 1)}(0)}.$$

We use the following result (stated in more general form than required here to later be applied to the critical case) proven in Appendix A.4 to conclude the proof.

Proposition 3.3. Let $(X_{i,n})_{i \leq n,n \in \mathbb{N}}$ be a triangular array of independent centered random variables with the probability density functions $f_{i,n}$ (w.r.t. Lebesgue measure on \mathbb{R}). Assume that

- 1. $f_{i,n} \in L^r(\mathbb{R})$ for some $r \in (1,2]$ (independent of i,n) such that $\sup_{i,n} ||f_{i,n}||_{L^r} \leq M$ for some M > 0:
- 2. for all $i \leq n, n \in \mathbb{N}$, $\sigma_{i,n}^2 := \operatorname{Var}[X_{i,n}] < \infty$ ordered such that $\sigma_{1,n}^2 \geq \ldots \geq \sigma_{n,n}^2$;
- 3. Lindeberg's condition is satisfied, that is, for any $\varepsilon > 0$,

$$\frac{\sum_{i\geq 1} \mathbb{E}[X_{i,n}^2 \mathbb{1}_{\{|X_{i,n}| > \varepsilon s_n\}}]}{s_n^2} \xrightarrow{n \to \infty} 0,$$

where $s_n^2 := \sum_{i=1}^n \sigma_{i,n}^2$.

4. there exist $\delta > 0$ (uniform), $K(n) \geq 1$, $l_*(n) \geq 1$, $n \geq l^*(n) \geq 2\lceil \frac{r}{r-1} \rceil$ such that for all n sufficiently large

$$(a) \frac{\sum_{i \geq l_{*}(n)} \sigma_{i,n}^{2}}{\sum_{i=1}^{n} \sigma_{i,n}^{2}} \geq \delta;$$

$$(b) \frac{\sum_{i \geq l_{*}(n)} \mathbb{E}[X_{i,n}^{2} \mathbb{1}_{\{|X_{i,n}| > K(n)\}}]}{\sum_{i \geq l_{*}(n)} \sigma_{i,n}^{2}} \leq \frac{1}{8};$$

$$(c) \frac{n - l^{*}(n)}{\sigma_{l^{*} n}^{2} \vee K(n)^{2}} \gg \log\left(\sum_{i=1}^{n} \sigma_{i,n}^{2}\right).$$

Then the relation

$$s_n f^{(n)}(s_n x) \xrightarrow{n \to \infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

holds uniformly over \mathbb{R} . Here, $f^{(n)}$ is the convolution of $f_{1,n},\ldots,f_{n,n}$, as well as the density function of $\sum_{i=1}^{n} X_{i,n}$. Note that $s_n f^{(n)}(s_n x)$ is the density of $\frac{\sum_{i=1}^{n} X_{i,n}}{s_n}$.

Remark 5. Note that the fourth assumption is redundant if $(\sigma_{i,n}^2)_{i,n}$ are uniformly bounded from above and away from zero.

We now check that the array $(Z_{i,u_n})_{i,u_n}$ (and fully analogously $(\lambda_i(Y_{i,n^d}^2-1))_{i,n^d})$ satisfies the required assumptions (up to reordering) with $l_*=1, l^*=2\lceil\frac{r}{r-1}\rceil$ and K large, but independent of n. It is clear that Z_{i,u_n} 's are centered and independent, $\sigma_{i,u_n}^2 := \operatorname{Var}[Z_{i,u_n}] =$ $2\mu_i^2 + 4\tilde{h}_i^2\mu_i \in (0,\infty)$. For the assumption 1., let f_{i,u_n} be the probability density function of Z_{i,u_n} , which is given by

$$f_{i,u_n}(t) = \frac{\left(f_Y\left(\frac{\sqrt{t + \mu_i + \tilde{h}_i^2} - \tilde{h}_i}{\sqrt{\mu_i}}\right) + f_Y\left(\frac{\sqrt{t + \mu_i + \tilde{h}_i^2} + \tilde{h}_i}{\sqrt{\mu_i}}\right)\right)}{2\sqrt{\mu_i}\sqrt{t + \mu_i + \tilde{h}_i^2}} \mathbb{1}_{\{t > -\mu_i - \tilde{h}_i^2\}},$$

where Y is a standard normal random variable. Hence, for any $r \in (1,2)$,

$$||f_{i,u_n}||_{L^r}^r = \int_0^\infty \frac{\left(f_Y\left(\frac{\sqrt{s}-\tilde{h}_i}{\sqrt{\mu_i}}\right) + f_Y\left(\frac{\sqrt{s}+\tilde{h}_i}{\sqrt{\mu_i}}\right)\right)^r}{2^r \mu_i^{r/2} s^{r/2}} \mathrm{d}s \le \int_0^\infty \frac{f_Y^r\left(\frac{\sqrt{s}-|\tilde{h}_i|}{\sqrt{\mu_i}}\right)}{\mu_i^{r/2} s^{r/2}} \mathrm{d}s$$

$$= \int_{0}^{\infty} \frac{f_{Y}^{r}\left(\sqrt{s} - \left|\frac{\tilde{h}_{i}}{\sqrt{\mu_{i}}}\right|\right)}{\mu_{i}^{r-1}s^{r/2}} ds = 2 \int_{0}^{\infty} \frac{f_{Y}^{r}\left(t - \left|\frac{\tilde{h}_{i}}{\sqrt{\mu_{i}}}\right|\right)}{\mu_{i}^{r-1}t^{r-1}} dt$$

$$\leq \frac{2}{\mu_{i}^{r-1}} \left(\int_{0}^{1} t^{1-r} dt + \int_{1}^{\infty} f_{Y}\left(t - \left|\frac{\tilde{h}_{i}}{\sqrt{\mu_{i}}}\right|\right) dt\right)$$

$$\leq \frac{2}{\mu_{i}^{r-1}} \left(\frac{1}{2-r} + 1\right) =: C(r) \frac{1}{\mu_{i}^{r-1}}$$

and so, $\sup_{i,n} \|f_{i,u_n}\|_{L^r}^r \leq C(r)/(\inf_i \mu_i)^{r-1} \leq C(r)(4d+2m^2)^{r-1}$ as follows from (2.9) (since $m_n^2 \xrightarrow{n \to \infty} m^2 > 0$). The latter, moreover, shows that for all $i \leq u_n$ uniformly in $n, \mu_i \in [(2m^2+4d)^{-1},2/m^2+L] =: [L_*,L^*]$. Besides that, we have seen above that there exist c,C>0 such that

$$\tilde{h}_i^2 \le \|h_{U^c}^{\sqrt{\beta}\gamma}\|_2^2 \le C(m^2, d) \|\gamma\|_2^2,$$

Therefore, $0 < 2L_*^2 \le \sigma_{i,u_n}^2 \le 2(L^*)^2 + 4C\|\gamma\|_2^2 L^* < \infty$ for all i, n uniformly (assumption 2. up to reordering). This implies that $2L_*^2 u_n \le s_n^2 \le (2(L^*)^2 + 4C\|\gamma\|_2^2 L^*)u_n$, so, $s_n^2 = \Theta(u_n)$. Let us check 4. (and 3.): (a) and (c) are obvious with our choice of l_*, l^* and K to be independent of n. For all K sufficiently large (greater than $4(L^* + C\|\gamma\|_2^2)$),

$$\{|Z_{i,u_n}| > K\}$$

$$= \left\{ (\tilde{h}_i + \sqrt{\mu_i} Y_{i,u_n})^2 > \underbrace{(\mu_i + \tilde{h}_i^2) + K}_{>K} \right\} \cup \left\{ (\tilde{h}_i + \sqrt{\mu_i} Y_{i,u_n})^2 < \underbrace{(\mu_i + \tilde{h}_i^2) - K}_{\leq L^* + C \|\gamma\|_2^2 - K < 0} \right\}$$

$$\subset \left\{ \sqrt{\mu_i} Y_{i,u_n} > \sqrt{K} - \tilde{h}_i \right\} \cup \left\{ \sqrt{\mu_i} Y_{i,u_n} < -\sqrt{K} - \tilde{h}_i \right\}$$

$$\subset \left\{ Y_{i,u_n} > (\sqrt{K} - \sqrt{C} \|\gamma\|_2) / \sqrt{L^*} \right\} \cup \left\{ Y_{i,u_n} < (-\sqrt{K} + \sqrt{C} \|\gamma\|_2) / \sqrt{L^*} \right\}$$

$$\subset \left\{ |Y_{i,u_n}| > \sqrt{K} / (2\sqrt{L^*}) \right\}.$$

We further have $Z_{i,u_n}^2 = \left(\mu_i(Y_{i,u_n}^2 - 1) + 2\tilde{h}_i\sqrt{\mu_i}Y_{i,u_n}\right)^2 \le (L^*)^2(Y_{i,u_n}^2 - 1)^2 + 4CL^*\|\gamma\|_2^2Y_{i,u_n}^2 + 4\sqrt{C}(L^*)^{3/2}\|\gamma\|_2|Y_{i,u_n}(Y_{i,u_n}^2 - 1)| \le \tilde{C}(Y_{i,u_n}^4 + |Y_{i,u_n}|^3 + Y_{i,u_n}^2 + |Y_{i,u_n}| + 1).$ Hence,

$$\frac{\mathbb{E}[Z_{i,u_n}^2 \mathbb{1}_{\{|Z_{i,u_n}| > K\}}]}{\sigma_{i,u_n}^2} \leq \tilde{C} \frac{\mathbb{E}\Big[(Y_{i,u_n}^4 + |Y_{i,u_n}|^3 + Y_{i,u_n}^2 + |Y_{i,u_n}| + 1) \mathbb{1}_{\{|Y_{i,u_n}| > \sqrt{K}/(2\sqrt{L^*})\}} \Big]}{\sigma_{i,u_n}^2} \\
\leq \frac{5\tilde{C}}{2L_*^2} \mathbb{E}\Big[Y_{i,u_n}^4 \mathbb{1}_{\{|Y_{i,u_n}| > \sqrt{K}/(2\sqrt{L^*})\}} \Big] = C' \mathbb{E}\Big[Y^4 \mathbb{1}_{\{|Y| > \sqrt{K}/(2\sqrt{L^*})\}} \Big].$$

and thus also, for any $\varepsilon > 0$, and all n sufficiently large,

$$\frac{\mathbb{E}[Z_{i,u_n}^2 \mathbb{1}_{\{|Z_{i,u_n}| > \varepsilon s_n\}}]}{\sigma_{i,u_n}^2} \le C' \mathbb{E}[Y^4 \mathbb{1}_{\{|Y| > \sqrt{\varepsilon s_n}/(2\sqrt{L^*})\}}].$$

The two estimates yield,

$$\frac{\sum_{i=1}^{u_n} \mathbb{E}[Z_{i,u_n}^2 \mathbb{1}_{\{|Z_{i,u_n}| > R\}}]}{s_n^2} \le C' \mathbb{E}[Y^4 \mathbb{1}_{\{|Y| > \sqrt{R}/(2\sqrt{L^*})\}}]$$

for $R \in \{K, \varepsilon s_n\}$. This can be made arbitrarily small by choosing K > 0 sufficiently large and for εs_n , the right-hand side clearly vanishes as $n \to \infty$.

This completes the verification of all the assumptions of Proposition 3.3 and accordingly implies that $s_n f_{\sum_{i=1}^{v_n} Z_{i,v_n}}(s_n x)$ and $\sigma_n f_{\sum_{i=1}^{n^d} \lambda_i(Y_{i,n^d}^2 - 1)}(\sigma_n x)$ converge to $f_{\mathcal{N}(0,1)}(x)$ uniformly over $x \in \mathbb{R}$, and herewith that $J_n \xrightarrow{n \to \infty} 1$ as desired.

3.2 The low-temperature regime

The current subsection proves the $\beta > \beta_c$ case of Theorem 3.1 and lays the groundwork for the critical case. In particular, we assume that $d \geq 3$ and $\beta \geq \beta_c$ and specify explicitly when we restrict to $\beta > \beta_c$. Recall that by Proposition 2.10,

Law
$$(\boldsymbol{\theta}) = \text{Law}\left((\boldsymbol{\gamma} + c\mathbf{1}_{\Lambda_n})/\sqrt{\beta} \mid ||\boldsymbol{\gamma} + c\mathbf{1}_{\Lambda_n}||_2 = \sqrt{\beta n^d}\right)$$

for an independent pair consisting of a zero-average GFF γ on Λ_n and a Lebesgue constant c. Our strategy will be based on the analysis of the joint law of (γ_U, c) conditioned on $\|\gamma + c \mathbf{1}_{\Lambda_n}\|_2 = \sqrt{\beta n^d}$ for a finite box $U \subset \mathbb{Z}^d$ (viewed as a subset of $\Lambda_n = \mathbb{T}_n^d$). More precisely, we will show that it converges to the product measure of the GFF ψ on \mathbb{Z}^d restricted to U and a Rademacher random variable X scaled by $\sqrt{\beta - \beta_c}$. Note that this would directly yield the desired local convergence in distribution of θ towards $\frac{1}{\sqrt{\beta}}\psi + \sqrt{\frac{\beta - \beta_c}{\beta}}X\mathbf{1}_{\mathbb{Z}^d}$.

Let $B \subset \mathbb{R}^U$ be a Borel set and $I = (a, b) \subset [0, \sqrt{\beta}]$. Observe that since γ is a zero-average GFF, $\langle \gamma, \mathbf{1}_{\Lambda_n} \rangle = 0$ almost surely. Therefore,

$$\mathbb{P}\left[\gamma_{U} \in B, c \in I \mid \|\gamma + c\mathbf{1}_{\Lambda_{n}}\|_{2} = \sqrt{\beta n^{d}}\right] = \mathbb{P}\left[\gamma_{U} \in B, c \in I \mid \|\gamma\|_{2}^{2} + c^{2}n^{d} = \beta n^{d}\right] \\
= \int_{B} dy \int_{I} dc_{0} \frac{f_{\gamma_{U}, \|\gamma\|_{2}^{2}}(y, n^{d}(\beta - c_{0}^{2}))}{f_{\|\gamma\|_{2}^{2} + c^{2}n^{d}}(\beta n^{d})} = \int_{B} dy f_{\gamma_{U}}(y) \frac{\int_{I} dc_{0} f_{\|\hat{\gamma}^{y}\|_{2}^{2}}(n^{d}(\beta - c_{0}^{2}))}{\int_{-\sqrt{\beta}}^{\sqrt{\beta}} d\tilde{c} f_{\|\gamma\|_{2}^{2}}(n^{d}(\beta - \tilde{c}^{2}))} \\
= \int_{B} dy f_{\gamma_{U}}(y) \frac{\int_{a^{2}}^{b^{2}} du \frac{1}{2\sqrt{u}} n^{d} f_{\|\hat{\gamma}^{y}\|_{2}^{2}}(n^{d}(\beta - u))}{2\int_{0}^{\beta} d\tilde{u} \frac{1}{2\sqrt{u}} \int_{0}^{b^{2}} du \frac{1}{\sqrt{u}} f_{\|\hat{\gamma}^{y}\|_{2}^{2}}(\beta - u)}{2\int_{0}^{\beta} d\tilde{u} \frac{1}{\sqrt{u}} \int_{0}^{b^{2}} du \frac{1}{\sqrt{u}} f_{\|\hat{\gamma}^{y}\|_{2}^{2}}(\beta - \tilde{u})} \\
= \int_{B} dy f_{\gamma_{U}}(y) \frac{\mathbb{E}\left[\frac{1}{\sqrt{\beta - \|\hat{\gamma}^{y}\|_{2}^{2}/n^{d}}} \mathbb{1}_{(\beta - b^{2}, \beta - a^{2})} \left(\frac{\|\hat{\gamma}^{y}\|_{2}^{2}}{n^{d}}\right)\right]}{2\mathbb{E}\left[\frac{1}{\sqrt{\beta - \|\hat{\gamma}^{y}\|_{2}^{2}/n^{d}}} \mathbb{1}_{[0,\beta)} \left(\frac{\|\hat{\gamma}^{y}\|_{2}^{2}}{n^{d}}\right)\right]} =: \int_{B} dy f_{\gamma_{U}}(y) R_{n}(y),$$

where $\hat{\gamma}^y$ is defined by $\hat{\gamma}_U^y \equiv y$ and

$$\hat{\gamma}_{U^{c}}^{y} \sim \mathcal{N}\left(G_{\Lambda_{n}}^{0\text{-avg.}}|_{U^{c}\times U}G_{\Lambda_{n}}^{0\text{-avg.}}|_{U^{2}}^{-1}y, G_{\Lambda_{n}}^{0\text{-avg.}}|_{U^{c}\times U^{c}} - G_{\Lambda_{n}}^{0\text{-avg.}}|_{U^{c}\times U}G_{\Lambda_{n}}^{0\text{-avg.}}|_{U^{2}}G_{\Lambda_{n}}^{0\text{-avg.}}|_{U\times U^{c}}\right)
=: \mathcal{N}(\nu(y), C) = \text{Law}(\gamma_{U^{c}}|\gamma_{U} = y).$$
(3.2)

Further note that we can assume that B is bounded since $\mathbb{E}[\|\boldsymbol{\gamma}_U\|_2^2 \mid \|\boldsymbol{\gamma}\|_2^2 + c^2 n^d = \beta n^d] = |U|\mathbb{E}[\gamma_0^2 \mid \|\boldsymbol{\gamma}\|_2^2 + c^2 n^d = \beta n^d] \leq |U|\beta$ by symmetries of the torus, and hence, by taking t > 0 large enough, $\mathbb{P}\left[\boldsymbol{\gamma}_U \notin B^{|U|}(0,t) \mid \|\boldsymbol{\gamma} + c\mathbf{1}_{\Lambda_n}\|_2 = \sqrt{\beta n^d}\right]$ can be made arbitrarily small.

Goal: Show that uniformly over bounded $B \ni y$,

$$R_{n}(y) = \frac{\mathbb{E}\left[\frac{1}{\sqrt{\beta - \|\hat{\gamma}^{y}\|_{2}^{2}/n^{d}}} \mathbb{1}_{(\beta - b^{2}, \beta - a^{2})} \left(\frac{\|\hat{\gamma}^{y}\|_{2}^{2}}{n^{d}}\right)\right]}{2\mathbb{E}\left[\frac{1}{\sqrt{\beta - \|\gamma\|_{2}^{2}/n^{d}}} \mathbb{1}_{[0, \beta)} \left(\frac{\|\gamma\|_{2}^{2}}{n^{d}}\right)\right]} \xrightarrow{n \to \infty} \frac{1}{2} \mathbb{1}_{I}(\sqrt{\beta - \beta_{c}}).$$
(3.3)

Note that by symmetry of c under the conditional law, we get the same result for -I. Altogether, this easily gives the desired statement about the limiting joint law.

To prove the goal for $\beta > \beta_c$ we proceed in three steps (two of which also hold for $\beta = \beta_c$):

Claim 1: $\|\gamma\|_2^2/n^d$ and $\|\hat{\gamma}^y\|_2^2/n^d$ converge almost surely towards β_c .

Claim 2: If $\beta > \beta_c$, there exist $C = C(\beta, d)$, $\hat{C} = \hat{C}(\beta, d) > 0$ such that for any $0 < \varepsilon < (\beta - \beta_c)/4$,

$$\mathbb{P}[S_n \ge \beta - \beta_c - \varepsilon] \le e^{-Cn^{d-2}}, \qquad \mathbb{P}[\hat{S_n}^y \ge \beta - \beta_c - \varepsilon] \le e^{-\hat{C}n^{d-2}}.$$

<u>Claim 3:</u> The density functions of $\|\gamma\|_2^2$ and $\|\hat{\gamma}^y\|_2^2$ are uniformly bounded. The latter also holds uniformly in y in a bounded Borel set B.

In view of (3.3) and due to expected high concentration of $\|\hat{\gamma}^y\|_2^2/n^d$ around its mean, the first step is self-explanatory. The latter two statements prove that the contribution of the regions $(\beta - \varepsilon, \beta)$ to the integrals in the denominator and numerator are negligible. One direct way to verify that would have been to show that the density function of $\|\hat{\gamma}^y\|_2^2/n^d$ is bounded on this interval, however, it is a highly non-trivial fact. So, instead we split $(\beta - \varepsilon, \beta)$ into further subintervals on one of which we could use that as expected the concentration away from β_c is extremely low, and on the remaining "infinitesimally" small interval bordering with β – integrability of $x \mapsto 1/\sqrt{\beta - x}$.

Combined the claims yield (3.3) in the low-temperature regime. Indeed, for $\varepsilon > 0$ small enough, since $\mathbb{R}_+ \ni x \mapsto 1/\sqrt{x}$ away from zero is bounded, by dominated convergence theorem and Claim 1 we get

$$\mathbb{E}\Big[\frac{1}{\sqrt{\beta-\|\hat{\gamma}^y\|_2^2/n^d}}\mathbb{1}_{(\beta-b^2,(\beta-a^2)\wedge(\beta-\varepsilon))}\Big(\frac{\|\hat{\gamma}^y\|_2^2}{n^d}\Big)\Big] \xrightarrow{n\to\infty} \frac{1}{\sqrt{\beta-\beta_c}}\mathbb{1}_{(a,b)}(\sqrt{\beta-\beta_c});$$

$$\mathbb{E}\Big[\frac{1}{\sqrt{\beta-\|\gamma\|_2^2/n^d}}\mathbb{1}_{[0,\beta-\varepsilon)}\Big(\frac{\|\gamma\|_2^2}{n^d}\Big)\Big] \xrightarrow{n\to\infty} \frac{1}{\sqrt{\beta-\beta_c}}.$$

Let us consider the remaining bits and show that they are vanishing in the limit. Let $\delta > 0$ be small,

$$0 \leq \mathbb{E}\left[\frac{1}{\sqrt{\beta - \|\boldsymbol{\gamma}\|_{2}^{2}/n^{d}}}\mathbb{1}_{[\beta - \varepsilon, \beta)}\left(\frac{\|\boldsymbol{\gamma}\|_{2}^{2}}{n^{d}}\right)\right]$$

$$\leq \frac{n^{d}}{\sqrt{\delta}} \mathbb{P}\left[\frac{\|\boldsymbol{\gamma}\|_{2}^{2}}{n^{d}} \in [\beta - \varepsilon, \beta - \delta/n^{2d})\right] + \sqrt{n^{d}} \mathbb{E}\left[\frac{1}{\sqrt{\beta n^{d} - \|\boldsymbol{\gamma}\|_{2}^{2}}}\mathbb{1}_{\beta n^{d} + [-\delta/n^{d}, 0)}(\|\boldsymbol{\gamma}\|_{2}^{2})\right].$$

$$\stackrel{\text{Claim 2}}{\leq} e^{-Cn^{d-2}} \stackrel{\text{Claim 3}}{\leq} \|f_{\|\boldsymbol{\gamma}\|_{2}^{2}}\|_{L^{\infty}} \int_{0}^{\delta/n^{d}} \mathrm{d}t/\sqrt{t} = 2\sqrt{\frac{\delta}{n^{d}}}$$

By first taking limit $n \to \infty$, we eliminate the first term, the second one can be then made arbitrarily small by adjusting δ . By replacing γ with $\hat{\gamma}^y$, we recover the same result for the latter. The claim concerning $R_n(y)$ follows immediately from the above two observations.

Let us now prove the claims.

Proof of Claim 1. For simplicity set $S_n := \frac{\|\gamma\|_2^2 - \mathbb{E}[\|\gamma\|_2^2]}{n^d} = \frac{\|\gamma\|_2^2 - n^d G^{0\text{-avg.}}(0,0)}{n^d}$ and $\hat{S}_n^y := \frac{\|\hat{\gamma}^y\|_2^2 - \mathbb{E}[\|\hat{\gamma}^y\|_2^2]}{n^d} = \frac{\|\hat{\gamma}_U^y\|_2^2 - \mathbb{E}[\|\hat{\gamma}_U^y\|_2^2]}{n^d}$. We will show that these random variables converge to zero almost surely, which in turn implies the claim. The latter remark is due to the fact that $G_{\Lambda_n}^{0\text{-avg.}}(0,0) \xrightarrow{n \to \infty} \beta_c$ and since $\frac{1}{n^d} \mathbb{E}[\|\hat{\gamma}_U^y\|_2^2] = \frac{1}{n^d} \mathrm{Trace}(C) + \frac{1}{n^d} \|\nu(y)\|_2^2 \xrightarrow{n \to \infty} \beta_c + 0$. Indeed, by Proposition 2.13, $G_{\Lambda_n}^{0\text{-avg.}}(x,y)$ converges uniformly over U^2 to $G_{\mathbb{Z}^d}(x,y)$ and so do the corresponding eigenvalues. The latter matrix is positive-definite, thus, there exists c > 0 (depending only on U) such that $0 < c \le \lambda_{\min}(G_{\Lambda_n}^{0\text{-avg.}}|_{U^2}) \le \lambda_{\max}(G_{\Lambda_n}^{0\text{-avg.}}|_{U^2}) \le \mathrm{Trace}(G_{\Lambda_n}^{0\text{-avg.}}|_{U^2}) \le 2|U|\beta_c$. Combined

with (2.13), this yields

$$\frac{1}{n^{d}} \|\nu(y)\|_{2}^{2} = \frac{1}{n^{d}} \|G_{\Lambda_{n}}^{0-\text{avg.}}|_{U^{c} \times U} G_{\Lambda_{n}}^{0-\text{avg.}}|_{U^{2}}^{-1} y\|_{2}^{2} \leq \frac{\|y\|_{2}^{2}}{c^{2} n^{d}} \sum_{x \in U^{c}, y \in U} G_{\Lambda_{n}}^{0-\text{avg.}}(x, y)^{2}
\leq \frac{\tilde{C}(U, d)}{n^{d}} \left(n \mathbb{1}_{\{d=3\}} + \log n \mathbb{1}_{\{d=4\}} + \mathbb{1}_{\{d \geq 5\}}\right) \frac{\|y\|_{2}^{2}}{c^{2}},$$
(3.4)

and fully analogously that $\frac{1}{n^d} \operatorname{Trace}(C) = \beta_c - o(1)$ with the error uniform in $y \in B$ (bounded). C is positive semi-definite as a covariance matrix, with eigenvalues $\mu_1 \geq \mu_2 \geq \ldots \geq \mu_{|U^c|} \geq 0$. We set $u_n \coloneqq |U^c|$ and observe that $\mu_{u_n} = 0$ with the unit eigenvector $q^{u_n} = \frac{1}{\sqrt{u_n}} \mathbf{1}_{U^c}$. Let $Q = (q^1, \ldots, q^{u_n})$ and P be the orthonormal matrices diagonalizing C and $G_{\Lambda_n}^{0\text{-avg}}$, respectively, i.e., $Q^T C Q = \operatorname{diag}((\mu_k)_{k=1}^{u_n})$ and $P^T G_{\Lambda_n}^{0\text{-avg}} P = \operatorname{diag}(0, (1/\eta_k)_{k=2}^{n^d})$, where $0 = \eta_1 < \eta_2 \leq \ldots \leq \eta_{n^d}$ are the eigenvalues of $-\Delta_{\Lambda_n}$. Let $(Y_i)_i$ be i.i.d. standard normal random variables. Then, if we set $h(y) \coloneqq Q^T \nu(y)$,

$$\|\gamma\|_{2}^{2} \stackrel{\text{law}}{=} \sum_{k=2}^{n^{d}} \frac{1}{\eta_{k}} Y_{k}^{2}; \qquad \|\hat{\gamma}_{U^{c}}^{y}\|_{2}^{2} \stackrel{\text{law}}{=} \sum_{k=1}^{u_{n}-1} (h_{k}(y) + \sqrt{\mu_{k}} Y_{k})^{2} + h_{u_{n}}(y)^{2}. \tag{3.5}$$

For any $\varepsilon > 0$ and all n sufficiently large, by Lemma 2.12 2., (2.15) and (3.4) we get

$$\begin{split} \mathbb{P}[|\hat{S}_{n}^{y}| > \varepsilon] &\leq \frac{\mathbb{E}[(\hat{S}_{n}^{y})^{2}]}{\varepsilon^{2}} \leq \frac{1}{n^{2d}\varepsilon^{2}} \sum_{k=1}^{u_{n}} (2\mu_{k}^{2} + 4\mu_{k}h_{k}^{2}(y)) \leq \frac{\mu_{1}\|\nu(y)^{2}\|_{2}}{n^{2d}\varepsilon^{2}} + \frac{2}{n^{2d}\varepsilon^{2}} \sum_{k=1}^{u_{n}-1} \frac{1}{\eta_{k+1}^{2}} \\ &\leq \frac{\tilde{C}_{d}}{\varepsilon^{2}} \left(\frac{1}{n^{2}} \mathbb{1}_{\{d=3\}} + \frac{\log(n)}{n^{4}} \mathbb{1}_{\{d=4\}} + \frac{1}{n^{d}} \mathbb{1}_{\{d\geq5\}} \right) \\ \mathbb{P}[|S_{n}| > \varepsilon] &\leq \frac{\mathbb{E}[S_{n}^{2}]}{\varepsilon^{2}} = \frac{1}{n^{2d}\varepsilon^{2}} \sum_{k=2}^{n^{d}} \frac{2}{\eta_{k}^{2}} \leq \frac{C_{d}}{\varepsilon^{2}} \left(\frac{1}{n^{2}} \mathbb{1}_{\{d=3\}} + \frac{\log(n)}{n^{4}} \mathbb{1}_{\{d=4\}} + \frac{1}{n^{d}} \mathbb{1}_{\{d\geq5\}} \right). \end{split}$$

for some C_d , $\tilde{C}_d > 0$ depending only on d. Since the estimates on the right-hand side are summable, Borel-Cantelli lemma implies almost sure convergence of \hat{S}_n^y and S_n towards zero.

Proof of Claim 2. The idea is to use the concentration inequality for Lipschitz functions of a standard normal vector \overline{Z} , [Mas07, Theorem 3.8, (3.10)], that is, if $f: \mathbb{R}^N \to \mathbb{R}$ is Lipschitz, x > 0, then $\mathbb{P}[f(\overline{Z}) \ge \mathbb{E}[f(\overline{Z})] + x] \le 2\frac{1}{\sqrt{2\pi}} \int_{x/\text{Lip}(f)}^{\infty} e^{-u^2/2} du \le e^{-\frac{(x/\text{Lip}(f))^2}{2}}$.

Let us define $f: \mathbb{R}^{n^d} \to \mathbb{R}$, $x \mapsto \|\sqrt{G_{\Lambda_n}^{0\text{-avg.}}}x\|_2$ and $\hat{f}^y: \mathbb{R}^{n^d} \to \mathbb{R}$, $x \mapsto \|\sqrt{C}x + \nu(y)\|_2$. Both these functions are Lipschitz with Lipschitz constants bounded by $\eta_2^{-1/2} = \Theta(n)$ and $\sqrt{\mu_1} = \Theta(n)$, respectively. We observe that for $\overline{Z} \sim \mathcal{N}(0, \mathrm{Id}_{n^d})$ and all n sufficiently large,

$$\begin{split} \mathbb{P}[S_n \geq \beta - \beta_c - \varepsilon] \leq \mathbb{P}[\|\gamma\|_2 \geq \sqrt{n^d} \sqrt{\beta - 2\varepsilon}] \\ \leq \mathbb{P}\big[f(\overline{Z}) \geq \mathbb{E}[f(\overline{Z})] + \sqrt{n^d} \sqrt{\beta - 2\varepsilon} - \underbrace{\mathbb{E}[f^2(\overline{Z})]]^{1/2}}_{\sqrt{n^d(\beta_c - o(1))}}\big] \\ \leq \exp\bigg(- \frac{1}{2}\Theta(n^{d-2}) \big(\sqrt{\beta - 2\varepsilon} - \sqrt{\beta_c - o(1)}\big)^2 \bigg) \leq e^{-Cn^{d-2}}; \end{split}$$

and fully analogously $\mathbb{P}[\hat{S}_n^y \ge \beta - \beta_c - \varepsilon] \le e^{-\hat{C}n^{d-2}}$.

Proof of Claim 3. Recall that $\|\gamma\|_2^2 \stackrel{\text{law}}{=} \sum_{k=2}^{n^d} \frac{1}{\eta_k} Y_k^2$ and $\|\hat{\gamma}^y\|_2^2 \stackrel{\text{law}}{=} \|y\|_2^2 + \sum_{k=1}^{u_n-1} (h_k(y) + \sqrt{\mu_k} Y_k)^2 + h_{u_n}(y)^2$. By Young's convolution inequality since $\|f\|_{L^1} = 1$ for a probability density function f, $\|f\|_{\hat{\gamma}^y}\|_{L^\infty} \leq \|f_{(h_l(y) + \sqrt{\mu_l} Y_l)^2 + (h_i(y) + \sqrt{\mu_l} Y_i)^2}\|_{L^\infty}$ and $\|f\|_{\gamma\|_2^2}\|_{L^\infty} \leq \|f_{Y_k^2/\eta_k + Y_j^2/\eta_j}\|_{L^\infty}$ for any $k \neq j, l \neq i$ (the choice will be specified later). Let us prove that there exists C > 0 uniform in y (here over all \mathbb{R}^U) such that $\|f_{(h_l(y) + \sqrt{\mu_l} Y_l)^2 + (h_i(y) + \sqrt{\mu_l} Y_i)^2}\|_{L^\infty} \leq C$. The remaining case can be easily concluded from this one. Let $x \in \mathbb{R}_+$ and write for simplicity $(h_l(y) + \sqrt{\mu_l} Y_l)^2 + (h_i(y) + \sqrt{\mu_l} Y_l)^2 = (a + cY)^2 + (b + dZ)^2$ (with Y, Z independent standard normal random variables). The density of $(a/c + Y)^2$ at x is given by $\frac{1}{2\sqrt{x}} (f_Y(\sqrt{x} - |a/c|) + f_Y(\sqrt{x} + |a/c|)) \leq \frac{1}{\sqrt{x}}$; therefore,

$$f_{(a+cY)^2+(b+dZ)^2}(x) = \int_0^x f_{c^2(a/c+Y)^2}(y) f_{d^2(b/d+Z)^2}(x-y) dy \le \frac{1}{cd} \int_0^1 \frac{1}{\sqrt{u(1-u)}} du = \frac{\pi}{cd}.$$

If c, d > 0 are uniformly bounded away from zero, we are done. In our setting we can for example pick $(l, i) = (\lceil u_n/2 \rceil, \lceil u_n/2 \rceil + 1) = (k, j)$, then by Lemma 2.12 2. and (2.15): $\mu_l, \mu_i, 1/\eta_k, 1/\eta_j = \Theta(1)$.

3.3 At the criticality

In this subsection we discuss the model at the critical temperature, $\beta=\beta_c$. Unlike the other two regimes we would need to treat the case d=3 slightly differently than the higher dimensions. Our proof for $d\geq 4$ combines the arguments of both non-critical regimes and makes use of Proposition 3.3 and (2.13). For d=3, we heavily rely on the fact that $\beta_c-G_{\Lambda_n}^{0\text{-avg.}}(0,0)=\Theta(1/n)>0$ (see (2.14)) and that the order of the variance of $\|\gamma\|_2^2$ is the square of the order of the largest eigenvalue of $G_{\Lambda_n}^{0\text{-avg.}}(0,0)$.

Let us start by recalling that up to the concluding argument and Claim 2, all the steps of the low-temperature regime are true for β_c too and our goal is to prove (3.3). In this case the latter amounts to showing that uniformly in y bounded,

$$R_{n} = \frac{\mathbb{E}\left[\frac{1}{\sqrt{\beta_{c} - \|\hat{\gamma}^{y}\|_{2}^{2}/n^{d}}} \mathbb{1}_{(\beta_{c} - b^{2}, \beta_{c} - a^{2})} \left(\frac{\|\hat{\gamma}^{y}\|_{2}^{2}}{n^{d}}\right)\right]}{2\mathbb{E}\left[\frac{1}{\sqrt{\beta_{c} - \|\gamma\|_{2}^{2}/n^{d}}} \mathbb{1}_{[0, \beta_{c})} \left(\frac{\|\gamma\|_{2}^{2}}{n^{d}}\right)\right]},$$

converges to 1/2 if a=0, and otherwise to zero. Note that this would imply that the conditional law of (γ_U, c) given $\|\gamma + c\mathbf{1}_{\Lambda_n}\|_2^2 = \beta_c n^d$ converges to $(\psi_U, 0)$, where ψ is the GFF on \mathbb{Z}^d . Note that if a>0, by Fatou's lemma applied to the denominator, dominated convergence theorem applied to the numerator and Claim 1 in the previous section, $\limsup_n R_n = 0$. This also yields that without loss of generality we can assume that $b^2 = \beta_c$. So, hereinafter let $a=0, b^2 = \beta_c$.

Let

$$X_{n} := \frac{\|\gamma\|_{2}^{2} - \mathbb{E}[\|\gamma\|_{2}^{2}]}{\sqrt{\text{Var}[\|\gamma\|_{2}^{2}]}}, \qquad \hat{X}_{n} := \frac{\|\hat{\gamma}^{y}\|_{2}^{2} - \mathbb{E}[\|\hat{\gamma}^{y}\|_{2}^{2}]}{\sqrt{\text{Var}[\|\hat{\gamma}^{y}\|_{2}^{2}]}};$$

$$C_{n} := \frac{\beta_{c}n^{d} - \mathbb{E}[\|\gamma\|_{2}^{2}]}{\sqrt{\text{Var}[\|\gamma\|_{2}^{2}]}}, \qquad \hat{C}_{n} := \frac{\beta_{c}n^{d} - \mathbb{E}[\|\hat{\gamma}^{y}\|_{2}^{2}]}{\sqrt{\text{Var}[\|\gamma^{y}\|_{2}^{2}]}}.$$

Then, using the fact (proven below) that

$$\operatorname{Var}[\|\hat{\gamma}^y\|_2^2] = \operatorname{Var}[\|\gamma\|_2^2] + \mathcal{O}\left(n^2\left(n\mathbb{1}_{\{d=3\}} + \log n\mathbb{1}_{\{d=4\}} + \mathbb{1}_{\{d\geq 5\}}\right)\right)$$

$$= (1 + o(1))\operatorname{Var}[\|\gamma\|_2^2] = \Theta\left(n^4\mathbb{1}_{\{d=3\}} + n^d\log n\mathbb{1}_{\{d=4\}} + n^d\mathbb{1}_{\{d\geq 5\}}\right),$$
(3.6)

we can rewrite R_n as

$$R_n = \frac{1 + o(1)}{2} \frac{\mathbb{E}\left[\frac{1}{\sqrt{\hat{C}_n - \hat{X}_n}} \mathbb{1}_{\{\hat{X}_n < \hat{C}_n\}}\right]}{\mathbb{E}\left[\frac{1}{\sqrt{C_n - X_n}} \mathbb{1}_{\{X_n < C_n\}}\right]}.$$

Note that since $\mathbb{E}[\|\hat{\gamma}^y\|_2^2] = G_{\Lambda_n}^{0\text{-avg.}}(0,0)u_n + o(n^{d/2})$, by (2.13) evaluated at y=0 and (2.14), we furthermore know that C_n and \hat{C}_n converge to the same finite limit C (depending on d): for $d \geq 4$, C=0; for d=3, C>0. As for \hat{X}_n and X_n , we have the following.

<u>Claim A:</u> X_n and \hat{X}_n converge in law to the same non-trivial (by symmetry, equivalent to finite non-zero) real random variable X. For $d \geq 4$, X is a standard normal random variable.

<u>Claim B:</u> The densities of \hat{X}_n and X_n are uniformly bounded.

Using these two observations and applying Portmanteau theorem to the upper/lower semicontinuous functions $x\mapsto 1/\sqrt{x}\mathbb{1}_{\{x\geq\varepsilon\}}$ and $x\mapsto 1/\sqrt{x}\mathbb{1}_{\{x>0\}}$, respectively, for $0<\varepsilon$ arbitrary small such that $\varepsilon< C(d=3)/2$, we obtain that

$$\lim_{n \to \infty} \mathbb{E}\left[\frac{1}{\sqrt{\hat{C}_n - \hat{X}_n}} \mathbb{1}_{\{\hat{X}_n < \hat{C}_n\}}\right] \le 2M\sqrt{\varepsilon} + \mathbb{E}\left[\frac{1}{\sqrt{C - X}} \mathbb{1}_{\{X < C - \varepsilon\}}\right];$$

$$\lim_{n \to \infty} \inf \mathbb{E}\left[\frac{1}{\sqrt{C_n - X_n}} \mathbb{1}_{\{X_n < C_n\}}\right] \ge \mathbb{E}\left[\frac{1}{\sqrt{C - X}} \mathbb{1}_{\{X < C\}}\right] \ge \mathbb{E}\left[\frac{1}{\sqrt{C - X}} \mathbb{1}_{\{X < C - \varepsilon\}}\right].$$

Here M>0 is the uniform bound on the densities of \hat{X}_n and X_n . Since X is non-trivial and of mean zero, the probability of it being non-positive is strictly positive. Hence, if d=3, since $C-\varepsilon>0$ in this case, there exists an absolute (independent of ε) constant p>0 such that $\mathbb{E}\left[\frac{1}{\sqrt{C-X}}\mathbb{1}_{\{X< C-\varepsilon\}}\right]>p$. Existence of such a constant for $d\geq 4$ even though $C(d\geq 4)=0$ is clear since X is a standard Gaussian. Thus, $R_n\leq \frac{1+o(1)}{2}\left(1+\frac{2M\sqrt{\varepsilon}}{p}\right)$. By taking $\varepsilon>0$ much smaller than p, we conclude that R_n is less or equal to 1/2 in the limit. By reversing the roles of the numerator and denominator, we derive that R_n is bounded from below by 1/2.

We proceed to discuss the remaining proofs of (3.6) and the claims. Our arguments for the latter will significantly differ for $d \ge 4$ and d = 3.

Proof of (3.6). It is an easy observation that $\text{Var}[\|\hat{\gamma}^y\|_2^2] = 2\sum_{k=1}^{u_n-1}(\mu_k^2 + 2h_k(y)^2\mu_k)$. By Lemma 2.12 2., (2.15) and (3.4), $\mu_1\|h(y)\|_2^2 \ll \Theta(n^4\mathbb{1}_{\{d=3\}} + n^d\log n\mathbb{1}_{\{d=4\}} + n^d\mathbb{1}_{\{d\geq 5\}}) = 2\sum_{k=1}^{u_n-1}\mu_k^2$ uniformly over y in a fixed bounded set, which further combined with (2.13) implies,

$$\frac{1}{2} \operatorname{Var}[\|\hat{\boldsymbol{\gamma}}^y\|_2^2] = \sum_{k=1}^{u_n - 1} \mu_k^2 + \mathcal{O}(n^2 (n\mathbb{1}_{\{d=3\}} + \log n\mathbb{1}_{\{d=4\}} + \mathbb{1}_{\{d \ge 5\}}));$$
$$\sum_{k=1}^{u_n - 1} \mu_k^2 = \operatorname{Trace}(C^2) = \frac{1}{2} \operatorname{Var}[\|\boldsymbol{\gamma}\|_2^2] + A_1 + A_2 + A_3,$$

where

$$A_{1} := \sum_{x,y \in U^{c}} G_{\Lambda_{n}}^{0-\text{avg.}}(x,y)^{2} - \frac{1}{2} \text{Var}[\|\boldsymbol{\gamma}\|_{2}^{2}] = -2 \sum_{\substack{x \in U^{c}, \\ z \in U}} G_{\Lambda_{n}}^{0-\text{avg.}}(x,z)^{2}$$
$$= \mathcal{O}((n\mathbb{1}_{\{d=3\}} + \log n\mathbb{1}_{\{d=4\}} + \mathbb{1}_{\{d\geq 5\}}));$$

$$\begin{split} A_2 &\coloneqq \sum_{\substack{z,z',\\w,w' \in U\\x,y \in U^c}} G_{\Lambda_n}^{0\text{-avg.}}(x,z) G_{\Lambda_n}^{0\text{-avg.}}|_{U^2}^{-1}(z,z') G_{\Lambda_n}^{0\text{-avg.}}(z',y) G_{\Lambda_n}^{0\text{-avg.}}(y,w) G_{\Lambda_n}^{0\text{-avg.}}|_{U^2}^{-1}(w,w') G_{\Lambda_n}^{0\text{-avg.}}(w',x) \\ &= \mathcal{O}\Big(\big(\int_1^n r^{4-2d} r^{d-1} \mathrm{d} r \big)^2 \Big) = \mathcal{O}\big(n^2 \mathbbm{1}_{\{d=3\}} + \log^2 n \mathbbm{1}_{\{d=4\}} + \mathbbm{1}_{\{d \geq 5\}} \big); \\ A_3 &\coloneqq -2 \sum_{\substack{z,z' \in U\\x,y \in U^c}} G_{\Lambda_n}^{0\text{-avg.}}|_{U^2}^{-1}(z,z') G_{\Lambda_n}^{0\text{-avg.}}(x,y) G_{\Lambda_n}^{0\text{-avg.}}(y,z) G_{\Lambda_n}^{0\text{-avg.}}(z',x) \\ &= \mathcal{O}\Big(\sum_{\substack{x \neq y \in \hat{\Lambda}_n \backslash \{0\}}} |x|^{2-d} |y|^{2-d} \mathrm{d}_{\mathbb{T}_n^d}(x,y)^{2-d} \Big) \\ &= \mathcal{O}\Big(2 \sum_{\substack{x \in \hat{\Lambda}_n \backslash \{0\}}} |x|^{4-2d} \sum_{\substack{y \in \hat{\Lambda}_n \backslash \{0,x\} \\ |y| \geq |x|}} \mathrm{d}_{\mathbb{T}_n^d}(x,y)^{2-d} \Big) \\ &= \mathcal{O}\Big(n^2 \sum_{\substack{x \in \hat{\Lambda}_n \backslash \{0\}}} |x|^{4-2d} \Big) = \mathcal{O}\Big(n^2 \Big(n \mathbbm{1}_{\{d=3\}} + \log n \mathbbm{1}_{\{d=4\}} + \mathbbm{1}_{\{d \geq 5\}} \Big) \Big). \end{split}$$

Altogether, as desired

$$\operatorname{Var}[\|\hat{\gamma}^y\|_2^2] = \operatorname{Var}[\|\gamma\|_2^2] + \mathcal{O}\left(n^2\left(n\mathbb{1}_{\{d=3\}} + \log n\mathbb{1}_{\{d=4\}} + \mathbb{1}_{\{d\geq 5\}}\right)\right)$$
$$= (1 + o(1))\operatorname{Var}[\|\gamma\|_2^2].$$

Proof of Claims A & B for $d \ge 4$: The key observation in this case is the following.

<u>Claim C:</u> The densities of X_n and \hat{X}_n^a converge uniformly towards the density function of a standard normal random variable.

The proof of this result is based on Proposition 3.3 (note that for d=3 even Lindeberg's condition fails) and the eigenvalue representation (3.5). But before we present it let us explain how it yields Claims A and B. Note that by the eigenvalue decomposition we can write \hat{X}_n as a sum of two independent random variables $P_n + T_n$, where P_n is the "good" part treated in Claim C and T_n is the sum of the remaining |U| terms. Then, by Claim C, P_n converges in law to a standard normal random variable and its density is uniformly bounded. Note that $T_n = \frac{\sum_{u_n - |U|}^{u_n} \mu_k(Y_k^2 - 1) + 2h_k(y)\sqrt{\mu_k}Y_k}{\sqrt{\text{Var}[\|\hat{\gamma}^y\|_2^2]}}$. Since these μ_k 's are uniformly bounded and $\|h(y)\|_2$ by (3.4) is growing slower than any polynomial, one can easily check that T_n possesses a density function and converges almost surely to zero. This implies that $T_n + P_n$ converges to a standard Gaussian variable (Claim A) and that the density of \hat{X}_n is uniformly bounded as a convolution of a bounded function and an integrable non-negative (with unit L^1 -norm) function (Claim B).

Proof of Claim A. We check that the required assumptions of Proposition 3.3 are satisfied by $\mathbf{Y}_n := ((h_k(y) + \sqrt{\mu_k}Y_k)^2 - \mu_k - h_k(y)^2)_{k=1}^{u_n - |U|-1}$ contributing to \hat{X}_n , and thus, also by $(\frac{1}{\eta_k}(Y_k^2 - 1))_{k=2}^{n^d}$ contributing to X_n , since the latter can be recovered from the former by setting $h(y) \equiv 0$ and $U = \emptyset$. More precisely, we consider independent families $(\mathbf{Y}_n)_n$ that

^aup to removing the last |U| terms in the eigenvalue decomposition (3.5) of \hat{X}_n containing μ_{u_n-q} for $1 \leq q \leq |U|$

form a triangular array. For better readability, we do not use double-indexing for the elements of each \mathbf{Y}_n .

The first assumption of the proposition can be directly deduced from the corresponding verification in the high-temperature regime since the eigenvalues $(\mu_k)_{k=1}^{u_n-|U|-1}$ of C and $(1/\eta_k)_{k=2}^{n^d}$ of $G_{\Lambda_n}^{0\text{-avg.}}$ are uniformly bounded away from zero. The next assumption (up to reordering which will be done for the verification of the fourth criterion) is clear since $\sigma_k^2 = 2\mu_k^2 + 4h_k(y)^2\mu_k < \infty$. From now on, index k would refer to the original order and \tilde{k} to the one according to decreasing variances. Recall that Lindeberg's condition follows from Lyapunov's condition, which in our setting for $\delta = 2$ appears as

$$\frac{1}{s_n^4} \sum_{k=1}^{u_n - |U| - 1} \mathbb{E} \left[\left((h_k(y) + \sqrt{\mu_k} Y_k)^2 - \mu_k - h_k(y)^2 \right)^4 \right] \xrightarrow{n \to \infty} 0,$$

where $s_n^2 = \sum_{k=1}^{u_n - |U| - 1} (2\mu_k^2 + 4h_k(y)^2 \mu_k) = 2(1 + o(1)) \sum_{k=1}^{u_n - |U| - 1} \mu_k^2$ by (3.4). We observe that this convergence indeed takes place since $\sum_{k=1}^{u_n - |U| - 1} \mathbb{E}\left[((h_k(y) + \sqrt{\mu_k}Y_k)^2 - \mu_k - h_k(y)^2)^4\right] \le C \sum_{k=1}^{u_n - |U| - 1} (\mu_k^4 + \mu_k^2 h_k(y)^4 + \mu_k^3 h_k(y)^2) \le C (\sum_{k=1}^{u_n - |U| - 1} \mu_k^4 + \mu_1^2 \|h(y)\|_2^4 + \mu_1^3 \|h\|_2^2)$, and by Lemma 2.12 2., (2.15) and (3.4), for $d \ge 4$, this is of order $\mathcal{O}(n^8 \mathbb{1}_{\{d \le 7\}} + n^d \log n \mathbb{1}_{\{d = 8\}} + n^d \mathbb{1}_{\{d \ge 9\}})$, but $s_n^4 = \Theta(n^8 (\log n)^2 \mathbb{1}_{\{d = 4\}} + n^{2d} \mathbb{1}_{\{d \ge 5\}})$. For the fourth criterion, consider the set $J = \{\sigma_k^2 : k \le u_n - |U| - 1$ such that $|h_k(y)| > 1/\log n\}$ – there are at most $(\log n) \|h^y\|_2^2$ such indices k. Notice that since $(\mu_k)_{k \le u_n - |U| - 1}$ are uniformly bounded away from zero and $\sigma_{k,u_n|U|-1}^2 = 2\mu_k^2 + 4\mu_k h_k(y)^2$, the order of the variances on the complement of J will be determined by the order of μ_k 's. Furthermore, elements of J after reordering can only be sent forward or remain at their place. Hence, for any $L \ge 1$,

$$\{\sigma_k^2: k \geq L\} \setminus J \quad \subset \ \{\sigma_{\tilde{k}}^2: \tilde{k} \geq L\} \setminus J \ \subset \quad \{\sigma_k^2: k \geq L - 2|J|\} \setminus J.$$

Let $\varepsilon \in (0,1)$ be very small but fixed. We choose required constants as follows:

- for $d \geq 5$, let $l^* = l_* := \inf\{l \geq \varepsilon n^d : l 2|J| \notin J, l \in \mathbb{Z}\}$, $K = K(\varepsilon)$ a large constant depending only on ε :
- for d=4, let $l^* := \inf\{l \ge \varepsilon n^d : l-2|J| \notin J, l \in \mathbb{Z}\}, l_* := \inf\{l \ge n^{d-\varepsilon} : l-2|J| \notin J, l \in \mathbb{Z}\}, K(n) = n^{\varepsilon}.$

Note that l^*, l_* can at most be equal to the reference value (either εn^d or $n^{d-\varepsilon}$) plus $|J| = o(n^{\delta})$ for any $\delta > 0$. Moreover, for $L \in \{l_*, l^*\}$,

$$\{\sigma_k^2: k \ge L\} \setminus J \subset \{\sigma_{\tilde{k}}^2: \tilde{k} \ge L\} \subset \{\sigma_k^2: k \ge L - 2|J|\}.$$

This implies that the l^* -th element of the reordered sequence $(\sigma_{\tilde{k}}^2)_{\tilde{k}}$ is smaller or equal to $\sigma_{l^*-2|J|}^2$ (of the original order). Let us verify the assumptions for $d \geq 5$. By Lemma 2.12 2. and (2.15), for $u_n - |U| - 1 \geq k \geq l_* - 2|J|, k \notin J$, $0 < c_* \leq \mu_k = \Theta(n^2/k^{2/d}) = \Theta(1) \leq \mu_{l_*-2|J|} \leq c^* < \infty$, $\sigma_k^2 = 2\mu_k^2 + 4\mu_k h_k(y)^2 = \Theta(1)$ uniformly. Thus, for $Z_k = (h_k(y) + \sqrt{\mu_k}Y_k)^2 - \mu_k - h_k(y)^2$, $Z \sim \mathcal{N}(0,1)$

$$\begin{split} \frac{\sum_{\tilde{k} \geq l_*}^{u_n - |U| - 1} \mathbb{E}[Z_{\tilde{k}}^2 \mathbb{1}_{\{|Z_{\tilde{k}}| > K\}}]}{\sum_{\tilde{k} \geq l_*}^{u_n - |U| - 1} \sigma_{\tilde{k}}^2} \leq \frac{\sum_{k \geq l_* - 2|J|, k \notin J}^{u_n - |U| - 1} \mathbb{E}[Z_k^2 \mathbb{1}_{\{|Z_k| > K\}}] + |J| \sigma_{l_* - 2|J|}^2}{\sum_{k \geq l_*, k \notin J}^{u_n - |U| - 1} \sigma_{\tilde{k}}^2} \\ \leq \frac{\sum_{k \geq l_*, k \notin J}^{u_n - |U| - 1} \mathbb{E}[Z_k^2 \mathbb{1}_{\{|Z_k| > K\}}] + 3|J| \sigma_{l_* - 2|J|}^2}{\sum_{k \geq l_*, k \notin J}^{u_n - |U| - 1} \sigma_{\tilde{k}}^2} \end{split}$$

$$\leq \frac{\mathbb{E}\left[(Z^2-1)^2\mathbb{1}_{\left\{\mu_{l_*-2|J|}Z^2+\frac{2\sqrt{\mu_{l_*-2|J|}}}{\log n}|Z|>K\right\}}\right]\sum_{k\geq l_*,k\notin J}^{u_n-|U|-1}\mu_k^2}{2\sum_{k\geq l_*,k\notin J}^{u_n-|U|-1}\mu_k^2} \\ + \frac{4\mu_1\|h(y)\|_2^2+\frac{1}{\log n}4\sqrt{3}\sum_{k\geq l_*,k\notin J}^{u_n-|U|-1}\mu_k^{3/2}+o(n)}{2\sum_{k\geq l_*,k\notin J}^{u_n-|U|-1}\mu_k^2} \\ \leq \frac{1}{2}\mathbb{E}\left[(Z^2-1)^2\mathbb{1}_{\left\{Z^2+|Z|>K/(2c^*)\right\}}\right]+o(n^{3-d})+\Theta\left(\frac{1}{\log n}\right)$$

can be made arbitrarily small by choosing $K > c^*$ sufficiently large. Furthermore,

$$\frac{\sum_{\tilde{k} \ge l_*}^{u_n - |U| - 1} \sigma_{\tilde{k}}^2}{s_n^2} \ge \frac{\sum_{k \ge l_*, k \notin J}^{u_n - |U| - 1} \sigma_{k}^2}{s_n^2} \ge \frac{\sum_{k = \varepsilon n^d + n}^{u_n - |U| - 1} 2\mu_{k}^2}{\Theta(n^d)}$$

$$\ge \frac{\Theta\left(n^4 \sum_{k = \varepsilon n^d + n^{d/2}}^{u_n - |U| - 1} k^{-\frac{4}{d}}\right)}{\Theta(n^d)} = (1 - \varepsilon^{\frac{d - 4}{d}})\Theta(1);$$

$$\frac{u_n - |U| - 1 - l^*}{\sigma_{l^* \text{ reorder}}^2 \vee K^2} \ge \frac{u_n - |U| - 1 - l^*}{\sigma_{l^* - 2|J|}^2 \vee K^2} = \Theta(n^d) \gg \log(n^d).$$

This completes the verification of all the assumptions for $d \ge 5$. Let d = 4, Fully analogously to the above, by Lemma 2.12 2. and (2.15),

$$\begin{split} \frac{\sum_{\tilde{k} \geq l_*}^{u_n - |U| - 1} \sigma_{\tilde{k}}^2}{s_n^2} &\geq \frac{\sum_{k \geq l_*, k \notin J}^{u_n - |U| - 1} \sigma_k^2}{s_n^2} \geq \frac{\Theta\left(\sum_{k = 2n^{d - \varepsilon}}^{u_n - |U| - 1} n^4 / k\right)}{\Theta(n^4 \log n)} = \frac{\Theta\left(\log\left(\frac{u_n}{n^{d - \varepsilon}}\right)\right)}{\Theta(\log n)} = \varepsilon\Theta(1); \\ &\frac{u_n - |U| - 1 - l^*}{\sigma_{l^*, \text{reorder.}}^2 \vee K(n)^2} \geq \frac{u_n - |U| - 1 - l^*}{\sigma_{l^* - 2|J|}^2 \vee K(n)^2} = \Theta\left(\frac{n^d}{(n^4 / (\varepsilon n^d)^{4/d}) \vee n^{2\varepsilon}}\right) \\ &= \Theta(n^{d - 2\varepsilon}) \gg \log(n^4 \log n); \\ &\frac{\sum_{\tilde{k} \geq l_*}^{u_n - |U| - 1} \mathbb{E}[Z_{\tilde{k}}^2 \mathbb{1}_{\{|Z_{\tilde{k}}| > K(n)\}}]}{\sum_{\tilde{k} \geq l_*}^{u_n - |U| - 1} \sigma_{\tilde{k}}^2} \leq \frac{\sum_{k \geq l_*, k \notin J}^{u_n - |U| - 1} \mathbb{E}[Z_k^2 \mathbb{1}_{\{|Z_k| > K(n)\}}] + 3|J|\sigma_{l_* - 2|J|}^2}{\sum_{k \geq l_*, k \notin J}^{u_n - |U| - 1} \sigma_{\tilde{k}}^2} \leq \frac{1}{2} \mathbb{E}\left[(Z^2 - 1)^2 \mathbb{1}_{\left\{Z^2 + |Z| > \frac{K(n)}{\log n} + |J|\sigma_{l_* - 2|J|}^2\right\}}\right] \\ &+ \frac{\Theta\left(n^2 ||h(y)||_2^2 + \frac{\sum_{k \geq l_*}^{u_n} \mu_k^{3/2}}{\log n} + |J|\sigma_{l_* - 2|J|}^2\right)}{\varepsilon\Theta(n^4 \log n)} \\ &= \frac{1}{2} \mathbb{E}\left[(Z^2 - 1)^2 \mathbb{1}_{\left\{Z^2 + |Z| > \frac{n^\varepsilon}{\Theta(n^\varepsilon/2)}\right\}}\right] + o(n^{-1}) + \Theta\left(\frac{1}{\log^2 n}\right). \end{split}$$

The latter is arbitrarily small.

Proof of Claims A & B for d = 3:

Proof of Claim B. The statement can be verified fully analogously to Claim 3 in the low-temperature regime (see Section 3.2) using eigenvalue representation (3.5) once we notice that for $\mu \in \{\mu_1, \mu_2, \eta_2^{-1}, \eta_3^{-1}\}$ and $v \in \{\operatorname{Var}[\|\hat{\gamma}^y\|_2^2], \operatorname{Var}[\|\boldsymbol{\gamma}\|_2^2]\}$: $\mu/\sqrt{v} = \Theta(1)$ uniformly in n. The latter is due to the above observations that $\mu, \eta^{-1} = \Theta(n^2)$ and the variances are of order $\Theta(n^4)$.

Proof of Claim A. Recall that if a sequence of random variables $(X_n)_n$ possesses moment generating functions (MGF) (M_n) that are finite on a common open interval around zero,

(-a,a) for a>0, i.e., $M_n(u)<\infty$ for all $n\in\mathbb{N}, u\in(-a,a)$, and there exists 0< b< a and a finite-valued function M defined on [-b,b] such that for any $u\in[-b,b]$, $M(b)=\lim_n M_n(b)$, then there exists a random variable X, which is the distributional limit of $(X_n)_n$, its MGF is finite on [-b,b] and coincides with M on this interval (cf. [Cur42, Theorem 3]). Using this fact it suffices to prove that the MGFs $(M_n)_n$ and $(\hat{M}_n)_n$ of $(X_n)_n$ and (\hat{X}_n) , respectively, converge pointwise for all $|x| \le b < \min\left(\frac{1}{2\sqrt{2}}, \frac{1}{4}\eta_2\sqrt{\mathrm{Var}[\|\gamma\|_2^2]}\right)$ (for all n sufficiently large) to some finite (on [-b,b]) function M, unequal to a constant function 1 (that is the limiting law is not that of a constant 0). Note that by Lemma 2.12 2., $\eta_2^2 \mathrm{Var}[\|\gamma\|_2^2] = 2\sum_{k=2}^{n^d} \eta_2^2/\eta_k^2 \sim C\sum_k k^{-4/3} \xrightarrow{n\to\infty} c > 0$, and therefore, we indeed can pick b>0 with the desired properties. Let $t\in[-b,b]$, then utilizing the eigenvalue representation (3.5) of \hat{X}_n and recalling that $\mathrm{Var}[\|\hat{\gamma}^y\|_2^2] = 2(1+o(n^{\varepsilon-1}))\sum_{k=1}^{u_n-1} \mu_k^2$ we obtain that

$$\hat{M}_{n}(t) = \exp\left(-t\sum_{k=1}^{u_{n}-1} \frac{\mu_{k}}{\sqrt{\operatorname{Var}[\|\hat{\gamma}^{y}\|_{2}^{2}]}} - \frac{1}{2}\sum_{k=1}^{u_{n}-1} \log\left(1 - \frac{2t\mu_{k}}{\sqrt{\operatorname{Var}[\|\hat{\gamma}^{y}\|_{2}^{2}]}}\right)\right)$$

$$\times \exp\left(t\sum_{k=1}^{u_{n}-1} \frac{h_{k}(y)^{2}}{\sqrt{\operatorname{Var}[\|\hat{\gamma}^{y}\|_{2}^{2}]}} \left(\left(1 - \frac{2t\mu_{k}}{\sqrt{\operatorname{Var}[\|\hat{\gamma}^{y}\|_{2}^{2}]}}\right)^{-1} - 1\right)\right)$$

$$|\cdot| \in \left[0.3b\|h(y)\|_{2}^{2}/\sqrt{\operatorname{Var}[\|\hat{\gamma}^{y}\|_{2}^{2}]}\right)^{\left(3.4\right)} \subset (0,o(n^{\varepsilon-1}))$$

$$= (1 + o(n^{\varepsilon-1})) \exp\left(\sum_{l=2}^{\infty} \frac{1}{2l} \left(\frac{\sqrt{2}t}{1 + o(n^{\varepsilon-1})}\right)^{l} \sum_{k=1}^{u_{n}-1} \left(\frac{\mu_{k}^{2}}{\sum_{p=1}^{u_{n}-1} \mu_{p}^{2}}\right)^{l/2}\right)$$

$$=: f_{n}(l)$$

for any $\varepsilon \in (0,1)$. Here we additionally used that the MGF of $(Z+\mu)^2$ with $Z \sim \mathcal{N}(0,1)$ is given by $\exp\left(\frac{\mu^2 x}{1-2x} - \frac{1}{2}\log(1-2x)\right)$ for all x < 1/2; and that $0 < \frac{2|t|\mu_k}{\sqrt{\text{Var}[\|\hat{\gamma}^y\|_2^2]}} \le \frac{(1+o(1))2b\mu_1}{\sqrt{\text{Var}[\|\hat{\gamma}\|^2]}} < \frac{1+o(1)}{2}\eta_2\mu_1 \le \frac{1+o(1)}{2} < 3/4$ by (2.15) and Lemma 2.12 2. for all n sufficiently large. Clearly, as $\mu_k^2/\sum_{p=1}^{u_n-1}\mu_p^2 < 1$ for all k and for l=2 the last sum in the exponential is just one, it is also bounded by one for all l>2. Therefore, uniformly in $t\in[-b,b], |f_n(l)|\le (3/4)^l/(2l)$, and the latter is summable. Therefore, by dominated convergence theorem,

$$\lim_{n \to \infty} \hat{M}_n(t) = \exp\left(\sum_{l=2}^{\infty} \frac{1}{2l} (\sqrt{2}t)^l \lim_{n \to \infty} \sum_{k=1}^{u_n - 1} \left(\frac{\mu_k^2}{\sum_{p=1}^{u_n - 1} \mu_p^2}\right)^{l/2}\right).$$

And fully analogously,

$$\lim_{n \to \infty} M_n(t) = \exp\left(\sum_{l=2}^{\infty} \frac{1}{2l} (\sqrt{2}t)^l \lim_{n \to \infty} \sum_{k=2}^{n^d} \left(\frac{1/\eta_k^2}{\sum_{p=2}^{n^d} 1/\eta_p^2}\right)^{l/2}\right).$$

It remains to show that for each fixed $l \geq 3$, $\lim_n \sum_k \left(\frac{\mu_k^2}{\sum_p \mu_p^2}\right)^{l/2} = \lim_n \sum_k \left(\frac{1/\eta_k^2}{\sum_p 1/\eta_p^2}\right)^{l/2}$ and that these limits are indeed well-defined. Note that for l=2, both sides of the latter expression are equal to one, which already suffices to conclude that the limiting distribution is non-trivial. Let us start by showing that $\lim_n \sum_k \left(\frac{1/\eta_k^2}{\sum_p 1/\eta_p^2}\right)^{l/2}$ is well-defined. First observe that

$$\sum_{k=2}^{n^d} \left(\frac{1/\eta_k^2}{\sum_{p=2}^{n^d} 1/\eta_p^2} \right)^{l/2} = \sum_{x \in [0,n) \cap \mathbb{Z}^d \backslash \{0\}} \left(\frac{1/\eta_x^2}{\sum_{y \in [0,n) \cap \mathbb{Z}^d \backslash \{0\}} 1/\eta_y^2} \right)^{l/2} = \sum_{x \in \hat{\Lambda}_n \backslash \{0\}} \left(\frac{1/\eta_x^2}{\sum_{y \in \hat{\Lambda}_n \backslash \{0\}} 1/\eta_y^2} \right)^{l/2},$$

where $\eta_x = 2\sum_{i=1}^3 (1 - \cos(2\pi x_i/n))$ and $\hat{\Lambda}_n = [-n/2, n/2)^3 \cap \mathbb{Z}^3$. Let f(n) be an arbitrary function slowly growing to infinity such that f(n) = o(n), e.g., $f(n) = \log(n)$, then using explicit form of η_x and Taylor expanding it, one can easily get that

$$\sum_{y \in \hat{\Lambda}_n \setminus \{0\}} \frac{1}{n^4 \eta_y^2} = \left(1 + \mathcal{O}\left(f(n)^{-1}\right)\right) \sum_{\substack{y \in \hat{\Lambda}_n \setminus \{0\} \\ |y| \le f(n)}} \frac{1}{(2\pi)^4 |y|^4} + o(1) \stackrel{n \to \infty}{\sim} \sum_{y \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{(2\pi)^4 |y|^4}.$$

And fully analogously,

$$\sum_{x \in \hat{\Lambda}_n \setminus \{0\}} \left(\frac{1}{n^2 \eta_x} \right)^l \overset{n \to \infty}{\sim} \sum_{x \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{(2\pi)^{2l} |x|^{2l}}.$$

Altogether,

$$\lim_{n \to \infty} \sum_{k} \left(\frac{1/\eta_k^2}{\sum_{p} 1/\eta_p^2} \right)^{l/2} = \sum_{x \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|x|^{2l}} / \left(\sum_{y \in \mathbb{Z}^3 \setminus \{0\}} \frac{1}{|y|^4} \right)^{l/2} < \infty.$$

It remains to prove that $\sum_k \left(\frac{\mu_k^2}{\sum_p \mu_p^2}\right)^{l/2} / \sum_k \left(\frac{1/\eta_k^2}{\sum_p 1/\eta_p^2}\right)^{l/2} \to 1$, which can further be reduced to showing $\sum_k \mu_k^l / \sum_k 1/\eta_k^l \to 1$ as n tends to infinity. The latter simplification is due to the proof of (3.6), which gives us that $\sum_p \mu_p^2 = \frac{1}{2} \mathrm{Var}[\|\gamma\|_2^2] + o(1) = \sum_p 1/\eta_p^2 + o(1)$. By Lemma 2.12 2. and the proof of (2.15) (and since the contribution of the smallest |U| eigenvalues μ_k to the sum is irrelevant), we furthermore know that $\sum_k \mu_k^l / \sum_k 1/\eta_k^l \le 1$ and for $k \ll n^d$, $\mu_k \ge \max(\lambda_k - \alpha_1, \lambda_{k+|U|})$, where λ_k is the k-th largest eigenvalue of $G' := G_{\Lambda_n}^{0\text{-avg.}}|_{U^c \times U^c}$ and α_1 is the largest eigenvalue of $A := G_{\Lambda_n}^{0\text{-avg.}}|_{U^c \times U}G_{\Lambda_n}^{0\text{-avg.}}|_{U \times U}G_{\Lambda_n}^{0\text{-avg.}}|_{U \times U^c}$. Notice that $0 < \alpha_1 \le \mathrm{Trace}(A) \le C(U) \sum_{x \in U^c, y \in U} G_{\Lambda_n}^{0\text{-avg.}}(x,y)^2 = \mathcal{O}((\log n)^{3d}n)$ by (2.12). Hence, since $\lambda_k \ge 1/\eta_{k+|U|+1} = \Theta(n^2/k^{2/3})$, for $k \le \log^3 n$: $\mu_k = \lambda_k (1-o(1))$ with o(1) uniform in this range of k. Combined with an argument analogous to our proof of the existence of the limit, this implies that

$$\frac{\sum_{k=1}^{u_n-1}(\mu_k/n^2)^l}{\sum_{k=2}^{n^d} 1/(n^2\eta_k)^l} \sim \frac{\sum_{k=1}^{\log^3 n}(\mu_k/n^2)^l}{\sum_{k=2}^{\log^3 n} 1/(n^2\eta_k)^l} \sim \frac{\sum_{k=1}^{\log^3 n}(\lambda_k/n^2)^l}{\sum_{k=2}^{\log^3 n} 1/(n^2\eta_k)^l} \sim \frac{\sum_{k=1}^{u_n}(\lambda_k/n^2)^l}{\sum_{k=2}^{n^d} 1/(n^2\eta_k)^l} = \frac{\operatorname{Trace}((G')^l)}{\operatorname{Trace}((G^{0\text{-avg.}})^l)}$$

as $n \to \infty$ and that $\operatorname{Trace}((G')^l) = \Theta(n^{2l})$. But now due to spacial homogeneity of zero-average Green's function (and periodicity as we are on the torus),

$$\operatorname{Trace}((G_{\Lambda_{n}}^{0-\operatorname{avg.}})^{l}) - \operatorname{Trace}((G')^{l}) = \sum_{k=1}^{l} \sum_{\substack{I \subset \{1, \dots, l\} \\ |I| = k}} \sum_{\substack{x_{i} \in U : i \in I; \\ x_{j} \in U^{c} : j \notin I}} \prod_{i=1}^{l} G_{\Lambda_{n}}^{0-\operatorname{avg.}}(x_{k}, x_{k+1}) \\
\leq \sum_{k=1}^{l} \binom{l}{k} \sum_{u \in U} \sum_{\substack{x_{i} \in \Lambda_{n} : i \neq 1}} G_{\Lambda_{n}}^{0-\operatorname{avg.}}(u, x_{2}) \left(\prod_{i=1}^{l-1} G_{\Lambda_{n}}^{0-\operatorname{avg.}}(x_{k}, x_{k+1}) \right) G_{\Lambda_{n}}^{0-\operatorname{avg.}}(x_{l}, u) \\
= \sum_{k=1}^{l} \binom{l}{k} \frac{|U|}{n^{3}} \operatorname{Trace}((G_{\Lambda_{n}}^{0-\operatorname{avg.}})^{l}) = (2^{l} - 1) \frac{|U|}{n^{3}} \operatorname{Trace}((G_{\Lambda_{n}}^{0-\operatorname{avg.}})^{l})$$

with the convention that $x_{l+1} = x_1$. This immediately yields that the ratio of interest $\operatorname{Trace}((G')^l)/\operatorname{Trace}((G_{\Lambda_n}^{0-\operatorname{avg.}})^l)$ converges to 1 as n tends to infinity, and herewith completes our proof.

3.4 Local correlation functions of the Spherical model

The key objective of this subsection is to prove Corollary 3.2, that is to show that

$$\nu_{\Lambda_n,\beta} \left[\prod_{x \in U} \theta_x^{i_x} \right] \xrightarrow{n \to \infty} \mathbb{E} \left[\prod_{x \in U} \alpha_x^{i_x} \right], \tag{3.7}$$

where $\sqrt{\beta}\alpha$ in correspondence with Theorem 3.1 is either an m^2 -massive GFF on \mathbb{Z}^d if $\beta < \beta_c$, a GFF on \mathbb{Z}^d if $\beta = \beta_c$ or a GFF on \mathbb{Z}^d plus an independent drift $\sqrt{\beta - \beta_c} X \mathbf{1}_{\mathbb{Z}^d}$ with a Rademacher random variable X if $\beta > \beta_c$. Note that since we already know that $\boldsymbol{\theta}_U$ converges in law towards $\boldsymbol{\gamma}_U$, to obtain (3.7), it suffices (see, for instance, [Bil95, Theorem 25.12]) to verify uniform integrability of variables $\left[\prod_{x \in U} \theta_x^{i_x}\right]_n$. The latter follows instantaneously from the following:

Lemma 3.4 (Moments of spins of the spherical model). Let θ be a configuration of the spherical model on torus $\Lambda_n = \mathbb{T}_n^d$ at inverse temperature $\beta > 0$. For any $p \in \mathbb{N}_0$, there exists a constant $C_p > 0$ independent of n such that

$$\nu_{\Lambda_n,\beta} [|\theta_0|^{2p}] \le C_p.$$

Indeed, since $\sup_n \mathbb{E}[|X_n|^2] < \infty$ implies uniform integrability of $(X_n)_n$ and as by generalized Hölder's inequality,

$$\nu_{\Lambda_n,\beta} \bigg[\prod_{x \in U} |\theta_x|^{2i_x} \bigg] \le \prod_{x \in U} \nu_{\Lambda_n,\beta} \bigg[|\theta_x|^{2i_x|U|} \bigg]^{1/|U|} \le \nu_{\Lambda_n,\beta} \bigg[|\theta_0|^{2|U| \max_x(i_x)} \bigg].$$

The last inequality follows from the fact that spins of the spherical model on torus are identically distributed.

We now proceed to the proof of the lemma, which is motivated by [Luk20, Lemma 3].

Proof. Recall that by Proposition 2.8, $\sqrt{\beta}\theta$ has the same law as any massive $(m^2 > 0)$ GFF φ on Λ_n conditioned on $\|\varphi\|_2^2 = \beta n^d$. Therefore,

$$\beta^p \nu_{\Lambda_n,\beta} \big[\theta_0^{2p} \big] = \mathbb{E} \big[\boldsymbol{\varphi}_0^{2p} \ \big| \ \| \boldsymbol{\varphi} \|_2^2 = \beta n^d \big].$$

Let $Q = (q_{x,w})_{x,w \in [0,n)^d \cap \mathbb{Z}^d}$ be an orthonormal matrix consisting of eigenvectors of G_{Λ_n,m^2} such that $Q^T G_{\Lambda_n,m^2} Q = \operatorname{diag}(((m^2 + \eta_w)^{-1})_w)$, where $(\eta_w)_{w \in [0,n)^d \cap \mathbb{Z}^d}$ are the eigenvalues of $-\Delta_{\Lambda_n}$. Let $(Z_w)_w$ be i.i.d. standard normal random variables. Then,

$$\varphi = (\varphi_x)_{x \in \Lambda_n} \stackrel{\text{law}}{=} \left(\sum_{w \in [0,n)^d \cap \mathbb{Z}^d} q_{x,w} \frac{1}{\sqrt{m^2 + \eta_w}} Z_w \right)_{x \in \Lambda_n};$$
$$\|\varphi\|_2^2 \stackrel{\text{law}}{=} \sum_{w \in [0,n)^d \cap \mathbb{Z}^d} \frac{1}{\eta_w + m^2} Z_w^2.$$

Using this we obtain on the one hand that

$$\beta^{p} = \frac{1}{n^{dp}} \mathbb{E} \left[\| \boldsymbol{\varphi} \|_{2}^{2p} \mid \| \boldsymbol{\varphi} \|_{2}^{2} = \beta n^{d} \right] = \frac{1}{n^{dp}} \sum_{\substack{w_{1}, \dots, w_{p} \\ \in [0, n)^{d} \cap \mathbb{Z}^{d}}} \mathbb{E} \left[\prod_{j=1}^{p} \frac{Z_{w_{j}}^{2}}{m^{2} + \eta_{w_{j}}} \mid \sum_{w} \frac{Z_{w}^{2}}{\eta_{w} + m^{2}} = \beta n^{d} \right].$$

And on the other hand, by Lemma 2.12,

$$\mathbb{E}\left[\boldsymbol{\varphi}_{0}^{2p} \mid \|\boldsymbol{\varphi}\|_{2}^{2} = \beta n^{d}\right] = \sum_{\substack{w_{1}, \dots, w_{2p} \\ \in [0, n)^{d} \cap \mathbb{Z}^{d}}} \left(\prod_{j=1}^{2p} q_{0, w_{j}}\right) \mathbb{E}\left[\prod_{j=1}^{2p} \frac{Z_{w_{j}}}{\sqrt{m^{2} + \eta_{w_{j}}}} \mid \sum_{w} \frac{Z_{w}^{2}}{\eta_{w} + m^{2}} = \beta n^{d}\right]$$

$$= \frac{1}{n^{dp}} \sum_{\substack{w_1, \dots, w_{2p} \\ \in [0, n)^d \cap \mathbb{Z}^d}} \mathbb{E} \left[\prod_{j=1}^{2p} \frac{Z_{w_j}}{\sqrt{m^2 + \eta_{w_j}}} \, \middle| \, \sum_{w} \frac{Z_w^2}{\eta_w + m^2} = \beta n^d \right].$$

Since conditional density of $(Z_w/\sqrt{m^2 + \eta_w})_w$ given $\sum_{w \in [0,n)^d \cap \mathbb{Z}^d} \frac{1}{\eta_w + m^2} Z_w^2 = \beta n^d$ depends in each coordinate only on its modulus, the conditional law of the new vector obtained by swapping a sign of a coordinate remains unchanged. Therefore, as long as there exists an odd power of some Z_w in the latter product, its conditional expectation vanishes, and hence,

$$\mathbb{E}\left[\varphi_0^{2p} \mid \|\varphi\|_2^2 = \beta n^d\right] \leq \frac{(2p)!}{n^{dp}} \sum_{\substack{w_1, \dots, w_p \\ \in [0, n)^d \cap \mathbb{Z}^d}} \mathbb{E}\left[\prod_{j=1}^p \frac{Z_{w_j}^2}{m^2 + \eta_{w_j}} \mid \sum_w \frac{Z_w^2}{\eta_w + m^2} = \beta n^d\right] = (2p)!\beta^p.$$

So,
$$\nu_{\Lambda_n,\beta}[\theta_0^{2p}] \leq (2p)! = C_p$$
 as desired.

4 The infinite spin-dimensionality distributional limit of the spin O(N) model

The goal of the present section is to understand the "local" infinite spin-dimensionality limit $(N \to \infty)$ of the spin O(N) model both on the finite domain and in the infinite volume limit. More precisely, we prove the following two theorems corresponding to these cases respectively that combined with Scheffé's lemma¹ yield Theorem 1.2 in the introduction.

Theorem 4.1 $(N \to \infty \text{ limit of spin } O(N) \text{ model on a finite torus). Let } \Lambda = \mathbb{T}_n^d \text{ and } \mathbf{S} = (S_x)_{x \in \Lambda} \in (\sqrt{N}\mathbb{S}^{N-1})^{|\Lambda|} \text{ be a configuration of the spin } O(N) \text{ model at inverse temperature } \beta > 0, i.e., \mathbf{S} \sim \mu_{\Lambda,N,\beta}. \text{ Then for any } M \in \mathbb{N}, \text{ the density function of } \overline{\mathbf{S}}^M \coloneqq (S_x^i)_{x \in \Lambda}^{i=1,\ldots,M} \text{ converges as } N \to \infty \text{ to that of an } M\text{-vectorial massive } GFF \text{ scaled by } 1/\sqrt{\beta} \text{ with the mass } m^2 \text{ depending on } \beta, d \text{ and } n \text{ in such a way that}$

$$G_{\Lambda m^2}(x,x) = \beta \text{ for all } x \in \Lambda,$$

or more explicitly, m² is the unique solution to

$$\frac{1}{n^d} \frac{1}{m^2} + \frac{1}{n^d} \sum_{x \in (\hat{\Lambda} + (\lfloor n/2 \rfloor, \dots, \lfloor n/2 \rfloor)) \setminus \{0\}} \frac{1}{m^2 + 2\sum_{k=1}^d (1 - \cos(2\pi x_k/n))} = \beta.$$
(4.1)

Theorem 4.2 ($N \to \infty$ limit of spin O(N) model when volume grows to infinity). For each $n, N \in \mathbb{N}$, let $\Lambda_n = \mathbb{T}_n^d$ and $[\mathbf{S}]_n^N = [(S_x)_{x \in \Lambda_n}]_n^N$ be a configuration of the spin O(N) model on the domain Λ_n at inverse temperature $\beta > 0$. Then, for any $M \in \mathbb{N}$ and a finite box $U \subset \mathbb{Z}^d$ considered for each n sufficiently large as a subset of Λ_n that is properly contained in it, as $N \to \infty$ followed by $n \to \infty$, $([\overline{\mathbf{S}}_U^M]_n^N)_n := [(S_x^i)_{x \in U}^{i=1,\dots,M}]_n^N)_{(n,N)}$ converges in law towards:

1. $\beta < \beta_c$: an M-vectorial massive GFF on \mathbb{Z}^d restricted to U scaled by $1/\sqrt{\beta}$ with the mass m^2 depending on β and d in such a way that

$$G_{\mathbb{Z}^d,m^2}(x,x) = \beta \text{ for all } x \in \mathbb{Z}^d;$$

2. $\beta = \beta_c$: an M-vectorial GFF on \mathbb{Z}^d restricted to U scaled by $1/\sqrt{\beta}$;

¹Scheffé's lemma states that if a sequence of integrable functions $(f_n)_n$ on a measure space $(X, \mathcal{F}, \lambda)$ converges almost everywhere to another integrable function f, then $L^1(\lambda)$ -convergence of (f_n) towards f holds if and only if $||f_n||_{L^1(\mu)} \to ||f||_{L^1(\mu)}$.

3. $\beta > \beta_c$: an M-vectorial GFF on \mathbb{Z}^d restricted to U scaled by $1/\sqrt{\beta}$ plus an independent constant (over U) random drift of the form $\sqrt{\frac{\beta-\beta_c}{\beta}}\overline{Z}^M\mathbf{1}_U$ with \overline{Z}^M being an M-dimensional standard normal vector.

4.1 On the finite torus

This subsection is concerned with the infinite spin-dimensionality limit of the spin O(N) model on a finite torus and in particular proves Theorem 4.1.

Before proceeding to the proof let us quickly discuss the strategy. First, by Proposition 2.9 we know that

$$\operatorname{Law}\left(\overline{\mathbf{S}}^{M}\right) = \operatorname{Law}\left(\overline{\mathbf{\Phi}}^{M}/\sqrt{\beta} \mid ||\Phi_{x}|| = \sqrt{\beta N}, \ \forall x \in \Lambda\right),$$

for any N-vectorial m^2 -massive GFF $\Phi := (\overline{\Phi}^M, \overline{\Phi}^{\text{rest}})^T$ such that $m^2 > 0$. Thus, for any $\overline{\gamma}^M := (\overline{\gamma}_x^M)_{x \in \Lambda} \in (\mathbb{R}^M)^{n^d}$, we have the following equality of densities w.r.t. Lebesgue measure on \mathbb{R}^{Mn^d} :

$$f_{\overline{\mathbf{S}}^{M}}(\overline{\gamma}^{M}) = f_{\overline{\mathbf{\Phi}}^{M}/\sqrt{\beta}} \Big|_{(\|\Phi_{x}\|)_{x} = (\sqrt{\beta}N)_{x}}(\overline{\gamma}^{M}) = \frac{f(\overline{\mathbf{\Phi}}^{M}/\sqrt{\beta}, (\|\Phi_{x}\|^{2})_{x})}{f(\|\Phi_{x}\|^{2})_{x}((\beta N)_{x})}$$

$$= f_{\overline{\mathbf{\Phi}}^{M}/\sqrt{\beta}}(\overline{\gamma}^{M}) \frac{f(\|\overline{\Phi_{x}}^{rest}\|^{2})_{x}}{f(\|\Phi_{x}\|^{2})_{x}((\beta N)_{x})}. \tag{4.2}$$

Since by the central limit theorem, $\left(\frac{\|\overline{\Phi_x}^{\mathrm{rest}}\|^2 - \mathbb{E}[\|\overline{\Phi_x}^{\mathrm{rest}}\|^2]}{\sqrt{N}}\right)_x = \left(\frac{\|\overline{\Phi_x}^{\mathrm{rest}}\|^2 - NG_{\Lambda,m^2}(0,0)}{\sqrt{N}}\right)_x$ converges to a centered n^d -dimensional Gaussian vector, we might expect that by choosing m^2 as in (4.1) we will be able to conclude that the above ratio converges to one, which would complete the proof. Motivated by this discussion, we state and prove (see A.3) the following result, which is a partial generalization of [GK68, Chp.8 §46 Theorem 1] from \mathbb{R} to \mathbb{R}^k .

Proposition 4.3. Let $(\mathbf{X}^i)_i$ be a sequence of i.i.d. centered random vectors with the probability density function (w.r.t. Lebesgue measure on \mathbb{R}^k) $f: \mathbb{R}^k \to \mathbb{R}$. Assume that

- 1. $f \in L^r(\mathbb{R}^k)$ for some $r \in (1,2]$;
- 2. all the entries of the covariance matrix C of $\mathbf{X}^1 = (X_j^1)_{j=1}^k$ are well-defined and finite, or equivalently, for all $1 \leq j \leq k$, $X_j^1 \in L^2(\mathbb{P})$;
- 3. C is positive definite.

Then the relation

$$n^{k/2} f^{(n)}(\sqrt{n} \boldsymbol{x}) \xrightarrow{n \to \infty} \frac{1}{(2\pi)^{k/2} \sqrt{\det(C)}} e^{-\frac{1}{2}(\boldsymbol{x}, C^{-1} \boldsymbol{x})}$$

holds uniformly over $\mathbf{x} \in \mathbb{R}^k$. Here, $f^{(n)}$ is the n-fold convolution of f, as well as the density function of $\sum_{i=1}^{n} \mathbf{X}^i$.

Proof of Theorem 4.1. Choose $m^2 > 0$ as in (4.1). Then, $\mathbb{E}[(\Phi_x^i)^2] = G_{\Lambda,m^2}(x,x) = \beta$. We start by checking that i.i.d. vectors $((\Phi^i)^2 - \beta := ((\Phi_x^i)^2 - \beta)_{x \in \Lambda})_i$ satisfy assumptions of

Proposition 4.3. Property 2. is obvious since we are working with squares of Gaussians. As for 3., as a covariance matrix, C is positive semi-definite. Furthermore, $\mathbf{x}^T C \mathbf{x} = 0$ iff $\langle \mathbf{x}, (\mathbf{\Phi}^i)^2 - \beta \rangle = 0$ almost surely, which in turn is true only for $\mathbf{x} = 0$. To verify 1., we derive a formula for the density function f of $((\Phi_x^1)^2 - \beta)_{x \in \Lambda}$ and show that it belongs to $L^r(\mathbb{R}^{n^d})$ for any $r \in [1, 2)$. Let f_{φ} be the density of an m^2 -massive GFF on Λ , then

$$f(t) = \left(\prod_{x} \frac{1}{2\sqrt{t_x + \beta}}\right) \sum_{(k_x)_x \in \{0,1\}^{n^d}} f_{\varphi} \left(\left((-1)^{k_x} \sqrt{t_x + \beta} \right)_x \right) \mathbb{1}_{\{\forall x: t_x + \beta > 0\}}.$$

Recall that the eigenvalues of $(-\Delta_{\Lambda} + m^2)$ belong to $[m^2, m^2 + 4d]$ (see Lemma 2.12). Therefore,

$$f(t) \le \frac{2^{n^d}}{(2\pi)^{n^d/2} \sqrt{\det G_{\Lambda,m^2}}} \prod_x \left(\frac{1}{2\sqrt{t_x + \beta}} e^{-\frac{1}{2}m^2(t_x + \beta)} \mathbb{1}_{\{t_x + \beta > 0\}} \right).$$

From this explicit formula we directly see that $f \in L^r(\mathbb{R}^{n^d})$ for any $r \in [1,2)$. Now that we have established that Proposition 4.3 holds in our setting, we return to the ratio (4.2) derived above.

$$\frac{f_{(\|\overline{\Phi_x}^{\mathrm{rest}}\|^2)_x}\left((\beta N - \beta \|\overline{\gamma}_x^M\|^2)_x\right)}{f_{(\|\Phi_x\|^2)_x}((\beta N)_x)} = \left(\frac{N}{N-M}\right)^{\frac{n^d}{2}} \times \frac{\left(N-M\right)^{\frac{n^d}{2}} f_{(\|\overline{\Phi_x}^{\mathrm{rest}}\|^2 - (N-M)\beta)_x}\left(\sqrt{N-M}(\frac{\beta M - \beta \|\overline{\gamma}_x^M\|^2}{\sqrt{N-M}})_x\right)}{N^{n^d/2} f_{(\|\Phi_x\|^2 - N\beta)_x}(\sqrt{N}(0)_x)} \times \frac{N \to \infty}{\Pr{\mathrm{op. 4.3}}} \xrightarrow{f_{\mathbf{Z}}(\mathbf{0})} = 1,$$

where **Z** is a centered Gaussian vector in \mathbb{R}^{n^d} with covariances $\mathbb{E}[Z_x Z_y] = \mathbb{E}[(\varphi_x^2 - \beta)(\varphi_y^2 - \beta)] = 2G_{\Lambda,m^2}^2(x,y)$.

4.2 The infinite-volume limit: $N \to \infty$ followed by $n \to \infty$

In this subsection we discuss the infinite spin-dimensionality and infinite volume limit and prove Theorem 4.2.

By first taking limit $N \to \infty$, we significantly simplify our work in light of Theorem 4.1. Indeed, the latter yields that for any $n \in \mathbb{N}$ fixed, $[\overline{\mathbf{S}}_U^M]_n^N$ converges in law as $N \to \infty$ to $\frac{1}{\sqrt{\beta}}[\overline{\mathbf{\Phi}}_U^M]_n \coloneqq \frac{1}{\sqrt{\beta}}[(\Phi_x^i)_{x \in U}^{i=1,\dots,M}]_n$, an M-vectorial m_n^2 -massive GFF on $\Lambda_n \coloneqq \mathbb{T}_n^d$ restricted to U. Now, since we are working with M independent identically distributed centered Gaussian vectors, to complete the proof we only have to verify that the covariance matrix of one of the coordinate processes $([\varphi_U]_n)_n$ converges to the covariance matrix of either an m^2 -massive GFF $\tilde{\psi}_U$ on \mathbb{Z}^d restricted to U if $\beta < \beta_c$, to a GFF ψ_U on \mathbb{Z}^d restricted to U if $\beta = \beta_c$, or if $\beta > \beta_c$, to the one of $\psi_U + \sqrt{\beta} - \beta_c Z \mathbf{1}_U$, where Z is an independent standard normal random variable. For this, recall that the critical inverse temperature $\beta_c \in (0, \infty]$ corresponds to $G_{\mathbb{Z}^d}(0,0)$. In particular, $\beta_c = \infty$ if d = 2, $\beta_c \in (0,\infty)$ for $d \geq 3$. Let us have a closer look at (4.1): using Riemann sum approximation, for $m^2 > 0$ if d = 2, or $m^2 \geq 0$ if $d \geq 3$, we get

$$\frac{1}{n^{d}} \sum_{x \in (\hat{\Lambda} + (\lfloor n/2 \rfloor, \dots, \lfloor n/2 \rfloor)) \setminus \{0\}} \frac{1}{m^{2} + 2 \sum_{k=1}^{d} (1 - \cos(2\pi x_{k}/n))}
\stackrel{n \to \infty}{\sim} \int_{(0,1)^{d}} \frac{1}{m^{2} + 2 \sum_{k=1}^{d} (1 - \cos(2\pi x_{k}))} dx = G_{\mathbb{Z}^{d}, m^{2}}(0,0).$$
(4.3)

This implies that if $\beta < \beta_c$, $(m_n^2)_n$, solutions to (4.1) for $\Lambda = \Lambda_n$, converge to $m^2 > 0$, the unique solution to $G_{\mathbb{Z}^d,m^2}(0,0) = \beta$. It remains to show that in this case $(\beta < \beta_c)$, $G_{\Lambda_n,m_n^2}(x,y) \to G_{\mathbb{Z}^d,m^2}(x,y)$ for all $x,y \in U$. But this has already been done in Section 2.2 right after Definition 2.5.

Suppose that $\beta > \beta_c$ $(d \ge 3)$. By (4.3) we see that in the limit the sum in (4.1) can be at most β_c . Hence, m_n^2 must be of order $O(1/n^d)$. Since all non-zero eigenvalues of $-\Delta_{\Lambda_n^d}$ are at least of order $\Omega(1/n^2)$,

$$\frac{1}{n^d} \sum_{x \in (\hat{\Lambda} + (\lfloor n/2 \rfloor, \dots, \lfloor n/2 \rfloor)) \setminus \{0\}} \frac{1}{m_n^2 + 2 \sum_{k=1}^d (1 - \cos(2\pi x_k/n))} \\
= \frac{1 - \mathcal{O}(n^{2-d})}{n^d} \sum_{x \in (\hat{\Lambda} + (\lfloor n/2 \rfloor, \dots, \lfloor n/2 \rfloor)) \setminus \{0\}} \frac{1}{2 \sum_{k=1}^d (1 - \cos(2\pi x_k/n))} \stackrel{n \to \infty}{\sim} \beta_c.$$

Therefore, $m_n^2 = \frac{1}{\beta - \beta_c} \frac{1}{n^d} (1 + o(1))$ with o(1) uniform in n large. Let u^1, \ldots, u^{n^d} be the orthonormal eigenvectors of $-\Delta_{\Lambda_n^d}$ corresponding to $0 = \eta_1 < \eta_2 \le \ldots \le \eta_{n^d}$; in particular, $u^1 = \frac{1}{\sqrt{n^d}} \mathbf{1}_{\Lambda_n}$. For any $x, y \in U$, we have

$$G_{\Lambda_{n}^{d},m_{n}^{2}}(x,y) = \sum_{k=2}^{n^{d}} u^{k}(x) \frac{1}{m_{n}^{2} + \eta_{k}} u^{k}(y) + u^{1}(x)u^{1}(y) \frac{1}{m_{n}^{2}}$$

$$= (1 + o(1)) \Big[\sum_{k=2}^{n^{d}} u^{k}(x) \frac{1}{\eta_{k}} u^{k}(y) + \frac{1}{n^{d}} (\beta - \beta_{c}) n^{d} \Big]$$

$$= (1 + o(1)) \Big[G_{\Lambda_{n}^{d}}^{0-\text{avg.}}(x,y) + (\beta - \beta_{c}) \Big]$$

$$\xrightarrow{\text{unif. over } x, y \in U} G_{\mathbb{Z}^{d}}(x,y) + (\beta - \beta_{c}),$$

$$(4.4)$$

where $G_{\Lambda_n^d}^{0\text{-avg.}}$ is the zero-average Green's function on torus. We conclude by noticing that $(G_{\mathbb{Z}^d}(x,y)+(\beta-\beta_c))_{x,y\in U}$ is the covariance matrix of $\psi_U+\sqrt{\beta-\beta_c}Z\mathbf{1}_U$.

Let $\beta = \beta_c$. From (2.5) and Remark 2 combined with the argument in the high temperature regime we easily see that for any $\varepsilon > 0$, there exists N sufficiently large such that for all $n \geq N$, $0 < m_n^2 < \varepsilon$. Thus, for any $\delta \in (0,1)$,

$$G_{\mathbb{Z}^d,\varepsilon}(x,y) \xleftarrow{n \to \infty}_{\text{unif. in } U} G_{\Lambda_n,\varepsilon}(x,y) \le G_{\Lambda_n,m_n^2}(x,y) \le G_{\Lambda_n,\min(m_n^2,n^{-2-\delta})}(x,y).$$

As $\varepsilon \to 0$, $G_{\mathbb{Z}^d,\varepsilon}(x,y)$ converges to $G_{\mathbb{Z}^d}(x,y)$ uniformly in $x,y \in U$. Moreover, analogously to (4.4),

$$\begin{split} G_{\Lambda_n,\min(m_n^2,n^{-2-\delta})}(x,y) &= \sum_{k=2}^{n^d} u^k(x) \frac{1}{\min(m_n^2,n^{-2-\delta}) + \eta_k} u^k(y) + \frac{1}{\min(m_n^2,n^{-2-\delta})n^d} \\ &= \sum_{k=2}^{n^d} u^k(x) \frac{1}{\eta_k} u^k(y) (1 - \mathcal{O}(n^{-\delta})) + \mathcal{O}(n^{2+\delta-d}) \\ &\quad + \left(\beta_c - \frac{1 - \mathcal{O}(m_n^2 n^2)}{n^d} \sum_{k=2}^{n^d} \frac{1}{\eta_k}\right) \mathbbm{1}_{\{m_n^2 \le n^{-2-\delta}\}} \\ &= (1 - o(1)) G_{\Lambda_n^d}^{0\text{-avg.}}(x,y) + o(1) \\ &\quad + \left(\beta_c - (1 - o(1)) G_{\Lambda_n^d}^{0\text{-avg.}}(0,0)\right) \mathbbm{1}_{\{m_n^2 \le n^{-2-\delta}\}} \end{split}$$

$$\xrightarrow[\text{unif. over } x,y\in U]{} G_{\mathbb{Z}^d}(x,y)$$

Hence, G_{Λ_n,m_n^2} converges uniformly in U^2 to $G_{\mathbb{Z}^d}$ as as desired.

A Appendix

A.1 Polynomial decay of zero-average Green's function

The goal of this subsection is to prove the following.

<u>Claim:</u> There exists C = C(d) > 0 such that for all n sufficiently large and $y \in \hat{\Lambda}$,

$$|G_{\Lambda}^{0\text{-avg.}}(0,y) - G_{\mathbb{Z}^d}(0,y)| \le Cn^{2-d}.$$

Proof. Recall that $\hat{\Lambda} = [-n/2, n/2)^d \cap \mathbb{Z}^d$ is the canonical projection of the torus $\Lambda = \mathbb{T}_n^d$ onto \mathbb{Z}^d and by Proposition 2.13,

$$2dG_{\Lambda}^{0\text{-avg.}}(0,y) = \int_{0}^{\infty} \left(\mathbb{P}_{\Lambda}^{0}[\overline{X}_{t} = y] - \frac{1}{n^{d}} \right) dt,$$

where $\overline{\mathbf{X}} = (\overline{X}_t)_{t \geq 0}$ is the continuous-time simple random walk on Λ viewed as a graph. To obtain the desired estimate we follow the strategy of the proof of [Abä17, Proposition 1.5] corresponding to our (2.12) and split the above integral approximately at the time $\overline{\mathbf{X}}$ reaches equilibrium (the uniform distribution). More precisely, as in the reference (with N=n in our setting), for $\lambda_* := \frac{1}{d}(1-\cos(2\pi/n)) \stackrel{n\to\infty}{\sim} \frac{2\pi^2}{dn^2}$ (the spectral gap of $\overline{\mathbf{X}}$) and $t_* := \frac{\log(n^d)}{\lambda_*}$ by [LP17, Theorem 20.6],

$$\int_{t_*}^{\infty} \left| \mathbb{P}_{\Lambda}^0[\overline{X}_t = y] - \frac{1}{n^d} \right| \mathrm{d}t \le \int_{t_*}^{\infty} e^{-\lambda_* t} \mathrm{d}t = \frac{1}{\lambda_* n^d} = \Theta(n^{2-d}).$$

The remaining part of the integral will be treated similarly to the referenced proof but with much greater precision. Note that

$$\begin{split} & \int_0^{t_*} \left(\mathbb{P}^0_{\Lambda}[\overline{X}_t = y] - \frac{1}{n^d} \right) \mathrm{d}t = \int_0^{t_*} \sum_{k=0}^{\infty} \mathbb{P}^0_{\Lambda}[N_t = k] \mathbb{P}^0_0[X_k = y] \mathrm{d}t - \frac{t_*}{n^d} \\ & \in \sum_{k=0}^{t_* + \sqrt{4t_* \log(n^d)}} \mathbb{P}^0_0[X_k = y] \int_0^{t_*} \frac{t^k e^{-t}}{k!} \mathrm{d}t - \frac{t_*}{n^d} + \left[0, \int_0^{t_*} \mathbb{P}^0_{\Lambda}[N_t > t_* + \sqrt{4t_* \log(n^d)}] \mathrm{d}t \right]. \end{split}$$

Here N_t is the number of jumps of $(\overline{X}_s)_s$ up to time t. Let us show that $\int_0^{t_*} \mathbb{P}^0_{\Lambda}[N_t > t_* + \sqrt{4t_* \log(n^d)}] dt = o(n^{2-d})$ as $n \to \infty$. For $t \in (0, t_*)$ by exponential Markov's inequality,

$$\mathbb{P}_{\Lambda}^{0}[N_{t} > t_{*} + \sqrt{4t_{*}\log(n^{d})}] \leq \inf_{\lambda > 0} \exp\left(t(e^{\lambda} - 1) - \lambda\left(t_{*} + \sqrt{4t_{*}\log(n^{d})}\right)\right)
\leq \exp\left(t_{*}\inf_{\lambda > 0}\left[\left(e^{\lambda} - 1\right) - \lambda\left(1 + \sqrt{4\lambda_{*}}\right)\right]\right)
= \exp\left(t_{*}\left[\sqrt{4\lambda_{*}} - \underbrace{\log\left(1 + \sqrt{4\lambda_{*}}\right)}_{\geq \sqrt{4\lambda_{*}} - 4\lambda_{*}/2}\left(1 + \sqrt{4\lambda_{*}}\right)\right]\right)
\leq \exp\left(-2t_{*}\lambda_{*}(1 - o(1))\right) \leq (n^{d})^{-3/2}.$$
(A.1)

And thus, $\int_0^{t_*} \mathbb{P}^0_{\Lambda}[N_t > t_* + \sqrt{4t_* \log(n^d)}] dt \le t_* n^{-3d/2} = \Theta(n^{2-3d/2} \log(n)) = o(n^{2-d})$ as desired. Furthermore, for $0 \le k \le t_* + \sqrt{4t_* \log(n^d)} =: t_*'$,

$$\mathbb{P}^{0}_{0}[X_{k} = y] = \sum_{\substack{v \in \mathbb{Z}^{d}: \\ d_{\mathbb{Z}^{d}}(0, y + nv) \le t'_{*}}} \mathbb{P}^{0, \mathbb{Z}^{d}}_{0}[X_{k} = y + nv].$$

We split the latter sum into two parts: $\sqrt{t'_*}C_d\sqrt{\log(n^d)} \leq d_{\mathbb{Z}^d}(0, y + nv) \leq t'_*$ and $d_{\mathbb{Z}^d}(0, y + nv) \leq \sqrt{t'_*}C_d\sqrt{\log(n^d)}$ for some $C_d > 0$ to be determined later. We show that the contribution of the former fragment of the sum to $\sum_{k=0}^{t'_*} \mathbb{P}_0^0[X_k = y] \int_0^{t_*} \frac{t^k e^{-t}}{k!} dt$ can be made as small as $o(n^{2-d})$ if C_d is chosen appropriately (the lower bound is clearly zero). For some appropriate absolute (depending only on the dimension) constants $c, c', r_0 > 0, R_0 < \infty$,

$$\sum_{k=0}^{t_{*}} \int_{0}^{t_{*}} \frac{t^{k} e^{-t}}{k!} dt \underbrace{\sum_{\substack{v \in \mathbb{Z}^{d}: \sqrt{t_{*}'} C_{d} \sqrt{\log(n^{d})} \leq \\ d_{\mathbb{Z}^{d}}(0, y + nv) \leq t_{*}' \\ \leq c(t_{*}'/n)^{d}}}_{\leq L_{*}} \leq c(t_{*}'/n)^{d}$$

$$\leq \mathbb{P}_{0}^{0, \mathbb{Z}^{d}} \left[\max_{k=0}^{\lfloor t_{*}' \rfloor} d_{\mathbb{Z}^{d}}(0, X_{k}) \geq \sqrt{t_{*}'} C_{d} \sqrt{\log(n^{d})} \right] \underbrace{t_{*} c(t_{*}'/n)^{d}}_{=\Theta(n^{2+d} \log^{d+1}(n))}$$

$$\leq c' n^{2+d} \log^{d+1}(n) \mathbb{P}_{0}^{0, \mathbb{Z}^{d}} \left[\max_{k=0}^{\lfloor t_{*}' \rfloor} |X_{k}| \geq (C_{d}/d) \sqrt{t_{*}' \log(n^{d})} \right]$$

$$\leq c' n^{2+d} \log^{d+1}(n) R_{0} e^{-r_{0}(C_{d}/d)^{2} \log(n^{d})} = \Theta(n^{2+d-r_{0}C_{d}^{2}/d} \log^{d+1}(n)).$$

The last inequality (and in particular, existence of r_0, R_0 as above) follows from [LL10, Proposition 2.1.2 (b)]. By choosing C_d such that $C_d > d\sqrt{2/r_0}$, say, $C_d = (2d/\sqrt{r_0}) \vee 1$, we get that the latter expression is $o(n^{2-d})$. It remains to treat

$$I_n := \sum_{k=0}^{t'_*} \sum_{\substack{v \in \mathbb{Z}^d : d_{\mathbb{Z}^d}(0, y + nv) \\ \leq C_d \sqrt{t'_* \log(n^d)}}} \mathbb{P}_0^{0, \mathbb{Z}^d} [X_k = y + nv] \int_0^{t_*} \frac{t^k e^{-t}}{k!} dt - \frac{t_*}{n^d}.$$

First observe that $\int_0^{t_*} \frac{t^k e^{-t}}{k!} dt = \mathbb{P}[\operatorname{Pois}(t_*) \geq k+1]$ (by definition of the upper incomplete gamma function and explicit expression of Poisson cumulative distribution function in terms of it). Therefore,

$$I_n \le \sum_{k=0}^{t'_*} \sum_{\substack{v \in \mathbb{Z}^d : d_{\mathbb{Z}^d}(0, y + nv) \\ \le C_d \sqrt{t'_* \log(n^d)}}} \mathbb{P}_0^{0, \mathbb{Z}^d} [X_k = y + nv] - \frac{t_*}{n^d},$$

but also

$$\begin{split} I_{n} &\geq \mathbb{P}[\operatorname{Pois}(t_{*}) \geq t_{*} - \sqrt{4t_{*} \log(n^{d})}] \\ &\times \sum_{k=0}^{t_{*} - \sqrt{4t_{*} \log(n^{d})}} \sum_{v \in \mathbb{Z}^{d} : d_{\mathbb{Z}^{d}}(0, y + nv)} \mathbb{P}_{0}^{0, \mathbb{Z}^{d}} [X_{k} = y + nv] - \frac{t_{*}}{n^{d}} \\ &\geq (1 - n^{-3d/2}) \sum_{k=0}^{t_{*} - \sqrt{4t_{*} \log(n^{d})}} \sum_{v \in \mathbb{Z}^{d} : d_{\mathbb{Z}^{d}}(0, y + nv)} \mathbb{P}_{0}^{0, \mathbb{Z}^{d}} [X_{k} = y + nv] - \frac{t_{*}}{n^{d}}. \end{split}$$

In the latter inequality we have used that fully analogously to (A.1), $\mathbb{P}[\operatorname{Pois}(t_*) < t_* - \sqrt{4t_*\log(n^d)}] \le n^{-3d/2}$. Define $p_k : \mathbb{R}^d \to \mathbb{R}$ by $p_k(x) \coloneqq 2\left(\frac{d}{2\pi k}\right)^{d/2}e^{-\frac{d|x|^2}{2k}}$ and write $x \leftrightarrow k$ if k and x have the same parity, i.e., if $k+x_1+\ldots+x_d$ is even. Set further $E_k(x) \coloneqq \left(\mathbb{P}_0^{0,\mathbb{Z}^d}[X_k = x] - p_k(x)\right)\mathbb{1}_{\{k\leftrightarrow x\}}$. By [Law12, Lemma 1.5.2], $|x|^\alpha \sum_{k\geq 0}|E_k(x)| \xrightarrow{|x|\to\infty} 0$ for every $\alpha < d$. Note that for $y \in \hat{\Lambda}$ and $v \in \mathbb{Z}^d \setminus \{0\}$, $|y+nv| = \Omega(n) \ge n/2$. Altogether, this implies that for $t \in \{t'_*, t_* - \sqrt{4t_*\log(n^d)}\}$ and $\alpha \in (d-2, d)$,

$$\sum_{k=0}^{t} \sum_{v \in \mathbb{Z}^{d}: d_{\mathbb{Z}^{d}}(0, y + nv) \atop \leq C_{d} \sqrt{t'_{*} \log(n^{d})}} \mathbb{P}_{0}^{0, \mathbb{Z}^{d}} [X_{k} = y + nv]$$

$$= \sum_{k=1}^{t} \sum_{v \in \mathbb{Z}^{d} \setminus \{0\}: d_{\mathbb{Z}^{d}}(0, y + nv)} p_{k}(y + nv) \mathbb{1}_{\{k \leftrightarrow y + nv\}}$$

$$\leq C_{d} \sqrt{t'_{*} \log(n^{d})}$$

$$=: J_{n}(t)$$

$$+ \sum_{k=0}^{t} \mathbb{P}_{0}^{0, \mathbb{Z}^{d}} [X_{k} = y] \pm \underbrace{\Theta((\log n)^{d}) o(n^{-\alpha})}_{=o(n^{2-d})}.$$

Claim 1:
$$0 \le 2dG_{\mathbb{Z}^d}(0, y) - \sum_{k=0}^t \mathbb{P}_0^{0, \mathbb{Z}^d} [X_k = y] \le o(n^{2-d}).$$

Claim 2: $|J_n(t) - \frac{t_*}{n^d}| = \mathcal{O}(n^{2-d}).$

Note that combined with the above arguments these claims would imply the desired result. Indeed, by the above and since $G_{\mathbb{Z}^d}(0,y)$ is finite for all $y \in \mathbb{Z}^d$,

$$-\mathcal{O}(n^{2-d}) + \underbrace{(1 - n^{-3d/2}) \Big(J_n + \sum_{k=0}^t \mathbb{P}_0^{0,\mathbb{Z}^d} [X_k = y] \Big) - \frac{t_*}{n^d}}_{\geq 2dG^{0-\text{avg.}}(0,y)} \leq 2dG^{0-\text{avg.}}(0,y)$$

$$\leq \mathcal{O}(n^{2-d}) + \underbrace{J_n - \frac{t_*}{n^d} + \sum_{k=0}^t \mathbb{P}_0^{0,\mathbb{Z}^d} [X_k = y]}_{\leq 2dG_{\mathbb{Z}^d}(0,y) + \mathcal{O}(n^{2-d})}.$$

Hence, $|G^{0\text{-avg.}}(0,y) - G_{\mathbb{Z}^d}(0,y)| = \mathcal{O}(n^{2-d}).$

Let us first prove Claim 1. Recall that $2dG_{\mathbb{Z}^d}(0,y) = \sum_{k\geq 0} \mathbb{P}_0^{0,\mathbb{Z}^d}[X_k=y]$. Therefore, it only remains to check that the tail $(k\geq t)$ of this sum decays at rate $o(n^{2-d})$. This follows almost immediately from the following three observations: $p_k(y) \leq C(d)r^{-\frac{d}{2}}, |E_k(y)| \leq O(k^{-\frac{d}{2}-1})$ by [Law12, Theorem 1.2.1 (1.10)] and $t = \Omega(n^2 \log n)$, which together imply that the tail is bounded by $\int_{Cn^2 \log n}^{\infty} (r^{-\frac{d}{2}-1} + r^{-\frac{d}{2}}) \leq O(n^{2-d}/(\log n)^{\frac{d}{2}-1}) = o(n^{2-d})$ as $d \geq 3$.

We turn to Claim 2 and first show that the contribution of $k < \varepsilon n^2$ terms (for any $0 < \varepsilon < \frac{1}{4}$) is $\mathcal{O}(n^{2-d})$. Indeed, since $\mathbb{R}_+ \ni r \mapsto p_r(z)$ is increasing on $(0, |z|^2)$ for any $z \in \mathbb{R}^d$ and $|y + nv|^2 \ge \frac{n^2}{4}$ for $v \in \mathbb{Z}^d \setminus \{0\}$,

$$\sum_{k=1}^{\varepsilon n^2} \sum_{v \in \mathbb{Z}^d \setminus \{0\} : d_{\mathbb{Z}^d}(0, y + nv)} p_k(y + nv) \le \sum_{v \in \mathbb{Z}^d \setminus \{0\} : d_{\mathbb{Z}^d}(0, y + nv) \atop \leq C_d \sqrt{t_*' \log(n^d)}} \int_1^{2\varepsilon n^2} 2\left(\frac{d}{2\pi r}\right)^{d/2} e^{-\frac{d|y + nv|^2}{2r}} dr$$

$$\leq \sum_{v \in \mathbb{Z}^d \setminus \{0\}: d_{\mathbb{Z}^d}(0, y + nv)} c_{d|y + nv|^2} d_{\frac{d|y + nv|^2}{4\varepsilon n^2}} s^{\frac{d}{2} - 2} e^{-s} ds$$

$$\leq \tilde{c}_d n^{2 - d} \varepsilon^{2 - \frac{d}{2}} \sum_{v \in \mathbb{Z}^d \setminus \{0\}: d_{\mathbb{Z}^d}(0, y + nv) \atop \leq C_d \sqrt{t'_* \log(n^d)}} \left| \frac{y}{n} + v \right|^{-2} e^{-\frac{d}{4\varepsilon} \left| \frac{y}{n} + v \right|^2}$$

$$\leq c'_d n^{2 - d} \varepsilon^{2 - \frac{d}{2}} \prod_{i=1}^d \left(\sum_{v_i = -2\tilde{C}_d \log n} e^{-\frac{d}{4\varepsilon} \left(\frac{y_i}{n} + v_i \right)^2} \right)$$

$$\leq c''_d n^{2 - d} \varepsilon^{2 - \frac{d}{2}} \left(\int_{\mathbb{R}} e^{-\frac{d}{4\varepsilon} x^2} dx + 1 \right)^d = \hat{c}_d n^{2 - d} \varepsilon^{2 - \frac{d}{2}}.$$

In the third line we used that for all x > 1,

$$\int_{x}^{\infty} s^{d/2} e^{-s} ds \le x^{d/2 - 2} e^{-x} \left(1 + \sum_{k=2}^{\lfloor d/2 \rfloor} \prod_{j=2}^{k} \left(\frac{d}{2} - j \right) \right).$$

Note also that there exists an absolute constant c>0 such that for all $|z| \leq 2C_d \sqrt{t'_* \log(n^d)}$, $t'_* \geq s \geq \varepsilon n^2$: $p_{\lfloor s \rfloor}(z)/p_s(z) \in \left(1-c\frac{\log^2 n}{\varepsilon^2 n^2}, 1+c\frac{\log^2 n}{\varepsilon^2 n^2}\right)$. Since $|y+nv| \leq d_{\mathbb{Z}^d}(0,y+nv) \leq d|y+nv|$, and $\frac{1}{2}p_k$ is the density function of $\mathcal{N}\left(0,\frac{k}{d}\mathrm{Id}_d\right)$, this yields that

$$J_n(t) \leq_a \left(1 \pm c \frac{\log^2 n}{\varepsilon^2 n^2}\right) \int_{\varepsilon n^2}^t ds \int_{B^d\left(\frac{-y}{n}, aC_d\sqrt{\frac{t'_* \log(n^d)}{n^2}}\right) \setminus [-\frac{1}{2}, \frac{1}{2})^d} \frac{p_s\left(n\left(\frac{y}{n} + \lfloor v \rfloor\right)\right)}{2} dv + \mathcal{O}(n^{2-d})$$

$$= \frac{n^2}{n^d} \left(1 \pm c \frac{\log^2 n}{\varepsilon^2 n^2}\right) \int_{\varepsilon}^{\frac{t}{n^2}} dr \int_{B^d\left(\frac{-y}{n}, aC_d\sqrt{\frac{t'_* \log(n^d)}{n^2}}\right) \setminus [-\frac{1}{2}, \frac{1}{2})^d} \frac{p_r\left(\frac{y}{n} + \lfloor v \rfloor\right)}{2} dv + \mathcal{O}(n^{2-d}),$$

where $\lfloor v \rfloor \in \mathbb{Z}^d$ stands for the point on the grid \mathbb{Z}^d such that $v \in \lfloor v \rfloor + \lfloor -1/2, 1/2 \rfloor^d$ and \lessgtr_a here means that $J_n(t) \leq \text{r.h.s.}$ with a=2, non-negative $\mathfrak{O}(n^{2-d})$ and + in $1 \pm c \frac{\log^2 n}{\varepsilon^2 n^2}$ and $J_n(t) \geq \text{r.h.s.}$ for a=1/(2d) with $\mathfrak{O}(n^{2-d})=0$ and - in $1 \pm c \frac{\log^2 n}{\varepsilon^2 n^2}$. Observe that

$$0 \le \frac{n^2}{n^d} \int_{\varepsilon}^{\frac{t}{n^2}} dr \frac{p_r(\frac{y}{n})}{2} \le C_d \frac{n^2}{n^d} \int_{\varepsilon}^{\frac{t}{n^2}} r^{-\frac{d}{2}} dr \le \mathcal{O}(n^{2-d}).$$

Therefore, we can add $\left[-\frac{1}{2},\frac{1}{2}\right]^d$ back to the integral by adding $-\mathfrak{O}(n^{2-d})$ to the lower bound of $J_n(t)$, for the upper bound other than that nothing changes. Using first less precise bounds, let us further show that

$$J_n(t) = \frac{n^2}{n^d} \int_{\varepsilon}^{\frac{t_*}{n^2}} dr \int_{\mathbb{R}^d} \frac{p_r(\frac{y}{n} + \lfloor v \rfloor)}{2} dv + \mathcal{O}(n^{2-d}).$$
 (A.2)

By the mean-value theorem, for any $v \in B^d(\frac{-y}{n}, 2C_d\sqrt{\frac{t'_*\log(n^d)}{n^2}})$, there exists $c_v \in [0, 1]$ such that

$$\begin{split} \left| p_r \left(\frac{y}{n} + \lfloor v \rfloor \right) - p_r \left(\frac{y}{n} + v \right) \right| \\ & \leq p_r \left(\frac{y}{n} + (1 - c_v)v + c_v \lfloor v \rfloor \right) \frac{d \left| \frac{y}{n} + (1 - c_v)v + c_v \lfloor v \rfloor \right|}{r} |v - \lfloor v \rfloor| \\ & \leq e^{\frac{d}{2r} c_v^2 |v - \lfloor v \rfloor|^2} p_r \left(\frac{1}{\sqrt{2}} \left(\frac{y}{n} + v \right) \right) d^{3/2} \frac{\left| \frac{y}{n} + v \right| + \sqrt{d}}{r} \end{split}$$

$$\leq c_d(\varepsilon)p_r\Big(\frac{1}{\sqrt{2}}\Big(\frac{y}{n}+v\Big)\Big)\frac{\left|\frac{y}{n}+v\right|+1}{r}.$$

Furthermore, for any C > 0, $0 < s \le t'_*$,

$$\mathbb{P}[\|\mathcal{N}(0, (s/d)\mathrm{Id}_d)\| \ge C\sqrt{t'_*\log(n^d)}] \le \mathbb{P}[\|\mathcal{N}(0, \mathrm{Id}_d)\| \ge C\sqrt{d\log(n^d)}] \\
\le 1 - \mathbb{P}[|\mathcal{N}(0, 1)| \le C\sqrt{\log(n^d)}]^d \le 1 - (1 - n^{-dC^2})^d = \mathcal{O}(n^{-dC^2}).$$

Hence, for $R := aC_d \sqrt{\frac{t'_* \log(n^d)}{n^2}}$,

$$\int_{B^d\left(\frac{-y}{n},R\right)} \frac{p_r\left(\frac{y}{n} + \lfloor v \rfloor\right)}{2} dv = \int_{B^d\left(0,R\right)} \frac{p_r(u)}{2} du \pm \frac{c_d(\varepsilon)}{r} \mathcal{O}\left(\int_{B^d\left(0,R\right)} \frac{p_r(u/\sqrt{2})(|u|+1)}{2} du\right)$$

$$= \mathbb{P}[\|\mathcal{N}(0,(r/d)\mathrm{Id}_d)\| \le R] \pm \frac{c_d(\varepsilon)}{r} \mathcal{O}\left(1 + \mathbb{E}[\|\mathcal{N}(0,(r/d)\mathrm{Id}_d)\|]\right)$$

$$= 1 - \mathcal{O}(n^{-dC_d^2}) \pm \frac{1}{\sqrt{r}} \mathcal{O}(1)$$

and analogously,

$$\int_{B^d \left(\frac{-y}{n}, R\right)^c} \frac{p_r\left(\frac{y}{n} + \lfloor v \rfloor\right)}{2} dv = \mathbb{P}[\|\mathcal{N}(0, (r/d)\mathrm{Id}_d)\| \ge R](1 + \mathcal{O}(1/r))$$

$$\pm \frac{1}{r} \mathcal{O}\left(\mathbb{E}[\|\mathcal{N}(0, (r/d)\mathrm{Id}_d)\|^2]^{1/2} \mathbb{P}[\|\mathcal{N}(0, (r/d)\mathrm{Id}_d)\| \ge R]^{1/2}\right)$$

$$= \mathcal{O}(n^{-dC_d^2/2}) \left(1 \pm \frac{1}{\sqrt{r}}\right)$$

where we additionally used Cauchy inequality. This in turn implies that

$$\frac{\log^2 n}{n^d} \int_{\varepsilon}^{\frac{t}{n^2}} dr \int_{B^d \left(\frac{-y}{n}, R\right)} \frac{p_r \left(\frac{y}{n} + \lfloor v \rfloor\right)}{2} dv = \mathcal{O}\left(\frac{t}{n^2} \frac{\log^2 n}{n^d}\right) = o(n^{2-d});$$

$$\frac{n^2}{n^d} \int_{\varepsilon}^{\frac{t}{n^2}} dr \int_{B^d \left(\frac{-y}{n}, R\right)^c} \frac{p_r \left(\frac{y}{n} + \lfloor v \rfloor\right)}{2} dv = \mathcal{O}\left(n^{-dC_d^2/2} \frac{t}{n^d}\right) = o(n^{2-d});$$

$$\frac{n^2}{n^d} \int_{\frac{t}{n^2}}^{\frac{t}{n^2}} dr \int_{B^d \left(\frac{-y}{n}, R\right)} \frac{p_r \left(\frac{y}{n} + \lfloor v \rfloor\right)}{2} dv = \mathcal{O}\left(\frac{t - t_*}{n^d}\right) = o(n^{2-d}).$$

Hence, (A.2). Note that the above arguments as well imply that $|J_n(t) - \frac{t_*}{n^d}| = \mathcal{O}(n^{2-d}\sqrt{\frac{t_*}{n^2}}) = \mathcal{O}(n^{2-d}\sqrt{\log n})$.

Let us now prove that $|J_n(t) - \frac{t_*}{n^d}| = \mathcal{O}(n^{2-d})$. By Taylor's theorem for multivariate functions, for $k \in \mathbb{Z}^d$, $x \coloneqq \frac{y}{n} + k$ and $u, w \in B \coloneqq [-1/2, 1/2)^d$:

$$p_r(x) = p_r(x+u) - u^T \nabla p_r(x) - u^T \int_0^1 (1-t)D^2 p_r(x+tu) dt u.$$

Note that the second term on the right-hand side is symmetric w.r.t. the origin as a function in u. Hence, if we integrate both sides w.r.t. u and w over B,

$$p_r(x) - \int_B p_r(x+u) du + \int_B \int_B \int_0^1 (1-t)u^T D^2 p_r(x+tu+w) u dt du dw$$

=
$$\int_B \int_B \int_0^1 (1-t)u^T (D^2 p_r(x+tu+w) - D^2 p_r(x+tu)) u dt du dw.$$

since |B| = 1. Observe that $(D^2 p_r(z))_{i,j} = \left(\frac{d}{r}\right)^2 p_r(z) \left(z_i z_j - \frac{r}{d} \delta_{i,j}\right)$ with the maximal eigenvalue $\left(\frac{d}{r}\right)^2 p_r(z) \left(|z|^2 - \frac{r}{d}\right)$ and the remaining ones equal to $-\frac{d}{r} p_r(z)$. Thus, since $r > \varepsilon > 0$,

$$\left| \int_{B} \int_{0}^{1} (1-t)u^{T} D^{2} p_{r}(z+tu) u \, dt du \right| \leq \int_{B} \int_{0}^{1} |u|^{2} \left(\frac{d}{r}\right)^{2} p_{r}(z+tu) \left(2|z|^{2}+2|u|^{2}+\frac{r}{d}\right) dt du
\leq \left(\frac{d}{r}\right)^{2} \frac{d}{d} \left(2|z|^{2}+\frac{d}{2}+\frac{r}{d}\right) \int_{B} \int_{0}^{1} e^{\frac{d}{2r}t^{2}|u|^{2}} p_{r}\left(\frac{z}{\sqrt{2}}\right) dt du
\leq \left(\frac{d}{r}\right)^{2} \left(2|z|^{2}+\frac{d}{2}+\frac{r}{d}\right) c_{d}(\varepsilon) p_{r}\left(\frac{z}{\sqrt{2}}\right).$$

The latter as a function in z is clearly integrable, hence, by Fubini's theorem

$$\sum_{x \in \frac{y}{n} + \mathbb{Z}^d} \left(\int_B p_r(x+u) du - \int_B \int_B \int_0^1 (1-t)u^T D^2 p_r(x+tu+w) u dt du dw \right)$$
$$= \int_{\mathbb{R}^d} p_r(z) dz - \int_B \int_0^1 (1-t)u^T \int_{\mathbb{R}^d} D^2 p_r(z+tu) dz u dt du = 2.$$

Here we also used that $\frac{1}{2}p_r(z)$ is a density function of $\mathcal{N}(0, \frac{r}{d}\mathrm{Id}_d)$ and so, $\int_{\mathbb{R}^d} p_r(z) \left(z_i z_j - \frac{r}{d}\delta_{i,j}\right) = 2\left(\frac{r}{d}\delta_{i,j} - \frac{r}{d}\delta_{i,j}\right) = 0$. Moreover, by the mean-value theorem, for any $i, j \leq d, z \in \mathbb{R}^d$,

$$\begin{split} &|\left(D^{2}p_{r}(z+w)-D^{2}p_{r}(z)\right)_{i,j}| \leq \sup_{c \in [0,1]} |\nabla(D^{2}p_{r})_{i,j}(z+cw)||w| \\ &= \left(\frac{d}{r}\right)^{3} \sup_{c \in [0,1]} p_{r}(z+cw) \left| \left(-\prod_{l \in \{i,j,k\}} (z+cw)_{l} + \frac{r}{d} \sum_{l \in \{i,j,k\}} (z+cw)_{l} \delta_{\{i,j,k\} \setminus \{l\}}\right)_{k} \right| |w| \\ &\leq \left(\frac{d}{r}\right)^{3} p_{r} \left(\frac{z+w}{\sqrt{2}}\right) \frac{\sqrt{d}}{2} \sup_{c \in [0,1]} e^{\frac{d}{2r}(1-c)^{2}|w|^{2}} \left(|z+cw|^{3} + 3\sqrt{d} \frac{r}{d}|z+cw|\right) \\ &\leq C(d,\varepsilon) \left(\frac{d}{r}\right)^{3} p_{r} \left(\frac{z+w}{\sqrt{2}}\right) \left(|z+w|^{3} + |z+w|^{2} + \frac{r}{d}|z+w| + |z+w| + \frac{r}{d} + 1\right) \\ &\leq \tilde{C}(d,\varepsilon) \left(\frac{d}{r}\right)^{3} p_{r} \left(\frac{z+w}{\sqrt{2}}\right) \left(|z+w|^{3} + \frac{r}{d}|z+w| + \frac{r}{d}\right). \end{split}$$

For us, of interest is z = x + tu. Note that by analogous estimation procedures we can further conclude that

$$p_r\left(\frac{z+w}{\sqrt{2}}\right)\left(|z+w|^3 + \frac{r}{d}|z+w| + \frac{r}{d}\right) \le C(d,\varepsilon)p_r\left(\frac{x+w}{2}\right)\left(|x+w|^3 + \frac{r}{d}|x+w| + \frac{r}{d}\right)$$

This then yields that

$$\begin{split} \sum_{x \in \frac{y}{n} + \mathbb{Z}^d} \int_B \left| \int_0^1 (1 - t) u^T \left(D^2 p_r(x + tu + w) - D^2 p_r(x + tu) \right) u \, \mathrm{d}t \mathrm{d}u \right| \mathrm{d}w \\ & \leq C'(d, \varepsilon) \left(\frac{d}{r} \right)^3 \sum_{x \in \frac{y}{n} + \mathbb{Z}^d} \int_B p_r \left(\frac{x + w}{2} \right) \left(|x + w|^3 + \frac{r}{d} |x + w| + \frac{r}{d} \right) \mathrm{d}w \\ & \leq \hat{C}(d, \varepsilon) \left(\frac{d}{r} \right)^3 \left(\mathbb{E} \left[\left\| \mathcal{N} \left(0, \frac{r}{d} \mathrm{Id}_d \right) \right\|^3 \right] + \frac{r}{d} \mathbb{E} \left[\left\| \mathcal{N} \left(0, \frac{r}{d} \mathrm{Id}_d \right) \right\| \right] + \frac{r}{d} \right) \\ & \leq \bar{C}(d, \varepsilon) r^{-3/2}. \end{split}$$

Altogether, this part combined with (A.2) show that

$$J_n(t) = \frac{n^2}{n^d} \int_{\epsilon}^{\frac{t_*}{n^2}} \left(1 \pm \mathcal{O}(1)r^{-3/2}\right) dr + \mathcal{O}(n^{2-d}) = \frac{t_*}{n^d} \pm \mathcal{O}(n^{2-d})$$

with O(1) uniform in $r > \varepsilon$. This concludes the proof of Claim 2 as desired.

A.2 Bound on $G_{\Lambda}^{0\text{-avg.}}(0,y)$ for $y \in \hat{U}$ in dimension three

Let d=3. Recall that $\hat{\Lambda} \subset [-n/2, n/2)^d \cap \mathbb{Z}^d$ is the canonical projection of Λ onto \mathbb{Z}^d , $\hat{U} := (\partial [-\lfloor n/2 \rfloor, \lfloor n/2 \rfloor]^d) \cap \mathbb{Z}^d$ and U is the corresponding preimage under the canonical projection. Let $y=(y_1,\ldots,y_d) \in U$.

Claim: In dimension d=3 for all $n\in\mathbb{N}$ sufficiently large, $n^{d-2}G_{\Lambda}^{0\text{-avg.}}(0,y)\leq c$ for $c<\frac{3^{2/3}}{2(4\pi)^{2/3}}\frac{a}{c}$.

Proof. Without loss of generality, due to the symmetries of torus we can assume that $y_1 \ge \ldots \ge y_d \ge 0$. In particular, since $y \in \hat{U}$, $y_1 = \lfloor \frac{n}{2} \rfloor$. In A.1 we showed that

$$2d n^{d-2}G_{\Lambda}^{0\text{-avg.}}(0,y) \leq \frac{d}{2\pi^2} + o(1) + c_d \varepsilon^{2-\frac{d}{2}} + \int_{\varepsilon}^{\frac{t_*}{n^2}} dr \left(\sum_{\substack{k \in \mathbb{Z}^3 \\ \forall i: |k_i| \leq C_d \log n}} q_r \left(\frac{y}{n} + k \right) - 1 \right),$$

where
$$t_* = \frac{d \log(n^d)}{2\pi^2} n^2$$
 and $q_r(x) = \left(\frac{d}{2\pi r}\right)^{d/2} e^{-\frac{d|x|^2}{2r}}$.

Let us first verify that even if n is odd, we can set $y_1 = n/2$, which would lead to an error of order at most o(1). Note that since $|k_1| \le C_d \log n$, $r > \varepsilon > 0$,

$$q_r\left(\frac{y}{n} + \left(\frac{1}{2n}, 0, \dots, 0\right) + k\right) = q_r\left(\frac{y}{n} + k\right)e^{-\frac{d}{2r}\left(\frac{1}{(2n)^2} + (k_1 + \frac{y_1}{n})\frac{1}{n}\right)} = q_r\left(\frac{y}{n} + k\right)\left(1 + O\left(\frac{\log n}{\varepsilon n}\right)\right).$$

Since we have already proven in A.1, that $\sum_{k\in\mathbb{Z}^3} q_r(\frac{y}{n}+k) \le 1+\mathcal{O}(1)r^{-3/2} \le c(\varepsilon)$, the above implies that

$$\mathcal{O}\left(\frac{\log n}{\varepsilon n}\right) \int_{\varepsilon}^{\frac{t_*}{n^2}} dr \sum_{\substack{k \in \mathbb{Z}^3 \\ \forall i : |k_i| \le C_d \log n}} q_r \left(\frac{y}{n} + k\right) = \mathcal{O}\left(\frac{\log^2 n}{n}\right) = o(1).$$

Therefore,

$$2d \, n^{d-2} G_{\Lambda}^{0\text{-avg.}}(0,y) \le \frac{d}{2\pi^2} + o(1) + c_d \varepsilon^{2-\frac{d}{2}} + \int_{\varepsilon}^{\frac{t_*}{n^2}} \mathrm{d}r \left(\sum_{k \in \mathbb{Z}^3} q_r \left(\frac{y^*}{n} + k \right) - 1 \right), \tag{A.3}$$

where $y_1^* = n/2$ and the remaining coordinates are the same as of y.

We further show that $\sum_{k\in\mathbb{Z}^3} q_r(\frac{y^*}{n}+k)$ under our previous assumptions on y^* is maximized at $(\frac{1}{2},0,\ldots,0)$. For this notice that $q_r(x)=\prod_{i=1}^d f_r(x_i)$ with f_r being a density function of a centered normally distributed random variable of variance r/d. Hence,

$$\sum_{k \in \mathbb{Z}^3} q_r \left(\frac{y^*}{n} + k \right) = \prod_{i=1}^d \left(\sum_{l \in \mathbb{Z}} f_r \left(\frac{y_i^*}{n} + l \right) \right) =: \prod_{i=1}^d F_r \left(\frac{y_i^*}{n} \right).$$

Note that F_r is a function on a one-dimensional continuous torus $\mathbb{T} = [-1/2, 1/2]/\sim$ (or alternatively, a periodic function on \mathbb{R}). Since furthermore it is in $L^2(\mathbb{T})$, by Carleson's theorem it coincides with its Fourier series almost everywhere, and thus by continuity, everywhere. We have

$$F_r\Big(\frac{y_i^*}{n}\Big) = \sum_{p \in \mathbb{Z}} e^{-\frac{r}{2d}(2\pi)^2 p^2} \cos\Big(2\pi p \frac{y_i^*}{n}\Big) \le \sum_{p \in \mathbb{Z}} e^{-\frac{r}{2d}(2\pi)^2 p^2}$$

^aThis bound would suffice to prove the second claim of point 8 of Proposition 2.14.

with equality reached at $y_i^* = 0$. So, from now on let $y^* = (\frac{1}{2}, 0, \dots, 0)$ (independently of the value of $y \in \hat{U}$).

Let us further investigate the last summand of (A.3). Using the aforementioned Fourier series and noticing that all of them are absolutely convergent and integrable as functions in r over $r > \varepsilon$, we get

$$\begin{split} I &\coloneqq \int_{\varepsilon}^{\frac{t_*}{n^2}} \mathrm{d}r \bigg(\sum_{k \in \mathbb{Z}^3} q_r \Big(\frac{y^*}{n} + k \Big) - 1 \bigg) \\ &= \int_{\varepsilon}^{\frac{t_*}{n^2}} \mathrm{d}r \bigg[\sum_{w \in \mathbb{Z}^2 \setminus \{0\}} e^{-\frac{r}{2d}(2\pi)^2 |w|^2} \bigg(\underbrace{1 - 2 \sum_{l=1}^{\infty} (-1)^{l-1} e^{-\frac{r}{2d}(2\pi)^2 l^2}}_{\in [0,1] \text{ unif. in } r > \varepsilon} \bigg) - 2 \sum_{l=1}^{\infty} (-1)^{l-1} e^{-\frac{r}{2d}(2\pi)^2 l^2} \bigg]. \end{split}$$

It is easy to see that $\int_{t_*/n^2}^{\infty} \sum_{l=1}^{\infty} e^{-\frac{r}{2d}(2\pi)^2 l^2} dr = \mathcal{O}(n^{-1})$. Therefore, the above is bounded by the integral of the integrand over (ε, ∞) plus $\mathcal{O}(n^{-1})$. By Fubini's theorem, the former integral equals to

$$\sum_{w \in \mathbb{Z}^2 \setminus \{0\}} \int_{\varepsilon}^{\infty} \mathrm{d} r e^{-\frac{r}{2d}(2\pi)^2 |w|^2} \left(1 - 2\sum_{l=1}^{\infty} (-1)^{l-1} e^{-\frac{r}{2d}(2\pi)^2 l^2}\right) - 2\sum_{l=1}^{\infty} (-1)^{l-1} \frac{2d}{(2\pi)^2 l^2} e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2}.$$

Note that for a fixed constant M > 0 (to be determined, but independent of ε),

$$\begin{split} & \sum_{\substack{w \in \mathbb{Z}^2 \backslash \{0\} \\ |w| > M/\sqrt{\varepsilon}}} \int_{\varepsilon}^{\infty} \mathrm{d} r e^{-\frac{r}{2d}(2\pi)^2 |w|^2} \bigg(\underbrace{1 - 2\sum_{l=1}^{\infty} (-1)^{l-1} e^{-\frac{r}{2d}(2\pi)^2 l^2}}_{\leq 1} \bigg) \leq \frac{2d}{(2\pi)^2} \sum_{\substack{w \in \mathbb{Z}^2 \backslash \{0\} \\ |w| > M/\sqrt{\varepsilon}}} \frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 |w|^2}}{|w|^2} \\ & \leq \frac{2d}{(2\pi)^2} \bigg(2\pi \int_{\frac{M}{\sqrt{\varepsilon}}}^{\infty} \mathrm{d} s \frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 s^2}}{s} + 4 \int_{\frac{M}{\sqrt{\varepsilon}}}^{\infty} \mathrm{d} s \frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 s^2}}{s^2} \bigg) \leq \frac{(2d)^2 e^{-\frac{M^2}{2d}(2\pi)^2}}{(2\pi)^4 M^2} \bigg(\pi + 2 \frac{\sqrt{\varepsilon}}{M} \bigg). \end{split}$$

Altogether,

$$\begin{split} I &\leq \mathcal{O}\Big(\frac{1}{n}\Big) + \frac{(2d)^2 e^{-\frac{M^2}{2d}(2\pi)^2}}{(2\pi)^4 M^2} \Big(\pi + 2\frac{\sqrt{\varepsilon}}{M}\Big) \\ &+ \frac{2d}{(2\pi)^2} \sum_{\substack{w \in \mathbb{Z}^2 \backslash \{0\} \\ \forall i: |w_i| \leq M/\sqrt{\varepsilon}}} e^{-\frac{\varepsilon}{2d}(2\pi)^2 |w|^2} \bigg[\frac{1}{|w|^2} - 2\sum_{\substack{l=1 \\ \text{odd}}}^{M/\sqrt{\varepsilon}} \left(\frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2}}{|w|^2 + l^2} - \frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 (l+1)^2}}{|w|^2 + (l+1)^2}\right)\bigg] \\ &- 2\frac{2d}{(2\pi)^2} \sum_{\substack{l=1 \\ \text{odd}}}^{\infty} \left(\frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2}}{l^2} - \frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 (l+1)^2}}{(l+1)^2}\right). \end{split}$$

We can write

$$\frac{1}{|w|^2} = 2 \frac{2 + \pi |w| \tanh\left(\frac{\pi |w|}{2}\right) - \pi |w| \coth\left(\frac{\pi |w|}{2}\right)}{4|w|^2} + \pi \frac{\coth\left(\frac{\pi |w|}{2}\right) - \tanh\left(\frac{\pi |w|}{2}\right)}{2|w|}$$

$$= 2 \sum_{k=1}^{\infty} \frac{4k - 1}{(|w|^2 + (2k - 1)^2)(|w|^2 + 4k^2)} + \pi \frac{1}{|w| \sinh(\pi |w|)}$$

$$= 2 \sum_{\substack{l=1 \text{odd}}}^{\infty} \left(\frac{1}{|w|^2 + l^2} - \frac{1}{|w|^2 + (l + 1)^2}\right) + \pi \frac{1}{|w| \sinh(\pi |w|)}$$

using the following series' representations (cf. [GR07, 1.217 1, 1.421 2, 1.421 4]) for the hyperbolic trigonometric functions

$$\tanh(x) = 8x \sum_{k=1}^{\infty} \frac{1}{\pi^2 (2k-1)^2 + 4x^2} \quad \forall \ x \in \mathbb{R};$$
$$\coth(x) = 2x \sum_{k=1}^{\infty} \frac{1}{\pi^2 k^2 + x^2} + \frac{1}{x} \quad \forall \ x \in \mathbb{R} \setminus \{0\}.$$

The proof of these expansions follows directly from [Dun09]. We have shown so far that

$$\begin{split} I &\leq C_{M}\sqrt{\varepsilon} + \frac{(2d)^{2}e^{-\frac{M^{2}}{2d}(2\pi)^{2}}}{(2\pi)^{4}M^{2}}\pi \\ &+ \frac{2d}{(2\pi)^{2}}\sum_{\substack{w \in \mathbb{Z}^{2}\backslash\{0\}\\|w| \leq M/\sqrt{\varepsilon}}} e^{-\frac{\varepsilon}{2d}(2\pi)^{2}|w|^{2}}2\Bigg[\sum_{\substack{l=1\\\text{odd}}}^{M/\sqrt{\varepsilon}} \left(\frac{1-e^{-\frac{\varepsilon}{2d}(2\pi)^{2}l^{2}}}{|w|^{2}+l^{2}} - \frac{1-e^{-\frac{\varepsilon}{2d}(2\pi)^{2}(l+1)^{2}}}{|w|^{2}+(l+1)^{2}}\right) \\ &+ \sum_{\substack{l=1+M/\sqrt{\varepsilon}\\\text{odd}}}^{\infty} \left(\frac{1}{|w|^{2}+l^{2}} - \frac{1}{|w|^{2}+(l+1)^{2}}\right)\Bigg] &=:A_{1} \\ &+ \frac{2d}{(2\pi)^{2}}\sum_{\substack{w \in \mathbb{Z}^{2}\backslash\{0\}\\|w| \leq M/\sqrt{\varepsilon}}} e^{-\frac{\varepsilon}{2d}(2\pi)^{2}|w|^{2}}\pi \frac{1}{|w|\sinh(\pi|w|)} \\ &=:A_{2} \\ &-2\frac{2d}{(2\pi)^{2}}\sum_{\substack{l=1\\\text{odd}}}^{\infty} \left(\frac{e^{-\frac{\varepsilon}{2d}(2\pi)^{2}l^{2}}}{l^{2}} - \frac{e^{-\frac{\varepsilon}{2d}(2\pi)^{2}(l+1)^{2}}}{(l+1)^{2}}\right) \\ &=:A_{3}. \end{split}$$

for an appropriate $C_M > 0$. For the term A_3 , notice that by Fatou's lemma

$$\liminf_{\varepsilon \to 0} \sum_{\substack{l=1 \text{odd}}}^{\infty} \left(\frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2}}{l^2} - \frac{e^{-\frac{\varepsilon}{2d}(2\pi)^2 (l+1)^2}}{(l+1)^2} \right) \ge \sum_{\substack{l=1 \text{odd}}}^{\infty} \left(\frac{1}{l^2} - \frac{1}{(l+1)^2} \right) = \frac{\pi^2}{12}.$$

Therefore, for all $\varepsilon > 0$ very small, there exists $c(\varepsilon) > 0$ which converges to 0 as ε does, such that $A_3 \leq -\frac{2d}{(2\pi)^2} \frac{\pi^2}{6} + c(\varepsilon)$. As for A_2 ,

$$\begin{split} \frac{(2\pi)^2}{2d} A_2 &\leq \sum_{w \in \mathbb{Z}^2 \setminus \{0\}} \frac{\pi}{|w| \sinh(\pi|w|)} \leq 2\pi \int_2^\infty \mathrm{d}s \frac{\pi s}{s \sinh(\pi s)} + \sum_{k=1}^\infty \frac{4\pi}{k \sinh(\pi k)} \\ &+ \frac{4\pi}{|(1,1)| \sinh(\pi|(1,1)|)} + \frac{4\pi}{|(2,2)| \sinh(\pi|(2,2)|)} + \frac{8\pi}{|(1,2)| \sinh(\pi|(1,2)|)} \\ &\leq -2\pi \log(\tanh(\pi)) + \frac{4\pi}{\sinh(\pi)} + \int_1^\infty \mathrm{d}s \frac{2\pi}{\sinh(\pi s)} \\ &+ 4\pi \left(\frac{1}{\sqrt{2} \sinh(\pi \sqrt{2})} + \frac{1}{2\sqrt{2} \sinh(\pi 2\sqrt{2})} + \frac{2}{\sqrt{5} \sinh(\pi \sqrt{5})} \right) \\ &= -2\pi \log(\tanh(\pi)) - 2\log(\tanh(\pi/2)) \\ &+ 4\pi \left(\frac{1}{\sinh(\pi)} + \frac{1}{\sqrt{2} \sinh(\pi \sqrt{2})} + \frac{1}{2\sqrt{2} \sinh(\pi 2\sqrt{2})} + \frac{2}{\sqrt{5} \sinh(\pi \sqrt{5})} \right). \end{split}$$

We have $\frac{(2\pi)^2}{2d}(A_2+A_3) \leq c'(\varepsilon) + 2\pi(0.2411-\pi/12) \leq c'(\varepsilon) - 0.04\pi$. It remains to estimate A_1 . We start by investigating $l \mapsto \frac{1-e^{-\frac{\varepsilon}{2d}(2\pi)^2l^2}}{|w|^2+l^2} =: g(l)$ – of interest to us are its regions of

monotonicity. For $l \geq 1$,

$$\frac{(|w|^2 + l^2)}{2l}g'(l) = e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2} \left(1 + \frac{\varepsilon}{2d} (2\pi)^2 (l^2 + |w|^2) \right) - 1$$

$$\geq \left(1 - \frac{\varepsilon}{2d} (2\pi)^2 l^2 \right) \left(1 + \frac{\varepsilon}{2d} (2\pi)^2 (l^2 + |w|^2) \right) - 1$$

$$= -\left(\frac{\varepsilon(2\pi)^2}{2d} \right)^2 \left(l^4 + l^2 |w|^2 - \frac{2d}{(2\pi)^2 \varepsilon} |w|^2 \right).$$

In particular, the subinterval of $[1, \infty)$ on which g(l) is decreasing ($\{l \geq 1 : g'(l) < 0\}$) is therefore included in the range of solutions of the following inequality

$$l^{4} + l^{2}|w|^{2} \ge \frac{2d}{(2\pi)^{2}\varepsilon}|w|^{2}$$

$$\updownarrow$$

$$l^{2} + \frac{|w|^{2}}{2} \ge |w|^{2}\sqrt{\frac{2d}{(2\pi)^{2}\varepsilon}}\frac{1}{|w|^{2}} + \frac{1}{4}$$

$$\updownarrow l \ge 1$$

$$l \ge |w|\sqrt{\sqrt{\frac{2d}{(2\pi)^{2}\varepsilon}}\frac{1}{|w|^{2}} + \frac{1}{4} - \frac{1}{2}} =: w_{*}.$$

Furthermore, observe that

$$g(l) - g(l+1) = \frac{(2l+1)(1 - e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2}) - (|w|^2 + l^2)(e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2} - e^{-\frac{\varepsilon}{2d}(2\pi)^2 (l+1)^2})}{(|w|^2 + l^2)(|w|^2 + (l+1)^2)}$$

$$\leq \frac{2l+1}{(|w|^2 + l^2)(|w|^2 + (l+1)^2)}.$$

Hence, since on $\{l \ge 1 : g'(l) \ge 0\}$, $g(l) - g(l+1) \le 0$ (up to a borderline value of l),

$$\begin{split} 2\sum_{l=1}^{M/\sqrt{\varepsilon}} \left(\frac{1-e^{-\frac{\varepsilon}{2d}(2\pi)^2 l^2}}{|w|^2+l^2} - \frac{1-e^{-\frac{\varepsilon}{2d}(2\pi)^2(l+1)^2}}{|w|^2+(l+1)^2}\right) + 2\sum_{l=1+M/\sqrt{\varepsilon}}^{\infty} \left(\frac{1}{|w|^2+l^2} - \frac{1}{|w|^2+(l+1)^2}\right) \\ &\leq 2\sum_{l=w_*-1}^{\infty} \frac{2l+1}{(|w|^2+l^2)(|w|^2+(l+1)^2)} \\ &\leq \int_{w_*-1}^{\infty} \frac{(2x+1)\mathrm{d}x}{(|w|^2+x^2)(|w|^2+(x+1)^2)} + \frac{C}{|w|^3} \mathbbm{1}_{\left\{|w| \geq \frac{c}{\sqrt{\varepsilon}}\right\}} \\ &= \frac{1}{|w|} \left[\arctan\left(\frac{w_*}{|w|}\right) - \arctan\left(\frac{w_*-1}{|w|}\right)\right] + \frac{C}{|w|^3} \mathbbm{1}_{\left\{|w| \geq \frac{c}{\sqrt{\varepsilon}}\right\}} \\ &\leq \frac{1}{|w|^2+(w_*-1)^2} + \frac{C}{|w|^3} \mathbbm{1}_{\left\{|w| \geq \frac{c}{\sqrt{\varepsilon}}\right\}} \end{split}$$

for appropriate constants c, C > 0. The second summand in the third line comes from the evaluation of the integrand at its local maximum point (which clearly has the leading term proportional to $|w|^2$) if it is greater or equal to $w_* - 1$. Note further that $w_* - 1 \ge w_*(1 - \sqrt{c}\varepsilon^{1/4})$ for some c > 0 and $|w|^2 + w_*^2(1 - \sqrt{c}\varepsilon^{1/4})^2 \ge |w|^2(1 + (1 - c\sqrt{\varepsilon})(\sqrt{\frac{1}{4} + \frac{2d}{(2\pi)^2\varepsilon}\frac{1}{|w|^2}} - \frac{1}{2})) \ge |w|^2(1 + (1 - c\sqrt{\varepsilon})(1 + c\sqrt{c}\varepsilon^{1/4})^2)$

²An explicit computation shows that the pre-coefficient of the leading term of the local maximum point is $1/\sqrt{3}$.

 $\frac{1}{2}(|w|^2 + 2\frac{1-c\sqrt{\varepsilon}}{1+c\sqrt{\varepsilon}}\sqrt{\frac{2d}{(2\pi)^2\varepsilon}}|w|)$. For simplicity set $a := 2\frac{1-c\sqrt{\varepsilon}}{1+c\sqrt{\varepsilon}}\sqrt{\frac{2d}{(2\pi)^2\varepsilon}} \approx 2\sqrt{\frac{2d}{(2\pi)^2\varepsilon}}$. It is now possible to estimate A_1 ,

$$\frac{(2\pi)^2}{2d} A_1 \leq \sum_{\substack{w \in \mathbb{Z}^2 \setminus \{0\} \\ |w| \leq M/\sqrt{\varepsilon}}} \left[\frac{2}{|w|^2 + a|w|} + \frac{C}{|w|^3} \mathbb{1}_{\{|w| \geq \frac{c}{\sqrt{\varepsilon}}\}} \right]
\leq 2\pi \int_1^{\frac{M}{\sqrt{\varepsilon}}} \frac{2r dr}{r^2 + ar} + 4 \int_1^{\infty} \frac{2dr}{r^2 + ar} + C' \int_{\frac{c}{\sqrt{\varepsilon}}}^{\infty} \frac{r dr}{r^3} + \frac{8}{1+a}
\leq C\sqrt{\varepsilon} + 4\pi \log\left(\frac{a + M/\sqrt{\varepsilon}}{1+a}\right) + 8 \frac{\log(a+1)}{a}
\leq 4\pi \log\left(1 + \frac{2\pi M}{2\sqrt{2d}}\right) + c(\varepsilon)$$

with $c(\varepsilon) \to 0$ as $\varepsilon \to 0$. To sum up, for all n sufficiently large,

$$n^{d-2}G_{\Lambda}^{0-\text{avg.}}(0,y) \le \frac{1 - 0.04\pi + 4\pi \log\left(1 + \frac{2\pi M}{2\sqrt{2d}}\right) + \pi \frac{2d}{(2\pi)^2 M^2} e^{-\frac{M^2}{2d}(2\pi)^2}}{(2\pi)^2} + c(\varepsilon) + c_d \varepsilon^{2-\frac{d}{2d}}(\varepsilon)$$

with $c(\varepsilon) > 0$ that can be made arbitrarily small by choosing $\varepsilon > 0$ sufficiently small (but non-zero). Note that since d = 3, the last summand is as well arbitrary small. Let us choose M > 0 such that $\frac{2\pi M}{\sqrt{2d}} = 1$. Then,

$$n^{d-2}G_{\Lambda}^{0-\text{avg.}}(0,y) \le \frac{1 - 0.04\pi + 4\pi \log(3/2) + \pi e^{-1}}{(2\pi)^2} + c(\varepsilon) < \frac{3^{2/3}}{2(4\pi)^{2/3}}.$$
 (A.4)

A.3 Proof of Proposition 4.3

Let us recall the result that we want to prove.

Proposition A.1. Let $(\mathbf{X}^i)_i$ be a sequence of i.i.d. centered random vectors with the probability density function (w.r.t. Lebesgue measure on \mathbb{R}^k) $f: \mathbb{R}^k \to \mathbb{R}$. Assume that

- 1. $f \in L^r(\mathbb{R}^k)$ for some $r \in (1,2]$;
- 2. all the entries of the covariance matrix C of $\mathbf{X}^1 = (X_j^1)_{j=1}^k$ are well-defined and finite, or equivalently, for all $1 \leq j \leq k$, $X_j^1 \in L^2(\mathbb{P})$;
- 3. C is positive definite.

Then the relation

$$n^{k/2} f^{(n)}(\sqrt{n} \boldsymbol{x}) \xrightarrow{n \to \infty} \frac{1}{(2\pi)^{k/2} \sqrt{\det(C)}} e^{-\frac{1}{2}(\boldsymbol{x}, C^{-1} \boldsymbol{x})}$$

holds uniformly w.r.t. \mathbf{x} in \mathbb{R}^k . Here, $f^{(n)}$ is the n-fold convolution of f, as well as the density function of $\sum_{i=1}^n \mathbf{X}^i$.

The following proof follows the lines of the one for the case k=1 in [GK68, Chp.8 §46 Theorem 1].

Proof. Let ψ denote the characteristic function of \mathbf{X}^1 , and ψ_n the characteristic function of $\sum_{i=1}^{n} \mathbf{X}^i$. In particular,

$$\psi(t) = \int_{\mathbb{R}^k} e^{it \cdot x} f(x) dx$$
 and $\psi_n(t) = \int_{\mathbb{R}^k} e^{it \cdot x} f^{(n)}(x) dx = (\psi(t))^n$.

By assumption f belongs to $L^r(\mathbb{R}^k)$ and to $L^1(\mathbb{R}^k)$ as a density function. Therefore, also $f \in L^u(\mathbb{R}^k)$ for any $u \in (1,r)$. This can be deduced from Hölder's inequality as follows. Let $\theta = \frac{r}{r-1} \frac{u-1}{u} \in (0,1), \ p = \frac{r-1}{u-1} \in (1,\infty)$ and p' = p/(p-1) such that $\frac{1}{p} + \frac{1}{p'} = 1$. Then by Hölder's inequality,

$$\int_{\mathbb{R}^k} |f|^u \mathrm{d}x \le \||f|^{u\theta}\|_{L^p} \||f|^{u(1-\theta)}\|_{L^{p'}} = \|f\|_{L^r}^{r/p} \|f\|_{L^1}^{1/p'} < \infty.$$

Let $\hat{f}(t) \coloneqq \int_{\mathbb{R}^k} e^{-2\pi i(\boldsymbol{x},t)} f(\boldsymbol{x}) d\boldsymbol{x}$ be the Fourier transform of f. Then by Hausdorff-Young inequality (see, e.g., [Gra14, Proposition 2.2.16]) we know that if $f \in L^p(\mathbb{R}^k)$ for some $p \in [1,2]$, then $\hat{f} \in L^{p'}(\mathbb{R}^k)$ for $p' = \frac{p}{p-1}$. Thus, the above observations about f and the fact that $\hat{f}(t) = \psi(-2\pi t)$ yield that ψ is in $L^q(\mathbb{R}^k)$ for any $q \ge \frac{r}{r-1}$. In particular, we get that for any $n \ge \frac{r}{r-1}$, ψ_n is integrable. Hence, by inversion formula theorem,

$$f^{(n)}(\boldsymbol{x}) = \frac{1}{(2\pi)^k} \int_{\mathbb{R}^k} e^{-i(\boldsymbol{x},\boldsymbol{t})} \psi_n(\boldsymbol{t}) d\boldsymbol{t},$$

and thus,

$$n^{k/2} f^{(n)}(\sqrt{n} \boldsymbol{x}) = \frac{1}{(2\pi)^k} \int_{\mathbb{R}^k} e^{-i(\boldsymbol{x}, \boldsymbol{t})} \psi_n(\boldsymbol{t}/\sqrt{n}) d\boldsymbol{t}.$$

On the other hand, we know that

$$\frac{1}{(2\pi)^{k/2}\sqrt{\det(C)}} e^{-\frac{1}{2}(\boldsymbol{x},C^{-1}\boldsymbol{x})} = \frac{1}{(2\pi)^k} \int_{\mathbb{R}^k} e^{-i(\boldsymbol{x},t) - \frac{1}{2}(\boldsymbol{t},C\boldsymbol{t})} d\boldsymbol{t}.$$

Therefore, to conclude the proof it suffices to show that

$$R_n(\boldsymbol{x}) \coloneqq \int_{\mathbb{R}^k} e^{-i(\boldsymbol{x},\boldsymbol{t})} \left(\psi_n(\boldsymbol{t}/\sqrt{n}) - e^{-\frac{1}{2}(\boldsymbol{t},C\boldsymbol{t})} \right) d\boldsymbol{t}$$

converges to zero as n tends to infinity uniformly over $\boldsymbol{x} \in \mathbb{R}^k$. By central limit theorem (multivariate version) and dominated convergence theorem ($\|\psi_n\|_{L^{\infty}} \leq 1$), for any r > 0 fixed,

$$\left| \int_{B^k(0,r)} e^{-i(\boldsymbol{x},\boldsymbol{t})} \left(\psi_n(\boldsymbol{t}/\sqrt{n}) - e^{-\frac{1}{2}(\boldsymbol{t},C\boldsymbol{t})} \right) d\boldsymbol{t} \right| \xrightarrow{n \to \infty} 0 \quad \text{uniformly over } \boldsymbol{x}.$$

Moreover, by taking r sufficiently large we can make $\left| \int_{|t|>r} e^{-i(x,t)-\frac{1}{2}(t,Ct)} dt \right|$ arbitrarily small. It is only left to treat the term

$$I := \int_{|\mathbf{t}| \searrow r} e^{-i(\mathbf{x}, \mathbf{t})} \psi_n(\mathbf{t}/\sqrt{n}) d\mathbf{t},$$

which we further split into two integrals I_1 and I_2 over $r < |\mathbf{t}| \le \varepsilon \sqrt{n}$ and $|\mathbf{t}| > \varepsilon \sqrt{n}$, respectively, for some $\varepsilon > 0$ small. Let us start with I_1 . By our assumptions, ψ is twice differentiable with

$$\nabla \psi(\mathbf{0}) = \mathbf{0}, \quad D^2 \psi(\mathbf{0}) = -C.$$

Hence, by Taylor's theorem, in the neighbourhood of 0,

$$\psi(t) - 1 = (\nabla \psi(\mathbf{0}))^T t + \frac{1}{2} t^T (D^2 \psi(\mathbf{0})) t + o(|t|^2) = -\frac{1}{2} t^T C t + o(|t|^2).$$

By assumption, C is positive definite, so it is possible to choose $\varepsilon > 0$ sufficiently small such that for all $|t| \le \varepsilon$, the remainder of the expansion $o(|t|^2)$ is smaller than $\frac{1}{4}t^TCt$. Then, $|\psi(t)| \le 1 - \frac{1}{4}t^TCt \le e^{-\frac{1}{4}t^TCt}$ and

$$|I_1| \le \int_{r < |\boldsymbol{t}| \le \varepsilon \sqrt{n}} |\psi_n(\boldsymbol{t}/\sqrt{n})| d\boldsymbol{t} \le \int_{r < |\boldsymbol{t}| \le \varepsilon \sqrt{n}} e^{-n\frac{1}{4n}\boldsymbol{t}^T C \boldsymbol{t}} d\boldsymbol{t} \le \int_{r < |\boldsymbol{t}|} e^{-\frac{1}{4}\boldsymbol{t}^T C \boldsymbol{t}} d\boldsymbol{t}.$$

Thus, by taking r sufficiently large we can make $|I_1|$ arbitrarily small. Let us finally consider I_2 . Recall that $\hat{f}(t) = \psi(-2\pi t)$, and thus by Riemann-Lebesgue lemma (see [Gra14, Proposition 2.2.17]), $|\psi(t)| \to 0$ as $|t| \to \infty$. Moreover, $|\psi(t)| < 1$ for $t \neq 0$. Indeed, suppose first that there exists a non-zero t such that $\psi(t) = 1$. Then the following must hold:

$$\int \underbrace{(1 - \cos((t, x)))}_{>0} f(x) dx = \int \underbrace{(1 - \sin((t, x)))}_{>0} f(x) dx = 0.$$

And hence for $f(\boldsymbol{x})d\boldsymbol{x}$ -a.e. $\boldsymbol{x}, 1-\cos((\boldsymbol{t},\boldsymbol{x}))=0$, which is clearly impossible unless \boldsymbol{t} is zero. Suppose now that $|\psi(\boldsymbol{t})|=1$, and consider $\mathbf{Z}:=\mathbf{X}^1-\mathbf{X}^2$. Then $\psi_Z(\boldsymbol{t})=1$, and we get that for $f_Z(\boldsymbol{x})d\boldsymbol{x}$ -a.e. $\boldsymbol{x}, 1-\cos((\boldsymbol{t},\boldsymbol{x}))=0$. This again implies that $\boldsymbol{t}=0$. Altogether, we proved that there exists c>0 such that $|\psi(\boldsymbol{t})|< e^{-c}$ for all $|\boldsymbol{t}|>\varepsilon$. Thus, for a fixed $q>\frac{r}{r-1}$,

$$|I_2| \le e^{-c(n-q)} \int_{|\boldsymbol{t}| > \varepsilon\sqrt{n}} |\psi(\boldsymbol{t}/\sqrt{n})|^q d\boldsymbol{t} = \underbrace{e^{-c(n-q)} n^{k/2}}_{n \to \infty} \underbrace{\int_{|\boldsymbol{t}| > \varepsilon} |\psi(\boldsymbol{t})|^q d\boldsymbol{t}}_{\le \infty}.$$

A.4 Proof of Proposition 3.3

We want to prove the following.

Proposition A.2. Let $(X_{i,n})_{i \leq n,n \in \mathbb{N}}$ be a triangular array of independent centered random variables with the probability density functions $f_{i,n}$ (w.r.t. Lebesgue measure on \mathbb{R}). Assume that

- 1. $f_{i,n} \in L^r(\mathbb{R})$ for some $r \in (1,2]$ (independent of i,n) such that $\sup_{i,n} ||f_{i,n}||_{L^r} \leq M$ for some M > 0;
- 2. for all $i \leq n, n \in \mathbb{N}$, $\sigma_{i,n}^2 := \operatorname{Var}[X_{i,n}] < \infty$ ordered such that $\sigma_{1,n}^2 \geq \ldots \geq \sigma_{n,n}^2$;
- 3. Lindeberg's condition is satisfied, that is, for any $\varepsilon > 0$,

$$\frac{\sum_{i\geq 1} \mathbb{E}[X_{i,n}^2 \mathbb{1}_{\{|X_{i,n}| > \varepsilon s_n\}}]}{s_n^2} \xrightarrow{n \to \infty} 0,$$

where $s_n^2 := \sum_{i=1}^n \sigma_{i,n}^2$.

4. there exist $\delta > 0$ (uniform), $K(n) \geq 1$, $l_*(n) \geq 1$, $n \geq l^*(n) \geq 2\lceil \frac{r}{r-1} \rceil$ such that for all n sufficiently large

$$(a) \frac{\sum_{i \ge l_*(n)} \sigma_{i,n}^2}{\sum_{i=1}^n \sigma_{i,n}^2} \ge \delta;$$

$$(b) \frac{\sum_{i \ge l_*(n)} \mathbb{E}[X_{i,n}^2 \mathbb{1}_{\{|X_{i,n}| > K(n)\}}]}{\sum_{i \ge l_*(n)} \sigma_{i,n}^2} \le \frac{1}{8};$$

$$(c) \ \frac{n-l^*(n)}{\sigma_{l^*,n}^2 \vee K(n)^2} \gg \log\Big(\sum_{i=1}^n \sigma_{i,n}^2\Big).$$

Then the relation

$$s_n f^{(n)}(s_n x) \xrightarrow{n \to \infty} \frac{1}{\sqrt{2\pi}} e^{-\frac{x^2}{2}}$$

holds uniformly over \mathbb{R} . Here, $f^{(n)}$ is the convolution of $f_{1,n},\ldots,f_{n,n}$, as well as the density function of $\sum_{i=1}^{n} X_{i,n}$. Note that $s_n f^{(n)}(s_n x)$ is the density of $\frac{\sum_{i=1}^{n} X_{i,n}}{s_n}$.

The following proof is a modification of the one given in A.3.

Proof. Let $\psi_{i,n}$ denote the characteristic function of $X_{i,n}$, and $\psi^{(n)}$ the characteristic function of $\sum_{i=1}^n X_{i,n}$, $\hat{g}(t) := \int_{\mathbb{R}} e^{-2\pi i x t} g(x) dx$ be the Fourier transform of $g: \mathbb{R} \to \mathbb{R}$ (for an appropriate g). With the same argument as in A.3, $\hat{f}_{i,n}, \psi_{i,n} \in L^q(\mathbb{R})$ for any $q \geq \frac{r}{r-1}$. In particular, by Hölder's inequality, for any fixed $n \geq \frac{r}{r-1}$,

$$\|\psi^{(n)}\|_{L^1} = \left\|\prod_{i=1}^n \psi_{i,n}\right\|_{L^1} \le \prod_{i=1}^n \|\psi_{i,n}\|_{L^n} < \infty.$$

Hence, by inversion formula theorem, $f^{(n)}(x) = \frac{1}{2\pi} \int_{\mathbb{R}} e^{-ixt} \psi^{(n)}(t) dt$. To conclude the proof it suffices to show that

$$D_n(\boldsymbol{x}) := \int_{\mathbb{R}} e^{-ixt} \left(\psi^{(n)}(t/s_n) - e^{-\frac{t^2}{2}} \right) dt$$

converges to zero as n tends to infinity uniformly over $x \in \mathbb{R}$. We first split D_n into two integrals over $|t| \leq r$ and |t| > r for some r > 0 (to be specified later). By Lindeberg-Feller central limit theorem and dominated convergence theorem ($\|\psi_n\|_{L^{\infty}} \leq 1$), the former converges to zero as n tends to infinity uniformly in $x \in \mathbb{R}$. By taking r sufficiently large we can also make $\left|\int_{|t|>r} e^{-ixt-\frac{t^2}{2}} \mathrm{d}t\right|$ arbitrarily small. The remaining part $\int_{|t|>r} e^{-ixt} \psi^{(n)}(t/s_n) \mathrm{d}t$ we further split into two integrals I_1 and I_2 over $r < |t| \leq \delta_n s_n$ and $|t| > \delta_n s_n$, respectively, for some $\delta_n > 0$ small (to be chosen later in dependence of K(n)). By our assumptions, $\psi'_{i,n}(0) = 0$ and $\psi''_{i,n}(0) = -\sigma^2_{i,n}$ for all i, n. Hence, by Taylor's theorem, in the neighbourhood of 0 ($|t| \leq \delta_n$),

$$\psi_{i,n}(t) = 1 - \frac{\sigma_{i,n}^2 t^2}{2} + R_{i,n}(t).$$

Since $e^{ix} = 1 + ix - \frac{x^2}{2} + R(x)$ with $|R(x)| \le \min(|x|^3/6, x^2)$, the remainder term $R_{i,n}(t)$ in the above expansion of $\psi_{i,n}$ satisfies

$$|R_{i,n}(t)| \leq \frac{1}{6} \mathbb{E}[|tX_{i,n}|^3 \mathbb{1}_{\{|X_{i,n}| \leq K\}}] + \mathbb{E}[|tX_{i,n}|^2 \mathbb{1}_{\{|X_{i,n}| \geq K\}}]$$

$$\leq \frac{K\delta_n}{6} t^2 \sigma_{i,n}^2 + t^2 \mathbb{E}[|X_{i,n}|^2 \mathbb{1}_{\{|X_{i,n}| \geq K\}}]$$

for any K>0. In particular, also for K(n) as in the fourth assumption. By choosing $\delta_n=\frac{3}{4K(n)}>0$, we obtain that $|\psi_{i,n}(t)|\leq 1-\frac{3}{8}\sigma_{i,n}^2t^2+t^2\mathbb{E}[|X_{i,n}|^2\mathbb{1}_{\{|X_{i,n}|\geq K(n)\}}]\leq \exp(-\frac{3}{8}\sigma_{i,n}^2t^2+t^2\mathbb{E}[|X_{i,n}|^2\mathbb{I}_{\{|X_{i,n}|\geq K(n)\}}]$

 $t^2 \mathbb{E}[|X_{i,n}|^2 \mathbb{1}_{\{|X_{i,n}| \ge K(n)\}}])$ for all $|t| \le \delta_n$. Hence,

$$|I_{1}| \leq \int_{r < |t| \leq \delta_{n} s_{n}} |\psi^{(n)}(t/s_{n})| dt \leq \int_{r < |t| \leq \delta_{n} s_{n}} \prod_{i=l_{*}(n)}^{n} e^{-\frac{3}{8} \frac{\sigma_{i,n}^{2}}{s_{n}^{2}} t^{2} + t^{2} \frac{\mathbb{E}[|X_{i,n}|^{2} \mathbb{I}_{\{|X_{i,n}| \geq K(n)\}}]}{s_{n}^{2}}} dt$$

$$\leq \int_{r < |t| \leq \delta_{n} s_{n}} \exp\left(-t^{2} \underbrace{\frac{\sum_{i \geq l_{*}(n)} \sigma_{i,n}^{2}}{s_{n}^{2}}}_{\geq \delta} \underbrace{\left[\frac{3}{8} - \frac{\sum_{i \geq l_{*}(n)} \mathbb{E}[|X_{i,n}|^{2} \mathbb{1}_{\{|X_{i,n}| \geq K(n)\}}]}{\sum_{i \geq l_{*}(n)} \sigma_{i,n}^{2}}\right]}\right) dt$$

$$\leq \int_{r < |t|} \exp(-t^{2} \delta/4) dt$$

By further increasing r, we can make $|I_1|$ arbitrarily small. Consider I_2 . Let $\mathcal{M}_{i,n}$ be the median of $f_{i,n}(X_{i,n})$, i.e., $\mathbb{P}[f_{i,n}(X_{i,n}) \geq \mathcal{M}_{i,n}] \geq 1/2$ and $\mathbb{P}[f_{i,n}(X_{i,n}) \leq \mathcal{M}_{i,n}] \geq 1/2$. By [BCG12, Theorem 2], there exist two absolute constants $c_1, c_2 > 0$ such that

$$|\psi_{i,n}(t)| < \begin{cases} 1 - c_1/(\mathcal{M}_{i,n}^2 \sigma_{i,n}^2), & |t| \ge \pi/(4\sigma_{i,n}); \\ 1 - c_2 t^2/\mathcal{M}_{i,n}^2, & 0 < |t| < \pi/(4\sigma_{i,n}). \end{cases}$$

Note that since $f_{i,n}(X_{i,n})$ is positive, by Markov's inequality for any $\lambda > 0$,

$$\mathbb{P}[f_{i,n}(X_{i,n}) \ge \lambda] \le \frac{\mathbb{E}[f_{i,n}(X_{i,n})^{r-1}]}{\lambda^{r-1}} = \frac{\int f_{i,n}^r(x) dx}{\lambda^{r-1}} \le \frac{M^r}{\lambda^{r-1}},$$

which directly implies that $0 < \mathcal{M}_{i,n} \le (2M^r)^{1/(r-1)} =: C^{1/2}$. Thus, for all $i \le n, n \in \mathbb{N}$:

$$|\psi_{i,n}(t)| < \begin{cases} 1 - c_1/(C\sigma_{i,n}^2), & |t| \ge \pi/(4\sigma_{i,n}); \\ 1 - c_2\delta_n^2/C, & \delta_n < |t| < \pi/(4\sigma_{i,n}); \end{cases}$$

and $\sup_{i\geq l^*(n)} |\psi_{i,n}(t)| < 1 - \min(c_1/(C\sigma_{l^*,n}^2), c_2\delta_n^2/C) \le e^{-c'\min(\sigma_{l^*,n}^{-2},\delta_n^2)}$ for some c'>0 and all $|t|\geq \delta_n$. Thus, for any fixed integer $2\lceil \frac{r}{r-1}\rceil > q > \frac{r}{r-1}$,

$$|I_{2}| \leq s_{n} \int_{|t| > \delta_{n}} \prod_{i=1}^{n} |\psi_{i,n}(t)| dt \leq e^{-(n-l^{*})c' \min(\sigma_{l^{*},n}^{-2},\delta_{n}^{2})} s_{n} \int_{|t| > \delta_{n}} \prod_{i=1}^{q} |\psi_{i,n}(t)| dt$$

$$\leq e^{-(n-l^{*})c' \min(\sigma_{l^{*},n}^{-2},\delta_{n}^{2})} s_{n} \prod_{i=1}^{q} \|\psi_{i,n}\|_{L^{q}} \leq e^{-(n-l^{*})c' \min(\sigma_{l^{*},n}^{-2},\delta_{n}^{2})} s_{n} \prod_{i=1}^{q} (2\pi)^{1/q} \|f_{i,n}\|_{L^{q/(q-1)}}$$

$$\leq e^{-(n-l^{*})c' \min(\sigma_{l^{*},n}^{-2},\delta_{n}^{2})} s_{n} 2\pi \prod_{i=1}^{q} \left[\|f_{i,n}\|_{L^{r}}^{r/(q(r-1))} \|f_{i,n}\|_{L^{1}}^{1-r/(q(r-1))} \right]$$

$$\leq 2\pi e^{-(n-l^{*})c' \min(\sigma_{l^{*},n}^{-2},\delta_{n}^{2})} s_{n} M^{r/(r-1)} \xrightarrow[\text{unif in } r]{n \to \infty} 0,$$

where we used Hölder's and Hausdorff-Young inequalities and the third assumption $(\delta_n = c/K(n))$.

References

- [Abä17] Angelo Abächerli. Local picture and level-set percolation of the Gaussian free field on a large discrete torus. Stochastic Processes and their Applications, 129, 07 2017.
- [AF02] David Aldous and James Allen Fill. Reversible Markov Chains and Random Walks on Graphs, 2002. Unfinished monograph, recompiled 2014, available at http://www.stat.berkeley.edu/\$\sim\$aldous/RWG/book.html.

- [AGS22] Juhan Aru, Christophe Garban, and Avelio Sepúlveda. Percolation for 2D classical Heisenberg model and exit sets of vector valued GFF. *ArXiv eprints*, 12 2022.
- [BCG12] Sergey Bobkov, Gennadiy Chistyakov, and Friedrich Götze. Bounds for characteristic functions in terms of quantiles and entropy. *Electronic Communications* in *Probability*, 17(none):1 9, 2012.
- [BD87] J.G. Brankov and D.M. Danchev. On the limit Gibbs states of the spherical model. *Journal of Physics A: Mathematical and General*, 20(14):4901, oct 1987.
- [BGM⁺13] J. M. Borwein, M. L. Glasser, R. C. McPhedran, J. G. Wan, and I. J. Zucker. Lattice Sums Then and Now. Encyclopedia of Mathematics and its Applications. Cambridge University Press, 2013.
- [Bil95] P. Billingsley. *Probability and Measure*. Wiley Series in Probability and Statistics. Wiley, 1995.
- [BK52] T. H. Berlin and M. Kac. The Spherical Model of a Ferromagnet. *Phys. Rev.*, 86:821–835, Jun 1952.
- [CSZ19] A. Crisanti, A. Sarracino, and M. Zannetti. Condensation versus ordering: From the spherical models to Bose-Einstein condensation in the canonical and grand canonical ensemble. *Phys. Rev. Res.*, 1:023022, Sep 2019.
- [Cur42] J. H. Curtiss. A Note on the Theory of Moment Generating Functions. The Annals of Mathematical Statistics, 13(4):430 433, 1942.
- [DH18] Sayan Das and Rajat Subhra Hazra. Extremal process of the zero-average Gaussian Free Field for $d \geq 3$, 2018.
- [Dun09] William Dunham. When Euler Met l'Hôpital. *Mathematics Magazine*, 82(1):16–25, 2009.
- [FV17] Sacha Friedli and Yvan Velenik. Statistical Mechanics of Lattice Systems: A Concrete Mathematical Introduction. Cambridge University Press, 2017.
- [GB68] J. D. Gunton and M. J. Buckingham. Condensation of the Ideal Bose Gas as a Cooperative Transition. *Phys. Rev.*, 166:152–158, Feb 1968.
- [GK68] B.V. Gnedenko and A.N. Kolmogorov. Limit Distributions for Sums of Independent Random Variables. Addison-Wesley Mathematical Series. Addison-Wesley, 1968.
- [GP93] J. Gough and J.V. Pulé. The spherical model and Bose-Einstein condensation. Helvetica Physica Acta, 66(1), 1993.
- [GR07] I.S. Gradshteyn and I.M. Ryzhik. 1 Elementary Functions. In Alan Jeffrey, Daniel Zwillinger, I.S. Gradshteyn, and I.M. Ryzhik, editors, *Table of Integrals, Series, and Products (Seventh Edition)*, pages 25–62. Academic Press, Boston, seventh edition edition, 2007.
- [Gra14] Loukas Grafakos. Classical Fourier analysis, volume Graduate texts in mathematics. Springer, New York, third edition edition, 2014.
- [HJ12] R.A. Horn and C.R. Johnson. *Matrix Analysis*. Cambridge University Press, 2012.

- [KT71] M. Kac and C.J. Thompson. Spherical model and the infinite spin dimensionality limit. *Physica Norvegica*, 5:163–168, 1971.
- [KT77] Mark Kac and Colin J. Thompson. Correlation functions in the spherical and mean spherical models. *Journal of Mathematical Physics*, 18(8):1650–1653, 08 1977.
- [Kup80] Antti J. Kupiainen. On the 1/n expansion. Communications in Mathematical Physics, 73(3):273 294, 1980.
- [Law12] G.F. Lawler. *Intersections of Random Walks*. Modern Birkhäuser Classics. Springer New York, 2012.
- [Lax55] Melvin Lax. Relation Between Canonical and Microcanonical Ensembles. *Phys. Rev.*, 97:1419-1420, Mar 1955.
- [LL10] Gregory F. Lawler and Vlada Limic. Random Walk: A Modern Introduction. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2010.
- [LP17] D.A. Levin and Y. Peres. *Markov Chains and Mixing Times*. MBK. American Mathematical Society, 2017.
- [Luk20] Jani Lukkarinen. Multi-state Condensation in Berlin–Kac Spherical Models. Communications in Mathematical Physics, 373:389–433, 2020.
- [LW52] H. W. Lewis and G. H. Wannier. Spherical Model of a Ferromagnet. *Phys. Rev.*, 88:682–683, Nov 1952.
- [LW53] H. W. Lewis and G. H. Wannier. Spherical Model of a Ferromagnet. *Phys. Rev.*, 90:1131–1131, Jun 1953.
- [Mad19] Erwin Madelung. Das elektrische Feld in Systemen von regelmäßig angeordneten Punktladungen. *Physikalische Zeitschrift*, 19:524–533, 1919.
- [Mas07] Pascal Massart. Concentration Inequalities and Model Selection, Ecole d'Eté de Probabilités de Saint-Flour XXXIII 2003. Lecture Notes in Mathematics Springer-verlag, 1896, 01 2007.
- [Pol75] A.M. Polyakov. Interaction of goldstone particles in two dimensions. Applications to ferromagnets and massive Yang-Mills fields. *Physics Letters B*, 59(1):79–81, 1975.
- [Rod17] Pierre-François Rodriguez. A 0-1 law for the massive Gaussian free field. *Probability Theory and Related Fields*, 169, 12 2017.
- [Shc88] M. V. Shcherbina. Spherical limit of n-vector correlations. *Theoretical and Mathematical Physics*, 77:1323–1331, 1988.
- [SM75] Yu.N. Sudarev S.A. Molchanov. Gibbs states in the spherical model. *Dokl. Akad. Nauk SSSR*, 224:536–539, 1975.
- [Sta68a] H. E. Stanley. Dependence of Critical Properties on Dimensionality of Spins. *Phys. Rev. Lett.*, 20:589–592, Mar 1968.
- [Sta68b] H. E. Stanley. Spherical Model as the Limit of Infinite Spin Dimensionality. *Phys. Rev.*, 176:718–722, Dec 1968.

- [Szn91] Alain-Sol Sznitman. Topics in propagation of chaos. In Paul-Louis Hennequin, editor, *Ecole d'Eté de Probabilités de Saint-Flour XIX* 1989, pages 165–251, Berlin, Heidelberg, 1991. Springer Berlin Heidelberg.
- [YW04] C. C. Yan and G. H. Wannier. Observations on the Spherical Model of a Ferromagnet. *Journal of Mathematical Physics*, 6(11):1833–1838, 12 2004.