HIGHER COMPLEX SOBOLEV SPACES ON COMPLEX MANIFOLDS

THAI DUONG DO¹ AND DUC-BAO NGUYEN²

ABSTRACT. We study higher complex Sobolev spaces and their corresponding functional capacities. In particular, we prove the Moser-Trudinger inequality for these spaces and discuss some relationships between these spaces and the complex Monge-Ampère equation.

Classification AMS 2020: 32Uxx, 32W20, 46E35.

Keywords: complex Sobolev space, plurisubharmonic function, Moser-Trudinger inequality, Monge-Ampère equation, positive closed current.

CONTENTS

| 1. Introduction | 1 |
|----------------------------------|----|
| 2. Higher complex Sobolev space | 4 |
| 3. Moser-Trudinger inequalities | 13 |
| 4. Complex Monge-Ampère equation | 26 |
| Acknowledgments | 33 |
| References | 33 |

1. INTRODUCTION

In this paper, we study higher complex Sobolev spaces introduced by Dinh which extend the notion of complex Sobolev space introduced earlier by himself and Sibony in [DS06a]. The complex Sobolev space has been systematically studied by Vigny in [Vig07]. The key observation is that this space takes into account the complex structure of the ambient space and is stable under holomorphic transformations; thus, one could view it as a tailored version of the classical Sobolev space to the complex setting. As a consequence, this functional space plays a key role in complex dynamics and it leads to many fruitful applications in this field (e.g., see [BD22, DKW21, DS06a, DV24, Vig15, Vu20]). Not limited to the theory of complex dynamics, the complex Sobolev spaces also find applications in other fields of mathematics. For example, in the study of the complex Monge-Ampère equation, this space has been used as a test space to obtain the necessary and sufficient conditions for whether a given measure is a Monge-Ampère measure with Hölder potentials (see [DKN20, DMV20]). Moreover, it has recently found applications in studying uniform diameter bound for Kähler metrics (see [Vu23]). Due to its importance, the complex Sobolev space has attracted a lot of attention from several mathematicians (see[DMV20, Vig07, VV23, Vu23]).

The classical Sobolev spaces $W^{n,p}$ are important tools in studying differential equations and are usually used as solution spaces for many fundamental equations. An important

Date: May 13, 2024.

fact is that when n and p go to infinity, the solution gains more regularity and becomes the ancient solution.

We study higher complex Sobolev spaces in both local (i.e., on bounded domains of \mathbb{C}^k) and global (i.e., on compact Kähler manifolds) settings. Let Ω be a bounded domain in \mathbb{C}^k , we denote by $PSH(\Omega)$ the set of plurisubharmonic (psh) functions in Ω . Let (X, ω) be a compact Kähler manifold of dimension k, we denote by QPSH(X) the set of quasi-plurisubharmonic (qpsh) functions on X. These functions are locally given as the sum of a smooth and a plurisubharmonic function. We also denote by $PSH(X, \omega)$ the set of ω -plurisubharmonic (ω -psh) functions on X, i.e., the set of functions $\varphi \in L^1(X, \mathbb{R} \cup \{-\infty\})$ such that φ qpsh and $\omega + dd^c \varphi \ge 0$.

Definition 1.1 (Higher complex Sobolev spaces). For $q \ge 1$, we define inductively *q*-complex Sobolev spaces W_q^* as follows

(1) (local setting) $W_q^*(\Omega)$ is the set of all functions $\varphi \in W^{1,2}(\Omega)$ such that $d\varphi \wedge d^c \varphi \leq dd^c \psi$ for some $\psi \in W_{q-1}^*(\Omega) \cap PSH(\Omega)$ ($\psi \in PSH(\Omega)$ when q = 1) satisfying

$$\int_\Omega dd^c\psi\wedge\omega^{k-1}<+\infty, ext{ where }\omega ext{ is the standard K}$$
ähler form on \mathbb{C}^k

(2) (global setting) $W_q^*(X)$ is the set of all functions $\varphi \in W^{1,2}(X)$ such that $d\varphi \wedge d^c \varphi \leq c_{\varphi} \omega + dd^c \psi$ for some constant $c_{\varphi} \geq 0$ and $\psi \in W_{q-1}^*(X) \cap QPSH(X)$ ($\psi \in QPSH(X)$ when q = 1).

We will introduce some notions and technical tools to study these functional spaces. In particular, using techniques in [DS06a, Vig07], we will build a quasinorm $\|\cdot\|_{*,q}$ on W_q^* which makes W_q^* a quasi-Banach space sharing many properties with W_1^* . After that, we will prove some basic properties of these spaces and consider some specific examples. Moreover, we also introduce a family of Vigny's functional capacities and show that all these capacities characterize pluripolar sets, similarly to how the original Vigny's capacity does.

In Section 3, we will prove the Moser-Trudinger inequalities for these higher complex Sobolev spaces. An important point in our results is that the exponent in the Moser-Trudinger inequalities goes to infinity when q goes to infinity. It follows that when qgrows, our spaces will gain more regularity and get closer to the bounded functions space.

Theorem 1.2 (Moser-Trudinger inequality in a local setting). Let $\alpha \in [1, 2^q)$ and K be a compact subset of the unit ball \mathbb{B} of \mathbb{C}^k . Let v_1, \ldots, v_k be psh functions which are Hölder continuous of Hölder exponent β for some $\beta \in (0, 1)$ on \mathbb{B} . Let $\varphi \in W_q^*(\mathbb{B})$. Assume that $\|v_j\|_{C^\beta} \leq 1$ for $1 \leq j \leq k$ and $\|\varphi\|_{*,q} \leq 1$. Then there exist strictly positive constants c_1 and c_2 depending on K, α , and β but independent of $\varphi, v_1, \ldots, v_k$ such that

$$\int_{K} e^{c_{1}|\varphi|^{\alpha}} dd^{c} v_{1} \wedge \dots \wedge dd^{c} v_{k} \leq c_{2}.$$

In particular, there exist strictly positive constants c_1 and c_2 depending on K and α such that for every $\varphi \in W_q^*(\mathbb{B})$ with $\|\varphi\|_{*,q} \leq 1$, there holds

$$\int_{K} e^{c_1 |\varphi|^{\alpha}} \omega^k \le c_2,$$

where ω is the standard Kähler form in \mathbb{C}^k .

Note that two functions in $W_q^*(\mathbb{B})$ are equal if they are equal almost everywhere. By [VV23, Theorem 1.1] (see also [DMV20]), for every function φ in $W_1^*(\mathbb{B})$ (and thus $W_q^*(\mathbb{B})$), all points are Lebesgue point except for points in some pluripolar set. So by considering the canonical values of φ at its Lebesgue points, the first integral in the theorem makes sense as $dd^cv_1 \wedge \cdots \wedge dd^cv_k$ has no mass on pluripolar sets. Throughout this paper, we always use the canonical values of φ as above.

The Moser-Trudinger inequality for W_1^* was proven in [DMV20] by using the slicing method. Recently, in [VV23], alongside the main goal of proving that the complement of the Lebesgue point set of functions in W_1^* is pluripolar, Vigny and Vu obtained a version of the Moser-Trudinger inequality for W_1^* which corresponds to the last assertion of Theorem 1.2 for q = 1. For $\varphi \in W_1^*(\mathbb{B})$, their strategy was to bound $|\varphi|^{\alpha}$ by some psh function which allows them to use Skoda's integrability theorem. The construction of the psh bound is motivated by the proof of Josefson's theorem (see [Jos78]). Following their strategy, we construct the psh bound for $|\varphi|^{\alpha}$ where $\varphi \in W_q^*(\mathbb{B})$ (see Theorem 3.11 below) and deduce Theorem 1.2 by using a singular version of Skoda's integrability theorem which has been obtained in [DNS10] (see also [Kau17]). We also obtain the following global version of Theorem 1.2.

Theorem 1.3 (Moser-Trudinger inequality in a global setting). Let (X, ω) be a compact Kähler manifold and $\alpha \in [1, 2^q)$. Let v_1, \ldots, v_k be ω -psh functions which are Hölder continuous of Hölder exponent β for some $\beta \in (0, 1)$. Let $\varphi \in W_q^*(X)$. Assume that $\|v_j\|_{C^{\beta}} \leq 1$ for $1 \leq j \leq k$ and $\|\varphi\|_{*,q} \leq 1$. Then there exist strictly positive constants c_1 and c_2 depending on X, ω, α and β but independent of $\varphi, v_1, \ldots, v_k$ such that

$$\int_X e^{c_1|\varphi|^{\alpha}} (\omega + dd^c v_1) \wedge \dots \wedge (\omega + dd^c v_k) \le c_2$$

In particular, there exist strictly positive constants c_1 and c_2 depending on X, α , and ω such that for every $\varphi \in W_q^*(X)$ with $\|\varphi\|_{*,q} \leq 1$, there holds

$$\int_X e^{c_1|\varphi|^{\alpha}} \omega^k \le c_2$$

In Section 4, we will discuss the connection of the higher complex Sobolev spaces to the theory of complex Monge-Ampère equation. In the global setting, the class $\mathcal{E}(X,\omega)$, introduced by Guedj-Zeriahi in [GZ07], is the largest class of ω -psh functions on which the complex Monge-Ampère operator is well defined and the comparison principle is valid. We will show that ω -psh functions with bounded $\|\cdot\|_{*,q}$ -norm belong to this space for every $q \ge 1$. Furthermore, among the subsets of $\mathcal{E}(X,\omega)$, the finite energy classes $\mathcal{E}^p(X,\omega)$ have important applications in the variational approach to complex Monge-Ampère equation (see [BBGZ13]). In [DGL20], the authors proved a Moser-Trudinger inequality for functions in $\mathcal{E}^p(X,\omega)$. The crucial point here is that the exponent in their Moser-Trudinger inequality is 1 + p/k which converges to infinity as p goes to infinity. This similar property with our space $W_q^*(X)$ motivated us to study if $W_q^*(X) \cap PSH(X,\omega)$ is contained in some $\mathcal{E}^{p(q)}$ with p(q) increases to infinity when q goes to infinity. It turns out we can choose p(q) = q - 1. **Theorem 1.4.** Let (X, ω) be a compact Kähler manifold. Then we have the following inclusions:

- (1) W^{*}_q(X) ∩ PSH(X, ω) ⊂ E(X, ω) for q ≥ 1,
 (2) W^{*}_q(X) ∩ PSH(X, ω) ⊂ E^{q-1}(X, ω) for q ≥ 2.

The proof of this theorem is by induction and relies on estimations around energy functionals.

In the local setting, the domain of the definition of the Monge-Ampère operator $\mathcal{D}(\Omega)$ has been well understood after the works of Cegrell ([Ceg04]) and Błocki ([Blo06]). It has been proved in [Blo04] that in \mathbb{C}^2 , the Monge-Ampère operator is well-defined for functions in $W_{\text{loc}}^{1,2}$ and thus in $W_{1,\text{loc}}^*$, where $W_{1,\text{loc}}^*$ is the set of functions that locally are functions in W_1^* . Our next theorem generalizes this fact by showing that the Monge-Ampère operator $(dd^c \cdot)^k$ is well-defined in the sense of Cegrell-Błocki for psh locally *q*-complex Sobolev functions when $q \ge k - 1$.

Theorem 1.5. $W^*_{a,loc}(\Omega) \cap PSH(\Omega) \subset \mathcal{D}(\Omega)$ for $q \geq k-1$.

We prove this theorem by defining the product of (1,1)-currents using an induction on the number of currents.

The paper is organized as follows. In Section 2, we consider some basic properties of W_a^* and corresponding Vigny's capacity. In Section 3, we construct a psh bound in order to prove Theorems 1.2 and 1.3. Finally, in Section 4, we study the relationship between W_q^* and the complex Monge-Ampère equation and prove Theorems 1.4 and 1.5.

2. HIGHER COMPLEX SOBOLEV SPACE

In this section, we prove some basic properties and give examples of our spaces. We also study their corresponding Vigny's capacities.

2.1. Quasinorms and compactness. We now define the quasinorm on W_q^* . First, we consider the local case. Let $\varphi_0 \in W_q^*(\Omega)$. Then, by definition, there exist psh functions $\varphi_1, \ldots, \varphi_q$ in Ω such that

$$\|dd^c\varphi_j\| < +\infty \text{ for all } j = 1, \ldots, q$$

and

(2.1)
$$d\varphi_{j-1} \wedge d^c \varphi_{j-1} \le dd^c \varphi_j \text{ for all } j = 1, \dots, q.$$

We call a sequence $(\varphi_1, \ldots, \varphi_q)$ satisfying (2.1) is a defining sequence of φ_0 in $W_q^*(\Omega)$. We define the q-star quasinorm for $W^*_q(\Omega)$ as follows

$$\|\varphi\|_{W_q^*(\Omega)} = \|\varphi\|_{L^2} + \min\Big\{\sum_{j=1}^q \|dd^c\varphi_j\|^{1/2^j} : (\varphi_1, \dots, \varphi_q) \text{ is a defining sequence of } \varphi\Big\}.$$

In the sequel, for simplicity, we will use the notation $\|\cdot\|_{*,q}$ instead of $\|\cdot\|_{W_a^*(\Omega)}$ when there is no possible confusion of domain.

Next, we consider the global case. Let $\varphi_0 \in W_a^*(X)$. Then, by definition, there exist qpsh functions $\varphi_1, \ldots, \varphi_q$ in X and constants $c_1, \ldots, c_q \ge 0$ such that

(2.2)
$$d\varphi_{j-1} \wedge d^c \varphi_{j-1} \le c_j \omega + dd^c \varphi_j \text{ for all } j = 1, \dots, q.$$

We call a sequence $((c_1, \varphi_1), \ldots, (c_q, \varphi_q))$ satisfying (2.2) is a *defining sequence* of φ_0 in $W_q^*(X)$. We define the *q*-star quasinorm for $W^*(X)$ as follows

$$\|\varphi\|_{*,q} = \|\varphi\|_{L^2} + \min\Big\{\sum_{j=1}^q c_j^{1/2^j} : \Big((c_1,\varphi_1),\ldots,(c_q,\varphi_q)\Big) \text{ is a defining sequence of }\varphi\Big\}.$$

In the global case, note that by Poincaré-Wirtinger inequality, if we replace L^2 in the definition of $\|\cdot\|_{*,q}$ by L^1 , we obtain an equivalent quasinorm.

Proposition 2.1. The function $\varphi \mapsto \|\varphi\|_{*,q}$ defines a quasinorm on $W_q^*(\Omega)$ (and $W_q^*(X)$ respectively).

Proof. Consider the local case. We first check the homogeneity. Let $\psi = \lambda \varphi$, then $(\varphi_1, \ldots, \varphi_q)$ is a defining sequence for φ iff $(|\lambda|^2 \varphi_1, \ldots, |\lambda|^{2^q} \varphi_q)$ is a defining sequence for ψ . Hence, $\|\psi\|_{*,q} = |\lambda| \|\varphi\|_{*,q}$. We now only need to check the quasi-triangle inequality. Let φ and ψ be functions in $W_q^*(\Omega)$ where $(\varphi_1, \ldots, \varphi_q)$ and (ψ_1, \ldots, ψ_q) are two corresponding defining sequences. Put $f = \varphi + \psi$. It is sufficient to prove that (f_1, \ldots, f_q) , with $f_j = \frac{4^{2^{j-1}}}{2}(\varphi_j + \psi_j)$, is a defining sequence of f satisfying

$$\sum_{j=1}^{q} \|dd^{c}f_{j}\|^{1/2^{j}} \leq 2^{1-1/2^{q}} \Big(\sum_{j=1}^{q} \|dd^{c}\varphi_{j}\|^{1/2^{j}} + \sum_{j=1}^{q} \|dd^{c}\psi_{j}\|^{1/2^{j}}\Big).$$

We prove it by induction on *q*. For q = 1, we have

$$df \wedge d^c f = d\varphi \wedge d^c \varphi + d\psi \wedge d^c \psi + (d\varphi \wedge d^c \psi + d\psi \wedge d^c \varphi).$$

It follows from Cauchy-Schwarz inequality that

$$d\varphi \wedge d^c \psi + d\psi \wedge d^c \varphi \le d\varphi \wedge d^c \varphi + d\psi \wedge d^c \psi.$$

This implies that

$$df \wedge d^c f \le 2(d\varphi \wedge d^c \varphi + d\psi \wedge d^c \psi) \le dd^c \left(2\varphi_1 + 2\psi_1\right).$$

Then, for $f_1 = 2(\varphi_1 + \psi_1)$, we have

$$df \wedge d^c f \le dd^c f_1,$$

and

$$\|dd^{c}f_{1}\|^{1/2} \leq \sqrt{2}(\|dd^{c}\varphi_{1}\|^{1/2} + \|dd^{c}\psi_{1}\|^{1/2}).$$

This finishes the proof for q = 1. Now we assume that the desired property is true for q-1 where $q \ge 2$. We have $f_{q-1} = \frac{4^{2^{q-2}}}{2}(\varphi_{q-1} + \psi_{q-1})$. It thus follows from Cauchy-Schwarz inequality and the definition of defining sequences that

$$df_{q-1} \wedge d^c f_{q-1} \leq \frac{4^{2^{q-1}}}{2} dd^c \left(\varphi_q + \psi_q\right).$$

Then, for $f_q = \frac{4^{2^{q-1}}}{2}(\varphi_q + \psi_q)$, we have

$$df_{q-1} \wedge d^c f_{q-1} \le dd^c f_q,$$

and

$$\|dd^c f_q\|^{1/2^q} \le 2^{1-1/2^q} \left(\|dd^c \varphi_q\|^{1/2^q} + \|dd^c \psi_q\|^{1/2^q} \right)$$

Thus, by the induction hypothesis, we have

$$\begin{split} \sum_{j=1}^{q} \|dd^{c}f_{j}\|^{1/2^{j}} &= \sum_{j=1}^{q-1} \|dd^{c}f_{j}\|^{1/2^{j}} + \|dd^{c}f_{q}\|^{1/2^{q}} \\ &\leq 2^{1-1/2^{q-1}} \Big(\sum_{j=1}^{q-1} \|dd^{c}\varphi_{j}\|^{1/2^{j}} + \sum_{j=1}^{q-1} \|dd^{c}\psi_{j}\|^{1/2^{j}} \Big) \\ &+ 2^{1-1/2^{q}} \Big(\|dd^{c}\varphi_{q}\|^{1/2^{q}} + \|dd^{c}\psi_{q}\|^{1/2^{q}} \Big) \\ &\leq 2^{1-1/2^{q}} \Big(\sum_{j=1}^{q} \|dd^{c}\varphi_{j}\|^{1/2^{j}} + \sum_{j=1}^{q} \|dd^{c}\psi_{j}\|^{1/2^{j}} \Big), \end{split}$$

as desired. The proof in the local case is thus completed.

Consider the global case. The homogeneity is clear. Now let φ and ψ be functions in $W_q^*(X)$ where $((\alpha_1, \varphi_1), \ldots, (\alpha_q, \varphi_q))$ and $((\beta_1, \psi_1), \ldots, (\beta_q, \psi_q))$ are corresponding defining sequences. Let $f = \varphi + \psi$. As in the local case, it is sufficient to prove that $((\gamma_1, f_1), \ldots, (\gamma_q, f_q))$, with $\gamma_j = \frac{4^{2^{j-1}}}{2}(\alpha_j + \beta_j)$ and $f_j = \frac{4^{2^{j-1}}}{2}(\varphi_j + \psi_j)$, is a defining sequence of f satisfying

$$\sum_{j=1}^{q} \gamma_j^{1/2^j} \le 2^{1-1/2^q} \Big(\sum_{j=1}^{q} \alpha_j^{1/2^j} + \sum_{j=1}^{q} \beta_j^{1/2^j} \Big).$$

We prove it by induction on q. For q = 1, the same computation as in the local case gives

$$df \wedge d^c f \le (2\alpha_1 + 2\beta_1)\,\omega + dd^c\,(2\varphi_1 + 2\psi_1)$$

Then, for $\gamma_1 = 2(\alpha_1 + \beta_1)$ and $f_1 = 2(\varphi_1 + \psi_1)$, we have

$$df \wedge d^c f \le \gamma_1 \omega + dd^c f_1,$$

and

$$\gamma_1^{1/2} \le \sqrt{2}(\alpha_1^{1/2} + \beta_1^{1/2}).$$

This finishes the proof for q = 1. Now we assume that the desired property is true for q - 1 where $q \ge 2$. We have $\gamma_{q-1} = \frac{4^{2^{q-2}}}{2}(\alpha_{q-1} + \beta_{q-1})$ and $f_{q-1} = \frac{4^{2^{q-2}}}{2}(\varphi_{q-1} + \psi_{q-1})$. It thus follows from Cauchy-Schwarz inequality and the definition of defining sequences that

$$df_{q-1} \wedge d^c f_{q-1} \leq \frac{4^{2^{q-1}}}{2} (\alpha_q + \beta_q) \omega + \frac{4^{2^{q-1}}}{2} dd^c (\varphi_q + \psi_q).$$

Then, for $\gamma_q = \frac{4^{2^{q-1}}}{2}(\alpha_q + \beta_q)$ and $f_q = \frac{4^{2^{q-1}}}{2}(\varphi_q + \psi_q)$, we have

$$df_{q-1} \wedge d^c f_{q-1} \le \gamma_q \omega + dd^c f_q,$$

and

$$\gamma_q^{1/2^q} \le 2^{1-1/2^q} \left(\alpha_q^{1/2^q} + \beta_q^{1/2^q} \right).$$

It thus follows from the induction hypothesis that

$$\begin{split} \sum_{j=1}^{q} \gamma_j^{1/2^j} &= \sum_{j=1}^{q-1} \gamma_j^{1/2^j} + \gamma_q^{1/2^q} \\ &\leq 2^{1-1/2^{q-1}} \Big(\sum_{j=1}^{q-1} \alpha_j^{1/2^j} + \sum_{j=1}^{q-1} \beta_j^{1/2^j} \Big) + 2^{1-1/2^q} \Big(\alpha_q^{1/2^q} + \beta_q^{1/2^q} \Big) \\ &\leq 2^{1-1/2^q} \Big(\sum_{j=1}^{q} \alpha_j^{1/2^j} + \sum_{j=1}^{q} \beta_j^{1/2^j} \Big), \end{split}$$

as desired. The proof is completed.

Remark 2.2. Note that when the domain Ω is nice (e.g., convex domain), we have $W_1^* = W^*$. This restriction is from the fact that we cannot solve the equation $dd^c\varphi = T$ in every domain. The spaces W_q^* is decreasing in q. Indeed, by definition, the q-star quasinorms are increasing in q, namely

$$\|\varphi\|_{*,q} \le \|\varphi\|_{*,q+1}$$
 for every $\varphi \in W_{q+1}^*$.

Next, we prove a compactness property of W_a^* .

Proposition 2.3. Let (φ_n) be a bounded sequence in $W_q^*(\Omega)$ $(W_q^*(X)$ respectively). Then there exists a subsequence (φ_{n_j}) and a function $\varphi \in W_q^*(\Omega)$ $(W_q^*(X)$ respectively) such that φ_{n_j} converges weakly to φ in $W^{1,2}$ and $\|\varphi\|_{*,q} \leq \lim_{j \to +\infty} \|\varphi_{n_j}\|_{*,q}$.

Proof. We prove the proposition for the local case only, as the proof for the global case merely requires a modification of this proof.

We proceed by induction on q. For q = 1, the proposition is a slight modification of [Vig07, Proposition 4]. Now, we assume that the desired property is true for q - 1 where q > 1. Let $\psi_n \in W_{q-1}^*$ such that $d\varphi_n \wedge d^c\varphi_n \leq dd^c\psi_n$ and $\|\psi_n\|_{*,q-1}$ is bounded. It follows from (φ_n) is bounded in W_q^* and $\|\varphi_n\|_{*,q-1} \leq \|\varphi_n\|_{*,q}$ that (φ_n) is also bounded in W_{q-1}^* . Thus, by the induction hypothesis, there are subsequences $(\varphi_{n_j}), (\psi_{n_j})$ of $(\varphi_n), (\psi_n)$ and functions $\varphi, \psi \in W_{q-1}^*$ such that $\varphi_{n_j}, \psi_{n_j}$ converge weakly to φ, ψ respectively in $W^{1,2}$ and $\|\psi\|_{*,q-1} \leq \lim_{j \to +\infty} \|\psi_{n_j}\|_{*,q-1}$. Hence, $dd^c\psi_{n_j}$ converges weakly to $dd^c\psi$. Moreover, as noted in the proof of [Vig07, Proposition 4], if Θ is a weak limit of $(d\varphi_{n_j} \wedge d^c\varphi_{n_j})$, then $d\varphi \wedge d^c\varphi \leq \Theta$. So, we have $d\varphi \wedge d^c\varphi \leq dd^c\psi$. Therefore, $\varphi \in W_q^*$. Let $(\psi_{0,1}, \ldots, \psi_{0,q-1})$ and $(\psi_{n_j,1}, \ldots, \psi_{n_j,q-1})$ be respectively the defining sequence of ψ and ψ_{n_j} such that

$$\|\psi\|_{*,q-1} = \|\psi\|_{L^2} + \sum_{t=1}^{q-1} \|dd^c\psi_{0,t}\|^{1/2^t},$$

and

$$\|\psi_{n_j}\|_{*,q-1} = \|\psi_{n_j}\|_{L^2} + \sum_{t=1}^{q-1} \|dd^c \psi_{n_j,t}\|^{1/2^t}.$$

We observe that

(2.3)
$$\|\varphi\|_{*,q} \le \|\varphi\|_{L^2} + \|dd^c\psi\|^{1/2} + \sum_{t=1}^{q-1} \|dd^c\psi_{0,t}\|^{1/2^{t+1}}$$

8

It follows from $\|\psi\|_{*,q-1} \leq \lim_{j \to +\infty} \|\psi_{n_j}\|_{*,q-1}$ and φ_{n_j} , ψ_{n_j} converge weakly to φ , ψ respectively in $W^{1,2}$ that

(2.4)
$$\|\varphi\|_{L^2} \leq \liminf_{j \to +\infty} \|\varphi_{n_j}\|_{L^2}$$

and

(2.5)
$$\|dd^c\psi\|^{1/2} \leq \liminf_{j \to +\infty} \|dd^c\psi_{n_j}\|^{1/2},$$

and

(2.6)
$$\sum_{t=1}^{q-1} \|dd^{c}\psi_{0,t}\|^{1/2^{t+1}} \leq \liminf_{j \to +\infty} \sum_{t=1}^{q-1} \|dd^{c}\psi_{n_{j},t}\|^{1/2^{t+1}}.$$

Combining inequalities (2.3), (2.4), (2.5), and (2.6) gives us

$$\|\varphi\|_{*,q} \leq \liminf_{j \to +\infty} \left(\|\varphi_{n_j}\|_{L^2} + \|dd^c \psi_{n_j}\|^{1/2} + \sum_{t=1}^{q-1} \|dd^c \psi_{n_j,t}\|^{1/2^{t+1}} \right).$$

Then, by the choice of ψ_{n_i} , we obtain

$$\|\varphi\|_{*,q} \le \liminf_{j \to +\infty} \|\varphi_{n_j}\|_{*,q}$$

By passing to a subsequence of (φ_{n_i}) , we complete the proof.

Now, we prove that with the quasinorm $\|\cdot\|_{*,q}$, W_q^* is a quasi-Banach space.

Proposition 2.4. W_q^* endowed with the quasinorm $\|\cdot\|_{*,q}$ is a quasi-Banach space.

Proof. The proof proceeds by induction on q. For q = 1, the proposition is a slight modification of [Vig07, Proposition 1]. Now, we assume that W_{q-1}^* is a quasi-Banach space endowed with the quasinorm $\|\cdot\|_{*,q-1}$ for $q \ge 2$. Let (φ_n) be a Cauchy sequence in W_q^* . Since $\|\varphi\|_{*,q-1} \le \|\varphi\|_{*,q}$, (φ_n) is also a Cauchy sequence in W_{q-1}^* . It thus follows from the induction hypothesis that this sequence converges to a function $\varphi \in W_{q-1}^*$. For every $\epsilon > 0$, there is an integer N such that for n and m greater than N, we have

$$d(\varphi_n - \varphi_m) \wedge d^c(\varphi_n - \varphi_m) \le dd^c \psi_{n,m}$$

where $\psi_{n,m}$ is a function with $\|\psi_{n,m}\|_{*,q-1} < \epsilon$. Let *n* go to infinity. Since $(\varphi_n - \varphi_m)_n$ converges in $W^{1,2}$ to $\varphi - \varphi_m$, $d(\varphi_n - \varphi_m) \wedge d^c(\varphi_n - \varphi_m)$ converges in L^1 to $d(\varphi - \varphi_m) \wedge d^c(\varphi - \varphi_m)$. By Proposition 2.3, we can find a subsequence of $(\psi_{n,m})_n$ which converges weakly in $W^{1,2}$ to a function ψ_m . This function satisfies

$$d(\varphi - \varphi_m) \wedge d^c(\varphi - \varphi_m) \le dd^c \psi_m$$

and $\|\psi_m\|_{*,q-1} \leq \epsilon$. Thus, $\varphi \in W_q^*$ and φ_n converges to φ in W_q^* . We complete our proof.

Remark 2.5. It is worth noting that in the case when q = 1, one can use a slight modification of the proof of [Vig07, Proposition 1] to prove that $\|\cdot\|_{*,1}$ is a norm and W_1^* is a Banach space with respect to this norm. It is also interesting to know if we can build a norm on W_q^* for q > 1. Note that by Aoki-Rolewicz's Theorem, we can pick some equivalent quasinorm $\|\cdot\|$ with $\|\cdot\|_{*,q}$, such that for some $0 , <math>\|x\|^p + \|y\|^p \le \|x + y\|^p$ for all $x, y \in W_q^*$ (see [KPR84]).

2.2. Examples. We now give specific examples for functions in W_q^* .

Example 2.6. Let φ be a psh function in Ω . Assume that φ satisfies the following condition

 $(2.7) d\varphi \wedge d^c \varphi \le r dd^c \varphi$

for some r > 0. Then it is clear by induction that $\varphi \in W_q^*(\Omega)$ for all q. The condition (2.7) was introduced in [DF83] and has many applications in studying the Bergman kernel of Ω (see for example [BC00, Blo13]).

Example 2.7. Let φ be an ω -psh function in a compact Kähler manifold (X, ω) . So $dd^c \varphi + \omega \ge 0$. If φ is bounded (assume for simplicity that $0 \le \varphi \le 1$), then we have

$$d\varphi \wedge d^c \varphi = dd^c (\varphi^2)/2 - \varphi dd^c \varphi \le \omega + dd^c (\varphi^2)/2,$$

$$d(\varphi^2/2) \wedge d^c (\varphi^2/2) = \varphi^2 d\varphi \wedge d^c \varphi \le d\varphi \wedge d^c \varphi \le \omega + dd^c (\varphi^2/2)$$

By choosing the defining sequence to be $((1, \varphi^2/2), \ldots, (1, \varphi^2/2))$, we deduce that $\varphi \in W_q^*(X)$ for all q. Thus, bounded ω -psh functions belong to the intersection of all $W_q^*(X)$. If φ is unbounded (assume for simplicity that $\varphi \leq -1$). Let $\psi = -\log(-\varphi)$. We have

$$d\psi \wedge d^c\psi = rac{d\varphi \wedge d^c\varphi}{|\varphi|^2}$$
 and
 $dd^c\psi = -rac{dd^c\varphi}{\varphi} + rac{d\varphi \wedge d^c\varphi}{|\varphi|^2}$

It follows that $d\psi \wedge d^c\psi = dd^c\psi + dd^c\varphi/\varphi \leq dd^c\psi + \omega$. Thus, by choosing the defining sequence to be $((1,\psi),\ldots,(1,\psi))$, we deduce that $\psi \in W_q^*(X)$ for all q.

Example 2.8. It follows from [Vig07, Section 2.3] (see also [DS06a, Proposition 4.1]) that if $f, g \in W_q^*$, then

$$d\max(f,g) \wedge d^c \max(f,g) \le dd^c(f_1 + g_1),$$

$$d\min(f,g) \wedge d^c \min(f,g) \le dd^c(f_1 + g_1),$$

where f_1, g_1 are functions in W_{q-1}^* such that $df \wedge d^c f \leq dd^c f_1$ and $dg \wedge d^c g \leq dd^c g_1$. So, both $\max(f, g)$ and $\min(f, g)$ belong to W_q^* . Moreover, there exists a constant c not depending on f and g such that

$$\|\max(f,g)\|_{*,q} \le c \left(\|f\|_{*,q} + \|g\|_{*,q}\right)$$
 and $\|\min(f,g)\|_{*,q} \le c \left(\|f\|_{*,q} + \|g\|_{*,q}\right)$.

Example 2.9. Let φ be the function defined by $-(-\log |z_1|^2)^{\alpha}$ in the unit ball \mathbb{B} of \mathbb{C}^k . Then,

$$i\partial\varphi\wedge\bar{\partial}\varphi = \frac{idz_1\wedge d\bar{z}_1}{|z_1|^2(-\log|z_1|^2)^{2-2\alpha}}.$$

Let $\psi = -(-\log|z_1|^2)^{2lpha}$. We have

$$\bar{\partial}\psi = 2\alpha(-\log|z_1|^2)^{2\alpha-1}\frac{1}{\bar{z_1}}d\bar{z_1}, \qquad i\partial\bar{\partial}\psi = 2\alpha(1-2\alpha)\frac{idz_1 \wedge d\bar{z_1}}{|z_1|^2(-\log|z_1|^2)^{2-2\alpha}}.$$

Thus,

$$i\partial\varphi\wedge\bar\partial\varphi=\frac{1}{2\alpha(1-2\alpha)}i\partial\bar\partial\psi$$

So, by induction on q and [Vig07, Example 2], we can prove that $\varphi \in W_q^*(\mathbb{B})$ if and only if $\alpha < \frac{1}{2q}$.

Example 2.10. Let φ be a subharmonic function in $W^{1,2}(U)$ where U is an open subset of \mathbb{C} . Then for any ball $B \subset U$, by Poincaré-Sobolev inequality, we have

$$\frac{1}{|B|} \int_{B} |\varphi - m_B(\varphi)| \le c \Big(\int_{B} |\nabla \varphi|^2 \Big)^{1/2},$$

where c is a constant not depending on φ and B. Let the radius of B go to zero, we deduce that φ is a VMO function. Thus, as pointed out in [BW24, Theorem 1.1], φ must have zero Lelong number at every point of U. Now, consider φ to be a psh function in $W_1^*(\Omega)$. By slicing method (see [DMV20]) and Siu's theorem (see [Dem, Chapter. III (7.13)]), we deduce that φ has zero Lelong number at every point of Ω . By remark after [Vig07, Proposition 6], we deduce that $W_q^*(\Omega)$ is a VMO space for every q > 1. The same result holds in the global setting. In fact, as we will see in Theorem 1.4, if $\varphi \in W_1^*(X) \cap PSH(X, \omega)$, then $\varphi \in \mathcal{E}(X, \omega)$. And it is well-known that functions in $\mathcal{E}(X, \omega)$ have zero Lelong numbers at every point.

2.3. **Density theorems.** We now prove density theorems for W_q^* . The approximate sequences have been constructed in [Vig07], and we use the same construction for our spaces.

First, we consider the local case. We have the following result.

Theorem 2.11. Let K be a relatively compact subset of Ω and $\varphi \in W_q^*(\Omega)$. Then there exists a sequence of smooth functions (φ_n) converges to φ in $W^{1,2}(K)$. Moreover, we have

$$\|\varphi\|_{W_q^*(K)} \le \lim_{n \to +\infty} \|\varphi_n\|_{W_q^*(K)} \le \|\varphi\|_{W_q^*(\Omega)}.$$

Proof. Take χ to be a non-negative smooth radial function with compact support in \mathbb{C}^k such that $\int_{\mathbb{C}^k} \chi = 1$ and define

$$\chi_{\epsilon}(z) = \epsilon^{-2k} \chi\left(\frac{z}{\epsilon}\right) \cdot$$

Put $\varphi_{\epsilon} = \varphi * \chi_{\epsilon}$, then φ_{ϵ} is well-defined in K when ϵ is small enough. Consider $\psi \in W_{q-1}^*(\Omega) \cap PSH(\Omega)$ such that $d\varphi \wedge d^c\varphi \leq dd^c\psi$. Let $(\epsilon_n)_n$ be a sequence decreasing to zero and define $\varphi_n = \varphi_{\epsilon_n}$, $\psi_n = \psi * \chi_{\epsilon_n}$. Then, by [Vig07, Lemma 5], we have $d\varphi_n \wedge d^c\varphi_n \leq dd^c\psi_n$. Now, let $(\varphi_1, \ldots, \varphi_q)$ be a defining sequence for φ . Define $\varphi_{n,j} = \varphi_j * \chi_{\epsilon_n}$ for $j = 1, \ldots, q$. Then by an induction, $(\varphi_{n,1}, \ldots, \varphi_{n,q})$ is a defining sequence for φ_n . Moreover, we also have

(2.8)
$$\lim_{n \to +\infty} \int_{K} dd^{c} \varphi_{n,j} \wedge \omega^{k-1} \leq \int_{\Omega} dd^{c} \varphi_{j} \wedge \omega^{k-1} \text{ for } j = 1, \dots, q.$$

It follows that (φ_n) is a bounded sequence in $W_q^*(K)$. Thus, by Proposition 2.3, there exists a subsequence (φ_{n_m}) converges to φ in $W^{1,2}(K)$ and satisfies the first inequality. By (2.8), this sequence also satisfies the second inequality. We complete the proof. \Box

We now consider the case of compact Kähler manifolds. We have the following result.

Theorem 2.12. Let $\varphi \in W_q^*(X)$. Then there exists a sequence of smooth functions (φ_n) such that φ_n converges to φ weakly in $W^{1,2}(X)$. Moreover, there is a constant c not depending on φ such that $\lim_{n \to +\infty} \|\varphi_n\|_{*,q} \leq c \|\varphi\|_{*,q}$.

Proof. Let (K_n^+) and (K_n^-) be two sequences of positive closed (k, k)-currents constructed in [DS04]. Recall that $(K_n^+ - K_n^-)$ converges to the current of integration on the diagonal

of $X \times X$. Define $K_n = K_n^+ - K_n^-$. Now consider $\varphi \in W_q^*(X)$ and put

$$\varphi_n(x) = \int_{y \in X} \varphi(y) K_n(x, y).$$

Let ψ be the function in $W^*_{q-1}(X)$ such that $d\varphi \wedge d^c\varphi \leq c(\omega + dd^c\psi)$, then we can bound $d\varphi_n \wedge d^c\varphi_n$ by

$$A\int_{y\in X} K_n^{\pm}(x,y) \wedge (\omega + dd^c\psi).$$

Here A is a constant depending on the manifold X and $\|\varphi\|_{*,q}$. This is a positive closed current (note that both K_n^+ and K_n^- are positive closed current) and for N big enough, we can bound this as the form $2A(\omega + dd^c\psi_n)$ where

$$\psi_n = \int_{y \in X} \psi(y) K_n^{\pm}(x, y).$$

As pointed out in [Vig07, Section 2.4] (see [DNS22, Appendix A] for more information), after iterating this convolution several times, we can make φ_n smooth. The result is thus followed by using [Vig07, Theorem 10] and induction in q.

2.4. Vigny's functional capacity. Consider the case when (X, ω) is a compact Kähler manifold. Following [Vig07], for a Borel set *E* in *X*, we define

 $L_q(E) = \left\{ \varphi \in W_q^* : \varphi \leq -1 \text{ a.e on some neighborhood of } E \text{ and } \varphi \leq 0 \text{ on } X \right\}.$

The corresponding Vigny's capacity for W_a^* can be defined as follows

$$\operatorname{Cap}_{q}(E) = \inf \left\{ \left\| \varphi \right\|_{*,q}^{2} : \varphi \in L_{q}(E) \right\}.$$

These capacities share similar properties with the original Vigny's capacity in [Vig07]. We list below some important properties. The proofs are modifications of Vigny's proofs in [Vig07].

Proposition 2.13. The capacity Cap_q satisfies the following properties

- (1) for $E \subset F \subset X$, $\operatorname{Cap}_q(E) \leq \operatorname{Cap}_q(F)$;
- (2) if (E_j) is a sequence of Borel sets in X, $\operatorname{Cap}_q(\cup_j E_j) \leq c \sum_j \operatorname{Cap}_q(E_j)$ for some

constant c not depending on (E_j) ;

- (3) $\operatorname{Cap}_q(X) = 1$ and $\operatorname{Cap}_q(E) \leq 1$ for any $E \subset X$;
- (4) if (K_j) is a decreasing sequence of compact sets, $\lim_{j \to +\infty} \operatorname{Cap}_q(K_j) = \operatorname{Cap}_q(\cap_n K_j);$
- (5) if (E_j) is an increasing sequence of Borel sets, $\operatorname{Cap}_q(\cup_j E_j) = \lim_{j \to +\infty} \operatorname{Cap}_q(E_j)$, that is, Cap_q is a Choquet capacity.

Proof. We note that, due to Example 2.8, in W_q^* , we also have the following inequality

$$\|\min(f,g)\|_{*,q}^{2} \le c \left(\|f\|_{*,q}^{2} + \|g\|_{*,q}^{2} \right).$$

Moreover, $||f||_{*,q} = 1$ where f is the constant function 1 on X. We can now follow the proofs of [Vig07, Proposition 27 and Theorem 30] to finish the proof.

 Cap_q also characterizes pluripolar sets in a manner analogous to the original Vigny's capacity.

Theorem 2.14. There exists a strictly positive constant B such that for all Borel subset E of X, we have

$$B^{-1}\operatorname{Cap}_{\omega}(E) \leq \operatorname{Cap}_{q}(E) \leq B(\operatorname{Cap}_{\omega}(E))^{\frac{1}{k^{2q-1}}}.$$

In particular, $\operatorname{Cap}_{a}(E) = 0$ if and only if E is pluripolar.

Recall that the capacity Cap_{ω} was defined by Kołodziej in [Kol03]. It is related to the well-known Bedford-Taylor capacity ([BT82]), and defined by

$$\operatorname{Cap}_{\omega}(E) = \sup \Big\{ \int_{E} (\omega + dd^{c}u)^{k} : \ u \in \operatorname{PSH}(X, \omega), \ -1 \le u \le 0 \Big\},\$$

where E is a Borel subset of X. We refer the reader to [GZ05] for more information on this capacity for local and global settings.

We also need the following notion of capacity introduced by Dinh and Sibony in [DS06b]. It is related to the capacities of Alexander in [Ale81] and of Siciak in [Sic62], see also [HL06] and [GZ05]. For a Borel subset E of X, we consider the function

$$V_E(x) = \sup \left\{ u(x) : u \in PSH(X, \omega) \text{ and } u \leq 0 \text{ on } E \right\}.$$

Then V_E is a non-negative ω -psh function. Define

$$\mathcal{J}(E) = \exp\left(-\sup_{X} V_E(x)\right).$$

Recall from [GZ05, Proposition 6.1] the following relation between Cap_{ω} and \mathcal{J} .

Theorem 2.15. There is a strictly positive constant A such that for all compact subset K of X, we have

$$\exp\left(-\frac{A}{\operatorname{Cap}_{\omega}(K)}\right) \leq \mathcal{J}(K) \leq e \cdot \exp\left(-\frac{1}{\operatorname{Cap}_{\omega}(K)^{1/k}}\right) \cdot$$

Now we can prove Theorem 2.14.

Proof of Theorem 2.14. We follow proofs in [Vig07, DKN20]. Due to Example 2.7, if φ is a qpsh function such that $\varphi < -1$, then the function $\psi = -\log(-\varphi)$ belongs to W_q^* for all q, and ψ has the same poles set as φ . We can now follow the proofs of [Vig07, Proposition 28] to see the first inequality.

We now consider the second inequality. Since both capacities are regular (see [GZ05, Theorem 4.2] and Proposition 2.13.(5)), we only need to show this inequality for any compact regular set $K \subset X$ instead of E. Let $M = \sup_{X} V_K(x)$. Without loss of generality, we may assume that $M < +\infty$, otherwise both capacities vanish. If $M \leq 1$, then by [DKN20, page 14], the desired inequality follows. Now we only consider the case when $M \geq 1$. Define

$$f_K(x) = \frac{V_K(x) - M}{M}$$

Then f_K is equal -1 on K with $-1 \le f_K \le 0$ and f_K is qpsh with $dd^c f_K + M^{-1}\omega \ge 0$. Since f_K is a bounded qpsh function, it follows from Example 2.7 that $f_K \in W_q^*$ for all q. We now compute $||f_K||_{*,q}$. Direct calculation gives us

$$df_K \wedge d^c f_K = -f_K dd^c f_K + \frac{1}{2} dd^c (f_K^2) \le \frac{\omega}{M} + \frac{1}{2} dd^c (f_K^2),$$

$$d(f_{K}^{2}/2) \wedge d^{c}(f_{K}^{2}/2) = (f_{K}^{2})df_{K} \wedge d^{c}f_{K} \leq df_{K} \wedge d^{c}f_{K} \leq \frac{\omega}{M} + \frac{1}{2}dd^{c}(f_{K}^{2}) \cdot$$

Hence, we can choose $((M^{-1}, f_K^2/2), \ldots, (M^{-1}, f_K^2/2))$ as a defining sequence. Now, as mentioned in the definition of $\|\cdot\|_{*,q}$, if we replace L^2 -norm by L^1 -norm, we obtain an equivalent norm. Since $\max_X f_K = 0$, we have

$$\|f_K\|_{L^1} = \int_X -f_K \omega^n \le \frac{A}{M}$$

for some constant A depending only on (X, ω) because the set of ω -psh function u such that $\max_{X} u = 0$ is a compact subset in $L^{1}(X)$. Thus, we get

$$||f_K||_{*,q} = ||f_K||_{L^1} + \sum_{j=1}^q \frac{1}{M^{2^j}} \le \frac{A}{M} + \sum_{j=1}^q \frac{1}{M^{1/2^j}} \le \frac{B'}{M^{1/2^q}}$$

for some constant B' > 0 since $M \ge 1$. So, by Theorem 2.15,

$$\operatorname{Cap}_{q}(U) \le \|f_{K}\|_{*,q}^{2} \le \frac{B^{\prime 2}}{M^{1/2^{q-1}}} \le B\left(\operatorname{Cap}_{\omega}(U)\right)^{\frac{1}{k^{2q-1}}}.$$

This finishes the proof.

Remark 2.16. Theorem 2.14 directly shows that the capacities Cap_q with $p \ge 1$, are equivalent capacities. Observe that the sequence $(\operatorname{Cap}_q)_{q\ge 1}$ is increasing and always bounded by 1. Given a Borel set E, it is an interesting question to study the behavior of $\operatorname{Cap}_q(E)$ as q goes to infinity and their relationship with the Lebesgue measure.

Remark 2.17. As in [Vig07, Remark 33], one can define Cap_q in the local case by the same method. It is also a Choquet capacity and the sets of zero capacity are exactly the pluripolar sets.

3. MOSER-TRUDINGER INEQUALITIES

In this section, we prove a sequence of estimations of L^m type. Then we construct the psh (qpsh) bound for functions in W_q^* to prove Theorems 1.2 and 1.3.

3.1. Estimations of L^m type. First, we need some auxiliary results. We begin by recalling the following version of Moser-Trudinger inequality for functions in W_1^* .

Theorem 3.1. [DMV20, Theorem 1.1] Let Ω be a domain in \mathbb{C}^k and K a compact subset of U. Let v_1, \ldots, v_k be psh functions which are Hölder continuous of Hölder exponent $\beta \in (0, 1)$ on Ω . Let $\varphi \in W_1^*(\Omega)$. Assume that $\|v_j\|_{C^\beta} \leq 1$ for $1 \leq j \leq k$ and $\|\varphi\|_{*,1} \leq 1$. Then there exist strictly positive constants α and c depending on U, K, β but independent of $\varphi, v_1, \ldots, v_k$ such that

$$\int_{K} e^{\alpha |\varphi|^{2}} dd^{c} v_{1} \wedge \dots \wedge dd^{c} v_{k} \leq c.$$

In particular, φ belongs to L_{loc}^p with respect to the measure $dd^cv_1 \wedge \cdots \wedge dd^cv_k$ for every $p \in [1, \infty)$.

The following result is a direct consequence of Theorem 3.1.

13

14

Corollary 3.2. Let K be a compact subset of Ω and $m \in \mathbb{N}$. There exists a strictly positive constant c_1 so that

$$\int_{K} |\varphi|^{m} \omega^{k} \le c_{1},$$

for every $\varphi \in W_1^*(\Omega)$ with $\|\varphi\|_{*,1} \leq 1$. In particular, the estimate holds for every $\varphi \in W_q^*(\Omega)$ with $\|\varphi\|_{*,q} \leq 1$.

Here, for the last assertion, we use that $\|\varphi\|_{*,1} \leq \|\varphi\|_{*,q}$ for $\varphi \in W_q^*$.

Next, let φ be a positive function in $W_q^*(\mathbb{B})$ and $(\varphi_1, \ldots, \varphi_q)$ be a defining sequence for φ such that

$$\|\varphi\|_{*,q} = \|\varphi\|_{L^2} + \sum_{j=1}^q \|dd^c\varphi_j\|^{1/2^j}.$$

We can assume that $\varphi_q \leq 0$.

Remark 3.3. Recall that if $(\varphi_1, \ldots, \varphi_q)$ is a defining sequence for φ , then $(\varphi_{1,\epsilon}, \ldots, \varphi_{q,\epsilon})$ is a defining sequence for φ_{ϵ} where φ_{ϵ} is the standard regularization of φ and $\varphi_{j,\epsilon}$ is the standard regularization of φ_j for $j = 1, \ldots, q$ (see Theorem 2.11).

We set

- $\phi_n = \max(\varphi_q, -n)$,
- $h_n = 1 + \phi_n/n \in \text{PSH}(\mathbb{B}, [0, 1]), h_n = 0 \text{ on } \{\varphi_q \le -n\},\$
- $T_n = dd^c (h_n^2/2)$. This is a positive closed (1, 1)-current which vanishes on $\{\varphi_q < -n\}$.

We have the following elementary lemma.

Lemma 3.4. Assume that φ_{q-1} and φ_q are smooth. Then

- (1) $dh_n \wedge d^c h_n \leq T_n$ and $h_n dd^c h_n \leq T_n$,
- (2) $d\varphi_{q-1} \wedge d^c \varphi_{q-1} \leq dd^c \phi_n$ on $\{h_n > 0\}$,
- (3) $d\varphi_{q-1} \wedge d^c \varphi_{q-1} \wedge T_n \leq dd^c \phi_{n+1} \wedge T_n$.

By (1), (2), and the definition of h_n , we have the following estimate.

Corollary 3.5. Assume that φ_{q-1} and φ_q are smooth. Then

$$h_n d\varphi_{q-1} \wedge d^c \varphi_{q-1} \le n h_n dd^c h_n \le n T_n$$

Now, we state and prove a sequence of estimates of L^m type.

Definition 3.6. Let ω be the canonical Kähler form on \mathbb{C}^k . For every $m \in \mathbb{N}$, $K \subseteq \mathbb{B}$, define for $0 \le p \le k$

$$I_{m,p,K} = \sup_{v_1,\dots,v_p \in \mathsf{PSH}(\mathbb{B},[0,1])} \int_K h_n^2 \varphi^{2m} dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge \omega^{k-p}$$

and for $0 \le p \le k-1$

$$J_{m,p,K} = \sup_{v_1,\dots,v_p \in \mathsf{PSH}(\mathbb{B},[0,1])} \int_K \varphi^{2m} T_n dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge \omega^{k-p-1}$$

As noted after Theorem 1.2, we use the canonical representative of φ here, and thus all the integrals make sense. In what follows, we use \lesssim or \gtrsim to denote \leq or \geq respectively modulo a multiplicative constant independent of n and φ provided $\|\varphi\|_{*,q} \leq 1$. Note that

all the definitions depend on n. However, we omit the index n since the estimates do not depend on n.

Lemma 3.7. There is a constant c = c(m, K) independent of φ and n such that

$$J_{m,0,K} \le cn^{\frac{m}{2^{q-1}}}$$

for every n.

Proof. By a standard regularization, we can assume that $\varphi, \varphi_1, \ldots, \varphi_q$ are smooth. Let χ be a smooth cut-off function such that $\chi \equiv 1$ on an open neighborhood of K, and χ is supported on \mathbb{B} . It is sufficient to prove that

$$J_m = \int_{\mathbb{B}} \chi^2 \varphi^{2m} T_n \wedge \omega^{k-1} \lesssim n^{\frac{m}{2q-1}}.$$

It follows from the definition of T_n and Stokes' formula that

$$J_{m} = \frac{1}{2} \int_{\mathbb{B}} \chi^{2} \varphi^{2m} dd^{c} h_{n}^{2} \wedge \omega^{k-1}$$

$$= -\int_{\mathbb{B}} \chi \varphi^{2m} d\chi \wedge d^{c} h_{n}^{2} \wedge \omega^{k-1} - m \int_{\mathbb{B}} \chi^{2} \varphi^{2m-1} d\varphi \wedge d^{c} h_{n}^{2} \wedge \omega^{k-1}$$

$$= -2 \Big(\int_{\mathbb{B}} \chi \varphi^{2m} h_{n} d\chi \wedge d^{c} h_{n} \wedge \omega^{k-1} + m \int_{\mathbb{B}} \chi^{2} \varphi^{2m-1} h_{n} d\varphi \wedge d^{c} h_{n} \wedge \omega^{k-1} \Big).$$

Let A_1 and A_2 be respectively the first and second integrals inside the brackets. It follows from Cauchy-Schwarz inequality, Corollary 3.2 and Lemma 3.4 that

$$\begin{aligned} A_1^2 &\leq \Big(\int_{\mathbb{B}} \chi^2 \varphi^{2m} dh_n \wedge d^c h_n \wedge \omega^{k-1}\Big) \Big(\int_{\mathbb{B}} \varphi^{2m} h_n^2 d\chi \wedge d^c \chi \wedge \omega^{k-1}\Big) \\ &\lesssim \Big(\int_{\mathbb{B}} \chi^2 \varphi^{2m} T_n \wedge \omega^{k-1}\Big) \Big(\int_{\mathrm{supp}(\chi)} \varphi^{2m} \wedge \omega^k\Big) \\ &\lesssim J_m, \end{aligned}$$

and

$$\begin{aligned} A_2^2 &\leq \Big(\int_{\mathbb{B}} \chi^2 \varphi^{2m} dh_n \wedge d^c h_n \wedge \omega^{k-1}\Big) \Big(\int_{\mathbb{B}} \chi^2 \varphi^{2m-2} h_n^2 d\varphi \wedge d^c \varphi \wedge \omega^{k-1}\Big) \\ &\leq \Big(\int_{\mathbb{B}} \chi^2 \varphi^{2m} T_n \wedge \omega^{k-1}\Big) \Big(\int_{\mathbb{B}} \chi^2 \varphi^{2m-2} h_n^2 dd^c \varphi_1 \wedge \omega^{k-1}\Big) \\ &\leq J_m B_{1,m-1}, \end{aligned}$$

where

$$B_{j,m} = \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dd^c \varphi_j \wedge \omega^{k-1} \text{ for } j = 1, \dots, q.$$

This implies $J_m \lesssim \sqrt{J_m} + \sqrt{J_m} \sqrt{B_{1,m-1}}$, and hence (3.1) $J_m \lesssim B_{1,m-1}$.

To estimate $B_{1,m-1}$, we first observe that, by Stokes' formula, for $j = 1, \ldots, q-1$,

$$B_{j,m} = -2\Big(\int_{\mathbb{B}} \chi \varphi^{2m} h_n^2 d\chi \wedge d^c \varphi_j \wedge \omega^{k-1} \\ + m \int_{\mathbb{B}} \chi^2 \varphi^{2m-1} h_n^2 d\varphi \wedge d^c \varphi_j \wedge \omega^{k-1} + \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n dh_n \wedge d^c \varphi_j \wedge \omega^{k-1}\Big)$$

tively. It follows from Cauchy-Schwarz inequality that

$$C_{j,1}^{2} \leq \left(\int_{\mathbb{B}} \varphi^{2m} h_{n}^{2} d\chi \wedge d^{c}\chi \wedge \omega^{k-1}\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge \omega^{k-1}\right)$$
$$\lesssim \left(\int_{\mathrm{supp}(\chi)} \varphi^{2m} \omega^{k}\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge \omega^{k-1}\right),$$
$$C_{j,2}^{2} \leq \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} h_{n}^{2} d\varphi \wedge d^{c}\varphi \wedge \omega^{k-1}\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge \omega^{k-1}\right)$$
$$\leq \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} h_{n}^{2} dd^{c}\varphi_{1} \wedge \omega^{k-1}\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge \omega^{k-1}\right)$$

and

$$C_{j,3}^{2} \leq \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} dh_{n} \wedge d^{c} h_{n} \wedge \omega^{k-1}\Big) \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge \omega^{k-1}\Big).$$

It thus follows from the definition of the defining sequence, Corollary 3.2, Lemma 3.4, Corollary 3.5 and inequality (3.1) that

$$\begin{split} C_{j,1}^2 \lesssim \begin{cases} B_{j+1,m} & \text{if } j < q-1 \\ nB_{1,m-1} & \text{if } j = q-1, \end{cases} & C_{j,2}^2 \lesssim \begin{cases} B_{1,m-1}B_{j+1,m} & \text{if } j < q-1 \\ nB_{1,m-1}^2 & \text{if } j = q-1, \end{cases} \\ & C_{j,3}^2 \lesssim \begin{cases} B_{1,m-1}B_{j+1,m} & \text{if } j < q-1 \\ nB_{1,m-1}^2 & \text{if } j = q-1. \end{cases} \end{split}$$

Note that we use (3.1) and the fact that $0 \le h_n \le 1$ when j = q - 1. Then, we have

$$B_{j,m}^2 \lesssim \begin{cases} B_{j+1,m} + B_{1,m-1}B_{j+1,m} & \text{if } j < q-1\\ nB_{1,m-1}^2 & \text{if } j = q-1, \end{cases}$$

which implies

$$\begin{cases} B_{j,m}^2 \lesssim B_{1,m-1} B_{j+1,m} \text{ for } j < q-1 \\ B_{q-1,m} \lesssim \sqrt{n} B_{1,m-1}, \end{cases}$$

and hence,

$$B_{1,m}^{2^{q-2}} \lesssim B_{q-1,m} B_{1,m-1}^{2^{q-2}-1} \lesssim \sqrt{n} B_{1,m-1}^{2^{q-2}}$$

It thus follows that

$$B_{1,m} \lesssim n^{\frac{1}{2^{q-1}}} B_{1,m-1},$$

 $B_{1,m} \lesssim n^{\frac{m}{2^{q-1}}}.$

and hence

(3.2)

Now, combining inequalities (3.1) and (3.2) gives us

$$J_m \lesssim n^{\frac{m}{2^{q-1}}},$$

as desired.

Lemma 3.8. There is a constant c = c(m, K) independent of φ and n such that $J_{m,p,K} \leq cn^{\frac{m}{2q-1}}$

for every n.

Proof. We prove the lemma by induction on p. If p = 0, the desired assertion is Lemma 3.7. Assume now that it is true for all p' with $p' \leq p - 1$. By a standard regularization, we can assume that $\varphi, \varphi_1, \ldots, \varphi_q$ are smooth. Let χ be a smooth cut-off function such that $\chi \equiv 1$ on an open neighborhood of K, and χ is supported on \mathbb{B} . It is sufficient to prove that

$$J_{m,p} = \sup_{v_1, \dots, v_p \in \mathsf{PSH}(\mathbb{B}, [0,1])} \int_{\mathbb{B}} \chi^2 \varphi^{2m} T_n dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge \omega^{k-p-1} \lesssim n^{\frac{m}{2q-1}}$$

We prove this inequality by induction on m (p now fixed). When m = 0, it is obvious. Assume that it is true for all m' with $m' \leq m - 1$. Consider $v_1, \ldots, v_p \in \text{PSH}(\mathbb{B}, [0, 1])$, we set $R = dd^c v_2 \wedge \cdots \wedge dd^c v_p \wedge T_n \wedge \omega^{k-p-1}$. It follows from Stokes' formula that

$$\int_{\mathbb{B}} \chi^2 \varphi^{2m} T_n dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge \omega^{k-p-1}$$

= $-2 \Big(\int_{\mathbb{B}} \chi \varphi^{2m} d\chi \wedge d^c v_1 \wedge R + m \int_{\mathbb{B}} \chi^2 \varphi^{2m-1} d\varphi \wedge d^c v_1 \wedge R \Big).$

Let D_1 and D_2 be the first and second integrals inside the brackets respectively. By the Cauchy-Schwarz inequality, the induction hypothesis on p, the induction hypothesis on m, and Lemma 3.4, we have

$$D_{1}^{2} \leq \left(\int_{\mathbb{B}} \varphi^{2m} d\chi \wedge d^{c}\chi \wedge R\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} dv_{1} \wedge d^{c}v_{1} \wedge R\right)$$
$$\lesssim \left(\int_{\mathrm{supp}(\chi)} \varphi^{2m} R \wedge \omega\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} dd^{c}(v_{1}^{2}) \wedge R\right)$$
$$\lesssim J_{m,p-1,\mathrm{supp}(\chi)} J_{m,p}$$
$$\lesssim n^{\frac{m}{2^{q-1}}} J_{m,p},$$

and

$$D_{2}^{2} \leq \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} d\varphi \wedge d^{c} \varphi \wedge R\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} dv_{1} \wedge d^{c} v_{1} \wedge R\right)$$
$$\lesssim \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} dd^{c} \varphi_{1} \wedge R\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} dd^{c} (v_{1}^{2}) \wedge R\right)$$
$$\lesssim E_{1,m-1} J_{m,p},$$

where

$$E_{j,m} = \sup_{v_2,\dots,v_p \in \mathsf{PSH}(\mathbb{B},[0,1])} \int_{\mathbb{B}} \chi^2 \varphi^{2m} dd^c \varphi_j \wedge R \text{ for } j = 1,\dots,q.$$

Note that since χ depends on K, the estimate here only depends on K. Then, by taking the supremum over all $v_1, \ldots, v_p \in \text{PSH}(\mathbb{B}, [0, 1])$, we observe that

(3.3)
$$J_{m,p} \lesssim n^{\frac{m}{2q-1}} + E_{1,m-1}.$$

To estimate $E_{1,m-1}$, we consider $v_2, \ldots, v_p \in PSH(\mathbb{B}, [0, 1])$. It follows from Stokes' formula that, for $j = 1, \ldots, q-1$,

$$\int_{\mathbb{B}} \chi^2 \varphi^{2m} dd^c \varphi_j \wedge R = -2 \Big(\int_{\mathbb{B}} \chi \varphi^{2m} d\chi \wedge d^c \varphi_j \wedge R + m \int_{\mathbb{B}} \chi^2 \varphi^{2m-1} d\varphi \wedge d^c \varphi_j \wedge R \Big).$$

Let $F_{j,1}$ and $F_{j,2}$ be the first and second integrals inside the brackets respectively. By the Cauchy-Schwarz inequality, the induction hypothesis on p, and the induction hypothesis

on m, we have

$$F_{j,1}^{2} \leq \left(\int_{\mathbb{B}} \varphi^{2m} d\chi \wedge d^{c}\chi \wedge R\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R\right)$$
$$\lesssim J_{m,p-1,\operatorname{supp}(\chi)} \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R\right)$$
$$\lesssim n^{\frac{m}{2^{q-1}}} \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R\right),$$

and

$$F_{j,2}^{2} \leq \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} d\varphi \wedge d^{c} \varphi \wedge R\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R\right)$$
$$\leq \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} dd^{c} \varphi_{1} \wedge R\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R\right)$$
$$\leq E_{1,m-1} \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R\right).$$

Then, by taking the supremum over all $v_1, \ldots, v_p \in PSH(\mathbb{B}, [0, 1])$, Lemma 3.4 and inequality (3.3), we have

$$E_{j,m}^2 \lesssim \begin{cases} E_{j+1,m} \left(E_{1,m-1} + n^{\frac{m}{2q-1}} \right) & \text{if } j < q-1 \\ n \left(n^{\frac{m}{2q-1}} \left(n^{\frac{m}{2q-1}} + E_{1,m-1} \right) + E_{1,m-1} \left(n^{\frac{m}{2q-1}} + E_{1,m-1} \right) \right) & \text{if } j = q-1. \end{cases}$$

It thus follows that

$$\begin{cases} E_{j,m}^2 \lesssim E_{j+1,m} \left(E_{1,m-1} + n^{\frac{m}{2q-1}} \right) \text{ if } j < q-1 \\ E_{q-1,m} \lesssim \sqrt{n} \left(E_{1,m-1} + n^{\frac{m}{2q-1}} \right). \end{cases}$$

Then, we have

$$E_{1,m}^{2^{q-2}} \le E_{q-1,m} \left(E_{1,m-1} + n^{\frac{m}{2^{q-1}}} \right)^{2^{q-2}-1} \lesssim \sqrt{n} \left(E_{1,m-1} + n^{\frac{m}{2^{q-1}}} \right)^{2^{q-2}},$$

and hence,

$$E_{1,m} \lesssim n^{\frac{1}{2^{q-1}}} \Big(E_{1,m-1} + n^{\frac{m}{2^{p-1}}} \Big).$$

Therefore, we obtain

(3.4)
$$E_{1,m} \lesssim n^{\frac{m+1}{2q-1}}$$

Combining inequalities (3.3) and (3.4) gives us

$$J_{m,p} \lesssim n^{\frac{m}{2^{q-1}}}$$

as desired.

Lemma 3.9. There is a constant c = c(m, K) independent of φ and n such that

$$I_{m,p,K} \le cn^{\frac{m}{2^{q-1}}}$$

for every n.

Proof. We argue similarly as in the proof of Lemma 3.8, by induction on p. If p = 0, the desired assertion follows from Corollary 3.2. We assume now that it is true for every p' with $p' \leq p - 1$. By a standard regularization, we can assume that $\varphi, \varphi_1, \ldots, \varphi_q$ are smooth. Let χ be a smooth cut-off function such that $\chi \equiv 1$ on an open neighborhood of K, and χ is supported on \mathbb{B} . To prove the desired assertion, it suffices to prove that

$$I_{m,p} = \sup_{v_1, \dots, v_p \in \mathsf{PSH}(\mathbb{B}, [0,1])} \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dd^c v_1 \wedge \dots \wedge dd^c v_p \wedge \omega^{k-p} \lesssim n^{\frac{m}{2q-1}}$$

We prove this inequality by induction on m (p now fixed). When m = 0, it is obvious. Assume that it is true for all m' with $m' \le m - 1$. Consider $v_1, \ldots, v_p \in \text{PSH}(\mathbb{B}, [0, 1])$, we set $R' = dd^c v_2 \land \cdots \land dd^c v_p \land \omega^{k-p}$. It follows from Stokes' formula that

$$\begin{split} \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dd^c v_1 \wedge R' &= -2 \Big(\int_{\mathbb{B}} \chi \varphi^{2m} h_n^2 d\chi \wedge d^c v_1 \wedge R' \\ &+ \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n dh_n \wedge d^c v_1 \wedge R' + m \int_{\mathbb{B}} \chi^2 \varphi^{2m-1} h_n^2 d\varphi \wedge d^c v_1 \wedge R' \Big). \end{split}$$

Let G_1, G_2 , and G_3 be the first, second, and third integrals inside the brackets respectively. By the Cauchy-Schwarz inequality, the induction hypothesis on p, the induction hypothesis on m, and Lemma 3.8, we have

$$G_{1}^{2} \leq \left(\int_{\mathbb{B}} h_{n}^{2} \varphi^{2m} d\chi \wedge d^{c}\chi \wedge R'\right) \left(\int_{\mathbb{B}} h_{n}^{2} \varphi^{2m} \chi^{2} dv_{1} \wedge d^{c}v_{1} \wedge R'\right)$$

$$\lesssim I_{m,p-1, \operatorname{supp}(\chi)} \left(\int_{\mathbb{B}} h_{n}^{2} \varphi^{2m} \chi^{2} dd^{c}(v_{1}^{2}) \wedge R'\right)$$

$$\lesssim n^{\frac{m}{2^{q-1}}} I_{m,p},$$

$$G_{2}^{2} \leq \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} dh_{n} \wedge d^{c}h_{n} \wedge R'\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} dv_{1} \wedge d^{c}v_{1} \wedge R'\right)$$

$$\leq \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} T_{n} \wedge R'\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} dd^{c}(v_{1}^{2}) \wedge R'\right)$$

$$\leq J_{m,p-1, \operatorname{supp}(\chi)} I_{m,p}$$

$$\lesssim n^{\frac{m}{2^{q-1}}} I_{m,p},$$

and

$$G_3^2 \leq \left(\int_{\mathbb{B}} \chi^2 \varphi^{2m-2} h_n^2 d\varphi \wedge d^c \varphi \wedge R'\right) \left(\int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dv_1 \wedge d^c v_1 \wedge R'\right)$$
$$\leq \left(\int_{\mathbb{B}} \chi^2 \varphi^{2m-2} h_n^2 dd^c \varphi_1 \wedge R'\right) \left(\int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dd^c (v_1^2) \wedge R'\right)$$
$$\leq H_{1,m-1} I_{m,p}$$

where

$$H_{j,m} = \sup_{v_2,\ldots,v_p \in \mathsf{PSH}(\mathbb{B},[0,1])} \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dd^c \varphi_j \wedge R' \text{ for } j = 1,\ldots,q.$$

Then, by taking the supremum over all $v_1, \ldots, v_p \in \text{PSH}(\mathbb{B}, [0, 1])$, we have (3.5) $I_{m,p} \lesssim n^{\frac{m}{2^{q-1}}} + H_{1,m-1}.$ To estimate $H_{1,m-1}$, we consider $v_2, \ldots, v_p \in PSH(\mathbb{B}, [0, 1])$. It follows from Stokes' formula that, for $j = 1, \ldots, q-1$,

$$\int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n^2 dd^c \varphi_j \wedge R' = -2 \Big(\int_{\mathbb{B}} \chi \varphi^{2m} h_n^2 d\chi \wedge d^c \varphi_j \wedge R' + \int_{\mathbb{B}} \chi^2 \varphi^{2m-1} h_n^2 d\varphi \wedge d^c \varphi_j \wedge R' + \int_{\mathbb{B}} \chi^2 \varphi^{2m} h_n dh_n \wedge d^c \varphi_j \wedge R' \Big).$$

Let $L_{j,1}, L_{j,2}$, and $L_{j,3}$ be the first, second, and third integrals inside the brackets respectively. It follows from Lemma 3.4, Lemma 3.8, Cauchy-Schwarz inequality and the induction hypothesis on p that

$$\begin{split} L_{j,1}^{2} &\leq \Big(\int_{\mathbb{B}} \varphi^{2m} h_{n}^{2} d\chi \wedge d^{c}\chi \wedge R'\Big) \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R'\Big) \\ &\leq \Big(\int_{\mathrm{supp}(\chi)} \varphi^{2m} h_{n}^{2} \wedge R' \wedge \omega\Big) \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R'\Big) \\ &\leq I_{m,p-1,\mathrm{supp}(\chi)} \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R'\Big) \\ &\lesssim n^{\frac{m}{2q-1}} \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R'\Big), \\ L_{j,2}^{2} &\leq \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} h_{n}^{2} d\varphi \wedge d^{c}\varphi \wedge R'\Big) \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c}\varphi_{j} \wedge R'\Big) \end{split}$$

$$\int \mathcal{L}^{2} = \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m-2} h_{n}^{2} dd^{c} \varphi_{1} \wedge R'\right) \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R'\right)$$
$$\lesssim H_{1,m-1} \left(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R'\right),$$

and

$$\begin{split} L_{j,3}^{2} &\leq \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} dh_{n} \wedge d^{c}h_{n} \wedge R'\Big) \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R'\Big) \\ &\leq \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} T_{n} \wedge R'\Big) \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R'\Big) \\ &\leq J_{m,p-1, \text{supp}(\chi)} \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R'\Big) \\ &\lesssim n^{\frac{m}{2q-1}} \Big(\int_{\mathbb{B}} \chi^{2} \varphi^{2m} h_{n}^{2} d\varphi_{j} \wedge d^{c} \varphi_{j} \wedge R'\Big). \end{split}$$

Then, by taking the supremum over all $v_2, \ldots, v_p \in PSH(\mathbb{B}, [0, 1])$, Corollary 3.5, Lemma 3.8, and inequality (3.5), we have

$$H_{j,m}^2 \lesssim \begin{cases} H_{j+1,m} \Big(H_{1,m-1} + n^{\frac{m}{2q-1}} \Big) \text{ if } j < q-1 \\ n \Big(H_{1,m-1} + n^{\frac{m}{2q-1}} \Big)^2 \text{ if } j = q-1. \end{cases}$$

It thus follows that

$$\begin{cases} H_{k,m}^2 \lesssim H_{k+1,m} \Big(H_{1,m-1} + n^{\frac{m}{2q-1}} \Big) \text{ if } k < q-1 \\ H_{q-1,m} \lesssim \sqrt{n} \Big(H_{1,m-1} + n^{\frac{m}{2q-1}} \Big), \end{cases}$$

Then, we

$$\begin{split} H_{1,m-1}^{2^{q-2}} &\lesssim H_{q-1,m} \Big(H_{1,m-1} + n^{\frac{m}{2^{q-1}}} \Big)^{2^{q-2}-1} \lesssim \sqrt{n} \Big(H_{1,m-1} + n^{\frac{m}{2^{q-1}}} \Big)^{2^{q-2}}. \end{split}$$
 have
$$H_{1,m} \lesssim n^{\frac{1}{2^{q-1}}} \Big(H_{1,m-1} + n^{\frac{m}{2^{q-1}}} \Big), \end{split}$$

which implies

(3.6)
$$H_{1,m} \lesssim n^{\frac{m+1}{2q-1}}$$

Combining inequalities (3.5) and (3.6) gives

$$I_{m,p} \lesssim n^{\frac{m}{2q-1}},$$

as desired.

We end this subsection by estimates of Bedford-Taylor's capacity of sub-level sets, which will be used in the sequel. We recall that, for a compact subset $K \subset \Omega$, the Bedford-Taylor's capacity of K in Ω is defined by the formula

$$\operatorname{Cap}(K,\Omega) = \sup \left\{ \int_{K} (dd^{c}u)^{n} : u \in \operatorname{PSH}(\Omega), -1 \le u \le 0 \right\}.$$

We refer the reader to [BT82, Kol98, Kol05] for more information on this capacity.

For every Borel set $E \subset \mathbb{B}$, the relative extremal function for E in \mathbb{B} is defined as

$$u_E = \sup \left\{ u \in \mathsf{PSH}(\mathbb{B}) : u \le 0 \text{ on } \mathbb{B}, u \le -1 \text{ on } E \right\}$$

Let u_E^* be the upper semicontinuous regularization of u_E . Then we have the following relationship between the relative extremal functions and Bedford-Taylor's capacity.

$$\operatorname{Cap}(E,\mathbb{B}) = \int_{\mathbb{B}} (dd^{c}u_{E}^{*})^{k} = \int_{\overline{E}} (dd^{c}u_{E}^{*})^{k},$$

see [BT82]. Now, fix a compact subset K of \mathbb{B} . Let $2^{2^{q-1}} < \lambda < 2^{2^q}$ be a constant. For $n \in \mathbb{N}$, we set

$$K_n = \Big\{ z \in K : \varphi(z) \ge 2^n, \varphi_q \ge -\lambda^n \Big\},$$

and $u_n = u_{K_n}$.

We have the following estimates for Bedford-Taylor's capacity of K_n in \mathbb{B} .

Lemma 3.10. For every $m \in \mathbb{N}$, there is a constant c_m independent of φ such that for all $n \in \mathbb{N}$

$$\int_{K_n} (dd^c u_n^*)^k \le \operatorname{Cap}(K_n, \mathbb{B}) \le c_m \left(\frac{\lambda}{2^{2^q}}\right)^{\frac{mn}{2^{q-1}}}.$$

Proof. The first inequality is clear. We only need to prove the second inequality. Let l be a positive integer and V_l be an open subset in \mathbb{B} so that $\operatorname{Cap}(V_l, \mathbb{B}) \leq l^{-1}$ and φ is continuous on $\mathbb{B} \setminus V_l$. We have

(3.7)
$$\operatorname{Cap}(K_n, \mathbb{B}) \leq \operatorname{Cap}(K_n \setminus V_l, \mathbb{B}) + \operatorname{Cap}(V_l, \mathbb{B}) \leq \int_{K_n \setminus V_l} (dd^c u_{K_n \setminus V_l}^*)^k + l^{-1}.$$

Since $h_{2\lambda^n} \ge 1/2$ on K_n (recall that $h_n = 1 + \max(\varphi_q, -n)/n$), we have

$$\int_{K_n \setminus V_l} (dd^c u^*_{K_n \setminus V_l})^k \le 4 \times 4^{-mn} \int_{K_n \setminus V_l} h_{2\lambda^n}^2 \varphi^{2m} (dd^c u^*_{K_n \setminus V_l})^k \lesssim 4^{-mn} (2\lambda^n)^{\frac{m}{2^{q-1}}} \lesssim \left(\frac{\lambda}{2^{2^q}}\right)^{\frac{mn}{2^{q-1}}}$$

by Lemma 3.9. This combined with inequality (3.7) gives

$$\operatorname{Cap}(K_n, \mathbb{B}) \le c_m \left(\frac{\lambda}{2^{2^q}}\right)^{\frac{mn}{2^{q-1}}} + l^{-1}$$

Let $l \to +\infty$, we complete the proof.

3.2. Psh bound (local version). Now we construct the psh bound for functions in W_q^* .

Theorem 3.11. Let $\varphi \in W_q^*(\mathbb{B})$ with $\|\varphi\|_{*,q} \leq 1$ and $\alpha \in [1, 2^q)$. Then for every compact subset K of \mathbb{B} , there exist a strictly positive constant C and a psh function u on \mathbb{B} such that

$$|\varphi|^{\alpha} \le -u$$

on K and $||u||_{L^1(K)} \leq C$.

Proof. Fix $\alpha \in [1, 2^q)$ and $\lambda \in (\max(2^{\alpha}, 2^{2^{q-1}}), 2^{2^q})$. Assume that $\varphi \in W_q^*(\mathbb{B})$ with $\varphi \ge 0$. Let $(\varphi_1, \ldots, \varphi_q)$ be a defining sequence for φ such that

$$\|\varphi\|_{*,q} = \|\varphi\|_{L^2} + \sum_{j=1}^q \|dd^c \varphi_j\|^{1/2^j}.$$

We can assume that $\varphi_q \leq 0$. Consider

$$u = \sum_{n=1}^{\infty} 2^{n\alpha} \left(u_n^* + \frac{\max(\varphi_q, -\lambda^n)}{\lambda^n} \right) \cdot$$

Similar to the proof of [VV23, Theorem 1.3], we need to prove the following claims.

Claim 1. $2^{\alpha}u + 1 \leq -\varphi^{\alpha}$ outside a pluripolar set.

Claim 2. *u* is not identically $-\infty$.

Claim 3. One can choose u so that L^1 -norm of u is bounded by a constant that not depends on φ .

Proof of Claim 1. Set

$$A_n = \left\{ x \in \mathbb{B} : \varphi(x) \in [2^n, 2^{n+1}), \varphi_q > -\lambda^n \right\}.$$

Let $x \in \mathbb{B} \setminus (\bigcup_{n=1}^{\infty} A_n)$. Thus, there exists n such that $\varphi(x) \in [2^n, 2^{n+1})$ and $\varphi_q \leq -\lambda^n$. Thus

$$u(x) \le 2^{n\alpha} \left(\frac{\max(\varphi_q, -\lambda^n)}{\lambda^n}\right) \le -2^{n\alpha}$$

This means

$$2^{\alpha}u + 1 \le -\varphi^{\alpha}$$

on $\mathbb{B} \setminus (\bigcup_{n=1}^{\infty} A_n)$. Note that $u_n^* = u_n$ outside some pluripolar set E_n , on $A_n \setminus E_n$, we have $u(x) \leq 2^{n\alpha} u_n(x) = -2^{n\alpha}$.

Hence,

$$2^{\alpha}u + 1 \le -\varphi^c$$

on $\left(\bigcup_{n=1}^{\infty} A_n\right) \setminus \left(\bigcup_{n=1}^{\infty} E_n\right)$ and the Claim 1 follows.

Proof of Claim 2. Let

$$B_n = \Big\{ 2^{n\alpha} u_n^* < \frac{1}{n^2} \rho \Big\},$$

where $\rho(z) = ||z||^2 - 1$. By the comparison principle (see [Kol05, Theorem 1.16]) and Lemma 3.10, we have, for every $m \ge 1$, there is a constant $c_m > 0$ depending only on msuch that

$$\int_{B_n} \omega^k = \int_{B_n} (dd^c \rho)^k \le n^{2k} 2^{nk\alpha} \int_{B_n} (dd^c u_n^*)^k \le c_m n^{2k} 2^{nk\alpha} \left(\frac{\lambda}{2^{2q}}\right)^{\frac{mn}{2q-1}}$$

Then for every $n_0 \ge 1$,

$$\sum_{n\geq n_0} \int_{B_n} \omega^k \leq c_m \sum_{n\geq n_0} n^{2k} \left(2^{2^{q-1}k\alpha} \left(\frac{\lambda}{2^{2^q}} \right)^m \right)^{\frac{n}{2^{q-1}}}.$$

Thus by choosing m large enough (precisely so that $2^{2^{q-1}k\alpha} \left(\frac{\lambda}{2^{2^q}}\right)^m < 1$), we see that

$$\sum_{n\geq n_0}\int_{B_n}\omega^k<\int_{\mathbb{B}}\omega^k,$$

for n_0 large enough (independent of φ and K). In particular, $\mathbb{B} \setminus (\bigcup_{n \ge n_0} B_n) \neq \emptyset$ and

$$0 \ge 2^{n\alpha} u_n^*(x) \ge \frac{1}{n^2} \rho(x), \text{ for all } n \ge n_0, \ x \in \mathbb{B} \setminus \big(\cup_{n \ge n_0} B_n \big).$$

Since $\mathbb{B} \setminus (\bigcup_{n \ge n_0} B_n)$ has a positive measure, it is a non-pluripolar set. It thus follows that we can choose $x_0 \in \mathbb{B} \setminus (\bigcup_{n \ge n_0} B_n)$ such that $\varphi_q(x_0) \neq -\infty$. Now we observe that

$$u(x_0) = \sum_{n \ge 1} 2^{n\alpha} \left(\frac{\max(\varphi_q(x_0), -\lambda^n)}{\lambda^n} \right) + \sum_{n \ge 1} 2^{n\alpha} u_n^*(x_0)$$

$$\ge \varphi_q(x_0) \sum_{n \ge 1} \left(\frac{2^{\alpha}}{\lambda} \right)^n + \sum_{1 \le n \le n_0} 2^{n\alpha} u_n^*(x_0) + \sum_{n > n_0} 2^{n\alpha} u_n^*(x_0)$$

$$\ge \varphi_q(x_0) \sum_{n \ge 1} \left(\frac{2^{\alpha}}{\lambda} \right)^n - \sum_{1 \le n \le n_0} 2^{n\alpha} + \sum_{n > n_0} \frac{1}{n^2} \rho(x_0) > -\infty.$$

This proves Claim 2.

Proof of Claim 3. Define

$$M_{\varphi} = \inf \left\{ \|u\|_{L^{1}(K)} : |\varphi|^{\alpha} \le -u, u \in \mathrm{PSH}(\mathbb{B}) \right\}.$$

Let

$$M = \sup\left\{M_{\varphi}, \varphi \in W_q^*, \varphi \ge 0, \|\varphi\|_{*,q} \le 1\right\}$$

Suppose by contradiction that $M = +\infty$. Hence, we can find a sequence $(\varphi_n)_n$ in $W_q^*(\mathbb{B})$ with $\|\varphi\|_{*,q} \leq 1$ such that $M_{\varphi_n} \geq 2^n$. Define

$$v = \sum_{n \ge 1} \frac{\varphi_n}{n^2} \cdot$$

Then, since W_q^* is a quasi-Banach space, $v \in W_q^*$. Hence, by Claim 2, we can find a negative psh function u on \mathbb{B} with $|u| \ge v^{\alpha}$. This deduce that $|u| \ge n^{-2\alpha}\varphi_n^{\alpha}$ and hence, $n^{2\alpha}|u| \ge \varphi_n^{\alpha}$. It follows that

$$M_{\varphi_n} \le \|n^{2\alpha}u\|_{L^1(K)} = n^{2\alpha}\|u\|_{L^1(K)} \ll 2^n,$$

when n is big enough. This is a contradiction since $M_{\varphi_n} \geq 2^n$. So $M < +\infty$ and the Claim 3 follows.

Now, for general φ , we decompose $\varphi = \max(\varphi, 0) + \min(\varphi, 0)$. By Example 2.8, we can rescale the function a little bit to get the answer. This completes the proof of Theorem 3.11.

Example 3.12. (Sharpness of the exponential α) As in [VV23, page 13], we consider the case when $\alpha > 2^q$ and k = 1. We choose the function φ to be $(-\log |z|^2)^{1/2^q-\delta}$ where $\delta \in (0, 1/2^q)$. Then by Example 2.9, $\varphi \in W_q^*$. Since $\alpha > 2^q$, we can choose δ small enough such that $\beta = \alpha(1/2^q - \delta) > 1$. Thus, $\varphi^{\alpha} = (-\log |z|^2)^{\beta}$ and then $e^{c\varphi^{\alpha}} \ge \frac{c}{|z|^2}$ which is not locally integrable at 0 in \mathbb{C} . Also, note that by arguments in [VV23], this function is not bounded from above by minus of a subharmonic function. So the exponential coefficient here as well as the one in Theorem 1.2 can not be greater than 2^q . It is natural to predict that the result still holds for $\alpha = 2^q$ (like in the case q = 1), but currently, we don't know how to prove that.

3.3. **Psh bound (global version).** The following result is a global version of Theorem 3.11.

Theorem 3.13. Let $\varphi \in W_q^*(X)$ with $\|\varphi\|_{*,q} \leq 1$ and $\alpha \in [1, 2^q)$. Then there exist a strictly positive constant C not depending on φ and a negative $C\omega$ -psh u on X such that

 $|\varphi|^{\alpha} \leq -u$ and $||u||_{L^{1}(X)} \leq C.$

Proof. We follow the proof of Theorem 3.11 almost line by line with the only additional consideration being the careful selection of u_n to ensure that it is $C\omega$ -psh for some uniform constant C. Recall that the capacity of a Borel set E in X with respect to a Kähler form η on X is defined as

$$\operatorname{Cap}_{\eta}(E) = \sup \Big\{ \int_{X} (\eta + dd^{c}v)^{k} : v \in \operatorname{PSH}(X, \eta), \ -1 \le v \le 0 \Big\}.$$

Following [GZ05, Proposition 3.1, Theorem 3.2], we have

$$\mathsf{Cap}_{\eta}(E) = \int_{X} -u_{E,\eta}^{*}(\eta + dd^{c}u_{E,\eta}^{*})^{k} = \int_{\overline{E}} -u_{E,\eta}^{*}(\eta + dd^{c}u_{E,\eta}^{*})^{k}$$

where

$$u_{E,\eta} = \sup \Big\{ u \in \mathsf{PSH}(X,\eta) : u \le 0 \text{ on } X \text{ and } u \le -1 \text{ on } E \Big\}.$$

Let $\alpha \in [1, 2^q)$ and take λ such that $2^{\alpha} < \lambda$. As in Theorem 3.11, it suffices to take $\varphi \in W_q^*(X)$ with $\varphi \ge 0$. Let $((c_1, \varphi_1), \ldots, (c_q, \varphi_q))$ be a defining sequence such that

$$\|\varphi\|_{*,q} = \|\varphi\|_{L^2} + \sum_{j=1}^q c_j^{1/2^j}.$$

We can assume that $\varphi_q \leq 0$. Consider

$$X_n = \Big\{ x \in X : \varphi(x) \in [2^n, 2^{n+1}), \ \varphi_q(x) > -\lambda^n \Big\}.$$

We set $u_n = u_{X_n, 3^{-n\alpha}\omega}$ and

$$u = \sum_{n=1}^{\infty} 2^{n\alpha} \left(u_n^* + \frac{\max(\varphi_q, -\lambda^n)}{\lambda^n} \right) \cdot$$

It follows from the proof of Lemma 3.10 that for every $m \in \mathbb{N}$, there exists a constant c_m independent of φ such that, for all $n \in \mathbb{N}$,

(3.8)
$$\operatorname{Cap}_{3^{-n\alpha}\omega}(X_n) \le c_m \left(\frac{\lambda}{2^{2^q}}\right)^{\frac{mn}{2^{q-1}}}$$

As before, we also have $2^{\alpha}u + 1 \leq -\varphi^{\alpha}$ outside a pluripolar set. Let

$$\eta = \sum_{n=1}^{\infty} 2^{n\alpha} (3^{-n\alpha}\omega) + 2^{n\alpha}\lambda^{-n}c_q\omega \le C\omega$$

for some C > 0 depending only on α and λ (note that $c_q \leq 1$). Since u_n^* is $3^{-n\alpha}\omega$ -psh and φ_q is $c_q \omega$ -psh, we have u is $C \omega$ -psh if $u \neq -\infty$. It remains to check that $u \neq -\infty$. We argue as in the proof of Theorem 3.11. Let

$$Y_n = \left\{ 2^{n\alpha} u_n^* < -\frac{1}{n^2} \right\} \cdot$$

It follows from the comparison principle (see [Kol05, Theorem 6.4]) and inequality (3.8) that, for any $m \in \mathbb{N}$,

$$\int_{Y_n} (3^{-n\alpha}\omega)^k \le \int_{Y_n} (3^{-n\alpha}\omega + dd^c u_n^*)^k \le 2^{n\alpha} n^2 \int_{Y_n} -u_n^* (3^{-n\alpha}\omega + dd^c u_n^*)^k \le 2^{n\alpha} n^2 \operatorname{Cap}_{3^{-n\alpha}\omega}(X_n) \le c_m 2^{n\alpha} n^2 \left(\frac{\lambda}{2^{2q}}\right)^{\frac{mn}{2q-1}}.$$

Hence, for every $n_0 \ge 1$, we have

$$\sum_{n \ge n_0} \int_{Y_n} \omega^k \le c_m \sum_{n \ge n_0} 2^{n\alpha} 3^{nk\alpha} n^2 \left(\frac{\lambda}{2^{2^q}}\right)^{\frac{mn}{2^{q-1}}} < c_m \sum_{n \ge n_0} n^2 \left(3^{(k+1)\alpha} \left(\frac{\lambda}{2^{2^q}}\right)^{\frac{m}{2^{q-1}}}\right)^n.$$

Since $\lambda < 2^{2^q}$, we can choose m large enough such that

$$3^{(k+1)\alpha} \left(\frac{\lambda}{2^{2^q}}\right)^{\frac{m}{2^{q-1}}} < 1$$

It thus follows that

$$\sum_{n \ge n_0} \int_{Y_n} \omega^k < 1 = \int_X \omega^k$$

for n_0 large enough (independent of φ). In particular, $X \setminus (\bigcup_{n \ge n_0} Y_n) \neq \emptyset$ and

$$0 \ge 2^{n\alpha} u_n^*(x) \ge -\frac{1}{n^2}, \text{ for all } n \ge n_0, \ x \in X \setminus \left(\bigcup_{n \ge n_0} Y_n \right).$$

Since $X \setminus (\bigcup_{n \ge n_0} Y_n)$ has a positive measure, it is a non-pluripolar set. It thus follows that we can choose $x_0 \in X \setminus (\bigcup_{n \ge n_0} Y_n)$ such that $\varphi_q(x_0) \neq -\infty$. Now we observe that

$$u(x_0) = \sum_{n \ge 1} 2^{n\alpha} \left(\frac{\max(\varphi_q(x_0), -\lambda^n)}{\lambda^n} \right) + \sum_{n \ge 1} 2^{n\alpha} u_n^*(x_0)$$

$$\ge \varphi_q(x_0) \sum_{n \ge 1} \left(\frac{2^{\alpha}}{\lambda} \right)^n + \sum_{1 \le n \le n_0} 2^{n\alpha} u_n^*(x_0) + \sum_{n > n_0} 2^{n\alpha} u_n^*(x_0)$$

$$\ge \varphi_q(x_0) \sum_{n \ge 1} \left(\frac{2^{\alpha}}{\lambda} \right)^n - \sum_{1 \le n \le n_0} 2^{n\alpha} - \sum_{n > n_0} \frac{1}{n^2} > -\infty.$$

The proof is thus completed.

4. COMPLEX MONGE-AMPÈRE EQUATION

In this section, we study the relationship between W_q^* and the complex Monge-Ampère operator.

4.1. Relationship between the classes $W_q^*(X)$, $\mathcal{E}(X,\omega)$ and $\mathcal{E}^p(X,\omega)$. Let (X,ω) be a compact Kähler manifold of dimension k. We recall some definitions from [GZ07]. Let φ be some unbounded ω -psh function on X and consider $\varphi_j = \max(\varphi, -j)$ be canonical approximation of φ by bounded ω -psh functions. By [BT87], we can define the Monge-Ampère measure $(\omega + dd^c \varphi_j)^k$. The sequence of measures

$$\mathbf{1}_{\{\varphi>-j\}}\left(\omega+dd^{c}\varphi_{j}\right)^{k}$$

is an increasing sequence and converges to the non-pluripolar Monge-Ampère measure μ_{φ} of φ . Its total mass $\mu_{\varphi}(X)$ can take any value in $\left[0, \int_{X} \omega^{k}\right]$. Define

$$\mathcal{E}(X,\omega) = \Big\{ \varphi \in \mathrm{PSH}(X,\omega) : \mu_{\varphi}(X) = \int_{X} \omega^{k} \Big\}.$$

Recall the following criterion for functions in $\mathcal{E}(X, \omega)$.

Lemma 4.1. [GZ07, Lemma 1.2] Fix $\varphi \in PSH(X, \omega)$ and define $\varphi_j = \max(\varphi, -j)$ for $j \in \mathbb{N}$. Let (s_j) be any sequence of real numbers converging to $+\infty$, such that $s_j \leq j$ for all $j \in \mathbb{N}$. The following conditions are equivalent

(1)
$$\varphi \in \mathcal{E}(X, \omega);$$

(2) $(\omega + dd^c \varphi_j)^k (\varphi \leq -j) \to 0;$
(3) $(\omega + dd^c \varphi_j)^k (\varphi \leq -s_j) \to 0.$

Now, we prove that any ω -psh complex Sobolev function belongs to $\mathcal{E}(X, \omega)$.

Proof of Theorem 1.4(1). Assume that φ is a negative ω -psh function in $W_1^*(X)$ and $\varphi_j = \max(\varphi, -j)$. We note that φ_j/j equals to -1 when $\varphi \leq -j$ and equals to $\varphi/j < 0$ when $\varphi > -j$. Let T_{φ} be a positive closed current on X such that $d\varphi \wedge d^c\varphi \leq T_{\varphi}$. By [Vig07] or

Example 2.8, we have $d\varphi_j \wedge d^c \varphi_j \leq T_{\varphi}$. By Stokes' formula, we have

$$\begin{split} \int_{\{\varphi \leq -j\}} (\omega + dd^c \varphi_j)^k &\leq -\int_X \frac{\varphi_j}{j} (\omega + dd^c \varphi_j)^k \\ &= \frac{1}{j} \Big(\int_X d\varphi_j \wedge d^c \varphi_j \wedge \sum_{m=0}^{k-1} (\omega + dd^c \varphi_j)^{k-1-m} \wedge \omega^m - \int_X \varphi_j \omega^k \Big) \\ &\leq \frac{1}{j} \Big(\int_X T_\varphi \wedge \sum_{m=0}^{k-1} (\omega + dd^c \varphi_j)^{k-1-m} \wedge \omega^m - \int_X \varphi \omega^k \Big) \cdot \end{split}$$

Since T_{φ} is closed, the first integral doesn't change if we replace the closed current $\sum_{m=0}^{k-1} (\omega + dd^c \varphi_j)^{k-1-m} \wedge \omega^m$ by a closed form in its de Rham cohomology class. We can replace it by $k\omega^{k-1}$ and obtain

$$\int_{\{\varphi \le -j\}} (\omega + dd^c \varphi_j)^k \le \frac{k \cdot \|T_\varphi\|}{j} - \frac{1}{j} \Big(\int_X \varphi \omega^k \Big) \to 0 \text{ as } j \to +\infty \cdot$$

This, combined with Lemma 4.1, finishes our proof.

Next, we recall the definition of finite energy classes in [GZ07]. For simplicity, assume that $\int_X \omega^k = 1$. Denote $\omega_{\varphi} = \omega + dd^c \varphi$. For bounded ω -psh functions, define the energy functional

$$E_p(\varphi) = -\frac{1}{k+1} \sum_{m=0}^k \int_X (-\varphi)^p (\omega_{\varphi})^m \wedge \omega^{k-m} \cdot$$

We can extend this functional for arbitrary ω -psh functions by canonical approximation as above

$$E_p(\varphi) = \lim_{j \to +\infty} E_p(\varphi_j).$$

So, we can define the finite energy class

$$\mathcal{E}^p(X,\omega) = \{\varphi \in \mathcal{E}(X,\omega) : E_p(\varphi) > -\infty\}.$$

To prove Theorem 1.4(2), we first recall some facts about finite energy classes that will be used in sequel. We only use the results for the class $\mathcal{E}^p(X,\omega)$ with $p \ge 1$. For more general classes, we refer the readers to [GZ07] and [DV22].

Proposition 4.2. Let φ and ψ be bounded non-positive ω -psh functions and $p \ge 1$. Then, for every positive closed current T of bi-dimension (1, 1), we have

$$0 \le \int_X (-\varphi)^p \omega_\psi \wedge T \le 2p \int_X (-\varphi)^p \omega_\varphi \wedge T + 2p \int_X (-\psi)^p \omega_\psi \wedge T.$$

Recall that $\omega_{\varphi} = \omega + dd^c \varphi$ and $\omega_{\psi} = \omega + dd^c \psi$. For a proof of this proposition, see [GZ07, Proposition 3.6].

Proposition 4.3. Let $\varphi_0, \ldots, \varphi_k$ be bounded non-positive ω -psh functions and $p \ge 1$. Then there exists a strictly positive constant C_p depending only on p such that

$$\int_X (-\varphi_0)^p \omega_{\varphi_1} \wedge \dots \wedge \omega_{\varphi_k} \le C_p \max_{0 \le m \le k} \Big(\int_X (-\varphi_m)^p \omega_{\varphi_m}^k \Big).$$

See [GZ07, Proposition 3.8] for a proof. We will use the following direct corollary.

Corollary 4.4. Let φ and ψ be non-positive functions in $\mathcal{E}^p(X, \omega)$ with $p \ge 1$. Then for $0 \le m \le k - 1$, there exists a strictly positive constant C_p depending only on p such that

$$\int_X (-\varphi)^p \omega_\psi \wedge \omega_\varphi^{k-m-1} \wedge \omega^m \le C_p \Big(\int_X (-\varphi)^p \omega_\varphi^k + \int_X (-\psi)^p \omega_\psi^k \Big)$$

Here the measure $\omega_{\psi} \wedge \omega_{\varphi}^{k-m-1} \wedge \omega^{m}$ is defined in the non-pluripolar sense (see [GZ17, Chapter 10.2.3].

Proof. Let $\varphi_j = \max(\varphi, -j)$ and $\psi_j = \max(\psi, -j)$. By [GZ17, Theorem 10.18],

$$\int_X (-\varphi_j)^p \omega_{\psi} \wedge \omega_{\varphi}^{k-m-1} \wedge \omega^m = \lim_{j' \to +\infty} \int_X (-\varphi_j)^p \omega_{\psi_{j'}} \wedge \omega_{\varphi_{j'}}^{k-m-1} \wedge \omega^m$$

By Proposition 4.3, we have

$$\begin{split} \int_X (-\varphi_j)^p \omega_{\psi_{j'}} \wedge \omega_{\varphi_{j'}}^{k-m-1} \wedge \omega^m &\leq C_p \Big(\int_X (-\varphi_j)^p \omega_{\varphi_j}^k + \int_X (-\varphi_{j'})^p \omega_{\varphi_{j'}}^k + \int_X (-\psi_{j'})^p \omega_{\psi_{j'}}^k \Big) \\ &\leq C_p \Big(\int_X (-\varphi)^p \omega_{\varphi}^k + \int_X (-\psi)^p \omega_{\psi}^k \Big). \end{split}$$

Let $j' \to +\infty$ and then let $j \to +\infty$ give us the result.

Proof of Theorem 1.4(2). We prove this by induction. First, consider the case when q = 2. Let ψ be the function in $W_1^*(X)$ such that $d\varphi \wedge d^c\varphi \leq C(\omega + dd^c\psi)$. By Theorem 1.4(1), we have $\varphi, \psi \in \mathcal{E}(X, \omega)$. Now, by using Stokes' formula, we have

$$-E_1(\varphi) = \frac{1}{k+1} \sum_{m=1}^k \sum_{l=0}^{m-1} \int_X d\varphi \wedge d^c \varphi \wedge \omega_{\varphi}^l \wedge \omega^{k-l-1}$$
$$\leq \frac{C}{k+1} \sum_{m=1}^k \sum_{l=0}^{m-1} \int_X \omega_{\psi} \wedge \omega_{\varphi}^l \wedge \omega^{k-l-1}$$
$$= \frac{C}{k+1} \sum_{m=1}^k \sum_{l=0}^{m-1} 1 < +\infty.$$

So, $\varphi \in \mathcal{E}^1(X, \omega)$. Assume now that our theorem is true for q-1 with q > 2. Let ψ be an ω -psh function in $W^*_{q-1}(X)$ such that $d\varphi \wedge d^c\varphi \leq C(\omega + dd^c\psi)$. By the induction hypothesis, we have $\varphi, \psi \in \mathcal{E}^{q-2}(X, \omega)$. We observe that

$$dd^{c}((-\varphi)^{q}/q) = (q-1)(-\varphi)^{q-2}d\varphi \wedge d^{c}\varphi - (-\varphi)^{q-1}dd^{c}\varphi.$$

Then, by using Stokes's formula, we can write

$$\begin{split} \int_X (-\varphi)^{q-1} dd^c \varphi \wedge T &= (q-1) \int_X (-\varphi)^{q-2} d\varphi \wedge d^c \varphi \wedge T - \int_X dd^c ((-\varphi)^q/q) \wedge T \\ &= (q-1) \int_X (-\varphi)^{q-2} d\varphi \wedge d^c \varphi \wedge T, \end{split}$$

for T is a sufficiently regular positive closed current of bi-dimension (1, 1). This deduces that

$$-E_{q-1}(\varphi) = \frac{1}{k+1} \sum_{m=0}^{k} \int_{X} (-\varphi)^{q-1} (\omega_{\varphi})^{m} \wedge \omega^{k-m}$$

$$= \int_{X} (-\varphi)^{q-1} \omega^{k} + \frac{q-1}{k+1} \sum_{m=1}^{k} \sum_{l=0}^{m-1} \int_{X} (-\varphi)^{q-2} d\varphi \wedge d^{c} \varphi \wedge (\omega_{\varphi})^{l} \wedge \omega^{k-l-1}$$

$$\leq \int_{X} (-\varphi)^{q-1} \omega^{k} + \frac{C(q-1)}{k+1} \sum_{m=1}^{k} \sum_{l=0}^{m-1} \int_{X} (-\varphi)^{q-2} \omega_{\psi} \wedge (\omega_{\varphi})^{l} \wedge \omega^{k-l-1}$$

$$< +\infty$$

by applying Corollary 4.4 for p = q - 2 and the induction hypothesis. So $\varphi \in \mathcal{E}^{q-1}(X, \omega)$ and we complete our proof.

Remark 4.5. The above theorem gives us a lower bound for p(q) by q-1. From Example 3.12 and [DGL20, Theorem 2.1], we see that p(q) has an upper bound by $k(2^q - 1)$. It is an interesting question to know what is the best choice for p(q). It has been noted in [Vig07] that $W_1^*(X)$ is a BMO space and hence by [JN61], there are constants c and A such that $\int_X e^{c|\varphi|} \omega^k \leq A$ for all φ in $W_1^*(X)$ with $\|\varphi\|_{*,1} \leq 1$. This fact can be used instead of Skoda's integrability theorem in the proof of Theorem 2.1 in [DGL20].

4.2. Relationship between the classes $W_q^*(\Omega)$ and $\mathcal{D}(\Omega)$. We will prove Theorem 1.5. First, we recall Błocki's criterion for functions in class $\mathcal{D}(\Omega)$.

Theorem 4.6. [Blo06, Theorem 1.1] Let Ω be a domain in \mathbb{C}^k and φ be a negative psh function on Ω , the following properties are equivalent

- (1) There exists a measure μ in Ω such that if $U \subset \Omega$ is open and a sequence $\varphi_n \in PSH(U) \cap C^{\infty}(U)$ is decreasing to u in U, then $(dd^c \varphi_n)^k$ tends weakly to μ in U;
- (2) For every open subset U of Ω and any sequence $\varphi_n \in PSH(U) \cap C^{\infty}(U)$ decreasing to φ in U, the sequence $(dd^c\varphi_n)^k$ is locally weakly bounded in U;
- (3) For every open subset U of Ω and any sequence $\varphi_n \in PSH(U) \cap C^{\infty}(U)$ decreasing to φ in U, the sequences

(4.1)
$$|\varphi_n|^{k-p-2} d\varphi_n \wedge d^c \varphi_n \wedge (dd^c \varphi_n)^p \wedge \omega^{k-p-1}, \qquad p = 0, \dots, k-2,$$

are locally weakly bounded in U;

(4) For every z ∈ Ω there exists an open neighborhood U of z in Ω and a sequence φ_n ∈ PSH(U) ∩ C[∞](U) decreasing to φ in U such that the sequences in (4.1) are locally weakly bounded in U.

The following result is the key point in our proof of Theorem 1.5.

Proposition 4.7. Let $1 \le p \le k-1$ and $0 \le m \le k-p$. Let q_1, \ldots, q_{p+1} be integers satisfying

(1)
$$q_{p+1} \ge \max_{1 \le j \le p} q_j$$
,
(2) $\sum_{j=1}^p q_j \ge p(p-1+m)$,

(3) $\max_{1 \le j \le p} q_j - \min_{1 \le j \le p} q_j \le 1.$

Let $\varphi_1, \ldots, \varphi_{p+1}$ be negative smooth psh functions on Ω such that, for $j = 1, \ldots, p+1$, $\|\varphi_j\|_{*,q_j} \leq 1$. Assume that there exists a defining sequence $(\varphi_{j,1}, \ldots, \varphi_{j,q_j})$ of φ_j such that $\varphi_{j,l}$ is smooth for $l = 1, \ldots, q_j$, $\|\varphi_{j,l}\|_{*,q_j-l} \leq 1$ for $l = 1, \ldots, q_j - 1$ and $\|\varphi_{j,q_j}\|_{L^1} \leq 1$. Then for every compact subset K of Ω , there is a constant C > 0 depending only on K, m and p such that

$$\int_{K} (-\varphi_{p+1})^{m} dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p} \wedge \omega^{k-p} \leq C.$$

Proof. We prove by induction on p and m. Fix a compact set K and a cut-off function χ on Ω such that $0 \le \chi \le 1$ and $\chi = 1$ on K.

Consider the case p = 1. If m = 0 then the desired property is trivial. Assume that the desired property is true for $0, \ldots, m-1$ for some $1 \le m \le k-p$. We now prove it for m. By hypothesis, $q_2 \ge q_1 \ge m$. Hence, there exist smooth psh functions ψ_1, ψ_2 satisfying $\|\psi_1\|_{*,m-1} \le 1$, $\|\psi_2\|_{*,m-1} \le 1$ such that

(4.2)
$$d\varphi_1 \wedge d^c \varphi_1 \leq dd^c \psi_1, \quad d\varphi_2 \wedge d^c \varphi_2 \leq dd^c \psi_2$$

By Stokes' formula, we have

(4.3)
$$\int_{K} (-\varphi_2)^m dd^c \varphi_1 \wedge \omega^{k-1} \lesssim \left| \int_{\Omega} \chi(-\varphi_2)^{m-1} d\varphi_2 \wedge d^c \varphi_1 \wedge \omega^{k-1} + \left| \int_{\Omega} (-\varphi_2)^m d\chi \wedge d^c \varphi_1 \wedge \omega^{k-1} \right|,$$

where \leq denotes \leq modulo a multiplicative constant depending only on *m* and *K*. For the first term, by using Cauchy-Schwarz inequality and inequalities (4.2), we can bound it from above by the square root of

$$\left(\int_{\Omega} \chi(-\varphi_2)^{m-1} dd^c \psi_1 \wedge \omega^{k-1}\right) \left(\int_{\Omega} \chi(-\varphi_2)^{m-1} dd^c \psi_2 \wedge \omega^{k-1}\right).$$

It follows from the induction hypothesis for m - 1 that both factors are bounded by a constant depending only on $supp(\chi)$ (and hence only on K). Hence, the first term of the RHS of inequality (4.3) is bounded by a constant depending only on K. For the second term of the RHS of inequality (4.3), by using Cauchy-Schwarz inequality and inequalities (4.2), we can bound it from above by the square root of

$$\left(\int_{\Omega} (-\varphi_2)^{m+1} d\chi \wedge d^c \chi \wedge \omega^{k-1}\right) \left(\int_{\operatorname{supp}(\chi)} (-\varphi_2)^{m-1} dd^c \psi_1 \wedge \omega^{k-1}\right).$$

We use the induction hypothesis for m - 1 for the second factor. The first factor can be bound by using Corollary 3.2. Hence, the second term of the RHS of inequality (4.3) is bounded by a constant depending only on K. The proof for the case p = 1 is thus complete.

Now, we consider the case when p > 1. Assume that the desired property is true for p - 1, where $2 \le p \le k - 1$. We prove that it is true for p. We can further assume that $q_p = \max_{1 \le j \le p} q_j$. Consider the case m = 0. We note that

$$dd^{c}\varphi_{1}\wedge\cdots\wedge dd^{c}\varphi_{p}=dd^{c}\left(\varphi_{p}dd^{c}\varphi_{1}\wedge\cdots\wedge dd^{c}\varphi_{p-1}\right).$$

Hence, by Stokes' formula, we have

$$\int_{K} dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p} \wedge \omega^{k-p} \leq \int_{\Omega} \chi dd^{c} (\varphi_{p} dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p-1}) \wedge \omega^{k-p} \\
= \int_{\Omega} (-\varphi_{p}) dd^{c} \chi \wedge dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p-1} \wedge \omega^{k-p} \\
\leq \int_{\mathrm{supp}(\chi)} (-\varphi_{p}) dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p-1} \wedge \omega^{k-p+1}.$$

We now only need to check if $\varphi_1, \ldots, \varphi_p$ satisfies the induction hypothesis for p-1 and m = 1. Indeed, condition (2) becomes

$$(4.4) q_1 + \dots + q_p \ge p(p-1)$$

Condition (3) thus becomes

(4.5)
$$q_p - q_j \le 1 \text{ for every } j = 1, \dots, p-1$$

Combining inequalities (4.4) and (4.5) gives us

$$\sum_{j=1}^{p-1} q_j = \frac{p-1}{p} \sum_{j=1}^{p-1} q_j + \frac{1}{p} \sum_{j=1}^{q-1} q_j \ge \frac{p-1}{p} \sum_{j=1}^{p-1} q_j + \frac{1}{p} \sum_{j=1}^{q-1} (q_p - 1)$$
$$= \frac{p-1}{p} \left(\sum_{j=1}^p q_j - 1 \right) \ge \frac{p-1}{p} \cdot p(p-1)$$
$$= (p-1)(p-1) = (p-1)(p-2+1).$$

It follows that q_1, \ldots, q_p satisfy the condition (2) for p-1 and m = 1. Obviously, q_1, \ldots, q_p satisfy the conditions (1) and (3) for p-1 and m = 1. Hence, by applying the induction hypothesis for p-1 and m = 1, we get the statement for m = 0.

Assume that the statement is true for $0, \ldots, m-1$, where $1 \le m \le k-p$. We now prove it for m. We can further assume that $q_p = \max_{1 \le j \le p} q_j$. Since $q_{p+1} \ge q_p$, there exist smooth psh functions ϕ_p, ϕ_{p+1} satisfying $\|\phi_p\|_{*,q_p-1} \le 1$, $\|\phi_{p+1}\|_{*,q_p-1} \le 1$ such that

(4.6)
$$d\varphi_p \wedge d^c \varphi_p \leq dd^c \phi_p, \quad d\varphi_{p+1} \wedge d^c \varphi_{p+1} \leq dd^c \phi_{p+1}.$$

We need to bound, for $\varphi = \varphi_{p+1}$,

$$\int_{K} \chi(-\varphi)^{m} dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p} \wedge \omega^{k-p}$$

By Stokes' formula, we have

(4.7)
$$\int_{K} \chi(-\varphi)^{m} dd^{c} \varphi_{1} \wedge \cdots \wedge dd^{c} \varphi_{p} \wedge \omega^{k-p}$$
$$\lesssim \left| \int_{\Omega} (-\varphi)^{m} d\chi \wedge d^{c} \varphi_{p} \wedge dd^{c} \varphi_{1} \wedge \cdots \wedge dd^{c} \varphi_{p-1} \wedge \omega^{k-p} \right|$$
$$+ \left| \int_{\Omega} \chi(-\varphi)^{m-1} d\varphi \wedge d^{c} \varphi_{p} \wedge dd^{c} \varphi_{1} \wedge \cdots \wedge dd^{c} \varphi_{p-1} \wedge \omega^{k-p} \right|$$

where \leq denotes \leq modulo a multiplicative constant depending only on *m* and *K*. For the second term of the RHS of inequality (4.7), by using Cauchy-Schwarz inequality and

inequalities (4.6), we can bound it from above by the square root of

$$\left(\int_{\Omega} \chi(-\varphi)^{m-1} dd^{c} \phi_{p} \wedge dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p-1} \wedge \omega^{k-p}\right) \\ \times \left(\int_{\Omega} \chi(-\varphi)^{m-1} dd^{c} \phi_{p+1} \wedge dd^{c} \varphi_{1} \wedge \dots \wedge dd^{c} \varphi_{p-1} \wedge \omega^{k-p}\right)$$

We note that $q_p - 1, q_1, \ldots, q_{p-1}, q_{p+1}$ satisfy the conditions (1), (2), and (3) for p and m-1. It follows from the induction hypothesis for m-1 that both factors are bounded by a constant depending only on $supp(\chi)$ (and hence only on K). Hence, the second term of the RHS of inequality (4.7) is bounded by a constant depending only on K. For the first term of the RHS of inequality (4.7), by using Cauchy-Schwarz inequality and inequalities (4.6), we can bound it from above by the square root of

$$\left(\int_{\Omega} (-\varphi)^{m+1} d\chi \wedge d^{c}\chi \wedge dd^{c}\varphi_{1} \wedge \dots \wedge dd^{c}\varphi_{p-1} \wedge \omega^{k-p}\right) \times \left(\int_{\mathrm{supp}(\chi)} (-\varphi)^{m-1} dd^{c}\phi_{p} \wedge dd^{c}\varphi_{1} \wedge \dots \wedge dd^{c}\varphi_{p-1} \wedge \omega^{k-p}\right).$$

To deal with the second factor, we note that $q_p - 1, q_1, \ldots, q_{p-1}, q_{p+1}$ satisfy the conditions (1), (2) and (3) for p and m - 1. Thus, we can bound the second factor by using the induction hypothesis for m - 1. To deal with the first factor, we first observe that $q_1, \ldots, q_{p-1}, q_{p+1}$ satisfy the conditions (1) and (3) for p - 1 and m + 1. For the condition (2), as in the case m = 0, we have

$$\sum_{j=1}^{p} q_j \ge p(p-1+m),$$

and

$$q_p - q_j \le 1$$
, for every $j = 1, ..., p - 1$.

It follows that

$$\sum_{j=1}^{p-1} q_j \ge \frac{p-1}{p} \left(\sum_{j=1}^p q_j - 1 \right) \ge \frac{p-1}{p} \cdot p(p-1+m) = (p-1)(p-1+m).$$

Hence $q_1, \ldots, q_{p-1}, q_{p+1}$ satisfy the conditions (2) for p-1 and m+1. Thus, we can bound the first factor by using the induction hypothesis for p-1 and m+1. Therefore, we can bound the first term of the RHS of inequality (4.7). The proof is complete.

We now prove that psh q-complex Sobolev functions belong to $\mathcal{D}(\Omega)$ for $q \ge k-1$.

End of proof of Theorem 1.5. Let $\varphi \in W^*_{k-1,\text{loc}}(\Omega) \cap \text{PSH}(\Omega)$. Since the problem is local, we can assume that $\varphi \in W^*_{k-1}(\Omega) \cap \text{PSH}(\Omega)$. We will prove that φ satisfies the condition (4) of Proposition 4.6. Indeed, we consider an open relatively compact subset U of Ω and let (φ_n) be the sequence of smooth psh functions constructed in the proof of Theorem 2.11. We note that $\|\varphi_n\|_{W^*_{k-1}(U)} \leq 2\|\varphi\|_{W^*_{k-1}(\Omega)}$ for every n. Let $(\varphi_{0,1}, \ldots, \varphi_{0,k-1})$ be a smooth defining sequence for φ . By the proof of Theorem 2.11, we can construct smooth defining sequence $(\varphi_{n,1}, \ldots, \varphi_{n,k-1})$ for φ_n such that $\|\varphi_{n,l}\|_{W^*_{k-1-l}(U)} \leq 2\|\varphi_{0,l}\|_{W^*_{k-1-l}(\Omega)}$ for $1 \leq l \leq 1$

k-2, $\|\varphi_{n,k-1}\|_{L^1(U)} \leq 2\|\varphi_{0,k-1}\|_{L^1(\Omega)}$, for every *n*. Since $d\varphi_n \wedge d^c\varphi_n \leq dd^c\varphi_{n,1}$, we have

$$\int_{U} |\varphi_{n}|^{k-p-2} d\varphi_{n} \wedge d^{c} \varphi_{n} \wedge (dd^{c} \varphi_{n})^{p} \wedge \omega^{k-p-1}$$

$$\leq \int_{U} |\varphi_{n}|^{k-p-2} dd^{c} \varphi_{n,1} \wedge (dd^{c} \varphi_{n})^{p} \wedge \omega^{k-p-1}.$$

Rescale φ if necessary, we can assume that, for every n, $\|\varphi_n\|_{W^*_{k-1}(U)} \leq 1$, $\|\varphi_{n,l}\|_{W^*_{k-1-l}(U)} \leq 1$ 1 for $1 \leq l \leq k-2$ and $\|\varphi_{n,k-1}\|_{L^1(U)} \leq 1$. It follows from Proposition 4.7 that the right-hand side is uniformly bounded by a constant depending only on U. Therefore, $|\varphi_n|^{k-p-2}d\varphi_n \wedge d^c\varphi_n \wedge (dd^c\varphi_n)^p \wedge \omega^{k-p-1}$ are locally weakly bounded in U for every $p = 0, \ldots, k-2$. The proof is thus complete. \Box

Acknowledgments

The first named author is supported by the MOE (Singapore) grant MOE-T2EP20120-0010. The second named author is supported by the Singapore International Graduate Award (SINGA). We would like to thank Tien-Cuong Dinh, Quôc Anh Ngô and Hoang-Son Tran for their help during the preparation of this paper.

REFERENCES

- [Ale81] H. Alexander, Projective capacity, Recent developments in several complex variables (Proc. Conf. Princeton Univ. Princeton, N. J., 1979), 3–27, Ann. of Math. Stud. 100, Princeton Univ. Press, Princeton, N.J. 1981.
- [AT84] H. Alexander, B. Taylor, Comparison of two capacities in \mathbb{C}^n , Math. Z. 186 (1984), 407–417.
- [BBGZ13] R.J. Berman, S. Boucksom, V. Guedj, A. Zeriahi, A variational approach to complex Monge-Ampère equations, Publ. Math. Inst. Hautes Études Sci. 117 (2013), 179–245.
- [BC00] B. Berndtsson, P. Charpentier, A Sobolev mapping property of the Bergman kernel, Math. Z. 235 (2000), no. 1, 1–10.
- [BD22] F. Bianchi, T.-C. Dinh, Equilibrium states of endomorphisms of \mathbb{P}^k : spectral stability and limit theorems, arXiv:2204.02856, to appear in Geom. Funct. Anal.
- [Blo04] Z. Błocki, On the definition of the Monge-Ampère operator in \mathbb{C}^2 , Math. Ann. **328** (2004), no. 3, 415–423.
- [Blo05] Z. Błocki, *Weak solutions to the complex Hessian equation*, Ann. Inst. Fourier (Grenoble) **55** (2005), 1735–1756.
- [Blo06] Z. Błocki, The domain of definition of the complex Monge–Ampère operator, Amer. J. Math. 128 (2006), 519–530.
- [Blo13] Z. Błocki, *Suita conjecture and the Ohsawa-Takegoshi extension theorem*, Invent. Math. **193** (2013), no. 1, 149–158.
- [BK07] Z. Błocki, S. Kołodziej, On regularization of plurisubharmonic functions on manifolds, Proc. Amer. Math. Soc. 135 (2007), no. 7, 2089–2093.
- [BT82] E. Bedford, B. A. Taylor, A new capacity for plurisubharmonic functions, Acta Math. 149 (1982), 1–41.
- [BT87] E. Bedford, B. A. Taylor, Fine topology, Silov boundary, and $(dd^c)^n$, J. Funct. Anal. 72 (1987), 225–251.
- [BW24] S. Biard, J. Wu, Equivalence between VMO functions and Zero Lelong numbers functions, arXiv:2403.03568, 2024.
- [Ceg98] U. Cegrell, *Pluricomplex energy*, Acta Math. 180 (1998), 187–217.
- [Ceg04] U. Cegrell, The general definition of the complex Monge–Ampère operator, Ann. Inst. Fourier (Grenoble) 54 (2004), 159–179.
- [Cho55] G. Choquet, Theory of capacities, Ann. Inst. Fourier. 5 (1955), 131-295.
- [Dar15] T. Darvas, The Mabuchi geometry of finite energy classes, Adv. Math. 285 (2015), 182–219.

- [Dem] J.-P. Demailly, *Complex analytic and differential geometry*, available at https://www-fourier.ujf-grenoble.fr/ demailly/manuscripts/agbook.pdf.
- [DV24] H. De Thélin, G. Vigny, Exponential mixing of all orders and CLT for generic birational maps of \mathbb{P}^k , arXiv:2402.01178, 2024.
- [DGL20] E. Di Nezza, V. Guedj, C. Lu, *Finite entropy vs finite energy*, Comment. Math. Helv. **96** (2021), 389–419.
- [DKW21] T.-C. Dinh, L. Kaufmann, H. Wu, *Products of random matrices: a dynamical point of view*, Pure Appl. Math. Q. **17** (2021), 933–969.
- [DKN20] T.-C. Dinh, S. Kołodziej, and N.C. Nguyen, *The Complex Sobolev Space and Hölder continuous solutions to Monge-Ampère equations*, Bull. Lond. Math. Soc. **54** (2022), 772–790.
- [DMV20] T.-C. Dinh, G. Marinescu, and D.-V. Vu, *Moser-Trudinger inequalities and complex Monge-Ampère equation*, Ann. Sc. Norm. Super. Pisa Cl. Sci. **24** (2023), 927–954.
- [DNS10] T.-C. Dinh, V.-A. Nguyên, N. Sibony, *Exponential estimates for plurisubharmonic functions and stochastic dynamics*, J. Differential Geom. **84** (2010), 465–488.
- [DNS22] T.-C. Dinh, V.-A. Nguyên, N. Sibony, *Unique ergodicity for foliations on compact Kähler surfaces*, Duke Math. J. **171** (2022), 2627–2698.
- [DS04] T.-C. Dinh, N. Sibony, *Regularization of currents and entropy*, Ann. Ecole Norm. Sup. **37** (2004), 959–971.
- [DS06a] T.-C. Dinh, N. Sibony, *Decay of correlations and the central limit theorem for meromorphic maps*, Comm. Pure Appl. Math. **59** (2006), 754–768.
- [DS06b] T.-C. Dinh, N. Sibony, Distribution des valeurs d'une suite de transformations méromorphes et applications, Comment. Math. Helv. **81** (2006), 221–258.
- [DV22] D.T. Do, D.-V. Vu, Complex Monge-Ampère equations with solutions in finite energy classes, Math. Res. Lett. 29 (2022), 1659–1683.
- [DF83] H. Donnelly, C. Fefferman, L²-cohomology and index theorem for the Bergman metric, Ann. of Math.
 118 (1983), 593–618.
- [Eva10] L.C. Evans, *Partial differential equations*, vol. 19 of Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, second ed., 2010.
- [FZ73] H. Federer, W. Ziemer, *The Lebesgue set of a function whose distribution derivatives are p-th power summable*, Indiana Univ. Math. J. **22** (1972/73), 139–158.
- [FS95] J.E. Fornæss, N. Sibony, Oka's inequality for currents and applications, Math. Ann. 301 (1995), 399–419.
- [GZ05] V. Guedj, A. Zeriahi, Intrinsic capacities on compact Kähler manifolds, J. Geom. Anal. 15 (2005), 607–639.
- [GZ07] V. Guedj, A. Zeriahi, *The weighted Monge-Ampère energy of quasiplurisubharmonic functions*, J. Funct. Anal. **250** (2007), 442–482.
- [GZ17] V. Guedj, A. Zeriahi, *Degenerate complex Monge-Ampère equations*, EMS Tracts Math, 26 European Mathematical Society (EMS), Zürich, 2017, xxiv+472 pp.
- [HL06] F.-R. Harvey, H.-B. Lawson, *Projective hulls and the projective Gelfand transform*, Asian J. Math. **10** (2006), 607–646.
- [JN61] F. John, L. Nirenberg, On functions of bounded mean oscillation, Comm. Pure. Appl. Math. 14 (1961), 415–426.
- [Jos78] B. Josefson, On the equivalence between locally polar and globally polar sets for plurisubharmonic functions on \mathbb{C}^n , Ark. Mat. 16 (1978), 109–115.
- [Kau17] L. Kaufmann, A Skoda-type integrability theorem for singular Monge-Ampère measures, Michigan Math. J. 66 (2017), 581–594.
- [Kol98] S. Kołodziej, The complex Monge-Ampère equation, Acta Math. 180 (1998), 69–117.
- [Kol03] S. Kołodziej, *The Monge-Ampère equation on compact Kähler manifolds*, Indiana Univ. Math. J. **52** (2003), 667–686.
- [Kol05] S. Kołodziej, *The complex Monge-Ampère equation and pluripotential theory*, Mem. Amer. Math. Soc. **178** (2005), no.840, x+64 pp.
- [KPR84] N.J. Kalton, N.T. Peck, James W. Roberts, An F-space sampler, London Math. Soc. Lecture Note Ser. 89, Cambridge University Press, Cambridge, 1984, xii+240 pp.

- [Mos70] J. Moser, A sharp form of an inequality by N. Trudinger, Indiana Univ. Math. J. 20 (1970/71), 1077–1092.
- [Sic62] J. Siciak, On some extremal functions and their applications in the theory of analytic functions of several complex variables, Trans. Amer. Math. Soc. **105** (1962), 322–357.
- [Siu74] Y.T. Siu, Analyticity of sets associated to Lelong numbers and the extension of positive closed currents, Invent. Math. **27** (1974), 53–156.
- [Tru67] N.S. Trudinger, On imbeddings into Orlicz spaces and some applications, J. Math. Mech. 17 (1967), 473–483.
- [Vig07] G. Vigny, Dirichlet-like space and capacity in complex analysis in several variables, J. Funct. Anal. 252 (2007), 247–277.
- [Vig15] G. Vigny, Exponential decay of correlations for generic regular birational maps of \mathbb{P}^k , Math. Ann. **362** (2015), 1033–1054.
- [VV23] G. Vigny, D.-V. Vu, *Lebesgue points of functions in the complex Sobolev space*, arXiv:2306.12332, to appear in Internat. J. Math.
- [Vu20] D.-V. Vu, Equilibrium measures of meromorphic self-maps on non-Kähler manifolds, Trans. Amer. Math. Soc. **373** (2020), 2229–2250.
- [Vu23] D.-V. Vu, Continuity of functions in complex Sobolev spaces, arXiv:2312.01635, 2023.
- [Wal68] J.B. Walsh, *Continuity of envelopes of plurisubharmonic functions*, J. Math. Mech. **18** (1968) 143–148.
- [Yau78] S.T. Yau, On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I, Comm. Pure Appl. Math. **31** (1978) 339–411.

¹Department of Mathematics, National University of Singapore - 10, Lower Kent Ridge Road - Singapore 119076³

Email address: drdo94@nus.edu.sg, duongdothai.vn@gmail.com

²Department of Mathematics, National University of Singapore - 10, Lower Kent Ridge Road - Singapore 119076

Email address: e1101880@u.nus.edu, meliodaspbc@gmail.com

³On leave from Institute of Mathematics, Vietnam Academy of Science and Technology