

The Morse property of limit functions appearing in mean field equations on surfaces with boundary

Zhengni Hu* Thomas Bartsch†

May 13, 2024

Abstract: In this paper, we study the Morse property for functions related to limit functions of mean field equations on a smooth, compact surface Σ with boundary $\partial\Sigma$. Given a Riemannian metric g on Σ we consider functions of the form

$$f_g(x) := \sum_{i=1}^m \sigma_i^2 R^g(x_i) + \sum_{\substack{i,j=1 \\ i \neq j}}^m \sigma_i \sigma_j G^g(x_i, x_j) + h(x_1, \dots, x_m),$$

where $\sigma_i \neq 0$ for $i = 1, \dots, m$, G^g is the Green function of the Laplace-Beltrami operator on (Σ, g) with Neumann boundary conditions, R^g is the corresponding Robin function, and $h \in C^2(\Sigma^m, \mathbb{R})$ is arbitrary. We prove that for any Riemannian metric g , there exists a metric \tilde{g} which is arbitrarily close to g and in the conformal class of g such that $f_{\tilde{g}}$ is a Morse function. Furthermore we show that, if all $\sigma_i > 0$, then the set of Riemannian metrics for which f_g is a Morse function is open and dense in the set of all Riemannian metrics.

Key Words: Green function, Robin function, Morse property, Transversality theorem

AMS subject classification: 35J08, 35J25, 53C21, 58J60

1 Introduction and main results

Let Σ be a smooth, compact surface with boundary $\partial\Sigma$. For a Riemannian metric g on Σ let $G^g : \Sigma \times \Sigma \rightarrow \mathbb{R}$ be the Green function for the Laplace-Beltrami operator $-\Delta_g$ with

*Supported by CSC No. 202106010046.

†Supported by DFG grant BA 1009/19-1.

Neumann boundary conditions and mean value 0; i.e. for each $\xi \in \Sigma$ the function $G^g(\cdot, \xi)$ is the unique solution of

$$\begin{cases} -\Delta_g G^g(\cdot, \xi) = \delta_\xi - \frac{1}{|\Sigma|_g} & \text{in } \overset{\circ}{\Sigma} := \Sigma \setminus \partial\Sigma, \\ \partial_{\nu_g} G^g(\cdot, \xi) = 0 & \text{on } \partial\Sigma, \\ \int_{\Sigma} G^g(\cdot, \xi) dv_g = 0. \end{cases}$$

Here ν_g is the unit outward normal on $\partial\Sigma$, dv_g is the area element of Σ , $|\Sigma|_g = \int_{\Sigma} dv_g$ and δ_ξ is the Dirac distribution on Σ concentrated at $\xi \in \Sigma$. Setting $\kappa(\xi) = 2\pi$ for $\xi \in \overset{\circ}{\Sigma}$ and $\kappa(\xi) = \pi$ for $\xi \in \partial\Sigma$, the Robin function $R^g : \Sigma \rightarrow \mathbb{R}$ is defined by

$$R^g(\xi) := \lim_{x \rightarrow \xi} \left(G^g(x, \xi) + \frac{1}{\kappa(\xi)} \log d_g(x, \xi) \right),$$

where d_g denotes the distance with respect to the metric g . Given integers $m \geq l \geq 0$ with $m \geq 1$ and real numbers $\sigma_i \neq 0$, $i = 1, \dots, m$, we set

$$X := \overset{\circ}{\Sigma}^l \times (\partial\Sigma)^{m-l} \quad \text{and} \quad \Delta_X := \{x \in X : x_i = x_j \text{ for some } i \neq j\},$$

and define, for an arbitrary given function $h \in \mathcal{C}^2(\Sigma^m, \mathbb{R})$:

$$(1.1) \quad f_g : X \setminus \Delta_X \rightarrow \mathbb{R}, \quad f_g(x) = \sum_{i=1}^m \sigma_i^2 R^g(x_i) + \sum_{\substack{i,j=1 \\ i \neq j}}^m \sigma_i \sigma_j G^g(x_i, x_j) + h(x_1, \dots, x_m).$$

We will prove that f_g is a Morse function for a “generic” metric g on Σ , in a sense that will be made precise below.

Functions of the form (1.1) play a crucial role in understanding the blow-up behavior of solutions of mean field equations like

$$(1.2) \quad \begin{cases} -\Delta_g u = \lambda \left(\frac{Ve^u}{\int_{\Sigma} Ve^u dv_g} - \frac{1}{|\Sigma|_g} \right) & \text{in } \overset{\circ}{\Sigma}, \\ \partial_{\nu_g} u = 0 & \text{on } \partial\Sigma, \\ \int_{\Sigma} u dv_g = 0. \end{cases}$$

For a compact surface without boundary it has been proven in [7] that a nondegenerate or more generally, an isolated stable critical point $x = (x_1, \dots, x_m)$ of f_g , with $l = m$ and $h(x) = 2 \sum_{i=1}^m \sigma_i \log V(x_i)$, gives rise to solutions (λ, u) of (1.2) with λ close to $8\pi m$ and such that u blows up as $\lambda \rightarrow 8\pi m$ precisely at $x_1, \dots, x_m \in \Sigma$. Similar results have been obtained in the case of mean field equations on a bounded domain in \mathbb{R}^2 with Dirichlet boundary conditions

in [8, 13], and for solutions of Gelfand's problem and steady states of the Keller-Segel system (see [2, 5, 6, 10] for instance).

Now we state our main results. We fix an integer $k \geq 0$ and $0 < \alpha < 1$, and let $\text{Riem}^{k+2,\alpha}(\Sigma)$ be the space of Riemannian metrics of class $\mathcal{C}^{k+2,\alpha}$ on Σ , i.e. the space of symmetric and positive definite sections $\Sigma \rightarrow (T\Sigma \otimes T\Sigma)^*$ of class $\mathcal{C}^{k+2,\alpha}$.

Theorem 1.1. *If $\sigma_i \neq 0$ for $i = 1, \dots, m$ and $\sigma_1 + \dots + \sigma_m \neq 0$ then for any $g \in \text{Riem}^{k+2,\alpha}(\Sigma)$ the set*

$$\mathcal{M}_g^{k+2,\alpha}(\Sigma) := \{\psi \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+) : f_{\psi g} \text{ is a Morse function}\}$$

is a residual subset of $\mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$.

As a consequence of Theorem 1.1 for any Riemannian metric g on Σ , there exists a metric \tilde{g} which is arbitrarily close to g and conformal to g , and such that $f_{\tilde{g}}$ is a Morse function.

Theorem 1.2. *If $\sigma_i > 0$ for all $i = 1, \dots, m$ then the set*

$$\text{Riem}_{\text{Morse}}^{k+2,\alpha}(\Sigma) := \{g \in \text{Riem}^{k+2,\alpha}(\Sigma) : f_g \text{ is a Morse function}\}$$

is an open and dense subset of $\text{Riem}^{k+2,\alpha}(\Sigma)$.

Remark 1.3. *In [1] the authors considered functions f_g of the form (1.1) on a surface Σ without boundary and with $h(x_1, \dots, x_m) = \sum_{i=1}^m \log V(x_i)$ where $V \in \mathcal{C}^2(\Sigma, \mathbb{R}_+)$. This is motivated by the mean field equation (1.2). They proved that f_g is a Morse function for V in an open and dense subset of $\mathcal{C}^2(\Sigma, \mathbb{R}_+)$. Also related is the paper [3], which deals with functions like f_g where G is the Green function of the Dirichlet Laplace operator on a bounded smooth domain $\Omega \subset \mathbb{R}^n$. In [3], it is proved that f_g is a Morse function for a generic domain Ω . The present paper seems to be the first investigating the Morse property of f_g as in (1.1) as a function of the Riemannian metric.*

2 Preliminaries

Riemann surfaces are locally conformally flat, so there exist isothermal coordinates where the metric is conformal to the Euclidean metric (see [4, 11, 15]). We modify the isothermal coordinates applied in [7, 9, 16] to a Riemann surface with boundary. For $\xi \in \Sigma$ there exists a local chart $y_\xi : \Sigma \supset U(\xi) \rightarrow B^\xi \subset \mathbb{R}^2$ transforming g to $e^{\varphi_\xi \circ y_\xi} \langle \cdot, \cdot \rangle_{\mathbb{R}^2}$. We may assume that $y_\xi(\xi) = 0$. For $\xi \in \overset{\circ}{\Sigma}$ we may also assume that $\overline{U(\xi)} \subset \overset{\circ}{\Sigma}$ and that the image of y_ξ is a disc

$B^\xi := \{y \in \mathbb{R}^2 : |y| < 2r_\xi\}$ of radius $2r_\xi > 0$. For $\xi \in \partial\Sigma$, by Lemma 4 in [16], there exists an isothermal coordinate system $(U(\xi), y_\xi)$ near ξ such that the image of y_ξ is a half disk $B_{2r_\xi}^+ := \{y \in \mathbb{R}^2 : |y| < 2r_\xi, y_2 \geq 0\}$ of radius $2r_\xi > 0$ and $y_\xi(U(\xi) \cap \partial\Sigma) = B_{2r_\xi}^+ \cap \partial\mathbb{R}_+^2$. For any $x \in y_\xi^{-1}(B_{2r_\xi}^+ \cap \partial\mathbb{R}_+^2)$, we have

$$(2.1) \quad (y_\xi)_*(\nu_g(x)) = -\exp\left(-\frac{\varphi_\xi(y)}{2}\right) \frac{\partial}{\partial y_2} \Big|_{y=y_\xi(x)}.$$

In this case, we take $B^\xi = B_{2r_\xi}^+$. For $\xi \in \Sigma$ and $0 < r \leq 2r_\xi$ we set

$$B_r^\xi := B^\xi \cap \{y \in \mathbb{R}^2 : |y| < r\} \quad \text{and} \quad U_r(\xi) := y_\xi^{-1}(B_r^\xi).$$

Recall that $\varphi_\xi : B^\xi \rightarrow \mathbb{R}$ is related to K , the Gaussian curvature of Σ , by the equation

$$-\Delta\varphi_\xi(y) = 2K(y_\xi^{-1}(y))e^{\varphi_\xi(y)} \quad \text{for all } y \in B^\xi.$$

We can assume that y_ξ and φ_ξ depend smoothly on ξ , and that $\varphi_\xi(0) = 0$ and $\nabla\varphi_\xi(0) = 0$.

Observe that we have for $\zeta \in U(\xi)$,

$$\lim_{x \rightarrow \zeta} \frac{d_g(x, \zeta)}{|\varphi_\xi(x) - \varphi_\xi(\zeta)|} = e^{\frac{1}{2}\varphi_\xi(\zeta)},$$

which implies

$$(2.2) \quad R^g(\zeta) = \lim_{x \rightarrow \zeta} \left(G^g(x, \zeta) + \frac{1}{\kappa(\zeta)} \log |y_\xi(x) - y_\xi(\zeta)| \right) + \frac{1}{2\kappa(\zeta)} \varphi_\xi(y_\xi(\zeta)),$$

and in particular, using $\varphi_\xi(y_\xi(\xi)) = \varphi_\xi(0) = 0$:

$$R^g(\xi) = \lim_{x \rightarrow \xi} \left(G^g(x, \xi) + \frac{1}{\kappa(\xi)} \log |y_\xi(x)| \right).$$

Now we construct a regular part $H^g(x, \xi)$ of the Green function $G^g(x, \xi)$ such that $H^g(\xi, \xi) = R^g(\xi)$. Let $\chi \in \mathcal{C}^\infty(\mathbb{R}, [0, 1])$ be such that

$$\chi(s) = \begin{cases} 1 & \text{if } |s| \leq 1, \\ 0 & \text{if } |s| \geq 2. \end{cases}$$

For $\xi \in \overset{\circ}{\Sigma}$ we choose $\delta_\xi = \min\{\frac{1}{2}r_\xi, \frac{1}{2}\text{dist}(x, \partial\Sigma)\}$, and for $\xi \in \partial\Sigma$ we set $\delta_\xi := \frac{1}{2}r_\xi$. Next we define the cut-off function $\chi_\xi \in \mathcal{C}^\infty(\Sigma, [0, 1])$ by

$$\chi_\xi(x) := \begin{cases} \chi(|y_\xi(x)|/\delta_\xi) & \text{if } x \in U(\xi) \\ 0 & \text{if } x \in \Sigma \setminus U(\xi) \end{cases}.$$

Then, for $\xi \in \Sigma$ the function $H_\xi^g := H^g(\cdot, \xi) : \Sigma \rightarrow \mathbb{R}$ is defined to be the unique solution of the Neumann problem

$$(2.3) \quad \begin{cases} \Delta_g H_\xi^g = \frac{1}{\kappa(\xi)} (\Delta_g \chi_\xi) \log |y_\xi| + \frac{2}{\kappa(\xi)} \langle \nabla^g \chi_\xi, \nabla^g \log |y_\xi| \rangle_g + \frac{1}{|\Sigma|_g} & \text{in } \overset{\circ}{\Sigma}, \\ \partial_{\nu_g} H_\xi^g = \frac{1}{\kappa(\xi)} (\partial_{\nu_g} \chi_\xi) \log |y_\xi| + \frac{1}{\kappa(\xi)} \chi_\xi \partial_{\nu_g} \log |y_\xi| & \text{on } \partial\Sigma, \\ \int_{\Sigma} H_\xi^g dv_g = \frac{1}{\kappa(\xi)} \int_{\Sigma} \chi_\xi \log |y_\xi| dv_g. \end{cases}$$

Lemma 2.1. *For $g \in \text{Riem}^{k+2,\alpha}(\Sigma)$ the function H^g is of class $\mathcal{C}^{k+3,\alpha}$ in any compact subsets of $\Sigma \times \overset{\circ}{\Sigma}$ and in $\Sigma \times \partial\Sigma$. Moreover, it satisfies*

$$G^g(x, \xi) = -\frac{1}{\kappa(\xi)} \chi_\xi(x) \cdot \log |y_\xi(x)| + H^g(x, \xi)$$

and $H^g(\xi, \xi) = R^g(\xi)$. Consequently R^g is of class $\mathcal{C}^{k+3,\alpha}$ in any compact subsets of $\overset{\circ}{\Sigma}$ and in $\partial\Sigma$.

Proof. First we observe that $e^{\varphi_\xi} \in \mathcal{C}^{k+2,\alpha}(B^\xi, \mathbb{R})$ because $g \in \text{Riem}^{k+2,\alpha}(\Sigma)$. For $\xi \in \overset{\circ}{\Sigma}$, by the choice of δ_ξ we have that $-\Delta_g H_\xi^g$ is of class $\mathcal{C}^{k+2,\alpha}$ class in Σ and $\partial_{\nu_g} H^g(x, \xi) \equiv 0$ on $\partial\Sigma$. Now the Schauder estimate for the Neumann problem (see [14, 16], for instance) implies that the solution of (2.3) uniquely exists in $\mathcal{C}^{k+4,\alpha}(\Sigma, \mathbb{R})$.

For $x \in U(\xi) \cap \partial\Sigma$, setting $y = y_\xi(x)$ we have:

$$\partial_{\nu_g} \log |y_\xi(x)| \stackrel{(2.1)}{=} -e^{-\frac{1}{2}\varphi_\xi(y)} \frac{\partial}{\partial y_2} \log |y| = -e^{-\frac{1}{2}\varphi_\xi(y)} \frac{y_2}{|y|^2} \equiv 0.$$

Clearly, $\partial_{\nu_g} \chi(|y_\xi|(x)) = 0$ for $x \in \partial\Sigma \cap U_{\delta_\xi}(\xi)$. It follows that $\partial_{\nu_g} H_\xi^g$ is of class $\mathcal{C}^{k+2,\alpha}$ on $\partial\Sigma$. Moreover $\Delta_g H_\xi^g$ is of class $\mathcal{C}^{k+2,\alpha}$ in Σ . Consequently (2.3) has a unique solution $H_\xi^g \in \mathcal{C}^{k+3,\alpha}(\Sigma, \mathbb{R})$ by the Schauder estimates.

Finally H_ξ^g is uniformly bounded in $\mathcal{C}^{k+3,\alpha}$ for ξ in any compact subsets of $\overset{\circ}{\Sigma}$ and in $\partial\Sigma$. Therefore $H^g(\xi, \xi)$ is in $\mathcal{C}^{k+3,\alpha}$ in any compact subsets of $\overset{\circ}{\Sigma}$ and in $\partial\Sigma$, and $H^g(\xi, \xi) = R^g(\xi)$ by (2.2). \square

For $g \in \text{Riem}^{k+2,\alpha}(\Sigma)$ we now consider the map

$$\mathcal{H}^g : \Sigma \times \Sigma \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+) \rightarrow \mathbb{R}, \quad \mathcal{H}^g(x, \xi, \psi) := H^{\psi g}(x, \xi).$$

Proposition 2.2. *The map \mathcal{H}^g is \mathcal{C}^1 in $\Sigma \times \mathring{\Sigma} \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$ and in $\Sigma \times \partial\Sigma \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$. Moreover, we have*

$$(2.4) \quad D_\psi \mathcal{H}^g(x, \xi, 1)[\theta] = -\frac{1}{|\Sigma|_g} \int_{\Sigma} (G^g(z, x) + G^g(z, \xi)) \theta(z) dv_g(z),$$

for any $\theta \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R})$.

Proof. For $g \in \text{Riem}^{k+2,\alpha}(\Sigma)$ and $\psi \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$ we clearly have $\psi g \in \text{Riem}^{k+2,\alpha}(\Sigma)$. By a direct calculation we obtain the following equations for $H_\xi^{\psi g} - H_\xi^g$:

$$\begin{cases} -\Delta_g(H_\xi^{\psi g} - H_\xi^g) = \frac{1}{|\Sigma|_g} - \frac{\psi}{|\Sigma|_{\psi g}} & \text{in } \mathring{\Sigma}, \\ \partial_{\nu_g}(H_\xi^{\psi g} - H_\xi^g) = 0 & \text{on } \partial\Sigma, \\ \int_{\Sigma} (H_\xi^{\psi g} - H_\xi^g) dv_g = - \int_{\Sigma} G_\xi^{\psi g}(\psi - 1) dv_g. \end{cases}$$

An expansion of $|\Sigma|_{\psi g} = \int_{\Sigma} dv_{\psi g}$ yields:

$$-\Delta_g(H_\xi^{\psi g} - H_\xi^g) = \frac{1 - \psi}{|\Sigma|_g} + \frac{\int_{\Sigma} (\psi - 1) dv_g}{|\Sigma|_g^2} + \mathcal{O}(\|\psi - 1\|_{\mathcal{C}^{k+2,\alpha}}^2) \quad \text{as } \|\psi - 1\|_{\mathcal{C}^{k+2,\alpha}} \rightarrow 0.$$

Recall that $G_\xi^{\psi g} = -\frac{4}{\sigma(\xi)}\chi(|y_\xi|) \log |y_\xi| + H_\xi^{\psi g}$, so the representation formula gives:

$$(2.5) \quad H_\xi^{\psi g}(x) - H_\xi^g(x) = \frac{1}{|\Sigma|_g} \int_{\Sigma} G^{\psi g}(z, \xi)(\psi(z) - 1) dv_g(z) - \frac{1}{|\Sigma|_{\psi g}} \int_{\Sigma} G^g(z, x)(\psi(z) - 1) dv_g(z).$$

By standard elliptic estimates (see [14, 16]) there exists a constant C such that

$$\|H_\xi^{\psi g} - H_\xi^g\|_{\mathcal{C}^{k+4,\alpha}} \leq C \cdot \|\psi - 1\|_{\mathcal{C}^{k+2,\alpha}},$$

thus

$$(2.6) \quad H_\xi^{\psi g} \rightarrow H_\xi^g \quad \text{in } \mathcal{C}^{k+4,\alpha} \text{ as } \psi \rightarrow 1 \text{ in } \mathcal{C}^{k+2,\alpha}.$$

According to the construction of $H^g(x, \xi)$, the convergence in (2.6) is uniform for ξ in any compact subset of $\mathring{\Sigma}$, and in $\partial\Sigma$. It follows that $\mathcal{H}_g(x, \xi, \cdot)$ is continuous at 1, uniformly for $x \in \Sigma$ and ξ in any compact subsets of $\mathring{\Sigma}$ or $\xi \in \partial\Sigma$. Using $\mathcal{H}_g(x, \xi, \psi) = \mathcal{H}_{\psi g}(x, \xi, 1)$ we see that $\mathcal{H}_g(x, \xi, \cdot)$ is continuous at every $\psi \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$.

Next we prove that $\mathcal{H}^g(x, \xi, \psi)$ is \mathcal{C}^1 with respect to ψ . We fix $\theta \in \mathcal{C}^{2+k,\alpha}(\Sigma, \mathbb{R})$ and consider the metric $(1 + t\theta)g$ with t sufficiently small so that $1 + t\theta > 0$. Then

$$(2.7) \quad w_\xi^t(x) := \frac{1}{t} (H^{(1+t\theta)g}(x, \xi) - H^g(x, \xi)) = \frac{1}{t} (\mathcal{H}^g(x, \xi, 1 + t\theta) - \mathcal{H}^g(x, \xi, 1))$$

satisfies the following equations as $t \rightarrow 0$:

$$(2.8) \quad \begin{cases} -\Delta_g w_\xi^t = \frac{1}{|\Sigma|_g^2} \int_\Sigma \theta \, dv_g - \frac{\theta}{|\Sigma|_g} + \mathcal{O}(t) & \text{in } \overset{\circ}{\Sigma}, \\ \partial_{\nu_g} w_t(x, \xi) = 0 & x \in \partial\Sigma, \\ \int_\Sigma w_\xi^t \, dv_g = - \int_\Sigma G_\xi^{(1+t\theta)g} \cdot \theta \, dv_g \end{cases}$$

where $\mathcal{O}(t)$ is defined with respect to the $\mathcal{C}^{k+2,\alpha}$ -norm. Applying the standard elliptic estimates, w_ξ^t converges as $t \rightarrow 0$ to some function $w_\xi^0(x) = D_\psi \mathcal{H}^g(x, \xi, 1)[\theta]$ in $\mathcal{C}^{k+4,\alpha}(\Sigma, \mathbb{R})$. Moreover, w_ξ^0 satisfies the equations

$$(2.9) \quad \begin{cases} -\Delta_g w_\xi^0 = \frac{1}{|\Sigma|_g^2} \int_\Sigma \theta \, dv_g - \frac{\theta}{|\Sigma|_g} & \text{in } \overset{\circ}{\Sigma}, \\ \partial_{\nu_g} w_\xi^0 = 0 & \text{on } \partial\Sigma, \\ \int_\Sigma w_\xi^0 \, dv_g = - \int_\Sigma G_\xi^g \cdot \theta \, dv_g. \end{cases}$$

The representation formula now implies (2.4):

$$D_\psi \mathcal{H}^g(x, \xi, 1)[\theta] = w_\xi^0(x) = -\frac{1}{|\Sigma|_g} \int_\Sigma (G^g(z, x) + G^g(z, \xi)) \cdot \theta(z) \, dv_g(z).$$

Replacing g by ψg and θ by $\frac{\theta}{\psi}$, we obtain the derivative of $\mathcal{H}_g(x, \xi, \psi)$ for arbitrary $\psi \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$:

$$(2.10) \quad \begin{aligned} D_\psi \mathcal{H}^g(x, \xi, \psi)[\theta] &= \lim_{t \rightarrow 0} \frac{1}{t} (H^{(\psi+t\theta)g}(x, \xi) - H^{\psi g}(x, \xi)) \\ &= -\frac{1}{|\Sigma|_{\psi g}} \int_\Sigma (G^{\psi g}(z, x) + G^{\psi g}(z, \xi)) \cdot \frac{\theta(z)}{\psi(z)} \, dv_{\psi g}(z) \\ &= -\frac{1}{|\Sigma|_{\psi g}} \int_\Sigma (G^{\psi g}(z, x) + G^{\psi g}(z, \xi)) \cdot \theta(z) \, dv_g(z). \end{aligned}$$

In order to see that $\mathcal{H}_g(x, \xi, \psi)$ is continuously Fréchet differentiable with respect to ψ it is sufficient to prove that $D_\psi \mathcal{H}_g(x, \xi, \psi)$ is continuous in ψ as a linear operator on $\mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$. For $\psi_1, \psi_2 \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$ there holds

$$\begin{aligned} |D_\psi \mathcal{H}_g(x, \xi, \psi_1)[\theta] - D_\psi \mathcal{H}_g(x, \xi, \psi_2)[\theta]| &= \left| \frac{1}{|\Sigma|_{\psi_1 g}} \int_\Sigma (G^{\psi_1 g}(z, x) + G^{\psi_1 g}(z, \xi)) \theta(z) \, dv_g(z) \right. \\ &\quad \left. - \frac{1}{|\Sigma|_{\psi_2 g}} \int_\Sigma (G^{\psi_2 g}(z, x) + G^{\psi_2 g}(z, \xi)) \theta(z) \, dv_g(z) \right| \\ &\leq C \cdot \|\theta\|_{\mathcal{C}^{k+2,\alpha}} \cdot \|\psi_1 - \psi_2\|_{\mathcal{C}^{k+2,\alpha}} \end{aligned}$$

where we applied (2.10); here $C > 0$ is a constant. Therefore \mathcal{H}^g is \mathcal{C}^1 in $\Sigma \times \overset{\circ}{\Sigma} \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$ and in $\Sigma \times \partial\Sigma \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$. \square

3 Proof of Theorem 1.1

The proof is based on the following transversality theorem [12, Theorem 5.4].

Theorem 3.1. *Let M, Ψ, N be Banach manifolds of class \mathcal{C}^r for some $r \in \mathbb{N}$, let $\mathcal{D} \subset M \times \Psi$ be open, let $\mathcal{F} : \mathcal{D} \rightarrow N$ be a \mathcal{C}^r map, and fix a point $z \in N$. Assume for each $(y, \psi) \in \mathcal{F}^{-1}(z)$ that:*

- (1) $D_y \mathcal{F}(y, \psi) : T_y M \rightarrow T_z N$ is semi-Fredholm with index $< r$;
- (2) $D\mathcal{F}(y, \psi) : T_y M \times T_\psi \Psi \rightarrow T_z N$ is surjective;
- (3) $\mathcal{F}^{-1}(z) \rightarrow \Psi$, $(y, \psi) \mapsto \psi$, is σ -proper.

Then

$$\Psi_{reg} := \{\psi \in \Psi : z \text{ is a regular value of } \mathcal{F}(\cdot, \psi)\}$$

is a residual subset of Ψ .

Proof of Theorem 1.1. We set $\Psi := \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$ and consider the functions $f_{\psi g} : X \setminus \Delta_X \rightarrow \mathbb{R}$ for $\psi \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$ first in a local isothermal chart. Let $y_{\xi_i} : \Sigma \supset U(\xi_i) \rightarrow B^{\xi_i} \subset \mathbb{R}^2$ be isothermal charts of Σ as in Section 2 with $\xi_1, \dots, \xi_l \in \overset{\circ}{\Sigma}$, $\xi_{l+1}, \dots, \xi_m \in \partial\Sigma$. For simplicity of notation we set $Y_i := y_{\xi_i} : U_i := U(\xi_i) \rightarrow B_i := B^{\xi_i} \subset \mathbb{R}^2$ for $i = 1, \dots, l$, and $Y_i := \pi_1 \circ y_{\xi_i} : U_i := U(\xi_i) \cap \partial\Sigma \rightarrow B_i \subset \mathbb{R}$ for $i = l+1, \dots, m$; here $\pi_1 : \mathbb{R}^2 \rightarrow \mathbb{R}$ is the projection onto the first component. For $i = l+1, \dots, m$ we thus have $y_{\xi_i}(x) = (Y_i(x), 0)$. Then

$$Y = Y_1 \times \cdots \times Y_m : X \supset U := U_1 \times \cdots \times U_m \rightarrow \mathbb{R}^{l+m}, \quad (x_1, \dots, x_m) \mapsto (Y_1(x_1), \dots, Y_m(x_m)),$$

is a chart of X . Set $M = N := \mathbb{R}^{l+m}$ and $V := Y(U \cap X \setminus \Delta_X) \subset \mathbb{R}^{l+m} = M$ so that $\mathcal{D} := V \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$ is an open subset of $\mathbb{R}^{l+m} \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)$.

We will apply Theorem 3.1 to prove that $0 \in \mathbb{R}^{l+m}$ is a regular value of

$$\nabla(f_{\psi g} \circ Y^{-1}) : \mathbb{R}^{l+m} \supset V = Y(U \cap X \setminus \Delta_X) \rightarrow \mathbb{R}^{l+m}$$

for ψ in a residual subset $\Psi_U \subset \Psi$. This implies that the restriction of $f_{\psi g}$ to $U \cap X \setminus \Delta_X$ is a Morse function for $\psi \in \Psi_U$. Then Theorem 1.1 follows because X is covered by finitely many neighborhoods U as above and because the intersection of finitely many residual sets is a residual set.

It remains to prove that the map

$$\mathcal{F}_g : \mathbb{R}^{l+m} \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+) \supset \mathcal{D} \rightarrow \mathbb{R}^{l+m}, \quad (y, \psi) \mapsto \nabla(f_{\psi g} \circ Y^{-1})(y),$$

satisfies the assumptions of Theorem 3.1 with $r = 1$. Concerning the differentiability it is clear that \mathcal{F}_g is \mathcal{C}^1 as a function of $y \in V$. In order to see that \mathcal{F}_g is also \mathcal{C}^1 in ψ , by Lemma 2.1 it is sufficient to prove that $\nabla_y(H^{\psi g}(Y_i^{-1}(\cdot), Y_j^{-1}(\cdot)))$ is \mathcal{C}^1 in ψ . We recall that w_ξ^t is defined by (2.7). Applying the representation formula of w_ξ^t and Lebesgue's dominated convergence theorem, we have for $(y, \psi) \in \mathcal{D}$ and $\theta \in \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R})$:

$$\begin{aligned} D_\psi|_{\psi=1} \nabla_y H^{\psi g}(Y_i^{-1}(y_i), Y_j^{-1}(y_j))[\theta] &= \lim_{t \rightarrow 0} \nabla_y \left(w_\xi^t(x) \Big|_{x=Y_i^{-1}(y_i), \xi=Y_j^{-1}(y_j)} \right) \\ &\stackrel{(2.5)}{=} \lim_{t \rightarrow 0} \left(-\frac{1}{|\Sigma|_g} \int_\Sigma \nabla_y G_{Y_j^{-1}(y_j)}^{(1+t\theta)g} \theta dv_g + \int_\Sigma \frac{1}{t} \left(\frac{1}{|\Sigma|_g} - \frac{1+t\theta}{|\Sigma|_{(1+t\theta)g}} \right) \nabla_y G_{Y_i^{-1}(y_i)}^g dv_g \right) \\ &= -\frac{1}{|\Sigma|_g} \int_\Sigma \nabla_y (G^g(z, Y_i^{-1}(y_i)) + G^g(z, Y_j^{-1}(y_j))) \cdot \theta(z) dv_g(z) \\ &= \nabla_y D_\psi \mathcal{H}^g(Y_i^{-1}(y_i), Y_j^{-1}(y_j), 1)[\theta], \end{aligned}$$

where we used Proposition 2.2. Since $D_\psi \mathcal{F}_g(y, \psi)[\theta] = D_\psi \mathcal{F}_{\psi g}(y, 1) \left[\frac{\theta}{\psi} \right]$ we obtain

$$D_\psi \mathcal{F}_g(y, \psi)[\theta] = -\frac{1}{|\Sigma|_\psi} \int_\Sigma \nabla_y (G^{\psi g}(z, Y_i^{-1}(y_i)) + G^{\psi g}(z, Y_j^{-1}(y_j))) \cdot \theta(z) dv_g(z).$$

As in the proof of Proposition 2.2 we deduce that $\mathcal{F}_g(y, \psi)$ is \mathcal{C}^1 on U .

Now we need to check the assumptions (1)-(3) of Theorem 3.1. Obviously $D_y \mathcal{F}_g(y, \psi) : \mathbb{R}^{l+m} \rightarrow \mathbb{R}^{l+m}$ is a Fredholm operator of index $0 < 1$, hence (1) holds. Also, (3) is easy to prove: For $j \in \mathbb{N}$ the set

$$M_j := \{Y(x) \in \mathbb{R}^{l+m} : x \in U, d_g(x, \Delta_X \cup \partial U) \geq 2^{-j}\} \subset Y(U) \subset \mathbb{R}^{l+m}$$

is compact as a continuous image of a compact set. Therefore the map

$$M_j \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+) \rightarrow \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+), \quad (y, \psi) \mapsto \psi,$$

is proper, hence its restriction to $\mathcal{F}_g^{-1}(0) \cap (M_j \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+))$ is proper. Since $V = \bigcup_{j=1}^\infty M_j$, so $\mathcal{D} = \bigcup_{j=1}^\infty (M_j \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+))$ it follows that the map

$$\mathcal{F}_g^{-1}(0) = \bigcup_{j=1}^\infty \left(\mathcal{F}_g^{-1}(0) \cap (M_j \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+)) \right) \rightarrow \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}_+), \quad (y, \psi) \mapsto \psi,$$

is σ -proper.

Finally we prove the surjectivity of the derivative $D\mathcal{F}_g(y, \psi) : \mathbb{R}^{l+m} \times \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}) \rightarrow \mathbb{R}^{l+m}$ at a point $(y, \psi) \in \mathcal{F}_g^{-1}(0)$. In fact, we shall prove that $D_\psi \mathcal{F}_g(y, \psi) : \mathcal{C}^{k+2,\alpha}(\Sigma, \mathbb{R}) \rightarrow \mathbb{R}^{l+m}$ is onto. Since

$$D_\psi \mathcal{F}_g(y, \psi)[\theta] = D_\psi \mathcal{F}_{\psi g}(y, 1)[\theta/\psi] \quad \text{for } \theta \in \mathcal{C}^{2+k,\alpha}(\Sigma, \mathbb{R})$$

it is sufficient to consider the case $\psi \equiv 1$. We observe for $\theta \in \mathcal{C}^{2+k,\alpha}(\Sigma, \mathbb{R})$:

$$D_\psi \Big|_{\psi=1} \nabla_y G^{\psi g}(Y_i^{-1}(y_i), Y_j^{-1}(y_j))[\theta] = D_\psi \Big|_{\psi=1} \nabla_y H^{\psi g}(Y_i^{-1}(y_i), Y_j^{-1}(y_j))[\theta].$$

Now Proposition 2.2 yields for $\theta \in \mathcal{C}^{2+k,\alpha}(\Sigma, \mathbb{R})$ with $\text{supp}(\theta) \subset \Sigma \setminus \{Y_1^{-1}(y_1), \dots, Y_m^{-1}(y_m)\}$:

$$\begin{aligned} D_\psi \mathcal{F}_g(y, 1)[\theta] &= \frac{d}{dt} \Big|_{t=0} \left(\sum_{i=1}^m \sigma_i^2 \nabla_y R^{(1+t\theta)g}(Y_i^{-1}(y_i)) \right. \\ &\quad \left. + \sum_{\substack{i,j=1 \\ i \neq j}}^m \sigma_i \sigma_j \nabla_y G^{(1+t\theta)g}(Y_i^{-1}(y_i), Y_j^{-1}(y_j)) + \nabla_y (h \circ Y^{-1})(y) \right) \\ &= -\frac{1}{|\Sigma|_g} \int_\Sigma \left(\sum_{i=1}^m 2\sigma_i^2 \nabla_y G^g(z, Y_i^{-1}(y_i)) \right. \\ &\quad \left. + \sum_{\substack{i,j=1 \\ i \neq j}}^m \sigma_i \sigma_j (\nabla_y G^g(z, Y_i^{-1}(y_i)) + \nabla_y G^g(z, Y_j^{-1}(y_j))) \right) \theta(z) dv_g(z). \end{aligned}$$

Consider an element $u = (u_1, \dots, u_m) \in \mathbb{R}^{l+m}$ with $u_1, \dots, u_l \in \mathbb{R}^2$, $u_{l+1}, \dots, u_m \in \mathbb{R}$, that is orthogonal to the range of $D_\psi \mathcal{F}_g(y, 1)$, i.e. it satisfies for every $\theta \in \mathcal{C}^{2+k,\alpha}(\Sigma, \mathbb{R})$ with $\text{supp}(\theta) \subset \Sigma \setminus \{Y_1^{-1}(y_1), \dots, Y_m^{-1}(y_m)\}$:

$$\begin{aligned} 0 &= \langle u, D_\psi \mathcal{F}_g(y, 1)[\theta] \rangle = \sum_{i=1}^m \langle u_i, (D_\psi \mathcal{F}_g(y, 1)[\theta])_i \rangle \\ &= -\sum_{i=1}^m \frac{1}{|\Sigma|_g} \left(2\sigma_i^2 + 2 \sum_{\substack{j=1 \\ j \neq i}}^m \sigma_i \sigma_j \right) \int_\Sigma \langle u_i, \nabla_{y_i} G^g(z, Y_i^{-1}(y_i)) \rangle \cdot \theta(z) dv_g(z) \end{aligned}$$

This implies, using $\sum_{j=1}^m \sigma_j \neq 0$:

$$(3.1) \quad \sum_{i=1}^m \sigma_i \langle u_i, \nabla_{y_i} G^g(z, Y_i^{-1}(y_i)) \rangle = 0 \quad \text{for } z \in \Sigma \setminus \{Y_1^{-1}(y_1), \dots, Y_m^{-1}(y_m)\}.$$

Setting $\kappa_i = 2\pi$ for $i = 1, \dots, l$ and $\kappa_i = \pi$ for $i = l+1, \dots, m$ we have

$$G^g(z, Y_i^{-1}(y_i)) = H^g(z, Y_i^{-1}(y_i)) - \frac{1}{\kappa_i} \chi(4|Y_i(z) - y_i|/\delta_{\xi_i}) \log |Y_i(z) - y_i|.$$

Now we define $z_i(t) := Y_i^{-1}(y_i + tu_i)$ for $i \in \{1, \dots, m\}$ and observe that

$$\nabla_{y_j} G^g(z_i(t), Y_j^{-1}(y_j)) = \mathcal{O}(1) \quad \text{as } t \rightarrow 0 \text{ for } j \neq i$$

whereas

$$\langle u_i, \nabla_{y_i} G^g(z_i(t), Y_i^{-1}(y_i)) \rangle \rightarrow \infty \quad \text{as } t \rightarrow 0.$$

Equation (3.1) implies $u_i = 0$ because $\sigma_i \neq 0$. This holds for all i , hence $u = 0$ and $D_\psi \mathcal{F}_g(y, \psi)$ must be onto. \square

4 Proof of Theorem 1.2

Theorem 1.2 follows easily from the following lemma.

Lemma 4.1. *If $\sigma_i > 0$ for $i = 1, \dots, m$ then $\lim_{x \rightarrow \partial(X \setminus \Delta_X)} |\nabla^g f_g(x)|_g = \infty$.*

Proof of Theorem 1.2. Lemma 4.1 implies that the set of critical points of f_g is compact. It follows that if f_g is a Morse function, then it has only finitely many critical points, and any C^2 -perturbation of f_g is also a Morse function. Therefore $\text{Riem}_{\text{Morse}}^{k+2,\alpha}(\Sigma)$ is an open subset of $\text{Riem}^{k+2,\alpha}(\Sigma)$. By Theorem 1.1 it is also a dense subset. \square

Proof of Lemma 4.1. Consider a sequence $x^n \in X \setminus \Delta_X$ such that

$$x^n \rightarrow x^\infty \in \partial(X \setminus \Delta_X) = \Delta_X \cup ((\partial\Sigma)^l \times (\partial\Sigma)^{m-l}) \subset \Sigma^l \times (\partial\Sigma)^{m-l}.$$

CASE 1. $x^\infty \in \Delta_X \subset \overset{\circ}{\Sigma}^l \times (\partial\Sigma)^{m-l}$.

Then, there exists $\xi \in \Sigma$ such that the set $I := \{i \in \{1, \dots, m\} : x_i^\infty = \xi\}$ contains at least two elements. Moreover, if $\xi \in \overset{\circ}{\Sigma}$, then $I \subset \{1, \dots, l\}$; if $\xi \in \partial\Sigma$, then $I \subset \{l+1, \dots, m\}$. In either case, we have that

$$|\nabla_{x_i}^g R^g(x_i^n)| = \mathcal{O}(1) \text{ and } |\nabla_{x_i}^g G^g(x_i^n, x_j^n)| = \mathcal{O}(1) \quad \text{as } n \rightarrow +\infty,$$

for $i \in I$ and $j \notin I$, and $|\nabla_{x_i}^g H^g(x_i^n, x_j^n)| = \mathcal{O}(1)$ for $i, j \in I$. Let $y_\xi : U(\xi) \rightarrow B^\xi \subset \mathbb{R}^2$ be the isothermal chart with $y_\xi(\xi) = 0$ introduced in Section 2. Then for $i \in I$, setting

$y_j^n := y_\xi(x_j^n) \in \mathbb{R}^2$ for $j \in I$ and assuming $y_i^n \neq 0$, we obtain as $n \rightarrow \infty$:

$$\begin{aligned}
|\nabla^g f_g(x^n)|_g &\geq |\nabla_{x_i}^g f_g(x^n)|_g = 2 \left| \nabla_{x_i}^g \sum_{j \in I \setminus \{i\}} \sigma_i \sigma_j G^g(x_i^n, x_j^n) \right|_g + \mathcal{O}(1) \\
&= \frac{2}{\kappa(\xi)} \left| \nabla_{x_i}^g \sum_{j \in I \setminus \{i\}} \sigma_i \sigma_j \log |y_\xi(x_i^n) - y_\xi(x_j^n)| \right|_g + \mathcal{O}(1) \\
&\geq \frac{2c}{\kappa(\xi)} \left| \nabla_{y_i} \sum_{j \in I \setminus \{i\}} \sigma_i \sigma_j \log |y_i^n - y_j^n| \right| + \mathcal{O}(1) \\
&= \frac{2c}{\kappa(\xi)} \left| \sum_{j \in I \setminus \{i\}} \sigma_i \sigma_j \frac{y_i^n - y_j^n}{|y_i^n - y_j^n|^2} \right| + \mathcal{O}(1) \\
&\geq \frac{2c}{\kappa(\xi)} \left| \sum_{j \in I \setminus \{i\}} \sigma_i \sigma_j \left\langle \frac{y_i^n - y_j^n}{|y_i^n - y_j^n|^2}, \frac{y_i^n}{|y_i^n|} \right\rangle \right| + \mathcal{O}(1)
\end{aligned}$$

The second inequality is a consequence of the fact that there exists $c > 0$ such that for any function $g : U(\xi) \rightarrow \mathbb{R}$:

$$|\nabla_x^g g(x)|_g \geq c |\nabla(g \circ y_\xi^{-1})(y_\xi(x))| \quad \text{for } x \in U(\xi).$$

Now for $n \in \mathbb{N}$ we choose $i(n) \in I$ with $|y_{i(n)}^n| \geq |y_j^n|$ for all $j \in I$. This implies $y_{i(n)}^n \neq 0$ and

$$\langle y_{i(n)}^n - y_j^n, y_{i(n)}^n \rangle \geq \frac{1}{2} |y_{i(n)}^n - y_j^n|^2 > 0 \quad \text{for } j \in I \setminus \{i(n)\},$$

hence

$$\left\langle \frac{y_{i(n)}^n - y_j^n}{|y_{i(n)}^n - y_j^n|^2}, \frac{y_{i(n)}^n}{|y_{i(n)}^n|} \right\rangle \geq \frac{1}{2|y_{i(n)}^n|} \quad \text{for } j \in I \setminus \{i(n)\}.$$

As a consequence we obtain, using $\sigma_i \sigma_j > 0$ for all $i, j \in I$:

$$|\nabla^g f_g(x^n)|_g \geq |\nabla_{x_{i(n)}}^g f_g(x^n)|_g \geq \frac{c}{\kappa(\xi)} \pi \sum_{j \in I \setminus \{i(n)\}} \sigma_{i(n)} \sigma_j \frac{1}{|y_{i(n)}^n|} + \mathcal{O}(1) \rightarrow \infty \quad \text{as } n \rightarrow \infty.$$

CASE 2. $x^\infty \notin \Delta_X$. Then there exists $i \in \{1, \dots, l\}$ such that $x_i^n \rightarrow \xi$ for some $\xi \in \partial\Sigma$.

We fix an isothermal chart $(y_\xi, U(\xi))$ around ξ as introduced in Section 2. For any $\zeta \in U_{r_\xi}(\xi)$, we decompose $G^g(\cdot, \zeta)$ as follows:

$$(4.1) \quad G^g(\cdot, \zeta) = \tilde{H}^g(\cdot, \zeta) - \frac{1}{\kappa(\zeta)} \chi(4|y_\xi(\cdot)| - y_\xi(\zeta)|/r_\xi) \log |y_\xi(\cdot) - y_\xi(\zeta)|.$$

Equation (2.2) implies $R^g(\zeta) = \tilde{H}^g(\zeta, \zeta) + \frac{1}{2\kappa(\zeta)} \varphi_\xi(y_\xi(\zeta))$. Let ∂_1, ∂_2 be the standard basis of \mathbb{R}^2 and define $\partial_{\zeta_1}, \partial_{\zeta_2} \in T_\zeta \Sigma$ be the corresponding basis of the tangent space of Σ at $\zeta \in U(\xi)$.

Fix $\zeta \in U_{r_\xi}(\xi) \cap \overset{\circ}{\Sigma}$ and set $\delta := y_\xi(\zeta)_2 > 0$. The representation formula for \tilde{H}^g yields:

$$\begin{aligned}
(4.2) \quad & \partial_{\zeta_1} \tilde{H}^g(\eta, \zeta)|_{\eta=\zeta} = \frac{1}{|\Sigma|_g} \int_{\Sigma} \partial_{\zeta_1} \tilde{H}^g(\cdot, \zeta) dv_g - \int_{\Sigma} G^g(\cdot, \zeta) \Delta_g \partial_{\zeta_1} \tilde{H}^g(\cdot, \zeta) dv_g \\
& + \int_{\partial\Sigma} G^g(\cdot, \zeta) \partial_{\nu_g} \partial_{\zeta_1} \tilde{H}^g(\cdot, \zeta) ds_g \\
& = \frac{1}{2\pi} \int_{\partial\Sigma} G^g(\zeta, x) \partial_{\nu_g} \partial_{\zeta_1} (\chi(4|y_\xi(x) - y_\xi(\zeta)|/r_\xi) \log |y_\xi(x) - y_\xi(\zeta)|) ds_g(x) + \mathcal{O}(1) \\
& = -\frac{1}{2\pi} \int_{B^\xi \cap \partial\mathbb{R}_+^2} G^g(\zeta, y_\xi^{-1}(y)) \chi(4|y - y_\xi(\zeta)|/r_\xi) \frac{2(y_1 - y_\xi(\zeta)_1)(y_2 - y_\xi(\zeta)_2)}{|y - y_\xi(\zeta)|^4} dy + \mathcal{O}(1) \\
& = -\frac{1}{2\pi\delta} \int_{-\varepsilon/\delta}^{\varepsilon/\delta} G^g(\zeta, y_\xi^{-1}((\delta s, 0) + (y_\xi(\zeta)_1, 0))) \frac{2s}{(s^2 + 1)^2} ds + \mathcal{O}(1).
\end{aligned}$$

as $\delta \rightarrow 0$. Here $\varepsilon \in (0, \frac{r_\xi}{16})$ is chosen sufficiently small. Decomposing G^g as in (4.1), we deduce for $\zeta \in U(\xi)$ with $y_\xi(\zeta)_2 < \frac{r_\xi}{16}$:

$$G^g(\zeta, y_\xi^{-1}((\delta s, 0) + (y_\xi(\zeta)_1, 0))) = \tilde{H}^g(\zeta, y_\xi^{-1}((\delta s, 0) + (y_\xi(\zeta)_1, 0))) - \frac{1}{\pi} \log |(\delta s, -\delta)|,$$

Now we apply the mean value theorem for \tilde{H}^g and obtain as $\delta s \rightarrow 0$:

$$(4.3) \quad \tilde{H}^g(\zeta, y_\xi^{-1}((\delta s, 0) + (y_\xi(\zeta)_1, 0))) = \tilde{H}^g(\zeta, y_\xi^{-1}(y_\xi(\zeta)_1, 0)) + \mathcal{O}(|\delta s| \sup_{x \in \partial\Sigma} \|\nabla_x \tilde{H}^g(\cdot, x)\|_{C(\Sigma)})$$

This implies as $\delta \rightarrow 0$:

$$\begin{aligned}
& \left| -\frac{1}{2\pi\delta} \int_{-\varepsilon/\delta}^{\varepsilon/\delta} G^g(\zeta, y_\xi^{-1}((\delta s, 0) + (y_\xi(\zeta)_1, 0))) \frac{2s}{(s^2 + 1)^2} ds \right| \\
& \leq \left| -\frac{1}{2\pi\delta} \tilde{H}^g(\zeta, y_\xi^{-1}(y_\xi(\zeta)_1, 0)) \int_{|s| \leq \varepsilon/\delta} \frac{2s}{(s^2 + 1)^2} ds \right| \\
& \quad + \left| \mathcal{O} \left(\int_{|s| \leq \varepsilon/\delta} \frac{2s^2}{(s^2 + 1)^2} ds \right) + \frac{1}{2\pi^2\delta} \int_{|s| \leq \varepsilon/\delta} \log(\delta\sqrt{s^2 + 1}) \frac{2s}{(s^2 + 1)^2} ds \right| \\
& \leq \mathcal{O}(1).
\end{aligned}$$

Now (4.2) yields $\partial_{\zeta_1} \tilde{H}^g(\zeta, \zeta) = \mathcal{O}(1)$ for $\zeta \in U_{r_\xi}(\xi) \cap \partial\Sigma$. Consequently, for $\zeta \in U_{r_\xi}(\xi)$ with $y_\xi(\zeta)_2 < \frac{r_\xi}{16}$, we have proven that:

$$(4.4) \quad \partial_{\zeta_1} R^g(\zeta) = \mathcal{O}(1) \quad \text{as } |y_\xi(\zeta)| \rightarrow 0.$$

The representation formula of \tilde{H}^g yields for $\zeta \in U_{r_\xi}(\xi) \cap \mathring{\Sigma}$ as $\delta \rightarrow 0$:

$$\begin{aligned}
\partial_{\zeta_2} \tilde{H}^g(\eta, \zeta)|_{\eta=\zeta} &= \frac{1}{|\Sigma|_g} \int_{\Sigma} \partial_{\zeta_2} \tilde{H}^g(\cdot, \zeta) dv_g - \int_{\Sigma} G^g(\cdot, \zeta) \Delta_g \partial_{\zeta_2} \tilde{H}^g(\cdot, \zeta) dv_g \\
&\quad + \int_{\partial\Sigma} G^g(\cdot, \zeta) \partial_{\nu_g} \partial_{\zeta_2} \tilde{H}^g(\cdot, \zeta) ds_g \\
&= \frac{1}{2\pi} \int_{\partial\Sigma} G^g(\zeta, x) \partial_{\nu_g} \partial_{\zeta_2} (\chi(4|y_\xi(x) - y_\xi(\zeta)|/r_\xi) \log |y_\xi(x) - y_\xi(\zeta)|) ds_g(x) + \mathcal{O}(1) \\
&\leq \frac{1}{2\pi} \int_{B_\xi \cap \partial\mathbb{R}_+^2} G^g(\zeta, y_\xi^{-1}(y)) \chi(4|y - y_\xi(\zeta)|/r_\xi) \frac{(y_1 - y_\xi(\zeta)_1)^2 - (y_2 - y_\xi(\zeta)_2)^2}{|y - y_\xi(\zeta)|^4} dy + \mathcal{O}(1) \\
&\leq \frac{1}{2\pi\delta} \int_{-\varepsilon/\delta}^{\varepsilon/\delta} G^g(\zeta, y_\xi^{-1}((\delta s, 0) + (y_\xi(\zeta)_1, 0))) \frac{s^2 - 1}{(s^2 + 1)^2} ds + \mathcal{O}(1) \\
&\stackrel{(4.3)}{\leq} \frac{1}{2\pi\delta} \tilde{H}^g(\zeta, y_\xi^{-1}(y_\xi(\zeta)_1, 0)) \int_{|s| \leq \varepsilon/\delta} \frac{s^2 - 1}{(s^2 + 1)^2} ds + \mathcal{O}\left(1 + \int_{|s| \leq \varepsilon/\delta} \frac{|s(s^2 - 1)|}{(s^2 + 1)^2} ds\right) \\
&\quad + \frac{\log(\delta^{-1})}{2\pi^2\delta} \int_{|s| \leq \varepsilon/\delta} \frac{s^2 - 1}{(s^2 + 1)^2} ds + \frac{1}{2\pi^2\delta} \int_{|s| \leq \varepsilon/\delta} \log\left(\frac{1}{\sqrt{s^2 + 1}}\right) \frac{s^2 - 1}{(s^2 + 1)^2} ds + \mathcal{O}(1) \\
&\leq \frac{1}{2\pi^2\delta} \int_{|s| \geq 1} \log\left(\frac{\sqrt{s^{-2} + 1}}{\sqrt{s^2 + 1}}\right) \frac{s^2 - 1}{(s^2 + 1)^2} ds - \frac{1}{2\pi^2\delta} \int_{|s| \geq \varepsilon/\delta} \log\left(\frac{1}{\sqrt{s^2 + 1}}\right) \frac{s^2 - 1}{(s^2 + 1)^2} ds \\
&\quad - \frac{\log(\delta^{-1})\varepsilon}{\pi^2(\delta^2 + \varepsilon^2)} + \mathcal{O}(\delta^{-1/2}) + \mathcal{O}(\log(\delta^{-1}) + 1) \\
&\leq -\frac{1}{2\pi^2\delta} \int_{|s| \geq 1} \log(s^2) \frac{s^2 - 1}{(s^2 + 1)^2} ds + \mathcal{O}(\delta^{-\frac{1}{2}}) \leq -\frac{1}{4\pi\delta}.
\end{aligned}$$

The last inequality used the identity

$$\int_{|s| \geq 1} \log(s^2) \frac{s^2 - 1}{(s^2 + 1)^2} ds = \pi.$$

From the above estimate we deduce for $\zeta \in \mathring{\Sigma} \cap U_{r_\xi}(\xi)$:

$$(4.5) \quad \partial_{\zeta_2} R^g(\zeta) \leq -\frac{1}{2\pi|y_\xi(\zeta)_2|} \quad \text{as } |y_\xi(\zeta)_2| \rightarrow 0.$$

Now if $I := \{i \in \{1, \dots, m\} : x_i^n \rightarrow \xi\}$ contains only a single element i then (4.5) yields

$$\begin{aligned}
|\nabla^g f_g(x^n)|_g &\geq |\nabla_{x_i}^g f_g(x^n)|_g \geq |\sigma_i^2 \nabla_{x_i}^g R^g(x_i^n) + 2\sigma_i \sum_{j \neq i} \nabla_{x_i}^g G^g(x_i^n, x_j^n) + \nabla_{x_i}^g h(x^n)|_g \\
&\geq \sigma_i^2 |\nabla_{x_i}^g R^g(x_i^n)| + \mathcal{O}(1) \geq \frac{c\sigma_i^2}{2\pi|y_\xi(x_i^n)_2|} + \mathcal{O}(1) \\
&\rightarrow \infty \quad \text{as } n \rightarrow \infty.
\end{aligned}$$

Here $c > 0$ is a constant such that for any function $F : U(\xi) \rightarrow \mathbb{R}$:

$$(4.6) \quad |\nabla_x^g F(x)|_g \geq c |\nabla(F \circ y_\xi^{-1})(y_\xi(x))| \quad \text{for } x \in U(\xi).$$

Next we consider the case that I contains at least two elements. Then $\xi_i^n \in U(\xi)$ for n large and $i \in I$. By a direct calculation we obtain for $j \in I \setminus \{i\}$, $\iota = 1, 2$ and $n \rightarrow \infty$:

$$\begin{aligned}
(4.7) \quad & \partial_{\zeta_\iota} G^g(x_j^n, \zeta)|_{\zeta=x_i^n} = \partial_{\zeta_\iota} \tilde{H}^g(x_j^n, \zeta)|_{\zeta=x_i^n} \\
& + \frac{1}{\kappa(x_i^n)} \log \frac{1}{|y_\xi(x_j^n) - y_\xi(x_i^n)|} \partial_{\zeta_\iota} \chi(4|y_\xi(x_j^n) - y_\xi(\zeta)|/r_\xi)|_{\zeta=x_i^n} \\
& + \frac{1}{\kappa(x_i^n)} \chi(4|y_\xi(x_j^n) - y_\xi(x_i^n)|/r_\xi) \partial_{\zeta_\iota} \log \frac{1}{|y_\xi(x_j^n) - y_\xi(\zeta)|} \Big|_{\zeta=x_i^n} \\
& = \partial_{\zeta_\iota} \tilde{H}^g(x_j^n, \zeta)|_{\zeta=x_i^n} - \frac{1}{\kappa(x_i^n)} \frac{(y_\xi(x_i^n) - y_\xi(x_j^n))_\iota}{|y_\xi(x_j^n) - y_\xi(x_i^n)|^2} + \mathcal{O}(1).
\end{aligned}$$

For $n \in \mathbb{N}$ we set

$$\varrho_n := \max \left(\{y_\xi(x_i^n)_2 : i \in I\} \cup \{|y_\xi(x_i^n)_1 - y_\xi(x_j^n)_1| : i, j \in I \text{ with } i \neq j\} \right).$$

If there exists $i(n) \in I$ such that $\varrho_n = y_\xi(x_{i(n)}^n)_2$, then $i(n) \in \{1, \dots, l\}$ satisfies

$$y_\xi(x_{i(n)}^n)_2 - y_\xi(x_j^n)_2 \geq 0 \text{ and } y_\xi(x_{i(n)}^n)_2 \geq |(y_\xi(x_i^n)_1 - y_\xi(x_j^n)_1)| \quad \text{for every } i, j \in I.$$

Given $\zeta = x_{i(n)}^n$ and $\eta = x_j^n$ with $j \in I \cap \{1, \dots, l\} \setminus \{i(n)\}$, we will calculate the upper bound of $\partial_{\zeta_2} \tilde{H}^g(\eta, \zeta)$ as $n \rightarrow 0$. Setting $a = y_\xi(\eta)_2 > 0$ and $b = y_\xi(\zeta)_1 - y_\xi(\eta)_1$ we have for $|b| \leq \varrho_n$ as $n \rightarrow \infty$:

$$\begin{aligned}
\partial_{\zeta_2} \tilde{H}^g(\eta, \zeta) &= \frac{1}{|\Sigma|_g} \int_{\Sigma} \partial_{\zeta_2} \tilde{H}^g(\cdot, \zeta) dv_g - \int_{\Sigma} G^g(\cdot, \eta) \Delta_g \partial_{\zeta_2} \tilde{H}^g(\cdot, \zeta) dv_g \\
&\quad + \int_{\partial\Sigma} G^g(\cdot, \eta) \partial_{\nu_g} \partial_{\zeta_2} \tilde{H}^g(\cdot, \zeta) ds_g \\
&= \mathcal{O}(1) + \frac{1}{2\pi} \int_{\partial\Sigma} G^g(\eta, x) \partial_{\nu_g} \partial_{\zeta_2} (\chi(4|y_\xi(x) - y_\xi(\zeta)|/r_\xi) \log |y_\xi(x) - y_\xi(\zeta)|) ds_g(x) \\
&\leq \mathcal{O}(1) + \frac{1}{2\pi} \int_{B^\xi \cap \partial\mathbb{R}_+^2} G^g(\eta, y_\xi^{-1}(y)) \chi(4|y - y_\xi(\zeta)|/r_\xi) \frac{(y_1 - y_\xi(\zeta)_1)^2 - (y_2 - y_\xi(\zeta)_2)^2}{|y - y_\xi(\zeta)|^4} dy \\
&\leq \mathcal{O}(1) + \frac{1}{2\pi \varrho_n} \int_{-\varepsilon/\varrho_n}^{\varepsilon/\varrho_n} G^g(\eta, y_\xi^{-1}((\varrho_n s, 0) + (y_\xi(\zeta)_1, 0))) \frac{s^2 - 1}{(s^2 + 1)^2} ds \\
&\stackrel{(4.3)}{\leq} \frac{1}{2\pi \varrho_n} \tilde{H}^g(\eta, y_\xi^{-1}(y_\xi(\zeta)_1, 0)) \int_{|s| \leq \varepsilon/\varrho_n} \frac{s^2 - 1}{(s^2 + 1)^2} ds + \mathcal{O} \left(1 + \int_{|s| \leq \varepsilon/\varrho_n} \frac{|s(s^2 - 1)|}{(s^2 + 1)^2} ds \right) \\
&\quad - \frac{\log(a)}{2\pi^2 \varrho_n} \int_{|s| \leq \varepsilon/\varrho_n} \frac{s^2 - 1}{(s^2 + 1)^2} ds + \frac{1}{2\pi^2 \varrho_n} \int_{|s| \leq \varepsilon/\varrho_n} \log \left(\frac{1}{\sqrt{1 + (\varrho_n s + b)^2/a^2}} \right) \frac{s^2 - 1}{(s^2 + 1)^2} ds \\
&\leq -\frac{1}{2\pi^2 \varrho_n} \int_{|s| \geq \varepsilon/\varrho_n} \left(\log(a^{-1}) + \log \left(\frac{1}{\sqrt{1 + (\varrho_n s + b)^2/a^2}} \right) \right) \frac{s^2 - 1}{(s^2 + 1)^2} ds \\
&\quad + \frac{1}{2\pi^2 \varrho_n} \int_{|s| \geq 1} \log \left(\frac{\sqrt{1 + (\varrho_n s^{-1} + b)^2/a^2}}{\sqrt{1 + (\varrho_n s + b)^2/a^2}} \right) \frac{s^2 - 1}{(s^2 + 1)^2} ds + \mathcal{O}(1 + \log(\varrho_n^{-1})) \\
&\leq \mathcal{O}(\log(\varrho_n^{-1}))
\end{aligned}$$

Here we used the inequalities:

$$\log \left(\frac{\sqrt{1 + (\varrho_n s^{-1} + b)^2/a^2}}{\sqrt{1 + (\varrho_n s + b)^2/a^2}} \right) \leq 0, \text{ for } |b| \leq \varrho_n, |s| \geq 1$$

and

$$0 \leq \log \sqrt{a^2 + (\varrho_n s + b)^2} \frac{s^2 - 1}{(s^2 + 1)^2} \leq C \frac{\varrho_n^{\frac{1}{2}}}{s^{\frac{3}{2}}}$$

for some constant $C > 0$, any $s \in \{s \in \mathbb{R} : |s| \geq \varepsilon/\varrho_n, a^2 + (\varrho_n s + b)^2 \geq 1\}$. We notice that we have for $j \in I \cap \{l+1, \dots, m\}$:

$$(4.8) \quad |\partial_{\zeta_2} \tilde{H}^g(x_j^n, \zeta)|_{\zeta=x_{i(n)}^n} = |\partial_{\zeta_2} \tilde{H}^g(\zeta, x_j^n)|_{\zeta=x_{i(n)}^n} \leq \mathcal{O}(1) \quad \text{as } n \rightarrow \infty;$$

and for $j \in I \cap \{1, \dots, l\} \setminus \{i(n)\}$, $\varrho_n = y_\xi(x_{i(n)}^n)$:

$$(4.9) \quad |\partial_{\zeta_2} \tilde{H}^g(x_j^n, \zeta)|_{\zeta=x_{i(n)}^n} \leq \mathcal{O}(\log(\varrho_n^{-1})) \quad \text{as } n \rightarrow \infty.$$

Now (4.5)-(4.9) imply for $n \rightarrow \infty$:

$$\begin{aligned} |\nabla^g f_g(x^n)|_g &\geq \left| \nabla_{x_{i(n)}}^g \left(\sigma_i^2 R^g(x_{i(n)}^n) + 2\sigma_{i(n)} \sum_{j \neq i(n)} \sigma_j G^g(x_{i(n)}^n, x_j^n) + h(x^n) \right) \right|_g \\ &\geq c \left| \partial_{(x_{i(n)})_2} \left(\sigma_{i(n)}^2 R^g(\xi_{i(n)}^n) + 2\sigma_{i(n)} \sum_{j \in I \setminus \{i(n)\}} \sigma_j G^g(\xi_{i(n)}, \xi_j) \right) \right| + \mathcal{O}(1) \\ &\geq \mathcal{O}(\log(\varrho_n^{-1})) + \sigma_{i(n)}^2 \frac{c}{2\pi \varrho_n} \\ &\rightarrow \infty. \end{aligned}$$

Here $c > 0$ is as in (4.6). If $\varrho_n > \max\{y_\xi(x_i^n) : i \in I\}$, then we take $i(n) \in I$ such that $y_\xi(x_{i(n)}^n)_1 = \max\{y_\xi(x_i^n) : i \in I\}$. The proof of Lemma 6 in [16] implies for $x \in U(\xi)$ and $y_\xi^*(x) := (y_\xi(x)_1, -y_\xi(x)_2)$:

$$\left\| G^g(\cdot, x) + \frac{1}{2\pi} \log |y_\xi(\cdot) - y_\xi(x)| + \frac{1}{2\pi} \log |y_\xi(\cdot) - y_\xi^*(x)| \right\|_{C^1(U_{r_\xi}(\xi))} \leq C$$

where $C > 0$ depends only on (Σ, g) and on ξ . Therefore we have for $j \in I \setminus \{i(n)\}$ as $n \rightarrow \infty$:

$$(4.10) \quad \partial_{\zeta_1} G(x_j^n, \zeta) \Big|_{\zeta=x_{i(n)}^n} \leq -\frac{1}{2\pi} \frac{y_\xi(x_{i(n)}^n)_1 - y_\xi(x_j^n)_1}{|y_\xi(x_{i(n)}^n) - y_\xi(x_j^n)|^2} - \frac{1}{2\pi} \frac{y_\xi(x_{i(n)}^n)_1 - y_\xi(x_j^n)_1}{|y_\xi(x_{i(n)}^n) - y_\xi^*(x_j^n)|^2} + \mathcal{O}(1).$$

We can take $i'(n) \in I \setminus \{i(n)\}$ such that $\varrho_n = y_\xi(x_{i(n)}^n)_1 - y_\xi(x_{i'(n)}^n)_1$ by the assumption. The inequality (4.10) yields

$$(4.11) \quad \partial_{\zeta_1} G(x_{i'(n)}^n, \zeta) \Big|_{\zeta=x_{i(n)}^n} \leq -\frac{1}{4\pi \varrho_n} \quad \text{as } n \rightarrow \infty.$$

From (4.4) together with (4.10) and (4.11) we derive the following estimate for the gradient of f_g as $n \rightarrow \infty$:

$$\begin{aligned}
|\nabla^g f_g(x^n)|_g &\geq \left| \nabla_{x_{i(n)}}^g \left(\sigma_i^2 R^g(x_{i(n)}^n) + 2\sigma_{i(n)} \sum_{j \neq i(n)} \sigma_j G^g(x_{i(n)}^n, x_j^n) + h(x^n) \right) \right|_g \\
&\geq c \left| \partial_{(x_{i(n)})_1} \left(\sigma_{i(n)}^2 R^g(\xi_{i(n)}^n) + 2\sigma_{i(n)} \sum_{j \in I \setminus \{i(n)\}} \sigma_j G^g(\xi_{i(n)}, \xi_j) \right) \right| + \mathcal{O}(1) \\
&\geq \mathcal{O}(1) + \sigma_{i(n)} \sigma(i'(n)) \frac{c}{2\pi\varrho_n} \\
&\rightarrow \infty
\end{aligned}$$

Again $c > 0$ is as in (4.6). This completes the proof of Lemma 4.1. \square

References

- [1] AHMEDOU, M., BARTSCH, T., AND FIERNKRAZ, T. Equilibria of vortex type Hamiltonians on closed surfaces. *Topol. Methods Nonlinear Anal.* 61, 1 (2023), 239–256.
- [2] BARAKET, S., AND PACARD, F. Construction of singular limits for a semilinear elliptic equation in dimension 2. *Calc. Var. Partial Differ. Equ.* 6, 1 (1997), 1–38.
- [3] BARTSCH, T., MICHELETTI, A. M., AND PISTOIA, A. The Morse property for functions of Kirchhoff-Routh path type. *Discrete Contin. Dyn. Syst.-S* 12, 7 (2019), 1867–1877.
- [4] CHERN, S.-S. An elementary proof of the existence of isothermal parameters on a surface. *Proc. Amer. Math. Soc.* 6 (1955), 771–782.
- [5] DEL PINO, M., KOWALCZYK, M., AND MUSSO, M. Singular limits in Liouville-type equations. *Calc. Var. Partial Differ. Equ.* 24, 1 (2005), 47–81.
- [6] DEL PINO, M., AND WEI, J. Collapsing steady states of the Keller–Segel system. *Nonlinearity* 19, 3 (2006), 661–684.
- [7] ESPOSITO, P., AND FIGUEROA, P. Singular mean field equations on compact Riemann surfaces. *Nonlinear Anal.* 111 (2014), 33–65.
- [8] ESPOSITO, P., GROSSI, M., AND PISTOIA, A. On the existence of blowing-up solutions for a mean field equation. *Ann. Inst. H. Poincaré Anal. Non Linéaire* 22, 2 (2005), 227–257.

- [9] FIGUEROA, P. Bubbling solutions for mean field equations with variable intensities on compact Riemann surfaces. *J. d'Analyse Math.*, DOI 10.1007/s11854-023-0303-2 (2023).
- [10] GLADIALI, F., AND GROSSI, M. Some results for the Gelfand's problem. *Commun. Partial Differ. Equ.* 29, 9-10 (2005), 1335–1364.
- [11] HARTMAN, P., AND WINTNER, A. On uniform Dini conditions in the theory of linear partial differential equations of elliptic type. *Amer. J. Math.* 77 (1955), 329–354.
- [12] HENRY, D. *Perturbation of the Boundary in Boundary-Value Problems of Partial Differential Equations*. London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 2005.
- [13] MA, L., AND WEI, J. Convergence for a Liouville equation. *Comment. Math. Helv.* 76, 3 (Sep. 2001), 506–514.
- [14] NARDI, G. Schauder estimate for solutions of Poisson's equation with Neumann boundary condition. *Enseign. Math.* 60, 3/4 (2014), 421–435.
- [15] VEKUA, I. N. The problem of reduction to canonical form of differential forms of elliptic type and the generalized Cauchy-Riemann system. *Dokl. Akad. Nauk SSSR (N.S.)* 100 (1955), 197–200.
- [16] YANG, Y., AND ZHOU, J. Blow-up analysis involving isothermal coordinates on the boundary of compact Riemann surface. *J. Math. Anal. Appl.* 504, 2 (2021), 125440.

THOMAS BARTSCH, ZHENGN NI HU
 Mathematisches Institut
 Universität Giessen
 Arndtstr. 2
 35392 Giessen, Germany
 Thomas.Bartsch@math.uni-giessen.de
 Zhengni.Hu@math.uni-giessen.de