

Classification of π_1 -injective proper maps between non-compact surfaces

Sumanta Das

Abstract

We show that almost all π_1 -injective proper maps between two non-compact surfaces, where surfaces are possibly of infinite type, can be properly homotoped to finite-sheeted covering maps.

1 Introduction

All manifolds will be assumed to be second countable and Hausdorff. A *surface* (resp. *bordered surface*) is a connected, orientable 2-dimensional manifold with an empty (resp. a non-empty) boundary. We say a connected 2-manifold with or without boundary is of *finite type* if its fundamental group is finitely generated; otherwise, we say it is of *infinite type*.

In 1927, Nielsen [21] proved that any π_1 -injective map between two compact surfaces is homotopic to a covering map. In dimension three, achieving an analog classification of π_1 -injective proper maps is also possible, subject to various constraints. For example, Waldhausen [29, Theorem 6.1.] proved that if $f: N \rightarrow M$ is π_1 -injective map between two connected, closed, orientable, irreducible 3-manifolds, where N is non-simply-connected and M is Haken, then f homotopic to a covering map. Brown and Tucker [1, Theorem 4.2] showed that if $f: \mathfrak{N} \rightarrow \mathfrak{M}$ is a π_1 -injective proper map between two connected, non-compact, orientable, irreducible, end-irreducible, boundaryless 3-manifolds such that $\pi_1(\mathfrak{N})$ is not isomorphic to the fundamental group of any compact surface, then f is properly homotopic to a finite-sheeted covering map.

In this note, we prove that most π_1 -injective proper maps between non-compact surfaces become finite-sheeted covering maps through proper homotopy. Our first theorem considers all surfaces except the plane \mathbb{C} and the punctured plane \mathbb{C}^* , where $\mathbb{C}^* := \mathbb{C} \setminus \{0\}$. The next two theorems consider these two excluded surfaces separately.

Theorem 1.1 Let Σ', Σ be two non-compact oriented surfaces such that Σ' is neither the plane nor the punctured plane. Suppose $f: \Sigma' \rightarrow \Sigma$ is a π_1 -injective proper map. Then f is properly homotopic to a d -sheeted covering map $p: \Sigma' \rightarrow \Sigma$ for some positive integer d . Thus, $\deg(f) = \pm d (\neq 0)$. Moreover, Σ' is of infinite type if and only if Σ is of infinite type.

Theorem 1.2 Suppose f is a π_1 -injective proper map from \mathbb{C}^* to a non-compact oriented surface Σ . Let $n := \deg(f)$. Then, we have the following.

- (i) If $n = 0$, then there exists a π_1 -injective, proper embedding $\iota: \mathbb{S}^1 \times [0, \infty) \hookrightarrow \Sigma$, along with a non-zero integer d , such that after a proper homotopy, f can be described by the proper map $\mathbb{S}^1 \times \mathbb{R} \ni (z, t) \mapsto \iota(z^d, |t|) \in \Sigma$. Thus, Σ has an isolated planar end, and given any compact subset K of Σ , there exists a proper map g properly homotopic to f such that $\text{im}(g) \subseteq \Sigma \setminus K$.
- (ii) If $n \neq 0$, then $\Sigma = \mathbb{C}^*$ and f is properly homotopic to the covering $\mathbb{C}^* \ni z \mapsto z^n \in \mathbb{C}^*$ if $n > 0$, and to the covering $\mathbb{C}^* \ni z \mapsto \bar{z}^{-n} \in \mathbb{C}^*$ if $n < 0$.

Theorem 1.3 Suppose f is a proper map from \mathbb{C} to a non-compact oriented surface Σ . Let $n := \text{deg}(f)$. Then, we have the following.

- (i) If $n = 0$, then for every compact subset K of Σ , there exists a proper map g properly homotopic to f such that $\text{im}(g) \subseteq \Sigma \setminus K$.
- (ii) If $n \neq 0$, then $\Sigma = \mathbb{C}$, and f is properly homotopic to the branched covering $\mathbb{C} \ni z \mapsto z^n \in \mathbb{C}$ if $n > 0$, and to the branched covering $\mathbb{C} \ni z \mapsto \bar{z}^{-n} \in \mathbb{C}$ if $n < 0$.

1.4 Idea of the proof

It might be possible to prove [Theorem 1.1](#) by modifying the approach used by Brown and Tucker [1, Theorem 4.2] as every non-compact surface admits exhaustion by essential compact bordered sub-surfaces. However, we will choose a different method to establish [Theorem 1.1](#). In fact, we will employ the strong topological rigidity theorem [3, Theorem 1], which states that if a homotopy equivalence between two non-compact surfaces is a proper map, then it is properly homotopic to a homeomorphism, provided that the surfaces are neither the plane nor the punctured plane. It is not difficult to see that the strong topological rigidity, along with the map lifting lemma of covering space theory, directly implies that the map f in [Theorem 1.1](#) is homotopic to a covering map. However, to prove that this covering map can be realized as finite-sheeted and that this homotopy can be realized as proper homotopy requires an argument based on the non-zero degree of the lifting map. See [Theorem 3.4](#) for a proof.

The proofs of the remaining two theorems require modifying the surgery techniques on a proper map between non-compact surfaces mentioned in [Theorem 3.6](#) and [Theorem 3.10](#). The necessity of these techniques can be described as follows: In order to prove [Theorem 1.3\(i\)](#), one may use [Theorem 2.2.2](#) to properly homotope f so that $\text{im}(f)$ is contained in the complement of a disk in Σ . However, [Theorem 2.2.2](#) does not indicate, given any compact subset K of Σ , whether f can be properly homotoped so that $\text{im}(f)$ is contained in $\Sigma \setminus K$. Here, we need to use a variation of [Theorem 3.6](#); see [Theorem 3.8](#) for proof. Now, let's consider the strategy for proving [Theorem 1.3\(ii\)](#). Since the domain of a non-zero degree map always contains 'more loops than its co-domain' (see [Theorem 3.9](#)), it's easy to see that $\Sigma = \mathbb{C}$. At this point, by applying [Theorem 2.2.2](#), we can at most expect to obtain a disk D such that $f^{-1}(D)$ is a pairwise-disjoint union of disks D_1, \dots, D_n , each mapped homeomorphically onto D . Now, if $n = 1$, then by first applying Cerf-Palais' disk theorem and then the Alexander trick, we can conclude the proof. However, when $n \geq 2$, we can't conclude in this manner. This is where we need to utilize those surgery techniques to transform $f^{-1}(D)$ into a single disk D' , such that $f|_{\partial D'} \rightarrow \partial D$ is an n -sheeted covering. See [Theorem 3.15](#) for proof.

To prove [Theorem 1.2\(i\)](#), one must apply these surgery techniques to a one-manifold \mathcal{A} that decomposes Σ into bordered sub-surfaces, with each complementary component being either a one-holed torus, a pair of pants, or a punctured disk. Therefore, after a proper homotopy, we

may assume f sends the whole \mathbb{C}^* into a punctured disk D_* , which appears as a complementary component of this decomposition. This is where we are using the fact that any two non-trivial circles on \mathbb{C}^* co-bound an annulus. If e_ℓ is the isolated planar end of Σ determined by D_* , then $\text{Ends}(f)$ sends both elements of $\text{Ends}(\mathbb{C}^*)$ to e_ℓ . Now, the same analogy that we used to prove [Theorem 1.3\(i\)](#) can be applied here to show that for every compact subset K of Σ , f can be properly homotoped such that $\text{im}(f)$ becomes disjoint from K . See [Theorem 3.20](#) for proof. On the other hand, unlike [Theorem 1.3\(ii\)](#), we can use the advantage of the non-triviality of the fundamental group of \mathbb{C}^* to prove [Theorem 1.2\(ii\)](#). Since any two subgroups of \mathbb{Z} of index n are the same, to prove [Theorem 1.2\(ii\)](#), all we need to show is that f is properly homotopic to a covering map, and any homeomorphism of \mathbb{C}^* is properly homotopic to either the identity or the complex conjugation. The former fact can be proved using the same analogy that we employed in proving [Theorem 1.1](#), while the latter follows from a result by Epstein [4, Theorem 5.7.]. See [Theorem 3.24](#) for proof.

2 Preliminaries

2.1 Proper maps

Let X and Y be topological spaces. A map from X to Y is called a *proper map* if the inverse image of each compact subset of Y is a compact subset of X . A sequence $\{x_n\}$ in X is said to *diverge to infinity* if, for every compact set $K \subseteq X$, there are at most finitely many values of n for which $x_n \in K$. If X is second countable Hausdorff, then a map $f: X \rightarrow Y$ is proper if and only if f takes sequences diverging to infinity in X to sequences diverging to infinity in Y [19, Propositions 4.92. and 4.93.(b)]. A proper map between manifolds is a closed map. More generally, a proper map from a space to a compactly generated Hausdorff space is a closed map [19, Theorem 4.95] [25].

If a homotopy $\mathcal{H}: X \times [0, 1] \rightarrow Y$ is a proper map, then we call \mathcal{H} a *proper homotopy*. Two proper maps from X to Y are said to be *properly homotopic* if there is a proper homotopy between them. It is possible that a homotopy $H: X \times [0, 1] \rightarrow Y$ through proper maps may fail to be a proper homotopy. For example, the homotopy $H: \mathbb{C} \times [0, 1] \ni (z, t) \mapsto tz^2 - z \in \mathbb{C}$ (the pre-image of 0 under H contains the pair $(n, 1/n)$ for each positive integer n). However, a homotopy through homeomorphisms of a manifold is proper homotopy. More generally, if X is a locally compact Hausdorff space and $\mathcal{H}: X \times [0, 1] \rightarrow X$ is a homotopy through homeomorphisms, then $X \times [0, 1] \ni (x, t) \mapsto (\mathcal{H}(x, t), t) \in X \times [0, 1]$ is a homeomorphism [2, Theorem 1.3], and thus \mathcal{H} is a proper map.

2.2 The degree of a proper map

Let M_1 and M_2 be two connected, oriented, topological n -manifolds. Denote the preferred generator of the n -th singular cohomology with compact support $H_c^n(M_j, \partial M_j; \mathbb{Z}) \cong \mathbb{Z}$ compatible with the orientation of M_j by $[M_j]$ for each $j = 1, 2$. If $f: (M_1, \partial M_1) \rightarrow (M_2, \partial M_2)$ is a proper map, then the *degree* of f is the unique integer $\text{deg}(f)$ that satisfies $H_c^n(f)([M_2]) = \text{deg}(f) \cdot [M_1]$. If manifolds are compact, then the notion of compactly supported cohomological degree agrees with the notion of the usual degree defined by singular cohomology. Note that if $f, g: (M_1, \partial M_1) \rightarrow (M_2, \partial M_2)$ are proper maps such that there is a proper homotopy $\mathcal{H}: M_1 \times [0, 1] \rightarrow M_2$ with $\mathcal{H}(\partial M_1 \times [0, 1]) \subseteq \partial M_2$ from f to g , then $\text{deg}(f) = \text{deg}(g)$. Moreover, the degree is multiplicative, i.e., the degree of the composition of two proper maps is the product of their degrees.

In the following two theorems, when we mention a disk D in a smooth n -manifold X , we mean that D is the image of $\{z \in \mathbb{R}^n : |z| \leq 1\}$ under a smooth embedding $\{z \in \mathbb{R}^n : |z| \leq 2\} \hookrightarrow X$. The first one is a useful tool to calculate the degree, and the second one geometrically realizes the (algebraic) notion of the degree.

Theorem 2.2.1 [5, Lemma 2.1b.] Let $f: M \rightarrow N$ be a proper map between two connected, oriented, smooth manifolds of the same dimension such that $f^{-1}(\partial N) = \partial M$. Suppose there exists a disk D in $\text{int}(N)$ with the property that $f^{-1}(D)$ is the pairwise disjoint union of disks D_1, \dots, D_k in $\text{int}(M)$ (k must be a non-negative integer as f is proper) such that f maps each $f^{-1}(D_i)$ homeomorphically onto D . For each i , let ε_i be $+1$ or -1 according as the homeomorphism $f|_{f^{-1}(D_i)}: f^{-1}(D_i) \rightarrow D$ is orientation-preserving or orientation-reversing. Then $\deg(f) = \sum_{i=1}^k \varepsilon_i$. In particular, if $f^{-1}(D) = \emptyset$, then $\deg(f) = 0$.

Theorem 2.2.2 [16, 17] [5, Theorems 3.1 and 4.1] Let $f: M \rightarrow N$ be a proper map between two connected, oriented, smooth manifolds of the same dimension such that $f^{-1}(\partial N) \subseteq \partial M$. Let $\ell := |\deg(f)|$. Then there is a proper map $g: M \rightarrow N$ with $g(\partial M) \subseteq \partial N$ and a homotopy $\mathcal{H}: M \times [0, 1] \rightarrow N$ with the following properties:

- There exists a compact subset $K \subseteq \text{int}(M)$ such that $\mathcal{H}(x, t) = f(x)$ for all $(x, t) \in (M \setminus K) \times [0, 1]$. In particular, \mathcal{H} is a proper homotopy relative to ∂M .
- There exists a disk $D \subseteq \text{int}(N)$ such that $g^{-1}(D)$ is the pairwise disjoint union of disks D_1, \dots, D_ℓ in $\text{int}(M)$ and g maps each $g^{-1}(D_i)$ homeomorphically onto D .

Therefore, if the degree of g , which is the same as the degree of f , is positive (resp. negative), then $g|_{g^{-1}(D_i)}: g^{-1}(D_i) \rightarrow D$ is an orientation-preserving (resp. orientation-reversing) homeomorphism for each $i = 1, \dots, \ell$. In particular, if $\ell = 0$, then f can be properly homotoped to a non-surjective map, relative to the complement of a compact subset of $\text{int}(M)$.

The following theorem is a well-known fact whose proof in the particular case when manifolds are boundaryless can be found in [3, Lemma 3.6.4.1].

Theorem 2.2.3 Let $f: (M, \partial M) \rightarrow (N, \partial N)$ be a proper map between two connected, oriented, smooth manifolds of the same dimension. If $\deg(f) \neq 0$, then f is surjective.

2.3 Ends of spaces

Let X be a connected topological manifold. Then there is a nested sequence $K_1 \subseteq K_2 \subseteq \dots$ of compact, connected subsets of X such that $\cup_i K_i = X$, $K_i \subseteq \text{int}(K_{i+1})$ for each i , $\cap_i (X \setminus K_i) = \emptyset$, and the closure of each component of any $X \setminus K_i$ is non-compact [12, Exercise 3.3.4]. We refer to such a nested sequence as an *efficient exhaustion by compacta* of X .

Let $\text{Ends}(X)$ be the set of all sequences (V_1, V_2, \dots) , where V_i is a component of $X \setminus K_i$ and $V_1 \supseteq V_2 \supseteq \dots$. Consider $X^\dagger := X \cup \text{Ends}(X)$ with the topology generated by the basis consisting of all open subsets of X , and all sets V_i^\dagger , where $V_i^\dagger := V_i \cup \{(V'_1, V'_2, \dots) \in \text{Ends}(X) | V'_i = V_i\}$. Then X^\dagger is a separable, compact, and metrizable space and X is an open dense subset of X^\dagger . That is, X^\dagger is a compactification of X , often referred to as the *Freudenthal compactification* of X . The subspace $\text{Ends}(X)$ of X^\dagger is a totally disconnected space. Thus, $\text{Ends}(X)$ is a closed subset of the Cantor set. Note that the space $\text{Ends}(X)$ is independent of the choice of efficient exhaustion of X by compacta.

Now, let Y be another connected topological manifold. Recall that every proper map $f: X \rightarrow Y$ induces a map $\text{Ends}(f): \text{Ends}(X) \rightarrow \text{Ends}(Y)$ that can be used to extend $f: X \rightarrow Y$ to a map $f^\dagger: X^\dagger \rightarrow Y^\dagger$ between the Freudenthal compactifications. This induced map between the spaces of ends can be defined as follows:

Let $f: X \rightarrow Y$ be a proper map. Choose efficient exhaustions by compacta $\{K_i\}$ and $\{L_i\}$ of X and Y , respectively. Consider an end $e = (V_1, V_2, \dots)$ of X , where V_i is a component of $X \setminus K_i$. Notice that each entry V_i of (V_1, V_2, \dots) determines all entries of the lower index; hence, every subsequence of entries of an end determines that end. By properness, for each i , there is a j_i such that $f(X \setminus K_{j_i}) \subseteq Y \setminus L_i$. Without loss of generality, we may assume $j_1 < j_2 < \dots$. By connectedness, f sends V_{j_i} into a component, say W_i , of $Y \setminus L_i$. Let $e' := (W_1, W_2, \dots)$, and $\text{Ends}(f)(e) := e'$. Now, define $f^\dagger: X^\dagger \rightarrow Y^\dagger$ as $f^\dagger|_X := f$ and $f^\dagger|_{\text{Ends}(X)} := \text{Ends}(f)$.

The map $f^\dagger: X^\dagger \rightarrow Y^\dagger$ thus defined is continuous [12, Proposition 3.3.12 (a)]. Also, one can show that the induced map of the identity is the identity, and the induced map of a composition of two proper maps is the composition of their induced maps. Moreover, if two proper maps $f_0, f_1: X \rightarrow Y$ are properly homotopic, then $\text{Ends}(f_0) = \text{Ends}(f_1)$ [12, Proposition 3.3.12 (b)].

2.4 Goldman's inductive procedure

For integers $g, b, p \geq 0$, denote the compact, connected, orientable 2-manifold of genus g with b boundary components by $S_{g,b}$. Let $S_{g,b,p}$ be the 2-manifold after removing p points from $\text{int}(S_{g,b})$. We will call $S_{0,0}, S_{0,2}, S_{0,3}, S_{1,1}, S_{1,2}$, and $S_{0,1,1}$ a disk, an annulus, a pair of pants, a one-holed torus, a two-holed torus, and a punctured disk, respectively. For brevity, $S_{g,0}$ will be denoted as S_g .

Now, let Σ be a non-compact surface. Goldman's inductive procedure [10, Section 2.6.] tells that there is an efficient exhaustion $S_1 \subseteq S_2 \subseteq \dots$ by compacta of Σ with the following properties:

- Each S_i is a smoothly embedded compact bordered sub-surface of Σ .
- For each i , the closure of each component of $\Sigma \setminus S_i$ intersects S_i in a single component of ∂S_i .
- S_1 is a disk, and $S_i \setminus \text{int}(S_{i-1})$ is either a pair of pants or an annulus or a two-holed torus for each $i \geq 2$.

Consider such an efficient exhaustion by compacta of Σ , and let $e := (V_1, V_2, \dots) \in \text{Ends}(\Sigma)$ be an end, where V_i is a component of $X \setminus S_i$. We say e is a *planar end* if V_i is embeddable in \mathbb{R}^2 for some positive integer i . An end is said to be *non-planar* if it is not planar. Notice that the set of all planar ends of Σ under the subspace topology is an open subset of $\text{Ends}(\Sigma)$. Define the *genus* of Σ as $g(\Sigma) := \sup g(S)$, where S is a compact bordered subsurface of Σ .

3 Proofs of the main theorems

3.1 Proof of Theorem 1.1

It is known [11, Page 8] that if $f: M \rightarrow N$ is a proper map between two connected, oriented, boundaryless, smooth n -manifolds of degree $d \neq 0$, then $\|N\| \cdot |d| \leq \|M\|$, where $\|\cdot\|$ denotes the Gromov simplicial volume. Also, recall that $\|S_g\| = \max\{0, -2\chi(S_g)\}$ [11, Page 9] [9, Corollary 7.5]. Thus, if $f: S_g \rightarrow S_h$ is a map of non-zero degree, then $g \geq h$ (every map from the sphere to the torus

is null-homotopic). However, the analog argument doesn't say the genus of the domain is higher than the genus of the co-domain when surfaces are of the infinite type because the simplicial volume of an infinite type surface is infinite [14, Proof of Proposition 4.2]. The following lemma states that the genus of the domain of a non-zero degree map is at least the genus of its co-domain, even if the surfaces have a finite genus but are of infinite type.

Lemma 3.1 Let $f: \Sigma' \rightarrow \Sigma$ be a proper map of non-zero degree between non-compact oriented surfaces. Then $g(\Sigma') \geq g(\Sigma)$.

Proof. If $g(\Sigma') = \infty$, then we are done. So assume $g(\Sigma') < \infty$. If possible, suppose $g(\Sigma) > g(\Sigma')$. Let h_1, h_2, \dots be a collection of pairwise-disjoint smoothly embedded copies of the one-holed torus in Σ such that $\Sigma \setminus \cup_i h_i$ is a planar surface. For $i \geq g(\Sigma') + 2$, identifying each h_i to a point in ∂h_i gives a quotient map $q: \Sigma \rightarrow \bar{\Sigma}$ from Σ to a non-compact surface $\bar{\Sigma}$. Orient $\bar{\Sigma}$. Notice that q is proper and $\deg(q) = \pm 1$ by [Theorem 2.2.1](#). Moreover, $g(\bar{\Sigma}) = g(\Sigma') + 1$. Thus, $qf: \Sigma' \rightarrow \bar{\Sigma}$ is a proper map of a non-zero degree between two finite genus non-compact surfaces. Consider the orientations on Σ'^{\dagger} and $\bar{\Sigma}^{\dagger}$ induced from Σ' and $\bar{\Sigma}$. By [Theorem 2.2.2](#) and [Theorem 2.2.1](#), the degree of $(qf)^{\dagger}: \Sigma'^{\dagger} \rightarrow \bar{\Sigma}^{\dagger}$ is the same as the degree of $qf: \Sigma' \rightarrow \bar{\Sigma}$. Thus, $(qf)^{\dagger}$ is a map of non-zero degree from the $\Sigma'^{\dagger} = S_{g(\Sigma')}$ to $\bar{\Sigma}^{\dagger} = S_{g(\Sigma')+1}$, a contradiction. \square

The following lemma states that when a map of non-zero degree exists between two oriented manifolds, the domain has more ends than the co-domain.

Lemma 3.2 Let $f: (M, \partial M) \rightarrow (N, \partial N)$ be a proper map between two connected, oriented, smooth manifolds of the same dimension. If $\deg(f) \neq 0$, then $\text{Ends}(f)$ is surjective.

Proof. Fix an efficient exhaustion $\{L_i\}$ by compacta of N . Suppose $\text{Ends}(f)$ is not surjective. Then $f^{\dagger}: M^{\dagger} \rightarrow N^{\dagger}$ is not surjective. Since M^{\dagger} is compact and N^{\dagger} is Hausdorff, the image of f^{\dagger} is closed. Thus, there exists a basic open set $W_i \cup \{(W'_1, W'_2, \dots) \in \text{Ends}(N) \mid W'_i = W_i\} \subseteq N^{\dagger} \setminus \text{im}(f^{\dagger})$, where W_i is a component of $N \setminus L_i$. But $(f^{\dagger})^{-1}(W_i) = f^{-1}(W_i) \neq \emptyset$ by [Theorem 2.2.3](#), a contradiction. \square

Let N be a connected topological n -manifold, and $p: \tilde{N} \rightarrow N$ a covering. By invariance of domain and covering space theory, \tilde{N} is a connected topological n -manifold such that p sends $\partial \tilde{N}$ onto ∂N and $\text{int}(\tilde{N})$ onto $\text{int}(N)$. If N is orientable, then \tilde{N} is also orientable because p is a local homeomorphism [7, Lemma 125.17. (a)]. If an orientation for N is given, we always consider that orientation of \tilde{N} for which p becomes orientation-preserving. The following lemma, whose proof is drawn upon ideas from [5, Theorem 3.1.], tells that if the degree of either the lifting or the map doesn't vanish, then the corresponding covering is finite-sheeted.

Lemma 3.3 Let $f: M \rightarrow N$ be a proper map between two connected, oriented, topological manifolds of the same dimension n such that $f(\partial M) \subseteq \partial N$, and let $p: \tilde{N} \rightarrow N$ be the covering corresponding to the subgroup $\text{im } \pi_1(f)$ of $\pi_1(N)$. Suppose $\tilde{f}: M \rightarrow \tilde{N}$ is a lift of f . Then, \tilde{f} is a proper map and $\tilde{f}(\partial M) \subseteq \partial \tilde{N}$. Moreover, if either $\deg(f) \neq 0$ or $\deg(\tilde{f}) \neq 0$, then p is a proper map.

To understand the various unlabeled maps and ι present in the four commutative diagrams of the proof of [Lemma 3.3](#), let's first recall the definition of singular cohomology with compact support. Consider a pair $A \subseteq X$ of topological space, and let n be a non-negative integer. Recall that the n -th singular cohomology with compact support $H_c^n(X, A; \mathbb{Z})$ is equal to the direct limit $\varinjlim H^n(X, A \cup (X \setminus K); \mathbb{Z})$, where K is a compact subset of X and various maps to define this direct system are inclusion induced maps. Hence, for a compact subset K of X , the definition of direct limit yields an obvious map $H^n(X, A \cup (X \setminus K); \mathbb{Z}) \rightarrow H_c^n(X, A; \mathbb{Z})$.

Proof of Lemma 3.3. Note that $p \circ \tilde{f} = f$ implies $\tilde{f}(\partial M) \subseteq \partial \tilde{N}$. Moreover, \tilde{f} is proper because f is proper.

Now, let either $\deg(f) \neq 0$ or $\deg(\tilde{f}) \neq 0$. We need to show that p is a proper map. On the contrary, suppose p is not proper, i.e., $p^{-1}(x)$ is infinite for some $x \in \text{int}(N)$. Define $C := f^{-1}(x) \subseteq \text{int}(M)$. So, the compact set $\tilde{f}(C)$ is contained in the discrete space $p^{-1}(x)$, i.e., we can write $\tilde{f}(C) = \{\tilde{x}_1, \dots, \tilde{x}_k\}$ for points $\tilde{x}_1, \dots, \tilde{x}_k \in p^{-1}(x) \subset \tilde{N}$. Since $p^{-1}(x)$ is infinite, we have an $\tilde{x} \in p^{-1}(x) \setminus \{\tilde{x}_1, \dots, \tilde{x}_k\}$. Thus, $\tilde{f}(M) \subseteq \tilde{N} \setminus \tilde{x}$.

Since \tilde{N} is orientable, $H^n(\tilde{N}, \tilde{N} \setminus \tilde{y}) \rightarrow H_c^n(\tilde{N}, \partial \tilde{N})$ is an isomorphism for every $y \in \text{int}(\tilde{N})$. Now, the following commutative diagram shows that $\deg(\tilde{f}) = 0$.

$$\begin{array}{ccc} H_c^n(\tilde{N}, \partial \tilde{N}) & \xrightarrow{\tilde{f}^*} & H_c^n(M, \partial M) \\ \cong \uparrow & & \uparrow \\ H^n(\tilde{N}, \tilde{N} \setminus \tilde{x}) & \xrightarrow{\tilde{f}^*} & H^n(M, M) = 0 \end{array}$$

With the contradiction method, we conclude that if $\deg(\tilde{f}) \neq 0$, then p is a proper map.

Now, to show that $\deg(f) = 0$, let's consider the outermost part of the following commutative diagram. Notice that it is sufficient to prove that the composition of the two blue arrows is zero, thereby implying $\deg(f) = 0$.

$$\begin{array}{ccc} H_c^n(N, \partial N) & \xrightarrow{f^*} & H_c^n(M, \partial M) \\ \cong \uparrow & & \uparrow \iota \\ H^n(N, N \setminus x) & \xrightarrow{f^*} & H^n(M, M \setminus C) \\ p^* \downarrow & \searrow \tilde{f}^* & \uparrow \tilde{f}^* \\ H^n(\tilde{N}, \tilde{N} \setminus p^{-1}(x)) & \longrightarrow & H^n(\tilde{N}, (\tilde{N} \setminus p^{-1}(x)) \cup \{\tilde{x}_1, \dots, \tilde{x}_k\}) \oplus \bigoplus_{\ell=1}^k H^n(\tilde{N}, \tilde{N} \setminus \tilde{x}_\ell) \end{array}$$

The following commutative diagram

$$\begin{array}{ccc} H^n(\tilde{N}, (\tilde{N} \setminus p^{-1}(x)) \cup \{\tilde{x}_1, \dots, \tilde{x}_k\}) & \xrightarrow{\tilde{f}^*} & H^n(M, M \setminus C) \\ & \searrow \tilde{f}^* & \uparrow \iota \\ & & H^n(M, M) = 0 \end{array}$$

tells that $H^n(\tilde{N}, (\tilde{N} \setminus p^{-1}(x)) \cup \{\tilde{x}_1, \dots, \tilde{x}_k\}) \xrightarrow{\tilde{f}^*} H^n(M, M \setminus C) \xrightarrow{\iota} H_c^n(M, \partial M)$ is zero map.

Also, for each $\ell = 1, \dots, k$, if we consider the following commutative diagram,

$$\begin{array}{ccc} H^n(\tilde{N}, \tilde{N} \setminus \tilde{x}_\ell) & \xrightarrow{\tilde{f}^*} & H^n(M, M \setminus C) \\ \cong \downarrow & & \downarrow \iota \\ H_c^n(\tilde{N}, \partial \tilde{N}) & \xrightarrow{\tilde{f}^*} & H_c^n(M, \partial M) \\ \cong \uparrow & & \uparrow \\ H^n(\tilde{N}, \tilde{N} \setminus \tilde{x}) & \xrightarrow{\tilde{f}^*} & H^n(M, M) = 0 \end{array}$$

then $\bigoplus_{\ell=1}^k H^n(\tilde{N}, \tilde{N} \setminus \tilde{x}_\ell) \xrightarrow{f^*} H^n(M, M \setminus C) \xrightarrow{\iota} H_c^n(M, \partial M)$ is zero map. With the contradiction method, we conclude that if $\deg(f) \neq 0$, then p is a proper map. \square

Now we are ready to give a proof of [Theorem 1.1](#).

Theorem 3.4 ([Theorem 1.1](#)) Let Σ', Σ be two non-compact oriented surfaces such that Σ' is neither the plane nor the punctured plane. Suppose $f: \Sigma' \rightarrow \Sigma$ is a π_1 -injective proper map. Then f is properly homotopic to a d -sheeted covering map $p: \Sigma' \rightarrow \Sigma$ for some positive integer d . Thus, $\deg(f) = \pm d (\neq 0)$. Moreover, Σ' is of infinite type if and only if Σ is of infinite type.

Proof. Let $p: \tilde{\Sigma} \rightarrow \Sigma$ be the covering corresponding to the subgroup $\text{im } \pi_1(f)$ of $\pi_1(\Sigma)$, and let $\tilde{f}: \Sigma' \rightarrow \tilde{\Sigma}$ be a lift of f with respect to p , i.e., $p \circ \tilde{f} = f$. Thus, $\text{im } \pi_1(p) = \text{im } \pi_1(f)$, and hence $\pi_1(\tilde{f}): \pi_1(\Sigma') \rightarrow \pi_1(\tilde{\Sigma})$ is an isomorphism because a covering map induces injection between fundamental groups. Note that if a non-compact surface has an infinite cyclic (resp. trivial) fundamental group, then it is homeomorphic to the punctured plane (resp. plane) [[3](#), [Theorem 3.1.9](#)]. Also, the properness of f implies the properness of \tilde{f} by [Lemma 3.3](#). Since non-compact surfaces are $K(\pi, 1)$ CW-complexes, by Whitehead theorem \tilde{f} is a homotopy equivalence. Thus, by [[3](#), [Theorem 1](#)], \tilde{f} is properly homotopic to a homeomorphism, which implies $\deg(\tilde{f}) = \pm 1$. Therefore, using [Lemma 3.3](#), we may assume that p is a d -sheeted covering for some positive integer d and $\tilde{\Sigma}$ is orientable. Fix an orientation of $\tilde{\Sigma}$. By [Theorem 2.2.1](#), $\deg(p) = \pm d$. Since $\deg(f) = \deg(p\tilde{f}) = (\pm d) \cdot \deg(\tilde{f}) = (\pm d) \cdot (\pm 1)$, we can conclude that $\deg(f) = \pm d (\neq 0)$, and f is properly homotopic to the d -sheeted covering map p .

Now, we need to show that Σ' is of infinite type if and only if Σ is of infinite type. At first, suppose Σ is of finite type. Therefore, $\pi_1(\Sigma)$ is a finitely generated free group and $\text{im } \pi_1(p)$ is a finite index subgroup of $\pi_1(\Sigma)$. By the Nielsen-Schreier index formula, $\pi_1(\tilde{\Sigma})$ is finitely generated. Since $\pi_1(\tilde{f})$ is an isomorphism, $\pi_1(\Sigma')$ is finitely generated. So, Σ' is of finite type. Next, suppose Σ is of infinite type. If Σ is an infinite genus surface, then [Lemma 3.1](#) tells that Σ' is an infinite type surface. On the other hand, if Σ has a finite genus, then $\text{Ends}(\Sigma)$ must be infinite, and thus by [Lemma 3.2](#), we can tell that Σ' is of infinite type. \square

3.2 Proof of [Theorem 1.3](#)

The remainder of this note is based on *surgery on a proper map between non-compact surfaces*. We will first recall a few terminologies and theorems.

Let S be a connected, orientable two-dimensional manifold with or without boundary. A *circle* on S is the image of an embedding of \mathbb{S}^1 into S . We say a circle \mathcal{C} on S is a *trivial circle* if \mathcal{C} bounds an embedded disk in S . A circle on S will be called *primitive* if it is not trivial. Note that if the image of an embedding $f: \mathbb{S}^1 \hookrightarrow S$ is a primitive circle on S , then $[f] \in \pi_1(S)$ is a primitive element [[4](#), [Theorems 1.7 and 4.2](#)]. A pairwise disjoint collection $\mathcal{A} = \{\mathcal{C}_\alpha\}$ of smoothly embedded circles on S is called a *locally finite curve system* (in short, LFCS) on S if for each compact subset K of Σ , $\mathcal{C}_\alpha \cap K = \emptyset$ for all but finitely many α . The theorem below tells that the transversal pre-image of an LFCS is again an LFCS.

Theorem 3.5 (Approximation and transversality in proper category) [[3](#), §3.2] Let $f: \Sigma' \rightarrow \Sigma$ be a proper map between non-compact surfaces, and let \mathcal{A} be an LFCS on Σ . Then f can be properly homotoped so that it becomes smooth as well as transverse to \mathcal{A} . Thus, after such a proper homotopy, for each component \mathcal{C} of \mathcal{A} , either $f^{-1}(\mathcal{C})$ is empty or a pairwise disjoint finite collection of smoothly embedded circles on Σ' ; in particular, $f^{-1}(\mathcal{A})$ is an LFCS on Σ' .

Let Σ be a non-compact surface. Suppose \mathcal{A} is an LFCS on Σ and $\{\Sigma_n\}$ is an at most countable collection of bordered sub-surfaces of Σ . We say \mathcal{A} *decomposes* Σ into bordered sub-surfaces, where *complementary components* are $\{\Sigma_n\}$ if each Σ_n is a closed subset of Σ , $\text{int}(\Sigma_n) \cap \text{int}(\Sigma_m) = \emptyset$ if $n \neq m$, $\cup_n \Sigma_n = \Sigma$, and $\cup_n \partial \Sigma_n = \cup \mathcal{A}$. For example, if $\Sigma \neq \mathbb{R}^2$, then there is an LFCS \mathcal{C} on Σ such that \mathcal{C} decomposes Σ into bordered sub-surfaces, and a complementary component of this decomposition is either a two-holed torus, a pair of pants, or a punctured disk. This follows from Goldman's inductive procedure [3, Theorem 3.1.5]. We call an LFCS \mathcal{A} on Σ a *preferred LFCS* if it satisfies either of the following:

- (i) \mathcal{A} is a finite collection of primitive circles on Σ .
- (ii) \mathcal{A} decomposes Σ into bordered sub-surfaces, and a complementary component of this decomposition is either a two-holed torus, a pair of pants, an annulus, or a punctured disk.

Now, we state our first theorem on surgery on a proper map.

Theorem 3.6 (Disk removal) [3, Theorem 3.3.5] Let $f: \Sigma' \rightarrow \Sigma$ be a smooth proper map between two non-compact surfaces, where $\Sigma' \neq \mathbb{R}^2 \neq \Sigma$. Suppose \mathcal{A} is a preferred LFCS on Σ such that f is transverse to \mathcal{A} . Then, f can be properly homotoped to remove all trivial components of the LFCS $f^{-1}(\mathcal{A})$ such that each primitive component of $f^{-1}(\mathcal{A})$ has an open neighborhood on which this proper homotopy is constant.

Remark 3.7 The only usage of the hypothesis $\Sigma' \neq \mathbb{R}^2$ to prove [Theorem 3.6](#) is to discard the possibility that there exist infinitely many components C'_1, C'_2, \dots of $f^{-1}(\mathcal{A})$ bounding the disks D'_1, D'_2, \dots , respectively such that C'_n is contained in the interior of D'_{n+1} for each n . Thus, if \mathcal{A} has only finitely many components, then the same conclusion of [Theorem 3.6](#) holds even if we consider \mathbb{R}^2 as the domain of f . However, note that the hypothesis $\Sigma \neq \mathbb{R}^2$ can never be dropped because to send every disk in Σ' bounded by a component of $f^{-1}(\mathcal{A})$ into a circle $C_\delta \subset \Sigma \setminus \cup \mathcal{A}$ using homotopy long exact sequence, C_δ must be a primitive circle parallel to a component of \mathcal{A} .

At this point, we are ready to prove the easy part of [Theorem 1.3](#).

Theorem 3.8 ([Theorem 1.3\(i\)](#)) Let f be a proper map from \mathbb{C} to a non-compact oriented surface Σ . Suppose $\deg(f) = 0$. Then, for every compact subset K of Σ , there exists a proper map g properly homotopic to f such that $\text{im}(g) \subseteq \Sigma \setminus K$.

Proof. First, assume $\Sigma = \mathbb{R}^2$. By [Theorem 2.2.2](#), properly homotope f so that the image of f misses a point $a \in \mathbb{R}^2$. Since any translation map of \mathbb{R}^2 is properly homotopic to the identity map of \mathbb{R}^2 , we may assume $a = 0$. Thus, there exists $r > 0$ such that $|f| \geq r$ because proper maps between manifolds are closed maps. Consider a compact subset K of \mathbb{R}^2 . Let n be a positive integer such that $K \subseteq \{z \in \mathbb{R}^2 : |z| \leq nr\}$. Since $1 + nt \geq 1$ for every $t \in [0, 1]$, the map $\mathbb{R}^2 \times [0, 1] \ni (z, t) \mapsto (1 + nt) \cdot f(z) \in \mathbb{R}^2$ is a proper homotopy from f to $g := (n + 1)f$. Certainly, $\text{im}(g) \cap K = \emptyset$. So, we are done when $\Sigma = \mathbb{R}^2$.

Now, suppose $\Sigma \neq \mathbb{R}^2$. Let K be a compact subset of Σ . Using Goldman's inductive procedure, find a compact bordered subsurface S of Σ such that $K \subseteq \text{int}(S)$ and each component of ∂S is a primitive circle on Σ . By [Theorem 3.5](#), we may assume f is smooth as well as transverse to ∂S . Therefore, $f^{-1}(\partial S)$ is a pairwise disjoint finite collection of smoothly embedded circles on \mathbb{R}^2 . Since $f^{-1}(\partial S)$ has finitely many components, each of which bounds a disk in \mathbb{R}^2 , by considering a procedure similar to that given in proof of [Theorem 3.6](#), we can properly homotope f to a proper

map $g: \mathbb{R}^2 \rightarrow \Sigma$ so that $g^{-1}(\partial S) = \emptyset$ (see [Remark 3.7](#)). Using continuity of g , either $\text{im}(g) \subset S$ or $\text{im}(g) \subseteq \Sigma \setminus S$. Since g is proper, the former can't happen, i.e., $\text{im}(g) \subseteq \Sigma \setminus S$, and hence $\text{im}(g) \cap K = \emptyset$. \square

Notice that, unlike the first part, in the second part of the above proof, we haven't utilized the geometric realization of the fact that $\deg(f) = 0$. This is because if f is a proper map from \mathbb{R}^2 to a non-compact oriented surface Σ of non-zero degree, then Σ must be \mathbb{R}^2 , by [Theorem 3.9](#) below. Note that [Theorem 3.9](#) appears as a corollary [[5](#), Corollary 3.4.] of a theorem [[5](#), Theorem 3.1.] by Epstein, which states that the absolute degree is the same as the (algebraic) degree up to sign when manifolds are orientable. Epstein credits Olum [[22](#)] for the proof of [[5](#), Corollary 3.4.]. Since [[5](#), Corollary 3.4.] comes without proof and follows from a general theory, we will give a proof of [Theorem 3.9](#) using [Lemma 3.3](#).

Theorem 3.9 Let $f: M \rightarrow N$ be a proper map between two connected, oriented, topological manifolds of the same dimension such that $f(\partial M) \subseteq \partial N$. If $\deg(f) \neq 0$, then the index $[\pi_1(N) : \text{im } \pi_1(f)]$ divides $\deg(f)$. In particular, if $\deg(f) = \pm 1$, then $\pi_1(f)$ is surjective.

Proof. Let $p: \tilde{N} \rightarrow N$ be the covering corresponding to the subgroup $\text{im } \pi_1(f)$ of $\pi_1(N)$. Thus, $\text{im } \pi_1(p) = \text{im } \pi_1(f)$. Consider a lift $\tilde{f}: \tilde{M} \rightarrow \tilde{N}$ of f with respect to the covering p , i.e., $f = p \circ \tilde{f}$. By [Lemma 3.3](#), \tilde{N} is a connected, orientable manifold such that $\dim \tilde{N} = \dim N$, $p(\partial \tilde{N}) = \partial N$, and $\tilde{f}(\partial \tilde{M}) \subseteq \partial \tilde{N}$. Furthermore, [Lemma 3.3](#) tells that p and \tilde{f} both are proper maps because $\deg(f) \neq 0$. Therefore, p is a d -sheeted covering map for some positive integer d . By covering space theory, $d = [\pi_1(N) : \text{im } \pi_1(p)] = [\pi_1(N) : \text{im } \pi_1(f)]$. Fix an orientation for \tilde{N} . Then, $\deg(p) = \pm d$. Therefore, $\deg(f) = \deg(p \circ \tilde{f}) = \deg(p) \deg(\tilde{f})$, i.e., $[\pi_1(N) : \text{im } \pi_1(f)]$ divides $\deg(f)$. \square

Another surgery technique is required to prove the remaining portion of [Theorem 1.3](#).

Theorem 3.10 (Level-preserving homotopy from degree-one map to homeomorphism) [[3](#), Theorem 3.4.3] Let $f: \Sigma' \rightarrow \Sigma$ be a smooth proper map between two non-compact surfaces, where $\Sigma' \neq \mathbb{R}^2 \neq \Sigma$. Suppose \mathcal{A} is a preferred LFCS on Σ such that f is transverse to \mathcal{A} . If f is a homotopy equivalence, then f can be properly homotoped to remove all trivial components of the $f^{-1}(\mathcal{A})$ as well as to map each primitive component of $f^{-1}(\mathcal{A})$ homeomorphically onto a component of \mathcal{A} . Moreover, after this proper homotopy, if C'_p and C are components of $f^{-1}(\mathcal{A})$ and \mathcal{A} , respectively, such that $f|_{C'_p} \rightarrow C$ is a homeomorphism, then C'_p (resp. C) has two one-sided tubular neighborhoods \mathcal{U}' and \mathcal{V}' (resp. \mathcal{U} and \mathcal{V}) with some identifications $(\mathcal{U}', C'_p) \equiv (C'_p \times [1, 2], C'_p \times 2) \equiv (\mathcal{V}', C'_p)$ (resp. $(\mathcal{U}, C) \equiv (C \times [1, 2], C \times 2) \equiv (\mathcal{V}, C)$) such that the following hold:

- (i) $\mathcal{U}' \cup \mathcal{V}'$ is a (two-sided) tubular neighborhood of C'_p ;
- (ii) $f|_{\mathcal{U}'} \rightarrow \mathcal{U}$ and $f|_{\mathcal{V}'} \rightarrow \mathcal{V}$ are homeomorphisms described by $C'_p \times [1, 2] \ni (z, t) \mapsto (f(z), t) \in C \times [1, 2]$.

Remark 3.11 Instead of assuming that f is a homotopy equivalence in [Theorem 3.10](#), if we assume that for a primitive component C'_p of $f^{-1}(\mathcal{A})$, there exists a component C of \mathcal{A} such that $f(C'_p) \subset C$ and the restriction $f|_{C'_p} \rightarrow C$ has degree $n \neq 0$, then f can be properly homotoped so that $f|_{C'_p} \rightarrow C$ becomes an n -sheeted covering map. Moreover, after this proper homotopy, f maps each small enough one-sided tubular neighborhood $C'_p \times [1, 2]$, $C'_p \times 2 \equiv C'_p$ of C'_p (on either side of C'_p) onto a small enough one-sided tubular neighborhood $C \times [1, 2]$, $C \times 2 \equiv C$ of C in the following manner: $C'_p \times [1, 2] \ni (z, t) \mapsto (f(z), t) \in C \times [1, 2]$.

Remark 3.12 Instead of assuming that f is a homotopy equivalence in [Theorem 3.10](#), if we assume that f is π_1 -injective, then f can be properly homotoped to remove all trivial components from $f^{-1}(\mathcal{A})$ as well as to send each component \mathcal{C}'_p of $f^{-1}(\mathcal{A})$ onto a component \mathcal{C} of \mathcal{A} such that the restriction $f|_{\mathcal{C}'_p} \rightarrow \mathcal{C}$ is a covering map. Moreover, after this proper homotopy, f maps each small enough one-sided tubular neighborhood $\mathcal{C}'_p \times [1, 2]$, $\mathcal{C}'_p \times 2 \equiv \mathcal{C}'_p$ of \mathcal{C}'_p (on either side of \mathcal{C}'_p) onto a small enough one-sided tubular neighborhood $\mathcal{C} \times [1, 2]$, $\mathcal{C} \times 2 \equiv \mathcal{C}$ of \mathcal{C} in the following manner: $\mathcal{C}'_p \times [1, 2] \ni (z, t) \mapsto (f(z), t) \in \mathcal{C} \times [1, 2]$.

To prove [Theorem 1.3\(ii\)](#), we also need the following lemmas, whose proofs are straightforward.

Lemma 3.13 If $\theta: \mathbb{S}^1 \times [0, 1] \rightarrow \mathbb{S}^1 \times [0, 1]$ is a map which sends $\mathbb{S}^1 \times \{0, 1\}$ into $\mathbb{S}^1 \times 0$, then φ can be homotoped rel. $\mathbb{S}^1 \times \{0, 1\}$ to send $\mathbb{S}^1 \times [0, 1]$ into $\mathbb{S}^1 \times 0$.

Lemma 3.14 Let $\varphi: \mathbb{S}^1 \times [1, 3] \rightarrow \mathbb{S}^1 \times [1, 2]$ be a map such that $\varphi(\mathbb{S}^1 \times r) \subseteq \mathbb{S}^1 \times r$ for $1 \leq r \leq 2$ and $\varphi(\mathbb{S}^1 \times [2, 3]) \subseteq \mathbb{S}^1 \times 2$. Then φ can be homotoped rel. $\mathbb{S}^1 \times \{1, 3\}$ to satisfy $\varphi^{-1}(\mathbb{S}^1 \times 2) = \mathbb{S}^1 \times 3$.

Theorem 3.15 ([Theorem 1.3\(ii\)](#)) Let f be a proper map from \mathbb{C} to a non-compact oriented surface Σ . Suppose $n := \deg(f)$ is non-zero. Then, $\Sigma = \mathbb{C}$, and f is properly homotopic to the branched covering $\mathbb{C} \ni z \mapsto z^n \in \mathbb{C}$ if $n > 0$, and to the branched covering $\mathbb{C} \ni z \mapsto \bar{z}^{-n} \in \mathbb{C}$ if $n < 0$.

Proof. By [Theorem 3.9](#), $\Sigma = \mathbb{C}$. At first, suppose $n > 0$. Let \mathcal{A} be the collection of all circles in \mathbb{C} centered at 0 with integer radii. Then \mathcal{A} is an LFCS on \mathbb{C} . By [Theorem 3.5](#), we may assume f is a smooth proper map of degree $n > 0$ as well as transverse to \mathcal{A} . Therefore, $f^{-1}(\mathcal{A})$ is an LFCS by [Theorem 3.5](#). Now, [Theorem 2.2.2](#) gives a proper map g properly homotopic to f with the following properties: the map g equals f outside a compact subset K of the form $\{z \in \mathbb{C} : |z| \leq r\}$ for some $r > 0$, and there exists a disk D in \mathbb{C} such that $g^{-1}(D)$ is the union of pairwise disjoint disks D_1, \dots, D_n in \mathbb{C} such that $g|_{D_j} \rightarrow D$ is an orientation-preserving homeomorphism for each $j = 1, \dots, n$. Without loss of generality, we may assume K contains $\cup_{j=1}^n D_j$. Consider the bordered surfaces $\mathbf{S}' := \mathbb{C} \setminus \cup_{j=1}^n \text{int}(D_k)$ and $\mathbf{S} := \mathbb{C} \setminus \text{int}(D)$. Notice that $g(\mathbf{S}') \subseteq \mathbf{S}$ and g sends each component of $\partial \mathbf{S}' = g^{-1}(\partial \mathbf{S})$ homeomorphically onto $\partial \mathbf{S}$. Denote the restriction map $g|_{\mathbf{S}'} \rightarrow \mathbf{S}$ by g_{res} . Since \mathcal{A} is an LFCS, there exists a component \mathcal{C} of \mathcal{A} such that $\mathcal{C} \cap (f(K) \cup g(K)) = \emptyset$ and the interior of \mathcal{C} (in \mathbb{C}) contains D . Since $f = g$ on $\mathbb{C} \setminus K$, we can say that $f^{-1}(\mathcal{C}) = g^{-1}(\mathcal{C}) = g_{\text{res}}^{-1}(\mathcal{C})$ doesn't intersect with K , g_{res} is smooth outside the compact subset $\mathbf{K}' := \mathbf{S}' \cap K$ of \mathbf{S}' , and g_{res} is transverse to \mathcal{C} . In particular, each component of $g_{\text{res}}^{-1}(\mathcal{C})$ is either a trivial circle or primitive circle on \mathbf{S}' lying in $\mathbf{S}' \setminus \mathbf{K}' = \mathbb{C} \setminus K = \{z \in \mathbb{C} : |z| > r\}$.

Now, we aim to properly homotope $g_{\text{res}}: \mathbf{S}' \rightarrow \mathbf{S}$ to a proper map $\widetilde{g_{\text{res}}}: \mathbf{S}' \rightarrow \mathbf{S}$ such that $\mathcal{C}' := \widetilde{g_{\text{res}}}^{-1}(\mathcal{C})$ is a single circle in $\text{int}(\mathbf{S}')$ and $\widetilde{g_{\text{res}}}|_{\mathcal{C}'} \rightarrow \mathcal{C}$ is an n -sheeted covering map, but here we want every proper homotopy to be relative to $\partial \mathbf{S}'$. So, here are the steps.

At first, applying [Theorem 2.2.2](#) on g_{res} , and then using [Theorem 2.2.1](#), we can tell $\deg(g_{\text{res}}) = \deg(g)$, i.e., in particular, $\deg(g_{\text{res}}) \neq 0$. Recall that a proper homotopy relative to the boundary doesn't change the degree. Therefore, g_{res} remains surjective after any proper homotopy rel. $\partial \mathbf{S}'$; see [Theorem 2.2.3](#).

Since \mathcal{C} is a primitive circle on \mathbf{S} , using a trick similar to the one given in the proof of [Theorem 3.6](#), after a proper homotopy \mathcal{H}_1 , we may assume each component of the non-empty set $g_{\text{res}}^{-1}(\mathcal{C})$ is a primitive circle on \mathbf{S}' . Note that the homotopy that appears in the proof of [Theorem 3.6](#) is, in fact, relative to the complement of every small enough neighborhood of the union of all disks bounded by trivial circles. Thus, we may assume \mathcal{H}_1 is relative to $\partial \mathbf{S}'$.

Therefore, for some positive integer m , we can write $g_{\text{res}}^{-1}(\mathcal{C}) = \mathcal{C}'_1 \sqcup \cdots \sqcup \mathcal{C}'_m$ (see Figure 1 for labeling of these circles), where each \mathcal{C}'_i is a primitive circle on \mathbf{S}' . Thus $g_{\text{res}}^{-1}(\mathcal{C})$ decomposes \mathbf{S}' into a copy of $S_{0,n+1}$, $(m-1)$ -copies $\mathcal{A}'_1, \dots, \mathcal{A}'_{m-1}$ of $S_{0,2}$ (see Figure 1 for labeling of these annuli), and a copy of $S_{0,1,1}$. Similarly, \mathcal{C} decomposes \mathbf{S} into a copy of $S_{0,2}$ and a copy of $S_{0,1,1}$.




Fig. 1: The description of $g_{\text{res}}: \mathbf{S}' \rightarrow \mathbf{S}$ after removing trivial components from $g_{\text{res}}^{-1}(\mathcal{C})$.

Restricting g_{res} on those copies, we get a map $\xi: S_{0,n+1} \rightarrow S_{0,2}$ such that $\xi|_{\partial D_j} \rightarrow \partial D$ is an orientation-preserving homeomorphism for each $j = 1, \dots, n$ and ξ sends the other boundary component $\partial S_{0,n+1} \setminus \cup_{j=1}^n \partial D_j$ into \mathcal{C} . The naturality of the homology long exact sequence and [13, Exercise 31 of Section 3.3] give the following commutative diagram

$$\begin{array}{ccc}
 H_2(S_{0,n+1}, \partial S_{0,n+1}) \cong \mathbb{Z} & \xrightarrow{1 \mapsto \bigoplus^{n+1} 1} & \bigoplus^{n+1} \mathbb{Z} \cong H_1(\partial S_{0,n+1}) \\
 \times \deg(\xi) \downarrow & & \downarrow \bigoplus^{n+1} 1 \mapsto n \oplus \ell \\
 H_2(S_{0,2}, \partial S_{0,2}) \cong \mathbb{Z} & \xrightarrow{1 \mapsto \bigoplus^2 1} & \bigoplus^2 \mathbb{Z} \cong H_1(\partial S_{0,2})
 \end{array}$$

where $\ell := \deg(\xi|_{\partial S_{0,n+1} \setminus \cup_{j=1}^n \partial D_j} \rightarrow \mathcal{C})$. The commutativity of the diagram tells $\deg(\xi) = n$ and $\ell = n$. Since any two components of $g_{\text{res}}^{-1}(\mathcal{C})$ co-bound an annulus in \mathbf{S}' and any two homotopic maps $\mathbb{S}^1 \rightarrow \mathbb{S}^1$ have the same degree, $\deg(g_{\text{res}}|_{\mathcal{C}'_i} \rightarrow \mathcal{C}) = \pm n$ for each $i = 1, \dots, m$ (the minus sign comes because two boundary components of an oriented annulus are oppositely oriented). Thus, using a trick similar to the one mentioned in the proof of Theorem 3.10, together with Remark 3.11, after a proper homotopy \mathcal{H}_2 , we may assume that $g_{\text{res}}|_{\mathcal{C}'_i} \rightarrow \mathcal{C}$ is an n -sheeted covering map for

each $i = 1, \dots, m$. Since the homotopy mentioned in the proof of [Theorem 3.10](#) is relative to the complement of a small neighborhood of $g_{\text{res}}^{-1}(\mathcal{C}) = \mathcal{C}'_1 \sqcup \dots \sqcup \mathcal{C}'_m$, one can ensure that \mathcal{H}_2 is relative to $\partial \mathbf{S}'$.

Now, we remove the components $\mathcal{C}'_2, \dots, \mathcal{C}'_m$ from $g_{\text{res}}^{-1}(\mathcal{C})$. First, consider the annulus \mathcal{A}'_1 . By continuity, $g_{\text{res}}(\mathcal{A}'_1)$ will be one of two sides of \mathcal{C} in \mathbf{S} , i.e., the compact set $g_{\text{res}}(\mathcal{A}'_1)$ is contained in a one-sided tubular neighborhood of \mathcal{C} in \mathbf{S} . Notice that if \mathbf{S}'' is a compact bordered subsurface of \mathbf{S}' such that $\mathbf{S}'' \cap \partial \mathbf{S}' = \emptyset$, then a homotopy of $g_{\text{res}}|_{\mathbf{S}''}$ rel. $\partial \mathbf{S}''$ can be extended to a proper homotopy of g_{res} rel. $\mathbf{S}' \setminus \text{int}(\mathbf{S}'') (\supset \partial \mathbf{S}')$. Thus, after a homotopy of $g_{\text{res}}|_{\mathcal{A}'_1}$ rel. $\partial \mathcal{A}'_1$ (see [Lemma 3.13](#)), we may assume $g_{\text{res}}(\mathcal{A}'_1) = \mathcal{C}$. Applying this argument to each of $\mathcal{A}'_2, \dots, \mathcal{A}'_{m-1}$, after a homotopy of $g_{\text{res}}|_{\mathcal{A}'_1 \cup \dots \cup \mathcal{A}'_{m-1}}$ rel. $\partial(\mathcal{A}'_1 \cup \dots \cup \mathcal{A}'_{m-1})$, we may assume $g_{\text{res}}(\mathcal{A}'_1 \cup \dots \cup \mathcal{A}'_{m-1}) = \mathcal{C}$. Further, the part 2. of [Theorem 3.10](#) tells that there exists a one-sided tubular neighborhood $\mathcal{C}'_m \times [1, 2] \equiv \mathcal{V}' \subset S_{0,1,1}$ of $\mathcal{C}'_m \equiv \mathcal{C}'_m \times 2$ and a one-sided tubular neighborhood $\mathcal{C} \times [1, 2] \equiv \mathcal{V}$ of $\mathcal{C} \equiv \mathcal{C} \times 2$ such that $g_{\text{res}}(z, r) = g_{\text{res}}(z) \times r$ for all $(z, r) \in \mathcal{C}'_m \times [1, 2]$. Applying [Lemma 3.14](#), after a homotopy of $g_{\text{res}}|_{(\mathcal{A}'_1 \cup \dots \cup \mathcal{A}'_{m-1}) \cup \mathcal{V}'}$ rel. $\mathcal{C}'_1 \sqcup (\mathcal{C}'_m \times 1)$, we have $g_{\text{res}}^{-1}(\mathcal{C}) = \mathcal{C}'_1$ and $g_{\text{res}}|_{\mathcal{C}'_1} \rightarrow \mathcal{C}$ is an n -sheeted covering map.

Since every proper homotopy of g_{res} has been done rel. $\partial \mathbf{S}'$, pasting $g|_{g^{-1}(D)} \rightarrow D$ with all those proper homotopies, we can say that $g: \mathbb{C} \rightarrow \mathbb{C}$ can be properly homotoped so that $g|_{g^{-1}(\mathcal{C})} \rightarrow \mathcal{C}$ becomes an orientation-preserving n -sheeted covering map from a (single) circle onto a circle. Since an n -sheeted orientation-preserving covering $\mathbb{S}^1 \rightarrow \mathbb{S}^1$ is of the form $\mathbb{S}^1 \ni z \mapsto h(z^n) \in \mathbb{S}^1$ for some orientation-preserving self-homomorphism $h: \mathbb{S}^1 \rightarrow \mathbb{S}^1$, by an application of [Theorem 3.1](#), after a proper homotopy, we may assume $g|_{g^{-1}(\mathbb{S}^1)} = \mathbb{S}^1 \ni z \mapsto z^n \in \mathbb{S}^1$. Now, applying [Proposition 3.16](#) and [Proposition 3.17](#), we can conclude that g , and consequently f as well, is properly homotopic to $\mathbb{C} \ni z \mapsto z^n \in \mathbb{C}$. This completes the proof of the case when $\deg(f) = n > 0$.

Now, assume $\deg(f) = n < 0$. Since the complex conjugation is an orientation-reversing self-homeomorphism of \mathbb{C} , the map $\bar{f}: \mathbb{C} \ni z \mapsto \bar{f}(\bar{z}) \in \mathbb{C}$ is orientation-preserving of degree $-n$. Thus, \bar{f} is properly homotopic to $\mathbb{C} \ni z \mapsto z^{-n} \in \mathbb{C}$ by the previous case. Therefore, f is properly homotopic to $\mathbb{C} \ni z \mapsto \bar{z}^{-n} \in \mathbb{C}$. \square

The famous Annulus Theorem says that if two orientation-preserving locally flat embeddings φ and ψ of the closed unit ball $\mathbb{B}^n := \{x \in \mathbb{R}^n : |x| \leq 1\}$ into \mathbb{R}^n satisfies $\varphi(\mathbb{B}^n) \subset \psi(\text{int}(\mathbb{B}^n))$, then $\psi(\mathbb{B}^n) \setminus \varphi(\text{int}(\mathbb{B}^n))$ is homeomorphic to $\mathbb{S}^{n-1} \times [0, 1]$. The annulus theorem was proved in dimension two by Radó [27], in dimension three by Moise [20, Theorem 1], in dimension four by Quinn [26, Page 506], and in dimensions at least five by Kirby [18, Page 576]. It is well known that the annulus theorem establishes the following theorem, which is the main tool for showing that the connected sum operation of topological n -manifolds is independent of the choice of embeddings of the n -balls. The analog of [Theorem 3.1](#) for smooth manifolds is called the Cerf-Palais' disk theorem [23, Theorem 5.5.] [24, Theorem B].

Theorem 3.1 [8, Proof of Theorem 4.12.] [7, Theorem 153.10.] [2, Theorem 1.3] Let M be an oriented, connected n -dimensional topological manifold without boundary. If $\varphi, \psi: \mathbb{B}^n \hookrightarrow M$ are two locally flat orientation-preserving embeddings of the closed unit balls into M , then there exists a homotopy $\mathcal{H}: M \times [0, 1] \rightarrow M$ through homeomorphisms such that $\mathcal{H}(-, 0) = \text{id}_M$ and $\mathcal{H}(-, 1) \circ \varphi = \psi$. In particular, \mathcal{H} is a proper homotopy.

In the proof of [Theorem 1.3\(ii\)](#), we have also used the following well-known facts, for which proof will be provided by modifying the proof of the Alexander trick [6, Lemma 2.1].

Proposition 3.16 Let $f: \mathbb{B}^2 \rightarrow \mathbb{B}^2$ be a map such that $f^{-1}(\mathbb{S}^1) = \mathbb{S}^1$. Then f can be homotoped rel. \mathbb{S}^1 to a map $g: \mathbb{B}^2 \rightarrow \mathbb{B}^2$ such that $g(0) = 0$ and $g(z) = |z| \cdot f\left(\frac{z}{|z|}\right)$ if $z \neq 0$.

Proof. Consider $H: \mathbb{B}^2 \times [0, 1] \rightarrow \mathbb{B}^2$ defined as follows

$$H(z, t) := \begin{cases} (1-t) \cdot f\left(\frac{z}{1-t}\right) & \text{if } 0 \leq |z| < 1-t, \\ |z| \cdot f\left(\frac{z}{|z|}\right) & \text{if } 1-t \leq |z| \leq 1 \text{ and } (z, t) \neq (0, 1), \\ 0 & \text{if } (z, t) = (0, 1). \end{cases}$$

Define $g := H(-, 1)$. □

Proposition 3.17 Let $\mathbb{B}_*^2 := \{z \in \mathbb{C} : 0 < |z| \leq 1\}$. Suppose $f: \mathbb{B}_*^2 \rightarrow \mathbb{B}_*^2$ be a proper map such that $f^{-1}(\mathbb{S}^1) = \mathbb{S}^1$. Then f can be properly homotoped rel. \mathbb{S}^1 to a proper map $g: \mathbb{B}_*^2 \rightarrow \mathbb{B}_*^2$ such that $g(z) = |z| \cdot f\left(\frac{z}{|z|}\right)$ for each $z \in \mathbb{B}_*^2$.

Proof. Define $\mathcal{H}: \mathbb{B}_*^2 \times [0, 1] \rightarrow \mathbb{B}_*^2$ by

$$\mathcal{H}(z, t) := \begin{cases} (1-t) \cdot f\left(\frac{z}{1-t}\right) & \text{if } 0 < |z| \leq 1-t, \\ |z| \cdot f\left(\frac{z}{|z|}\right) & \text{if } 1-t < |z| \leq 1. \end{cases}$$

Define $g := \mathcal{H}(-, 1)$. Now, we prove \mathcal{H} is a proper map. So let $\{(z_n, t_n)\}$ is a sequence in $\mathbb{B}_*^2 \times [0, 1]$ with $z_n \rightarrow 0$. We need to show that $\mathcal{H}(z_n, t_n) \rightarrow 0$. Define $N_1 := \{n \in \mathbb{N} : 1 - t_n < |z_n|\}$ and $N_2 := \{n \in \mathbb{N} : |z_n| \leq 1 - t_n\}$. Then $\mathbb{N} = N_1 \cup N_2$. Therefore, it is enough to show $\{\mathcal{H}(z_n, t_n) : n \in N_k\} \rightarrow 0$ whenever N_k is infinite for each $k = 1, 2$. If N_1 is infinite, then $\{\mathcal{H}(z_n, t_n) : n \in N_k\} \rightarrow 0$ by the definition of \mathcal{H} .

Next, assume N_2 is infinite. To prove $\{\mathcal{H}(z_n, t_n) : n \in N_2\} \rightarrow 0$, pick any $\varepsilon > 0$, and let $N_{2\varepsilon} := \{n \in N_2 : 1 - t_n < \varepsilon\}$. By the definition, $|\mathcal{H}(z_n, t_n)| \leq (1 - t_n) < \varepsilon$ for all $n \in N_{2\varepsilon}$. Moreover, if $N_2 \setminus N_{2\varepsilon}$ is infinite, then $\{z_n/(1 - t_n) : n \in N_2 \setminus N_{2\varepsilon}\} \rightarrow 0$, which implies $\{f(z_n/(1 - t_n)) : n \in N_2 \setminus N_{2\varepsilon}\} \rightarrow 0$ (as f is proper), and thus $|\mathcal{H}(z_n, t_n)| \leq |f(z_n/(1 - t_n))| < \varepsilon$ for all but finitely many $n \in N_2 \setminus N_{2\varepsilon}$. Therefore, $|\mathcal{H}(z_n, t_n)| < \varepsilon$ for all but finitely many $n \in N_2$. So, we are done. □

3.3 Proof of Theorem 1.2

Now, we will prove Theorem 1.2. At first, a couple of observations. Let Σ be a non-compact surface. We call a bordered subsurface S of Σ *essential* if the inclusion-induced map $\pi_1(S) \rightarrow \pi_1(\Sigma)$ is injective. Note that an end $e = (V_1, V_2, \dots)$ of Σ is an isolated planar end if and only if there exists an integer n_0 such that \overline{V}_n is a punctured disk for all $n \geq n_0$. Thus, a non-compact surface Σ has an isolated planar end if and only if Σ contains an essential punctured disk.

Lemma 3.18 Let f be a π_1 -injective proper map from \mathbb{C}^* to a non-compact oriented surface Σ . Suppose $\deg(f) = 0$. Then Σ contains an essential punctured disk D_* such that f can be properly homotoped so that $f(\mathbb{C}^*) \subseteq D_*$.

Proof. Notice that $\Sigma \neq \mathbb{R}^2$ because $\pi_1(f)$ is injective. So, there exists an LFCS \mathcal{A} on Σ such that \mathcal{A} decomposes Σ into bordered sub-surfaces, and a complementary component of this decomposition is homeomorphic to either $S_{1,1}$, $S_{0,3}$, or $S_{0,1,1}$. By Theorem 3.5, we may assume f is smooth as well

as transverse to \mathcal{A} . Therefore, $f^{-1}(\mathcal{A})$ is an LFCS on \mathbb{C}^* by [Theorem 3.5](#). Since each component of \mathcal{A} is a primitive circle on Σ , and f is π_1 -injective, by [Remark 3.12](#), after a proper homotopy, we may further assume that each component of $f^{-1}(\mathcal{A})$, if any, is a primitive circle on \mathbb{C}^* , and f maps for every component of $f^{-1}(\mathcal{A})$ onto a component of \mathcal{A} by a covering map. Moreover, any two primitive circles on \mathbb{C}^* co-bound an annulus in \mathbb{C}^* , but no two distinct components of \mathcal{A} co-bound an annulus in Σ together imply either $f^{-1}(\mathcal{A}) = \emptyset$ or there exists a component \mathcal{C} of \mathcal{A} such that $f^{-1}(\mathcal{A}) = f^{-1}(\mathcal{C}) \neq \emptyset$.

At first, suppose $f^{-1}(\mathcal{A}) = \emptyset$. Then $f(\mathbb{C}^*)$, being non-compact, must be contained in a punctured disk D_* that appears as a complementary component of the decomposition of Σ by \mathcal{A} . Hence, the proof for this case is complete.

From now onwards we will assume $f^{-1}(\mathcal{A}) = f^{-1}(\mathcal{C}) \neq \emptyset$. If $f^{-1}(\mathcal{C})$ has more than one component, then we want to properly homotope f so that $f^{-1}(\mathcal{C})$ becomes a single circle and $f|_{f^{-1}(\mathcal{C})} \rightarrow \mathcal{C}$ is a covering map. The main tool for all those proper homotopies is the observation that if S' is a compact bordered subsurface of \mathbb{C}^* , then a homotopy of $f|_{S'}$ rel. $\partial S'$ can be extended to a proper homotopy of f .

Notice that any two distinct components of $f^{-1}(\mathcal{C})$, being primitive circles on \mathbb{C}^* , co-bound an annulus in \mathbb{C}^* . So, let \mathcal{A}' be an annulus co-bounded by two components of $f^{-1}(\mathcal{C})$. The continuity of f tells that in the decomposition of Σ by \mathcal{A} , $f(\mathcal{A}')$ lies within one of the two complementary components that share \mathcal{C} as a common boundary. Thus, we can always find a compact bordered subsurface S of Σ such that $f|_{\mathcal{A}'} \rightarrow S$. Since $f|_{\mathcal{A}'} \rightarrow S$ is π_1 -injective and $f|\partial\mathcal{A}' \rightarrow \mathcal{C}$ is a local homeomorphism, after a homotopy of $f|_{\mathcal{A}'} \rightarrow S$ rel. $\partial\mathcal{A}'$, we may assume $f(\mathcal{A}') \subseteq \mathcal{C}$ [[29](#), Lemma 1.4.3]. Let $\mathcal{A}'_{\text{out}}$ denote the union of all annuli co-bounded by two components of $f^{-1}(\mathcal{C})$. Then, after a homotopy of $f|_{\mathcal{A}'_{\text{out}}} \rightarrow \Sigma$ rel. $\partial\mathcal{A}'_{\text{out}}$, we may assume $f(\mathcal{A}'_{\text{out}}) \subseteq \mathcal{C}$. Denote the boundary components of $\mathcal{A}'_{\text{out}}$ by \mathcal{C}' and \mathcal{C}'' . Now, [Remark 3.12](#) tells that there exists a one-sided tubular neighborhood $\mathcal{C}' \times [1, 2] \equiv \mathcal{V}'$ of $\mathcal{C}' \equiv \mathcal{C}' \times 2$ and a one-sided tubular neighborhood $\mathcal{C} \times [1, 2] \equiv \mathcal{V}$ of $\mathcal{C} \equiv \mathcal{C} \times 2$ such that $f(z, r) = f(z) \times r$ for all $(z, r) \in \mathcal{C}' \times [1, 2]$. Applying [Lemma 3.14](#), after a homotopy of $f|_{\mathcal{A}'_{\text{out}} \cup \mathcal{V}'}$ rel. $\mathcal{C}'' \sqcup (\mathcal{C}' \times 1)$, we can assume that $f^{-1}(\mathcal{C}) = \mathcal{C}''$ and $f|_{\mathcal{C}''} \rightarrow \mathcal{C}$ is an n -sheeted covering map.

At this point, two cases arise depending on whether Σ is equal to \mathbb{C}^* or not. Let's begin by assuming $\Sigma \neq \mathbb{C}^*$. Thus, if S_1 and S_2 are two distinct, complementary components of the decomposition of Σ by \mathcal{A} such that S_1 is a punctured disk and $\partial S_1 \cap \partial S_2 \neq \emptyset$, then S_2 must be compact. Since f is proper, $f(\mathbb{C}^*)$ is non-compact. Now, upon considering the decomposition of Σ by \mathcal{A} , we can tell that $f(\mathbb{C}^*)$ is contained in one of the two complementary components that share \mathcal{C} as a common boundary. Hence, there exists a punctured disk D_* appears as a complementary component of the decomposition of Σ by \mathcal{A} such that $\partial D_* = \mathcal{C}$ and $f(\mathbb{C}^*) \subseteq D_*$.

Next, assume $\Sigma = \mathbb{C}^*$, and let $D_*^{(1)}$ and $D_*^{(2)}$ be the punctured disks appear as complementary components of the decomposition of \mathbb{C}^* by \mathcal{C} . Note that in this case, $f(\mathbb{C}^*)$ cannot be equal to \mathbb{C}^* ; otherwise, applying [Proposition 3.17](#) on each $f|_{f^{-1}(D_*^{(1)})} \rightarrow D_*^{(1)}$ and $f|_{f^{-1}(D_*^{(2)})} \rightarrow D_*^{(2)}$, would yield $\deg(f) = \pm n \neq 0$. So, $f(\mathbb{C}^*)$ must be contained in one of $D_*^{(1)}$ or $D_*^{(2)}$. So, we are done. \square

Notice that, except in the last paragraph of the previous proof, we haven't utilized the fact that $\deg(f) = 0$. This is because of the following proposition.

Proposition 3.19 Let f be a π_1 -injective proper map from \mathbb{C}^* to a non-compact oriented surface Σ such that $\deg(f) \neq 0$. Then Σ must be to \mathbb{C}^* .

Proof. Since $\pi_1(f)$ is injective, $\Sigma \neq \mathbb{R}^2$. Now, note that the fundamental group of a non-compact surface other than the plane and the punctured plane is a free group of rank at least two. Thus, if $\Sigma \neq \mathbb{C}^*$, then the index $[\pi_1(\Sigma) : \text{im } \pi_1(f)]$ must be finite by [Theorem 3.9](#). Thus, by the Nielsen–Schreier index formula, $\text{im } \pi_1(f)$ is a free group of rank at least 2, a contradiction. \square

Now, we are ready to prove the first part of [Theorem 1.2](#).

Theorem 3.20 ([Theorem 1.2\(i\)](#)) Let f be a π_1 -injective proper map from $\mathbb{S}^1 \times \mathbb{R}$ to a non-compact oriented surface Σ . Suppose $\deg(f) = 0$. Then there exists a π_1 -injective, proper embedding $\iota: \mathbb{S}^1 \times [0, \infty) \hookrightarrow \Sigma$, along with a non-zero integer d , such that after a proper homotopy, f can be described by the proper map $\mathbb{S}^1 \times \mathbb{R} \ni (z, t) \mapsto \iota(z^d, |t|) \in \Sigma$. Thus, Σ has an isolated planar end, and given any compact subset K of Σ , there exists a proper map g properly homotopic to f such that $\text{im}(g) \subseteq \Sigma \setminus K$.

Proof. Notice that $\Sigma \neq \mathbb{R}^2$ because f is π_1 -injective. By [Lemma 3.18](#), we may assume that $f(\mathbb{S}^1 \times \mathbb{R}) \subseteq D_*$ for some essential punctured disk $D_* \subset \Sigma$. Let e be the (isolated planar) end of Σ determined by D_* . Consider a locally-finite, pairwise-disjoint collection $\mathcal{A} := \{C_i : i = 1, 2, \dots\}$ of smoothly embedded primitive circles on Σ such that each C_i is contained in $\text{int}(D_*)$. Observe that if a proper map g is properly homotopic to f , then $g^{-1}(\mathcal{A}) \neq \emptyset$ because $\text{Ends}(g) = \text{Ends}(f)$ sends both elements of $\text{Ends}(\mathbb{S}^1 \times \mathbb{R})$ to e . By [Theorem 3.5](#), we may assume f is smooth as well as transverse to \mathcal{A} . Since each component of \mathcal{A} is a primitive circle on Σ , and f is π_1 -injective, by [Remark 3.12](#), after a proper homotopy, we may further assume that each component of the non-empty LFCS $f^{-1}(\mathcal{A})$ is a primitive circle on $\mathbb{S}^1 \times \mathbb{R}$ and f maps for every component of $f^{-1}(\mathcal{A})$ onto a component of \mathcal{A} by a covering map. Thus, there exists a positive integer n_0 such that $f^{-1}(C_{n_0}) \neq \emptyset$ and f sends every component of $f^{-1}(C_{n_0})$ onto C_{n_0} by a covering map. Now, by an argument similar to what is given in the proof of [Lemma 3.18](#), after a proper homotopy, we may assume that $C'_{n_0} := f^{-1}(C_{n_0})$ is a single circle and $f|_{C'_{n_0}} \rightarrow C_{n_0}$ is a finite-sheeted covering. Let \mathcal{D}_* be the essential punctured disk contained in D_* such that $\partial \mathcal{D}_* = C_{n_0}$. Notice that $f(\mathbb{S}^1 \times \mathbb{R}) = D_*$.

Now, choose $r \neq 0$ so that $\mathbb{S}^1 \times r$ doesn't intersect C'_{n_0} . Since $\mathbb{S}^1 \times r$ co-bounds an annulus with each of the circles $\mathbb{S}^1 \times 0$ and C'_{n_0} , the isotopy extension theorem [[6](#), [Proposition 1.11](#)] [[15](#), [Theorem 1.3](#).] gives two homotopies $\mathcal{H}_1, \mathcal{H}_2: (\mathbb{S}^1 \times \mathbb{R}) \times [0, 1] \rightarrow \mathbb{S}^1 \times \mathbb{R}$ through homeomorphisms such that $\mathcal{H}_1(C'_{n_0}, 1) = \mathbb{S}^1 \times r$ and $\mathcal{H}_1(\mathbb{S}^1 \times r, 1) = \mathbb{S}^1 \times 0$. By [[2](#), [Theorem 1.3](#)], \mathcal{H}_1 and \mathcal{H}_2 are proper homotopies. Thus, there exists a homeomorphism of $\mathbb{S}^1 \times \mathbb{R}$ properly homotopic to the identity, which sends C'_{n_0} onto $\mathbb{S}^1 \times 0$. Therefore, after a proper homotopy, we may assume that $f(\mathbb{S}^1 \times \mathbb{R}) = D_*$ and $f|_{\mathbb{S}^1 \times 0} \rightarrow \partial \mathcal{D}_* = C_{n_0}$ is a finite-sheeted covering. Now, by the classification of finite-sheeted covering maps of the circle and together with an application of [Proposition 3.17](#) on each of $f|_{\mathbb{S}^1 \times (-\infty, 0]} \rightarrow \mathcal{D}_*$ and $f|_{\mathbb{S}^1 \times [0, \infty)} \rightarrow \mathcal{D}_*$, we can conclude that there exists a π_1 -injective, proper embedding $\iota: \mathbb{S}^1 \times [0, \infty) \hookrightarrow \Sigma$ with $\text{im}(\iota) = \mathcal{D}_*$ such that after a proper homotopy, f can be described by the proper map $\mathbb{S}^1 \times \mathbb{R} \ni (z, t) \mapsto \iota(z^d, |t|) \in \Sigma$ for some non-zero integer d . Moreover, by an argument similar to that given in the first part of the proof of [Theorem 3.8](#), we can conclude that given any compact subset K of Σ , there exists a proper map g properly homotopic to f such that $\text{im}(g) \subseteq \Sigma \setminus K$. \square

An argument similar to the one given in the proof of [Theorem 3.20](#) provides the following:

Proposition 3.21 Let $f: \mathbb{S}^1 \times \mathbb{R} \rightarrow \mathbb{S}^1 \times \mathbb{R}$ be a π_1 -injective proper map of degree 0. Then f is properly homotopic to $\mathbb{S}^1 \times \mathbb{R} \ni (z, t) \mapsto (z^d, |t|) \in \mathbb{S}^1 \times \mathbb{R}$ for some integer $d \neq 0$.

Now, we are ready to prove the second part of [Theorem 1.2](#). For that, we need the following lemma.

Lemma 3.22 Let $\varphi: \mathbb{C}^* \rightarrow \mathbb{C}^*$ be a proper map of non-zero degree. If φ is a homotopy equivalence, then it is properly homotopic to a homeomorphism.

Remark 3.23 The non-zero degree assumption in [Lemma 3.22](#) can't be dropped. For example, $f: \mathbb{S}^1 \times \mathbb{R} \ni (z, t) \mapsto (z, |t|) \in \mathbb{S}^1 \times \mathbb{R}$ is a self-homotopy equivalence of \mathbb{C}^* which is not properly homotopic to any homeomorphism of \mathbb{C}^* because $\deg(f) = 0$.

Proof. Pick a smoothly embedded primitive circle \mathcal{C} on \mathbb{C}^* . Thus, there are punctured disks $D_*^{(1)}$ and $D_*^{(2)}$ such that $\mathbb{C}^* = D_*^{(1)} \cup D_*^{(2)}$ and $\mathcal{C} = D_*^{(1)} \cap D_*^{(2)}$. By [Theorem 3.5](#), after a proper homotopy, we may assume φ is smooth as well as transverse to \mathcal{C} . Now, [Theorem 2.2.3](#) tells that φ remains surjective even after a proper homotopy. Therefore, $\varphi^{-1}(\mathcal{C})$ is a non-empty, pairwise-disjoint, finite collection of smoothly embedded circles on \mathbb{C}^* . Further, after a proper homotopy, similar to what we have done in the proof of [Theorem 3.15](#), we may assume $\mathcal{C}' := \varphi^{-1}(\mathcal{C})$ is a primitive circle on \mathbb{C}^* and $\varphi|_{\mathcal{C}'} \rightarrow \mathcal{C}$ is a homeomorphism. This is possible as φ is a homotopy equivalence. Let $D_*^{(1)'}$ and $D_*^{(2)'}$ be the punctured disks such that $\mathbb{C}^* = D_*^{(1)'} \cup D_*^{(2)'}$ and $\mathcal{C}' = D_*^{(1)'} \cap D_*^{(2)'}$. Since φ is surjective, possibly after re-indexing, we may assume $\varphi^{-1}(D_*^{(j)}) = D_*^{(j)'}$ for $j = 1, 2$. Finally, to conclude apply [Proposition 3.17](#) on $\varphi|_{D_*^{(j)'}} \rightarrow D_*^{(j)}$ for $j = 1, 2$. \square

Theorem 3.24 ([Theorem 1.2\(ii\)](#)) Let f be a π_1 -injective proper map from \mathbb{C}^* to a non-compact oriented surface Σ . Suppose $n := \deg(f)$ is non-zero. Then, $\Sigma = \mathbb{C}^*$ and f is properly homotopic to the covering $\mathbb{C}^* \ni z \mapsto z^n \in \mathbb{C}^*$ if $n > 0$, and to the covering $\mathbb{C}^* \ni z \mapsto \bar{z}^{-n} \in \mathbb{C}^*$ if $n < 0$

Proof. By [Proposition 3.19](#), $\Sigma = \mathbb{C}^*$. Let $p: \tilde{\Sigma} \rightarrow \mathbb{C}^*$ be the covering corresponding the subgroup $\text{im } \pi_1(f)$ of $\pi_1(\mathbb{C}^*)$, and let $\tilde{f}: \mathbb{C}^* \rightarrow \tilde{\Sigma}$ be a lift of f w.r.t. p , i.e., $p \circ \tilde{f} = f$. Thus, $\text{im } \pi_1(p) = \text{im } \pi_1(f)$, and hence $\pi_1(\tilde{f})$ is an isomorphism because a covering map induces injection between fundamental groups. In particular, the fundamental group of $\tilde{\Sigma}$ is infinite cyclic. Hence, $\tilde{\Sigma} = \mathbb{C}^*$. The properness of f implies the properness of \tilde{f} by [Lemma 3.3](#). Since non-compact surfaces are $K(\pi, 1)$ CW-complexes, by Whitehead theorem \tilde{f} is a homotopy equivalence. By [Lemma 3.3](#), we may assume p is a d -sheeted covering for some positive integer d and $\tilde{\Sigma}$ is orientable. Fix an orientation of $\tilde{\Sigma}$. By [Theorem 2.2.1](#), $\deg(p) = \pm d$. Now, $\deg(\tilde{f}) \neq 0$ because $0 \neq n = \deg(f) = \deg(p\tilde{f}) = (\pm d) \cdot \deg(\tilde{f})$. Thus, $\tilde{f}: \mathbb{C}^* \rightarrow \mathbb{C}^*$ is a homotopy equivalence and $\deg(\tilde{f}) \neq 0$. So \tilde{f} is properly homotopic to a homeomorphism by [Lemma 3.22](#), i.e., $\deg(\tilde{f}) = \pm 1$. Hence, $n = \pm d$, and f is properly homotopic to a d -sheeted covering.

At first, suppose $n > 0$. By the previous paragraph, without loss of generality, we may assume f is an n -sheeted covering map. Now, covering space theory [[13](#), [Proposition 1.37](#).] gives a self-homeomorphism h of \mathbb{C}^* such that $f(z) = h(z^n)$ for all $z \in \mathbb{C}^*$. Certainly, h is orientation-preserving. By [Proposition 3.25](#) below, we may assume $f(1) = 1 = h(1)$. Now, [[4](#), [Theorem 5.7](#).] tells that there exists a level-preserving homeomorphism $\mathcal{H}: \mathbb{C}^* \times [0, 1] \rightarrow \mathbb{C}^* \times [0, 1]$ which agrees with h on $\mathbb{C}^* \times 0$ and with $\text{Id}_{\mathbb{C}^*}$ on $\mathbb{C}^* \times 1$. The projection $\mathbb{C}^* \times [0, 1] \rightarrow \mathbb{C}^*$ is proper implies h is properly homotopic to $\text{Id}_{\mathbb{C}^*}$. So we are done when $\deg(f) = n > 0$.

Now, assume $\deg(f) = n < 0$. Since the complex conjugation is an orientation-reversing self-homeomorphism of \mathbb{C} , the map $\bar{f}: \mathbb{C}^* \ni z \mapsto \bar{f}(\bar{z}) \in \mathbb{C}^*$ is orientation-preserving of degree $-n$. Thus, \bar{f} is properly homotopic to $\mathbb{C}^* \ni z \mapsto z^{-n} \in \mathbb{C}^*$ by the previous case. Therefore, f is properly homotopic to $\mathbb{C}^* \ni z \mapsto \bar{z}^{-n} \in \mathbb{C}^*$. \square

It is known that every connected, boundaryless manifold M is homogeneous, i.e., for any two points x and y of M , there exists a homeomorphism h of M sending x to y . Moreover, one can also show that the homeomorphism h can be chosen so that it is homotopic to the identity map of M [28, Lemma 6.4]. In case M is non-compact, [Theorem 3.1](#) tells us that the homeomorphism h can be chosen so that it becomes properly homotopic to the identity of M .

Proposition 3.25 Let M be an orientable, connected topological manifold without boundary. If x and y are two points of M , then there exists a self-homeomorphism $\varphi: M \rightarrow M$ properly homotopic to the identity of M such that $\varphi(x) = y$.

Acknowledgments

This note is a chapter of my PhD thesis under the guidance of Siddhartha Gadgil. I thank him for his encouragement throughout this study and for several helpful conversations. This work is supported by a scholarship from the National Board for Higher Mathematics.

References

- [1] E. M. Brown and T. W. Tucker. On proper homotopy theory for noncompact 3-manifolds. *Trans. Amer. Math. Soc.*, 188:105–126, 1974.
- [2] R. H. Crowell. Invertible isotopies. *Proc. Amer. Math. Soc.*, 14:658–664, 1963.
- [3] Sumanta Das. Strong Topological Rigidity of Non-Compact Orientable Surfaces. *Algebr. Geom. Topol. (to appear)*, 2023. Available at [arXiv:2111.11194v3](https://arxiv.org/abs/2111.11194v3).
- [4] D. B. A. Epstein. Curves on 2-manifolds and isotopies. *Acta Math.*, 115:83–107, 1966.
- [5] D. B. A. Epstein. The degree of a map. *Proc. London Math. Soc. (3)*, 16:369–383, 1966.
- [6] Benson Farb and Dan Margalit. *A primer on mapping class groups*, volume 49 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 2012.
- [7] Stefan Friedl. *Topology (Course notes)*. AMS Open Math Notes, July 2023. Reference OMN:202307.111368. Available at <https://www.ams.org/open-math-notes/omn-view-listing?listingId=111368>.
- [8] Stefan Friedl, Matthias Nagel, Patrick Orson, and Mark Powell. A survey of the foundations of four-manifold theory in the topological category, 2024. [arXiv:1910.07372v3](https://arxiv.org/abs/1910.07372v3).
- [9] Roberto Frigerio. *Bounded cohomology of discrete groups*, volume 227 of *Mathematical Surveys and Monographs*. American Mathematical Society, Providence, RI, 2017.
- [10] Martin E. Goldman. An algebraic classification of noncompact 2-manifolds. *Trans. Amer. Math. Soc.*, 156:241–258, 1971.
- [11] Michael Gromov. Volume and bounded cohomology. *Inst. Hautes Études Sci. Publ. Math.*, 56:5–99 (1983), 1982.
- [12] Craig R. Guilbault. Ends, shapes, and boundaries in manifold topology and geometric group theory. In *Topology and geometric group theory*, volume 184 of *Springer Proc. Math. Stat.*, pages 45–125. Springer, [Cham], 2016.

- [13] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002. Available at <https://pi.math.cornell.edu/hatcher/AT/AT.pdf>.
- [14] Nicolaus Heuer and Clara Löh. The spectrum of simplicial volume of non-compact manifolds. *Geom. Dedicata*, 215:243–253, 2021.
- [15] Morris W. Hirsch. *Differential topology*, volume 33 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1994. Corrected reprint of the 1976 original.
- [16] Heinz Hopf. Zur Topologie der Abbildungen von Mannigfaltigkeiten Erster Teil. neue Darstellung der Theorie des Abbildungsgrades für topologische Mannigfaltigkeiten. *Math. Ann.*, 100(1):579–608, 1928.
- [17] Heinz Hopf. Zur Topologie der Abbildungen von Mannigfaltigkeiten Zweiter Teil Klasseninvarianten von Abbildungen. *Math. Ann.*, 102(1):562–623, 1930.
- [18] Robion C. Kirby. Stable homeomorphisms and the annulus conjecture. *Ann. of Math. (2)*, 89:575–582, 1969.
- [19] John M. Lee. *Introduction to topological manifolds*, volume 202 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 2000.
- [20] Edwin E. Moise. Affine structures in 3-manifolds. II. Positional properties of 2-spheres. *Ann. of Math. (2)*, 55:172–176, 1952.
- [21] Jakob Nielsen. Untersuchungen zur Topologie der geschlossenen zweiseitigen Flächen. *Acta Math.*, 50:189–358, 1927.
- [22] Paul Olum. Mappings of manifolds and the notion of degree. *Ann. of Math. (2)*, 58:458–480, 1953.
- [23] Richard S. Palais. Natural operations on differential forms. *Trans. Amer. Math. Soc.*, 92:125–141, 1959.
- [24] Richard S. Palais. Extending diffeomorphisms. *Proc. Amer. Math. Soc.*, 11:274–277, 1960.
- [25] Richard S. Palais. When proper maps are closed. *Proc. Amer. Math. Soc.*, 24:835–836, 1970.
- [26] Frank Quinn. Ends of maps. III. Dimensions 4 and 5. *J. Differential Geometry*, 17(3):503–521, 1982.
- [27] T. Radó. Über den Begriff der Riemannschen Fläche. *Acta Litt. Sci. Szeged*, 2:101–121, 1925. Available at <https://www.maths.ed.ac.uk/v1ranick/papers/rado.pdf>.
- [28] James W. Vick. *Homology theory*, volume 145 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, second edition, 1994. An introduction to algebraic topology.
- [29] Friedhelm Waldhausen. On irreducible 3-manifolds which are sufficiently large. *Ann. of Math. (2)*, 87:56–88, 1968.

DEPARTMENT OF MATHEMATICS, INDIAN INSTITUTE OF SCIENCE, BANGALORE 560012, INDIA

E-mail address: sumantadas@iisc.ac.in