

# MANIFOLDS WITH TRIVIAL CHERN CLASSES II: MANIFOLDS ISOGENOUS TO A TORUS PRODUCT, COFRAMED MANIFOLDS AND A QUESTION BY BALDASSARRI.

FABRIZIO CATANESE

**ABSTRACT.** Motivated by a general question addressed by Mario Baldassarri in 1956, we discuss characterizations of the Pseudo-Abelian Varieties introduced by Roth, and we introduce a first new notion, of Manifolds Isogenous to a  $k$ -Torus Product: the latter have the last  $k$  Chern classes trivial in rational cohomology and vanishing Chern numbers.

We show that in dimension 2 the latter class is the correct substitute for some incorrect assertions by Enriques, Dantoni, Roth and Baldassarri: these are the surfaces with  $K_X$  nef and  $c_2(X) = 0 \in H^4(X, \mathbb{Z})$ .

We observe in the last section, using a construction by Chad Schoen, that such a simple similar picture does not hold in higher dimension.

We discuss then, as a class of solutions to Baldassarri's question, manifolds isogenous to projective (respectively: Kähler) manifolds whose tangent bundle or whose cotangent bundle has a trivial subbundle of positive rank.

We see that the class of 'partially framed' projective manifolds (that is, whose tangent bundle has a trivial subbundle) consists, in the case where  $K_X$  is nef, of the Pseudo-Abelian varieties of Roth; while the class of 'partially co-framed' projective manifolds is not yet fully understood in spite of the new results that we are able to show here: and we formulate some open questions and conjectures.

In the course of the paper we address also the case of more general compact complex Manifolds, introducing the new notions of suspensions over parallelizable Manifolds, of twisted hyperelliptic Manifolds, and we describe the known results under the Kähler assumption.

*In memory of Mario Baldassarri (1920-1964).*

## CONTENTS

1. Introduction.	2
2. Surfaces with second Chern class equal to zero, and a Chern class characterization of Hyperelliptic surfaces.	3
3. Pseudo Abelian Varieties and Varieties Isogenous to a $k$ -Torus-Product	9
4. On the non Kähler case	14
5. Partially framed and co-framed manifolds	17

---

*Date:* May 2, 2024.

AMS Classification: 14F, 14K, 14C25

6. Mathematical and Historical comments on Baldassarri's paper	21
6.1. Historical comments	21
6.2. Mathematical comments	23
7. Manifolds with vanishing Chern numbers	25
7.1. The Schoen Calabi-Yau threefolds $X$ with $c_3(X) = 0$	26
References	28

## 1. INTRODUCTION.

Our first motivation was to understand a question posed in Mario Baldassarri's paper [Bald56], which aimed at determining the complex projective Manifolds whose last Chern classes are zero in rational cohomology, namely  $c_i(X) = 0 \in H^{2i}(X, \mathbb{Q})$  for  $i \geq k + 1$ . He made the wrong claim that these manifolds are exactly the Pseudo-Abelian varieties introduced by Roth [Roth54], hence his work also suggested to have a better understanding of the possible characterizations of the Pseudo-Abelian varieties.

Baldassarri's claim arose from the attempt to generalize to higher dimensions an incorrect assertion concerning algebraic surfaces (deriving from a minor mistake in the Castelnuovo-Enriques classification theorem of algebraic surfaces).

After clarifying the surface case, we explain here the state of the art in all dimensions, prove some new results, define new classes of compact complex Manifolds, and pose some new questions.

A first basic principle that Baldassarri and his predecessors missed to consider (as the use of topology was not sufficiently established at the time) was the following:

**Remark 1.1. (Isogeny principle):** If we have a finite unramified map between two compact complex Manifolds  $f : Z \rightarrow X$ , then  $c_i(Z) = 0 \in H^{2i}(Z, \mathbb{Q})$  if and only if  $c_i(X) = 0 \in H^{2i}(X, \mathbb{Q})$ .

Defining **isogeny** between Manifolds as the equivalence relation generated by the existence of such finite unramified maps, we see that the set of Manifolds which are solutions to Baldassarri's question consists of a union of isogeny classes.

Baldassarri's problem has been solved in the extremal case where  $k = 0$ : the characterization of the compact Kähler Manifolds with all Chern classes zero in rational (or real) cohomology was solved in 1978, thanks to Yau's celebrated theorem [Yau78] on the existence of Kähler-Einstein metrics on Manifolds with  $c_1(X) = 0 \in H^2(X, \mathbb{Q})$ . It follows that these Manifolds are the ones isogenous to complex tori (they are the so-called Hyperelliptic Manifolds, quotients  $X = T/G$  of a complex torus  $T$  by a finite group  $G$  acting freely on  $T$ ).

More precisely, we have:

**Theorem 1.2.** *(Yau) A compact Kähler manifold  $X$  such that  $c_1(X) = 0$ ,  $c_2(X) = 0$ , in  $H^*(X, \mathbb{R})$ , is a Hyperelliptic manifold.*

What happens for  $k \geq 1$ ?

The first significant case is the case where  $n := \dim(X) = 2, k = 1$ .

Already to describe this case we need some definition: a compact Kähler Manifold  $X$  is said to be a **torus product** iff  $X \cong T \times Y$ , where  $T$  is a complex torus of dimension  $\geq 1$ .

Whereas one says that  $X$  is **strongly isogenous to a torus product** if  $X = (T \times Y)/G$ , where  $G$  acts on  $T, Y$  and freely by its diagonal action on  $T \times Y$  (we shall see later that, if  $X$  is a compact Kähler non uniruled Manifold, the above two notions are equivalent).

The case of Roth's **Pseudo Abelian Varieties** is the special case where  $X = (T \times Y)/G$ ,  $X$  is projective, and moreover  $G$  acts on  $T$  as a group of translations.

In the case  $n = 2, k = 1$  we have a complete answer (see Theorem 2.8 for more details):

**Theorem 1.3.** *Let  $X$  be a compact Kähler surface with  $c_2(X) = 0 \in H^4(X, \mathbb{Z})$ . Then:*

- (1) *if  $K_X$  is nef, then  $X$  is strongly isogenous to a torus product, namely  $X = (T \times Y)/G$ , where  $G$  acts freely via a diagonal action on  $T \times Y$ ;*
- (2) *if  $X$  is a minimal surface, but  $K_X$  is not nef, then  $X$  is a  $\mathbb{P}^1$ -bundle over an elliptic curve  $E$ ;*
- (3) *if  $X$  is non minimal, then  $X$  is a blow up of a  $\mathbb{P}^1$ -bundle over a curve  $C$  of genus  $g \geq 2$ .*

*In the first two cases  $c_1(X)^2 = c_2(X) = 0$ , while in the last case  $\chi(X) = 1 - g$ , hence  $c_1(X)^2 = K_X^2 < 0$ .*

*If  $c_1(X) = 0 \in H^2(X, \mathbb{R})$ , then  $X$  is either a complex torus, or a hyperelliptic surface.*

*If  $c_1(X) = 0 \in \text{Pic}(X)$ , then  $X$  is a complex torus.*

There are three features in our above theorem for  $n = 2$ :

- (I): in case (1), where  $K_X$  is nef,  $X$  need not be a Pseudo-Abelian variety;
- (II): since  $\chi(\mathcal{O}_X) \leq 0$ , then  $X$  always possesses a non zero holomorphic 1-form  $\omega \in H^0(\Omega_X^1)$ ;
- (III): if moreover  $X$  is minimal, then  $\omega$  is everywhere non vanishing;
- (IV): if  $X$  is minimal, then  $X$  is birationally covered by a trivial family of (positive dimensional) complex tori, that is, there exists a dominant rational map  $\psi : T \times Y \dashrightarrow X$ .

The first feature (I) contradicts results of Dantoni [Dant43] and Enriques [Enr05b], (indeed Enriques' error is also reproduced in the classification theorem of Castelnuovo and Enriques [CastEn15]). These papers put Baldassarri and Roth on the wrong track.

The second feature (II) is the one which, as we shall see, fails to hold true in dimension  $n \geq 3$  as a consequence of vanishing of the top Chern class; while, if (II) holds true, since  $(-1)^n c_n(X)$  is the expected number of zeros of a holomorphic 1-form, then (III) looks plausible.

If one takes (III) as an assumption, that there exists a holomorphic 1-form without zeros, then Baldassarri's claim holds at least in dimension 3, as shown by [HS21b] (see also a similar result in Theorem 5.5).

The failure of (II) for  $n \geq 3$  is due to work of Chad Schoen [Schoen88], which shows that in dimension 3 there are simply connected manifolds, actually Calabi Yau manifolds ( $K_X$  trivial in  $Pic^0(X)$ ) with  $c_3(X) = 0$ .

Concerning the failure of feature (IV) for  $n \geq 3$ , our result here is that the Schoen threefolds are not birationally covered by a family of isomorphic Abelian surfaces, see Proposition 7.2, and that they do not admit a fibration onto a surface with general fibres isomorphic to a fixed elliptic curve  $E$ , see Proposition 7.3. Hence that they are also birationally far away from being isogenous to a torus product.

The first conclusion that we reach is therefore that, in higher dimensions, the right assumption to make is feature (III).

And the guiding observation is that an easy class of solutions to Baldassarri's question is provided by the manifolds  $X$  which are isogenous **to a partially framed or to a partially co-framed** manifold  $Z$ .

Our definition here is that  $Z$  is partially (tangentially) framed if the tangent bundle  $\Theta_Z$  admits a maximal trivial subbundle  $\mathcal{O}_X^k$ , with  $k > 0$ .

Whereas  $Z$  is said to be partially co-framed (cotangentially framed) if the cotangent bundle  $\Omega_Z^1$  admits such a maximal trivial subbundle  $\mathcal{O}_X^k$ , with  $k > 0$ .

The first definition brings back into play Roth's Pseudo-Abelian Varieties, see Theorem 5.3, which contains results also in the non projective case:

**Theorem 1.4.** *A tangentially  $k$ -framed projective Manifold with  $K_X$  nef is a pseudo-Abelian variety, in particular  $\Theta_X \cong \mathcal{O}_X^k \oplus \mathcal{F}$ .*

The previous classical Theorem was essentially proven by Roth and, thanks to the work of Fujiki [Fuj78] and Lieberman [Li78], the class of partially (tangentially) framed projective manifolds is understood. We also observe, using results of [Li78], [AMN12] (which also contains an excellent survey of the theory of framed Kähler manifolds), that the picture becomes more complicated if we enlarge our consideration to the wider realm of cKM (compact Kähler Manifolds), where more complicated constructions such as Seifert fibrations, principal torus bundles and torus suspensions enter into the picture (see section 4).

We also consider in Section 3 the non Kähler case, where we define a more general notion of suspension over a parallelizable Manifold, for which a splitting of the type  $\Theta_X \cong \mathcal{O}_X^k \oplus \mathcal{F}$  holds true.

And we consider all the Manifolds isogenous to a parallelizable Manifold, calling them **twisted Hyperelliptic Manifolds** since they also enjoy the

property of having all the real Chern classes equal to zero (but a general converse result is missing).

Passing to the second class, of partially co-framed manifolds, we see how this interesting class is more mysterious and presents some intriguing questions (see the discussion following Proposition 5.2 and Theorem 5.5).

A new result which we prove here is the following

**Theorem 1.5.** *Assume that  $X$  is a  $k$ -coframed compact Kähler manifold, such that either*

*(a)  $q := h^0(\Omega_X^1) = k$ ,*

*or more generally*

*(b) the framing is given by  $f^*(\Omega_A^1)$ , where  $f : X \rightarrow A$  is holomorphic and  $A$  is a complex torus of dimension  $k$  ( $f$  is the Albanese map  $a_X$  in the first case).*

*(i) Then  $f$  is a differentiable fibre bundle.*

*(ii) If  $X$  is projective, and  $K_X$  is nef, then  $X$  is a pseudo-Abelian variety provided one of the following assumptions is satisfied:*

*(1)  $n - k \leq 3$ ,*

*(2)  $K_X$  is pseudoample (there is a positive integer  $m$  such that  $|mK_X|$  is base point free),*

*(3) the fibres of  $f$  have a pseudoample canonical system,*

*(4)  $k \leq 5$ .*

A partial analogue of (1) holds also for compact Kähler manifolds, in view of [C-H-P16].

For the proof of the previous Theorem 5.5 we use a recent result by Taji [Taj23], proving birational isotriviality under the more general assumption that the fibres admit a good minimal model.

In the projective case with  $K_X$  nef we do not have yet more examples of  $k$ -coframed Manifolds than the class of the Pseudo-Abelian varieties.

Observe that hypothesis (2) holds if the so called Abundance conjecture is true, hence in particular it is reasonable to pose the following

**Conjecture 1.6.** *Assume that  $X$  is a  $k$ -coframed projective manifold with  $q = k$  and with  $K_X$  nef: then  $X$  is a pseudo-Abelian variety.*

In fact, we ask the more general

**Question 1.7.** *Are the  $k$ -coframed projective manifolds with  $K_X$  nef just the Pseudo-Abelian varieties?*

Finally, while for  $k$ -framed Manifolds and for  $k$ -coframed Manifolds the last  $k$  Chern classes are zero (in the Chow ring if the Manifolds are projective), we observe that more generally for the Manifolds isogenous to a  $k$ -Torus product (in view of Remark 1.1) not only the last  $k$  Chern classes are zero in rational cohomology, but also all the Chern numbers of  $X$  are equal to zero.

Hence in the last section we briefly discuss Manifolds with  $K_X$  nef and with vanishing Chern numbers.

For  $n = 2$ , as we already illustrated, there are only manifolds isogenous to a torus product; but, in dimension  $n = 3$ , in spite of a recent result by Hao-Schreieder [HS21a], describing threefolds with  $c_1c_2 = 0$ , and Kodaira dimension 2, as being birationally isogenous to a torus product, we have also the Schoen threefolds, which do not exhibit this feature.

Section 6 contains also some historical comments which may be useful in order to relate the present results to the classical literature.

## 2. SURFACES WITH SECOND CHERN CLASS EQUAL TO ZERO, AND A CHERN CLASS CHARACTERIZATION OF HYPERELLIPTIC AND ABELIAN SURFACES

In this section we establish Theorem 1.3 mentioned in the Introduction: this is done rerunning the proof of the classification of surfaces.

The classification theorem by Castelnuovo and Enriques [CastEn15] was extended by Kodaira [Kod68], and a crucial result is the following (see [Bea78], Chapter 6, Theorem VI.13, Theorem 1.4 of [CatLi19], and especially the crucial Theorem of [Cat22]):

**Theorem 2.1.** *Let  $S$  be a compact smooth complex surface, minimal in the strong sense that  $K_S$  is nef, and such that  $\chi(S) = 0$  (which implies that  $K_S^2 = 0$  and the topological Euler number  $e(S) = c_2(S) = 0$ ). Equivalently, assume that  $K_S$  is nef, and that  $e(S) = c_2(S) = 0$ .*

*Then  $X$  is strongly isogenous to a torus product, namely  $X = (T \times Y)/G$ , where  $T$  is a torus of positive dimension (1 or 2) and  $G$  acts freely via a diagonal action on  $T \times Y$ .*

More precisely,

**Theorem 2.2.** *The surfaces strongly isogenous to a torus product are classified as follows:*

- 1)  $p_g(S) = 1$ ,  $q(S) = 2$ , and  $S$  is a complex torus  $A$  (a **hyperelliptic surface of grade 1**), or
- 2)  $p_g(S) = p$ ,  $q(S) = p + 1$  and  $S$  is **isogenous** to an elliptic product, i.e.  $S$  is the quotient  $(C_1 \times C_2)/G$  of a product of curves of genera

$$g_1 := g(C_1) = 1, g_2 := g(C_2) \geq 1,$$

*by a free action of a finite group of product type (that is,  $G$  acts faithfully on  $C_1, C_2$  and we take the diagonal action  $g(x, y) := (gx, gy)$ ), such that, if we denote by  $g'_j = g(C_j/G)$ , then*

$$g'_1 + g'_2 = p + 1.$$

*Case 2) bifurcates into two subcases:*

(2.1,p) **pseudo-elliptic case:**  $g'_1 = 1$  (hence  $G$  acts on  $C_1$  by translations),  $C_2/G$  has genus  $p$ , and we assume <sup>1</sup>, for  $p = 1$ , that  $g_2 \geq 2$ ; or  
 (2.0,p)  $g'_1 = 0$  (hence  $C_1/G \cong \mathbb{P}^1$ ),  $C_2/G$  has genus  $p + 1 = q(S)$ , and we assume <sup>2</sup>, for  $p = 0$ , that  $g_2 \geq 2$ ; here the image of Albanese map  $\alpha : S \rightarrow \text{Alb}(S)$  equals  $C_2/G \subset \text{Alb}(S)$ .

Case (2.1,0) with  $g_2 = 1$  is the case where  $S$  is a properly hyperelliptic (bielliptic) surface (a **hyperelliptic surface of grade  $\geq 2$** ):

$S = (E_1 \times C_2)/G$ , where  $E_1, C_2$  are elliptic curves, and  $G$  acts via an action of product type, such that  $G$  acts on  $E_1$  via translations, and faithfully on  $C_2$  with  $C_2/G \cong \mathbb{P}^1$ .

In this case all the fibres of the Albanese map are isomorphic to  $C_2$ ,  $P_{12}(S) = 1$ , and  $S$  admits also an elliptic fibration  $\psi : S \rightarrow C_2/G \cong \mathbb{P}^1$ .

In the other cases (2.0,p), (2.1,p), for  $p \geq 1$ , (2.1,0) with  $g_2 \geq 2$ ,  $S$  is **isogenous to a higher genus elliptic product**, this means that  $C_2$  has genus  $g_2 \geq 2$ . Here  $S$  is properly elliptic and  $P_{12}(S) \geq 2$ .

The cases are distinguished mainly by the geometric genus  $p_g(S) = p$ .

In the case of the torus and of the hyperelliptic surfaces  $K_S$  is numerically equivalent to zero, whereas in the other cases  $K_S$  is not numerically equivalent to zero.

Moreover, the three cases are also distinguished (notice that  $c_1(S)$  is the class of the divisor  $-K_S$ ) by

- $K_S = 0 \in \text{Pic}(S)$  for the case 1) of a complex torus,
- $c_1(S) = 0 \in H^2(S, \mathbb{Z})$  but  $K_S \neq 0 \in \text{Pic}(S)$  in the case of properly hyperelliptic surfaces,
- $c_1(S) \neq 0 \in H^2(S, \mathbb{Q})$  in the other cases where  $S$  is isogenous to a higher genus elliptic product.

*Proof.* The part which is not contained in the cited sources concerns how to distinguish the several cases.

Everything is straightforward, except the assertion that for hyperelliptic surfaces (case (2.1,0) with  $g_1 = 1$ ) the first integral Chern class is zero: but this is a special case of our result on Bagnera de Franchis manifolds, Theorem 3.1 of [Cat23].

□

The following can be readily checked:

**Proposition 2.3.** *In case 1)  $A$  acts transitively and freely on  $A$ , so  $\text{Aut}^0(S)$  has dimension 2.*

*In case (2.1,p),  $\text{Aut}^0(S)$  has dimension 1:  $C_1 = E_1$  acts on  $S$ , transitively on the orbit closures, but with stabilizer  $H \subset G \subset E_1$  for the classes of points  $(x, y) \in E_1 \times C_2$  such that  $Hy = y$ .*

---

<sup>1</sup>to exclude that we are in case 1)

<sup>2</sup>to exclude that we are in case (2.1,0)



Finally, in case  $(2.0, p)$ ,  $\text{Aut}^0(S)$  has dimension 0 : there is no action of  $C_1$  on  $S$ , but the general fibres of the Albanese map  $\alpha : S \rightarrow C_2/G$  are isomorphic to  $C_1$  (a finite number shall only be isogenous to  $C_1$ ).

With a weaker notion of minimality we have a counterexample to the previous Theorem 2.1, as we shall now see.

**Proposition 2.4.** *If the smooth surface satisfies  $c_1^2 = c_2 = 0$ , that is  $K_S^2 = e(S) = 0$ , then  $S$  is minimal.*

*Proof.* By the Noether's formula, from  $K_S^2 = 0$  and  $e(S) = c_2(S) = 0$  follows  $\chi(\mathcal{O}_S) = 0$ .

And  $\chi(\mathcal{O}_S)$  is a birational invariant.

If  $S$  is not minimal, then  $S$  is the blow-up of a minimal surface  $S'$  with  $e(S') < 0$ . By Castelnuovo's theorem  $S'$  is a ruled surface with  $q(S') \geq 2$ . But then  $0 > \chi(\mathcal{O}_{S'}) = \chi(\mathcal{O}_S) = 0$ , a contradiction.  $\square$

**Corollary 2.5.** *Consider the minimal surfaces  $S$  with Chern numbers  $c_1^2 = c_2 = 0$ , that is with  $K_S^2 = e(S) = 0$ .*

*If  $K_S$  is nef,  $S$  is isogenous to a product  $Y \times A$ , with a torus  $A$  of dimension  $\geq 1$ .*

*If  $K_S$  is not nef, then  $S$  is a  $\mathbb{P}^1$ -bundle over an elliptic curve.*

*Proof.* By Theorem 2.1 there remains only to consider the case where  $K_S$  is not nef, hence  $S$  is ruled. Therefore, since the case of  $S = \mathbb{P}^2$  yields  $e(S) = 3$ , we must have a  $\mathbb{P}^1$ -bundle over a curve  $C$ . In this case, since  $0 = e(S) = 4(1 - g(C))$ , we get that  $C$  has genus  $g(C) = 1$ .  $\square$

**Remark 2.6.** If we take an elliptic curve  $B$ , and a vector bundle  $V$  of rank 2 on  $B$ , say  $V = L \oplus M$ , where  $\deg(L) = \deg(M)$ , then  $\mathbb{P}(V)$  is minimal, but the group of automorphisms of  $\mathbb{P}(V)$  consist of  $\mathbb{C}^*$  for general choice of  $L, M$  (see [Mar71], also Theorem 7.3 of [CatLiu21]).

Hence the action is not transitive on the orbit closures, because the two sections of  $\mathbb{P}(V)$  are left invariant by the automorphism group.

For completeness we show that:

**Proposition 2.7.** *Surfaces in the class  $(2.0, 0)$  do exist.*

*Proof.* Let  $G := (\mathbb{Z}/2)^3$ , and make it first act on an elliptic curve  $C_1$  as the group of transformations

$$z \mapsto \pm z + \eta, \quad 2\eta = 0,$$

so that  $G$  has generators  $\eta_1, \eta_2, \epsilon$ , where  $\epsilon(z) = -z$ .

To get a second action on  $C_2$  such that  $C_2/G =: E_2$  is an elliptic curve, we take  $E_2$  to be an elliptic curve,  $\mathcal{B} = \{x_1, x_2\}$  a branch set, so that

$$\pi_1(E_2 \setminus \mathcal{B}) = \langle \alpha, \beta, \gamma_1, \gamma_2 \mid \gamma_1 \gamma_2 = [\alpha, \beta] \rangle,$$



hence  $H_1(E_2 \setminus \mathcal{B}) = \mathbb{Z}\alpha \oplus \mathbb{Z}\beta \oplus \mathbb{Z}\gamma_1$ .

Define  $\mu : H_1(E_2 \setminus \mathcal{B}) \rightarrow G$  by:

$$\mu(\alpha) = \epsilon, \mu(\beta) = \eta_1, \mu(\gamma_1) = \eta_2 \Rightarrow \mu(\gamma_2) = \eta_2.$$

We want to prove that the product action of  $G$  on  $C_1 \times C_2$  is free.

To this purpose we observe that  $\eta_2$  has eight fixed points on  $C_2$ , and, since  $C_2$  has genus 5,  $E'_2 := C_2/\langle \eta_2 \rangle$  is an elliptic curve, so that  $E'_2 \rightarrow E_2$  is étale, that is,  $G/\langle \eta_2 \rangle$  acts freely on  $E'_2$ . The conclusion is that the only element acting on  $C_2$  with fixed points is  $\eta_2$ ; since  $\eta_2$  acts freely on  $C_1$ , the product action is free. □

We end this section observing that if  $S$  is a minimal surface with  $e(S) = 0$ , then either  $S$  is a  $\mathbb{P}^1$  bundle over an elliptic curve, or  $K_S$  is nef. In the latter case it must be  $K_S^2 = 0$ , since for  $K_S^2 < 0$   $S$  is ruled, hence  $K_S$  is not nef, and for  $K_S^2 > 0$   $S$  is of general type, and then  $e(S) > 0$ .

The following is the correction of the theorem of Dantoni [Dant43]; it shows that if  $S$  is not a torus then the surface is ‘elliptic’ only in the weak sense that it has a rational map with fibres elliptic curves, and not in the strong sense that a torus of dimension at least 1 should act on  $S$ , as we saw in Proposition 2.3 and in the previous remark 2.6 (it was also stated as Theorem 1.3 in the Introduction with only slightly different wording).

**Theorem 2.8.** *A surface  $S$  with  $e(S) = 0$  is either the blow up of a  $\mathbb{P}^1$ -bundle over a curve of genus at least 2, or it is minimal, and then it is either a complex torus, or a hyperelliptic surface, or it is isogenous to a higher genus elliptic product, or it is a  $\mathbb{P}^1$ -bundle over an elliptic curve.*

*In the last three cases  $S$  contains a 1-dimensional family of isomorphic elliptic curves whose union is dense in  $S$ .*

*If the canonical divisor is numerically trivial, then  $S$  is either a complex torus, or a hyperelliptic surface.*

### 3. PSEUDO ABELIAN VARIETIES AND VARIETIES ISOGENOUS TO A $k$ -TORUS-PRODUCT

The first aim of this section is to show how to derive from Roth’s original definition of the Pseudo Abelian Varieties the modern definition which we adopt in this article; and how Roth’s definition leads, in the case of compact Kähler Manifolds, to the notion of Seifert fibrations.

Later on we dwell on other notions, for instance we give a new definition, of suspension over a parallelizable Manifold.

Leonard Roth [Roth54], [Roth55] defined the Psudo-Abelian varieties as follows:

**Definition 3.1.** (Roth)

A projective variety  $X$  of dimension  $n$  is said to be Pseudo Abelian of order  $k$  if

i)  $\text{Aut}^0(X)$  contains a complex torus  $T$ , of maximal dimension  $= k$ , with the property that its orbits are all of dimension  $k$ .

Recall in fact the following theorem by Fujiki and Lieberman [Fuj78], [Li78]

**Theorem 3.2.** *If  $X$  is a compact Kähler manifold (or more generally in the Fujiki class) then there is an exact sequence of groups*

$$1 \rightarrow L \rightarrow \text{Aut}^0(X) \rightarrow T_X \rightarrow 1$$

where  $L$  is a linear algebraic group, and  $T_X$  is a complex torus.

The Lie Algebra of  $\text{Aut}^0(X)$  is  $\mathfrak{A} := H^0(\Theta_X)$ , while the Lie Algebra of  $L$  is the space  $\mathfrak{L}$  of vector fields which admit zeros ( $\mathfrak{L}$  is an ideal in the Lie Algebra  $\mathfrak{A}$ ).

If  $X$  is not uniruled, then  $L = \{1\}$ .

We explain now how to derive from Roth's definition a simpler one.

We have an action  $T \times X \rightarrow X$ , and we denote by  $T_x$  the orbit of  $x$ , image of  $T \times \{x\}$ .

Hence the orbits  $T_x$  of  $T$  give a variety  $\mathcal{V}$  of dimension  $n - k$  in the Hilbert scheme  $\mathcal{H}$  (Douady space) of  $X$ , and the restriction of the universal family to  $\mathcal{V}$ ,  $\phi : \mathcal{U} \rightarrow \mathcal{V}$ , yields  $\mathcal{U}$  which maps isomorphically to  $X$ . In fact,  $\dim(\mathcal{U}) = n = \dim(X)$ , and the identity of  $X$  factors as:

$$X \rightarrow \mathcal{U} \rightarrow \mathcal{V} \times X \rightarrow X,$$

where  $x \in X \mapsto (T_x, x)$ .

Hence we may write:  $\phi : X \rightarrow \mathcal{V}$ , and since the action of  $T$ ,

$$a : T \times X \rightarrow X$$

commutes with the projection over  $\mathcal{V}$ , and is effective, then the general fibre of  $\phi$  is isomorphic to  $T$ .

The other fibres are instead of the form  $T/G'$ , where  $G'$  is a finite subgroup of  $T$ . Since the general fibres are isomorphic, over an open set  $\mathcal{V}'$  of  $\mathcal{V}$  we have a holomorphic bundle (for instance, as a consequence of Kuranishi's theorem).

Because of the action of  $T$  on the fibres the monodromy of the bundle centralizes the group of translations hence the monodromy transformations consist of translations, and we have a principal bundle.

If  $X$  is projective, then the monodromy is finite, hence we get a finite group  $G \subset T$ .

The fibration is isotrivial, hence there exists a finite Galois base change  $f : Y \rightarrow \mathcal{V}$  with group  $G$  such that the fibre product is birational to a product

$$Y \times_{\mathcal{V}} X \sim Y \times T.$$

At each point of  $\mathcal{V}$  the local monodromy  $G'$  is a subgroup of  $G$ , hence the fibre product  $Y \times_{\mathcal{V}} X$  yields an unramified covering of  $X$  (since  $G'$  yields an unramified covering of the corresponding fibre).

Hence the fibre product  $Y \times_{\mathcal{V}} X$  is smooth, is birational to  $Y \times T$ , and all the fibres are isomorphic to  $T$ : therefore,  $Y \times_{\mathcal{V}} X \cong Y \times T$ , compatibly with the projection onto  $Y$ .

This motivates an equivalent definition, and some related definitions:

**Definition 3.3.** (I) A complex manifold  $X$  of dimension  $n$  is said to be a  $k$ -Pseudo-Torus (or Pseudo-Torus of order  $k$ ) if there is a torus  $T$  of dimension  $k$ , a manifold  $Y$ , and a finite Abelian group  $G$  acting on  $T$  faithfully via translations, acting faithfully on  $Y$ , such that the quotient of the product action is isomorphic to  $X$ :

$$X = (Y \times T)/G.$$

(II) If moreover  $X$  is projective, we shall say that  $X$  is a Pseudo-Abelian Variety of order  $k$  in the strong sense.

(III) A compact Kähler Manifold  $X$  is said to be Seifert fibred, cf. [Li78], if there is a finite abelian unramified covering  $Z \rightarrow X$  with Abelian Galois group  $G$  (hence  $X = Z/G$ ) such that  $Z$  is a principal bundle  $Z \rightarrow Y$ , with fibre a (positive dimensional) complex torus  $T$ , and where the action of  $G$  commutes with the action of  $T$  on  $Z$ .

(IV) A compact complex manifold  $X$  is called ([AMN12]) a suspension over a complex torus  $T = \mathbb{C}^k/\Lambda$  if there is a compact complex manifold  $Y$  and a homomorphism  $\rho : \Lambda \rightarrow \text{Aut}(Y)$  such that

$$X = (\mathbb{C}^k \times Y)/\Lambda, \lambda(z, y) := (z + \lambda, \rho(\lambda)(y)).$$

**Remark 3.4.** (a) The reason for definition (III) is that if  $X$  is only a cKM, and it has an action of a complex torus  $T$ , with all the orbits tori of the same dimension, then the global monodromy is not necessarily finite. But the local monodromies around the points corresponding to multiple fibres are finite subgroups of  $T$ , and since we have a finite number of them, the local monodromies generate a finite subgroup  $G$  of  $T$ . Associated to this subgroup there is an covering  $Y \rightarrow \mathcal{V}$  which is a principal  $T$ -bundle.

(b) In the case of definition (IV) of a suspension over a torus,  $\mathbb{C}^k$  acts on  $X$  via translations on the first summand, but in general there is no complex torus acting on  $X$ .

Indeed the  $\mathbb{C}^k$ -orbits are the projections  $\pi(\mathbb{C}^k \times \{y\})$ , which are isomorphic to  $\mathbb{C}^k/\text{Stab}_y$ , hence they do not need be compact.

The suspension over a torus is a Pseudo-torus if and only if the subgroup  $\text{Im}(\rho) \subset \text{Aut}(Y)$  is finite. Equivalently, if and only if all orbits are tori (which amounts to  $\text{Stab}_y$  being a subgroup of finite index in  $\Lambda$  for all  $y \in Y$ ): since then  $\text{Ker}(\rho)$  has finite index.

In particular the suspension is Pseudo-Abelian if and only if it is a Pseudo-torus.

(c) Observe that for such a suspension we have a splitting of the tangent bundle

$$\Theta_X \cong \mathcal{O}_X^k \oplus \mathcal{F},$$

hence also of the cotangent bundle.

In [AMN12] Example 2.4, page 1005, it is observed that a suspension  $X$  as above is Kähler if and only if  $Y$  is Kähler and a finite index subgroup  $\Lambda' \subset \Lambda$  maps to  $\text{Aut}^0(Y)$ .

In this case, taking  $G := \Lambda/\Lambda'$ , there is a Galois covering  $Z$  of  $X$  with group  $G$ . If moreover  $\text{Aut}^0(Y)$  is trivial, then  $Z = T' \times Y$  and we have a pseudo-torus.

If  $\text{Aut}^0(Y)$  is non-trivial, one cannot conclude that  $Z$  is a principal torus bundle, since the action of  $\Lambda'$  on  $Y$  need not be properly discontinuous.

(d) If  $X$  is Seifert fibred, then  $X = Z/G$ , and we have an exact sequence corresponding to the principal bundle  $Z \rightarrow Y$ :

$$0 \rightarrow \mathcal{O}_Z^k \rightarrow \Theta_Z \rightarrow \mathcal{F}_Z \rightarrow 0.$$

Since the action of  $G$  commutes with the action of  $T$ , the exact sequence descends to  $X$ , and we have

$$0 \rightarrow \mathcal{O}_X^k \rightarrow \Theta_X \rightarrow \mathcal{F} \rightarrow 0.$$

As we shall soon see, the above notions of Seifert-fibred manifold, or of pseudo-torus, and of suspension over a torus are not general enough in order to deal with manifolds with Chern classes trivial in real cohomology, hence we give another definition (important for the case of projective varieties):

**Definition 3.5.** A compact complex Manifold  $X$  of dimension  $n$  is said to be

- (i) **Isogenous to a k-Torus product** (MITP of order  $k$ ), if  $X$  is isogenous to a product  $Y \times T$ , where  $T$  is a torus of dimension  $k > 0$ ,  $Y$  is a compact complex manifold. If  $k$  is maximal with this property we say that  $X$  is a maximal MITP of order  $k$ .
- (ii) **Strongly Isogenous to a k-Torus product** if there is a torus  $T$  of dimension  $k > 0$ , a compact complex manifold  $Y$ , and a finite group  $G$  acting on  $T$ , and  $Y$  such that  $G$  acts freely on the product  $Y \times T$  via the induced diagonal action  $(g(t, y) := (g(t), g(y)))$  and moreover:

$$X = (Y \times T)/G.$$

If  $k$  is maximal with this property we say that  $X$  is a maximal SITP of order  $k$ .

If  $X$  is projective, we shall say that it is a Variety Isogenous to a k-Torus product.

**Proposition 3.6.** *If  $X$  is not uniruled, and  $X$  is a compact Kähler Manifold, the two notions (i) and (ii) are equivalent in the maximal case.*

*Proof.* Clearly (ii) implies (i), hence it suffices to show that (i) implies (ii), and we shall do it by induction on the number  $h$  of finite unramified maps which make  $X$  isogenous to such a product  $Y \times T$ .

**Step (a(1))** For  $h = 1$ , assume that there is a finite unramified map  $f : X \rightarrow Y \times T$ .

Then, taking the Galois closure  $Z$ , and letting  $G$  be the Galois group, we see that  $Z$  is associated to an epimorphism  $\psi : \pi_1(Y) \times \pi_1(T) \rightarrow G$ , and we denote  $G_1 := \psi(\pi_1(Y))$ ,  $G_2 := \psi(\pi_1(T))$ .

Then  $G$  is a quotient of  $G_1 \times G_2$  by the normal subgroup  $G_1 \cap G_2$ .

To the epimorphism  $\phi : \pi_1(Y) \times \pi_1(T) \rightarrow (G_1 \times G_2)$  corresponds a product Manifold  $Y_1 \times T_1$ , and clearly  $Z = (Y_1 \times T_1)/G_1 \cap G_2$ , hence  $X$  is as desired, since  $G_1 \times G_2$  acts on  $T_1$  diagonally (and on  $T_1$  via translations. Hence  $X$  is also a quotient of  $(Y_1 \times T_1)$  by another subgroup of  $G_1 \times G_2$ .

**Step (b(1))** For  $h = 1$ , assume that there is a finite unramified map  $f : (Y \times T) \rightarrow X$ . If  $f$  is not Galois, take the Galois closure  $Z$ , and denote by  $\Gamma$  the Galois group of  $Z \rightarrow X$ , and by  $G$  the subgroup which is the Galois group of  $Z \rightarrow (Y \times T)$ . Arguing as in Step (a(1)), we find a Galois cover  $(Y_1 \times T_1) \rightarrow (Y \times T)$  with group  $G_1 \times G_2$ .

We claim that the unramified cover  $(Y_1 \times T_1) \rightarrow X$  is Galois; to establish this claim it suffices to show that the elements  $\gamma \in \Gamma$  admit lifts to  $(Y_1 \times T_1)$ . Now,  $Z$  has a split tangent bundle  $\Theta_Z = \mathcal{F} \oplus \mathcal{O}_Z^k$ , and since  $Z$  is not uniruled, it follows from the Theorem of Fujiki and Lieberman, and by the maximality of  $k$ , that  $H^0(\Theta_Z) = H^0(\mathcal{O}_Z^k)$ .

Hence the splitting is uniquely determined, and the group  $T_2$  maps onto  $\text{Aut}^0(Z)$ . The fibration  $Z \rightarrow (Z/T_2)$  is then isotrivial, and  $(Y_1 \times T_1)$  is the canonical product pull back under  $Y_1 \rightarrow (Z/T_2)$ .

Therefore the lifting property holds true.

**Step (a(h))** Assume that  $X_2 = (Y_2 \times T_2)/G_2$ , as in (ii), and that there is a finite unramified map  $f : X \rightarrow X_2$ .

Then, taking a component  $X'$  of the fibre product  $X \times_{X_2} (Y_2 \times T_2)$ , and applying (a(1)) we get that  $X'$ , hence also  $X$ , admits  $(Y_1 \times T_1)$  as an unramified covering, hence we can conclude because of Step (b(1)).

**Step (b(h))** Assume then that  $X_2 = (Y_2 \times T_2)/G_2$ , as in (ii), and that there is a finite unramified map  $f : X_2 \rightarrow X$ .

Then taking the composition we get an unramified map  $(Y_2 \times T_2) \rightarrow X$ , and again we are done by Step (b(1)).

□

The following are easy important properties of such Manifolds and Varieties Isogenous to a Torus product.

**Proposition 3.7.** *Let the complex Manifold  $X$  of dimension  $n$  be Isogenous to a  $k$ -Torus product  $Y \times T$  where  $\dim(T) = k > 0$ .*

*Then the integral Chern classes  $c_i(Y \times T) \in H^*(Y \times T, \mathbb{Z})$  vanish for  $i \geq n - k + 1$ , and the rational Chern classes  $c_i(X) \in H^*(X, \mathbb{Q})$  vanish for  $i \geq n - k + 1$ .*

*Moreover all the Chern numbers of  $X$  vanish.*

If moreover  $X$  is (respectively is isogenous to) a pseudo-torus, or more generally the suspension over a torus, or is Seifert fibred, then all the integral Chern classes  $c_i(X) \in H^*(X, \mathbb{Z})$  vanish for  $i \geq n - k + 1$  (respectively the corresponding rational Chern classes vanish).

*Proof.* Since the tangent bundle of  $Y \times T$  is the direct sum of the pull back of the tangent bundle of  $Y$  with the pull back of the tangent bundle of  $T$  which is trivial, the total Chern class of  $Y \times T$  is the pull back of the total Chern class of  $Y$ , and this proves the first assertion.

The second assertion follows since the Chern classes of  $X$  pull back to the Chern classes of the product  $Y \times T$ , and  $H^*(X, \mathbb{Q}) = H^*(Y \times T, \mathbb{Q})^G$ .

The third assertion follows since any isobaric polynomial of weight  $n$  in the Chern classes of  $X$  is a rational multiple of the same isobaric polynomial of weight  $n$  in the Chern classes of  $Y$ , and we are assuming  $k > 0$ .

The last assertion follows since the tangent bundle of  $X$  either splits as  $\Theta_X = \mathcal{O}_X^k \oplus \mathcal{F}$ , or has a bundle exact sequence

$$0 \rightarrow \mathcal{O}_X^k \rightarrow \Theta_X \rightarrow \mathcal{F} \rightarrow 0.$$

□

#### 4. ON THE NON KÄHLER CASE

In the realm of compact complex manifolds, the Manifolds  $X$  with a holomorphically trivial tangent bundle  $\Theta_X \cong \mathcal{O}_X^n$  are called the **parallelizable Manifolds**.

By a Theorem of Wang [Wang54] every parallelizable Manifold is a quotient  $X = \mathcal{G}/\Lambda$ , where  $\mathcal{G}$  is a complex Lie group and  $\Lambda$  is a cocompact discrete subgroup.

Since every Manifold isogenous to a parallelizable Manifold has trivial real Chern classes, it would be interesting to study this class in more detail (in the Kähler case the parallelizable Manifolds are the complex tori, and the Manifolds isogenous to a torus are the Hyperelliptic Manifolds).

**Proposition 4.1.** *Every Manifold  $Y$  isogenous to a parallelizable Manifold  $X' = \mathcal{G}/\Lambda'$  is obtained from another parallelizable Manifold  $X = \mathcal{G}/\Lambda$  dividing by the action of a finite group  $G$  acting freely on  $X$ .*

*$G$  is a group of affine transformations of  $X$ , of the form  $f(x) = gF(x)$  where  $g \in \mathcal{G}$  and  $F$  has a fixed point  $x_0$  and has derivative  $DF$  determined by the value of the derivative at  $x_0$ .*

*The quotient Manifold  $Y := X/G$  will be called a **twisted hyperelliptic Manifold**.*

*Proof.* If  $Y$  is isogenous to  $X'$ , then  $Y$  has also  $\mathcal{G}$  as universal cover, as it is trivial to see.

Hence we can write  $Y = \mathcal{G}/\Gamma$ , where  $\Gamma$  acts freely on  $\mathcal{G}$  and with compact quotient.

Observe that any isogeny amounts, up to isomorphism, to a finite index containment between fundamental groups, viewed as discrete groups of automorphisms of  $\mathcal{G}$  (since  $\mathcal{G}$  is simply connected).

It is a general fact that the intersection  $H \cap K$  of two finite index subgroups  $H, K$  of a group  $\Gamma'$  is again of finite index in  $\Gamma'$  (it is the stabilizer of the orbit of  $H \times K$  on  $\Gamma'/H \times \Gamma'/K$ <sup>3</sup>).

Hence, if  $\Gamma_2$  is of finite index in  $\Gamma_1$  and  $\Gamma_3$ , and  $\Gamma_4$  is of finite index in  $\Gamma_3$  and  $\Gamma_5$ , then  $\Gamma_2 \cap \Gamma_4$  is of finite index in  $\Gamma_3$ , hence also in  $\Gamma_1$  and  $\Gamma_5$ .

Therefore, since we can always assume that the number of direct relations yielding the isogeny is even (introducing an equality step), we can reduce to the case where  $\Gamma$  and  $\Lambda'$  contain a finite index subgroup in both of them, call it  $\Lambda''$ . By replacing  $X'' = \mathcal{G}/\Lambda''$  by the Galois closure of  $X'' \rightarrow Y$ , we obtain  $Y = X/G$  as desired.

Now, if  $f \in G$ , there exists  $g \in \mathcal{G}$  such that  $F := g^{-1}f$  has a fixed point  $x_0 := \Lambda$  on  $X$ . The derivative of  $F$  is a holomorphic section of the trivial bundle  $DF \in H^0(\text{End}(\mathcal{O}_X^n))$  gotten by the action on the space of left invariant vector fields: since  $X$  is compact,  $DF$  is constant, hence  $f(x) = gF(x)$  is affine.

□

Of course each such automorphism  $F$  lifts to an automorphism  $\tilde{F}$  of  $\mathcal{G}$  which must have the property of fixing the identity and of sending  $\Lambda$  to  $\Lambda$ , since we must have

$$\tilde{F}(g\lambda) = \tilde{F}(g)\pi_1(F)(\lambda) \Rightarrow \tilde{F}(\lambda) = \pi_1(F)(\lambda),$$

this is exactly as it happens for automorphisms of complex tori.

We give now a simple example of such Manifolds.

**Proposition 4.2.** *Consider the Iwasawa Manifold  $X = \mathcal{G}/\Lambda$ , where  $\mathcal{G}$  is the complex Lie group of unipotent  $3 \times 3$  complex matrices, that is, upper triangular matrices and with diagonal entries equal to 1.*

*And let  $\Lambda$  be the subgroup of matrices with upper-diagonal entries belonging to the ring  $\mathbb{Z}[i]$  of Gaussian integers.*

*Then we define the following automorphism of  $X$ , such that*

$$r[(z_{12}, z_{21}, z_{13})] := [(iz_{12}, z_{21} + 1/4, iz_{13})].$$

*Then  $r$  has order 4, and generates a group  $G \cong \mathbb{Z}/4$  acting freely on  $X$ . The quotient is a twisted Hyperelliptic Manifold with  $H^0(\Omega_Y^1)$  of dimension 1 and generated by a closed holomorphic 1-form..*

*Proof.* The automorphism is well defined, since the product  $z\lambda$  has coordinates

$$(z_{12} + \lambda_{12}, z_{21} + \lambda_{21}, z_{13} + \lambda_{13} + z_{12}\lambda_{21}),$$

hence we infer  $[r(z)] = [r(z\lambda)]$  since

---

<sup>3</sup>see for instance: stack Overflow



$$\begin{aligned}
[r(z\lambda)] &= [(iz_{12} + i\lambda_{12}, z_{21} + \lambda_{21} + 1/4, iz_{13} + i\lambda_{13} + iz_{12}\lambda_{21})] = \\
&= [(iz_{12}, z_{21} + \lambda_{21} + 1/4, iz_{13} + iz_{12}\lambda_{21})] = [r(z)\lambda'],
\end{aligned}$$

where  $\lambda' = (0, \lambda_{21}, 0)$ .

That the action is free follows since for the second coordinate

$$z_{21} \mapsto z_{21} + 1/4.$$

It is known that  $H^0(\Omega_X^1)$  is generated by

$$\omega_1 := dz_{12}, \omega_2 := dz_{21}, \omega_3 := dz_{13} - z_{21}dz_{12},$$

which are mapped by  $r$  to  $i\omega_1, \omega_2, i\omega_3 - (i/4)\omega_1$ .

Hence  $H^0(\Omega_X^1)^G$  is generated by  $\omega_1$ .  $\square$

We leave aside here the general discussion of the structure of twisted Hyperelliptic Manifolds, but we raise a question.

**Question 4.3.** The twisted Hyperelliptic Manifolds, being isogenous to parallelizable Manifolds, are examples of compact complex manifolds with all the real Chern classes  $c_{i,\mathbb{R}}(X) = 0, \forall i$ .

Are there other examples ?

We can also define another general class of Manifolds, which we call Suspensions over a parallelizable Manifold.

**Definition 4.4.** (S-P-M) A compact complex manifold  $X$  is called a **suspension over a parallelizable Manifold**  $Z = \mathcal{G}/\Lambda$  if there is a compact complex manifold  $Y$  and a homomorphism  $\rho : \Lambda \rightarrow \text{Aut}(Y)$  such that

$$X = (\mathcal{G} \times Y)/\Lambda, \lambda(z, y) := (z\lambda, y\rho(\lambda)).$$

**Proposition 4.5.** *If  $X$  is a suspension over a parallelizable Manifold  $Z$ , then we have a splitting of the tangent bundle*

$$\Theta_X = \mathcal{O}_X^k \oplus \mathcal{F},$$

where  $k = \dim(Z)$ .

*Proof.* The splitting follows right away since  $\Lambda$  has a product action. Moreover, the first summand is trivial because  $Z = \mathcal{G}/\Lambda$  is parallelizable and the first summand is just the pull-back of the tangent bundle  $\Theta_Z$  for the projection  $X \rightarrow Z$ , with fibres isomorphic to  $Y$ .  $\square$

**Remark 4.6.** In the case where  $\rho$  has finite image, we get that,  $\Lambda'$  being defined as  $\Lambda' := \ker(\rho)$ , then  $X$  is isogenous to  $Z' \times Y$ , where  $Z' := \mathcal{G}/\Lambda'$ . And we get then triviality of the Chern numbers of  $X$ .

In the next section we shall show how, assuming that  $X$  is Kähler, the occurrence of such a splitting  $\Theta_X = \mathcal{O}_X^k \oplus \mathcal{F}$  is understood as the structure of a suspension over a torus.

It is interesting to see whether a similar characterization of suspensions over a parallelizable Manifold holds also for compact complex manifolds.

For this purpose one has to take  $k$  maximal, and the first key point would be to see whether  $H^0(\mathcal{O}_X^k) \subset H^0(\Theta_X)$  is a Lie subalgebra  $\mathfrak{G}$ : then we would have an action on  $X$  of the simply connected Lie group  $\mathcal{G}$  associated to  $\mathfrak{G}$ .

## 5. PARTIALLY FRAMED AND CO-FRAMED MANIFOLDS

We begin with a simple definition yielding complex Manifolds with all the last  $k$  integral Chern classes equal to zero.

**Definition 5.1.** A complex manifold  $X$  of dimension  $n$  is said to be  $k$ -tangentially framed, or simply  $k$ -framed, if the holomorphic tangent bundle  $\Theta_X$  admits a trivial subbundle  $\cong \mathcal{O}_X^k$ , and where  $k > 0$  is maximal.

A complex manifold  $X$  of dimension  $n$  is said to be  $k$ -cotangentially framed, or simply  $k$ -co-framed, if the holomorphic cotangent bundle  $\Omega_X^1$  admits a trivial subbundle  $\cong \mathcal{O}_X^k$ , and where  $k > 0$  is maximal.

In the rest of the section we shall use the results of [Li78], [Fuj78], [AMN12], often for simplicity we might refer to the exposition given in the last paper.

**Proposition 5.2.** *Assume that  $X$  is a  $k$ -coframed compact Kähler manifold and let  $W \subset h^0(\Omega_X^1)$  be the corresponding maximal vector subspace consisting entirely of nowhere vanishing holomorphic 1-forms (of dimension  $k \geq 1$ ). Then  $W$  determines an analytically integrable foliation with trivial normal bundle.*

*i) If moreover the subspace  $W$  corresponds to a quotient torus  $T'$  of  $\text{Alb}(X) = H^0(\Omega_X^1)^\vee / (H_1(X, \mathbb{Z})/\text{Tors})$ , then the foliation is algebraically integrable, consisting of the fibres of  $\Psi : X \rightarrow T'$ , which is a differentiable fibre bundle.*

*ii) If we have a splitting  $\Omega_X^1 \cong \mathcal{O}_X^k \oplus \mathcal{F}^\vee$ , then  $\Psi$  is a holomorphic fibre bundle with fibre  $Y$ .*

*iii) If moreover  $Y$  has finite automorphism group, then  $X$  is a  $k$ -Pseudo-Torus product  $X = (Y \times T)/G$  with  $\dim(T) = k > 0$ .*

*Proof.* A subspace  $W$  which is maximal with the property that all forms  $\omega \in W \setminus \{0\}$  are nowhere vanishing, has a basis  $\omega_1, \dots, \omega_k$  such that the forms  $\omega_j$  are linearly independent at each point, hence they generate a trivial rank  $k$  subbundle of  $\Omega_X^1$ .

The associated foliation is analytically integrable, because  $X$  is a cKM, and holomorphic 1-forms are closed, hence the distribution induced by  $W$  is integrable, and spans a trivial subbundle of the cotangent bundle.

The foliation on  $X$  is induced by a foliation on  $\text{Alb}(X)$ , corresponding to the annihilator of  $W$ . This foliation is algebraically integrable if, as we assume, it corresponds to the projection onto a quotient torus  $T'$ .

The composed map  $\Psi : X \rightarrow T'$  has fibres of dimension  $k$ , which are therefore union of leaves. Since the fibres are smooth, if they are not connected, we would get by the Stein factorization on unramified covering of  $T'$ , which is again a quotient of  $Alb(X)$  by the universal property of the Albanese variety.

Hence we may assume that the fibres of  $\Phi$  are connected, and we have a differentiable fibre bundle.

The splitting  $\Omega_X^1 \cong \mathcal{O}_X^k \oplus \mathcal{F}^\vee$  guarantees that the Kodaira-Spencer map for the family is identically zero, hence by Kuranishi's theorem we have a holomorphic bundle.

If this is a holomorphic bundle, with fibre  $Y$ , and  $Aut(Y)$  is finite, there is an unramified covering  $T \rightarrow T'$  with group  $G$  such that the pull back is a product. Therefore  $X = (T \times Y)/G$ , where  $G$  acts on  $T$  via translations.  $\square$

**Theorem 5.3.** *Let  $X$  be a compact Kähler complex Manifold of dimension  $n$ : then  $X$  is the suspension over a torus  $T$  with  $\dim(T) = k > 0$  if and only if there is a  $k$ -framing yielding a partial tangential splitting*

$$\Theta_X \cong \mathcal{O}_X^k \oplus \mathcal{F}.$$

*A  $k$ -framing of  $X$  yields the structure of a Seifert fibration on  $X$  in the case where  $h^0(\Theta_X) = k$ .*

*Moreover, a  $k$ -framed projective manifold  $X$  is a suspension over an Abelian Variety and is indeed a Pseudo-Abelian variety if  $K_X$  is nef.*

*Proof.* We have already seen that for a suspension over a torus we have such a partial tangential splitting.

Conversely, recall that  $H^0(\Theta_X)$  is the Lie algebra of the Lie group  $Aut(X)$ . The Lie Algebra  $H^0(\Theta_X) =: \mathfrak{A}_X$  contains the Lie ideal  $\mathcal{H}_X^1$  of the vector fields admitting zeros, and there is (see [AMN12], page 1002), a direct sum

$$\mathfrak{A}_X = \mathcal{H}_X^1 \oplus \mathcal{A},$$

where  $\mathcal{A}$  is a maximal Abelian subalgebra generated by nowhere vanishing vector fields.

By Theorem 3.14 of [Li78],  $\mathcal{H}_X^1$  is precisely the subspace of  $\mathcal{H}_X$  yielding the zero flow on  $Alb(X)$ .

If  $\mathcal{H}_X^1 = 0$  then the trivial subbundle yields  $k$  everywhere linearly independent vector fields, which, see [Fuj78], [Li78], and also Theorem 1.2 of [AMN12], generate the action of a  $k$ -dimensional complex torus  $T$  with smooth orbits  $T_x$ , quotients of  $T$  by a finite group  $H_x$ .

Hence  $X$  is Seifert fibred, as we have discussed earlier, cf. Theorem 4.9 by Lieberman [Li78].

If we have a tangential splitting  $\Theta_X \cong \mathcal{O}_X^k \oplus \mathcal{F}$ , we get a corresponding cotangent splitting,

$$\Omega_X^1 \cong \mathcal{O}_X^k \oplus \mathcal{F}^\vee.$$

It suffices, by the preceding proposition, to show that the coframing defines a subspace  $W \subset H^0(\Omega_X^1)$  corresponding to a quotient torus  $T'$  of  $\text{Alb}(X)$ . Since every automorphism of  $X$  yields an affine action on  $\text{Alb}(X)$ , the action of the torus  $T$ , which spans  $W^\vee$ , yields a subtorus  $A$  of  $\text{Alb}(X)$ . Then we define  $T' := \text{Alb}(X)/A$ , and we get  $\Psi : X \rightarrow T'$  which is a holomorphic bundle, with parallel transport given by the action of  $T$ : hence  $X$  is the suspension over a torus.

Finally, if  $X$  is projective,  $T'$  is an Abelian variety. We defer the reader to Theorem 0.3 of [AMN12] for the proof of the last assertions, see also [Li77].  $\square$

**Remark 5.4.** Theorem 0.3 of [AMN12] proves the following very interesting result: if  $X$  is a compact Kähler manifold which is  $k$ -framed, then  $X$  admits a small deformation which is a suspension over a  $k$ -dimensional torus, and which is a  $k$ -Pseudo-Torus if  $Kod(X) \geq 0$ .

If  $X$  is projective and  $k$ -framed, as we saw, they show that  $X$  is Pseudo-Abelian.

**Theorem 5.5.** *Assume that  $X$  is a  $k$ -coframed compact Kähler manifold, such that either*

(a)  $q := h^0(\Omega_X^1) = k$ ,

*or more generally*

(b) *the framing is given by  $f^*(\Omega_A^1)$ , where  $f : X \rightarrow A$  is holomorphic and  $A$  is a complex torus of dimension  $k$  ( $f$  is the Albanese map  $a_X$  in the first case).*

(i) *Then  $f$  is a differentiable fibre bundle.*

(ii) *If  $X$  is projective, and  $K_X$  is nef, then  $X$  is a pseudo-Abelian variety provided one of the following assumptions is satisfied:*

(1)  $n - k \leq 3$ ,

(2)  $K_X$  is pseudoample (there is a positive integer  $m$  such that  $|mK_X|$  is base point free),

(3) the fibres of  $f$  have a pseudoample canonical system,

(4)  $k \leq 5$ .

*Proof.* The first assertion follows from i) of Proposition 5.2.

We prove now as a warm up the second assertion (ii) in the case where the fibres of  $f$  have dimension 1, hence they are smooth curves of genus  $g \geq 1$ . Since the fibration induces a holomorphic map to the Teichmüller space  $\mathcal{T}_g$  which is biholomorphic to a bounded domain, this map is constant and we have a holomorphic fibre bundle.

Since  $X$  is projective, the monodromy is finite, so there exists a finite unramified map  $A' \rightarrow A := \text{Alb}(X)$  such that the pull back is a product, hence  $X$  is a Pseudo-Abelian variety.

Since  $K_X$  restricts to the canonical bundle on the fibres, the fibres have nef canonical divisor, and clearly (2) implies (3).

Also (1) implies (3): if the fibres have dimension  $\leq 3$ , then the abundance conjecture is true by results of Miyaoka and Kawamata, as explained in [Flips92], that is, (3) holds.

We are reduced to proving the result under assumption (3), which implies that the fibres have a good minimal model.

We can apply then Theorems 1.1, 1.2 and 1.3 of [Taj23] implying that we have birational isotriviality, namely, there exists a base change  $\phi : W \rightarrow A$  such that the pull-back

$$\phi^*(X) := W \times_A X$$

of  $f : X \rightarrow A$  is birational to a product. Given a general smooth fibre  $F$  of  $f$ , we can assume that  $\phi^*(X)$  is birational to  $W \times F$ , and that this birational map induces the identity on  $F$ .

Hence there is a Zariski open set  $U$  of  $A$  such that the fibres over  $U$  are isomorphic to  $F$ .

We apply now the results of Wavrik[Wav69] in order to conclude that, for each fibre  $F_a$ , the Kuranishi space  $Def(F_a)$  is a local moduli space.

The hypothesis needed is that the dimension of the space of holomorphic vector fields  $h^0(F_a, \Theta_{F_a})$  is constant. Since the fibres  $F_a$  have nef canonical divisor, they do not admit vanishing holomorphic vector fields, see Theorem 3.2, hence their space of vector fields is the Lie algebra of a complex torus  $Aut^0(F_a)$ , hence the fibres, being projective with nef canonical divisor, are pseudo-Abelian. And a deformation of a  $k$ -pseudo Abelian variety is  $k$ -pseudo Abelian: this shows that the dimension  $h^0(F_a, \Theta_{F_a})$ , which is upper semicontinuous by standard results, is also lower semicontinuous, whence it is constant and we can apply Wavrik's result.

We have now a holomorphic map of a neighbourhood  $\mathcal{U}$  of  $a \in A$  into  $Def(F_a)$ , and this map is constant on a dense subset: hence this holomorphic map is constant, and we have shown that  $f$  is a holomorphic bundle.

Since  $X$  is projective, we conclude as above that  $X$  is a pseudo Abelian variety.

Under assumption (4) we can instead apply [Taj21] to obtain birational isotriviality, and proceed similarly.

□

**Remark 5.6.** (I) Isotriviality of such a fibration was first shown by Kovacs in the case where the fibres have ample canonical divisor [Kov97], and this result was extended in [W-W23] who proved birational isotriviality in the more general case of fibres of general type.

**Example 5.7.** This example concerns  $k$ -coframed varieties.

(1) A smooth fibration onto an Abelian variety need not be isotrivial, and not even a holomorphic bundle.

Let in fact  $A$  be an elliptic curve and assume that we have an embedding  $j : A \rightarrow S$ , where  $S$  is a surface of general type, say with  $K_S$  ample.

Let  $\Gamma \subset A \times S$  be the graph of  $j$ , and take  $X$  to be the blow up of  $A \times S$  with centre the smooth curve  $\Gamma$ .

Then the differentiable fibre bundle  $f : X \rightarrow A$  is not a holomorphic bundle, since  $\text{Aut}(S)$  is finite, and the pairs  $(S, p_1)$  and  $(S, p_2)$  are not isomorphic for general  $P_1, P_2 \in j(A)$ .

(2) In this case  $K_X$  is not nef.

We pose now the following general questions, (b) being a generalization of Conjecture 1.6:

**Question 5.8.** Assume that  $X$  is a  $k$ -coframed projective manifold. Does there exist a coframing  $V$  (a subbundle  $V \cong \mathcal{O}_X^k$  of  $\Omega_X^1$ ) such that i) of Proposition 5.2 holds, namely the subspace  $W = H^0(V) \subset H^0(\Omega_X^1)$  defines a quotient Abelian variety  $A'$ ?

(b) If  $X$  is projective with  $K_X$  nef and it admits a fibration  $f : X \rightarrow A'$  onto an Abelian variety  $A'$  with all fibres smooth, is then  $f$  a holomorphic bundle (and hence  $X$  is a pseudo-Abelian variety)?

**Remark 5.9.** Question (a) asks, in the case where  $X$  is projective, whether there is a choice of the subspace  $W \subset H^0(\Omega_X^1)$  corresponding to an Abelian subvariety of  $\text{Alb}(X)$ , equivalently, whether  $\Lambda^{2k}(W \oplus \bar{W})$  is a point defined over  $\mathbb{Q}$  in the Grassmann manifold  $\text{Grass}(2k, H^1(X, \mathbb{C})) \subset \mathbb{P}(\Lambda^{2k}(H^1(X, \mathbb{C})))$ . One may reduce to the case where  $X$  is defined over an algebraic extension of  $\mathbb{Q}$ .

## 6. MATHEMATICAL AND HISTORICAL COMMENTS ON BALDASSARRI'S PAPER

**6.1. Historical comments.** Baldassarri [Bald56] wanted to characterize the smooth projective manifolds  $X$  whose first  $h$  canonical systems

$$K_0(X), \dots, K_{h-1}(X)$$

have degree zero.

At that time, even if Baldassarri in his *Ergebnisse* book [Bald56book] (the book was rather influential, it was for instance translated in Russian by Manin) was exposing the new methods in the theory of Algebraic Varieties, the concept of canonical systems of all dimension was based on geometric approaches.

The canonical systems of a manifold (see [Roth56]), defined in a geometric way by Todd, Eger and later in a simpler way by Beniamino Segre [Seg52], [Seg54]<sup>4</sup> after proposals made by Severi, were shown in 1955 by Nakano [Nak55] to be the so called Chern classes of the cotangent bundle (see [At98] for an historical account and [Ful84] as a general reference): more precisely, up to sign, to the system  $K_h(X)$  corresponds the Chern class  $c_{n-h}(X)$ , where  $n$  is the dimension of  $X$  (and we can consider the Chern classes either as elements of the Chow ring of  $X$ , or as integral cohomology classes).

---

<sup>4</sup>using the embedding covariants for the case of the diagonal  $\Delta_X \subset X \times X$ : this approach was also later followed by Grothendieck.

For this reason we rephrase the questions raised by Baldassarri in terms of Chern classes: Baldassarri dealt with the question of characterizing all the projective varieties  $X$  such that the rational Chern classes  $c_i(X) \in H^*(X, \mathbb{Q})$  vanish for  $n - k + 1 \leq i \leq n$ , but not for  $i = n - k$ .

This question is still wide open, except for the case  $k = n$  (cf. Theorem 1.2). As we have shown, already in the surface case even the condition that all the Chern numbers are zero (this means  $K_X^2 = e(X) = 0$ ) does not imply that  $X$  is a complex torus or a hyperelliptic surface (in the old notation a torus was also called a hyperelliptic surface, of grade, or rank, 1). It implies (see Theorem 2.8) that the surface is either a torus or it is birationally (but not necessarily biregularly) covered by a 1-dimensional family of isomorphic elliptic curves.

Baldassarri's question also suggests (see question (IV) below) to classify, if possible, the varieties (respectively, the cKM)  $X$  whose integral Chern classes  $c_i(X) \in H^*(X, \mathbb{Z})$  vanish for  $n - k + 1 \leq i \leq n$ .

Again, this question is still open, even for the case  $k = n$ , as we saw in Part I [Cat23].

The paper [Bald56] by Baldassarri, suggesting that the Pseudo Abelian Varieties of Roth might be the varieties with such vanishing top real Chern classes, motivated other more general questions. But, in view of what we have seen already in the case of surfaces, one has to include first of all a condition of minimality of  $X$  in a strong sense, for instance that  $K_X$  is nef. Or require only a birational isomorphism with a product (because a  $\mathbb{P}^1$ -bundle over an elliptic curve  $T$  is only birational to  $\mathbb{P}^1 \times T$ ), or that  $X$  be only birational to  $(Y \times T)/G$ , where the action is only free in codimension 1 (as in [HS21a]).

Todd's review of [Bald56] pointed out a wrong intermediate result in the paper, admitting as counterexample the blow up  $X$  of  $\mathbb{P}^3$  with centre a smooth curve of genus 3, which is a regular manifold  $X$  having  $c_3 = 0$ .

Todd's counterexample could somehow be quickly dismissed as only pointing out the need to assume that  $X$  is a minimal manifold, say with  $K_X$  nef, as we already mentioned.

We observed however here in this paper that the flaw is not simply a technical problem, the main claim by Baldassarri that the solutions are Roth's Pseudo-Abelian varieties is incorrect also for minimal manifolds. Because for instance the class of Manifolds isogenous to a  $k$ -torus product, which is a class of solutions to Baldassarri's problem, is (see section 2) a larger class than the class of Roth's Pseudo-Abelian varieties.

Todd saw however that the crucial flaw in Baldassarri's argument was the attempt to show (in dimension at least 3) that manifolds with top Chern class equal to zero possess non zero holomorphic 1-forms.

What is historically interesting for us is to observe that the 'original sin' of [Bald56] was to try to extend to higher dimension some wrong results by Enriques, Dantoni and Roth (indeed the error of Enriques is also reproduced in the classification theorem of Castelnuovo and Enriques [CastEn15]).



Baldassarri and Roth were inspired by a paper by Dantoni [Dant43], devoted to the minimal surfaces  $X$  with  $c_2(X) = 0$ . Dantoni uses surface classification, especially an article by Enriques of 1905 [Enr05b], claiming that the non ruled surfaces with these properties are the hyperelliptic manifolds and the ‘elliptic’ surfaces. But ‘elliptic’ for Enriques here does not have the same standard meaning introduced later by Kodaira and others: Enriques requires the action of a fixed elliptic curve on  $X$ , with all orbits of dimension 1.

Hence Enriques and Dantoni in their classification omit to consider the case of quotients  $X = (E \times C)/G$  where the action of the finite group  $G$  is free, of product type, but  $G$  does not act on the elliptic curve  $E$  via translations, and moreover  $C$  is a curve of genus  $g \geq 2$ , such that the quotient  $C/G$  is an elliptic curve (see section 2, and for instance [CatLi19] or [Bea78] for the special case  $p_g = 0$ , and [CB] or [Cat22] for the general case). Indeed, in this latter case the automorphism group of  $X$  has dimension zero.

Dantoni’s paper inspired Roth [Roth54] [Roth55] who defined the Pseudo-Abelian varieties as the manifolds admitting the action of a complex torus of positive dimension  $= k$  having all orbits of dimension  $k$ , and such that  $k$  is maximal with this property.

But for instance, in [Roth53], Roth does not realize about the existence of Hyperelliptic threefolds with automorphism groups of dimension zero, and believes that Hyperelliptic threefolds can only be Pseudo-Abelian varieties with  $k \geq 1$ .

The first conclusion is simple: the ‘original sin’ was to consider only quotients  $X = (T \times Y)/G$  where  $T$  is a torus,  $G$  acts freely via a product action, and  $G$  acts on  $T$  via translations. Obviously under these assumptions  $T$  acts on  $X$  and the orbits have all the same dimension  $k = \dim(T)$ !

**6.2. Mathematical comments.** Baldassarri’s paper suggested the following other questions:

**Question 6.1.** (I) What can be said about a projective Manifold (respectively, a compact Kähler manifold)  $X$  with  $K_X$  nef and such that all its Chern numbers are equal to zero?

(II) Is a projective Manifold (respectively, a compact Kähler manifold)  $X$  with  $K_X$  nef and such that its rational Chern classes  $c_i(X) \in H^*(X, \mathbb{R})$  vanish for  $k + 1 \leq i \leq n$  isogenous to a  $k$ -Torus Product (respectively, isogenous to a  $k$ -framed or  $k$ -coframed manifold)?

(III) Same question for a projective Manifold (or compact Kähler manifold)  $X$  with  $K_X$  nef and with all the Chern numbers equal to zero.

(IV) What can we say about a projective Manifold (or cKM)  $X$  whose integral Chern classes  $c_i(X) \in H^*(X, \mathbb{Z})$  vanish for  $k + 1 \leq i \leq n$ ?

We shall see in the next section that work of Chad Schoen [Schoen88] gives a negative answer to Questions (II) and (III).

**Remark 6.2.** a) We have mentioned that Baldassarri's claim is wrong already for surfaces.

b) The claim is also wrong in  $\dim = 3$  since here, as shown in Prop. 1.5 of Part I, [Cat23], there are Hyperelliptic Threefolds whose group of Automorphisms is discrete, hence they are not Pseudo-Abelian. Indeed, if one looks at the paper by Roth [Roth53] on Hyperelliptic threefolds, one sees that Roth does not consider the case of Hyperelliptic Threefolds for which  $\text{Aut}^0(X)$  consists only of the Identity.

c) In the same paper Roth calls, following Enriques, [Enr05a], [Enr05b], 'the elliptic surfaces' the surfaces such that  $\text{Aut}^0(X)$  is an elliptic curve.

Here the modern terminology, introduced by Kodaira, differs: an elliptic surface is a surface admitting a fibration  $f : S \rightarrow C$  with fibres elliptic curves; in general it does not possess non-trivial automorphisms.

As we saw in Part I [Cat23], however, there are Hyperelliptic Threefolds and Varieties (hence for them  $c_n(X) = 0$ ) which are regular ( $H^1(\mathcal{O}_X) = 0$ ).

Hence the crucial fact that Baldassarri wants to use, that for  $c_n(X) = 0$  we have an irregular variety possessing a holomorphic 1-form without zeros has to be taken as an assumption.

To give an idea of the difficulty of the type of questions considered by Baldassarri, let us notice that for instance the investigation of varieties such that there is a holomorphic 1-form  $\omega \in H^0(\Omega_X^1)$  without zeros has been taken up (before our present investigations) in the last years by Schreieder and Hao ([Schre21] [HS21b]), and a classification has been achieved only in dimension 3.

What is more interesting is that, under this much stronger assumption of the existence of such a form  $\omega$ , the results confirms, in a special case, the result claimed by Baldassarri (see also our Theorem 5.5).

**Theorem 6.3. (Hao-Schreieder [HS21b])**

*Let  $X$  be a smooth projective threefold satisfying property*

*(A) : admitting a holomorphic 1-form  $\omega \in H^0(\Omega_X^1)$  without zeros.*

*Then the minimal model program for  $X$  yields a birational morphism  $\sigma : X \rightarrow X^{\min}$  blowing up smooth elliptic curves which are not contracted by the Albanese map, and such that*

*(2) there is a smooth morphism  $\pi : X^{\min} \rightarrow A$  to an Abelian variety of positive dimension;*

*(3) If the Kodaira dimension is non negative, then a finite étale cover of  $X^{\min}$  is a product  $X' \cong A' \times S'$ , where  $S'$  is smooth projective, and the composite map  $A' \cong A' \times \{s'\} \rightarrow A$  is finite and étale.*

*(4) If the Kodaira dimension is equal to  $-\infty$ , then either*

*(4a)  $X^{\min}$  has a smooth Del Pezzo fibration over an elliptic curve, or*

*(4b)  $X^{\min}$  has a conic bundle fibration  $f : X^{\min} \rightarrow S$  over a smooth surface  $S$  satisfying property (A). Moreover, either  $f$  is smooth, or  $A$  is smooth and the degeneracy locus of  $f$  is a finite union of elliptic curves which are étale over  $A$ .*

**Corollary 6.4.** *If  $X$  is a smooth projective threefold satisfying property (A) of admitting a holomorphic 1-form  $\omega \in H^0(\Omega_X^1)$  without zeros, and the Kodaira dimension of  $X$  is non negative, then  $X^{\min}$  is a Pseudo-Abelian variety.*

*Proof.* We are in case (3), and obviously  $A'$  acts on the product  $A' \times S'$ . Since the map  $A' \rightarrow A$  is finite and unramified, it is a quotient map, with Galois group  $G$ , a finite Abelian group of translations of  $A'$ . We have  $A' \times S' \rightarrow X^{\min} \rightarrow A$ , and since  $A' \times S' \rightarrow A'$  is a smooth fibration with fibre  $S'$ , hence also  $X^{\min} \rightarrow A$  is a smooth fibration with fibre  $S'$ . Since the pull back of  $X^{\min} \rightarrow A$  to  $A'$  is isomorphic to the product  $A' \times S'$ , we conclude that  $A'$  acts on  $X^{\min}$  and  $X^{\min} = (A' \times S')/G$ , where  $G$  acts on  $A'$  by translations.  $\square$

**Remark 6.5.** Let  $X = (A_1 \times A_2 \times C)/G$ , where  $A_1, A_2$  are elliptic curves,  $G$  acts on  $A_1, A_2, C$ , with a free action on  $A_1$ , while  $A_2/G$  and  $C/G$  have genus 0.

Then  $X$  is a MITP of order  $k = 2$ , but it is a Pseudo-Abelian Variety of order 1, such that  $h^0(\Omega_X^1) = 1$ , and property (A) is satisfied.

The historical conclusion that we can draw is that Baldassarri's paper, even if vitiated by the 'original sin' (trying to extend results which were not correct already in small dimension  $n = 2, 3$ ) poses some problems which, in spite of the tremendous substantial and technical progress which took place in the last 60 or more years, are still open and challenging.

The typical example is the question of describing the  $k$ -co-framed manifolds  $X$  (see Proposition 5.2 and Example 5.7).

## 7. MANIFOLDS WITH VANISHING CHERN NUMBERS

The title of this section is on purpose ambiguous: one may ask about Manifolds for which certain Chern numbers vanish, or all the way consider Manifolds for which all the Chern numbers are equal to zero.

We have given the example of MITP, Manifolds  $X$  Isogenous to a  $k$ -Torus Product, as a prototype of manifolds with all the Chern numbers equal to zero.

We have also observed that, at least in dimension 2, all the surfaces with all the Chern numbers  $c_1^2 = c_2 = 0$ , or the minimal surfaces with  $c_2 = 0$ , in view of Theorem 2.8, are MITP's, if  $K_S$  is nef, or birational to a MITP if  $S$  is ruled.

Because if  $S$  is minimal and not elliptic ruled, there exists a Galois étale covering  $S' \rightarrow S$  such that  $S' \cong T \times Y$ , where  $T$  is a complex torus with  $\dim(T) > 0$ .

More generally, we have the class of manifolds  $X$  isogenous to a partially cotangentially framed manifold  $X$ .

If we go up to dimension 3, there are three Chern numbers,  $c_1^3, c_3$ , and  $c_1 c_2 = \chi(\mathcal{O}_X)$ .

A recent result by Hao and Schreieder goes in the direction of answering the question, [HS21a] in the special case where the Kodaira dimension is  $n - 1$  (extending part (1) of Theorem 1.3):

**Theorem 7.1. (Hao-Schreieder)** *Let  $X$  be a minimal model with  $\dim(X) = n$  and Kodaira dimension  $n - 1$ .*

*Then  $c_1^{n-2}c_2(X) = 0$  if and only if  $X$  is birational to a quotient  $Z = (E \times Y)/G$ , where*

- (1)  $Z$  has canonical singularities;*
- (2)  $E$  is an elliptic curve and  $Y$  is a normal projective variety with  $K_Y$  ample;*
- (3)  $G$  acts diagonally, faithfully on each factor, and freely in codimension two on  $E \times Y$ .*

Now, the case where  $X$  is a threefold of general type (in this case it can be  $c_1c_2 = 0$ , as shown by Ein and Lazarsfeld, [EL97]), is excluded if  $X$  is minimal, since then  $c_1^3 > 0$ .

Hence the missing cases are the cases of Kodaira dimension 0 and 1. For Kodaira dimension 0,  $c_1(X) = 0 \in H^2(X, \mathbb{Q})$ , hence remains to see what happens for  $c_3 = 0$ . Some examples of simply connected Calabi-Yau threefolds with  $c_3 = 0$  have been constructed by Chad Schoen [Schoen88] and other examples were later found by Volker Braun [Bra12] (the latter as hypersurfaces in a toric fourfold).

**7.1. The Schoen Calabi-Yau threefolds  $X$  with  $c_3(X) = 0$ .** We want to discuss now the former examples by Schoen, and show some partial results which seem to indicate that they should not be birational to a quotient of a torus product.

These examples are constructed as small resolutions of fibre products  $X = S_1 \times_{\mathbb{P}^1} S_2$  where  $f_i : S_i \rightarrow \mathbb{P}^1$  is a rational elliptic surface with a section. Let  $f : X \rightarrow \mathbb{P}^1$  be the fibre product of  $f_1$  and  $f_2$ .

We are interested in the special case where the critical values of  $f_1, f_2$  are different; then the fibre product  $X$  is smooth, and all the fibres  $F$  of  $f$  are either a product of two elliptic curves, or the product of an elliptic curve with a degenerate fibre. Hence all the fibres  $F$  have Euler number  $e(F) = 0$ , and  $c_3(X) = e(X) = 0$ .

The canonical divisor of  $X$  is trivial, for instance we can take the  $f_i = \frac{F_i}{G_i}$  to be given by a pencil of plane cubics with simple base points: then  $X$  is the small resolution of a hypersurface of bidegree  $(3, 3)$   $X' \subset \mathbb{P}^2 \times \mathbb{P}^2$ ,

$$X' = \{(x, y) | F_1(x)G_2(y) = F_2(y)G_1(x)\}$$

hence  $X'$  and  $X$  have trivial canonical divisor (see [Schoen88] page 181). Moreover, essentially by the hyperplane theorem of Lefschetz,  $X$  is simply connected (see at any rate (2.1) of [Schoen88], page 181).

Thus  $X$  is a Calabi-Yau threefold with  $c_3(X) = 0$ .

**Proposition 7.2.** *The Schoen threefolds  $X$  (with  $K_X$  trivial and  $c_3(X) = 0$ ) cannot be birationally covered by a 1-dimensional family of subvarieties which are isomorphic to a fixed Abelian surface  $T$ .*

*Proof.* Assume that  $X$  is birationally covered by a family  $T \times C$  (where by the way  $C = \mathbb{P}^1$  since  $q(X) = 0$ ).

For a general  $b \in C$ ,  $T_b := T \times \{b\}$  cannot map to a fibre of  $f : X \rightarrow \mathbb{P}^1$ , since  $f$  is the fibre product of  $f_1$  and  $f_2$  and we may assume that the fibrations  $f_i$  do not have constant moduli.

Hence  $T_b$ , which is a subvariety of  $X$ , dominates  $\mathbb{P}^1$  through the morphism  $f$ .

But a linear system of dimension one on an Abelian surface yields a morphism to  $\mathbb{P}^1$  only if the system is not ample, that is, the fibre is a union of translates of an elliptic curve  $E \subset T$ . Varying  $b$ , the elliptic curve  $E$  is fixed (since it corresponds to a subgroup of the first homology group of  $T$ ), therefore all the fibres of  $f$  contain an elliptic curve isomorphic to  $E$ . This is a contradiction, as the general fibres of  $f_1$  and  $f_2$  are not even isogenous to a fixed elliptic curve  $E$ . □

**Proposition 7.3.** *The Schoen threefolds  $X$  do not admit a fibration  $\psi : X \rightarrow S$  onto a surface  $S$  such that the general fibre is isomorphic to a fixed elliptic curve  $E$ .*

*Proof.* Since  $X$  is simply connected, and the general fibre of  $\psi$  is connected, it follows that  $S$  is also simply connected.

Let  $D$  be the divisorial part of the set of critical values of  $\psi$ , and let  $D^*$  be its smooth locus: define

$$S^* := (S \setminus \text{Sing}(D)), \quad X^* = \psi^{-1}(S^*).$$

Then we have an exact sequence

$$\pi_1(E) \rightarrow \pi_1(X^*) \rightarrow \pi_1(S^*) \rightarrow 1,$$

and we observe that also  $X^*, S^*$  are simply connected, since we have removed a subvariety of real codimension 4.

The image of  $\pi_1(E) \rightarrow \pi_1(X^*)$  is the quotient of  $\pi_1(E)$  by the local monodromies around the irreducible components  $D_j$  of  $D^*$ .

To understand these local monodromies, take a general curve section  $C$  of  $S$ . The inverse image of  $C$  is an elliptic fibration  $\Sigma \rightarrow C$  over  $C$  such that all the smooth fibres are isomorphic to  $E$ .

The fibration is isotrivial, hence  $\Sigma = (C' \times E)/G$ , where  $C' \rightarrow C$  is Galois with group  $G$ .

$G$  acts on  $C' \times E$  via a product action. If there are fixpoints for the action of  $G$  on  $C'$ , then, since the quotient is smooth, it follows that the isotropy subgroups act freely on  $E$ , hence by translations. So the local holomorphic monodromies are translations by torsion points, and the monodromy acts trivially on the first homology of  $E$ .

The conclusion is that the image of  $\pi_1(E) \rightarrow \pi_1(X^*)$  is an infinite group, and this is a contradiction since  $\pi_1(X^*)$  is trivial.  $\square$

The construction of Chad Schoen [Schoen88] applied to other elliptic fibrations leads to threefolds  $X$  with Kodaira dimension 1, and  $c_3(X) = 0$ . We have not yet investigated whether we can achieve with this construction trivial Chern number  $c_1(X)c_2(X) = 0$ .

**Acknowledgements:** I would like to thank Francesco Baldassarri for bringing the paper [Bald56] by Mario Baldassarri to our attention, thus raising my interest in these questions; he, Ciro Ciliberto and Flaminio Flamini provided me with the text of [Dant43]. Thanks to Ciliberto for a quick but useful discussion, and to Thomas Peternell for mentioning the article by Braun.

Many thanks to Matthias Schütt for bringing the examples of [Schoen88] to my attention and explaining their key features.

Thanks to Pierre Deligne for answering an email query (see Remark 5.9).

Thanks to Adriano Tomassini for pointing out Wang's reference [Wang54].

## REFERENCES

- [AMN12] JAUME AMORÓS, MÒNICA MANJARÍN, MARCEL NICOLAU: Deformations of Kähler manifolds with nonvanishing holomorphic vector fields. *J. Eur. Math. Soc. (JEMS)* 14, No. 4, 997–1040 (2012).
- [Ap55] MADHUMALATI APTE: Sur certaines classes caractéristiques des variétés kählériennes compactes. *C. R. Acad. Sci., Paris* 240, 149–151 (1955).
- [At98] FRANCIS ATIYAH: Obituary: John Arthur Todd. *Bull. Lond. Math. Soc.* 30, No. 3, 305–316 (1998).
- [BdF08] GIUSEPPE BAGNERA, MICHELE DE FRANCHIS: Le superficie algebriche le quali ammettono una rappresentazione parametrica mediante funzioni iperellittiche di due argomenti, *Mem. di Mat. e di Fis. Soc. It. Sc. (3)* 15, 253–343 (1908).
- [Bald56] MARIO BALDASSARRI: Una caratterizzazione delle varietà abeliane e pseudoabeliane. *Ann. Mat. Pura Appl. (4)* 42 (1956), 227–252.
- [Bald56book] MARIO BALDASSARRI: *Algebraic Varieties*. *Ergebnisse der Mathematik und ihrer Grenzgebiete*, vol.12, Springer Berlin (1956).
- [BCF15] INGRID BAUER, FABRIZIO CATANESE, DAVIDE FRAPPORTI: Generalized Burniat type surfaces and Bagnera-De Franchis varieties. *J. Math. Sci. Univ. Tokyo* 22 (2015), no. 1, 55–111.
- [Bea78] ARNAUD BEAUVILLE: Surfaces algébriques complexes. *Astérisque* 54, 1–172 (1978).
- [Bra12] VOLKER BRAUN: The 24-cell and Calabi-Yau threefolds with Hodge numbers (1,1). *J. High Energy Phys.* 2012, no. 5, 101, front matter+19 pp.
- [C-H-P16] FRÉDÉRIC CAMPANA, ANDREAS HÖRING, THOMAS PETERNELL: Abundance for Kähler threefolds. *Ann. Sci. Éc. Norm. Supér. (4)* 49, No. 4, 971–1025 (2016).
- [CastEn15] GUIDO CASTELNUOVO, FEDERIGO ENRIQUES: Die algebraischen Flächen vom Gesichtspunkte der birationalen Transformation aus. *Enz. d. Math. Wiss. III C* 6 b, 674–768 (Band III 2, Heft 6) (1915) (*Memorie Scelte* vol. 3, pp. 197–296).
- [CatCil93] FABRIZIO CATANESE, CIRO CILIBERTO: On the irregularity of cyclic coverings of algebraic surfaces. *Geometry of complex projective varieties (Cetraro, 1990)*, 89–115, *Sem. Conf.*, 9, Mediterranean, Rende, (1993).



- [Cat08] FABRIZIO CATANESE: Differentiable and deformation type of algebraic surfaces, real and symplectic structures. CIME Course: Symplectic 4-manifolds and algebraic surfaces, Springer Lecture Notes in Math. 1938 (2008) 55–167.
- [CB] FABRIZIO CATANESE, INGRID BAUER: *ETH Lectures on algebraic surfaces*, preliminary version, 160 pages, (2009).
- [Cat22] FABRIZIO CATANESE: The  $P_{12}$ -Theorem: the Classification of surfaces and its historical development, Chapter I of the book : Curves and Surfaces, a history of Shapes (2022).
- [Cat23] FABRIZIO CATANESE: Manifolds with trivial Chern classes I: Hyperelliptic Manifolds and a question by Severi (2023), to appear in the Annali Scuola Normale Sup. Pisa.
- [CC17] FABRIZIO CATANESE, PIETRO CORVAJA: Teichmüller spaces of generalized hyperelliptic manifolds. Complex and symplectic geometry, 39-49, Springer INdAM Ser., 21, Springer, Cham (2017).
- [CatLi19] FABRIZIO CATANESE, BINRU LI: Enriques’ classification in characteristic  $p > 0$ : the  $P_{12}$ -theorem. Nagoya Math. J. 235, 201–226 (2019).
- [CatLiu21] FABRIZIO CATANESE, WENFEI LIU: On topologically trivial automorphisms of compact Kähler manifolds and algebraic surfaces. Atti Accad. Naz. Lincei, Cl. Sci. Fis. Mat. Nat., IX. Ser., Rend. Lincei, Mat. Appl. 32, No. 2, 181–211 (2021).
- [Dant43] GIOVANNI DANTONI: Determinazione delle superficie con serie di Severi di ordine nullo o negativo. Atti Accad. Italia, Mem., Cl. Sci. Fis. Mat. Nat. 14, 39–49 (1943).
- [EL97] LAWRENCE EIN, ROBERT LAZARSFELD: Singularities of theta divisors and the birational geometry of irregular varieties. J. Am. Math. Soc. 10, No. 1, 243–258 (1997).
- [Enr05a] FEDERIGO ENRIQUES Sulle superficie algebriche che ammettono un gruppo continuo di trasformazioni birazionali in se stesse. Palermo Rend. 20, 61–72 (1905).
- [Enr05b] FEDERIGO ENRIQUES Sulle superficie algebriche di genere geometrico zero. Palermo Rend. 20, 1–33 (1905).
- [EnrSev09] FEDERIGO ENRIQUES, FRANCESCO SEVERI Mémoire sur les surfaces hyperelliptiques. Acta Math. 32 (1909), no. 1, 283–392, and Acta Math. 33 (1910), no. 1, 321–403.
- [Fuj78] AKIRA FUJIKI: On automorphism groups of compact Kähler manifolds. Invent. Math. 44 (1978), no. 3, 225–258.
- [Ful84] WILLIAM FULTON: Intersection theory. Ergebnisse der Mathematik und ihrer Grenzgebiete, 3. Folge, Bd. 2. Berlin etc.: Springer-Verlag. xi, 470 p. (1984).
- [HS21a] FENG HAO, STEFAN SCHREIEDER: Equality in the Bogomolov-Miyaoka-Yau inequality in the non-general type case. J. Reine Angew. Math. 775 (2021), 87–115.
- [HS21b] FENG HAO, STEFAN SCHREIEDER: Holomorphic one-forms without zeros on threefolds, Geom. Topol. 25 (2021), 409–444.
- [Kob87] SHOSHICHI KOBAYASHI: Differential geometry of complex vector bundles. Publications of the Mathematical Society of Japan, 15; Kano Memorial Lectures, 5. Princeton, NJ: Princeton University Press; Tokyo: Iwanami Shoten Publishers. xi, 304 p. (1987).
- [Kod68] KUNIIHIKO KODAIRA: On the structure of compact complex analytic surfaces. III. Amer. J. Math. 88 (1968) 55–83.
- [Flips92] KOLLÁR, JÁNOS ET AL. Flips and abundance for algebraic threefolds. A summer seminar at the University of Utah, Salt Lake City, 1991. Astérisque. 211. Paris: Société Mathématique de France, 258 p. (1992).
- [Kov97] SÁNDOR J. KOVÁCS: Families over a base with a birationally nef tangent bundle. Math. Ann. 308, No. 2, 347–359 (1997).



- [Lai11] C.-J. LAI: Varieties fibered by good minimal models, *Math. Ann.* 350 (2011), no. 3, 533–547.
- [La01] HERBERT LANGE: Hyperelliptic varieties. *Tohoku Math. J. (2)* 53 (2001), no. 4, 491–510.
- [Li78] DAVID I. LIEBERMAN: Compactness of the Chow scheme: applications to automorphisms and deformations of Kähler manifolds. *Fonctions de plusieurs variables complexes, III (Sém. Francois Norguet, 1975â–1977)*, pp. 140–186, *Lecture Notes in Math.*, 670, Springer, Berlin, 1978.
- [Li77] DAVID I. LIEBERMAN: Holomorphic vector fields on projective varieties. *Several complex Variables, Proc. Symp. Pure Math. 30, Part 1, Williamstown 1975*, 273–276 (1977).
- [Lu-Ta18] STEVEN LU, BEHROUZ TAJI: A characterization of finite quotients of abelian varieties. *Int. Math. Res. Not.* 2018, No. 1, 292–319 (2018).
- [Mar71] MASAKI MARUYAMA: On automorphism groups of ruled surfaces. *J. Math. Kyoto Univ.* 11 (1971), 89–112.
- [Miya87] YOICHI MIYAOKA: The Chern classes and Kodaira dimension of a minimal variety. *Algebraic geometry, Proc. Symp., Sendai/Jap. 1985, Adv. Stud. Pure Math.* 10, 449–476 (1987).
- [Mum70] DAVID MUMFORD: Abelian varieties. *Studies in Mathematics. Tata Institute of Fundamental Research 5. London: Oxford University Press.* VIII, 242 p. (1970), with notes by C. P. Ramanujam.
- [Nak55] SHIGEO NAKANO: Tangential vector bundle and Todd canonical systems of an algebraic variety. *Mem. Coll. Sci. Univ. Kyoto, Ser. A* 29, 145–149 (1955).
- [Roth53] LEONARD ROTH: Hyperelliptic threefolds. *Proc. Camb. Philos. Soc.* 49, 397–409 (1953).
- [Roth54] LEONARD ROTH: Pseudo-Abelian varieties. *Proc. Camb. Philos. Soc.* 50, 360–371 (1954).
- [Roth55] LEONARD ROTH: Some properties of pseudo-Abelian varieties. *Ann. Mat. Pura Appl., IV. Ser.* 38, 281–302 (1955).
- [Roth56] LEONARD ROTH: Sistemi canonici ed anticanonici. *Univ. Genova, Pubbl. Ist. Mat.* 21, 118 p. (1956).
- [Schre21] STEFAN SCHREIEDER: Zeros of holomorphic one-forms and topology of Kähler manifolds. *Int. Math. Res. Not. IMRN* 2021, no. 8, 6169–6183.
- [Schoen88] CHAD SCHOEN: On fiber products of rational elliptic surfaces with section. *Math. Z.* 197, No. 2, 177–199 (1988).
- [Seg52] BENIAMINO SEGRE: Variétés covariantes d ’ immersion et variétés canoniques sur une variété algébrique ou topologique. *C. R. Acad. Sci., Paris* 234, 1731–1733 (1952).
- [Seg54] BENIAMINO SEGRE: Dilatazioni e varietà canoniche sulle varietà algebriche. *Ann. Mat. Pura Appl., IV. Ser.* 37, 139–155 (1954).
- [Sev51] FRANCESCO SEVERI: Fondamenti per la geometria sulle varietà algebriche. II. *Ann. Mat. Pura Appl., IV. Ser.* 32, 1–81 (1951).
- [Som74] ANDREW JOHN SOMMESE: Holomorphic vector-fields on compact Kähler manifolds. *Math. Ann.* 210, 75–82 (1974).
- [Taj21] BEHROUZ TAJI: On the Kodaira dimension of base spaces of families of manifolds. *J. Pure Appl. Algebra* 225, No. 11, Article ID 106729, 21 p. (2021).
- [Taj23] BEHROUZ TAJI: Birational geometry of smooth families of varieties admitting good minimal models. *Eur. J. Math.* 9, No. 4, Paper No. 88, 44 p. (2023).
- [Yau78] SHING-TUNG YAU: On the Ricci curvature of a compact Kähler manifold and the complex Monge-Ampère equation. I. *Commun. Pure Appl. Math.* 31, 339–411 (1978).
- [Wang54] HSIEN-CHUNG WANG: Complex parallelisable manifolds. *Proc. Am. Math. Soc.* 5, 771–776 (1954).

- [Wav69] WAVRIK, JOHN J. Obstructions to the existence of a space of moduli. Global Analysis, Papers in Honor of K. Kodaira, Princeton University Press, 403–414 (1969).
- [W-W23] CHUANHAO WEI, LEI WU: Isotriviality of smooth families of varieties of general type. Manuscr. Math. 172, No. 1-2, 139–168 (2023).

MATHEMATISCHES INSTITUT DER UNIVERSITÄT BAYREUTH, NW II, UNIVERSITÄTSSTR.  
30, 95447 BAYREUTH  
*Email address:* `fabrizio.catanese@uni-bayreuth.de`

KOREA INSTITUTE FOR ADVANCED STUDY, HOEGIRO 87, SEOUL, 133–722.