

ON THE FOLKLORE SET AND DIRICHLET SPECTRUM FOR MATRICES

MUMTAZ HUSSAIN, JOHANNES SCHLEISCHITZ, AND BENJAMIN WARD

ABSTRACT. We study the Folklore set of Dirichlet improvable matrices in $\mathbb{R}^{m \times n}$ which are neither singular nor badly approximable. We prove the non-emptiness for all positive integer pairs m, n apart from $\{m, n\} = \{1, 1\}$ and $\{m, n\} = \{2, 3\}$ in a constructive way. For a wide range of integer pairs (m, n) we construct subsets of the Folklore set with an exact prescribed Dirichlet constant (in some right neighbourhood of 0). This enables us to provide information on the Dirichlet Spectrum of matrices. The key technique of our construction is to build certain 2-dimensional simultaneously approximable very singular vectors, and then ‘lift’ these vectors to higher dimensions by ‘glueing’ them to a large set of well-behaved matrices. Our technique is also applicable to arbitrary norms. As a corollary we obtain lower bounds on the Hausdorff dimension of these sets. These statements complement previous results of the middle-named author (Selecta Math. 2023), Beresnevich et. al. (Adv. Math. 2023), and Das et. al. (Adv. Math. 2024).

Keywords: Dirichlet spectrum, Folklore set, Uniform approximation, Singular sets, linear forms, Hausdorff dimension.

Math Subject Classification 2010: 11J13, 11J82, 11J83, 11K60

1. UNIFORM APPROXIMATION AND THE FOLKLORE SET

1.1. Dual approximation. Dirichlet’s approximation theorem for dual linear forms states that for any $\boldsymbol{\xi} = (\xi_1, \dots, \xi_n) \in \mathbb{R}^n$ and any $N \in \mathbb{N}$ the system

$$(1) \quad \begin{aligned} |\hat{\mathbf{b}} \cdot \boldsymbol{\xi} + b_{n+1}| &:= |b_1 \xi_1 + \dots + b_n \xi_n + b_{n+1}| < N^{-n} \\ \|\hat{\mathbf{b}}\| &:= \max_{1 \leq i \leq n} |b_i| \leq N \end{aligned}$$

has non-zero integer solution $\mathbf{b} = (\hat{\mathbf{b}}, b_{n+1}) \in \mathbb{Z}^{n+1}$. For fixed $0 < c < 1$ the set of *c-Dirichlet improvable numbers*, denoted as $Di_n(c)$, are those for which the right-hand side of (1) can be “improved” by c . That is,

$$Di_n(c) := \left\{ \boldsymbol{\xi} \in \mathbb{R}^n : \begin{cases} |\hat{\mathbf{b}} \cdot \boldsymbol{\xi} + b_{n+1}| < cN^{-n} \\ \|\hat{\mathbf{b}}\| \leq N \end{cases} \text{ is soluble for all } N \geq N_0(c, \boldsymbol{\xi}) \right\}.$$

Generally $\boldsymbol{\xi} \in \mathbb{R}^n$ is said to be *Dirichlet improvable*, if there exists some $0 < c < 1$ for which $\boldsymbol{\xi} \in Di_n(c)$, and $\boldsymbol{\xi} \in \mathbb{R}^n$ is said to be *Singular*, if it is *c-Dirichlet improvable* for all $c > 0$. Concisely

$$Di_n := \bigcup_{0 < c < 1} Di_n(c), \quad \text{and} \quad Sing_n := \bigcap_{0 < c < 1} Di_n(c).$$

Trivially, for any $0 < c < 1$, we have that $Sing_n \subseteq Di_n(c) \subseteq Di_n$.

In another direction, from Dirichlet's approximation theorem it is immediate that for any $\boldsymbol{\xi} \in \mathbb{R}^n$

$$(2) \quad |\hat{\mathbf{b}} \cdot \boldsymbol{\xi} + b_{n+1}| < \|\hat{\mathbf{b}}\|^{-n}$$

has infinitely many integer solutions $\mathbf{b} \in \mathbb{Z}^{n+1}$. The set of *Badly approximable vectors*, denoted as Bad_n , consists of the set of $\boldsymbol{\xi} \in \mathbb{R}^n$ such that the right-hand side of (2) cannot be "improved" by an arbitrarily small constant, otherwise $\boldsymbol{\xi}$ is called well approximable. That is,

$$Bad_n := \left\{ \boldsymbol{\xi} \in \mathbb{R}^n : \text{There exists } \varepsilon > 0 : |\hat{\mathbf{b}} \cdot \boldsymbol{\xi} + b_{n+1}| \geq \varepsilon \|\hat{\mathbf{b}}\|^{-n} \text{ for all } \mathbf{b} \in \mathbb{Z}^{n+1} \setminus \{\mathbf{0}\} \right\}.$$

We have the relations

$$Bad_n \subseteq Di_n \quad \text{and} \quad Bad_n \cap Sing_n = \emptyset.$$

The left was shown by Davenport and Schmidt [14, Theorem 2], the right is obvious. It is well known that all of the sets highlighted above are n -dimensional Lebesgue nullsets. This can be proven through a variety of methods, see for example [5] for an overview. So to give any meaningful notion of size to these sets we consider the Hausdorff dimension, denoted \dim_H . See [15] for the definition and properties of the Hausdorff dimension. The following results on Di_n , $Sing_n$, and Bad_n have been proven:

$$\dim_H Bad_n = n, \quad \text{for } n = 1 \text{ in [17] and for } n \geq 2 \text{ in [29],}$$

$$\dim_H Di_n = n, \quad [29] \text{ combined with [14, Theorem 2],}$$

$$\dim_H Sing_n = \frac{n^2}{n+1}, \quad \text{for } n = 2 \text{ in [8] and for } n \geq 2 \text{ in [9].}$$

The dimensions of $Di_n(c)$ are not yet established, but given $n \geq 2$ and any constant $C > 0$, for $c > 0$ sufficiently small and for any real number $t > n$ the bounds

$$\frac{n^2}{n+1} + c^t \leq \dim_H Di_n(c) \leq \frac{n^2}{n+1} + C \cdot c^{n/2}$$

were proven in [9, Theorem 1.3]. See also [9, Corollary 6.12] for an improvement on the lower bound.

The reader will notice that the dimension results for $Sing_n$ and $Di_n(c)$ hold only for $n \geq 2$. For $n = 1$ it was proven by Khintchine [18] that $Sing_1 = \mathbb{Q}$, in fact $Di_1(c) = \mathbb{Q}$ for any $c < \frac{1}{2}$. By combining this with Theorem 1 of Davenport-Schmidt [14], we obtain

$$Di_1 = \mathbb{Q} \cup Bad_1.$$

In higher dimensions much less was known, until the recent paper of Beresnevich et. al. [6], who showed that, for $n \geq 2$, the *Folklore set*

$$FS_n := Di_n \setminus (Bad_n \cup Sing_n)$$

has a continuum of points. They mention that this question was first asked in [31] (see the remark immediately following [31, Theorem 4.6]) and furthermore conjecture that, for $n \geq 2$, we should have $\dim_H FS_n = n$ [6, Problem 3.1].

In the latest (published) version of the paper by Das et al. [12], a variational principle was developed that can be applied to obtain a lower bound

$$(3) \quad \dim_H FS_n \geq \dim_H \text{Sing}_n = n - 1 + \frac{1}{n+1}.$$

See [12, Theorem 3.15]. The previous iteration of the variational principle had not been immediately applicable to the Folklore set, thus not yet implying [6], see [6, Remark 4.4]. At the time of preparing this manuscript the authors were not aware of the recent advances in [12]. Our proof technique is different and more constructive in nature. In particular we are able to prescribe exact Dirichlet constants, even for arbitrary norms, in certain cases. We have attempted to make clear throughout where the bound (3), and more generally (7) below, improve on our statements and where our results are new. Note that (3) is still not the conjectured (full) dimension, which so far remains elusive.

In this article, we prove a range of results that extend the theorems given in [6], and complement [12], mainly in two different ways:

- In Section 2 we consider the Folklore set in the dual approximation setting and add an additional condition that the point has some specific Dirichlet constant c . We prove this set has Hausdorff dimension at least $n - 2$ for all constants c in some right neighbourhood of 0. In fact we prove a refined version involving arbitrary norms. In particular the Dirichlet spectrum (to be defined below) in such a setting contains a proper interval starting at 0.
- In Section 3 we consider the Folklore set in the setting of systems of linear forms. In almost all cases of m, n , we prove non-emptiness (Theorem 3.1) and in most of these cases positive Hausdorff dimension (Theorems 3.3, 3.4, 3.6, 3.7). Again, in several cases we can describe any small enough positive exact Dirichlet constant, even when considering a setup with arbitrary norms.

Prior to stating these results we introduce some more notation, formulated in the general matrix setting and for arbitrary norms.

1.2. Systems of linear forms and arbitrary norms. Let $n, m \in \mathbb{N}$, and let $\|\cdot\|_1$ be any norm on \mathbb{R}^n and $\|\cdot\|_2$ any norm on \mathbb{R}^m . For a given real $m \times n$ matrix $\Omega \in \mathbb{R}^{m \times n}$ and a parameter $t \geq 1$ define the piecewise constant approximation function

$$\psi_\Omega^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t) = \min_{\substack{\mathbf{b}=(\hat{\mathbf{b}}, \tilde{\mathbf{b}}) \in \mathbb{Z}^{n+m}: \\ 1 \leq \|\hat{\mathbf{b}}\|_1 \leq t}} \left\| \Omega \cdot \hat{\mathbf{b}} + \tilde{\mathbf{b}} \right\|_2 = \min_{\substack{\mathbf{b}=(\hat{\mathbf{b}}, \tilde{\mathbf{b}}) \in \mathbb{Z}^{n+m}: \\ 1 \leq \|\hat{\mathbf{b}}\|_1 \leq t}} \left\| \sum_{i=1}^n \Omega_i b_i + \tilde{\mathbf{b}} \right\|_2,$$

where Ω_i denotes the i th column of the matrix Ω , and

$$(4) \quad \hat{\mathbf{b}} = (b_1, \dots, b_n), \quad \tilde{\mathbf{b}} = (b_{n+1}, \dots, b_{n+m}), \quad \mathbf{b} = (\hat{\mathbf{b}}, \tilde{\mathbf{b}})$$

interpreted as column vectors. See works of Lagarias [21, 22] and the survey articles [24, 10] for the history and further details on best approximations in dimensions greater than 1.

Define the Dirichlet constant of a matrix as

$$\Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) = \limsup_{t \rightarrow \infty} t^{n/m} \psi_{\Omega}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t),$$

and further let

$$\tilde{\Theta}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) = \liminf_{t \rightarrow \infty} t^{n/m} \psi_{\Omega}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t).$$

The values $\Theta^{m \times n}$ and $\tilde{\Theta}^{m \times n}$ compare the approximation quality with the standard Dirichlet function $t^{-n/m}$ in the uniform and asymptotic settings, respectively. By equivalence of norms and Dirichlet's Theorem, we derive

$$(5) \quad D = D^{m \times n}(\|\cdot\|_1, \|\cdot\|_2) := \sup_{\Omega \in \mathbb{R}^{m \times n}} \Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) < \infty.$$

For $\|\cdot\|_1$ and $\|\cdot\|_2$ the standard maximum norms, we have $D = 1$, with the upper estimate coming from Minkowski's Theorem for systems of linear forms and this value being attained for Lebesgue almost all matrices by well-known metrical results, see [2, 27] for further details on the Dirichlet spectrum. For $0 \leq c \leq D$, define the following sets

$$\begin{aligned} Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) &:= \{\Omega \in \mathbb{R}^{m \times n} : \Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) \leq c\}, \\ Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2) &:= \{\Omega \in \mathbb{R}^{m \times n} : \Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) < D\} \\ &= \bigcup_{0 < c < D} Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c), \\ Sing_{m,n} = Sing_{m,n}(\|\cdot\|_1, \|\cdot\|_2) &:= \{\Omega \in \mathbb{R}^{m \times n} : \Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) = 0\} \\ &= \bigcap_{0 < c < D} Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c), \\ Bad_{m,n} = Bad_{m,n}(\|\cdot\|_1, \|\cdot\|_2) &:= \{\Omega \in \mathbb{R}^{m \times n} : \tilde{\Theta}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) > 0\}. \end{aligned}$$

Indeed, the singular and bad sets do not depend on the norms by equivalence of norms. Moreover it is known and follows from Mahler's Dual Convex Body Theory that

$$Bad_{m,n} = Bad_{n,m}, \quad \text{and} \quad Sing_{m,n} = Sing_{n,m}.$$

The Hausdorff dimensions of $Bad_{m,n}$ and $Di_{m,n}$ are again full [30, 14], while $\dim_H(Sing_{m,n}) = mn(1 - \frac{1}{m+n})$, as was recently proven in [12, Theorem 1.1].

Additional to these sets, for any $0 \leq c < D$, define the pairwise disjoint sets

$$DI_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) := \{\Omega \in \mathbb{R}^{m \times n} : \Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) = c\}.$$

Then $DI_{m,n}(\|\cdot\|_1, \|\cdot\|_2, 0) = Sing_{m,n}$. Note further that

$$(6) \quad DI_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \subseteq Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \subseteq Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2).$$

Given this setup we define the Folklore set on $m \times n$ matrices with arbitrary norms as

$$FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2) = Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2) \setminus (Bad_{m,n} \cup Sing_{m,n})$$

and, for any $0 < c < D$, the smaller set

$$FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) = DI_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \setminus Bad_{m,n}.$$

When the norms are the maximum norms we just denote the sets by $FS_{m,n}$ and $FS_{m,n}(c)$ respectively. In this setting, generalizing (3), it is proved in [12, Theorem 3.15] that

$$(7) \quad \dim_H FS_{m,n} \geq \dim_H \text{Sing}_{m,n} = mn - \frac{mn}{m+n}.$$

Prior to stating our metrical results we give the following lemma, which gives a nice relationship between a matrix of $FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c)$ and its transpose.

Lemma 1.1. *Given m, n , there exists $\delta = \delta_{m,n} > 0$ so that if $\Omega \in FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c)$ for some $0 < c < \delta$, then its transpose Ω^T satisfies $\Omega^T \in FS_{n,m}(\|\cdot\|_2, \|\cdot\|_1)$.*

We will provide a proof at the end in Section 9 but should note it is a simple consequence of German's transference principle [16, Theorem 7]. Note when considering the transpose we lose control over the precise Dirichlet constant.

It is worth mentioning that a different setup for defining normed Dirichlet improvability, with respect to a single arbitrary norm on \mathbb{R}^{m+n} , has been recently studied, see for example [4, 20]. It is unclear to us how the two setups are precisely related, in particular, if our results remain valid in the single norm setting.

2. ON THE FOLKLORE SET IN THE DUAL APPROXIMATION SETTING

2.1. Maximum norm. Let us, for now, return to our previous setting of classical dual approximation and maximum norm, that is $m = 1$, $\|\cdot\|_1 = \|\cdot\|_\infty = \max_{1 \leq i \leq n} |\cdot|$, and $\|\cdot\|_2 = |\cdot|$. For ease of notation, for $c \in (0, 1)$ we write

$$DI_n(c) := DI_{1,n}(\|\cdot\|_\infty, |\cdot|, c), \quad FS_n(c) := DI_n(c) \setminus (\text{Bad}_n \cup \text{Sing}_n).$$

It was shown in [26] by the second name author that the set $FS_n(c)$ is non-empty for any $c \in (0, 1)$, refining the aforementioned result from [6] and also complementing [12].

Our new result for the vector case and maximum norm is the following.

Theorem 2.1. *For any natural number $n \geq 4$, there exists explicitly computable $c_n \in (0, 1)$ such that for any $c \in (0, c_n)$ we have*

$$\dim_H FS_n(c) \geq n - 2.$$

Remark 1. We believe the claim holds for $n = 3$ as well, probably improving on the bound that can be obtained from the methods in [27, 26]. However, problems occur in our proof in this case stemming from auxiliary results from [25], leaving this case open.

Remark 2. In [27] a lower bound of weaker order $(3/8)n + o(n)$ as $n \rightarrow \infty$ is shown for the corresponding sets with respect to simultaneous approximation for any $c \in (0, 1)$. In [26] it is shown that $FS_n(c)$ for a linear form as defined above is not empty for any $c \in (0, 1)$, and reference to [27] is given on how to obtain metrical results in a similar way. However, again these bounds will turn out considerably weaker than in Theorem 2.1 above. The bound of Theorem 2.1 can in fact be further improved a little with some effort, see Remark 15 in Section 5.1.

While the method of the proof of [12, Theorem 3.15] shows that any set

$$Di_n(c) \setminus (Bad_n \cup Sing_n) \supseteq FS_n(c), \quad c \in (0, 1)$$

has Hausdorff dimension bounded from below as in (3) no matter how small c is, the Dirichlet constant cannot be fixed in their work. So our Theorem 2.1 that permits control over the precise Dirichlet constant, at least in some interval, is new. We may take

$$c_n = \frac{n-3}{8(n-2)^{3/2}} \cdot \sqrt{\pi} \cdot \frac{\Gamma(n-\frac{1}{2})}{\Gamma(n)}$$

with Γ the Euler Gamma function. As $c \rightarrow 0^+$, for the Hausdorff dimension we cannot hope for anything better than $n-1 + \frac{1}{n+1}$ the value in (3), as Cheung and Chevallier [9] showed this as the asymptotics for the Hausdorff dimension of the larger set $Di_n(c) \supseteq FS_n(c)$. Hence we consider our bound reasonably good. Similar to [27, 26] the expected increase of the Hausdorff dimension of $FS_n(c)$ as a function of $c \in [0, 1]$ is not reflected in Theorem 2.1.

Using [6, Proposition 1.1 and Lemma 4.9] we infer the following corollary towards the original conjecture [6, Problem 3.1]. Denote the Folklore set for simultaneous approximation to n numbers by $FS_n^* = FS_{n,1}^*$ using notation of Section 1.2.

Corollary 2.2. *Let $n \geq 4$. Then we have*

$$\dim_H FS_n \geq n-2, \quad \dim_H FS_n^* \geq n-2.$$

This result is stated more for sake of completeness as the metrical bound is weaker than (3) proved in [12]. However our bound is obtained by a constructive method and does not use the variational principle.

Remark 3. We are not able to deduce an equivalent claim for any set $FS_n^*(c) := FS_{n,1}^*(c)$ with $c > 0$. The reason is that the sets $FS_n^*(c)$ do not coincide with $FS_n(c)$ for $c > 0$, indeed Lemma 1.1 is too weak. On the other hand, similar to [6] or [26, Section 11] it allows for deducing non-emptiness (Hausdorff dimension $\geq n-2$ again) for sets

$$Di_n^*(c) \setminus (Bad_n^* \cup Di_n^*(c')) = Di_{n,1}(c) \setminus (Bad_{n,1} \cup Di_{n,1}(c'))$$

where suitable $c' \in (0, c)$ can be explicitly computed as a function of $n \geq 2, c \in (0, 1)$. An analogous claim, with stronger Hausdorff dimension estimate as in (7) and a different (most likely smaller) constant c' , can be obtained from the uniform version of the variational principle in [12] as well.

2.2. Arbitrary norms. Given any norm $\|\cdot\| = \|\cdot\|_1$ on \mathbb{R}^n and $c > 0$, let us write

$$DI_n(\|\cdot\|, c) := DI_{1,n}(\|\cdot\|, |\cdot|, c), \quad FS_n(\|\cdot\|, c) := DI_n(\|\cdot\|, c) \setminus (Bad_n \cup Sing_n).$$

Our second result extends Theorem 2.1 to arbitrary norms.

Theorem 2.3. *Let $n \geq 4$ and $\|\cdot\|$ be any norm on \mathbb{R}^n . Then there is $c_n = c_n(\|\cdot\|) > 0$ such that for any $c \in [0, c_n]$ we have*

$$\dim_H FS_n(\|\cdot\|, c) \geq n-2.$$

In particular the Dirichlet spectrum

$$\mathcal{D}_n(\|\cdot\|) := \{\Theta(\|\cdot\|, \boldsymbol{\xi}) : \boldsymbol{\xi} \in \mathbb{R}^n\} := \{\Theta^{1 \times n}(\|\cdot\|, |\cdot|, \boldsymbol{\xi}) : \boldsymbol{\xi} \in \mathbb{R}^n\} \subseteq [0, D]$$

with D from (5), contains some right neighborhood of 0.

Clearly this result is again independent from the work of Das et al [12].

Remark 4. Since $FS_n(\|\cdot\|, c) \subseteq FS_n(\|\cdot\|)$ the lower bound on the Hausdorff dimension applies to $FS_n(\|\cdot\|)$ as well, which is however weaker than (3).

Remark 5. For $n = 3$ and a general norm, even non-emptiness of the set $FS_n(\|\cdot\|, c)$ for a given $c \in (0, D)$, with D as in (5), remains unknown. For certain norms, this can be obtained from the method in [26]. Again we expect the Hausdorff dimension to rise as a function of $c \in [0, D]$, not reflected in the theorem.

Remark 6. For results on the Dirichlet spectrum with respect to the maximum norm see [27, Appendix] and more completely [26]. In [26] it is proven that the spectrum is the entire unit interval, i.e. $\mathcal{D}_n(\|\cdot\|_\infty) = [0, 1]$. Theorem 2.3 tells us the weaker statement that the spectrum contains some small interval starting at the origin, however, our result holds for any norm. Notice Theorem 2.3 is for the dual approximation setting. For results on the Dirichlet spectrum in the simultaneous approximation setting for various norms on \mathbb{R}^2 see the recent articles [1, 19] for the current state of the art. In short, it was shown in [19, Corollary 1.2] that the corresponding upper bound of $D = D^{2 \times 1}(|\cdot|, \|\cdot\|_2)$, as defined in (5), is an accumulation point for any norm on \mathbb{R}^2 , and in [1] it is proven that for any arbitrary norm the Dirichlet spectrum contains an interval starting at the origin, see [1, Corollary 1]. Both of these results are for simultaneous approximation on \mathbb{R}^2 . We should stress that results for specific norms were known prior to the statements of [1, 19], see for example [2, 3, 27].

Remark 7. By Lemma 1.1, from Theorem 2.3 we may deduce an analogous metrical result to Corollary 2.2 on the simultaneous approximation Folklore set for an arbitrary norm and $m \geq 4$. This will be covered by the more general Theorem 3.7 below.

3. THE FOLKLORE SET FOR A SYSTEM OF LINEAR FORMS

We now come to the matrix setting $\min\{m, n\} > 1$.

3.1. Two fundamental results for maximum norm. In this section we state some results for the maximum norm that can be deduced from our more general results following in the subsequent sections. Our first result below on the Folklore set is (mostly) weaker than the claims in [12]. However, we still want to highlight it, as it will be a consequence of our considerably more general results below that indeed do not follow from work of Das et al [12].

Theorem 3.1. *Let m, n be positive integers satisfying*

$$\{m, n\} \notin \{\{1, 1\}, \{2, 3\}\}.$$

Then

$$FS_{m,n} = FS_{m,n}(\|\cdot\|_\infty, \|\cdot\|_\infty) \neq \emptyset.$$

In fact

$$\dim_H FS_{m,n} \geq mn - 2 \max\{m, n\},$$

and all claims hold when we choose the involved matrices of full \mathbb{R} -rank $\min\{m, n\}$.

Remark 8. Regarding the metrical claim on the dimension appearing in Theorem 3.1, we state in advance that it may be conjectured that

$$\dim_H FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2) = mn,$$

and more generally that

$$\dim_H FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) = mn - o(1)$$

as $c \rightarrow D^-$ for D from (5), for any pair of norms. However, as $c \rightarrow 0^+$, we cannot expect the limit of the Hausdorff dimension of $Di_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \supseteq FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c)$ to exceed the value from (7).

Our result follows from a combination of Theorems 3.5, 3.7 below combined with the cases $m = 1, n \geq 2$ and $m \geq 2, n = 1$ already settled in [27, 26]. The metrical result is non-trivial as soon as $\min\{m, n\} \geq 3$.

We stress again that Theorem 3.1 as such is, up to the claim on the rank, implied by the stronger bound (7) from [12] which also applies to the cases $\{m, n\} = \{2, 3\}$ we miss. However, we emphasize that considerably refined information is provided by the partial results formulated below. The refinements will provide us with information on one or more of the following properties: stronger Hausdorff dimension results in several cases; arbitrary norms; more information on the Dirichlet spectrum, ideally imposing an exact Dirichlet constant akin to Theorem 2.3; and \mathbb{Q} -linear independence of matrix entries together with $\{1\}$. The exact claims we obtain depend on the dimensions m, n of the matrix.

Regarding the full rank claim, since the Hausdorff dimension of rank deficient real $m \times n$ matrices equals $mn - (\max\{m, n\} - \min\{m, n\} + 1)$, it seems only implied by (7) if

$$(8) \quad mn - \frac{mn}{m+n} > mn - (\max\{m, n\} - \min\{m, n\} + 1).$$

Asymptotically this is equivalent to

$$(9) \quad \frac{\max\{m, n\}}{\min\{m, n\}} > \frac{\sqrt{5} + 1}{2} - o(1), \quad m, n \rightarrow \infty,$$

so roughly speaking, the matrices should not be close to being square matrices.

Secondly, we want to directly state results on the Dirichlet spectrum

$$(10) \quad \mathcal{D}_{m,n} := \{\Theta^{m \times n}(\|\cdot\|_\infty, \|\cdot\|_\infty, \Omega) : \Omega \in \mathbb{R}^{m \times n}\} \subseteq [0, 1].$$

As a reference point, we recall that a direct consequence of [26, 27] is that

$$(11) \quad \mathcal{D}_{m,1} = \mathcal{D}_{1,n} = [0, 1], \quad m \geq 2, n \geq 2.$$

The first claim below verifies for asymptotically 3/4 of the cases of m, n (with respect to maximum norm on \mathbb{N}^2 up to given height) that $\mathcal{D}_{m,n}$ contains some proper interval starting at 0; the other claims provide some refined information in special cases.

Theorem 3.2. *Let k, m, n be positive integers.*

- If $\frac{m}{n} \in (0, \frac{1}{2}] \cup (1, \infty)$, $(m, n) \notin \{(3, 2), (5, 2), (7, 2), \dots\}$, then $\mathcal{D}_{m,n}$ contains some right neighborhood $[0, c_{m,n}]$ of 0.
- If $m|n$ or $n|m$ and $(m, n) \neq (1, 1)$ we have $\mathcal{D}_{m,n} = [0, 1]$.
- Moreover

$$(12) \quad \mathcal{D}_{m,n} \subseteq \mathcal{D}_{km, kn}.$$

Remark 9. The first claim holds more generally for any pair of expanding norms, see Section 3.2 for a definition. In most cases $\|\cdot\|_1$ can even be arbitrary.

These results are new, in particular, independent of [12]. The first claim follows directly from a combination of Theorem 3.4 and Theorem 3.6 below, where more general norms are considered, apart from the vector cases $\min\{m, n\} = 1$ which are implied by the second claim of the theorem. The last claim (12) is Corollary 8.2 below, and the second claim follows in turn via (11) settled in [27, 26] and since $D = 1$ for maximum norms. Note that (12) implies that we can restrict to coprime m, n for the study of the Dirichlet spectrum with respect to maximum norms.

Towards the proof of Theorems 3.1 and 3.2, we consider two cases. Firstly when $\frac{m}{n} \in [\frac{1}{2}, 2]$, and secondly when $\frac{m}{n} \in (0, \frac{1}{2}) \cup (2, \infty)$. Within these cases, we again split into further two subcases depending which variable is larger and transition by using Lemma 1.1. For example in the first case we begin with $\frac{m}{n} \in [1, 2]$ and then use Lemma 1.1 to obtain results for when $\frac{m}{n} \in [\frac{1}{2}, 1]$.

3.2. The case $m/n \in [1/2, 2]$. We first treat the case $m = n$. Here we keep restricting to maximum norms.

Theorem 3.3. *For $m = n \geq 2$, the set $FS_{n,n} = FS_{n,n}(\|\cdot\|_\infty, \|\cdot\|_\infty)$ satisfies*

$$\dim_H FS_{n,n} \geq (n-1)^2 + 1.$$

Moreover, we can choose the matrices of full \mathbb{R} -rank n (i.e. invertible).

Again, up to the rank claim, this is implied by the result from [12]. The rank claim indeed appears not to be covered by [12] as (8) is easily checked to fail. The construction of our proof gives very limited control of the exact Dirichlet constant here. Even 0 being an accumulation point for the Dirichlet spectrum needs some additional conjectural assumption, which on the other hand can be inferred from [12, Theorem 3.15] and its proof method.

Now let us study the case $m \neq n$. Following [27], let us call a norm $\|\cdot\|$ on \mathbb{R}^k expanding if $\|\mathbf{x}\| \geq \|\pi_j(\mathbf{x})\|$ for any $\mathbf{x} \in \mathbb{R}^k$ and every $1 \leq j \leq k$, where

$$\pi_j : (x_1, \dots, x_k) \rightarrow (0, \dots, 0, x_j, 0, \dots, 0)$$

is the projection to the j -th coordinate axis. As remarked in [27] this is a rather mild condition that in particular includes the maximum norm and all p -norms $\|\mathbf{x}\| = (\sum |x_i|^p)^{1/p}$, $p \geq 1$. Let us again call

$$\mathcal{D}_{m,n}(\|\cdot\|_1, \|\cdot\|_2) = \{\Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) : \Omega \in \mathbb{R}^{m \times n}\} \subseteq [0, D]$$

with $D > 0$ as in (5), the Dirichlet spectrum in dimensions m, n . Denote further by $[x]$ the smallest integer $\geq x$. We show the following.

Theorem 3.4. *Assume m, n are positive integers with*

$$1 < \frac{m}{n}, \quad (m, n) \notin \{(3, 2), (5, 2), (7, 2), \dots\}.$$

Let $\|\cdot\|_1$ be an arbitrary norm on \mathbb{R}^n and $\|\cdot\|_2$ be any expanding norm on \mathbb{R}^m , unless if $n = 2$ assume both norms $\|\cdot\|_1$ and $\|\cdot\|_2$ are expanding.

Then there exists $c_{m,n} = c_{m,n}(\|\cdot\|_1, \|\cdot\|_2) > 0$ so that for any $c \in [0, c_{m,n}]$, we have

$$FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \neq \emptyset$$

in fact unless $n = 2$ or $(m, n) = (4, 3)$ we have

$$\dim_H FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \geq m \cdot (n - 1) - \left\lceil \frac{m}{n} \right\rceil,$$

and for $(m, n) = (4, 3)$ we still have the weaker bound

$$\dim_H FS_{4,3}(\|\cdot\|_1, \|\cdot\|_2, c) \geq 4.$$

In particular the Dirichlet spectrum $\mathcal{D}_{m,n}(\|\cdot\|_1, \|\cdot\|_2)$ contains a right neighborhood of 0. Moreover, in these claims we can choose the matrices of full \mathbb{R} -rank n .

Remark 10. We can improve the Hausdorff dimension formulas in Theorem 3.4 a little. For $\|\cdot\|_2$ the maximum norm, this works via applying [27, Theorem 3.1] to estimate from below $\dim_H FS_{\ell,1}(c)$ for $c \in [0, 1]$ where $\ell = \lceil m/n \rceil$. This positive value can be added given the method of our proof below and using (20). For general expanding $\|\cdot\|_2$ a similar argument applies see [27, Remark 2]. This resembles Remark 2. In particular in the cases $(m, 2)$ for $m \geq 4$ even, we can settle a positive Hausdorff dimension.

Remark 11. The condition of $\|\cdot\|_2$ being expanding can be relaxed to asking that for some space spanned by $\ell := \lceil m/n \rceil$ canonical base vectors $\mathbf{e}_{i_1}, \dots, \mathbf{e}_{i_\ell}$ the projected norm onto it is expanding. For example, for $n < m \leq 2n$ and $i_1 = 1, i_2 = 2$ this means the norm $\|(x_1, x_2)\|_2' := \|(x_1, x_2, 0, \dots, 0)\|_2$ on \mathbb{R}^2 is expanding.

As we prescribe exact Dirichlet constant, our Theorem 3.4 is not implied by [12]. We remark that full \mathbb{Q} -rank of columns, a weaker statement, would be equivalent to the matrix giving rise to an infinite (non-terminating) sequence of best approximations.

For Theorem 3.1 we only need the case $m/n \in (1, 2]$, however for the full claim of Theorem 3.2 we require the full statement of Theorem 3.4.

By equivalence of norms, for arbitrary norms, we have the according result that the Folklore set is non-empty and the Dirichlet spectrum accumulates at 0.

We now restrict to the case $m/n \in (1, 2]$ as the implications below from other cases will be superseded from results in Section 3.3. By Lemma 1.1 we may deduce an analogous

result to Theorem 3.4, apart from notably not fixing exact Dirichlet constant c , in the cases

$$n \geq 4, \quad \frac{1}{2} \leq \frac{m}{n} < 1.$$

Together with the case $m = n$ from Theorem 3.3, we may summarize

Theorem 3.5. *Let m, n be positive integers with*

$$\frac{m}{n} \in \left[\frac{1}{2}, 2\right], \quad \{m, n\} \notin \{\{1, 1\}, \{2, 3\}\}.$$

Then

$$FS_{m,n}(\|\cdot\|_\infty, \|\cdot\|_\infty) \neq \emptyset,$$

in fact

$$(13) \quad \dim_H FS_{m,n}(\|\cdot\|_\infty, \|\cdot\|_\infty) \geq mn - 2 \max\{m, n\}.$$

Moreover, we can choose the matrices of full \mathbb{R} -rank $\min\{m, n\}$. If additionally $m \neq n$, then more generally for any norms $\|\cdot\|_1$ on \mathbb{R}^n and $\|\cdot\|_2$ on \mathbb{R}^m

$$FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2) \neq \emptyset, \quad \dim_H FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2) \geq mn - 2 \max\{m, n\},$$

and $\mathcal{D}_{m,n}(\|\cdot\|_1, \|\cdot\|_2)$ accumulates at 0, in particular, it is not finite.

By contrast to Theorem 3.4, this theorem is again, apart from the rank statement in certain cases, implied by [12, Theorem 3.15] and its proof. Regarding the rank claim, by (9) essentially the cases $\max\{m, n\}/\min\{m, n\} \in [1, (1 + \sqrt{5})/2 = 1.61\dots]$ seem not covered by [12].

We remark that (13) can be improved by our method unless $\{m, n\} = \{2, 4\}$ where the right hand side vanishes. In our constructions of all Theorems 3.3, 3.4, 3.5 the real matrices in our construction have some 0 entries. In particular the entries are not \mathbb{Q} -linearly independent as in Theorems 3.6, 3.7 below, which would be desirable.

3.3. The case $m/n \notin (1/2, 2)$. Now assume $n \geq 2m$. In this case, we establish the following generalization of Theorem 2.3.

Theorem 3.6. *Let m, n be positive integers satisfying*

$$(14) \quad \frac{m}{n} \leq \frac{1}{2}, \quad n \neq m + 2.$$

Let $\|\cdot\|_1$ be any norm on \mathbb{R}^n and $\|\cdot\|_2$ be any norm on \mathbb{R}^m . Then there exists $c_{m,n} = c_{m,n}(\|\cdot\|_1, \|\cdot\|_2) > 0$ such that for any $c \in [0, c_{m,n}]$ we have

$$\dim_H FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \geq (n - 2)m.$$

Analogous claims hold when we restrict to $\Omega \in FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c)$ so that

- (i) Ω has full \mathbb{R} -rank m
- (ii) the set of all its entries $\Omega_{i,j}$ together with 1 is linearly independent over \mathbb{Q} .

In particular, $\mathcal{D}_{m,n}(\|\cdot\|_1, \|\cdot\|_2)$ contains some right neighborhood of 0.

Due to fixing the Dirichlet constant, none of the claims are implied by the work in [12].

Theorem 2.3 is the special case $m = 1$ when identifying $\|\cdot\|$ with $\|\cdot\|_1$ and $\|\cdot\|_2 = |\cdot|$ with the absolute value on \mathbb{R} . Note that when $m = 1$ conditions (i), (ii) are automatically satisfied for non-singular vectors. We will use Theorem 2.3 to prove Theorem 3.6.

Conditions (14) on m, n can be stated equivalently as

$$n \geq 2m, \quad (m, n) \notin \{(1, 3), (2, 4)\}.$$

Note that in assumption (i), we impose independence of rows over \mathbb{R} , not \mathbb{Q} .

In the case $(m, n) = (1, 3)$ where the results of Theorem 3.6 are not applicable, avoiding results from [12], we can still deduce that the larger set

$$(15) \quad Di_{1,3}(\|\cdot\|_1, \|\cdot\|_2, c) \setminus Bad_{1,3} \supseteq FS_{1,3}(\|\cdot\|_1, \|\cdot\|_2, c)$$

is uncountable for arbitrarily small $c > 0$, since by [26] we have $FS_{1,3}(\|\cdot\|_\infty, \|\cdot\|_\infty, c) \neq \emptyset$ for any $c \in [0, 1]$ and by the equivalence of norms. By [12], in fact, the Hausdorff dimension bound (7) applies for the left-hand side set in (15).

Similar to Section 3.2, using Lemma 1.1 for the reverse case $m \geq 2n$, we get the following corollary which prior to the latest version of [12] was new if $\min\{m, n\} > 1$ even for maximum norms.

Theorem 3.7. *Let m, n be positive integers satisfying*

$$(16) \quad \frac{m}{n} \notin \left(\frac{1}{2}, 2\right), \quad (m, n) \notin \{(1, 3), (3, 1), (2, 4), (4, 2)\}.$$

Let $\|\cdot\|_1$ be any norm on \mathbb{R}^n and $\|\cdot\|_2$ be any norm on \mathbb{R}^m . Then

$$\dim_H FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2) \geq (\max\{m, n\} - 2) \cdot \min\{m, n\} = mn - 2 \cdot \min\{m, n\}.$$

Moreover we may again assume (i), (ii), and the Dirichlet spectrum $\mathcal{D}_{m,n}(\|\cdot\|_1, \|\cdot\|_2)$ accumulates at 0, in particular it is not finite.


By contrast to Theorem 3.6, since we lose our grip on the exact Dirichlet constant and (9) is implied by (16), all but the claim (ii) are directly implied by Das et al [12]. Since exceptions to (ii) form a countable union of rational hyperplanes thus of Hausdorff dimension $mn - 1$ which is not exceeded by the value from (7) as soon as $\min\{m, n\} > 1$, (ii) is indeed not implied unless in the vector cases.

Note also that for the cases $\{m, n\} = \{2, 4\}$ and $\{m, n\} = \{1, 3\}$ excluded in (16), we have results via Theorem 3.4 and [27, 26] respectively.

Remark 12. It may be true that $Di_{m,n} = Di_{n,m}$ for every pair m, n . If $\min\{m, n\} = 1$, one inclusion of the set identity was proved by Davenport and Schmidt [13]. Assuming this property would give a shortcut avoiding Lemma 1.1 for the proofs of Theorems 3.5, 3.7.

4. BRIEF OUTLINE OF PROOFS

In summary, the claims above are connected in the following way (see below for Corollary 8.2):



Theorem 2.1 is proven in Sections 5.1 and 5.2, and Theorem 2.3 is proven in Section 6. The dotted arrow in the above figure between Theorem 2.3 and Theorem 3.6 indicates that the methodology of proof is similar, but by no means does Theorem 2.3 immediately imply Theorem 3.6. The same applies for the arrow between [27, Theorem 2.2] and Theorem 3.4, where besides Lemma 7.1 also, the rather easy Lemma 8.1 below is a crucial ingredient. The complete proof of Theorem 3.6 is given in Section 7. In Section 8, Theorems 3.3 and 3.4 are proven. All other implications are rather immediate. Thus the main claims that remain to be proved below are those highlighted in bold font. Lemma 1.1 is proven in Section 9.

All proofs of claims from Sections 2 and 3.3 are based on constructions for the case $m = 1, n = 2$ of one linear form in two variables. For Theorem 2.1, we can use a construction from [26], however for the results in arbitrary norms (Theorem 2.3) we follow essentially a different construction from [28], with some refinements. Then, for either result, similar as in [28], we use a method of Moshchevitin [25] to extend the claim to larger n . In Theorem 3.6 when $m > 1$, there is some additional variational argument needed to extend the $m = 1, n = 2$ case first to arbitrary m and $n = 2$ before we use Moshchevitin’s argument to be able to increase n as well arbitrarily. As an artefact of the method from [25], we lose the cases $m = 1, n = 3$ and $m = 2, n = 4$, on the other

hand, $n \geq 2m$ is essentially required for the first step $m = 1, n = 2$. Note that we gain back the case $m = 2, n = 4$ via transference from Section 3.2.

Concerning Section 3.2, for Theorem 3.4 we apply a result from [27] on $m = 2, n = 1$. We also again employ Moshchevitin's construction [25] here. We lose the case $(m, n) = (3, 2)$ in Theorem 3.1 and the cases $(m, 2)$ for $m \geq 3$ odd in Theorem 3.2 due to problems originating in [27]. The special case $m = n \geq 2$ of Theorem 3.3 is not covered by the argument, indeed it does not need either of these prerequisite results [25, 27] but uses a different, self-contained construction. Its proof is considerably shorter and easier than for the other main claims in bold font.

5. PROOF OF THEOREM 2.1

5.1. Simplifying the problem. In short, the proof combines the special two-dimensional case of [26] with an idea of Moshchevitin [25]. We first recall the result on linear forms in two variables. Throughout this section fix $n \in \mathbb{N}_{\geq 2}$ and let $\|\cdot\|_1 = \|\cdot\|_\infty$ on \mathbb{R}^n and $\|\cdot\|_2 = |\cdot|$ on \mathbb{R} . For $\zeta \in \mathbb{R}^n$ and $t \geq 1$ for ease of notation write

$$\psi_\zeta^{1 \times n}(\|\cdot\|_1, |\cdot|, t) = \psi_\zeta^{1 \times n}(t).$$

Theorem 5.1 ([26]). *Let $n \geq 2$ and $c \in [0, 1]$. Then there exist \mathbb{Q} -linearly independent with 1 real vectors $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2$ satisfying*

$$(17) \quad \limsup_{t \rightarrow \infty} t^n \psi_\xi^{1 \times 2}(t) = c,$$

and moreover

$$(18) \quad \liminf_{t \rightarrow \infty} t^n \psi_\xi^{1 \times 2}(t) = 0.$$

Remark 13. When $n \geq 3$, the liminf statement (18) follows from the limsup claim (17), for example by Marnat and Moshchevitin [23]. The linear independence over \mathbb{Q} of $(\xi, 1)$ is also implied by (17), so it is just stated for completeness.

Observe that for $n > 2$ this is not the natural order of decay as in Dirichlet's theorem, but considerably faster, so ξ is singular. This rate of decay will be required to increase the dimension to n .

The claim is a special case of [26, Theorem 1.1] for $n = 2$ and which treats "exact uniform approximation" with respect to more general decay functions Φ . As the proof of this result was only sketched in [26], we briefly recall the construction for convenience of the reader, with $\Phi(t) = ct^{-n}$. Define an integer sequence $(a_j)_{j \geq 1}$ as follows: Take the initial two terms a_1, a_2 large enough with $a_1 | a_2$. Now iteratively, for $k \geq 1$ having constructed the first $2k$ terms a_1, \dots, a_{2k} , let the next two terms are given by the recursion

$$a_{2k+1} = a_{2k}^{M_k}$$

and

$$a_{2k+2} = a_{2(k+1)} = a_{2k+1} \cdot [c^{-1} a_{2k+1}^{n-1}] \in (a_{2k+1}^n, \infty),$$

with integers $M_k \rightarrow \infty$ that tend to be infinitely fast enough. It turns out that $\xi = (\xi_1, \xi_2) \in \mathbb{R}^2$ with

$$\xi_j = \sum_{k=0}^{\infty} a_{2k+j}^{-1}, \quad j = 1, 2,$$

has the desired properties of Theorem 5.1. An alternative construction for c small enough can be derived from the method in [28], we will need the latter for our results for arbitrary norms.

Now we want to blow up the vectors in Theorem 5.1 from \mathbb{R}^2 to \mathbb{R}^n by adding components ξ_3, \dots, ξ_n . Let us call the new vector(s) $\zeta \in \mathbb{R}^n$. Then the following is an easy observation.

Proposition 1. *Take any $\xi \in \mathbb{R}^2$ as in Theorem 5.1. Any $\zeta \in \mathbb{R}^n$ with first two coordinates $\xi_1 = \zeta_1, \xi_2 = \zeta_2$, that is any vector $\zeta \in \{\xi\} \times \mathbb{R}^{n-2} \subseteq \mathbb{R}^n$, satisfies $\zeta \in Di_n(c) \setminus Bad_n$.*

Proof. Let us write $\xi^* = (\xi_1, \xi_2, 1)$ and similarly $\zeta^* = (\zeta_1, \dots, \zeta_n, 1)$. Denote $e_i = (0, 0, \dots, 0, 1, 0, \dots, 0)$ the i -th canonical base vector of \mathbb{R}^{n+1} .

Our claim follows from the fact that

$$(19) \quad \psi_{\zeta}^{1 \times n}(t) \leq \psi_{\xi}^{1 \times 2}(t), \quad t > 0.$$

Indeed, any t induces a best approximation $\mathbf{b} = (b_1, b_2, b_3) \in \mathbb{Z}^3$ with respect to ξ that realizes $\psi_{\xi}(t)$. Then the blown-up vector

$$\tilde{\mathbf{b}} := b_1 \mathbf{e}_1 + b_2 \mathbf{e}_2 + b_3 \mathbf{e}_{n+1} = (b_1, b_2, 0, \dots, 0, b_3) \in \mathbb{Z}^{n+1}$$

induces the same maximum norm and approximation quality

$$\|\hat{\mathbf{b}}\| = \|\tilde{\mathbf{b}}\|, \quad |\tilde{\mathbf{b}} \cdot \zeta^*| = |\mathbf{b} \cdot \xi^*| = \psi_{\xi}^{1 \times 2}(t).$$

so the minimum over all integer points defining $\psi_{\zeta}(t)$ cannot be larger. Trivially (18) and (19) implies

$$\liminf_{t \rightarrow \infty} t^n \psi_{\zeta}^{1 \times n}(t) = 0,$$

so $\zeta \notin Bad_n$. Similarly (17) and (19) implies

$$\limsup_{t \rightarrow \infty} t^n \psi_{\zeta}^{1 \times n}(t) \leq c,$$

so $\zeta \in Di_n(c)$. □

Proposition 1 does not imply that $\zeta \notin Sing_n$. For example, if $n \geq 3$ taking $\zeta = (\xi_1, \xi_2, \xi_1 + \xi_2, 0, \dots, 0) \in \mathbb{R}^n$ then $\zeta \in Sing_n$. However, the following Theorem enables us to say that this does not happen for a significantly large subset of \mathbb{R}^{n-2} of “additional coordinates”. In fact, we can ensure the induced Dirichlet constant is not smaller than the one of ξ for generic vectors of \mathbb{R}^{n-2} .

Theorem 5.2. *Let $n \geq 4$ be an integer. Take any $\xi \in \mathbb{R}^2$ as in Theorem 5.1 derived from any small enough $c \in (0, c(n))$ as in Theorem 2.1. Then for a set*

$$\mathcal{Y} = \mathcal{Y}_{n, \xi} \subseteq \mathbb{R}^{n-2}$$

of positive $n - 2$ dimensional Lebesgue measure consisting of ζ_3, \dots, ζ_n , the vector $\zeta = (\xi_1, \xi_2, \zeta_3, \dots, \zeta_n) \in \mathbb{R}^n$ satisfies

$$\Theta^{1 \times n}(\zeta) = \limsup_{t \rightarrow \infty} t^n \psi_{\zeta}^{1 \times n}(t) = \limsup_{t \rightarrow \infty} t^n \psi_{\xi}^{1 \times 2}(t) = c,$$

or equivalently

$$\zeta \in DI_n(c).$$

As for Theorem 2.1, we again believe the claim should hold for $n = 3$ as well.

Remark 14. Presumably the $(n - 2)$ -dimensional Lebesgue measure of the set \mathcal{Y} is full, however, this is not required for our purposes.

By combining Proposition 1 and Theorem 5.2 it is clear that

$$\zeta \in DI_n(c) \setminus Bad_n = FS_n(c).$$

Let \mathcal{Z}_c denote the set of $\xi \in \mathbb{R}^2$ satisfying Theorem 5.1. Since $\lambda_n(\mathcal{Y}) > 0$, for any $c \in (0, c_n)$ we have that

$$\dim_H FS_n \geq \dim_H FS_n(c) \geq \dim_H \mathcal{Z}_c \times \mathcal{Y} \geq \dim_H(\mathcal{Z}_c) + \dim_H \mathcal{Y} \geq n - 2.$$

The second to last inequality follows from a standard result on the Hausdorff dimension of Cartesian products

$$(20) \quad \dim_H A \times B \geq \dim_H A + \dim_H B$$

for A, B arbitrary sets in Euclidean spaces, see [32]. Hence Theorem 2.1 is proven modulo Theorem 5.2.

Remark 15. Notice that in the above calculation, we simply use $\dim_H \mathcal{Z}_c \geq 0$. It is here where our lower bound could be improved. For $\tau > 2$ a parameter, define

$$Sing_2(\tau) := \left\{ \xi \in \mathbb{R}^2 : \limsup_{t \rightarrow \infty} t^\tau \psi_{\xi}^{1 \times 2}(t) = 0 \right\}.$$

It can be deduced from [11, Theorem 1.4-1.9] by a change of variables that for any $\tau > 2 + \sqrt{2}$ we have

$$\dim_H Sing_2(\tau) = \frac{2}{\tau}.$$

Clearly $\mathcal{Z}_c \subseteq Sing_2(n - \varepsilon)$ for any $\varepsilon > 0$, and so

$$\dim_H \mathcal{Z}_c \leq \frac{2}{n - \varepsilon}.$$

Thus for $n \geq 4 > 2 + \sqrt{2}$ the best we could hope for via this method is

$$\dim_H FS_n(c) \geq n - 2 + \frac{2}{n}.$$

On the other hand, a positive lower bound for $\dim_H(\mathcal{Z}_c)$ can be obtained by a similar variational method as in [28, Theorem 2.1], we do not carry it out.

Theorem 5.2 remains to be proved, for which we reserve the next section.

5.2. Proof of Theorem 5.2. We have to show the Dirichlet constant of many ζ is at least c , the rest is obvious. We follow ideas from the proof of Moshchevitin [25, Theorem 12] with some twists. Write $\zeta = (\xi_1, \xi_2, \zeta_3, \dots, \zeta_n)$ and restrict to $(\zeta_3, \dots, \zeta_n)$ in the $n-2$ -dimensional unit ball $B_{n-2}(\mathbf{0}, 1) \subseteq \mathbb{R}^{n-2}$ with respect to the Euclidean norm (this will be slightly easier to handle than for the maximum norm). Denote by

$$\mathbf{b}_v = (b_{1,v}, b_{2,v}, b_{3,v}) \in \mathbb{Z}^3, \quad v \geq 1,$$

the best approximations with respect to $\xi = (\xi_1, \xi_2)$ and derive

$$M_v = \max_{j=1,2} |b_{j,v}| = \|\hat{\mathbf{b}}_v\|, \quad v \geq 1$$

and the minimal values

$$L_v = |b_{1,v}\xi_1 + b_{2,v}\xi_2 + b_{3,v}|, \quad v \geq 1.$$

Clearly $(L_v)_{v \geq 1}$ is a strictly decreasing sequence and $(M_v)_{v \geq 1}$ is strictly increasing. Further, note that we have

$$\psi_{\xi}^{1 \times 2}(t) = L_v, \quad t \in [M_v, M_{v+1}),$$

hence

$$(21) \quad \limsup_{t \rightarrow \infty} t^n \psi_{\xi}^{1 \times 2}(t) = \limsup_{v \rightarrow \infty} L_v M_{v+1}^n.$$

Denote $\mathbf{e}_i = (0, 0, \dots, 0, 1, 0, \dots, 0)$ the i -th canonical base vector of \mathbb{R}^{n+1} . To finish the proof, it suffices to show the following:

Lemma 5.3. *Let $n \geq 4$ an integer and $\xi = (\xi_1, \xi_2)$ as in Theorem 5.1. For a positive measure set $\mathcal{Y} \subseteq B_{n-2}(0, 1) \subseteq \mathbb{R}^{n-2}$ consisting of $\gamma = (\xi_3, \dots, \xi_n)$, we have that for infinitely many integers $v \geq 1$, letting $T_v := M_{v+1} - 1/2$, the integer best approximation for $\zeta = (\xi, \gamma)$ at T_v is given as*

$$b_{1,v}\mathbf{e}_1 + b_{2,v}\mathbf{e}_2 + b_{3,v}\mathbf{e}_3 = (b_{1,v}, b_{2,v}, 0, \dots, 0, b_{3,v}) \in \mathbb{Z}^{n+1},$$

thus the best approximation function at values T_v equals

$$\psi_{\zeta}^{1 \times n}(T_v) = \psi_{\xi}^{1 \times 2}(T_v) = L_v.$$

Observe that since T_v is of the same order as M_{v+1} , for ζ as in the lemma, (21) and the corresponding property for ζ readily implies

$$\Theta^{1 \times n}(\zeta) \geq \limsup_{v \rightarrow \infty} L_v T_v^n = \limsup_{t \rightarrow \infty} t^n \psi_{\xi}^{1 \times 2}(t),$$

the reverse inequality is clear by Proposition 1. Hence, assuming the lemma is true, the claim of Theorem 5.2 follows.

We finally prove the preceding lemma as in [25].

Proof of Lemma 5.3. We show that for a positive portion of $B_{n-2}(\mathbf{0}, 1) \subseteq \mathbb{R}^{n-2}$ of vectors $\gamma = (\zeta_3, \dots, \zeta_n)$, we have for infinitely many v that

$$(22) \quad \min \left| x_1 \xi_1 + x_2 \xi_2 + \sum_{i=3}^n x_i \zeta_i - y \right| \geq L_v$$

with minimum taken over all integer points

$$\mathbf{z} = (x_1, \dots, x_n, y) \in \mathbb{Z}^{n+1} \setminus \{\mathbf{0}\}$$

with

$$(23) \quad 0 < \max_{1 \leq j \leq n} |x_j| \leq T_v.$$

For $v \geq 1$, denote by \mathcal{S}_v the set of such vectors of \mathbb{R}^{n-2} taken over all integer points x_i, y obeying (23), i.e.

$$\mathcal{S}_v := \left\{ \gamma \in B_{n-2}(\mathbf{0}, 1) : \min_{\substack{(\mathbf{x}, y) \in \mathbb{Z}^{n+1}: \\ 0 < \max_{1 \leq i \leq n} |x_i| \leq T_v}} \left| x_1 \xi_1 + x_2 \xi_2 + \sum_{i=3}^n x_i \zeta_i - y \right| \geq L_v \right\}.$$

Then our set is the limsup set of the \mathcal{S}_v , or its complement is the set of all vectors that lie only in finitely many \mathcal{S}_v , that is,

$$(24) \quad B_{n-2}(\mathbf{0}, 1) \setminus \mathcal{Y} \subseteq \bigcup_{i \geq 1} \bigcap_{j \geq i} \mathcal{S}_j^c \subseteq \bigcup_{i \geq 1} \mathcal{S}_i^c.$$

Assume that any complement

$$\mathcal{U}_v := \mathcal{S}_v^c = \left\{ \gamma \in B_{n-2}(\mathbf{0}, 1) : \min_{\substack{(\mathbf{x}, y) \in \mathbb{Z}^{n+1}: \\ 0 < \max_{1 \leq i \leq n} |x_i| \leq T_v}} \left| x_1 \xi_1 + x_2 \xi_2 + \sum_{i=3}^n x_i \zeta_i - y \right| < L_v \right\}, \quad v \geq 1,$$

has $n-2$ dimensional Lebesgue measure less than some δ strictly smaller than the volume of $B_{n-2}(\mathbf{0}, 1)$. Then this will also be true for $B_{n-2}(\mathbf{0}, 1) \setminus \mathcal{Y}$ by the following easy result.

Proposition 2. *Let λ be a measure on a space \mathcal{T} . Assume H_i are increasing subsets of \mathcal{T} , i.e. $H_1 \subseteq H_2 \subseteq \dots$. Then*

$$\lambda \left(\bigcup_{k=1}^{\infty} H_k \right) = \lim_{k \rightarrow \infty} \lambda(H_k).$$

We will require the estimate $\lambda \left(\bigcup_{k=1}^{\infty} H_k \right) \leq \lim_{k \rightarrow \infty} \lambda(H_k)$ only.

Proof. Writing $G_j = H_j \setminus H_{j-1}$ with $H_0 = \emptyset$, we see

$$\begin{aligned} \lambda \left(\bigcup_{k=1}^{\infty} H_k \right) &= \lambda \left(\bigcup_{k=1}^{\infty} G_k \right) \\ &= \sum_{k=1}^{\infty} \lambda(G_k) \\ &= \lim_{N \rightarrow \infty} \sum_{k=1}^N \lambda(G_k) \\ &= \lim_{N \rightarrow \infty} \lambda \left(\bigcup_{k=1}^N H_k \right) \\ &= \lim_{N \rightarrow \infty} \lambda(H_N). \end{aligned}$$

□

In the sequel denote by λ the $(n-2)$ -dimensional Lebesgue measure of a set $A \subseteq \mathbb{R}^{n-2}$. As indicated before, we apply Proposition 2 to

$$H_i = \bigcap_{j \geq i} \mathcal{S}_j^c = \bigcap_{j \geq i} \mathcal{U}_j \subseteq \mathcal{U}_i, \quad i \geq 1,$$

to see via (24) that

$$\lambda(B_{n-2}(\mathbf{0}, 1) \setminus \mathcal{Y}) \leq \lim_{v \rightarrow \infty} \lambda(\mathcal{U}_v).$$

Assume we have shown that for some uniform Δ we have

$$(25) \quad \lambda(\mathcal{U}_v) \leq \Delta < \lambda(B_{n-2}(\mathbf{0}, 1)), \quad v \geq 1.$$

Then we conclude that

$$\lambda(B_{n-2}(\mathbf{0}, 1) \setminus \mathcal{Y}) \leq \Delta < \lambda(B_{n-2}(\mathbf{0}, 1))$$

and thus

$$\lambda(\mathcal{Y}) = \lambda(B_{n-2}(\mathbf{0}, 1)) - \lambda(B_{n-2}(\mathbf{0}, 1) \setminus \mathcal{Y}) \geq \lambda(B_{n-2}(\mathbf{0}, 1)) - \Delta > 0,$$

as desired.

To verify (25), let us estimate $\lambda(\mathcal{U}_v)$ for fixed v . For any $\boldsymbol{\gamma} = (\zeta_3, \dots, \zeta_n) \in \mathcal{U}_v$ it follows that

$$\min_{\substack{(\mathbf{x}, y) \in \mathbb{Z}^{n+1}: \\ 0 < \max_{1 \leq i \leq n} |x_i| \leq T_v}} \left| x_1 \xi_1 + x_2 \xi_2 + \sum_{i=3}^n x_i \zeta_i - y \right| < L_v.$$

Equivalently

$$x_3 \zeta_3 + \dots + x_n \zeta_n \in J_v(x_1, x_2, y)$$

where $J_v(\cdot)$ is given as the interval

$$J_v(x_1, x_2, y) = (-y - x_1 \xi_1 - x_2 \xi_2 - L_v, -y - x_1 \xi_1 - x_2 \xi_2 + L_v).$$

Let

$$\Omega_v(x_1, \dots, x_n, y) = \left\{ (\zeta_3, \dots, \zeta_n) \in B_{n-2}(0, 1) \subseteq \mathbb{R}^{n-2} : \sum_{i=3}^n x_i \zeta_i \in J_v(x_1, x_2, y) \right\}.$$

Observe that $\Omega_v(x_1, \dots, x_n, y)$ is empty if $|y| > |x_1 \xi_1| + |x_2 \xi_2| + \sum_{i=3}^n |x_i \zeta_i| + L_v$. Fix for now $\varepsilon > 0$. Since $L_v \rightarrow 0$ as $v \rightarrow \infty$ and as we may take ξ_1, ξ_2 arbitrarily small positive numbers (the ξ from Theorem 5.1 are easily seen to be dense in \mathbb{R}), say smaller than $\varepsilon/8$, and $(\zeta_3, \dots, \zeta_n) \in B_{n-2}(\mathbf{0}, 1)$, Cauchy-Schwarz inequality yields for any $\varepsilon > 0$ this is true if

$$|y| > (\sqrt{n-2} + \varepsilon/2)T_v$$

as then

$$\begin{aligned} |y| &> (\sqrt{n-2} + \varepsilon/2)T_v \\ &\geq (\varepsilon/4) \cdot T_v + \left(\sum_{i=3}^n |x_i|^2 \right)^{1/2} \left(\sum_{i=3}^n |\zeta_i|^2 \right)^{1/2} + L_v \\ &\geq |\xi_1| |x_1| + |\xi_2| |x_2| + \left(\sum_{i=3}^n |x_i|^2 \right)^{1/2} \left(\sum_{i=3}^n |\zeta_i|^2 \right)^{1/2} + L_v \\ &\geq |x_1 \xi_1| + |x_2 \xi_2| + \sum_{i=3}^n |x_i \zeta_i| + L_v, \quad v \geq v_0(\varepsilon). \end{aligned}$$

The above shows that if we let

$$(26) \quad \Omega_v = \bigcup_{x_1, \dots, x_n} \bigcup_y \Omega_v(x_1, \dots, x_n, y),$$

with unions are taken over integers pairs x_1, x_2 as in (23) and integers y with $|y| \leq (\sqrt{n-2} + \varepsilon/2)T_v$, then

$$\mathcal{U}_v \subseteq \Omega_v, \quad v \geq v_0(\varepsilon).$$

The distance between the parallel planes defining $\Omega_v(x_1, \dots, x_n, y)$ equals $2(x_3^2 + \dots + x_n^2)^{-1/2}$. Thus each $\Omega_v(x_1, \dots, x_n, y)$ is contained in a cylinder of radius one and height $2L_v \cdot (x_3^2 + \dots + x_n^2)^{-1/2}$. Denoting by λ' the $(n-3)$ -dimensional Lebesgue measure, this has volume

$$(27) \quad \begin{aligned} \lambda(\Omega_v(x_1, \dots, x_n, y)) &= 2\lambda'(B_{n-3}(\mathbf{0}, 1)) \cdot L_v (x_3^2 + \dots + x_n^2)^{-1/2} \\ &\leq 2L_v \cdot \lambda'(B_{n-3}(\mathbf{0}, 1)) \cdot \frac{1}{\max_{3 \leq i \leq n} |x_i|}. \end{aligned}$$

Since, assuming $T_v \geq \varepsilon^{-1}$, there are at most

$$2T_v(\sqrt{n-2} + \varepsilon/2) + 1 \leq T_v(2\sqrt{n-2} + 2\varepsilon)$$

integers y in the inner union of (26), in total we get

$$\begin{aligned} \lambda(\mathcal{U}_v) &\leq \lambda(\Omega_v) \\ &\leq \sum_{x_1, x_2} \sum_y \sum_{x_3, \dots, x_n} \lambda(\Omega_v(x_1, \dots, x_n, y)) \\ &\leq 2(2\sqrt{n-2} + 2\varepsilon) L_v T_v \lambda'(B_{n-3}(\mathbf{0}, 1)) \cdot \sum_{x_1, x_2} \sum_{x_3, \dots, x_n} \frac{1}{\max_{3 \leq i \leq n} |x_i|} \end{aligned}$$

for $v \geq v_0(\varepsilon)$. The outer sum over x_1, x_2 gives a factor $[T_v]^2 \leq T_v^2$ by (23). We split the inner sum into sets where $\max_{3 \leq i \leq n} |x_i| = k \in [1, [T_v]]$ is constant, which correspond to $2(n-2) = 2n-4$ faces on a hypercube for each k according to which variable equals $\pm k$, giving in total $(2n-4)k^{n-3}$ integer points for each k to sum $1/k$ over. Hence we may further estimate

$$\begin{aligned} \lambda(\mathcal{U}_v) &\leq 2(2n-4)(2\sqrt{n-2} + 2\varepsilon) \lambda'(B_{n-3}(\mathbf{0}, 1)) \cdot L_v T_v^3 \cdot \sum_{k=1}^{[T_v]} k^{n-4} \\ &\leq 2(2n-4)(2\sqrt{n-2} + 2\varepsilon) \lambda'(B_{n-3}(\mathbf{0}, 1)) \left(\frac{1}{n-3} + \varepsilon_2 \right) L_v T_v^n, \quad v \geq v_1(\varepsilon, \varepsilon_2). \end{aligned}$$

Here the error term $\varepsilon_2 > 0$ tends to 0 as $v \rightarrow \infty$, and we used $n \neq 3$ to avoid a logarithmic term appearing in the last sum. We may assume $\varepsilon_2 < \varepsilon$ for all large v . So if $L_v < cT_v^{-n}$ for any $c \in (0, c_n - \varepsilon)$ for $\varepsilon > 0$ some small manipulation of ε and with c_n satisfying

$$\lambda(B_{n-2}(\mathbf{0}, 1)) > c_n \frac{4(2n-4)\sqrt{n-2} \lambda'(B_{n-3}(\mathbf{0}, 1))}{n-3},$$

or equivalently

$$c_n < \frac{n-3}{4(2n-4)\sqrt{n-2}} \cdot \frac{\lambda(B_{n-2}(\mathbf{0}, 1))}{\lambda'(B_{n-3}(\mathbf{0}, 1))} = \frac{n-3}{8(n-2)^{3/2}} \cdot \sqrt{\pi} \cdot \frac{\Gamma(n-\frac{1}{2})}{\Gamma(n)},$$

indeed (25) holds for some $\Delta < \lambda(B_{n-2}(\mathbf{0}, 1))$, depending on c_n , and all $v \geq v_0(\Delta) = v_0(c_n)$. Hence the complement contained in \mathcal{Y} has positive $(n-2)$ -dimensional Lebesgue measure. Since we can choose $\varepsilon > 0$ and thus $\varepsilon > 0$ arbitrarily small, we may just assume that $c < c_n$ and the proof of Lemma 5.3 is complete. \square

Remark 16. Improvements of c_n can be made by sharpening the contribution $2\sqrt{n-2}+1$ for the number of y . Indeed, for most y the obtained intersections with the unit ball have a considerably smaller volume than $2L_v \cdot (x_3^2 + \dots + x_n^2)^{-1/2}$, so it seems we can gain some factor by more concise estimates.

6. PROOF OF THEOREM 2.3

6.1. Outline. The method of proof is similar to that of Theorem 2.1. Namely, the proof is done in two steps:

Step I: For small enough c , construct $\boldsymbol{\xi} = (\xi_1, \xi_2)$ such that

$$\limsup_{t \rightarrow \infty} t^n \psi_{\boldsymbol{\xi}}^{1 \times 2}(\|\cdot\|, t) = c.$$

Step II: Show that for a positive measure set (ξ_3, \dots, ξ_n) with respect to $(n-2)$ -dimensional Lebesgue measure, the extended vector $\zeta = (\xi_1, \dots, \xi_n)$ satisfies

$$\Theta^{1 \times n}(\zeta) = \limsup_{t \rightarrow \infty} t^n \psi_\zeta^{1 \times n}(\|\cdot\|, t) = c$$

as well.

6.2. Proof of Step I. This is the main substance of the proof. For this, we use a different construction than for the maximum norm where we employed the construction from [26]. Rather we follow the proof of [28, Theorem 2.1] but with some notable twists.

Write $\xi^* = (\xi, 1) \in \mathbb{R}^3$ and use a similar meaning of star for other quantities. Conversely denote by a hat over a vector in \mathbb{R}^3 its restriction to the first 2 coordinates by chopping off the last coordinate, i.e. $\widehat{\xi^*} = \xi$.

We construct our real vector. Let $n < \tau < \mu$ be parameters to be chosen later related by the identity

$$(28) \quad \tau\mu - 1 = n\mu \quad \iff \quad \mu = \frac{1}{\tau - n}.$$

Let $\alpha_j, \beta_j, \gamma_j, \delta_j$ be strictly increasing positive integer sequences and a real number $r = r(\|\cdot\|) > 0$ all to be fixed later. Derive integers of the form

$$A_j = 2^{\alpha_j} 3^{\gamma_j}, \quad B_j = 5^{\beta_j} 7^{\delta_j},$$

satisfying as $j \rightarrow \infty$

$$(29) \quad A_{j+1} = (1 + o(1))rB_j^\tau, \quad B_j = (1 + o(1))A_j^\mu.$$

We first show that such choices are possible. For A_j , we use that $\{\log 2, \log 3\}$ is a \mathbb{Q} -linearly independent set (easily seen by applying exponential map to a putative vanishing linear form and using unique prime factorization to disprove it) and the following easy Proposition 3. Similarly for B_j we use that $\{\log 5, \log 7\}$ is \mathbb{Q} -linearly independent.

Proposition 3. *Let a, b, c be real numbers with a, b linearly independent over \mathbb{Q} . Then ordering the set $\{k_1 a + k_2 b + c : k_i \in \mathbb{N}\}$ increasingly and denoting it by $(z_i)_{i \geq 1}$ we have $|z_{i+1} - z_i| \rightarrow 0$.*

Proof. Clearly, we may assume $c = 0$ and suppose without loss of generality that $b < a$. We may then equivalently consider

$$\mathcal{Z} = \{k_1 + k_2 \frac{b}{a} : k_i \in \mathbb{N}\}$$

instead. Since $\frac{b}{a}$ is irrational, Kronecker's Theorem (see e.g. [7, Section 3.5 Theorem IV]) shows that the set is dense in the unit interval when considering all $(k_1, k_2) \in \mathbb{Z} \times \mathbb{N}$. Let $\varepsilon = 1/K > 0$ with a large integer K and partition $[0, 1]$ into K intervals

$$I_t = [(t-1)\varepsilon, t\varepsilon], \quad 1 \leq t \leq K$$

of length $1/K$. Then for any I_t there exist $(k'_1(t), k'_2(t)) \in \mathbb{Z} \times \mathbb{N}$ so that $k'_1(t) + k'_2(t) \frac{b}{a} \in I_t$. Take $T_\varepsilon := \max_t |k'_1(t)|$ and consider the set

$$\mathcal{Z}_{T_\varepsilon} = \{(k_1(t) + N) + k_2(t) \frac{b}{a} : 1 \leq t \leq K, N \in \mathbb{N}_{\geq T_\varepsilon}\} \subset \mathcal{Z}.$$

It is easy to see that for any $M \in \mathbb{N}_{\geq T_\varepsilon}$ the interval $M + I_t$ contains a point from $\mathcal{Z}_{T_\varepsilon}$. This shows that \mathcal{Z} is ε dense on some interval $[3T_\varepsilon, \infty)$. Since $\varepsilon > 0$ is arbitrary the claim is proved. \square

We can now apply the proposition alternatingly to

$$a = \log 2, b = \log 3, c = \log r, \quad a = \log 5, b = \log 7, c = 0$$

respectively to construct iteratively our sequences α_j, γ_j , which correspond to $k_1 = \alpha_{j+1} - \alpha_j \geq 0$ and $k_2 = \gamma_{j+1} - \gamma_j \geq 0$, resp. $k_1 = \beta_{j+1} - \beta_j \geq 0$ and $k_2 = \delta_{j+1} - \delta_j \geq 0$. Thereby, having chosen all $\alpha, \beta, \gamma, \delta$. up to index j , we construct first pairs $\alpha_{j+1}, \gamma_{j+1}$ and then $\beta_{j+1}, \delta_{j+1}$, and the iterative construction is complete. So, having chosen r , we may assume all the above is defined.

Note that (29) implies

$$(30) \quad A_{j+1} = (1 + o(1))rA_j^{\tau_\mu}, \quad B_{j+1} = (1 + o(1))r^\mu B_j^{\tau_\mu}.$$

Moreover, again by (29), upon changing initial terms if necessary, we can assume

$$(31) \quad 1 < A_1 < B_1 < A_2 < B_2 < \dots$$

Finally let $\xi = (\xi_1, \xi_2)$ with

$$\xi_1 = \sum_{j \geq 1} A_j^{-1}, \quad \xi_2 = \sum_{j \geq 1} B_j^{-1}.$$

We claim that it satisfies the assertions of the theorem upon choosing r appropriately.

For $j \geq 1$ put

$$\sigma_j := A_1^{-1} + \dots + A_j^{-1}, \quad \eta_j := B_1^{-1} + \dots + B_j^{-1}.$$

We clearly have $\sigma_j \rightarrow \xi_1$ and $\eta_j \rightarrow \xi_2$ as $j \rightarrow \infty$, and

$$F_j = A_j \sigma_j \in \mathbb{N}, \quad G_j = B_j \eta_j \in \mathbb{N}.$$

Then since the sequences $(\alpha_j), (\beta_j), (\gamma_j), (\delta_j)$ are strictly increasing we have $F_j \equiv 1 \pmod{2 \cdot 3}$ and $G_j \equiv 1 \pmod{5 \cdot 7}$, which imply the coprimality assertions

$$(32) \quad (A_j, F_j) = (F_j, 6) = 1, \quad (B_j, G_j) = (G_j, 35) = 1.$$

Let

$$\mathbf{v}_j = (A_j, 0, -F_j), \quad \mathbf{w}_j = (0, B_j, -G_j).$$

Note that $\hat{\mathbf{v}}_j = (A_j, 0)$ and $\hat{\mathbf{w}}_j = (0, B_j)$. Hence their norms $\|\cdot\|$ satisfy

$$(33) \quad \|\hat{\mathbf{v}}_j\| = \|\mathbf{e}_1\| A_j = d_1 A_j, \quad d_1 := \|\mathbf{e}_1\| = \|(1, 0)\|,$$

and

$$(34) \quad \|\hat{\mathbf{w}}_j\| = \|\mathbf{e}_2\| B_j = d_2 B_j \stackrel{(29)}{=} (1 + o(1)) d_2 A_j^\mu, \quad d_2 := \|\mathbf{e}_2\| = \|(0, 1)\|,$$

Thus by (31) clearly

$$(35) \quad \|\hat{\mathbf{v}}_j\| < \|\hat{\mathbf{w}}_j\| < \|\hat{\mathbf{v}}_{j+1}\|, \quad j \geq j_0.$$

Here j_0 depends on d_1, d_2 , thus on the chosen norm. Then by (30) moreover

$$(36) \quad \begin{aligned} |\mathbf{v}_j \cdot \boldsymbol{\xi}^*| &= |A_j \cdot \xi_1 + 0 \cdot \xi_2 - F_j| = A_j(A_{j+1}^{-1} + A_{j+2}^{-1} + \cdots) \\ &= A_j A_{j+1}^{-1} (1 + o(1)) \stackrel{(30)}{=} (1 + o(1)) r^{-1} A_j^{-(\tau\mu-1)} \end{aligned}$$

and

$$(37) \quad \begin{aligned} |\mathbf{w}_j \cdot \boldsymbol{\xi}^*| &= |0 \cdot \xi_1 + B_j \cdot \xi_2 - G_j| = B_j(B_{j+1}^{-1} + B_{j+2}^{-1} + \cdots) \\ &= (1 + o(1)) B_j B_{j+1}^{-1} \stackrel{(30)}{=} (1 + o(1)) r^{-\mu} \cdot B_j^{-(\tau\mu-1)}, \end{aligned}$$

are small linear forms for $j \geq 1$. We next show

Lemma 6.1. *As soon as r is large enough, all best approximations for $\boldsymbol{\xi}$ of large norm have up to sign one of the following forms*

$$\mathbf{v}_j, \quad \mathbf{w}_j, \quad \mathbf{v}_j + \mathbf{w}_j, \quad \mathbf{w}_j - \mathbf{v}_j, \quad \mathbf{w}_j + \mathbf{v}_{j+1}, \quad \mathbf{v}_{j+1} - \mathbf{w}_j.$$

Proof. Let \mathbf{b} be any best approximation. Then by (35) there is an index j such that either $\|\hat{\mathbf{v}}_j\| \leq \|\hat{\mathbf{b}}\| < \|\hat{\mathbf{w}}_j\|$ or $\|\hat{\mathbf{w}}_j\| \leq \|\hat{\mathbf{b}}\| < \|\hat{\mathbf{v}}_{j+1}\|$. We show that in the first case $\mathbf{b} = \pm \mathbf{v}_j$ or $\pm(\mathbf{v}_j \pm \mathbf{w}_j)$, and in the latter case $\mathbf{b} = \pm \mathbf{w}_j$ or $\pm(\mathbf{w}_j \pm \mathbf{v}_{j+1})$. Assume the first case, so

$$(38) \quad \|\hat{\mathbf{v}}_j\| \leq \|\hat{\mathbf{b}}\| < \|\hat{\mathbf{w}}_j\|.$$

the latter works very similarly by symmetry. First, observe that since \mathbf{b} is the best approximation of norm at least $\|\hat{\mathbf{v}}_j\|$, we know that

$$(39) \quad |\mathbf{b} \cdot \boldsymbol{\xi}^*| \leq |\mathbf{v}_j \cdot \boldsymbol{\xi}^*|.$$

We distinguish two cases.

Case 1: \mathbf{b} lies in the two-dimensional subspace of \mathbb{R}^3 spanned by $\mathbf{v}_j, \mathbf{w}_j$, i.e. $\mathbf{b} \in \langle \mathbf{v}_j, \mathbf{w}_j \rangle_{\mathbb{R}} \cap \mathbb{Z}^3$. The special form of A_j, B_j and (32) imply the following crucial result on integer vectors in the two-dimensional lattices $\langle \mathbf{v}_j, \mathbf{w}_j \rangle_{\mathbb{R}} \cap \mathbb{Z}^3$.

Proposition 4. *For $\mathbf{v}_j, \mathbf{w}_j$ as above, if a linear combination $g\mathbf{v}_j + h\mathbf{w}_j$ is an integer vector, then in fact $g \in \mathbb{Z}$ and $h \in \mathbb{Z}$. In other words, $\langle \mathbf{v}_j, \mathbf{w}_j \rangle_{\mathbb{R}} \cap \mathbb{Z}^3 = \langle \mathbf{v}_j, \mathbf{w}_j \rangle_{\mathbb{Z}}$.*

Proof. Clearly, we must have $g, h \in \mathbb{Q}$. If we write $(p_1/q_1)\mathbf{v}_j + (p_2/q_2)\mathbf{w}_j$ with p_i/q_i in lowest terms, then it is clear that q_1 must consist of exclusively non-negative integer powers of 2 and 3 and q_2 of non-negative integer power of 5 and 7 to make the first two coordinates $(p_1/q_1)A_j = (p_1/q_1)2^{\alpha_j}3^{\gamma_j}$ resp. $(p_2/q_2)B_j = (p_2/q_2)5^{\beta_j}7^{\delta_j}$ of $g\mathbf{v}_j + h\mathbf{w}_j$ integers. But then by (32) clearly the third coordinate $(p_1/q_1)F_j + (p_2/q_2)G_j$ is not an integer unless $q_1 = q_2 = 1$. \square

We first show that any integer linear combination $\mathbf{b} = g\mathbf{v}_j + h\mathbf{w}_j$ representing the best approximation as above must have $|g| \leq 1$. Assume $|g| \geq 2$. It is easily seen that for any norm on \mathbb{R}^2 there is a constant $\theta = \theta(\|\cdot\|)$ such that $\|\mathbf{w}\| \leq W$ for $\mathbf{w} = (w_1, w_2) \in \mathbb{R}^2$ and $W > 0$ implies $|w_i| \leq \theta W$, $i = 1, 2$. Hence (38) and the special form of $\mathbf{v}_j, \mathbf{w}_j$ imply

$$|g| \leq \theta, \quad |h| \leq \theta$$

for some θ independent of j . On the other hand, since r, μ are fixed and $A_j \rightarrow \infty$, it is clear from (36), (37) and (29) that

$$|\mathbf{w}_j \cdot \boldsymbol{\xi}^*| = o(|\mathbf{v}_j \cdot \boldsymbol{\xi}^*|), \quad j \rightarrow \infty.$$

Combination implies

$$\begin{aligned} |\mathbf{b} \cdot \boldsymbol{\xi}^*| &= |(g\mathbf{v}_j + h\mathbf{w}_j) \cdot \boldsymbol{\xi}^*| \geq |g| \cdot |\mathbf{v}_j \cdot \boldsymbol{\xi}^*| - |h| \cdot |\mathbf{w}_j \cdot \boldsymbol{\xi}^*| \geq 2|\mathbf{v}_j \cdot \boldsymbol{\xi}^*| - |h| \cdot |\mathbf{w}_j \cdot \boldsymbol{\xi}^*| \\ &\geq 2|\mathbf{v}_j \cdot \boldsymbol{\xi}^*| - \theta \cdot |\mathbf{w}_j \cdot \boldsymbol{\xi}^*| > |\mathbf{v}_j \cdot \boldsymbol{\xi}^*|, \quad j \geq j_0, \end{aligned}$$

contradicting (39). Hence $|g| \leq 1$. Now if $|h| \geq 2$ then for any $|g| \leq 1$ using (35) we get

$$\|\hat{\mathbf{b}}\| = \|g\hat{\mathbf{v}}_j + h\hat{\mathbf{w}}_j\| \geq |h| \cdot \|\hat{\mathbf{w}}_j\| - \|\hat{\mathbf{v}}_j\| \geq 2 \cdot \|\hat{\mathbf{w}}_j\| - \|\hat{\mathbf{v}}_j\| > \|\hat{\mathbf{w}}_j\|$$

contradicting (38). Thus both g, h are among $\{-1, 0, 1\}$. Hence we are left with the vectors of the lemma.

Case 2: \mathbf{b} does not lie in the space spanned by $\mathbf{v}_j, \mathbf{w}_j$. For this case, we use an easy consequence of Minkowski's Second Convex Body Theorem.

Lemma 6.2. *There exists a constant $c > 0$ such that for any $\boldsymbol{\xi} \in \mathbb{R}^2$ and any parameter $Q \geq 1$, the system*

$$|b_1| \leq Q, \quad |b_2| \leq Q, \quad |b_1\xi_1 + b_2\xi_2 + b_3| < cQ^{-2}$$

does not have three linearly independent solutions in integer vectors $\mathbf{b} = (b_1, b_2, b_3)$.

Proof. Consider the integer lattice \mathbb{Z}^3 and the box of $(x_1, x_2, x_3) \in \mathbb{R}^3$ with coordinates

$$|x_1| \leq Q, \quad |x_2| \leq Q, \quad |\xi_1 x_1 + \xi_2 x_2 + x_3| \leq cQ^{-2}.$$

It has volume $8c$, independent of Q and ξ_1, ξ_2 . Hence, by Minkowski's Second Convex Body Theorem, the product of the induced successive minima is $\ll c$, hence choosing c small enough the third successive minimum is smaller than 1. This means there cannot be three linearly independent integer points within the box, which in turn is equivalent to the claim. \square

We first notice that both \mathbf{v}_j and \mathbf{w}_j induce approximations of order greater than two. By (34), (36) and as our choice $(\tau\mu - 1)/\mu = n > 2$, we get

$$(40) \quad |\mathbf{v}_j \cdot \boldsymbol{\xi}^*| = (1 + o(1))r^{-1}A_j^{-(\tau\mu-1)} \ll r^{-1}B_j^{-n} \ll r^{-1}\|\hat{\mathbf{w}}_j\|^{-n} < r^{-1}\|\hat{\mathbf{w}}_j\|^{-2}$$

and for \mathbf{w}_j by (37) we have a stronger estimate that also yields

$$(41) \quad |\mathbf{w}_j \cdot \boldsymbol{\xi}^*| = o(1) \cdot B_j^{-(\tau\mu-1)} = o(\|\hat{\mathbf{w}}_j\|^{-n}) = o(\|\hat{\mathbf{w}}_j\|^{-2}), \quad j \rightarrow \infty.$$

Combined with (35), (38), (39), we have

$$\max\{\|\hat{\mathbf{b}}\|, \|\hat{\mathbf{v}}_j\|, \|\hat{\mathbf{w}}_j\|\} = \|\hat{\mathbf{w}}_j\|, \quad \max\{|\mathbf{b} \cdot \boldsymbol{\xi}^*|, |\mathbf{v}_j \cdot \boldsymbol{\xi}^*|, |\mathbf{w}_j \cdot \boldsymbol{\xi}^*|\} \ll r^{-1}\|\hat{\mathbf{w}}_j\|^{-2}.$$

By the assumptions of Case 2, the three vectors $\mathbf{v}_j, \mathbf{w}_j, \mathbf{b}$ are linearly independent. So we get a contradiction to Lemma 6.2 for r sufficiently large and $Q = \|\hat{\mathbf{w}}_j\|$, as soon as $\|\hat{\mathbf{w}}_j\|$ is sufficiently large. Hence, in total, Case 2 provides only finitely many best approximations, of small norm, so Lemma 6.1 is proved. \square

For studying $\psi_{\xi}^{1 \times 2}$, we may restrict to integer vectors of the form appearing in Lemma 6.1. Now $|\mathbf{w}_j \cdot \xi^*|$ is much larger than $|\mathbf{v}_{j+1} \cdot \xi^*|$, and $|\mathbf{v}_j \cdot \xi^*|$ is much larger than $|\mathbf{w}_j \cdot \xi^*|$ by (29), (36), (37). Hence by the triangle inequality $\mathbf{v}_j \pm \mathbf{w}_j$ induce the same approximation quality as \mathbf{v}_j up to a factor $1 + o(1)$, and similarly, $\mathbf{w}_j \pm \mathbf{v}_{j+1}$ induce the same approximation quality up to $1 + o(1)$ as \mathbf{w}_j , i.e

$$(42) \quad |(\mathbf{v}_j \pm \mathbf{w}_j) \cdot \xi^*| = (1 + o(1)) \cdot |\mathbf{v}_j \cdot \xi^*|, \quad |(\mathbf{v}_{j+1} \pm \mathbf{w}_j) \cdot \xi^*| = (1 + o(1)) \cdot |\mathbf{w}_j \cdot \xi^*|.$$

Moreover as the norm of $\hat{\mathbf{w}}_j$ is much larger than the norm of $\hat{\mathbf{v}}_j$ and the norm of $\hat{\mathbf{v}}_{j+1}$ is much larger than the norm of $\hat{\mathbf{w}}_j$, by (29) and (33), we have by the triangle inequality that

$$(43) \quad \|\hat{\mathbf{v}}_j \pm \hat{\mathbf{w}}_j\| = (1 + o(1))\|\hat{\mathbf{w}}_j\|, \quad \|\hat{\mathbf{v}}_{j+1} \pm \hat{\mathbf{w}}_j\| = (1 + o(1))\|\hat{\mathbf{v}}_{j+1}\|.$$

By (42), (43) and since we have freedom up to a factor $1 + o(1)$ in our claim to be proved, we may assume that all best approximations are among $\mathbf{v}_j, \mathbf{w}_j$. Now we choose r appropriately. To finish the proof we show that choosing τ, μ, r suitably we have

$$(44) \quad |\mathbf{v}_j \cdot \xi^*| = (1 + o(1)) \cdot c\|\hat{\mathbf{w}}_j\|^{-n}, \quad |\mathbf{w}_j \cdot \xi^*| = o(1) \cdot \|\hat{\mathbf{v}}_{j+1}\|^{-n}.$$

Assume this is true. Then indeed partitioning the interval $[\|\mathbf{v}_1\|, \infty)$ into consecutive disjoint intervals of the form $[\|\mathbf{v}_j\|, \|\mathbf{w}_j\|)$ and $[\|\mathbf{w}_j\|, \|\mathbf{v}_{j+1}\|)$ we see

$$t^n \psi_{\xi}^{1 \times 2}(\|\cdot\|, t) \leq c(1 + o(1)) \quad t \rightarrow \infty.$$

On the other hand, since there is no better approximation than the one induced by \mathbf{v}_j for any $t < \|\mathbf{w}_j\|$, for $t_j := \|\mathbf{w}_j\| - 1$ we have the reverse inequality

$$t_j^n \psi_{\xi}^{1 \times 2}(\|\cdot\|, t_j) \geq c(1 + o(1)) \quad j \rightarrow \infty.$$

Combining proves the equality

$$\limsup_{t \rightarrow \infty} t^n \psi_{\xi}^{1 \times 2}(\|\cdot\|, t) = c.$$

For the left identity of (44) equivalently by (34), (36) we need

$$(1 + o(1))r^{-1}A_j^{-(\tau\mu-1)} = c \cdot d_2^{-n}A_j^{-n\mu},$$

so we pick

$$r = c^{-1}d_2^n A_j^{n\mu - \tau\mu + 1} \stackrel{(28)}{=} c^{-1}d_2^n.$$

Similarly, by (33), (37) for the right identity of (44) it suffices to have

$$B_j^{-(\tau\mu-1)} = o(d_1^{-n}A_{j+1}^{-n\tau}).$$

Since $A_{j+1}^{-n\tau} \asymp B_j^{-n\tau^2}$ by (29) and d_1 is constant, it suffices to notice that upon (28) we may choose $\tau < \mu$ so that $n\tau^2 < \tau\mu - 1$. Indeed it suffices to choose τ just slightly larger than n so that $\mu = 1/(\tau - n) > \tau^2$ which is equivalent to $n\tau^2 < \tau\mu - 1$. We leave the short calculation to the reader. We have completed step I. Assuming c is small enough, we apply Step II below.

6.3. Proof of Step II. The proof of step II works analogously to the special case of maximum norm in Section 5.2. Our norm $\|\cdot\|$ on \mathbb{R}^n induces a projected norm $\|\cdot\|'$ on \mathbb{R}^2 via

$$(45) \quad \|(x_1, x_2)\|' := \|(x_1, x_2, 0, \dots, 0)\|.$$

Then for this norm $\|\cdot\|'$ we apply step I to get $\boldsymbol{\xi} = (\xi_1, \xi_2)$ with Dirichlet constant c , i.e.

$$\limsup_{t \rightarrow \infty} t^n \psi_{\boldsymbol{\xi}}^{1 \times 2}(\|\cdot\|', t) = c.$$

Let $\mathbf{b}_v = (b_{1,v}, b_{2,v}, b_{3,v})$ be the integer vector sequence constructed in Step I.

By construction of $\|\cdot\|'$, it is again clear by the same argument as in Proposition 1 that any vector $\boldsymbol{\zeta} = (\zeta_1, \dots, \zeta_n) \in \mathbb{R}^n$ extending $\boldsymbol{\xi}$ has Dirichlet constant at most c with respect to $\|\cdot\|$, i.e.

$$\limsup_{t \rightarrow \infty} t^n \psi_{\boldsymbol{\zeta}}^{1 \times n}(\|\cdot\|, t) \leq \limsup_{t \rightarrow \infty} t^n \psi_{\boldsymbol{\xi}}^{1 \times 2}(\|\cdot\|', t) = c.$$

Indeed, we can choose the embedded integer vectors $\iota(\mathbf{b}_v)$ via

$$\iota : (y_1, y_2, y_3) \rightarrow (y_1, y_2, 0, \dots, 0, y_3)$$

from \mathbb{R}^3 to \mathbb{R}^{n+1} , which by (45) induce the same heights and approximation quality

$$\|\widehat{\iota(\mathbf{b}_v)}\| = \|\widehat{\mathbf{b}_v}\|', \quad |\iota(\mathbf{b}_v) \cdot \boldsymbol{\zeta}^*| = |\mathbf{b}_v \cdot \boldsymbol{\xi}^*|$$

for any \mathbf{b}_v , with hat notation according to (4) on the respective spaces. For the non-trivial lower estimate $\geq c$ with respect to $\|\cdot\|$, define

$$L_v = |\iota(\mathbf{b}_v) \cdot \boldsymbol{\zeta}^*| = |\mathbf{b}_v \cdot \boldsymbol{\xi}^*| = |b_{1,v}\xi_1 + b_{2,v}\xi_2 + b_{3,v}|, \quad M_v = M_v(\|\cdot\|) = \|\widehat{\mathbf{b}_v}\|' = \|\widehat{\iota(\mathbf{b}_v)}\|$$

and $T_v = M_v - 1/2$ as in the proof for the maximum norm but where M_v, T_v are now defined with respect to our norm $\|\cdot\|$. Then by the equivalence of norms on \mathbb{R}^n , for any integer vector of norm $\|(x_1, \dots, x_n)\| \leq T_v$ we still have

$$\max_{1 \leq i \leq n} |x_i| \leq C(\|\cdot\|) \cdot T_v$$

for some constant C depending on the norm only. We replace (23) by this modified inequality and follow the proof of the special case Lemma 5.3 where the maximum norm was treated. The twist by the constant C will result in some altered value of c_n , depending on the chosen norm, for the same conclusion. We omit its explicit calculation in dependence of C .

7. PROOF OF THEOREM 3.6

7.1. Proof excluding linear independence claims. We proceed as in § 6.2 to construct the first two elements of the first line of our matrix Ω by $\boldsymbol{\xi} = (\xi_1, \xi_2) = (\Omega_{1,1}, \Omega_{1,2})$, upon modifying the construction so that (28) is replaced by

$$\tau\mu - 1 = \frac{n}{m}\mu \stackrel{(14)}{\geq} 2\mu.$$

Define $\psi_{\xi}^{1 \times 2}(\|\cdot\|'_1, t)$ with respect to the restriction $\|\cdot\|'_1$ of $\|\cdot\|_1$ to \mathbb{R}^2 in the integer vector \mathbf{b} , as in (45). Then by the same arguments of the proof of Theorem 2.3 we will have

$$(46) \quad \limsup_{t \rightarrow \infty} t^{n/m} \psi_{\xi}^{1 \times 2}(\|\cdot\|'_1, t) = c.$$

Hereby we use our assumption $n/m \geq 2$ for Case 2, based on Minkowski's Second Convex Body Theorem, to work analogously.

Now take the matrix $V \in \mathbb{R}^{m \times 2}$ with all m lines equal to ξ , i.e.

$$V_{j,1} = \xi_1, \quad V_{j,2} = \xi_2, \quad 1 \leq j \leq m.$$

This matrix gives rise to an approximation function

$$\psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t)$$

with norms defined as above. First, assume for simplicity that $\|\cdot\|_2 = \|\cdot\|_{\infty}$ is the maximum norm on \mathbb{R}^m . We claim that

$$(47) \quad \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) = \psi_{\xi}^{1 \times 2}(\|\cdot\|'_1, t), \quad t > 0.$$

Indeed to see

$$(48) \quad \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) \geq \psi_{\xi}^{1 \times 2}(\|\cdot\|'_1, t)$$

for all t , note that for arbitrary $\mathbf{b} = (b_1, \dots, b_{m+2}) \in \mathbb{Z}^{m+2}$ and for $V^* \in \mathbb{R}^{m \times (m+2)}$ obtained by glueing V to the $m \times m$ identity matrix to the right, the induced value

$$\|V^* \cdot \mathbf{b}\|_2 = \|V \cdot \hat{\mathbf{b}} + \tilde{\mathbf{b}}\|_2 = \|V \cdot (b_1, b_2) + (b_3, \dots, b_{m+2})\|_2$$

with vectors interpreted as column vectors, is at least as large as the modulus of any entry, since we are considering $\|\cdot\|_2 = \|\cdot\|_{\infty}$ for now. But for the first entry this equals $|(b_1, b_2) \cdot \xi + b_3| = |\xi^* \cdot (b_1, b_2, b_3)|$. Taking the minimum over \mathbf{b} as in the definition of $\psi_V^{m \times 2}$ shows the inequality (48). Conversely, if we extend arbitrary $\mathbf{b} = (b_1, b_2, b_3) \in \mathbb{Z}^3$ into \mathbb{Z}^{m+2} via $\eta(\mathbf{b}) := (b_1, b_2, b_3, \dots, b_3)$ with last m coordinates equal to b_3 , by choice of maximum norm, we have $\|V^* \cdot \eta(\mathbf{b})\|_2 = |\xi^* \cdot (b_1, b_2, b_3)|$, so passing to the minimum over \mathbf{b} again we get the reverse inequality

$$(49) \quad \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) \leq \psi_{\xi}^{1 \times 2}(\|\cdot\|'_1, t).$$

Combining (46), (47) we get

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) = \limsup_{t \rightarrow \infty} t^{n/m} \psi_{\xi}^{1 \times 2}(\|\cdot\|'_1, t) = c.$$

To finish the proof, we show that for a positive measure set of $B \in \mathbb{R}^{m \times (n-2)}$ inducing $\Omega = (V, B)$ via putting it to the right of V , we have

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_{\Omega}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t) = \limsup_{t \rightarrow \infty} t^{n/m} \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t).$$

Again the inequality

$$(50) \quad \limsup_{t \rightarrow \infty} t^{n/m} \psi_{\Omega}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t) \leq \limsup_{t \rightarrow \infty} t^{n/m} \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t)$$

is easy to check for arbitrary $\|\cdot\|_2$ and extensions $\Omega \in \mathbb{R}^{m \times n}$ of V by an arbitrary matrix $B \in \mathbb{R}^{m \times (n-2)}$. Indeed, for a sufficient estimate

$$\psi_{\Omega}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t) \leq \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t), \quad t > 0$$

we embed the integer best approximations of V by adding 0 coordinates via

$$\sigma : \mathbf{b} = (b_1, b_2, \dots, b_{m+2}) \rightarrow (b_1, b_2, 0, \dots, 0, b_3, \dots, b_{m+2}) \in \mathbb{Z}^{m+n}.$$

These induce the same norm $\|\widehat{\sigma(\mathbf{b})}\|_1 = \|\widehat{\mathbf{b}}\|'_1$ where

$$\widehat{\sigma(\mathbf{b})} = (b_1, b_2, \dots, b_n) = (b_1, b_2, 0, \dots, 0), \quad \widehat{\mathbf{b}} = (b_1, b_2),$$

with hat again as in (4) on the respective spaces. With Ω^* derived from Ω likewise as for V^* , they further induce the same approximation qualities $\|\Omega^* \cdot \sigma(\mathbf{b})\|_2 = \|V^* \cdot \mathbf{b}\|_2$ since in fact $\Omega^* \cdot \sigma(\mathbf{b}) = V^* \cdot \mathbf{b}$, proving (50). For the reverse estimate

$$(51) \quad \limsup_{t \rightarrow \infty} t^{n/m} \psi_{\Omega}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t) \geq \limsup_{t \rightarrow \infty} t^{n/m} \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) = c,$$

we use a generalisation of Step II from § 6.3 for $m > 1$ as in Moshchevitin [25, Theorem 12]. We obtain that for small enough c and a positive measure set of remaining matrices $B \in \mathbb{R}^{m \times (n-2)}$ with respect to $m(n-2)$ dimensional Lebesgue measure when added to the right of V to obtain Ω , we have (51). Hereby we require $n \neq m+2$ for similar reasons as $n \geq 4$ was previously needed in our proof for one linear form, see [25]. Moreover, the assumption $n \geq 2m$ is vital for the volume estimates. See Lemma 7.1 below for a generalisation. The proof for $\|\cdot\|_2$ the maximum norm is complete.

Now let the norm $\|\cdot\|_2$ be arbitrary. Again we prove (48), (49), (50), (51). The critical estimates where the norm $\|\cdot\|_2$ is relevant for the argument are (48), (49). First notice that by the special form of V , if \mathbf{b} is not of the form $\mathbf{b} = (b_1, b_2, b_3, b_3, \dots, b_3) \in \mathbb{Z}^{m+2}$ then some entry of $V^* \cdot \mathbf{b}$ will be of modulus at least $1/2$, so $\|V^* \cdot \mathbf{b}\|_{\infty} \geq 1/2$, hence by equivalence of norms, we deduce

$$\|V^* \cdot \mathbf{b}\|_2 \gg 1$$

with some absolute constant depending on the norm $\|\cdot\|_2$ only. Hence we only need to take into account vectors of the form $\mathbf{b} = (b_1, b_2, b_3, b_3, \dots, b_3)$ when studying $\psi_V^{m \times 2}$. But then the vector $V^* \cdot \mathbf{b} \in \mathbb{R}^m$ lies on the line $\lambda \cdot (1, 1, \dots, 1) \in \mathbb{R}^m$ and has norm $|\lambda| \cdot \|(1, 1, \dots, 1)\|_2$ proportional to $|\lambda|$. Hence we have to minimize $|\lambda|$. In § 6.2, see in particular Lemma 6.1, we showed that this is achieved when the subvector (b_1, b_2, b_3) is essentially of the form $\mathbf{v}_j = (A_j, 0, -F_j)$ and $\mathbf{w}_j = (0, B_j, -G_j)$ for A_j, B_j, F_j, G_j defined in the same section (as shown in § 6.2, for the other possible vectors of Lemma 6.1 we may only improve the approximation quality by a negligible factor $1 + o(1)$ as $j \rightarrow \infty$). These arguments show that the best integer approximations for V are essentially of the form

$$(52) \quad \mathbf{c}_j := (A_j, 0, -F_j, \dots, -F_j), \quad \mathbf{d}_j := (0, B_j, -G_j, \dots, -G_j)$$

so that for $\mathbf{v}_j, \mathbf{w}_j \in \mathbb{Z}^3$ as before, we get error vectors

$$V^* \cdot \mathbf{a}_j = V \cdot \begin{pmatrix} A_j \\ 0 \end{pmatrix} - F_j \cdot \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} = \mathbf{v}_j \cdot \boldsymbol{\xi}^* \cdot \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}^m$$

and

$$V^* \cdot \mathbf{b}_j = V \cdot \begin{pmatrix} 0 \\ B_j \end{pmatrix} - G_j \cdot \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} = \mathbf{w}_j \cdot \boldsymbol{\xi}^* \cdot \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \end{pmatrix} \in \mathbb{R}^m.$$

Then for $\Gamma := \|(1, 1, \dots, 1)\|_2$ they have norms

$$\|V^* \cdot \mathbf{c}_j\|_2 = |\mathbf{v}_j \cdot \boldsymbol{\xi}^*| \cdot \Gamma, \quad \|V^* \cdot \mathbf{d}_j\|_2 = |\mathbf{w}_j \cdot \boldsymbol{\xi}^*| \cdot \Gamma.$$

Thus we have to adjust the proof of the special case by the constant factor Γ in the calculation, so that in order to finalize Step I we now need to pick

$$r = c^{-1} \Gamma^{-1} d_2^m, \quad d_2 = \|\mathbf{e}_2\|'_1 = \|(0, 1)\|'_1$$

to get (49). The reverse inequality (48) holds as well as we noticed that the vectors in (52) are (essentially) the integer best approximations. We observed that (50) holds independent of $\|\cdot\|_2$.

Finally, Step II is to again show the reverse inequalities (51) hold. This is done analogously to Lemma 5.3 but in higher dimension, as in Moshchevitin [25, Theorem 12]. Again essentially irrespective of the norm $\|\cdot\|_2$. We state the according analogue of Lemma 5.3 that we will prove analogously but without effort to make constants effective.

Lemma 7.1. *Let m, n be positive integers with $n \geq 2$ and $n \neq m + 2$. Let $\|\cdot\|_1$ and $\|\cdot\|_2$ be any norms on \mathbb{R}^n and \mathbb{R}^m and $\|\cdot\|'_1$ be the restriction of $\|\cdot\|_1$ to \mathbb{R}^2 as in (45). Further, let $V \in \mathbb{R}^{m \times 2}$ be any matrix with*

$$(53) \quad \limsup_{t \rightarrow \infty} t^{n/m} \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) = c$$

for small enough $c > 0$, depending on m, n . Let $\mathbf{b}_v \in \mathbb{Z}^{m+2}$ be the associated sequence of best approximations, and $\hat{\mathbf{b}}_v = (b_{v,1}, b_{v,2})$ of norms and approximation qualities

$$M_v := \|\hat{\mathbf{b}}_v\|'_1, \quad L_v := \|V^* \cdot \mathbf{b}_v\|_2.$$

Then, for a positive measure set $\mathcal{Y} \subseteq B_{m(n-2)}(\mathbf{0}, 1) \subseteq \mathbb{R}^{m(n-2)}$ consisting of matrices $B \in \mathbb{R}^{m \times (n-2)}$, we have that for infinitely many integers $v \geq 1$, letting $T_v := M_{v+1} - 1/2$, the integer best approximation for $\Omega = (V, B) \in \mathbb{R}^{m \times n}$ at T_v is given as

$$b_{1,v} \mathbf{e}_1 + b_{2,v} \mathbf{e}_2 + b_{3,v} \mathbf{e}_{n+1} + \dots + b_{m+2,v} \mathbf{e}_{n+m} = (b_{1,v}, b_{2,v}, 0, \dots, 0, b_{3,v}, \dots, b_{m+2,v}) \in \mathbb{Z}^{n+m},$$

thus the best approximation function at values T_v equals

$$\psi_\Omega^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, T_v) = \psi_V^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, T_v) = L_v.$$

Proof. Fix $V \in \mathbb{R}^{m \times 2}$. As in the proof of Lemma 5.3 we construct the set \mathcal{Y} to be composed of matrices $B \in \mathcal{Y}$ such that the matrices $\Omega = (V, B) \in \mathbb{R}^{m \times n}$ satisfy

$$\psi_{\Omega}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, T_v) = \psi_V^{m \times 2}(\|\cdot\|_1', \|\cdot\|_2, T_v) = L_v.$$

By the equivalence of norms we have that there exists $C(\|\cdot\|_1) > 0$ dependent only on $\|\cdot\|_1$ such that

$$\|\hat{\mathbf{x}}\|_1 \leq T_v \quad \implies \quad \|\hat{\mathbf{x}}\|_{\infty} \leq C(\|\cdot\|_1)T_v.$$

Thus if again $\Omega^* = (V, B)^* \in \mathbb{R}^{m \times (m+n)}$ is obtained by putting the $m \times m$ identity matrix to the right of Ω and we let

$$S_v := \left\{ B \in B_{m(n-2)}(\mathbf{0}, 1) : \min_{\substack{\mathbf{x} \in \mathbb{Z}^{n+m} \\ 0 < \|\hat{\mathbf{x}}\|_{\infty} \leq C(\|\cdot\|_1)T_v}} \|(V, B)^* \cdot \mathbf{x}\|_2 \geq L_v \right\},$$

then

$$\mathcal{Y} \supseteq \limsup_{v \rightarrow \infty} S_v.$$

Hence if we can show that the set

$$(54) \quad \lambda_{m(n-2)} \left(\liminf_{v \rightarrow \infty} S_v^c \right) \leq \delta < \lambda_{m(n-2)} (B_{m(n-2)}(\mathbf{0}, 1))$$

then clearly $\lambda_{m(n-2)}(\mathcal{Y}) > 0$. Furthermore, by showing that

$$(55) \quad \lambda_{m(n-2)}(S_v^c) \leq \delta \quad v \geq 1,$$

and applying Proposition 2 we have (54), so proving (55) is sufficient to verify $\lambda_{m(n-2)}(\mathcal{Y}) > 0$.

Let $z = (z_3, \dots, z_n)$, $y = (x_{n+1}, \dots, x_{n+m})$ and

$$X_v(z, x_1, x_2, y) = \left\{ \gamma \in B_{m(n-2)}(\mathbf{0}, 1) : \begin{pmatrix} z_3 \gamma_{1,1} + \dots + z_n \gamma_{1,n-2} \\ \vdots \\ z_3 \gamma_{m,1} + \dots + z_n \gamma_{m,n-2} \end{pmatrix} \in J_v(x_1, x_2, y) \right\}$$

where

$$J_v(x_1, x_2, y) := \prod_{i=1}^m (-x_1 \xi_1 - x_2 \xi_2 - x_{n+i} - C(\|\cdot\|_2)L_v, -x_1 \xi_1 - x_2 \xi_2 - x_{n+i} + C(\|\cdot\|_2)L_v).$$

Here we have again used equivalence of norms, namely that there exists $C(\|\cdot\|_2) > 0$ dependent only on $\|\cdot\|_2$ so that for any $X \in \mathbb{R}^m$

$$C(\|\cdot\|_2)^{-1} \|X\|_{\infty} \leq \|X\|_2 \leq C(\|\cdot\|_2) \|X\|_{\infty}.$$

Considering one row of $X_v(z, x_1, x_2, y)$ at a time (denoted $X_v^{(i)}(z, x_1, x_2, y)$ for $1 \leq i \leq m$) we have the same calculation as in (27), that is

$$\lambda_{n-2} (X_v^{(i)}(z, x_1, x_2, y)) \leq 2C(\|\cdot\|_2)L_v \lambda_{n-3}(B_{n-3}(\mathbf{0}, 1)) \frac{1}{\|z\|_{\infty}}.$$

Hence

$$\begin{aligned} \lambda_{m(n-2)}(X_v(z, x_1, x_2, y)) &\leq 2^m C(\|\cdot\|_2)^m L_v^m \lambda_{n-3}(B_{n-3}(\mathbf{0}, 1))^m \frac{1}{\|z\|_\infty^m} \\ &= C(m, n, \|\cdot\|_1, \|\cdot\|_2) L_v^m \frac{1}{\|z\|_\infty^m}. \end{aligned}$$

Using similar calculations as appearing in Lemma 5.3 we have

$$\begin{aligned} \lambda_{m(n-2)}(S_v^c) &\leq \sum_{0 < \max_{i=1,2} |x_i| \leq C(\|\cdot\|_1) T_v} \sum_{\|y\|_\infty \leq C(\|\cdot\|_1) T_v} \sum_{\|z\|_\infty \leq C(\|\cdot\|_1) T_v} \lambda_{m(n-2)}(X_v(z, x_1, x_2, y)) \\ &\leq C(m, n, \|\cdot\|_1, \|\cdot\|_2) L_v^m T_v^{m+2} \sum_{\|z\|_\infty \leq C(\|\cdot\|_1) T_v} \frac{1}{\|z\|_\infty^m} \\ &\leq C'(m, n, \|\cdot\|_1, \|\cdot\|_2) L_v^m T_v^{m+2} \sum_{k=1}^{[C(\|\cdot\|_1) T_v]} k^{n-3-m} \\ &\stackrel{(n \neq m+2)}{\leq} C''(m, n, \|\cdot\|_1, \|\cdot\|_2) L_v^m T_v^{m+2} T_v^{1+n-3-m} \\ &\leq C''(m, n, \|\cdot\|_1, \|\cdot\|_2) L_v^m T_v^n. \end{aligned}$$

Thus, for any $0 < \delta < \lambda_{m(n-2)}(B_{m(n-2)}(\mathbf{0}, 1))$, if $L_v < c T_v^{-\frac{n}{m}}$ then for any $c \in (0, C'''(m, n, \|\cdot\|_1))$ with

$$C'''(m, n, \|\cdot\|_1, \|\cdot\|_2) < \left(\frac{\delta}{C''(m, n, \|\cdot\|_1, \|\cdot\|_2)} \right)^{\frac{1}{m}},$$

we have that

$$\lambda_{m(n-2)}(S_v^c) < \delta$$

as required. \square

There is nothing essential about V having two columns, the lemma can readily be generalized. The condition $n \neq m + 2$ entered in the proof for similar reasons as the requirement $n \geq 4$ in Lemma 5.3, see also [25]. (This condition needs to be adapted to $n \neq m + m_0$ when V has m_0 columns.) All claims of Theorem 3.6 apart from the linear independence statements are proved.

7.2. Proof of linear independence claims. The first claim (i) on linear independence of rows follows from Step II. It suffices to prove linearly independent rows can be arranged for the submatrix B of $\Omega = (V, B)$. Now since $n - 2 \geq m$ by (14), the matrix B does not have fewer columns than rows, so if B does not have the property it is rank deficient. However, the set of rank-deficient matrices has zero Lebesgue measure as all its $m \times m$ minors must have determinant 0, so essentially at least one coordinate is a function of the others, leading to positive codimension. On the other hand, we showed that a positive Lebesgue measure set of matrices B gives rise to Ω as in the theorem. Hence removing such rank-deficient matrices still leaves us with a set of positive $m \times (n - 2)$ -dimensional Lebesgue measure from which we can choose suitable B .

For the latter claim (ii) on \mathbb{Q} linearly independent entries, note that in the construction of Section 7.1 certain entries of V (thus Ω) are identical, so the condition does not hold.

To fulfil this additional property, let us modify our construction by varying signs. First construct the submatrix $V \in \mathbb{R}^{m \times 2}$ as above by taking for A_j, B_j as before

$$\Omega_{k,1} = V_{k,1} = \sum_{j=1}^{\infty} \delta_{k,j} A_j^{-1}, \quad \Omega_{k,2} = V_{k,2} = \sum_{j=1}^{\infty} \delta_{k,j}^* B_j^{-1}, \quad (1 \leq k \leq m)$$

with sequences $\delta_{k,j} \in \{-1, 1\}$, $\delta_{k,j}^* \in \{-1, 1\}$ to be chosen later. When $\delta_{k,j} = \delta_{k,j}^* = 1$ for all k, j we are in the standard case of the previous sections.

When $m = 1$, all arising matrices/vectors $(V_{1,1}, V_{1,2}) = (\xi_1, \xi_2)$ share the properties of the standard case above. For $m > 1$, we get more general error vectors than in (52), now of the form

$$(56) \quad \tilde{\mathbf{c}}_j := V \cdot \begin{pmatrix} A_j \\ 0 \end{pmatrix} - \begin{pmatrix} F_{j,1} \\ F_{j,2} \\ \vdots \\ F_{j,m} \end{pmatrix}, \quad \tilde{\mathbf{d}}_j := V \cdot \begin{pmatrix} 0 \\ B_j \end{pmatrix} - \begin{pmatrix} G_{j,1} \\ G_{j,2} \\ \vdots \\ G_{j,m} \end{pmatrix}, \quad j \geq 1,$$

where

$$F_{j,k} = A_j \sum_{u=1}^j \delta_{k,u} A_u^{-1}, \quad G_{j,k} = B_j \sum_{u=1}^j \delta_{k,u}^* B_u^{-1}, \quad (1 \leq k \leq m),$$

depending on the choice of $\delta_{k,j}, \delta_{k,j}^*$. They again satisfy congruence properties

$$F_j \equiv \pm 1 \pmod{6} \quad \text{and} \quad G_j \equiv \pm 1 \pmod{35}$$

for the same reasons, sufficient for coprimality with 6 respectively 35. Thus the entries of the remainders $\tilde{\mathbf{c}}_j, \tilde{\mathbf{d}}_j$ are given as

$$c_{j,k} = A_j \sum_{u=j+1}^{\infty} \delta_{k,u} A_u^{-1}, \quad d_{j,k} = B_j \sum_{u=j+1}^{\infty} \delta_{k,u}^* B_u^{-1}, \quad (1 \leq k \leq m).$$

However, since $A_j/A_{j+1} \rightarrow 0$ and $B_j/B_{j+1} \rightarrow 0$ we again see

$$c_{j,k} = \delta_{k,j+1} \cdot A_j A_{j+1}^{-1} (1 + o(1)), \quad d_{j,k} = \delta_{k,j+1}^* \cdot B_j B_{j+1}^{-1} (1 + o(1)), \quad (1 \leq k \leq m).$$

Hence, by the equivalence of norms, it is clear that the error vectors in (56) all have $\|\cdot\|_2$ norm of order respectively

$$(57) \quad \|\tilde{\mathbf{c}}_j\|_2 = A_j A_{j+1}^{-1} \Gamma_j (1 + o(1)), \quad \|\tilde{\mathbf{d}}_j\|_2 = B_j B_{j+1}^{-1} \tilde{\Gamma}_j (1 + o(1)), \quad j \rightarrow \infty,$$

with $\Gamma_j, \tilde{\Gamma}_j$ of one of the numbers

$$\Gamma_j, \tilde{\Gamma}_j \in \{\|(\pm 1, \pm 1, \dots, \pm 1)\|_2\}, \quad j \geq 1,$$

for any possible sign choice. The values $\tilde{\Gamma}_j$ do not matter much as the according evaluations will be of negligible order by the analogue of (44). Now if we let

$$\Gamma := \max\{\|(\pm 1, \pm 1, \dots, \pm 1)\|_2\}$$

with maximum overall 2^m sign choices, we can use the argument of Section 7.1. Hereby we may assume that the maximizing sign choice occurs for infinitely many $(\delta_{1,j}, \dots, \delta_{m,j})$,

as otherwise we redefine Γ via the maximum among those m term sign sequences occurring infinitely often. Moreover, due to (57), the arguments of Section 6.2 applied to each of the m coordinates separately still apply very similarly and yield that the vectors $(A_j, 0, -F_{j,1}, \dots, -F_{j,m})$ and $(0, B_j, -G_{j,1}, \dots, -G_{j,m})$ essentially form the sequence of best approximations for the induced V . We leave the details to the reader. Then very similar arguments as in the classical case including Lemma 7.1 show that the Dirichlet constant $\Theta^{m \times n}(\Omega)$ of many such $\Omega = \Omega(\delta, \delta^*)$ equals c , as before.

Finally we show that for many choices of $\delta_{k,j}, \delta_{k,j}^*$ all entries of $\Omega \in \mathbb{R}^{m \times n}$ are linearly independent over \mathbb{Q} together with $\{1\}$. Let us first show this for the submatrix $V \in \mathbb{R}^{m \times 2}$. Label the entries $V_{i,j}$ of V by v_1, \dots, v_{2m} with $v_{2h} = V_{h,2}$ and $v_{2h-1} = V_{h,1}$, $1 \leq h \leq m$. Now the first line of V is \mathbb{Q} -linearly independent for any choice of $\delta_{1,j}, \delta_{1,j}^*$ sequences as otherwise the sequence of best approximations for $(V_{1,1}, V_{1,2})$ would terminate, but the proof above shows that this is not the case. Hence $\{1, v_1, v_2\}$ is \mathbb{Q} -linearly independent. Then we proceed inductively. Having constructed the first $\ell \geq 2$ elements v_1, \dots, v_ℓ , we choose the δ or δ^* sequence for $v_{\ell+1}$ so that we avoid \mathbb{Q} -linear dependence of $v_{\ell+1}$ with the already chosen $1, v_1, \dots, v_\ell$. This is possible since the \mathbb{Q} -span of $\{1, v_1, \dots, v_\ell\}$ is countable, but we have uncountably many choices of δ or δ^* sequences. These clearly induce uncountably many pairwise distinct numbers $v_{\ell+1}$, the injectivity of the corresponding maps $\varphi : \{-1, 1\}^{\mathbb{N}} \rightarrow \mathbb{R}$ resp. $\varphi^* : \{-1, 1\}^{\mathbb{N}} \rightarrow \mathbb{R}$ follows from the rapid increase of the sequences A_j resp. B_j . Repeating this for $\ell = 2, 3, \dots, 2m-1$, finally all entries of V will be independent over \mathbb{Q} with $\{1\}$, as desired.

Now Step II of the proof above combined with the following easy argument show that we can transition from V to Ω .

Proposition 5. *Let g, h be positive integers. Assume given real numbers x_1, \dots, x_g are \mathbb{Q} -linearly independent together with 1. Then only for a set of Hausdorff dimension $h-1$ of vectors $\mathbf{y} = (y_1, \dots, y_h)$ in \mathbb{R}^h , the joint vector $\mathbf{z} = (x_1, \dots, x_g, y_1, \dots, y_h)$ is \mathbb{Q} -linearly dependent with 1.*

Proof. Since the x_j are independent, if $\{1, \mathbf{z}\}$ is dependent over \mathbb{Q} then there are integers M_i, N_i, N with

$$N + \sum_{i=1}^g M_i x_i + \sum_{i=1}^h N_i y_i = 0$$

and not all N_i are 0. But this means that \mathbf{y} lies in a countable union of affine hyperplanes of \mathbb{R}^h . Hence we see that the set of \mathbf{y} inducing dependency over \mathbb{Q} indeed has Hausdorff dimension at most $h-1$. \square

It suffices to take $g = 2m, h = (n-2)m$ and to choose x_i the entries of V (first two columns of Ω) and y_i the remaining entries of Ω (entries of B) in the proposition to conclude that a full dimensional set of possible matrices $B \in \mathbb{R}^{m \times (n-2)} \cong \mathbb{R}^{m(n-2)}$ remains (in fact of positive $m(n-2)$ dimensional Hausdorff measure).

8. PROOFS OF THEOREMS 3.3 AND 3.4

Theorems 3.3 admits a rather easy proof, as a consequence of the following simple lemma. We consider maximum norms on both \mathbb{R}^n and \mathbb{R}^m and simply write $\psi_\Omega^{m \times n}(t)$ for $\psi_\Omega^{m \times n}(\|\cdot\|_\infty, \|\cdot\|_\infty, t)$.

Lemma 8.1. *Let $\Omega \in \mathbb{R}^{m \times n}$ be a block matrix consisting of rectangular blocks $\Omega_1, \dots, \Omega_k$ of sizes (m_j, n_j) , $1 \leq j \leq k$ along the diagonal, $m = \sum m_j$, $n = \sum n_j$. Then*

$$\psi_\Omega^{m \times n}(t) = \min_{1 \leq j \leq k} \psi_{\Omega_j}^{m_j \times n_j}(t), \quad t > 0.$$

Proof. First observe that by the form of Ω , the problem decomposes into k decoupled linear form approximation problems. Fix t and let $\mathbf{b}_1, \dots, \mathbf{b}_k$ in $\mathbb{Z}^{m_j + n_j}$ be the best approximation integer vectors inducing $\psi_{\Omega_j}^{m_j \times n_j}(t)$ for $1 \leq j \leq k$, i.e. $\mathbf{b} = \mathbf{b}_j$ minimizes $\|\Omega_j^* \mathbf{b}\|_\infty = \|\Omega_j \hat{\mathbf{b}} - \tilde{\mathbf{b}}\|_\infty$ among integer vectors \mathbf{b} with $0 < \|\mathbf{b}\|_\infty \leq t$. The inequality

$$\psi_\Omega^{m \times n}(t) \leq \min_{1 \leq j \leq k} \psi_{\Omega_j}^{m_j \times n_j}(t)$$

follows by taking the integer vector $\mathbf{b} \in \mathbb{Z}^{m+n}$ with entries of \mathbf{b}_{j_0} at the according places, with j_0 inducing the minimum of $\|\Omega_j^* \mathbf{b}_j\|_\infty$ over j , and 0 entries at all other places. By choice of the maximum norm, this will have the same norm

$$\|\Omega^* \cdot \mathbf{b}\|_\infty = \|\Omega_{j_0}^* \cdot \mathbf{b}_{j_0}\|_\infty = \psi_{\Omega_{j_0}}^{m_{j_0} \times n_{j_0}}(t) = \min_{1 \leq j \leq k} \psi_{\Omega_j}^{m_j \times n_j}(t).$$

Conversely, since some coordinate of the best approximation vector for given t is not 0 and the system is decoupled and we use maximum norm, we must have

$$\psi_\Omega^{m \times n}(t) \geq \min_{1 \leq j \leq k} \psi_{\Omega_j}^{m_j \times n_j}(t).$$

□

The deduction of the case $m = n$ works as follows.

Proof of Theorem 3.3. As in Lemma 8.1, take a matrix of the form $\Omega = \text{diag}(\Omega_1, \Omega_2)$ with diagonal blocks $\Omega_1 \in \text{Bad}_{n-1, n-1}$ and $\Omega_2 = \{\eta\} \notin \text{Bad}_{1,1} \cup \mathbb{Q}$ are arbitrary. Keep in mind that as $n/n = (n-1)/(n-1) = 1/1 = 1$, the Dirichlet exponent (remark: not constant) is 1 for all Ω_1, η and Ω , for all arguments below.

By Lemma 8.1 we have

$$(58) \quad \Theta^{n \times n}(\Omega) = \limsup_{t \rightarrow \infty} t \cdot \psi_\Omega^{n \times n}(t) = \limsup_{t \rightarrow \infty} t \cdot \min\{\psi_{\Omega_1}^{(n-1) \times (n-1)}(t), \psi_\eta^{1 \times 1}(t)\}.$$

Now since $\Omega_1 \in \text{Bad}_{n-1, n-1} \subseteq \text{Di}_{n-1, n-1}$ we have

$$\Theta^{(n-1) \times (n-1)}(\Omega_1) = \limsup_{t \rightarrow \infty} t^{\frac{n-1}{n-1}} \cdot \psi_{\Omega_1}^{(n-1) \times (n-1)}(t) = \limsup_{t \rightarrow \infty} t \cdot \psi_{\Omega_1}^{(n-1) \times (n-1)}(t) < 1$$

so $\Omega \in \text{Di}_{n, n}$ as well by (58). Moreover since

$$\limsup_{t \rightarrow \infty} t \psi_\eta^{1 \times 1}(t) \geq 1/2 > 0$$

for any irrational η by Khintchine [18] and

$$\liminf_{t \rightarrow \infty} t \psi_{\Omega_1}^{(n-1) \times (n-1)}(t) > 0$$

since $\Omega_1 \in \text{Bad}_{n-1, n-1}$, again by (58) we have $\Theta^{n \times n}(\Omega) > 0$ and thus $\Omega \notin \text{Sing}_{n, n}$. Finally again by Lemma 8.1

$$\tilde{\Theta}^{n \times n}(\Omega) = \liminf_{t \rightarrow \infty} t^{n/n} \psi_{\Omega}(t) \leq \liminf_{t \rightarrow \infty} t^{1/1} \psi_{\eta}(t) = 0$$

since $\eta \notin \text{Bad}_{1,1}$, so $\Omega \notin \text{Bad}_{n, n}$. Hence $\Omega \in FS_{n, n}$.

It is well-known that $\text{Bad}_{n-1, n-1}$ has full Hausdorff dimension $(n-1)^2$ and that the complement of $\text{Bad}_{1,1}$ has full 1-dimensional Lebesgue measure, see Section 1. Removing the countable set \mathbb{Q} clearly has no effect on the Lebesgue measure. In particular, we can choose regular (full \mathbb{R} -rank) matrices B as the complementary set has strictly lower Hausdorff dimension, and by the diagonal form and $\eta \neq 0$ the same will be true for any arising Ω . This argument together with (20) also shows that the dimension of the set of matrices Ω constructed above is at least $(n-1)^2 + 1$. \square

We want to formulate another consequence of Lemma 8.1 for maximum norms that is used in the deduction of some claims of Theorem 3.2.

Corollary 8.2. *With $\mathcal{D}_{m, n}$ as in (10), for any positive integers m, n, k and $c > 0$ we have*

$$\mathcal{D}_{m, n} \subseteq \mathcal{D}_{km, kn},$$

and

$$\dim_H FS_{m, n} \leq \dim_H FS_{km, kn}, \quad \dim_H FS_{m, n}(c) \leq \dim_H FS_{km, kn}(c).$$

Proof. For any $m \times n$ matrix Ω , by Lemma 8.1 the $km \times kn$ -matrix $\Omega' := \text{diag}(\Omega, \dots, \Omega)$ with k diagonal blocks Ω induces the same approximation function

$$\psi_{\Omega}^{m \times n}(t) = \psi_{\Omega'}^{km \times kn}(t), \quad t > 0.$$

Since the Dirichlet exponents coincide via $(kn)/(km) = n/m$, the claims follow directly. \square

For the proof of Theorem 3.4 we combine Lemmas 8.1, 7.1 with the main result from [27] on simultaneous approximation $n = 1$.

Proof of Theorem 3.4. Let us introduce

$$\ell := \left\lceil \frac{m}{n} \right\rceil \geq 2$$

so that

$$(59) \quad (\ell - 1)n < m \leq \ell n.$$

For simplicity, first we treat the case $\ell = 2$ (that suffices for the sake of Theorem 3.1).

Case $\ell = 2$: Then

$$(60) \quad \frac{1}{2} \leq \frac{n}{m} < 1.$$

Moreover we can assume $m \geq 4$ and $n \geq 2$, as the otherwise our exclusion of $(m, n) = (3, 2)$ forces $n = 1$, and this case is a trivial consequence of [27]. Let us first consider maximum norms

$$(61) \quad \|\cdot\|_1 = \|\cdot\|_2 = \|\cdot\|_\infty.$$

Then [27, Theorem 2.2] in the case of simultaneous approximation to two numbers can be applied to $\Phi(t) = ct^{-n/m}$ for any $c \in (0, 1)$, as the conditions (d1)-(d3) of [27] are satisfied by our choice of Φ and (60). This yields $\xi \in \mathbb{R}^{2 \times 1}$ with

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_\xi^{2 \times 1}(t) = c, \quad \text{and} \quad \liminf_{t \rightarrow \infty} t^{n/m} \psi_\xi^{2 \times 1}(t) = 0.$$

Now let $\zeta \in \text{Bad}_{m-2,1}$ arbitrary unless if $m = 4$ then $\zeta = \xi$. Since $m - 2 \geq 2$ by assumption, it is easily seen that in either case

$$\psi_\xi^{2 \times 1}(t) \leq \psi_\zeta^{(m-2) \times 1}(t), \quad t \geq t_0.$$

Hence by Lemma 8.1 the matrix $U = \text{diag}(\xi, \zeta) \in \mathbb{R}^{m \times 2}$ satisfies

$$(62) \quad \limsup_{t \rightarrow \infty} t^{n/m} \psi_U^{m \times 2}(t) = c, \quad \liminf_{t \rightarrow \infty} t^{n/m} \psi_U^{m \times 2}(t) = 0,$$

as well.

If $n = 2$ then we must have $m = 4$ due to our assumptions $\frac{n}{m} \geq \frac{1}{2}$ and $m \geq 4$, and then we can just take $\Omega = U$. Note also that in this case our claimed bound for Hausdorff dimension becomes zero, so there is nothing more to prove.

If $n \geq 3$, then in Lemma 7.1 take $V = U \in \mathbb{R}^{m \times 2}$. Note that the problematic case $n = m + 2$ of the lemma is excluded here via (60). Notice further V satisfies (53) by (62). Hence, assuming $c > 0$ is small enough, there exists a set of positive $m(n-2)$ -dimensional Lebesgue measure $\mathcal{Y} \subseteq \mathbb{R}^{m \times (n-2)}$ of matrices so that for any $B \in \mathcal{Y}$, the $\mathbb{R}^{m \times n}$ matrix $\Omega = (B, U)$ satisfies

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_\Omega^{m \times n}(t) = c.$$

Finally, we show Ω is not badly approximable. By (62) there exists a sequence of best approximations such that on the sequence $(t_j)_{j \geq 1}$ we have

$$\lim_{j \rightarrow \infty} t_j^{n/m} \psi_U^{m \times 2}(t_j) = 0.$$

Considering Ω , take the same sequence of the best approximations and place a zero in all entries not related to U (that is starting from the third) to obtain the same statement, i.e.

$$\lim_{j \rightarrow \infty} t_j^{n/m} \psi_\Omega^{m \times n}(t_j) = 0.$$

Combining these two claims we infer $\Omega \in FS_{m,n}(c)$. Using (20) the dimension statement follows since we have shown that (when $m \geq 5$)

$$\dim_H FS_{m,n}(c) \geq \dim_H \mathcal{Y} + \dim_H \text{Bad}_{m-2,1} = m(n-2) + m - 2,$$

where we used that $\text{Bad}_{m-2,1}$ has full Hausdorff dimension and \mathcal{Y} has positive Lebesgue measure. In the case $(m, n) = (4, 3)$ we obtain

$$\dim_H FS_{4,3}(c) \geq \dim_H \mathcal{Y} = m(n-2) = 4.$$

We remark that for $(m, n) = (4, 2)$ the according estimate holds as well, but the right hand side vanishes. This completes the proofs for the maximum norm, i.e. (61).

Now let $\|\cdot\|_1$ be arbitrary and still $\|\cdot\|_2 = \|\cdot\|_\infty$. Let $\|\cdot\|'_1$ be the projected norm

$$\|(x_1, x_2)\|'_1 := \|(x_1, x_2, 0, \dots, 0)\|_1.$$

Let

$$\gamma = \|e_1\|'_1$$

the norm of the first base vector of \mathbb{R}^2 . By [27, Theorem 2.2] and again (60), for small enough $c > 0$ we may pick $\boldsymbol{\xi} \in \mathbb{R}^{2 \times 1}$ such that

$$(63) \quad \limsup_{t \rightarrow \infty} t^{n/m} \psi_{\boldsymbol{\xi}}^{2 \times 1}(t) = c_1 := c\gamma^{n/m}, \quad \liminf_{t \rightarrow \infty} t^{n/m} \psi_{\boldsymbol{\xi}}^{2 \times 1}(t) = 0.$$

Suppose we can show that, for any $U = \text{diag}(\boldsymbol{\xi}, \boldsymbol{\zeta}) \in \mathbb{R}^{m \times 2}$ as above, which we can identify with elements in $\{\boldsymbol{\xi}\} \times \text{Bad}_{m-2,1} \subseteq \mathbb{R}^m$, we have that

$$(64) \quad \psi_U^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) = \psi_{\boldsymbol{\xi}}^{2 \times 1}(|\cdot|, \|\cdot\|_\infty, \gamma^{-1}t), \quad t \geq t_0.$$

If (64) holds, then indeed

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_U^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) = \gamma^{-n/m} \limsup_{t \rightarrow \infty} t^{n/m} \psi_{\boldsymbol{\xi}}^{2 \times 1}(|\cdot|, \|\cdot\|_\infty, \gamma^{-1}t) = \gamma^{-n/m} c_1 = c,$$

and

$$\liminf_{t \rightarrow \infty} t^{n/m} \psi_U^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) = \gamma^{-n/m} \liminf_{t \rightarrow \infty} t^{n/m} \psi_{\boldsymbol{\xi}}^{2 \times 1}(|\cdot|, \|\cdot\|_\infty, \gamma^{-1}t) = 0.$$

Hence for c small enough we again we may again apply Lemma 7.1, to find some set $\mathcal{Y} \subseteq \mathbb{R}^{m \times (n-2)}$ of positive Lebesgue measure such that for any $B \in \mathcal{Y}$ we have

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_{(U,B)}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t) = c, \quad \liminf_{t \rightarrow \infty} t^{n/m} \psi_{(U,B)}^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, t) = 0.$$

So $(U, B) \in FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c)$, and analogously to the maximum norm by (20) we conclude

$$\dim_H FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \geq \dim_H \mathcal{Y} + \dim_H \text{Bad}_{m-2,1} = m(n-2) + m - 2,$$

for $m \geq 5$, and accordingly for $(m, n) = (4, 3)$ we get the bound

$$\dim_H FS_{4,3}(\|\cdot\|_1, \|\cdot\|_2, c) \geq m(n-2) = 4.$$

Since $[m/n] = 2$ by (60), we indeed recognize these as special cases of our claim. Thus to complete the proof for $\ell = 2$ we need to show (64) holds.

For the inequality

$$(65) \quad \psi_U^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) \leq \psi_{\boldsymbol{\xi}}^{2 \times 1}(|\cdot|, \|\cdot\|_\infty, \gamma^{-1}t), \quad t \geq t_0$$

observe that any $(x, \mathbf{w}) \in \mathbb{Z} \times \mathbb{Z}^2$ with $|x| \leq \gamma^{-1}t$ gives rise to $\mathbf{z} = (x, 0) = x\mathbf{e}_1 \in \mathbb{Z}^2$ of norm $\|\mathbf{z}\|'_1 = |x|\gamma \leq t$. Moreover since

$$(66) \quad U \cdot \mathbf{z} - (\mathbf{w}, 0, \dots, 0) = (\boldsymbol{\xi} \cdot x - \mathbf{w}, 0, \dots, 0) \in \mathbb{R}^m,$$

with the same first two coordinates followed by $m-2$ zeros, we have the identity

$$(67) \quad \|U \cdot \mathbf{z} - (\mathbf{w}, 0, \dots, 0)\|_\infty = \|\boldsymbol{\xi} \cdot x - \mathbf{w}\|_\infty,$$

where we used $\|\cdot\|_2$ being the maximum norm. As t is arbitrary (65) is implied.

Now consider the reverse inequality

$$(68) \quad \psi_U^{m \times 2}(\|\cdot\|'_1, \|\cdot\|_2, t) \geq \psi_{\xi}^{2 \times 1}(\|\cdot\|, \|\cdot\|_{\infty}, \gamma^{-1}t), \quad t \geq t_0.$$

Begin with $(m, n) \neq (4, 2)$. Note that if $\mathbf{z} = (z_1, z_2) \in \mathbb{Z}^2$ with $\|\mathbf{z}\|'_1 \leq t$ does not lie in the space $z_2 = 0$, then by equivalence of norms for any $\mathbf{y} \in \mathbb{Z}^m$

$$(69) \quad \|U \cdot \mathbf{z} - \mathbf{y}\|_{\infty} \gg \|\mathbf{z}\|_{\infty}^{-1/(m-2)} \gg \|\mathbf{z}\|_1'^{-1/(m-2)} \geq t^{-1/(m-2)}$$

where the most left estimate holds since the system is decoupled and $\zeta \in \text{Bad}_{m-2,1}$. As we noticed above that $m \geq 4$ and $1/2 \leq n/m$, in our case $(m, n) \neq (4, 2)$ we have the strict inequality $1/(m-2) < n/m$. Then, in view of (63), approximation quality (69) is beaten by $\psi_{\xi}^{2 \times 1}(\|\cdot\|, \|\cdot\|_{\infty}, \gamma^{-1}t)$, so extending accordingly the involved integer best approximation approximations to \mathbb{Z}^n by adding 0 entries leads to the same smaller value. So we can exclude such vectors. But for vectors \mathbf{z} in the linear space $z_2 = 0$, the integer vector \mathbf{y} minimizing $\|U \cdot \mathbf{z} - \mathbf{y}\|_{\infty}$ for given \mathbf{z} is easily seen to be of the form $\mathbf{y} = (\mathbf{w}, 0, \dots, 0)$ with $\mathbf{w} \in \mathbb{Z}^2$, as otherwise some entry of $U \cdot \mathbf{z} - \mathbf{y}$ is of modulus at least 1. Then the output $U \cdot \mathbf{z} - \mathbf{y} \in \mathbb{R}^m$ again satisfies (66) and hence (67), settling (68) via a short calculation. For $m = 4, n = 2$ a very similar argument applies using $\zeta = \xi$, however we need $\|\cdot\|_1$ to be expanding to exclude better approximations outside the space $z_2 = 0$. If the expanding property is assumed, then it is again easy to see that we may restrict \mathbf{z} to this space and the claims follows easily. We leave the details to the reader.

Finally, let $\|\cdot\|_2$ be any expanding norm on \mathbb{R}^m and consider again the induced projected norm $\|\cdot\|'_2$ on \mathbb{R}^2 by

$$\|(x_1, x_2)\|'_2 := \|(x_1, x_2, 0, \dots, 0)\|_2.$$

It is easy to see that $\|\cdot\|'_2$ is an expanding norm on \mathbb{R}^2 . A minor adaption of [27, Corollary 2] to more general Φ satisfying (d1)-(d3) of [27] can be proved analogously. When applied with respect to simultaneous approximation of two reals, and as (d1)-(d3) hold for our $\Phi(t) = ct^{-n/m}$ in view of (60) and $c \leq 1$, it yields the following: For any small enough $c > 0$, we can similarly arrange some $\xi \in \mathbb{R}^{2 \times 1}$ to satisfy

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_{\xi}^{2 \times 1}(\|\cdot\|, \|\cdot\|'_2, t) = c_1 := c\gamma^{n/m},$$

and

$$\liminf_{t \rightarrow \infty} t^{n/m} \psi_{\xi}^{2 \times 1}(\|\cdot\|, \|\cdot\|'_2, t) = 0.$$

We can then copy the proof above replacing $\|\cdot\|_{\infty}$ by $\|\cdot\|'_2$ whenever considering $\psi_{\xi}^{2 \times 1}$. Again by the same argument, it suffices to show the accordingly altered variant of (64). For the estimate (65), the only place where we needed $\|\cdot\|_2$ to be the maximum norm is to deduce that the projected vectors satisfy (67). However, in view of (66), the defining property of $\|\cdot\|'_2$ means that the according equality still holds. The proof of the reverse inequality (68) is analogous to the above, only with different implied constants in (69). The case $\ell = 2$ is done.

Case $\ell \geq 3$: Then by (59) similarly to (60) we have

$$(70) \quad \frac{1}{\ell} \leq \frac{n}{m} < \frac{1}{\ell - 1}.$$

Now again [27, Theorem 2.2] can be applied to $\Phi(t) = ct^{-n/m}$ for any $c \in (0, 1)$, as its conditions (d1)-(d3) are satisfied by (70). This yields $\xi \in \mathbb{R}^{\ell \times 1}$ so that

$$\limsup_{t \rightarrow \infty} t^{n/m} \psi_{\xi}^{\ell \times 1}(t) = c, \quad \text{and} \quad \liminf_{t \rightarrow \infty} t^{n/m} \psi_{\xi}^{\ell \times 1}(t) = 0.$$

Now let $\zeta \in \text{Bad}_{m-\ell, 1}$ arbitrary unless if $m = 2\ell$ then $\zeta = \xi$. We again consider $U = \text{diag}(\xi, \zeta) \in \mathbb{R}^{m \times 2}$ as above. As soon as strict inequality

$$(71) \quad m > 2\ell$$

holds, we have $1/(m - \ell) < n/m$ so that the analogue of (69) similarly implies that we can restrict to the space $z_2 = 0$. It can be checked that the analogous arguments of the case $\ell = 2$ above apply. Notably when transitioning to any expanding norm $\|\cdot\|_2$, here we apply a variant of [27, Corollary 2] with respect to simultaneous approximation to ℓ numbers.

Finally we need to discuss the separate cases when (71) is satisfied, and evaluate the Hausdorff dimensions. Recall we have assumed $\ell \geq 3$, moreover we can assume $n \geq 2$ as $n = 1$ was done in [26]. Via (59), condition (71) is readily checked to be satisfied if $n \geq 3$. Thus we are left with the subset of cases $n = 2$. The cases $n = 2, m \geq 4$ even, leading to $m = 2\ell$, can be dealt with similarly to the case $(m, n) = (4, 2)$ for $\ell = 2$ above. This indeed leaves only the cases $n = 2$, and m odd, as excluded in the theorem.

By construction, and the repeatedly used observation that $\text{Bad}_{m-\ell, 1}$ has full Hausdorff dimension $m - \ell$, we conclude that unless $m = 2\ell$, we get Hausdorff dimension bound

$$\dim_H FS_{m,n}(c) \geq m \cdot (n - 2) + (m - \ell) = m(n - 1) - \left\lfloor \frac{m}{n} \right\rfloor, \quad m \geq 2\ell + 1,$$

of the theorem. The remaining case $m = 2\ell = 2\lceil m/n \rceil$ is readily checked to lead precisely to the cases $n = 2$ and m even or $(m, n) = (4, 3)$. When $(m, n) = (4, 3)$ we have $\ell = 2$, which is already treated above. It should be remarked that for $n = 2$ our Hausdorff dimension bound is just $m(n - 2) = 0$, thus of no interest. \square

9. PROOF OF LEMMA 1.1

For matrix $\Omega \in \mathbb{R}^{m \times n}$ and real numbers $A, B \in \mathbb{R}_+$ define the convex bodies

$$M_{A,B}(\Omega) := \left\{ \mathbf{z} \in \mathbb{R}^{n+m} : \max_{1 \leq j \leq m} \left| \sum_{i=1}^n \Omega_{j,i} z_i + z_{n+j} \right| \leq A, \quad \max_{1 \leq i \leq n} |z_i| \leq B \right\},$$

$$\widehat{M}_{A,B}(\Omega) := \left\{ \mathbf{z} \in \mathbb{R}^{n+m} : \max_{1 \leq i \leq n} \left| z_i - \sum_{j=1}^m \Omega_{i,j} z_{n+j} \right| \leq B, \quad \max_{1 \leq j \leq m} |z_{n+j}| \leq A \right\}.$$

The following theorem proven by German is crucial.

Theorem 9.1. [16, Theorem 7] *Let $\Omega \in \mathbb{R}^{m \times n}$ and $A, B \in \mathbb{R}_+$. If*

$$M_{A,B}(\Omega) \cap \mathbb{Z}^{n+m} \setminus \{\mathbf{0}\} \neq \emptyset,$$

then

$$\widehat{M}_{A^*, B^*}(\Omega) \cap \mathbb{Z}^{n+m} \setminus \{\mathbf{0}\} \neq \emptyset$$

for

$$A^* = C(A^n B^{1-n})^{\frac{1}{n+m-1}} \quad \text{and} \quad B^* = C(A^{1-m} B^m)^{\frac{1}{n+m-1}},$$

with $C > 0$ constant dependent only on n and m .

Equipped with Theorem 9.1 we can prove Lemma 1.1.

Proof. Since $\Omega \in FS_{m,n}(\|\cdot\|_1, \|\cdot\|_2, c) \subseteq \mathbb{R}^{m \times n}$ then we have that

$$\Theta^{m \times n}(\|\cdot\|_1, \|\cdot\|_2, \Omega) = c.$$

By the equivalence of norms we have that the same is true up to some constant for maximum norms, so

$$\Theta^{m \times n}(\|\cdot\|_\infty, \|\cdot\|_\infty, \Omega) \asymp c.$$

Hence, for all sufficiently large $t \in \mathbb{R}$ and some $\rho > 0$ there is an integer

$$\mathbf{b} \in M_{\rho t^{-n/m}, t}(\Omega) \cap \mathbb{Z}^{n+m} \setminus \{\mathbf{0}\}.$$

Hence, by Theorem 9.1, for all sufficiently large $u \in \mathbb{R}$ there exists integer

$$\mathbf{b} \in \widehat{M}_{u, \delta u^{-m/n}}(\Omega) \cap \mathbb{Z}^{n+m} \setminus \{\mathbf{0}\},$$

for

$$\delta = C^{\frac{m+n}{n}} \rho^{\frac{m}{n(n+m-1)}}.$$

Hence $\Omega^T \in Di_{n,m}(\delta)$, and by equivalence of norms there exists some $\delta^* > 0$ such that $\Omega^T \in Di_{n,m}(\|\cdot\|_2, \|\cdot\|_1, \delta^*)$. If $D^* = D^{n \times m}(\|\cdot\|_1, \|\cdot\|_2)$ is the Dirichlet constant from (5) for the transpose problem, for small enough δ and thus $\delta^* < D^*$ this will induce a Dirichlet improvable matrix Ω^T . Conversely, for any $0 < \gamma < c$ there exists an unbounded sequence of $t \in \mathbb{R}$ for which

$$\widehat{M}_{t, \gamma t^{-n/m}}(\Omega^T) \cap \mathbb{Z}^{n+m} \setminus \{\mathbf{0}\} = \emptyset,$$

otherwise $\Omega \in Di_{m,n}(\gamma)$, which is clearly false since $\Omega \in DI_{m,n}(c)$. Applying the contrapositive of Theorem 9.1 we obtain that there exist certain arbitrarily large $u \in \mathbb{R}$ and a constant $C^* > 0$ for which

$$M_{C^* u^{-n/m}, u}(\Omega^T) \cap \mathbb{Z}^{n+m} \setminus \{\mathbf{0}\} = \emptyset.$$

Hence $\Omega^T \notin Sing_{n,m}$. Furthermore, one can similarly show that $\Omega^T \notin Bad_{n,m}$, and so

$$\Omega^T \in Di_{n,m}(\|\cdot\|_2, \|\cdot\|_1, \delta^*) \setminus (Sing_{n,m} \cup Bad_{n,m}).$$

As previously mentioned $Sing_{n,m}$ and $Bad_{n,m}$ are independent of norms, and so $\Omega^T \in FS_{n,m}(\|\cdot\|_2, \|\cdot\|_1)$ as required. \square

Acknowledgements. The research of Mumtaz Hussain and Benjamin Ward is supported by the Australian Research Council Discovery Project (200100994). Some of this work was carried out when Johannes Schleisitz visited La Trobe University supported by an International Visitor Program of the Sydney Mathematical Research Institute (SMRI) and La Trobe University, we thank them for their generous support. Finally, we thank Tushar Das and David Simmons for pointing out some relevant results from their paper [12].

REFERENCES

- [1] A. Agin. Constructing displacement vectors. *arXiv preprint arXiv:2308.03049*, 2023.
- [2] R. K. Akhunzhanov and N. G. Moshchevitin. A note on Dirichlet spectrum. *Mathematika*, 68(3):896–920, 2022.
- [3] R. K. Akhunzhanov and D. O. Shatskov. On Dirichlet spectrum for two-dimensional simultaneous Diophantine approximation. *Mosc. J. Comb. Number Theory*, 3(3-4):5–23, 2013.
- [4] N. Andersen and W. Duke. On a theorem of Davenport and Schmidt. *Acta Arith.*, 198(1):37–75, 2021.
- [5] V. Beresnevich, S. Datta, A. Ghosh, and B. Ward. Bad is null. <https://arxiv.org/abs/2307.10109>, preprint 2023.
- [6] V. Beresnevich, L. Guan, A. Marnat, F. Ramírez, and S. Velani. Dirichlet is not just bad and singular. *Adv. Math.*, 401:Paper No. 108316, 57, 2022.
- [7] J. W. S. Cassels. *An introduction to Diophantine approximation*. Cambridge Tracts in Mathematics and Mathematical Physics, No. 45. Cambridge University Press, New York, 1957.
- [8] Y. Cheung. Hausdorff dimension of the set of singular pairs. *Ann. of Math. (2)*, 173:no. 1, 127–167, 2011.
- [9] Y. Cheung and N. Chevallier. Hausdorff dimension of singular vectors. *Duke Math. J.*, 165(12):2273–2329, 2016.
- [10] N. Chevallier. Best simultaneous Diophantine approximations and multidimensional continued fraction expansions. *Mosc. J. Comb. Number Theory*, 3(1):3–56, 2013.
- [11] T. Das, L. Fishman, D. Simmons, and M. Urbański. A variational principle in the parametric geometry of numbers, with applications to metric Diophantine approximation. *C. R. Math. Acad. Sci. Paris*, 355(8):835–846, 2017.
- [12] T. Das, L. Fishman, D. Simmons, and M. Urbański. A variational principle in the parametric geometry of numbers. *Adv. Math.*, 437:Paper No. 109435, 130, 2024.
- [13] H. Davenport and W. M. Schmidt. Dirichlet’s theorem on diophantine approximation. II. *Acta Arith.*, 16:413–424, 1969/70.
- [14] H. Davenport and W. M. Schmidt. Dirichlet’s theorem on diophantine approximation. In *Symposia Mathematica, Vol. IV (INDAM, Rome, 1968/69)*, pages 113–132. Academic Press, London, 1970.
- [15] K. Falconer. *Fractal geometry*. John Wiley & Sons, Ltd., Chichester, third edition, 2014. Mathematical foundations and applications.
- [16] O. N. German. On Diophantine exponents and Khintchine’s transference principle. *Mosc. J. Comb. Number Theory*, 2(2):22–51, 2012.
- [17] V. Jarník. Diophantische Approximationen und Hausdorffsches Mass. *Mat. Sb.*, 36:371–382, 1929.
- [18] A. Khinchin. Über eine Klasse linearer diophantischer Approximationen. *Rend. Circ. Mat. Palermo*, 50:170–195, 1926.
- [19] D. Kleinbock and A. Rao. Abundance of Dirichlet-improvable pairs with respect to arbitrary norms. *Mosc. J. Comb. Number Theory*, 11(1):97–114, 2022.
- [20] D. Kleinbock and A. Rao. Weighted uniform Diophantine approximation of systems of linear forms. *Pure Appl. Math. Q.*, 18(3):1095–1112, 2022.
- [21] J. C. Lagarias. Best simultaneous Diophantine approximations. I. Growth rates of best approximation denominators. *Trans. Amer. Math. Soc.*, 272(2):545–554, 1982.
- [22] J. C. Lagarias. Best simultaneous Diophantine approximations. II. Behavior of consecutive best approximations. *Pacific J. Math.*, 102(1):61–88, 1982.
- [23] A. Marnat and N. G. Moshchevitin. An optimal bound for the ratio between ordinary and uniform exponents of Diophantine approximation. *Mathematika*, 66(3):818–854, 2020.
- [24] N. G. Moshchevitin. Best diophantine approximations: the phenomenon of degenerate dimension. *London Math. Soc. Lecture Note Ser.*, 338:158–182, 2007.
- [25] N. G. Moshchevitin. Singular Diophantine systems of A. Ya. Khinchin and their application. *Uspekhi Mat. Nauk*, 65(3(393)):43–126, 2010.

- [26] J. Schleischitz. Dirichlet spectrum for one linear form. *Bull. Lond. Math. Soc.*, 55(3):1330–1339, 2023.
- [27] J. Schleischitz. Exact uniform approximation and Dirichlet spectrum in dimension at least two. *Selecta Math. (N.S.)*, 29(5):Paper No. 86, 2023.
- [28] J. Schleischitz. Metrical results on the geometry of best approximations for a linear form. *arXiv preprint arXiv:2302.08403*, 2023.
- [29] W. Schmidt. A metrical theorem in diophantine approximation. *Canadian Journal of Mathematics*, 12:619–631.
- [30] W. M. Schmidt. Badly approximable systems of linear forms. *J. Number Theory*, 1:139–154, 1969.
- [31] F. Süess. Simultaneous diophantine approximation on affine subspaces and dirichlet improvability. *PhD Thesis, University of York*, 2017.
- [32] C. Tricot, Jr. Two definitions of fractional dimension. *Math. Proc. Cambridge Philos. Soc.*, 91(1):57–74, 1982.

MUMTAZ HUSSAIN, DEPARTMENT OF MATHEMATICAL AND PHYSICAL SCIENCES, LA TROBE UNIVERSITY, BENDIGO 3552, AUSTRALIA.

Email address: `m.hussain@latrobe.edu.au`

JOHANNES SCHLEISCHITZ, MIDDLE EAST TECHNICAL UNIVERSITY, NORTHERN CYPRUS CAMPUS, KALKANLI, GÜZELYURT

Email address: `johannes@metu.edu.tr`; `jschleischitz@outlook.com`

BENJAMIN WARD, DEPARTMENT OF MATHEMATICAL AND PHYSICAL SCIENCES, LA TROBE UNIVERSITY, BENDIGO 3552, AUSTRALIA.

Email address: `Ben.Ward@latrobe.edu.au`