

Spherically symmetric configurations in the quadratic $f(R)$ gravity

V. I. Zhdanov,^{1,2,*} O. S. Stashko,^{3,4,†} and Yu. V. Shtanov^{5,1,‡}

¹*Taras Shevchenko National University of Kyiv, Kyiv 01601, Ukraine*

²*Igor Sikorsky Kyiv Polytechnic Institute, Kyiv 03056, Ukraine*

³*Department of Physics, Princeton University, Princeton, USA*

⁴*Goethe Universität, Max-von-Laue Str. 1, Frankfurt am Main, 60438, Germany*

⁵*Bogolyubov Institute for Theoretical Physics, Metrologichna St. 14-b, Kyiv 03143, Ukraine*

We study spherically symmetric configurations of the quadratic $f(R)$ gravity in the Einstein frame. In case of a purely gravitational system, we have determined the global qualitative behavior of the metric and the scalaron field for all static solutions satisfying the conditions of asymptotic flatness. These solutions are proved to be regular everywhere except for a naked singularity at the center; they are uniquely determined by the total mass \mathfrak{M} and the “scalar charge” Q characterizing the strength of the scalaron field at spatial infinity. The case $Q = 0$ yields the Schwarzschild solution, but an arbitrarily small $Q \neq 0$ leads to the appearance of a central naked singularity having a significant effect on the neighboring region, even when the space-time metric in the outer region is practically insensitive to the scalaron field. Approximation procedures are developed to derive asymptotic relations near the naked singularity and at spatial infinity, and the leading terms of the solutions are presented. We investigate the linear stability of the static solutions with respect to radial perturbations satisfying the null Dirichlet boundary condition at the center and numerically estimate the range of parameters corresponding to stable/unstable configurations. In particular, the configurations with sufficiently small Q turn out to be linearly unstable.

Keywords: modified gravity, $f(R)$ gravity, scalar fields, naked singularities

I. INTRODUCTION

Modification of General Relativity (GR) by a Lagrangian in the form of a nonlinear function $f(R)$ of the scalar curvature R is, perhaps, the simplest one and has long been the subject of numerous studies and applications (see [1–3] for reviews). Compared to GR, such $f(R)$ gravity theory contains one extra scalar degree of freedom, which can be used for modelling a wide variety of phenomena, from the early inflationary regime [4–7], consistent with current observations [8], to dark energy and dark matter at later epoch [9–19].

Natural questions arise about the possible effects of modified gravity in models of isolated systems that can simulate real astrophysical objects. Isolated configurations have received considerable attention in both GR and modified theories of gravity due to various aspects ranging from the black hole no-hair theorems to naked singularities, stability properties, etc. (see, e.g., [20–34] and references therein).

In the present paper, we investigate asymptotically flat spherically symmetric (SS) systems and their stability in the framework of the quadratic $f(R)$ gravity. The key point is that we use the transition to a convenient Einstein frame, that enables us to use previous results on

global properties of the SS systems with scalar field, especially of [35]. In particular, it was shown [35] that for fairly general scalar field potentials there exist a naked singularity (NS) at the center of the SS configuration. Systems with NS, if they really exist in nature, could be extremely interesting. Indeed, in the modern cosmological epoch, the effects of the modified gravity are expected to be very weak in astrophysical objects, but they can manifest themselves in the extreme conditions near classical NS. On the other hand, there is a widespread opinion that systems with NS do not exist in the universe according to the Penrose’s Cosmic Censorship hypothesis [36, 37]. This hypothesis has never been proven, and its discussion shifts to issues of stability and fine-tuning of the input data [38–42]. Apparently, the existence and stability of NS depends on the types of configurations involved and, perhaps, on the specific parameter regions in concrete models [43–46].

Static SS solutions in $f(R)$ gravity were studied earlier in [25], where the existence of NS at the center was pointed out. However, the consideration of [25] is essentially based on numerical computations, and some of its assumptions may be questioned or at least require rigorous justification. First of all, one must be sure that regular solutions with all possible configuration parameters satisfying the assumption of asymptotic flatness do exist for all positive values of the radial variable r . This is a non-trivial question in case of a nonlinear system; indeed, there are examples of static SS configurations where “spherical” singularities arise at finite values of $r > 0$ [35, 47]. Another important issue, that requires proper justification, concerns the asymptotic properties of solutions necessary for the linear stability problems considered in this work.

Our paper is organized as follows. In Section II, we

* valery.zhdanov@knu.ua

† alexander.stashko@gmail.com

‡ shtanov@bitp.kyiv.ua

review the general relations of $f(R)$ gravity as regards the transition to the Einstein frame. Section III concentrates on the quadratic $f(R)$ model. Field equations in the Einstein frame are written in Section IV for the case of a non-stationary SS metric. Section V deals with the purely gravitational system; we prove that *all* SS solutions with nonzero scalar field satisfying the asymptotic-flatness conditions, exist for all $r > 0$ and may have singularity only at $r = 0$. Also in this Section, asymptotic formulas near NS are obtained which are further used in numerical calculations. In Section VI we write down equations for perturbations. In Section VII we use numerical calculations to illustrate the static solutions and to study their linear stability. In Section VIII, we discuss our results. The details concerning the proof of the existence and uniqueness and justification of approximation methods are given in appendices.

II. BASIC EQUATIONS AND NOTATION

In $f(R)$ gravity, the standard Lagrangian of General Relativity in the gravitational action is replaced by a more general function¹ of the scalar curvature R :

$$S_g = -\frac{1}{2\kappa} \int d^4x \sqrt{-g} f(R),$$

where $\kappa = 8\pi G$, and G is Newton's gravitational constant. The corresponding dynamical equations for the physical metric $g_{\mu\nu}$ (Jordan frame) are [1–3]

$$f'(R)R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}f(R) + (\square - \nabla_\mu \nabla_\nu) f'(R) = \kappa T_{\mu\nu}, \quad (1)$$

where $T_{\mu\nu} = T_{\nu\mu}$ is the energy-momentum tensor of non-gravitational fields satisfying the covariant conservation law

$$\nabla_\nu T^{\mu\nu} = 0. \quad (2)$$

There is a well-known procedure [1, 2] to represent the equations of the $f(R)$ gravity, written for the physical metric $g_{\mu\nu}$ (Jordan frame), in the form of the usual Einstein equations for a conformally transformed metric $\hat{g}_{\mu\nu}$ (Einstein frame)

$$g_{\mu\nu} = e^{-2\xi} \hat{g}_{\mu\nu}, \quad e^{2\xi} = f'(R), \quad (3)$$

accompanied by equations for an additional canonically normalized scalar field (SF) $\phi = \sqrt{6/\kappa} \xi$. In this paper, we will describe this scalar field (the scalaron) by the dimensionless scalar ξ .

The self-interaction potential $W(\xi)$ of the scalar field ξ can be introduced parametrically as follows:

$$e^{2\xi} = f'(u), \quad W(\xi) = \frac{1}{2} e^{-4\xi} [f(u) - f'(u)u]. \quad (4)$$

The extrema of this potential are located at the values of ξ corresponding to the values of u satisfying the equation $uf'(u) = 2f(u)$. The quadratic part of the potential around its absolute minimum determines the scalaron mass μ :

$$\mu^2 = \frac{1}{3} \left[\frac{1}{f''(u)} - \frac{u}{f'(u)} \right],$$

where the right-hand side is evaluated at the minimum of the potential. Non-observation [48, 49] of the scalaron-induced Yukawa forces [50] between non-relativistic masses leads to a lower bound on the scalaron mass (see also [11, 12, 51]):

$$\mu \geq 2.7 \text{ meV} \quad \text{at 95\% C.L.} \quad (5)$$

This lower bound leads to a rather small length scale $l_\mu = 1/\mu$.

Denote

$$\hat{T}_{\mu\nu} = e^{-2\xi} T_{\mu\nu}. \quad (6)$$

To avoid confusion due to the presence of two metrics, we emphasize that, from this point on, ‘‘hats’’ correspond to the Einstein frame [1, 2], in which covariant differentiation, raising and lowering indexes are performed with the metric tensor $\hat{g}_{\mu\nu}$. On the account of (3) and (4), equation (1) leads to the following equations in the Einstein frame:

$$\hat{R}_{\mu\nu} - \frac{1}{2} \hat{g}_{\mu\nu} \hat{R} = \hat{T}_{\mu\nu}^{(\xi)} + \kappa \hat{T}_{\mu\nu}, \quad (7)$$

where

$$\hat{T}_{\mu\nu}^{(\xi)} = 6\partial_\mu \xi \partial_\nu \xi - \hat{g}_{\mu\nu} [3\xi_{,\alpha} \xi_{,\beta} \hat{g}^{\alpha\beta} - W(\xi)]. \quad (8)$$

Equations (7) are supplemented by the equation for the scalaron ξ :

$$\hat{\nabla}_\alpha \hat{\nabla}^\alpha \xi = -\frac{1}{6} \frac{\partial W}{\partial \xi} + \frac{1}{6} \kappa \hat{T}, \quad \hat{T} = \hat{g}^{\mu\nu} \hat{T}_{\mu\nu}. \quad (9)$$

The system of equations (7), (9) is equivalent to the fourth-order equations (1).

The covariant conservation law (2) also must be rewritten in the Einstein frame. For the energy-momentum tensor, we have

$$\hat{\nabla}_\mu \hat{T}^\mu{}_\nu = -\hat{T} \partial_\nu \xi, \quad \hat{T} = \hat{g}_{\alpha\beta} \hat{T}^{\alpha\beta}. \quad (10)$$

¹ We use the units in which $c = \hbar = 1$, the metric signature is $(+, -, -, -)$, and the curvature conventions are $R^\alpha{}_\beta\gamma\delta = \partial_\gamma \Gamma^\alpha{}_{\beta\delta} - \dots$, $R_{\mu\nu} = R^\alpha{}_\mu\alpha\nu$.

III. QUADRATIC $f(R)$ GRAVITY

In what follows, we deal with the simplest model, in which²

$$f(R) = R - \frac{R^2}{6\mu^2}, \quad (11)$$

where μ is the scalaron mass. In the original inflationary model due to Starobinsky [4], described by this Lagrangian, the scalaron mass has relatively large value $\mu \approx 3 \cdot 10^{13}$ GeV [8].

The scalaron potential (4) for model (11) is

$$W(\xi) = \frac{3}{4}\mu^2 (1 - e^{-2\xi})^2. \quad (12)$$

It is monotonically increasing with ξ in the domain $\xi \geq 0$ asymptotically rising to a plateau. In the case of an asymptotically flat space-time in the initial (Jordan) frame, we assume that the scalar curvature vanishes at spatial infinity; correspondingly, $\xi = \frac{1}{2} \ln[f'(R)] \rightarrow 0$.

For most of astrophysical systems with continuous distributions of matter, we expect ξ to be very small. For $|\xi| \ll 1$, we have

$$W(\xi) \approx 3\mu^2\xi^2. \quad (13)$$

In a static case, we deal with metric $g_{\mu\nu} \equiv g_{\mu\nu}(\mathbf{r})$; $g_{0i} \equiv 0$, yielding

$$\hat{\Delta}\xi \equiv -\frac{1}{\sqrt{-\hat{g}}}\frac{\partial}{\partial x^i}\left[\sqrt{-\hat{g}}\hat{g}^{ij}\frac{\partial\xi}{\partial x^j}\right] = \mu^2\xi - \frac{\kappa}{6}\hat{T}, \quad (14)$$

where $i, j = 1, 2, 3$.

As noted above, the value of the scalaron mass μ is bounded from below by (5) so as not to contradict the existing observations. A somewhat stronger lower bound, $\mu \gtrsim 4.4$ meV, arises in the consideration of the scalaron as a dark-matter candidate [18, 19]. Such a mass corresponds to the length scale $l_\mu = \mu^{-1} \lesssim 4.5 \cdot 10^{-3}$ cm, which is a very small value in the astrophysical realm. In the case of an astrophysical object with mass \mathfrak{M} and gravitational radius $r_g = 2G\mathfrak{M}$, we typically deal with a very large dimensionless quantity $\mu r_g \gg 1$. This allows us to make a general estimate for the value of ξ inside a sufficiently smooth distribution of \hat{T} . Indeed, equation (14) can be written in a form that may be considered as a source of asymptotic approximations:

$$\xi = \frac{\kappa}{6\mu^2}\hat{T} - \frac{1}{\mu^2}\hat{\Delta}\xi, \quad (15)$$

with the last term on the right-hand side regarded as a small perturbation. This gives us a simple approximate formula

$$\xi \approx \frac{\kappa}{6\mu^2}\left(\hat{T} - \frac{1}{\mu^2}\hat{\Delta}\hat{T}\right). \quad (16)$$

This formula is applicable if $\mu^2\hat{T} \gg \hat{\Delta}\hat{T}$, which, however, may fail to be satisfied at the boundary of the body, where (16) must be replaced by a more exact relation.

Further in this paper we focus primarily on mathematical properties of the model and allow for arbitrary values of μ where possible.

IV. SPHERICALLY SYMMETRIC CONFIGURATIONS

A. Field equations in the Einstein frame

The metric of a spherically symmetric space-time in the Schwarzschild (curvature) coordinates can be written as

$$ds^2 = e^\alpha dt^2 - e^\beta dr^2 - r^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (17)$$

where $r > 0$, $\alpha \equiv \alpha(t, r)$, $\beta \equiv \beta(t, r)$. The nontrivial Einstein equations (7) in this case are

$$\begin{aligned} \frac{\partial}{\partial r}[r(e^{-\beta} - 1)] &= -\kappa r^2 \hat{T}_0^0 \\ -r^2 \left[3e^{-\alpha} \left(\frac{\partial\xi}{\partial t}\right)^2 + 3e^{-\beta} \left(\frac{\partial\xi}{\partial r}\right)^2 + W(\xi) \right] &, \end{aligned} \quad (18)$$

$$\begin{aligned} r e^{-\beta} \frac{\partial\alpha}{\partial r} + e^{-\beta} - 1 &= -\kappa r^2 \hat{T}_1^1 \\ + r^2 \left[3e^{-\alpha} \left(\frac{\partial\xi}{\partial t}\right)^2 + 3e^{-\beta} \left(\frac{\partial\xi}{\partial r}\right)^2 - W(\xi) \right] &, \end{aligned} \quad (19)$$

$$\frac{\partial\beta}{\partial t} = -\kappa r e^\beta \hat{T}_0^1 + 6r \frac{\partial\xi}{\partial t} \frac{\partial\xi}{\partial r}, \quad (20)$$

where $\xi \equiv \xi(t, r)$, and the structure of $\hat{T}_{\mu\nu}$ corresponds to the spherically symmetric case.

Equation (9) yields

$$\begin{aligned} e^{-\frac{\alpha+\beta}{2}} \left[\frac{\partial}{\partial t} \left(e^{\frac{\beta-\alpha}{2}} \frac{\partial\xi}{\partial t} \right) - \frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 e^{\frac{\alpha-\beta}{2}} \frac{\partial\xi}{\partial r} \right) \right] \\ = -\frac{1}{6} \left[W'(\xi) - \kappa \hat{T} \right]. \end{aligned} \quad (21)$$

Note that, for $\xi \equiv 0$ and $\hat{T} = 0$, we get the Schwarzschild solution.

B. Static SS solutions

In the case of a static isolated regular configuration, the quantities α , β and ξ depend only on r . We assume that, for $r < R_0$, we have a continuous matter distribution with non-zero regular $\hat{T}_{\mu\nu}$, whereas $\hat{T}_{\mu\nu} = 0$ in an outer region $r > R_0$.

² We neglect the presence of a small cosmological constant.

In what follows, we consider purely gravitational case with $\hat{T} = 0$. For a static case, Eqs. (18), (19) yield

$$\frac{d}{dr}(\alpha + \beta) = 6r \left(\frac{d\xi}{dr} \right)^2, \quad (22)$$

$$\frac{d}{dr}(\alpha - \beta) = -\frac{2}{r} + \frac{2e^\beta}{r} [1 - r^2 W(\xi)]. \quad (23)$$

The system of equations with respect to α , β , and ξ is closed by adding the equation for the SF:

$$\frac{d}{dr} \left[r^2 e^{\frac{\alpha-\beta}{2}} \frac{d\xi}{dr} \right] = \frac{r^2}{6} e^{\frac{\alpha+\beta}{2}} W'(\xi). \quad (24)$$

In the case of an asymptotically flat static space-time, we assume

$$\lim_{r \rightarrow \infty} [r\alpha(r)] = -r_g, \quad \lim_{r \rightarrow \infty} [r\beta(r)] = r_g, \quad (25)$$

where $r_g = 2G\mathfrak{M}$, and $\mathfrak{M} > 0$ is the configuration mass.

In the asymptotically flat configuration, we assume $\xi(r) \rightarrow 0$ as $r \rightarrow \infty$, and Eq. (24) can be approximated by the equation for a free massive scalar field on the Schwarzschild background [52–54]:

$$\frac{d}{dr} \left[r(r - r_g) \frac{d\xi}{dr} \right] = r^2 \mu^2 \xi. \quad (26)$$

We discard unbounded solutions of (26) as $r \rightarrow \infty$ leaving only those with exponentially decaying $\xi(r)$. Taking into account the results of [52–55] on the asymptotic behavior at infinity, we assume

$$\lim_{r \rightarrow \infty} \left[\left(\frac{r}{r_g} \right)^{1+M\mu} e^{\mu r} \xi(r) \right] = Q. \quad (27)$$

Here and below, $M = r_g/2$, and the constant Q measures the strength of the scalaron field at spatial infinity and will be dubbed as the “scalar charge.”

For given $\mu > 0$, $\mathfrak{M} > 0$ and Q , we claim that *there is $r_0 > R_0$ such that solution $\alpha(r) \in C^1$, $\beta(r) \in C^1$, $\xi(r) \in C^2$ of Eqs. (22)–(24) exists for $r \geq r_0$ and is uniquely defined by conditions (25), (27) regardless of the interior solution and structure of the energy-momentum tensor.* This statement is physically quite understandable; however, its rigorous proof requires some analytical work, which we present in Appendix A.

Our next step is to show that, if $\hat{T}_{\mu\nu} \equiv 0$ for all $r > 0$, then one can put $r_0 = 0$; in this case, the parameters M and Q completely define the static SS configuration. This is the subject of the next Section V.

The numerical investigations, which we perform below, require more detailed information on the asymptotic properties at large r . This is also considered in Appendix A, where we present a convergent approximation method to justify asymptotics of decaying static solutions $\xi(r)$ for $\mu r \gg 1$, $r \gg r_g$. The leading terms of this asymptotics for $|\xi| \ll 1$ are

$$\xi(r) = Q \left(\frac{r_g}{r} \right)^{1+M\mu} \left(1 + \frac{b_1}{r} + \frac{b_2}{r^2} \right) e^{-\mu r}, \quad (28)$$

where

$$b_1 = \frac{M}{2}(1 + 3M\mu), \quad (29)$$

$$b_2 = \frac{M}{8\mu}(2 + 5M\mu + 16M^2\mu^2 + 9M^3\mu^3). \quad (30)$$

Note that (28) differs by a power-law factor from the usual Yukawa asymptotics in flat space. In view of the consideration in Appendix A, an asymptotic relation for $d\xi/dr$ can be obtained by formal differentiation of (28).

V. GLOBAL BEHAVIOR OF STATIC SOLUTIONS AND ASYMPTOTIC PROPERTIES NEAR NS

Now we consider a purely gravitational SS system, that is $\hat{T}_{\mu\nu} \equiv 0$. In this Section we essentially use the method of [35].

According to the previous results, there exists an asymptotically flat solution of Eqs. (22)–(24) on $[r_0, \infty)$ for some $0 < r_0 < \infty$. The set of all such r_0 has infimum $r_0^* = \inf r_0 < \infty$, and the solution exists for $r \in (r_0^*, \infty)$. We prove that $r_0^* = 0$.

Suppose (on the contrary) that $r_0^* > 0$ and consider the solution on (r_0^*, ∞) .

The SF potential (12) has the property

$$\xi W'(\xi) > 0, \quad \xi \neq 0. \quad (31)$$

Using this equation, it is easy to see that

$$\frac{d\Gamma}{dr} = r^2 e^{\frac{\alpha-\beta}{2}} \xi'^2 + \frac{r^2}{6} e^{\frac{\alpha+\beta}{2}} \xi W'(\xi), \quad (32)$$

where $\Gamma(r) \equiv r^2 e^{\frac{\alpha-\beta}{2}} \xi \xi'$, $\xi' \equiv d\xi/dr$.

The right-hand side of (32) is positive for any nontrivial $\xi(r)$, and the function $\Gamma(r)$ is strictly increasing. For the solutions satisfying (27), we have $\Gamma(r) \rightarrow 0$ as $r \rightarrow \infty$. In a nontrivial case, this is possible if and only if $\Gamma(r) < 0$, so that $\forall r > r_0^*$ the functions $\xi(r)$ and $\xi'(r)$ are nonzero and have opposite signs.

Now we will prove that, in the case of $\hat{T}_{\mu\nu} \equiv 0$, solutions of (22)–(24) with asymptotic conditions (25), (27) can be regularly extended to the left of r_0^* .

Similar to [35], we introduce variables $x = (\alpha + \beta)/2$ and

$$X = e^x, \quad Y = \left(\frac{r}{r_g} \right) e^{\frac{\alpha-\beta}{2}}, \quad Z = -rY \frac{d\xi}{dr}. \quad (33)$$

Then, using (23) and (22), we obtain an equivalent first-order system in terms of the dimensionless variables X ,

Y , Z , and ξ :

$$\frac{dX}{dr} = \frac{3XZ^2}{rY^2}, \quad (34a)$$

$$\frac{dY}{dr} = \frac{X}{r_g} X [1 - r^2 W(\xi)], \quad (34b)$$

$$\frac{dZ}{dr} = -\frac{r^2}{6r_g} X W'(\xi), \quad (34c)$$

$$\frac{d\xi}{dr} = -\frac{Z}{rY}, \quad (34d)$$

It is sufficient to show that the right-hand sides of system (34) are bounded as $r \rightarrow r_0^*$. We shall use the monotonicity properties of $X(r)$, $Z(r)$, and $\xi(r)$ following directly from (34a), (34c), (34d).

Consider first the case $\xi(r) > 0$. In the domain $\xi > 0$, where $W(\xi)$ and $W'(\xi)$ are bounded, Eq. (34a) implies that $X(r) > 0$ is monotonically increasing and bounded for $r \rightarrow r_0^*$. Therefore, the right-hand sides of (34b), (34c) are bounded and integrable as $r \rightarrow r_0^* + 0$ and there exist finite limits $Z(r) \rightarrow Z_0$ and $Y(r) \rightarrow Y_0$.

Evidently, $Z_0 > 0$ because $Z(r)$ is a strictly decreasing function as a consequence of (34c).

In order to estimate the value of Y_0 from below, we consider an interval $(r_0^*, r_1]$ for some $r_1 < \infty$. Taking into account that $X(r)$ is a monotonically increasing function, we can choose it as an independent variable. After division of (34b) by (34a), we have, for $r_0^* < r < r_1$,

$$-\frac{d}{dX} \left(\frac{r_g}{Y} \right) = \frac{r}{3Z^2(r)} [1 - r^2 W(\xi(r))] < \frac{r_1}{3Z^2(r_1)},$$

where we have taken into account $Z(r) > Z(r_1)$. After integration from $X(r)$ to $X(r_1)$, we obtain

$$\frac{1}{Y(r)} < \frac{1}{Y(r_1)} + \frac{r_1 X(r_1)}{3r_g Z^2(r_1)}. \quad (35)$$

This excludes the case $Y_0 = 0$ yielding $Y_0 > 0$.

Now we see that the right-hand sides of (34a), (34d) are also bounded and integrable and there exist finite limits $X(r)$ and $\xi(r)$ for $r \rightarrow r_0^* > 0$.

Thus, the whole system (34) is regular for $r \rightarrow r_0^*$ and, according to the local existence and uniqueness theorems, it can be extended to the left neighbourhood of this point. The contrary assumption is false and we must put $r_0^* = 0$.

Now we can repeat considerations concerning $Y(r), Z(r)$ on $(0, \infty)$ yielding

$$Z_0 = \lim_{r \rightarrow 0^+} Z(r) > 0 \quad (36)$$

and

$$Y_0 = \lim_{r \rightarrow 0^+} Y(r) > 0. \quad (37)$$

This completes the consideration of the case $\xi(r) > 0$.

The case of $\xi(r) < 0$ differs from that considered above due to the exponential behavior of $W(\xi)$, $W'(\xi)$ for $\xi < 0$.

However, here we also can show that the right-hand side of (34c) is bounded by using the same reasoning as in [35] (Lemma 4); see Appendix B. Further consideration is similar to the case of positive $\xi(r)$.

Finally, we summarize that *solution* $\alpha(r) \in C^1$, $\beta(r) \in C^1$, $\xi(r) \in C^2$ of Eqs. (22)–(24) satisfying (25), (27) exists for all $r > 0$ and is unique. Moreover, there exist the limits from the right (36), (37).

Using the estimates of Y_0 and Z_0 , we infer a logarithmic behavior of α , β and ξ corresponding to a power-law behavior of $X(r)$ and $Y(r)$ as $r \rightarrow 0$:

$$x \sim \eta \ln \left(\frac{r}{r_g} \right), \quad \xi(r) \sim -\zeta \ln \left(\frac{r}{r_g} \right), \quad (38)$$

$$\alpha(r) \sim (\eta - 1) \ln \left(\frac{r}{r_g} \right), \quad \beta(r) \sim (\eta + 1) \ln \left(\frac{r}{r_g} \right), \quad (39)$$

where $\zeta = Z_0/Y_0 > 0$ and $\eta = 3\zeta^2$. These constants can be related with asymptotic parameters \mathfrak{M} and Q at infinity by means of numerical methods.

Relations (38), (39) justify the choice of zero approximation for the iteration procedure described in Appendix C, which enables us to obtain more detailed asymptotic relations for $r \rightarrow 0$. Here we present the resulting leading orders of the metric coefficients,

$$e^\alpha(r) \approx X_0 Y_0 \left(\frac{r}{r_g} \right)^{\eta-1} \left[1 - \sigma \frac{\eta-1}{(\eta+1)^2} \left(\frac{r}{r_g} \right)^{\eta+1} \right], \quad (40)$$

$$e^\beta(r) \approx \sigma \left(\frac{r}{r_g} \right)^{\eta+1} \left[1 - \sigma \frac{3\eta+1}{(\eta+1)^2} \left(\frac{r}{r_g} \right)^{\eta+1} \right], \quad (41)$$

and of the scalaron field,

$$\xi(r) \approx -\zeta \ln \left(\frac{r}{r_g} \right) + \xi_0 + \frac{\zeta \sigma}{(\eta+1)^2} \left(\frac{r}{r_g} \right)^{\eta+1}, \quad (42)$$

where $\zeta = Z_0/Y_0$, $\sigma = X_0/Y_0$, $\eta = 3\zeta^2$, and $X_0 > 0$ is the constant arising in (C7). In case of asymptotically flat configurations these constants are related to the parameters Q and $M\mu$. These dependencies can be obtained numerically; the example for $\eta = \eta(M\mu)$ is presented on Fig. 3.

For the original metric in the Jordan frame, we have

$$ds^2 = A(r)dt^2 - B(r)dr^2 - R^2(r) (d\theta^2 + \sin^2 \theta d\varphi^2), \quad (43)$$

where according to (3)

$$A(r) = e^{\alpha-2\xi}, \quad B(r) = e^{\beta-2\xi}, \quad R(r) = r e^{-\xi}. \quad (44)$$

The Kretschmann scalar behaves as

$$K = R_{\alpha\beta\gamma\delta} R^{\alpha\beta\gamma\delta} \sim \left(\frac{r_g}{r} \right)^{6(1+\zeta^2)+4\zeta}. \quad (45)$$

Using (44), it is easy to verify that radial null geodesics emanating from the origin can reach the external observer in finite time. Thus, for arbitrarily small SF $\xi \neq 0$, there is a naked singularity in the center.³

Transition from (43) to the curvature coordinates yields

$$ds^2 = \tilde{A}dt^2 - \tilde{B}dR^2 - R^2(d\theta^2 + \sin^2\theta d\varphi^2), \quad (46)$$

where the leading terms as $R \rightarrow 0$ are

$$\tilde{A}(R) = X_0 Y_0 e^{3\xi_0(\zeta-1)} \left(\frac{R}{r_g}\right)^t \left[1 + \mathcal{O}\left(\frac{R}{r_g}\right)^s\right],$$

$$\tilde{B}(R) = \frac{\sigma}{(\zeta+1)^2} e^{s\xi_0} \left(\frac{R}{r_g}\right)^s \left[1 + \mathcal{O}\left(\frac{R}{r_g}\right)^s\right],$$

$$t = 3\zeta - 1, \quad s = \frac{3\zeta^2 + 1}{\zeta + 1}, \quad \zeta \neq -1.$$

The integer values of s, t found in [25] can be obtained in part with $\zeta = -2/3, \pm 1/3, 1, 3$. The cases $(t, s) = (0, 0), (1, -1)$ indicated in [25] turn out to be impossible in our analysis of asymptotically flat configurations with $\zeta \neq 0$. Note that taking into account the next order terms from (40)–(42) shows that metric coefficients in the general case cannot be represented in the form of the Frobenius expansion, as assumed in [25] (cf. also [56]); this is true only for special values of ζ .

VI. SPHERICAL PERTURBATIONS

Our aim is to find a region of parameters μ, Q for which static SS configurations described by Eqs. (18)–(21) with $\hat{T}_{\mu,\nu} \equiv 0$ are unstable against small perturbations. To check for stability/instability issues, we consider linear perturbations, which can be expressed as superposition of functions with the time dependence⁴ $\sim \exp(-i\Omega t)$, $\Omega \neq 0$. The boundary conditions for the spatial dependence of the perturbations will be given below.

There is an extensive literature on the linear perturbations against spherically symmetric background (see, e.g., [57, 58]). The perturbations can be separated into axial and polar modes [57, 58], which can be treated independently. However, in order to show that the system is unstable, it is sufficient to show that there exists at least one unstable mode. Correspondingly, we will restrict ourselves to the case of radial perturbations. In this case, the treatment of our problem follows the same scheme as in [43].

³ Here, the center of the spherically symmetric system is defined as the origin of the curvature coordinates for metric (17).

⁴ Strictly speaking, we consider a class of time-dependent perturbations that satisfy certain boundary conditions as functions of r (see below) and can be expressed in the form of a Laplace/Fourier transform as functions of t .

Based on the exact equations (18)–(21) in vacuum ($\hat{T}_{\mu\nu} \equiv 0$), we perturb the static SS background $\alpha(r), \beta(r), \xi(r)$ by considering $\alpha(r) + \alpha_1(t, r), \beta(r) + \beta_1(t, r), \xi(r) + \xi_1(t, r)$, where α_1, β_1, ξ_1 represent small perturbations.

After linearization, Eq. (20) yields

$$\dot{\beta}_1 = 6r\dot{\xi}_1\xi'.$$

Neglecting the static additive, we have

$$\beta_1 = 6r\xi_1\xi'. \quad (47)$$

The linearized sum of Eqs. (18)–(19) yields

$$\frac{\alpha'_1 - \beta'_1}{2} = \xi_1 e^{\beta} \{6\xi' [1 - r^2 W(\xi)] - rW'(\xi)\}. \quad (48)$$

Eqs. (47), (48) allow us to express all perturbations through ξ_1 . Then we perform linearization of (21) taking into account this equation for the background values, and substitute $\xi_1 = r^{-1}\Phi$ to obtain the master equation in the form

$$\frac{\partial^2 \Phi}{\partial t^2} - e^{\frac{\alpha-\beta}{2}} \frac{\partial}{\partial r} \left[e^{\frac{\alpha-\beta}{2}} \frac{\partial \Phi}{\partial r} \right] + U_{\text{eff}} \Phi = 0, \quad (49)$$

where

$$U_{\text{eff}}(r) = \frac{e^{\alpha-\beta}}{r} \frac{\alpha' - \beta'}{2} + \frac{1}{6} e^{\alpha} W''(\xi) - e^{\alpha} \xi' [6\xi' (1 - r^2 W(\xi)) - 2rW'(\xi)]. \quad (50)$$

Using (23) and (34d), we get

$$U_{\text{eff}}(r) = -\frac{e^{\alpha-\beta}}{r^2} + \frac{e^{\alpha}}{r^2} [1 - r^2 W(\xi)] \left(1 - 6\frac{Z^2}{Y^2}\right) - 2e^{\alpha} \left[\frac{Z}{Y} W'(\xi) - \frac{1}{12} W''(\xi)\right]. \quad (51)$$

Typical examples of U_{eff} are shown in Fig. 4.

The initial value problem related to equation (49) would be incomplete without a boundary condition at the center and at the infinity. While behavior at infinity is treated in a standard way, we have no physical idea of what happens near a naked singularity. On the other hand, the problem must be mathematically correct in order to provide a unique solution to the linearized equations [43, 44, 46, 59–61]. Our analysis below is analogous to that of [43], which takes into account that we deal with small perturbations.

For a single mode $\Phi \sim \exp(-i\Omega t)$ we have from (49)

$$e^{\frac{\alpha-\beta}{2}} \frac{\partial}{\partial r} \left[e^{\frac{\alpha-\beta}{2}} \frac{\partial \Phi}{\partial r} \right] + [\Omega^2 - U_{\text{eff}}] \Phi = 0, \quad (52)$$

The asymptotic behavior of $U_{\text{eff}}(r)$ as $r \rightarrow 0$ is dominated by the term

$$\frac{e^{\alpha-\beta}}{r^2} \approx \frac{K_0}{r^4} \left[1 + \frac{2\sigma}{\eta+1} \left(\frac{r}{r_g}\right)^{\eta+1} \right], \quad (53)$$

where $K_0 = Y_0^2 r_g^2$ is not contained in the general asymptotic solution of (52)

$$\Phi(r) = C_1 r + C_2 r \ln(r/r_g), \quad r \rightarrow 0. \quad (54)$$

In order that this solution and its derivatives be regular at the origin, we must put $C_2 = 0$; otherwise, the perturbation cannot be considered small. Moreover, if $C_2 \neq 0$, then the perturbed Kretschmann scalar turns out to be much larger than (45) as $r \rightarrow 0$ (cf. considerations in [44]).

Therefore, we consider solutions of (52) with regular first derivative at $r = 0$ corresponding to

$$\Phi(r) = C_1 r, \quad C_2 = 0, \quad r \rightarrow 0. \quad (55)$$

The boundary condition at spatial infinity is

$$\Phi(r) \rightarrow 0, \quad r \rightarrow \infty. \quad (56)$$

In terms of the ‘‘tortoise’’ radial variable r_* defined as

$$\frac{dr_*}{dr} = e^{(\beta-\alpha)/2}, \quad r_*(0) = 0, \quad (57)$$

Eq. (52) can be written in the familiar form

$$\frac{\partial^2 \Phi}{\partial r_*^2} - U_{\text{eff}}[r(r_*)] \Phi = -\Omega^2 \Phi. \quad (58)$$

Thus we have the problem of finding an eigenpair $(\Omega^2, \Phi \in C^2)$ for the symmetric operator on the left-hand side of (58) with the null Dirichlet boundary condition at the center and condition (56) at infinity⁵. To be precise, we work in the space of C^2 -functions, which are square integrable on $(0, \infty)$. Obviously, the eigenvalues Ω^2 of this problem are real; for brevity, we call solutions ‘‘unstable’’ if $\Omega^2 < 0$ and ‘‘stable’’ otherwise.

For large values of the radial variable, taking into account (28) and (13), we have

$$W_{\text{eff}}''(r) \approx \left(1 - \frac{r_g}{r}\right) \mu^2, \quad r \rightarrow \infty. \quad (59)$$

The corresponding asymptotic for large r is

$$\Phi(r) \sim D_1(\Omega) e^{kr} + D_2(\Omega) e^{-kr}, \quad k = \sqrt{\mu^2 - \Omega^2}.$$

Therefore, a necessary condition for (56) to be satisfied is $\mu^2 > \Omega^2$. The equation for the eigenvalues is

$$D_1(\Omega) = 0, \quad (60)$$

and it ensures condition (56).

VII. NUMERICAL SOLUTIONS

A. Static SS solutions

We performed backward numerical integration of (23), (24) with respect to α , β , and ξ starting from some sufficiently large value of the radial variable $r_{\text{init}} \gg r_g$ and $\mu r_{\text{init}} \gg 1$, where ξ can be assumed sufficiently small and one can use the asymptotic relations (28) to specify the initial conditions.⁶ We typically opted $r_{\text{init}} \sim 100 r_g$ for moderate μ . Note that, in this case, the backward integration is more convenient than using the shooting method, which involves several shooting parameters.

Numerical investigations show that $e^{\alpha(r)}$ is monotonically increasing, while there is a maximum of $e^{\beta(r)}$ at some r_{max} so that this function is monotonically increasing in the interval $(0, r_{\text{max}})$ and decreasing for $r > r_{\text{max}}$. Near the origin, $e^{\alpha(r)} > e^{\beta(r)}$, but, at some $r = r_c$, the sign of the inequality changes and we have $e^{\alpha(r)} < 1 < e^{\beta(r)}$ for $r > r_c$.

For fixed M, Q , as μ increases, the maximum of $e^{\beta(r)}$ grows and becomes very sharp, whereas, to the left of this maximum, the graphs of $e^{\alpha(r)}$, $e^{\beta(r)}$ are pressed to the abscissa axis. The same situation is observed with fixed μ as Q decreases and becomes small. To the right of the maximum, we have $e^{\alpha(r)} < e^{\beta(r)}$; as $r \rightarrow \infty$, these functions tend to the asymptotic value equal to unity. The larger is μ , the faster the asymptotic values are reached.

There are different modes of behavior of $Y(r)$. For fixed M and μ , and for a sufficiently small Q , we have $r^2 W(\xi(r)) \leq 1$ for all r , and $Y(r)$ is monotonically increasing with the asymptotics $Y(r) \approx r$ for large r . For sufficiently large $|Q|$, we have an interval with negative right-hand side of (34b), where $Y(r)$ is a decreasing function. This interval is bounded because $\xi(r)$ is an exponentially decreasing function and, for sufficiently large r , again $r^2 W(\xi(r)) < 1$, and $Y(r)$ is monotonically increasing. This behavior is typical in case of large $M\mu$; in this case we have a plateau of almost constant $Y(r), Z(r)$ near the center. Far from the center, SF tends to zero exponentially according to Eq. (28) and (D2) leading to $e^\alpha \approx 1 - r_g/r \approx e^{-\beta}$.

For $M\mu \gg 1$ we used the approximation (D2) of Appendix D, which enables us to get the initial conditions for the backwards numerical integration with, e.g., $r_{\text{init}} \sim (5 \div 10) r_g$ and $\xi(r_{\text{init}}) \sim 0.01$. Here we practically have the Schwarzschild metric for $r > r_0$. Note that the case $M\mu \gg 1$ is quite difficult to process on a laptop due to the rapid change of solutions near the region where the condition of smallness of ξ begins to be violated.

Figs. 1 and 2 show the resulting metric coefficients $\tilde{A}(R)$ and $\tilde{B}(R)$ recalculated to the curvature coordinates of the Jordan frame (46). Their behavior is qualitatively

⁵ For numerical investigations, it is more convenient to deal directly with initial equation (52).

⁶ Asymptotic formulas for $d\xi/dr$ are obtained by formal differentiation of (28).

similar to the case of the Einstein frame. The values (Y_0, Z_0) are derived numerically according to (37), (36); this enables us to obtain the asymptotic parameters ζ and $\eta = 3\zeta^2$. Figure 3 shows the dependence $\eta(M\mu)$.




FIG. 1. Typical behavior of $\tilde{A}(R)$ in (46) for different values of $M\mu$ and for $Q = 10$.




FIG. 2. Typical behavior of $\tilde{B}(R)$ in (46) for different values of $M\mu$ and for $Q = 10$.

B. (In)stability against radial perturbations

Here we present the results of numerical analysis on linear stability/instability. We imposed null Dirichlet conditions in the center, because otherwise perturbations cannot be considered small. The algorithm of calculations involved two main stages.

- We chose the initial conditions for static SS (background) solutions according to the asymptotic for-




FIG. 3. Behavior of $\eta = 3\zeta^2$ as a function of $M\mu$ for different values of Q .

mula (28) at sufficiently large r and perform backward integration to get these solutions according to subsection VII A.

- In order to check stability of the static SS solution described by α, β, ξ , we considered numerical solutions of the initial-value problem for (52) with initial data according to the asymptotic formula (55) for sufficiently small r . We used the shooting method with the single shooting parameter Ω^2 to check condition (60) yielding the correct asymptotic behavior (56) at infinity. This procedure was carried out for a set of different parameters $(M\mu, Q)$.

In particular, we found that the static SS configurations considered above are linearly unstable at least in some region of parameters, which includes, for fixed $M\mu$, sufficiently small values of Q . This is enough to state that corresponding configurations are unstable; at the same time, the existence of a white region in Fig. 6 does not mean stability with respect to another types of perturbations.

Fig. 5 demonstrates typical dependencies of the eigenvalues Ω as functions of Q . The domains of $(M\mu, Q)$, which correspond to linearly stable and unstable SS solutions against radial perturbations are shown in Fig. 6. Transition to the Jordan frame preserves the time dependence $\sim e^{-i\Omega t}$ and, therefore, the stability/instability domains in Fig. 6.

VIII. DISCUSSION

We studied spherically symmetric configurations of the quadratic $f(R)$ gravity [4]. Transition to the Einstein frame made possible purely analytical treatment of static global solutions using the method of [35]. As follows from our findings, static asymptotically flat SS solutions with





FIG. 4. Typical examples of the effective potential $U_{\text{eff}}[r(r_*)]$ for radial perturbations in the case of $M\mu = 1$. The upper/lower panel shows the case of positive/negative Q . In both cases, $U_{\text{eff}} \rightarrow \mu^2$ as $r_* \rightarrow \infty$.

zero $T_{\mu\nu}$ are regular outside the center for any scalaron mass and for the global constants $\mathfrak{M} > 0$, $Q \neq 0$; they do not have spherical singularities, which can arise, e.g., in configurations with nonlinear scalar-field potentials [47].

In the case of an isolated regular SS structure with a continuous mass-energy distribution in the central region, taking into account the experimental and theoretical bounds on the scalaron mass, the contribution of the scalaron field to the observable effects are expected to be very small. In this case, the gravitational field is determined by the energy-momentum tensor inside the body. Outside the SS body, the field is completely determined by two parameters: the mass \mathfrak{M} and the “scalar charge” Q .

Returning to the case of a purely gravitational configuration without ordinary matter ($T_{\mu\nu} \equiv 0$) we state that the parameters \mathfrak{M} and Q act as the only characteristics of the static spherically symmetric system, and




FIG. 5. Typical dependencies of $\Omega^2(Q)$. Black curves: $Q > 0$; blue curves: $Q < 0$.




FIG. 6. The grey region corresponds to linear instability against radial perturbations of the static SS configurations defined by the parameters $M\mu$ and Q . The boundary line corresponds to zero eigenvalues satisfying (60).

there is necessarily a naked singularity at the center if $Q \neq 0$. In this regard, we confirm the statement of [25] concerning the occurrence of NS at the center (although some provisions of this work need adjustment). The only exception is the Schwarzschild solution that corresponds to zero scalaron field ($Q = 0$). It is important to note that, whatever small nonzero scalar charge could be, it strongly affects the solution of the quadratic $f(R)$ gravity [4] in the interior region ($r \lesssim r_g$) that is visible to an external observer and permeable to external signals moving towards the singularity. Apparently, this situation is typical for many isolated systems with a scalar field, ranging from the linear massless scalar field [62, 63] to configurations with more complicated potentials [35].

No matter how small the scalar field is, it has a strong impact on the properties of spacetime in this region. At the same time, for large values of Q , the deviations from the Schwarzschild metric can be noticeable at distances as large as, say, $r \sim 5\text{--}10 r_g$. In any case, it is reasonable to defer detailed discussion of the corresponding observational effects until stability issues are resolved.

Our results on the global properties of static SS solutions can be generalized to a wider class of $f(R)$ gravity models. Most restrictive assumption of Section V is the monotonicity requirement (31). From (4) we have

$$W'(\xi) = e^{-4\xi} \left[u \frac{df}{du} - 2f(u) \right].$$

Taking into account that $\text{sgn } \xi = \text{sgn } (f' - 1)$, we have that (31) is satisfied if

$$\left(\frac{df}{du} - 1 \right) \left(u \frac{df}{du} - 2f \right) > 0.$$

Another restriction deals with the behavior of $|f(u)|$ as $|u| \rightarrow \infty$; it must be not faster than $\sim u^2$.

In this paper, we were not able to propose a completely analytic treatment of stability, and we used numerical methods. We derived equations for linear perturbations and studied the asymptotic properties of their solutions near the center and for large r . As it is pointed out in Section VI, the physical conditions near the naked singularity are unknown; for example, the singularity may be radiating, absorbing or something else. However, we are limited to the assumption of small perturbations, which leads us to the restriction (55) and corresponds to the null Dirichlet condition at the center. With this condition we found exponentially growing modes of perturbations (satisfying boundary conditions at the center and at infinity) for some $M\mu, Q$ shown in the grey region of Fig. 6. The static configurations with μ, Q from this region are unstable with respect to linear perturbations. Although our numerical simulations involve moderate values of Q and $M\mu$, they suggest that the domain of instability becomes larger for large $M\mu$ so as to include astrophysically interesting values.

On the other hand, there exists the region of parameters, where linear perturbations are bounded (the white region in Fig. 6). This could indicate the existence of stable configurations, if not for the following circumstances: (i) in case of a nonlinear system, the existence of bounded linear perturbations does not always mean stability, and nonlinear corrections must be studied (cf., e.g., [44]); (ii) other types of linear perturbations (polar and axial ones) are also possible. In fact, there are general results on the linear stability against axial perturbations leading to a Schrödinger-type equation [64]. One can show that the wave potential in this equation appears to be positive-definite in case of our problem; this can be used to prove stability with respect to axial linear perturbations. However, investigation of polar perturbations is mandatory.

Acknowledgements. O.S. is grateful to Igor Klebanov for kind hospitality at Princeton University. V.I.Z. and Yu.V.S. acknowledge partial support under the scientific program ‘‘Astronomy and space physics’’ of the Taras Shevchenko National University of Kyiv. The work of Yu.V.S. was supported by the Simons Foundation and by the National Academy of Sciences of Ukraine under project 0121U109612.

Appendix A: Asymptotic properties as $r \rightarrow \infty$

Here we prove the existence of the unique solutions of (22)–(24) with conditions (25), (27); at the same time, our aim is to justify formula (28). The proof uses the standard techniques of the theory of differential equations used in considerations of the conditional stability [65]. We will isolate the linear part of the equations using specific properties of the system, separate the nonlinear terms and reduce the problem to application of the contraction mapping principle. The specific feature of our treatment is that we reduce the problem to integral equations so that to have expressions like (28) at the first iteration.

Further we use a new field variable $\psi = r\xi$; also it is natural to rewrite (22), (23) using the variables

$$x = \frac{\alpha + \beta}{2}, \quad y = \frac{\beta - \alpha}{2} + \ln(1 - r_g/r). \quad (\text{A1})$$

This gives

$$\frac{dx}{dr} = F_2(r, \psi, d\psi/dr), \quad (\text{A2})$$

$$\frac{dy}{dr} + \frac{x + y}{r - r_g} = F_3(r, x, y, \psi, d\psi/dr), \quad (\text{A3})$$

where the nonlinear terms (starting from the second order in x, y, ψ , and $d\psi/dr$), are collected into F_2 and F_3 . Due to (25), we have

$$x(\infty) = y(\infty) = 0. \quad (\text{A4})$$

Using (23), we write Eq. (24) for static SF in the form

$$\frac{d^2\psi}{dr^2} + \left(\frac{d\psi}{dr} - \frac{\psi}{r} \right) \left[\frac{1}{r} (1 - e^\beta) + r e^\beta W(\xi) \right] = \frac{r}{6} e^\beta W'(\xi). \quad (\text{A5})$$

Here, in the first order, we have $e^\beta = 1 - r_g/r$ and $W'(\xi) = 6\mu^2\xi$. After some rearrangement, we collect the linear part of (A5) into the left-hand side:

$$\frac{d^2\psi}{dr^2} - \left(\frac{d\psi}{dr} - \frac{\psi}{r} \right) \frac{r_g}{r(r - r_g)} - \frac{\mu^2\psi}{1 - r_g/r} = F_1(r, \alpha, \beta, \psi, d\psi/dr), \quad (\text{A6})$$

where F_1 contains the nonlinear terms in x, y, ψ , and $d\psi/dr$.

For arbitrary $n = 1, 2, \dots$, one can find an asymptotic solution⁷ of (A6) in the form

$$v(r, \mu) = e^{-\mu r} \left(\frac{r_g}{r} \right)^m \left(1 + \sum_{k=1}^n \frac{b_k(\mu)}{r^k} \right), \quad (\text{A7})$$

$m \equiv m(\mu) = M\mu$, which satisfies the *exact* relation

$$\frac{d^2 v}{dr^2} - \left(\frac{dv}{dr} - \frac{v}{r} \right) \frac{r_g}{r(r-r_g)} - \frac{\mu^2 v}{1-r_g/r} = \frac{p_n(r, \mu)v}{r^{n+2}}, \quad (\text{A8})$$

where $p_n(r, \mu)$ is a rational function of r , bounded as $r \rightarrow \infty$; the coefficients b_k are determined recursively. In particular, in the case of formula (28), we have $n = 2$; the validity of (A8) can be checked by direct substitution.

Denote

$$\psi_1(r) = v(r, \mu), \quad \psi_2(r) = v(r, -\mu) \sim e^{\mu r} \left(\frac{r}{r_g} \right)^m. \quad (\text{A9})$$

The equation for ψ_2 is obtained from (A8) by the change $\mu \rightarrow -\mu$ leading also to $m \rightarrow -m$. Now we have the exponentially decaying approximate solution of (A6) and exponentially growing one. The Wronskian of ψ_1, ψ_2 is calculated directly to be $w_{12} = 2\mu [1 + \mathcal{O}(r^{-1})]$.

We use the method of variation of constants by setting

$$\psi(r) = K_1(r)\psi_1(r) + K_2(r)\psi_2(r), \quad (\text{A10a})$$

$$\frac{d\psi}{dr} = K_1(r)\frac{d\psi_1}{dr} + K_2(r)\frac{d\psi_2}{dr}, \quad (\text{A10b})$$

where the ‘‘variable constants’’ are K_1 and K_2 . Unlike the usual method of variation of constants, functions ψ_1, ψ_2 are not exact solutions of the linear part of (A6), but satisfy (A8) with different signs of μ .

Let $Q_* = Qr_g$. In order to have the asymptotics $\psi \sim Q_*\psi_1(r)$ as $r \rightarrow \infty$, corresponding to (28), we require

$$K_1(\infty) = Q_*, \quad K_2(\infty) = 0. \quad (\text{A11})$$

A consequence of (A10) is

$$\frac{dK_1}{dr}\psi_1(r) + \frac{dK_2}{dr}\psi_2(r) = 0. \quad (\text{A12})$$

Now we differentiate (A10b) and substitute the result into (A6) using (A8) and (A10). We obtain

$$\frac{dK_1}{dr} \frac{d\psi_1}{dr} + \frac{dK_2}{dr} \frac{d\psi_2}{dr} = -G(r, K_1, K_2) + F_1, \quad (\text{A13})$$

where

$$G(r, K_1, K_2) = \frac{p_n(r, \mu)K_1\psi_1 + p_n(r, -\mu)K_2\psi_2}{r^{n+2}}. \quad (\text{A14})$$

Combination of (A12) and (A13) yields separate equations for dK_1/dr and dK_2/dr and then, by virtue of (A11), leads to integral equations

$$K_1(r) = Q_* - \int_r^\infty \frac{G(t) - F_1(t)}{w_{12}(t)} \psi_2(t) dt, \quad (\text{A15})$$

$$K_2(r) = \int_r^\infty \frac{G(t) - F_1(t)}{w_{12}(t)} \psi_1(t) dt. \quad (\text{A16})$$

Equations (A2), (A3) with conditions (A4) lead to the integral equations

$$x(r) = - \int_r^\infty F_2(t) dt, \quad (\text{A17})$$

$$y(r) = - \int_r^\infty \left[\frac{(t-r)F_2(t) + (t-r_g)F_3(t)}{r-r_g} \right] dt. \quad (\text{A18})$$

The integral equations (A15), (A16), (A17), (A18) can be presented in terms of new variables $u \equiv \{u_j\}$, $j = 1, \dots, 4$, by means of the following substitutions:

$$K_1(r) = u_1(r), \quad K_2(r) = u_2(r) \frac{\psi_1(r)}{\psi_2(r)}, \quad (\text{A19})$$

and

$$x(r) = u_3(r)\psi_1(r), \quad y(r) = u_4(r)\psi_1(r). \quad (\text{A20})$$

Accordingly, we have

$$\psi(r) = (u_1 + u_2)\psi_1(r),$$

$$\tilde{G}(r, u) \equiv G(r, K_1, K_2) = \frac{p_n(r, \mu)u_1 + p_n(r, -\mu)u_2}{r^{n+2}}\psi_1,$$

and the functions

$$\tilde{F}_j(r, u) \equiv F_j \left(r, x, y, \psi, \frac{d\psi}{dr} \right), \quad j = 1, 2, 3,$$

should also be expressed in terms of u by means of (A19)–(A20).

⁷ In fact, we look for an asymptotic solution of the linear homogeneous part of (A6) defined by the left-hand side. The nonlinear corrections due to the right-hand side of (A6), which are exponentially smaller than the terms of (A7) as $r \rightarrow \infty$, are ‘‘automatically’’ taken into account in the iteration process below.

Introduce the operator $\mathfrak{A} = \{\mathfrak{A}_1, \mathfrak{A}_2, \mathfrak{A}_3, \mathfrak{A}_4\}$, where

$$\mathfrak{A}_1(r, u) = Q_* - \int_r^\infty \frac{\tilde{G}(t, u) - \tilde{F}_1(t, u)}{w_{12}(t)} \psi_2(t) dt, \quad (\text{A21})$$

$$\mathfrak{A}_2(r, u) = \frac{\psi_2(r)}{\psi_1(r)} \int_r^\infty \frac{\tilde{G}(t, u) - \tilde{F}_1(t, u)}{w_{12}(t)} \psi_1(t) dt, \quad (\text{A22})$$

$$\mathfrak{A}_3(r, u) = -\frac{1}{\psi_1(r)} \int_r^\infty \tilde{F}_2(r, u) dt, \quad (\text{A23})$$

$$\mathfrak{A}_4(r, u) = -\int_r^\infty \frac{(t-r)\tilde{F}_2(t, u) + (t-r_g)\tilde{F}_3(t, u)}{\psi_1(r)(r-r_g)} dt. \quad (\text{A24})$$

With this definition, the operator equation

$$u(r) = \mathfrak{A}(r, u), \quad (\text{A25})$$

is equivalent to system (A6), (A2), (A3) with conditions (27) and (A4) and, therefore, to the system of equations (22)–(24) with conditions (25), (27). Equation (A25) will be considered on interval $[r_0, \infty)$, where $r_0 > r_g$ will be taken sufficiently large.

Note that

$$|\psi(r)| = |(u_1 + u_2)\psi| \leq |u(r)|_4 \psi_1(r), \quad (\text{A26})$$

$$\left| \frac{d\psi}{dr} \right| \leq \mu [1 + \mathcal{O}(r^{-1})] |u(r)|_4 \psi_1(r), \quad (\text{A27})$$

$$|\tilde{G}(r, u)| \leq \frac{C_1}{r^{n+2}} \psi_1(r) |u|_4, \quad C_1 = \mathcal{O}(1), \quad (\text{A28})$$

$|u|_4 = \sum_{i=1}^4 |u_i|$, where we used the explicit expressions (A7), (A9) and representations (A10). Further, we use that, for sufficiently large r ,

$$\begin{aligned} \psi_1(r)\psi_2(r) &= 1 + \mathcal{O}(r^{-1}), \\ \int_r^\infty \psi_1^2(t) dt &= \frac{\psi_1^2(r)}{2\mu} [1 + \mathcal{O}(r^{-1})]. \end{aligned}$$

After lengthy calculations, we obtain that the functions \tilde{F}_j are Lipschitz continuous:

$$|\tilde{F}_j(r, u) - \tilde{F}_j(r, u')| \leq H(u, u') |u - u'|_4, \quad (\text{A29})$$

where $j = 1, 2, 3$,

$$H(u, u') \leq C_2 \max\{|u|_4, |u'|_4\} \psi_1^2(r), \quad C_2 = \mathcal{O}(1).$$

Note that we have two small parameters: $\psi_1(r)$ and $(r_g/r)^{n+1}$; the contribution of the first one to estimates (arising due to the nonlinear terms F_j) is exponentially

smaller for $r \geq r_0$. Then, for sufficiently large r_0 , we have

$$\varepsilon(r_0) = \max\{(r_g/r_0)^{n+1}, \psi_1(r_0)\} = (r_g/r_0)^{n+1}.$$

Denote $\|f\|_a^b \equiv \sup\{f(t), t \in [a, b]\}$. Consider the Banach space \mathcal{B} of bounded continuous functions on $[r_0, \infty)$ with the norm $\|f\|_{r_0}^\infty$ and a domain $\mathcal{D} \subset \mathcal{B} : \forall u \in \mathcal{D}, |u| \leq 2|Q_*|$. The operator \mathfrak{A} is well defined on \mathcal{D} ; all integrals are convergent: in (A21), the integrand is estimated as $\sim r^{-(n+2)}$ due to (A28) and (A29); in (A22)–(A24), we have exponentially decaying expressions in the integrands.

By virtue of the above estimates, we have, for $u \in \mathcal{D}$,

$$|\mathfrak{A}(r, u)| \leq Q_* + C_3 \varepsilon(r_0) \|u\|_r^\infty, \quad (\text{A30})$$

$$|\mathfrak{A}(r, u) - \mathfrak{A}(r, v)| \leq C_1 \varepsilon(r_0) \|u - v\|_{r_0}^\infty, \quad (\text{A31})$$

where constants $C_1, C_3 = \mathcal{O}(1)$. Then from (A30) we have that (for sufficiently large r_0) $\mathfrak{A}(\mathcal{B}) \subset \mathcal{B}$ and, from (A31), we infer the contraction property. Therefore, equation (A25) has a unique solution.

This means that there exists a unique solution $\alpha(r) \in C^1$, $\beta(r) \in C^1$, $\xi(r) \in C^2$ of Eqs. (22)–(24) with conditions (25), (27).

From (A21), (A22) we infer

$$\begin{aligned} |\psi(r) - Q_* \psi_1(r)| &= |u_1(r) - Q_* + u_2(r)| \psi_1(r) \\ &\leq \mathcal{O}(r^{-(n+1)}) \psi_1(r). \end{aligned}$$

This confirms formula (28). Also, using (A23), (A24), we have

$$|u_3(r)| + |u_4(r)| \lesssim \psi_1(r),$$

whence, on the account of (A1),

$$\begin{aligned} \left| \alpha(r) - \ln\left(1 - \frac{r_g}{r}\right) \right| + \left| \beta(r) + \ln\left(1 - \frac{r_g}{r}\right) \right| \\ \lesssim \mathcal{O}[\psi_1^2(r)]. \quad (\text{A32}) \end{aligned}$$

Formula (28) is the result of the first iteration of (A25). The next iterations add higher-order corrections in powers of $1/r^{n+1}$ and $\exp(-\mu r)$, although the latter are useful only if very accurate results are required. In our study, the first iteration yielding (28) and (A32) was quite sufficient to impose the initial conditions for backward numerical integration described in subsection VII A.

Appendix B: The case of $\xi < 0$

Here we use the method of [35]. The key question is to prove that the right-hand sides of (34b) and (34c) are bounded as $r \rightarrow 0$.

Analogously to Section V, consider $r < r_1$ for some $r_1 < \infty$. If $\xi < 0$ for $r \leq r_1$, then $0 < W(\xi) < (3/4)\mu^2 \exp(4|\xi|)$, $|W'(\xi)| < 3\mu^2 \exp(4|\xi|)$, and $|\xi(r)| <$

monotonically decreasing due to (34d). For both right-hand sides (34b) and (34c), it suffices to estimate

$$P(r) \equiv r^2 X(r) e^{4|\xi|} = r_1^2 X_1 e^{4|\xi_1|} e^{-2L+4d-(x_1-x)}, \quad (\text{B1})$$

where $x \equiv x(r) \equiv \ln[X(r)]$, $r < r_1$, and we denoted $x_1 = x(r_1)$, $X_1 = X(r_1) = \exp(x_1)$, $\xi_1 = \xi(r_1)$, $d(r) = |\xi(r)| - |\xi_1| > 0$, $L \equiv L(r) = -\ln(r/r_1) > 0$.

We use the inequality

$$\frac{[\xi(r) - \xi(r_1)]^2}{\ln(r_1/r)} \leq \int_r^{r_1} t \left(\frac{d\xi}{dt} \right)^2 dt, \quad r < r_1,$$

which is valid for any continuously differentiable function ξ (see Eq. (31) of [35]). By virtue of (22),

$$x_1 - x = 3 \int_r^{r_1} t \left(\frac{d\xi}{dt} \right)^2 dt \geq \frac{3}{L} d^2.$$

Then in (B1)

$$\begin{aligned} -2L + 4d - (x_1 - x) &\leq -2L + 4d - \frac{3}{L} d^2 \\ &= -\frac{3}{L} \left(d - \frac{2}{3}L \right)^2 - \frac{2}{3}L < 0, \end{aligned}$$

whence

$$P(r) \leq r_1^2 X_1 (r/r_1)^{2/3} e^{4|\xi_1|}.$$

Therefore, the exponent $e^{4|\xi|}$ is suppressed by $X(r)$; the right hand sides of (34b) and (34c) are bounded. This is sufficient to infer the existence of the finite limits Z_0 and Y_0 for $r \rightarrow 0$. The considerations leading to the lower bounds $|Z_0| > 0$ and $Y_0 > 0$ are similar to the corresponding arguments in Section V for positive $\xi(r)$.

Appendix C: Asymptotic behavior near NS

Here we formulate an iteration procedure to find the asymptotic expansion near NS in curvature coordinates. Taking into account (36) and (37), we have from (34b), (34c)

$$Y(r) = Y_0 + Y_1(r), \quad Z(r) = Z_0 + Z_1(r), \quad (\text{C1})$$

where

$$Y_1(r) = \int_0^r X(t) [1 - t^2 W(\xi(t))] \frac{dt}{r_g} = \mathcal{O}[(r/r_g)^{\eta+1}], \quad (\text{C2})$$

$$Z_1(r) = -\frac{1}{6} \int_0^r t^2 X(t) W'(\xi(t)) \frac{dt}{r_g} = \mathcal{O}[(r\mu)^2 (r/r_g)^{\eta+2\zeta+1}]. \quad (\text{C3})$$

Now we can separate the logarithmic terms in x , ξ by setting

$$x(r) = \eta \ln \left(\frac{r}{r_g} \right) + x_1^*(r), \quad (\text{C4a})$$

$$\xi(r) = -\zeta \ln \left(\frac{r}{r_g} \right) + \xi_1^*(r), \quad (\text{C4b})$$

obtaining equations with smooth right-hand sides:

$$\frac{dx_1^*}{dr} = \frac{3}{r} D_1(r), \quad D_1(r) = \frac{Z^2}{Y^2} - \frac{Z_0^2}{Y_0^2}, \quad (\text{C5})$$

$$\frac{d\xi_1^*}{dr} = \frac{D_2(r)}{r}, \quad D_2(r) = \frac{Z_0}{Y_0} - \frac{Z}{Y}. \quad (\text{C6})$$

On account of (C2), (C3), we have $D_1(r) = \mathcal{O}(r^{\eta+1})$ and $D_2(r) = \mathcal{O}(r^{\eta+1})$.

Then

$$x_1^*(r) = \ln X_0 + 3 \int_0^r \frac{dt}{t} D_1(t), \quad (\text{C7})$$

$$\xi_1^*(r) = \xi_0 + \int_0^r \frac{dt}{t} D_2(t), \quad (\text{C8})$$

$X_0 > 0$ and ξ_0 being free constants.

Equations (C2), (C3), (C7), (C8) with notation (C1), (C4a), (C4b) form a system of integral equations for the functions Y_1 , Z_1 , χ_1 , and ξ_1 . This system is ready for successive approximations dealing with well-defined continuous expressions at each step of the iterative procedure that may be used to obtain the asymptotic solution of four equations (18)–(21) with four constants Y_0 , Z_0 , X_0 , and ξ_0 .

The results of the first iterations can be written as follows

$$Y(r) = Y_0 + \frac{X_0}{\eta+1} \left(\frac{r}{r_g} \right)^{\eta+1} [1 + \mathcal{O}(\mu^2 r^2)], \quad (\text{C9})$$

$$\begin{aligned} x(r) &= \eta \ln \left(\frac{r}{r_g} \right) + \ln X_0 \\ &\quad - \frac{2\eta}{(\eta+1)^2} \frac{X_0}{Y_0} \left(\frac{r}{r_g} \right)^{\eta+1} [1 + \mathcal{O}(\mu^2 r^2)], \end{aligned} \quad (\text{C10})$$

$$\begin{aligned} \xi(r) &= -\zeta \ln \left(\frac{r}{r_g} \right) + \xi_0 \\ &\quad + \frac{\zeta}{(\eta+1)^2} \frac{X_0}{Y_0} \left(\frac{r}{r_g} \right)^{\eta+1} [1 + \mathcal{O}(\mu^2 r^2)], \end{aligned} \quad (\text{C11})$$

where $\zeta = Z_0/Y_0$, $\eta = 3\zeta^2$.

In case of a spatially flat configuration, the constants Y_0 , Z_0 , X_0 , and ξ_0 are functions of the pair M , Q that uniquely defines the configuration. Having the numerical solutions of Sec. VII, we get Z_0 of (36), Y_0 of (37). We note a good numerical convergence to these limits.

Returning to the initial variables e^α , e^β , we obtain the asymptotic formulas (40)–(42).

Appendix D: Approximate SF for large $M\mu$ and small ξ

For $r > r_g$, as r increases, we have $\xi \rightarrow 0$ and the metric reaches the Schwarzschild asymptotics. Moreover, as it was pointed out in Sec. III, the realistic values of $M\mu$ must be very large, so the asymptotic behavior occurs very quickly. Then we can use the WKB method in case of small $\xi \ll 1$, using (13) and (26). Substituting $\xi = e^S$

into (26) gives

$$\frac{d^2 S}{dr^2} + \left(\frac{dS}{dr}\right)^2 + \frac{dS}{dr} \frac{2r - r_g}{r(r - r_g)} = \frac{r\mu^2}{r - r_g}. \quad (\text{D1})$$

Then we set $S = S_0 + S_1 + \dots$, where the first two orders $|S_0| \sim \mathcal{O}(\mu) \gg |S_1| \sim \mathcal{O}(1)$ of the WKB method applied to (D1) yield

$$\xi(r) = \frac{Q_1 \exp\left(-\mu r \sqrt{1 - r_g/r}\right)}{(1 - r_g/r)^{1/4} \left(1 + \sqrt{1 - r_g/r}\right)^{\mu r_g}} \left(\frac{r_g}{r}\right)^{1+M\mu}, \quad (\text{D2})$$

where we must put $Q_1 = Q(4/e)^{M\mu}$ so as to have correspondence with (28) for $r \gg r_g$. Approximation (D2) is effective for $\xi(r) \ll 1$, $\mu\sqrt{r(r - r_g)} \gg \ln(r/r_g)$. Note that formula (28), unlike (D2), cannot be used for r comparable with r_g .

-
- [1] T. P. Sotiriou and V. Faraoni, $f(R)$ theories of gravity, *Reviews of Modern Physics* **82**, 451 (2010), [arXiv:0805.1726 \[gr-qc\]](#).
- [2] A. De Felice and S. Tsujikawa, $f(r)$ theories, *Living Reviews in Relativity* **13**, 3 (2010), [arXiv:1002.4928 \[gr-qc\]](#).
- [3] S. Nojiri, S. D. Odintsov, and V. K. Oikonomou, Modified gravity theories on a nutshell: Inflation, bounce and late-time evolution, *Phys. Rept.* **692**, 1 (2017), [arXiv:1705.11098 \[gr-qc\]](#).
- [4] A. A. Starobinsky, A new type of isotropic cosmological models without singularity, *Phys. Lett. B* **91**, 99 (1980).
- [5] A. Vilenkin, Classical and quantum cosmology of the Starobinsky inflationary model, *Phys. Rev. D* **32**, 2511 (1985).
- [6] Y. Shtanov, V. Sahni, and S. S. Mishra, Tabletop potentials for inflation from $f(R)$ gravity, *J. Cosmol. Astroparticle Phys.* **03**, 023 (2023), [arXiv:2210.01828 \[gr-qc\]](#).
- [7] S. D. Odintsov, V. K. Oikonomou, I. Giannakoudi, F. P. Fronimos, and E. C. Lympieriadou, Recent advances in inflation, *Symmetry* **15**, 1701 (2023), [arXiv:2307.16308 \[gr-qc\]](#).
- [8] Y. Akrami *et al.* (Planck Collaboration), Planck 2018 results. X. Constraints on inflation, *Astron. Astrophys.* **641**, A10 (2020), [arXiv:1807.06211 \[astro-ph.CO\]](#).
- [9] S. Capozziello, V. F. Cardone, and A. Troisi, Dark energy and dark matter as curvature effects, *J. Cosmol. Astroparticle Phys.* **08**, 001 (2006), [arXiv:astro-ph/0602349](#).
- [10] S. Nojiri and S. D. Odintsov, Dark energy, inflation and dark matter from modified $F(R)$ gravity, *TSPU Bulletin* **N8(110)**, 7 (2011), [arXiv:0807.0685 \[hep-th\]](#).
- [11] J. A. R. Cembranos, Dark matter from R^2 gravity, *Phys. Rev. Lett.* **102**, 141301 (2009), [arXiv:0809.1653 \[hep-ph\]](#).
- [12] J. A. R. Cembranos, Modified gravity and dark matter, *J. Phys. Conf. Ser.* **718**, 032004 (2016), [arXiv:1512.08752 \[hep-ph\]](#).
- [13] C. Corda, H. J. Mosquera Cuesta, and R. Lorduy Gomez, High-energy scalarons in R^2 gravity as a model for Dark Matter in galaxies, *Astropart. Phys.* **35**, 362 (2012), [arXiv:1105.0147 \[gr-qc\]](#).
- [14] T. Katsuragawa and S. Matsuzaki, Dark matter in modified gravity?, *Phys. Rev. D* **95**, 044040 (2017), [arXiv:1610.01016 \[gr-qc\]](#).
- [15] T. Katsuragawa and S. Matsuzaki, Cosmic history of chameleonic dark matter in $F(R)$ gravity, *Phys. Rev. D* **97**, 064037 (2018), [Erratum: *Phys. Rev. D* **97**, 129902 (2018)], [arXiv:1708.08702 \[gr-qc\]](#).
- [16] B. K. Yadav and M. M. Verma, Dark matter as scalaron in $f(R)$ gravity models, *J. Cosmol. Astroparticle Phys.* **10**, 052 (2019), [arXiv:1811.03964 \[gr-qc\]](#).
- [17] N. Parbin and U. D. Goswami, Scalarons mimicking dark matter in the Hu–Sawicki model of $f(R)$ gravity, *Mod. Phys. Lett. A* **36**, 2150265 (2021), [arXiv:2007.07480 \[gr-qc\]](#).
- [18] Y. Shtanov, Light scalaron as dark matter, *Physics Letters B* **820**, 136469 (2021), [arXiv:2105.02662 \[hep-ph\]](#).
- [19] Y. Shtanov, Initial conditions for the scalaron dark matter, *J. Cosmol. Astroparticle Phys.* **10**, 079 (2022), [arXiv:2207.00267 \[astro-ph.CO\]](#).
- [20] J. D. Bekenstein, Transcendence of the law of baryon-number conservation in black-hole physics, *Phys. Rev. Lett.* **28**, 452 (1972).
- [21] J. D. Bekenstein, Nonexistence of baryon number for static black holes, *Phys. Rev. D* **5**, 1239 (1972).
- [22] J. D. Bekenstein, Novel “no-scalar-hair” theorem for black holes, *Phys. Rev. D* **51**, R6608 (1995).
- [23] B. Holdom and J. Ren, Not quite a black hole, *Phys. Rev. D* **95**, 084034 (2017), [arXiv:1612.04889 \[gr-qc\]](#).
- [24] R. Kase and S. Tsujikawa, Neutron stars in $f(R)$ gravity and scalar-tensor theories, *J. Cosmol. Astroparticle Phys.* **09**, 054 (2019), [arXiv:1906.08954 \[gr-qc\]](#).
- [25] E. Hernández-Lorenzo and C. F. Steinwachs, Naked singularities in quadratic $f(R)$ gravity, *Phys. Rev. D* **101**, 124046 (2020), [arXiv:2003.12109 \[gr-qc\]](#).
- [26] D. D. Doneva and S. S. Yazadjiev, No-hair theorems for noncanonical self-gravitating static multiple scalar fields, *Phys. Rev. D* **102**, 10.1103/physrevd.102.084055 (2020).

- [27] J.-P. Hong, M. Suzuki, and M. Yamada, Charged black holes in non-linear Q-clouds with $O(3)$ symmetry, *Physics Letters B* **803**, 135324 (2020), [arXiv:1907.04982 \[gr-qc\]](#).
- [28] J.-P. Hong, M. Suzuki, and M. Yamada, Spherically symmetric scalar hair for charged black holes, *Phys. Rev. Lett.* **125**, 111104 (2020), [arXiv:2004.03148 \[gr-qc\]](#).
- [29] A. Bakopoulos and T. Nakas, Analytic and asymptotically flat hairy (ultra-compact) black-hole solutions and their axial perturbations, *Journal of High Energy Physics* **04**, 096 (2022), [arXiv:2107.05656 \[gr-qc\]](#).
- [30] A. De Felice and S. Tsujikawa, Stability of Schwarzschild black holes in quadratic gravity with Weyl curvature domination, *J. Cosmol. Astroparticle Phys.* **10**, 004 (2023), [arXiv:2307.06490 \[gr-qc\]](#).
- [31] A. De Felice and S. Tsujikawa, Excluding static and spherically symmetric black holes in Einsteinian cubic gravity with unsuppressed higher-order curvature terms, *Physics Letters B* **843**, 138047 (2023), [arXiv:2305.07217 \[gr-qc\]](#).
- [32] X. Ye and S.-W. Wei, Distinct topological configurations of equatorial timelike circular orbit for spherically symmetric (hairy) black holes, *J. Cosmol. Astroparticle Phys.* **07**, 049 (2023), [arXiv:2301.04786 \[gr-qc\]](#).
- [33] I. Bogush, D. Gal'tsov, G. Gylchev, K. Kobialko, P. Nedkova, and T. Vetsov, Photon surfaces, shadows, and accretion disks in gravity with minimally coupled scalar field, *Phys. Rev. D* **106**, 024034 (2022), [arXiv:2205.01919 \[gr-qc\]](#).
- [34] H. K. Nguyen, Beyond Schwarzschild–de Sitter spacetimes. III. A perturbative vacuum with nonconstant scalar curvature in $R + R^2$ gravity, *Phys. Rev. D* **107**, 104009 (2023), [arXiv:2211.07380 \[gr-qc\]](#).
- [35] V. I. Zhdanov and O. S. Stashko, Static spherically symmetric configurations with N nonlinear scalar fields: Global and asymptotic properties, *Phys. Rev. D* **101**, 064064 (2020), [arXiv:1912.00470 \[gr-qc\]](#).
- [36] R. Penrose, Gravitational collapse and space-time singularities, *Phys. Rev. Lett.* **14**, 57 (1965).
- [37] R. Penrose, Gravitational collapse: The role of general relativity, *General Relativity and Gravitation* **34**, 1141 (2002).
- [38] D. Christodoulou, Violation of cosmic censorship in the gravitational collapse of a dust cloud, *Communications in Mathematical Physics* **93**, 171 (1984).
- [39] A. Ori and T. Piran, Naked singularities in self-similar spherical gravitational collapse, *Phys. Rev. Lett.* **59**, 2137 (1987).
- [40] P. S. Joshi and I. H. Dwivedi, Naked singularities in spherically symmetric inhomogeneous Tolman-Bondi dust cloud collapse, *Phys. Rev. D* **47**, 5357 (1993).
- [41] P. S. Joshi, D. Malafarina, and R. Narayan, Distinguishing black holes from naked singularities through their accretion disc properties, *Classical and Quantum Gravity* **31**, 015002 (2013).
- [42] Y. C. Ong, Space–time singularities and cosmic censorship conjecture: A Review with some thoughts, *International Journal of Modern Physics A* **35**, 2030007 (2020).
- [43] M. A. Clayton, L. Demopoulos, and J. Légaré, The dynamical stability of the static real scalar field solutions to the Einstein-Klein-Gordon equations revisited, *Physics Letters A* **248**, 131 (1998).
- [44] G. W. Gibbons, S. A. Hartnoll, and A. Ishibashi, On the Stability of Naked Singularities with Negative Mass, *Progress of Theoretical Physics* **113**, 963 (2005), [arXiv:hep-th/0409307 \[hep-th\]](#).
- [45] R. J. Gleiser and G. Dotti, Instability of the negative mass Schwarzschild naked singularity, *Classical and Quantum Gravity* **23**, 5063 (2006), [arXiv:gr-qc/0604021 \[gr-qc\]](#).
- [46] G. Dotti, Linear stability of black holes and naked singularities, *Universe* **8**, 38 (2022).
- [47] O. Stashko and V. I. Zhdanov, Singularities in static spherically symmetric configurations of general relativity with strongly nonlinear scalar fields, *Galaxies* **9**, 72 (2021), [arXiv:2109.01931 \[gr-qc\]](#).
- [48] D. J. Kapner, T. S. Cook, E. G. Adelberger, J. H. Gundlach, B. R. Heckel, C. D. Hoyle, and H. E. Swanson, Tests of the gravitational inverse-square law below the dark-energy length scale, *Phys. Rev. Lett.* **98**, 021101 (2007), [arXiv:hep-ph/0611184](#).
- [49] E. G. Adelberger, B. R. Heckel, S. A. Hoedl, C. D. Hoyle, D. J. Kapner, and A. Upadhye, Particle-physics implications of a recent test of the gravitational inverse-square law, *Phys. Rev. Lett.* **98**, 131104 (2007), [arXiv:hep-ph/0611223](#).
- [50] K. S. Stelle, Classical gravity with higher derivatives, *Gen. Rel. Grav.* **9**, 353 (1978).
- [51] L. Perivolaropoulos and L. Kazantzidis, Hints of modified gravity in cosmos and in the lab?, *Int. J. Mod. Phys. D* **28**, 1942001 (2019), [arXiv:1904.09462 \[gr-qc\]](#).
- [52] R. A. Asanov, Static scalar and electric fields in Einstein's theory of relativity, *Soviet Journal of Experimental and Theoretical Physics* **26**, 424 (1968).
- [53] R. A. Asanov, Point source of massive scalar field in gravitational theory, *Theoretical and Mathematical Physics* **20**, 667 (1974).
- [54] D. J. Rowan and G. Stephenson, The massive scalar meson field in a Schwarzschild background space, *Journal of Physics A Mathematical General* **9**, 1261 (1976).
- [55] O. Stashko and V. Zhdanov, Spherically symmetric configurations in general relativity in the presence of a linear massive scalar field: Separation of a distribution of test body circular orbits, *Ukrainian Journal of Physics* **64**, 189 (2019).
- [56] H. Lü, A. Perkins, C. N. Pope, and K. S. Stelle, Spherically symmetric solutions in higher-derivative gravity, *Phys. Rev. D* **92**, 124019 (2015).
- [57] S. Chandrasekhar, *The Mathematical Theory of Black Holes* (Oxford University Press, Oxford, UK, 1998).
- [58] E. Berti, V. Cardoso, and A. O. Starinets, Quasinormal modes of black holes and black branes, *Classical and Quantum Gravity* **26**, 163001 (2009), [arXiv:0905.2975 \[gr-qc\]](#).
- [59] R. M. Wald, Dynamics in nonglobally hyperbolic, static space-times, *Journal of Mathematical Physics* **21**, 2802 (1980).
- [60] A. Ishibashi and A. Hosoya, Who's afraid of naked singularities? Probing timelike singularities with finite energy waves, *Phys. Rev. D* **60**, 104028 (1999).
- [61] A. Ishibashi and R. M. Wald, Dynamics in non-globally-hyperbolic static spacetimes: II. General analysis of prescriptions for dynamics, *Classical and Quantum Gravity* **20**, 3815 (2003), [arXiv:gr-qc/0305012 \[gr-qc\]](#).
- [62] I. Z. Fisher, Scalar mesostatic field with regard for gravitational effects, *Zh. Exp. Theor. Phys.*, **18**, 636 (1948), [arXiv:gr-qc/9911008 \[gr-qc\]](#).
- [63] A. I. Janis, E. T. Newman, and J. Winicour, Reality of the Schwarzschild singularity, *Phys. Rev. Lett.* **20**, 878

- (1968).
- [64] O. S. Stashko, O. V. Savchuk, and V. I. Zhdanov, Quasi-normal modes of naked singularities in presence of non-linear scalar fields, *Phys. Rev. D* **109**, 024012 (2024), [arXiv:2307.04295 \[gr-qc\]](#).
- [65] E. A. Coddington and N. Levinson, *Theory of Ordinary Differential Equations* (McGraw-Hill, New York, 1955).