PWEXP: An **R** Package Using Piecewise Exponential Model for Study Design and Event/Timeline Prediction

Tianchen Xu

Bristol Myers Squibb

Rachael Wen Bristol Myers Squibb

April 30, 2024

Abstract

Parametric assumptions such as exponential distribution are commonly used in clinical trial design and analysis. However, violation of distribution assumptions can introduce biases in sample size and power calculations. Piecewise exponential (PWE) hazard model partitions the hazard function into segments each with constant hazards and is easy for interpretation and computation. Due to its piecewise property, PWE can fit a wide range of survival curves and accurately predict the future number of events and analysis time in event-driven clinical trials, thus enabling more flexible and reliable study designs. Compared with other existing approaches, the PWE model provides a superior balance of flexibility and robustness in model fitting and prediction.

The proposed **PWEXP** package is designed for estimating and predicting PWE hazard models for right-censored data. By utilizing well-established criteria such as AIC, BIC, and cross-validation log-likelihood, the **PWEXP** package chooses the optimal number of change-points and determines the optimal position of change-points. With its particular goodness-of-fit, the **PWEXP** provides accurate and robust hazard estimation, which can be used for reliable power calculation at study design and timeline prediction at study conduct. The package also offers visualization functions to facilitate the interpretation of survival curve fitting results.

Keywords: piecewise exponential, clinical trial, PWEXP, Change-points, timeline prediction, event prediction, ${\bf R}$

1 Introduction

Survival analysis is a statistical method used to analyze time-to-event data, such as the time from a subject's randomization until he/she experiences an event (e.g., disease progression or death). One of the most commonly parametric models in survival analysis is the exponential hazard model, which assumes a constant hazard rate over time. However, in reality, the hazard rate is barely constant over time. Abrupt or gradual changes in the hazard rates can be observed due to environmental factors, specific maintenance activity, and so on. The change of hazard rates over time could happen multiple times (Qian and Zhang, 2013). A more flexible model is needed and the piecewise exponential hazard model is a popular alternative. The piecewise exponential (PWE) model partitions the hazard over time into several time intervals, assuming the hazard rates are constant within each time interval, but can vary between intervals. By selecting appropriate

			${f R}$ Package		
	PWEXP	pch	muhaz	eventTrack	Piecewise- Changepoint
Method	MLE/OLS	MLE	MLE	PML	Bayesian
Model estimation with					
Given change-points	\checkmark	\checkmark	×	×	×
Unknown change-points	\checkmark	$\times^{(a)}$	$\times^{(b)}$	$\checkmark^{(c)}$	\checkmark
Partial known change-points	\checkmark	×	×	×	×
Tail robustness adjustment	$\checkmark^{(d)}$	×	×	×	×
Inference					
CI/Asymptotic variance	\checkmark	\checkmark	×	\checkmark	$\checkmark^{(e)}$
Change-point test	×	×	×	\checkmark	×
Model summary					
AIC, BIC	\checkmark	\checkmark	×	×	×
PML, WAIC ^{(f)}	×	×	×	×	\checkmark
Likelihood	\checkmark	\checkmark	×	\checkmark	\checkmark
CV likelihood	\checkmark	×	×	×	×

Table 1: Comparison between the **PWEXP** package and other alternatives for PWE model estimation. The symbol \checkmark indicates the feature is available, and \times indicates it is not available. Note: (a) The **pch** package uses empirical quantiles as change-points; (b) The **muhaz** package uses even length intervals for change-points; (c) The **eventTrack** uses general-purpose optimizer "Nelder-Mead Simplex" method to estimate change-points; (d) The **PWEXP** provides several approaches to avoid an unstable tail (See Section 2.6); (e) Credible interval; (f) PML: Pseudo-Marginal Likelihood, WAIC: Widely Applicable Information Criterion.

change-points to create closely-spaced boundaries and dividing the time axis into smaller intervals where the hazard changes rapidly and wider intervals where the hazard changes slowly, the hazard rate within each time interval can be estimated and their estimations can be used to approximate the Kaplan-Meier (KM) survival curves. Such approximation capability is due to its inherent flexibility which allows it to effectively capture a wide range of hazard rate patterns (Matthews and Farewell, 1982).

Estimated piecewise hazard rates from from historical data can provide assumptions for clinical trial design, including sample size power calculation, interim analysis (IA) plan, trial duration estimation, and resource allocation, etc. Commonly used packages for clinical trial design such as **rpact** (Wassmer and Pahlke, 2024), **gsDesign** (Anderson, 2023) can utilize piecewise exponential models, which will be illustrated in later sections.

Over the past decades, significant progress has been made in developing mature theories regarding the asymptotic properties of estimators in PWE models. Friedman (1982); Yao (1986) demonstrated that the maximum likelihood (ML) change-point estimator is consistent under certain conditions. Henderson (1990) derived exact critical values for likelihood ratio test for a single change-point and the corresponding piecewise exponential hazard rates. Goodman et al. (2011) proposed a Wald-type test statistic to test the existence of multiple change-points in a PWE model.

However, despite the significant progress in PWE model inference, there are still limited approaches available to accurately estimate the change-points in real datasets. For a PWE model with one single changepoint, it is recommended to employ grid search or solve the score equations using numerical algorithms like

	R Package			
_	PWEXP	eventTrack	gestate	eventPred
Sub-model				
Accural model	uniform/user defined	user defined	user defined	(piecewise) Poisson/time- decay/B-spline
Event model	$\exp/pwexp$	KM+exp tail	exp/Weibull/log- normal	various ^(a)
Drop-out model	$\exp/pwexp$	$\begin{array}{llllllllllllllllllllllllllllllllllll$		$various^{(a)}$
Prediction			,	
Event prediction	\checkmark	\checkmark	\checkmark	\checkmark
Timeline prediction	\checkmark	×	×	\checkmark
Confidence interval	\checkmark	×	\checkmark	$\checkmark^{(c)}$

Table 2: Comparison between the **PWEXP** package and other alternatives for event/timeline prediction. The symbol \checkmark indicates the feature is available, and \times indicates it is not available. Note: (a) The **eventPred** includes these event/drop-out models: exponential, Weibull, log-logistic, log-normal, piecewise exponential (with known change-points), model averaging, spline; (b) These drop-out distributions can be used in the **gestate** package as known models (since the package cannot estimate them from data): generalized Gamma, Gompertz, log-logistic, mixture exponential, mixture Weibull, piecewise exponential; (c) The **eventPred** package does not account for the uncertainty from model estimation.

Newton-Raphson (Dupuy, 2006; Li et al., 2013). For a PWE model with multiple change-points, Goodman et al. (2011) maximum (profile) likelihood estimation with the Nelder-Mead Simplex optimization algorithm is applicable. However, since the log-likelihood function is highly non-smooth, the recommended method heavily depends on the initial values and is often very slow and unstable in practice. There are also several Bayesian methods available, including stochastic approximation Monte Carlo algorithm(Kim et al., 2020), reversible jump MCMC method and some variants (Chapple et al., 2020; Cooney and White, 2021a).

The proposed **PWEXP** offers several ways to estimate PWE models ranging from a single change-point to multiple change-points, with the number and positions of change-points know or not known. Compared with available packages on CRAN for PWE model estimation (Frumento, 2021; Hess and Robert, 2021; Rufibach, 2022; Cooney and White, 2021b), **PWEXP** stands out among these packages by goodness-of-fit, robust estimation of varied survival curves, and offering a comprehensive toolset (Table 1).

Another primary focus of **PWEXP** is the event/timeline prediction in clinical trials. In event-driven clinical trials, accurately predicting future analysis time or events by using interim data is crucial for resource allocation, informed decision-making, and effective program development. Following the classic event prediction framework (Bagiella and Heitjan, 2001; Ying and Heitjan, 2008), at the current calendar time, t_0 , we are interested in predicting the conditional expectation of the number of events $ED(t_0, t')$ that will have been observed by time t' ($t' > t_0$): $ED(t_0, t) = D(t_0) + Q(t_0, t') + R(t_0, t')$, where $D(t_0)$ is the total number of subjects with events by time t_0 ; $Q(t_0, t')$ is the expected number of subjects who are in the study with no events by time t_0 and will have events by time t'; and $R(t_0, t')$ is the expected number of subjects who are in the study who will enroll and have events between times t_0 and t'. If the trial has finished enrollment by t_0 , then $R(t_0, t') = 0$. Among the three components, $D(t_0)$ is always known; $Q(t_0, t')$ and $R(t_0, t')$ requires modeling

on the distribution of events and loss to follow-up (drop-out) (e.g., using PWE models). For $R(t_0, t')$, an additional accrual prediction model is also required. Table 2 lists existing **R** packages for event/timeline prediction. **PWEXP** package offers both event and timeline prediction along with confidence intervals. The **eventPred** can also take PWE model as event and drop-out model, but it cannot estimate a model without known change-points.

In the remainder of this article, Section 2 introduces the theories and estimation approaches of PWE model, Section 3 focuses on the event prediction framework, Section 4 demonstrates the **PWEXP** package, followed with Section 5 presents a general analysis workflow of event prediction by using simulated data, Section 6 illustrates how to use **PWEXP** for study design with a real data example, and the last section is for summary and conclusion.

2 Piecewise Exponential (PWE) Model

2.1 PWE Distribution and Survival Model

This section begins with a brief introduction to the Piecewise Exponential (PWE) distribution, followed by an overview of the PWE model for right censoring data.

When the hazard rates of a random variable T is piecewise constant with a total of r change-points d_k $(1 \le k \le r)$:

$$h(t) = \begin{cases} \lambda_1, & t < d_1 \\ \lambda_2, & d_1 \le t < d_2 \\ \dots & \\ \lambda_{r+1}, & t \ge d_r \end{cases}$$

,

then T follows a piecewise exponential (PWE) distribution.

The density function and survival function (1-cumulative density function) of T are given by:

$$f(t) = h(t)e^{-H(t)} = \begin{cases} \lambda_1 e^{-\lambda_1 t}, & t < d_1 \\ \lambda_2 e^{(\lambda_2 - \lambda_1)d_1 - \lambda_2 t}, & d_1 \le t < d_2 \\ \dots \\ \lambda_{r+1} e^{\left[\sum_{i=1}^r (\lambda_{i+1} - \lambda_i)d_i\right] - \lambda_{r+1} t}, & t \ge d_r \end{cases}$$
$$S(t) = e^{-H(t)} = \begin{cases} e^{-\lambda_1 t}, & t < d_1 \\ e^{(\lambda_2 - \lambda_1)d_1 - \lambda_2 t}, & d_1 \le t < d_2 \\ \dots \\ e^{\left[\sum_{i=1}^r (\lambda_{i+1} - \lambda_i)d_i\right] - \lambda_{r+1} t}, & t \ge d_r. \end{cases}$$

The dpwexp() and ppwexp() functions in the **PWEXP** package can be used to calculate these values.

A PWE random variable T can be generated with inverse transform method, which relies on the quantile

function Q(p):

$$Q(p) = \begin{cases} \frac{-\log(1-p)}{\lambda_1} & p < 1 - e^{-\lambda d_1} \\ \frac{(\lambda_2 - \lambda_1)d_1 - \log(1-p)}{\lambda_2} & 1 - e^{-\lambda d_1} \le p < 1 - e^{(\lambda_2 - \lambda_1)d_1 - \lambda_2 d_2} \\ \dots \\ \frac{\left[\sum_{i=1}^r (\lambda_{i+1} - \lambda_i)d_i\right] - \log(1-p)}{\lambda_{r+1}} & p \ge 1 - e^{\left[\sum_{i=1}^r (\lambda_{i+1} - \lambda_i)d_i\right] - \lambda_{r+1} d_r}. \end{cases}$$

The qpwexp() function can calculate quantile values and rpwexp() function can generate PWE random variables from given parameters.

From a survival clinical trial, we observe *n* pair of iid random variables $(T_1, \delta_1), (T_2, \delta_2), \dots, (T_n, \delta_n)$, where δ_i indicates whether PWE distributed X_i (i.e. an event of interest such as death) is observed ($\delta_i = 1$) or not (non-informative right censored, $\delta_i = 0$). Following Klein et al. (2003, Chpater 3.5), the log-likelihood of the dataset can be constructed as:

$$\begin{split} \log(L) &= \sum_{j \in D_1} \left[\log(\lambda_1) - \lambda_1 t_j \right] + \sum_{j \in C_1} \left[-\lambda_1 t_j \right] \\ &+ \sum_{j \in D_2} \left[\log(\lambda_2) + (\lambda_2 - \lambda_1) d_1 - \lambda_2 t_j \right] + \sum_{j \in C_2} \left[(\lambda_2 - \lambda_1) d_1 - \lambda_2 t_j \right] \\ &+ \cdots \\ &+ \sum_{j \in D_{r+1}} \left\{ \log(\lambda_{r+1}) + \sum_{i=1}^r \left[(\lambda_{i+1} - \lambda_i) d_i \right] - \lambda_{r+1} t_j \right\} + \sum_{j \in C_{r+1}} \left\{ \sum_{i=1}^r \left[(\lambda_{i+1} - \lambda_i) d_i \right] - \lambda_{r+1} t_j \right\} \end{split}$$

where D_k or C_k represent the index set of event times or censoring times that fall into kth piece of the PWE distribution (i.e., $D_k = \{i \mid \delta_i = 1, d_{k-1} \leq T_i < d_k\}$, $C_k = \{i \mid \delta_i = 0, d_{k-1} \leq T_i < d_k\}$; here we let $d_0 = 0, d_{k+1} = +\infty$ for simpler notation). Obviously, the log-likelihood function $\log(L)$ is not smooth at $d_k = T_i$, resulting in maximum likelihood estimation infeasible using traditional optimization methods such as Newton-Raphson.

In the subsequent sections, we will first introduce the estimation of hazard rates when the positions of change-points are known (Section 2.2). Then we will move on to cases where we need to estimate both hazard rates and change-points without prior knowledge of positions of change-points (Section 2.3). Finally, we will cover the estimation of hazard rates and change-points when partial change-points are known (Section 2.4).

2.2 Parameter Estimation with Known Change-points

In this section, we assume all change-points $d_1, d_2, \dots d_r$ are already known. The log-likelihood function $\log(L)$ becomes smooth with respect to parameter parameter λ , allowing us to take a derivative of $\log(L)$

wrt λ :

$$\begin{aligned} \frac{\partial \log(L)}{\partial \lambda_1} &= \frac{n_{D_1}}{\lambda_1} - \sum_{j \in C_1} t_j - n_{2+} d_1 \\ \frac{\partial \log(L)}{\partial \lambda_2} &= \frac{n_{D_2}}{\lambda_2} - \sum_{j \in D_2, C_2} (t_j - d_1) - n_{3+} (d_2 - d_1) \\ & \cdots \\ \frac{\partial \log(L)}{\partial \lambda_r} &= \frac{n_{D_r}}{\lambda_r} - \sum_{j \in D_r, C_r} (t_j - d_{r-1}) - n_{r+1+} (d_r - d_{r-1}) \\ \frac{\partial \log(L)}{\partial \lambda_{r+1}} &= \frac{n_{D_{r+1}}}{\lambda_{r+1}} - \sum_{j \in D_{r+1}, C_{r+1}} (t_j - d_r) \end{aligned}$$

where n_{D_k} is the number of events that fall into the kth piece (i.e., $n_{D_k} = |D_k|$); n_{k^+} is the number of events and censoring that fall into the kth to the last pieces (i.e., $n_{k^+} = \sum_{l=k}^{r+1} |D_l| + |C_l|$).

Setting all derivatives equal to 0, we obtain the maximum likelihood estimators of hazard rates:

$$\begin{split} \hat{\lambda}_1 = & \frac{n_{D_1}}{\sum_{j \in C_1} t_j + n_{2^+} d_1} \\ \hat{\lambda}_2 = & \frac{n_{D_2}}{\sum_{j \in D_2, C_2} (t_j - d_1) + n_{3^+} (d_2 - d_1)} \\ & \cdots \\ \hat{\lambda}_r = & \frac{n_{D_r}}{\sum_{j \in D_r, C_r} (t_j - d_{r-1}) + n_{r+1^+} (d_r - d_{r-1})} \\ \hat{\lambda}_{r+1} = & \frac{n_{D_{r+1}}}{\sum_{j \in D_{r+1}, C_{r+1}} (t_j - d_r)}. \end{split}$$

2.3 Parameter Estimation with Unknown Change-points

The log-likelihood function $\log(L)$ is differentiable with respect to d_k as long as d_k is not equal to any event time or censoring time $(d_k \neq T_i)$. Under this assumption, we take the derivative of $\log(L)$ wrt d_k :

$$\frac{\partial \log(L)}{\partial d_k} = \sum_{j \in D_{k+1}} [\lambda_{k+1} - \lambda_k] + \sum_{j \in C_{k+1}} [\lambda_{k+1} - \lambda_k]$$
$$= n_{k+1} + (\lambda_{k+1} - \lambda_k).$$

Since $\lambda_{k+1} \neq \lambda_k$, then the expression above cannot be zero when d_k falls between any two consecutive sample time points. This fact implies that only when d_k is equal to any of the sample values T_i , the log-likelihood function achieves the maximum value (Loubert, 1986). Based on these findings, we propose three estimation approaches in the following sections.

2.3.1 MLE by Brute-force Search

Inspired by Qian and Zhang (2013)'s least squared approach with grid search, we propose employing a brute-force search to calculate the log-likelihood values for all potential change-point combinations. For each

Algorithm 1 Brute-force Search

procedure BFS(times T, censoring status C, number of change-points *nbreak*, max number of combinations max_set) $T_sub \leftarrow sub_sample(T, nbreak, max_set)$ candidate $\leftarrow combn(T_sub, nbreak)$ $candidate \leftarrow candidate[, sample(ncol(candicate, max_set))]$ for $chq_pt = each column in candidate do$ estimate hazard rates by the formula in Section 2.2 if any hazard rate is 0 then skip to next end if calculate log-likelihood L by the formula in Section 2.1 end for change-points chg_pt and hazard rates with largest L is the MLE end procedure function SUB_SAMPLE(T, nbreak, max_set) initialize $nl \leftarrow 1$ $nr \leftarrow length(T)$ repeat $NN \leftarrow choose(floor((nl+nr)/2), nbreak)$ if $NN > max_set$ then $nr \leftarrow floor((nl+nr)/2)$

else $nl \leftarrow floor((nl + nr)/2)$ end if until nr - nl < 1.5return $T_sub \leftarrow sample(T, nr)$ end function

change-point combination candidate, hazard rates can be estimated using the formula outlined in Section 2.2 and thus the log-likelihood can be obtained. The change-points combination and hazard rates with the largest log-likelihood value are selected as the maximum likelihood estimator. The full brute-force search method guarantees to find the optimal solution, but it comes with a high computational burden.

Especially, when the number of samples or the number of change-points are relatively large, the combination of change-points becomes prohibitively large for a brute-force search to be feasible. In such instances, we draw random sub-samples first and then conduct an exhaustive search based on these sub-samples. The 'max_set' argument in the pwexp.fit() function controls the maximum number of combinations to be explored. We summarize the entire procedure in pseudo-algorithm 1. This brute-force search with a subsampling method reduces computational intensity but introduces a risk of missing the optimal solution due to the sub-sampling process (i.e., the randomly selected sub-samples missed or do not cover the 'optimal' change-points).

2.3.2 OLS on Survival Function

The second method for estimating change-points relies on the survival curve. By taking a logarithmic transformation of the PWE survival function, we obtain a piecewise linear function:

$$\log(S(t)) = \begin{cases} -\lambda_1 t, & t < d_1 \\ (\lambda_2 - \lambda_1)d_1 - \lambda_2 t, & d_1 \le t < d_2 \\ \cdots \\ [\sum_{i=1}^r (\lambda_{i+1} - \lambda_i)d_i] - \lambda_{r+1} t, & t \ge d_r \\ = -\lambda_1 t + (\lambda_1 - \lambda_2)(t - d_1)_+ \cdots + (\lambda_r - \lambda_{r+1})(t - d_r)_+. \end{cases}$$

Following Küchenhoff (1996) and Muggeo (2003), we can fit a piecewise linear regression on log survival function $\log(S(t))$ with the **segmented** package and obtain the OLS estimation of change-points. Once the change-points are determined, hazard rates can be estimated using the formula described in Section 2.2. We summarize the entire procedure in pseudo-algorithm 2. Note that the OLS method does not obtain maximum likelihood estimators.

Algorithm 2 OLS on Survival Function
procedure $OLS(times T, censoring status C, number of change-points nbreak)$
$S \leftarrow survfit(surv(T,C) \sim 1)$
$chg_pt \leftarrow segmented(lm(log(S) \sim T))$
estimate hazard rates by the formula in Section 2.2
calculate log-likelihood L by the formula in Section 2.1
end procedure

2.3.3 Hybrid Method

The change-points estimated in the OLS method are typically close to, but not exactly, the maximum likelihood estimators. To obtain the true maximum likelihood estimator, we propose a hybrid method that combines the brute-force search with the OLS method.

Specifically, instead of randomly drawing sub-samples in brute-force search, we draw sub-samples from the values near the estimated change-points from the OLS method. For example, we may consider samples within some percentage of the confidence intervals around the change-points. Subsequently, we conduct an exhaustive search based on these sub-samples. The 'max_set' argument in the pwexp.fit() function controls the maximum number of combinations to be explored. This hybrid method is significantly more efficient and has a high probability of finding the optimal solution in practice. Based on our experience, this approach fits most datasets well and overcomes previous limitations of both Brute-force Search and OLS methods. The entire procedure of hybrid method is summarized in pseudo-algorithm 3.

2.4 Parameter Estimation with Partial Known Change-points

The proposed **PWEXP** package also accepts partial known change-points when some of the change-points are specified by the 'breakpoint' argument in the pwexp.fit() function and the total number of change-points

Algorithm 3 Hybrid Method

```
procedure HB(times T, censoring status C, number of change-points nbreak, max number of combina-
tions max\_set)
   S \leftarrow survfit(surv(T, C) \sim 1)
   chg with SE \leftarrow segmented(lm(log(S) \sim T))
   chg\_candidate_1 \leftarrow T \text{ in } 95\% \text{ CI of } chg_1
   chg\_candidate_{nbreak} \leftarrow T \text{ in } 95\% \text{ CI of } chg_{nbreak}
   candidate \leftarrow expand.grid(chg\_candidate_1, \cdots, chg\_candidate_{nbreak})
   candidate \leftarrow candidate[sample(nrow(candicate, max\_set)),]
   for chq_pt = each row in candidate do
       estimate hazard rates by the formula in Section 2.2
       if any hazard rate is 0 then
           skip to next
       end if
       calculate log-likelihood L by the formula in Section 2.1
   end for
   change-points chg_pt and hazard rates with largest L are returned
end procedure
```

('nbreak') is larger than the length of 'breakpoint'. This feature is particularly useful when the user wants to manually include certain change-points in the fitted model while retaining the flexibility of the PWE model. For example, if the survival curve is expected to change at the start of a treatment switch, we can manually specify a change-point at this time point. To achieve this:

- For the brute-force search method, the user will conduct an exhaustive search on the combinations of 'length(breakpoint)' pre-specified change-points along with ('nbreak length(breakpoint)') candidate change-points from data to obtain the maximum likelihood estimator.
- For the OLS on survival function method, when fitting a piecewise linear regression on log survival function log(S(t)) with the segmented() function from the **segmented** package, the user can set the 'npsi' argument to the number of unknown change-points ('npsi = nbreak length(breakpoint)') and the 'fixed.psi' argument to the pre-specified change-points ('fixed.psi = breakpoint'). Then the fitted result will have a total of 'nbreak' change-points including all pre-specified change-points.
- For the hybrid method, the user first follows the approach used in the OLS on survival function method to identify the remaining ('nbreak length(breakpoint)') change-points. Subsequently, the user can fix the known change-points (specified in 'breakpoint') and draw sub-samples from the values near the estimated change-points from the OLS method. Finally, the user can conduct an exhaustive search based on these sub-samples to obtain the maximum likelihood estimator.

2.5 Parameter Estimation Summary

The pwexp.fit() function is designed for PWE model estimation. Figure 1 illustrates the process for estimating a PWE model: The pwexp.fit() function first validates the 'breakpoint' argument (if not 'NULL'). Any change-points that are too early or too late such that there are no events before or after them are deleted with a warning. If two adjacent change-points are too close to each other such that there are no events

between them, they are combined by taking their average. Subsequently, based on the 'optimizer' argument, the function performs parameter estimation as described in the preceding sections.



Figure 1: Diagram of the process for estimating a PWE model using the pwexp.fit() function

To quantify the uncertainty of estimated model parameters, we can employ the boot.pwexp.fit() function to conduct bootstrapping. This technique involves resampling from the original dataset with replacement to generate multiple bootstrap samples. For each resampled dataset, we fit the PWE model and record the parameter estimates. By analyzing the distribution of parameter estimates obtained from the bootstrap samples, we can construct confidence intervals to capture the uncertainty associated with the model parameters.

2.6 Tail Robustness

In a real-world dataset, some 'abnormal' patterns such as sudden drops may occur in the tail of a survival curve. For example, in Figure 2, the gray Kaplan-Meier curve exhibits a sudden drop at t = 32. If this abrupt change is not properly addressed and a piecewise exponential model is fitted directly to the data, it will result in the blue curve, which over-estimates the hazard rate in the tail.

To mitigate this issue, several strategies can be employed:

1. One strategy is to ensure that a sufficient number of events are used in estimating the hazard rate for the tail of the survival curve. This can be achieved by setting a minimum threshold for the number of events required in the estimation of the last exponential piece. The 'min_pt_tail' argument (default:



Figure 2: Survival function of an example dataset. This is an illustration of different approaches to adjust tail.

'min_pt_tail = 5') in the pwexp.fit() function is used to specify this minimum number. For example, setting 'min_pt_tail = 10', we will have at least 10 events to be used to estimate the tail, resulting in a more robust estimation demonstrated by the orange curve in Figure 2.

- 2. Another strategy is to constrain the position of the last change-point to prevent it from occurring too late. This actually increases the number of events used in estimating the last exponential piece as well. The 'exclude_int' argument in the pwexp.fit() function can be used to exclude an interval from having any change-points. For example, by setting 'exclude_int = c(23, Inf)', we exclude any change-points after time t = 23, leading to the purple curve in Figure 2.
- 3. The third strategy involves cutting (re-censoring) the data prior to the occurrence of the sudden drop by using the cut_dat() function. Subsequently, a PWE model is fitted to the truncated data. For example, here we re-censor the data at t = 32 and red curve is the fitted model.

These strategies offer practical solutions to address 'abnormal' tails in survival data and enhance the reliability of hazard rate estimations.

2.7 Determination of Change-points Numbers

In the preceding sections, the number of change-points is assumed to be predetermined. However, in practice, the optimal number of change-points is usually unknown. One common approach is to plot the KM curve and fit a series of PWE models, starting from 0 change-points and gradually increasing the number of change-points. By performing a virtual comparison between the KM curve and the fitted PWE models, to ascertain the adequacy of the number of change-points in the PWE model for accurately fitting the KM curve.

Furthermore, statistical criteria such as Akaike Information Criterion (AIC) (from the pwexp.fit() function), Bayesian Information Criterion (BIC) (from the pwexp.fit() function), and cross-validation loglikelihood (from the cv.pwexp.fit() function) can assist in determining the optimal number of change-points. These statistical metrics provide a more objective basis for assessing 'goodness-of-fit' of the model, and help to select the most appropriate model configuration.

3 Event Prediction with PWE

In this section, we will discuss the process of conducting event prediction using the PWE model. Prior to delving into the details, we need some preliminary knowledge outlined in Section 3.1. Following this, we will elaborate on event prediction methodologies.

3.1 Conditional PWE Distribution

Suppose random variable T follows a PWE distribution as described in Section 2.1, then the conditional survival function (1-cumulative density function) and cumulative distribution function of T given T > R is given by

$$\begin{split} S(t|t>R) &= \frac{S(t)}{S(R)}, \quad \text{where } S(t), S(R) \text{ as defined in Section 2.1} \\ F(t|t>R) &= 1 - \frac{S(t)}{S(R)}, \quad \text{where } S(t), S(R) \text{ as defined in Section 2.1.} \end{split}$$

The conditional_ppwexp() function in the **PWEXP** package can be used to calculate these values.

The corresponding conditional PWE random variable can be generated with inverse transform method, which relies on the conditional quantile function Q(p|t > R):

$$Q(p|t > R) = \begin{cases} \frac{\lambda_1 R - \log(1-p)}{\lambda_1}, & p < F(d_1|t > R), R < d_1\\ \frac{\left[\sum_{i=1}^{k-1} (\lambda_{i+1} - \lambda_i) d_i\right] + \lambda_1 R - \log(1-p)}{\lambda_k}, & F(d_{k-1}|t > R) \le p < F(d_k|t > R), R < d_1\\ \\ \cdots\\ \frac{\left[\sum_{i=m}^{k-1} (\lambda_{i+1} - \lambda_i) d_i\right] + \lambda_m R - \log(1-p)}{\lambda_k}, & F(d_{k-1}|t > R) \le p < F(d_k|t > R), d_{m-1} \le R < d_m \end{cases}$$

The conditional_qpwexp() function can calculate conditional quantile values and the conditional_rpwexp() function can generate the conditional PWE random variables from given parameters.

3.2 Event Prediction

As discussed in Section 1, following the classic event prediction framework (Bagiella and Heitjan, 2001; Ying and Heitjan, 2008), at the current calendar time, t_0 , the conditional expectation of the number of events $ED(t_0, t')$ by time t' ($t' > t_0$) is:

$$ED(t_0, t') = D(t_0) + Q(t_0, t') + R(t_0, t'),$$

where $D(t_0)$ is the observed events by t_0 ; $Q(t_0, t')$ is the expected number of subjects who are in the study with no events by time t_0 and will have events by t'; and $R(t_0, t')$ is the expected number of subjects who will enroll and have events between t_0 and t'.

Assume that the PWE distributed event times T_i are independent, and that the PWE distributed censoring times C_i are also independent of each other and of the event times.

Suppose R is the index set of the subjects who will enroll at $U_i > t_0$. For an individual subject, the probability of observing an event before t' is:

$$P_i^R = \Pr(Y_i + U_i \le t', T_i < C_i) = \Pr(\min(T_i, C_i) + U_i \le t', T_i < C_i).$$

Define random variable $V_l = I(\min(T_l, C_l) + U_i \le t, T_l < C_l)$, where T_l and C_l can be generated as described in Section 2.1 using the rpwexp() function. Then $R(t_0, t')$ can be approximately simulated as

$$R(t_0, t') = \sum_{i \in R} P_i^R \approx \sum_{i \in R} \left\{ \frac{1}{N} \sum_{l=1}^N V_l \right\},\,$$

where N is the number of simulation iterations controlled by the 'n_each' argument in the predict() function of the **PWEXP** package.

Suppose Q is the index set of the subjects who are in the study with no events at time t_0 . For an individual subject, the probability of observing an event before t' is:

$$P_i^Q = \Pr(Y_i + U_i \le t', T_i < C_i | Y_i + U_i > t_0)$$

= $\Pr(\min(T_i, C_i) + U_i \le t', T_i < C_i | \min(T_i, C_i) + U_i > t_0)$
= $\Pr(\min(T_i, C_i) + U_i \le t', T_i < C_i | T_i + U_i > t_0, C_i + U_i > t_0).$

Define $W_l = I(\min(T_l, C_l) + U_i \le t', T_l < C_l)|T_l > t_0 - U_i, C_l > t_0 - U_i$, where the conditional T_l and C_l can be generated as described in Section 3.1 using the conditional_rpwexp() function. Then $Q(t_0, t')$ can be approximately simulated as

$$Q(t_0, t') = \sum_{i \in Q} P_i^Q \approx \sum_{i \in Q} \left\{ \frac{1}{N} \sum_{l=1}^N W_l \right\},$$

where N is the number of simulation iterations controlled by the 'n_each' argument in the predict() function.

Bringing all these elements together, the expectation of the number of events $ED(t_0, t')$ can be simulated as

$$ED(t_0, t') \approx D(t_0) + \sum_{i \in Q} \left\{ \frac{1}{N} \sum_{l=1}^N W_l \right\} + \sum_{i \in R} \left\{ \frac{1}{N} \sum_{l=1}^N V_l \right\}.$$

Similarly, we can define the predictive number of events $PED(t_0, t')$ as the result of a single generation:

$$PED(t_0, t') = D(t_0) + \sum_{i \in R} V_i + \sum_{i \in Q} W_i.$$

Both $ED(t_0, t')$ and $PED(t_0, t')$ can be estimated using the plot_event() function.

3.3 Confidence Interval

We will adopt percentile bootstrap to compute the confidence interval of the expected number of events $ED(t_0, t')$. The procedure comprises several steps. First we perform case resampling (with replacement) to obtain 'nsim' samples. Then we conduct 'nsim' PWE model estimations to get 'nsim' sets of parameters. For each set of parameters, we estimate the bootstrapped $ED(t_0, t')^*$ following the method outlined in the preceding section. Finally,

$$\left(ED(t_0, t')^*_{(\alpha/2)}, ED(t_0, t')^*_{(1-\alpha/2)}\right)$$

represents the $1 - \alpha/2$ percentile of the bootstrapped $ED(t_0, t')^*$.

For a predictive interval of the number of events $PED(t_0, t')$, we will follow a similar procedure. We conduct case resampling (with replacement) to obtain 'nsim' samples and conduct PWE model estimations to obtain 'nsim' sets of parameters. For each set of parameters, we repeatedly generate N (the 'n_each' argument in the predict() function) predictive number of event $PED(t_0, t')^*$ as described in the last section. Finally,

$$\left(PED(t_0, t')^*_{(\alpha/2)}, PED(t_0, t')^*_{(1-\alpha/2)}\right)$$

denotes the $1 - \alpha/2$ percentile of the bootstrapped $PED(t_0, t')^*$. The predictive interval accounts for both the uncertainty stemming from model estimation and the variability of event occurrence. As a result, it is always wider than the confidence interval of the expected number of events $ED(t_0, t')$. The plot_event() function can provide both types of confidence intervals.

4 Description of the PWEXP Package

4.1 Installation

The **PWEXP** package is developed in **R** and available from CRAN at https://CRAN.R-project.org/package=PWEXP. Hence, it can be installed in any computer with **R** with the command:

R> install.packages("PWEXP")

The development version of the package can be installed from GitHub using **devtools** with the following command:

R> devtools::install_github("zjph602xtc/PWEXP")

These commands also import **fastmatch** package and **segmented** package. Packages **parallel**, **doSNOW** and **foreach** are needed if multi-core calculation is enabled for bootstrapping and cross-validation. Several vignettes have been prepared using **knitr**. An online package tutorial is available at https://zjph602xtc.github.io/PWEXP/.

4.2 Software Architecture and Main Functions

	Simulate Data	Split Data	Fit PWE Model & Model Selection	Bootstrap CI	Prediction		
Description	 simulate TTE data with censoring and death multiple arms and stratifications custom distributions for TTE, censor, death 	 split dataset into training and testing sets recensor training data at the cut off time 	 fit piece exponential model with fixed, partial fixed or unknown breakpoints obtain AIC, BIC, CV likelihood 	 use bootstrap to obtain the CI of estimated parameters 	 predict timelines or number of events obtain the Cl of predicted results 		
Main function	simdata()	cut_dat()	<pre>pwexp.fit() cv.pwexp.fit()</pre>	<pre>boot.pwexp.fit()</pre>	<pre>predict()</pre>		
Visualization	<pre>plot_survival() plot_event()</pre>		<pre>plot_survival()</pre>	<pre>plot_survival()</pre>	<pre>plot_event()</pre>		
Auxiliary function	Study design/plan:	sim_ioiiowup()	pweyn() dpweyn() :	rnweyn()			
Auxiliary function	<pre>Piecewise exponential dist: dpwexp(), ppwexp(), qpwexp(), rpwexp() Conditional piecewise exp dist: ppwexp_conditional(), qpwexp_conditional(), rpwexp_conditional()</pre>						

Figure 3: Main functions and structure of the **PWEXP** package

The **PWEXP** package follows a software architecture based on functional programming. Each main function returns an instance of an 'S3' class, which implements relevant 'S3' methods to facilitate downstream analyses. The **PWEXP** package offers a comprehensive suite of tools tailored for analyzing survival data with a piecewise exponential distribution. It facilitates various tasks related to survival analysis and event prediction based on the fitted model. Specifically, the package comprises several components designed for different tasks as shown in Figure 3:

• Survival data simulation

- the simdata() function generates simulated survival dataset with:
 - * randomization/enrollment time defined by randomization/enrollment rate (number of subjects per month) or randomization/enrollment curve (number of subjects in each month);
 - * multiple treatment groups with stratification by user-defined allocation ratio;
 - * primary endpoint (event), drop-out and death with exponential distribution or user-defined distributions (e.g., piecewise exponential, mixture distribution, etc.) for each stratification in each group.
- the plot_survival() function visualizes KM estimator for right censored data;
- the plot_event() function plots the number of cumulative events.
- Data cut-off

- the cut_dat() function truncates the data by a clinical cut-off date (DCOD), only retaining subjects randomized before this time and re-censoring the data. It is useful to create a training data set.

- PWE model estimation
 - the pwexp.fit() function fits the piecewise exponential model to right censored data with:
 - * pre-specified change-points;
 - * partially pre-specified change-points;
 - * unknown change-points (to be fitted from the data).

- the pwexp.fit() function also calculates model AIC, BIC;
- the cv.pwexp.fit() conducts model cross-validation and obtains the CV log-likelihood;
- the plot_survival() plots the fitted survival curve for 'pwexp.fit' object.

• Model bootstrapping

- the boot.pwexp.fit() conducts bootstrapping for an existing model;
- the plot_survival() plots fitted survival curve and its confidence interval for 'boot.pwexp.fit' object.
- Events/timeline prediction
 - the predict() function predicts the number of events and timeline based on the event and drop-out model (optional);
 - the predict() function provides the CI of the predicted number of events or timeline for a bootstrapping model;
 - the plot_event() plots predicted number of events (for 'predict.pwexp.fit' object) and its CI (if applicable, for 'predict.boot.pwexp.fit' object).
- Auxiliary functions
 - the auxiliary function sim_followup() estimates follow-up time and number of events at design stage;
 - the auxiliary functions dpwexp(), ppwexp(), qpwexp(), rpwexp() are the PDF, CDF, quantile function, generator function for PWE distribution;
 - the auxiliary functions ppwexp_conditional(), qpwexp_conditional(), rpwexp_conditional() are the CDF,
 quantile function, generator function for conditional PWE distribution.

In the following sections, we will present two examples to illustrate the usage of the **PWEXP** package. The first example focuses on event prediction using simulated data, while the second example demonstrates a study design based on a real case study.

5 Event/timeline Prediction Example

In this section, we will present a comprehensive workflow for event and timeline prediction. The workflow comprises three steps. First, we simulate survival data and truncate the dataset at the time t_0 when 80% of the subjects are randomized. Subsequently, we fit a PWE model to the truncated dataset. Finally, we perform event and timeline prediction for time t' with $t' > t_0$ by using the estimators from the truncated dataset and compare the predicted events/timelines with the 'real' data from simulation.

5.1 Simulate PWE Data

Here we utilize the simdata() function to create a simple example dataset with the following characteristics:

- randomization rate is defined as 20 subjects per month ('rand_rate = 20'), and total sample size is 1000 ('total_sample = 1000');
- the primary endpoint (event) follows a PWE distribution with monthly hazard rates of 0.1 for t < 5 months, 0.01 for $5 \le t < 14$ months, 0.2 for $t \ge 14$ months (defined by 'myevent_dist' function);

- the drop-out follows an exponential distribution with a drop-out probability of 3%/month ('drop_rate = 0.03')(equivalently, a monthly drop-out hazard rate is $-\log(1 0.03) = 0.0304$);
- the 'add_column' argument requests additional variables to be included in the dataset.

```
R> myevent_dist <- function(n)rpwexp(n, rate = c(0.1, 0.01, 0.2),
+ breakpoint = c(5,14))
R> dat <- simdata(rand_rate = 20, total_sample = 1000, drop_rate = 0.03,
+ advanced_dist = list(event_dist = myevent_dist),
+ add_column = c('censor_reason', 'event', 'followT', 'followT_abs'))
R> head(dat)
```

	ID	randT	eventT	dropT	$\verb"censor_reason"$	event	followT	followT_abs
1	1	18.95190	0.9741066	47.025224	<na></na>	1	0.9741066	19.92601
2	2	18.38245	3.1487676	1.336703	drop_out	0	1.3367033	19.71915
3	3	38.76302	14.3753607	35.737477	<na></na>	1	14.3753607	53.13838
4	4	27.59118	3.2444541	12.549571	<na></na>	1	3.2444541	30.83563
5	5	48.42573	16.1987174	24.916275	<na></na>	1	16.1987174	64.62445
6	6	35.62735	14.1660874	62.585078	<na></na>	1	14.1660874	49.79344

In the generated dataset 'dat', all subjects are followed for infinite time and we will truncate it by a clinical cut-off date (DCOD) with the cut_dat() function in the next step. These variables are included in dataset 'dat' (See Appendix A for more details):

- 'randT' is the randomization time for each subject;
- 'eventT' and 'dropT' are the event time and drop-out time, respectively;
- 'followT' is the follow-up time, which is the minimum value of 'eventT', 'dropT'. In real-world datasets, this is the observation time;
- 'followT_abs' is the sum of 'randT' and 'followT';
- 'event' indicates whether the primary event occurred at the end of follow-up with 0 for censoring and 1 for occurrence of event. If a subject is censored, 'censor_reason' shows the type of censoring (i.e., 'drop_out', 'death' or 'never_event' (when 'followT == Inf')).

We can visualize the survival curve and cumulative number of events using the plot_survival(), plot_event() functions respectively. Figure 4(a) displays the KM curve of the simulated events. Notably, the curve drops faster before t = 5 and after t = 14 with hazard rates being 0.1 and 0.2, respectively, which are larger than the hazard rate of 0.01 in the middle time interval. Figure 4(b) is the cumulative number of events over time. It is important to note that the x-axis represents the time from randomization of each subject, allowing us to observe changes in the event rate due to the piecewise exponential distribution (by setting argument 'abs_time = F' in the the plot_event() function). If we were to plot the event curve from the beginning of the trial, the figure would be Figure 10.

```
R> plot_survival(dat$eventT, dat$event, conf.int = F, mark.time = T,
+ xlim = c(0, 25), main = 'Survival Function of the Simulation Data')
```

```
R> plot_event(dat$eventT, abs_time = F, dat$event, xlim = c(0, 25),
+ main = 'Number of Cumulative Events of the Simulation Data')
```



Figure 4: (a) KM curve for the simulated dataset; (b) Event curve for the simulated dataset.

To compare the prediction results with the ground truth, we need to cut the complete dataset 'dat' at the interim analysis time. For instance, we can cut the dataset using the time of the clinical cut-off date (DCOD) when 80% of subjects are randomized. This operation retains only data available by DCOD in the returned data 'train'. Additionally, the returned data will be re-censored at the DCOD.

```
R> cut <- quantile(dat$randT, probs = 0.8)
R> cut
   80%
39.99107
R> train <- cut_dat(cut = cut, data = dat, var_randT = 'randT',
                   var_followT = 'followT', var_followT_abs = 'followT_abs',
+
                   var_event = 'event', var_censor_reason = 'censor_reason')
+
R> head(train)
  ID
        randT
                  eventT
                              dropT censor_reason event
                                                           followT followT_abs
  1 18.95190
               0.9741066 47.025224
                                             <NA>
                                                       1 0.9741066
                                                                      19.92601
1
2
  2 18.38245
               3.1487676 1.336703
                                         drop_out
                                                       0 1.3367033
                                                                      19.71915
  3 38.76302 14.3753607 35.737477
                                                       0 1.2280545
                                                                      39.99107
3
                                              cut
```

4 27.59118 3.2444541 12.549571

4

<NA>

1 3.2444541

30.83563

6	6 35.62735	14.1660874 62.585078	cut	0 4.3637258	39.99107
7	7 13.92114	0.9511499 85.247486	<na></na>	1 0.9511499	14.87229

5.2 PWE Model Estimation

In this section, we aim to determine the optimal piecewise exponential model for the training dataset 'train'. Given that the position and number of change-points are unknown in practice, we will fit models 'fit_b0' to 'fit_b4' with 0 to 4 unknown change-points. The 'nbreak' argument is the total number of change-points. It is worth noting that the 'fit_b0' model is actually an exponential model since there are no change-points.

```
R> fit_b0 <- pwexp.fit(train$followT, train$event, nbreak = 0)
R> fit_b1 <- pwexp.fit(train$followT, train$event, nbreak = 1)
R> fit_b2 <- pwexp.fit(train$followT, train$event, nbreak = 2)
R> fit_b3 <- pwexp.fit(train$followT, train$event, nbreak = 3)
R> fit_b4 <- pwexp.fit(train$followT, train$event, nbreak = 4)</pre>
```

As indicated in Section 2.7, the initial step is to visualize the fitted curves alongside the real KM curve. Now let's proceed by plotting the fitted curves:

```
R> plot_survival(train$followT, train$event, xlim = c(0, 40),
+ main = 'Fitted Models with Different Number of Breakpoints')
R> plot_survival(fit_b0, col = 'green', lwd = 2)
R> plot_survival(fit_b1, col = 'blue', lwd = 3, show_breakpoint = F)
R> plot_survival(fit_b2, col = 'red', lwd = 3,
+ breakpoint_par = list(col = 'grey50', lty = 2))
R> plot_survival(fit_b3, col = 'orange', lwd = 3, show_breakpoint = F)
R> legend('topright', c('training data', '0 breakpoints', '1 breakpoint',
+ '2 breakpoints', '3 breakpoints'), lwd = 3,
+ col = c('black', 'green', 'blue', 'red', 'orange'))
```

Figure 5 shows the four fitted models. It is clear that the exponential model does not adequately capture the observed survival patterns. However, the PWE model with one change-point, located around t = 14, demonstrated a much closer alignment with the actual KM curve. Moreover, the PWE model with two change-points exhibits a perfect fit. Increasing the number of change-points beyond two does not yield significant improvement in the model fit.

Visualization method can be combined with statistical criteria such as AIC, BIC, and CV Log likelihood to aid in the process of model selection. For illustrative purposes, we fit two additional models:

- 1. 'fit_a0' with two pre-specified change-points at t = 5 and t = 14;
- 2. 'fit_a1' with one pre-specified change-point at t = 14, while the total number of change-points is set to 2. The position of the other change-point will be estimated from the training data.

```
R> fit_a0 <- pwexp.fit(train$followT, train$event, breakpoint = c(5,14))
R> fit_a1 <- pwexp.fit(train$followT, train$event, nbreak = 2, breakpoint = c(14))</pre>
```



Figure 5: Visualization of PWE models with different number of change-points

Note that models 'fit_a0' and 'fit_a1' are solely for demonstration purposes and will not be considered for subsequent analysis, as the actual positions of change-points will not be known in practice.

The BIC values for the six models are plotted in Figure 6. The model 'fit_a0' with two pre-specified true known change-points has the smallest BIC and serves as a baseline. Among the models without pre-specified change-points, we observe that 'fit_b2', which has the correct number of change-points, exhibits the lowest BIC value.

```
R> model_name <- c('2 fixed breakpoints', '1 fixed + 1 unknown bkp',
                   '0 unknown bkp (exp model)', '1 unknown bkp', '2 unknown bkp',
+
                   '3 unknown bkp', '4 unknown bkp')
+
  model_ind <- c("a0", "a1", "b0", "b1", "b2", "b3", "b4")</pre>
R>
R> col <- c("#FBB4AE", "#B3CDE3", "#C7E9C0", "#A1D99B", "#74C476", "#41AB5D", "#238B45")
  barplot(c(fit_a0$BIC, fit_a1$BIC, fit_b0$BIC, fit_b1$BIC, fit_b2$BIC,
R>
             fit_b3$BIC, fit_b4$BIC), xlab = 'Model',
+
          ylab = 'BIC', ylim = c(2700, 3400), xpd = F, col = col,
+
          names.arg = model_ind, main = 'BIC of Fitted Models')
R> legend('topleft', model_name, fill = col, ncol = 2, text.width = 3.5)
```

Cross-validation (CV) log-likelihood can be used to further assist in model selection. The cv.pwexp.fit() function conducts cross-validation either from an estimated PWE model or directly from the data. Additionally, the 'parallel' argument enables parallel computing to reduce the computation time.

R> fit_a0_cv <- cv.pwexp.fit(fit_a0, nsim = 100)</pre>



Figure 6: BIC of fitted PWE models

```
R> fit_a1_cv <- cv.pwexp.fit(fit_a1, nsim = 100)
R> fit_b0_cv <- cv.pwexp.fit(fit_b0, nsim = 100)
R> fit_b1_cv <- cv.pwexp.fit(fit_b1, nsim = 100)
R> fit_b2_cv <- cv.pwexp.fit(fit_b2, nsim = 100, parallel = TRUE, mc.core = 10)
R> fit_b3_cv <- cv.pwexp.fit(fit_b3, nsim = 100, parallel = TRUE, mc.core = 10)
R> fit_b4_cv <- cv.pwexp.fit(train$followT, train$event, nbreak = 4, nsim = 100)</pre>
```

The returned object 'cv.pwexp.fit' is a vector of 'nsim' CV log-likelihoods. We utilize a boxplot to visualize the CV log-likelihoods in Figure 7.

```
R> boxplot(cv ~ model_ind, data.frame(cv = c(fit_a0_cv, fit_a1_cv, fit_b0_cv, fit_b1_cv,
+ fit_b2_cv, fit_b3_cv, fit_b4_cv),
+ model_ind = rep(model_ind, each = 100)),
+ ylab = 'CV log likelihood', xlab = 'Model', ylim = c(-320,-250),
+ col = col, main = 'CV Log Likelihood of Fitted Models')
R> legend('topleft', model_name, fill = col, ncol = 2, text.width = 3)
```

Consistent with the BIC results, the model 'fit_a0' demonstrates the largest CV log-likelihood and serves as a baseline. Among the models without pre-specified change-points, 'fit_b2' continues to exhibit the largest CV log-likelihood.

Therefore, we select model 'fit_b2' as our final optimal model for downstream analysis. The model details are stored in the object attributes list 'attributes(fit_b2)'. Specifically, we can extract the estimated hazard rates and change-points using 'attr(fit_b2, 'lam')' and 'attr(fit_b2, 'brk')', respectively:



Figure 7: Cross validation log-likelihood of fitted PWE models

R> fit_b2

brk1	brk2	lam1	lam2	lam3	likelihood	AIC	BIC
5.077201	13.95023	0.08955893	0.009611296	0.1907673	-1393.633	2797.266	2820.689
R> attr(1	fit_b2, ']	Lam')					
[1] 0.089	9558930 0.	.009611296 (0.190767295				
R> attr(1	fit_b2, 't	ork')					
[1] 5.077	7201 13.95	50226					

We can utilize the boot.pwexp.fit() function to bootstrap this model and estimate the variance of the estimated parameters. The returned object includes a data frame with 'nsim' rows, each corresponding to estimated parameters in each bootstrapping sample. Similar to the 'pwexp.fit' object, the bootstrapping model details are also restored in the object attributes list 'attributes(fit_b2_boot)'. While bootstrapping the model is optional, it provides confidence intervals for event and timeline predictions, enhancing the reliability of our results. Figure 8 illustrates the fitted model with a 95% confidence interval.

R> fit_b2_boot <- boot.pwexp.fit(fit_b2, nsim = 100, parallel = TRUE, mc.core = 10)
R> plot_survival(train\$followT, train\$event, xlim = c(0,40),

+ main = 'Fitted Model for Primary Events with 95% CI')
R> plot_survival(fit_b2_boot, col = 'red', alpha = 0.05, CI_par = list(col = '#ff9896'))
R> brk_ci <- apply(attr(fit_b2_boot, 'brk'), 2, function(x)quantile(x, c(0.025, 0.975)))
R> abline(v = brk_ci, col = 'grey', lwd = 2, lty = 3)



Figure 8: Visualization of the final model with 95% CI

5.3 Event/timeline Prediction

As described in Section 3, for event/timeline prediction, a censoring model for drop-out and death (if present) can be estimated together with the event model. The censoring model is similar to the event model described in the previous section, but the response of the model is drop-out and death (if present) rather than event of interest. The censoring model is discretionary in scenarios where the drop-out (and death) rate is negligible. Nevertheless, in situations where the drop-out (and death) rates exhibit significance, constructing a censoring model becomes imperative to ensure precise event/timeline prediction.

Here, we first create a variable 'drop_indicator' to be a time-to-event endpoint that represents the time of drop-out (and we do not have death in the simulated data), and it will be used as the response variable in the censoring model. We fit an exponential censoring model with bootstrapping.

```
R> fit_censor_boot <- boot.pwexp.fit(train$followT, drop_indicator,
+ nbreak = 0, nsim = 100)
R> plot_survival(fit_censor_boot, col = 'red', alpha = 0.01,
+ CI_par = list(col='#ff9896'))
```



Figure 9: Visualization of the fitted censoring model for drop-out with 95% CI

Finally, we use the predict() function to predict the number of future events. The 'future_rand' argument defines the randomization curve of the remaining subjects. In this example, the future randomization rate remains 20 subjects per month ('rand_rate = 20'), and the total sample to be enrolled after the analysis time is the planned number of samples ('1000') minus the number of randomized subjects at the analysis time ('NROW(train)'). The returned object from predict() contains predicted event curve functions, which can be used with the plot_event() function to calculate the estimated events and plot the curve.

```
R> predicted_boot <- predict(fit_b2_boot, analysis_time = cut,
+ censor_model_boot = fit_censor_boot,
+ future_rand = list(rand_rate = 20, total_sample = 1000 - NROW(train)),
+ n_each = 30)
```

We use plot_event() function to plot the event curve with confidence interval ('type = 'confidence') and predictive interval ('type = 'predictive''):

```
R> plot_event(dat$followT_abs, abs_time = T, event = dat$event, ylim = c(0, 800),
+ main = 'Predicted Number of Events')
```



Time from first randomization

Figure 10: Predicted number of future events

```
R> plot_event(train$followT_abs, abs_time = T, event = train$event, add = T, col='blue')
R> pred_event_confidence <- plot_event(predicted_boot, eval_at = seq(45, 90, 5),</pre>
+
                                        type = 'confidence')
R> pred_event_predictive <- plot_event(predicted_boot, eval_at = seq(45, 90, 5),
                                      type = 'predictive', CI_par = list(col = 'purple'))
+
R> legend('bottomright', c('data used to train model', 'true event curve',
                            'predicted event curve', 'confidence interval',
+
                            'predictive interval'), lwd = 2,
+
                col=c('blue', 'black', 'red', '#ff9896', 'purple'), lty = c(1,1,1,2,2))
+
R> head(pred_event_confidence)
     time n_event 5% n_event 95% n_event
[1,]
       45 494.3127
                     489.2371
                                  500.9097
[2,]
       50 565.9164
                     557.6987
                                  577.3319
[3,]
       55 614.9518
                     604.7011
                                  627.8907
[4,]
       60 647.3800
                     637.2311
                                  660.7467
[5,]
       65 676.7368
                     666.0635
                                  689.8273
[6,]
       70 692.2490
                     680.6740
                                  705.5199
R> head(pred_event_predictive)
     time n_event 5% n_event 95% n_event
```



Cumulative events

Figure 11: Predicted timeline for given number of future events

[1,]	45 494.4746	482.0519	507.2512
[2,]	50 566.0482	549.3727	584.1406
[3,]	55 614.8935	595.8857	634.8421
[4,]	60 647.5956	628.1870	666.9981
[5,]	65 677.5232	657.7572	696.6753
[6,]	70 693.0376	673.1747	711.9456

The returned data frame contains the number of predicted events and its corresponding 95% confidence/predictive interval at the specified times (by the 'eval_at' argument in the plot_event() function).

By setting argument 'xyswitch = TRUE', we can use the $plot_event()$ function to obtain the timeline for a given number of events in the future.

```
R> plot_event(dat$followT_abs, xlim = c(0, 800), event = dat$event,
              xyswitch = T, main = 'Predicted Timeline')
+
R> plot_event(train$followT_abs, abs_time = T, event=train$event, add = T,
              xyswitch = T, col = 'blue')
+
R> pred_time_confidence <- plot_event(predicted_boot, xyswitch = T,</pre>
                                       eval_at = seq(500, 700, 50), type = 'confidence')
+
R> pred_time_predictive <- plot_event(predicted_boot, xyswitch = T,</pre>
                                       eval_at = seq(500, 700, 50), type = 'predictive',
+
                                       CI_par = list(col = 'purple'))
+
R> legend('bottomright', c('data used to train model', 'true timeline',
                            'predicted timeline', 'confidence interval',
+
```

```
+ 'predictive interval'), lwd = 2,
+ col = c('blue', 'black', 'red', '#ff9896', 'purple'), lty = c(1,1,1,2,2))
R> head(pred_time_confidence)
```

	$\texttt{n_event}$	time	5% time	95% time
[1,]	500	45.37890	44.94330	45.79609
[2,]	550	48.85392	48.17232	49.38727
[3,]	600	53.05977	51.75140	54.30394
[4,]	650	60.41130	58.20409	62.13599
[5,]	700	77.63385	67.49335	NA

R> head(pred_time_predictive)

	n_event	time	5% time	95% time
[1,]	500	45.37776	44.53543	46.30582
[2,]	550	48.84987	47.72773	50.05325
[3,]	600	53.06970	51.16706	55.60011
[4,]	650	60.40710	57.27906	63.61008
[5,]	700	78.28629	65.73887	NA

6 Study Design Example

In this section, we will demonstrate how to use the **PWEXP** package to assist in study design, such as sample size and study duration calculation, with one example.

Assume we are planning a phase 3 trial design with target population be patients newly diagnosed with diffuse large B-cell lymphoma and an international prognostic index (IPI) score of 4-5. The primary objective of the trial is to demonstrate the superiority of the experimental treatment compared to the control arm in terms of overall survival (OS) endpoint. Additional characteristics for the study design are listed as below:

- subjects are randomized in a 1:1 ratio;
- a two-sided log-rank test will be employed, ensuring 90% power at a significance level of 5% (or one-sided at 2.5%);
- the target hazard ratio for the primary endpoint OS, is set to 0.6;
- a monthly dropout probability of 1% is assumed;
- interim analyses are scheduled at 40% and 70% of total information;
- alpha-spending version of O'Brien-Fleming boundary for efficacy; no futility interim;
- a total of 660 subjects will be recruited;
- recruitment begins at a rate 15 pt/month during the first 12 months; subsequently, increase of the number of sites and ramp up of recruitment by +6 pt/month each month until a maximum of 45 pt/month.

6.1 Obtaining Model Assumption

The initial step for study design involves deriving assumptions for the OS rate in the control arm by analyzing available historical data or literature such as Ruppert et al. (2020). To estimate the OS rate for the target population, we digitize the OS curves for IPI 4-5 (red curve) from Figure 2A using WebPlotDigitizer (Rohatgi, 2024). Subsequently, we reconstruct individual subject data (IPD) using the **IPDfromKM** package (Liu et al., 2021). The resulting OS curve for IPI 4-5 patients is represented by the gray curve in Figure 12.



Figure 12: OS curve for IPI 4-5 subjects (Ruppert et al., 2020). Colored curves are fitted parametric models.

We explore various parametric models, including the exponential, Weibull, Gamma, and log-normal models, in an attempt to fit the data. However, these models prove inadequately capturing the KM curve. As a result, we consider employing a piecewise exponential model as a potential solution.

```
R> dat <- read.csv(paste0("https://raw.githubusercontent.com/zjph602xtc/PWEXP/",
                   "main/docs/ruppert%20ipd.csv"))
+
R> PWE <- pwexp.fit(dat$Survival.time, dat$Status, nbreak = 2)</pre>
R> PWE
brk1
       brk2
                lam1
                             lam2
                                         lam3 likelihood
                                                               AIC
                                                                        BIC
14.716 29.85 0.023956 0.009931584 0.004189957 -1065.844 2141.689 2161.984
R> PWEXP::ppwexp(12 * 1:4, rate = attr(PWE, 'lam'), breakpoint = attr(PWE, 'brk'),
          lower.tail = F)
+
```

[1] 0.7501575 0.6409900 0.5894241 0.5605208

The red curve in Figure 12 represents the fitted PWE model, which closely matches the actual KM curve. Based on the the fitted PWE model, the IPI 4-5 population group exhibits monthly hazard rates of 0.0240, 0.00993, and 0.00419 for the time intervals of $(0, \le 14.7 \text{ months})$, $(> 14.7 \text{ months}, \le 29.9 \text{ months})$, and (> 29.9 months), respectively. These hazard rates result in a PFS rate of 75.0%, 64.1%, 58.9%, and 56.1% at year 1, year 2, year 3, and year 4, respectively.

6.2 Integration with Design Package

Once OS rates were estimated by the fitted piecewise model 'PWE', they can be readily utilized in various clinical trial design packages. The model details are stored in the object attributes list 'attributes(PWE)'. Specifically, we can extract the estimated hazard rates and change-points using 'attr(PWE, 'lam')' and 'attr(PWE, 'brk')', respectively. As an illustration, we will demonstrate the integration with **rpact** package for study design:

```
R> design <- getDesignGroupSequential(
     sided = 1, alpha = 0.025, beta = 0.1,
+
     informationRates = c(0.4, 0.7, 1),
+
     typeOfDesign = "asOF")
+
R> accrualTime <- getAccrualTime(accrualTime = c(0, 12:16),
                                  accrualIntensity = c(15, 21, 27, 33, 39, 45),
+
                                  maxNumberOfSubjects = 660)
+
R> y <- getSampleSizeSurvival(</pre>
     design = design,
+
+
     lambda2 = attr(PWE, 'lam'),
     dropoutRate1 = 0.01, dropoutRate2 = 0.01, dropoutTime = 1,
+
     allocationRatioPlanned = 1,
+
     accrualTime = accrualTime,
+
+
     piecewiseSurvivalTime = c(0,attr(PWE,'brk')),
     hazardRatio = 0.6)
+
R> summary(y)
Sample size calculation for a survival endpoint
Sequential analysis with a maximum of 3 looks (group sequential design), overall
significance level 2.5% (one-sided).
The results were calculated for a two-sample logrank test,
HO: hazard ratio = 1,
H1: hazard ratio = 0.6, piecewise survival distribution,
piecewise survival time = c(0, 14.716, 29.85),
control lambda(2) = c(0.024, 0.01, 0.004), maximum number of subjects = 660,
accrual time = c(12, 13, 14, 15, 16, 24),
```

accrual intensity = c(15, 21, 27, 33, 39, 45), dropout rate(1) = 0.01, dropout rate(2) = 0.01, dropout time = 1, power 90%.

Stage	1	2	3
Information rate	40%	70%	100%
Efficacy boundary (z-value scale)	3.357	2.445	2.001
Overall power	0.0981	0.6139	0.9000
Number of subjects	536.2	660.0	660.0
Expected number of subjects under H1	647.8		
Cumulative number of events	65.3	114.3	163.4
Analysis time	21.248	27.089	35.146
Expected study duration	29.6		
Cumulative alpha spent	0.0004	0.0074	0.0250
One-sided local significance level	0.0004	0.0073	0.0227
Efficacy boundary (t)	0.436	0.633	0.731
Exit probability for efficacy (under HO)	0.0004	0.0070	
Exit probability for efficacy (under H1)	0.0981	0.5157	

Legend:

(t): treatment effect scale

Based on the results, we will conduct two interim analyses at 21.2 and 27.1 months, with the final analysis scheduled for 35.1 months to achieve an overall 90% power. A total of 65.3, 114.3, 163.4 cumulative events are expected to occur at each analysis time.

6.3 Event/timeline and Follow-up Simulation

The **PWEXP** package also offers simulation capabilities for study design purposes. The sim_followup() function can estimate the number of events at any specified time ('type = 'calendar"). Additionally, it provides summary statistics of the follow-up time. In the example below, the follow-up time is defined as the duration from randomization to censoring (either drop-out or end of the trial, whichever occurs first), specified by 'follow_up_endpoint = c('drop_out', 'cut')' (Betensky, 2015). We can define summary statistics for the follow-up time. For instance, here we define 'prop_12' as the proportion of subjects whose follow-up time is larger than 12 months. Moreover, we request statistics such as the 'mean', 'median', and 'sum' of the follow-up time.

+	drop_rate = 0.01, by_group = T,
+	n_rand = c(rep(15, 12), 21, 27, 33, 39, rep(45, 8)),
+	<pre>advanced_dist = list(event_dist = c(pfs_trt, pfs_con)),</pre>
+	<pre>rep = 1000, stat = c(mean, median, sum, prop_12),</pre>
+	<pre>follow_up_endpoint = c('cut', 'drop_out'))</pre>

\$T_all

	at	analysis_time	event	subjects	mean	median	sum	prop_12
1	21.248	21.248	65.422	536.18	7.376394	5.662857	3954.951	0.2292295
2	27.089	27.089	114.560	660.00	11.076918	9.670736	7310.766	0.3550924
3	35.146	35.146	163.726	660.00	17.951402	17.106201	11847.925	0.8340242

\$T_by_group

	at	group	$analysis_time$	event	subjects	mean	median	sum	prop_12
1	21.248	con	21.248	40.165	268.285	7.379129	5.666560	1979.673	0.2293759
4	21.248	trt	21.248	25.257	267.895	7.373329	5.662474	1975.278	0.2290497
2	27.089	con	27.089	69.915	330.000	11.082933	9.680618	3657.368	0.3553515
5	27.089	trt	27.089	44.645	330.000	11.070902	9.661505	3653.398	0.3548333
3	35.146	con	35.146	99.260	330.000	17.949422	17.114265	5923.309	0.8343364
6	35.146	trt	35.146	64.466	330.000	17.953382	17.106924	5924.616	0.8337121

The output displays the expected number of events at interim analyses (IAs) and final analysis (FA) as 65.4, 114.6, and 163.7, respectively. These figures closely align with the theoretical calculations performed using the **rpact** package.

If we plot the expected number of events according to theoretical calculations from the **rpact** package and simulation results from the **PWEXP** for each month in Figure 13, the simulation result is very accurate throughout the whole study duration. Furthermore, the simulation results provide summary statistics of follow-up for each arm at specified times. For example, at the time of the final analysis (35.1 months), 83.4% of subjects will have a follow-up duration longer than 1 year. This information is crucial for trial sponsors and health authorities in evaluating trial's conduct and performance.

The sim_followup() function can also estimate the timeline based on a specified number of randomized subjects or number of events ('type = 'sample" or 'type = 'event"). The following example demonstrates the analysis time for every 10% increase in information fraction from 10% to 100%:

\$T_all

	at	analysis_time	event	subjects	mean	median	sum
1	16	11.03759	16	166.952	5.280129	5.174119	894.3154



Figure 13: Expected number of events according to theoretical calculations from the **rpact** package and simulation results from the **PWEXP** package

2	33	15.90078	33	299.653	6.350680	5.413976	1906.5474
3	49	18.83120	49	427.435	6.744884	5.033877	2891.7742
4	65	21.16049	65	532.298	7.359293	5.650851	3929.0303
5	82	23.23549	82	620.501	8.093451	6.535725	5034.8399
6	98	25.09083	98	656.938	9.317254	7.851815	6124.7430
7	114	27.04290	114	659.976	11.028731	9.625848	7278.7666
8	131	29.35984	131	660.000	13.060754	11.770090	8620.0980
9	147	31.95780	147	660.000	15.279185	14.168758	10084.2622
10	163	35.37393	163	660.000	18.095337	17.312156	11942.9226

7 Summary and Discussion

Survival analysis plays a crucial role in various fields, where understanding the time until an event of interest occurs is the primary goal. The **PWEXP** package presented in this paper offers a comprehensive suite of tools tailored specifically for survival data with a piecewise exponential distribution (PWE). The **PWEXP** package follows a modular and user-friendly design, making it accessible to both novice and experienced researchers. The package is able to automatically fit the change-points, assess model performance using statistical criteria like AIC, BIC, and cross-validation log-likelihood, and estimate parameter uncertainty through bootstrapping. In addition to model estimation, the package supports event/timeline simulation and prediction. These tools empower researchers to explore complex survival datasets, derive meaningful

insights, and make informed decisions. Additionally, the package's integration with other statistical tools, such as **rpact**, **gsDesign**, enables researchers to perform comprehensive trial design simulations, including power calculations, interim analyses, and timeline estimations.

There are several future development directions worth studying. For example, current event/timeline prediction methods assume timely and complete reporting of data, which may not always align with real-world scenarios. Incorporating event report lag considerations, as explored in previous studies (Wang et al., 2012), could enhance prediction accuracy. Additionally, accounting for factors like cross-over (e.g., switch from control to treatment arm) and treatment switches (e.g., receive subsequent therapy) would broaden the package's applicability in diverse research contexts.

Computational details

The results in this paper were obtained using \mathbf{R} 4.3.0 with the **PWEXP** 0.5.0 package. \mathbf{R} itself and all packages used are available from the Comprehensive \mathbf{R} Archive Network (CRAN) at https://CRAN. R-project.org/.

References

Anderson, K. (2023). gsDesign: Group Sequential Design. R package version 3.6.0.

- Bagiella, E. and D. F. Heitjan (2001). Predicting analysis times in randomized clinical trials. Statistics in medicine 20(14), 2055–2063.
- Betensky, R. A. (2015). Measures of follow-pp in time-to-event studies: Why provide them and what should they be? *Clinical Trials* 12(4), 403–408.
- Chapple, A. G., T. Peak, and A. Hemal (2020). A novel bayesian continuous piecewise linear log-hazard model, with estimation and inference via reversible jump markov chain monte carlo. *Statistics in Medicine* 39(12), 1766–1780.
- Cooney, P. and A. White (2021a). Change-point detection for piecewise exponential models. arXiv preprint arXiv:2112.03962.
- Cooney, P. and A. White (2021b). Change-point detection for piecewise exponential models. arXiv, 1–21.
- Dupuy, J.-F. (2006). Estimation in a change-point hazard regression model. Statistics & Probability Letters 76(2), 182–190.
- Friedman, M. (1982). Piecewise exponential models for survival data with covariates. The Annals of Statistics 10(1), 101–113.
- Frumento, P. (2021). *pch:* Piecewise Constant Hazard Models for Censored and Truncated Data. **R** package version 2.0.

- Goodman, M. S., Y. Li, and R. C. Tiwari (2011). Detecting multiple change points in piecewise constant hazard functions. *Journal of Applied Statistics* 38(11), 2523–2532.
- Henderson, R. (1990). A problem with the likelihood ratio test for a change-point hazard rate model. Biometrika 77(4), 835–843.
- Hess, K. and G. Robert (2021). *muhaz:* Hazard Function Estimation in Survival Analysis. R package version 1.2.6.4.
- Kim, J., S. Cheon, and Z. Jin (2020). Bayesian multiple change-points estimation for hazard with censored survival data from exponential distributions. *Journal of the Korean Statistical Society* 49, 15–31.
- Klein, J. P., M. L. Moeschberger, et al. (2003). Survival Analysis: Techniques for Censored and Truncated Data, Volume 1230. Springer.
- Küchenhoff, H. (1996). An exact algorithm for estimating breakpoints in segmented generalized linear models.
- Li, Y., L. Qian, and W. Zhang (2013). Estimation in a change-point hazard regression model with long-term survivors. Statistics & Probability Letters 83(7), 1683–1691.
- Liu, N., Y. Zhou, and J. J. Lee (2021). Ipdfromkm: Reconstruct individual patient data from published kaplan-meier survival curves. *BMC medical Research Methodology* 21(1), 111.
- Loubert, S. K. (1986). Inference Procedures for the Piecewise Exponential Model When the Data Are Arbitrarily Censored. Iowa State University.
- Matthews, D. E. and V. T. Farewell (1982). On testing for a constant hazard against a change-point alternative. *Biometrics*, 463–468.
- Muggeo, V. M. (2003). Estimating regression models with unknown break-points. Statistics in Medicine 22(19), 3055–3071.
- Qian, L. and W. Zhang (2013). Multiple change-point detection in piecewise exponential hazard regression models with long-term survivors and right censoring. In *Contemporary Developments in Statistical Theory:* A Festschrift for Hira Lal Koul, pp. 289–304. Springer.
- Rohatgi, A. (2024). Webplotdigitizer: Version 4.7.
- Rufibach, K. (2022). eventTrack: Event Prediction for Time-to-Event Endpoints. R package version 1.0.2.
- Ruppert, A. S., J. G. Dixon, G. Salles, A. Wall, D. Cunningham, V. Poeschel, C. Haioun, H. Tilly, H. Ghesquieres, M. Ziepert, et al. (2020). International prognostic indices in diffuse large b-cell lymphoma: A comparison of ipi, r-ipi, and nccn-ipi. *Blood, The Journal of the American Society of Hematology* 135(23), 2041–2048.
- Wang, J., C. Ke, Q. Jiang, C. Zhang, and S. Snapinn (2012). Predicting analysis time in event-driven clinical trials with event-reporting lag. *Statistics in Medicine* 31(9), 801–811.

- Wassmer, G. and F. Pahlke (2024). *rpact: Confirmatory Adaptive Clinical Trial Design and Analysis.* R package version 3.5.1.
- Yao, Y.-C. (1986). Maximum likelihood estimation in hazard rate models with a change-point. *Communications in Statistics-Theory and Methods* 15(8), 2455–2466.
- Ying, G.-s. and D. F. Heitjan (2008). Weibull prediction of event times in clinical trials. Pharmaceutical Statistics: The Journal of Applied Statistics in the Pharmaceutical Industry 7(2), 107–120.



A Generating Synthetic Survival Data with simdata()

Figure 14: Relationship between variables in the generated dataset

The simdata() function can generate very flexible synthetic survival dataset with:

- randomization/enrollment time defined by randomization/enrollment rate (number of subjects per month) with 'rand_rate' argument, or randomization/enrollment curve (number of subjects in each month) with 'n_rand' argument;
- multiple treatment groups with stratification by user-defined allocation ratio with 'group', 'strata' and 'allocation' arguments;
- primary endpoint (event), drop-out and death with exponential distribution with 'event_lambda', 'drop_rate', 'death_lambda' arguments, or user-defined distributions (e.g., piecewise exponential, mixture distribution, etc.) with 'advanced_dist' argument for each stratification in each group.

In the generated dataset, all subjects are followed for infinite time and we can truncate it by a clinical cut-off date (DCOD) with the cut_dat() function later. The following variables are included the generated dataset (some are requested by the 'add_column' argument):

- 'randT' is the randomization time for each subject;
- 'eventT', 'dropT', 'deathT' are the event time, drop-out time and death time, respectively;

- 'followT' is the follow-up time, which is the minimum value of 'eventT', 'dropT' and 'deathT';
- 'eventT_abs', 'dropT_abs', 'deathT_abs' and 'followT_abs' are the corresponding time from the beginning of the trial (e.g., 'eventT_abs' is the sum of 'randT' and 'eventT', etc.);
- 'event' indicates whether the primary event occurred at the end of follow-up with value 1 for occurrence of event. 'censor' indicates whether the censoring (including drop-out and death) occurred at the end of follow-up with value 1 for censoring. If a subject is censored, 'censor_reason' shows the type of censoring (i.e., 'drop_out', 'death' or 'never_event' (when 'followT == Inf')).

Figure 14 shows some examples of the relationship between these variables.