
NON-EXPLOSION SOLUTIONS FOR A CLASS OF STOCHASTIC PHYSICAL DIFFUSION OSCILLATORS

A PREPRINT

Amrane . Houas

Laboratory of Applied Mathematics,
University Mohamed Khider, Biskra, Algeria.
houas.amrane@univ-biskra.dz

Fateh. Merahi

Department of Mathematics
University of Batna II, Algeria.
f.merahi@univ-batna2.dz

Mustafa. Moumni

Laboratory of Photonic Physics and Nano-Materials,
University of Biskra, Algeria,
LPRIM, Department of Physics, University of Batna I, Algeria.
m.moumni@univ-batna.dz

May 1, 2024

ABSTRACT

In this work, we are interested in problems that are related to the physical phenomena of diffusion. We will focus on the theoretical aspect of the study, such as existence, uniqueness and non-explosive solutions. We will weaken the conditions imposed on the coefficients of the stochastic differential equations (SDE) that model some diffusion phenomena of mechanics. The work will be based on a general non-explosion criterion and we will obtain sufficient conditions so that the solution for a certain class of diffusions does not explode. We will construct Lyapunov functions that ensure the non-explosion of the solutions. Two important oscillators, namely the Duffing and the Van Der Pol oscillators, belong to this class. The Euler-Maruyama method is applied to these two oscillators to give us a simulation solution for them.

Keywords SDE · Non-explosion · Oscillator · Euler-Maruyama method · AMS: · 60F99 · 60G07 · 37H05

1 Introduction

Differential equations for stochastic processes arise when studying many problems in physics and engineering (Sobczyk [2013], Thygesen [2023], Särkkä and Solin [2019], Øksendal [2022]). They are typically of two basic types:

$$\begin{cases} \dot{X}_t = A(t)X_t + B(t)X_{t_0} = c & (I) \end{cases}$$

$$\begin{cases} \dot{X}_t = f(t, X_t, \eta_t)X_{t_0} = c & (II) \end{cases}$$

where $A(t)$, $B(t)$ and η_t are random functions and the initial condition c is also a random variable in both cases. If the random functions are sufficiently regular, we can consider the problems (I) and (II) as a family of problems corresponding to different realisations. In this case we can treat them with the classical methods of differential equations. The situation is different when the random functions are of the white noise type. White noise is perceived as a Gaussian stochastic process with zero mean and constant spectral density along the entire real axis Karatzas and Shreve [2012]. Such a process does not really exist because its covariance function is the Dirac function and therefore its variance is infinite. However, white noise is a very important mathematical idealisation for S.D.E. random influences that vary rapidly and whose values are virtually uncorrelated at different times Øksendal [2003].

The analysis of stochastic dynamical systems leads to differential equations of the following form

$$\dot{X}_t = f(t, X_t) + G(t, X_t)\xi_t$$

Here ξ_t is the White noise, G (respectively f) is a function defined from $[t_0, t] \times \mathbb{R}^d$ (respectively $[t_0, t] \times \mathbb{R}^d$) with values in $Mat_{\mathbb{R}}(d, d)$ (respectively \mathbb{R}^d).

Some difficulties arise in studying these equations and they are due to the fact that white noise does not exist as a stochastic process in the classical sense. Other difficulties arise from the multitude of possible interpretations of this type of equation Ikeda and Watanabe [2014].

The challenges associated with this particular equation have been significantly mitigated due to the contributions of Ito, Bernstein, and Guikhman Ikeda and Watanabe [2014]. Independently, the first two authors developed methods for addressing this type of equation (Ito et al. [1951], Gikhman et al. [2007]). The solutions to these equations are Markovian processes with continuous trajectories and serve as probabilistic theoretical models for physical diffusion phenomena with random influences. A notable example of a diffusion process is the Wiener or Brownian motion process, as elaborated upon in the reference Burdzy [2014].

Non-linear stochastic oscillators have numerous important applications in various fields of science and engineering (Kyzioł and Okniński [2015], Pikovsky et al. [2003], Bommundt [1993]). This is because they correspond more closely to natural phenomena, as linearity is often just an approximation of reality that only serves to facilitate the theoretical study of real phenomena. In this paper, our focus lies on the exploration of non-explosive solutions within the realm of a specific class of diffusion mechanics, which can be mathematically represented as::

$$\ddot{x}_t + bf(x_t, \dot{x}_t) + g(x_t) = \sigma(x_t, \dot{x}_t) \dot{W}_t$$

Considering the solution as a vector in the phase space, the coefficients of drift and diffusion of the associated SDE do not satisfy the global conditions of existence and uniqueness of the solution. This implies that these solutions can escape to infinity in a finite time (known as an exploding SDE). To prevent this divergence, we establish sufficient conditions for f , g , and σ based on non-explosion criteria in the general case Khasminskii [2011].

Our focus is on the Duffing-Van der Pol oscillator (Van der Pol [1920], Van der Pol and Van Der Mark [1927], Duffing [1918], Guckenheimer and Holmes [2013]), a special case of these oscillators. This oscillator plays a crucial role in various fields, including physics, chemistry, biology, electronics, and engineering (Szemplińska-Stupnicka and Rudowski [1997], Chudzik et al. [2011], Bold et al. [2003], Kudryashov [2021])

2 Motivation

Modelling the dynamic behaviour of structures such as large suspension bridges and oil platforms, which are subjected to random oscillations caused by aerodynamic effects and earthquakes, leads to second-order equations of the following type:

$$\ddot{x}(t) + f(x(t), \dot{x}(t)) = \sigma(x(t), \dot{x}(t)) \dot{W}(t) \quad (1)$$

Where :

- $x(t) \in \mathbb{R}^n$ with $t \geq 0$
- f is a function defined from $\mathbb{R}^n \times \mathbb{R}^n$ to \mathbb{R}^n
- σ is a function defined from $\mathbb{R}^n \times \mathbb{R}^n$ to $Mat_{\mathbb{R}}(n, m)$
- $\dot{W}(t)$ is a white noise of dimension m

The first step consists of representing equation (1) as Itô stochastic differential equation.

The classical method of approaching equations of type (1) is to place the system in phase space by considering the process $z(t) = (x(t), \dot{x}(t))$ and to represent (1) as Itô SDE on \mathbb{R}^{2n} .

So if we set $y(t) = \dot{x}(t)$, the equation (1) is transformed to :

$$\begin{cases} dx(t) = y(t)dt \\ dy(t) = -f(x(t), y(t)) + \sigma(x(t), y(t))dW(t) \end{cases} \quad (2)$$

The aim of this study is to investigate the existence and uniqueness of solutions to the equation (2) that do not explode in finite time.

Due to the importance of Duffing-Van der Pol oscillators in the theory of stability and bifurcations for stochastic dynamical systems (Baxendale [2004], Arnold and Khas' minskii [1993], Arnold et al. [1996], Keller and Ochs [1999], Schenck-Hoppé [1996a], Li et al. [2023], Benedetti et al. [2022]), this study relies on two significant examples:

Duffing oscillator :

$$\ddot{x}(t) + 2\alpha\omega_0\dot{x}(t) + \omega_0^2(x(t) + \lambda x(t)^3) = \sigma \dot{W}(t) \quad (3)$$

Van Der Pol oscillator

$$\ddot{x}(t) + 2\xi\omega_0(x(t)^2 - 1)\dot{x}(t) + \omega_0^2(x(t) + \gamma x(t)^3) = \sigma \dot{W}(t) \quad (4)$$

3 Non-Explosion Criteria

Let's consider the SDE :

$$\begin{cases} dx(t) = b(x(t))dt + \sigma(x(t))dW(t), t > 0 \\ x(0) = x_0 p.s \end{cases} \quad (5)$$

With

- $x(t) \in \mathbb{R}^n$ with $t \geq 0$
- b is a function defined from \mathbb{R}^n to \mathbb{R}^n
- σ is a function defined from \mathbb{R}^n to $Mat_{\mathbb{R}}(n, n)$

Theorem 1. (Non-Explosion Criteria) (Khasminskii [2011])

Under the hypothesis :

H1) $\forall R > 0, \exists K_R > 0 : |b(x) - b(y)| + |\sigma(x) - \sigma(y)| \leq K_R |x - y|, \forall |x| \leq R, \forall |y| \leq R$

H2) $x(0)$ independent to $W = \{W(t), t \geq 0\}$ and $E|x_0|^2 < \infty$

H3) There exist a function $x \mapsto V(x)$ from \mathbb{R}^n to \mathbb{R}_+ of C^2 class, named the Lyapounov function and it verifies

$$\begin{cases} \forall x \in \mathbb{R}^n LV(x) \leq cV(x), c > 0 \\ V(x) \rightarrow +\infty \text{ if } |x| \rightarrow +\infty \end{cases}$$

With

$$\begin{cases} L = \sum_{i=1}^n b_i(x) \frac{\partial}{\partial x_i} + \frac{1}{2} \sum_{i,j=1}^n a_{ij}(x) \frac{\partial^2}{\partial x_i \partial x_j}; \\ a(x) = \sigma(x)\sigma^T(x) \end{cases}$$

being the differential generator associated with the diffusion process solution of SDE

The SDE given by (5) has a unique solution that does not explode in a finite time and it is a diffusion process.

4 The main results

4.1 Diffusion of mechanics

Consider the oscillator

$$\ddot{x}(t) + b(x(t), \dot{x}(t)) + g(x(t)) = \sigma(x(t), \dot{x}(t))\dot{W}(t) \quad (6)$$

Where

- $x(t) \in \mathbb{R}^n$, with $t \geq 0$
- $b : \mathbb{R}^{2n} \rightarrow \mathbb{R}^n$
- $g : \mathbb{R}^n \rightarrow \mathbb{R}^n$
- $\dot{W}(t)$ is a white noise
- $\sigma : \mathbb{R}^n \times \mathbb{R}^n \rightarrow Mat_{\mathbb{R}}(n, n)$

If we put $y(t) = \dot{x}(t)$, then equation 6 will write as :

$$\begin{cases} dx(t) = y(t)dt \\ dy(t) = -[b(x(t), y(t)) + g(x(t))]dt + \sigma(x(t), y(t))dW(t) \end{cases} \quad (7)$$

The differential generator associated with the equation (7) is :

$$L = \langle y, \nabla_x \rangle - \langle b(x, y) + g(x), \nabla_y \rangle + \frac{1}{2} Tr[\sigma(x, y)\sigma^T(x, y)D_y^2]$$

with $\nabla_x = (\frac{\partial}{\partial x_1}, \dots, \frac{\partial}{\partial x_n})$, $\nabla_y = (\frac{\partial}{\partial y_1}, \dots, \frac{\partial}{\partial y_n})$ and $D_y^2 = ((\frac{\partial}{\partial x_i \partial y_j})_{ij})$, $\langle ., . \rangle$ is the scalar product

Theorem 2. Let's consider the system

$$\begin{cases} dx(t) = y(t)dt \\ dy(t) = -[b(x(t), y(t)) + g(x(t))]dt + \sigma(x(t), y(t))dW(t) \\ x(0) = x_0, y(0) = y_0 \end{cases} \quad (8)$$

Under the conditions :

- 1) b is locally Lipschitzian,
- 2) The function g derives from a potential i.e $g(x) = \nabla_x G(x)$ where $G : \mathbb{R}^n \rightarrow \mathbb{R}$ of C^2 class such that $G(x) \rightarrow +\infty$ if $|x| \rightarrow +\infty$
- 3) $\exists K_1 > 0, \exists c > 0$ and

$$\frac{1}{2} \text{Tr}[\sigma(x, y)\sigma^T(x, y)] \leqslant \langle y, b(x, y) \rangle + c[G(x) + \frac{1}{2}|y|^2] + K_1$$

- 4) The random variable (x_0, y_0) satisfies the hypothesis **H2** of the theorem 1

The system 8 admits a unique solution that does not explode in a finite time

Proof. We will demonstrate that all the hypotheses of Theorem 1 are satisfied.

Firstly, it should be noted that b is locally Lipschitzian. Secondly, G is of C^2 class, which means that g is of C^1 class. Additionally, considering condition 03, the coefficients of the system in 8 are locally Lipschitzian (**H1** is satisfied).

Now consider the function $V(x, y) = G(x, y) + \frac{1}{2}|y|^2 + K$, where K is a positive constant that will be chosen further. Since $G(x) \rightarrow +\infty$ if $|x| \rightarrow +\infty$, then $\exists K_2 > 0 : G(x) + K_2 \geqslant 0 \forall x \in \mathbb{R}^n$ and so, we can choose the constant K sufficiently large to ensure that

$$V(x, y) \geqslant 0 \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^n \quad V(x, y) \rightarrow +\infty \text{ if } R = |x|^2 + |y|^2 \rightarrow +\infty$$

Moreover, since G is of C^2 class, V is also of C^2 class.

Applying the differential generator associated with 8 to the function V yields

$$\begin{aligned} LV(x, y) &= \langle y, g(x) \rangle - \langle y, b(x, y) + g(x) \rangle + \frac{1}{2} \text{Tr}[\sigma(x, y)\sigma^T(x, y)] \\ &= -\langle y, b(x, y) \rangle + \frac{1}{2} \text{Tr}[\sigma(x, y)\sigma^T(x, y)] \\ &\leqslant c[G(x) + \frac{1}{2}|y|^2] + K_1 \quad (\text{condition 03}) \\ &\leqslant cV(x, y) - cK + K_1 \end{aligned}$$

We choose $K \geqslant \frac{K_1}{c}$ such that $LV(x, y) \leqslant cV(x, y), \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^n$

Therefore, the Hypothesis H3 of Theorem 1 is satisfied, which implies that the system (8) has a unique solution that does not explode in a finite time. \square

Corollary 3. Under the conditions 1), 2) and 4) and if $\sigma(x, y) = \sigma$ is a constant and b verifies

$$\exists \alpha \geqslant 0 : \langle b(x, y), y \rangle \geqslant -\alpha|y|^2 \quad \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^n,$$

the result of the theorem remains valid.

Proof. We have

$$[G(x) + \frac{1}{2}|y|^2] + \langle y, b(x, y) \rangle + K \geqslant cG(x) + (\frac{c}{2} - \alpha)|y|^2 + K, \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^n.$$

Because $\text{Tr}[\sigma(x, y)\sigma^T(x, y)] \geqslant 0$, we have to choose the constant K sufficiently large and $c > 2\alpha$ such that $cG(x) + (\frac{c}{2} - \alpha)|y|^2 + K \geqslant \frac{1}{2}\text{Tr}[\sigma(x, y)\sigma^T(x, y)]$; hence the result follows. \square

4.2 Change of the variable

The energy function of the system (8) serves as a Lyapunov function in Theorem 2. The function is defined as follows: $V(x, y) = G(x, y) + \frac{1}{2} |y|^2 < y, y > + K$. However, it is necessary to assume that $G(x) \rightarrow +\infty$ if $|x| \rightarrow +\infty$; This is the first restriction.

A second restriction arises from the fact that if we apply the differential generator L to this function, we have for $y = 0$: $LV(x, y) = \frac{1}{2} \text{Tr}[\sigma(x, y)\sigma^T(x, y)]$ and we cannot have $LV(x, y) \rightarrow -\infty$ if $|x|^2 + |y|^2 \rightarrow +\infty$, which is a necessary condition to ensure the existence of a stationary solution.

To overcome these difficulties, we can apply a change of variables in the starting system and search for another Lyapunov function for the new equivalent system.

So, we assume that the system (6) has the form:

$$\ddot{x}_i(t) + f_i(x_i(t))\dot{x}_i(t) + \frac{\partial G}{\partial x_i}(x(t)) = \sum_{j=1}^n \sigma_{ij}(x(t), \dot{x}(t))\dot{W}_j, i = \overline{1, n} \quad (9)$$

where f_i ($1 \leq i \leq n$), is an application from \mathbb{R} to \mathbb{R} and G is an application from \mathbb{R}^n to \mathbb{R} .

Let $\dot{x}(t) = y(t) = (y_1(t), \dots, y_n(t))$.

In phase space, the oscillator (9) is written

$$\begin{cases} dx_i(t) = y_i(t) \\ dy_i(t) = -[f_i(x_i(t)) + \frac{\partial G}{\partial x_i}(x(t))]dt + \sum_{j=1}^n \sigma_{ij}(x(t), y(t))\dot{W}_j, i = \overline{1, n} \end{cases} \quad (10)$$

Now, consider the application

$$\begin{cases} \Phi : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}^n \times \mathbb{R}^n \\ (x, y) \mapsto (x_1, \dots, x_n, y_1 + F(x_1), \dots, y_n + F(x_n)) \end{cases} \quad (11)$$

where $F_i(x_i) = \int_0^{x_i} f_i(s)ds$

If we suppose that the functions f_i are of class C^1 , then the application defined by (11) is a C^2 diffeomorphism from $\mathbb{R}^n \times \mathbb{R}^n$ to $\mathbb{R}^n \times \mathbb{R}^n$.

Let's apply Itô's formula to the process

$$\begin{cases} \Phi_i(x, y) = x_i \\ \Phi_{i+n}(x, y) = y_i + F_i(x_i), i = \overline{1, n} \end{cases} \quad (12)$$

We obtain

$$\begin{cases} d\Phi_i = (\Phi_{i+n} - F_i(\Phi_i))dt \\ d\Phi_{i+n} = -\frac{\partial G}{\partial x_i}dt + \sum_{j=1}^n \sigma_{ij}(\varphi, \varphi - F(\varphi))dW_j(t), i = \overline{1, n} \end{cases} \quad (13)$$

where $\varphi = (\Phi_1, \dots, \Phi_n)$, $F(\varphi) = (F_1(\Phi_1), \dots, F_n(\Phi_n))$

Going back to the notation (x, y) in (13), the system (10) is therefore equivalent to the following system

$$\begin{cases} dx_i = (y_i - F_i(x_i))dt \\ dy_i = -\frac{\partial G}{\partial x_i}dt + \sum_{j=1}^n \sigma_{ij}(x, y - F(x))dW_j(t), i = \overline{1, n} \end{cases} \quad (14)$$

where $F(x) = (F_1(x_1), \dots, F_n(x_n))$

Let

$$H(x) = \sum_{i=1}^n \int_0^{x_i} F_i(s)ds \quad (15)$$

then, the differential generator associated with the system (14) is given by

$$L = < y - \nabla H(x), \nabla x > - < \nabla G(x), \nabla y > + \frac{1}{2} \text{Tr}[\sigma(x, y - F(x))\sigma^T(x, y - F(x))D_y^2] \quad (16)$$

Now we are interested in studying the non-explosive criteria for the solution of the system (14) under weaker hypotheses than those of Theorem 2.

4.2.1 Scalar case

In this case, equation 9 is written as

$$\ddot{x}(t) + f(x(t))\dot{x}(t) + g(x(t)) = \sigma(x(t), y(t))\ddot{W}(t) \quad (17)$$

According to (14), it is equivalent to the system

$$\begin{cases} dx(t) = (y(t) - F(x(t)))dt \\ dy(t) = -g(x(t))dt + \sigma(x(t), y(t) - F(x(t)))dW(t) \end{cases} \quad (18)$$

where $F(x) = \int_0^x f(s)ds$

Theorem 4. Suppose that

- 01)** f and g are C^1 class and σ is locally Lipschitzian,
- 02)** $\exists c_1 > 0, \exists c_2 > 0; x[F(x) + g(x)] \geq c_1 x^2, \forall |x| \geq c_2$,
- 03)** $\exists K_1 > 0 : \frac{1}{2}\sigma^2(x, y - F(x)) \geq \frac{1}{2}(y - F(x))^2 + F(x)[\frac{1}{2}F(x) + g(x)] + K_1, \forall (x, y) \in \mathbb{R}^2$,

then for each initial condition that verifies the hypothesis **H2** of Theorem 1, the system (18) has a unique solution that does not explode in a finite time.

Remark 5. This theorem shows that the following assumption

$$\exists c_1 > 0, \exists c_2 > 0; xg(x) \geq c_1 x^2, \forall |x| \geq c_2$$

previously thought to be necessary in precedent works (Schenk-Hoppé [1996b]), is in fact not necessary.

Proof. According to hypothesis **01**, the coefficients of system (18) satisfy the condition of local Lipschitz. Now, let us consider the function

$$V(x, y) = \int_0^x [F(s) + g(s)]ds + \frac{1}{2}y^2 + K$$

V is C^2 class and the condition **02** implies that $V(x, y) \geq 0$ and $V(x, y) \rightarrow +\infty$ if $x^2 + y^2 \rightarrow +\infty$. Applying the differential generator associated with equation (18) to V yields:

$$\begin{aligned} LV(x, y) &= (y - F(x))[F(x) + g(x)] - g(x)y + \frac{1}{2}\sigma^2(x, y - F(x)) \\ &= yF(x) - F(x)[F(x) + g(x)] + \frac{1}{2}\sigma^2(x, y - F(x)) \\ &\quad - \frac{1}{2}(y - F(x))^2 - F(x)[\frac{1}{2}F(x) + g(x)] + \frac{1}{2}y^2 + \frac{1}{2}\sigma^2(x, y - F(x)) \end{aligned}$$

and using hypothesis **03** gives

$$LV(x, y) \leq \frac{1}{2}y^2 + K_1, \forall (x, y) \in \mathbb{R}^2$$

If we choose $K \geq K_1$, then

$$LV(x, y) \leq \frac{1}{2}y^2 + K \leq V(x, y)$$

Therefore, this verifies the Hypothesis **H3** of Theorem 1, and as a result, the system (18) has a unique solution that does not explode in a finite time. \square

4.2.2 Vectorial case

Theorem 6. Assume that

- 01** $f_i, \frac{\partial G}{\partial x_i}$, are C^1 class, for $i = 1, \dots, n$

- 02** $\lim_{|x| \rightarrow +\infty} im [H(x) + G(x)] = +\infty$

$$\mathbf{03} \quad \exists K_2 \geq 0 : Tr[\sigma(x, y - F(x))\sigma^T(x, y - F(x))] \leqslant \langle y, y \rangle + \langle \nabla H(x), \nabla H(x) + 2\nabla G(x) \rangle + K_2, \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^n$$

then for each initial condition that verifies **H2** of Theorem 1, the system 14 admits a unique solution that does not explode in a finite time

Proof. The condition **01** ensures that the hypothesis **H1** is verified. Now consider the function

$$V(x, y) = H(x) + \frac{1}{2} \langle y, y \rangle + G(x) + K$$

where $K > 0$ is a constant.

Because $f_i, \frac{\partial G}{\partial x_i}$ are C^1 class for $i = 1, \dots, n$, then V is C^2 class.

The condition **02**) implies that we have $V(x, y) \geq 0, \forall (x, y) \in \mathbb{R}^n \times \mathbb{R}^n$ and $V(x, y) \rightarrow +\infty$ if $R = |x|^2 + |y|^2 \rightarrow +\infty$. Let us apply the differential generator associated with V ; we get:

$$\begin{aligned} LV(x, y) &= \langle y - \nabla H, \nabla H + \nabla G \rangle - \langle \nabla G, y \rangle + \frac{1}{2} Tr[\sigma(x, y - F(x))\sigma^T(x, y - F(x))] \\ &= V(x, y) - \frac{1}{2} \langle y, y \rangle - \frac{1}{2} \langle \nabla H, \nabla H \rangle - \langle \nabla H, \nabla G \rangle + \frac{1}{2} Tr[\sigma(x, y - F(x))\sigma^T(x, y - F(x))] \end{aligned}$$

Using the condition **03**) and choosing $K \geq K_2$ we get $LV(x, y) \leq V(x, y)$, then the system (14) admits a unique solution that does not explode at finite time. \square

5 Applications

It will now be shown that both the Duffing and the Vander-Pol oscillators belong to the analyzed class of oscillators and that their solutions do not explode.

5.1 Scalar case

Example 1 (Duffing Oscillator)

Let's consider the oscillator

$$\ddot{x}(t) + 2\alpha\omega_0\dot{x}(t) + \omega_0^2(x(t) + \lambda x(t)^3) = \sigma \dot{W}(t) \quad (19)$$

where α, λ and ω_0 are constants in \mathbb{R}_+^* and σ is a constant. In this case, we have

$$yb(x, y) = 2\alpha\omega_0 y^2, G(x) = \omega_0^2(\frac{x^2}{2} + \lambda \frac{x^4}{4})$$

then $yb(x, y) \leq -y^2$ and $G(x) \rightarrow +\infty$ if $x \rightarrow +\infty$, therefore, corollary 1 applies in this case, which means that the solutions of the equation (19) do not explode in a finite amount of time.

Example 2 (Vander-Pol Oscillator)

We have the system:

$$\ddot{x}(t) + 2\xi\omega_0(x(t)^2 - 1)\dot{x}(t) + \omega_0^2(x(t) + \gamma x(t)^3) = \sigma \dot{W}(t) \quad (20)$$

where ξ, γ and ω_0 are constants in \mathbb{R}_+^* and σ is a constant. In this case, we have

$$yb(x, y) = \xi\omega_0(x^2 - 1)y^2, G(x) = \omega_0^2(\frac{x^2}{2} + \gamma \frac{x^4}{4})$$

Here it is clear that corollary 1 is satisfied, meaning that solutions of (refeq20) do not explode in finite time.

Example 3 (Duffing-Van Der Pol Oscillator)

Considering the oscillator

$$\ddot{x}(t) + [\sum_{j=1}^{2m} \xi_j x(t)^j] \dot{x}(t) + [\sum_{j=1}^{2n} a_j x(t)^{j+1}] = \sigma(x(t), \dot{x}(t)) \dot{W}(t) \quad (21)$$

with $\xi_{2m} > 0, a_{2n} < 0$, and $\xi_j \in \mathbb{R}$ for $j = \overline{1, 2m-1}$, $a_j \in \mathbb{R}$ for $j = \overline{1, 2n-1}$, and $m > n \geq 1$. The application σ satisfies the condition (03) of theorem 3.

In this case, we have:

$$F(x) = \sum_{j=1}^{2m} \frac{\xi_j}{j+1} x^{j+1}, g(x) = \sum_{j=1}^{2n} a_j x^{j+1} \text{ and}$$

$$\exists \alpha > 0, \exists c_1 \geq 0 : x[F(x) + G(x)] \geq \alpha x^2, \text{ if } |x| \geq c_1.$$

This verifies the conditions of Theorem 3, showing that solutions of (21) do not explode in finite time.

5.2 Vectorial case

Example 1 (Duffing Oscillator)

We study the oscillator :

$$\ddot{X}(t) + B\dot{X}(t) + g(X(t)) = \sigma(X(t), \dot{X}(t))\dot{W}(t) \quad (22)$$

where $X(t) \in \mathbb{R}^n, \forall t \geq 0, B = (b_{ij})$ is a positive matrix of rang $n \times n$.

Suppose that

$$g(x) = Ax + \sum_{i=1}^n K_{ii}x_i^3 e_i, K_{ii} > 0, \forall i = \overline{1, n}$$

where $A = (a_{ij})$ is a symmetric positive matrix of rang $n \times n$ and $\{e_i\}_{i=\overline{1, n}}$ is the canonical base of \mathbb{R}^n , then

$$G(x) = \frac{1}{2} \langle x, Ax \rangle + \frac{1}{4} \sum_{i=1}^n K_{ii}x_i^4 \geq \frac{1}{2} \langle x, Ax \rangle + \frac{\alpha}{2} |x|^2, (\alpha > 0),$$

and so, $G(x) \rightarrow +\infty$ if $|x| \rightarrow +\infty$

In addition we have

$$\langle y, B(x, y) \rangle = \langle y, By \rangle \geq 0.$$

If the matrix $\sigma(x, y)$ verifies :

$$\exists K_1 > 0, \exists \lambda \geq 0 : \text{Tr}[\sigma(x, y)\sigma^T(x, y)] \leq \lambda[G(x) + y^2] + K_1$$

then theorem 2 is applicable and the solutions of (22) are non-explosive in a finite time.

Example 2

Let be the oscillator :

$$\ddot{x}_i(t) + \xi_i(x_i^{2n_1}(t))\dot{x}_i(t) + \frac{\partial G}{\partial x_i}(x(t)) = \sum_{j=1}^n \sigma_{ij}(x_i(t), \dot{x}_i(t))dW_j(t), i = \overline{1, n} \quad (23)$$

with $G(x) = -\sum_{j=1}^n (a_j x_j^{2n_2+2} + \nu x_1 x_i)$ where $\xi_i > 0, a_i > 0$ and ν are real numbers and $n_1 > n_2 > 0$ are natural numbers.

In this case, we have

$$f_i(x) = \xi_i x_i^{2n_1}, H(x) = \sum_{i=1}^n \frac{\xi_i}{(2n_1+1)(2n_1+2)} x_i^{2n_1+2}$$

The matrix $\sigma(x, y)$ satisfies condition (03) of theorem 4 and, so, the solutions of system 23 do not explode in a finite time.

6 Simulation of Duffing and Vander-Pol oscillators with Euler-Maruyama method

This section provides a simulation of the solution for Duffing and Vanderpol oscillators, expressed as a stochastic differential equation, using the Euler-Maruyama method.

The Euler-Maruyama method is a numerical technique used to simulate solutions to stochastic differential equations (SDEs). It is an extension of the classical Euler method for ordinary differential equations (ODEs) and provides an approximation of the trajectories of stochastic processes Milstein and Tretyakov [2004], particularly those involving random fluctuations like Brownian motion. This method is particularly useful when closed-form analytical solutions are difficult to obtain. The Euler-Maruyama method has various applications in different fields. The method is commonly used to study the dynamics of physical systems under random perturbations, simulate financial models with stochastic components, analyze biological processes influenced by noise, and more (Higham et al. [2002], Lang and Motschan-Armen [2023], Wang et al. [2020], Jin [2023], Hiderah [2023]).

6.1 Simulation of Duffing oscillator

We have already shown the existence, uniqueness and non-explosivity of the Duffing oscillator solution in 19

$$\ddot{x}(t) + 2\alpha\omega_0\dot{x}(t) + \omega_0^2(x(t) + \lambda x(t)^3) = \sigma \dot{W}(t)$$

where α , λ , ω_0 , and σ are constants in \mathbb{R}_+^* .

We are going to do a simulation of the solution for (19) by means of the Euler-Maruyama method.

6.1.1 Discretization steps with Euler-Maruyama method

Starting with the given SDE (19), we perform the following steps

- 1) Select a small time step size Δt to discretize the continuous-time equation. This step size determines the granularity of the simulation.
- 2) Set initial conditions for $x(0)$ and $v(0)$, where $v(t) = \dot{x}(t)$. Also, set up arrays to store the values of x , v , and time t at each step.
- 3) Generate random numbers following a normal distribution, scaled by $\sqrt{\Delta t}$, to represent the increments of the Wiener process (Brownian motion) at each time step.
- 4) Iterate over time steps from $t = 0$ to $t = T$ (where T is the total simulation time) with steps of Δt .
- 5) For each step:

- a) Calculate the deterministic term a and the stochastic term b based on the SDE (19)

$$\begin{aligned} a &= -2\alpha\omega v(t) - \omega^2 x(t) - \omega^2 \lambda x(t)^3 \\ b &= \sigma \dot{W}(t) \end{aligned}$$

- b) Update x and v using the Euler-Maruyama update rule:

$$\begin{aligned} x(t + \Delta t) &= x(t) + v(t)\Delta t \\ v(t + \Delta t) &= v(t) + a\Delta t + b \end{aligned}$$

- 6) After the simulation is complete, we plot the values of x and v over time t to visualize the behavior of the Duffing oscillator.

Figure 1 shows the graph of the simulation solution of position and velocity for the following values of the Duffing Oscillator Parameters: $\alpha = 0.5$, $\lambda = 3$, $\omega_0 = 1$, and $\sigma = 2$.

6.2 Simulation of Vander-Pol oscillator

In (20), we have demonstrated the existence, uniqueness, and no-explosion of the solution of the Vander-Pol oscillator

$$\ddot{x}(t) + 2\xi\omega_0(x(t)^2 - 1)\dot{x}(t) + \omega_0^2(x(t) + \gamma x(t)^3) = \sigma \dot{W}(t)$$

where ξ , γ , ω_0 , and σ are constants in \mathbb{R}_+^* . So, by means of the Euler-Maruyama method, we give a simulation of a solution for the Vander-Pol oscillator for some values of its parameters.

6.2.1 Discretization steps with Euler-Maruyama method

Starting with the given SDE (20), we follow the same steps as for the Duffing case. Based on the equation (20) the deterministic and stochastic terms are given by:

$$\begin{aligned} a &= 2\xi\omega x(t)^2 v(t) - 2\xi\omega v(t) + \omega^2 x(t) + \omega^2 \gamma x(t)^3 \\ b &= \sigma \dot{W}(t) \end{aligned}$$

Figure 2 shows the graph of the simulated solution for $\xi = 0.1$, $\gamma = 0.25$, $\omega = 1$, and $\sigma = 0.1$




Figure 1: Duffing oscillator with Brownian movement: position and velocity




Figure 2: Vander-Pol oscillator with Brownian movement: position and velocity

7 Conclusion

This work focuses on the theoretical study of stochastic differential equations, particularly on the conditions for the existence, uniqueness, and non-explosion of solutions. The study is centred on problems related to the physical phenomena of diffusion. The results show that it is possible to obtain sufficient non-explosive conditions. Sufficient conditions for constructing Lyapunov functions ensuring nonexplosive solutions are presented for a class of diffusion phenomena, including Duffing and Vander-Pol oscillators in scalar and vectorial cases. These reduced criteria has the potential to be applied to a wider range of physical cases. Finally, a simulation of Duffing and Vander-Pol oscillators using the Euler-Maruyama method is provided to demonstrate the non-explosive nature of the solutions.

Declaration of Conflict of interest statement All authors contributed to the study's conception and design. Material preparation, data collection, and analysis were performed by [Amrane Houas] and [Fateh Merahi], and [mustapha Moumni]. The first draft of the manuscript to submission was written by [Amrane Houas] and the co-authors [Fateh Merahi] and [mustapha Moumni] commented on this version of the manuscript and approved it. The authors did not receive support from any organization for the submitted work.

References

- Kazimierz Sobczyk. *Stochastic differential equations: with applications to physics and engineering*, volume 40. Springer Science & Business Media, 2013.
- Uffe Høgsbro Thygesen. *Stochastic Differential Equations for Science and Engineering*. CRC Press, 2023.
- Simo Särkkä and Arno Solin. *Applied stochastic differential equations*, volume 10. Cambridge University Press, 2019.
- Bernt Øksendal. Space-time stochastic calculus and white noise. In *Mathematics Going Forward: Collected Mathematical Brushstrokes*, pages 629–649. Springer, 2022.
- Ioannis Karatzas and Steven Shreve. *Brownian motion and stochastic calculus*, volume 113. Springer Science & Business Media, 2012.
- B Øksendal. Stochastic differential equations: An introduction with applications, 2003, 2003.
- Nobuyuki Ikeda and Shinzo Watanabe. *Stochastic differential equations and diffusion processes*, volume 24. Elsevier, 2014.
- Kiyosi Ito, Kiyosi Itô, Kiyosi Itô, Japon Mathématicien, Kiyosi Itô, and Japan Mathematician. *On stochastic differential equations*, volume 4. American Mathematical Society New York, 1951.
- Iosif Ilyich Gikhman, Anatoli Vladimirovich Skorokhod, Iosif Ilyich Gikhman, and Anatoli Vladimirovich Skorokhod. *Stochastic differential equations*. Springer, 2007.
- Krzysztof Burdzy. Brownian motion. In *Brownian Motion and its Applications to Mathematical Analysis*, pages 1–10. Springer, 2014.
- J Kyzioł and A Okniński. The duffing–van der pol equation: Metamorphoses of resonance curves. *Nonlinear Dynamics and Systems Theory*, page 25, 2015.
- Arkady Pikovsky, Michael Rosenblum, and Jürgen Kurths. *Synchronization: a universal concept in nonlinear sciences*, volume 12. Cambridge University Press, 2003.
- E Brømmundt. Awrejcewicz, j., bifurcation and chaos in coupled oscillators. singapore etc., world scientific publishing co. 1991. viii, 245 pp. isbn 981-02-0579-1. *ZAMP-Journal of Applied Mathematics and Mechanics/Zeitschrift für Angewandte Mathematik und Mechanik*, 73(1):46–46, 1993.
- Rafail Khasminskii. *Stochastic stability of differential equations*, volume 66. Springer Science & Business Media, 2011.
- Balthasar Van der Pol. theory of the amplitude of free forced triode vibrations. *Radio review*, 1:701–710, 1920.
- Balth Van der Pol and Jan Van Der Mark. Frequency demultiplication. *Nature*, 120(3019):363–364, 1927.
- Georg Duffing. *Erzwungene Schwingungen bei veränderlicher Eigenfrequenz und ihre technische Bedeutung*. Vieweg, 1918.
- John Guckenheimer and Philip Holmes. *Nonlinear oscillations, dynamical systems, and bifurcations of vector fields*, volume 42. Springer Science & Business Media, 2013.
- W Szemplińska-Stupnicka and J Rudowski. The coexistence of periodic, almost-periodic and chaotic attractors in the van der pol-duffing oscillator. *Journal of Sound and Vibration*, 199(2):165–175, 1997.

- Agnieszka Chudzik, Przemyslaw Perlikowski, Andrzej Stefanski, and Tomasz Kapitaniak. Multistability and rare attractors in van der pol–duffing oscillator. *International Journal of Bifurcation and Chaos*, 21(07):1907–1912, 2011.
- Katherine Bold, Chantal Edwards, John Guckenheimer, Sabyasachi Guharay, Kathleen Hoffman, Judith Hubbard, Ricardo Oliva, and Warren Weckesser. The forced van der pol equation ii: Canards in the reduced system. *SIAM Journal on Applied Dynamical Systems*, 2(4):570–608, 2003.
- Nikolay A Kudryashov. The generalized duffing oscillator. *Communications in Nonlinear Science and Numerical Simulation*, 93:105526, 2021.
- Peter H Baxendale. Stochastic averaging and asymptotic behavior of the stochastic duffing–van der pol equation. *Stochastic Processes and Their Applications*, 113(2):235–272, 2004.
- Ludwig Arnold and Rafail Z Khas’ minskii. *Stability index for nonlinear stochastic differential equations*. Universität Bremen. Fachbereiche Mathematik/Informatik, Elektrotechnik/Physik. Forschungsschwerpunkt Dynamische Systeme, 1993.
- Ludwig Arnold, N Sri Namachchivaya, and Klaus R Schenk-Hoppé. Toward an understanding of stochastic hopf bifurcation: a case study. *International Journal of Bifurcation and Chaos*, 6(11):1947–1975, 1996.
- Hannes Keller and Gunter Ochs. Numerical approximation of random attractors. *Stochastic dynamics*, pages 93–115, 1999.
- Klaus Reiner Schenk-Hoppé. Bifurcation scenarios of the noisy duffing-van der pol oscillator. *Nonlinear dynamics*, 11(3):255–274, 1996a.
- Yajie Li, Yongtao Sun, Ying Hao, Xiangyun Zhang, Feng Wang, Heping Shi, and Bin Wang. Stochastic stability of the fractional and tri-stable van der vol oscillator with time-delay feedback driven by gaussian white noise. *Thermal Science*, 27(3 Part A):2155–2164, 2023.
- Kaio CB Benedetti, Paulo Batista Gonçalves, Stefano Lenci, and Giuseppe Rega. Global analysis of stochastic nonlinear dynamical systems. part 2: Influence of uncertainties and noise on basins/attractors topology and integrity. *International Journal of Non-Linear Mechanics*, 159:104594, 11 2022.
- Klaus Reiner Schenk-Hoppé. Deterministic and stochastic duffing-van der pol oscillators are non-explosive. *Zeitschrift für angewandte Mathematik und Physik ZAMP*, 47(5):740–759, 1996b.
- Grigori N Milstein and Michael V Tretyakov. *Stochastic numerics for mathematical physics*, volume 39. Springer, 2004.
- Desmond J Higham, Xuerong Mao, and Andrew M Stuart. Strong convergence of euler-type methods for nonlinear stochastic differential equations. *SIAM journal on numerical analysis*, 40(3):1041–1063, 2002.
- Annika Lang and Ioanna Motschan-Armen. Euler-maruyama approximations of the stochastic heat equation on the sphere. *arXiv preprint arXiv:2307.07564*, 2023.
- Zhen Wang, Yu Miao, et al. Euler-maruyama scheme for sdes with dini continuous coefficients. *arXiv preprint arXiv:2009.12791*, 2020.
- Diancong Jin. Central limit theorem for temporal average of backward euler–maruyama method. *arXiv preprint arXiv:2307.04181*, 2023.
- Kamal Hiderah. The truncated euler–maruyama method of one-dimensional stochastic differential equations involving the local time at point zero. *Random Operators and Stochastic Equations*, 31(2):141–152, 2023.