CP Violation Problem

Nicolai Popov,^{1,*} William J. Briscoe,^{2,†} and Igor Strakovsky^{2,‡}

¹Sektion Physik der Ludwig-Maximilians-Universität, 4 Schellingstrasse, 80799 München, Germany ²Institute for Nuclear Studies, Department of Physics, The George Washington University, Washington, DC 20052, USA

(Dated: May 1, 2024)

The strong CP violation problem has a long history emanating from its discovery 60 years ago in the decay of neutral kaons and the subsequent experimental and theoretical studies over several decades. We review herein experimental data that observe indirect CP violation of the order of ~ 10^{-3} , as well as the discovery of direct CP violation of the order of ~ 10^{-6} . The verification of CP violation in the decay of charged kaons was also observed. Data reflecting CP violation in the decays of B and D mesons have become very important and are also discussed in this review.

I. INTRODUCTION

New particles called K-mesons, or kaons, were discovered in cosmic rays in the late 1940s and continued to be studied intensively at charged-particle accelerators beginning in 1954. These are K^{\pm} (which are a particle/antiparticle pair) and a K^0 meson which is its own antiparticle. They have zero spin and a mass approximately 970 times that of an electron. Participating in strong interactions, they are hadrons, but with zero baryon charge, also having a new quantum number, strangeness S, namely S = 1 for K^0 and S = -1 for \bar{K}^0 , characterizing their behavior in strong interaction processes. Until 1956, the law of conservation of parity was considered fundamental as was the law of conservation of energy and momentum. However, Lee and Yang's suggestion of spatial parity non-conservation in weak interactions [1] was confirmed in 1957 by Wu *et al.* in studying the beta decay of Co⁶⁰ nuclei at NBS (now NIST) [2].

^{*} science.popov@googlemail.com

[†] briscoe@gwu.edu

[‡] igor@gwu.edu

An important result of the study of K^0 mesons was the discovery of CP parity violation, or T parity violation, according to the CPT theorem [3, 4]. Here, P is a spatial inversion - parity, that is, a mirror image of the coordinates, replacing the with opposite ones for polar vectors $\vec{r} \to -\vec{r}$, as well as momentum $\vec{p} \to -\vec{p}$, while for pseudovectors, angular momentum and spin, do not change. C – charge conjugation parity, with the replacement of particles by antiparticles, *i.e.*, a change in the electric charge, hyper-charge, baryon and lepton charges and, as a consequence, a change in the sign of the magnetic moment. In this case, the spatial coordinates, momentum and spin of the particle do not change. T – time reversal, $t \to -t$, that is, reversibility of movement, with a change in the signs of momentum and spins. These parities are conserved in electromagnetic and strong interactions, but are violated in the weak interaction.

II. CP PARITY

Neither K^0 nor \overline{K}^0 has a certain CP parity, but do have negative spatial parity P. The action of the operators C, P, and CP on the wave functions K^0 and \overline{K}^0 can be written in the form:

$$C|K^{0}\rangle = -|\bar{K}^{0}\rangle, \qquad C|\bar{K}^{0}\rangle = -|K^{0}\rangle,$$
$$P|K^{0}\rangle = -|\bar{K}^{0}\rangle, \qquad P|\bar{K}^{0}\rangle = -|\bar{K}^{0}\rangle,$$
$$CP|K^{0}\rangle = -|\bar{K}^{0}\rangle, \qquad CP|\bar{K}^{0}\rangle = -|K^{0}\rangle,$$

where it is possible that antiparticles K^0 and \bar{K}^0 are transitioning from one to the other. Thus, from the states K^0 and \bar{K}^0 it is possible to construct linear combinations having definite CP parity values. These are

$$|K_1^0\rangle = \sqrt{\frac{1}{2}} \left(|K^0\rangle + |\bar{K}^0\rangle \right),\,$$

with CP parity +1, which can decay into two pions, and the orthogonal superposition

$$|K_2^0\rangle = \sqrt{\frac{1}{2}} \left(|K^0\rangle - |\bar{K}^0\rangle \right),$$

has CP parity -1, that can decay into two pions only as a result of a CP violating process such as the weak interaction. K_1^0 and K_2^0 are not a particle and an antiparticle and therefore may have different decay channels. Thus, the CP parities of K_1^0 and K_2^0 are:

$$CP|K_1^0\rangle = +|K_1^0\rangle$$
,
 $CP|K_2^0\rangle = -|K_2^0\rangle$.

Let us consider the conservation of combined CP-parity in the decays of K-mesons into 2 and 3 pions. 2π and 3π systems with orbital momentum l = 0 are eigenstates of the CPoperator. The operator P is equivalent to replacing $\pi^+\pi^-$ with the acquisition of the wave function of the multiplier $(-1)^l$.

$$P|\pi^{+}\pi^{-}\rangle = P|\pi^{+}\rangle P|\pi^{-}\rangle (-1)^{l} = +|\pi^{+}\pi^{-}\rangle.$$

The operator C turns π^+ into π^- and π^- into π^+ , that is, it is also equivalent to replacement $\pi^+\pi^-$ mesons

$$C|\pi^{+}\pi^{-}\rangle = (-1)^{l} |\pi^{+}\pi^{-}\rangle = +|\pi^{+}\pi^{-}\rangle$$

for l = 0. And the eigenvalue of the CP operator of the 2π system is equal to +1:

$$CP|\pi^+\pi^-\rangle = (-1)^{2l} |\pi^+\pi^-\rangle = +|\pi^+\pi^-\rangle.$$

The CP parity of a system of three pion state depends on the orbital state \vec{L} of the 3π which is the sum of the orbital momentum vectors \vec{l} of $\pi^+\pi^-$ pair and the orbital momentum $\vec{l'}$ of π^0 relative to the center of mass of this pion pair, that is, $\vec{L} = \vec{l} + \vec{l'} = 0$ ($\vec{l'} = -\vec{l}$).

$$P|\pi^{+}\pi^{-}\pi^{0}\rangle = P|\pi^{+}\rangle P|\pi^{-}\rangle P|\pi^{0}\rangle (-1)^{L} = (-1)|\pi^{+}\pi^{-}\pi^{0}\rangle,$$
$$C|\pi^{+}\pi^{-}\pi^{0}\rangle = |\pi^{+}\pi^{-}\pi^{0}\rangle (-1)^{L} = (+1)|\pi^{+}\pi^{-}\pi^{0}\rangle,$$
$$CP|\pi^{+}\pi^{-}\pi^{0}\rangle = (-1)|\pi^{+}\pi^{-}\pi^{0}\rangle$$

for L = 0. In the result, the eigenvalue of the CP operator of the 3π system is equal to -1. Since 2π and 3π systems are eigenvalues of the CP operator at zero orbital momentum, we assume in above relations that this momentum is equal zero.

The states K_1^0 and K_2^0 have certain CP parity values, without, however, having a certain strangeness. We can write down:

$$|K^{0}\rangle = \sqrt{\frac{1}{2}} \left(|K_{1}^{0}\rangle + |K_{2}^{0}\rangle \right),$$

$$\left|\bar{K}^{0}\right\rangle = \sqrt{\frac{1}{2}} \left(\left|K_{1}^{0}\right\rangle - \left|K_{2}^{0}\right\rangle\right).$$

That is K^0 and \bar{K}^0 are a superposition of the states K_1^0 and K_2^0 . Since $CP(K_1^0) = +1$, while conservation the combined parity, the average lifetime of which splits into 2π and $\tau(K_1^0) \approx 0.9 \times 10^{-10}$ s. Since $CP(K_2^0) = -1$ and K_2^0 break down into 3π , with a useful life $\tau(K_2^0) \approx 5 \times 10^{-8}$ s due to the smaller phase volume of the decay products. Due to the low energy release in 3π decays, their probabilities are approximately three orders of magnitude lower than the probabilities of 2π decays. Since 50% of the K^0 meson consists of the K_1^0 component, its decays into 2π are observed near the target. At a greater distance from the target, decays of the K_2^0 component are observed in 3π mesons.

The interconversion of K^0 and \bar{K}^0 mesons in a vacuum is their oscillation $K^0 \leftrightarrow \bar{K}^0$. Being a process with a change in strangeness $\Delta S = 2$, since S = 1 for the K^0 meson and S = -1 for the \bar{K}^0 meson. It occurs because of weak interactions between the quarks, from which K^0 mesons come together: $K^0 = \bar{s}d \leftrightarrow s\bar{d} = \bar{K}^0$. As a result of these transformations, K^0 and \bar{K}^0 do not have a certain mass and a certain lifetime. States with a certain mass and a certain lifetime are short-lived K^0_S and long-lived K^0_L mesons.

The difference between the masses of K_L^0 and K_S^0 is due to the weak interaction causing $K^0 \leftrightarrow \bar{K}^0$ transitions, and is very small [5]:

$$\Delta m_K = m(K_L^0) - m(K_S^0) = (3.510 \pm 0.018) \times 10^{-12} \text{ MeV}.$$

In the presence of CP invariance, we can consider: $|K_S^0\rangle \equiv |K_1^0\rangle$ and $|K_L^0\rangle \equiv |K_2^0\rangle$.

Let at the initial moment t = 0 there are pure K^0 mesons in vacuum, obtained, for example, in the reaction $\pi^- p \to K^0 \Lambda$ in the absence of \bar{K}^0 mesons. Then after time t there will be the following beam:

$$\frac{1}{2} \left[(K^0 + \bar{K}^0) \ e^{-i(m_1 - \Gamma_1/2)t} + (K^0 - \bar{K}^0) \ e^{-i(m_2 - \Gamma_2/2)t} \right]$$

If K_1^0 and K_2^0 are stable, $\Gamma_1 = \Gamma_2 = 0$, the probabilities of finding

$$P_{K^{0}}(t) = \frac{1}{4} |e^{-im_{1}t} + e^{-im_{2}t}|^{2} = \cos^{2}\left(\frac{\Delta m_{K}}{2}t\right)$$
$$P_{\bar{K}^{0}}(t) = \frac{1}{4} |e^{-im_{1}t} - e^{-im_{2}t}|^{2} = \sin^{2}\left(\frac{\Delta m_{K}}{2}t\right)$$

and after time $\tau = \pi (m_2 - m_1) = \pi / \Delta m_K$ (oscillation period) there would be no K^0 mesons in beam, since:

$$P_{K^0}(\tau) = \cos^2\left(\frac{\Delta m_K}{2}\tau\right) = \cos^2\frac{\pi}{2} = 0,$$

$$P_{\bar{K}^0}(\tau) = \sin^2\left(\frac{\Delta m_K}{2}\tau\right) = \sin^2\frac{\pi}{2} = 1.$$

Such oscillations, which are superimposed by exponential decay due to $\Gamma_{1,2} \neq 0$, are observed experimentally. It is possible to observe them, for example, in semileptonic decays of neutral kaons that satisfy the rule $\Delta Q = \Delta S$. This rule is due to the fact that during the transition $s \rightarrow u$ the strangeness and electric charge of the quark increase by one.

$$K^{0}\left(\frac{d}{\bar{s}}\right) \to \pi^{-}\left(\frac{d}{\bar{u}}\right) l^{+}\nu_{l} , \quad K^{0}\left(\frac{d}{\bar{s}}\right) \nrightarrow \pi^{+}\left(\frac{u}{\bar{d}}\right) l^{-}\bar{\nu}_{l} ,$$

$$\bar{K}^{0}\left(\frac{s}{\bar{d}}\right) \to \pi^{+}\left(\frac{u}{\bar{d}}\right) l^{-}\bar{\nu}_{l} , \quad \bar{K}^{0}\left(\frac{s}{\bar{d}}\right) \nrightarrow \pi^{-}\left(\frac{d}{\bar{u}}\right) l^{+}\nu_{l} .$$

Here $l \equiv e, \mu$. The sign of the charged lepton indicates the strangeness of K^0 or \bar{K}^0 . Oscillations can be measured by recording the number of electrons and positrons arising from K_{e3} decays. Such experiments make it possible to determine Δm_K . Reactions with \neq are prohibited by the rule $\Delta Q = \Delta S$ and have not been observed experimentally.

The K^0 -meson, arising as a result of the strong interaction, at some distance from the point of birth is practically transformed due to the weak interaction into \bar{K}^0 and therefore turns out to be capable of causing nuclear reactions, for example, $\bar{K}^0 p \to \Lambda \pi^+$ characteristic of \bar{K}^0 and prohibited for K^0 due to the preservation of strangeness in strong interaction. In 1955, Pais and Piccioni predicted oscillations in a beam of neutral K mesons within the framework of the regeneration effect [6]. Regeneration consisted of the use of an absorber, positioned at a distance determined by τ_S and τ_L to demonstrate the superposition of K^0 and \bar{K}^0 . Let us take a pure beam K^0 in vacuum (equal of parts K_S^0 and K_L^0). After, *e.g.*, $t \approx 10 \tau_S$, the K_S^0 intensity down by factor of $e^{-t/\tau_S} = e^{-10} \approx 4.5 \times 10^{-5}$ (0.5 m for 1 GeV kaon) The 6.2 GeV proton synchrotron at Lawrence Berkeley National Laboratory (LBNL) with a π^- beam of 1.1 GeV was used for the experiment.

A beam of neutral K-mesons decays in flight so that short-lived K_S^0 disappears, leaving a beam of long-lived pure K_L^0 . If this beam passes through matter, K^0 and \bar{K}^0 interact with nuclei in different ways. With K^0 , quasi-elastic scattering on nucleons occurs, while \bar{K}^0 can produce hyperons. Due to the different interactions of these components, the quantum coherence between the two particles is lost. The resulting beam contains various linear superpositions of K^0 and \bar{K}^0 . Each superposition is a mixture of K_S^0 and K_L^0 , so that K_S^0 is restored when a beam of K-mesons passes through the substance of the regenerator , as was observed in the above experiment. Anomalous regeneration of K_S^0 was obtained in Ref. [7]. The problem of time reversal in quantum mechanics has been discussed since the thirties [8]. In 1964, Christenson, Cronin, Fitch, and Turlay discovered CP violation in an experiment at the Brookhaven National Laboratory (BNL) AGS accelerator [9] through the decay of the neutral K_L^0 meson to 2π mesons in spite of the main decay channel to 3π . In fact, there are two K-mesons: $K_1^0 \to \pi^+\pi^-$ with CP = +1 (while CP = +1 for the state), that is,, the even state, and $K_2^0 \to \pi^+\pi^-\pi^0$ with CP = -1, *i.e.*, the odd state.

The long-lived K_L^0 has a probability of approximately 0.2% decaying into 2π with CP parity violation, which is small, but is, however, an important discovery for fundamental physics [5]:

$$\frac{\Gamma(K_L^0 \to \pi^+ \pi^-)}{\Gamma(K_L^0 \to all)} = (2.03 \pm 0.04) \times 10^{-3} \,.$$

CP violation in mixtures of neutral mesons $K^0 \leftrightarrow \bar{K}^0$ is an indirect violation of CP parity, in contrast to direct violation at different decay rates of a particles and antiparticles and interference between direct and indirect violation of CP parity. There is no strict identification of K_S^0 with K_1^0 and K_L^0 with K_2^0 . Instead, the states K_L^0 and K_S^0 are defined as follows:

$$|K_L^0\rangle = \left(1 + |\epsilon|^2\right)^{-1/2} \left(|K_2^0\rangle + \epsilon |K_1^0\rangle\right) = \left[2\left(1 + |\epsilon|^2\right)\right]^{-1/2} \left[(1 + \epsilon)|K^0\rangle - (1 - \epsilon)|\bar{K}^0\rangle\right],$$

$$|K_S^0\rangle = \left(1 + |\epsilon|^2\right)^{-1/2} \left(|K_1^0\rangle + \epsilon |K_2^0\rangle\right) = \left[2\left(1 + |\epsilon|^2\right)\right]^{-1/2} \left[(1 + \epsilon)|K^0\rangle + (1 - \epsilon)|\bar{K}^0\rangle\right].$$

The laws of Nature would be the same for matter and for antimatter. Most phenomena are C- and P-symmetric, and also CP-symmetric for electromagnetic, strong, and gravitational interactions. The weak interactions violate the C - and P-symmetric as well as some CP-symmetry of unknown nature with the supposition of superweak interaction. At the same time, the Universe is made chiefly of matter, rather than consisting of equal parts of matter and antimatter as might be expected.

III. MATTER-ANTIMATTER PROBLEM

In 1967, Sakharov showed that CP violation was one of the necessary conditions for the almost complete destruction of antimatter at the early stage of the development of the Universe [10, 11]. It can be shown that to create this imbalance between matter and antimatter, the Sakharov conditions must be satisfied, namely, baryon number violation, C symmetry and CP symmetry violated, and the interaction is outside what is allowed by thermal equilibrium. Baryon number violation is a necessary condition for producing an excess of baryons over antibaryons. One of its conditions is the violation of CP symmetry in the extreme conditions of the first seconds after the Big Bang. Since our Universe is not a sea of photons without other matter, after the Big Bang different physical laws were in effect for matter and antimatter: that is, CP symmetry was violated. However, breaking Csymmetry is also necessary so that interactions that produce more baryons than antibaryons are not balanced by interactions that produce more antibaryons than baryons. Breaking of CP symmetry is also necessary, since otherwise an equal number of left baryons and right antibaryons would be produced, as well as an equal number of right baryons and left antibaryons. The interactions should be outside the limitations of thermal equilibrium, otherwise, CPT symmetry would compensate for the processes of decreasing or increasing the baryon number.

IV. CP VIOLATION IN STANDARD MODEL

Possible CP violation is associated with complex weak coupling constants owing to the CP-odd interaction of participating particles. The CP violation appears naturally in united gauge theories with spontaneous symmetry breaking. This mechanism is possible through the complex vacuum expectation value of the Higgs field (Fig. 1). The simplest model of CP breaking through the Higgs field coupling results in a complex phase in the Cabibbo-Kobayashi-Maskawa mass matrix (CKM) [12, 13].

The interaction of quarks in the Standard Model with charged gauge bosons W^{\pm} is described by:

$$L = -\frac{g}{2\sqrt{2}} \bar{u}_i \gamma^{\mu} (1 - \gamma_5) V_{ij} d_j W^{\pm}_{\mu} + h.c. ,$$

where g is the weak interaction constant, u_i and d_j – the fields of quark states of three generations, W^{\pm} – the fields of vector charged bosons, V_{ij} is the quark mixing matrix, that is, the CKM matrix:

$$V = \begin{pmatrix} c_{12}c_{13} & s_{12}c_{13} & s_{13}e^{-i\delta} \\ -s_{12}c_{23} - c_{12}s_{23}s_{13}e^{i\delta} & c_{12}c_{23} - s_{12}s_{23}s_{13}e^{i\delta} & s_{23}c_{13} \\ s_{12}s_{23} - c_{12}c_{23}s_{13}e^{i\delta} & -c_{12}s_{23} - s_{12}c_{23}s_{13}e^{i\delta} & c_{23}c_{13} \end{pmatrix}$$



FIG. 1. The Standard Model of elementary particles, taken from Ref. [5].

here $c_{ij} = \cos \theta_{ij}$, $s_{ij} = \sin \theta_{ij}$ and one phase δ in the interval $[0, 2\pi]$ [5]:

$$\theta_{12} = (13.04 \pm 0.05)^{\circ} ,$$

$$\theta_{13} = (0.201 \pm 0.011)^{\circ} ,$$

$$\theta_{23} = (2.38 \pm 0.06)^{\circ} ,$$

$$\delta = (0.201 \pm 0.011)^{\circ} .$$

In the case of two generations of quarks, the mixing matrix is parameterized by a single Cabibbo angle $\theta_c \approx 13^{\circ}$:

$$V = \begin{pmatrix} \cos \theta_c & \sin \theta_c \\ -\sin \theta_c & \cos \theta_c \end{pmatrix}$$

which cannot provide a mechanism for CP violation. The presence of a third generation of quarks, for example, the t quark, is a necessary condition for violation of the CP parity.

Within the framework of the Standard Model using the CMK-matrix, calculations of CP violating effects were carried out in the kaon system as well as in systems with B and D

mesons, where these effects are greater and were also discovered experimentally. Violation of the CP symmetry in decays of B mesons was observed in 2001 in the BaBar experiment at SLAC and in experiments at the KEK facility.

As for baryon number, there is no experimental evidence of the violation of this number. The violation of such conservation in the Standard Model is possible outside the framework of perturbation theory. Baryon number breaking is possible within the framework of supersymmetry or Grand Unification Theory (GUT).



V. DISCOVERY OF INDIRECT CP VIOLATION

FIG. 2. Plan view of the detector arrangement, taken from Ref. [9].

Figure 2, taken from Ref. [9], presents a general overview of the detector system of the experiment performed at Brookhave National Laboratory. Spark chambers were located in each arm of a two-arm spectrometer; these record the trajectories of particles before and after the deflecting magnet. Cherenkov scintillation counters triggered the spark chambers when a coincidence was detected and the resulting tracks were photographed and recorded on film. The detector system was installed in the beam of K^0 meson beam line at the Brookhaven synchrotron at a distance such that the K_1^0 mesons had already decayed so that only K_2^0 mesons remained in the beam. The angle between the arms was chosen for

optimal determination of the kaons that decay into two charged pions. Helium gas filled the volume where this decay occurred to minimize the detection of three-particle decays of K_2^0 mesons through rescattering. In this case, the vector sum of the momenta of the two detected particles must be oriented in the direction of the beam of K_2^0 mesons, and the missing mass of the decay particle, calculated from the data on the decay products, must coincide with the mass of the K-meson. 56 desired events were recorded with a background of 11 events. From these data it follows that the fraction of the decay of K_2^0 into two pions in relation to all decay modes involving charged particles is equal to 2×10^{-3} . This was the first evidence of CP violation of parity in the kaon system. Almost six months were spent searching for an alternative CP violation, but to no avail. Thus their conclusion that they obtained indirect CP violation (see Fig. 2, taken from Ref. [9]) in the mixing of neutral kaon states ($K^0 \leftrightarrow \bar{K}^0$).

VI. DIRECT AND INDIRECT CP VIOLATION

In the agreement with Wolfenstein superposition [14], postulated in 1964, any superweak interaction could exist. A direct consequence of this manifests itself only in the mixing $\Delta S = 2$ between K^0 and \bar{K}^0 with the absence in $\Delta S = 1$. A consequence of this assumption could be correlations between K^0 decay amplitudes. Let

$$\eta^{+-} = \frac{A(K_L^0 \to \pi^+ \pi^-)}{A(K_S^0 \to \pi^+ \pi^-)} , \quad \eta^{00} = \frac{A(K_L^0 \to \pi^0 \pi^0)}{A(K_S^0 \to \pi^0 \pi^0)} .$$

In this case the decay amplitude of the CP-odd combination K_2^0 combination to a 2π final state might differ from zero, corresponding to a direct CP violation ϵ . However, $\eta^{+-} = \eta^{00} = \epsilon$ in the case of indirect CP violation. There are two decay amplitudes, one is the I = 0 and the second in the I = 2 states, the and final states. As the Clebsch-Gordan coefficients projecting, the I = 0 and I = 2 states for the $\pi^+\pi^-$ and $2\pi^0$ final states are different, we have:

$$\epsilon' = i\sqrt{1/2} \ Im\left(\frac{A_2}{A_0}\right) \ e^{i(\delta_2 - \delta_0)}$$

here $A_{0,2}$ and $\delta_{0,2}$ are the amplitudes and the strong phases of the 2π final isospin states I = 0, 2. Then, there exists direct CP violation, determined by complex parameter ϵ' , which states that there is possible $K_2^0 \to 2\pi$ in the absence above equality, *i.e.*:

$$\eta^{+-} = \epsilon + \epsilon' , \quad \eta^{00} = \epsilon - 2\epsilon' ,$$

Modern experimental data, taken from Refs. [5], are the following:

 $\epsilon = (2.228 \pm 0.011) \times 10^{-3},$ $\epsilon + \epsilon' = (2.232 \pm 0.011) \times 10^{-3},$ $\epsilon - 2\epsilon' = (2.221 \pm 0.011) \times 10^{-3},$ $Re(\epsilon'/\epsilon) = (1.65 \pm 0.26) \times 10^{-3},$ $\phi_{+-} = (43.51 \pm 0.05)^{\circ},$ $\phi_{00} = (43.7 \pm 0.8)^{\circ}.$

These phases are for $\eta^{+-} = |\eta^{+-}|e^{i\phi_{+-}}$ and $\eta^{00} = |\eta^{00}|e^{i\phi_{00}}$. An estimate of the phases ϕ_{+-} and ϕ_{00} was given by Bell and Steinberger in 1965 [15] indicating their equality to the phase ϕ_{ϵ} and taking into account the CPT theorem with the above-mentioned Δm_K and $\Gamma_S = 1.11 \times 10^{10} \text{ s}^{-1}$ they are equal $\phi_{\epsilon} = 43.5^{\circ}$ in agreement with the experimental values mentioned above. Experimental confirmation of the approximate equality $\eta_{+-} \approx \eta_{00}$ was obtained in the 1970s [16–18].

$$Re(\epsilon'/\epsilon) = \frac{1}{6}(1-|\frac{\eta^{00}}{\eta^{+-}}|^2) = \frac{1}{6}(1-R)$$

where R is the so-called double ratio:

$$R = \frac{\Gamma(K_L^0 \to \pi^0 \pi^0)}{\Gamma(K_S^0 \to \pi^0 \pi^0)} / \frac{\Gamma(K_L^0 \to \pi^+ \pi^-)}{\Gamma(K_S^0 \to \pi^+ \pi^-)} \,.$$

After discovery of indirect CP violation [9], further experiments were carried for search direct CP violation (see Fig. 3). There are two possible manners of violation of the CP. In the first, the long-lived kaon is a mixture of two quantum states, mainly CP-odd with just a small amount of CP-even and in the second, CP violation in decay of K^0 when the actual quark reactions underlying the particle transformations. In the decay of neutral kaon, a strange quark disappears with creation of pions composed of only up and down quarks. CP violation via this route is called direct one.



FIG. 3. The indirect and direct modes of CP violation.

VII. CERN AND FERMILAB EXPERIMENTS

In the NA31 experiment performed at CERN [19, 20], the decay channels of the neutral and charged kaons were simultaneously recorded both near the dipping volume and far from it (K_L^0) . This made it possible to avoid the background from the interaction of neutrons in the regenerator and to minimize the difference in the geometries of the decay lengths of which differ by approximately three orders of magnitude. The energies of neutral and charged particles were measured by two calorimeters filled with liquid argon. A detailed description of the detectors in this experiment is given in Refs. [19, 20]. Short-lived and long-lived kaons were born in collisions of protons of the 450 GeV Super Proton Synchrotron (SPS) with two targets, one of which $(K_L^0 \text{ target})$ was located at a large distance (about 100 m) from decay volume, and the other $(K_S^0 \text{ target})$ close to it. The trigger for launching the selection process to isolate decays into charged pions was triggered on the basis of data from the scintillation hodoscope located between the second wire chamber and the electromagnetic calorimeter. The trigger for isolating decays into neutral pions was launching by a hodoscope placed in liquid argon. The main problems of this experiment were related to the lack of a spectrometer to measure charged tracks. In this case, about 40% of the decays to 2π were lost.

A characteristic of another experiment E731 (Fermilab) [21, 22] was the simultaneous use of two almost parallel K_L^0 beams, one of which, before the decay volume, passed through a B_4C regenerator, where (due to the difference in the scattering of the nuclei K^0 and \bar{K}^0) coherent regeneration of the K_L^0 component occurred. A magnetic spectrometer was used to record tracks of charged particles. Neutral decays were recorded by an electromagnetic calorimeter containing 804 glass cells. Photomultipliers recorded Cherenkov light from electrons and positrons generated in glass by photons or high-energy electrons. The trigger of charged decays was launched as a result of coincidences of signals in two opposite quadrants of the scintillation hodoscope. Compared with the NA31 experiment, it is clear that in the E731 experiment, the background from $K_L^0 \rightarrow 3\pi^0$ in the neutral decay mode is significantly lower due to the closer location of the decay volume to the detectors. There is also less background in charged mode from a detector with better resolution in measuring the momentum of charged particles. At the same time, the result of the E731 experiment strongly depended on the correct accounting of the efficiency of decay detection performed by Monte Carlo simulation. The results of these experiments are the following: NA31 (CERN) $Re(\epsilon'/\epsilon) = (2.30 \pm 0.65) \times 10^{-3}$ and E731 $Re(\epsilon'/\epsilon) = (0.74 \pm 0.61) \times 10^{-3}$. From a comparison of these experiments it is clear that they do not agree with each other (with a confidence level of less than 10%). Experiment NA31 indicates the possibility of direct CP violation $\approx 10^{-3}$, while the result of experiment E731 is compatible with zero. Therefore, there was no clear conclusion about a direct violation of the CP invariance at that time, and further more accurate measurements were required.

The theoretical calculations obtained for ϵ'/ϵ up to the 80s ranged from 0.002 to 0.02 [23–

25]. Theoretical consideration of this problem continued in the 1990s with a decrease in the calculated ϵ'/ϵ to 10^{-4} [26–28] with an indication of the reduction of the so-called loop "penguin" diagrams [29]. However, these diagrams are highly sensitive to possible contributions from unknown particles and therefore provide excellent probes for new sources of CP violation. The known source of CP violation, which is governed by the CKM matrix in the quark sector, is insufficient to account for the huge excess of matter over antimatter in the Universe; new sources of CP violation are necessary to resolve this problem.

The NA48 experiment was significantly improved compared to the previous NA31 experiment with the participation of new institutions in the collaboration. The aim was to increase the precision to obtain ϵ'/ϵ to 10^{-4} . Significant changes were made to the beam system and the detector. The NA48 experiment at CERN [30, 31] had made separate runs with long-lived and short-lived kaon, K_L^0 and K_S^0 , produced in two different targets by protons from the same CERN SPS beam. All four modes of decays in beams K_L^0 and K_S^0 to reduce systematic uncertainties in the measurement of decay probabilities were output into the same decay volume simultaneously and almost collinearly. The K_L^0 beam is produced by a ten times more 450 GeV proton beam than for NA31, colliding with a beryllium target. The protons for the K_S^0 beam are derived from the free protons in the K_L^0 target colliding with a silicon mono-crystal. Unlike NA31, a magnetic spectrometer was used for the measurement of charged decays with better accuracy. Although the generation of decays $K_L^0 \rightarrow 2\pi$ required an intense initial beam of protons, a much less intense beam of protons was sufficient to create the beam K_S^0 when creating two-pion decays, which is approximately 5 orders of magnitude lower than the intensity of the beam for generation K_L^0 , which approximately corresponds to the square of the indirect violation parameter $|\eta^{+-}|$.

The simultaneous generation of two kaon beams ensured their equal sensitivity to variations in the intensity of the primary proton beam within 10%. Free protons (without their interaction with the target K_L^0) were directed to a curved silicon crystal below the beam axis K_L^0 . This was a small fraction of protons ($\approx 10^{-5}$) due to its collision near the decay volume with the target K_S^0 . The two-pion decays of this beam were decays due to the large difference in the lifetimes K_L^0 and K_S^0 and the low probability of such decays for K_L^0 . The decay volume was 6 m from the target K_S^0 . Here are the beams K_S^0 and K_L^0 exited from the two apertures of the last collimator into a general decay region. To identify decays in the beam K_S^0 a "beam tagging station" consisting of different scintillators was used. The triggers for recording decays $K^0 \to 2\pi^0$ worked based on the analog summation of signals from the group of cells of a liquid krypton calorimeter.

The world average result is the following [5, 32]:

$$\begin{aligned} Re(\epsilon'/\epsilon) &= (1.68 \pm 0.20) \times 10^{-3} ,\\ \frac{\Gamma(K^0 \to \pi^+ \pi^-) - \Gamma(\bar{K}^0 \to \pi^+ \pi^-)}{\Gamma(\bar{K}^0 \to \pi^+ \pi^-) + \Gamma(\bar{K}^0 \to \pi^+ \pi^-)} &= 2Re(\epsilon') = (5.3 \pm 0.6) \times 10^{-6} ,\\ \frac{\Gamma(K^0 \to \pi^0 \pi^0) - \Gamma(\bar{K}^0 \to \pi^0 \pi^0)}{\Gamma(\bar{K}^0 \to \pi^0 \pi^0) + \Gamma(\bar{K}^0 \to \pi^0 \pi^0)} &= -4Re(\epsilon') = (-10.6 \pm 1.02) \times 10^{-6} . \end{aligned}$$

Let us summarize the above. The NA48 experiment at CERN had made separate runs with long-lived and short-lived kaons. Both experiments in CERN used simultaneous beams of long- and short-lived kaons and taking data on charged and neutral pion production by both kaon beams at the same time. Installed in CERN's highest intensity proton beamline it uses a large and sophisticated detector. Some of the CERN protons go to make long-lived kaons, and the remaining particles are bent by a crystal and used to make a parallel beam of short-lived kaons. In this way the two varieties of kaons are clearly differentiated, even though they eventually decay in the same way.

A detailed description of the selection of events in this and previous experiments is given in [32, 33]. These reviews describe the beams, detectors, and analysis methods used in CERN experiments, which made significant contributions to their results and established new standards for precision measurements.

The final measurements of ϵ'/ϵ made by the most precise experiments, KTEV, NA48, NA31, and E731, together with the average value are given in Fig. 4, taken from [5]. In contrast to the significant improvements of the NA48 experiment over the previous NA31 experiment at CERN, the KTeV experiment [34, 35] followed the approach of the previous E731 experiment, also at Fermilab. Both beams were formed by the collision of protons with an energy of 800 MeV with a common target. The magnets removed charged particles from the beams. The common absorber increased the proportion of neutral kaons relative to neutrons in neutral beams. A plastic regenerator that promotes the formation of K_S^0 in one of the beams was moved to block the other neutral beam. The beams diverged from each other and were completely separated in the decay volume. The spectrometer, consisting of four drift chambers, including sensitive wires and a large dipole magnet, was filled with helium. The electromagnetic calorimeter consisted of 3100 cells of cesium iodine



FIG. 4. The comparison of the results of different experiments, taken from Ref. [5].

(CsI) crystals. The trigger system included scintillation hodoscope signals for decays into charged pions and summation of the energy release in the calorimeter during decays into neutral pions. The analysis of selected events included fitting the obtained data by modeling the Monte Carlo method.

Using a subset of data collected in 1996–1997, they found [36, 37]:

$$Re(\epsilon'/\epsilon) = (28.0 \pm 4.1) \times 10^{-4}$$

The final result of the KTeV experiment is it follows [38]:

$$Re(\epsilon'/\epsilon) = (19.2 \pm 2.1) \times 10^{-4}$$

A detailed description of the experiments NA31 and NA48 at CERN and KTeV at Fermilab is given in Refs. [32, 33]. The data on ϵ'/ϵ obtained in these experiments made it possible to unconditionally establish for the first time the presence of direct CP violation in the decays of neutral kaons. Given the most accurate result of the NA48 experiment (with first indicating of direct CP violation in experiment NA31), the contribution of other experiments is extremely important for a complete picture of the discovered phenomenon. Detailed reviews are devoted to the violation of CP in neutral kaon decays [11, 32, 33, 39–42].

Along with obtaining new experimental results, theoretical research into the problem continued. The topics of Lorentz and CPT violation was discussed in Refs. [43–45]. They referenced the possibility of spontaneous CPT breaking that arises in string theory considering the standard model extension (SME) taking into account this effect in the neutral meson system [43, 44]. However, Lorenz symmetry breaking, although leading to the CPT violating terms, cannot lead to the CPT violation in experimentally interesting $K - \bar{K}$ and analogous systems [45]. At the same time, any indication on CPT violation is absent up to now.

VIII. ALTERNATIVE RESEARCH

The decay $(K_S^0 \to \mu^+ \mu^-)$ was investigated in [46] as a sensitive probe of the short-distance region. They updated the Standard Model predictions of the short-distance contribution by a few present. As the CP violating effect is small, ($\approx 10^{-3}$) also small corrections should also be taken into account for accurate extraction from experimental data.

In Ref. [47] was considered CP violating asymmetry in the energy of the charged pions in the decay in term of two parameters. They may be determined by studying the corresponding time-dependent asymmetry in the decay of the neutral kaons with tagged. Initial strangeness. The measure of the CP conserving asymmetry will provide a test of the Chiral Perturbation Theory and give information on the final-state interaction of the three pions.

Higher-order electroweak contributions to indirect CP violation in neutral kaons were calculated in [48]. The authors wrote that the theoretical calculation of ϵ_K had historically large perturbative errors arising from charm-quark corrections. These errors were larger than expected higher-order electroweak corrections. Recently, a simple reparametrization of the effective Hamiltonian reduced perturbation errors, making higher-order electroweak calculations justified, They presented the O(1%) leading logarithm electroweak contribution to ϵ_K .

In 1973, trying to find an explanation for CP violation in the decay of neutral kaons and starting from the idea of Cabibbo about the mixing of two generations of quarks, Kobayashi and Maskawa predicted the existence of a third generation. Indeed, the bottom b quark was discovered in 1977, and the t quark in 1995.

The differences in the properties of B and B mesons predicted by Kobayashi and Maskawa theory, including CP violation, were discovered in the quark sector on the BaBar experiment at the Stanford Linear Accelerator Center (SLAC) and Belle experiments at the High Energy Accelerator Research Organization (KEK) in 2002–2007. CP violation for quarks with a charge of -1/3 corresponds to a strange s-quark that is part of kaons and a beauty *b*-quark that is part of *B*-mesons. However, CP violation was also predicted for quarks with charge of +2/3, corresponding to the charm *c* quark that is part of the *D*-meson. The weak force appears to act not upon a quark state, as identified by the flavor of quark, but on a quantum mixture of two types of quark for indirect CP violation contrary to direct CP violation in pure quark state.

A large number of CP violating processes in *B*-meson and *D* meson decays have been discovered starting in 2001 in *B*-factory experiments, including the BaBar experiment at SLAC [49] and the Belle experiment at the High Energy Accelerator Research Organization (KEK) [50] also by the Large Hadron Collider beauty Collaboration (LHCb) at CERN [51]. In 2010, D0 Collaboration at Fermilab's Tevatron Particle Collider, an international team of 500 physicists indicated a 1% difference between the production of pairs of muons and pairs of antimuons in the decay of B mesons produced in high-energy collisions [52].

The CP violating processes observed so far fit well into the framework of the Standard Model. However, surprises cannot be ruled out when studying such processes with the need to search for alternative sources of CP violation with access to new physics. This problem is discussed in the Ref. [53]. The consideration via a loop transition, *i.e.*, penguin decays, is very sensitive to possible contributions from unknown particles, provided any probes for new sources of CP violation [54].

A measurement of time-dependent CP violation in the decays of the B^0 and \bar{B}^0 mesons to the final states $J/\psi(\rightarrow \mu^+\mu^-)K_S^0$, $\psi(2S)(\rightarrow \mu^+\mu^-)K_S^0$, and $J/\psi(\rightarrow e^+e^-)K_S^0$ with $K_S^0 \rightarrow \pi^+\pi^-$ (the J/ψ meson was discovered in 1974 by two groups of experimentalists at SLAC and BNL, confirming the existence of charm quark c) was considered in Ref. [55]. In the agreement with their statement the obtained CP violation parameters represent the most prestige single measurement of the CKM angle β to date and was more precise than the current world average.

Fleischer gave a review [56] on CP violation in decays B: recent development and future

perspectives. He indicated that these processes provide a powerful tool to probe New Physics beyond the Standard Model. Also the review of recent data in B decays is given by Malami in Ref. [57].

CP violation was also predicted for D mesons, which include a representative of the second type of quarks with a charge of +2/3, *i.e.*, a charmed *c*-quark [58]. The latest publication of the LHCb Collaboration report that for the first time it was possible to accurately establish CP violation for D mesons. The asymmetry A_{CP} of the decay of the D meson relative to the \overline{D} was studied in two different channels, on $\pi^+\pi^-$ and K^+K^- . But this difference also depends on the unequal determination of the decay products in these two channels. However, the difference in asymmetries in the two decay channels reduces this detection effect, leaving only CP violation. The result was symmetry violation [59]:

$$\Delta A_{CP} = (-15.4 \pm 2.9) \times 10^{-4}$$

Starting with 2012 KOTO Collaboration (KEK) investigated rare decay in the J-PARC 30 GeV Main Ring due to direct CP violation [60]. They obtained the upper limit of $BR(K_L^0 \to \pi^0 \nu \bar{\nu}) < 5.1 \times 10^{-8}$ (90% *CL*). This mode directly breaks the CP symmetry and is highly suppressed in the Standard Model. Moreover, the theoretical uncertainty of this decay is only a few percent. These features make this decay one of the best probes to search for new physics beyond the Standard Model. They concluded that the number of observed events is statistically consistent with the background expectation. The agreement of theoretical prediction in frame of Standard Model with this experimental result could be indication on absence any necessity for new physics.

The analogous search was carried out by the NA62 Collaboration in CERN in the first search for rare $K^+ \to \pi^+ \nu \bar{\nu}$ using the decay-in-flight technique [61]. They received the upper limit of

$$BR_{exp}(K^+ \to \pi^+ \nu \bar{\nu}) < 14 \times 10^{-10} \ (95\% \ CL)$$
.

This value agrees, within error, with the prediction of the Standard Model,

$$BR_{theor}(K^+ \to \pi^+ \nu \bar{\nu}) < 10 \times 10^{-10} \ (95\% \ CL)$$

limiting the search for theories beyond it. The kaons were born in a collision with a beryllium target of a photon beam with an energy of 450 GeV from the CERN Super Proton Synchrotron (SPS). The 270-meter-long detector, using a system of triggers, made it possible to select kaon decays necessary for calculating the background and searching for the desired decay channel. This ultra rare decay channel K was detected for the first time with a statistical accuracy of 3.5σ . The current theoretical and experimental situation for the golden modes $K \to \pi \nu \bar{\nu}$, taken from [5], is summarized in Fig. 5.



FIG. 5. Summary of current situation for the golden modes $K \to \pi \nu \bar{\nu}$, taken from Ref. [5]. The red ellipse shows the 1σ Standard Model prediction with input from CKMfitter [62]; the green (yellow) region corresponds to the NA62 (BNL787/949) 1σ measurement; and the dashed orange line marks the 90% CL KOTO upper bound [63]. The black shaded region shows the Grossman and Nir [64] exclusion.

The existing experiments indicate that CP violation is too weak compared to what is required to explain the matter-antimatter asymmetry. The magnitude of the CP violation effect will be proportional to the function of all three mixing angles and the δ known as the Jarlskog invariant:

$$J_{CP}^{CKM} = c_{12} c_{13}^2 c_{23} s_{12} s_{13} s_{23} \sin \delta \approx 3 \times 10^{-5}$$

The quark sector cannot explain the baryon asymmetry of the Universe while mixing angles are small.

The recent results are reported at "The 2024 International Workshop on Future Tau

Charm Facilities," Shanghai, China, January 14-18, 2024,

https://indico.pnp.ustc.edu.cn/event/91/ and at "22nd Conference on Flavor Physics and CP Violation," Bangkok, Thailand, May 27-31, 2024, https://indico.cern.ch/event/1291023/.

An open question remains to understand the participation of heavy quarks during CP violation in the weak interaction, while spatial parity violation (P-violation) is characteristic of all weak interaction processes. In contrast to P parity violation in weak interaction. CP parity violation does not exceed a fraction of a percent. Finally, further explaining the CP violation beyond the Standard Model, that is, within the framework of new physics, is an important task in studies of this process.

One of the fundamental problems of modern physics is the explanation of the causes of CP in the weak interaction.

ACKNOWLEDGMENTS

One of us, N.P., thanks Dr. Wilhelm Czaplinski for collaboration. This work was supported in part by the U. S. Department of Energy, Office of Science, Office of Nuclear Physics, under Award No. DE–SC0016583.

- T. D. Lee and C. N. Yang, "Question of Parity Conservation in Weak Interactions," Phys. Rev. 104, 254 (1956).
- [2] C. S. Wu, E. Ambler, R. W. Hayward, D. D. Hoppes, and R. P. Hudson, "Experimental Test of Parity Conservation in β Decay," Phys. Rev. 105, 1413 (1957).
- [3] G. Luders, "On the equivalence of invariance under time reversal and under particleantiparticle conjugation for relativistic field theories," Kong. Dan. Vid. Sel. Mat. Fys. Med. 28N5, 1 (1954).
- [4] W. Pauli, "Niels Bohr and the development of physics," (Pergamon Press, London, 1955).
- [5] R. L. Workman *et al.* [Particle Data Group], "Review of Particle Physics," Prog. Theor. Exp. Phys. **2022**, 083C01 (2022).
- [6] A. Pais and O. Piccioni, "Note on the decay and absorption of the Theta0," Phys. Rev. 100, 1487 (1955).

- [7] L. B. Leipuner, W. Chinowsky, R. Crittenden, R. Adair, B. Musgrave, and F. T. Shively,
 "Anomalous regeneration of K-10 mesons from K-20 mesons," Phys. Rev. 132, 2285 (1963).
- [8] E. Wigner, "Grupprntheorie und ihre Anwendungen auf die Quantenmechanik der Atomspekten," (Friedr Vieweg und Sohn, Baunschweig, 1931); Nachr. Gesell. Der Wissen, Math.-Phys. Klasse 5, 546 (1932).
- [9] J. H. Christenson, J. W. Cronin, V. L. Fitch, and R. Turlay, "Evidence for the 2π decay of the K₂⁰ meson," Phys. Rev. Lett. 13, 138 (1964).
- [10] A. D. Sakharov, "Violation of CP Invariance, C asymmetry, and baryon asymmetry of the universe," Pisma Zh. Eksp. Teor. Fiz. 5 (1967), 32 (1967) [JETP Lett. 5, 24 (1967)]; Usp. Fiz. Nauk 161, 61 (1991) [Sov. Phys. Usp. 34, 392 (1991)].
- [11] N. Popov and I. Strakovsky, "Neutrino Mass: Past, Present, and Future," (NOVA Science Publishers, 2021).
- [12] N. Cabibbo, "Unitary symmetry and leptonic decays," Phys. Rev. Lett. 10, 531 (1963).
- [13] M. Kobayashi and T. Maskawa, "CP violation in the renormalizable theory of weak interaction," Prog. Theor. Phys. 49, 652 (1973).
- [14] L. Wolfenstein, "Violation of CP invariance and the possibility of very weak interactions," Phys. Rev. Lett. 13, 562 (1964).
- [15] J. S. Bell and J. Steinberger, "Weak interactions of kaons," in Proceedings of Oxford Int. Conf. on Elementary Particles, September 1965, Eds.T. R. Walsh, A. E. Taylor, R. G. Moorhousde and B. Southworth (Ruserford High Energy Lab., Chilton, Didicot, 1966) pg. 195.
- [16] M. Holder *et al.* "On the decay $K_L \to \pi^0 \pi^0$," PhysLett. B **40**, 141 (1972).
- [17] M. Banner, J. W. Cronin, C. M. Hoffman, B. C. Knapp, and M. J. Shochet, "Measurement of eta00/eta+-," Phys. Rev. Lett. 28, 1597 (1972).
- [18] J. H. Christenson, J. H. Goldman, E. Hummel, S. D. Roth, T. W. L. Sanford, and J. Sculli, "Measurement of the phase and magnitude of Eta00," Phys. Rev. Lett. 43, 1209 (1979).
- [19] H. Burkhardt *et al.* [NA31 Collaboration], "First evidence for direct CP violation," Phys. Lett. B 206, 169 (1988).
- [20] G. D. Barr et al. [NA31 Collaboration], "A New measurement of direct CP violation in the neutral kaon system," Phys. Lett. B 317, 233 (1993)
- [21] L. K. Gibbons *et al.* [E731 Collaboration], "Measurement of the CP violation parameter Re(ε'/ε)," Phys. Rev. Lett. **70**, 1203 (1993).

- [22] L. K. Gibbons *et al.* [E731 Collaboration], "CP and CPT symmetry test from the two pion decays of the neutral kaon with the Fermilab E731 detector," Phys. Rev. D 55, 6625 (1997).
- [23] J. R. Ellis, M. K. Gaillard, and D. V. Nanopoulos, "Lefthanded Currents and CP Violation," Nucl. Phys. B 109, 213 (1976).
- [24] F. J. Gilman and M. B. Wise, "The delta I = 1/2 rule and violation of CP in the six quark model," Phys. Lett. B 83, 83 (1979).
- [25] B. Guberina and R. D. Peccei, "Quantum Chromodynamic effects and CP violation in the Kobayashi-Maskawa Model," Nucl. Phys. B 163, 289 (1980).
- [26] G. Buchalla, A. J. Buras, and M. K. Harlander, "The anatomy of epsilon-prime / epsilon in the Standard Model," Nucl. Phys. B 337, 313 (1990).
- [27] M. Ciuchini, E. Franco, G. Martinelli, L. Reina, and L. Silvestrini, "An Upgraded analysis of epsilon-prime epsilon at the next-to-leading order," Z. Phys. C 68, 239 (1995).
- [28] A. J. Buras, M. Jamin, and M. E. Lautenbacher, "A 1996 analysis of the CP violating ratio epsilon-prime / epsilon," Phys. Lett. B 389, 749 (1996).
- [29] J. M. Flynn and L. Randall, "The electromagnetic penguin contribution to epsilon-prime / epsilon for large top quark mass," Phys. Lett. B 224, 221 (1989) [erratum: Phys. Lett. B 235412 (1990)].
- [30] J. R. Batley et al. [NA48 Collaboration], "A precision measurement of direct CP violation in the decay of neutral kaons into two pions," Phys. Lett. B 544, 97 (2002), and references therein.
- [31] A. Lai *et al.* [NA48 Collaboration], "A precise measurement of the direct CP violation parameter Re(ε'/ε)," Eur. Phys. J. C 22, 231 (2001).
- [32] L. Iconomidou-Fayard and D. Fournier, "The Discovery of Direct CP Violation," Adv. Ser. Direct. High Energy Phys. 23, 205 (2015).
- [33] V. D. Kekelidze and D. T. Madigozhin, "On the observation of direct CP violation in neutral kaon decays," Fizika Elementarnykh Chastits i Atomnogo Yadra 38, 1163 (2007) [Phys. Part. Nucl. 38, 610 (2007)].
- [34] A. Alavi-Harati *et al.* [KTeV Collaboration], "Observation of direct CP violation in $K_{S,L} \to \pi\pi$ decays," Phys. Rev. Lett. **83**, 22 (1999).
- [35] A. Alavi-Harati *et al.* [KTeV Collaboration], "Measurements of direct CP violation, CPT symmetry, and other parameters in the neutral kaon system," Phys. Rev. D 67, 012005 (2003)

[erratum: Phys. Rev. D **70**, 079904 (2004)]

- [36] Y. B. Hsiung [for KTeV Collaboration], "An Observation of direct CP violation epsilon prime/epsilon result from KTeV," Nucl. Phys. B Proc. Suppl. 86, 312 (2000).
- [37] A. Glazov, "Measurement of direct CP violation in K meson decays at KTeV," Nucl. Phys. A 675, 116C (2000).
- [38] E. Abouzaid *et al.* [KTeV Collaboration], "Precise Measurements of Direct CP Violation, CPT Symmetry, and Other Parameters in the Neutral Kaon System," Phys. Rev. D 83, 092001 (2011).
- [39] T. D. Lee and C. S. Wu, "Weak Interactions," Ann. Rev. Nucl. Part. Sci. 15, 381 (1965).
- [40] L. B. Okun, "Leptons and Quarks," (North-Holland, Amsterdam, N.Y., Oxford, 1982).
- [41] I. I. Bigi and A. I. Sando, "CP Violation," (Cambridge Monographs on Particle Physics, Nuclear Physics and Cosmology, 2016).
- [42] N. Popov, "Muon Physics," (Cambridge Scholars Publishing, 2019).
- [43] D. Colladay and V. A. Kostelecky, "CPT violation and the standard model," Phys. Rev. D 55, 6760 (1997).
- [44] V. A. Kostelecky, "Topics in Lorentz and CPT violation," [arXiv:1010.4559 [hep-ph]].
- [45] G. Cvetic, S. K. Kang, C. S. Kim, and K. M. Lee, "No CPT violation from tilted brane in neutral meson - anti-meson systems," Phys. Rev. D 62, 057901 (2000).
- [46] J. Brod and E. Stamou, "Impact of indirect CP violation on $Br(K_S \to \mu^+ \mu^-)_{l=0}$," JHEP **05**, 155 (2023).
- [47] F. Buccella, O. Pisanti, and F. Sannino, "Energy charge correlation in the pi+ pi- pi0 decay of K(L) and of tagged neutral kaons," Z. Phys. C 66, 135 (1995).
- [48] J. Brod, S. Kvedaraite, Z. Polonsky, and A. Youssef, "Higher-Order Electroweak Contributions to Indirect CP Violation in Neutral Kaons," PoS DISCRETE2022, 041 (2024).
- [49] B. Aubert *et al.* [BaBar Collaboration], "Measurement of CP violating asymmetries in B⁰ decays to CP eigenstates," Phys. Rev. Lett. 86, 2515 (2001).
- [50] K. Abe *et al.* [Belle Collaboration], "Observation of large CP violation in the neutral B meson system," Phys. Rev. Lett. 87, 091802 (2001).
- [51] R. Aaij et al. [LHCb Collaboration], "First observation of CP violation in the decays of B⁰_s mesons," Phys. Rev. Lett. **110**, 221601 (2013).

- [52] V. M. Abazov *et al.* [D0 Collaboration], "Search for the rare decay $B_s^0 \to \mu^+ \mu^-$," Phys. Lett. B **693**, 539 (2010).
- [53] Y. Nir, "Exploring new physics with CP violation in neutral D and B decays," Nuovo Cim. A 109, 991 (1996).
- [54] M. Z. Barel, K. De Bruyn, R. Fleischer, and E. Malami, "Penguin Effects in $B_d^0 \to J/\psi K_S^0$ and $B_s^0 \to J/\psi \phi$," PoS **CKM2021**, 111 (2023).
- [55] R. Aaij *et al.* [LHCb Collaboration], "Measurement of CP violation in $B^0 \to \psi (\to l^+ l^-) K_S^0 (\to \pi^+\pi^-)$ decays," Phys. Rev. Lett. **132**, 021801 (2024).
- [56] R. Fleischer, "CP violation in B decays: Recent developments and future perspectives," [arXiv:2402.00710 [hep-ph]].
- [57] E. Malami, "Theoretical highlights of CP violation in *B* decays," [arXiv:2402.16976 [hep-ph]].
- [58] T. Feldmann, S. Nandi, and A. Soni, "Repercussions of Flavour Symmetry Breaking on CP Violation in D-Meson Decays," JHEP 06, 007 (2012).
- [59] R. Aaij et al. [LHCb Collaboration], "Observation of CP violation in charm decays," Phys. Rev. Lett. 122, 211803 (2019).
- [60] J. K. Ahn *et al.* [KOTO Collaboration], "Study of the $K_L \to \pi^0 \nu \bar{\nu}$ decay at the J-PARC KOTO experiment," Phys. Rev. Lett. **126**, 121801 (2021).
- [61] E. Cortina Gil *et al.* [NA62 Collaboration], "First search for $K^+ \to \pi^+ \nu \bar{\nu}$ using the decay-inflight technique," Phys. Lett. B **791**, 156 (2019).
- [62] J. Charles *et al.* "Predictions of selected flavour observables within the Standard Model," Phys. Rev. D 84, 033005 (2011).
- [63] J. K. Ahn *et al.* [KOTO Collaboration], "Search for the $K_L \to \pi^0 \nu \overline{\nu}$ and $K_L \to \pi^0 X^0$ decays at the J-PARC KOTO experiment," Phys. Rev. Lett. **122**, 021802 (2019).
- [64] Y. Grossman and Y. Nir, " $K_L \to \pi^0 \nu \bar{\nu}$ beyond the standard model," Phys. Lett. B **398**, 163 (1997).