

EQUIVALENCE BETWEEN VALIDITY OF THE p -POINCARÉ INEQUALITY AND FINITENESS OF THE STRICT p -CAPACITARY INRADIUS.

A.-K. GALLAGHER 

ABSTRACT. It is shown that the p -Poincaré inequality holds on an open set Ω in \mathbb{R}^n if and only if the strict p -capacitary inradius of Ω is finite. To that end, new upper and lower bounds for the infimum of the associated nonlinear Rayleigh quotients are derived.

1. INTRODUCTION

Let Ω be an open set in \mathbb{R}^n , and $\|\cdot\|_{p,\Omega}$ the L^p -norm on Ω for $1 < p < \infty$. The classical p -Poincaré inequality is said to hold on Ω if there exists a constant $C > 0$ such that

$$(1.1) \quad \|f\|_{p,\Omega} \leq C \|\nabla f\|_{p,\Omega} \quad \forall f \in C_c^\infty(\Omega),$$

where $C_c^\infty(\Omega)$ denotes the space of smooth functions with compact support in Ω . That is, (1.1) holds if the infimum of the so-called nonlinear Rayleigh quotient

$$\lambda_{1,p}(\Omega) = \inf \left\{ \|\nabla f\|_{p,\Omega}^p / \|f\|_{p,\Omega}^p : f \in C_c^\infty(\Omega) \setminus \{0\} \right\},$$

is positive. If $\lambda_{1,p}(\Omega)$ is positive and attained at some non-zero function in $W_0^{1,p}(\Omega)$, i.e., the closure of $C_c^\infty(\Omega)$ with respect to the $L^p(\Omega)$ -Sobolev 1-norm, then it is the smallest generalized eigenvalue for the p -Laplacian, with Dirichlet boundary condition, on Ω in the distributional sense.

In this note, we employ the concept of strict p -capacitary inradius of Ω , originally introduced in the case of $p = 2$ in [6] for $n = 2$ and in [5] for $n \geq 3$, and derive the equivalence between its finiteness and the positivity of $\lambda_{1,p}(\Omega)$. In the definition of the strict p -capacitary inradius, we use the notion of Sobolev p -capacity, $C_p(K)$, which, for a compact set $K \subset \mathbb{R}^n$, is defined as

$$C_p(K) = \inf \left\{ \|u\|_{p,\mathbb{R}^n}^p + \|\nabla u\|_{p,\mathbb{R}^n}^p : u \in \mathcal{S}, u \geq 1 \text{ on } K \right\},$$

where \mathcal{S} denotes the space of Schwartz functions. The strict p -capacitary inradius of Ω is defined as

$$(1.2) \quad \rho_p(\Omega) = \sup \left\{ r > 0 : \forall \epsilon > 0 \exists x \in \mathbb{R}^n \text{ such that } C_p(\overline{\mathbb{B}_r(x)} \cap \Omega^c) < \epsilon \right\},$$

where $\mathbb{B}_r(x)$ denotes the open ball in \mathbb{R}^n of radius $r > 0$ with center $x \in \mathbb{R}^n$. Roughly speaking, finiteness of $\rho_p(\Omega)$ means that the complement of Ω is somewhat evenly distributed in \mathbb{R}^n . In fact, if $\rho_p(\Omega)$ is finite and $R > \rho_p(\Omega)$, then there exists a $\delta > 0$ such that within R -units of any point in \mathbb{R}^n , we may find a set in the complement of Ω whose Sobolev p -capacity is at least δ .

Date: May 1, 2024.

2020 Mathematics Subject Classification. 35P30, 35J70, 31C45.

Key words and phrases. p -Poincaré inequality, capacitary inradius, p -Laplacian.

Finiteness of $\rho_p(\Omega)$ is a necessary condition for (1.1) to hold on an open set Ω . In fact, our first result yields a sharp upper bound for $\rho_p(\Omega)$ in terms of $\lambda_{1,p}(\Omega)$.

Theorem 1.3. *Let $\Omega \subset \mathbb{R}^n$ be an open set, $1 < p < \infty$. Suppose $\lambda_{1,p}(\Omega) > 0$.*

(i) *Then*

$$(1.4) \quad \left(\rho_p(\Omega)\right)^p \leq \lambda_{1,p}(\mathbb{B}_1(0))/\lambda_{1,p}(\Omega).$$

(ii) *If Ω is connected and bounded, then equality in (1.4) holds if and only if*

$$\Omega = \mathbb{B}(x, \rho_p(\Omega)) \setminus K$$

for some $x \in \mathbb{R}^n$ and some compact set K with $C_p(K) = 0$. Furthermore, if Ω is connected and unbounded such that $\rho_p(\Omega) = \rho_p(\Omega \cap \mathbb{B}_R(0))$ for some $R > 0$, then equality in (1.4) cannot hold.

The heart of the matter of the proof of Theorem 1.3 is a continuity result for $\lambda_{1,p}$. That is, suppose $\{K_j\}_{j \in \mathbb{N}}$ is a sequence contained in the closure of some bounded domain D with smooth boundary. If $C_p(K_j)$ tends to zero as $j \rightarrow \infty$ and, for each $j \in \mathbb{N}$, $D \setminus K_j$ is an open set with smooth boundary, then $\lambda_{1,p}(D \setminus K_j)$ tends to $\lambda_{1,p}(D)$. This was originally proved in [6] for the case $n = 2$, $p = 2$, $D = \mathbb{B}_1(0)$, and the logarithmic capacity in place of C_p . Regularity results for the Dirichlet problem and for first eigenfunctions of the p -Laplacian, Δ_p , on open, bounded sets with smooth boundary as well as the theory of p -harmonic functions allow this continuity result to be extended to the case of the Sobolev p -capacity with $1 < p < \infty$, see Lemma 3.1.

To state the sufficiency of the finiteness of $\rho_p(\Omega)$, we define the scalar

$$(1.5) \quad \delta_R(\Omega) := \sup \left\{ \delta \geq 0 : C_p(\overline{\mathbb{B}_R(x)} \cap \Omega^c) \geq \delta \forall x \in \mathbb{R}^n \right\}$$

for any $R > 0$. Note that $\delta_R(\Omega)$ is positive whenever $\rho_p(\Omega)$ is finite and $R > \rho_p(\Omega)$.

Theorem 1.6. *Let $\Omega \subset \mathbb{R}^n$ be an open set, $1 < p < \infty$. Suppose $\rho_p(\Omega) < \infty$ and $R > \rho_p(\Omega)$.*

(i) *Then,*

$$(1.7) \quad \frac{\delta_R(\Omega)}{R^n \cdot \|E_R\|_{op}^p} \leq \lambda_{1,p}(\Omega),$$

for any bounded, linear extension operator $E_R : W^{1,p}((0, R)^n) \rightarrow W^{1,p}(\mathbb{R}^n)$ satisfying $(E_R f)|_{(0, R)^n} = f$ for $f \in W^{1,p}((0, R)^n)$.

(ii) *There exist constants $C = C(n, p) > 0$ and $\gamma_R(\Omega) \in (0, 1]$ such that*

$$(1.8) \quad C\gamma_R(\Omega) \cdot R^{-p} \leq \lambda_{1,p}(\Omega).$$

Part (i) of Theorem 1.6 is a direct consequence of a Poincaré-type inequality for any function in $W^{1,p}(\Omega)$ which has a representative that vanishes on a set of positive p -Sobolev capacity in Ω , where Ω is a bounded extension domain. This inequality originates in the work of Meyers [16], and was completed by Adams, see Theorem 8.3.3 and the notes in Section 8.3 on pg. 231 in [1]. For the proof of Theorem 1.6, one simply writes \mathbb{R}^n as a union of closed cubes with mutually disjoint interiors and of side length larger than the strict p -capacitary inradius of the open set under consideration. Then, one applies this Poincaré-type inequality to each cube. This kind of proof is contained in the works of Maz'ya–Shubin

[13] and Souplet [17]. Souplet also splits \mathbb{R}^n into cubes and then uses a Poincaré-type inequality for functions which vanish on a set of positive Lebesgue measure. Maz'ya–Shubin derive a Poincaré-type inequality in [13, Lemma 3.1], similar to the one used in this note, but for balls which forces them take the multiplicity of coverings by balls into account in order to obtain a global estimate.

Theorems 1.3 and 1.6 may be summarized in a qualitative manner as follows.

Corollary 1.9. *Let $\Omega \subset \mathbb{R}^n$ be an open set, $1 < p < \infty$. Then,*

$$\lambda_{1,p}(\Omega) > 0 \Leftrightarrow \rho_p(\Omega) < \infty.$$

It is well-known that the finiteness of the inradius, $\mathfrak{R}(\Omega)$, of an open set Ω , i.e., the supremum of the radii of all balls contained in Ω , is a necessary condition for (1.1). This can be seen by a simple scaling argument, see, e.g., Souplet's proof of [17, Prop. 2.1 (i)]. Souplet additionally shows in [17, Prop. 2.1] with $p \in [1, \infty)$, that this condition is also sufficient as long as Ω is a domain which satisfies a uniform exterior cone condition; in the case of $p = 2$ this appears to go back to work of Agmon [2]. In [17], Souplet introduces a measure-theoretic inradius, which yields a sufficient condition for the validity of (1.1) without any regularity assumptions on the boundary of the domain Ω . Souplet's formulation of inradius actually inspired the notion of the capacitary inradius in (1.2), originally introduced in [6]. A sufficient condition, similar to Souplet's, was previously obtain by Lieb in [9, Corollary 2]. We note that the assumption of finiteness of either of these conditions is a stronger assumption than the one of finiteness of the strict capacitary inradius defined in (1.2). The reason for that is that these measure-theoretic inradii do not take into account sets in the complement which are of Lebesgue measure zero but of positive Sobolev p -capacity. However, for any pair (p', p) with $1 < p' < p \leq n$, there exists a set $E \subset \mathbb{R}^n$ of Lebesgue measure zero, such that $C_{p'}(E) = 0$ while $C_p(E) > 0$, see [1, Theorem 5.5.1].

A complete description in the flavor of Corollary 1.9 was first given by Maz'ya–Shubin in [13] in the case of $p = 2$ and $n \geq 3$. The authors of [13] use different notions of capacity and of capacitary inradius than presented in this note; see Lemma 2.14 on how their capacitary inradius relates to the one defined in (1.2). For $p = 2$ and $n \geq 3$, estimate (1.4) is an improvement over the upper bound for $\lambda_{1,2}(\Omega)$ provided in [13] while (1.8) and the lower bound given in [13] are similar. We also point to the work of Vitolo [19] in which he shows that if $p > n$ and Ω is a domain with finite inradius, $\mathfrak{R}(\Omega)$, then $\lambda_{1,p}(\Omega) > 0$. Note that if $p > n$, then singletons have positive p -Sobolev capacity, so that $\mathfrak{R}(\Omega) = \rho_p(\Omega)$, i.e., Theorem 1.6 rediscovers Vitolo's result.

Together with Lebl and Ramachandran, we considered the problem of describing the validity of the Poincaré inequality in the case of $n = 2, p = 2$ in potential-theoretic terms in [6]. Originally, we intended to investigate which potential-theoretic conditions yield the L^2 -closed range property of (the weak maximal extension of) the Cauchy–Riemann operator which constitutes an open problem in several complex variables. This closed range property turns out to be equivalent to (1.1), and in fact, we showed both that Corollary 1.9 holds for $p = 2$ and that (1.1) is equivalent to the existence of a smooth, bounded function on Ω such that its Laplacian has a positive lower bound on Ω . These results were later shown to hold for $p = 2, n \geq 3$ with the Newtonian capacity in place of C_p in [5] by the author of this note. We show in Lemma 2.11 that the strict p -capacitary inradius

defined in (1.2) does not depend on the choice of p -capacity as long as the sets of zero p -capacity are the same as the sets of for which C_p is zero as well. In particular, the strict capacity inradii defined in [6, 5] are the same as the one defined in (1.2) for $p = 2$, see the paragraph subsequent to the proof of Lemma 2.11.

This note is structured as follows. The notions of capacity, strict p -capacity inradius, and $\lambda_{1,p}$ and their basic properties are recalled in Section 2. The proofs of Theorem 1.3 and Theorem 1.6 are given in Section 3 and 4, respectively.

Acknowledgement. I am very grateful to Carlo Morpurgo for his insights he shared with me while completing this project.

2. PRELIMINARIES

2.1. Sobolev p -capacity.

Definition 2.1. Let K be a compact set in \mathbb{R}^n , $1 < p < \infty$. Then

$$C_p(K) := \inf \left\{ \|u\|_{p,\mathbb{R}^n}^p + \|\nabla u\|_{p,\mathbb{R}^n}^p : u \in \mathcal{S}, u \geq 1 \text{ on } K \right\}.$$

This definition may be extended to open sets $U \subset \mathbb{R}^n$ by setting

$$(2.2) \quad C_p(U) = \sup \{ C_p(K) : K \subset U, K \text{ compact} \}.$$

It then follows that

$$C_p(K) = \inf \{ C_p(U) : K \subset U, U \text{ open} \},$$

which can be proven analogously to [1, Proposition 2.2.3]. The definition of C_p may now be extended to arbitrary sets by setting

$$(2.3) \quad C_p(E) := \inf \{ C_p(U) : E \subset U, U \text{ open} \}$$

for $E \subset \mathbb{R}^n$. A set $E \subset \mathbb{R}^n$ is called p -polar, if $C_p(E) = 0$. Moreover, two functions are said to equal p -quasi everywhere if they equal outside a p -polar set.

Next, we present some standard properties of C_p .

Lemma 2.4. Let $E \subset \mathbb{R}^n$. Then

- (i) If $E' \subset E$, then $C_p(E') \leq C_p(E)$.
- (ii) If $x \in \mathbb{R}^n$, then $C_p(E + x) = C_p(E)$.
- (iii) If $s > 0$, then $C_p(sE) \leq s^n \max\{1, s^{-p}\} C_p(E)$.
- (iv) If $\{E_i\}_{i \in \mathbb{N}} \subset \mathbb{R}^n$ such that $E = \bigcup_{i=1}^{\infty} E_i$, then $C_p(E) \leq \sum_{i=1}^{\infty} C_p(E_i)$.
- (v) Any Borel set E is capacitable, i.e.,

$$C_p(E) = \sup \{ C_p(K) : K \subset E, K \text{ compact} \} = \inf \{ C_p(U) : E \subset U, U \text{ open} \}.$$

Proof. The proofs of (i)–(iv) for arbitrary sets follow from (2.2) and (2.3) once (i)–(iii) have been established for compact sets. For compact sets, (i), (ii) and (iv) follow directly from Definition 2.1 while (iii) follows from a change of variable argument yielding

$$C_p(sK) = \inf \left\{ s^n \|u\|_{p,K}^p + \frac{s^n}{s^p} \|\nabla u\|_{p,K}^p : u \geq 1 \text{ on } K, u \in \mathcal{S} \right\}.$$

For the proof of (v), see Propositions 2.3.12 and 2.3.13 as well as Theorem 2.3.11 and the succeeding remark in [1]. \square

2.2. The strict p -capacitary inradius. In a slight deviation of (1.2), we define the strict p -capacitary inradius as follows.

Definition 2.5. Let $\Omega \subset \mathbb{R}^n$ be an open set, $1 < p < \infty$. Then, the strict p -capacitary inradius of Ω is defined as

$$(2.6) \quad \rho_p(\Omega) = \sup \left\{ r > 0 : \forall \epsilon > 0 \exists x \in \mathbb{R}^n \text{ such that } C_p(\mathbb{B}_r(x) \cap \Omega^c) < \epsilon \right\}.$$

We show first that this definition of ρ_p agrees with (1.2), although C_p is not invariant under taking closures.

Lemma 2.7. Let $\Omega \subset \mathbb{R}^n$ be an open set, $1 < p < \infty$. Then

$$(2.8) \quad \rho_p(\Omega) = \sup \left\{ r > 0 : \forall \epsilon > 0 \exists x \in \mathbb{R}^n \text{ such that } C_p(\overline{\mathbb{B}_r(x)} \cap \Omega^c) < \epsilon \right\}.$$

Proof. Let us denote the right hand side of (2.8) by $\hat{\rho}_p(\Omega)$. By the monotonicity of the Sobolev p -capacity, it is immediate that $\hat{\rho}_p(\Omega) \leq \rho_p(\Omega)$. Now, suppose that $0 < R < \rho_p(\Omega)$. Then, for all $\epsilon > 0$ there exists an $x \in \mathbb{R}^n$ such that $C_p(\mathbb{B}_R(x) \cap \Omega^c) < \epsilon$, and hence

$$C_p(\overline{\mathbb{B}_{R-\delta}(x)} \cap \Omega^c) < \epsilon \quad \forall \delta \in (0, R).$$

Therefore, $R - \delta < \hat{\rho}_p(\Omega)$ for all $\delta \in (0, R)$, which implies $\rho_p(\Omega) \leq \hat{\rho}_p(\Omega)$. \square

In the following, we collect some basic properties of the strict p -capacitary inradius. To do so, we recall that the inradius, $\mathfrak{R}(\Omega)$, of an open set $\Omega \subset \mathbb{R}^n$ is defined as

$$\mathfrak{R}(\Omega) = \sup \{ r > 0 : \exists x \in \mathbb{R}^n \text{ such that } \mathbb{B}_r(x) \subset \Omega \}.$$

We also define the p -capacitary inradius, $r_p(\Omega)$, by

$$r_p(\Omega) = \sup \left\{ r > 0 : \exists x \in \mathbb{R}^n \text{ such that } C_p(\mathbb{B}_r(x) \cap \Omega^c) = 0 \right\}.$$

Lemma 2.9. Let $\Omega \subset \mathbb{R}^n$ be an open set, $1 < p < \infty$.

- (i) If $x \in \mathbb{R}^n$ and $s > 0$, then $\rho_p(\Omega + x) = \rho_p(\Omega)$ and $\rho_p(s\Omega) = s\rho_p(\Omega)$.
- (ii) If $\Omega \subset \Omega'$ is an open set, the $\rho_p(\Omega) \leq \rho_p(\Omega')$. If additionally, $\Omega' \setminus \Omega$ is p -polar, then $\rho_p(\Omega) = \rho_p(\Omega')$.
- (iii) $\rho_p(\Omega) \geq r_p(\Omega) \geq \mathfrak{R}(\Omega)$, and equality holds if $p > n$.
- (iv) If Ω is bounded, then $\rho_p(\Omega) = r_p(\Omega)$. Moreover, there exists an $x^\circ \in \mathbb{R}^n$, such that

$$C_p(\mathbb{B}_{\rho_p(\Omega)}(x^\circ) \cap \Omega^c) = 0,$$

i.e., $r_p(\Omega)$ is attained.

- (v) $\rho_p(\Omega) = \lim_{R \rightarrow \infty} \rho_p(\Omega \cap \mathbb{B}_R(0))$.

Proof. The translation invariance of ρ_p holds because it holds for C_p , see (ii) of Lemma 2.4. To check the linearity under dilations we first note that

$$\mathbb{B}_r(x) \cap \Omega^c = \frac{1}{s} (\mathbb{B}_{sr}(xs) \cap (s\Omega)^c).$$

Thus, if for a given $r > 0$ and $\epsilon > 0$ there exists an $x \in \mathbb{R}^n$ such that

$$C_p(\mathbb{B}_{sr}(xs) \cap (s\Omega)^c) < \epsilon,$$

then

$$C_p(\mathbb{B}_r(x) \cap \Omega^c) \leq s^{-n} \max\{1, s^p\} \epsilon,$$

by (iii) of Lemma 2.4. It then follows that $r < \rho_p(\Omega)$ whenever $sr < \rho_p(s\Omega)$, and hence, $s\rho_p(\Omega) \leq \rho_p(s\Omega)$ for any $s > 0$. We now may repeat this argument with $t = \frac{1}{s}$ in place of s and $t^{-1}\Omega$ in place of Ω to obtain

$$t\rho_p(t^{-1}\Omega) \leq \rho_p(\Omega) \Rightarrow s^{-1}\rho_p(s\Omega) \leq \rho_p(\Omega),$$

which yields

$$s\rho_p(\Omega) \leq \rho_p(s\Omega) \leq s\rho_p(\Omega),$$

hence, the proof of (i) is complete.

The first part of (ii) follows from the definition. The second part follows after observing

$$C_p(\mathbb{B}_r(x) \cap \Omega^c) \leq C_p(\mathbb{B}_r(x) \cap (\Omega')^c)$$

by (iv) of Lemma 2.4 and the fact that $\mathbb{B}_r(x) \cap (\Omega' \setminus \Omega)$ is p -polar.

The set of inequalities in (iii) follows directly from the definitions of the inradii. Equality holds if $p > n$, because $C_p(\{x\}) > 0$ for $x \in \mathbb{R}^n$. To wit, if $R > \mathfrak{R}(\Omega)$, then $\mathbb{B}_R(x) \cap \Omega^c$ is non-empty for all $x \in \mathbb{R}^n$ so that $C_p(\mathbb{B}_R(x) \cap \Omega^c) > C_p(\{0\})$ for all $x \in \mathbb{R}^n$. Hence, $R > \rho_p(\Omega)$, so that $\mathfrak{R}(\Omega) = \rho_p(\Omega)$ follows.

For the proof of (iv), suppose Ω is bounded. Then, by definition of $\rho_p(\Omega)$, there exists a sequence $\{(r_j, x_j)\}_{j \in \mathbb{N}}$ in $\mathbb{R}_0^+ \times \mathbb{R}^n$ such that $\{r_j\}_{j \in \mathbb{N}}$ is an increasing sequence which converges to $\rho_p(\Omega)$, and

$$C_p(\mathbb{B}_{r_j}(x_j) \cap \Omega^c) < \frac{1}{j}.$$

Since Ω is a bounded set, it follows that $\{x_j\}$ is a bounded sequence, thus, has a convergent subsequence. For ease of notation, let us denote the subsequence by $\{x_j\}_{j \in \mathbb{N}}$. Write x° for the limit point. It suffices to prove that

$$(2.10) \quad C_p(\mathbb{B}_{\rho_p(\Omega)}(x^\circ) \cap \Omega^c) = 0$$

holds. To prove (2.10), let $\delta \in (0, \rho_p(\Omega))$ and choose $j_0 \in \mathbb{N}$ such that $|x^\circ - x_j| < \frac{\delta}{2}$ for all $j \geq j_0$. Then, choose $j_1 \geq j_0$ such that $\rho_p(\Omega) < r_j + \frac{\delta}{2}$ for all $j \geq j_1$. It follows that

$$\overline{\mathbb{B}_{\rho_p(\Omega)-\delta}(x^\circ)} \subset \mathbb{B}_{r_j}(x_j) \quad \forall j \geq j_1,$$

and, therefore,

$$C_p(\overline{\mathbb{B}_{\rho_p(\Omega)-\delta}(x^\circ)} \cap \Omega^c) \leq C_p(\mathbb{B}_{r_j}(x_j) \cap \Omega^c) < \frac{1}{j} \quad \forall j \geq j_1.$$

Letting $j \rightarrow \infty$ then yields

$$C_p(\overline{\mathbb{B}_{\rho_p(\Omega)-\delta}(x^\circ)} \cap \Omega^c) = 0 \quad \forall \delta \in (0, \rho_p(\Omega)),$$

and, hence, by part (v) of Lemma 2.4, the claimed (2.10) follows.

Part (v) follows directly from the monotonicity property in (ii). \square

We now can prove that the strict p -capacitary inradius does not depend on the choice of p -capacity.

Lemma 2.11. *Let $\Omega \subset \mathbb{R}^n$ be open, $1 < p < \infty$. Let $\Gamma_p : \mathcal{P}(\mathbb{R}^n) \rightarrow \mathbb{R}_0^+ \cup \{\infty\}$ be such that*

- (a) $\Gamma_p(\emptyset) = 0$,
- (b) $E \subset E' \Rightarrow \Gamma_p(E) \leq \Gamma_p(E')$,
- (c) *all Borel sets are capacitable with respect to Γ_p .*

Suppose $\Gamma_p(E) = 0$ iff $C_p(E) = 0$ for all bounded Borel sets $E \subset \mathbb{R}^n$. Then

$$(2.12) \quad \rho_p(\Omega) = \sup \left\{ r > 0 : \forall \epsilon > 0 \exists x \in \mathbb{R}^n \text{ such that } \Gamma_p(\mathbb{B}_r(x) \cap \Omega^c) < \epsilon \right\}.$$

Proof. Note first that properties (a)–(c) ensure that (iv) and (v) of Lemma 2.9 hold for the capacity inradius, $\rho_p^\Gamma(\Omega)$, defined by the right hand side of (2.12).

Next, write Ω_R for $\Omega \cap \mathbb{B}_R(0)$ for $R > 0$. Then, by (v) and (iv) of Lemma 2.9, it follows that

$$(2.13) \quad \begin{aligned} \rho_p(\Omega) &= \lim_{R \rightarrow \infty} \rho_p(\Omega_R) = r_p(\Omega_R) \\ &= \sup \left\{ r > 0 : \exists x \in \mathbb{R}^n \text{ such that } \Gamma_p(\mathbb{B}_r(x) \cap (\Omega_R)^c) = 0 \right\}, \end{aligned}$$

where the last step follows from the assumption that $\Gamma_p(E) = 0$ iff $C_p(E) = 0$ for all bounded Borel set $E \subset \mathbb{R}^n$. Since (iv)–(v) of Lemma 2.9 hold for the (strict) capacity inradius with respect to Γ_p , it follows

$$\sup \left\{ r > 0 : \exists x \in \mathbb{R}^n \text{ such that } \Gamma_p(\mathbb{B}_r(x) \cap (\Omega_R)^c) = 0 \right\} = \lim_{R \rightarrow \infty} \rho_p^\Gamma(\Omega_R) = \rho_p^\Gamma(\Omega),$$

i.e., ρ_p is invariant under the choice of Γ_p . \square

Both the logarithmic capacity for $n = 2$ and the Newtonian capacity for $n \geq 3$ satisfy (a)–(c) of Lemma 2.11. Moreover, their bounded polar Borel sets are equal to the bounded Borel sets which are polar with respect to C_2 by Theorem 1 ($m = 1$, $p = 2$) in [12], see also Theorems A and B ($\alpha = 0$, $m = 2$) in [20]. As a consequence, the strict capacity inradius ρ_2 with respect to C_2 is the same as the one defined in [6] with respect to the logarithmic capacity for $n = 2$ and the one defined in [5] with respect to the Newtonian capacity for $n \geq 3$.

Maz'ya and Shubin used a different notion of inradius, formulated in terms of the Wiener capacity, in their work [13], for $p = 2$ and $n \geq 3$. To wit, they defined the *interior capacity radius*, $r_{\Omega, \gamma}$, of an open set $\Omega \subset \mathbb{R}^n$ for $\gamma \in (0, 1)$ by

$$r_{\Omega, \gamma} = \sup \left\{ r > 0 : \exists x \in \mathbb{R}^n \text{ such that } C'_2(\overline{\mathbb{B}_r(x)} \setminus \Omega) \leq \gamma C'_2(\overline{\mathbb{B}_r(0)}) \right\},$$

where the Wiener capacity C'_p , $1 < p < \infty$, is defined by

$$C'_p(K) = \inf \left\{ \|\nabla u\|_{p, \mathbb{R}^n}^p : u \geq 1 \text{ on } K, u \in \mathcal{S} \right\}$$

for $K \subset \mathbb{R}^n$ compact.

This interior capacity radius relates to ρ_2 , for $n \geq 3$, as follows.

Lemma 2.14. *Let $\Omega \subset \mathbb{R}^n$, $n \geq 3$, be an open set. Then $\rho_2(\Omega) = \inf\{r_{\Omega, \gamma} : \gamma \in (0, 1)\}$.*

Proof. Note first that the Wiener capacity is equivalent to the Newtonian capacity, see, e.g., pg. 4 in [13] for a sketch of the proof. That means in particular that the Wiener capacity satisfies the hypotheses of Lemma 2.11, and hence, it suffices to show that $\rho'_2(\Omega) = \inf\{r_{\Omega, \gamma} : \gamma \in (0, 1)\}$, where

$$\rho'_2(\Omega) = \sup \{ r > 0 : \forall \epsilon > 0 \exists x \in \mathbb{R}^n \text{ such that } C'_2(\mathbb{B}_r(x) \cap \Omega^c) < \epsilon \}.$$

Moreover, the arguments supplied in the proof of (2.8) let us work with this alternative formulation for $\rho'_2(\Omega)$:

$$\rho'_2(\Omega) = \sup \{ r > 0 : \forall \epsilon > 0 \exists x \in \mathbb{R}^n \text{ such that } C'_2(\overline{\mathbb{B}_r(x)} \cap \Omega^c) < \epsilon \}.$$

Now, let $\gamma \in (0, 1)$ be given. Let $R < \rho'_2(\Omega)$, and choose $\epsilon = \gamma C'_2(\overline{\mathbb{B}_R(0)})$. Then, by definition of $\rho'_2(\Omega)$, there exists an $x \in \mathbb{R}^n$ such that

$$C'_2(\overline{\mathbb{B}_R(x)} \cap \Omega^c) < \epsilon = \gamma C'_2(\overline{\mathbb{B}_R(0)}).$$

Thus, $R < r_{\Omega, \gamma}$ for any given $\gamma \in (0, 1)$, and hence, $\rho'_2(\Omega) \leq \inf\{r_{\Omega, \gamma} : \gamma \in (0, 1)\}$.

Next, let $R > \rho'_2(\Omega)$. Then, there exists an $\epsilon > 0$ such that $C'_2(\overline{\mathbb{B}_R(x)} \cap \Omega^c) \geq \epsilon$ for all $x \in \mathbb{R}^n$. Let $\gamma_0 = \min\{1, \epsilon/C'_2(\overline{\mathbb{B}_R(0)})\}$. It then follows that

$$C'_2(\overline{\mathbb{B}_R(x)} \cap \Omega^c) > \gamma C'_2(\overline{\mathbb{B}_R(0)}) \quad \forall \gamma \in (0, \gamma_0)$$

for all $x \in \mathbb{R}^n$. That is, $R > r_{\Omega, \gamma}$ for all $\gamma \in (0, \gamma_0)$. Since $r_{\Omega, \gamma}$ is increasing in γ , we obtain that $R > \inf\{r_{\Omega, \gamma} : \gamma \in (0, 1)\}$. Hence, $\rho'_2(\Omega) \geq \inf\{r_{\Omega, \gamma} : \gamma \in (0, 1)\}$. \square

Understanding the p -capacitary inradius as limit of Maz'ya–Shubin-like inradii as in Lemma 2.14 comes in handy in the proof of part (ii) of Theorem 1.6. We define

$$r_{p, \gamma}(\Omega) := \sup\{r > 0 : \exists x \in \mathbb{R}^n \text{ such that } C_p(\mathbb{B}_r(x) \cap \Omega^c) \leq \gamma C_p(\mathbb{B}_r(0))\}$$

for $\gamma \in (0, 1)$ and $1 < p < \infty$. A proof similar to the one given in Lemma 2.14 then yields the following.

Corollary 2.15. *Let $\Omega \subset \mathbb{R}^n$ and $1 < p < \infty$. Then $\rho_p(\Omega) = \inf\{r_{p, \gamma}(\Omega) : \gamma \in (0, 1)\}$.*

2.3. Infimum of the nonlinear Rayleigh quotient for p -Laplacian.

Definition 2.16. *Let $\Omega \subset \mathbb{R}^n$ be an open set, $1 < p < \infty$. Then*

$$\lambda_{1, p}(\Omega) := \inf \left\{ \frac{\|\nabla u\|_{p, \Omega}^p}{\|u\|_{p, \Omega}} : u \in C_c(\Omega) \setminus \{0\} \right\}.$$

We collect some elementary properties of $\lambda_{1, p}$ in the following lemma.

Lemma 2.17. *Let $\Omega \subset \mathbb{R}^n$ be an open set. Then*

- (i) *If $\Omega' \subset \Omega$ is an open set, then $\lambda_{1, p}(\Omega) \leq \lambda_{1, p}(\Omega')$.*
- (ii) *If $x \in \mathbb{R}^n$, then $\lambda_{1, p}(\Omega + x) = \lambda_{1, p}(\Omega)$.*
- (iii) *If $s > 0$, then*

$$\lambda_{1, p}(s\Omega) = \lambda_{1, p}(\Omega)s^{-p}.$$

- (iv) *If $\Omega' \subset \Omega$ is an open set such that $\Omega \setminus \Omega'$ is p -polar, then $\lambda_{1, p}(\Omega) = \lambda_{1, p}(\Omega')$.*

Proof. Parts (i)–(iii) follow from the definition, the translation invariance of the Lebesgue measure, and a change of variable argument, respectively. For part (iv), we note that if $\Omega \setminus \Omega'$ is p -polar, then $W_0^{1, p}(\Omega) = W_0^{1, p}(\Omega')$, see, for instance, [8, Theorem 2.43]. Hence, by definition, $\lambda_{1, p}(\Omega) = \lambda_{1, p}(\Omega')$. \square

If $\lambda_{1, p}(\Omega)$ is positive and attained at some $u \in W_0^{1, p}(\Omega)$, then u is a weak solution to

$$(2.18) \quad \begin{cases} \operatorname{div}(|\nabla u|^{p-2} \nabla u) + \lambda |u|^{p-2} u = 0 & \text{in } \Omega \\ u = 0 & \text{on } \partial\Omega \end{cases}$$

for $\lambda = \lambda_{1, p}(\Omega)$. Such a function u is called a first eigenfunction of Δ_p , where

$$\Delta_p u := \operatorname{div}(|\nabla u|^{p-2} \nabla u).$$

Here, u being a weak solution to (2.18) means that

$$(2.19) \quad \int_{\Omega} |\nabla u|^{p-2} \nabla u \circ \nabla \chi \, dm = \lambda \int_{\Omega} |u|^{p-2} u \chi \, dm$$

for all $\chi \in C_c^\infty(\Omega)$.

It is well-known that if $\Omega \Subset \mathbb{R}^n$ is a smoothly bounded domain, then $\lambda_{1,p}(\Omega)$ is positive and attained at some non-zero $u \in W_0^{1,p}(\Omega) \cap C(\overline{\Omega})$. Moreover, such u may be assumed to be positive in Ω . A comprehensive resource for standard results on the first eigenvalue and eigenfunctions for Δ_p are the lecture notes by Lindqvist [11]. The regularity result for the first eigenfunctions is due to Gariepy–Ziemer in [7] for $1 < p \leq n$; in the case of $p > n$, it is well-known that, under these conditions on Ω , a representative of $u \in W^{1,p}(\Omega)$ is in $C(\overline{\Omega})$, see, e.g., Theorem 5 in §5.6.2 in [4].

3. PROOF OF THEOREM 1.3

The following lemma is crucial for the proof of Theorem 1.3.

Lemma 3.1. *Let $D \Subset \mathbb{R}^n$ be a smoothly bounded domain. Let $K_j \subset \overline{D}$, $j \in \mathbb{N}$, be compact sets. Suppose that $\lim_{j \rightarrow \infty} C_p(K_j) = 0$ and $D_j := D \setminus K_j$ is an open set with smooth boundary for all $j \in \mathbb{N}$. Then, $\lim_{j \rightarrow \infty} \lambda_{1,p}(D_j) = \lambda_{1,p}(D)$.*

Proof. Note first that if $p > n$, then $\lim_{j \rightarrow \infty} C_p(K_j) = 0$ implies that K_j is empty for all j sufficiently large, and hence, the conclusion holds trivially.

Since D is a bounded domain, $\lambda_{1,p}(D) > 0$, and there exists a weak solution, $\varphi \in W_0^{1,p}(D) \cap C(\overline{D})$, to (2.18) which is positive on D , see Section 2.3 for references. After rescaling, we may further assume that $0 < \varphi \leq 1$ on D . Next, for each $j \in \mathbb{N}$, there exists a weak solution $h_j \in W^{1,p}(D_j)$ to the Dirichlet problem for Δ_p with boundary data φ on D_j , i.e.,

$$\begin{cases} \Delta_p h_j = 0 & \text{in } D_j \\ h_j = \varphi & \text{on } bD_j \end{cases}'$$

see, e.g., [10, Th. 2.16]. It follows from work by Maz'ya [14], see also [10, Th. 2.16], that $h_j \in C(\overline{D_j})$, and hence $h_j = 0$ on $bD_j \cap bD$ and $0 < h_j \leq 1$ on $bD_j \setminus bD$. Finally, set $\psi_j = \varphi - h_j$ on D_j . Then, $\psi_j \in W_0^{1,p}(D_j) \cap C(\overline{D_j})$ and $\Delta_p \psi_j = \Delta_p \varphi$ holds weakly on D_j . Using (2.19), we compute

$$\begin{aligned} \|\nabla \psi_j\|_{p,D_j}^p &= \int_{D_j} |\nabla \psi_j|^{p-2} \nabla \psi_j \circ \nabla \psi_j \, dm \\ &= \int_{D_j} |\nabla \varphi|^{p-2} \nabla \varphi \circ \nabla \psi_j \, dm \\ &= \lambda_{1,p}(D) \int_{D_j} |\varphi|^{p-2} \varphi \cdot \psi_j \, dm \leq \lambda_{1,p}(D) \|\varphi\|_{p,D_j}^{p-1} \cdot \|\psi_j\|_{p,D_j}, \end{aligned}$$

where the last step follows from Hölder inequality. Using this estimate, in conjunction with the definition of $\lambda_{1,p}(D_j)$, yields

$$\frac{1}{\lambda_{1,p}(D_j)} \geq \frac{\|\psi_j\|_{p,D_j}^p}{\|\nabla \psi_j\|_{p,D_j}^p} \geq \frac{\|\psi_j\|_{p,D_j}^{p-1}}{\lambda_{1,p}(D) \cdot \|\varphi\|_{p,D_j}^{p-1}},$$

and therefore,

$$\left(\frac{\lambda_{1,p}(D)}{\lambda_{1,p}(D_j)}\right)^{\frac{1}{p-1}} \geq \frac{\|\psi_j\|_{p,D_j}}{\|\varphi\|_{p,D_j}} \geq \frac{\|\varphi\|_{p,D_j} - \|h_j\|_{p,D_j}}{\|\varphi\|_{p,D_j}} \geq 1 - \frac{\|h_j\|_{p,D_j}}{\|\varphi\|_{p,D_j}}.$$

Since φ is bounded on D and $\lim_{j \rightarrow \infty} C_p(D \setminus D_j) = 0$, so that $\lim_{j \rightarrow \infty} m(D \setminus D_j) = 0$ as well, it follows from the Dominated Convergence Theorem that

$$\lim_{j \rightarrow \infty} \|\varphi\|_{p,D_j} = \|\varphi\|_{p,D}.$$

Hence, it suffices to show that $\lim_{j \rightarrow \infty} \|h_j\|_{p,D_j} = 0$.

For that, define

$$U_j = \{u \in \mathcal{S} : u \equiv 1 \text{ near } K_j, 0 \leq u \leq 1\}$$

for each $j \in \mathbb{N}$. Temporarily fix $j \in \mathbb{N}$ and $u \in U_j$. Then, we may find a $g_j \in W^{1,p}(D_j) \cap C(\overline{D_j})$ which is a weak solution to

$$(3.2) \quad \begin{cases} \Delta_p g_j = 0 & \text{in } D_j \\ g_j = u & \text{on } bD_j \end{cases}.$$

Note that $h_j \leq g_j$ holds on D_j by the comparison principle for p -harmonic functions, see, e.g., [10, Th. 2.15], and the fact that $\varphi \leq u$ on bD_j . Thence,

$$(3.3) \quad \|h_j\|_{p,D_j} \leq \|g_j\|_{p,D_j} \leq \|g_j - u\|_{p,D_j} + \|u\|_{p,D_j}.$$

By (3.2), $g_j - u \in W_0^{1,p}(D_j)$. Thus, since $D_j \subset D$, Poincaré inequality is applicable to the first term on the right hand side of (3.3) with a uniform constant, i.e.,

$$\|g_j - u\|_{p,D_j} \leq C \|\nabla g_j - \nabla u\|_{p,D_j}$$

with $C = (\lambda_{1,p}(D))^{-1/p}$. Therefore, we have

$$\|h_j\|_{p,D_j} \leq C \|\nabla g_j\|_{p,D_j} + C \|\nabla u\|_{p,D_j} + \|u\|_{p,D_j}.$$

Next, note that (3.2) implies that g_j is a quasi-minimizer, i.e.,

$$\|\nabla g_j\|_{p,D_j} \leq \|\nabla f\|_{p,D_j}$$

for any $f \in W^{1,p}(D_j)$ with $g_j - f \in W_0^{1,p}(D_j)$, in particular, for $f = u$, see, e.g., [10, Th. 2.15]. Thus,

$$\|h_j\|_{p,D_j} \leq 2C \|\nabla u\|_{p,D_j} + \|u\|_{p,D_j}.$$

Therefore,

$$\begin{aligned} \|h_j\|_{p,D_j}^p &\leq 2^{p-1}(2C+1) \cdot \inf \left\{ \|u\|_{p,D_j}^p + \|\nabla u\|_{p,D_j}^p : u \in U_j \right\} \\ &= 2^{p-1}(2C+1)C_p(K_j), \end{aligned}$$

where the last step follows from the fact that $C_p(K_j) = \inf \left\{ \|u\|_{p,D_j}^p + \|\nabla u\|_{p,D_j}^p : u \in U_j \right\}$. This can be proved the same was as (ii) in §2.2.1 in [15]. This concludes the proof since $\lim_{j \rightarrow \infty} C_p(K_j) = 0$ by hypothesis. \square

The following is a slight variation of [5, Proposition 6.1].

Lemma 3.4. *Let $K \subset \mathbb{R}^n$ be a compact set such that $\mathbb{B}_1(0) \cap K \neq \emptyset$. Then, for any $\epsilon > 0$, there exists a compact set $K_\epsilon \subset \overline{\mathbb{B}_1(0)}$ such that*

- (i) $K \cap \overline{\mathbb{B}_1(0)} \subset K_\epsilon$,
- (ii) $C_p(K_\epsilon) \leq C_p(K) + \epsilon$,
- (iii) $\mathbb{B}_1(0) \setminus K_\epsilon$ has smooth boundary.

The above lemma differs from [5, Proposition 6.1] in two ways. On the one hand K_ϵ is a compact subset of $\overline{\mathbb{B}_1(0)}$ instead of a relatively compact subset of $\mathbb{B}_1(0)$, and on the other hand, the Sobolev p -capacity is used instead of the Newtonian capacity. That the latter change is acceptable is due to the fact that only the properties of monotonicity, countable subadditivity and outer regularity are used. To ensure that the former change is also correct, one only needs to set $K_\epsilon = \overline{\Omega} \cap \overline{\mathbb{B}_1(0)}$ in the last paragraph of the proof of [5, Proposition 6.1].

Now we are set to prove part (i) of Theorem 1.3.

Proof of part (i) of Theorem 1.3. Suppose that $\lambda_{1,p}(\Omega) > 0$ and

$$(\rho_p(\Omega))^p > \lambda_{1,p}(\mathbb{B}_1(0))/\lambda_{1,p}(\Omega)$$

hold. Then, we may choose a positive $R < \rho_p(D)$ such that

$$(3.5) \quad R^p > \lambda_{1,p}(\mathbb{B}_1(0))/\lambda_{1,p}(\Omega).$$

Let us first consider the case $p > n$. Then, $\rho_p(\Omega) = \mathfrak{R}(\Omega)$, and hence, there exists an $x \in \mathbb{R}^n$ such that $\mathbb{B}_R(x) \subset \Omega$. Using (3.5) as well as (i) and (iii) of Lemma 2.17, we then obtain

$$\lambda_{1,p}(\mathbb{B}_R(x)) \geq \lambda_{1,p}(\Omega) > \lambda_{1,p}(\mathbb{B}_1(0))R^{-p} = \lambda_{1,p}(\mathbb{B}_R(0)).$$

This is a contradiction, and hence, $\mathfrak{R}(\Omega) \leq \lambda_{1,p}(\mathbb{B}_1(0))/\lambda_{1,p}(\Omega)$.

For the case of $p \in (1, n]$, it follows from the definition of $\rho_p(\Omega)$ and (3.5) that for all $j \in \mathbb{N}$ there exists a $x_j \in \mathbb{R}^n$ with

$$C_p(\overline{\mathbb{B}_R(x_j)} \cap \Omega^c) < \frac{1}{2j^\gamma},$$

where $\gamma := R^{-n} \max\{1, R^p\}$. For each $j \in \mathbb{N}$, define $\mathfrak{A}_j \subset \overline{\mathbb{B}_1(0)}$ by setting

$$\mathfrak{A}_j = \{x \in \mathbb{R}^n : Rx + x_j \in \overline{\mathbb{B}_R(x_j)} \cap \Omega^c\}.$$

Then, it follows from parts (ii) and (iii) of Lemma 2.4 that $C_p(\mathfrak{A}_j) < 1/(2j)$. Now, for each $j \in \mathbb{N}$, we may apply Lemma 3.4 to the pair $(\mathfrak{A}_j, 1/(2j))$ to obtain a compact set A_j such that $C_p(A_j) \leq 1/j$, $\Omega_j := \mathbb{B}_1(0) \setminus A_j$ is an open set with smooth boundary, and $\mathfrak{A}_j \cap \overline{\mathbb{B}_1(0)} \subset A_j$. The latter implies that

$$R\Omega_j + x_j \subset \Omega.$$

It then follows from parts (i)–(iii) of Lemma 2.17

$$\lambda_{1,p}(\Omega_j) = R^p \lambda_{1,p}(R\Omega_j + x_j) \geq R^p \lambda_{1,p}(\Omega).$$

Hence, by (3.5), there is some $\epsilon > 0$ such that

$$\lambda_{1,p}(\Omega_j) > \lambda_{1,p}(\mathbb{B}_1(0)) + \epsilon$$

for all $j \in \mathbb{N}$. This is a contradiction to Lemma 3.1, with $D = \mathbb{B}_1(0)$ and $K_j = A_j$, and it follows that

$$\left(\rho_p(\Omega)\right)^p \leq \lambda_{1,p}(\mathbb{B}_1(0))/\lambda_{1,p}(\Omega).$$

□

The second part of Theorem 1.3 is proved by using the following lemma.

Lemma 3.6. *Let $\Omega_1 \subset \Omega_2$ be open sets in \mathbb{R}^n , Ω_2 connected and bounded. If $\lambda_{1,p}(\Omega_1) = \lambda_{1,p}(\Omega_2)$, then $\Omega_2 \setminus \Omega_1$ is p -polar.*

The necessity of the connectedness assumption in Lemma 3.6 can be seen by considering the example of $\Omega_1 = \mathbb{B}_1(0)$ and $\Omega_2 = \mathbb{B}_1(0) \cup \mathbb{B}_1(x)$ for some $x \notin \overline{\mathbb{B}_2(0)}$. In this case $\lambda_{1,p}(\Omega_1) = \lambda_{1,p}(\Omega_2)$ while $\Omega_2 \setminus \Omega_1$ is p -polar.

Although, this result is presumably well-known, at least for $p = 2$, we present a proof of Lemma 3.6 here for the sake of completeness.

Proof. Suppose that $\lambda_{1,p}(\Omega_1) = \lambda_{1,p}(\Omega_2)$. Since Ω_1 is bounded, $\lambda_{1,p}(\Omega_1)$ is positive and there exists a $\varphi \in W_0^{1,p}(\Omega_1)$ which is an eigenfunction of Δ_p with eigenvalue $\lambda_{1,p}(\Omega_1)$. Then, there exists $\tilde{\varphi} \in W_0^{1,p}(\mathbb{R}^n)$ such that $\tilde{\varphi} = \varphi$ almost everywhere on Ω_1 and $\tilde{\varphi} = 0$ p -quasi everywhere on \mathbb{R}^n , see, e.g., [8, Theorem 4.5]. It follows that $\tilde{\varphi}$ is not zero p -quasi everywhere on Ω_2 and $\tilde{\varphi} \in W_0^{1,p}(\Omega_2)$ since $\Omega_1 \subset \Omega_2$. Hence, $\tilde{\varphi}$ is an eigenfunction with eigenvalue $\lambda_{1,p}(\Omega_1)$ for Δ_p on Ω_2 . The assumption $\lambda_{1,p}(\Omega_1) = \lambda_{1,p}(\Omega_2)$ then implies that $\tilde{\varphi}$ is a first eigenfunction for Δ_p on Ω_2 . Since Ω_2 is connected and bounded, $\tilde{\varphi}$ is non-zero almost everywhere on Ω_2 . This is a contradiction to $\tilde{\varphi} = 0$ p -quasi everywhere on Ω_2 unless $\Omega_2 \setminus \Omega_1$ is p -polar. \square

Proof of part (ii) of Theorem 1.3. Suppose Ω is bounded. Then, $\lambda_{1,p}(\Omega) > 0$, hence $\rho_p(\Omega) < \infty$ by part (i) of Theorem 1.3. Moreover, by (iv) of Lemma 2.9, there exists an x° such that

$$C_p(\mathbb{B}_{\rho_p(\Omega)}(x^\circ) \cap \Omega^c) = 0.$$

This means that there exists a set K with $C_p(K) = 0$, which is relatively closed in $\mathbb{B}_{\rho_p(\Omega)}(x^\circ)$, such that

$$(\mathbb{B}_{\rho_p(\Omega)}(x^\circ) \setminus K) \subset \Omega.$$

By hypothesis,

$$\lambda_{1,p}(\Omega) = \lambda_{1,p}(\mathbb{B}_{\rho_p(\Omega)}(0)) = \lambda_{1,p}(\mathbb{B}_{\rho_p(\Omega)}(x^\circ) \setminus K).$$

Lemma 3.6 is now applicable with $\Omega_1 = \mathbb{B}_{\rho_p(\Omega)}(x^\circ) \setminus K$ and $\Omega_2 = \Omega$. Thus, we obtain that $\Omega \setminus (\mathbb{B}_{\rho_p(\Omega)}(x^\circ) \setminus K)$ is p -polar which concludes the proof of (ii) for the bounded case.

Next, suppose Ω is unbounded such $\rho_p(\Omega) = \rho_p(\Omega \cap \mathbb{B}_R(0))$ for some $R > 0$, and

$$\lambda_{1,p}(\mathbb{B}_1(0))(\rho_p(\Omega))^{-p} = \lambda_{1,p}(\Omega).$$

For ease of notation, write Ω_R in place of $\Omega \cap \mathbb{B}_R(0)$. Without loss of generality, we may assume that $\rho_p(\Omega_R) < R$. Part (i) of Lemma 2.17 and (1.4) for Ω_R yield

$$\lambda_{1,p}(\Omega) \leq \lambda_{1,p}(\Omega_R) \leq \lambda_{1,p}(\mathbb{B}_1(0))(\rho_p(\Omega_R))^{-p}.$$

Thus, it follows from the hypothesis and the two preceding estimates that

$$(3.7) \quad \lambda_{1,p}(\Omega_R) = \lambda_{1,p}(\mathbb{B}_1(0))(\rho_p(\Omega_R))^{-p}.$$

Denote by $\{Z_j\}_{j \in \mathbb{N}}$ the set of connected components of Ω_R , and note that

$$(3.8) \quad \begin{aligned} \rho_p(\Omega_R) &= \sup \{ \rho_p(Z_j) : j \in \mathbb{N} \}, \\ \lambda_{1,p}(\Omega_R) &= \inf \{ \lambda_{1,p}(Z_j) : j \in \mathbb{N} \}. \end{aligned}$$

Since Ω_R is bounded and $\rho_p(\Omega_R) > 0$, it follows that $\rho_p(\Omega_R)$ is attained at $\rho_p(Z_{j_0})$ for some $j_0 \in \mathbb{N}$. Then, (1.4) yields

$$\begin{aligned} 0 < \lambda_{1,p}(\Omega_R) &\leq \lambda_{1,p}(Z_{j_0}) \leq \lambda_{1,p}(\mathbb{B}_1(0)) \left(\rho_p(Z_{j_0})\right)^{-p} \\ &= \lambda_{1,p}(\mathbb{B}_1(0)) \left(\rho_p(\Omega_R)\right)^{-p} \end{aligned}$$

Thus, by (3.7) and the choice of j_0 , it follows that $\lambda_{1,p}(\Omega_R) = \lambda_{1,p}(Z_{j_0})$. We now may apply part (ii) of Theorem 1.3 for the bounded case to Z_{j_0} . That is, there exists an $x \in \mathbb{R}^n$ such that Z_{j_0} is equal to $\mathbb{B}_{\rho_p(\Omega_R)}(x)$ modulo a p -polar set K . Now, either the boundary of $\mathbb{B}_{\rho_p(\Omega_R)}(x)$ meets the boundary of $\mathbb{B}_R(0)$ tangentially or the closure of $\mathbb{B}_{\rho_p(\Omega_R)}(x)$ is contained in $\mathbb{B}_R(0)$. In the latter case, it follows that Z_{j_0} is actually a connected component of Ω , and hence $\Omega = Z_{j_0}$ which is a contradiction to Ω being unbounded. In the former case, it follows from the openness of Ω and $\rho_p(\Omega_R) < R$ that Z_{j_0} is in fact a connected component of Ω which, again, is a contradiction. Thus, equality in (1.4) cannot hold. \square

4. PROOF OF THEOREM 1.6

The next theorem is the essence of Theorem 1.6. It is a special case of Theorem 8.3.3 in [1] which is formulated for $(1, p)$ -extension domains. A domain Ω is of this class, if there exists a linear, bounded operator $E_\Omega : W^{1,p}(\Omega) \rightarrow W^{1,p}(\mathbb{R}^n)$ such that $(E_\Omega f)|_\Omega = f$ for $f \in W^{1,p}(\Omega)$.

Theorem 4.1. *Let $\Omega \subset \mathbb{R}^n$ be a bounded $(1, p)$ -extension domain, and suppose that $f \in W^{1,p}(\Omega)$, $1 < p < \infty$. Let K be a closed subset of Ω such that $C_p(K) > 0$. Suppose that $f|_K = 0$, then*

$$(4.2) \quad \|f\|_{p,\Omega} \leq (m(\Omega))^{\frac{1}{p}} \cdot \|E_\Omega\|_{\text{op}} \cdot \frac{\|\nabla f\|_{p,\Omega}}{(C_p(K))^{1/p}},$$

where $m(\Omega)$ is the Lebesgue measure of Ω and $\|E_\Omega\|_{\text{op}}$ is the operator norm of a bounded, linear extension operator $E_\Omega : W^{1,p}(\Omega) \rightarrow W^{1,p}(\mathbb{R}^n)$ such that $(E_\Omega f)|_\Omega = f$ for $f \in W^{1,p}(\Omega)$.

Proof. This follows straightforwardly from Theorem 8.3.3 and Lemma 8.3.2 in [1]. First, consider Lemma 8.3.2, with $m = 1$, $\sigma = 0$, so that $a_0 = \int_\Omega f \, d\mu_0$ and

$$\|L f\|_{W^{1,p}(\Omega)} = a_0 \cdot (m(\Omega))^{1/p}.$$

Then, use Theorem 8.3.3, with $m = 1$, $\beta = 0$, and $\|G_1 * \mu_0\|_{p'} = C_p(K)^{-1/p}$. Collecting the constants determining A in (8.3.7), we get $A = (m(\Omega))^{1/p} \cdot \|E_\Omega\|_{\text{op}}$. \square

Proof of Theorem 1.6. It follows from Calderón's work in [3, Theorem 12] that any open square in \mathbb{R}^n is a $(1, p)$ -extension domain. Hence, for given $R > 0$, there exist bounded, linear operators from $W^{1,p}((0, R)^n)$ to $W^{1,p}(\mathbb{R}^n)$ which are the identity operator on $(0, R)^n$. Let E_R be such an operator.

Suppose $\rho_p(\Omega)$ is finite. Let $R > \rho_p(\Omega)$ and $\delta_R(\Omega)$ be defined as in (1.5). We note that arguments analogous to the ones used in the proof of Lemma 2.7 yield

$$\delta_R(\Omega) = \sup\{\delta > 0 : C_p(\mathbb{B}_R(x) \cap \Omega^c) \geq \delta \, \forall x \in \mathbb{R}^n\}.$$

Now, for each $m \in \mathbb{Z}^n$, set

$$Q_m = \{x \in \mathbb{R}^n : x_j \in (Rm_j, R(m_j + 1)) \quad \forall j \in \{1, \dots, n\}\}.$$

Since

$$\int_{\mathbb{R}^n} |\cdot|^p dm = \sum_{m \in \mathbb{Z}^n} \int_{Q_m} |\cdot|^p dm,$$

it suffices to show that

$$\|f\|_{p, Q_m} \leq \frac{R^{n/p} \|E_R\|_{\text{op}}}{(\delta_R(\Omega))^{1/p}} \cdot \|\nabla f\|_{p, Q_m} \quad \forall f \in C_c^\infty(\Omega)$$

holds for each $m \in \mathbb{Z}^n$.

For that, let $\eta \in (0, \delta_R(\Omega))$ be fixed. Then, by our choice of R , we may choose a compact set $K_m \subset Q_m \cap \Omega^c$ such that $C_p(K_m) \geq \eta$ for each $m \in \mathbb{Z}^n$. Since $f \in C_c^\infty(\Omega)$ implies that $f \in C^\infty(\overline{Q_m})$ with $f|_{K_m} = 0$ for all $m \in \mathbb{Z}^n$, Theorem 4.1 is applicable here and yields

$$\|f\|_{p, Q_m} \leq \frac{R^{n/p} \|E_R\|_{\text{op}}}{\eta^{1/p}} \|\nabla f\|_{p, Q_m} \quad \forall f \in C_c^\infty(\Omega)$$

for all $m \in \mathbb{Z}^n$. As this holds for any $\eta \in (0, \delta_R(\Omega))$, the proof of part (i) of Theorem 1.6, i.e.,

$$\frac{\delta_R(\Omega)}{R^n \|E_R\|_{\text{op}}^p} \leq \lambda_{1,p}(\Omega)$$

is complete.

To prove part (ii) of Theorem 1.6, again, suppose $\rho_p(\Omega) < \infty$ and $R > \rho_p(\Omega)$. Set $\gamma_R(\Omega) = \delta_R(\Omega)/C_p(\mathbb{B}_R(0))$, i.e.,

$$\gamma_R(\Omega) = \sup\{\gamma \in (0, 1) : C_p(\mathbb{B}_R(x) \cap \Omega^c) \geq \gamma C_p(\mathbb{B}_R(0)) \quad \forall x \in \mathbb{R}^n\}.$$

This means that $r_{p,\gamma}(\Omega) < R$ for all $\gamma \in (0, \gamma_R(\Omega))$, see also Corollary 2.15. Furthermore, there exists a constant $c = c(n, p)$ such that

$$C_p(\mathbb{B}_R(0)) \geq cR^{n-p},$$

see Section 2.12 in [8]. Hence,

$$c\gamma_R(\Omega) \cdot R^{n-p} \leq \delta_R(\Omega).$$

Next, we may choose an extension operator E_R for $(0, R)^n$ such that $\|E_R\|_{\text{op}}$ only depends on n and p , and not on R , see Stein's construction of extension operators in [18, Theorem 5], in particular, note Theorem 5' and (c) on pg. 190 therein in regards to the operator norm dependencies. It follows that there exists a constant $C = C(n, p) > 0$ such that

$$(4.3) \quad C\gamma_R(\Omega) \cdot R^{-p} \leq \lambda_{1,p}(\Omega).$$

□

Remark. Estimate (4.3) is in line with the estimate obtained by Maz'ya–Shubin in case of $p = 2$, $n \geq 3$, in (3.19) of [13], since $r_{p,\gamma_R(\Omega)}(\Omega) = R$.

REFERENCES

- [1] ADAMS, D. R., AND HEDBERG, L. I. *Function spaces and potential theory*, vol. 314 of *Grundlehren Math. Wiss.* Berlin: Springer-Verlag, 1995.
- [2] AGMON, S. Lectures on elliptic boundary value problems. Prepared for publication by B. Frank Jones jun. with the assistance of George W. Batten jun. Van Nostrand Mathematical Studies. 2. Princeton, N.J.-Toronto-New York-London: D. van Nostrand Company, Inc. v, 291 p. (1965)., 1965.
- [3] CALDERÓN, A. P. Lebesgue spaces of differentiable functions and distributions. *Proc. Sympos. Pure Math.* 4, 33-49 (1961)., 1961.
- [4] EVANS, L. C. *Partial differential equations*, vol. 19 of *Graduate Studies in Mathematics*. American Mathematical Society, 1998.
- [5] GALLAGHER, A.-K. On the Poincaré inequality on open sets in \mathbb{R}^n , to appear in *Computational Methods and Function Theory*.
- [6] GALLAGHER, A.-K., LEBL, J., AND RAMACHANDRAN, K. The closed range property for the $\bar{\partial}$ -operator on planar domains. *J. Geom. Anal.* 31, 2 (2021), 1646–1670.
- [7] GARIEPY, R., AND ZIEMER, W. P. A regularity condition at the boundary for solutions of quasilinear elliptic equations. *Arch. Ration. Mech. Anal.* 67 (1977), 25–39.
- [8] HEINONEN, J., KILPELÄINEN, T., AND MARTIO, O. *Nonlinear potential theory of degenerate elliptic equations. With a new preface, corrigenda, and epilogue consisting of other new material*, reprint of the 2006 edition ed. Dover Books Math. Mineola, NY: Dover Publications, 2018.
- [9] LIEB, E. H. On the lowest eigenvalue of the Laplacian for the intersection of two domains. *Invent. Math.* 74 (1983), 441–448.
- [10] LINDQVIST, P. *Notes on the p -Laplace equation*, vol. 102 of *Rep., Univ. Jyväskylä, Dep. Math. Stat.* Jyväskylä: University of Jyväskylä, Department of Mathematics and Statistics, 2006.
- [11] LINDQVIST, P. A nonlinear eigenvalue problem. In *Topics in mathematical analysis*. Hackensack, NJ: World Scientific, 2008, pp. 175–203.
- [12] LITTMAN, W. A connection between α -capacity and $m - p$ polarity. *Bull. Am. Math. Soc.* 73 (1967), 862–866.
- [13] MAZ'YA, V., AND SHUBIN, M. Can one see the fundamental frequency of a drum? *Letters in Mathematical Physics* 74 (2005), 135–1151.
- [14] MAZ'YA, V. G. On continuity in a boundary point of solutions of quasilinear elliptic equations. *Vestn. Leningr. Univ., Mat. Mekh. Astron.* 25, 3 (1970), 42–55.
- [15] MAZ'YA, V. G. *Sobolev spaces. With applications to elliptic partial differential equations. Transl. from the Russian by T. O. Shaposhnikova*, 2nd revised and augmented ed. ed., vol. 342 of *Grundlehren Math. Wiss.* Berlin: Springer, 2011.
- [16] MEYERS, N. G. Integral inequalities of Poincaré and Wirtinger type. *Arch. Ration. Mech. Anal.* 68 (1978), 113–120.
- [17] SOUPLLET, P. Geometry of unbounded domains, Poincaré inequalities and stability in semilinear parabolic equations. *Comm. Partial Differential Equations* 24, 5-6 (1999), 951–973.
- [18] STEIN, E. M. *Singular integrals and differentiability properties of functions*, vol. 30 of *Princeton Math. Ser.* Princeton University Press, Princeton, NJ, 1970.
- [19] VITOLO, A. $H^{1,p}$ -eigenvalues and L^∞ -estimates in quasicylindrical domains. *Commun. Pure Appl. Anal.* 10, 5 (2011), 1315–1329.
- [20] WALLIN, H. A connection between α -capacity and L^p -classes of differentiable functions. *Ark. Mat.* 5 (1965), 331–341.

GALLAGHER TOOL & INSTRUMENT LLC, REDMOND, WA 98052, USA
 Email address: anne.g@gallagherti.com