

STARSHAPED COMPACT HYPERSURFACES IN WARPED PRODUCT MANIFOLDS I: PRESCRIBED CURVATURE EQUATIONS

BIN WANG

ABSTRACT. In [Starshaped compact hypersurfaces with prescribed Weingarten curvature in warped product manifolds. *Calc. Var. Partial Differential Equations* **57** (2018), no.2, Paper No. 42.], Chen-Li-Wang obtained global curvature estimates for $(n-1)$ -convex starshaped compact hypersurfaces Σ in warped product manifolds, satisfying the prescribed $(n-1)$ -curvature equation

$$\sigma_{n-1}(\kappa(X)) = \Psi(X, \nu) \quad \text{for all } X \in \Sigma.$$

In this note, we prove the same result for $(n-2)$ -convex starshaped compact hypersurfaces satisfying the prescribed $(n-2)$ -curvature equation

$$\sigma_{n-2}(\kappa(X)) = \Psi(X, \nu) \quad \text{for all } X \in \Sigma.$$

The main ingredient of our proof is a new test function and a notable feature of which is that it also works for the $n-1$ case. Furthermore, the proof happens to be simpler than that of Chen-Li-Wang, because we have synthesized techniques from several authors, the most inspiring of which is the recent breakthrough made by Guan, Ren and Wang in [Global C^2 -estimates for convex solutions of curvature equations. *Comm. Pure Appl. Math.* 68 (2015), no. 8, 1287–1325.].

As an application of our method, we obtain global C^2 estimates for semi-convex and $(k+1)$ -convex solutions of the general k -curvature equation

$$\sigma_k(\kappa(X)) = \Psi(X, \nu) \quad \text{for all } X \in \Sigma.$$

We also establish the existence of k -convex starshaped compact hypersurfaces satisfying the prescribed curvature measure type equation

$$\sigma_k(\kappa(X)) = \langle X, \nu \rangle^p \psi(X) \quad \text{for all } X \in \Sigma$$

where $p \in (-\infty, 0) \cup (0, 1]$. In particular, the C^1 estimate for this equation holds for all $p \neq 0$ without any barrier conditions.

CONTENTS

1. Introduction	2
Acknowledgement	6
2. Preliminaries	6
3. A Model Computation	10
4. The Prescribed Curvature Equation in General	16
4.1. When $k = n-1, n \geq 3$ or $k = n-2, n \geq 5$	18
4.2. When Σ is semi-convex or $(k+1)$ -convex	18
5. The Prescribed Curvature Measure Type Equation	20
5.1. The C^0 Estimate	21
5.2. The C^1 Estimate for all $p \in \mathbb{R}$	21
5.3. The C^2 Estimate for $p \leq 1$	25
References	30

2020 *Mathematics Subject Classification.* Primary 53C21; Secondary 35J60.

Key words and phrases. Starshaped compact hypersurfaces, prescribed Weingarten curvature equations, warped product manifolds, curvature estimates, the prescribed curvature measure problem.

1. INTRODUCTION

It is evident that the study of the σ_k equation has become a central objective in global differential geometry. A few classical examples include: the Minkowski problem [2, 11, 30, 41, 42, 45, 46], the Alexandrov-Chern problem [1, 12, 13], and the problem of finding closed strictly convex hypersurfaces whose Weingarten curvature is prescribed on $\mathbb{S}^n$ in terms of its inverse Gauss map [18]. More recent examples include: the prescribed curvature measure problem [19, 21, 54], the σ_k -Nirenberg problem [34–36], and the Christoffel-Minkowski problem [22–24].

Here we are interested in a natural variation of the aforementioned problems stated as follows¹: under what conditions a given sufficiently smooth, positive function $\Phi(X, \nu)$ is the k -th mean curvature $\sigma_k(\kappa(X))$ of a smooth closed hypersurface Σ embedded in a space form of constant sectional curvature as a graph over a sphere?

Formulating it as a PDE problem, we are concerned with the existence of star-shaped compact solutions Σ in space forms to the curvature equation

$$(1.1) \quad \sigma_k(\kappa(X)) = \Psi(X, \nu) \quad \text{for all } X \in \Sigma,$$

for which establishing a priori estimates is a crucial step. The C^0 and C^1 estimates have been derived long ago under barrier conditions; see for example, [6] or [50]. However, the C^2 estimates had remained open due to the appearance of a gradient term on the right-hand side and can only be obtained in some special cases.

For $k = 1$, the equation is quasi-linear and the required estimates follow from classical theory of quasi-linear PDEs; see [4, 5, 51]. While for $k = n$, the equation is of Monge-Ampère type and the estimates follow from Caffarelli-Nirenberg-Spruck's seminal work [7] on the Monge-Ampère equation; see [16, 33, 43, 44].

For $2 \leq k \leq n - 1$, if Ψ is independent of ν i.e. $\Psi(X, \nu) = \psi(X)$, then the problem has been completely settled by Caffarelli-Nirenberg-Spruck [8] in $\mathbb{R}^{n+1}$, by Jin-Li [29] in $\mathbb{H}^{n+1}$, by Li-Oliker [37] in $\mathbb{S}^{n+1}$ and by De Andrade-Barbosa-De Lira [15] in warped product manifolds. On the other hand, if the right-hand side only depends on ν i.e. $\Psi(X, \nu) = \varphi(\nu)$, this problem is solved in $\mathbb{R}^{n+1}$ for convex solutions by Guan-Guan in [18]. Ivochkina [27, 28] considered the Dirichlet problem for (1.1) in bounded domains in $\mathbb{R}^n$ and obtained the C^2 estimates under extra conditions on the dependence of Ψ on ν .

This longstanding problem of obtaining C^2 estimates for equation (1.1) when $2 \leq k \leq n - 1$ and when the right-hand side depends on ν has recently been raised again by Guan, Li and Li in [19, Remark 3.5]: suppose that there is an a priori C^1 bound of Σ . Can one conclude there is an a priori C^2 bound of Σ in terms of the C^1 norm of Σ , Ψ , n and k ?

In 2015, a major breakthrough was made by Guan, Ren and Wang [25]: they established global C^2 estimates for

- convex hypersurfaces satisfying $\sigma_k(\kappa(X)) = \Psi(X, \nu)$ and
- 2-convex hypersurfaces satisfying $\sigma_2(\kappa(X)) = \Psi(X, \nu)$.

Their method was to employ a new test function

$$(1.2) \quad Q = \begin{cases} \frac{1}{2} \log \sum_l \kappa_l^2 - N \log \langle X, \nu \rangle & \text{for convex hypersurfaces} \\ \log \log \sum_l e^{\kappa_l} - (1 + \varepsilon) \log \langle X, \nu \rangle + \frac{a}{2} |X|^2 & \text{for 2-convex hypersurfaces} \end{cases}$$

which is nonlinear in principal curvatures but with sufficiently good convexity properties; this helped them overcome difficulties caused by allowing Ψ to depend on ν . See also the simpler proofs by Chu [14] and Spruck-Xiao [50] for both results respectively; Chen-Li-Wang [10] later generalized these two results to warped-product manifolds.

¹This question is a part of one of Yau's problems [55, problem 59], which concerns the existence of closed hypersurfaces of prescribed genus.

By using a modified version of the above test function

$$(1.3) \quad Q = \log \log \sum_l e^{\kappa_l} - N \log \langle X, \nu \rangle,$$

Ren and Wang [47, 48] solved, in a rather powerful manner, the same problem in $\mathbb{R}^{n+1}$ when $k = n - 1$ and $k = n - 2$. Our first result is a simpler proof for their C^2 estimates and we extend them to warped product manifolds. In particular, the $k = n - 2$ case in e.g. the hyperbolic space $\mathbb{H}^{n+1}$ is new.

Theorem 1.1. *Let $k = n - 1, n \geq 3$ or $k = n - 2, n \geq 5$. Suppose that Σ is a closed strictly star-shaped k -convex hypersurface in a warped product manifold $\bar{M}$ satisfying the curvature equation (1.1) for some positive function $\Psi(X, \nu) \in C^2(\Gamma)$, where Γ is an open neighborhood of the unit normal bundle of Σ in $\bar{M} \times \mathbb{S}^n$. Then there exists some constant $C > 0$ depending only on $n, k, \|\Sigma\|_{C^1}, \inf \Psi, \|\Psi\|_{C^2}$ and the curvature $\bar{R}$ of $\bar{M}$ such that*

$$\max_{X \in \Sigma} \kappa_{\max}(X) \leq C.$$

Several remarks are in order.

Remark 1.2. First, we shall explicitly point out that the $k = n - 1$ part of the theorem in warped product manifolds has already been obtained by Chen-Li-Wang in [10, Section 6]. Their method was based on the ground-breaking paper of Guan-Ren-Wang [25] and also on how Ren-Wang solved the $k = n - 1$ case in $\mathbb{R}^{n+1}$ [47].

It is then plausible to think that, since the $k = n - 2$ case has also been solved by Ren-Wang in $\mathbb{R}^{n+1}$ [48], so based on their proof and by imitating the method of Chen-Li-Wang, one can readily derive the C^2 estimates for the σ_{n-2} equation in warped product manifolds as well. Although this result has not been written and published (as far as we know), this method of proof should have already been known to those authors.

Remark 1.3. Apart from providing a written proof for the $k = n - 2$ case, the main contributions of our theorem 1.1 are two-fold: on one hand, we approach the C^2 estimate by a different method and our proof is simpler than the ones in [10, 47, 48], due to our synthesis of techniques from several other authors. On the other hand, our method (inherited from Lu's paper [39] with minor simplifications) is robust and hence is also applicable in other situations concerning curvature estimates for prescribed curvature equations; see theorem 1.8 and theorem 1.13 below. We believe that this method could be applied to more equations of similar forms.

Remark 1.4. Note that the results of Guan-Ren-Wang [25] and Ren-Wang [47, 48] were all obtained in the Euclidean space $\mathbb{R}^{n+1}$. It is then worth to wonder if the estimates could still hold when the ambient space is e.g. the hyperbolic space $\mathbb{H}^{n+1}$. Due to the nature of $\mathbb{H}^{n+1}$ having negative curvature, obtaining the C^2 estimate in this case is usually more difficult: there would exist an extra negative term $-h_{11}$ which would not exist in the $\mathbb{R}^{n+1}$ case.

As demonstrated in the work of Spruck-Xiao [50], this problematic term can be handled by applying the maximum principle to $\log \kappa_{\max}$, instead of the more complicated terms in (1.2) and (1.3). Our proof follows the exact same spirit: we synthesize the test functions of Spruck-Xiao [50] and Ren-Wang [47, 48] i.e. (1.5) and (1.3), to get a new one

$$(1.4) \quad Q = \log \kappa_{\max} - N \log \langle X, \nu \rangle + \alpha \Phi$$

which utilizes the largest principal curvature and gives more "good" third order terms; and we apply a standard perturbation argument as in [14] to resolve the issue that $\kappa_{\max}$ may have multiplicity more than one; proceeding to follow Lu's derivation in [39, 40], we can remove the large negative term $-\frac{h_{11}^2}{h_{11}^2}$ by invoking Ren-Wang's concavity inequality in [47, 48]; finally, it is the elegant paper

[50] of Spruck and Xiao that made us realize that our proof can be readily carried over to space forms and later to warped product manifolds after ² reading Chen-Li-Wang's paper [10].

Remark 1.5. In [10, Theorem 4.1], Chen-Li-Wang used the same test function of Spruck-Xiao i.e. (1.5), to generalize the $k = 2$ case from space forms to warped product manifolds, but they did not seem to realize the potential use of the new test function (1.4). Because in that paper, Chen-Li-Wang used

$$Q = \begin{cases} \frac{1}{2} \log \sum_l \kappa_l^2 - N \log \langle X, \nu \rangle + \alpha \Phi & \text{for convex hypersurfaces} \\ \log \log \sum_l e^{\kappa_l} - N \log \langle X, \nu \rangle + \alpha \Phi & \text{for } (n-1)\text{-convex hypersurfaces} \end{cases}$$

which is in the same form of Guan-Ren-Wang's test function (1.2). We note that applying the maximum principle to the largest principal curvature is the key to making our proof simpler than theirs ³.

Since the $k = n - 1$ case in warped product manifolds is contained in Chen-Li-Wang's work [10], we shall re-state the theorem for the $k = n - 2$ case alone for emphasis, and again, this result is new in hyperbolic space $\mathbb{H}^{n+1}$.

Theorem 1.6. *Let $n \geq 3$. Suppose that Σ is a closed strictly star-shaped $(n-2)$ -convex hypersurface in a warped product manifold $\overline{M}$ satisfying the curvature equation*

$$\sigma_{n-2}(\kappa(X)) = \Psi(X, \nu) \quad \text{for all } X \in \Sigma$$

for some positive function $\Psi(X, \nu) \in C^2(\Gamma)$, where Γ is an open neighborhood of the unit normal bundle of Σ in $\overline{M} \times \mathbb{S}^n$. Then there exists some constant $C > 0$ depending only on $n, \|M\|_{C^1}, \inf \Psi, \|\Psi\|_{C^2}$ and the curvature $\overline{R}$ of $\overline{M}$ such that

$$\max_{X \in \Sigma} \kappa_{\max}(X) \leq C.$$

Remark 1.7. When $n = 3$, it reduces to the prescribed mean curvature equation which has been solved long ago; when $n = 4$, it becomes the σ_2 equation in dimension four which has been solved by Guan-Ren-Wang [25, Theorem 1.4] in $\mathbb{R}^{n+1}$, by Spruck-Xiao [50, Theorem 2.1] in space forms and by Chen-Li-Wang [10, Theorem 4.1] in warped product manifolds.

Therefore, here we only need to prove theorem 1.6 for $n \geq 5$; this is also the applicable range of Ren-Wang's inequality, see lemma 2.11. Since the lower order estimates and the existence theorem (once C^2 estimates are obtained) are already established in [50, Theorem 3.3] and [10, Theorem 1.1] (under some barrier conditions), we do not list them here again.

Next, we state our second result: by using the test function of Spruck-Xiao in [50]

$$(1.5) \quad Q = \log \kappa_{\max} - \log(\langle X, \nu \rangle - a) + \alpha \Phi,$$

we can prove C^2 estimates for k -convex solutions to the prescribed curvature measure type equation i.e.

$$(1.6) \quad \sigma_k(\kappa(X)) = \langle X, \nu \rangle^p \psi(X) \quad \text{for all } X \in \Sigma.$$

When $p = 1$, this is exactly the equation for the prescribed curvature measure problem whose significance is discussed in Guan-Li-Li and Yang's papers [19, 54], in which C^2 estimates have been obtained in $\mathbb{R}^{n+1}$ and $\mathbb{H}^{n+1}$, respectively. Following Lu, who obtained C^2 -estimates for the wider range $p \in (-\infty, 0) \cup (0, 1]$ in $\mathbb{H}^{n+1}$ [39, Theorem 1.2], we extend their results not only to warped product manifolds and also to the same wider range of p 's.

²This note was initially written for curvature estimates in space forms. Later, Lu pointed out the paper of Chen-Li-Wang and suggested the generalization to warped product manifolds.

³This has also been observed by Chu [14], and this is how he provided a simple proof for Guan-Ren-Wang's result.

Theorem 1.8. *Let $1 \leq k \leq n$ and $p \in (-\infty, 0) \cup (0, 1]$. Suppose that Σ is a closed strictly star-shaped k -convex hypersurface in a warped product manifold $\overline{M}$ satisfying the prescribed curvature measure type equation (1.6) for some positive function $\psi(X) \in C^2(\Sigma)$. Then there exists some constant $C > 0$ depending only on $n, k, p, \|M\|_{C^1}, \inf \psi, \|\psi\|_{C^2}$ and the curvature $\overline{R}$ of $\overline{M}$ such that*

$$\max_{X \in \Sigma} \kappa_{\max}(X) \leq C.$$

Remark 1.9. We shall mention that in [9], Chen proved theorem 1.8 for $k = 2$ in $\mathbb{R}^{n+1}$ based on the study of optimal concavity of the σ_2 operator; and Huang-Xu [26] proved theorem 1.8 for all $2 \leq k \leq n$ in $\mathbb{R}^n$ by generalizing the method of Guan-Li-Li [19].

By extending the lower order estimates in [21, 26, 54] to warped product manifolds, the existence theorem follows.

Theorem 1.10. *If either (i) $1 \leq k \leq n-1, p \in (-\infty, 0) \cup (0, 1]$, or (ii) $k = n, p \in (-\infty, 0) \cup (0, 1)$, then there exists a unique k -convex star-shaped $C^{3,\alpha}$ hypersurface Σ in $\overline{M}$ satisfying (1.6).*

For $k = n$ and $p = 1$, the same existence result would hold with the additional assumption that $\min_M \psi > 1$.

Remark 1.11. In particular, this generalizes the main result [54, Theorem 1.1 and Theorem 5.1] of Yang from $p = 1$ to $p \in (-\infty, 0) \cup (0, 1]$ in hyperbolic space. But still, our proof is highly inspired by that of Yang.

Remark 1.12. The extra condition $\min \psi > 1$ is required only when deriving the C^0 estimate in the case that $k = n$ and $p = 1$. On the other hand, our C^1 estimate for (1.6) holds for all $p \neq 0$ without any barrier conditions; previously, this result was only obtained in $\mathbb{R}^{n+1}$ [26, Lemma 2.3] and hence it is new in e.g. $\mathbb{H}^{n+1}$.

The ultimate goal is to know whether theorem 1.1 holds for $3 \leq k \leq n-3$; see remark 3.5 in [19]. This is a rather difficult task and it still remains open up to this date.

Recall that in their groundbreaking paper, Guan-Ren-Wang proved this for convex solutions and they claimed in a remark [25, Remark 4.7] that their proof would also work for semi-convex and $(k+1)$ -convex solutions, by modifying their test function (1.2). We provide a simpler proof for this claim using our test function (1.4) which will make a concavity inequality of Lu in [39] applicable and we extend this result to warped product manifolds.

Theorem 1.13. *Let $3 \leq k \leq n-3$. Suppose that Σ is a closed strictly star-shaped k -convex hypersurface in a warped product manifold $\overline{M}$ satisfying the curvature equation (1.1) for some positive function $\Psi(X, \nu) \in C^2(\Gamma)$, where Γ is an open neighborhood of the unit normal bundle of Σ in $\overline{M} \times \mathbb{S}^n$.*

If either one of the following conditions

- (a) Σ is semi-convex i.e. there exists some $\eta > 0$ such that for all $X \in M$ and all $1 \leq i \leq n$, we have $\kappa_i(X) \geq -\eta$.
- (b) Σ is $(k+1)$ -convex i.e. $\kappa(X) \in \Gamma_{k+1}$.

holds, then there exists some constant $C > 0$ depending only on $n, k, \eta, \|M\|_{C^1}, \inf \Psi, \|\Psi\|_{C^2}$ and the curvature $\overline{R}$ of $\overline{M}$ such that

$$\max_{X \in \Sigma} \kappa_{\max}(X) \leq C.$$

Remark 1.14. Lu [39] proved the semi-convex case in $\mathbb{H}^{n+1}$ based on the method of Yang [54] and that of Guan-Ren-Wang [25]; our proof closely follows theirs, and in fact, it is simply a modest extension of Lu's proof with minor simplifications.

Note also that Lu's proof would work for the $(k+1)$ -convex case as well and that was how we proved the $(k+1)$ -convex case in our initial draft. Later, Lu pointed out to us that, according to

a lemma of Li-Ren-Wang [31], $(k+1)$ -convexity implies semi-convexity. Therefore, it is sufficient to solve the semi-convex case only. We also include a version of the Li-Ren-Wang lemma in the context of our curvature equation; see lemma 2.13.

Finally, we would like to mention that, in a follow-up note [52], we will derive global C^2 estimates for another class of curvature equations:

$$\frac{\sigma_k(\eta[\Sigma])}{\sigma_l(\eta[\Sigma])} = \Psi(X, \nu), \quad \forall X \in \Sigma$$

where $\eta[\Sigma]$ is the $(0, 2)$ tensor η defined by

$$\eta_{ij} = Hg_{ij} - h_{ij}$$

i.e. the first Newton transformation of h_{ij} with respect to g_{ij} . In that article, we will also obtain purely interior C^2 estimates, the Pogorelov type C^2 estimates, the Liouville theorem and solve the Dirichlet problem on Riemannian manifolds, all for the corresponding Hessian equations:

$$\frac{\sigma_k(\eta[u])}{\sigma_l(\eta[u])} = f(x, u, Du), \quad \forall x \in \Omega \subseteq \mathbb{R}^n$$

where

$$\eta[u] = (\Delta u)I - D^2u.$$

The rest of this note is organized as follows. In section 2, we list auxiliary facts such as geometric formulas about hypersurfaces in warped product manifolds and properties of the σ_k operator. In section 3, we perform a formal computation for a combined test function, which will be quoted in later sections. In section 4, we prove theorem 1.1, theorem 1.6 and theorem 1.13. Finally in section 5, we derive C^0 , C^1 and C^2 estimates for the prescribed curvature measure type equation (1.10), which will imply theorem 1.8 and theorem 1.10.

Acknowledgement. We would like to thank Professor Siyuan Lu for introducing the papers [10, 47, 48] of Ren-Wang and Chen-Li-Wang to us. This work is also inspired by Lu's recent two papers [39, 40]. We would also like to thank him for pointing out a lemma due to Li-Ren-Wang in [31] i.e. our lemma 2.13, which simplifies our proof further.

2. PRELIMINARIES

In this section, we collect facts and formulas about hypersurfaces in warped product manifolds, which can all be found in [10] or [15]; here we are merely listing these information for completeness; also to fix notations.

Let (M, g') be a compact Riemannian manifold and let I be an interval in $\mathbb{R}$. For a smooth positive function $\phi : I \rightarrow \mathbb{R}$, we define a warped product manifold $\overline{M} = I \times_\phi M$ endowed with the metric

$$\overline{g} = ds^2 = dr^2 + \phi(r)^2 g', \quad r \in I$$

which will also be simply denoted by $\langle \cdot, \cdot \rangle$.

Remark 2.1. Note that when $M = \mathbb{S}^n$,

$$\overline{M} = \begin{cases} \mathbb{R}^{n+1}, & \text{if } \phi(r) = r \text{ and } I = [0, \infty) \\ \mathbb{H}^{n+1}, & \text{if } \phi(r) = \sinh(r) \text{ and } I = [0, \infty) \\ \mathbb{S}^{n+1}, & \text{if } \phi(r) = \sin(r) \text{ and } I = [0, \pi/2) \end{cases}$$

The Riemannian connection on $\overline{M}$ will be denoted by $\overline{\nabla}$ and the connection on M will be denoted by ∇' . The same convention applies to the curvature tensors $\overline{R}$ and R' .

Let $\{e_1, \dots, e_{n-1}\}$ be an orthonormal frame in M and let $\{\theta_i\}$ be the associated dual frame. An orthonormal frame in $\overline{M}$ can then be defined as $\bar{e}_i := (1/\phi)e_i$, $1 \leq i \leq n-1$ and $\bar{e}_0 := \partial/\partial r$; the associated dual frame is then $\bar{\theta}_i := \phi\theta_i$, $1 \leq i \leq n-1$ and $\bar{\theta}_0 = dt$.

We may represent a starshaped hypersurface Σ as a radial graph of a differentiable function $r : M \rightarrow I$ over M i.e.

$$\Sigma = \{X(z) = (r(z), z) : z \in M\},$$

whose tangent space is spanned at each point by the vectors

$$X_i = \phi\bar{e}_i + r_i\bar{e}_0,$$

where r_i 's are the components of the differential $dr = r_i\theta^i$.

The unit outward ⁴ normal is given by

$$\nu = \frac{1}{\sqrt{\phi^2 + |\nabla' r|^2}} \left(\phi\bar{e}_0 - \sum_i r^i\bar{e}_i \right);$$

here $|\nabla' r|^2 = r^i r_i$ is the squared norm of $\nabla' r = r^i e_i$.

The induced metric on Σ is given by

$$g_{ij} = \langle X_i, X_j \rangle = \phi^2 \delta_{ij} + r_i r_j \quad \text{with inverse} \quad g^{ij} = \frac{1}{\phi^2} \left(\delta_{ij} - \frac{r^i r^j}{\phi^2 + |\nabla' r|^2} \right)$$

The second fundamental form of Σ is given by

$$h_{ij} = \langle \bar{\nabla}_{X_j} X_i, \nu \rangle = \frac{1}{\sqrt{\phi^2 + |\nabla' r|^2}} (-\phi r_{ij} + 2\phi' r_i r_j + \phi^2 \phi' \delta_{ij}),$$

where r_{ij} are the components of the Hessian $\nabla'^2 r = \nabla' dr$ of r in M .

Lemma 2.2. *Let X_0 be a point of Σ and let $\{E_0 = \nu, E_1, \dots, E_n\}$ be an adapted frame field such that each E_i is a principal direction and the associated dual frame satisfies $\omega_i^k = 0$ at X_0 . Then at X_0 , we have*

$$\nabla_k h_{ij} = \nabla_j h_{ik} + \bar{R}_{0ijk}$$

and

$$h_{ii11} - h_{11ii} = h_{11} h_{ii}^2 - h_{11}^2 h_{ii} + 2(h_{ii} - h_{11})\bar{R}_{i1i1} + h_{11}\bar{R}_{i0i0} - h_{ii}\bar{R}_{1010} + \bar{R}_{i1i0;1} - \bar{R}_{1i10;i}.$$

Remark 2.3. The frame field E_i may be obtained from the adapted frame $\nu, X_1, \dots, X_n$ by the Gram-Schmidt procedure. Since this last frame depends only on r and $\nabla' r$, we may conclude that the components of $\bar{R}$ and $\bar{\nabla}\bar{R}$ calculated in terms of the frame E_i depend only on r and $\nabla' r$.

Proof. See lemma 2.1 in [15]. □

$\overline{M}$ naturally comes with a conformal Killing position vector field $V = \phi(r)\partial_r$ and we denote by $\nu(V)$ the outward unit normal. That is, we will study the prescribed curvature equation in the following form:

$$(2.1) \quad \sigma_k(\kappa[\Sigma]) = \Psi(V, \nu).$$

We also define two auxiliary quantities: the support function $u := \langle V, \nu \rangle$ and

$$\Phi(r) := \int_0^r \phi(\rho) d\rho.$$

⁴In [10], Chen-Li-Wang used the inward normal; the reader shall note the sign difference.

Lemma 2.4.

$$\begin{aligned}\nabla_{E_i}\Phi &= \phi\langle\bar{e}_0, E_i\rangle E_i \\ \nabla_i u &= g^{kl}h_{ik}\nabla_{E_l}\Phi \\ \nabla_{E_i, E_j}^2\Phi &= \phi'g_{ij} - uh_{ij} \\ \nabla_{ij}u &= g^{kl}\left(\nabla_{E_k}h_{ij} - \bar{R}_{0ijk}\right)\nabla_{E_l}\Phi + \phi'h_{ij} - ug^{kl}h_{ik}h_{jl}\end{aligned}$$

Proof. See proposition 2.4 in [20], lemma 4.1 in [15] and lemma 2.3 in [10]. $\square$

Before we end this section, we list a few useful properties of the σ_k operator. The reader is also referred to [49, 53] and [38, Chapter XV] for more.

The k -th elementary symmetric polynomial $\sigma_k : \mathbb{R}^n \rightarrow \mathbb{R}$ is a smooth symmetric function of n variables, defined by

$$\sigma_k(\kappa) = \sum_{i_1 < i_2 < \dots < i_k} \kappa_{i_1} \cdots \kappa_{i_k}$$

and the Gårding cones are defined by

$$\Gamma_k = \{\kappa \in \mathbb{R}^n : \sigma_j(\kappa) > 0 \quad \forall 1 \leq j \leq k\}.$$

Definition 2.5. An embedded hypersurface Σ in $\bar{M}$ is said to be a k -convex or k -admissible solution to the curvature equation

$$\sigma_k(\kappa[\Sigma]) = \Psi(V, \nu)$$

if its principal curvatures $\kappa = (\kappa_1, \dots, \kappa_n)$ belong to the k th Gårding cone Γ_k at every point.

Notation 2.6. Observe that

$$\frac{\partial \sigma_k}{\partial \kappa_i} = \sigma_{k-1}(\kappa_1, \dots, \kappa_{i-1}, 0, \kappa_i, \dots, \kappa_n) = \sigma_{k-1}(\kappa) \Big|_{\kappa_i=0}$$

and we introduce the notation

$$\sigma_k^{ii}, \quad \sigma_{k-1}(\kappa|i), \quad \text{or} \quad \sigma_{k-1;i}(\kappa)$$

to mean the same thing.

Similarly, for the second order derivatives, we use the following notations

$$\sigma_k^{ii,jj} = \sigma_{k-2}(\kappa|ij) = \sigma_{k-2;ij}(\kappa) = \frac{\partial^2 \sigma_k}{\partial \kappa_i \partial \kappa_j}.$$

Lemma 2.7. For $\kappa \in \mathbb{R}^n$, we have the following

$$\sigma_k(\kappa) = \kappa_i \sigma_{k-1}(\kappa|i) + \sigma_k(\kappa|i), \quad \sum_{i=1}^n \kappa_i \sigma_{k-1}(\kappa|i) = k \sigma_k, \quad \sum_{i=1}^n \sigma_{k-1}(\kappa|i) = (n-k+1) \sigma_{k-1}$$

Lemma 2.8. For $\kappa \in \Gamma_k$ and $1 \leq l < k$, we have

$$\sigma_l(\kappa) > \kappa_1 \cdots \kappa_l$$

and

$$\sigma_k(\kappa) \leq C \kappa_1 \cdots \kappa_k.$$

Proof. For the first inequality, see lemma 12 in [48]; an explicit proof is given in the preprint version of that paper. For the second inequality, see lemma A.1 in [32]. $\square$

Lemma 2.9. For $\kappa = (\kappa_1, \dots, \kappa_n) \in \Gamma_k$, if $\kappa_i \leq 0$, then

$$-\kappa_i < \frac{n-k}{k} \kappa_1.$$

Proof. See lemma 11 in [48]. $\square$

Lemma 2.10. *Let $1 \leq k \leq n$. If $\kappa = (\kappa_1, \dots, \kappa_n) \in \Gamma_k$ is ordered as $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_n$, then we have*

$$\sum_{i=1}^n \kappa_i^2 \sigma_k^{ii} \geq \frac{k}{n} \kappa_1 \sigma_k.$$

Proof. When $k = 1$, the inequality follows by noting that $\sigma_1^{ii} = 1$ and $\kappa_1 \geq \frac{1}{n} \sigma_1$. When $2 \leq k \leq n$, by lemma 2.3 in [40], we have $\sum \kappa_i^2 \sigma_k^{ii} \geq \frac{k}{n} \sigma_1 \sigma_k$. Noting that $\sigma_1 - \kappa_1 = \frac{\partial \sigma_2}{\partial \kappa_i} > 0$ for $\kappa \in \Gamma_k \subseteq \Gamma_2$ yields the inequality. $\square$

Finally, we have the concavity inequalities due to Ren-Wang and Lu.

Lemma 2.11. *Let $k = n - 1, n \geq 3$ or $k = n - 2, n \geq 5$, and let $\xi \in \mathbb{R}^n$ be an arbitrary vector. Suppose that $\kappa \in \Gamma_k$ is ordered as*

$$\kappa_1 > \kappa_2 \geq \kappa_3 \geq \dots \geq \kappa_n.$$

Suppose also that $N_0 \leq \sigma_k(\kappa) \leq N_1$. If κ_1 is sufficiently large, then there exists some $\beta > 0$ such that

$$\kappa_1 \left[\beta \left(\sum_{i=1}^n \sigma_k^{ii} \xi_i \right)^2 - \sum_{p \neq q} \sigma_k^{pp,qq} \xi_p \xi_q \right] - \sigma_k^{11} \xi_1^2 + \sum_{i \neq 1} \frac{2\kappa_1}{\kappa_1 - \kappa_i} \sigma_k^{ii} \xi_i^2 \geq 0.$$

Proof. We first prove for $k = n - 2, n \geq 5$. By theorem 4 in [48], we have that

$$\kappa_1 \left[\beta \left(\sum_{i=1}^n \sigma_k^{ii} \xi_i \right)^2 - \sum_{p \neq q} \sigma_k^{pp,qq} \xi_p \xi_q \right] - \sigma_k^{11} \xi_1^2 + \sum_{i \neq 1} a_i \xi_i^2 \geq 0$$

where

$$a_i = \sigma_k^{ii} + (\kappa_1 + \kappa_i) \sigma_k^{11,ii}.$$

Recall that

$$\sigma_k^{11,ii} = \frac{\sigma_k^{ii} - \sigma_k^{11}}{\kappa_1 - \kappa_i}$$

and so we have

$$\begin{aligned} a_i &= \sigma_k^{ii} + (\kappa_1 + \kappa_i) \sigma_k^{11,ii} = \sigma_k^{ii} + \frac{\kappa_1 + \kappa_i}{\kappa_1 - \kappa_i} (\sigma_k^{ii} - \sigma_k^{11}) \\ &\leq \sigma_k^{ii} + \frac{\kappa_1 + \kappa_i}{\kappa_1 - \kappa_i} \sigma_k^{ii} \\ &= \frac{2\kappa_1}{\kappa_1 - \kappa_i} \sigma_k^{ii}. \end{aligned}$$

The result then follows. Note that we have used lemma 2.9 with $2k > n$ to get

$$\frac{\kappa_1 + \kappa_i}{\kappa_1 - \kappa_i} \geq 0.$$

For $k = n - 1$ with $n \geq 3$, we apply theorem 11 in [47] to have that

$$\kappa_1 \left[\beta \left(\sum_{i=1}^n \sigma_{n-1}^{ii} \xi_i \right)^2 - \sum_{p \neq q} \sigma_{n-1}^{pp,qq} \xi_p \xi_q \right] - \sigma_{n-1}^{11} \xi_1^2 + (1 + \varepsilon) \sum_{i \neq 1} \sigma_{n-1}^{ii} \xi_i^2 \geq 0$$

where $\varepsilon > 0$ is arbitrary and β depends on ε .

By lemma 2.9 again, we can choose ε small enough e.g. $\varepsilon = [(2n - 2)/n] - 1$ so that

$$\frac{2\kappa_1}{\kappa_1 - \kappa_i} \geq \frac{2n - 2}{n} = 1 + \varepsilon, \quad n \geq 3 \quad \text{and} \quad i \neq 1.$$

The result then follows. $\square$

Lemma 2.12. *Suppose $\kappa = (\kappa_1, \dots, \kappa_n) \in \Gamma_k$ is ordered as $\kappa_1 \geq \dots \geq \kappa_n$. Let $\xi \in \mathbb{R}^n$ be an arbitrary vector and $1 \leq l < k$. Given $\varepsilon, \delta, \delta_0 \in (0, 1)$, we can find some $\delta' > 0$ such that if $\kappa_l \geq \delta\kappa_1$ and $\kappa_{l+1} \leq \delta'\kappa_1$, then*

$$(2.2) \quad -\sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} \geq (1 - \varepsilon) \frac{\xi_1^2}{\kappa_1^2} - \delta_0 \sum_{i > l} \frac{\sigma_k^{ii} \xi_i^2}{\kappa_1 \sigma_k}.$$

Proof. See lemma 3.1 in [39]. □

Finally, we demonstrate the fact that $(k+1)$ -convexity implies semi-convexity, which is a lemma proved by Li-Ren-Wang in [31], for solutions of the Hessian equation $\sigma_k(D^2u) = f(x, u, Du)$. Here we prove the corresponding version for our curvature equation (1.1).

Lemma 2.13. *Suppose that Σ is a $(k+1)$ -convex solution to equation (1.1). Then there exists some constant $\eta > 0$ depending on known data of Σ and Ψ such that*

$$\kappa_i(X) \geq -\eta \quad \text{for all } 1 \leq i \leq n \text{ and all } X \in \Sigma.$$

Proof. The proof is adapted from [31, Lemma 7]. We choose η such that

$$\left(\frac{\eta}{n}\right)^k \geq \sup_{X \in \Sigma} \Psi(X, \nu).$$

Let $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_n$ denote the principal curvatures of Σ . Since Σ is $(k+1)$ -convex, we have that $\kappa = (\kappa_1, \dots, \kappa_n) \in \Gamma_{k+1}$. Hence by using lemma 2.8, we have that

$$\sigma_k(\kappa) \geq \kappa_1 \cdots \kappa_k \geq \kappa_k^k.$$

It follows that

$$\kappa_k \leq \frac{\eta}{n}.$$

Also, using $\kappa \in \Gamma_k$, we have that the sum of any $(n - k + 1)$ components of κ is positive. In particular, we have

$$\sum_{i=k}^n \kappa_i > 0$$

which implies that

$$0 < \kappa_k + \kappa_{k+1} + \dots + \kappa_n \leq (n - k) \kappa_k + \kappa_n \leq \frac{n - k}{n} \eta + \kappa_n$$

i.e.

$$\kappa_n + \eta \geq 0.$$

□

Remark 2.14. The proof also shows the trivial fact that if $k = n$ then $\kappa_n > 0$ i.e. $\kappa \in \Gamma_n$ and Σ is strictly convex.

3. A MODEL COMPUTATION

In this section, we establish preparatory work for obtaining the following curvature estimate, which will imply theorem 1.1, theorem 1.6, theorem 1.8, and theorem 1.13.

Theorem 3.1. *Let $\Sigma = \{(r(z), z) : z \in M\}$ be a closed k -convex hypersurface in $\overline{M}$ which is strictly star-shaped with respect to the origin and satisfies equation (2.1) for some positive function $\Psi(V, \nu) \in C^2(\Gamma)$, where Γ is an open neighborhood of the unit normal bundle of Σ in $\overline{M} \times \mathbb{S}^n$.*

Suppose that we have uniform control $0 < r_1 < r(z) < r_2 < b$ and $|r|_{C^1} \leq r_3$. If additionally either of the following holds:

- $k = n - 1, n \geq 3$ or $k = n - 2, n \geq 5$,

- Σ is semi-convex i.e. there exists some $\eta > 0$ such that for all $X \in \Sigma$ and all $1 \leq i \leq n$, we have $\kappa_i(X) \geq -\eta$.
- Σ is $(k+1)$ -convex i.e. $\kappa \in \Gamma_{k+1}$, or
- $\Phi(V, \nu) = \langle V, \nu \rangle^p \psi(V)$ for $p \in (-\infty, 0) \cup (0, 1]$ and $\psi \in C^2(\Sigma)$,

then there exists some $C > 0$ depending on $n, k, p, r_1, r_2, r_3, \inf \Psi, \|\Psi\|_{C^2}$ and $\bar{R}$ such that

$$\max_{z \in M} |\kappa_{\max}(z)| \leq C.$$

The proof for different cases will be different. More specifically, for the first three cases, we will use the test function

$$Q = \log \kappa_{\max} - N \log u + \alpha \Phi$$

with $N, \alpha > 0$ being some large numbers, in order to make the concavity inequalities in lemma 2.11 and lemma 2.12 applicable.

In contrast, for the fourth case, we will be working with the test function

$$Q = \log \kappa_{\max} - \log(u - a) + \alpha \Phi$$

without the large number N and where $a > 0$ satisfies $u \geq 2a$. This is because N would have produced a negative term with the coefficient $-N$ which are difficult to handle without a handy concavity inequality, if we would like N to be large. In this case, we have to make delicate use of the special structure of the right-hand side $\Psi(X, \nu) = u^p \psi(X)$ with the help of the gauge term a .

However, a large part of the derivation will be almost exactly the same in all the four cases. Therefore, in this section, we will do a "model computation" for the combined test function

$$Q = \log \kappa_{\max} - N \log(u - a) + \alpha \Phi.$$

The purpose for doing so is to reduce the labor and shorten the proof. More precisely, we will substitute $a = 0$ in section 4 and $N = 1$ in section 5, then we can proceed from there and thus redundant calculations will be eliminated.

Now we begin the calculations. Define

$$Q(z, \xi) = \log \mathbb{I}(\xi, \xi) - N \log(u - a) + \alpha \Phi$$

where ξ is a unit tangent vector to Σ at $X = (r(z), z)$ and $\mathbb{I}$ is the second fundamental form. Suppose that Q attains its maximum at $X_0 = (r(z_0), z_0)$. We may choose a local orthonormal frame $\{E_1, \dots, E_n\}$ such that at this point we have $\xi = E_1$ and $h_{ij} = \kappa_i \delta_{ij}$, where the principal curvatures are ordered as $\kappa_1 \geq \kappa_2 \geq \dots \geq \kappa_n$.

Remark 3.2. Technically speaking, our operation here is not really valid; because the point X_0 and hence the local orthonormal frame $\{e_i\}$ depend on the choice of the test function Q .

Here we are merely doing a symbolic computation formally! This is for not to repeat this part of calculation in section 4 and section 5.

Now, note that if κ_1 has multiplicity more than one i.e.

$$\kappa_1 = \kappa_2 = \dots = \kappa_m > \kappa_{m+1} \geq \dots \geq \kappa_n \quad \text{for some } m > 1,$$

then Q is not smooth at X_0 . To resolve this issue, we apply a standard perturbation argument following Chu [14]. Let g be the first fundamental form of Σ . Near X_0 , we define a new tensor B by

$$B(V_1, V_2) = g(V_1, V_2) - g(V_1, E_1)g(V_2, E_1)$$

for tangent vectors V_1 and V_2 . Denote by $B_{ij} = B(E_i, E_j)$ and it is clear that $B_{ij} = \delta_{ij}(1 - \delta_{1i})$. We now define a new matrix by $\tilde{h}_{ij} := h_{ij} + B_{ij}$ whose eigenvalues are ordered as

$$\tilde{\kappa}_1 \geq \tilde{\kappa}_2 \geq \dots \geq \tilde{\kappa}_n.$$

Note that $\kappa_1 \geq \tilde{\kappa}_1$ near X_0 and

$$\tilde{\kappa}_i = \begin{cases} \kappa_1, & i = 1 \\ \kappa_i + 1, & i > 1 \end{cases}$$

at X_0 . It then follows that $\tilde{\kappa}_1$ has multiplicity one and hence that the perturbed test function

$$\tilde{Q} := \log \tilde{\kappa}_1 - N \log(u - a) + \alpha \Phi$$

is smooth at X_0 .

Remark 3.3. In [14], Chu defined $\tilde{\kappa}_i = \kappa_i - 1$ for $i > 1$. Here we change the constant to be +1 in order to have that ⁵ $\tilde{\kappa} = \kappa + (0, 1, 1, \dots, 1) \in \Gamma_k$. This change is crucial as to satisfy the hypotheses of lemma 2.11 and lemma 2.12.

We will proceed to perform a lengthy and tedious computation at the point X_0 ; the reader may start from (3.13).

Notation 3.4. In our derivation below, the symbol C or $C_{\dots}$ with some index will denote some controlled quantity whose magnitude depends on some known data. Its value may change from line to line without explicitly saying so, but it will still be denoted by the same symbol. This would not cause confusion because its magnitude is not of relevance in our analysis.

Also, whenever we say the vague phrase " κ_1 is sufficiently large", it would mean that given any known constant $C > 0$, we must have $\kappa_1 > C$ otherwise the desired curvature bound $\kappa_1 \leq C$ would have been obtained already.

Notation 3.5. We may view σ_k as an operator taking symmetric matrices as its argument; in our case the matrix is h_{ij} and we may write $F(h_{ij})$ to denote the operator.

When we view it as a function taking the eigenvalues of matrices i.e. an n -dimensional vector as its argument, in our case the vector is the principal curvatures $\kappa = (\kappa_1, \dots, \kappa_n)$ and we write $f(\kappa)$ to denote the function.

We also denote

$$F^{ij}(h) = \frac{\partial F}{\partial h_{ij}}, \quad F^{ij,kl} = \frac{\partial^2 F}{\partial h_{ij} \partial h_{kl}}$$

and we have

$$F^{ij}(h) = f_i(\kappa) \delta_{ij}$$

when h is diagonalized and where f_i is the i -th partial derivative of f .

Using $\tilde{h}_{ij} = h_{ij} + B_{ij}$, we have at X_0 that

$$\begin{aligned} (\tilde{\kappa}_1)_i &= \tilde{h}_{11i} = h_{11i} \\ (\tilde{\kappa}_1)_{ii} &= \tilde{h}_{11ii} + 2 \sum_{p \neq 1} \frac{\tilde{h}_{1pi}^2}{\tilde{\kappa}_1 - \tilde{\kappa}_p} = h_{11ii} + 2 \sum_{p \neq 1} \frac{h_{1pi}^2}{\kappa_1 - \tilde{\kappa}_p} \end{aligned}$$

and so

$$(3.1) \quad 0 = \frac{(\tilde{\kappa}_1)_i}{\tilde{\kappa}_1} - N \frac{u_i}{u - a} + \alpha \Phi_i = \frac{h_{11i}}{\kappa_1} - N \frac{u_i}{u - a} + \alpha \Phi_i$$

$$\begin{aligned} (3.2) \quad 0 &\geq \frac{(\tilde{\kappa}_1)_{ii}}{\tilde{\kappa}_1} - \frac{(\tilde{\kappa}_1)_i^2}{\tilde{\kappa}_1^2} - N \frac{u_{ii}}{u - a} + N \frac{u_i^2}{(u - a)^2} + \alpha \Phi_{ii} \\ &= \frac{h_{11ii}}{\kappa_1} + 2 \sum_{p \neq 1} \frac{h_{1pi}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_p)} - \frac{h_{11i}^2}{\kappa_1^2} - N \frac{u_{ii}}{u - a} + N \frac{u_i^2}{(u - a)^2} + \alpha \Phi_{ii} \end{aligned}$$

⁵Because Γ_k is an open symmetric convex cone.

Contracting (3.2) with $F = \sigma_k$, we obtain that

$$(3.3) \quad \begin{aligned} 0 \geq & \sum_i \frac{F^{ii} h_{11ii}}{\kappa_1} + 2 \sum_i \sum_{p \neq 1} \frac{F^{ii} h_{1pi}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_p)} - \sum_i \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\ & - N \sum_i \frac{F^{ii} u_{ii}}{u - a} + N \sum_i \frac{F^{ii} u_i^2}{(u - a)^2} + \alpha \sum_i F^{ii} \Phi_{ii}. \end{aligned}$$

By the commutator formula (see lemma 2.2),

$$\nabla_{11} h_{ii} = \nabla_{ii} h_{11} + h_{11} h_{ii}^2 - h_{11}^2 h_{ii} + 2(h_{ii} - h_{11}) \bar{R}_{i1i1} + h_{11} \bar{R}_{i0i0} - h_{ii} \bar{R}_{1010} + \bar{R}_{i1i0;1} - \bar{R}_{1i10;i}.$$

from which it follows that for some $C > 0$ depending on $k, \bar{R}$ and Ψ

$$(3.4) \quad F^{ii} h_{11ii} \geq F^{ii} h_{ii11} - \kappa_1 \sum_i F^{ii} \kappa_i^2 + \kappa_1^2 k \Psi - C \kappa_1 \sum_i F^{ii} - C.$$

where we have used lemma 2.7 to get $\sum F^{ii} \kappa_i = kF = k\Psi$.

Also, for the second term in (3.3), we have

$$(3.5) \quad \begin{aligned} 2 \sum_i \sum_{p \neq 1} \frac{F^{ii} h_{1pi}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_p)} & \geq 2 \sum_{p \neq 1} \frac{F^{pp} h_{1pp}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_p)} + 2 \sum_{p \neq 1} \frac{F^{11} h_{1p1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_p)} \\ & = 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)}. \end{aligned}$$

Now, by inserting (3.4) and (3.5) into (3.3), we have

$$(3.6) \quad \begin{aligned} 0 \geq & \sum_i \frac{F^{ii} h_{ii11}}{\kappa_1} - \sum_i \kappa_i^2 F^{ii} + \kappa_1 k \Psi - C \sum_i F^{ii} - C \\ & + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_i \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\ & - N \sum_i \frac{F^{ii} u_{ii}}{u - a} + N \sum_i \frac{F^{ii} u_i^2}{(u - a)^2} + \alpha \sum_i F^{ii} \Phi_{ii} \end{aligned}$$

Now let us evaluate the third line. By lemma 2.4, we have

$$\begin{aligned} u_{ii} &= \sum_k (h_{iik} - \bar{R}_{0iik}) \Phi_k + \phi' \kappa_i - \kappa_i^2 u \\ \Phi_{ii} &= \phi' - u \kappa_i \end{aligned}$$

and hence (3.6) becomes

$$\begin{aligned}
(3.7) \quad 0 &\geq \sum_i \frac{F^{ii} h_{ii11}}{\kappa_1} - \sum_i F^{ii} \kappa_i^2 + \kappa_1 k \Psi - C \sum F^{ii} - C + N \sum_i \frac{F^{ii} u_i^2}{(u-a)^2} \\
&\quad + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_i \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\
&\quad - \frac{N}{u-a} \sum_i F^{ii} \left[\sum_k (h_{iik} - \bar{R}_{0iik}) \Phi_k + \phi' \kappa_i - \kappa_i^2 u \right] + \alpha \sum_i F^{ii} (\phi' - u \kappa_i) \\
&= \sum_i \frac{F^{ii} h_{ii11}}{\kappa_1} - \sum_i \kappa_i^2 f_i + \kappa_1 k \Psi - C \sum F^{ii} - C + N \sum_i \frac{F^{ii} u_i^2}{(u-a)^2} \\
&\quad + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_i \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + \frac{N}{u-a} \sum F^{ii} \bar{R}_{0iik} \Phi_k \\
&\quad - \frac{N}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right) - \frac{N k \phi'}{u-a} \Psi + \frac{N u}{u-a} \sum_i F^{ii} \kappa_i^2 + \alpha \phi' \sum_i F^{ii} - \alpha k u \Psi \\
&\geq \left(\frac{N u}{u-a} - 1 \right) \sum_i F^{ii} \kappa_i^2 + (\alpha \phi' - C) \sum_i F^{ii} + \kappa_1 k \Psi - C \alpha - C \\
&\quad + \sum_i \frac{F^{ii} h_{ii11}}{\kappa_1} - \frac{N}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right) + N \sum_i \frac{F^{ii} u_i^2}{(u-a)^2} \\
&\quad + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_i \frac{F^{ii} h_{11i}^2}{\kappa_1^2}
\end{aligned}$$

where we have used lemma 2.7 for $\sum_i \kappa_i f_i$.

Next, we proceed to handle the h_{ii11} term. By differentiating (2.1) twice, we obtain that

$$(3.8) \quad F^{ii} h_{iik} = \Psi_V (\nabla_{E_k} V) + h_{ks} \Psi_\nu (E_s)$$

$$(3.9) \quad \text{and } F^{ii} h_{ii11} + F^{ij,kl} h_{ij1} h_{kl1} = \nabla_{11} \Psi$$

Thus, we have

$$\begin{aligned}
(3.10) \quad 0 &\geq \left(\frac{N u}{u-a} - 1 \right) \sum_i F^{ii} \kappa_i^2 + (\alpha \phi' - C) \sum_i f_i + k F \kappa_1 - C \alpha - C \\
&\quad - \sum_{i,j,k,l} \frac{F^{ij,kl} h_{ij1} h_{kl1}}{\kappa_1} + \frac{\Psi_{11}}{\kappa_1} - \frac{N}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right) \\
&\quad + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_i \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + N \sum_i \frac{F^{ii} u_i^2}{(u-a)^2}.
\end{aligned}$$

Next, by an inequality due to Andrews [3] and Gerhard [17], we have

$$\begin{aligned}
- \sum_{i,j,k,l} \frac{F^{ij,kl} h_{ij1} h_{kl1}}{\kappa_1} &= - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + \sum_{p \neq q} \frac{F^{pp,qq} h_{pq1}^2}{\kappa_1} \\
&\geq - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{i > m} \frac{1}{\kappa_1} \frac{F^{ii} - F^{11}}{\kappa_1 - \tilde{\kappa}_i} h_{11i}^2
\end{aligned}$$

and so (3.10) becomes

$$\begin{aligned}
 (3.11) \quad 0 \geq & \left(\frac{Nu}{u-a} - 1 \right) \sum_i F^{ii} \kappa_i^2 + (\alpha\phi' - C) \sum_i F^{ii} + kF\kappa_1 - C\alpha - C \\
 & - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{i > m} \frac{1}{\kappa_1} \frac{F^{ii} - F^{11}}{\kappa_1 - \tilde{\kappa}_i} h_{11i}^2 \\
 & + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_i \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + N \sum_i \frac{F^{ii} u_i^2}{(u-a)^2} \\
 & + \frac{\Psi_{11}}{\kappa_1} - \frac{N}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right)
 \end{aligned}$$

Regrouping terms, we get

$$\begin{aligned}
 (3.12) \quad 0 \geq & \left(\frac{Nu}{u-a} - 1 \right) \sum_i F^{ii} \kappa_i^2 + (\alpha\phi' - C) \sum_i F^{ii} + kF\kappa_1 - C\alpha - C \\
 & - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + N \frac{F^{11} u_1^2}{(u-a)^2} \\
 & + 2 \sum_{i > m} \frac{1}{\kappa_1} \frac{F^{ii} - F^{11}}{\kappa_1 - \tilde{\kappa}_i} h_{11i}^2 + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_{i \neq 1} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + N \sum_{i \neq 1} \frac{F^{ii} u_i^2}{(u-a)^2} \\
 & + \frac{\Psi_{11}}{\kappa_1} - \frac{N}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right)
 \end{aligned}$$

For the third line, we have

$$\begin{aligned}
 & 2 \sum_{i > m} \frac{1}{\kappa_1} \frac{F^{ii} - F^{11}}{\kappa_1 - \tilde{\kappa}_i} h_{11i}^2 + 2 \sum_{i \neq 1} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_{i \neq 1} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\
 = & 2 \sum_{i > m} \frac{F^{ii} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + 2 \sum_{1 < i \leq m} \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \sum_{i \neq 1} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \\
 \geq & \sum_{i \in I} \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \quad I := \{i > m : \kappa_1 + \tilde{\kappa}_i < 0\}
 \end{aligned}$$

because for $1 < i \leq m$,

$$2 \frac{F^{11} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \frac{F^{ii} h_{11i}^2}{\kappa_1^2} = (2\kappa_1 - 1) \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \geq 0;$$

and for $i > m$,

$$2 \frac{F^{ii} h_{11i}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \frac{F^{ii} h_{11i}^2}{\kappa_1^2} = \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \geq 0 \quad \text{if } \kappa_1 + \tilde{\kappa}_i \geq 0.$$

Therefore, we arrive at

$$\begin{aligned}
(3.13) \quad 0 \geq & \left(\frac{Nu}{u-a} - 1 \right) \sum_i F^{ii} \kappa_i^2 + (\alpha\phi' - C) \sum_i F^{ii} + kF\kappa_1 - C\alpha - C \\
& - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + N \frac{F^{11} u_1^2}{(u-a)^2} \\
& + \sum_{i \in I} \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + N \sum_{i \in I} \frac{F^{ii} u_i^2}{(u-a)^2} + \frac{\Psi_{11}}{\kappa_1} - \frac{N}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right).
\end{aligned}$$

It is from here we shall continue the derivation based upon different conditions listed in theorem 3.1.

4. THE PRESCRIBED CURVATURE EQUATION IN GENERAL

In this section, we prove theorem 3.1 for the first three cases i.e.

$$(4.1) \quad \text{Case 1: } k = n-1, n \geq 3 \quad \text{or} \quad k = n-2, n \geq 5$$

$$(4.2) \quad \text{Case 2: } M \text{ is semi-convex}$$

$$(4.3) \quad \text{Case 3: } M \text{ is } (k+1)\text{-convex}$$

For these cases, we shall employ the test function

$$Q = \log \kappa_{\max} - N \log u + \alpha \Phi.$$

By proceeding exactly as in section 3, or simply by substituting $a = 0$ in (3.13) we obtain that

$$\begin{aligned}
(4.4) \quad 0 \geq & (N-1) \sum_i F^{ii} \kappa_i^2 + (\alpha\phi' - C) \sum_i F^{ii} + kF\kappa_1 - C\alpha - C \\
& - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + N F^{11} \frac{u_1^2}{u^2} \\
& + \sum_{i \in I} \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + N \sum_{i \in I} \frac{F^{ii} u_i^2}{u^2} + \frac{\Psi_{11}}{\kappa_1} - \frac{N}{u} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right)
\end{aligned}$$

Let us deal with the third line first. For the first term, we claim that

Claim 4.1. Under either conditions (4.1) or (4.2), we have

$$\sum_{i \in I} \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} \geq 0.$$

Proof of the Claim. It suffices to prove the numerator of the quotient

$$\frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i}$$

is positive when $\kappa_i < 0$.

If k and n satisfy condition (4.1), then by lemma 2.9 we have that

$$\kappa_1 + \tilde{\kappa}_i = \kappa_1 + \kappa_i + 1 > \kappa_1 - \frac{n-k}{k} \kappa_1 + 1 = \frac{2k-n}{k} \kappa_1 + 1 > 0.$$

If M is semi-convex i.e. $\kappa_i \geq -\eta$, then

$$\kappa_1 + \tilde{\kappa}_i \geq \kappa_1 - \eta + 1 \geq 0$$

by assuming that κ_1 is sufficiently large. □

With this claim and by throwing away excessive positive terms, we have that

$$\begin{aligned}
 0 &\geq (N-1) \sum_i F^{ii} \kappa_i^2 + (\alpha\phi' - C) \sum_i F^{ii} - C\alpha - C \\
 &\quad - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} \\
 &\quad + \frac{\Psi_{11}}{\kappa_1} - \frac{N}{u} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right)
 \end{aligned} \tag{4.5}$$

Next, we estimate the remaining two terms in the third line.

Claim 4.2.

$$\frac{\Psi_{11}}{\kappa_1} - \frac{N}{u} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right) \geq -C\kappa_1 - C\alpha - CN - C$$

for some $C > 0$ depending on $\bar{R}, \|\Psi\|_{C^2}, \|\Phi\|_{C^1}$ and $\|\Sigma\|_{C^1}$.

Proof of Claim 4.2. The term Φ_{11} can have the following rough estimate:

$$\begin{aligned}
 \Phi_{11} &= d_V^2 \Psi(\nabla_{E_1} V, \nabla_{E_1} V) + d_V \Psi(\nabla_{E_1, E_1}^2 V) \\
 &\quad + 2d_V d_\nu \Psi(\nabla_{E_1} V, \nabla_{E_1} V) + d_\nu^2 \Psi(\nabla_{E_1} \nu, \nabla_{E_1} \nu) \\
 &\quad + d_\nu \Psi(\nabla_{E_1, E_1}^2 \nu) \\
 &\geq \sum_k h_{1k1} (d_\nu \Psi)(E_k) - Ch_{11}^2 - Ch_{11} - C \\
 &= \sum_k (h_{11k} + \bar{R}_{01k1}) (d_\nu \Psi)(E_k) - Ch_{11}^2 - Ch_{11} - C \\
 &\geq \sum_k h_{11k} (d_\nu \Psi)(E_k) - C\kappa_1^2 - C\kappa_1 - C
 \end{aligned}$$

for some $C > 0$ depending on $\bar{R}$ and $\|\Psi\|_{C^2}$. Note that

$$\begin{aligned}
 &\sum_k \frac{h_{11k}}{\kappa_1} (d_\nu \Psi)(E_k) - \frac{N}{u} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right) \\
 &= \sum_k \frac{h_{11k}}{\kappa_1} (d_\nu \Psi)(E_k) - N \sum_k F_k \frac{\Phi_k}{u} \\
 &= \sum_k \left(N \frac{u_k}{u} - \alpha \Phi_k \right) (d_\nu \Psi)(E_k) - N \sum_k [h_{kk} (d_\nu \Psi)(E_k) + (d_V \Psi)(\nabla_{E_k} V)] \frac{\Phi_k}{u} \\
 &= N \sum_k \frac{h_{kk} \Phi_k}{u} (d_\nu \Psi)(E_k) - N \sum_k h_{kk} (d_\nu \Psi)(E_k) \frac{\Phi_k}{u} \\
 &\quad - \alpha \sum_k d_\nu(\Psi)(E_k) - N \sum_k (d_V \Psi)(\nabla_{E_k} V) \frac{\Phi_k}{u} \\
 &\geq -C\alpha - CN,
 \end{aligned}$$

where the second equality is due to (3.1) and (3.8), and the third equality is due to lemma 2.4.

The result follows. $\square$

With this claim, it follows that

$$(4.6) \quad \begin{aligned} 0 \geq & (N-1) \sum_i \kappa_i^2 f_i + (\alpha\phi' - C) \sum_i f_i - C(\kappa_1 + \alpha + N + 1) \\ & - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} \end{aligned}$$

Now we can proceed to eliminate the second line by applying the concavity inequalities due to Ren-Wang and Lu i.e. lemma 2.11 and lemma 2.12.

4.1. **When $k = n - 1, n \geq 3$ or $k = n - 2, n \geq 5$.**

In this case, by applying lemma 2.11 to $\xi_i = h_{ii1}$ and divide by κ_1^2 , it follows that

$$- \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} \geq -\frac{\beta F_1^2}{\kappa_1}$$

for some $\beta > 0$ depending on $\inf \Psi$ and $\|\Psi\|_{C^0}$. Since

$$F_1^2 = [h_{11}(d_\nu \Psi)(E_1) + (d_V \Psi)(\nabla_{E_1} V)]^2,$$

we have that

$$- \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} \geq -C\beta\kappa_1$$

for some $C > 0$ depending on $\|\Psi\|_{C^1}$, $\|\Phi\|_{C^1}$ and $\|\Sigma\|_{C^1}$. Thus, (4.6) becomes:

$$0 \geq (N-1) \sum_i \kappa_i^2 f_i + (\alpha\phi' - C_1) \sum_i f_i - C(\kappa_1 + \alpha + \beta\kappa_1 + N + 1).$$

Since on Σ , ϕ' is bounded from below by a positive controlled constant and so by choosing α large enough, we may assume that the coefficient $\alpha\phi' - C_1 \geq 0$ and hence the second term is non-negative.

Finally, by lemma 2.10, the inequality reduces to

$$0 \geq (N-1) \frac{k}{n} \kappa_1 \sigma_k - C\kappa_1 - C.$$

Choosing N large enough yields the estimate.

4.2. **When Σ is semi-convex or $(k+1)$ -convex.**

For cases 2 and 3, this time we do not have a concavity inequality as handy as that of Ren-Wang to handle the second line of (4.6). However, we could overcome this difficulty by employing Lu's concavity inequality (2.2) which is weaker but more generic, along with an iteration argument which is inspired by Guan-Ren-Wang [25] and Yang [54].

Remark 4.3. Recall that according to lemma 2.13, it is sufficient to prove the semi-convex case only.

Let $\varepsilon > 0$ be some small number which will be chosen later and let $\delta_0 = \frac{1}{2}$. Pick an arbitrary $\delta_1 \in (0, 1)$, say $\delta_1 = \frac{1}{3}$, then trivially we have $\kappa_1 \geq \delta_1 \kappa_1$. Now by lemma 2.12, there exists some $\delta_2 > 0$ and for this δ_2 , if $\kappa_2 \leq \delta_2 \kappa_1$, then Lu's inequality (2.2) holds for $l = 1$.

We will soon deal with the situation in which Lu's inequality holds; here the key argument is that if the reverse happens i.e. $\kappa_2 > \delta_2 \kappa_1$, then we continue to pick some $\delta_3 > 0$ by lemma 2.12 and for this $\delta_3 > 0$, we do the same analysis as above: either the process stops here and we have Lu's inequality to be analyzed, or we have $\kappa_3 > \delta_3 \kappa_1$ and the process goes on.

Suppose that the process goes on until $\kappa_k > \delta_k \kappa_1$ i.e. we have $\kappa_i > \delta_i \kappa_1$ for $1 \leq i \leq k$. Then we immediately have the following by assuming κ_1 is sufficiently large:

$$\begin{aligned} \sigma_k(\kappa) &\geq \kappa_1 \cdots \kappa_k - C_{n,k} \cdot \kappa_1 \cdots \kappa_{k-1} \cdot \eta \\ &= \kappa_1 \cdots \kappa_{k-1} (\kappa_k - C\eta) \\ &\geq \kappa_1 \cdot (\delta_2 \kappa_1) \cdots (\delta_{k-1} \kappa_1) \cdot (\delta_k \kappa_1 - C\eta) \\ &\geq C\delta_2 \cdots \delta_k \kappa_1^k \quad \text{if } \Sigma \text{ is semi-convex i.e. } \kappa_i \geq -\eta. \end{aligned}$$

which implies the desired bound for κ_1 .

Now suppose that the process stops at some index $m \leq l < k$ and then Lu's inequality (2.2) holds for l :

$$-\sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} \geq (1 - \varepsilon) \frac{\xi_1^2}{\kappa_1^2} - \frac{1}{2} \sum_{i > l} \frac{\sigma_k^{ii} \xi_i^2}{\kappa_1 \sigma_k}.$$

Taking $F = \sigma_k$, $\xi_i = h_{ii1}$ and multiply the inequality by $\frac{\sigma_k}{\kappa_1}$, we have that

$$-\sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} \geq (1 - \varepsilon) \frac{h_{111}^2}{\kappa_1^3} \sigma_k - \frac{1}{2} \sum_{i > l} \frac{F^{ii} h_{ii1}^2}{\kappa_1^2} - \frac{F_1^2}{\kappa_1 \sigma_k}.$$

The second line of (4.6) can now be estimated as follows:

$$\begin{aligned} &-\sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} \\ (4.7) \quad &\geq -\frac{F_1^2}{\kappa_1 \sigma_k} + \frac{h_{111}^2}{\kappa_1^2} \left(\frac{1 - \varepsilon}{\kappa_1} \sigma_k - \sigma_k^{11} \right) + \sum_{i > l} \left[\frac{2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} - \frac{1}{2\kappa_1^2} \right] F^{ii} h_{ii1}^2 \\ &\geq -C\kappa_1 + (1 - \varepsilon) \sigma_k(\kappa|1) \frac{h_{111}^2}{\kappa_1^3} - \varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} \end{aligned}$$

where we have used

$$\sigma_k(\kappa) = \kappa_1 \sigma_{k-1}(\kappa|1) + \sigma_k(\kappa|1)$$

and

$$\frac{2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} - \frac{1}{2\kappa_1^2} = \frac{3\kappa_1 + \tilde{\kappa}_i}{2\kappa_1^2 (\kappa_1 - \tilde{\kappa}_i)} \geq 0 \quad \text{by repeating the proof of claim 4.1.}$$

Substituting (4.7) into (4.6), we obtain that

$$\begin{aligned} (4.8) \quad 0 &\geq (N - 1) \sum_i \kappa_i^2 F^{ii} - \varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} \\ &\quad + (\alpha\phi' - C) \sum_i f_i + (1 - \varepsilon) \sigma_k(\kappa|1) \frac{h_{111}^2}{\kappa_1^3} \\ &\quad - C\kappa_1 - C\alpha - CN - C. \end{aligned}$$

We proceed to show how to deal with the terms involving ε . By the critical equation (3.1), we have that

$$\begin{aligned} -\varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} &= -\varepsilon F^{11} \left(N \frac{u_1}{u} - \alpha \Phi_1 \right)^2 \\ &\geq -C\varepsilon N^2 F^{11} \kappa_1^2 - C\varepsilon \alpha^2 F^{11} \\ &\geq -C\varepsilon N^2 F^{11} \kappa_1^2 \end{aligned}$$

and

$$\begin{aligned}
(1 - \varepsilon)\sigma_k(\kappa|1)\frac{h_{111}^2}{\kappa_1^3} &\geq -C\kappa_2 \cdots \kappa_k \cdot \eta \cdot \frac{1}{\kappa_1} \cdot \left(N\frac{u_1}{u} - \alpha\Phi_1\right)^2 \\
&\geq -CN^2\kappa_1 \cdots \kappa_k - C\alpha^2 \frac{\kappa_2 \cdots \kappa_k}{\kappa_1} \\
&\geq -CN^2\kappa_1 \cdots \kappa_k \quad \text{if } \kappa_i \geq -\eta.
\end{aligned}$$

Recall that on Σ , ϕ' is bounded from below by a positive controlled constant and so by choosing α large enough, we may assume that the coefficient $\alpha\phi' - C_1$ is positive. Therefore, by lemma 2.7, lemma 2.8 and choosing α large enough, we have that

$$\begin{aligned}
&(\alpha\phi' - C_1) \sum_i f_i + (1 - \varepsilon)\sigma_k(\kappa|1)\frac{h_{111}^2}{\kappa_1^3} \\
&= (\alpha\phi' - C_1)(n - k + 1)\sigma_{k-1} + (1 - \varepsilon)\sigma_k(\kappa|1)\frac{h_{111}^2}{\kappa_1^3} \\
&\geq C\alpha\kappa_1 \cdots \kappa_{k-1} - CN^2\kappa_1 \cdots \kappa_k \\
&= (\kappa_1 \cdots \kappa_{k-1}) \cdot (C\alpha - CN^2\kappa_k)
\end{aligned}$$

If $C\alpha - CN^2\kappa_k \geq 0$, then we are done dealing with this term. Otherwise, we would have $\kappa_k \geq CN$ by choosing $\alpha = N^3$, which will then imply that

$$\begin{aligned}
\sigma_k(\kappa) &\geq \kappa_1 \cdots \kappa_k - C\kappa_1 \cdots \kappa_{k-1} \cdot \eta \\
&= \kappa_1 \cdots \kappa_{k-1}(\kappa_k - C\eta) \\
&\geq \kappa_1 \cdots \kappa_{k-1}(CN - C\eta) \\
&\geq C\kappa_1
\end{aligned}$$

by choosing N sufficiently large. Therefore, we may assume that $C\alpha - CN^2\kappa_k \geq 0$ and remove the second line of (4.8) to have that

$$\begin{aligned}
0 &\geq (N - 1) \sum_i \kappa_i^2 F^{ii} - \varepsilon \frac{F^{11} h_{111}^2}{\kappa_1^2} - C\kappa_1 - C\alpha - CN - C \\
&\geq (N - 1) \sum_i \kappa_i^2 F^{ii} - C\varepsilon N^2 F^{11} \kappa_1^2 - C\kappa_1 - C\alpha - CN - C \\
&\geq (N - C) \sum_i \kappa_i^2 F^{ii} - C\kappa_1 - C\alpha - CN - C \\
&\geq CN\kappa_1\sigma_k - C\kappa_1 - C\alpha - CN - C \\
&\geq (CN - C)\kappa_1 - C
\end{aligned}$$

where we have chosen $\varepsilon = \frac{1}{N^2}$ and applied lemma 2.10. The desired estimate for κ_1 follows by choosing N large enough.

5. THE PRESCRIBED CURVATURE MEASURE TYPE EQUATION

In this section, we prove theorem 1.10 by deriving C^0, C^1 and C^2 estimates for the prescribed curvature measure type equation

$$\sigma_k(\kappa[\Sigma]) = \langle X, \nu \rangle^p \psi(X).$$

5.1. The C^0 Estimate.

Lemma 5.1. *Let (i) $1 \leq k \leq n-1, p \in (-\infty, 0) \cup (0, 1]$, or (ii) $k = n, p \in (-\infty, 0) \cup (0, 1)$. If $\Sigma = \{(r(z), z) : z \in M\}$ is a closed k -convex hypersurface in $\overline{M}$ which is strictly star-shaped with respect to the origin and satisfies (1.6) for some positive $\psi \in C^2(\Sigma)$, then there exist $C_1, C_2 > 0$ depending only on $n, k, \min_M \psi, \max_M \psi$ such that*

$$C_1 \leq \min r \leq \max r \leq C_2.$$

For $k = n$ and $p = 1$, the same result holds under the additional assumption that $\min_M \psi > 1$.

Proof. Our proof follows that of Guan-Lin-Ma [21, Lemma 2.2] and Yang [54, Theorem 3.2]. Observe that the equation can be written as

$$\sigma_k(h_j^i) = \langle X, \nu \rangle^p \overline{\psi}(X)$$

where

$$\langle X, \nu \rangle = \frac{\phi^2}{\sqrt{\phi^2 + |\nabla' r|^2}}, \quad \overline{\psi} = \phi^{-(n+1)} \psi;$$

see (2.12) in [21]. Since Σ is compact, r attains its maximum at some point z_0 . Since at this point, $\nabla' r = 0$, it follows that

$$\begin{aligned} h_j^i &= \frac{1}{\phi^2 \sqrt{\phi^2 + |\nabla' r|^2}} \left(\delta_{ik} - \frac{r^i r^k}{\phi^2 + |\nabla' r|^2} \right) (-\phi r_{kj} + 2\phi' r_k r_j + \phi^2 \phi' \delta_{kj}) \\ &= \frac{1}{\phi^3} (-\phi r_{ij} + \phi^2 \phi' \delta_{ij}) \\ &\geq \frac{\phi'}{\phi} \delta_{ij} \end{aligned}$$

where we have used the fact that $r_{ij} \leq 0$ at z_0 . Hence, by ellipticity of the equation operator,

$$\psi = \phi^{n+1-p} \cdot \sigma_k(h_j^i) \geq \phi^{n+1-p} \cdot \sigma_k\left(\frac{\phi'}{\phi} \delta_{ij}\right) = \phi^{n+1-p} \cdot \binom{n}{k} \left(\frac{\phi'}{\phi}\right)^k = \binom{n}{k} (\phi')^k \phi^{n+1-p-k}.$$

Similarly, at the point where r attains its minimum, we have that

$$\psi \leq \binom{n}{k} (\phi')^k \phi^{n+1-p-k}.$$

This proof works fine unless $p = 1$ and $k = n$, for which we would need the additional condition that $\min \psi > 1$. □

5.2. The C^1 Estimate for all $p \in \mathbb{R}$.

Theorem 5.2. *If $\Sigma = \{(r(z), z) : z \in M\}$ is a closed k -convex hypersurface in $\overline{M}$ which is strictly star-shaped with respect to the origin and satisfies (1.6) for some positive $\psi \in C^2(\Sigma)$, then there exist $C > 0$ depending only on $n, k, p, \min_M \psi, |\psi|_{C^1}$ and the curvature $\overline{R}$ of $\overline{M}$ such that*

$$\max_M |\nabla' r| \leq C.$$

Remark 5.3. A remarkable feature of this result is that it holds for all $p \neq 0$ without any barrier conditions. See [10, Lemma 3.1], [25, Lemma 5.1] and [50, Lemma 3.2] for the C^1 estimate of the general curvature equation under barrier conditions.

Proof. Our proof employs the test function of Chen-Li-Wang [10, Lemma 3.1] but follows the derivation of Huang-Xu [26, Lemma 2.3]. We first observe that

$$\langle V, \nu \rangle = \frac{\phi(r)^2}{\sqrt{\phi(r)^2 + |\nabla' r|^2}}$$

and so this gradient bound is equivalent to $\langle V, \nu \rangle \geq C > 0$, due to the C^0 estimate.

Consider the test function

$$P = \gamma(\Phi) - \log u$$

where γ is to be determined later. Suppose that P attains its maximum at $z_0 \in \Sigma$. If V is parallel to the normal direction ν at z_0 , then $\langle V, \nu \rangle = |V|$ and we are done. Assume this is not the case, then we can choose a local orthonormal frame $\{E_1, \dots, E_n\}$ such that $\langle V, E_1 \rangle \neq 0$ and $\langle V, E_i \rangle = 0$ for $i \geq 2$. Note also that $V = \langle V, E_1 \rangle E_1 + \langle V, \nu \rangle \nu$.

Then at z_0 , we have that

$$\begin{aligned} (5.1) \quad 0 &= P_i = \gamma' \Phi_i - \frac{u_i}{u} \\ 0 &\geq P_{ii} = \gamma'' \Phi_i^2 + \gamma' \Phi_{ii} - \frac{u_{ii}}{u} + \frac{u_i^2}{u^2} \\ &= \gamma'' \Phi_i^2 + \gamma' \Phi_{ii} - \frac{u_{ii}}{u} + (\gamma')^2 \Phi_i^2 \\ (5.2) \quad &= \gamma' \Phi_{ii} - \frac{u_{ii}}{u} + [\gamma'' + (\gamma')^2] \Phi_i^2 \end{aligned}$$

By (5.1), we have that

$$(5.3) \quad h_{11} = u\gamma' \quad \text{and} \quad h_{i1} = 0 \quad \forall i \geq 2.$$

We can then rotate the coordinate system such that $\{E_1, \dots, E_n\}$ are the principal curvature directions of the second fundamental form so that $h_{ij} = \kappa_i \delta_{ij}$.

Contracting (5.2) with F^{ii} and apply lemma 2.4, we have that

$$\begin{aligned} 0 &\geq \gamma' \sum F^{ii} \Phi_{ii} - \frac{1}{u} \sum F^{ii} u_{ii} + [\gamma'' + (\gamma')^2] \sum F^{ii} \Phi_i^2 \\ &= \gamma' \sum F^{ii} (\phi' - u\kappa_i) - \frac{1}{u} \sum F^{ii} \left[\sum_k (h_{iik} - \bar{R}_{0iik}) \Phi_k + \phi' \kappa_i - \kappa_i^2 u \right] \\ &\quad + [\gamma'' + (\gamma')^2] \sum F^{ii} \Phi_i^2 \\ &\geq \gamma' \phi' \sum F^{ii} - u\gamma' \cdot k\Psi - \frac{1}{u} \sum F^{ii} h_{ii1} \Phi_1 + \frac{1}{u} \sum F^{ii} \bar{R}_{0ii1} \Phi_1 - \frac{\phi'}{u} \cdot k\Psi + \sum F^{ii} \kappa_i^2 \\ &\quad + [\gamma'' + (\gamma')^2] F^{11} \Phi_1^2 \end{aligned}$$

Differentiating (1.10), we have that

$$F^{ii} h_{ii1} = pu^{p-1} u_1 \cdot \psi + u^p \cdot \psi_1.$$

Substituting this and $\Psi = u^p \psi$ into the above inequality, we obtain that

$$\begin{aligned} 0 &\geq \gamma' \phi' \sum F^{ii} - k\gamma' u^{p+1} \psi - \frac{\Phi_1}{u} u^p \left(p\psi \frac{u_1}{u} + \psi_1 \right) \\ &\quad - \frac{\Phi_1}{u} \sum F^{ii} \bar{R}_{0ii1} - k\phi' u^{p-1} \psi + \sum F^{ii} \kappa_i^2 + [\gamma'' + (\gamma')^2] F^{11} \Phi_1^2 \end{aligned}$$

where we have also used the skew symmetry property $\bar{R}_{0ii1} = -\bar{R}_{0i1i}$.

By (5.1), we have that

$$\frac{u_1}{u} = \gamma' \Phi_1.$$

The inequality then becomes

$$\begin{aligned} 0 &\geq \gamma' \phi' \sum F^{ii} - k\gamma' u^{p+1} \psi - pu^{p-1} \gamma' \psi \Phi_1^2 - \Phi_1 u^{p-1} \psi_1 \\ &\quad - \frac{\Phi_1}{u} \sum F^{ii} \bar{R}_{0i1i} - k\phi' u^{p-1} \psi + \sum F^{ii} \kappa_i^2 + [\gamma'' + (\gamma')^2] F^{11} \Phi_1^2 \end{aligned}$$

Now, using

$$|V|^2 = \langle V, E_1 \rangle^2 + \langle V, \nu \rangle^2 \quad \text{i.e.} \quad \Phi_1^2 = |V|^2 - u^2,$$

we have

$$\begin{aligned} (5.4) \quad 0 &\geq [\gamma'' + (\gamma')^2] [|V|^2 - u^2] F^{11} + \gamma' \phi' \sum F^{ii} - k\gamma' \psi u^{p+1} - [|V|^2 - u^2] p\gamma' \psi u^{p-1} \\ &\quad - \Phi_1 \psi_1 u^{p-1} - k\phi' \psi u^{p-1} + \sum F^{ii} \kappa_i^2 - \frac{\Phi_1}{u} \sum F^{ii} \bar{R}_{0i1i}. \end{aligned}$$

Rearranging the terms in (5.4), we have that

$$\begin{aligned} (5.5) \quad &[\gamma'' + (\gamma')^2] F^{11} u^2 + (k-p)\gamma' \psi u^{p+1} \\ &+ u^{p-1} [|V|^2 p\gamma' \psi + \Phi_1 \psi_1 + k\phi' \psi] \\ &\geq [\gamma'' + (\gamma')^2] \cdot |V|^2 F^{11} + \gamma' \phi' \sum F^{ii} \\ &+ \sum F^{ii} \kappa_i^2 - \frac{\Phi_1}{u} \sum F^{ii} \bar{R}_{0i1i} \end{aligned}$$

Now, choose

$$\gamma(s) = \frac{\alpha p}{s} \quad \text{i.e.} \quad \gamma'(s) = -\frac{\alpha p}{s^2} \quad \text{and} \quad \gamma''(s) = \frac{2\alpha p}{s^3},$$

where $\alpha > 0$ is some possibly large constant depending on the given value of p . We claim that

Claim 5.4. $(k-p)\gamma' \psi u^{p+1} + u^{p-1} [|V|^2 p\gamma' \psi + \Phi_1 \psi_1 + k\phi' \psi] \leq 0$

Proof. If $p < 0$ or $p > t$, then

$$\begin{aligned} &(k-p)\gamma' \psi u^{p+1} + p|V|^2 \gamma' \psi u^{p-1} \\ &= \gamma' \psi u^{p-1} [(k-p)u^2 + p|V|^2] \\ &= -\alpha p \frac{\psi}{\Phi^2} u^{p-1} [(k-p)u^2 + p|V|^2] \\ &\leq -\frac{\alpha p^2 \psi}{2\Phi^2} |V|^2 u^{p-1}, \end{aligned}$$

where we have assumed that

$$u^2 \leq \frac{p}{2(p-k)} |V|^2.$$

The desired inequality follows by choosing α large enough.

If $0 < p \leq k$, then we immediately have that

$$(k-p)\gamma' \psi u^{p+1} \leq 0$$

and

$$|V|^2 p\gamma' \psi + \Phi_1 \psi_1 + k\phi' \psi = -\frac{\alpha p^2}{\Phi^2} |V|^2 \psi + \Phi_1 \psi_1 + k\phi' \psi \leq 0$$

by choosing α large enough. □

Claim 5.5. $\sum F^{ii} \kappa_i^2 - \frac{\Phi_1}{u} \sum F^{ii} \bar{R}_{0i1i} \geq -C \sum F^{ii}$.

Proof. Note that in [10], they applied (5.3) to have that $\sum F^{ii} \kappa_i^2 \geq F^{11} \kappa_1^2 = u^2 (\gamma')^2 F^{11}$, but this would cancel out the important term in $[\gamma'' + (\gamma')^2] F^{11} u^2$. So here we just throw it away by positivity i.e. $\sum F^{ii} \kappa_i^2 \geq 0$.

For the second term, since $V = \langle V, E_1 \rangle E_1 + \langle V, \nu \rangle \nu$, it follows that $V \perp \text{Span}\{E_2, \dots, E_n\}$. Since we also have $E_1, \nu \perp \text{Span}\{E_2, \dots, E_n\}$, we can choose a coordinate system such that $\bar{e}_1 \perp \text{Span}\{E_2, \dots, E_n\}$, which implies that the pair $\{V, \bar{e}_1\}$ and $\{\nu, E_1\}$ would lie in the same plane, and

$$\text{Span}\{E_2, \dots, E_n\} = \text{Span}\{\bar{e}_2, \dots, \bar{e}_n\}.$$

Thus, by choosing $E_i = \bar{e}_i$ for $2 \leq i \leq n$, we may write

$$\begin{aligned} \nu &= \langle \nu, \bar{e}_0 \rangle \bar{e}_0 + \langle \nu, \bar{e}_1 \rangle \bar{e}_1 = \frac{u}{\phi} \bar{e}_0 + \langle \nu, \bar{e}_1 \rangle \bar{e}_1, \\ E_1 &= \langle E_1, \bar{e}_0 \rangle \bar{e}_0 + \langle E_1, \bar{e}_1 \rangle \bar{e}_1. \end{aligned}$$

The curvature tensor can then be computed as

$$\begin{aligned} \bar{R}_{0i1i} &= \bar{R}(\nu, E_i, E_1, E_i) \\ &= \frac{u}{\phi} \langle E_1, \bar{e}_0 \rangle \bar{R}(\bar{e}_0, \bar{e}_i, \bar{e}_0, \bar{e}_i) + \langle \nu, \bar{e}_1 \rangle \langle E_1, \bar{e}_1 \rangle \bar{R}(\bar{e}_1, \bar{e}_i, \bar{e}_1, \bar{e}_i) \\ &= \frac{u}{\phi} \langle E_1, \bar{e}_0 \rangle \bar{R}(\bar{e}_0, \bar{e}_i, \bar{e}_0, \bar{e}_i) - \frac{u \langle \nu, \bar{e}_1 \rangle^2}{\langle E_1, V \rangle} \bar{R}(\bar{e}_1, \bar{e}_i, \bar{e}_1, \bar{e}_i) \\ &= u \left(\frac{\langle E_1, \bar{e}_0 \rangle}{\phi} \bar{R}(\bar{e}_0, \bar{e}_i, \bar{e}_0, \bar{e}_i) - \frac{\langle \nu, \bar{e}_1 \rangle^2}{\langle E_1, V \rangle} \bar{R}(\bar{e}_1, \bar{e}_i, \bar{e}_1, \bar{e}_i) \right), \end{aligned}$$

where the second equality follows from $0 = \bar{R}_{ijk0}$ [10, Lemma 2.1] and the third equality is due to $0 = \langle V, \bar{e}_1 \rangle$. This cancels out the $\frac{1}{u}$ factor in the front and the result follows. $\square$

With these claims, the inequality (5.5) is reduced to

$$(5.6) \quad [\gamma'' + (\gamma')^2] F^{11} u^2 \geq [\gamma'' + (\gamma')^2] \cdot |V|^2 F^{11} + (\gamma' \phi' + C) \sum F^{ii}.$$

Finally, we can finish the derivation by one more claim.

Claim 5.6. $[\gamma'' + (\gamma')^2] \cdot |V|^2 F^{11} + (\gamma' \phi' + C) \sum F^{ii} \geq C F^{11}$.

Proof. By our choice of γ , we have that

$$\begin{aligned} &[\gamma'' + (\gamma')^2] \cdot |V|^2 F^{11} + \gamma' \phi' \sum F^{ii} \\ &= \left[\frac{2\alpha p}{\Phi^2} + \frac{\alpha^2 p^2}{\Phi^4} \right] \cdot |V|^2 F^{11} - \alpha p \frac{\phi'}{\Phi^2} \sum F^{ii}. \end{aligned}$$

If $p < 0$, the first term is still positive by choosing α large enough and the second term will immediately imply that

$$-\alpha p \frac{\phi'}{\Phi^2} \sum F^{ii} \geq C F^{11}.$$

If $p \geq 0$, then by (5.3) we have that $h_{11} = u\gamma' \leq 0$. Hence, we have that

$$F^{11} = \sigma_{k-1}(\kappa|1) = \sigma_{k-1}(\kappa) - \kappa_1 \sigma_{k-2}(\kappa|1) \geq \sigma_{k-1}(\kappa) \geq c(n, k) \sum F^{ii}$$

by lemma 2.7. The result then follows by choosing α large enough. $\square$

With this claim, the inequality (5.1) yields

$$[\gamma'' + (\gamma')^2] F^{11} u^2 \geq C F^{11}$$

and the bound $u \geq C$ is obtained by choosing α large enough so that $\gamma'' + (\gamma')^2 > 0$. $\square$

5.3. The C^2 Estimate for $p \leq 1$.

Finally, we derive the C^2 estimate for the prescribed curvature measure type equation i.e.

$$\sigma_k(\kappa(X)) = \langle X, \nu \rangle^p \psi(X) \quad \text{where } p \in (-\infty, 0) \cup (0, 1].$$

Proof of Theorem 1.8. For this case, we will use the test function of Spruck-Xiao [50]

$$Q = \log \kappa_{\max} - \log(u - a) + \alpha \Phi$$

where $u := \langle X, \nu \rangle$ and $u \geq 2a$, and $\alpha > 0$ is a constant to be chosen later.

By proceeding exactly as in section 3, or simply by substituting $N = 1$ in (3.13), we obtain that

$$(5.7) \quad \begin{aligned} 0 \geq & \left(\frac{u}{u-a} - 1 \right) \sum_i F^{ii} \kappa_i^2 + (\alpha \phi' - C) \sum_i F^{ii} + k F \kappa_1 - C \alpha - C \\ & - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} - \frac{F^{11} h_{111}^2}{\kappa_1^2} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} + \frac{F^{11} u_1^2}{(u-a)^2} \\ & + \sum_{i \in I} \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + \sum_{i \in I} \frac{F^{ii} u_i^2}{(u-a)^2} + \frac{\Psi_{11}}{\kappa_1} - \frac{1}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right). \end{aligned}$$

Recall that the index set I is defined as

$$I = \{i > m : \kappa_1 + \tilde{\kappa}_i < 0\}.$$

There are two major obstacles here: (1) unlike in claim 4.1, this time it is not clear if the quotient

$$\frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i}$$

is positive without extra conditions; and (2) it is difficult to handle the large negative term

$$-\frac{F^{11} h_{111}^2}{\kappa_1^2};$$

we have to make full use of all the positive terms in (5.7) combined with the special structure of the right-hand side $\Psi = u^p \psi$.

The first trick is to split the $\sum_i \kappa_i^2 F^{ii}$ term into

$$(5.8) \quad \begin{aligned} \frac{a}{u-a} \sum_i \kappa_i^2 F^{ii} &= \left(\frac{n-1}{n} \frac{a}{u-a} + \frac{1}{n} \frac{a}{u-a} \right) \sum_{i=1}^n F^{ii} \kappa_i^2 \\ &\geq \frac{n-1}{n} \frac{a}{u-a} \sum_{i=1}^n F^{ii} \kappa_i^2 + \frac{1}{n} \frac{a}{u-a} F^{11} \kappa_1^2 + \frac{1}{n} \frac{a}{u-a} \sum_{i \in I} F^{ii} \kappa_i^2 \end{aligned}$$

We claim that

Claim 5.7.

$$\frac{1}{n} \frac{a}{u-a} \sum_{i \in I} \kappa_i^2 f_i + \sum_{i \in I} \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + \sum_{i \in I} \frac{F^{ii} u_i^2}{(u-a)^2} \geq 0$$

and

$$\frac{1}{n} \frac{a}{u-a} F^{11} \kappa_1^2 - \frac{F^{11} h_{111}^2}{\kappa_1^2} + \frac{F^{11} u_1^2}{(u-a)^2} \geq 0,$$

where $I = \{i > m : \kappa_1 + \tilde{\kappa}_i < 0\}$.

Remark 5.8. The proof for this claim is in fact independent of this $\frac{1}{n}$ coefficient. The reason for the splitting (5.8) is that even after we have used parts of the sum to prove claim 5.7, we are still left with a complete sum $\frac{n-1}{n} \frac{a}{u-a} \sum F^{ii} \kappa_i$ or simply the term $F^{11} \kappa_1^2$, which are both of the same order as κ_1 and will crucially be used in the subsequent derivation. That is, the most important is not the magnitude of its coefficient but merely its presence.

proof of the claim. Observe the trivial fact that

$$\frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \geq -1$$

and recall from the first order critical equation (3.1) that

$$\frac{h_{11i}^2}{\kappa_1^2} = \left(\frac{u_i}{u-a} - \alpha \Phi_i \right)^2.$$

Also, note that for $i \in I$, we have $|\kappa_i| \geq \kappa_1$. It is then straightforward to have that

$$\begin{aligned} & \frac{1}{n} \frac{a}{u-a} \sum_{i \in I} \kappa_i^2 f_i + \sum_{i \in I} \frac{\kappa_1 + \tilde{\kappa}_i}{\kappa_1 - \tilde{\kappa}_i} \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + \sum_{i \in I} \frac{F^{ii} u_i^2}{(u-a)^2} \\ & \geq \sum_{i \in I} \frac{1}{n} \frac{a}{u-a} F^{ii} \kappa_i^2 - \frac{F^{ii} h_{11i}^2}{\kappa_1^2} + \frac{F^{ii} u_i^2}{(u-a)^2} \\ & = \sum_{i \in I} \frac{1}{n} \frac{a}{u-a} F^{ii} \kappa_i^2 - F^{ii} \left(\frac{u_i}{u-a} - \alpha \Phi_i \right)^2 + \frac{F^{ii} u_i^2}{(u-a)^2} \\ & = \sum_{i \in I} \frac{1}{n} \frac{a}{u-a} F^{ii} \kappa_i^2 + 2\alpha F^{ii} \frac{u_i}{u-a} \Phi_i - \alpha^2 F^{ii} \Phi_i^2 \\ & \geq \sum_{i \in I} \frac{1}{n} \frac{a}{u-a} F^{ii} \kappa_i^2 - C\alpha F^{ii} |\kappa_i| - C\alpha^2 F^{ii} \\ & \geq 0 \quad \text{by assuming } \kappa_1 \text{ is sufficiently large.} \end{aligned}$$

The proof for the second inequality is very similar:

$$\begin{aligned} & \frac{1}{n} \frac{a}{u-a} F^{11} \kappa_1^2 - \frac{F^{11} h_{111}^2}{\kappa_1^2} + \frac{F^{11} u_1^2}{(u-a)^2} \\ & \geq \frac{1}{n} \frac{a}{u-a} F^{11} \kappa_1^2 - C\alpha F^{11} |\kappa_1| - C\alpha^2 F^{11} \\ & \geq 0 \quad \text{by assuming } \kappa_1 \text{ is sufficiently large.} \end{aligned}$$

□

With this claim, the inequality (5.7) implies that

$$\begin{aligned} (5.9) \quad 0 & \geq \frac{n-1}{n} \frac{a}{u-a} \sum_{i=1}^n F^{ii} \kappa_i^2 + (\alpha \phi' - C) \sum_{i=1}^n F^{ii} \\ & \quad - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} \\ & \quad + \frac{\Psi_{11}}{\kappa_1} - \frac{1}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right) + kF \kappa_1 - C\alpha - C. \end{aligned}$$

We can proceed to invoke the special structure of Ψ to handle the first two terms in the third line of (5.9). We first find, by lemma 2.4, the first order equation (3.1) and the expression $F = \Psi = u^p \psi$,

that

$$\begin{aligned}
 \Psi_{11} &= (u^p \psi)_{11} \\
 &= \psi \left[pu^{p-1} u_{11} + p(p-1) u^{p-2} u_1^2 \right] + 2\psi_1 \cdot pu^{p-1} u_1 + u^p \cdot \psi_{11} \\
 &= \psi \cdot pu^{p-1} \left[\sum_k (h_{11k} - \bar{R}_{011k}) \Phi_k + \phi' \kappa_1 - \kappa_1^2 u \right] \\
 &\quad + \psi \cdot p(p-1) u^{p-2} u_1^2 + 2p\psi_1 u^p \frac{u_1}{u} + u^p \psi_{11} \\
 &\geq \psi \cdot pu^{p-1} \sum_k h_{11k} \Phi_k - \psi \cdot pu^p \kappa_1^2 + \psi \cdot p(p-1) u^{p-2} u_1^2 - C\kappa_1 - C \\
 &\geq \frac{pF}{u} \sum_k h_{11k} \Phi_k - pF\kappa_1^2 + p(p-1)F \frac{u_1^2}{u^2} - C\kappa_1 - C.
 \end{aligned}$$

for some C depending on $p, \bar{R}, \|\Sigma\|_{C^1}$ and $\|\psi\|_{C^2}$. Then, by noting that

$$\begin{aligned}
 & - \frac{1}{u-a} \sum_k \left(\sum_i f_i h_{iik} \Phi_k \right) + \frac{pF}{u} \sum_k \frac{h_{11k}}{\kappa_1} \Phi_k \\
 &= - \sum_k \frac{F_k \Phi_k}{u-a} + \frac{pF}{u} \sum_k \frac{h_{11k}}{\kappa_1} \Phi_k \\
 &= - \sum_k \left(\psi \cdot pu^{p-1} u_k + u^p \cdot \psi_k \right) \frac{\Phi_k}{u-a} + \frac{pF}{u} \sum_k \left(\frac{u_k}{u-a} - \alpha \Phi_k \right) \Phi_k \\
 &= - \frac{pF}{u} \sum_k \frac{u_k \Phi_k}{u-a} + \frac{pF}{u} \sum_k \frac{u_k \Phi_k}{u-a} - u^p \sum_k \frac{\psi_k \Phi_k}{u-a} - \alpha \frac{pF}{u} \sum_k \Phi_k^2 \\
 &\geq -C\alpha - C,
 \end{aligned}$$

it follows from (5.9) that

$$\begin{aligned}
 (5.10) \quad 0 &\geq \frac{n-1}{n} \frac{a}{u-a} \sum_{i=1}^n F^{ii} \kappa_i^2 + (\alpha\phi' - C) \sum_{i=1}^n F^{ii} \\
 &\quad - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} \\
 &\quad + (k-p)F\kappa_1 + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} - C\alpha - C.
 \end{aligned}$$

It remains to handle the second line and the term

$$p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2},$$

which can be negative if $p \in (0, 1)$.

5.3.1. *Case A:* $\kappa_2 \cdots \kappa_{k-1} \geq \alpha^{-\frac{k-2}{n}}$.

By concavity of the operator $\sigma_k^{1/k}$ i.e.

$$\frac{1}{k} \sigma_k^{\frac{1}{k}-1} \sum_{p \neq q} \sigma_k^{pp,qq} h_{pp1} h_{qq1} + \frac{1}{k} \left(\frac{1}{k} - 1 \right) \sigma_k^{\frac{1}{k}-2} \sum_{p,q} \sigma_k^{pp} h_{pp1} \sigma_k^{qq} h_{qq1} \leq 0,$$

we are able to obtain that

$$-\sum_{p \neq q} F^{pp,qq} h_{pp1} h_{qq1} \geq -\frac{k-1}{k} \frac{F_1^2}{F} \geq -C\kappa_1^2.$$

For the sum $\sum F^{ii}$, we use lemma 2.7 and lemma 2.8 to have that

$$\sum_{i=1}^n F^{ii} = (n-k+1)\sigma_{k-1}(\kappa) \geq c(n,k)\kappa_1 \cdots \kappa_{k-1}.$$

Hence,

$$\begin{aligned} & -\sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + (\alpha\phi' - C_1) \sum_{i=1}^n F^{ii} + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \\ & \geq -C\kappa_1 + (\alpha\phi' - C_1)c(n,k)\kappa_1 \cdots \kappa_{k-1} + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \\ & \geq -C\kappa_1 + C\alpha^{\frac{n-k+2}{n}} \kappa_1 \\ & \geq 0 \quad \text{by choosing } \alpha \text{ sufficiently large,} \end{aligned}$$

where the estimate $p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \geq -C\kappa_1$ is given in remark 5.9. Now, the inequality (5.10) along with lemma 2.10 yields

$$\begin{aligned} 0 & \geq \frac{n-1}{n} \frac{a}{u-a} \sum_{i=1}^n F^{ii} \kappa_i^2 + (k-p)F\kappa_1 - C\alpha - C \\ & \geq CF\kappa_1 - C\alpha - C \end{aligned}$$

as desired.

5.3.2. Case B: $\kappa_2 \cdots \kappa_{k-1} \leq \alpha^{-\frac{k-2}{n}}$.

For this case, we do not have a large enough lower bound for $\sum F^{ii}$ and this is where we have to restrict the range of p . We will proceed by the same iteration argument as in section 4.2 and apply Lu's inequality (2.2). Let $\delta = 1/3$, $\delta_0 = \min\{2k/n, 1/2\}$ and let $\varepsilon > 0$ to be chosen later. We consider two sub-cases according to lemma 2.12.

Case B.1. For each $2 \leq l \leq k$, there exists some $\delta_l > 0$ such that $\kappa_l \geq \delta_l \kappa_1$.

Then we immediately have that

$$\alpha^{-\frac{k-2}{n}} \geq \kappa_2 \cdots \kappa_{k-1} \geq \delta_2 \cdots \delta_{k-1} \kappa_1^{k-1}$$

and the estimate for κ_1 follows.

Case B.2. There exists $m \leq l < k$ such that (2.2) holds.

That is, we have

$$-\sum_{p \neq q} \frac{\sigma_k^{pp,qq} \xi_p \xi_q}{\sigma_k} + \frac{(\sum_i \sigma_k^{ii} \xi_i)^2}{\sigma_k^2} \geq (1-\varepsilon) \frac{\xi_1^2}{\kappa_1^2} - \delta_0 \sum_{i>l} \frac{\sigma_k^{ii} \xi_i^2}{\kappa_1 \sigma_k}.$$

Substituting $\xi_i = h_{ii1}$, the second line in (5.10) then becomes

$$\begin{aligned} & -\sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} \\ & \geq (1-\varepsilon) \sigma_k \frac{h_{111}^2}{\kappa_1^3} + \sum_{i>l} \left[\frac{2}{\kappa_1 (\kappa_1 - \tilde{\kappa}_i)} - \frac{\delta_0}{\kappa_1^2} \right] F^{ii} h_{ii1}^2 - \frac{F_1^2}{\kappa_1 \sigma_k}. \end{aligned}$$

By lemma 2.9 and our choice of δ_0 , we have that

$$\frac{2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} - \frac{\delta_0}{\kappa_1^2} \geq 0.$$

Therefore, by lemma 2.10, the first order equation (3.1) and the expression $F = \Psi = u^p \psi$, we have that

$$\begin{aligned} & \frac{n-1}{n} \frac{a}{u-a} \sum_{i=1}^n F^{ii} \kappa_i^2 - \sum_{p \neq q} \frac{F^{pp,qq} h_{pp1} h_{qq1}}{\kappa_1} + 2 \sum_{i \neq 1} \frac{F^{ii} h_{ii1}^2}{\kappa_1(\kappa_1 - \tilde{\kappa}_i)} + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \\ & \geq \frac{n-1}{n} \frac{a}{u-a} \sum_{i=1}^n F^{ii} \kappa_i^2 + (1-\varepsilon) \sigma_k \frac{h_{111}^2}{\kappa_1^3} - \frac{F_1^2}{\kappa_1 \sigma_k} + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \\ & = CF\kappa_1 + (1-\varepsilon) \frac{F}{\kappa_1} \left(\frac{u_1}{u-a} - \alpha \Phi_1 \right)^2 - \frac{(\psi \cdot pu^{p-1}u_1 + u^p \psi_1)^2}{\kappa_1 \sigma_k} + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \\ & = CF\kappa_1 + (1-\varepsilon) \frac{F}{\kappa_1} \left(\frac{u_1^2}{(u-a)^2} - 2\alpha \frac{u_1 \Phi_1}{u-a} + \alpha^2 \Phi_1^2 \right) \\ & \quad - \frac{1}{\kappa_1 \sigma_k} \left(\psi^2 p^2 u^{2p-2} u_1^2 + 2\psi pu^{2p-1} u_1 \psi_1 + u^{2p} \psi_1^2 \right) + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \\ & \geq CF\kappa_1 + \left[(1-\varepsilon) \frac{F}{\kappa_1} \frac{u_1^2}{(u-a)^2} - C\alpha \right] - p^2 \frac{F}{\kappa_1} \frac{u_1^2}{u^2} - C + p(p-1) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \\ & = CF\kappa_1 + \left[(1-\varepsilon) \frac{F}{\kappa_1} \frac{u_1^2}{(u-a)^2} - p \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \right] + \left[-p^2 \frac{F}{\kappa_1} \frac{u_1^2}{u^2} + p^2 \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \right] - C\alpha - C \\ & = CF\kappa_1 - \varepsilon \frac{F}{\kappa_1} \frac{u_1^2}{(u-a)^2} + \left[\frac{F}{\kappa_1} \frac{u_1^2}{(u-a)^2} - p \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \right] - C\alpha - C \\ & \geq CF\kappa_1 - C\varepsilon F\kappa_1 + (1-p) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} - C\alpha - C \end{aligned}$$

With this estimate, (5.10) then implies that

$$(5.11) \quad 0 \geq (C - C\varepsilon)F\kappa_1 + (1-p) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} + (k-p)F\kappa_1 - C\alpha - C.$$

If $p \leq 1$, then the proof is complete by choosing a sufficiently small $\varepsilon > 0$. □

Remark 5.9. It can be clearly seen that a sufficient condition for the estimate to hold is

$$(5.12) \quad (1-p) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} + (k-p)F\kappa_1 \geq 0.$$

Note that

$$u_1^2 = \kappa_1^2 \Phi_1^2 = \kappa_1^2 (|V|^2 - u^2),$$

so it follows that

$$(1-p) \frac{F}{\kappa_1} \frac{u_1^2}{u^2} \geq (1-p) \left(\frac{1}{\varepsilon_0} - 1 \right) F\kappa_1,$$

where

$$\varepsilon_0 := \inf_{\Sigma} \frac{u^2}{|V|^2}.$$

Hence, the inequality (5.12) holds if

$$(1-p)\left(\frac{1}{\varepsilon_0}-1\right)+(k-p)\geq 0,$$

which is true if and only if

$$p\leq 1+\varepsilon_0(k-1).$$

This shows that our estimate, in fact, holds for a wider range $p\leq 1+C$ for some $C>0$ depending on k and $\|\Sigma\|_{C^1}$. Taking $k=2$ will recover Chen's upper bound for p in [9, (4.32)-(4.33)].

However, since this constant is not universal, we shall not state it explicitly in our theorem.

REFERENCES

- [1] A. D. Aleksandrov, *Uniqueness theorems for surfaces in the large. I*, Vestnik Leningrad. Univ. **11** (1956), no. 19, 5–17. MR86338
- [2] A. Alexandroff, *Zur theorie der gemischten volumina von konvexen körpern. ii. neue ungleichungen zwischen den gemischten volumina und ihre anwendungen*, Rec. Math. [Mat. Sbornik] N.S. **2(44)** (1937), 1205–1238.
- [3] Ben Andrews, *Contraction of convex hypersurfaces in Euclidean space*, Calc. Var. Partial Differential Equations **2** (1994), no. 2, 151–171. MR1385524
- [4] I. Ja. Bakel'man and B. E. Kantor, *Estimates of the solutions of quasilinear elliptic equations that are connected with problems of geometry "in the large"*, Mat. Sb. (N.S.) **91(133)** (1973), 336–349, 471. MR333441
- [5] ———, *Existence of a hypersurface homeomorphic to the sphere in Euclidean space with a given mean curvature*, Geometry and topology, No. 1 (Russian), 1974, pp. 3–10. MR423266
- [6] J. Lucas M. Barbosa, Jorge H. S. Lira, and Vladimir I. Oliker, *A priori estimates for starshaped compact hypersurfaces with prescribed m th curvature function in space forms*, Nonlinear problems in mathematical physics and related topics, I, 2002, pp. 35–52. MR1970603
- [7] L. Caffarelli, L. Nirenberg, and J. Spruck, *The Dirichlet problem for nonlinear second-order elliptic equations. I. Monge-Ampère equation*, Comm. Pure Appl. Math. **37** (1984), no. 3, 369–402. MR739925
- [8] ———, *Nonlinear second order elliptic equations. IV. Starshaped compact Weingarten hypersurfaces*, Current topics in partial differential equations, 1986, pp. 1–26. MR1112140
- [9] ChuanQiang Chen, *Optimal concavity of some Hessian operators and the prescribed σ_2 curvature measure problem*, Sci. China Math. **56** (2013), no. 3, 639–651. MR3017945
- [10] Daguang Chen, Haizhong Li, and Zhizhang Wang, *Starshaped compact hypersurfaces with prescribed Weingarten curvature in warped product manifolds*, Calc. Var. Partial Differential Equations **57** (2018), no. 2, Paper No. 42, 26. MR3764587
- [11] Shiu Yuen Cheng and Shing Tung Yau, *On the regularity of the solution of the n -dimensional Minkowski problem*, Comm. Pure Appl. Math. **29** (1976), no. 5, 495–516. MR423267
- [12] Shiing-shen Chern, *Integral formulas for hypersurfaces in Euclidean space and their applications to uniqueness theorems*, J. Math. Mech. **8** (1959), 947–955. MR114170
- [13] ———, *Differential geometry and integral geometry*, Proc. Internat. Congress Math. 1958, 1960, pp. 441–449. MR120601
- [14] Jianchun Chu, *A simple proof of curvature estimate for convex solution of k -Hessian equation*, Proc. Amer. Math. Soc. **149** (2021), no. 8, 3541–3552. MR4273155
- [15] Francisco J. de Andrade, João Lucas M. Barbosa, and Jorge H. S. de Lira, *Closed Weingarten hypersurfaces in warped product manifolds*, Indiana Univ. Math. J. **58** (2009), no. 4, 1691–1718. MR2542970
- [16] Ph. Delanoë, *Plongements radiaux $S^n \hookrightarrow \mathbf{R}^{n+1}$ à courbure de Gauss positive prescrite*, Ann. Sci. École Norm. Sup. (4) **18** (1985), no. 4, 635–649. MR839688
- [17] Claus Gerhardt, *Closed Weingarten hypersurfaces in Riemannian manifolds*, J. Differential Geom. **43** (1996), no. 3, 612–641. MR1412678
- [18] Bo Guan and Pengfei Guan, *Convex hypersurfaces of prescribed curvatures*, Ann. of Math. (2) **156** (2002), no. 2, 655–673. MR1933079
- [19] Pengfei Guan, Junfang Li, and Yanyan Li, *Hypersurfaces of prescribed curvature measure*, Duke Math. J. **161** (2012), no. 10, 1927–1942. MR2954620
- [20] Pengfei Guan, Junfang Li, and Mu-Tao Wang, *A volume preserving flow and the isoperimetric problem in warped product spaces*, Trans. Amer. Math. Soc. **372** (2019), no. 4, 2777–2798. MR3988593
- [21] Pengfei Guan, Changshou Lin, and Xi-Nan Ma, *The existence of convex body with prescribed curvature measures*, Int. Math. Res. Not. IMRN **11** (2009), 1947–1975. MR2507106
- [22] Pengfei Guan, Changshou Lin, and Xi-nan Ma, *The Christoffel-Minkowski problem. II. Weingarten curvature equations*, Chinese Ann. Math. Ser. B **27** (2006), no. 6, 595–614. MR2273800

- [23] Pengfei Guan and Xi-Nan Ma, *The Christoffel-Minkowski problem. I. Convexity of solutions of a Hessian equation*, Invent. Math. **151** (2003), no. 3, 553–577. MR1961338
- [24] Pengfei Guan, Xi-Nan Ma, and Feng Zhou, *The Christoffel-Minkowski problem. III. Existence and convexity of admissible solutions*, Comm. Pure Appl. Math. **59** (2006), no. 9, 1352–1376. MR2237290
- [25] Pengfei Guan, Changyu Ren, and Zhizhang Wang, *Global C^2 -estimates for convex solutions of curvature equations*, Comm. Pure Appl. Math. **68** (2015), no. 8, 1287–1325. MR3366747
- [26] Yong Huang and Lu Xu, *Two problems related to prescribed curvature measures*, Discrete Contin. Dyn. Syst. **33** (2013), no. 5, 1975–1986. MR3002739
- [27] N. M. Ivochkina, *Solution of the Dirichlet problem for equations of m th order curvature*, Mat. Sb. **180** (1989), no. 7, 867–887, 991. MR1014618
- [28] ———, *The Dirichlet problem for the curvature equation of order m* , Algebra i Analiz **2** (1990), no. 3, 192–217. MR1073214
- [29] Qinian Jin and Yanyan Li, *Starshaped compact hypersurfaces with prescribed k -th mean curvature in hyperbolic space*, Discrete Contin. Dyn. Syst. **15** (2006), no. 2, 367–377. MR2199434
- [30] Hans Lewy, *On differential geometry in the large. I. Minkowski’s problem*, Trans. Amer. Math. Soc. **43** (1938), no. 2, 258–270. MR1501942
- [31] Ming Li, Changyu Ren, and Zhizhang Wang, *An interior estimate for convex solutions and a rigidity theorem*, J. Funct. Anal. **270** (2016), no. 7, 2691–2714. MR3464054
- [32] Yan Yan Li, *Interior gradient estimates for solutions of certain fully nonlinear elliptic equations*, J. Differential Equations **90** (1991), no. 1, 172–185. MR1094454
- [33] ———, *Group invariant convex hypersurfaces with prescribed Gauss-Kronecker curvature*, Multidimensional complex analysis and partial differential equations (São Carlos, 1995), 1997, pp. 203–218. MR1447225
- [34] YanYan Li, Han Lu, and Siyuan Lu, *On the σ_2 -Nirenberg problem on $\mathbb{S}^2$* , J. Funct. Anal. **283** (2022), no. 10, Paper No. 109606, 50. MR4458997
- [35] YanYan Li, Luc Nguyen, and Bo Wang, *The axisymmetric σ_k -Nirenberg problem*, J. Funct. Anal. **281** (2021), no. 9, Paper No. 109198, 60. MR4291513
- [36] ———, *On the σ_k -Nirenberg problem*, Amer. J. Math. **146** (2024), no. 1, 241–276. MR4691488
- [37] Yanyan Li and Vladimir I. Oliker, *Starshaped compact hypersurfaces with prescribed m -th mean curvature in elliptic space*, J. Partial Differential Equations **15** (2002), no. 3, 68–80. MR1927805
- [38] Gary M. Lieberman, *Second order parabolic differential equations*, World Scientific Publishing Co., Inc., River Edge, NJ, 1996. MR1465184
- [39] Siyuan Lu, *Curvature estimates for semi-convex solutions of Hessian equations in hyperbolic space*, Calc. Var. Partial Differential Equations **62** (2023), no. 9, Paper No. 257, 23. MR4659885
- [40] ———, *On the asymptotic Plateau problem in hyperbolic space*, Proc. Amer. Math. Soc. **151** (2023), no. 12, 5443–5451. MR4648938
- [41] Hermann Minkowski, *Volumen und Oberfläche*, Math. Ann. **57** (1903), no. 4, 447–495. MR1511220
- [42] Louis Nirenberg, *The Weyl and Minkowski problems in differential geometry in the large*, Comm. Pure Appl. Math. **6** (1953), 337–394. MR58265
- [43] V. I. Oliker, *Hypersurfaces in $\mathbb{R}^{n+1}$ with prescribed Gaussian curvature and related equations of Monge-Ampère type*, Comm. Partial Differential Equations **9** (1984), no. 8, 807–838. MR748368
- [44] Vladimir I. Oliker, *The Gauss curvature and Minkowski problems in space forms*, Recent developments in geometry (Los Angeles, CA, 1987), 1989, pp. 107–123. MR1034975
- [45] A. V. Pogorelov, *Regularity of a convex surface with given Gaussian curvature*, Mat. Sbornik N.S. **31/73** (1952), 88–103. MR52807
- [46] Aleksey Vasil’ yevich Pogorelov, *The Minkowski multidimensional problem*, Scripta Series in Mathematics, V. H. Winston & Sons, Washington, DC; Halsted Press [John Wiley & Sons], New York-Toronto-London, 1978. Translated from the Russian by Vladimir Oliker, Introduction by Louis Nirenberg. MR478079
- [47] Changyu Ren and Zhizhang Wang, *On the curvature estimates for Hessian equations*, Amer. J. Math. **141** (2019), no. 5, 1281–1315. MR4011801
- [48] ———, *The global curvature estimate for the $n - 2$ Hessian equation*, Calc. Var. Partial Differential Equations **62** (2023), no. 9, Paper No. 239, 50. MR4646879
- [49] Joel Spruck, *Geometric aspects of the theory of fully nonlinear elliptic equations*, Global theory of minimal surfaces, 2005, pp. 283–309. MR2167264
- [50] Joel Spruck and Ling Xiao, *A note on star-shaped compact hypersurfaces with prescribed scalar curvature in space forms*, Rev. Mat. Iberoam. **33** (2017), no. 2, 547–554. MR3651014
- [51] Andrejs E. Treibergs and S. Walter Wei, *Embedded hyperspheres with prescribed mean curvature*, J. Differential Geom. **18** (1983), no. 3, 513–521. MR723815
- [52] Bin Wang, *Starshaped compact hypersurfaces of prescribed curvature in warped product manifolds ii*, forthcoming.
- [53] Xu-Jia Wang, *The k -Hessian equation*, Geometric analysis and PDEs, 2009, pp. 177–252. MR2500526

- [54] Fengrui Yang, *Prescribed curvature measure problem in hyperbolic space*, Comm. Pure Appl. Math. **77** (2024), no. 1, 863–898. MR4666637
- [55] Shing Tung Yau, *Problem section*, Seminar on Differential Geometry, 1982, pp. 669–706. MR645762

DEPARTMENT OF MATHEMATICS, THE CHINESE UNIVERSITY OF HONG KONG, SHATIN, N.T., HONG KONG.
Email address: `bwang@math.cuhk.edu.hk`