

Photon propagator for inflation in the general covariant gauge

Silvije Domazet, Dražen Glavan,^a Tomislav Prokopec^b

^a*CEICO, FZU — Institute of Physics of the Czech Academy of Sciences,
Na Slovance 1999/2, 182 21 Prague 8, Czech Republic*

^b*Institute for Theoretical Physics, Spinoza Institute & EMMEΦ, Utrecht University,
Princetonplein 5, 3584 CC Utrecht, The Netherlands*

E-mail: sdomazet@mail.com, glavan@fzu.cz, t.prokopec@uu.nl

ABSTRACT: Photon propagator for power-law inflation is considered in the general covariant gauges within the canonical quantization formalism. Photon mode functions in covariant gauges are considerably more complicated than their scalar counterparts, except for the special choice of the gauge-fixing parameter we call the simple covariant gauge. We explicitly construct the position space photon propagator in the simple covariant gauge, and find the result considerably more complicated than its scalar counterpart. This is because of the need for explicitly inverting the Laplace operator acting on the scalar propagator, which results in Appell's fourth function. Our propagator correctly reproduces the de Sitter and flat space limits. We use this propagator to compute two simple observables: the off-coincident field strength-field strength correlator and the energy-momentum tensor, both of which yield consistent results. As a spinoff of our computation we also give the exact expression for the Coulomb gauge propagator in power-law inflation in arbitrary dimensions.

1 Introduction

The rapid expansion of primordial inflation quickly dilutes the Universe. Therefore, for most of inflation, it is the coupling of fields to the expanding background that controls the dynamics of their fluctuations. Massless vector fields, such as the electromagnetic field, couple conformally to gravity and thus do not directly sense the expansion of the Universe. Only when their conformality is broken can the effects of the expansion be communicated to them. This happens in instances when the vector field couples to another background field (condensate). Such is the case of symmetry-breaking scalar electrodynamics, where the scalar condensate generates a possibly time-dependent effective mass for the gauge vector. Some of the notable early Universe models that exhibit gauge field amplification due to this mechanism are Higgs inflation (see e.g. [1–11]) and axion inflation (see e.g. [12–18]). The Ratra model [19] is another case of conformality breaking, where a time-dependent scalar condensate couples to the vector kinetic term, thus amplifying vector perturbations in inflation. However, arguably more intriguing are examples where the breaking of conformality is due to loop effects from interactions with non-conformal fluctuations.

Due to the effect of gravitational particle production [20–22], infrared fluctuations of non-conformally coupled fields, such as scalars [23] and gravitons [24], are copiously produced by the almost exponential rapid expansion of the primordial inflationary Universe. Coupling of vectors to such fluctuations can lead to large secular loop corrections for vectors in inflation. This is true in inflationary scalar electrodynamics [25–32], where the photon can develop a mass gap [33–36] due to large infrared fluctuations of the charged scalar, which in turn exponentially damps its spatial correlations. A nonperturbative analysis [37, 38], based on Starobinsky’s stochastic formalism [39, 40], shows the generated photon mass to be nonperturbatively large. Coupling to the charged scalar fluctuations can also induce generation of magnetic fields during inflation [41–45] which could be observable at late times (see e.g. [46] for a review). For electromagnetism interacting with quantum gravity, loops of inflationary gravitons [47–57] engender secular corrections to the Coulomb potential, and the electric field of the propagating photon.

Thus far most of the loop computations have been performed in rigid de Sitter space where the Universe expands exactly exponentially. This idealization is very often a good approximation for inflationary spacetime, and it typically leads to considerable computational simplifications due to enhanced symmetry. However, de Sitter space is not always the most appropriate description for the inflating spacetime. In fact, observations [58] are good enough to decisively point at a small deviation from the exponential expansion of de Sitter space, characterized by slow-roll parameters which are a measure of the rate of decrease in the expansion rate. These slow-roll parameters are some of the key inflationary observables that discriminate between the models. We are interested in corrections that slow-roll parameters can engender

in inflationary loop corrections, especially in the infrared. This is generally not a mathematically tractable problem, but given that slow-roll parameters are small and evolve slowly, deviations from de Sitter expansion can often be treated adiabatically. Here we consider power-law inflation [59, 60] as an analytically tractable model.¹

Power-law inflation is characterized by a constant non-vanishing principal slow-roll parameter. Even though a constant slow-roll parameter is excluded by data [58], it is closer to the realistic slow-roll inflation than a rigid de Sitter space, and allows to capture the effects of non-vanishing principal slow-roll parameter. This should also be a good approximation when the evolution of the slow-roll parameter can be considered adiabatic, which indeed is the case in realistic inflationary models. On a technical side power-law inflation is considered almost as tractable as de Sitter space, because of the experience with scalar fields whose mode functions retain the same functional form as in de Sitter space, and scalar propagators that also retain the same functional form as in de Sitter, being expressible in terms of the hypergeometric function. Some of the notable works dealing with questions of quantum loop corrections in inflating spacetimes with a non-vanishing slow-roll parameter are [62–72].

In order to perform dimensionally regulated loop computations involving vector fields in power-law inflation we need to have the necessary two-point functions (propagators) in D dimensions. While the propagator for the massive vector in the Abelian Higgs model has been worked out in the unitary gauge [73], the massless vector propagator has not been worked out. That is why in this paper we compute the propagator for the massless vector field in power-law inflation. Massless vector field A_μ , that we henceforth refer to as the photon, is a $U(1)$ gauge field, and working out its two-point functions requires fixing a gauge. Perhaps the most natural gauge choice is the general covariant gauge, characterized by adding the covariant gauge-fixing term to the action,

$$S_{\text{gf}}[A_\mu] = \int d^Dx \sqrt{-g} \left[-\frac{1}{2\xi} (\nabla^\mu A_\mu)^2 \right], \quad (1.1)$$

that contains one free gauge-fixing parameter ξ , where $\nabla^\mu A_\mu$ is the covariant derivative of the vector potential with respect to the space time metric $g_{\mu\nu}$. This gauge in general preserves all of the spacetime isometries. However, even in de Sitter space where this is expected to be quite beneficial, the covariant gauges cannot preserve all de Sitter symmetries [74, 75] of the two-point functions, except in the exact transverse gauge [76]. Even though de Sitter invariant solutions to propagator equations of motion do exist [77, 78], they predict spurious behaviour in the infrared [80, 81] due to not respecting the necessary Ward-Takahashi identity [79]. Nonetheless the covariant gauge (1.1) still seems as a reasonable gauge choice to consider in power-law

¹Fermions also couple conformally to gravity, and experience the expansion only by coupling to non-conformally coupled fields. The effects of quantum corrections in power-law inflation with fermions running in the loops can be studied using the propagator worked out in [61].

inflation, and we construct the photon propagator in this gauge. Our computations yield the propagator not as practical as expected. While in de Sitter space the covariant gauge two-point functions are generally expressible in terms of the corresponding scalar two-point functions, in power-law inflation this is no longer the case, which complicates matter significantly.

Determining the photon propagator in momentum space amounts to computing the photon mode functions. We compute them for an arbitrary gauge-fixing parameter ξ in (1.1). Their solutions in power-law inflation is considerably more complicated than the corresponding solutions in de Sitter space. However, we identify a particular choice for the gauge-fixing parameter, that we refer to as the simple covariant gauge, for which the mode functions take on a simple form and are expressible in terms of scalar mode functions and their derivatives. We then use these mode functions to compute the position space photon propagator in the simple covariant gauge. Despite the relative simplicity of the mode functions, this propagator in power-law inflation is still much more complicated than its de Sitter counterpart, and cannot be expressed in terms of a finite number of derivatives of a scalar propagator. Rather, the propagator involves complicated Appell's fourth functions which are two-variable generalizations of the hypergeometric function. This complication appears because of the necessity to explicitly invert the Laplace operator acting on the scalar propagator. This inversion is also necessary to obtain the Coulomb gauge propagator, the result for which we report as well.

Following the introductory section that is now concluding, Sec. 2 collects some of the definitions and results for the scalar mode functions in power-law inflation, and the corresponding two-point functions, that are used throughout. Section 3 summarizes the canonical quantization procedure in the general covariant gauge, giving equations of motion for the field operators and the constraints. The photon mode functions are given in the general covariant gauge in Sec. 4, and in Sec. 5 the photon two-point function is computed in the simple covariant gauge. In Sec. 6 the flat space and the de Sitter space limits are worked out and compared with the literature. Section 7 gives results for two simplest observables. The concluding section summarizes and discusses the main results. A considerable amount of mathematical details, alternative derivations, and checks are collected in three sizable appendices A–C.

2 Preliminaries

Here we collect definitions of the background power-law inflation spacetime, and the results for quantum scalar fields in power-law inflation that are used throughout.

2.1 Power-law inflation

The geometry of the D -dimensional spatially flat Friedman-Lemaître-Robertson-Walker (FLRW) spacetime in conformal coordinates is described by the line element,

$$ds^2 = a^2(\eta)(-d\eta^2 + d\vec{x}^2), \quad (2.1)$$

that defines the conformally flat metric $g_{\mu\nu} = a^2(\eta) \text{diag}(-1, 1, \dots, 1)$, where a is the scale factor that encodes the dynamics of the expansion. The rate of expansion is conveniently captured by the conformal Hubble rate,

$$\mathcal{H} = \frac{1}{a} \frac{da}{d\eta}, \quad (2.2)$$

which is related to the physical Hubble rate $H = \mathcal{H}/a$. The acceleration of the expansion is captured by the principal slow-roll parameter,

$$\epsilon = -\frac{1}{H^2} \frac{dH}{dt} = 1 - \frac{1}{\mathcal{H}^2} \frac{d\mathcal{H}}{d\eta}, \quad (2.3)$$

that is connected to the deceleration parameter $q = \epsilon - 1$. In our conventions the Riemann tensor is defined as $R^\alpha_{\mu\beta\nu} = \partial_\beta \Gamma^\alpha_{\mu\nu} - \partial_\nu \Gamma^\alpha_{\mu\beta} + \Gamma^\rho_{\mu\nu} \Gamma^\alpha_{\beta\rho} - \Gamma^\rho_{\mu\beta} \Gamma^\alpha_{\nu\rho}$, where the Christoffel symbol is $\Gamma^\alpha_{\mu\nu} = \frac{1}{2} g^{\alpha\beta} (\partial_\mu g_{\nu\beta} + \partial_\nu g_{\mu\beta} - \partial_\beta g_{\mu\nu})$. Thus, the curvature tensors in FLRW spacetime are,

$$R_{\mu\nu\rho\sigma} = 2H^2 g_{\mu[\rho} g_{\sigma]\nu} + 4\epsilon H^2 (a^2 \delta^0_{[\mu} g_{\nu][\sigma} \delta^0_{\rho]}), \quad (2.4)$$

$$R_{\mu\nu} \equiv R^\alpha_{\mu\alpha\nu} = (D-1-\epsilon)H^2 g_{\mu\nu} + (D-2)\epsilon H^2 (a^2 \delta^0_\mu \delta^0_\nu), \quad (2.5)$$

$$R \equiv g^{\mu\nu} R_{\mu\nu} = (D-1)(D-2\epsilon)H^2. \quad (2.6)$$

If the principal slow-roll parameter is smaller than one, $\epsilon < 1$, the expansion is accelerating, which is the case in primordial slow-roll inflation where $0 < \epsilon \ll 1$. The special case of inflation is the power-law inflation [59, 60] characterized by a constant principal slow-roll parameter,

$$\epsilon = \text{const.} \ll 1 \quad \implies \quad \mathcal{H} = \frac{H_0}{1 - (1-\epsilon)H_0(\eta - \eta_0)} = H_0 a^{1-\epsilon}, \quad a(\eta) = \left(\frac{\mathcal{H}}{H_0}\right)^{\frac{1}{1-\epsilon}}, \quad (2.7)$$

where η_0 is the initial time at which $a(\eta_0) = 1$ and $\mathcal{H}(\eta_0) = H(\eta_0) = H_0$. The conformal time coordinate then ranges on an interval $\eta \in (-\infty, \bar{\eta})$, where $\bar{\eta} = \eta_0 + 1/[(1-\epsilon)H_0]$, while spatial coordinates cover $(D-1)$ -dimensional Euclidean space. The special case of power-law inflation for $\epsilon = 0$ is the de Sitter space, that has a constant physical Hubble rate $H = H_0$.

When expressing bi-local quantities, dependent on two spacetime points x and x' , such as the photon two-point functions, it is useful to employ bi-local variables that respect cosmological symmetries. The convenient choice for these variables is,

$$y(x; x') = (1-\epsilon)^2 \mathcal{H} \mathcal{H}' \Delta x^2, \quad u(x; x') = (1-\epsilon) \ln(aa'), \quad v(x; x') = (1-\epsilon) \ln(a/a'), \quad (2.8)$$

where henceforth primed quantities always refer to the primed coordinate, and where $\Delta x^2 = \|\vec{x} - \vec{x}'\|^2 - (\eta - \eta')^2$ is the Lorentz invariant distance between points. In order to represent the two-point functions as distributional limits of analytic bi-local functions, the bi-local variables above will acquire infinitesimal imaginary parts. Different prescriptions for these imaginary parts are defined in the two following subsections.

2.2 Scalar mode functions

The equation of motion for scalar modes is ubiquitous in cosmology,

$$\left[\partial_0^2 + k^2 - \left(\lambda^2 - \frac{1}{4} \right) (1-\epsilon)^2 \mathcal{H}^2 \right] \mathcal{U}_\lambda(\eta, \vec{k}) = 0, \quad (2.9)$$

where \mathcal{U}_λ is the suitably rescaled mode function of a non-minimally coupled scalar field, and where λ is a constant that can be related to the non-minimal coupling. The general solution in power-law inflation, where $\epsilon = \text{const.}$, is a linear combination,

$$\mathcal{U}(\eta, \vec{k}) = \alpha(\vec{k}) U_\lambda(\eta, k) + \beta(\vec{k}) U_\lambda^*(\eta, k), \quad (2.10)$$

of the positive-frequency Chernikov-Tagirov-Bunch-Davies (CTBD) mode function [82, 83],

$$U_\lambda(\eta, k) = e^{\frac{i\pi}{4}(2\lambda+1)} e^{\frac{-ik}{(1-\epsilon)H_0}} \sqrt{\frac{\pi}{4(1-\epsilon)\mathcal{H}}} H_\lambda^{(1)}\left(\frac{k}{(1-\epsilon)\mathcal{H}}\right), \quad (2.11)$$

and its complex conjugate, where $H_\lambda^{(1)}$ is the Hankel function of the first kind. The Wronskian for these two linearly independent solutions is,

$$\text{Im}\left[U_\lambda(\eta, k) \partial_0 U_\lambda^*(\eta, k) \right] = \frac{1}{2}. \quad (2.12)$$

In flat space the CTBD mode function reduces to,

$$U_\lambda(\eta, k) \stackrel{H_0 \rightarrow 0}{\sim} \frac{e^{-ik(\eta-\eta_0)}}{\sqrt{2k}} \left\{ 1 + \frac{i}{2} \left(\lambda^2 - \frac{1}{4} \right) \frac{(1-\epsilon)H_0}{k} - \frac{1}{8} \left(\lambda^2 - \frac{1}{4} \right) \left[\lambda^2 - \frac{9}{4} - 4ik(\eta-\eta_0) \right] \frac{(1-\epsilon)^2 H_0^2}{k^2} + \dots \right\}, \quad (2.13)$$

which is closely related to the UV limit $k/\mathcal{H} \gg 1$. On the other hand the IR expansion, i.e. the small momentum expansion $k/\mathcal{H} \ll 1$ of the CTBD mode function is

$$U_\lambda(\eta, k) \stackrel{k/\mathcal{H} \ll 1}{\sim} e^{\frac{i\pi}{4}(2\lambda+1)} e^{\frac{-ik}{(1-\epsilon)H_0}} \frac{\Gamma(\lambda) \Gamma(1-\lambda)}{\sqrt{4\pi(1-\epsilon)\mathcal{H}}} \left\{ \sum_{n=0}^{\infty} \frac{i(-1)^{n+1}}{n! \Gamma(n+1-\lambda)} \left[\frac{k}{2(1-\epsilon)\mathcal{H}} \right]^{2n-\lambda} - e^{-i\lambda\pi} \sum_{n=0}^{\infty} \frac{i(-1)^{n+1}}{n! \Gamma(n+1+\lambda)} \left[\frac{k}{2(1-\epsilon)\mathcal{H}} \right]^{2n+\lambda} \right\}. \quad (2.14)$$

We make frequent use of recurrence relations between contiguous scalar mode functions (scalar mode functions whose indices differ by one),

$$\left[\partial_0 + \left(\frac{1}{2} + \lambda \right) (1 - \epsilon) \mathcal{H} \right] U_\lambda = -ik U_{\lambda+1}, \quad \left[\partial_0 + \left(\frac{1}{2} - \lambda \right) (1 - \epsilon) \mathcal{H} \right] U_\lambda = -ik U_{\lambda-1}, \quad (2.15)$$

which follow from recurrence relations for Hankel functions (c.f. 10.6.2 in [84, 85]). These allow to express the Wronskian (2.12) as,

$$\text{Re} \left[U_\lambda(\eta, k) U_{\lambda+1}^*(\eta, k) \right] = \frac{1}{2k}. \quad (2.16)$$

We will also use the following identity,

$$\begin{aligned} & \left[\partial_0^2 + k^2 - \left(\lambda^2 - \frac{1}{4} \right) (1 - \epsilon)^2 \mathcal{H}^2 \right] \left(\frac{a^2}{\mathcal{H}} U_{\lambda+1} \right) \\ & = 2a^2 \mathcal{H} \left[2[\lambda(1 - \epsilon) + 1] U_{\lambda+1} - (1 + \epsilon) \frac{ik}{\mathcal{H}} U_\lambda \right], \end{aligned} \quad (2.17)$$

that follows from the equation of motion (2.9) and the recurrence relations in (2.15).

2.3 Scalar two-point functions

The positive-frequency Wightman two-point function for the scalar field in power-law inflation can be constructed as a sum-over-modes over its mode functions that satisfy the equation of motion (2.9),

$$i[-\Delta^+]_\lambda(x; x') = (aa')^{-\frac{D-2}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k} \cdot (\vec{x} - \vec{x}')} \mathcal{U}_\lambda(\eta, k) \mathcal{U}_\lambda^*(\eta', k), \quad (2.18)$$

where different states are captured by different Bogolyubov coefficients in mode functions (2.10). Implicit in this construction is the analytic continuation $\eta \rightarrow \eta - i\delta/2$ and $\eta' \rightarrow \eta' + i\delta/2$ (applied after the complex conjugation of the mode function), that preserves the property under complex conjugation $\{i[-\Delta^+]_\lambda(x; x')\}^* = i[-\Delta^+]_\lambda(x'; x)$. Nonequilibrium quantum field theory [86–92] requires the use of three other two-point functions (see [93, 94] for an introduction). The complex conjugate of (2.18) is the negative-frequency Wightman function, $i[+\Delta^-]_\lambda(x; x') = \{i[-\Delta^+]_\lambda(x; x')\}^*$. The Feynman propagator is then constructed from the two Wightman functions,

$$i[+\Delta^+]_\lambda(x; x') = \theta(\eta - \eta') i[-\Delta^+]_\lambda(x; x') + \theta(\eta' - \eta) i[+\Delta^-]_\lambda(x; x'), \quad (2.19)$$

and its complex conjugate is the Dyson propagator, $i[-\Delta^-]_\lambda(x; x') = \{i[+\Delta^+]_\lambda(x; x')\}^*$. It then follows, from the definitions and from the mode equation (2.9), that the two-point functions satisfy the following equation of motion,

$$\left[\square - (1 - \epsilon)^2 H^2 \left(\left[\frac{D-1-\epsilon}{2(1-\epsilon)} \right]^2 - \lambda^2 \right) \right] i[{}^{\text{a}}\Delta^{\text{b}}]_\lambda(x; x') = \text{S}^{\text{ab}} \frac{i\delta^D(x-x')}{\sqrt{-g}}, \quad (2.20)$$

where the scalar d'Alembertian is given by $\square = \nabla^\mu \nabla_\mu = -a^{-2} [\partial_0^2 + (D-2)\mathcal{H}\partial_0 - \nabla^2]$, and $\nabla^2 = \partial_i \partial_i$ is the Laplacian. Henceforth, the typewriter font indices denote the Schwinger-Keldysh polarity indices $\mathbf{a}, \mathbf{b} = \pm$, and we introduce an accompanying sign symbol $\mathbf{S}^{\mathbf{ab}}$ defined by,

$$\mathbf{S}^{++} = -\mathbf{S}^{--} = 1, \quad \mathbf{S}^{-+} = \mathbf{S}^{+-} = 0. \quad (2.21)$$

Note that, in addition to the equation of motion (2.20) with respect to the unprimed coordinate x , we have another independent equation with respect to the primed coordinate x' ,

$$\left[\square' - (1-\epsilon)^2 H'^2 \left(\left[\frac{D-1-\epsilon}{2(1-\epsilon)} \right]^2 - \lambda^2 \right) \right] i[\mathbf{a}\Delta^{\mathbf{b}}]_\lambda(x; x') = \mathbf{S}^{\mathbf{ab}} \frac{i\delta^D(x-x')}{\sqrt{-g}}. \quad (2.22)$$

In general the scalar two-point functions that respect cosmological symmetries of spatial homogeneity and isotropy can be expressed in terms of bi-local variables (2.8) with appropriate infinitesimal complex parts $\propto i\delta$ appended. This allows the two-point functions to be expressed as distributional limits of analytic functions. The complex parts appended to bi-local variables depend on the type of the two-point function, and different prescriptions are labeled by the Schwinger-Keldysh polarity indices. For the first variable,

$$y_{\mathbf{ab}} = (1-\epsilon)^2 \mathcal{H}\mathcal{H}' \Delta x_{\mathbf{ab}}^2, \quad (2.23)$$

the $i\delta$ is in the Lorentz invariant distance,²

$$\Delta x_{-+}^2 = \|\Delta \vec{x}\|^2 - (\Delta\eta - i\delta)^2, \quad \Delta x_{++}^2 = \|\Delta \vec{x}\|^2 - (|\Delta\eta| - i\delta)^2, \quad (2.24)$$

and $\Delta x_{+-}^2 = (\Delta x_{-+}^2)^*$, and $\Delta x_{--}^2 = (\Delta x_{++}^2)^*$. The complexified second variable is given by,

$$u_{-+} = u - \frac{1}{2}(1-\epsilon)(\mathcal{H} - \mathcal{H}')i\delta, \quad u_{++} = u - \frac{1}{2}(1-\epsilon)|\mathcal{H} - \mathcal{H}'|i\delta, \quad (2.25)$$

and $u_{+-} = (u_{-+})^*$, $u_{--} = (u_{++})^*$, while the third is given by,

$$v_{-+} = v - \frac{1}{2}(1-\epsilon)(\mathcal{H} + \mathcal{H}')i\delta \quad v_{++} = v - \frac{1}{2}(1-\epsilon)(\mathcal{H} + \mathcal{H}')\text{sgn}(\Delta\eta)i\delta, \quad (2.26)$$

and $v_{+-} = (v_{-+})^*$ and $v_{--} = (v_{++})^*$. For the sake of the flat space limit it is useful to define the $i\delta$ -prescription for the time difference as well,

$$\Delta\eta_{-+} = \Delta\eta - i\delta, \quad \Delta\eta_{++} = |\Delta\eta| - i\delta, \quad (2.27)$$

²The time dependent factor multiplying the Lorentz invariant distance in (2.23) strictly speaking also has complexified parts as does the u variable in (2.25), though it is irrelevant in all the expressions we consider and we thus omit it.

and the accompanying complex conjugates, $\Delta\eta_{+-} = (\Delta\eta_{-+})^*$ and $\Delta\eta_{--} = (\Delta\eta_{++})^*$. All of the prescriptions essentially follow from the one for the Wightman function (2.18).

The scalar two-point functions in power-law inflation can differ qualitatively depending on the index λ that they inherit from the mode equation (2.9). The preferred mode function is the CTBD one (2.11) which corresponds to the state that minimizes energy mode-per-mode in the asymptotic past. While this mode function is defined for any λ , it is clear from its IR behaviour (2.14) that for $\lambda \geq (D-1)/2$ this leads to an IR divergent sum-over-modes (2.18) [95–97], and that the Bogolyubov coefficients in the mode function (2.10) need to be chosen differently [95, 98]. We consider the two cases separately in the two following subsections.

2.3.1 IR finite scalar two-point functions

When the mode function index $\lambda < (D-1)/2$ there is no obstruction to choosing the mode function in the sum-over-modes (2.18) to be the CTBD one (2.11),

$$i[-\Delta^+]_{\lambda}(x; x') = (aa')^{-\frac{D-2}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot(\vec{x}-\vec{x}')} U_{\lambda}(\eta, k) U_{\lambda}^*(\eta', k). \quad (2.28)$$

This integral representation evaluates to [99, 102],

$$i[-\Delta^+]_{\lambda}(x; x') = i\Delta_{\lambda}(y_{-+}, u) = e^{-\frac{(D-2)\epsilon}{2(1-\epsilon)}u} \mathcal{F}_{\lambda}(y_{-+}), \quad (2.29)$$

where y_{-+} is the $i\delta$ -regulated distance function appropriate for the positive-frequency Wightman function, as introduced in (2.24) together with u , and the rescaled propagator function is expressed in terms of the hypergeometric function,

$$\begin{aligned} \mathcal{F}_{\lambda}(y) &= \frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\frac{D-1}{2} + \lambda) \Gamma(\frac{D-1}{2} - \lambda)}{(4\pi)^{\frac{D}{2}} \Gamma(\frac{D}{2})} \\ &\quad \times {}_2F_1\left(\left\{\frac{D-1}{2} + \lambda, \frac{D-1}{2} - \lambda\right\}, \left\{\frac{D}{2}\right\}, 1 - \frac{y}{4}\right), \end{aligned} \quad (2.30)$$

that satisfies the hypergeometric equation in disguise,

$$\left[(4y - y^2) \frac{\partial^2}{\partial y^2} + D(2 - y) \frac{\partial}{\partial y} + \lambda^2 - \left(\frac{D-1}{2}\right)^2 \right] \mathcal{F}_{\lambda}(y) = 0. \quad (2.31)$$

The function in (2.30) admits a power series representation around $y=0$,

$$\begin{aligned} \mathcal{F}_{\lambda}(y) &= \frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\frac{D-2}{2})}{(4\pi)^{\frac{D}{2}}} \left\{ \left(\frac{y}{4}\right)^{-\frac{D-2}{2}} + \frac{\Gamma(\frac{4-D}{2})}{\Gamma(\frac{1}{2} + \lambda) \Gamma(\frac{1}{2} - \lambda)} \sum_{n=0}^{\infty} \right. \\ &\quad \left. \times \left[\frac{\Gamma(\frac{3}{2} + \lambda + n) \Gamma(\frac{3}{2} - \lambda + n)}{\Gamma(\frac{6-D}{2} + n) (n+1)!} \left(\frac{y}{4}\right)^{n - \frac{D-4}{2}} - \frac{\Gamma(\frac{D-1}{2} + \lambda + n) \Gamma(\frac{D-1}{2} - \lambda + n)}{\Gamma(\frac{D}{2} + n) n!} \left(\frac{y}{4}\right)^n \right] \right\}, \end{aligned} \quad (2.32)$$

that we will make use of. We will also use recurrence relations between propagator functions with contiguous indices,

$$2\frac{\partial\mathcal{F}_\lambda}{\partial y} = (2-y)\frac{\partial\mathcal{F}_{\lambda+1}}{\partial y} - \left(\frac{D-3}{2} - \lambda\right)\mathcal{F}_{\lambda+1}, \quad (2.33)$$

$$2\frac{\partial\mathcal{F}_\lambda}{\partial y} = (2-y)\frac{\partial\mathcal{F}_{\lambda-1}}{\partial y} - \left(\frac{D-3}{2} + \lambda\right)\mathcal{F}_{\lambda-1}. \quad (2.34)$$

that follow from Gauss' recursion relations for hypergeometric functions (see (9.137) in [100]).

The solution for the scalar Feynman propagator, that is expressed in terms of the Wightman function as in (2.19) is then given by,

$$i[{}^{+\Delta^+}]_\lambda(x; x') = i\Delta_\lambda(y_{++}, u) = e^{-\frac{(D-2)\epsilon}{2(1-\epsilon)}u}\mathcal{F}_\lambda(y_{++}), \quad (2.35)$$

where y_{++} is complexified according to the prescription in (2.24). The remaining two-point functions are obtained from the definitions, which in the end amounts to simply changing the polarity indices on the y variable, denoting the appropriate prescription. Henceforth we suppress denoting explicitly the Schwinger-Keldysh polarity indices on variables, as they are either defined by the corresponding indices of the object being computed, or otherwise should be clear from the context. The expressions without any indices are meant to be valid in general.

2.3.2 IR divergent scalar two-point functions

When the index of the CTBD mode function is $\lambda \geq (D-1)/2$ the integral in (2.28) is divergent in the IR [95, 98]. This implies that (2.11) is not a legitimate choice for the mode functions in the IR sector. Rather, one should resort to choosing Bogolyubov coefficients in (2.10) such that the sum-over-modes representation (2.18) of the two-point function is well defined and convergent in the IR. This is guaranteed by fixing initial conditions at some time η_0 ,

$$\mathcal{U}_\lambda(\eta_0, k) = \bar{U}(k), \quad \partial_0\mathcal{U}_\lambda(\eta_0, k) = \partial_0\bar{U}(k), \quad (2.36)$$

such that $\bar{U}(k) \stackrel{k \rightarrow 0}{\propto} k^p$, where $p \geq -(D-1)/2$. This in turn determines the Bogolyubov coefficients in (2.10),

$$\alpha(k) = kU_{\lambda+1}^*(\eta_0, k)\bar{U}(k) + iU_\lambda^*(\eta_0, k)\left[\partial_0\bar{U}(k) + \left(\lambda + \frac{1}{2}\right)(1-\epsilon)H_0\bar{U}(k)\right], \quad (2.37a)$$

$$\beta(k) = kU_{\lambda+1}(\eta_0, k)\bar{U}(k) - iU_\lambda(\eta_0, k)\left[\partial_0\bar{U}(k) + \left(\lambda + \frac{1}{2}\right)(1-\epsilon)H_0\bar{U}(k)\right]. \quad (2.37b)$$

The evolution cannot generate IR divergences in (2.18) once they are absent at the initial time [95]. There are many possible choices for defining an IR finite state,

but since we are interested in studying loop corrections due to interactions, the appropriate choice for Bogolyubov coefficients in (2.10) is the one that minimizes the sensitivity to the details of regulating the IR. This we accomplish by assuming the full mode function to be the CTBD one all the way down to some deep IR scale $k_0 \ll H_0$,

$$k > k_0 \quad : \quad \bar{U}(k) = U_\lambda(\eta_0, k), \quad \text{and} \quad \partial_0 \bar{U}(k) = \partial_0 U_\lambda(\eta_0, k), \quad (2.38)$$

while the suppression for long wavelength modes $k < k_0$ is implemented via Bogolyubov coefficients (2.37) in some reasonable way. But given that $k_0/H_0 \ll 1$ we can neglect all positive powers of k_0 that appear in the two-point function, and the modes below k_0 can generate only such contributions. In accelerating spaces this approximation only becomes better at later times, as the infrared scale always appears as a decaying ratio k_0/\mathcal{H} . Thus, keeping only the negative powers of the deep IR scale k_0 , the two-point function is very well approximated by effectively introducing an IR cutoff k_0 into the integral (2.28),

$$i[-\Delta^+]_\lambda(x; x') = (aa')^{-\frac{D-2}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot(\vec{x}-\vec{x}')} \theta(k-k_0) U_\lambda(\eta, k) U_\lambda^*(\eta', k). \quad (2.39)$$

The two-point function can now be computed analytically by splitting it into two parts, according to the split of the step function $\theta(k-k_0) = 1 - \theta(k_0-k)$, and compute each part separately. However, since now each of the two parts will be given by an integral with zero as the lower limit of integration, both integrals will be IR divergent on their own. Despite this, their sum is finite because they come from splitting a manifestly finite quantity (2.39). This is why we are allowed to compute each individual part by regulating the IR by dimensional regularization, and after adding them together we are guaranteed to obtain a result that is correct as the two errors cancel each other out.³ We denote the split as,

$$i[-\Delta^+]_\lambda(x; x') = e^{-\frac{(D-2)\epsilon}{2(1-\epsilon)}u} \left[\mathcal{F}_\lambda(y) + \mathcal{W}_\lambda(y, u, v) \right] = i\Delta_\lambda(y, u, v). \quad (2.40)$$

where the first (bulk) part corresponds to the rescaled propagator function (2.30), and the IR part is given by,

$$\mathcal{W}_\lambda(y, u, v) = -(aa')^{-\frac{(D-2)(1-\epsilon)}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot(\vec{x}-\vec{x}')} \theta(k_0-k) U_\lambda(\eta, k) U_\lambda^*(\eta', k). \quad (2.41)$$

We evaluate this integral by only keeping terms that can potentially contribute negative powers of the effective IR cutoff k_0 .

³Dimensional regularization automatically subtracts any power-law divergences, whether UV or IR. While for the UV this is a convenient property, as it reduces the labor needed to renormalize a theory, it should not be employed to subtract divergences in the IR, as they are not universal but rather are symptomatic of unphysical properties of a chosen state, and need to be approached with due care [98].

To evaluate the IR part we first integrate over the angular coordinates. This is accomplished as in [102], by first writing out the volume element in angular coordinates, $d^{D-1}k = k^{D-2}dk d\Omega_{D-2}$, where the surface element of the $(D-2)$ -sphere, $d\Omega_{D-2} = \sin^{D-3}(\theta_{D-3})d\theta_{D-3} \sin^{D-4}(\theta_{D-4})d\theta_{D-4} \dots d\varphi$, is parametrized by $(D-2)$ angles $\theta_{D-3}, \theta_{D-4}, \dots, \varphi$. Then using the expression for the surface of the $(D-3)$ -sphere, and the representation of the Bessel function 8.411.7 from [100],

$$\int d\Omega_{D-3} = \frac{2\pi^{\frac{D-2}{2}}}{\Gamma(\frac{D-2}{2})}, \quad J_\lambda(z) = \frac{\left(\frac{z}{2}\right)^\lambda}{\sqrt{\pi}\Gamma(\frac{1}{2}+\lambda)} \int_0^\pi d\theta e^{\pm iz \cos(\theta)} \sin^{2\lambda}(\theta), \quad (2.42)$$

we arrive at the one-dimensional integral expression,

$$\mathcal{W}_\lambda = -\frac{(aa')^{-\frac{(D-2)(1-\epsilon)}{2}}}{(2\pi)^{\frac{D-1}{2}} \|\Delta\vec{x}\|^{\frac{D-3}{2}}} \int_0^{k_0} dk k^{\frac{D-1}{2}} J_{\frac{D-3}{2}}(k\|\Delta\vec{x}\|) U_\lambda(\eta, k) U_\lambda^*(\eta', k). \quad (2.43)$$

Next we expand the mode functions in the small argument limit, keeping only terms potentially generating negative powers of k_0 , and collecting the same powers of k ,

$$\begin{aligned} \mathcal{W}_\lambda = & -\frac{(aa')^{-\frac{(D-2)(1-\epsilon)}{2}} 2^{2\lambda-1} \Gamma^2(\lambda) \Gamma^2(1-\lambda)}{(2\pi)^{\frac{D+1}{2}} (1-\epsilon) \sqrt{\mathcal{H}\mathcal{H}'} \|\Delta\vec{x}\|^{\frac{D-3}{2}}} \\ & \times \sum_{N=0}^{\infty} \sum_{n=0}^N \frac{\left(-\frac{1}{4}\right)^N}{n!(N-n)! \Gamma(n+1-\lambda) \Gamma(N-n+1-\lambda)} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{N-2n} \\ & \times \int_0^{k_0} dk k^{\frac{D-1}{2}} J_{\frac{D-3}{2}}(k\|\Delta\vec{x}\|) \left[\frac{k^2}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'}\right]^{N-\lambda}. \end{aligned} \quad (2.44)$$

The integral over a single Bessel function and a power can be found in 1.8.1.1. of [101],

$$\int_0^z dz' z'^\rho J_\sigma(z') = \frac{z^{\rho+\sigma+1}}{2^\sigma(\rho+\sigma+1)\Gamma(\sigma+1)} {}_1F_2\left(\left\{\frac{\rho+\sigma+1}{2}\right\}, \left\{\frac{\rho+\sigma+3}{2}, \sigma+1\right\}, -\frac{z^2}{4}\right), \quad (2.45)$$

so that after some rearrangements, and after applying the Legendre duplication formula for the gamma function in the overall factor, the IR series takes the form,

$$\begin{aligned} \mathcal{W}_\lambda = & -\frac{[(1-\epsilon)H_0]^{D-2}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(\lambda)\Gamma(2\lambda)}{\Gamma(\frac{1}{2}+\lambda)\Gamma(\frac{D-1}{2})} \sum_{N=0}^{\infty} \sum_{n=0}^N \frac{1}{\left(\frac{D-1}{2}+N-\lambda\right)} \\ & \times \frac{\left(-\frac{1}{4}\right)^N \Gamma^2(1-\lambda)}{n!(N-n)! \Gamma(n+1-\lambda) \Gamma(N-n+1-\lambda)} \left[\frac{k_0^2}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'}\right]^{\frac{D-1}{2}+N-\lambda} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{N-2n} \\ & \times {}_1F_2\left(\left\{\frac{D-1}{2}+N-\lambda\right\}, \left\{\frac{D+1}{2}+N-\lambda, \frac{D-1}{2}\right\}, -\frac{(k_0\|\Delta\vec{x}\|)^2}{4}\right). \end{aligned} \quad (2.46)$$

Finally, restricting to the domain where $k_0\|\Delta\vec{x}\|\ll 1$, expanding the hypergeometric function above in a power series, and rearranging the triple series gives,

$$\begin{aligned} \mathcal{W}_\lambda(y, u, v) = & -\frac{[(1-\epsilon)H_0]^{D-2}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(\lambda)\Gamma(2\lambda)}{\Gamma(\frac{1}{2}+\lambda)\Gamma(\frac{D-1}{2})} \sum_{N=0}^{\lfloor\lambda-\frac{D-1}{2}\rfloor} \sum_{n=0}^N \sum_{\ell=0}^{N-n} \frac{c_{Nn\ell}}{(\frac{D-1}{2}+N-\lambda)} \\ & \times \left[\frac{k_0^2 e^{-u}}{(1-\epsilon)^2 H_0^2} \right]^{\frac{D-1}{2}-\lambda+N} \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right]^n \operatorname{ch}[(N-n-2\ell)v], \end{aligned} \quad (2.47)$$

where the coefficients are,

$$c_{Nn\ell} = \frac{\left(-\frac{1}{4}\right)^N}{n! \ell! (N-n-\ell)!} \frac{\Gamma^2(1-\lambda)\Gamma(\frac{D-1}{2})}{\Gamma(\frac{D-1}{2}+n)\Gamma(\ell+1-\lambda)\Gamma(N-n-\ell+1-\lambda)}, \quad (2.48)$$

and where the sum over N is cut by the floor function in the upper limit to $N \leq \lambda - (D-1)/2$, which guarantees that only the negative powers of k_0 are kept, while positive ones are neglected. This corresponds to the result obtained in [102], where the infrared regulation was implemented by considering the spatial sections of the spacetime to be flat torii with a large radius. Here, however, it is advantageous to consider the IR regulation to be implemented by the Bogolyubov coefficients defining the initial state, in order to avoid having to deal with linearization instability [103–113] that appears in gauge theories on compact manifolds. The fact that the two expressions for the two-point function agree regardless of the difference in the method for regulating the IR is due to the expressions in actuality being valid only for spatial separations $k_0\|\Delta\vec{x}\|\ll 1$. However, since $k_0 \ll H_0$ is chosen to be smaller than any physical scale of interest, that limitation is of no real significance.

Writing out the equation of motion (2.20) off-coincidence in terms of the bi-local variables reads,

$$\begin{aligned} & \left[(4y-y^2) \frac{\partial^2}{\partial y^2} + D(2-y) \frac{\partial}{\partial y} + 2(2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{(D-2)\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right. \\ & \left. - \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{D-1-\epsilon}{1-\epsilon} \right) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} \right) + \lambda^2 - \left(\frac{D-1-\epsilon}{2(1-\epsilon)} \right)^2 \right] i\Delta_\lambda(y, u, v) = 0. \end{aligned} \quad (2.49)$$

This is a generalization of Eq. (2.31) to the case when the CTBD mode function cannot be used due to an IR divergence. In this case there is another independent equation descending from the primed equation of motion (2.22), that is obtained by substituting $v \rightarrow -v$ everywhere inside the square brackets in (2.49). The two equations can be combined into another two equations of well defined parity that

have to be satisfied independently. Namely, the even equation is,

$$\begin{aligned} & \left[(4y-y^2) \frac{\partial^2}{\partial y^2} + D(2-y) \frac{\partial}{\partial y} - 2 \left[y + 4 \operatorname{sh}^2 \left(\frac{v}{2} \right) \right] \left(\frac{\partial}{\partial u} + \frac{(D-2)\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right. \\ & \left. - \left(\frac{\partial}{\partial u} + \frac{D-1-\epsilon}{1-\epsilon} \right) \frac{\partial}{\partial u} + \left(4 \operatorname{sh}(v) \frac{\partial}{\partial y} - \frac{\partial}{\partial v} \right) \frac{\partial}{\partial v} + \lambda^2 - \left(\frac{D-1-\epsilon}{2(1-\epsilon)} \right)^2 \right] i\Delta_\lambda(y, u, v) = 0, \end{aligned} \quad (2.50)$$

and the odd equation reads,

$$\begin{aligned} & \left[\left(\left[y + 4 \operatorname{sh}^2 \left(\frac{v}{2} \right) \right] \frac{\partial}{\partial y} + \frac{\partial}{\partial u} + \frac{D-1-\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial v} \right. \\ & \left. - 2 \operatorname{sh}(v) \left(\frac{\partial}{\partial u} + \frac{(D-2)\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right] i\Delta_\lambda(y, u, v) = 0. \end{aligned} \quad (2.51)$$

We can also derive generalizations of recurrence relations for rescaled propagator functions (2.33) and (2.34) that apply for the full scalar two-point functions. These are derived from recurrence relations between mode functions (2.15) that imply the following two reflection identities for time derivatives,

$$\begin{aligned} & \left[\partial_0 - \left(\lambda - \frac{D-3+\epsilon}{2(1-\epsilon)} \right) (1-\epsilon) \mathcal{H} \right] i[{}^{\text{a}}\Delta^{\text{b}}]_{\lambda+1}(x; x') \\ & = - \left[\partial'_0 + \left(\lambda + 1 + \frac{D-3+\epsilon}{2(1-\epsilon)} \right) (1-\epsilon) \mathcal{H}' \right] i[{}^{\text{a}}\Delta^{\text{b}}]_\lambda(x; x'), \end{aligned} \quad (2.52\text{a})$$

$$\begin{aligned} & \left[\partial'_0 - \left(\lambda - \frac{D-3+\epsilon}{2(1-\epsilon)} \right) (1-\epsilon) \mathcal{H}' \right] i[{}^{\text{a}}\Delta^{\text{b}}]_{\lambda+1}(x; x') \\ & = - \left[\partial_0 + \left(\lambda + 1 + \frac{D-3+\epsilon}{2(1-\epsilon)} \right) (1-\epsilon) \mathcal{H} \right] i[{}^{\text{a}}\Delta^{\text{b}}]_\lambda(x; x'). \end{aligned} \quad (2.52\text{b})$$

Then writing these out in terms of bi-local variables yields the desired generalized recurrence relations,

$$\begin{aligned} & \left[2 \frac{\partial}{\partial y} - \frac{1}{\operatorname{sh}(v)} \frac{\partial}{\partial v} \right] i\Delta_\lambda(y, u, v) \\ & = \left[(2-y) \frac{\partial}{\partial y} - \left(\frac{D-3+\epsilon}{2(1-\epsilon)} - \lambda \right) - \frac{\partial}{\partial u} - \frac{\operatorname{ch}(v)}{\operatorname{sh}(v)} \frac{\partial}{\partial v} \right] i\Delta_{\lambda+1}(y, u, v), \end{aligned} \quad (2.53\text{a})$$

$$\begin{aligned} & \left[2 \frac{\partial}{\partial y} - \frac{1}{\operatorname{sh}(v)} \frac{\partial}{\partial v} \right] i\Delta_\lambda(y, u, v) \\ & = \left[(2-y) \frac{\partial}{\partial y} - \left(\frac{D-3+\epsilon}{2(1-\epsilon)} + \lambda \right) - \frac{\partial}{\partial u} - \frac{\operatorname{ch}(v)}{\operatorname{sh}(v)} \frac{\partial}{\partial v} \right] i\Delta_{\lambda-1}(y, u, v), \end{aligned} \quad (2.53\text{b})$$

that we make frequent use of in the remainder of the paper.

3 Photon in FLRW

This section is devoted to a brief recap of the procedure for imposing the multiplier gauge for electromagnetism that corresponds to the general covariant gauge-fixing term (1.1), and to the canonical quantization of the resulting gauge-fixed theory. The details of this procedure, applicable to more general gauges and general FLRW spacetimes, are given in [79].

3.1 Classical photon in FLRW

Electromagnetism in D -dimensional curved spacetime is given by the covariantized Maxwell action,

$$S[A_\mu] = \int d^D x \sqrt{-g} \left[-\frac{1}{4} g^{\mu\rho} g^{\nu\sigma} F_{\mu\nu} F_{\rho\sigma} \right], \quad (3.1)$$

that is invariant under $U(1)$ gauge transformations, $A_\mu(x) \rightarrow A_\mu(x) + \partial_\mu \Lambda(x)$. In FLRW cosmological spaces the corresponding canonical formulation of electromagnetism is given by the canonical action [79],

$$\mathcal{S}[A_0, \Pi_0, A_i, \Pi_i, \ell] = \int d^D x \left[\Pi_0 \partial A_0 + \Pi_i \partial_0 A_i - \mathcal{H} - \ell \Psi_1 \right], \quad (3.2)$$

where (A_0, Π_0) and (A_i, Π_i) are canonical pairs of vector potentials and their conjugate momenta, whose Poisson brackets are determined by the symplectic part of the action, with the non-vanishing ones being,

$$\{A_0(\eta, \vec{x}), \Pi_0(\eta, \vec{x}')\} = \delta^{D-1}(\vec{x} - \vec{x}'), \quad \{A_i(\eta, \vec{x}), \Pi_j(\eta, \vec{x}')\} = \delta_{ij} \delta^{D-1}(\vec{x} - \vec{x}'). \quad (3.3)$$

The canonical Hamiltonian is given by,

$$\mathcal{H} = \frac{a^{4-D}}{2} \Pi_i \Pi_i - A_0 \partial_i \Pi_i + \frac{a^{D-4}}{4} F_{ij} F_{ij}, \quad (3.4)$$

where repeated lower spatial indices are summed over, and ℓ is a Lagrange multiplier that generates the primary constraint,

$$\Psi_1 = \Pi_0 \approx 0, \quad (3.5)$$

that vanishes on-shell, and whose conservation in turn generates a secondary constraint,

$$0 \approx \partial_0 \Psi_1 \approx \Psi_2 = \partial_i \Pi_i. \quad (3.6)$$

that also vanishes on-shell. The conservation of the secondary constraint generates no further constraints, and thus the two form a complete set of first-class constraints, $\{\Psi_1, \Psi_2\} = 0$. The fact that these two formulations — configuration space one given by (3.1) and the canonical one given by (3.2) — are equivalent can be confirmed by checking that the two generate equivalent sets of equations of motion.

The multiplier gauge is imposed by promoting the Lagrange multiplier in the canonical action (3.2) to a function of canonical variables, $\ell \rightarrow \bar{\ell}(A_0, \Pi_0, A_i, \Pi_i)$. This way of fixing the gauge does not allow to reduce the phase space (such as the Coulomb gauge would), but rather we work with all the components of the vector potential. On the other hand, multiplier gauges are much more conducive to preserving the symmetries of the gauge-invariant action. The particular choice for the multiplier that we consider here,

$$\ell \longrightarrow \bar{\ell} = -\frac{\xi}{2}a^{4-D}\Pi_0 + \partial_i A_i - (D-2)\mathcal{H}A_0. \quad (3.7)$$

is precisely motivated by this observation. This choice defines the gauge-fixed canonical action,

$$\mathcal{S}_\star[A_0, \Pi_0, A_i, \Pi_i] \equiv \mathcal{S}[A_0, \Pi_0, A_i, \Pi_i, \ell \rightarrow \bar{\ell}] = \int d^Dx \left[\Pi_0 \partial A_0 + \Pi_i \partial A_i - \mathcal{H}_\star \right], \quad (3.8)$$

with the gauge-fixed Hamiltonian,

$$\mathcal{H}_\star = \frac{a^{4-D}}{2}(\Pi_i \Pi_i - \xi \Pi_0 \Pi_0) + \Pi_i \partial_i A_0 + \Pi_0 \partial_i A_i - (D-2)\mathcal{H}\Pi_0 A_0 + \frac{a^{D-4}}{2}F_{ij}F_{ij}. \quad (3.9)$$

The the corresponding configuration space formulation of the gauge-fixed theory is consequently given by,

$$S_\star[A_\mu] = \int d^Dx \sqrt{-g} \left[-\frac{1}{4}g^{\mu\rho}g^{\nu\sigma}F_{\mu\nu}F_{\rho\sigma} - \frac{1}{2\xi}(\nabla^\mu A_\mu)^2 \right], \quad (3.10)$$

which is precisely the gauge-invariant action (3.1) with the general covariant gauge-fixing term (1.1) added. The gauge-fixed action now defines the gauge-fixed dynamics, in the sense that the ambiguity present in the original equations of motion are now fixed. For the gauge-fixed equations,

$$\partial_0 A_0 = -\xi a^{4-D}\Pi_0 + \partial_i A_i - (D-2)\mathcal{H}A_0, \quad (3.11a)$$

$$\partial_0 \Pi_0 = \partial_i \Pi_i + (D-2)\mathcal{H}\Pi_0, \quad (3.11b)$$

$$\partial_0 A_i = a^{4-D}\Pi_i + \partial_i A_0, \quad (3.11c)$$

$$\partial_0 \Pi_i = \partial_i \Pi_0 + a^{D-4}\partial_j F_{ji}, \quad (3.11d)$$

the initial value problem is well defined once the initial conditions are specified.

However, the gauge-fixed action is not a complete gauge-fixed description of the system, as it makes no reference to the first-class constraints (3.5) and (3.6) that are present in the original gauge-invariant system. In fact, the set of equations (3.11) admits many more solutions than the original gauge-invariant system of equations, regardless of the gauge freedom. That is why the two constraints (3.5) and (3.6)

have to be required to hold as subsidiary conditions on the initial value surface, i.e. on the initial conditions that are propagated by the evolution equations in (3.11),

$$\Psi_1(\eta_0, \vec{x}) = \Pi_0(\eta_0, \vec{x}) = 0, \quad \Psi_2(\eta_0, \vec{x}) = \partial_i \Pi_i(\eta_0, \vec{x}) = 0. \quad (3.12)$$

Equations of motion preserve these conditions so that they are valid for all times once imposed initially. Therefore, it is the gauge-fixed action (3.8) *and* the constraints (3.12) that define the gauge-fixed version of the theory.

3.2 Quantized photon in FLRW

The gauge-fixing procedure for the classical theory described in the preceding section lends itself well to canonical quantization in the Heisenberg picture. The gauge-fixed dynamics is quantized straightforwardly by promoting vector potential components and their conjugate momenta into field operators, $A_0 \rightarrow \hat{A}_0$, $\Pi_0 \rightarrow \hat{\Pi}_0$, $A_i \rightarrow \hat{A}_i$, $\Pi_i \rightarrow \hat{\Pi}_i$, and their nonvanishing Poisson brackets (3.3) to canonical commutation relations,

$$[\hat{A}_0(\eta, \vec{x}), \hat{\Pi}_0(\eta, \vec{x}')] = i\delta^{D-1}(\vec{x} - \vec{x}'), \quad [\hat{A}_i(\eta, \vec{x}), \hat{\Pi}_j(\eta, \vec{x}')] = \delta_{ij} i\delta^{D-1}(\vec{x} - \vec{x}'). \quad (3.13)$$

The equations of motion the field operators satisfy are just the same as the classical equations (3.11), with fields promoted to field operators. The algebra of field operators is assumed represented on some space of states. However, this space cannot be the usual Fock space because of the subsidiary conditions (3.12) that need to be quantized as well.

The generalization of the subsidiary condition to the quantized theory firstly requires that the expectation values of Hermitian operators associated to classical constraints, $\hat{\Psi}_1 = \hat{\Pi}_0$ and $\hat{\Psi}_2 = \partial_i \hat{\Pi}_i$, vanish at initial time on account of the correspondence principle,

$$\langle \Omega | \hat{\Psi}_1(\eta_0, \vec{x}) | \Omega \rangle = 0, \quad \langle \Omega | \hat{\Psi}_2(\eta_0, \vec{x}) | \Omega \rangle = 0. \quad (3.14)$$

Secondly, all the correlators of constraints have to likewise vanish, e.g. two-point correlators,

$$\langle \Omega | \hat{\Psi}_1(\eta_0, \vec{x}) \hat{\Psi}_1(\eta_0, \vec{x}') | \Omega \rangle = 0, \quad (3.15a)$$

$$\langle \Omega | \hat{\Psi}_1(\eta_0, \vec{x}) \hat{\Psi}_2(\eta_0, \vec{x}') | \Omega \rangle = 0, \quad (3.15b)$$

$$\langle \Omega | \hat{\Psi}_2(\eta_0, \vec{x}) \hat{\Psi}_2(\eta_0, \vec{x}') | \Omega \rangle = 0, \quad (3.15c)$$

which ensures no fluctuations of constraints can be measured. The evolution equations then ensure that these correlators then vanish for all times. The conditions in (3.14) and (3.15) are naturally implemented as conditions on the state vectors forming the space of states, in the form of an operator annihilating the state,

$$\hat{K}(\vec{x}) | \Omega \rangle = 0, \quad \forall \vec{x}. \quad (3.16)$$

Just as in the classical theory, the subsidiary conditions cut out a subspace of physically allowed initial conditions. In the quantized theory this condition is meant to cut out a subset of physically allowed states from the vector space of states. In order for the condition in (3.16) to be consistent with canonical commutation relations in (3.13) the operator \hat{K} cannot be Hermitian. Rather, it is some non-Hermitian linear combination of Hermitian constraint operators,

$$\hat{K}(\vec{x}) = \int d^{D-1}x' \left[f_1(\eta_0; \vec{x}, \vec{x}') \hat{\Psi}_1(\eta_0, \vec{x}) + f_2(\eta_0; \vec{x}, \vec{x}') \hat{\Psi}_2(\eta_0, \vec{x}) \right]. \quad (3.17)$$

In fact, the equations of motion guarantee that such a time-independent linear combination can be formed from Hermitian constraints at any point in time,

$$\hat{K}(\vec{x}) = \int d^{D-1}x' \left[f_1(\eta; \vec{x}, \vec{x}') \hat{\Psi}_1(\eta, \vec{x}) + f_2(\eta; \vec{x}, \vec{x}') \hat{\Psi}_2(\eta, \vec{x}) \right], \quad (3.18)$$

where the time-dependent coefficient functions satisfy their own equations of motion,

$$\partial_0 f_1 = -\nabla^2 f_2 - (D-2)\mathcal{H}f_1, \quad \partial_0 f_2 = -f_1, \quad (3.19)$$

There is a great deal of freedom in choosing initial conditions, and hence in choosing the subsidiary constraint operator, and particular choices can lead to more favourable expressions.

4 Field operator dynamics

The gauge-fixed dynamics of the linear theory is completely determined by solving the operator equations of motion, which are just the classical equations (3.11) with fields promoted to field operators, and by requiring the solutions to respect canonical commutation relations (3.13). This task is best accomplished if we first employ a couple of decompositions of the field operators, after which some of the equations decouple, and some can be brought into the form of the scalar mode equation (2.9). This section is devoted to introducing the necessary decompositions and to solving the resulting equations of motion.

Firstly, it is convenient to split the spatial components of field operators into transverse and longitudinal parts,

$$\hat{A}_i = \hat{A}_i^T + \hat{A}_i^L, \quad \hat{\Pi}_i = \hat{\Pi}_i^T + \hat{\Pi}_i^L, \quad (4.1)$$

such that $\partial_i \hat{A}_i^T = 0 = \partial_i \hat{\Pi}_i^T$. This is best accomplished with the use of transverse and longitudinal projectors,

$$\mathbb{P}_{ij}^T = \delta_{ij} - \frac{\partial_i \partial_j}{\nabla^2}, \quad \mathbb{P}_{ij}^L = \frac{\partial_i \partial_j}{\nabla^2}. \quad (4.2)$$

These projectors are orthogonal, $\mathbb{P}_{ij}^T \mathbb{P}_{jk}^L = \mathbb{P}_{ij}^L \mathbb{P}_{jk}^T = 0$, and idempotent, $\mathbb{P}_{ij}^T \mathbb{P}_{jk}^T = \mathbb{P}_{ik}^T$, and the field operator components are then projected out with correct properties, $\hat{A}_i^T = \mathbb{P}_{ij}^T \hat{A}_j$, $\hat{\Pi}_i^T = \mathbb{P}_{ij}^T \hat{\Pi}_j$, $\hat{A}_i^L = \mathbb{P}_{ij}^L \hat{A}_j$, $\hat{\Pi}_i^L = \mathbb{P}_{ij}^L \hat{\Pi}_j$. After this split the equations of motion for the transverse operators decouple from the rest.

Secondly, given the isotropy of spatially flat cosmological spaces, it is convenient to expand the field operators in Fourier modes of the comoving momentum space,

$$\hat{A}_0(\eta, \vec{x}) = a^{-\frac{D-2}{2}} \int \frac{d^{D-1}k}{(2\pi)^{\frac{D-1}{2}}} e^{i\vec{k}\cdot\vec{x}} \hat{\mathcal{A}}_0(\eta, \vec{k}), \quad (4.3a)$$

$$\hat{\Pi}_0(\eta, \vec{x}) = a^{\frac{D-2}{2}} \int \frac{d^{D-1}k}{(2\pi)^{\frac{D-1}{2}}} e^{i\vec{k}\cdot\vec{x}} \hat{\pi}_0(\eta, \vec{k}), \quad (4.3b)$$

$$\hat{A}_i^L(\eta, \vec{x}) = a^{-\frac{D-2}{2}} \int \frac{d^{D-1}k}{(2\pi)^{\frac{D-1}{2}}} e^{i\vec{k}\cdot\vec{x}} \frac{(-i)k_i}{k} \hat{\mathcal{A}}_L(\eta, \vec{k}), \quad (4.3c)$$

$$\hat{\Pi}_i^L(\eta, \vec{x}) = a^{\frac{D-2}{2}} \int \frac{d^{D-1}k}{(2\pi)^{\frac{D-1}{2}}} e^{i\vec{k}\cdot\vec{x}} \frac{(-i)k_i}{k} \hat{\pi}_L(\eta, \vec{k}), \quad (4.3d)$$

$$\hat{A}_i^T(\eta, \vec{x}) = a^{-\frac{D-4}{2}} \int \frac{d^{D-1}k}{(2\pi)^{\frac{D-1}{2}}} e^{i\vec{k}\cdot\vec{x}} \sum_{\sigma=1}^{D-2} \varepsilon_i^T(\sigma, \vec{k}) \hat{\mathcal{A}}_{T,\sigma}(\eta, \vec{k}), \quad (4.3e)$$

$$\hat{\Pi}_i^T(\eta, \vec{x}) = a^{\frac{D-4}{2}} \int \frac{d^{D-1}k}{(2\pi)^{\frac{D-1}{2}}} e^{i\vec{k}\cdot\vec{x}} \sum_{\sigma=1}^{D-2} \varepsilon_i^T(\sigma, \vec{k}) \hat{\pi}_{T,\sigma}(\eta, \vec{k}), \quad (4.3f)$$

where the powers of the scale factor have been factored out for later convenience. Here we introduced transverse polarization tensors with the following properties,

$$k_i \varepsilon_i(\sigma, \vec{k}) = 0, \quad [\varepsilon_i(\sigma, \vec{k})]^* = \varepsilon_i(\sigma, -\vec{k}), \quad (4.4)$$

$$\varepsilon_i^*(\sigma, \vec{k}) \varepsilon_i(\sigma', \vec{k}) = \delta_{\sigma\sigma'}, \quad \sum_{\sigma=1}^{D-2} \varepsilon_i^*(\sigma, \vec{k}) \varepsilon_j(\sigma, \vec{k}) = \delta_{ij} - \frac{k_i k_j}{k^2}, \quad (4.5)$$

where $k = \|\vec{k}\|$. The Fourier transforms of field operators in (4.3) are themselves Hermitian, which in momentum space implies they behave under conjugation as $\hat{\mathcal{O}}^\dagger(\vec{k}) = \hat{\mathcal{O}}(-\vec{k})$. The canonical commutators of the momentum space field operators are now,

$$[\hat{\mathcal{A}}_0(\eta, \vec{k}), \hat{\pi}_0(\eta, \vec{k}')] = [\hat{\mathcal{A}}_L(\eta, \vec{k}), \hat{\pi}_L(\eta, \vec{k}')] = i\delta^{D-1}(\vec{k} + \vec{k}'), \quad (4.6a)$$

$$[\hat{\mathcal{A}}_{T,\sigma}(\eta, \vec{k}), \hat{\pi}_{T,\sigma'}(\eta, \vec{k}')] = i\delta_{\sigma\sigma'} \delta^{D-1}(\vec{k} + \vec{k}'). \quad (4.6b)$$

The equations of motion for the transverse sector are

$$\partial_0 \hat{\mathcal{A}}_{T,\sigma} = \hat{\pi}_{T,\sigma} + \frac{1}{2}(D-4)\mathcal{H}\hat{\mathcal{A}}_{T,\sigma}, \quad (4.7)$$

$$\partial_0 \hat{\pi}_{T,\sigma} = -k^2 \hat{\mathcal{A}}_{T,\sigma} - \frac{1}{2}(D-4)\mathcal{H}\hat{\pi}_{T,\sigma}. \quad (4.8)$$

while the equations of motion for the scalar sector, including longitudinal and temporal components, are

$$\partial_0 \hat{\mathcal{A}}_0 = -\xi a^2 \hat{\pi}_0 + k \hat{\mathcal{A}}_L - \frac{1}{2}(D-2)\mathcal{H}\hat{\mathcal{A}}_0, \quad (4.9)$$

$$\partial_0 \hat{\pi}_0 = k \hat{\pi}_L + \frac{1}{2}(D-2)\mathcal{H}\hat{\pi}_0, \quad (4.10)$$

$$\partial_0 \hat{\mathcal{A}}_L = a^2 \hat{\pi}_L - k \hat{\mathcal{A}}_0 + \frac{1}{2}(D-2)\mathcal{H}\hat{\mathcal{A}}_L, \quad (4.11)$$

$$\partial_0 \hat{\pi}_L = -k \hat{\pi}_0 - \frac{1}{2}(D-2)\mathcal{H}\hat{\pi}_L. \quad (4.12)$$

4.1 Transverse sector

The equations of motion (4.7) and (4.8) of the transverse sector combine into a single second order equation,

$$\left[\partial_0^2 + k^2 - \left(\nu_T^2 - \frac{1}{4} \right) (1-\epsilon)^2 \mathcal{H}^2 \right] \hat{\mathcal{A}}_{T,\sigma} = 0, \quad (4.13)$$

$$\hat{\pi}_{T,\sigma} = \left[\partial_0 - \left(\nu_T + \frac{1}{2} \right) (1-\epsilon) \mathcal{H} \right] \hat{\mathcal{A}}_{T,\sigma}, \quad (4.14)$$

where we define the transverse sector index,

$$\nu_T = \frac{D-3-\epsilon}{2(1-\epsilon)}. \quad (4.15)$$

Equation (4.13) is easily recognized as the scalar mode equation (2.9) whose solution is then given by (2.10), upon which Eq. (4.14) can be recognized as the recurrence relation from (2.15). The solutions for transverse field operators are thus,

$$\hat{\mathcal{A}}_{T,\sigma}(\eta, k) = U_{\nu_T}(\eta, k) \hat{b}_T(\sigma, \vec{k}) + U_{\nu_T}^*(\eta, k) \hat{b}_T^\dagger(\sigma, -\vec{k}), \quad (4.16)$$

$$\hat{\pi}_{T,\sigma}(\eta, k) = -ik U_{\nu_T-1}(\eta, k) \hat{b}_T(\sigma, \vec{k}) + ik U_{\nu_T-1}^*(\eta, k) \hat{b}_T^\dagger(\sigma, -\vec{k}), \quad (4.17)$$

where the non-Hermitian operators $\hat{b}_T(\vec{k}, \sigma)$ can be seen as initial conditions. Their commutation relations follow from the canonical commutation relations (4.6). The non-vanishing ones are,

$$[\hat{b}_T(\sigma, \vec{k}), \hat{b}_T^\dagger(\sigma', \vec{k}')] = \delta_{\sigma\sigma'} \delta^{D-1}(\vec{k} - \vec{k}'), \quad (4.18)$$

which are just the commutation relations for creation and annihilation operators. Therefore the transverse sector of the space of states is naturally constructed as a Fock space. It is natural to consider the vacuum state of this sector to be the state that minimizes the energy mode-per-mode at an asymptotic past, by analogy with the CTBD state for the scalar field. Here this state corresponds to the one that is annihilated by the annihilation operator given in the solutions (4.16) and (4.17),

$$\hat{b}_T(\sigma, \vec{k})|\Omega\rangle = 0, \quad \forall \vec{k}, \sigma. \quad (4.19)$$

This is the state that we shall consider in Sec. 5 when computing the photon two-point function.

4.2 Scalar sector

The second and the fourth equations (4.10) and (4.12) of the scalar sector decouple from the rest, and combine into a single second order one,

$$\left[\partial_0^2 + k^2 - \left(\nu^2 - \frac{1}{4} \right) (1-\epsilon)^2 \mathcal{H}^2 \right] \hat{\pi}_L = 0, \quad (4.20)$$

$$\hat{\pi}_0 = -\frac{1}{k} \left[\partial_0 + \left(\nu + \frac{1}{2} \right) (1-\epsilon) \mathcal{H} \right] \hat{\pi}_L, \quad (4.21)$$

where we introduce the index of the scalar sector,

$$\nu = \frac{D-3+\epsilon}{2(1-\epsilon)}. \quad (4.22)$$

According to considerations in Sec. 2.2 the solutions are immediately written as,

$$\hat{\pi}_L(\eta, \vec{k}) = k U_\nu(\eta, k) \hat{b}_P(\vec{k}) + k U_\nu^*(\eta, k) \hat{b}_P^\dagger(-\vec{k}), \quad (4.23)$$

$$\hat{\pi}_0(\eta, \vec{k}) = ik U_{\nu+1}(\eta, k) \hat{b}_P(\vec{k}) - ik U_{\nu+1}^*(\eta, k) \hat{b}_P^\dagger(-\vec{k}), \quad (4.24)$$

where the non-Hermitian operators $\hat{b}_P(\vec{k})$ play the role of initial conditions/integration constants. The remaining two equations (4.9) and (4.11) combine into a sourced second-order equation,

$$\left[\partial_0^2 + k^2 - \left(\nu^2 - \frac{1}{4} \right) (1-\epsilon)^2 \mathcal{H}^2 \right] \hat{\mathcal{A}}_0 = (1-\xi) a^2 k \hat{\pi}_L - 2\xi a^2 \mathcal{H} \hat{\pi}_0, \quad (4.25)$$

$$\hat{\mathcal{A}}_L = \frac{1}{k} \left[\partial_0 + \left(\nu + \frac{1}{2} \right) (1-\epsilon) \mathcal{H} \right] \hat{\mathcal{A}}_0 + \frac{\xi a^2}{k} \hat{\pi}_0, \quad (4.26)$$

We can write the solutions to these equations as,

$$\begin{aligned} \hat{\mathcal{A}}_0(\eta, \vec{k}) &= U_\nu(\eta, k) \hat{b}_H(\vec{k}) + U_\nu^*(\eta, k) \hat{b}_H^\dagger(-\vec{k}) \\ &\quad + v_0(\eta, k) \hat{b}_P(\vec{k}) + v_0^*(\eta, k) \hat{b}_P^\dagger(-\vec{k}), \end{aligned} \quad (4.27)$$

$$\begin{aligned} \hat{\mathcal{A}}_L(\eta, \vec{k}) &= -i U_{\nu+1}(\eta, k) \hat{b}_H(\vec{k}) + i U_{\nu+1}^*(\eta, k) \hat{b}_H^\dagger(-\vec{k}) \\ &\quad - i v_L(\eta, k) \hat{b}_P(\vec{k}) + i v_L^*(\eta, k) \hat{b}_P^\dagger(-\vec{k}), \end{aligned} \quad (4.28)$$

where the homogeneous parts depend on operators $\hat{b}_H(\vec{k})$ standing for integration constants, while the particular parts depend on the integration constants for canonical momentum operators introduced in solutions (4.23) and (4.24). The particular mode functions v_0 and v_L in the solutions (4.27) and (4.28) satisfy equations descending from (4.25) and (4.26),

$$\begin{aligned} &\left[\partial_0^2 + k^2 - \left(\nu^2 - \frac{1}{4} \right) (1-\epsilon)^2 \mathcal{H}^2 \right] v_0 \\ &= \frac{-i\xi k a^2 \mathcal{H}}{[\nu(1-\epsilon)+1]} \left[2[\nu(1-\epsilon)+1] U_{\nu+1} - (1+\epsilon) \frac{ik}{\mathcal{H}} U_\nu \right] + \left(1 - \frac{\xi}{\xi_s} \right) k^2 a^2 U_\nu, \end{aligned} \quad (4.29)$$

$$v_L = \frac{i}{k} \left[\partial_0 + \left(\nu + \frac{1}{2} \right) (1-\epsilon) \mathcal{H} \right] v_0 - \xi a^2 U_{\nu+1}. \quad (4.30)$$

where we introduced the parameter,

$$\xi_s = \frac{\nu(1-\epsilon)+1}{\nu(1-\epsilon)-\epsilon} = \frac{D-1+\epsilon}{D-3-\epsilon}, \quad (4.31)$$

that we refer to as the *simple covariant gauge*. This name is suggestive of the simplification that happens when the gauge fixing parameter takes this value, $\xi = \xi_s$. The second part of the source in (4.29) drops out, and the identity (2.17) immediately gives the solutions,

$$v_0 \xrightarrow{\xi \rightarrow \xi_s} \frac{-ik}{(D-3-\epsilon)} \left[\frac{a^2}{\mathcal{H}} U_{\nu+1} - \frac{1}{H_0} U_\nu \right], \quad (4.32)$$

$$v_L \xrightarrow{\xi \rightarrow \xi_s} \frac{-ik}{(D-3-\epsilon)} \left[\frac{a^2}{\mathcal{H}} U_\nu - \frac{1}{H_0} U_{\nu+1} \right]. \quad (4.33)$$

These solutions for particular mode functions are chosen to satisfy the Wronskian-like relation,

$$\text{Re} \left(v_0 U_{\nu+1}^* + v_L U_\nu^* \right) = 0, \quad (4.34)$$

in addition to reducing to the corresponding de Sitter space solutions found in [74]. The Wronskian relations (2.16) and (4.34) are in fact sufficient to compute the commutation relations between the momentum space operators introduced appearing in solutions for the field operators, without solving for the particular mode functions. They follow from the canonical commutation relations (4.6),

$$[\hat{b}_H(\vec{k}), \hat{b}_H^\dagger(\vec{k}')] = [\hat{b}_P(\vec{k}), \hat{b}_P^\dagger(\vec{k}')] = 0, \quad [\hat{b}_H(\vec{k}), \hat{b}_P^\dagger(\vec{k}')] = -\delta^{D-1}(\vec{k} - \vec{k}'), \quad (4.35)$$

These commutators are not canonical, in the sense that they are not the ones of creation/annihilation operators. Nevertheless, these operators are used to construct a basis of the space of states [79].

Finding particular mode functions for $\xi \neq \xi_s$ is considerably more complicated and Appendix A is devoted to the detailed derivation. Requiring that the mode functions reduce to the de Sitter solutions from [74] for an arbitrary gauge-fixing parameter,

$$v_0 \xrightarrow{\epsilon \rightarrow 0} \frac{-i\xi k}{2(\nu_0+1)H_0} \left[\frac{\mathcal{H}}{H_0} U_{\nu_0+1} - U_{\nu_0} \right] - \left(1 - \frac{\xi}{\xi_s^0} \right) \frac{ik}{2H_0} \left[\frac{ik}{\nu_0 H_0} \frac{\partial U_{\nu_0}}{\partial \nu_0} + U_{\nu_0} \right], \quad (4.36a)$$

$$v_L \xrightarrow{\epsilon \rightarrow 0} \frac{-i\xi k}{2(\nu_0+1)H_0} \left[\frac{\mathcal{H}}{H_0} U_{\nu_0} - U_{\nu_0+1} \right] - \left(1 - \frac{\xi}{\xi_s^0} \right) \frac{ik}{2H_0} \left[\frac{ik}{\nu_0 H_0} \frac{\partial U_{\nu_0+1}}{\partial \nu_0} + \frac{\mathcal{H}}{\nu_0 H_0} U_{\nu_0} + U_{\nu_0+1} \right], \quad (4.36b)$$

where,

$$\nu \xrightarrow{\epsilon \rightarrow 0} \frac{D-3}{2} \equiv \nu_0, \quad \xi_s \xrightarrow{\epsilon \rightarrow 0} \frac{D-1}{D-3} \equiv \xi_s^0, \quad (4.37)$$

and that in the flat space limit they reduce directly to,

$$v_0 \xrightarrow{H_0 \rightarrow 0} \frac{1}{4} \left[(1+\xi) + 2(1-\xi)ik(\eta-\eta_0) \right] \frac{e^{-ik(\eta-\eta_0)}}{\sqrt{2k}}, \quad (4.38a)$$

$$v_L \xrightarrow{H_0 \rightarrow 0} \frac{1}{4} \left[-(1+\xi) + 2(1-\xi)ik(\eta-\eta_0) \right] \frac{e^{-ik(\eta-\eta_0)}}{\sqrt{2k}}, \quad (4.38b)$$

in addition to satisfying (4.34), essentially uniquely fixes the solutions to be,

$$v_0(\eta, k) = \frac{-i\xi k}{2[\nu(1-\epsilon)+1]} \left[\frac{a^2}{\mathcal{H}} U_{\nu+1}(\eta, k) - \frac{1}{H_0} U_\nu(\eta, k) \right] + \left(1 - \frac{\xi}{\xi_s}\right) \left[\mathcal{Q}(\eta, k) U_\nu(\eta, k) - \tilde{\mathcal{Q}}(\eta, k) U_\nu^*(\eta, k) \right], \quad (4.39)$$

$$v_L(\eta, k) = \frac{-i\xi k}{2[\nu(1-\epsilon)+1]} \left[\frac{a^2}{\mathcal{H}} U_\nu(\eta, k) - \frac{1}{H_0} U_{\nu+1}(\eta, k) \right] + \left(1 - \frac{\xi}{\xi_s}\right) \left[\mathcal{Q}(\eta, k) U_{\nu+1}(\eta, k) + \tilde{\mathcal{Q}}(\eta, k) U_{\nu+1}^*(\eta, k) \right]. \quad (4.40)$$

The coefficient functions here are given as linear combinations,

$$\mathcal{Q}(\eta, k) = \frac{i}{4\pi} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \left[-2 \cos(\pi\nu) \mathcal{J}_2\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) + \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) + \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, -\nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) \right] + \mathcal{A}, \quad (4.41)$$

$$\tilde{\mathcal{Q}}(\eta, k) = \frac{e^{\frac{-2ik}{(1-\epsilon)H_0}}}{4\pi} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \left[-2 \mathcal{J}_2\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) + e^{-i\pi\nu} \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) + e^{i\pi\nu} \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, -\nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) \right] + \mathcal{B}, \quad (4.42)$$

of special functions defined in terms of generalized hypergeometric functions,

$$\mathcal{J}_1(\rho, \lambda; z) = \frac{-\Gamma^2(-\lambda)z^{\rho+2\lambda}}{4^\lambda(\rho+2\lambda)} {}_2F_3\left(\left\{\frac{\rho}{2}+\lambda, \frac{1}{2}+\lambda\right\}, \left\{\frac{2+\rho}{2}+\lambda, 1+\lambda, 1+2\lambda\right\}, -z^2\right), \quad (4.43)$$

$$\mathcal{J}_2(\rho, \lambda; z) = \frac{\Gamma(\lambda)\Gamma(-\lambda)z^\rho}{\rho} {}_2F_3\left(\left\{\frac{\rho}{2}, \frac{1}{2}\right\}, \left\{\frac{2+\rho}{2}, 1+\lambda, 1-\lambda\right\}, -z^2\right), \quad (4.44)$$

and where the two integration constants are given by,

$$\mathcal{A} = \frac{-ik}{2(1+\epsilon)H_0} + \frac{i\Gamma\left(\frac{-\epsilon}{1-\epsilon}\right)\Gamma\left(\frac{1}{2}+\frac{\epsilon}{1-\epsilon}\right)\Gamma\left(\frac{-\epsilon}{1-\epsilon}+\nu\right)\Gamma\left(\frac{-\epsilon}{1-\epsilon}-\nu\right)}{4\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}+\nu\right)\Gamma\left(\frac{1}{2}-\nu\right)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}}, \quad (4.45)$$

$$\mathcal{B} = \frac{\Gamma\left(\frac{-\epsilon}{1-\epsilon}\right)\Gamma\left(\frac{-\epsilon}{1-\epsilon}+\nu\right)\Gamma\left(\frac{-\epsilon}{1-\epsilon}-\nu\right)}{4\Gamma\left(\frac{1}{2}\right)\Gamma\left(\frac{1}{2}-\frac{\epsilon}{1-\epsilon}\right)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} e^{\frac{-2ik}{(1-\epsilon)H_0}} e^{\frac{-i\pi\epsilon}{(1-\epsilon)}}. \quad (4.46)$$

The huge difference in the complexity of particular mode functions in the general covariant gauge versus the simple covariant gauge $\xi = \xi_s$ makes the former case rather intractable to work with. This is why from Sec. 5 on we consider only the simple covariant gauge when computing the position-space two-point functions and the observables.

4.3 Constraints

The subsidiary condition on the space of states (3.16) takes an analogous expression in momentum space,

$$\hat{\mathcal{K}}(\vec{k})|\Omega\rangle = 0, \quad \forall \vec{k}, \quad (4.47)$$

where the non-Hermitian constraint operator is given as a local linear combination,

$$\hat{\mathcal{K}}(\vec{k}) = c_1(\eta, k)\hat{\psi}_1(\eta, \vec{k}) + c_2(\eta, k)\hat{\psi}_2(\eta, \vec{k}), \quad (4.48)$$

of the momentum space Hermitian constraint operators,

$$\hat{\psi}_1(\eta, \vec{k}) = \hat{\pi}_0(\eta, \vec{k}), \quad \hat{\psi}_2(\eta, \vec{k}) = \hat{\pi}_L(\eta, \vec{k}). \quad (4.49)$$

The conservation of (4.48) implies equations for the coefficients,

$$\partial_0 c_1 - k c_2 + \frac{1}{2}(D-2)\mathcal{H}c_1 = 0, \quad \partial_0 c_2 + k c_1 - \frac{1}{2}(D-2)\mathcal{H}c_2 = 0. \quad (4.50)$$

They combine into a second order equation,

$$\left[\partial_0^2 + k^2 - \left(\nu^2 - \frac{1}{4} \right) (1-\epsilon)^2 \right] c_1 = 0, \quad (4.51)$$

$$c_2 = \frac{1}{k} \left[\partial_0 + \left(\nu + \frac{1}{2} \right) (1-\epsilon)\mathcal{H} \right] c_1, \quad (4.52)$$

that is the scalar mode equation (2.9) with the general solution given in (2.10). The general solution for the non-Hermitian constraint respecting the symmetries of the background,

$$\hat{\mathcal{K}}(\vec{k}) = e^{i\theta(k)} \left(e^{-i\varphi(k)} \text{ch}[\rho(k)] \hat{b}_P(\vec{k}) + e^{i\varphi(k)} \text{sh}[\rho(k)] \hat{b}_P^\dagger(\vec{k}) \right), \quad (4.53)$$

can be parametrized by three real functions θ , φ , and ρ of the momentum modulus. The space of states has to admit at least one vector annihilated by the operator above. To fully specify this state we need another operator that will annihilate it,

$$\hat{\mathcal{B}}(\vec{k})|\Omega\rangle = 0, \quad \forall \vec{k}, \quad (4.54)$$

and the choice preserving commutation relations (4.35),

$$[\hat{\mathcal{K}}(\vec{k}), \hat{\mathcal{K}}^\dagger(\vec{k}')] = [\hat{\mathcal{B}}(\vec{k}), \hat{\mathcal{B}}^\dagger(\vec{k}')] = 0, \quad [\hat{\mathcal{K}}(\vec{k}), \hat{\mathcal{B}}^\dagger(\vec{k}')] = -\delta^{D-1}(\vec{k} - \vec{k}') \quad (4.55)$$

is given by,

$$\hat{\mathcal{B}}(\vec{k}) = e^{i\theta(k)} \left(e^{-i\varphi(k)} \text{ch}[\rho(k)] \hat{b}_H(\vec{k}) + e^{i\varphi(k)} \text{sh}[\rho(k)] \hat{b}_H^\dagger(\vec{k}) \right). \quad (4.56)$$

The choice consistent with de Sitter symmetries [74] in the limit $\epsilon \rightarrow 0$, and with Poincaré symmetries [114, 115] in the flat space limit $H_0 \rightarrow 0$ would be for all the three real functions θ, φ , and ρ to vanish, and that would be the natural choice to consider here. Nevertheless, anticipating issues with IR convergence of the two-point function, we take for the Bogolyubov-like coefficients of the scalar sector to be,

$$\rho(k) = \theta(k) = \varphi(k) = 0, \quad k \geq k_0, \quad (4.57)$$

where $k_0 \ll H_0$ is some deep IR scale, which implies that the physical state has to satisfy,

$$\hat{b}_P(\vec{k})|\Omega\rangle = 0, \quad \hat{b}_H(\vec{k})|\Omega\rangle = 0, \quad k \geq k_0. \quad (4.58)$$

Below the IR scale k_0 we keep the general form of the Bogolyubov coefficients in (4.53) and (4.56), but with an additional assumption. They are chosen such that the contribution of deep IR modes to the two-point function contains only terms that vanish in the limit $k_0 \rightarrow 0$. That means their contribution to the two-point function can safely be neglected if $k_0/H_0 \ll 1$ is assumed small enough. The logic behind such a choice mirrors the one from Sec. 2.3.2 for the scalar two-point functions.

5 Two-point function

The two-point functions of a free field theory are the building blocks of loop expansions in quantum field theory. In this section we use the solutions for field operators found in the preceding section to evaluate the expectation values that define the two-point function. We first recount the general properties that two-point functions must satisfy regardless of the quantum state, and then proceed to define the natural Gaussian state with respect to which we compute the two-point function as a sum-over modes. Due to the complicated form of the scalar sector mode functions we only evaluate the sum-over-modes in the simple covariant gauge (4.31) in which the mode functions are tractable. Nonetheless, the final covariantized expression for the two-point functions we find is rather complicated, and does not exhibit a simple structure of being composed of scalar two-point functions and their derivatives. This owes to the necessity of explicitly evaluating the inverse Laplace operator acting on the scalar-two point function which results in a combination of Appell's fourth functions, as shown in Appendix B.

5.1 Generalities

The Schwinger-Keldysh formalism for nonequilibrium quantum field theory [86–92] (see e.g. [93, 94] for introductions) necessitates the use of several different two-point

functions in perturbative (loop) computations. The positive-frequency Wightman function for the photon field is defined as the expectation value of the off-coincident product of two field operators,

$$i[{}_{\mu}^{-}\Delta_{\nu}^{+}](x; x') = \langle \Omega | \hat{A}_{\mu}(x) \hat{A}_{\nu}(x') | \Omega \rangle, \quad (5.1)$$

while its complex conjugate, $i[{}_{\mu}^{+}\Delta_{\nu}^{-}](x; x') = \{i[{}_{\mu}^{-}\Delta_{\nu}^{+}](x; x')\}^*$, is the negative-frequency Wightman function. The expectation value of the time-ordered product is the Feynman propagator,

$$\begin{aligned} i[{}_{\mu}^{+}\Delta_{\nu}^{+}](x; x') &= \langle \Omega | \mathcal{T}(\hat{A}_{\mu}(x) \hat{A}_{\nu}(x')) | \Omega \rangle \\ &= \theta(\eta - \eta') i[{}_{\mu}^{-}\Delta_{\nu}^{+}](x; x') + \theta(\eta' - \eta) i[{}_{\mu}^{+}\Delta_{\nu}^{-}](x; x'), \end{aligned} \quad (5.2)$$

and its complex conjugate, $i[{}_{\mu}^{-}\Delta_{\nu}^{-}](x; x') = \{i[{}_{\mu}^{+}\Delta_{\nu}^{+}](x; x')\}^*$, is the Dyson propagator. The position space equations of motion (3.11) satisfied by field operators can be written in the more familiar covariant form,

$$\mathcal{D}_{\mu}{}^{\nu} \hat{A}_{\nu}(x) = 0, \quad \mathcal{D}_{\mu}{}^{\nu} = \nabla_{\rho} \nabla^{\rho} \delta_{\mu}^{\nu} - \left(1 - \frac{1}{\xi}\right) \nabla_{\mu} \nabla^{\nu} - R_{\mu}{}^{\nu}. \quad (5.3)$$

These equations of motion are inherited by the two-point functions,

$$\mathcal{D}_{\mu}{}^{\rho} i[{}_{\rho}^{\text{a}}\Delta_{\nu}^{\text{b}}](x; x') = \mathbf{S}^{\text{ab}} g_{\mu\nu} \frac{i\delta^D(x-x')}{\sqrt{-g}}, \quad (5.4)$$

where the sign symbol \mathbf{S}^{ab} is defined in (2.21). The local source in the equations for the Feynman and Dyson propagators descends from time-ordering in the definition (5.2), and from the canonical commutation relations (3.13). In multiplier gauges the equations of motion are not the only equations that two-point functions must satisfy. There are also subsidiary conditions (3.15) that descend from the classical first-class constraints. These subsidiary conditions are all captured by the Ward-Takahashi identity,

$$\nabla^{\mu} i[{}_{\mu}^{\text{a}}\Delta_{\nu}^{\text{b}}](x; x') = -\xi \partial'_{\nu} i[{}^{\text{a}}\Delta^{\text{b}}]_{\nu+1}(x; x'), \quad (5.5)$$

where the quantity on the right-hand-side is the massless, minimally coupled scalar (MMCS) two-point function,

$$\square i[{}^{\text{a}}\Delta^{\text{b}}]_{\nu+1}(x; x') = \mathbf{S}^{\text{ab}} \frac{i\delta^D(x-x')}{\sqrt{-g}}, \quad (5.6)$$

discussed in Sec. 2. We compute the photon two-point functions in the following section using the sum-over-modes representation with the mode functions worked out in sections 4.1 and 4.2. In appendix C the results for two-point functions are checked to satisfy the appropriate equation of motion (5.4) and the Ward-Takahashi identity (5.5).

5.2 Evaluating sum-over-modes

In Sec. 4 we have found solutions (4.16), (4.27), and (4.28) for the vector field operators in momentum space, and have specified the quantum state that we consider in (4.19) and (4.58). Together with the commutation relations (4.18) and (4.35), this is all we need to express the two-point function (5.1) as a sum-over-modes,

$$i[\bar{0}\Delta_0^+](x; x') = (aa')^{\frac{2-D}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot\Delta\vec{x}} \theta(k-k_0) \times \left[-U_\nu(\eta, k)v_0^*(\eta', k) - v_0(\eta, k)U_\nu^*(\eta', k) \right], \quad (5.7)$$

$$i[\bar{0}\Delta_i^+](x; x') = (aa')^{\frac{2-D}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot\Delta\vec{x}} \theta(k-k_0) \times \frac{k_i}{k} \left[U_\nu(\eta, k)v_L^*(\eta', k) + v_0(\eta, k)U_{\nu+1}^*(\eta', k) \right], \quad (5.8)$$

$$i[\bar{i}\Delta_0^+](x; x') = (aa')^{\frac{2-D}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot\Delta\vec{x}} \theta(k-k_0) \times \frac{k_i}{k} \left[U_{\nu+1}(\eta, k)v_0^*(\eta', k) + v_L(\eta, k)U_\nu^*(\eta', k) \right], \quad (5.9)$$

$$i[\bar{i}\Delta_j^+](x; x') = (aa')^{\frac{4-D}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot\Delta\vec{x}} \left(\delta_{ij} - \frac{k_i k_j}{k^2} \right) U_{\nu_T}(\eta, k)U_{\nu_T}^*(\eta', k) \quad (5.10) \\ - (aa')^{\frac{2-D}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} e^{i\vec{k}\cdot\Delta\vec{x}} \theta(k-k_0) \frac{k_i k_j}{k^2} \left[U_{\nu+1}(\eta, k)v_L^*(\eta', k) + v_L(\eta, k)U_{\nu+1}^*(\eta', k) \right].$$

Note that the reason why step functions $\theta(k-k_0)$ appear is given in Sec. 4.3. Because not all sum-over modes of the scalar sector will be IR finite for a natural definition of the state, we need to consider Bogolyubov coefficients in the IR that ameliorate this problem and regulate the behaviour of the mode function. We choose to implement this in a way that is least sensitive to this regulating procedure and can be parametrized effectively by a deep IR scale $k_0 \ll H_0$ that effectively manifests itself as an IR cutoff on the sums-over-modes descending from the scalar sector. There is no need for such regulation of the transverse sector. Note also that the $i\delta$ -prescriptions are implicit in (5.7)–(5.10) in the same way as for the scalar propagator (2.18).

In the following subsections we evaluate the integrals in (5.7)–(5.10) for the case of simple covariant gauge $\xi = \xi_s$ defined in (4.31), for which the solutions for the particular mode functions (4.32) and (4.33) of the scalar sector are tractable. Upon plugging in these mode functions we can utilize recurrence relations (2.15) to recognize that the components of the photon two-point function can all be expressed in terms of certain operators acting on sums-over-modes (2.39) that represent the

scalar two-point functions,

$$i[{}_{0}^{-}\Delta_0^+](x; x') = \frac{-1}{(D-3-\epsilon)} \left[\frac{a^2}{\mathcal{H}} \partial_0 + \frac{a'^2}{\mathcal{H}'} \partial'_0 + (D-2)(a^2 + a'^2) \right] i[{}^{-}\Delta^+]_{\nu}(x; x'), \quad (5.11)$$

$$i[{}_{0}^{-}\Delta_i^+](x; x') = \frac{\partial'_i}{(D-3-\epsilon)} \left[\frac{a^2}{\mathcal{H}} i[{}^{-}\Delta^+]_{\nu+1}(x; x') - \frac{a'^2}{\mathcal{H}'} i[{}^{-}\Delta^+]_{\nu}(x; x') \right], \quad (5.12)$$

$$i[{}_i^{-}\Delta_0^+](x; x') = \frac{\partial_i}{(D-3-\epsilon)} \left[\frac{a'^2}{\mathcal{H}'} i[{}^{-}\Delta^+]_{\nu+1}(x; x') - \frac{a^2}{\mathcal{H}} i[{}^{-}\Delta^+]_{\nu}(x; x') \right], \quad (5.13)$$

$$i[{}_i^{-}\Delta_j^+](x; x') = \delta_{ij} a a' i[{}^{-}\Delta^+]_{\nu_T}(x; x') + \frac{a a' \partial_i \partial'_j}{(1-\epsilon)^2 \mathcal{H} \mathcal{H}'} i[{}^{-}\Xi^+]_{\nu_T}(x; x') \\ - \frac{1}{(D-3-\epsilon)} \frac{\partial_i \partial'_j}{(1-\epsilon)^2 \mathcal{H} \mathcal{H}'} \left[\frac{a'^2}{\mathcal{H}'} \partial_0 + \frac{a^2}{\mathcal{H}} \partial'_0 + (D-3+\epsilon) \left(\frac{a^2 \mathcal{H}'}{\mathcal{H}} + \frac{a'^2 \mathcal{H}}{\mathcal{H}'} \right) \right] i[{}^{-}\Xi^+]_{\nu}(x; x'). \quad (5.14)$$

Thus we have accomplished expressing the photon two-point function in terms of derivatives acting on scalar two-point functions, and on what we call Laplace-inverted two-point functions that result from inverting the Laplace operator on the scalar two-point function,

$$i[{}^{-}\Xi^+]_{\lambda}(x; x') \equiv \frac{(1-\epsilon)^2 \mathcal{H} \mathcal{H}'}{\nabla^2} i[{}^{-}\Delta^+]_{\lambda}(x; x'), \quad (5.15)$$

where the time-dependent factor on the right-hand-side is chosen for convenience.

In the de Sitter limit the combination of temporal derivatives in the brackets of the second line in (5.14) conspire to produce a Laplacian that eliminates its inverse [74]. However, here we are not as fortunate, and have to consider computing (5.15) explicitly. This is accomplished in a form of an asymptotic series of derivative operators acting on the scalar two-point function in the next subsection, while in Appendix B a detailed analysis of the sum-over-modes representation of (5.15) is given and a closed form solution in terms of Appell's fourth function is derived.

For the remainder of this subsection it is not necessary to evaluate (5.15). It is sufficient to note that the Laplace-inverted two-point function inherits from the scalar two-point function the symmetry under the exchange of coordinates, and the cosmological symmetries, and that it can be considered as a function of bi-local variables, $i[{}^{\text{a}\Xi^{\text{b}}}]_{\lambda}(x; x') = i\Xi_{\lambda}(y_{\text{ab}}, u_{\text{ab}}, v_{\text{ab}})$, just as for the scalar two-point function. Thus we act explicitly with temporal derivatives in (5.11–5.14) to write the components of the two-point functions as,

$$i[{}_{0}^{-}\Delta_0^+](x; x') = \frac{\mathcal{H} \mathcal{H}' e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T H_0^2} \left[2 \operatorname{ch}\left(\frac{v}{1-\epsilon}\right) \left((2-y) \frac{\partial}{\partial y} - \frac{\partial}{\partial u} - 2\nu - 1 \right) \right. \\ \left. - 4 \operatorname{ch}\left(\frac{\epsilon v}{1-\epsilon}\right) \frac{\partial}{\partial y} - 2 \operatorname{sh}\left(\frac{v}{1-\epsilon}\right) \frac{\partial}{\partial v} \right] i\Delta_{\nu}(y, u, v), \quad (5.16)$$

$$i[{}_{0}^{-}\Delta_i^+](x; x') = \frac{(\partial'_i y) \mathcal{H} e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T (1-\epsilon) H_0^2} \left[e^{\frac{\epsilon v}{1-\epsilon}} \frac{\partial}{\partial y} i\Delta_{\nu+1}(y, u, v) - e^{\frac{-v}{1-\epsilon}} \frac{\partial}{\partial y} i\Delta_{\nu}(y, u, v) \right], \quad (5.17)$$

$$i[{}_i^{-}\Delta_0^+](x; x') = \frac{(\partial_i y) \mathcal{H}' e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T (1-\epsilon) H_0^2} \left[e^{\frac{-\epsilon v}{1-\epsilon}} \frac{\partial}{\partial y} i\Delta_{\nu+1}(y, u, v) - e^{\frac{v}{1-\epsilon}} \frac{\partial}{\partial y} i\Delta_{\nu}(y, u, v) \right], \quad (5.18)$$

$$i[\bar{i}\Delta_j^+](x; x') = \frac{\mathcal{H}\mathcal{H}'}{H_0^2} \delta_{ij} e^{\frac{\epsilon u}{1-\epsilon}} i\Delta_{\nu_T}(y, u, v) + \frac{e^{\frac{\epsilon u}{1-\epsilon}} \partial_i \partial'_j}{2\nu_T(1-\epsilon)^2 H_0^2} \left\{ 2\nu_T i\Xi_{\nu_T}(y, u, v) \right. \\ \left. + \left[2 \operatorname{ch}\left(\frac{\epsilon v}{1-\epsilon}\right) \left((2-y) \frac{\partial}{\partial y} - \frac{\partial}{\partial u} - 2\nu \right) - 4 \operatorname{ch}\left(\frac{v}{1-\epsilon}\right) \frac{\partial}{\partial y} + 2 \operatorname{sh}\left(\frac{\epsilon v}{1-\epsilon}\right) \frac{\partial}{\partial v} \right] i\Xi_\nu(y, u, v) \right\}. \quad (5.19)$$

Even though we cannot fully eliminate the complicated Laplace-inverted function, it is nonetheless advantageous to apply some simplifications. Introducing the notation $I[f(y)] \equiv \int^y dy' f(y')$ for the primitive function with respect to variable y , we can make use of the two identities for Laplace operators acting on some arbitrary bi-scalar,

$$I\left[\nabla^2 f(y, u, v)\right] = \nabla^2 I[f(y, u, v)] - 4(1-\epsilon)^2 \mathcal{H}\mathcal{H}' f(y, u, v), \quad (5.20)$$

$$\nabla^2 I[f(y, u, v)] = 4(1-\epsilon)^2 \mathcal{H}\mathcal{H}' \left(\left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right] \frac{\partial}{\partial y} + \frac{(D-1)}{2} \right) f(y, u, v), \quad (5.21)$$

that are proven by explicitly acting the derivatives, to derive another useful identity,

$$\left(\left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right] \frac{\partial}{\partial y} + \frac{D-3}{2} \right) i\Xi_\lambda(y, u, v) = \frac{1}{4} I[i\Delta_\lambda(y, u, v)]. \quad (5.22)$$

Applying it to the (ij) component in (5.19) results in,

$$i[\bar{i}\Delta_j^+](x; x') = \frac{\mathcal{H}\mathcal{H}'}{H_0^2} \delta_{ij} e^{\frac{\epsilon u}{1-\epsilon}} i\Delta_{\nu_T}(y, u, v) \\ + \frac{e^{\frac{\epsilon u}{1-\epsilon}} \partial_i \partial'_j}{2\nu_T(1-\epsilon)^2 H_0^2} \left[-\frac{1}{2} \operatorname{ch}\left(\frac{\epsilon v}{1-\epsilon}\right) I[i\Delta_\nu(y, u, v)] + I[i\Upsilon(y, u, v)] \right], \quad (5.23)$$

where for notational simplicity in the remainder of the paper it is useful to define

$$i\Upsilon(y, u, v) \equiv 2\nu_T \frac{\partial}{\partial y} i\Xi_{\nu_T}(y, u, v) - 2 \operatorname{ch}\left(\frac{\epsilon v}{1-\epsilon}\right) \left(\frac{\partial}{\partial u} + 2\nu - \frac{D-3}{2} \right) \frac{\partial}{\partial y} i\Xi_\nu(y, u, v) \\ - 2 \operatorname{sh}\left(\frac{\epsilon v}{1-\epsilon}\right) \left(2 \operatorname{sh}(v) \frac{\partial}{\partial y} - \frac{\partial}{\partial v} \right) \frac{\partial}{\partial y} i\Xi_\nu(y, u, v). \quad (5.24)$$

This form (5.23) of the (ij) component is useful because it makes manifest that in de Sitter limit all the complicated functions all cancel since $i\Upsilon \xrightarrow{\epsilon \rightarrow 0} 0$; this will be discussed further in Sec. 6.1.

It is also interesting to note down the expression for the two-point function of just the transverse components of the spatial vector potential,

$$i[\bar{i}\Delta_j^+]^{\text{ph}}(x; x') = \langle \Omega | \hat{A}_i^T(x) \hat{A}_j^T(x') | \Omega \rangle = e^{\frac{u}{1-\epsilon}} \left(\delta_{ij} + \frac{\partial_i \partial'_j}{\nabla^2} \right) i\Delta_{\nu_T}(y, u, v) \\ = e^{\frac{u}{1-\epsilon}} \left[\delta_{ij} i\Delta_{\nu_T}(y, u, v) + \partial_i \partial'_j \frac{i\Xi_{\nu_T}(y, u, v)}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} \right]. \quad (5.25)$$

This can rightfully be called the physical two-point function, as it corresponds to the two-point function in the Coulomb gauge, in which $\partial_i \hat{A}_i = 0$ and $\hat{A}_0 = 0$ stand as an operator equalities.

5.3 Laplace-inverted two-point function

The Laplace-inverted two-point function in (5.15) is a formal solution of the Poisson equation with the scalar Wightman function as the source. There are in fact such equations for all four combinations of Schwinger-Keldysh polarities,

$$\nabla^2 \frac{i[\mathbf{a}\Xi^{\mathbf{b}}]_{\lambda}(x; x')}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} = i[\mathbf{a}\Delta^{\mathbf{b}}]_{\lambda}(x; x'), \quad (5.26)$$

determined by different scalar two-point functions appearing as sources on the right-hand side. However, this is not the only equation that Laplace-inverted two-point functions have to satisfy. Using the equation of motion (2.20) for the scalar two-point function it follows that the Laplace-inverted two-point functions satisfy a fourth order equation,

$$\left[\square - (1-\epsilon)^2 H^2 \left(\left[\frac{D-1-\epsilon}{2(1-\epsilon)} \right]^2 - \lambda^2 \right) \right] \nabla^2 \frac{i[\mathbf{a}\Xi^{\mathbf{b}}]_{\lambda}(x; x')}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} = \mathbf{S}^{\mathbf{ab}} \frac{i\delta^D(x-x')}{\sqrt{-g}}. \quad (5.27)$$

Given that the Laplacian commutes with the scalar d'Alembertian, and that the Poisson equation for a point charge has a unique solution, it follows that the Laplace-inverted two-point functions satisfy another second order equation,

$$\begin{aligned} & \left[\square - (1-\epsilon)^2 H^2 \left(\left[\frac{D-1-\epsilon}{2(1-\epsilon)} \right]^2 - \lambda^2 \right) \right] \frac{i[\mathbf{a}\Xi^{\mathbf{b}}]_{\lambda}(x; x')}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} \\ &= \mathbf{S}^{\mathbf{ab}} \nabla^{-2} \frac{i\delta^D(x-x')}{\sqrt{-g}} = \frac{-\mathbf{S}^{\mathbf{ab}} \Gamma\left(\frac{D-3}{2}\right) i\delta(\eta-\eta')}{4\pi^{\frac{D-1}{2}} \|\Delta\vec{x}\|^{D-3} \sqrt{-g}}. \end{aligned} \quad (5.28)$$

Naturally, Laplace-inverted two-point functions also satisfies the primed equation obtained by replacing $\square \rightarrow \square'$ and $H \rightarrow H'$ in the equation above. Also, using recurrence relations (2.15) for the mode functions, Laplace-inverted two-point functions can be seen to satisfy their own raising and lowering equations, respectively,

$$\begin{aligned} & \left[\partial_0 + \left(\frac{D-1-\epsilon}{2(1-\epsilon)} + \lambda \right) (1-\epsilon) \mathcal{H} \right] \left[\partial'_0 + \left(\frac{D-1-\epsilon}{2(1-\epsilon)} + \lambda \right) (1-\epsilon) \mathcal{H}' \right] \frac{i[\mathbf{a}\Xi^{\mathbf{b}}]_{\lambda}(x; x')}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} \\ &= -i[\mathbf{a}\Delta^{\mathbf{b}}]_{\lambda+1}(x; x') - \frac{\mathbf{S}^{\mathbf{ab}} \Gamma\left(\frac{D-3}{2}\right) i\delta(\eta-\eta')}{4\pi^{\frac{D-1}{2}} \|\Delta\vec{x}\|^{D-3} a^{D-2}}, \end{aligned} \quad (5.29)$$

$$\begin{aligned} & \left[\partial_0 + \left(\frac{D-1-\epsilon}{2(1-\epsilon)} - \lambda \right) (1-\epsilon) \mathcal{H} \right] \left[\partial'_0 + \left(\frac{D-1-\epsilon}{2(1-\epsilon)} - \lambda \right) (1-\epsilon) \mathcal{H}' \right] \frac{i[\mathbf{a}\Xi^{\mathbf{b}}]_{\lambda}(x; x')}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} \\ &= -i[\mathbf{a}\Delta^{\mathbf{b}}]_{\lambda-1}(x; x') - \frac{\mathbf{S}^{\mathbf{ab}} \Gamma\left(\frac{D-3}{2}\right) i\delta(\eta-\eta')}{4\pi^{\frac{D-1}{2}} \|\Delta\vec{x}\|^{D-3} a^{D-2}}. \end{aligned} \quad (5.30)$$

Different two-point functions with different Schwinger-Keldysh polarities can be represented by one function of bi-local variables (2.8) with different $i\delta$ -prescriptions,

$$i[\mathbf{a}\Xi^{\mathbf{b}}]_{\lambda}(x; x') = i\Xi_{\lambda}(y_{\mathbf{ab}}, u_{\mathbf{ab}}, v_{\mathbf{ab}}). \quad (5.31)$$

while the function on the right-hand-side then satisfies equations without local source terms. The first one follows from the defining Poisson equation (5.26),

$$\left(4\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]\frac{\partial}{\partial y}+2(D-1)\right)\frac{\partial}{\partial y}i\Xi_\lambda(y,u,v)=i\Delta_\lambda(y,u,v). \quad (5.32)$$

The second equation (5.28) and its primed counterpart are best given as two linear combinations, producing an even equation,

$$\begin{aligned} &\left[(4y-y^2)\frac{\partial^2}{\partial y^2}+D(2-y)\frac{\partial}{\partial y}-2\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]\left(\frac{\partial}{\partial u}+\frac{D\epsilon-2}{2(1-\epsilon)}\right)\frac{\partial}{\partial y}\right. \\ &\quad \left.-\left(\frac{\partial}{\partial u}+\frac{D-3+\epsilon}{1-\epsilon}\right)\frac{\partial}{\partial u}+\left(4\operatorname{sh}(v)\frac{\partial}{\partial y}-\frac{\partial}{\partial v}\right)\frac{\partial}{\partial v}+\lambda^2-\left(\frac{D-3+\epsilon}{2(1-\epsilon)}\right)^2\right]i\Xi_\lambda=0, \end{aligned} \quad (5.33)$$

and an odd equation,

$$\left[\left(\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]\frac{\partial}{\partial y}+\frac{\partial}{\partial u}+\frac{D-3+\epsilon}{2(1-\epsilon)}\right)\frac{\partial}{\partial v}-2\operatorname{sh}(v)\left(\frac{\partial}{\partial u}+\frac{D\epsilon-2}{2(1-\epsilon)}\right)\frac{\partial}{\partial y}\right]i\Xi_\lambda=0. \quad (5.34)$$

Lastly, the raising and lowering equations (5.29) and (5.30) take the following form,

$$\left[\left(\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]\frac{\partial}{\partial y}+\frac{\partial}{\partial u}+\frac{D-3+\epsilon}{2(1-\epsilon)}+\lambda\right)^2-\left(2\operatorname{sh}(v)\frac{\partial}{\partial y}-\frac{\partial}{\partial v}\right)^2\right]i\Xi_\lambda=-i\Delta_{\lambda+1}, \quad (5.35)$$

$$\left[\left(\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]\frac{\partial}{\partial y}+\frac{\partial}{\partial u}+\frac{D-3+\epsilon}{2(1-\epsilon)}-\lambda\right)^2-\left(2\operatorname{sh}(v)\frac{\partial}{\partial y}-\frac{\partial}{\partial v}\right)^2\right]i\Xi_\lambda=-i\Delta_{\lambda-1}. \quad (5.36)$$

Solving for $i\Xi_\lambda$ is best done by considering the first equation (5.32) out of the five above. From it we can readily solve for a derivative of $i\Xi_\lambda$ as a power series, by considering the expression in parentheses on the left-hand side of (5.32) as an expansion parameter,

$$\frac{\partial}{\partial y}i\Xi_\lambda(y,u,v)=\sum_{n=0}^{\infty}\left(\frac{-2}{D-1}\right)^n\left(\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]\frac{\partial}{\partial y}\right)^ni\Delta_\lambda(y,u,v). \quad (5.37)$$

It is convenient to commute the derivatives so they act only on $i\Delta_\lambda$, so that the operator is written as,

$$\left(\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]\frac{\partial}{\partial y}\right)^n=\sum_{\ell=1}^n\left\{\begin{matrix} n \\ \ell \end{matrix}\right\}\left[y+4\operatorname{sh}^2\left(\frac{v}{2}\right)\right]^\ell\frac{\partial^\ell}{\partial y^\ell}, \quad (5.38)$$

where the coefficients $\left\{\begin{matrix} n \\ \ell \end{matrix}\right\}$ in this expansion satisfy the following recurrence relations,

$$\left\{\begin{matrix} n+1 \\ \ell \end{matrix}\right\}=\ell\left\{\begin{matrix} n \\ \ell \end{matrix}\right\}+\left\{\begin{matrix} n \\ \ell-1 \end{matrix}\right\}, \quad \left\{\begin{matrix} n \\ n \end{matrix}\right\}=\left\{\begin{matrix} n \\ 1 \end{matrix}\right\}=1, \quad n\geq\ell\geq 1. \quad (5.39)$$

Therefore they are recognized to be Stirling numbers of the second kind (c.f. §26.8 of [84, 85]), that admit an explicit sum representation,

$$\left\{ \begin{matrix} n \\ \ell \end{matrix} \right\} = \sum_{j=0}^{\ell} \frac{(-1)^{\ell-j} j^n}{j! (\ell-j)!}. \quad (5.40)$$

Plugging this expansion back into the series solution (5.37), and reorganizing the series by grouping together all derivatives of the same order,

$$\frac{\partial}{\partial y} i\Xi_{\lambda} = \frac{i\Delta_{\lambda}}{2(D-1)} + \sum_{\ell=1}^{\infty} \left[\sum_{n=\ell}^{\infty} \left(\frac{-2}{D-1} \right)^n \left\{ \begin{matrix} n \\ \ell \end{matrix} \right\} \right] \times \left[y + 4 \operatorname{sh}^2 \left(\frac{v}{2} \right) \right]^{\ell} \frac{\partial^{\ell}}{\partial y^{\ell}} \frac{i\Delta_{\lambda}}{2(D-1)}. \quad (5.41)$$

The series over n can be recognized as the generating function for the Stirling numbers of the second kind (Eq. 26.8.11 from [84, 85]), that has a closed form expression,

$$\sum_{n=\ell}^{\infty} \left\{ \begin{matrix} n \\ \ell \end{matrix} \right\} z^{-n} = \frac{(-1)^{\ell} \Gamma(1-z)}{\Gamma(1-z+\ell)}. \quad (5.42)$$

This now yields,

$$\frac{\partial}{\partial y} i\Xi_{\lambda}(y, u, v) = \sum_{n=0}^{\infty} \frac{(-1)^n \Gamma\left(\frac{D-1}{2}\right)}{4 \Gamma\left(\frac{D+1}{2} + n\right)} \left[y + 4 \operatorname{sh}^2 \left(\frac{v}{2} \right) \right]^n \frac{\partial^n}{\partial y^n} i\Delta_{\lambda}(y, u, v). \quad (5.43)$$

Determining this derivative of $i\Xi_{\lambda}$ is sufficient for the photon two-point function due to the two spatial derivatives in (5.19). Nonetheless, we can use the identity that is proved by simple partial integration,

$$\sum_{n=0}^{\infty} c_n (y+\alpha)^n \frac{\partial^n}{\partial y^n} I[f(y)] = I \left[\sum_{n=0}^{\infty} \left(c_n + (n+1)c_{n+1} \right) (y+\alpha)^n \frac{\partial^n}{\partial y^n} f(y) \right], \quad (5.44)$$

to obtain the final expression for the Laplace-inverted two-point function,

$$i\Xi_{\lambda}(y, u, v) = \sum_{n=0}^{\infty} \frac{(-1)^n \Gamma\left(\frac{D-3}{2}\right)}{4 \Gamma\left(\frac{D-1}{2} + n\right)} \left[y + 4 \operatorname{sh}^2 \left(\frac{v}{2} \right) \right]^n \frac{\partial^n}{\partial y^n} I[i\Delta_{\lambda}(y, u, v)]. \quad (5.45)$$

Here the primitive function of the scalar two-point function with respect to y is defined as a term-by-term integral of the power series (2.32) and (2.47),

$$I[i\Delta_{\lambda}(y, u, v)] = e^{-\frac{(D-2)\epsilon}{2(1-\epsilon)}u} \left(I[\mathcal{F}_{\lambda}(y)] + I[\mathcal{W}(y, u, v)] \right), \quad (5.46)$$

such that no y -independent constants of integration are generated, neither in the bulk part,

$$I[\mathcal{F}_{\lambda}(y)] = -4 \frac{[(1-\epsilon)H_0]^{D-2}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma\left(\frac{D-3}{2} + \lambda\right) \Gamma\left(\frac{D-3}{2} - \lambda\right)}{\Gamma\left(\frac{D-2}{2}\right)} \times {}_2F_1 \left(\left\{ \frac{D-3}{2} + \lambda, \frac{D-3}{2} - \lambda \right\}, \left\{ \frac{D-2}{2} \right\}, 1 - \frac{y}{4} \right). \quad (5.47)$$

nor in the IR part,

$$\begin{aligned}
I[\mathcal{W}_\lambda(y, u, v)] &= -\frac{[(1-\epsilon)H_0]^{D-2}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(\lambda)\Gamma(2\lambda)}{\Gamma(\frac{1}{2}+\lambda)\Gamma(\frac{D-1}{2})} \sum_{N=0}^{\lfloor \lambda - \frac{D-1}{2} \rfloor} \sum_{n=0}^N \sum_{\ell=0}^{N-n} \frac{c_{Nn\ell}}{(\frac{D-1}{2}+N-\lambda)} \\
&\times \left[\frac{k_0^2 e^{-u}}{(1-\epsilon)^2 H_0^2} \right]^{\frac{D-1}{2}-\lambda+N} \frac{1}{(n+1)} \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right]^{n+1} \operatorname{ch}[(N-n-2\ell)v]. \quad (5.48)
\end{aligned}$$

Note that in deriving expression (5.45) we had not relied on the specific form of the source in Eq. (5.45), and thus it is valid for an arbitrary source for which the sum in (5.31) converges for some region of parameters. In principle there is a question of the homogeneous part of the solution, which is missed by the iterative method that generates the series solution. However, this contribution is not permitted as it would correspond to the Coulomb potential-like contribution that would generate a source which is not there. Checking that the solution in (5.45) also satisfies the remaining equations (5.33)–(5.36) is a tedious, but straightforward task of acting derivatives and applying equations for the scalar two-point function from Sec. 2.3.2. Note, however, that the series solution in (5.45) is asymptotic and that its practical utility is limited to small spatial separations. This corresponds to an implicit assumption that $\left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right] = (1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|$ is used as an expansion parameter when solving (5.32) iteratively. In general one should use the closed form solutions for the Laplace-inverted two-point function expressed in terms of Appell's fourth function, that we derive in Appendix B.

It is very useful to note that some of the equations that the Laplace-inverted two-point function satisfies are inherited by $i\Upsilon$ defined in (5.24). Firstly, it follows from Eq. (5.32) that $i\Upsilon$ satisfies,

$$\begin{aligned}
&\left(4 \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right] \frac{\partial}{\partial y} + 2(D-1) \right) i\Upsilon(y, u, v) \\
&= 2\nu_T i\Delta_{\nu_T}(y, u, v) - 2 \operatorname{ch}\left(\frac{\epsilon v}{1-\epsilon}\right) \left(\frac{\partial}{\partial u} + 2\nu - \frac{D-3}{2} \right) i\Delta_\nu(y, u, v) \\
&\quad - 2 \operatorname{sh}\left(\frac{\epsilon v}{1-\epsilon}\right) \left(2 \operatorname{sh}(v) \frac{\partial}{\partial y} - \frac{\partial}{\partial v} \right) i\Delta_\nu(y, u, v). \quad (5.49)
\end{aligned}$$

The remaining four equations (5.33)–(5.36) combine with the one above to form three additional equations that $i\Upsilon$ satisfies. The even equation (5.33) can be regarded as giving rise to another even equation,

$$\begin{aligned}
&\left[(4y-y^2) \frac{\partial^2}{\partial y^2} + (D+2)(2-y) \frac{\partial}{\partial y} - 2 \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right] \left(\frac{\partial}{\partial u} + \frac{D\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right. \\
&\quad \left. - \left(\frac{\partial}{\partial u} + \frac{D-1-\epsilon}{1-\epsilon} \right) \frac{\partial}{\partial u} + \left(4 \operatorname{sh}(v) \frac{\partial}{\partial y} - \frac{\partial}{\partial v} \right) \frac{\partial}{\partial v} + \frac{3-D(2-\epsilon)}{(1-\epsilon)^2} \right] i\Upsilon
\end{aligned}$$

$$\begin{aligned}
&= \frac{\text{sh}\left(\frac{\epsilon v}{1-\epsilon}\right)}{1-\epsilon} \left[2\epsilon \text{sh}(v) \frac{\partial}{\partial y} i\Delta_\nu - (1+\epsilon) \frac{\partial}{\partial v} i\Delta_\nu \right] + \frac{\text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right)}{1-\epsilon} \left[-2\epsilon \frac{\partial}{\partial y} i\Delta_{\nu+1} \right. \\
&\quad \left. - 2\epsilon \text{ch}(v) \frac{\partial}{\partial y} i\Delta_\nu + \left(\frac{\partial}{\partial u} + \frac{(D-2)\epsilon}{2(1-\epsilon)} \right) i\Delta_\nu + (1-\epsilon)\nu i\Delta_\nu \right] - \frac{\nu_T i\Delta_{\nu_T}}{1-\epsilon}, \quad (5.50)
\end{aligned}$$

the odd equation (5.34) as giving rise to another odd equation,

$$\begin{aligned}
&\left[\left(\left[y + 4 \text{sh}^2\left(\frac{v}{2}\right) \right] \frac{\partial}{\partial y} + \frac{\partial}{\partial u} + \frac{D-1-\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial v} - 2 \text{sh}(v) \left(\frac{\partial}{\partial u} + \frac{D\epsilon-2}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right] i\Upsilon \\
&= - \left[\frac{\epsilon \text{sh}(v)}{1-\epsilon} \text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right) + \text{ch}(v) \text{sh}\left(\frac{\epsilon v}{1-\epsilon}\right) \right] \frac{\partial}{\partial y} i\Delta_\nu \\
&\quad + \frac{\epsilon \text{sh}\left(\frac{\epsilon v}{1-\epsilon}\right)}{2(1-\epsilon)} \left[\frac{1}{\text{sh}(v)} \frac{\partial}{\partial v} i\Delta_{\nu+1} - \frac{\text{ch}(v)}{\text{sh}(v)} \frac{\partial}{\partial v} i\Delta_\nu - \nu i\Delta_\nu \right], \quad (5.51)
\end{aligned}$$

and the rising and lowering equations (5.35) and (5.35) as producing

$$\begin{aligned}
&\left(\frac{\partial}{\partial u} + \frac{(D-2)\epsilon}{2(1-\epsilon)} - \nu_T \right) \left(\frac{\partial}{\partial u} + \frac{(D-2)\epsilon}{2(1-\epsilon)} \right) i\Upsilon \\
&= \frac{1}{2} \left(\frac{\partial}{\partial u} + \frac{(D-2)\epsilon}{2(1-\epsilon)} - \nu_T \right) \left\{ \text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right) \left[\frac{1}{\text{sh}(v)} \frac{\partial}{\partial v} i\Delta_{\nu+1} - \frac{\text{ch}(v)}{\text{sh}(v)} \frac{\partial}{\partial v} i\Delta_\nu \right. \right. \\
&\quad \left. \left. + \left(\frac{\partial}{\partial u} + \frac{(D-2)\epsilon}{2(1-\epsilon)} \right) i\Delta_\nu \right] - \text{sh}\left(\frac{\epsilon v}{1-\epsilon}\right) \frac{\partial}{\partial v} i\Delta_\nu \right\}. \quad (5.52)
\end{aligned}$$

The association of the three equations above to their antecedents (5.33)–(5.36) is only qualitative since each of the new equations is a combination of essentially all of the antecedent ones. Additionally, in deriving the last equation (5.52) we used that,

$$\frac{\partial}{\partial u} i\Delta_{\nu_T} = -\frac{(D-2)\epsilon}{2(1-\epsilon)} i\Delta_{\nu_T}, \quad \frac{\partial}{\partial v} i\Delta_{\nu_T} = 0. \quad (5.53)$$

owing to the transverse sector index being $\nu_T < (D-1)/2$.

5.4 Covariantization

The components of the photon two-point function found in Sec. 5.2 can be written in a more systematic covariantized form,

$$\begin{aligned}
i[\mu^- \Delta_\nu^+](x; x') &= (\partial_\mu \partial'_\nu y) \mathcal{C}_1(y, u, v) + (\partial_\mu y) (\partial'_\nu y) \mathcal{C}_2(y, u, v) \\
&\quad + \left[(\partial_\mu y) (\partial'_\nu u) + (\partial_\mu u) (\partial'_\nu y) \right] \mathcal{C}_3(y, u, v) + \left[(\partial_\mu y) (\partial'_\nu u) - (\partial_\mu u) (\partial'_\nu y) \right] \bar{\mathcal{C}}_3(y, u, v) \\
&\quad + (\partial_\mu u) (\partial'_\nu u) \mathcal{C}_4(y, u, v), \quad (5.54)
\end{aligned}$$

where the elements of the appropriate bi-tensor basis for this form are constructed out of independent derivatives of the bi-scalar variables (2.8), and are multiplied by

five scalar structure functions. This is the tensor basis from [73] supplemented by the odd tensor structure with an accompanying structure function $\bar{\mathcal{C}}_3$. We need to determine the structure functions by matching the form (5.54) to the results in (5.16)–(5.18) and (5.23). This is facilitated by writing out explicitly the components of the covariantized form,

$$i[\bar{i}\Delta_j^+](x; x') = 2\delta_{ij}(1-\epsilon)^2\mathcal{H}\mathcal{H}'\left\{-\mathcal{C}_1 + I[\mathcal{C}_2]\right\} + \partial_i\partial'_j I^2[\mathcal{C}_2], \quad (5.55)$$

$$i[\bar{0}\Delta_i^+](x; x') = (1-\epsilon)\mathcal{H}(\partial'_iy)\left\{\mathcal{C}_1 - [2-y-2e^{-v}]\mathcal{C}_2 + \mathcal{C}_3 - \bar{\mathcal{C}}_3\right\}, \quad (5.56)$$

$$i[\bar{i}\Delta_0^+](x; x') = (1-\epsilon)\mathcal{H}'(\partial_iy)\left\{\mathcal{C}_1 - [2-y-2e^v]\mathcal{C}_2 + \mathcal{C}_3 + \bar{\mathcal{C}}_3\right\}, \quad (5.57)$$

$$i[\bar{0}\Delta_0^+](x; x') = (1-\epsilon)^2\mathcal{H}\mathcal{H}'\left\{-[2-y-4\text{ch}(v)]\mathcal{C}_1 - [4y-y^2-8+4(2-y)\text{ch}(v)]\mathcal{C}_2 - 2[2-y-2\text{ch}(v)]\mathcal{C}_3 - 4\text{sh}(v)\bar{\mathcal{C}}_3 + \mathcal{C}_4\right\}. \quad (5.58)$$

A straightforward comparison then yields the structure functions,

$$\mathcal{C}_1 = \frac{e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T(1-\epsilon)^2H_0^2}\left\{-\nu_T i\Delta_{\nu_T} - \frac{1}{2}\text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right)i\Delta_\nu + i\Upsilon\right\}, \quad (5.59)$$

$$\mathcal{C}_2 = \frac{e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T(1-\epsilon)^2H_0^2}\left\{-\frac{1}{2}\text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right)\frac{\partial}{\partial y}i\Delta_\nu + \frac{\partial}{\partial y}i\Upsilon\right\}, \quad (5.60)$$

$$\mathcal{C}_3 = \frac{e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T(1-\epsilon)^2H_0^2}\left\{\frac{\nu_T}{2}i\Delta_{\nu_T} + \frac{1}{2\text{sh}(v)}\text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right)\left[\frac{\partial}{\partial v}i\Delta_{\nu+1} - \text{ch}(v)\frac{\partial}{\partial v}i\Delta_\nu\right] - \frac{(D-3)}{4}\text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right)i\Delta_\nu - \frac{1}{2}\text{sh}\left(\frac{\epsilon v}{1-\epsilon}\right)\frac{\partial}{\partial v}i\Delta_\nu + \frac{(D-3)}{2}i\Upsilon\right\}, \quad (5.61)$$

$$\bar{\mathcal{C}}_3 = \frac{e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T(1-\epsilon)^2H_0^2}\left\{-\text{sh}\left(\frac{\epsilon v}{1-\epsilon}\right)\frac{\partial}{\partial y}\left[i\Delta_{\nu+1} + \text{ch}(v)i\Delta_\nu\right] - 2\text{sh}(v)\frac{\partial}{\partial y}i\Upsilon\right\}, \quad (5.62)$$

$$\mathcal{C}_4 = \frac{e^{\frac{\epsilon u}{1-\epsilon}}}{2\nu_T(1-\epsilon)^2H_0^2}\left\{\text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right)\left[\frac{\partial}{\partial u} + \frac{[2-y-2\text{ch}(v)]}{2\text{sh}(v)}\frac{\partial}{\partial v}\right]i\Delta_{\nu+1} + \frac{1}{2}\text{ch}\left(\frac{\epsilon v}{1-\epsilon}\right)\left[(2-y)\left(\frac{\partial}{\partial u} + 2\nu - D + 3\right) - \frac{[(2-y)\text{ch}(v)-2]}{\text{sh}(v)}\frac{\partial}{\partial v}\right]i\Delta_\nu - \text{sh}\left(\frac{\epsilon v}{1-\epsilon}\right)\frac{\partial}{\partial v}\left[(2-y)i\Delta_\nu + 2i\Delta_{\nu+1}\right] + \left[(4y-y^2)\frac{\partial}{\partial y} + (D-2)(2-y)\right]i\Upsilon\right\}, \quad (5.63)$$

where in simplifying the final expressions we made use of generalized recurrence relations (2.53b) and (2.53a) for scalar two-point functions, the reduction formula (5.49), and addition formulas for hyperbolic functions. These structure functions are the main result of our paper, as they determine the photon two-point function in the simple covariant gauge (4.31) in the covariant representation (5.54). The remaining

two-point functions defined in Sec. 5.1 are obtained from this result by changing the $i\delta$ -prescription for the bi-local variables in the structure functions and the tensor basis to the one appropriate according to (2.24) and (2.26).

Explicitly checking that our result for the photon two-point function (5.54) with solutions for the structure functions above satisfies both the equation of motion (5.4) and the Ward-Takahashi identity (5.5) is important, particularly in view of recent findings [79] that in de Sitter space a number of the results from the literature fails to satisfy the latter condition. Our result indeed satisfies all the necessary equations; the details necessary to perform the checks are given in Appendix C.

6 Various limits

In this section we derive several limits of the covariant form of the two-point function (5.54) with structure functions (5.59)–(5.63), and compare them to the literature where possible.

6.1 De Sitter limit

In the de Sitter limit the bi-local variables (2.8) simply take their values with $\epsilon = 0$, and accordingly so do the basis tensor introduced in (5.54). All the limits of the scalar two-point functions and derived quantities are manifestly finite in the de Sitter limit, due to the IR sum. Therefore, all the terms in structure functions (5.59)–(5.63) multiplied by ϵ automatically vanish in the de Sitter limit. The few terms that remain have simple limits. The mode function indices (4.15) and (4.22) from the transverse and scalar sectors become degenerate,

$$\nu, \nu_T \xrightarrow{\epsilon \rightarrow 0} \nu_0 = \frac{D-3}{2}. \quad (6.1)$$

The scalar two-point functions carrying those indices reduce to one and the same de Sitter invariant scalar two-point function,

$$i\Delta_\nu(y, u, v), i\Delta_{\nu_T}(y, u, v) \xrightarrow{\epsilon \rightarrow 0} \mathcal{F}_{\nu_0}(y), \quad (6.2)$$

Their derivatives with respect to y persist, but derivatives with respect to u and v vanish. For the Laplace-inverted two-point function (5.43) it is only relevant that the derivative with respect to u vanishes,

$$\frac{\partial}{\partial u} \frac{\partial}{\partial y} i\Xi_\nu(y, u, v) \xrightarrow{\epsilon \rightarrow 0} 0, \quad (6.3)$$

as this implies that all of them cancel completely in (5.24),

$$i\Upsilon(y, u, v) \xrightarrow{\epsilon \rightarrow 0} 0. \quad (6.4)$$

The limits above encompass all the terms appearing in (5.59)–(5.63), except for few instances where the scalar propagator with the index $\nu+1$ appears. Because in de Sitter limit we have for the index $\nu+1 \xrightarrow{\epsilon \rightarrow 0} \nu_0+1 = (D-1)/2$, the two-point function will reduce to the MMCS two-point function, which is known not to be de Sitter invariant. All instances when it appears include derivatives, which are evaluated using the form given in Sec. 2.3.2,

$$\frac{\partial}{\partial u} i\Delta_{\nu+1}(y, u, v) \xrightarrow{\epsilon \rightarrow 0} \frac{H_0^{D-2}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(D-1)}{\Gamma(\frac{D}{2})}, \quad \frac{\partial}{\partial v} i\Delta_{\nu+1}(y, u, v) \xrightarrow{\epsilon \rightarrow 0} 0. \quad (6.5)$$

Finally, taking the de Sitter value of the simple covariant gauge,

$$\xi_s \xrightarrow{\epsilon \rightarrow 0} \xi_s^0 = \frac{\nu_0+1}{\nu_0} = \frac{D-1}{D-3}, \quad (6.6)$$

leads to the de Sitter limit of structure functions,

$$\begin{aligned} \mathcal{C}_1 &\xrightarrow{\epsilon \rightarrow 0} \frac{1}{2\nu_0 H_0^2} \left[-\left(\nu_0 + \frac{1}{2}\right) \mathcal{F}_{\nu_0}(y) \right], & \mathcal{C}_2 &\xrightarrow{\epsilon \rightarrow 0} \frac{1}{2\nu_0 H_0^2} \left[-\frac{1}{2} \frac{\partial}{\partial y} \mathcal{F}_{\nu_0}(y) \right], \\ \mathcal{C}_3 &\xrightarrow{\epsilon \rightarrow 0} 0, & \bar{\mathcal{C}}_3 &\xrightarrow{\epsilon \rightarrow 0} 0, & \mathcal{C}_4 &\xrightarrow{\epsilon \rightarrow 0} \xi_s^0 \times \frac{H_0^{D-4}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(D-1)}{(D-1)\Gamma(\frac{D}{2})}. \end{aligned} \quad (6.7)$$

These correctly reproduce the de Sitter limit, including the de Sitter breaking non-vanishing structure function \mathcal{C}_4 recently found in [74, 75]. The latter is here a consequence of the first expression in (6.5) not vanishing, that is due to the non-existence of IR finite de Sitter invariant two-point functions for MMCS in the CTBD state [96, 97].

While the covariant gauge two-point function has a relatively simple de Sitter limit, compared to its form in power-law inflation, no such simplifications happen for the physical two-point function (5.25),

$$i[\bar{\Delta}_j^+]^{\text{ph}}(x; x') \xrightarrow{\epsilon \rightarrow 0} e^u \left[\delta_{ij} i\Delta_{\nu_0}(y, u, v) + \partial_i \partial'_j \frac{i\bar{\Xi}_{\nu_0}(y, u, v)}{\mathcal{H}\mathcal{H}'} \right], \quad (6.8)$$

since $i\bar{\Xi}_{\nu_\tau}$ retains its form (5.45) without any simplifications. It is really the transverse and the longitudinal contributions to the (ij) component of the two-point function that conspire so that the complicated parts cancel between them. In practice this is due to two non-commensurate mode function indices reducing to one and the same in the de Sitter limit (6.1).

6.2 Flat space limit

In the Minkowski space limit, $H_0 \rightarrow 0$, the three bi-local variables (2.8) reduce to,

$$y_{\text{ab}} \stackrel{H_0 \rightarrow 0}{\sim} (1-\epsilon)^2 H_0^2 \Delta x_{\text{ab}}^2, \quad u_{\text{ab}} \stackrel{H_0 \rightarrow 0}{\sim} (1-\epsilon) H_0 (\eta + \eta' - 2\eta_0), \quad v_{\text{ab}} \stackrel{H_0 \rightarrow 0}{\sim} (1-\epsilon) H_0 \Delta \eta_{\text{ab}}, \quad (6.9)$$

and accordingly the tensor structures reduce to,

$$(\partial_\mu \partial'_\nu y) \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} -2[(1-\epsilon)H_0]^2 \eta_{\mu\nu}, \quad (6.10a)$$

$$(\partial_\mu y)(\partial'_\nu y) \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} -4[(1-\epsilon)H_0]^4 \Delta x_\mu \Delta x_\nu, \quad (6.10b)$$

$$\left[(\partial_\mu y)(\partial'_\nu u) \pm (\partial_\mu u)(\partial'_\nu y) \right] \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} 2[(1-\epsilon)H_0]^3 (\Delta x_\mu \delta_\nu^0 \mp \delta_\mu^0 \Delta x_\nu), \quad (6.10c)$$

$$(\partial_\mu u)(\partial'_\nu u) \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} [(1-\epsilon)H_0]^2 \delta_\mu^0 \delta_\nu^0, \quad (6.10d)$$

where $\Delta x_\mu = x_\mu - x'_\mu$. For derivatives appearing in the scalar structure functions we have,

$$\frac{\partial}{\partial y} \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} \frac{1}{(1-\epsilon)^2 H_0^2} \frac{\partial}{\partial(\Delta x^2)}, \quad \frac{\partial}{\partial v} \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} \frac{1}{(1-\epsilon)H_0} \frac{\partial}{\partial \Delta \eta}. \quad (6.11)$$

The scalar two-point functions all reduce to the massless scalar two-point function,

$$i\Delta_\lambda(y, u, v) \stackrel{H_0 \rightarrow 0}{\longrightarrow} \frac{\Gamma(\frac{D-2}{2})}{4\pi^{\frac{D}{2}} (\Delta x^2)^{\frac{D-2}{2}}} \equiv i\check{\Delta}(\Delta x^2), \quad (6.12)$$

and its derivatives with respect to remaining bi-local variables reduce to,

$$\frac{\partial}{\partial u} i\Delta_\lambda(y, u, v) \stackrel{H_0 \rightarrow 0}{\longrightarrow} -\frac{(D-2)\epsilon}{2(1-\epsilon)} i\check{\Delta}(\Delta x^2), \quad \frac{\partial}{\partial v} i\Delta_\lambda(y, u, v) \stackrel{H_0 \rightarrow 0}{\longrightarrow} 0. \quad (6.13)$$

Note that here we additionally have to assume that $k_0 \rightarrow 0$, such that $k_0/H_0 \ll 1$ remains satisfied. Using these we can also infer the flat space limit of (5.43),

$$\begin{aligned} \frac{\partial}{\partial y} i\Xi_\lambda(y, u, v) &\stackrel{H_0 \rightarrow 0}{\longrightarrow} \sum_{n=0}^{\infty} \frac{(-1)^n \Gamma(\frac{D-1}{2})}{4\Gamma(\frac{D+1}{2}+n)} (\Delta x^2 + \Delta \eta^2)^n \frac{\partial^n}{\partial(\Delta x^2)^n} i\check{\Delta}(\Delta x^2) \\ &= \sum_{n=0}^{\infty} \frac{\Gamma(\frac{D-1}{2}) \Gamma(\frac{D-2}{2}+n)}{4\Gamma(\frac{D+1}{2}+n)} \frac{(\Delta x^2 + \Delta \eta^2)^n}{4\pi^{\frac{D}{2}} (\Delta x^2)^{\frac{D-2}{2}+n}} \\ &= \frac{i\check{\Delta}(\Delta x^2)}{2(D-1)} \times {}_2F_1\left(\left\{1, \frac{D-2}{2}\right\}, \left\{\frac{D+1}{2}\right\}, 1 + \frac{\Delta \eta^2}{\Delta x^2}\right) \equiv \frac{\partial}{\partial(\Delta x^2)} i\check{\Xi}(\Delta x^2, \Delta \eta), \end{aligned} \quad (6.14)$$

and of its derivative with respect to u ,

$$\frac{\partial}{\partial u} \frac{\partial}{\partial y} i\Xi_\lambda(y, u, v) \stackrel{H_0 \rightarrow 0}{\longrightarrow} -\frac{(D-2)\epsilon}{2(1-\epsilon)} \frac{\partial}{\partial(\Delta x^2)} i\check{\Xi}(\Delta x^2, \Delta \eta). \quad (6.15)$$

It is then straightforward to derive the equation that (6.14) satisfies,

$$\left[2(\Delta \eta)^2 \frac{\partial}{\partial(\Delta x^2)} - \Delta \eta \frac{\partial}{\partial \Delta \eta} + 1 \right] \frac{\partial}{\partial(\Delta x^2)} i\check{\Xi}(\Delta x^2, \Delta \eta) = \frac{1}{2} i\check{\Delta}(\Delta x^2), \quad (6.16)$$

which, together with identities (6.14) and (6.16), implies the flat space limit of (5.24),

$$i\Upsilon(y, u, v) \xrightarrow{H_0 \rightarrow 0} \frac{-\epsilon}{(1-\epsilon)} i\check{\Delta}(\Delta x^2). \quad (6.17)$$

Given all the listed flat space limits, it is only the first two structure functions out of (5.59)–(5.63) that contribute to the flat space limit,

$$i[{}_{\mu}^{-}\Delta_{\nu}^{+}](x; x') \xrightarrow{H_0 \rightarrow 0} \left[\eta_{\mu\nu} - \frac{(1-\xi_s)}{2} \left(\eta_{\mu\nu} - (D-2) \frac{\Delta x_{\mu} \Delta x_{\nu}}{\Delta x^2} \right) \right] i\check{\Delta}(\Delta x^2), \quad (6.18)$$

that can be written in a form more often encountered,

$$i[{}_{\mu}^{-}\Delta_{\nu}^{+}](x; x') \xrightarrow{H_0 \rightarrow 0} \left[\eta_{\mu\nu} - (1-\xi_s) \frac{\partial_{\mu} \partial_{\nu}}{\partial^2} \right] i\check{\Delta}(\Delta x^2), \quad (6.19)$$

where the gauge-fixing parameter corresponding to the simple covariant gauge defined in (4.31). The two-point function in (6.18) of course contains the correct $i\delta$ -prescription [116], inherited from the bi-local variables adapted for FLRW.

The flat-space limit of the physical two-point function (5.25) is inferred from the flat space limits of the scalar two-point function (6.12) and of the Laplace-inverted two-point function (B.30),

$$i[{}_i^{-}\Delta_j^{+}]^{\text{ph}}(x; x') \xrightarrow{H_0 \rightarrow 0} \frac{\Gamma(\frac{D-2}{2})}{4\pi^{\frac{D}{2}}} \left[\frac{\delta_{ij}}{(\Delta x^2)^{\frac{D-2}{2}}} - \frac{(-\Delta\eta^2)^{\frac{4-D}{2}}}{2(D-3)(D-4)} \times \right. \\ \left. \times \partial_i \partial'_j {}_2F_1 \left(\left\{ \frac{D-3}{2}, \frac{D-4}{2} \right\}, \left\{ \frac{D-1}{2} \right\}, 1 + \frac{\Delta x^2}{\Delta\eta^2} \right) \right]. \quad (6.20)$$

It corresponds to the Coulomb gauge two-point function in D -dimensional Minkowski space. Of course, in four spacetime dimensions, due to conformal invariance, the Coulomb gauge two-point function reduces to the flat space expression [117]. In position space this expression reads,

$$i[{}_i^{-}\Delta_j^{+}]^{\text{ph}}(x; x') \xrightarrow{D \rightarrow 4} \frac{1}{4\pi^2} \left\{ \frac{\delta_{ij}}{\Delta x^2} + \frac{\partial_i \partial'_j}{2} \left[\ln(\mu^2 \Delta x^2) + \frac{\Delta\eta}{\|\Delta\vec{x}\|} \ln \left(\frac{\Delta\eta + \|\Delta\vec{x}\|}{\Delta\eta - \|\Delta\vec{x}\|} \right) \right] \right\}. \quad (6.21)$$

Note that the seeming $1/(D-4)$ divergence is removed by the spatial derivatives in the second line of (6.20). Also note that the dependence on an arbitrary scale μ drops out of the expression due to spatial derivatives, and its role is just to make the argument of the logarithm dimensionless.

7 Simple observables

Our propagator in the simple covariant gauge given in (5.54) with structure functions (5.59)–(5.63) can now be used in loop computations. As a simple consistency check here we consider two simplest observables: the tree-level field strength correlators, and the one-loop energy-momentum tensor.

7.1 Field strength correlator

The tree-level correlator of the field stress tensor,

$$\langle \Omega | \hat{F}_{\mu\nu}(x) \hat{F}_{\rho\sigma}(x') | \Omega \rangle = 4(\delta_{[\mu}^{\alpha} \partial_{\nu]}) (\delta_{[\rho}^{\beta} \partial'_{\sigma]}) i [{}_{\alpha}^{-} \Delta_{\beta}^{+}] (x; x') \quad (7.1)$$

is gauge independent and constitutes an observable. It can thus serve as a simple consistency check for the photon two-point function we found in Sec 5. Acting the derivatives in (7.1) onto the covariantized representation of the two-point function (5.54) organizes itself in the following tensor basis [73],

$$\begin{aligned} \langle \Omega | \hat{F}_{\mu\nu}(x) \hat{F}_{\rho\sigma}(x') | \Omega \rangle &= (\partial_{\mu} \partial'_{\rho} y) (\partial'_{\sigma} \partial_{\nu} y) \mathcal{G}_1 + (\partial_{[\mu} y) (\partial_{\nu]} \partial'_{[\sigma} y) (\partial'_{\rho]} y) \mathcal{G}_2 \\ &+ \left[(\partial_{[\mu} y) (\partial_{\nu]} \partial'_{[\sigma} y) (\partial'_{\rho]} u) + (\partial_{[\mu} u) (\partial_{\nu]} \partial'_{[\sigma} y) (\partial'_{\rho]} y) \right] \mathcal{G}_3 \\ &+ \left[(\partial_{[\mu} y) (\partial_{\nu]} \partial'_{[\sigma} y) (\partial'_{\rho]} u) - (\partial_{[\mu} u) (\partial_{\nu]} \partial'_{[\sigma} y) (\partial'_{\rho]} y) \right] \bar{\mathcal{G}}_3 \\ &+ (\partial_{[\mu} u) (\partial_{\nu]} \partial'_{[\sigma} y) (\partial'_{\rho]} u) \mathcal{G}_4 + (\partial_{[\mu} y) (\partial_{\nu]} u) (\partial'_{[\rho} y) (\partial'_{\sigma]} u) \mathcal{G}_5, \end{aligned} \quad (7.2)$$

where the structure functions are expressed in terms of structure functions (5.59)–(5.63) of the photon two-point function,

$$\begin{aligned} \mathcal{G}_1 &= 4 \left(\frac{\partial \mathcal{C}_1}{\partial y} - \mathcal{C}_2 \right), & \mathcal{G}_2 &= \frac{\partial \mathcal{G}_1}{\partial y}, & \mathcal{G}_3 &= \frac{\partial \mathcal{G}_1}{\partial u}, & \bar{\mathcal{G}}_3 &= -\frac{\partial \mathcal{G}_1}{\partial v}, & (7.3) \\ \mathcal{G}_4 &= 4 \left(\frac{\partial^2 \mathcal{C}_1}{\partial u^2} - \frac{\partial^2 \mathcal{C}_1}{\partial v^2} - 2 \frac{\partial \mathcal{C}_3}{\partial u} - 2 \frac{\partial \bar{\mathcal{C}}_3}{\partial v} + \frac{\partial \mathcal{C}_4}{\partial y} \right), & \mathcal{G}_5 &= \frac{\partial \mathcal{G}_4}{\partial y} - \frac{\partial^2 \mathcal{G}_1}{\partial u^2} + \frac{\partial^2 \mathcal{G}_1}{\partial v^2}. \end{aligned}$$

Evaluating these structure functions is accomplished by judiciously applying equations (5.49)–(5.52) for the $i\Upsilon$ function defined in (5.24), as well as equations of motion (2.50) and (2.51), and recurrence relations (2.53) for the scalar two-point functions,

$$\mathcal{G}_1 = \frac{2 e^{-\frac{(D-4)\epsilon}{2(1-\epsilon)} u}}{(1-\epsilon)^2 H_0^2} \left[-\frac{\partial \mathcal{F}_{\nu_T}}{\partial y} \right], \quad (7.4a)$$

$$\mathcal{G}_2 = \frac{2 e^{-\frac{(D-4)\epsilon}{2(1-\epsilon)} u}}{(1-\epsilon)^2 H_0^2} \left[-\frac{\partial^2 \mathcal{F}_{\nu_T}}{\partial y^2} \right], \quad (7.4b)$$

$$\mathcal{G}_3 = \frac{2 e^{-\frac{(D-4)\epsilon}{2(1-\epsilon)} u}}{(1-\epsilon)^2 H_0^2} \left[\frac{(D-4)\epsilon}{2(1-\epsilon)} \frac{\partial \mathcal{F}_{\nu_T}}{\partial y} \right], \quad (7.4c)$$

$$\bar{\mathcal{G}}_3 = 0, \quad (7.4d)$$

$$\mathcal{G}_4 = \frac{2 e^{-\frac{(D-4)\epsilon}{2(1-\epsilon)} u}}{(1-\epsilon)^2 H_0^2} \left[\frac{(D-4)\epsilon}{2(1-\epsilon)} \left(1 - \frac{(D-4)\epsilon}{2(1-\epsilon)} \right) \mathcal{F}_{\nu_T} \right], \quad (7.4e)$$

$$\mathcal{G}_5 = \frac{2 e^{-\frac{(D-4)\epsilon}{2(1-\epsilon)} u}}{(1-\epsilon)^2 H_0^2} \left[\frac{(D-4)\epsilon}{2(1-\epsilon)} \frac{\partial \mathcal{F}_{\nu_T}}{\partial y} \right]. \quad (7.4f)$$

The end result reveals that the only contributions come from the transverse sector. That means our two-point function produces a gauge independent result for this observable, as it should. Furthermore, the $D \rightarrow 4$ limit reduces to the flat space vacuum result,

$$\langle \Omega | \hat{F}_{\mu\nu}(x) \hat{F}_{\rho\sigma}(x') | \Omega \rangle \xrightarrow{D \rightarrow 4} \frac{2}{\pi^2 (\Delta x^2)^2} \left[\eta_{\mu[\rho} \eta_{\sigma]\nu} - 4 \eta_{\alpha[\mu} \eta_{\nu][\sigma} \eta_{\rho]\beta} \frac{\Delta x^\alpha \Delta x^\beta}{\Delta x^2} \right]. \quad (7.5)$$

This is a manifestation of the conformal coupling of the photon to gravity in $D=4$, which allows for a natural choice of the photon vacuum state that does not sense the expansion. The limit (7.5) confirms that we have chosen precisely such a state for the photon.

7.2 Energy-momentum tensor

The energy-momentum tensor can be given by two different definitions: either as a variation of the gauge invariant Maxwell action (3.1), or as a variation of the gauge-fixed action (3.10). This is the case both in the classical and the quantum theory. However, it is sufficient to consider just the former definition,

$$T_{\mu\nu}(x) = \frac{-2}{\sqrt{-g}} \frac{\delta S}{\delta g^{\mu\nu}(x)} = \left(\delta_\mu^\rho \delta_\nu^\sigma - \frac{1}{4} g_{\mu\nu} g^{\rho\sigma} \right) g^{\alpha\beta} F_{\rho\alpha}(x) F_{\sigma\beta}(x), \quad (7.6)$$

as the latter differs by the contribution of the gauge-fixing part of the action that is guaranteed to vanish on-shell. In the classical theory this is a simple consequence of the two constraints (3.12) vanishing on-shell. In the quantized theory the contribution of the gauge-fixing part can be seen to vanish on-shell in two ways: (i) either due to the proper operator ordering and to the subsidiary condition (3.16) [79], or (ii) due to the cancellations between the gauge-fixing part and the Faddeev-Popov ghost part that has to be added if all the operators are Weyl-ordered [75, 118, 119].

Defining an operator associated to (7.6) is straightforward, since when expressed in terms of the canonical fields all the terms are composed either solely of transverse fields, or solely of constraints. Therefore, we may define the operator to be Weyl-ordered, and the expectation value essentially reduces to the coincident field strength correlator,

$$\langle \Omega | \hat{T}_{\mu\nu}(x) | \Omega \rangle = \left(\delta_\mu^\rho \delta_\nu^\sigma - \frac{1}{4} g_{\mu\nu} g^{\rho\sigma} \right) g^{\alpha\beta} \langle \Omega | \hat{F}_{\rho\alpha}(x) \hat{F}_{\sigma\beta}(x) | \Omega \rangle. \quad (7.7)$$

Computing the coincident correlator amounts to computing a dimensionally regulated coincidence limit of (7.1). This is best done in the tensor basis (7.2) where the only non-vanishing tensor structures in this limit are the first and the fourth one,

$$(\partial_\mu \partial'_{[\rho} y) (\partial'_{\sigma]} \partial_\nu y) \xrightarrow{x' \rightarrow x} 4 [(1-\epsilon)H]^4 g_{\mu[\rho} g_{\sigma]\nu}, \quad (7.8)$$

$$(\partial_{[\mu} u) (\partial_{\nu]} \partial'_{[\sigma} y) (\partial'_{\rho]} u) \xrightarrow{x' \rightarrow x} -2 [(1-\epsilon)H]^4 (a^2 \delta_{[\mu}^0 g_{\nu][\sigma} \delta_{\rho]}^0), \quad (7.9)$$

and where the dimensionally regulated coincident limits of the corresponding structure functions in (7.2) are inferred from the power-series representation (2.32),

$$\mathcal{G}_1 \xrightarrow{x' \rightarrow x} \frac{[(1-\epsilon)H]^{D-4}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(\frac{D+1}{2} + \nu_T) \Gamma(\frac{D+1}{2} - \nu_T) \Gamma(\frac{2-D}{2})}{\Gamma(\frac{1}{2} + \nu_T) \Gamma(\frac{1}{2} - \nu_T) (-D)} \xrightarrow{D \rightarrow 4} \frac{1}{32\pi^2(1-\epsilon)}, \quad (7.10)$$

$$\begin{aligned} \mathcal{G}_4 \xrightarrow{x' \rightarrow x} & \frac{[(1-\epsilon)H]^{D-4}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(\frac{D-1}{2} + \nu_T) \Gamma(\frac{D-1}{2} - \nu_T)}{\Gamma(\frac{1}{2} + \nu_T) \Gamma(\frac{1}{2} - \nu_T)} \\ & \times \left[1 - \frac{(D-4)\epsilon}{2(1-\epsilon)} \right] \frac{4\epsilon \Gamma(\frac{6-D}{2})}{(D-2)(1-\epsilon)} \xrightarrow{D \rightarrow 4} 0. \end{aligned} \quad (7.11)$$

These limits are finite in $D=4$, and thus no counterterms are necessary for renormalization. Plugging the coincident correlator into (7.7) and performing the remaining contractions finally gives a vanishing result for the gauge-invariant energy-momentum tensor,

$$\langle \Omega | \hat{T}_{\mu\nu}(x) | \Omega \rangle = 0. \quad (7.12)$$

There is still the conformal anomaly contributing to the energy-momentum tensor at one loop [120–122], that is not captured by the result (7.12). In fact, for conformally coupled fields in conformally flat backgrounds the conformal anomaly contribution to the energy-momentum tensor is the only nonvanishing one [123], and its unambiguous part is gauge independent [124]. In that sense the result in (7.12) is a consistency check of the two-point function we computed. The conformal anomaly contribution, however, does not appear due to divergences in the one-loop diagram corresponding to the source for the graviton tadpole, which is the energy-momentum tensor. As a matter of fact, there are no logarithmic divergences in that diagram, as we have shown in this section. The conformal anomaly contribution appears due to divergences found in other one-loop diagrams of the theory, and to recover it one would need to renormalize the effective action, rather than just the particular diagram corresponding to the observable we consider here.

8 Discussion

Understanding how large are the effects of quantum loop corrections to inflationary observables is of indisputable importance. In this work we were concerned with the building blocks necessary for quantifying these corrections in theories containing gauge vector fields in power-law inflation. Two-point functions of free quantum fields — propagators — are the basic ingredients for computing Feynman diagrams representing loop corrections. This motivated us to consider the photon propagator for power-law inflation in general covariant gauges. The main result of this work is the D -dimensional position space two-point function for the photon in power-law inflation. We have presented it in the covariantized form (5.54) with structure

functions given in (5.59)–(5.63). We have computed this two-point function in the simple covariant gauge (4.31) because only for this particular choice are the photon two-point functions (4.32) and (4.33) simple enough for the evaluation of the Fourier integrals in (5.7)–(5.10) to be feasible. Our two-point function satisfies both the equations of motion (5.4), and the Ward-Takahashi identity (5.5) that follows from the canonical quantization recalled in Sec. 3 and Sec. 4. The detailed checks of these are given in Appendix C. Furthermore, our result correctly reproduces the flat space limit, and the de Sitter space limit, including the de Sitter breaking term [74, 75]. Both of these limits are worked out in Sec. 6.

The complexity of the final result for the photon propagator is somewhat unexpected. The experience with non-minimally coupled massless scalar fields suggests that scalar mode functions, and consequently scalar two-point functions are just as complicated in power-law inflation as they are in de Sitter. The expectation was the same for photons, for which this turns out not to be the case. The complications descend from the fact that equations of motion couple the components of the vector potential in a way that in general no longer yields simple CTBD scalar mode functions and their derivatives as solutions. This is seen in Sec. 4.2 where we give the mode function solutions for the scalar sector of the vector potential, that are worked out in the accompanying appendix A. Only for a special choice of the gauge-fixing parameter $\xi = \xi_s$ in (4.31), that we dubbed the simple covariant gauge, do the mode functions simplify to the well known CTBD ones given in (2.11) and their derivatives. In this special case the momentum space two-point function of the photon retains the level of complexity it has in de Sitter.

In position space, on the other hand, even the simple covariant gauge two-point function takes on a considerably more complex form in power-law inflation than it does in the de Sitter space limit [74]. Here one is required to explicitly evaluate the inverse Laplace operator acting on the scalar two-point function. In Sec. 5.3 we have evaluated this Laplace-inverted two-point function in terms of a series (5.45) of higher and higher derivatives acting in the scalar two-point function, that is appropriate for sub-Hubble separations. In the accompanying appendix B the closed form expressions for this object are found in terms of Appell’s fourth function. This function is a particular instance of a non-factorizable double hypergeometric series, and its analytical structure is much more complicated than the structure of the hypergeometric function that describes scalar propagators. Incidentally, evaluating the Laplace-inverted two-point function explicitly is also necessary to compute the Coulomb gauge photon two-point function in position space, and we also report this result in (5.25).

The penultimate section of the main text is devoted to checking that our two-point function correctly reproduces two simple observables, namely the field strength correlator (7.1), and the energy-momentum tensor (7.7). It is confirmed that only the transverse sector contributes to these observables, and that the gauge sector

completely drops out. It is straightforward to demonstrate such manifest gauge independence in these two simple examples. This is because the lowest orders of the two observables are composed of a single photon two-point function only. In general this is not the case for more complicated loop computations. Our result passes all the tests before attempting these. The practical problem, however, might be the considerable complexity that our power-law inflation two-point function exhibits, compared to its de Sitter counterpart. Another issue is the absence of a free gauge-fixing parameter in the simple covariant gauge, which precludes manifest checks of gauge-independence of the final results when computing observables. However, gauge independence *can* be checked when combined with computation using the propagator in another gauge, such as the Coulomb gauge two-point function we also worked out in (5.25).

In conclusion, one can use our photon two-point function in the simple covariant gauge for loop computations in power-law inflation. However, it might be advisable to first examine whether different linear non-covariant gauges [79] lead to more tractable photon two-point functions, and consequently to simpler loop computations. It is also worth pointing out that similar complications as seen here are expected if attempting to construct the graviton propagator in covariant gauges in power-law inflation, for which no results are known (cf. [99]).

Acknowledgements

We are grateful to Richard P. Woodard for explaining to us the origins of conformal anomaly in dimensional regularization; to Igor Khavkine for helpful discussions about the analytic structure of Appell's functions; and to José L. López for looking into asymptotic expansions of Appell's first function and sending us preliminary results. DG was supported by the European Union and the Czech Ministry of Education, Youth and Sports (Project: MSCA Fellowship CZ FZU I — CZ.02.01.01/00/22_010/0002906). This work is part of the Delta ITP consortium, a program of the Netherlands Organisation for Scientific Research (NWO) that is funded by the Dutch Ministry of Education, Culture and Science (OCW) — NWO project number 24.001.027.

A Particular mode functions

Here we solve equations (4.29) and (4.30) for the particular mode functions for an arbitrary gauge-fixing parameter ξ . After shifting the mode functions,

$$v_0 = \frac{-i\xi k}{2[\nu(1-\epsilon)+1]} \left[\frac{a^2}{\mathcal{H}} U_{\nu+1} - \frac{1}{H_0} U_\nu \right] + \left(1 - \frac{\xi}{\xi_s}\right) w_0, \quad (\text{A.1})$$

$$v_L = \frac{-i\xi k}{2[\nu(1-\epsilon)+1]} \left[\frac{a^2}{\mathcal{H}} U_\nu - \frac{1}{H_0} U_{\nu+1} \right] + \left(1 - \frac{\xi}{\xi_s}\right) w_L, \quad (\text{A.2})$$

and applying identities (2.15) and (2.17), these equations read,

$$\left[\partial_0^2 + k^2 - \left(\nu^2 - \frac{1}{4}\right)(1-\epsilon)^2 \mathcal{H}^2 \right] w_0 = a^2 k^2 U_\nu, \quad (\text{A.3})$$

$$w_L = \frac{i}{k} \left[\partial_0 + \left(\nu + \frac{1}{2}\right)(1-\epsilon) \mathcal{H} \right] w_0. \quad (\text{A.4})$$

We solve the first equation using the retarded Green's function,

$$G_R(\eta; \eta') = \theta(\eta - \eta') G(\eta; \eta'), \quad G(\eta; \eta') = i \left[U_\nu(\eta, k) U_\nu^*(\eta', k) - U_\nu^*(\eta, k) U_\nu(\eta', k) \right], \quad (\text{A.5})$$

so that the solution for w_0 takes the form,

$$w_0(\eta, k) = k^2 \int_{\eta_0}^{\eta} d\eta' G(\eta; \eta') a^2(\eta') U_\nu(\eta', k) + \mathcal{A}(\epsilon, H_0, k) U_\nu(\eta, k) - \mathcal{B}(\epsilon, H_0, k) U_\nu^*(\eta, k) = \mathcal{Q}(\eta, k) U_\nu(\eta, k) - \tilde{\mathcal{Q}}(\eta, k) U_\nu^*(\eta, k), \quad (\text{A.6})$$

where the coefficient functions are,

$$\mathcal{Q}(\eta, k) = ik^2 \int_{\eta_0}^{\eta} d\eta' a^2(\eta') U_\nu^*(\eta', k) U_\nu(\eta', k) + \check{\mathcal{A}}(\epsilon, H_0, k), \quad (\text{A.7})$$

$$\tilde{\mathcal{Q}}(\eta, k) = ik^2 \int_{\eta_0}^{\eta} d\eta' a^2(\eta') U_\nu(\eta', k) U_\nu(\eta', k) + \check{\mathcal{B}}(\epsilon, H_0, k), \quad (\text{A.8})$$

with $\check{\mathcal{A}}$ and $\check{\mathcal{B}}$ arbitrary constants of integration. The solution for w_L follows from acting the derivatives in (A.4) onto the solution (A.6) for w_0 ,

$$w_L(\eta, k) = \mathcal{Q}(\eta, k) U_{\nu+1}(\eta, k) + \tilde{\mathcal{Q}}(\eta, k) U_{\nu+1}^*(\eta, k), \quad (\text{A.9})$$

using the fact that $G(\eta; \eta) = 0$. Therefore, the task of computing the particular mode functions is reduced to evaluating the two integrals from (A.7) and (A.8), and choosing the accompanying constants of integration. This is accomplished by making use of the integral 1.8.3.1. from [101],

$$\int_0^z dz' z'^{\rho-1} J_\lambda(z') J_\mu(z') = \frac{z^{\rho+\lambda+\mu}}{2^{\lambda+\mu} (\rho+\lambda+\mu) \Gamma(1+\lambda) \Gamma(1+\mu)} \times {}_3F_4 \left(\left\{ \frac{1+\lambda+\mu}{2}, \frac{2+\lambda+\mu}{2}, \frac{\rho+\lambda+\mu}{2} \right\}, \left\{ 1+\lambda, 1+\mu, 1+\lambda+\mu, \frac{2+\rho+\lambda+\mu}{2} \right\}, -z^2 \right), \quad (\text{A.10})$$

that is valid for $\text{Re}(\rho+\mu+\lambda) > 0$. In fact we need only two special cases of that result,

$$\begin{aligned} \int_0^z dz' z'^{\rho-1} J_\lambda(z') J_\lambda(z') &= \frac{z^{\rho+2\lambda}}{4^\lambda (\rho+2\lambda) \Gamma^2(1+\lambda)} \\ &\times {}_2F_3\left(\left\{\frac{\rho}{2}+\lambda, \frac{1}{2}+\lambda\right\}, \left\{\frac{2+\rho}{2}+\lambda, 1+\lambda, 1+2\lambda\right\}, -z^2\right) \\ &\equiv \frac{\mathcal{J}_1(\rho, \lambda; z)}{\Gamma(1+\lambda) \Gamma(1-\lambda) \Gamma(\lambda) \Gamma(-\lambda)}, \end{aligned} \quad (\text{A.11})$$

$$\begin{aligned} \int_0^z dz' z'^{\rho-1} J_\lambda(z') J_{-\lambda}(z') &= \frac{z^\rho}{\rho \Gamma(1+\lambda) \Gamma(1-\lambda)} {}_2F_3\left(\left\{\frac{\rho}{2}, \frac{1}{2}\right\}, \left\{\frac{2+\rho}{2}, 1+\lambda, 1-\lambda\right\}, -z^2\right) \\ &\equiv \frac{\mathcal{J}_2(\rho, \lambda; z)}{\Gamma(1+\lambda) \Gamma(1-\lambda) \Gamma(\lambda) \Gamma(-\lambda)}. \end{aligned} \quad (\text{A.12})$$

The integrals above are defined for a limited range of parameters only, the one in (A.11) for $\text{Re}(\rho+2\lambda) > 0$, and the one in (A.12) for $\text{Re}(\rho) > 0$, because of zero being the lower limit of integration. Nonetheless, we take the right-hand-sides to define the two functions \mathcal{J}_1 and \mathcal{J}_2 on an unlimited range of parameters. This detail is essentially immaterial for our purposes since the coefficient functions in (A.7) and (A.8) come with two integration constants. Using the results in (A.11) and (A.12), and the expression (2.11) for the mode function we can evaluate the coefficient functions,

$$\begin{aligned} \mathcal{Q}(\eta, k) &= \frac{i}{4\pi} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \left[-2 \cos(\pi\nu) \mathcal{J}_2\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) \right. \\ &\quad \left. + \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) + \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, -\nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) \right] + \mathcal{A}(\epsilon, H_0, k), \end{aligned} \quad (\text{A.13})$$

$$\begin{aligned} \tilde{\mathcal{Q}}(\eta, k) &= \frac{e^{\frac{-2ik}{(1-\epsilon)H_0}}}{4\pi} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \left[-2 \mathcal{J}_2\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) \right. \\ &\quad \left. + e^{-i\pi\nu} \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) + e^{i\pi\nu} \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, -\nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) \right] + \mathcal{B}(\epsilon, H_0, k), \end{aligned} \quad (\text{A.14})$$

up to the arbitrary integration constants, in which we have absorbed time independent contributions from the lower limit of integration in (A.7) and (A.8), and relabeled them to \mathcal{A} and \mathcal{B} . The two solutions for the coefficient functions above define our particular mode functions once the integration constants are chosen. In the following we fix them by considering three conditions: (i) the Wronskian-like relation (4.34), (ii) the de Sitter limit (4.36) computed in [74], and (iii) the Minkowski limit (4.38).

Wronskian relation. The Wronskian-like relation (4.34) for the shifted particular mode functions (A.1) and (A.2) reads,

$$\text{Re}\left(w_0 U_{\nu+1}^* + w_L U_\nu^*\right) = 0. \quad (\text{A.15})$$

Given the Wronskian in (2.16) this condition requires $\text{Re}[\mathcal{Q}(\eta, k)] = 0$, which consequently implies $\text{Re}[\mathcal{A}(\epsilon, H_0, k)] = 0$.

De Sitter limit. The de Sitter limit (4.36) we require implies that the shifted particular mode functions in (A.6) and (A.9) in the limit $\epsilon \rightarrow 0$ reduce to,

$$w_0(\eta, k) \xrightarrow{\epsilon \rightarrow 0} -\frac{ik}{2H_0} \left[\frac{ik}{\nu_0 H_0} \frac{\partial U_{\nu_0}}{\partial \nu_0} + U_{\nu_0} \right], \quad (\text{A.16a})$$

$$w_L(\eta, k) \xrightarrow{\epsilon \rightarrow 0} -\frac{ik}{2H_0} \left[\frac{ik}{\nu_0 H_0} \frac{\partial U_{\nu_0+1}}{\partial \nu_0} + \frac{a}{\nu_0} U_{\nu_0} + U_{\nu_0+1} \right]. \quad (\text{A.16b})$$

Demonstrating that this limit is correctly reproduced, and determining what are the conditions this imposes on the constants of integration in (A.6) and (A.9) is not straightforward. This is due to the parametric derivative of the mode function appearing in (A.16), that has to be tied to the special functions appearing in the coefficient functions (A.13) and (A.14). We start by working out the de Sitter limit of these special functions introduced in (A.11) and (A.12),

$$\begin{aligned} \mathcal{J}_1\left(\frac{-2\epsilon}{1-\epsilon}, \pm\nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) &\xrightarrow{\epsilon \rightarrow 0} \mp \frac{\Gamma^2(\mp\nu_0)}{2\nu_0} \left(\frac{k}{2\mathcal{H}}\right)^{\pm 2\nu_0} \times \\ &\times {}_2F_3\left(\left\{\pm\nu_0, \frac{1}{2}\pm\nu_0\right\}, \left\{1\pm\nu_0, 1\pm\nu_0, 1\pm 2\nu_0\right\}, -\frac{k^2}{\mathcal{H}^2}\right) \equiv \mathcal{J}_1^0\left(\pm\nu_0; \frac{k}{\mathcal{H}}\right), \end{aligned} \quad (\text{A.17})$$

$$\begin{aligned} \mathcal{J}_2\left(\frac{-2\epsilon}{1-\epsilon}, \nu; \frac{k}{(1-\epsilon)\mathcal{H}}\right) &\xrightarrow{\epsilon \rightarrow 0} -\frac{(1-\epsilon)\Gamma(\nu)\Gamma(-\nu)}{2\epsilon} + \Gamma(\nu_0)\Gamma(-\nu_0) \times \\ &\times \left[\ln\left(\frac{k}{\mathcal{H}}\right) - \frac{1}{(1-\nu_0^2)} \left(\frac{k}{2\mathcal{H}}\right)^2 {}_3F_4\left(\left\{1, 1, \frac{3}{2}\right\}, \left\{2, 2, 2+\nu_0, 2-\nu_0\right\}, -\frac{k^2}{\mathcal{H}^2}\right) \right] \\ &\equiv -\frac{(1-\epsilon)\Gamma(\nu)\Gamma(-\nu)}{2\epsilon} + \mathcal{J}_2^0\left(\nu_0; \frac{k}{\mathcal{H}}\right). \end{aligned} \quad (\text{A.18})$$

Thus, the coefficient functions (A.13) and (A.14) in the de Sitter limit are,

$$\begin{aligned} \mathcal{Q}(\eta, k) &\xrightarrow{\epsilon \rightarrow 0} \frac{i}{4\pi} \left(\frac{k}{H_0}\right)^2 \left[\mathcal{J}_1^0\left(\nu_0; \frac{k}{\mathcal{H}}\right) + \mathcal{J}_1^0\left(-\nu_0; \frac{k}{\mathcal{H}}\right) - 2\cos(\pi\nu_0) \mathcal{J}_2^0\left(\nu_0; \frac{k}{\mathcal{H}}\right) \right] \\ &\quad - \frac{i(1-\epsilon)\cot(\pi\nu)}{4\epsilon\nu} \left[\frac{k}{(1-\epsilon)H_0}\right]^{\frac{2}{1-\epsilon}} + \mathcal{A}(\epsilon \rightarrow 0, H_0, k), \end{aligned} \quad (\text{A.19})$$

$$\begin{aligned} \tilde{\mathcal{Q}}(\eta, k) &\xrightarrow{\epsilon \rightarrow 0} \frac{e^{-\frac{2ik}{H_0}}}{4\pi} \left(\frac{k}{H_0}\right)^2 \left[e^{-i\pi\nu_0} \mathcal{J}_1^0\left(\nu_0; \frac{k}{\mathcal{H}}\right) + e^{i\pi\nu_0} \mathcal{J}_1^0\left(-\nu_0; \frac{k}{\mathcal{H}}\right) - 2\mathcal{J}_2^0\left(\nu_0; \frac{k}{\mathcal{H}}\right) \right] \\ &\quad - \frac{e^{-\frac{2ik}{(1-\epsilon)H_0}}(1-\epsilon)}{4\epsilon\nu \sin(\pi\nu)} \left[\frac{k}{(1-\epsilon)H_0}\right]^{\frac{2}{1-\epsilon}} + \mathcal{B}(\epsilon \rightarrow 0, H_0, k). \end{aligned} \quad (\text{A.20})$$

Note that in the four expressions above ν still depends on ϵ , as opposed to ν_0 that does not; we could have expanded that dependence as well, but it is more convenient to keep it implicit. Recognizing the parametric derivatives of the mode functions in

the two expressions above is accomplished with the help of the result in Eq. (2.2) from [125] for the parametric derivative of the Bessel function of the first kind,

$$\begin{aligned} \frac{\partial}{\partial \lambda} J_\lambda(z) &= J_{-\lambda}(z) \frac{\Gamma(-\lambda)}{2\Gamma(1+\lambda)} \left(\frac{z}{2}\right)^{2\lambda} {}_2F_3\left(\left\{\lambda, \frac{1}{2}+\lambda\right\}, \left\{1+\lambda, 1+\lambda, 1+2\lambda\right\}, -z^2\right) \\ &- J_\lambda(z) \left[\frac{1}{2\lambda} + \psi(\lambda) - \ln\left(\frac{z}{2}\right) + \frac{z^2}{4(1-\lambda^2)} {}_3F_4\left(\left\{1, 1, \frac{3}{2}\right\}, \left\{2, 2, 2+\lambda, 2-\lambda\right\}, -z^2\right)\right]. \end{aligned} \quad (\text{A.21})$$

It is helpful to first express this result in terms of functions defined in (A.17) and (A.18),

$$\frac{\partial}{\partial \lambda} J_\lambda(z) = \frac{-\mathcal{J}_1^0(\lambda; z)}{\Gamma(\lambda)\Gamma(-\lambda)} J_{-\lambda}(z) - \left[\frac{1}{2\lambda} + \psi(\lambda) + \ln(2) - \frac{\mathcal{J}_2^0(\lambda; z)}{\Gamma(\lambda)\Gamma(-\lambda)}\right] J_\lambda(z). \quad (\text{A.22})$$

This then allows us to express the parametric derivative of the CTBD mode function in (2.11) in terms of the de Sitter limit of special functions given in (A.17) and (A.18),

$$\begin{aligned} \frac{\partial}{\partial \lambda} U_\lambda(\eta, k) &= U_\lambda(\eta, k) \left\{ -\frac{i \cot(\pi\lambda)}{2} [\psi(\lambda) + \psi(-\lambda) + 2 \ln(2)] \right. \\ &\quad \left. + \frac{i\lambda}{2\pi} \left[\mathcal{J}_1^0\left(\lambda; \frac{k}{\mathcal{H}}\right) + \mathcal{J}_1^0\left(-\lambda; \frac{k}{\mathcal{H}}\right) - 2 \cos(\pi\lambda) \mathcal{J}_2^0\left(\lambda; \frac{k}{\mathcal{H}}\right) \right] \right\} \\ &- U_\lambda^*(\eta, k) e^{-\frac{2ik}{\mathcal{H}_0}} \left\{ -\frac{1}{2 \sin(\pi\lambda)} [i\pi + \psi(\lambda) + \psi(-\lambda) + 2 \ln(2)] \right. \\ &\quad \left. + \frac{\lambda}{2\pi} \left[e^{-i\pi\lambda} \mathcal{J}_1^0\left(\lambda; \frac{k}{\mathcal{H}}\right) + e^{i\pi\lambda} \mathcal{J}_1^0\left(-\lambda; \frac{k}{\mathcal{H}}\right) - 2 \mathcal{J}_2^0\left(\lambda; \frac{k}{\mathcal{H}}\right) \right] \right\}, \end{aligned} \quad (\text{A.23})$$

where we have used the reflection formula for the digamma function, $\psi(1-\lambda) - \psi(\lambda) = \pi \cot(\pi\lambda)$. Furthermore, using (A.21) and the recurrence relation (2.15) we derive the analogous expression for a contiguous mode function,

$$\begin{aligned} \frac{\partial}{\partial \lambda} U_{\lambda+1}(\eta, k) &= \frac{i\mathcal{H}}{k} U_\lambda(\eta, k) + U_{\lambda+1}(\eta, k) \left\{ -\frac{i \cot(\pi\lambda)}{2} [\psi(\lambda) + \psi(-\lambda) + 2 \ln(2)] \right. \\ &\quad \left. + \frac{i\lambda}{2\pi} \left[\mathcal{J}_1^0\left(\lambda; \frac{k}{\mathcal{H}}\right) + \mathcal{J}_1^0\left(-\lambda; \frac{k}{\mathcal{H}}\right) - 2 \cos(\pi\lambda) \mathcal{J}_2^0\left(\lambda; \frac{k}{\mathcal{H}}\right) \right] \right\} \\ &+ U_{\lambda+1}^*(\eta, k) e^{-\frac{2ik}{\mathcal{H}_0}} \left\{ -\frac{1}{2 \sin(\pi\lambda)} [i\pi + \psi(\lambda) + \psi(-\lambda) + 2 \ln(2)] \right. \\ &\quad \left. + \frac{\lambda}{2\pi} \left[e^{-i\pi\lambda} \mathcal{J}_1^0\left(\lambda; \frac{k}{\mathcal{H}}\right) + e^{i\pi\lambda} \mathcal{J}_1^0\left(-\lambda; \frac{k}{\mathcal{H}}\right) - 2 \mathcal{J}_2^0\left(\lambda; \frac{k}{\mathcal{H}}\right) \right] \right\}, \end{aligned} \quad (\text{A.24})$$

where we made use of derivative properties,

$$\frac{\partial}{\partial z} \mathcal{I}_1^0(\lambda; z) = \Gamma(1+\lambda) \Gamma(1-\lambda) \Gamma(\lambda) \Gamma(-\lambda) \frac{J_\lambda(z) J_\lambda(z)}{z}, \quad (\text{A.25a})$$

$$\frac{\partial}{\partial z} \mathcal{I}_2^0(\lambda; z) = \Gamma(1+\lambda) \Gamma(1-\lambda) \Gamma(\lambda) \Gamma(-\lambda) \frac{J_\lambda(z) J_{-\lambda}(z)}{z}, \quad (\text{A.25b})$$

that follow from the definitions (A.11) and (A.12), and de Sitter limits (A.17) and (A.18). Finally, this allows us to write the de Sitter limit of the shifted particular mode functions (A.6) and (A.9) as,

$$\begin{aligned} w_0 \stackrel{\epsilon \rightarrow 0}{\sim} & \frac{k^2}{2\nu_0 H_0^2} \frac{\partial U_{\nu_0}}{\partial \nu_0} + \left\{ \frac{ik^2 \cot(\pi\nu_0)}{4\nu_0 H_0^2} \left[\psi(\nu_0) + \psi(-\nu_0) + 2 \ln(2) \right] \right. \\ & \left. - \frac{i(1-\epsilon) \cot(\pi\nu)}{4\epsilon\nu} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} + \mathcal{A}(\epsilon \rightarrow 0, H_0, k) \right\} U_{\nu_0} \\ & - \left\{ \frac{e^{-\frac{2ik}{H_0} k^2}}{4\nu_0 H_0^2 \sin(\pi\nu_0)} \left[i\pi + \psi(\nu_0) + \psi(-\nu_0) + 2 \ln(2) \right] \right. \\ & \left. - \frac{e^{\frac{-2ik}{(1-\epsilon)H_0} (1-\epsilon)}}{4\epsilon\nu \sin(\pi\nu)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} + \mathcal{B}(\epsilon \rightarrow 0, H_0, k) \right\} U_{\nu_0}^*, \quad (\text{A.26}) \end{aligned}$$

$$\begin{aligned} w_L \stackrel{\epsilon \rightarrow 0}{\sim} & \frac{-ik}{2\nu_0 H_0} \left[\frac{ik}{H_0} \frac{\partial U_{\nu_0+1}}{\partial \nu_0} + a U_{\nu_0} \right] + \left\{ \frac{ik^2 \cot(\pi\nu_0)}{4\nu_0 H_0^2} \left[\psi(\nu_0) + \psi(-\nu_0) + 2 \ln(2) \right] \right. \\ & \left. - \frac{i(1-\epsilon) \cot(\pi\nu)}{4\epsilon\nu} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} + \mathcal{A}(\epsilon \rightarrow 0, H_0, k) \right\} U_{\nu_0+1} \\ & + \left\{ \frac{e^{-\frac{2ik}{H_0} k^2}}{4\nu_0 H_0^2 \sin(\pi\nu_0)} \left[i\pi + \psi(\nu_0) + \psi(-\nu_0) + 2 \ln(2) \right] \right. \\ & \left. - \frac{e^{\frac{-2ik}{(1-\epsilon)H_0} (1-\epsilon)}}{4\epsilon\nu \sin(\pi\nu)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} + \mathcal{B}(\epsilon \rightarrow 0, H_0, k) \right\} U_{\nu_0+1}^*. \quad (\text{A.27}) \end{aligned}$$

Requiring these expressions to match (A.16) finally gives the constants of integration in the de Sitter limit,

$$\begin{aligned} \mathcal{A} \stackrel{\epsilon \rightarrow 0}{\sim} & -\frac{ik}{2H_0} + \frac{i(1-\epsilon) \cot(\pi\nu)}{4\epsilon\nu} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \\ & - \frac{ik^2 \cot(\pi\nu_0)}{4\nu_0 H_0^2} \left[\psi(\nu_0) + \psi(-\nu_0) + 2 \ln(2) \right] + \mathcal{O}(\epsilon), \quad (\text{A.28}) \end{aligned}$$

$$\begin{aligned} \mathcal{B} \stackrel{\epsilon \rightarrow 0}{\sim} & \frac{e^{\frac{-2ik}{(1-\epsilon)H_0} (1-\epsilon)}}{4\epsilon\nu \sin(\pi\nu)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \\ & - \frac{e^{-\frac{2ik}{H_0} k^2}}{4\nu_0 H_0^2 \sin(\pi\nu_0)} \left[i\pi + \psi(\nu_0) + \psi(-\nu_0) + 2 \ln(2) \right] + \mathcal{O}(\epsilon), \quad (\text{A.29}) \end{aligned}$$

where \mathcal{A} is purely imaginary as required by the Wronskian condition.

Flat space limit. The flat space limit (4.38) implies that the shifted particular mode functions in the limit $H_0 \rightarrow 0$ have to reduce to

$$w_0 \xrightarrow{H_0 \rightarrow 0} \frac{1}{4} \left[1 + 2ik(\eta - \eta_0) \right] \frac{e^{-ik(\eta - \eta_0)}}{\sqrt{2k}}, \quad w_L \xrightarrow{H_0 \rightarrow 0} \frac{1}{4} \left[-1 + 2ik(\eta - \eta_0) \right] \frac{e^{-ik(\eta - \eta_0)}}{\sqrt{2k}}. \quad (\text{A.30})$$

To show that this limit is reproduced by (A.6) and (A.9) we need the asymptotic series for the generalized hypergeometric functions in the limit of large argument that are given in §16.11 of [84, 85]. In particular we need two special cases of that asymptotic series,

$$\mathcal{I}_1(\rho, \pm\lambda, z) \xrightarrow{z \rightarrow \infty} \Gamma(\lambda) \Gamma(-\lambda) \Gamma(1+\lambda) \Gamma(1-\lambda) \left\{ \frac{\Gamma(\frac{1-\rho}{2}) \Gamma(\frac{\rho \pm \lambda}{2})}{2 \Gamma(\frac{1}{2}) \Gamma(\frac{2-\rho}{2}) \Gamma(\frac{2-\rho}{2} \pm \lambda)} - \frac{z^{\rho-1}}{\pi} \left[\frac{1}{(1-\rho)} + \frac{\cos(2z \mp \pi\lambda)}{2z} + \mathcal{O}(z^{-2}) \right] \right\}, \quad (\text{A.31})$$

$$\mathcal{I}_2(\rho, \lambda, z) \xrightarrow{z \rightarrow \infty} \Gamma(\lambda) \Gamma(-\lambda) \Gamma(1+\lambda) \Gamma(1-\lambda) \left\{ \frac{\Gamma(\frac{\rho}{2}) \Gamma(\frac{1-\rho}{2})}{2 \Gamma(\frac{1}{2}) \Gamma(\frac{2-\rho}{2} + \lambda) \Gamma(\frac{2-\rho}{2} - \lambda)} - \frac{z^{\rho-1}}{\pi} \left[\frac{\cos(\pi\lambda)}{(1-\rho)} + \frac{\cos(2z)}{2z} + \mathcal{O}(z^{-2}) \right] \right\}, \quad (\text{A.32})$$

which imply the following flat space limits for the coefficient functions (A.13) and (A.14),

$$\mathcal{Q}(\eta, k) \xrightarrow{H_0 \rightarrow 0} \frac{1}{2} \left[\frac{ik}{(1+\epsilon)H_0} + ik(\eta - \eta_0) \right] \left[1 + \mathcal{O}(H_0^2) \right] + \mathcal{A}(\epsilon, H_0 \rightarrow 0, k), \quad (\text{A.33})$$

$$- \frac{i \Gamma(\frac{-\epsilon}{1-\epsilon}) \Gamma(\frac{1}{2} + \frac{\epsilon}{1-\epsilon}) \Gamma(\frac{-\epsilon}{1-\epsilon} + \nu) \Gamma(\frac{-\epsilon}{1-\epsilon} - \nu)}{4 \Gamma(\frac{1}{2}) \Gamma(\frac{1}{2} + \nu) \Gamma(\frac{1}{2} - \nu)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}},$$

$$\tilde{\mathcal{Q}}(\eta, k) \xrightarrow{H_0 \rightarrow 0} - \frac{e^{-2ik(\eta - \eta_0)}}{4} \left[1 + \mathcal{O}(H_0) \right] + \mathcal{B}(\epsilon, H_0 \rightarrow 0, k) \quad (\text{A.34})$$

$$- e^{\frac{-2ik}{(1-\epsilon)H_0}} e^{\frac{-i\pi\epsilon}{1-\epsilon}} \frac{\Gamma(\frac{-\epsilon}{1-\epsilon}) \Gamma(\frac{-\epsilon}{1-\epsilon} + \nu) \Gamma(\frac{-\epsilon}{1-\epsilon} - \nu)}{4 \Gamma(\frac{1}{2}) \Gamma(\frac{1}{2} - \frac{\epsilon}{1-\epsilon})} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}}.$$

Thus the flat space limit of the shifted particular mode functions is given by,

$$w_0 \xrightarrow{H_0 \rightarrow 0} \frac{1}{4} \left[1 + 2ik(\eta - \eta_0) \right] \frac{e^{-ik(\eta - \eta_0)}}{\sqrt{2k}} + \left\{ \frac{ik}{2(1+\epsilon)H_0} + \mathcal{A}(\epsilon, H_0 \rightarrow 0, k) - \frac{i \Gamma(\frac{-\epsilon}{1-\epsilon}) \Gamma(\frac{1}{2} + \frac{\epsilon}{1-\epsilon}) \Gamma(\frac{-\epsilon}{1-\epsilon} + \nu) \Gamma(\frac{-\epsilon}{1-\epsilon} - \nu)}{4 \Gamma(\frac{1}{2}) \Gamma(\frac{1}{2} + \nu) \Gamma(\frac{1}{2} - \nu)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \right\} \frac{e^{-ik(\eta - \eta_0)}}{\sqrt{2k}}$$

$$+ \left\{ e^{\frac{-2ik}{(1-\epsilon)H_0}} e^{\frac{-i\pi\epsilon}{1-\epsilon}} \frac{\Gamma(\frac{-\epsilon}{1-\epsilon}) \Gamma(\frac{-\epsilon}{1-\epsilon} + \nu) \Gamma(\frac{-\epsilon}{1-\epsilon} - \nu)}{4 \Gamma(\frac{1}{2}) \Gamma(\frac{1}{2} - \frac{\epsilon}{1-\epsilon})} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} - \mathcal{B}(\epsilon, H_0 \rightarrow 0, k) \right\} \frac{e^{ik(\eta - \eta_0)}}{\sqrt{2k}}, \quad (\text{A.35})$$

$$\begin{aligned}
w_L \xrightarrow{H_0 \rightarrow 0} & \frac{1}{4} \left[-1 + 2ik(\eta - \eta_0) \right] \frac{e^{-ik(\eta - \eta_0)}}{\sqrt{2k}} + \left\{ \frac{ik}{2(1+\epsilon)H_0} + \mathcal{A}(\epsilon, H_0 \rightarrow 0, k), \right. \\
& \left. - \frac{i \Gamma\left(\frac{-\epsilon}{1-\epsilon}\right) \Gamma\left(\frac{1}{2} + \frac{\epsilon}{1-\epsilon}\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} + \nu\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} - \nu\right)}{4 \Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2} + \nu\right) \Gamma\left(\frac{1}{2} - \nu\right)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \right\} \frac{e^{-ik(\eta - \eta_0)}}{\sqrt{2k}} \\
& - \left\{ e^{\frac{-2ik}{(1-\epsilon)H_0}} e^{\frac{-i\pi\epsilon}{1-\epsilon}} \frac{\Gamma\left(\frac{-\epsilon}{1-\epsilon}\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} + \nu\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} - \nu\right)}{4 \Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2} - \frac{\epsilon}{1-\epsilon}\right)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}} \right. \\
& \left. - \mathcal{B}(\epsilon, H_0 \rightarrow 0, k) \right\} \frac{e^{ik(\eta - \eta_0)}}{\sqrt{2k}}. \tag{A.36}
\end{aligned}$$

In order for these expressions to match (A.30) the integration constants in the flat space limit must satisfy,

$$\mathcal{A} \xrightarrow{H_0 \rightarrow 0} \frac{-ik}{2(1+\epsilon)H_0} + \frac{i \Gamma\left(\frac{-\epsilon}{1-\epsilon}\right) \Gamma\left(\frac{1}{2} + \frac{\epsilon}{1-\epsilon}\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} + \nu\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} - \nu\right)}{4 \Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2} + \nu\right) \Gamma\left(\frac{1}{2} - \nu\right)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}}, \tag{A.37}$$

$$\mathcal{B} \xrightarrow{H_0 \rightarrow 0} e^{\frac{-2ik}{(1-\epsilon)H_0}} e^{\frac{-i\pi\epsilon}{1-\epsilon}} \frac{\Gamma\left(\frac{-\epsilon}{1-\epsilon}\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} + \nu\right) \Gamma\left(\frac{-\epsilon}{1-\epsilon} - \nu\right)}{4 \Gamma\left(\frac{1}{2}\right) \Gamma\left(\frac{1}{2} - \frac{\epsilon}{1-\epsilon}\right)} \left[\frac{k}{(1-\epsilon)H_0} \right]^{\frac{2}{1-\epsilon}}. \tag{A.38}$$

These two conditions, together with the two conditions (A.28) and (A.29) found by considering the de Sitter limit, constrain the choice for the constants of integration that we can make. In fact, it is most convenient to promote the flat space limit conditions (A.37) and (A.38) to the full choice for the constants. It is straightforward to check that this choice satisfies the de Sitter limit requirements automatically.

B Inverting Laplacian

The Laplace-inverted two-point function introduced in (5.15) can also be evaluated in a closed form, rather than in the series form (5.45) given in Sec. 5.3. This is accomplished by first reverting to the sum-over-modes representation,

$$i \left[{}^{-}\Xi^+ \right]_{\lambda}(x; x') = -(aa')^{\frac{2-D}{2}} \int \frac{d^{D-1}k}{(2\pi)^{D-1}} \frac{(1-\epsilon)^2 \mathcal{H} \mathcal{H}'}{k^2} e^{i\vec{k} \cdot \Delta \vec{x}} \theta(k - k_0) U_{\lambda}(\eta, k) U_{\lambda}^*(\eta', k), \tag{B.1}$$

where the step function effectively enforces the mode function suppression in the IR by introducing a cutoff $k_0 \ll H_0$. The UV convergence for all ranges of coordinates x and x' is ensured by the same $i\delta$ -prescription as for the scalar two-point function in (2.18). In this appendix we invert the Laplacian acting on the Wightman function, with the particular $i\delta$ -prescription implicit in all expressions,

$$\mathcal{H} \rightarrow \mathcal{H}_{-+} = \mathcal{H} \left[1 - \frac{(1-\epsilon)\mathcal{H}i\delta}{2} \right], \quad \mathcal{H}' \rightarrow \mathcal{H}'_{-+} = \mathcal{H}' \left[1 + \frac{(1-\epsilon)\mathcal{H}'i\delta}{2} \right]. \tag{B.2}$$

Generalization to other two-point functions is straightforward. We should also note it is more convenient to use \mathcal{H} , \mathcal{H}' , and $\|\Delta\vec{x}\|$ as variables for the two-point functions that in this appendix, instead of the bi-local variables (2.8) employed in the main text.

The integral in (B.1) can be evaluated exactly. Integrating over the angular coordinates involves an integral over a $(D-2)$ -sphere and an integral representation of the Bessel function of the first kind, given in (2.42), and it produces,

$$i[-\Xi^+]_{\lambda}(x; x') = -\frac{(aa')^{-\frac{D-2}{2}}(1-\epsilon)^2\mathcal{H}\mathcal{H}'}{(2\pi)^{\frac{D-1}{2}}\|\Delta\vec{x}\|^{\frac{D-3}{2}}}\int_{k_0}^{\infty} dk k^{\frac{D-5}{2}} J_{\frac{D-3}{2}}(k\|\Delta\vec{x}\|)U_{\lambda}(\eta, k)U_{\lambda}^*(\eta', k). \quad (\text{B.3})$$

We evaluate this resulting integral analogously to evaluating the integral for the scalar two-point functions in Sec. 2.3.2. This entails first writing out the integration over modes as the bulk (infinite) range and subtracting from it the infrared range, $\int_{k_0}^{\infty} = \int_0^{\infty} - \int_0^{k_0}$. Each part is then computed individually by dimensionally regulating their respective infrared behaviour. This procedure would be incorrect to apply on actually infrared divergent integrals. However, the integral in (B.3) is not infrared divergent to start with, and the errors we are making by dimensionally regulating the infrared of the two individual parts cancel each other out. Therefore, it is legitimate to split the integral in (B.3) into two contributions (cf. Eq. (2.40)),

$$i[-\Xi^+]_{\lambda}(x; x') = (aa')^{-\frac{(D-2)\epsilon}{2}}\left[\mathcal{M}_{\lambda}(x; x') + \mathcal{X}_{\lambda}(x; x')\right]. \quad (\text{B.4})$$

that we compute separately in the following subsections.

B.1 Bulk part

The bulk part of (B.4) is given by the integral representation,

$$\begin{aligned} \mathcal{M}_{\lambda}(x; x') &= -\frac{(aa')^{-\frac{(D-2)(1-\epsilon)}{2}}(1-\epsilon)\sqrt{\mathcal{H}\mathcal{H}'}}{8(2\pi)^{\frac{D-3}{2}}\|\Delta\vec{x}\|^{\frac{D-3}{2}}} \\ &\times \int_0^{\infty} dk k^{\frac{D-5}{2}} J_{\frac{D-3}{2}}(k\|\Delta\vec{x}\|)H_{\lambda}^{(1)}\left(\frac{k}{(1-\epsilon)\mathcal{H}}\right)H_{\lambda}^{(2)}\left(\frac{k}{(1-\epsilon)\mathcal{H}'}\right). \end{aligned} \quad (\text{B.5})$$

The convergence for all ranges of coordinates is guaranteed by the implicit $i\delta$ -prescription (B.2). We first evaluate the integral for the range of coordinates where the $i\delta$ -prescriptions are not necessary, and then analytically extended to the remaining ranges of coordinates. The result we use is an integral over three Bessel functions and a power that is given in 6.578.1. of [100], or in 2.12.42.5. of [101], and is originally due to Rice [126] and Bailey [127],

$$\begin{aligned} \int_0^{\infty} dz z^{\rho-1} J_{\sigma}(cz)J_{\lambda}(az)J_{\mu}(bz) &= \frac{2^{\rho-1}\Gamma\left(\frac{\lambda+\mu+\rho+\sigma}{2}\right)a^{\lambda}b^{\mu}c^{-\lambda-\mu-\rho}}{\Gamma(1+\lambda)\Gamma(1+\mu)\Gamma\left(1-\frac{\lambda+\mu+\rho-\sigma}{2}\right)} \\ &\times F_4\left(\frac{\lambda+\mu+\rho+\sigma}{2}, \frac{\lambda+\mu+\rho-\sigma}{2}; 1+\lambda, 1+\mu; \frac{a^2}{c^2}, \frac{b^2}{c^2}\right). \end{aligned} \quad (\text{B.6})$$

It holds when the following conditions are met,

$$\operatorname{Re}(\lambda + \mu + \rho + \sigma) > 0, \quad \operatorname{Re}(\rho) < \frac{5}{2}, \quad a, b, c > 0, \quad c > a + b. \quad (\text{B.7})$$

The function F_4 in (B.6) is the Appell's fourth function, defined inside the region $|\sqrt{X}| + |\sqrt{Y}| < 1$ by the convergent double power series,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) = \frac{\Gamma(\gamma) \Gamma(\gamma')}{\Gamma(\alpha) \Gamma(\beta)} \sum_{m=0}^{\infty} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+m+n) \Gamma(\beta+m+n)}{\Gamma(\gamma+m) \Gamma(\gamma'+n)} \frac{X^m Y^n}{m! n!}, \quad (\text{B.8})$$

and outside of it by its analytical extension. This definition implies symmetry properties,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) = F_4(\beta, \alpha; \gamma, \gamma'; X, Y) = F_4(\alpha, \beta; \gamma', \gamma; Y, X), \quad (\text{B.9})$$

and also that,

$$F_4(0, \beta; \gamma, \gamma'; X, Y) = 1. \quad (\text{B.10})$$

Performing either of the two sums in the definition (B.8) expresses it as a single series over hypergeometric functions,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) = \frac{\Gamma(\gamma')}{\Gamma(\alpha) \Gamma(\beta)} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+n) \Gamma(\beta+n)}{\Gamma(\gamma'+n) n!} Y^n {}_2F_1\left(\{\alpha+n, \beta+n\}, \{\gamma\}, X\right), \quad (\text{B.11})$$

which is particularly useful for defining the analytic continuation to the entire range of complex coordinates X and Y . In particular, it makes manifest the two branch points at $X=1$ and $Y=1$ that the Appell's fourth function possesses. We need only two special cases of the result in (B.6),

$$\begin{aligned} & \int_0^{\infty} dz z^{\rho-1} J_{\rho}(cz) J_{\lambda}(az) J_{\lambda}(bz) \\ &= \frac{2^{\rho-1} \Gamma(\rho+\lambda) c^{-\rho}}{\Gamma^2(1+\lambda) \Gamma(1-\lambda)} \left(\frac{ab}{c^2}\right)^{\lambda} F_4\left(\rho+\lambda, \lambda; 1+\lambda, 1+\lambda; \frac{a^2}{c^2}, \frac{b^2}{c^2}\right), \end{aligned} \quad (\text{B.12})$$

$$\int_0^{\infty} dz z^{\rho-1} J_{\rho}(cz) J_{\lambda}(az) J_{-\lambda}(bz) = \frac{2^{\rho-1} \Gamma(\rho) c^{-\rho}}{\Gamma(1+\lambda) \Gamma(1-\lambda)} \left(\frac{a}{b}\right)^{\lambda}, \quad (\text{B.13})$$

so that (B.5) evaluates to,

$$\begin{aligned} \mathcal{M}_{\lambda}(x; x') &= - \frac{[(1-\epsilon)H_0]^{D-2}}{16\pi^{\frac{D+1}{2}}} \frac{\Gamma\left(\frac{D-3}{2}\right) \Gamma(\lambda) \Gamma(-\lambda)}{[(1-\epsilon)^2 \mathcal{H} \mathcal{H}' \|\Delta\vec{x}\|^2]^{\frac{D-3}{2}}} \left[e^{i\pi\lambda} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{\lambda} + e^{-i\pi\lambda} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{-\lambda} \right] \\ &+ \mathcal{F}_{\lambda}\left((1-\epsilon)\mathcal{H}\|\Delta\vec{x}\|, (1-\epsilon)\mathcal{H}'\|\Delta\vec{x}\|\right) + \mathcal{F}_{-\lambda}\left((1-\epsilon)\mathcal{H}\|\Delta\vec{x}\|, (1-\epsilon)\mathcal{H}'\|\Delta\vec{x}\|\right), \end{aligned} \quad (\text{B.14})$$

where we have defined a shorthand notation for the two-variable function,

$$\mathcal{F}_\lambda(X, Y) = \frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\frac{D-3}{2} + \lambda) \Gamma(-\lambda)}{16\pi^{\frac{D+1}{2}} \lambda} (XY)^{-\frac{D-3}{2} - \lambda} \times F_4\left(\frac{D-3}{2} + \lambda, \lambda; 1 + \lambda, 1 + \lambda; \frac{1}{X^2}, \frac{1}{Y^2}\right). \quad (\text{B.15})$$

This result is obtained for the range of coordinates on which the result (B.6) is valid, which according to (B.7) translates into the condition $y(x; x') > 4$ on the bi-local variable. The result is then extended to the full range of coordinates by analytic continuation provided by the implicit $i\delta$ -prescriptions in time coordinates/conformal Hubble rates (B.2). It is interesting to note that the bulk solution (B.14), when written in terms of bi-local variables (2.8), depends only on y and v , while the sole dependence on u is in the overall factor taken out in (B.4).

The representation (B.14) for the integral defined in (B.5) is correct on the entire range of coordinates. However, in practice this representation is best adapted to the super-Hubble regime for which the arguments of Appell's fourth function take small values. We can derive an alternative representation adapted to the sub-Hubble regime, by making use of the identity 16.16.10. from [84, 85],

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) = \frac{\Gamma(\gamma') \Gamma(\beta - \alpha)}{\Gamma(\gamma' - \alpha) \Gamma(\beta)} (-Y)^{-\alpha} F_4\left(\alpha, \alpha - \gamma' + 1; \gamma, \alpha - \beta + 1; \frac{X}{Y}, \frac{1}{Y}\right) + \frac{\Gamma(\gamma') \Gamma(\alpha - \beta)}{\Gamma(\gamma' - \beta) \Gamma(\alpha)} (-Y)^{-\beta} F_4\left(\beta, \beta - \gamma' + 1; \gamma, \beta - \alpha + 1; \frac{X}{Y}, \frac{1}{Y}\right), \quad (\text{B.16})$$

that is valid for $|\text{Arg}(-Y)| < \pi$. This turns the solution (B.14) into,

$$\mathcal{M}_\lambda(x; x') = e^{\frac{i\pi(D-1)}{2}} \left[e^{i\pi\lambda} \overline{\mathcal{F}}_\lambda\left(\frac{\mathcal{H}}{\mathcal{H}'}, (1-\epsilon)\mathcal{H}\|\Delta\vec{x}\|\right) + e^{-i\pi\lambda} \overline{\mathcal{F}}_{-\lambda}\left(\frac{\mathcal{H}}{\mathcal{H}'}, (1-\epsilon)\mathcal{H}\|\Delta\vec{x}\|\right) \right] + e^{-\frac{i\pi(D-1)}{2}} \left[e^{-i\pi\lambda} \overline{\mathcal{F}}_\lambda\left(\frac{\mathcal{H}'}{\mathcal{H}}, (1-\epsilon)\mathcal{H}'\|\Delta\vec{x}\|\right) + e^{i\pi\lambda} \overline{\mathcal{F}}_{-\lambda}\left(\frac{\mathcal{H}'}{\mathcal{H}}, (1-\epsilon)\mathcal{H}'\|\Delta\vec{x}\|\right) \right], \quad (\text{B.17})$$

where we define another shorthand notation for a different two-variable function,

$$\overline{\mathcal{F}}_\lambda(X, Y) = \frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\frac{D-3}{2} + \lambda) \Gamma(-\lambda)}{16\pi^{\frac{D+1}{2}} (D-3)} X^{\frac{D-3}{2} + \lambda} \times F_4\left(\frac{D-3}{2} + \lambda, \frac{D-3}{2}; 1 + \lambda, \frac{D-1}{2}; X^2, Y^2\right). \quad (\text{B.18})$$

Note that we had applied the transformation (B.16) to (B.14) in a fashion that preserves manifest invariance under simultaneous complex conjugation and interchange $x \leftrightarrow x'$ that the two-point function must respect.

It is also possible represent the solution (B.14) in terms of Appell's first function instead of the fourth one. This is accomplished by first applying the argument transformation formula (3.1) from [128],

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) = F_4(\alpha, \beta; \gamma-1, \gamma'; X, Y) - \frac{\alpha\beta X}{\gamma(\gamma-1)} F_4(\alpha+1, \beta+1; \gamma+1, \gamma'; X, Y), \quad (\text{B.19})$$

followed by the transformation formula (4.1) from [127] between different Appell's functions,

$$F_4\left(\alpha, \beta; \gamma, \beta; \frac{-X}{(1-X)(1-Y)}, \frac{-Y}{(1-X)(1-Y)}\right) = (1-X)^\alpha (1-Y)^\alpha F_1(\alpha; \gamma-\beta, 1+\alpha-\gamma; \gamma; X, XY). \quad (\text{B.20})$$

While more is known about Appell's first function than the fourth one, we do not pursue the representation resulting from (B.19) and (B.20) above, as we did not find immediate advantages in employing it compared to the two representations in (B.14) and (B.17). Further representations and transformations of Appell's fourth function can be derived from transformations of the hypergeometric function [128] that might be useful when examining different ranges of coordinates.

B.2 Infrared part

The infrared part of the two-point function in (B.4) is given by,

$$\mathcal{X}_\lambda(x; x') = \frac{(aa')^{-\frac{(D-2)(1-\epsilon)}{2}} (1-\epsilon)^2 \mathcal{H}\mathcal{H}'}{(2\pi)^{\frac{D-1}{2}} \|\Delta\vec{x}\|^{\frac{D-3}{2}}} \int_0^{k_0} dk k^{\frac{D-5}{2}} J_{\frac{D-3}{2}}(k\|\Delta\vec{x}\|) U_\lambda(\eta, k) U_\lambda^*(\eta', k). \quad (\text{B.21})$$

We evaluate this integral by following the procedure of Sec. 2.3.2. This entails first expanding the CTBD mode functions in the low momentum limit (2.14), keeping only the terms that can generate negative powers of the IR cutoff k_0 ,

$$\begin{aligned} \mathcal{X}_\lambda(x; x') &= \frac{(aa')^{-\frac{(D-2)(1-\epsilon)}{2}} (1-\epsilon) \sqrt{\mathcal{H}\mathcal{H}'} \Gamma^2(\lambda) \Gamma^2(1-\lambda)}{2^{1-2\lambda} (2\pi)^{\frac{D+1}{2}} \|\Delta\vec{x}\|^{\frac{D-3}{2}}} \\ &\times \sum_{N=0}^{\infty} \sum_{n=0}^N \frac{(-\frac{1}{4})^N}{n!(N-n)! \Gamma(n+1-\lambda) \Gamma(N-n+1-\lambda)} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{N-2n} \\ &\times \int_0^{k_0} dk k^{\frac{D-5}{2}} J_{\frac{D-3}{2}}(k\|\Delta\vec{x}\|) \left[\frac{k^2}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'}\right]^{N-\lambda}. \end{aligned} \quad (\text{B.22})$$

Evaluating the resulting integrals according to (2.45) then gives,

$$\mathcal{X}_\lambda(x; x') = \frac{[(1-\epsilon)H_0]^{D-2}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(\lambda)\Gamma(2\lambda)}{\Gamma(\frac{1}{2}+\lambda)\Gamma(\frac{D-1}{2})} \sum_{N=0}^{\infty} \sum_{n=0}^N \frac{1}{(\frac{D-3}{2}+N-\lambda)}$$

$$\begin{aligned} & \times \frac{\left(-\frac{1}{4}\right)^N \Gamma^2(1-\lambda)}{n!(N-n)! \Gamma(n+1-\lambda) \Gamma(N-n+1-\lambda)} \left[\frac{k_0^2}{(1-\epsilon)^2 \mathcal{H} \mathcal{H}'}\right]^{\frac{D-3}{2}+N-\lambda} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{N-2n} \\ & \times {}_1F_2\left(\left\{\frac{D-3}{2}+N-\lambda\right\}, \left\{\frac{D-1}{2}+N-\lambda, \frac{D-1}{2}\right\}, -\frac{(k_0 \|\Delta\vec{x}\|)^2}{4}\right), \quad (\text{B.23}) \end{aligned}$$

where we have used the Legendre duplication formula to simplify the gamma functions in the overall factor. Expanding the hypergeometric function for small arguments $k_0 \|\Delta\vec{x}\| \ll 1$, and reorganizing the series produces the final expression,

$$\begin{aligned} \mathcal{X}_\lambda(x; x') &= \frac{[(1-\epsilon)H_0]^{D-2}}{(4\pi)^{\frac{D}{2}}} \frac{\Gamma(\lambda)\Gamma(2\lambda)}{\Gamma(\frac{1}{2}+\lambda)\Gamma(\frac{D-1}{2})} \left[\frac{k_0^2}{(1-\epsilon)^2 \mathcal{H} \mathcal{H}'}\right]^{\frac{D-3}{2}-\lambda} \quad (\text{B.24}) \\ & \times \sum_{N=0}^{\lfloor \lambda - \frac{D-3}{2} \rfloor} \sum_{n=0}^N \sum_{\ell=0}^{N-n} \frac{c_{Nn\ell}}{\left(\frac{D-3}{2}+N-\lambda\right)} \left[\frac{k_0^2}{(1-\epsilon)^2 \mathcal{H} \mathcal{H}'}\right]^N \left[(1-\epsilon)^2 \mathcal{H} \mathcal{H}' \|\Delta\vec{x}\|^2\right]^n \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{N-n-2\ell}, \end{aligned}$$

where the coefficients $c_{Nn\ell}$ are given in (2.48), and where the floor function truncates the external sum at $N \leq (D-3)/2$, thus keeping only negative powers of k_0 .

B.3 Various limits

Closed form solutions (B.14) and (B.14) for the Laplace-inverted two-point function greatly simplify in certain limits. We derive several of these limits here, including rederiving limits from Sec. 6 starting directly from closed form solutions given in terms of Appell's fourth function.

de Sitter limit. Reproducing the de Sitter limit (6.1) does not rely on the particular representation of the inverted Laplacian (5.15). Rather, it hinges on showing that (6.4) holds, which is true for any representation.

Flat space limit. When examining particular limits, different representations of the Appell's fourth function are useful. For the flat space limit of Sec. 6.2 we need to derive the asymptotic behaviour of (B.18) around the point $(X, Y) = (1, 0)$, to be applied in (B.17). This is because the arguments of Appell's functions in (B.17) in the flat space limit reduce to,

$$(1-\epsilon)\mathcal{H}\|\Delta\vec{x}\| \stackrel{H_0 \rightarrow 0}{\sim} (1-\epsilon)H_0\|\Delta\vec{x}\|, \quad (1-\epsilon)\mathcal{H}'\|\Delta\vec{x}\| \stackrel{H_0 \rightarrow 0}{\sim} (1-\epsilon)H_0\|\Delta\vec{x}\|, \quad (\text{B.25a})$$

$$\frac{\mathcal{H}^2}{\mathcal{H}'^2} \stackrel{H_0 \rightarrow 0}{\sim} 1 + 2(1-\epsilon)H_0\Delta\eta, \quad \frac{\mathcal{H}'^2}{\mathcal{H}^2} \stackrel{H_0 \rightarrow 0}{\sim} 1 - 2(1-\epsilon)H_0\Delta\eta. \quad (\text{B.25b})$$

where the $i\delta$ -prescriptions are implied. Therefore, the limit $(X, Y) \rightarrow (1, 0)$ has to be taken simultaneously because the particular ratio of arguments is finite and non-vanishing,

$$X \rightarrow 1, \quad Y \rightarrow 0, \quad \frac{Y}{(1-X)^2} \rightarrow Z, \quad 0 < Z < \infty. \quad (\text{B.26})$$

This can be accomplished as in [129], by first applying the transformation formula for the hypergeometric function 9.131.2. from [100],

$${}_2F_1\left(\{\alpha, \beta\}, \{\gamma\}, X\right) = \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} {}_2F_1\left(\{\alpha, \beta\}, \{\alpha+\beta-\gamma+1\}, 1-X\right) \quad (\text{B.27})$$

$$+ \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)} (1-X)^{\gamma-\alpha-\beta} {}_2F_1\left(\{\gamma-\alpha, \gamma-\beta\}, \{\gamma-\alpha-\beta+1\}, 1-X\right),$$

valid for $|\text{Arg}(1-X)| < \pi$, to the series representation of Appell's fourth function in (B.11), resulting in,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) = \frac{\Gamma(\gamma)\Gamma(\gamma')}{\Gamma(\alpha)\Gamma(\beta)} \sum_{n=0}^{\infty} \left\{ \frac{\Gamma(\alpha+n)\Gamma(\beta+n)\Gamma(\gamma-\alpha-\beta-2n)}{\Gamma(\gamma-\alpha-n)\Gamma(\gamma-\beta-n)\Gamma(\gamma'+n)n!} Y^n \right.$$

$$\times {}_2F_1\left(\{\alpha+n, \beta+n\}, \{\alpha+\beta-\gamma+1+2n\}, 1-X\right) + \frac{\Gamma(\alpha+\beta-\gamma+2n)}{\Gamma(\gamma'+n)n!} (1-X)^{\gamma-\alpha-\beta}$$

$$\times \left[\frac{Y}{(1-X)^2} \right]^n {}_2F_1\left(\{\gamma-\alpha-n, \gamma-\beta-n\}, \{\gamma-\alpha-\beta+1-2n\}, 1-X\right) \left. \right\}. \quad (\text{B.28})$$

Then applying the limit (B.26) for the variables simplifies this expression,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) \stackrel{(\text{B.26})}{\rightsquigarrow} \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)}$$

$$+ \frac{\Gamma(\gamma)\Gamma(\gamma')}{\Gamma(\alpha)\Gamma(\beta)} (1-X)^{\gamma-\alpha-\beta} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+\beta-\gamma+2n)}{\Gamma(\gamma'+n)n!} \left[\frac{Y}{(1-X)^2} \right]^n$$

$$= \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta)}{\Gamma(\gamma-\alpha)\Gamma(\gamma-\beta)} + \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)} (1-X)^{\gamma-\alpha-\beta}$$

$$\times {}_2F_1\left(\left\{\frac{\alpha+\beta-\gamma+1}{2}, \frac{\alpha+\beta-\gamma}{2}\right\}, \{\gamma'\}, \frac{4Y}{(1-X)^2}\right). \quad (\text{B.29})$$

We apply this result to functions (B.18), and plug them into (B.17) taking the flat space limit,

$$\mathcal{M}_\lambda(x; x') \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} \frac{[(1-\epsilon)H_0]^2}{(4\pi)^{\frac{D-1}{2}}} \frac{\Gamma(D-4)}{\Gamma(\frac{D-1}{2})} \times e^{-\frac{i\pi(D-2)}{2}} (\Delta\eta)^{4-D}$$

$$\times {}_2F_1\left(\left\{\frac{D-3}{2}, \frac{D-4}{2}\right\}, \left\{\frac{D-1}{2}\right\}, 1 + \frac{\Delta x^2}{\Delta\eta^2}\right). \quad (\text{B.30})$$

Note that this expression is divergent in $D \rightarrow 4$ by itself, but that it always appears in photon two-point function with derivatives acting on it that remove the divergence. The flat space limit we derived in Sec. 6.2 relies on demonstrating that the limit in (6.14) holds. Checking this limit here entails taking the derivative of (B.30) with respect to y , which in the flat space limit reduces to,

$$\frac{\partial}{\partial y} \stackrel{H_0 \rightarrow 0}{\rightsquigarrow} \frac{1}{(1-\epsilon)^2 H_0^2} \frac{\partial}{\partial(\Delta x^2)}, \quad (\text{B.31})$$

and then applying the second transformation formula 9.131.1 from [100],

$${}_2F_1\left(\{\alpha, \beta\}, \{\gamma\}, X\right) = (1-X)^{-\beta} {}_2F_1\left(\{\gamma-\alpha, \beta\}, \{\gamma\}, \frac{X}{X-1}\right), \quad (\text{B.32})$$

valid for $|\text{Arg}(1-X)| < \pi$. This finally produces

$$\frac{\partial}{\partial y} \mathcal{M}_\lambda(x; x') \xrightarrow{H_0 \rightarrow 0} \frac{\Gamma(D-2)}{4\Gamma\left(\frac{D+1}{2}\right)} \frac{(\Delta x^2)^{-\frac{D-2}{2}}}{(4\pi)^{\frac{D-1}{2}}} {}_2F_1\left(\left\{1, \frac{D-2}{2}\right\}, \left\{\frac{D+1}{2}\right\}, 1 + \frac{\Delta\eta^2}{\Delta x^2}\right). \quad (\text{B.33})$$

Upon applying some simple identities for gamma function, this is the same as expression (6.14), up to an overall factor $(aa')^{-\frac{(D-2)\epsilon}{2}}$ that was factored out in (B.4).

Spatial coincidence limit for the bulk part. The limit $\|\Delta\vec{x}\| \rightarrow 0$ is straightforwardly obtained from the second representation for the bulk part (B.17). First we need to consider the appropriate limit of the function defined in (B.18), that is obtained by first using the limit of Appell's fourth function for one vanishing argument,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) \xrightarrow{Y \rightarrow 0} {}_2F_1\left(\{\alpha, \beta\}, \{\gamma\}, X\right), \quad (\text{B.34})$$

followed by the quadratic transformation formula 15.8.21 from [84, 85],

$$\begin{aligned} & {}_2F_1\left(\{\alpha, \beta\}, \{\alpha-\beta+1\}, X^2\right) \\ &= (1+X)^{-2\alpha} {}_2F_1\left(\left\{\alpha, \alpha-\beta+\frac{1}{2}\right\}, \left\{2\alpha-2\beta+1\right\}, \frac{4X}{(1+X)^2}\right) \end{aligned} \quad (\text{B.35})$$

and the linear transformation 9.132.2 from [100],

$$\begin{aligned} {}_2F_1\left(\{\alpha, \beta\}, \{\gamma\}, X\right) &= \frac{\Gamma(\gamma)\Gamma(\beta-\alpha)}{\Gamma(\beta)\Gamma(\gamma-\alpha)} (-X)^{-\alpha} {}_2F_1\left(\{\alpha, \alpha+1-\gamma\}, \{\alpha+1-\beta\}, \frac{1}{X}\right) \\ &+ \frac{\Gamma(\gamma)\Gamma(\alpha-\beta)}{\Gamma(\alpha)\Gamma(\gamma-\beta)} (-X)^{-\beta} {}_2F_1\left(\{\beta, \beta+1-\gamma\}, \{\beta+1-\alpha\}, \frac{1}{X}\right), \end{aligned} \quad (\text{B.36})$$

that is valid for $|\text{Arg}(-X)| < \pi$. This sequence transforms the function in (B.18) into,

$$\begin{aligned} \overline{\mathcal{F}}_\lambda(X, Y) &\xrightarrow{Y \rightarrow 0} \frac{-[(1-\epsilon)H_0]^{D-2}}{(4\pi)^{\frac{D}{2}}(D-3)\sin(\pi\lambda)} \left[\frac{\Gamma\left(\frac{4-D}{2}\right)\Gamma\left(\frac{D-3}{2}+\lambda\right)}{2\Gamma\left(\frac{5-D}{2}+\lambda\right)} [e^{i\text{Arg}(-X)}]^{-\frac{D-3}{2}-\lambda} \right. \\ &\quad \times {}_2F_1\left(\left\{\frac{D-3}{2}+\lambda, \frac{D-3}{2}-\lambda\right\}, \left\{\frac{D-2}{2}\right\}, \frac{(1+X)^2}{4X}\right) \\ &\quad \left. + \frac{\Gamma\left(\frac{D-4}{2}\right)}{2} [e^{i\text{Arg}(-X)}]^{-\frac{1}{2}-\lambda} \left[\frac{(1+X)^2}{4X} \right]^{\frac{4-D}{2}} {}_2F_1\left(\left\{\frac{1}{2}+\lambda, \frac{1}{2}-\lambda\right\}, \left\{\frac{6-D}{2}\right\}, \frac{(1+X)^2}{4X}\right) \right]. \end{aligned} \quad (\text{B.37})$$

Applying this result to (B.17), keeping track of $i\delta$ -prescriptions, and applying the identity for gamma functions,

$$\frac{\Gamma(\alpha+\lambda)}{\Gamma(1-\alpha+\lambda)} - \frac{\Gamma(\alpha-\lambda)}{\Gamma(1-\alpha-\lambda)} = -2 \sin(\pi\lambda) \frac{\Gamma(\alpha+\lambda) \Gamma(\alpha-\lambda)}{\Gamma(\frac{1}{2}+\alpha) \Gamma(\frac{1}{2}-\alpha)}, \quad (\text{B.38})$$

then yields the sought spatial coincidence limit,

$$\begin{aligned} \mathcal{M}_\lambda(x; x') &= \frac{-2}{D-3} \times \frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\frac{D-3}{2}+\lambda) \Gamma(\frac{D-3}{2}-\lambda)}{(4\pi)^{\frac{D}{2}} \Gamma(\frac{D-2}{2})} \\ &\times {}_2F_1\left(\left\{\frac{D-3}{2}+\lambda, \frac{D-3}{2}-\lambda\right\}, \left\{\frac{D-2}{2}\right\}, 1+\text{sh}^2\left(\frac{v}{2}\right)\right). \end{aligned} \quad (\text{B.39})$$

This is precisely the spatial coincidence limit one obtains from the series representation (5.45) in Sec. 5.3, that receives contribution only from the $n=0$ term since $[y+4\text{sh}^2(\frac{v}{2})] = (1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|^2$.

Large spatial separations limit for the bulk part. The limit $\|\Delta\vec{x}\| \rightarrow \infty$ is best inferred from the first representation for the bulk solution (B.14). It involves taking the large argument limit of the function (B.15) appearing there,

$$\mathcal{F}_\lambda(X, Y) \stackrel{X, Y \rightarrow \infty}{\sim} \frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\frac{D-3}{2}+\lambda) \Gamma(-\lambda)}{16\pi^{\frac{D+1}{2}} \lambda} (XY)^{-\frac{D-3}{2}-\lambda}, \quad (\text{B.40})$$

after which the limit of (B.14) follows,

$$\begin{aligned} \mathcal{M}_\lambda(x; x') &= \frac{[(1-\epsilon)H_0]^{D-2}}{16\pi^{\frac{D+1}{2}}} \left\{ -\frac{\Gamma(\frac{D-3}{2}) \Gamma(\lambda) \Gamma(-\lambda)}{[(1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|^2]^{\frac{D-3}{2}}} \left[e^{i\pi\lambda} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^\lambda + e^{-i\pi\lambda} \left(\frac{\mathcal{H}}{\mathcal{H}'}\right)^{-\lambda} \right] \right. \\ &\left. + \frac{\Gamma(\frac{D-3}{2}+\lambda) \Gamma(-\lambda)}{\lambda [(1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|^2]^{\frac{D-3}{2}+\lambda}} - \frac{\Gamma(\frac{D-3}{2}-\lambda) \Gamma(\lambda)}{\lambda [(1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|^2]^{\frac{D-3}{2}-\lambda}} \right\}. \end{aligned} \quad (\text{B.41})$$

Deriving this result from the series representation (5.45) might be prohibitively difficult and whenever large spatial separations are called for it is better to appeal to the closed form solution (B.14). Note that, unlike the spatial coincidence limit (B.39), the limit of large spatial separation above exhibits a Coulomb-like tail that is usually generated as a response to a point charge. The same behaviour is exhibited in the Coulomb gauge in flat space (6.21) where the Coulomb-like contribution is present for large spatial separations (with both real and imaginary parts), but disappears in the limit of small separations due to the logarithm that multiplies it.

Temporal coincidence limit for the bulk part. The behaviour close to temporal coincidence $\eta' \rightarrow \eta$ of the bulk part of the Laplace-inverted two point function is best inferred starting from the second representation in (B.17). However, taking this limit

directly is impeded by the first argument tending to one, $Y \rightarrow 1$, where one of the poles of Appell's fourth function is located. The location of this pole descends from the corresponding pole of the hypergeometric equation, which is revealed by examining the single series representation (B.11) applied to Appell's fourth function in (B.18),

$$F_4(\alpha, \beta; \gamma, \gamma'; X^2, Y^2) = \frac{\Gamma(\gamma')}{\Gamma(\alpha)\Gamma(\beta)} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+n)\Gamma(\beta+n)}{\Gamma(\gamma'+n)n!} Y^{2n} \times {}_2F_1\left(\{\alpha+n, \beta+n\}, \{\gamma\}, X^2\right). \quad (\text{B.42})$$

The singular behaviour around the pole at $X \rightarrow 1$ is isolated by applying formula (B.27) for transforming the argument of the hypergeometric function,

$$F_4(\alpha, \beta; \gamma, \gamma'; X^2, Y^2) = \frac{\Gamma(\gamma')}{\Gamma(\alpha)\Gamma(\beta)} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+n)\Gamma(\beta+n)}{\Gamma(\gamma'+n)n!} Y^{2n} \times \left\{ \frac{\Gamma(\gamma)\Gamma(\gamma-\alpha-\beta-2n)}{\Gamma(\gamma-\alpha-n)\Gamma(\gamma-\beta-n)} {}_2F_1\left(\{\alpha+n, \beta+n\}, \{\alpha+\beta-\gamma+1+2n\}, 1-X^2\right) + \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma+2n)}{\Gamma(\alpha+n)\Gamma(\beta+n)} (1-X^2)^{\gamma-\alpha-\beta-2n} \times {}_2F_1\left(\{\gamma-\alpha-n, \gamma-\beta-n\}, \{\gamma-\alpha-\beta+1-2n\}, 1-X^2\right) \right\}. \quad (\text{B.43})$$

Both hypergeometric functions in the resulting expression above are now well behaved in the limit $X \rightarrow 1$. However, the second one is multiplied by more and more negative powers of a small quantity. These powers descend from the behaviour around the pole. We keep all such leading terms in the second series, and resum the series to obtain control over the singular behaviour,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) \stackrel{X \sim 1}{\simeq} \frac{\Gamma(\gamma)\Gamma(\gamma')}{\Gamma(\alpha)\Gamma(\beta)} (1-X^2)^{\gamma-\alpha-\beta} \sum_{n=0}^{\infty} \frac{\Gamma(\alpha+\beta-\gamma+2n)}{\Gamma(\gamma'+n)n!} \left[\frac{Y}{1-X^2}\right]^{2n} = \frac{\Gamma(\gamma)\Gamma(\alpha+\beta-\gamma)}{\Gamma(\alpha)\Gamma(\beta)} (1-X^2)^{\gamma-\alpha-\beta} {}_2F_1\left(\left\{\frac{\alpha+\beta-\gamma}{2}, \frac{\alpha+\beta-\gamma+1}{2}\right\}, \{\gamma'\}, \left[\frac{2Y}{1-X^2}\right]^2\right). \quad (\text{B.44})$$

This result should be followed by another another transformation formula (B.36) that inverts the argument of the hypergeometric function,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) \stackrel{X \sim 1}{\simeq} \frac{\Gamma(\gamma)\Gamma(\gamma')}{2\Gamma(\alpha)\Gamma(\beta)} (1-X^2)^{\gamma-\alpha-\beta} \times \left\{ \frac{\Gamma\left(\frac{\alpha+\beta-\gamma}{2}\right)}{\Gamma\left(\gamma'-\frac{\alpha+\beta-\gamma}{2}\right)} \left(-\left[\frac{Y}{1-X^2}\right]^2\right)^{\frac{\gamma-\alpha-\beta}{2}} \times {}_2F_1\left(\left\{\frac{\alpha+\beta-\gamma}{2}, \frac{\alpha+\beta-\gamma}{2}+1-\gamma'\right\}, \left\{\frac{1}{2}\right\}, \left[\frac{1-X^2}{2Y}\right]^2\right) \right\}$$

$$\begin{aligned}
& - \frac{\Gamma\left(\frac{\alpha+\beta-\gamma+1}{2}\right)}{\Gamma\left(\gamma'-\frac{\alpha+\beta-\gamma+1}{2}\right)} \left(-\left[\frac{Y}{1-X^2}\right]^2\right)^{\frac{\gamma-\alpha-\beta-1}{2}} \\
& \times {}_2F_1\left(\left\{\frac{\alpha+\beta-\gamma+1}{2}, \frac{\alpha+\beta-\gamma+3}{2}-\gamma'\right\}, \left\{\frac{3}{2}\right\}, \left[\frac{1-X^2}{2Y}\right]^2\right). \quad (\text{B.45})
\end{aligned}$$

The last step consists of taking the vanishing argument limit of the hypergeometric functions above, in which they reduce to unity,

$$\begin{aligned}
F_4(\alpha, \beta; \gamma, \gamma'; X, Y) & \stackrel{X \sim 1}{\simeq} \frac{\Gamma(\gamma)\Gamma(\gamma')}{2\Gamma(\alpha)\Gamma(\beta)} (1-X^2)^{\gamma-\alpha-\beta} \\
& \times \left\{ \frac{\Gamma\left(\frac{\alpha+\beta-\gamma}{2}\right)}{\Gamma\left(\gamma'-\frac{\alpha+\beta-\gamma}{2}\right)} \left(-\left[\frac{Y}{1-X^2}\right]^2\right)^{\frac{\gamma-\alpha-\beta}{2}} - \frac{\Gamma\left(\frac{\alpha+\beta-\gamma+1}{2}\right)}{\Gamma\left(\gamma'-\frac{\alpha+\beta-\gamma+1}{2}\right)} \left(-\left[\frac{Y}{1-X^2}\right]^2\right)^{\frac{\gamma-\alpha-\beta-1}{2}} \right\}. \quad (\text{B.46})
\end{aligned}$$

Applying this limit to (B.18) now gives,

$$\begin{aligned}
\overline{\mathcal{F}}_\lambda(X, Y) & \stackrel{X \sim 1}{\simeq} \frac{[(1-\epsilon)H_0]^{D-2}}{32\pi^{\frac{D-1}{2}} \sin(\pi\lambda)} X^{\frac{D-3}{2}+\lambda} (1-X^2)^{4-D} \\
& \times \left\{ \frac{\Gamma\left(\frac{D-3}{2}\right)}{2} \left(-\left[\frac{Y}{1-X^2}\right]^2\right)^{\frac{3-D}{2}} - \frac{\Gamma\left(\frac{D-4}{2}\right)}{\sqrt{\pi}} \left(-\left[\frac{Y}{1-X^2}\right]^2\right)^{\frac{4-D}{2}} \right\}. \quad (\text{B.47})
\end{aligned}$$

Finally, applying this to the representation in (B.17) gives

$$\mathcal{M}_\lambda(x; x') \stackrel{\eta' \rightarrow \eta}{\simeq} \frac{[(1-\epsilon)H_0]^{D-2}}{8\pi^{\frac{D}{2}}} \left[\frac{\sqrt{\pi} \Gamma\left(\frac{D-3}{2}\right) i v}{[(1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|^2]^{\frac{D-3}{2}}} - \frac{\Gamma\left(\frac{D-4}{2}\right)}{[(1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|^2]^{\frac{D-4}{2}}} \right]. \quad (\text{B.48})$$

The imaginary part of this result accounts for the local terms in equations (5.28)–(5.30) for the Laplace-inverted two-point function in the (+) prescription. Namely, two time derivatives acting close to coincidence produce

$$\begin{aligned}
\partial_0 \partial_0 \frac{i^{[+\Xi^+]}}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} \lambda(x; x') & = \partial_0 \partial_0 \left[\theta(\eta-\eta') \frac{i^{[-\Xi^+]}}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} \lambda(x; x') + \theta(\eta'-\eta) \frac{i^{[+\Xi^-]}}{(1-\epsilon)^2 \mathcal{H}\mathcal{H}'} \lambda(x; x') \right] \\
\stackrel{x' \rightarrow x}{\simeq} \partial_0 \partial_0 \left[\frac{\Gamma\left(\frac{D-3}{2}\right)}{8\pi^{\frac{D-1}{2}} \|\Delta\vec{x}\|^{D-3} a^{D-2}} \frac{i|v|}{(1-\epsilon)\sqrt{\mathcal{H}\mathcal{H}'}} \right] & = \frac{\Gamma\left(\frac{D-3}{2}\right) i \delta(\eta-\eta')}{4\pi^{\frac{D-1}{2}} \|\Delta\vec{x}\|^{D-3} a^{D-2}}. \quad (\text{B.49})
\end{aligned}$$

Moreover, the result in (B.48) also allows us to infer the behaviour close to coincidence of the function $i\Upsilon$ defined in (5.24),

$$i\Upsilon \stackrel{\eta' \rightarrow \eta}{\simeq} - \frac{\epsilon}{(1-\epsilon)} \times \frac{[(1-\epsilon)^2 \mathcal{H}\mathcal{H}']^{\frac{D-2}{2}} \Gamma\left(\frac{D-2}{2}\right)}{4\pi^{\frac{D}{2}} [(1-\epsilon)^2 \mathcal{H}\mathcal{H}' \|\Delta\vec{x}\|^2]^{\frac{D-2}{2}}}, \quad (\text{B.50})$$

where the real part of (B.48) contributes, while the contributions of the imaginary part end up canceling.

B.4 Series representation

Here we derive the series solution (5.43) for the Laplace-inverted two-point function starting from the closed form solutions (B.14) and (B.17) for the bulk part, and (B.24) for the infrared part.

Bulk part. The starting point is formula (3.3) from [128] for transforming the parameters of the Appell's fourth function,

$$F_4(\alpha, \beta; \gamma, \gamma'; X, Y) = \frac{-\beta}{\alpha-\beta} F_4(\alpha, \beta+1; \gamma, \gamma'; X, Y) + \frac{\alpha}{\alpha-\beta} F_4(\alpha+1, \beta; \gamma, \gamma'; X, Y). \quad (\text{B.51})$$

Iterating this identity $(N+1)$ times for the special combination of parameters appearing in (B.15) gives,

$$F_4(\alpha, \beta-1; \beta, \beta; X, Y) = \sum_{n=0}^N \frac{(1-\beta) \Gamma(\alpha-\beta+1) \Gamma(\alpha+n)}{\Gamma(\alpha) \Gamma(\alpha-\beta+2+n)} F_4(\alpha+n, \beta; \beta, \beta; X, Y) + \frac{\Gamma(\alpha-\beta+1) \Gamma(\alpha+1+N)}{\Gamma(\alpha) \Gamma(\alpha-\beta+2+N)} F_4(\alpha+1+N, \beta-1; \beta, \beta; X, Y). \quad (\text{B.52})$$

In the end we want to take the limit $N \rightarrow \infty$. Appell's functions in the series part of (B.52) are all in a particular form that can be reduced to rescale propagator functions (2.30) and their derivatives. We do this by modifying the example from Appendix A of [78]. The first step is to apply the reduction formula 9.182.7 from [100],

$$F_4\left(\alpha, \beta; \beta, \beta; \frac{-X}{(1-X)(1-Y)}, \frac{-Y}{(1-X)(1-Y)}\right) = (1-X)^\alpha (1-Y)^\alpha {}_2F_1\left(\{\alpha, \alpha-\beta+1\}, \{\beta\}, XY\right), \quad (\text{B.53})$$

that reduces them all to hypergeometric functions. Then we apply a sequence of three transformation identities for hypergeometric functions; first the quadratic transformation formula (B.35), then the argument transformation formula (B.32), followed by the argument inversion formula (B.36). This sequence transforms the reduction formula (B.53) into,

$$F_4(\alpha, \beta; \beta, \beta; X, Y) = \frac{\Gamma(2\beta-1)}{\Gamma(\beta-\frac{1}{2})} (4\sqrt{XY})^{-\alpha} \times \left\{ \frac{\Gamma(\beta-\alpha-\frac{1}{2})}{\Gamma(2\beta-\alpha-1)} {}_2F_1\left(\{\alpha, \alpha-2\beta+2\}, \{\alpha-\beta+\frac{3}{2}\}, \frac{(\sqrt{X}+\sqrt{Y})^2-1}{4\sqrt{XY}}\right) + \frac{\Gamma(\alpha-\beta+\frac{1}{2})}{\Gamma(\alpha)} \left[\frac{1-(\sqrt{X}+\sqrt{Y})^2}{4\sqrt{XY}} \right]^{\beta-\alpha-\frac{1}{2}} \times {}_2F_1\left(\{\beta-\frac{1}{2}, \frac{3}{2}-\beta\}, \{\beta-\alpha+\frac{1}{2}\}, \frac{(\sqrt{X}+\sqrt{Y})^2-1}{4\sqrt{XY}}\right) \right\}. \quad (\text{B.54})$$

Applying this reduction formula to all the terms in the series in identity (B.52), that we apply to Appell's functions in (B.14) gives after some tedious algebra,

$$\begin{aligned} \mathcal{M}_\lambda(x; x') &= \sum_{n=0}^N \frac{(-1)^n \Gamma(\frac{D-3}{2})}{4 \Gamma(\frac{D-1}{2} + n)} \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right]^n \frac{\partial^n}{\partial y^n} I[\mathcal{F}_\lambda(y)] \\ &\quad + \operatorname{rem}_\lambda^N \left((1-\epsilon) \mathcal{H} \|\Delta \vec{x}\|, (1-\epsilon) \mathcal{H}' \|\Delta \vec{x}\| \right), \end{aligned} \quad (\text{B.55})$$

where we have recognized that

$$\begin{aligned} &\frac{[(1-\epsilon)H_0]^2}{(4\pi)^{\frac{D}{2}}} \left(\frac{-1}{4}\right)^{n-1} \frac{\Gamma(\frac{D-3}{2} + \lambda + n) \Gamma(\frac{D-3}{2} - \lambda + n)}{\Gamma(\frac{D-2}{2} + n)} \\ &\quad \times {}_2F_1\left(\left\{\frac{D-3}{2} + \lambda + n, \frac{D-3}{2} - \lambda + n\right\}, \left\{\frac{D-2}{2} + n\right\}, 1 - \frac{y}{4}\right) = \frac{\partial^n}{\partial y^n} I[\mathcal{F}_\lambda(y)], \end{aligned} \quad (\text{B.56})$$

which follows from the definition of the rescaled propagator function (2.30), and where the remainder in the second line of (B.55) is

$$\begin{aligned} \operatorname{rem}_\lambda^N(X, Y) &= \frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\frac{D-3}{2}) \Gamma(\lambda) \Gamma(-\lambda)}{16\pi^{\frac{D+1}{2}} (XY)^{\frac{D-3}{2}}} \left[-e^{i\pi\lambda} \left(\frac{X}{Y}\right)^\lambda - e^{-i\pi\lambda} \left(\frac{X}{Y}\right)^{-\lambda} \right. \\ &\quad + \frac{\Gamma(\frac{D-1}{2} + \lambda + N) (XY)^{-\lambda}}{\Gamma(\frac{D-1}{2} + N) \Gamma(1 + \lambda)} F_4\left(\frac{D-1}{2} + \lambda + N, \lambda; 1 + \lambda, 1 + \lambda; \frac{1}{X^2}, \frac{1}{Y^2}\right) \\ &\quad \left. + \frac{\Gamma(\frac{D-1}{2} - \lambda + N) (XY)^\lambda}{\Gamma(\frac{D-1}{2} + N) \Gamma(1 - \lambda)} F_4\left(\frac{D-1}{2} - \lambda + N, -\lambda; 1 - \lambda, 1 - \lambda; \frac{1}{X^2}, \frac{1}{Y^2}\right) \right]. \end{aligned} \quad (\text{B.57})$$

It is now clear that the first line of (B.55) reproduces the series representation (5.45) from Sec. 5.3 upon taking the limit $N \rightarrow \infty$. This would imply that the expression in the second line of (B.55) has to vanish in the same limit.

Showing the remainder (B.57) vanishes would be straightforward if asymptotic expansions of Appell's fourth functions for large parameters were readily available. Unfortunately, that is not the case, and there seem to be no readymade results in the literature. Large parameter asymptotic expansions are known for some cases for Appell's first function F_1 [130, 131], so one might hope that expressing the remainder in terms of F_1 using (B.19) and (B.20) would help. Unfortunately, thus obtained representation in terms of F_1 is not covered by the asymptotic expansions in [130, 131]. The necessary expansions are currently work in progress [132].

IR part. Here we show that the IR part (B.24) of the Laplace-inverted two-point function corresponds to the series representation (5.43) and (5.45). We start by writing (B.24) in terms of bi-local variables (2.8), and rewriting it as a primitive

function with respect to y ,

$$\mathcal{X}_\lambda(x; x') = I \left[\frac{[(1-\epsilon)H_0]^{D-2} \Gamma(\lambda) \Gamma(2\lambda)}{(4\pi)^{\frac{D}{2}} \Gamma(\frac{D-1}{2}) \Gamma(\frac{1}{2} + \lambda)} \sum_{N=0}^{\infty} \sum_{n=0}^N \sum_{\ell=0}^{N-n} \frac{(n+1)c_{(N+1)(n+1)\ell}}{(\frac{D-1}{2} + N - \lambda)} \right. \\ \left. \times \left[\frac{k_0^2 e^{-u}}{(1-\epsilon)^2 H_0^2} \right]^{\frac{D-1}{2} - \lambda + N} \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right]^n \operatorname{ch}[(N-n-2\ell)v] \right]. \quad (\text{B.58})$$

Next we derive from the definition of the coefficients (2.48) that,

$$(n+1)c_{(N+1)(n+1)\ell} = -\frac{c_{Nn\ell}}{2(D-1+2n)}. \quad (\text{B.59})$$

If we expand the denominator formally in powers of $2n$,

$$(n+1)c_{(N+1)(n+1)\ell} = -\frac{1}{2(D-1)} \sum_{m=0}^{\infty} \left(\frac{-2}{D-1}\right)^m n^m c_{Nn\ell} \quad (\text{B.60})$$

we can plug it into the triple series, and write the coefficient n^m in terms of derivatives,

$$n^m \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right]^n = \left(\left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right] \frac{\partial}{\partial y} \right)^m \left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right]^n. \quad (\text{B.61})$$

Then, upon recognizing the last two rows as the IR part of the scalar two-point function (2.47),

$$\mathcal{X}_\lambda(y, u, v) = I \left[\sum_{m=0}^{\infty} \left(\frac{-2}{D-1}\right)^m \left(\left[y + 4 \operatorname{sh}^2\left(\frac{v}{2}\right) \right] \frac{\partial}{\partial y} \right)^m \frac{\mathcal{W}_\lambda(y, u, v)}{2(D-1)} \right]. \quad (\text{B.62})$$

This is precisely the IR part written in the intermediate form (5.37). It is written in the final form (5.43) following the procedure given in (5.38)–(5.42).

C Checks for two-point function

Checking explicitly that our result for the photon two-point function given in the covariant tensor basis (5.54) with the structure functions (5.59)–(5.63) satisfies the equations from Sec. 5.1 is important. This is facilitated by the following expressions for derivatives of the bi-local variables,

$$\nabla_\mu(\partial_\nu y) = g_{\mu\nu}(1-\epsilon)H^2(2-y-2\epsilon e^{-v}) - \frac{\epsilon}{1-\epsilon} \left[(\partial_\mu y)(\partial_\nu u) + (\partial_\mu u)(\partial_\nu y) \right], \quad (\text{C.1})$$

$$\nabla_\mu(\partial_\nu u) = -g_{\mu\nu}(1-\epsilon)H^2 - \frac{1+\epsilon}{1-\epsilon} (\partial_\mu u)(\partial_\nu u), \quad \nabla_\mu(\partial'_\nu u) = 0, \quad (\text{C.2})$$

and by the contraction identities for basis tensors given in Table 1. Additionally,

$g^{\mu\nu}(\partial_\mu y)(\partial_\nu y)$	$(1-\epsilon)^2 H^2(4y-y^2)$
$g^{\rho\sigma}(\partial'_\rho y)(\partial'_\sigma y)$	$(1-\epsilon)^2 H'^2(4y-y^2)$
$g^{\mu\nu}(\partial_\mu y)(\partial_\nu u)$	$(1-\epsilon)^2 H^2(2-y-2e^{-v})$
$g^{\rho\sigma}(\partial'_\rho y)(\partial'_\sigma u)$	$(1-\epsilon)^2 H'^2(2-y-2e^v)$
$g^{\mu\nu}(\partial_\mu u)(\partial_\nu u)$	$-(1-\epsilon)^2 H^2$
$g^{\rho\sigma}(\partial'_\rho u)(\partial'_\sigma u)$	$-(1-\epsilon)^2 H'^2$
$g^{\mu\nu}(\partial_\mu y)(\partial_\nu \partial'_\rho y)$	$(1-\epsilon)^2 H^2(2-y)(\partial'_\rho y)$
$g^{\rho\sigma}(\partial_\mu \partial'_\rho y)(\partial'_\sigma y)$	$(1-\epsilon)^2 H'^2(2-y)(\partial_\mu y)$
$g^{\mu\nu}(\partial_\mu u)(\partial_\nu \partial'_\rho y)$	$-(1-\epsilon)^2 H^2[(\partial'_\rho y) + 2e^{-v}(\partial'_\rho u)]$
$g^{\rho\sigma}(\partial_\mu \partial'_\rho y)(\partial'_\sigma u)$	$-(1-\epsilon)^2 H'^2[(\partial_\mu y) + 2e^v(\partial_\mu u)]$
$g^{\mu\nu}(\partial_\mu \partial'_\rho y)(\partial_\nu \partial'_\sigma y)$	$(1-\epsilon)^2 H^2[4(1-\epsilon)^2 H'^2 g'_{\rho\sigma} - (\partial'_\rho y)(\partial'_\sigma y)]$
$g^{\rho\sigma}(\partial_\mu \partial'_\rho y)(\partial_\nu \partial'_\sigma y)$	$(1-\epsilon)^2 H'^2[4(1-\epsilon)^2 H^2 g_{\mu\nu} - (\partial_\mu y)(\partial_\nu y)]$

Table 1. Contractions of tensor structures (table adopted from [73]).

equations from Sec. 2.3.2 satisfied by scalar two-point functions, and equations from Sec. 5.3 satisfied by the Laplace-inverted two-point functions need to be used. These checks are greatly facilitated by computer algebra programs such as *Cadabra* [133–135], and *Wolfram Mathematica* that we made extensive use of here.

Ward-Takahashi identity. We can expand the left-hand-side of the Ward-Takahashi identity (5.5) in the basis of two vectors,

$$\nabla^\mu i[\mathbf{a}_\mu \Delta_\nu^{\mathbf{b}}](x; x') = (1-\epsilon)^2 H^2 \left[(\partial'_\nu y) \mathcal{Z}_1(y, u, v) + (\partial'_\nu u) \mathcal{Z}_2(y, u, v) \right], \quad (\text{C.3})$$

where the two structure functions are expressed in terms of the structure functions of the photon two-point function,

$$\begin{aligned} \mathcal{Z}_1 = & \left[(2-y) \frac{\partial}{\partial y} - \frac{\partial}{\partial u} - \frac{\partial}{\partial v} - D - \frac{(D-2)\epsilon}{(1-\epsilon)} \right] (\mathcal{C}_1 + \mathcal{C}_3 - \bar{\mathcal{C}}_3) - 2e^{-v} \frac{\partial}{\partial y} (\mathcal{C}_3 - \bar{\mathcal{C}}_3) \\ & + \left[(4y-y^2) \frac{\partial}{\partial y} + (D+1)(2-y) + (2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{(D-2)\epsilon}{(1-\epsilon)} \right) \right] \mathcal{C}_2, \end{aligned} \quad (\text{C.4})$$

$$\begin{aligned} \mathcal{Z}_2 = & -2e^{-v} \left[\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{(D-2)\epsilon}{(1-\epsilon)} \right] \mathcal{C}_1 - 2e^{-v} (\mathcal{C}_3 - \bar{\mathcal{C}}_3) \\ & + \left[(4y-y^2) \frac{\partial}{\partial y} + D(2-y) + (2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{(D-2)\epsilon}{(1-\epsilon)} \right) \right] (\mathcal{C}_3 + \bar{\mathcal{C}}_3) \\ & + \left[(2-y-2e^{-v}) \frac{\partial}{\partial y} - \frac{\partial}{\partial u} - \frac{\partial}{\partial v} - (D-1) - \frac{(D-2)\epsilon}{(1-\epsilon)} \right] \mathcal{C}_4. \end{aligned} \quad (\text{C.5})$$

Upon plugging in the photon structure functions (5.59)–(5.63), and applying the equation (5.49) satisfied by $i\Upsilon$, and the recurrence relations (2.53) for the scalar two-point functions, and the equations of motion (2.50) and (2.51) they satisfy, the result simplifies to,

$$(1-\epsilon)^2 H^2 \mathcal{Z}_1 = -\xi_s \frac{\partial}{\partial y} i\Delta_{\nu+1}, \quad (1-\epsilon)^2 H^2 \mathcal{Z}_2 = -\xi_s \left(\frac{\partial}{\partial u} - \frac{\partial}{\partial v} \right) i\Delta_{\nu+1}, \quad (\text{C.6})$$

which exactly reproduces the right-hand side of the Ward-Takahashi identity (5.5).

Equation of motion. The Ward-Takahashi identity that we just checked also simplifies the equation of motion (5.4) satisfied by the two-point functions in the simple covariant gauge,

$$\left(\square \delta_\mu^\rho - R_\mu^\rho \right) i \left[{}_\rho^{\text{a}} \Delta_\nu^{\text{b}} \right] (x; x') = \mathbf{S}^{\text{ab}} g_{\mu\nu} \frac{i\delta^D(x-x')}{\sqrt{-g}} + (1-\xi_s) \partial_\mu \partial'_\nu i \left[{}^{\text{a}} \Delta^{\text{b}} \right]_{\nu+1} (x; x'). \quad (\text{C.7})$$

The left-hand-side can be expanded off-coincidence in the appropriate tensor basis,

$$\begin{aligned} \left(\square \delta_\mu^\rho - R_\mu^\rho \right) i \left[{}_\rho^{\text{a}} \Delta_\nu^{\text{b}} \right] (x; x') &= (1-\epsilon)^2 H^2 \left[(\partial_\mu \partial'_\nu y) \mathcal{E}_1(y, u, v) + (\partial_\mu y) (\partial'_\nu y) \mathcal{E}_2(y, u, v) \right. \\ &\quad + \left[(\partial_\mu y) (\partial'_\nu u) + (\partial_\mu u) (\partial'_\nu y) \right] \mathcal{E}_3(y, u, v) + \left[(\partial_\mu y) (\partial'_\nu u) - (\partial_\mu u) (\partial'_\nu y) \right] \bar{\mathcal{E}}_3(y, u, v) \\ &\quad \left. + (\partial_\mu u) (\partial'_\nu u) \mathcal{E}_4(y, u, v) \right], \end{aligned} \quad (\text{C.8})$$

where the structure functions are found to be,

$$\begin{aligned} \mathcal{E}_1 &= \left[(4y-y^2) \frac{\partial^2}{\partial y^2} + D(2-y) \frac{\partial}{\partial y} + 2(2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{(D-4)\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right. \\ &\quad \left. - \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{D-1-3\epsilon}{1-\epsilon} \right) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} \right) - \frac{D-2\epsilon}{1-\epsilon} \right] \mathcal{C}_1 + \frac{2}{1-\epsilon} \left[(2-y-2\epsilon e^{-v}) \mathcal{C}_2 - \mathcal{C}_3 + \bar{\mathcal{C}}_3 \right], \end{aligned} \quad (\text{C.9})$$

$$\begin{aligned} \mathcal{E}_2 &= \left[(4y-y^2) \frac{\partial^2}{\partial y^2} + (D+4)(2-y) \frac{\partial}{\partial y} + 2(2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{(D-2)\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right. \\ &\quad \left. - \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{D+1-5\epsilon}{1-\epsilon} \right) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} \right) - \frac{2(D-3\epsilon)}{1-\epsilon} \right] \mathcal{C}_2 - \frac{2}{1-\epsilon} \frac{\partial}{\partial y} (\mathcal{C}_1 + \mathcal{C}_3 - \bar{\mathcal{C}}_3), \end{aligned} \quad (\text{C.10})$$

$$\begin{aligned} \mathcal{E}_3 &= - \left[\frac{\epsilon}{1-\epsilon} (2-y+2e^{-v}) \frac{\partial}{\partial y} + \frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{(D-2\epsilon)\epsilon}{(1-\epsilon)^2} \right] \mathcal{C}_1 \\ &\quad - \left[\frac{\epsilon}{1-\epsilon} \left((4y-y^2) \frac{\partial}{\partial y} + D(2-y) \right) - (2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{(1+(D-3)\epsilon)\epsilon}{(1-\epsilon)^2} \right) \right] \mathcal{C}_2 \\ &\quad + \left[(4y-y^2) \frac{\partial^2}{\partial y^2} + (D+2)(2-y) \frac{\partial}{\partial y} + 2(2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{1-(D-3)\epsilon}{2(1-\epsilon)} \right) \frac{\partial}{\partial y} \right. \end{aligned}$$

$$\begin{aligned}
& - \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{D-4\epsilon}{1-\epsilon} \right) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} \right) - \frac{(D-1)-(2D+1)\epsilon+4\epsilon^2}{(1-\epsilon)^2} \Big] \mathcal{C}_3 \\
& + \left[\frac{1+\epsilon}{1-\epsilon} (2-y-2e^{-v}) \frac{\partial}{\partial y} + \frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{(D-1-\epsilon)\epsilon}{(1-\epsilon)^2} - \frac{1+\epsilon}{1-\epsilon} \right] \bar{\mathcal{C}}_3 - \frac{1}{1-\epsilon} \frac{\partial \mathcal{C}_4}{\partial y}, \quad (\text{C.11})
\end{aligned}$$

$$\begin{aligned}
\bar{\mathcal{E}}_3 = & \left[\frac{\epsilon}{(1-\epsilon)} (2-y-2e^{-v}) \frac{\partial}{\partial y} + \frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{(D-2\epsilon)\epsilon}{(1-\epsilon)^2} \right] \mathcal{C}_1 + \left[\frac{\epsilon}{(1-\epsilon)} (4y-y^2) \frac{\partial}{\partial y} \right. \\
& - (2-y+2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{(D-3+\epsilon)\epsilon}{(1-\epsilon)^2} \right) + 2(2-y) \frac{(D-1-\epsilon)\epsilon}{(1-\epsilon)^2} \Big] \mathcal{C}_2 \\
& + \left[\frac{1+\epsilon}{1-\epsilon} (2-y-2e^{-v}) \frac{\partial}{\partial y} + \frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{(D-1-\epsilon)\epsilon}{(1-\epsilon)^2} - \frac{1+\epsilon}{1-\epsilon} \right] \mathcal{C}_3 - \frac{1}{1-\epsilon} \frac{\partial \mathcal{C}_4}{\partial y} \\
& + \left[(4y-y^2) \frac{\partial^2}{\partial y^2} + (D+2)(2-y) \frac{\partial}{\partial y} + 2(2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{1-(D-3)\epsilon}{2(1-\epsilon)} \right) \right] \frac{\partial}{\partial y} \\
& - \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{D-4\epsilon}{1-\epsilon} \right) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} \right) - \frac{(D-1)-(2D+1)\epsilon+4\epsilon^2}{(1-\epsilon)^2} \Big] \bar{\mathcal{C}}_3, \quad (\text{C.12})
\end{aligned}$$

$$\begin{aligned}
\mathcal{E}_4 = & - \left[\frac{2\epsilon}{1-\epsilon} (4y-y^2) \frac{\partial}{\partial y} - 2(2-y) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{(D-2\epsilon)\epsilon}{(1-\epsilon)^2} \right) \right] (\mathcal{C}_3 + \bar{\mathcal{C}}_3) \\
& - 4e^{-v} \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{2-(D-4)\epsilon}{2(1-\epsilon)} \right) (\mathcal{C}_3 - \bar{\mathcal{C}}_3) + \frac{2(D-2)(1+\epsilon)\epsilon}{(1-\epsilon)^2} e^{-v} (\mathcal{C}_1 + \mathcal{C}_3 + \bar{\mathcal{C}}_3) \\
& + \left[(4y-y^2) \frac{\partial^2}{\partial y^2} + D(2-y) \frac{\partial}{\partial y} + 2(2-y-2e^{-v}) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} - \frac{2-(D-4)\epsilon}{2(1-\epsilon)} \right) \right] \frac{\partial}{\partial y} \\
& - \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} + \frac{D-1-3\epsilon}{1-\epsilon} \right) \left(\frac{\partial}{\partial u} + \frac{\partial}{\partial v} \right) + \frac{2(D-1-\epsilon)\epsilon}{(1-\epsilon)^2} \Big] \mathcal{C}_4. \quad (\text{C.13})
\end{aligned}$$

Upon plugging in the photon structure functions (5.59)–(5.63), and applying judiciously equations (5.49)–(5.52) satisfied by the function $i\Upsilon$ defined in (5.24), and equations (2.50)–(2.53) satisfied by the scalar two-point functions the expressions above evaluate to,

$$(1-\epsilon)^2 H^2 \mathcal{E}_1 = (1-\xi_s) \frac{\partial}{\partial y} i\Delta_{\nu+1}, \quad (\text{C.14a})$$

$$(1-\epsilon)^2 H^2 \mathcal{E}_2 = (1-\xi_s) \frac{\partial^2}{\partial y^2} i\Delta_{\nu+1}, \quad (\text{C.14b})$$

$$(1-\epsilon)^2 H^2 \mathcal{E}_3 = (1-\xi_s) \frac{\partial^2}{\partial y \partial u} i\Delta_{\nu+1}, \quad (\text{C.14c})$$

$$(1-\epsilon)^2 H^2 \bar{\mathcal{E}}_3 = - (1-\xi_s) \frac{\partial^2}{\partial y \partial v} i\Delta_{\nu+1}, \quad (\text{C.14d})$$

$$(1-\epsilon)^2 H^2 \mathcal{E}_4 = (1-\xi_s) \left(\frac{\partial^2}{\partial u^2} - \frac{\partial^2}{\partial v^2} \right) i\Delta_{\nu+1}. \quad (\text{C.14e})$$

These precisely account for the right-hand-side of the equation of motion (C.7) off-coincidence.

In order to check that the local terms on the right-hand-side of (C.7) are also correctly reproduced for the case of the Feynman (and consequently Dyson) propagator

it is sufficient to consider only the most singular terms from each photon structure function (5.59)–(5.63). For scalar propagators these are isolated easily using the power-series representation (2.32) for the bulk part, as the IR series (2.47) does not contribute. From there we find that the most singular term is just the conformal two-point function,

$$i\Delta_\nu(y, u, v) \stackrel{x' \rightarrow x}{\sim} i\Delta_{1/2}(y, u) = e^{-\frac{(D-2)\epsilon}{2(1-\epsilon)}u} \mathcal{F}_{1/2}(y) = \frac{[(1-\epsilon)^2 H H']^{\frac{D-2}{2}} \Gamma(\frac{D-2}{2})}{(4\pi)^{\frac{D}{2}}} \left(\frac{y}{4}\right)^{-\frac{D-2}{2}}. \quad (\text{C.15})$$

The leading singular behaviour for the function $i\Upsilon$ the leading singular can be inferred from the result in (B.50) by substituting $(1-\epsilon)^2 \mathcal{H} \mathcal{H}' \|\Delta \vec{x}\|^2 \rightarrow y$ in the denominator,

$$i\Upsilon(y, u, v) \stackrel{x' \rightarrow x}{\sim} -\frac{\epsilon}{1-\epsilon} i\Delta_{1/2}(y, u), \quad (\text{C.16})$$

which can be confirmed by showing that this result satisfies equations (5.49)–(5.52) close to coincidence. The leading singular behaviour of the photon structure functions is therefore captured by

$$\mathcal{C}_1 \stackrel{x' \rightarrow x}{\sim} \frac{(D-2)}{2(1-\epsilon)\nu_T} \mathcal{C}, \quad \mathcal{C}_2 \stackrel{x' \rightarrow x}{\sim} \frac{(1+\epsilon)}{2(1-\epsilon)\nu_T} \frac{\partial}{\partial y} \mathcal{C}, \quad (\text{C.17a})$$

$$\mathcal{C}_3 \stackrel{x' \rightarrow x}{\sim} \frac{(D-2)\epsilon}{4(1-\epsilon)\nu_T} \mathcal{C}, \quad \bar{\mathcal{C}}_3 \stackrel{x' \rightarrow x}{\sim} 0 \times v \mathcal{C}, \quad \mathcal{C}_4 \stackrel{x' \rightarrow x}{\sim} 0 \times \mathcal{C}, \quad (\text{C.17b})$$

where,

$$\mathcal{C}(y, u) = -\frac{i\Delta_{1/2}(y, u)}{2(1-\epsilon)^2 H H'}. \quad (\text{C.18})$$

We can then apply a simple derivative identity,

$$\begin{aligned} (\partial_\mu y) (\partial'_\nu y) f(y, u) &= \partial_\mu \partial'_\nu I^2[f(y, u)] - (\partial_\mu \partial'_\nu y) I[f(y, u)] \\ &- \left[(\partial_\mu y) (\partial'_\nu u) + (\partial_\mu u) (\partial'_\nu y) \right] \frac{\partial}{\partial u} I[f(y, u)] - (\partial_\mu u) (\partial'_\nu u) \frac{\partial^2}{\partial u^2} I^2[f(y, u)], \end{aligned} \quad (\text{C.19})$$

to the second tensor structure to rewrite the most singular piece of the photon propagator in a particularly convenient form,

$$\begin{aligned} i[\mu^+ \Delta_\nu^+] (x; x') \stackrel{x' \rightarrow x}{\sim} & \frac{(1+\epsilon)}{2(1-\epsilon)\nu_T} \partial_\mu \partial'_\nu I[\mathcal{C}] + (\partial_\mu \partial'_\nu y) \mathcal{C} \\ & + \left[(\partial_\mu y) (\partial'_\nu u) + (\partial_\mu u) (\partial'_\nu y) \right] \frac{\epsilon \mathcal{C}}{1-\epsilon}, \end{aligned} \quad (\text{C.20})$$

where we have neglected an irrelevant piece where the fourth tensor structure is multiplied by $y^{-\frac{D-4}{2}}$, which is not sufficiently singular to produce a local contribution in the equation of motion. We then use the identities which generate the temporal delta function,

$$\partial_0 \partial_0 y_{++} = \partial_0 \partial_0 y + 4(1-\epsilon)^2 \mathcal{H}^2 \times i\delta \times \delta(\eta - \eta'), \quad (\text{C.21a})$$

$$\partial_0 \partial'_0 y_{++} = \partial_0 \partial'_0 y - 4(1-\epsilon)^2 \mathcal{H}^2 \times i\delta \times \delta(\eta - \eta'), \quad (\text{C.21b})$$

and the covariant D -dimensional delta function,

$$-4(1-\epsilon)^2 H^2 \times i\delta \times \delta(\eta-\eta') \times \frac{\partial}{\partial y} i\Delta_{1/2}(y_{++}, u_{++}) = \frac{i\delta^D(x-x')}{\sqrt{-g}}, \quad (\text{C.22})$$

to compute the action of the photon kinetic operator from (5.3) onto the most singular part of the propagator (C.20),

$$\mathcal{D}_\mu{}^\rho \left\{ \partial_\rho \partial'_\nu I[\mathcal{C}](y_{++}, u_{++}) \right\} \overset{x' \rightarrow x}{\sim} -\frac{2}{\xi_s} (a^2 \delta_\mu^0 \delta_\nu^0) \frac{i\delta^D(x-x')}{\sqrt{-g}}, \quad (\text{C.23})$$

$$\mathcal{D}_\mu{}^\rho \left\{ (\partial_\rho \partial'_\nu y) \mathcal{C}(y_{++}, u_{++}) \right\} \overset{x' \rightarrow x}{\sim} \left[g_{\mu\nu} + \left(1 - \frac{1}{\xi_s}\right) (a^2 \delta_\mu^0 \delta_\nu^0) \right] \frac{i\delta^D(x-x')}{\sqrt{-g}}, \quad (\text{C.24})$$

$$\mathcal{D}_\mu{}^\rho \left\{ \left[(\partial_\rho y_{++}) (\partial'_\nu u_{++}) + (\partial_\rho u_{++}) (\partial'_\nu y_{++}) \right] \mathcal{C}(y_{++}, u_{++}) \right\} \overset{x' \rightarrow x}{\sim} 0. \quad (\text{C.25})$$

When combined, these relations correctly reproduce the local terms in the Feynman propagator equation of motion (5.4).

References

- [1] F. L. Bezrukov and M. Shaposhnikov, “The Standard Model Higgs boson as the inflaton,” *Phys. Lett. B* **659** (2008), 703-706 [arXiv:0710.3755 [hep-th]].
- [2] D. P. George, S. Mooij and M. Postma, “Effective action for the Abelian Higgs model in FLRW,” *JCAP* **11** (2012), 043 [arXiv:1207.6963 [hep-th]].
- [3] F. Bezrukov, “The Higgs field as an inflaton,” *Class. Quant. Grav.* **30** (2013), 214001 [arXiv:1307.0708 [hep-ph]].
- [4] D. P. George, S. Mooij and M. Postma, “Quantum corrections in Higgs inflation: the real scalar case,” *JCAP* **02** (2014), 024 [arXiv:1310.2157 [hep-th]].
- [5] I. Obata, T. Miura and J. Soda, “Dynamics of Electroweak Gauge Fields during and after Higgs Inflation,” *Phys. Rev. D* **90** (2014) no.4, 045005 [arXiv:1405.3091 [hep-th]].
- [6] F. Bezrukov, J. Rubio and M. Shaposhnikov, “Living beyond the edge: Higgs inflation and vacuum metastability,” *Phys. Rev. D* **92** (2015) no.8, 083512 [arXiv:1412.3811 [hep-ph]].
- [7] D. P. George, S. Mooij and M. Postma, “Quantum corrections in Higgs inflation: the Standard Model case,” *JCAP* **04** (2016), 006 [arXiv:1508.04660 [hep-th]].
- [8] Y. Ema, R. Jinno, K. Mukaida and K. Nakayama, “Violent Preheating in Inflation with Nonminimal Coupling,” *JCAP* **02** (2017), 045 [arXiv:1609.05209 [hep-ph]].
- [9] P. Adshead, J. T. Giblin and Z. J. Weiner, “Non-Abelian gauge preheating,” *Phys. Rev. D* **96** (2017) no.12, 123512 [arXiv:1708.02944 [hep-ph]].

- [10] E. I. Sfakianakis and J. van de Vis, “Preheating after Higgs Inflation: Self-Resonance and Gauge boson production,” *Phys. Rev. D* **99** (2019) no.8, 083519 [arXiv:1810.01304 [hep-ph]].
- [11] P. Adshead, J. T. Giblin, R. Grutkoski and Z. J. Weiner, “Gauge preheating with full general relativity,” *JCAP* **03** (2024), 017 [arXiv:2311.01504 [astro-ph.CO]].
- [12] E. Pajer and M. Peloso, “A review of Axion Inflation in the era of Planck,” *Class. Quant. Grav.* **30** (2013), 214002 [arXiv:1305.3557 [hep-th]].
- [13] P. Adshead, J. T. Giblin, T. R. Scully and E. I. Sfakianakis, “Gauge-preheating and the end of axion inflation,” *JCAP* **12** (2015), 034 [arXiv:1502.06506 [astro-ph.CO]].
- [14] P. Adshead, J. T. Giblin, T. R. Scully and E. I. Sfakianakis, “Magnetogenesis from axion inflation,” *JCAP* **10** (2016), 039 [arXiv:1606.08474 [astro-ph.CO]].
- [15] V. Domcke and K. Mukaida, “Gauge Field and Fermion Production during Axion Inflation,” *JCAP* **11** (2018), 020 [arXiv:1806.08769 [hep-ph]].
- [16] E. V. Gorbar, K. Schmitz, O. O. Sobol and S. I. Vilchinskii, “Gauge-field production during axion inflation in the gradient expansion formalism,” *Phys. Rev. D* **104** (2021) no.12, 123504 [arXiv:2109.01651 [hep-ph]].
- [17] A. Caravano, E. Komatsu, K. D. Lozanov and J. Weller, “Lattice simulations of Abelian gauge fields coupled to axions during inflation,” *Phys. Rev. D* **105** (2022) no.12, 123530 [arXiv:2110.10695 [astro-ph.CO]].
- [18] D. G. Figueroa, J. Lizarraga, A. Urrio and J. Urrestilla, “Strong Backreaction Regime in Axion Inflation,” *Phys. Rev. Lett.* **131** (2023) no.15, 151003 [arXiv:2303.17436 [astro-ph.CO]].
- [19] B. Ratra, “Cosmological ‘seed’ magnetic field from inflation,” *Astrophys. J. Lett.* **391** (1992), L1-L4
- [20] L. Parker, “Particle creation in expanding universes,” *Phys. Rev. Lett.* **21** (1968), 562-564.
- [21] L. Parker, “Quantized fields and particle creation in expanding universes. 1.,” *Phys. Rev.* **183** (1969), 1057-1068.
- [22] L. Parker, “Quantized fields and particle creation in expanding universes. 2.,” *Phys. Rev. D* **3** (1971), 346-356 [erratum: *Phys. Rev. D* **3** (1971), 2546-2546].
- [23] V. F. Mukhanov and G. V. Chibisov, “Quantum Fluctuations and a Nonsingular Universe,” *JETP Lett.* **33** (1981), 532-535 [*Pisma Zh. Eksp. Teor. Fiz.* **33** (1981) 549-553]
- [24] A. A. Starobinsky, “Spectrum of relict gravitational radiation and the early state of the universe,” *JETP Lett.* **30** (1979), 682-685 [*Pisma Zh. Eksp. Teor. Fiz.* **30** (1979) 719-723]
- [25] E. O. Kahya and R. P. Woodard, “Charged scalar self-mass during inflation,” *Phys. Rev. D* **72** (2005), 104001 [arXiv:gr-qc/0508015 [gr-qc]].

- [26] E. O. Kahya and R. P. Woodard, “One Loop Corrected Mode Functions for SQED during Inflation,” *Phys. Rev. D* **74** (2006), 084012 [arXiv:gr-qc/0608049 [gr-qc]].
- [27] T. Prokopec, N. C. Tsamis and R. P. Woodard, “Two Loop Scalar Bilinears for Inflationary SQED,” *Class. Quant. Grav.* **24** (2007), 201-230 [arXiv:gr-qc/0607094 [gr-qc]].
- [28] X. Chen, Y. Wang and Z. Z. Xianyu, “Standard Model Background of the Cosmological Collider,” *Phys. Rev. Lett.* **118** (2017) no.26, 261302 [arXiv:1610.06597 [hep-th]].
- [29] F. Fabián González and T. Prokopec, “Renormalization group approach to scalar quantum electrodynamics on de Sitter,” [arXiv:1611.07854 [gr-qc]].
- [30] X. Chen, Y. Wang and Z. Z. Xianyu, “Standard Model Mass Spectrum in Inflationary Universe,” *JHEP* **04** (2017), 058 [arXiv:1612.08122 [hep-th]].
- [31] F. K. Popov, “Debye mass in de Sitter space,” *JHEP* **06** (2018), 033 [arXiv:1711.11010 [hep-th]].
- [32] D. Glavan and G. Rigopoulos, “One-loop electromagnetic correlators of SQED in power-law inflation,” *JCAP* **02** (2021), 021 [arXiv:1909.11741 [gr-qc]].
- [33] T. Prokopec, O. Tornkvist and R. P. Woodard, “Photon mass from inflation,” *Phys. Rev. Lett.* **89** (2002), 101301 [arXiv:astro-ph/0205331 [astro-ph]].
- [34] T. Prokopec, O. Tornkvist and R. P. Woodard, “One loop vacuum polarization in a locally de Sitter background,” *Annals Phys.* **303** (2003), 251-274 [arXiv:gr-qc/0205130 [gr-qc]].
- [35] T. Prokopec and R. P. Woodard, “Vacuum polarization and photon mass in inflation,” *Am. J. Phys.* **72** (2004), 60-72 [arXiv:astro-ph/0303358 [astro-ph]].
- [36] T. Prokopec and E. Puchwein, “Photon mass generation during inflation: de Sitter invariant case,” *JCAP* **04** (2004), 007 [arXiv:astro-ph/0312274 [astro-ph]].
- [37] T. Prokopec, N. C. Tsamis and R. P. Woodard, “Stochastic Inflationary Scalar Electrodynamics,” *Annals Phys.* **323** (2008), 1324-1360 [arXiv:0707.0847 [gr-qc]].
- [38] T. Prokopec, N. C. Tsamis and R. P. Woodard, “Two loop stress-energy tensor for inflationary scalar electrodynamics,” *Phys. Rev. D* **78** (2008), 043523 [arXiv:0802.3673 [gr-qc]].
- [39] A. A. Starobinsky, “STOCHASTIC DE SITTER (INFLATIONARY) STAGE IN THE EARLY UNIVERSE,” *Lect. Notes Phys.* **246** (1986), 107-126
- [40] A. A. Starobinsky and J. Yokoyama, “Equilibrium state of a selfinteracting scalar field in the De Sitter background,” *Phys. Rev. D* **50** (1994), 6357-6368 [arXiv:astro-ph/9407016 [astro-ph]].
- [41] M. S. Turner and L. M. Widrow, “Inflation Produced, Large Scale Magnetic Fields,” *Phys. Rev. D* **37** (1988), 2743

- [42] E. A. Calzetta, A. Kandus and F. D. Mazzitelli, “Primordial magnetic fields induced by cosmological particle creation,” *Phys. Rev. D* **57** (1998), 7139-7144 [arXiv:astro-ph/9707220 [astro-ph]].
- [43] M. Giovannini and M. E. Shaposhnikov, “Primordial magnetic fields from inflation?,” *Phys. Rev. D* **62** (2000), 103512 [arXiv:hep-ph/0004269 [hep-ph]].
- [44] T. Prokopec, “Cosmological magnetic fields from photon coupling to fermions and bosons in inflation,” [arXiv:astro-ph/0106247 [astro-ph]].
- [45] A. C. Davis, K. Dimopoulos, T. Prokopec and O. Tornkvist, “Primordial spectrum of gauge fields from inflation,” *Phys. Lett. B* **501** (2001), 165-172 [arXiv:astro-ph/0007214 [astro-ph]].
- [46] R. Durrer and A. Neronov, “Cosmological Magnetic Fields: Their Generation, Evolution and Observation,” *Astron. Astrophys. Rev.* **21** (2013), 62 [arXiv:1303.7121 [astro-ph.CO]].
- [47] B. L. Hu and K. Shiokawa, “Wave propagation in stochastic space-times: Localization, amplification and particle creation,” *Phys. Rev. D* **57** (1998), 3474-3483 [arXiv:gr-qc/9708023 [gr-qc]].
- [48] D. A. R. Dalvit, F. D. Mazzitelli and C. Molina-Paris, “One loop graviton corrections to Maxwell’s equations,” *Phys. Rev. D* **63** (2001), 084023 [arXiv:hep-th/0010229 [hep-th]].
- [49] K. E. Leonard, T. Prokopec and R. P. Woodard, “Covariant Vacuum Polarizations on de Sitter Background,” *Phys. Rev. D* **87** (2013) no.4, 044030 [arXiv:1210.6968 [gr-qc]].
- [50] K. E. Leonard and R. P. Woodard, “Graviton Corrections to Vacuum Polarization during Inflation,” *Class. Quant. Grav.* **31** (2014), 015010 [arXiv:1304.7265 [gr-qc]].
- [51] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, “Electrodynamic Effects of Inflationary Gravitons,” *Class. Quant. Grav.* **31** (2014), 175002 [arXiv:1308.3453 [gr-qc]].
- [52] C. L. Wang and R. P. Woodard, “Excitation of Photons by Inflationary Gravitons,” *Phys. Rev. D* **91** (2015) no.12, 124054 [arXiv:1408.1448 [gr-qc]].
- [53] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, “Graviton Loop Corrections to Vacuum Polarization in de Sitter in a General Covariant Gauge,” *Class. Quant. Grav.* **32** (2015) no.19, 195014 [arXiv:1504.00894 [gr-qc]].
- [54] C. L. Wang and R. P. Woodard, “One-loop quantum electrodynamic correction to the gravitational potentials on de Sitter spacetime,” *Phys. Rev. D* **92** (2015), 084008 [arXiv:1508.01564 [gr-qc]].
- [55] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, “One loop graviton corrections to dynamical photons in de Sitter,” *Class. Quant. Grav.* **34** (2017) no.8, 085002 [arXiv:1609.00386 [gr-qc]].

- [56] S. P. Miao, T. Prokopec and R. P. Woodard, “Scalar enhancement of the photon electric field by the tail of the graviton propagator,” *Phys. Rev. D* **98** (2018) no.2, 025022 [arXiv:1806.00742 [gr-qc]].
- [57] D. Glavan, S. P. Miao, T. Prokopec and R. P. Woodard, “Explaining large electromagnetic logarithms from loops of inflationary gravitons,” *JHEP* **08** (2023), 195 [arXiv:2307.09386 [gr-qc]].
- [58] Y. Akrami *et al.* [Planck], “Planck 2018 results. X. Constraints on inflation,” *Astron. Astrophys.* **641** (2020), A10 [arXiv:1807.06211 [astro-ph.CO]].
- [59] F. Lucchin and S. Matarrese, “Power Law Inflation,” *Phys. Rev. D* **32** (1985), 1316.
- [60] D. La and P. J. Steinhardt, “Extended Inflationary Cosmology,” *Phys. Rev. Lett.* **62** (1989), 376 [erratum: *Phys. Rev. Lett.* **62** (1989), 1066].
- [61] J. F. Koksma and T. Prokopec, “Fermion Propagator in Cosmological Spaces with Constant Deceleration,” *Class. Quant. Grav.* **26** (2009), 125003 [arXiv:0901.4674 [gr-qc]].
- [62] A. Bilandzic and T. Prokopec, “Quantum radiative corrections to slow-roll inflation,” *Phys. Rev. D* **76** (2007), 103507 [arXiv:0704.1905 [astro-ph]].
- [63] T. M. Janssen, S. P. Miao, T. Prokopec and R. P. Woodard, “The Hubble Effective Potential,” *JCAP* **05** (2009), 003 [arXiv:0904.1151 [gr-qc]].
- [64] M. Herranen, T. Markkanen and A. Tranberg, “Quantum corrections to scalar field dynamics in a slow-roll space-time,” *JHEP* **05** (2014), 026 [arXiv:1311.5532 [hep-ph]].
- [65] S. P. Miao and R. P. Woodard, “Fine Tuning May Not Be Enough,” *JCAP* **09** (2015), 022 [arXiv:1506.07306 [astro-ph.CO]].
- [66] T. Markkanen, “Light scalars on cosmological backgrounds,” *JHEP* **01** (2018), 116 [arXiv:1711.07502 [gr-qc]].
- [67] M. B. Fröb, “One-loop quantum gravitational backreaction on the local Hubble rate,” *Class. Quant. Grav.* **36** (2019) no.9, 095010 [arXiv:1806.11124 [gr-qc]].
- [68] S. P. Miao, S. Park and R. P. Woodard, “Ricci Subtraction for Cosmological Coleman-Weinberg Potentials,” *Phys. Rev. D* **100** (2019) no.10, 103503 [arXiv:1908.05558 [gr-qc]].
- [69] A. Kyriazis, S. P. Miao, N. C. Tsamis and R. P. Woodard, “Inflaton effective potential for general ϵ ,” *Phys. Rev. D* **102** (2020) no.2, 025024 [arXiv:1908.03814 [gr-qc]].
- [70] S. Katuwal, S. P. Miao and R. P. Woodard, “Inflaton effective potential from photons for general ϵ ,” *Phys. Rev. D* **103** (2021) no.10, 105007 [arXiv:2101.06760 [gr-qc]].
- [71] W. C. C. Lima, “Graviton backreaction on the local cosmological expansion in slow-roll inflation,” *Class. Quant. Grav.* **38** (2021) no.13, 135015 [arXiv:2007.04995 [gr-qc]].

- [72] D. Glavan and T. Prokopec, “When tadpoles matter: one-loop corrections for spectator Higgs in inflation,” *JHEP* **10** (2023), 063 [arXiv:2306.11162 [hep-ph]].
- [73] D. Glavan, A. Marunović, T. Prokopec and Z. Zahraee, “Abelian Higgs model in power-law inflation: the propagators in the unitary gauge,” *JHEP* **09** (2020), 165 [arXiv:2005.05435 [gr-qc]].
- [74] D. Glavan and T. Prokopec, “Photon propagator in de Sitter space in the general covariant gauge,” *JHEP* **05** (2023), 126 [arXiv:2212.13982 [gr-qc]].
- [75] D. Glavan and T. Prokopec, “Even the photon propagator must break de Sitter symmetry,” *Phys. Lett. B* **841** (2023), 137928 [arXiv:2212.13997 [hep-th]].
- [76] N. C. Tsamis and R. P. Woodard, “A Maximally symmetric vector propagator,” *J. Math. Phys.* **48** (2007), 052306 [arXiv:gr-qc/0608069 [gr-qc]].
- [77] B. Allen and T. Jacobson, “Vector Two Point Functions in Maximally Symmetric Spaces,” *Commun. Math. Phys.* **103** (1986), 669.
- [78] M. B. Fröb and A. Higuchi, “Mode-sum construction of the two-point functions for the Stueckelberg vector fields in the Poincaré patch of de Sitter space,” *J. Math. Phys.* **55** (2014), 062301 [arXiv:1305.3421 [gr-qc]].
- [79] D. Glavan, “Photon quantization in cosmological spaces,” *Phys. Rev. D* **109** (2024) no.8, 085014 [arXiv:2212.13975 [hep-th]].
- [80] A. Youssef, “Infrared behavior and gauge artifacts in de Sitter spacetime: The photon field,” *Phys. Rev. Lett.* **107** (2011), 021101 [arXiv:1011.3755 [gr-qc]].
- [81] N. Rendell, “Large-distance behavior of the massless vector two-point function in de Sitter spacetime,” *Int. J. Mod. Phys. D* **27** (2018) no.11, 1843005 [arXiv:1802.00687 [gr-qc]].
- [82] N. A. Chernikov and E. A. Tagirov, “Quantum theory of scalar fields in de Sitter space-time,” *Ann. Inst. H. Poincaré Phys. Theor. A* **9** (1968), 109.
- [83] T. S. Bunch and P. C. W. Davies, “Quantum Field Theory in de Sitter Space: Renormalization by Point Splitting,” *Proc. Roy. Soc. Lond. A* **360** (1978), 117-134.
- [84] F. W. J. Olver, D. W. Lozier, R. F. Boisvert and C. W. Clark, eds., “NIST Handbook of Mathematical Functions,” (Cambridge University Press, Cambridge, 2010).
- [85] F. W. J. Olver, A. B. Olde Daalhuis, D. W. Lozier, B. I. Schneider, R. F. Boisvert, C. W. Clark, B. R. Miller, and B. V. Saunders, eds., “NIST Digital Library of Mathematical Functions,” <https://dlmf.nist.gov/>, Release 1.1.10 of 2023-06-15.
- [86] J. S. Schwinger, “Brownian motion of a quantum oscillator,” *J. Math. Phys.* **2** (1961), 407-432.
- [87] K. T. Mahanthappa, “Multiple production of photons in quantum electrodynamics,” *Phys. Rev.* **126** (1962), 329-340.

- [88] P. M. Bakshi and K. T. Mahanthappa, “Expectation value formalism in quantum field theory. 1.,” *J. Math. Phys.* **4** (1963), 1-11.
- [89] P. M. Bakshi and K. T. Mahanthappa, “Expectation value formalism in quantum field theory. 2.,” *J. Math. Phys.* **4** (1963), 12-16.
- [90] L. V. Keldysh, “Diagram technique for nonequilibrium processes,” *Zh. Eksp. Teor. Fiz.* **47** (1964), 1515-1527; *Sov. Phys. JETP* **20** (1965), 1018.
- [91] R. D. Jordan, “Effective Field Equations for Expectation Values,” *Phys. Rev. D* **33** (1986), 444-454.
- [92] E. Calzetta and B. L. Hu, “Closed Time Path Functional Formalism in Curved Space-Time: Application to Cosmological Back Reaction Problems,” *Phys. Rev. D* **35** (1987), 495.
- [93] J. Berges, “Introduction to nonequilibrium quantum field theory,” *AIP Conf. Proc.* **739** (2004) no.1, 3-62 [arXiv:hep-ph/0409233 [hep-ph]].
- [94] D. Glavan and T. Prokopec, “A pedestrian introduction to non-equilibrium QFT,” <https://webpace.science.uu.nl/proko101/LecturenotesNonEquilQFT.pdf>.
- [95] L. H. Ford and L. Parker, “Infrared Divergences in a Class of Robertson-Walker Universes,” *Phys. Rev. D* **16** (1977), 245-250.
- [96] B. Allen, “Vacuum States in de Sitter Space,” *Phys. Rev. D* **32** (1985), 3136
- [97] B. Allen and A. Folacci, “The Massless Minimally Coupled Scalar Field in De Sitter Space,” *Phys. Rev. D* **35** (1987), 3771.
- [98] T. M. Janssen and T. Prokopec, “Regulating the infrared by mode matching: A Massless scalar in expanding spaces with constant deceleration,” *Phys. Rev. D* **83** (2011), 084035 [arXiv:0906.0666 [gr-qc]].
- [99] T. Janssen and T. Prokopec, “A Graviton propagator for inflation,” *Class. Quant. Grav.* **25** (2008), 055007 [arXiv:0707.3919 [gr-qc]].
- [100] I. S. Gradshteyn and I. M. Ryzhik, “Table of integrals, series, and products,” seventh edition, edited by A. Jeffrey and D. Zwillinger, (Elsevier/Academic Press, Amsterdam, Netherlands, 2007).
- [101] A. P. Prudnikov, Yu. A. Brychkov and O. I. Marichev, “Integrals and Series, Volume 2: Special Functions,” (Gordon and Breach, United States, 1986).
- [102] T. M. Janssen, S. P. Miao, T. Prokopec and R. P. Woodard, “Infrared Propagator Corrections for Constant Deceleration,” *Class. Quant. Grav.* **25** (2008), 245013 [arXiv:0808.2449 [gr-qc]].
- [103] V. Moncrief, “Space-Time Symmetries and Linearization Stability of the Einstein Equations. 2.,” *J. Math. Phys.* **17** (1976), 1893-1902.
- [104] J. M. Arms, “LINEARIZATION STABILITY OF GRAVITATIONAL AND GAUGE FIELDS,” *J. Math. Phys.* **20** (1979), 443-453.

- [105] A. Higuchi, “Quantum linearization instabilities of de Sitter space-time. 1,” *Class. Quant. Grav.* **8** (1991), 1961-1981.
- [106] A. Higuchi, “Quantum linearization instabilities of de Sitter space-time. 2,” *Class. Quant. Grav.* **8** (1991), 1983-2004
- [107] N. C. Tsamis and R. P. Woodard, “The Structure of perturbative quantum gravity on a De Sitter background,” *Commun. Math. Phys.* **162** (1994), 217-248
- [108] D. Marolf, “Refined algebraic quantization: Systems with a single constraint,” [arXiv:gr-qc/9508015 [gr-qc]].
- [109] D. Marolf and I. Morrison, “Group Averaging of massless scalar fields in 1+1 de Sitter,” *Class. Quant. Grav.* **26** (2009), 035001 [arXiv:0808.2174 [gr-qc]].
- [110] D. Marolf and I. A. Morrison, “Group Averaging for de Sitter free fields,” *Class. Quant. Grav.* **26** (2009), 235003 [arXiv:0810.5163 [gr-qc]].
- [111] S. P. Miao, N. C. Tsamis and R. P. Woodard, “Transforming to Lorentz Gauge on de Sitter,” *J. Math. Phys.* **50** (2009) 122502 [arXiv:0907.4930 [gr-qc]].
- [112] J. Gibbons and A. Higuchi, “Removing the Faddeev-Popov zero modes from Yang-Mills theory in spacetimes with compact spatial sections,” *Phys. Rev. D* **91** (2015) no.2, 024006 [arXiv:1410.7830 [hep-th]].
- [113] E. Altas, E. Kilicarslan and B. Tekin, “Einstein–Yang–Mills theory: gauge invariant charges and linearization instability,” *Eur. Phys. J. C* **81** (2021) no.7, 648 [arXiv:2105.11744 [hep-th]].
- [114] S. N. Gupta, “Theory of longitudinal photons in quantum electrodynamics,” *Proc. Phys. Soc. A* **63** (1950) 681.
- [115] K. Bleuler, “A New method of treatment of the longitudinal and scalar photons,” *Helv. Phys. Acta* **23** (1950) 567.
- [116] F. Falceto, “Canonical quantization of the electromagnetic field in arbitrary ξ -gauge,” [arXiv:2211.16870 [hep-th]].
- [117] I. I. Cotaescu and C. Crucean, “The Quantum theory of the free Maxwell field on the de Sitter expanding universe,” *Prog. Theor. Phys.* **124** (2010), 1051-1066 [arXiv:0806.2515 [gr-qc]].
- [118] A. Belokogne and A. Folacci, “Stueckelberg massive electromagnetism in curved spacetime: Hadamard renormalization of the stress-energy tensor and the Casimir effect,” *Phys. Rev. D* **93** (2016) no.4, 044063 [arXiv:1512.06326 [gr-qc]].
- [119] A. Belokogne, A. Folacci and J. Queva, “Stueckelberg massive electromagnetism in de Sitter and anti-de Sitter spacetimes: Two-point functions and renormalized stress-energy tensors,” *Phys. Rev. D* **94** (2016) no.10, 105028 [arXiv:1610.00244 [gr-qc]].
- [120] D. M. Capper and M. J. Duff, “Trace anomalies in dimensional regularization,” *Nuovo Cim. A* **23** (1974), 173-183.

- [121] L. S. Brown and J. P. Cassidy, “Stress Tensor Trace Anomaly in a Gravitational Metric: General Theory, Maxwell Field,” *Phys. Rev. D* **15** (1977), 2810.
- [122] S. L. Adler, J. Lieberman and Y. J. Ng, “Regularization of the Stress Energy Tensor for Vector and Scalar Particles Propagating in a General Background Metric,” *Annals Phys.* **106** (1977), 279.
- [123] L. S. Brown and J. P. Cassidy, “Stress Tensors and their Trace Anomalies in Conformally Flat Space-Times,” *Phys. Rev. D* **16** (1977), 1712.
- [124] R. Endo, “Gauge Dependence of the Gravitational Conformal Anomaly for the Electromagnetic Field,” *Prog. Theor. Phys.* **71** (1984), 1366.
- [125] Yu. A. Brychkov, “Higher derivatives of the Bessel functions with respect to the order,” *Integral Transforms Spec. Funct.* **27** (2016), 566.
- [126] S. O. Rice, “On Contour Integrals for the Product of Two Bessel Functions,” *The Quarterly Journal of Mathematics* os-6:1 (1935), 52-64.
- [127] W. N. Bailey, “Some Infinite Integrals Involving Bessel Functions,” *Proc. London Math. Soc.* s2-40 (1936), 37-48.
- [128] Yu. A. Brychkov and Nasser Saad, “On some formulas for the Appell function $F_4(a,b;c,c';w,z)$,” *Integral Transforms Spec. Funct.* 28:9 (2017), 629-644.
- [129] H. Exton, “On the system of partial differential equations associated with Appell’s function F_4 ,” *J. Phys. A: Math. Gen.* **28** (1995), 631.
- [130] J. L. López, P. Pagola and E. Pérez Sinusía, “Asymptotics of the first Appell function F_1 with large parameters,” *Integral Transforms and Special Functions* 24:9 (2013), 715-733.
- [131] J. L. López, P. Pagola and E. Pérez Sinusía, “Asymptotics of the first Appell function F_1 with large parameters II,” *Integral Transforms and Special Functions* 24:12 (2013), 982-999.
- [132] J. L. López et al., *in progress*.
- [133] K. Peeters, “Introducing Cadabra: A Symbolic computer algebra system for field theory problems,” [arXiv:hep-th/0701238 [hep-th]].
- [134] K. Peeters, “A Field-theory motivated approach to symbolic computer algebra,” *Comput. Phys. Commun.* **176** (2007), 550-558 [arXiv:cs/0608005 [cs.SC]].
- [135] K. Peeters, “Cadabra2: computer algebra for field theory revisited,” *J. Open Source Softw.* **3** (2018) no.32, 1118.