Large deviations of current for the symmetric simple exclusion process on a semi-infinite line and on an infinite line with a slow bond

Kapil Sharma,^{1,*} Soumyabrata Saha,^{1,†} Sandeep Jangid,^{1,‡} and Tridib Sadhu^{1,§}

¹Department of Theoretical Physics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India

(Dated: May 14, 2024)

Two of the most influential exact results in classical one-dimensional diffusive transport are about current statistics for the symmetric simple exclusion process in the stationary state on a finite line coupled with two unequal reservoirs at the boundary, and in the non-stationary state on an infinite line. We present the corresponding result for the intermediate geometry of a semi-infinite line coupled with a single reservoir. This result is obtained using the fluctuating hydrodynamics approach of macroscopic fluctuation theory and confirmed by rare event simulations using a cloning algorithm. Our exact result enables us to address the corresponding problem on an infinite line in the presence of a slow bond and several related problems.

Keywords: large deviations; symmetric simple exclusion process; slow bond

Current fluctuations in non-equilibrium transport have long been a subject of interest in both classical and quantum contexts [1-7]. The major interest in these investigations lies in the full counting statistics in terms of large deviations of current. Apart from being a contender for extension of free energy for non-equilibrium systems, large deviation function (ldf) in general can characterize various peculiarities of non-equilibrium conditions, such as non-local response, emergent symmetries, and low-dimensional phase transitions [3, 8–14]. However, estimating ldf poses a major challenge, often requiring specialized integrability techniques [4, 15, 16] tailored to curated models or clever numerical sampling schemes of rare-probability events [17–21]. Understandably, exact results about ldf play an important role in the landscape of non-equilibrium physics, providing a benchmark for affirming qualitative predictions of approximate methods. Our work in this *Letter* presents a non-trivial addition to this list of exact results, fostering solutions to certain relevant problems [22–24].

Among the widely studied stochastic models of classical transport, is the symmetric simple exclusion process (SSEP) [3, 25–27]. The SSEP, along with its driven variants, has attained the status of a paradigmatic model in non-equilibrium statistical physics [16, 28]. Two celebrated exact results for the SSEP concern the large deviations of current on a finite lattice coupled with two unequal reservoirs, and on an infinite lattice starting with a non-stationary state. The two geometries represent distinct non-equilibrium scenarios: a stationary state for finite systems, and a time-dependent state relaxing towards an asymptotic equilibrium state for infinite systems. They illuminate crucial differences in their fluctuations. The finite system, in the long run, holds no memory of the initial state, while the infinite system exhibits an unusual dependence on the initial state even at long times [29, 30].

The large deviations of current in these two geometries were obtained using the additivity conjecture [31-

33] and integrability methods, such as the diagonalization of tilted matrix [34] or the matrix product states [4, 35] for the finite lattice, and the Bethe ansatz [36] for the infinite lattice. These microscopic results were subsequently verified [2, 37, 38] using a fluctuating hydrodynamics framework [36, 39, 40]. This framework, now famously known as the macroscopic fluctuation theory (MFT) [2, 41], emerged in the early 2000s from the seminal works of Bertini, De Sole, Gabrielli, Jona-Lasinio, and Landim, which presented a general approach for characterizing non-equilibrium fluctuations of diffusive systems. MFT has successfully led to exact results of large deviations in exclusion processes [2, 37–39, 42] and related transport models [39, 43–45].

In this *Letter*, we consider the intermediate scenario: the SSEP on a semi-infinite lattice [46, 47] coupled to a boundary reservoir with a density that is different from the initial bulk density of the system. This elucidates a non-equilibrium regime evolving into an asymptotic state in equilibrium with the reservoir, although the dynamical quantities remain sensitive to the initial conditions [48]. Only limited results [48–51] are known for this intermediate geometry, particularly because extending the aforementioned integrability methods for this geometry proves challenging. Even a solution via MFT has remained elusive.

We overcame these difficulties of the semi-infiniteproblem by uncovering a novel mapping to the infiniteline problem that led to an exact result for the ldf of current for SSEP on the semi-infinite geometry, which is our main result. Additionally, this exact result enables us to characterize current fluctuations in the presence of a slow bond, both in the semi-infinite and infinite geometries. The mapping to the infinite-problem is possible due to certain symmetries of the Euler-Lagrange equations and the boundary conditions of the associated variational problem within MFT. These techniques, however, could not be obviously exploited within microscopic approaches. Our exact result is a testament to the power



FIG. 1. The SSEP on a semi-infinite lattice coupled with a reservoir of density ρ_a with a coupling strength $\gamma > 0$.

of MFT for non-equilibrium systems that are otherwise formidable to approach.

Model and main result: The SSEP on a semi-infinite lattice (see Fig. 1), with sites indexed by $i = 1, 2, \cdots$, is composed of continuous-time hard-core random walkers hopping to adjacent sites with a unit rate provided the target site is empty. At the boundary site i = 1, particles are injected following exclusion with rate $\gamma \rho_a$ and removed at rate $\gamma(1-\rho_a)$, which models [3] coupling with a reservoir of density ρ_a . For each site *i* at a given time τ , we assign a binary occupation number $n_i(\tau)$ that takes values 0 and 1, depending on whether the site is empty or occupied, respectively. Initially, sites are filled following Bernoulli distribution with a uniform average density ρ_b . Our main result concerns the time-integrated current, Q_T , which represents the total flux of particles from the reservoir into the system over a time-period T. In the hydrodynamic description [52], expressed in terms of coarse-grained density $\rho(i/\sqrt{T}, \tau/T) \simeq n_i(\tau)$, the flux

$$Q_T = \sqrt{T} \int_0^\infty \mathrm{d}x \left[\rho(x,1) - \rho(x,0)\right] \tag{1}$$

measures the net change in the number of particles in the system. In the large T limit, its generating function has the asymptotics

$$\langle e^{\lambda Q_T} \rangle \sim e^{\sqrt{T}\mu_{\rm si}(\lambda,\rho_a,\rho_b)}.$$
 (2a)

Here, the scaled cumulant generating function (scgf) $\mu_{\rm si}(\lambda, \rho_a, \rho_b)$ (subscript 'si' denotes semi-infinite) is given by

$$\mu_{\rm si}(\lambda,\rho_a,\rho_b) = \int_{-\infty}^{\infty} \frac{\mathrm{d}k}{2\pi} \log\left[1 + 4\,\omega\,(1+\omega)\,\mathrm{e}^{-k^2}\right] \quad (2\mathrm{b})$$

where ω is a function of the parameters defined as [30, 36]

$$\omega(\lambda,\rho_a,\rho_b) = \rho_a(1-\rho_b)(\mathrm{e}^{\lambda}-1) + \rho_b(1-\rho_a)(\mathrm{e}^{-\lambda}-1).$$
(2c)

This dependence of scgf on λ , ρ_a , and ρ_b through a single function ω arises from a symmetry of the underlying dynamics of the SSEP [34, 36, 53]. The expression (2b) is consistent with our earlier [48] result in the low-density limit. It is instructive to compare (2b) with the corresponding result for the infinite line [36]. The scgf (2b) encapsulates all cumulants of Q_T in the large T limit. While the first three cumulants were initially reported in [48], we present here the fourth cumulant for $\rho_a = \rho_b = \rho$

$$\frac{\langle Q_T^4 \rangle_c}{\sqrt{T}} \simeq \frac{4}{\sqrt{\pi}} \rho \left(1 - \rho\right) \left[1 - 12\left(\sqrt{2} - 1\right) \rho \left(1 - \rho\right)\right].$$
(3)

Similar to the infinite line case [29], the scgf (2b) also admits the Gallavotti-Cohen-type fluctuation symmetry

$$\mu_{\rm si}(\lambda,\rho_a,\rho_b) = \mu_{\rm si}\bigg(\log\frac{\rho_b\left(1-\rho_a\right)}{\rho_a\left(1-\rho_b\right)} - \lambda,\rho_a,\rho_b\bigg).$$
(4)

Remarkably, the expression (2b) admits a Fredholm determinant representation

$$\mu_{\rm si}(\lambda,\rho_a,\rho_b) = \frac{1}{2\pi} \log \det \left[\mathbb{1} + 4\omega(1+\omega)K(x,y)\right], \quad (5)$$

with the kernel $K(x, y) = \delta(x - y) \exp(\partial_x^2)$. Analogous Fredholm determinant structure is observed in related contexts for the current distribution in the TASEP [54] and the ASEP [55, 56]. This representation underscores a connection between TASEP and random matrix theory, as emphasized by Johansson [54]. However, a similar relationship for the SSEP is not apparent. Interestingly, an alternative representation of (2b),

$$\mu_{\rm si}(\lambda, \rho_a, \rho_b) = \frac{1}{2\sqrt{\pi}} \sum_{n=1}^{\infty} \frac{(-1)^{n+1}}{n^{3/2}} \left[4\,\omega\,(1+\omega)\right]^n \quad (6)$$

bears a striking resemblance [57] to the scqf for the number of surviving particles in an assembly of annihilating random walkers (see eq. (9.106) in [15]).

Numerical verification: For confirming our result (2b) based on the hydrodynamic description of SSEP, we have independently generated the scgf using numerical simulation based on a continuous-time cloning algorithm [17, 18, 52] with 10^5 clones and measured over duration T = 250. The simulation result, plotted in Fig. 2, shows a good agreement with our theoretical result (2b) for a reasonably large value of λ . Deviations emerging at larger values of λ are a consequence of finite-size effects [52].

LDF: The scaling in (2a) corresponds to a large deviation asymptotics of the probability

$$P(Q_T = j\sqrt{T}) \sim e^{-\sqrt{T}\phi_{\rm si}(j)}$$
(7)

where $\phi_{\rm si}(j)$ is the ldf, related to $\mu_{\rm si}(\lambda)$ (reference to $\rho_{a(b)}$ is ignored) by a Legendre-Fenchel transformation [11] $\phi_{\rm si}(j) = \max_{\lambda} [j\lambda - \mu_{\rm si}(\lambda)]$. Using the transformation, it is immediate that (4) reflects the symmetry

$$\phi_{\rm si}(j) - \phi_{\rm si}(-j) = j \log \frac{\rho_b (1 - \rho_a)}{\rho_a (1 - \rho_b)}.$$
 (8)



FIG. 2. Solid red line represents the theoretical result of scgf (2b) for $\rho_{a(b)} = 0.5$, with blue dots representing the corresponding simulation result obtained by the cloning algorithm. The results closely match in the range $-1.5 \le \lambda \le 1.5$, with the deviations shown in the inset. For comparison, the corresponding scgf for infinite line μ_{inf} [36] is shown in black dashed line.

Similarly, the asymptotics $\mu_{\rm si}(\lambda, \rho_a, \rho_b) \simeq \frac{4\sqrt{2}}{3\pi} (\log \omega)^{3/2}$ of (2b) for large positive λ correspond [36] to the asymptotics of the ldf

$$\phi_{\rm si}(j) \simeq \frac{\pi^2 j^3}{24} - j \log \left(\rho_a \left(1 - \rho_b\right)\right)$$
 (9)

for large positive j. A similar analysis for large negative j gives asymptotics (9) with $j \rightarrow -j$ and $\rho_a \leftrightarrow \rho_b$. Notably, for the non-equilibrium condition $\rho_a \neq \rho_b$, the ldf is skewed.

The symmetry relation (8) and the asymptotics (9) are confirmed in the plot of the ldf in Fig. 3, generated by numerically evaluating the Legendre-Fenchel transformation of the scgf (2b).

Derivation: In the following, we outline our derivation of (2b) within the fluctuating hydrodynamics framework of MFT. The crucial idea behind MFT is to recognize the relevant hydrodynamic modes for a coarse-grained description of the dynamics and characterize the probability of their evolution in terms of an Action, which is analogous to the Martin-Siggia-Rose-Janssen-De Dominicis (MSRJD) Action [58–61] of the associated fluctuating hydrodynamics equation. For SSEP, the relevant hydrodynamic mode is the locally conserved density $\rho(x, t)$ evolving by [48, 52]

$$\partial_t \rho = \partial_x^2 \rho + \frac{1}{T^{1/4}} \,\partial_x \left(\sqrt{\sigma(\rho)} \,\eta \right) \tag{10}$$

where $\sigma(\rho) = 2\rho(1-\rho)$, and $\eta(x,t)$ is a delta-correlated Gaussian white noise with unit covariance. Correspond-



FIG. 3. The ldf indicated in solid red line, with the super-Gaussian asymptotes (9) for large |j| indicated in blue dotdashed lines. The black dashed line denotes the Gaussian ldf with the appropriate mean $\langle Q_T \rangle$ and variance $\langle Q_T^2 \rangle_c$. The plots are for $\rho_a = 0.6$ and $\rho_b = 0.3$. The inset confirms the symmetry (8).

ing MSRJD-Action on the semi-infinite line is [48, 52]

$$S_0 = \mathcal{F} + \int_0^1 \mathrm{d}t \int_0^\infty \mathrm{d}x \left[\widehat{\rho} \,\partial_t \rho - \frac{\sigma(\rho) \,(\partial_x \widehat{\rho})^2}{2} + \partial_x \rho \,\partial_x \widehat{\rho} \right]$$
(11)

where $\hat{\rho}$ is the response field and \mathcal{F} incorporates contributions from fluctuations in the initial state [29, 48]. Within this description, the generating function $\langle e^{\lambda Q_T} \rangle = \int \mathcal{D}[\rho, \hat{\rho}] e^{-\sqrt{T}S_{\lambda}[\rho, \hat{\rho}]}$ with $S_{\lambda} = S_0 - \lambda Q_T / \sqrt{T}$ and Q_T in (1).

For large T, the path-integral is dominated by a saddle point, leading to (2a) with

$$\mu_{\rm si}(\lambda) \simeq -\min_{\rho,\widehat{\rho}} S_{\lambda}\left[\rho,\widehat{\rho}\right] \equiv -S_{\lambda}\left[q_{\rm si}, p_{\rm si}\right]$$
(12)

(reference to $\rho_{a(b)}$ is ignored) where $(q_{\rm si}, p_{\rm si})$ is the least-Action path for $(\rho, \hat{\rho})$. This way, the problem reduces [29, 48] to solving the corresponding Euler-Lagrange equations

$$\partial_t q_{\rm si} = \partial_x^2 q_{\rm si} - \partial_x [\sigma(q_{\rm si}) \partial_x p_{\rm si}], \quad \text{and}$$
(13a)

$$\partial_t p_{\rm si} = -\partial_x^2 p_{\rm si} - \frac{\sigma(q_{\rm si})}{2} (\partial_x p_{\rm si})^2 \tag{13b}$$

in the semi-infinite domain x > 0, with the temporal boundary conditions

$$p_{\rm si}(x,0) = \lambda + \int_{\rho_b}^{q_{\rm si}(x,0)} \frac{2\,\mathrm{d}r}{\sigma(r)} \quad \text{and} \quad p_{\rm si}(x,1) = \lambda, \ (14)$$

where the integral in $p_{\rm si}(x, 0)$ is the contribution [29] from \mathcal{F} for the initial state of the SSEP with Bernoulli measure. Additional spatial boundary conditions,

$$q_{\rm si}(0,t) = \rho_a \quad \text{and} \quad p_{\rm si}(0,t) = 0$$
 (15)

are due to the "fast-coupling" [38, 52] with reservoir.

This variational problem is reminiscent of the corresponding problem on the infinite line [29] for a Bernoullimeasured initial state with an average density ρ_a for x < 0 and ρ_b for x > 0. The only differences between the two problems are the domain x and the boundary conditions. The infinite line problem was formulated in [29] within MFT, which was recently solved in [37] by identifying an ingenious mapping to the classical integrable system and employing the inverse scattering method. Despite having small differences, the semi-infinite case poses a new nontrivial problem [62], which is incredibly difficult to solve.

A remarkable simplification arises for the special choice of initial density pair $(\rho_a, \rho_b) \equiv (1/2, 0)$ for the semiinfinite line problem. For this choice of densities, there are no fluctuations in the initial state, which amounts to setting $\mathcal{F} = 0$ in (11) with the condition

$$q_{\rm si}(x,0) = 0. \tag{16}$$

The Euler-Lagrange equations for the corresponding variational problem (12) remains the same as in (13). The only difference comes in the initial condition in (14), which is now replaced by (16). This initial condition corresponds to the quenched averaging [29, 38].

We now show that this quenched semi-infinite line problem has a direct mapping to the quenched infinite line problem with $(\rho_a, \rho_b) \equiv (1, 0)$ and fugacity $\tilde{\lambda} = 2\lambda$. For the latter problem, the Euler-Lagrange equation is the same [29] as in (13), but now on the entire real line x with the temporal boundary conditions

$$q_{\inf}(x,0) = \theta(-x)$$
 and $p_{\inf}(x,1) = \lambda \theta(x)$ (17)

(subscript 'inf' denotes infinite). It is trivial to check that the solution admits the symmetry

$$q_{\inf}(x,t) = 1 - q_{\inf}(-x,t)$$
 (18a)

$$p_{\inf}(x,t) = \tilde{\lambda} - p_{\inf}(-x,t)$$
(18b)

which fixes the value of the fields at the origin

$$q_{\inf}(0,t) = \frac{1}{2}$$
 and $p_{\inf}(0,t) = \frac{\tilde{\lambda}}{2}$ (19)

at all times 0 < t < T. This conclusion is an essential part of our observation, as now, the solution $q_{inf}(x, t)$ on positive line x satisfies the boundary conditions (15, 16) of the semi-infinite line problem with $(\rho_a, \rho_b) \equiv (1/2, 0)$. For a similar correspondence of the response field, we define

$$\tilde{p}(x,t) = p_{\inf}(x,t) - \frac{\tilde{\lambda}}{2}$$
(20)

which too now replicates the boundary conditions $\tilde{p}(x, 1) = \lambda$ and $\tilde{p}(0, t) = 0$ of the semi-infinite line problem with $\tilde{\lambda} = 2\lambda$. The fields $(q_{\text{inf}}, \tilde{p})$ also satisfy the same Euler-Lagrange equations (13). Consequently, the least-Action path $(q_{\rm si}, p_{\rm si})$ for the semi-infinite line problem with $(\rho_a, \rho_b) \equiv (1/2, 0)$ and fugacity λ is related to the corresponding path $(q_{\rm inf}, p_{\rm inf})$ for the infinite line problem with $(\rho_a, \rho_b) \equiv (1, 0)$ and fugacity 2λ by

$$q_{\rm si}(x,t) = q_{\rm inf}(x,t)$$
 and $p_{\rm si}(x,t) = p_{\rm inf}(x,t) - \lambda$ (21)

for $x \ge 0$ at all times. This correspondence relates the least-Action of the two problems resulting [52] in our crucial observation:

$$\mu_{\rm si}\left(\lambda, \frac{1}{2}, 0\right) = \frac{1}{2}\,\mu_{\rm inf}(2\lambda, 1, 0) \tag{22}$$

where the pre-factor 1/2 comes from the half-domain of integration of x in (11). The latter scgf is known from the seminal work [36] of Derrida and Gerschenfeld, which culminates in

$$\mu_{\rm si}\left(\lambda, \frac{1}{2}, 0\right) = \int_{-\infty}^{\infty} \frac{\mathrm{d}k}{2\pi} \log\left[1 + (\mathrm{e}^{2\lambda} - 1)\mathrm{e}^{-k^2}\right].$$
(23)

The result (23) is for the specific initial density pair $(\rho_a, \rho_b) \equiv (1/2, 0)$. For extending the result for other densities, we invoke a well-known rotational symmetry [29, 52, 53] of the least-Action (12). Essentially, the least-Action paths for two sets of parameters $(\lambda, \rho_a, \rho_b)$ and $(\lambda', \rho_a', \rho_b')$ are related under a canonical transformation [29, 52, 53]. The symmetry results [36, 52, 53] in a dependence of the scgf on $(\lambda, \rho_a, \rho_b)$ through a single parameter

$$\mu_{\rm si}(\lambda,\rho_a,\rho_b) = R_{\rm si}\big(\omega(\lambda,\rho_a,\rho_b)\big) \tag{24}$$

where ω is defined in (2c). This dependence enables us [52] to deduce the function $R_{\rm si}(\omega)$ from the result (23), leading to the expression (2b).

Slow bond: Recent interests [33, 38, 48, 63–67] in studying the effects of the coupling strength with reservoir can also be addressed in the semi-infinite line problem. The result (2b) is independent [48] of the coupling strength γ (see Fig. 1) as long as it is larger than $\mathcal{O}(T^{-1/2})$. This is seen from the corresponding result for slow coupling $\gamma = \Gamma/\sqrt{T}$, where the boundaryfluctuations are significant, modifying the scgf (2b) to

$$\mu_{\rm si}^{\rm slow}(\lambda,\rho_a,\rho_b) = \min_{z} \left[\Gamma \sinh^2 (z-u) + R_{\rm si}(\sinh^2 z) \right]$$
(25)
with $\sinh^2 u = \omega(\lambda,\rho_a,\rho_b)$. In the $\Gamma \to \infty$ limit, (2b) is
recovered from (25).

The expression (25) is obtained following an additivity argument [33], where contributions in (25) are separately from the single bond joining the reservoir and the system, and the system itself, optimised over the density at their common site [52]. This construction is very similar to the discussion in [33] for a related context, except for a crucial distinction that unlike the latter example, there is



FIG. 4. The solid lines denote the scgf (25) for $\rho_{a(b)} = 0.5$ and different values of Γ (green for $\Gamma = 1$, blue for $\Gamma = 2$, and red for $\Gamma \to \infty$). The markers denote the corresponding numerical results obtained by the cloning algorithm with T = 500 and $N_c = 10^4$

no quasi-stationarity for the semi-infinite problem. Nevertheless, the additivity conjecture gives the correct result (25), as verified in Fig. 4.

A similar additivity argument helps to solve the problem of current fluctuation across a single slow bond on the infinite lattice [24, 68, 69]. For the hopping rate $\gamma = \Gamma/\sqrt{T}$ across the slow bond, and unity for rest of the lattice, the generating function of current $\langle e^{\lambda Q_T} \rangle \sim e^{\sqrt{T}\mu_{inf}^{slow}(\lambda,\rho_a,\rho_b)}$, for large T, with the scgf

$$\mu_{\inf}^{\text{slow}}(\lambda, \rho_a, \rho_b) = \min_{z_a, z_b} \left[R_{\text{si}} \left(\sinh^2 z_a \right) + \Gamma \sinh^2 \left(z_a + z_b - u \right) + R_{\text{si}} \left(\sinh^2 z_b \right) \right]$$
(26)

where $\sinh^2 u = \omega(\lambda, \rho_a, \rho_b)$. In the $\Gamma \to \infty$ limit, the celebrated infinite line result [36, 37] is recovered [52] from (26).

Other applications: Several other interesting conclusions can be drawn from our result. A known symmetry [29] of the MFT-Action extends our exact results to models [70–72] with a quadratic mobility $\sigma(\rho) = 2A\rho(B-\rho)$, where A and B are arbitrary constants, culminating in $\mu_{\sigma}(\lambda, \rho_{a}, \rho_{b}) = (1/A) \mu_{\text{ssep}}(\omega(AB\lambda, \rho_{a}/B, \rho_{b}/B))$.

Our semi-infinite line result provides immediate solutions for two closely related problems. The first problem [23] concerns an infinite line SSEP with fast injection at a single site, a scenario which gained prominence in related contexts [73–75]. At long-times t, the net injection of particles N_t on an empty lattice follows the asymptotics $\langle e^{\lambda N_t} \rangle \sim \exp \left[2 \sqrt{t} \, \mu_{\rm si}(\lambda, 1, 0)\right]$ with (2b), which results from the statistical independence of the two halves due to fast injection. The second problem concerns the survival probability $P_t(k)$ of the $k^{\rm th}$ particle (tracer) up to time t in a fully packed SSEP on \mathbb{Z}^+ in the presence of an absorbing site at the origin. The asymptotics (7) and the relation $P_t(k) = P(Q_t > k)$ with the cumulative probability of integrated current result in a stretched exponential decay $P_t(k) \sim \exp(-\sqrt{t/\tau_s})$ for $k \leq 2\sqrt{t/\pi}$ at long-times t, with $\tau_s = [\phi(-k/\sqrt{t})]^{-2}$.

Similar stretched exponential decays were noted in kinetically constrained models [22, 76–78], such as the energy-conserving spin-flip dynamics [22], where spin auto-correlation decays as $e^{-\sqrt{t/\tau}}$, with τ remaining undetermined. For the spin-model, it was realized [78] that the domain-wall dynamics is equivalent to the infiniteline SSEP at a uniform density ρ . This correspondence relates [79] the spin-auto-correlation to the generating function of current in SSEP leading [52] to an exact result $\tau = [\mu(i\pi, \rho, \rho)]^{-2}$, where $\mu(\lambda, \rho_a, \rho_b)$ is the scqf of the infinite line problem [36]. Similar correspondence holds in the presence of slow bond in the spin-flip dynamics, relating to (26).

Conclusions: There remain several related open problems of immediate interest. Most prominent among them is the scgf for fixed (quenched) initial states at arbitrary densities. The only available non-trivial result [29] is for the infinite line SSEP at half-filling, obtained by establishing a relation with the fluctuating (annealed) initial state. A similar mapping for the semi-infinite line yields the quenched scgf

$$\mu_{\rm si}^{\rm que}\left(\lambda, \frac{1}{2}, \frac{1}{2}\right) = \int_{-\infty}^{\infty} \frac{\mathrm{d}k}{\sqrt{8}\pi} \log\left[1 + \sinh^2\left(\lambda\right) \mathrm{e}^{-k^2}\right]. \tag{27}$$

Another pressing open question is the least-Action path at arbitrary densities for the semi-infinite line. In broader contexts, an extension of the problem in higher dimensions would be interesting, where only limited results are available [80]. From a practical point of view, the emergence of the infinite line SSEP in quantum circuits [7] is exciting, and a semi-infinite analog would be worth exploring. Along similar lines, extensions of our results for quantum analogues of the SSEP [81, 82] or for integrable models using generalized hydrodynamics [83, 84] would be timely.

We acknowledge the financial support of the Department of Atomic Energy, Government of India, under Project Identification No. RTI 4002. TS thanks Bernard Derrida for insightful discussions on the problem. The importance of semi-infinite line SSEP and the additivity ansatz for slow bond originated from those discussions. TS also thanks Mustansir Barma for introducing him to the works on stretched exponential decays, particularly the open problem in [22].

^{*} kapil.sharma@tifr.res.in

[†] soumyabrata.saha@tifr.res.in

[‡] sandeep.jangid@tifr.res.in

[§] tridib@theory.tifr.res.in

- M. Prähofer and H. Spohn, Current fluctuations for the totally asymmetric simple exclusion process, in *In and Out of Equilibrium: Probability with a Physics Flavor*, Vol. 51, edited by V. Sidoravicius (Birkhäuser Boston, 2002) p. 185–204.
- [2] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, Current fluctuations in stochastic lattice gases, Phys. Rev. Lett. **94**, 030601 (2005); Non equilibrium current fluctuations in stochastic lattice gases, J. Stat. Phys. **123**, 237–276 (2006).
- [3] B. Derrida, Non-equilibrium steady states: Fluctuations and large deviations of the density and of the current, J. Stat. Mech. 2007, P07023 (2007).
- [4] A. Lazarescu, The physicist's companion to current fluctuations: One-dimensional bulk-driven lattice gases, J. Phys. A: Math. Theor. 48, 503001 (2015).
- [5] T. Banerjee, S. N. Majumdar, A. Rosso, and G. Schehr, Current fluctuations in noninteracting run-and-tumble particles in one dimension, Phys. Rev. E 101, 052101 (2020).
- [6] H. Lee, L. S. Levitov, and A. Y. Yakovets, Universal statistics of transport in disordered conductors, Phys. Rev. B 51, 4079 (1995).
- [7] E. McCulloch, J. De Nardis, S. Gopalakrishnan, and R. Vasseur, Full counting statistics of charge in chaotic many-body quantum systems, Phys. Rev. Lett. 131, 210402 (2023).
- [8] T. Bodineau and B. Derrida, Distribution of current in nonequilibrium diffusive systems and phase transitions, Phys. Rev. E 72, 066110 (2005).
- [9] T. Bodineau, B. Derrida, V. Lecomte, and F. van Wijland, Long range correlations and phase transitions in non-equilibrium diffusive systems, J. Stat. Phys. 133, 1013–1031 (2008).
- [10] G. Jona-Lasinio, From fluctuations in hydrodynamics to nonequilibrium thermodynamics, Prog. Theor. Phys. Supp. 184, 262–275 (2010).
- [11] H. Touchette, The large deviation approach to statistical mechanics, Phys. Rep. 478, 1–69 (2009).
- [12] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, Towards a nonequilibrium thermodynamics: A self-contained macroscopic description of driven diffusive systems, J. Stat. Phys. **135**, 857–872 (2009).
- [13] P. I. Hurtado, C. Pérez-Espigares, J. J. del Pozo, and P. L. Garrido, Symmetries in fluctuations far from equilibrium, PNAS 108, 7704–7709 (2011).
- [14] Y. Baek, Y. Kafri, and V. Lecomte, Dynamical phase transitions in the current distribution of driven diffusive channels, J. Phys. A: Math. Theor. 51, 105001 (2018).
- [15] G. M. Schütz, Exactly solvable models for many-body systems far from equilibrium, in *Phase Transitions and Critical Phenomena*, Vol. 19, edited by C. Domb and J. L. Lebowitz (Academic Press, 2001) p. 1–251.
- [16] K. Mallick, The exclusion process: A paradigm for nonequilibrium behaviour, Physica A 418, 17–48 (2015).
- [17] C. Giardinà, J. Kurchan, and L. Peliti, Direct evaluation of large-deviation functions, Phys. Rev. Lett. 96, 120603 (2006).
- [18] V. Lecomte and J. Tailleur, A numerical approach to large deviations in continuous time, J. Stat. Mech. 2007, P03004 (2007).
- [19] C. Giardinà, J. Kurchan, V. Lecomte, and J. Tailleur, Simulating rare events in dynamical processes, J. Stat.

Phys. 145, 787–811 (2011).

- [20] C. Pérez-Espigares and P. I. Hurtado, Sampling rare events across dynamical phase transitions, Chaos 29, 083106 (2019).
- [21] A. K. Hartmann, Big Practical Guide to Computer Simulations, 2nd ed. (World Scientific, 2015).
- [22] H. Spohn, Stretched exponential decay in a kinetic Ising model with dynamical constraint, Commun. Math. Phys. 125, 3–12 (1989).
- [23] P. L. Krapivsky, Symmetric exclusion process with a localized source, Phys. Rev. E 86, 041103 (2012).
- [24] T. Franco, P. Gonçalves, and A. Neumann, Phase transition in equilibrium fluctuations of symmetric slowed exclusion, Stoch. Process. Their Appl. **123**, 4156 (2013); Corrigendum to "Phase transition in equilibrium fluctuations of symmetric slowed exclusion" [Stochastic Process. Appl. **123**(12) (2013) 4156–4185], Stoch. Process. Their Appl. **126**, 3235 (2016).
- [25] H. Spohn, Large Scale Dynamics of Interacting Particles (Springer Berlin, Heidelberg, 1991).
- [26] C. Kipnis and C. Landim, Scaling Limits of Interacting Particle Systems (Springer Berlin, Heidelberg, 1999).
- [27] T. M. Liggett, Stochastic Interacting Systems: Contact, Voter and Exclusion Processes (Springer Berlin, Heidelberg, 1999).
- [28] T. Chou, K. Mallick, and R. K. P. Zia, Non-equilibrium statistical mechanics: from a paradigmatic model to biological transport, Rep. Prog. Phys. 74, 116601 (2011).
- [29] B. Derrida and A. Gerschenfeld, Current fluctuations in one dimensional diffusive systems with a step initial density profile, J. Stat. Phys. 137, 978–1000 (2009).
- [30] P. L. Krapivsky and B. Meerson, Fluctuations of current in nonstationary diffusive lattice gases, Phys. Rev. E 86, 031106 (2012).
- [31] T. Bodineau and B. Derrida, Current fluctuations in nonequilibrium diffusive systems: An additivity principle, Phys. Rev. Lett. **92**, 180601 (2004).
- [32] T. Bodineau and B. Derrida, Cumulants and large deviations of the current through non-equilibrium steady states, C. R. Physique 8, 540–555 (2007).
- [33] B. Derrida, O. Hirschberg, and T. Sadhu, Large deviations in the symmetric simple exclusion process with slow boundaries, J. Stat. Phys. 182, 15 (2021).
- [34] B. Derrida, B. Douçot, and P. E. Roche, Current fluctuations in the one-dimensional symmetric exclusion process with open boundaries, J. Stat. Phys. 115, 717–748 (2004).
- [35] M. Gorissen, A. Lazarescu, K. Mallick, and C. Vanderzande, Exact current statistics of the asymmetric simple exclusion process with open boundaries, Phys. Rev. Lett. **109**, 170601 (2012).
- [36] B. Derrida and A. Gerschenfeld, Current fluctuations of the one dimensional symmetric simple exclusion process with step initial condition, J. Stat. Phys. 136, 1–15 (2009).
- [37] K. Mallick, H. Moriya, and T. Sasamoto, Exact solution of the macroscopic fluctuation theory for the symmetric exclusion process, Phys. Rev. Lett. 129, 040601 (2022); Exact solutions to macroscopic fluctuation theory through classical integrable systems (2024), arXiv:2404.16434.
- [38] S. Saha and T. Sadhu, Large deviations in the symmetric simple exclusion process with slow boundaries: A hydrodynamic perspective (2023), arXiv:2310.11350v2.

- [39] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, Macroscopic fluctuation theory, Rev. Mod. Phys. 87, 593 (2015).
- [40] G. Jona-Lasinio, Review article: Large fluctuations in non-equilibrium physics, Nonlin. Processes Geophys. 30, 253–262 (2023).
- [41] L. Bertini, A. De Sole, D. Gabrielli, G. Jona-Lasinio, and C. Landim, Fluctuations in stationary nonequilibrium states of irreversible processes, Phys. Rev. Lett. 87, 040601 (2001); Macroscopic fluctuation theory for stationary non-equilibrium states, J. Stat. Phys. 107, 635-675 (2002).
- [42] J. Tailleur, J. Kurchan, and V. Lecomte, Mapping nonequilibrium onto equilibrium: The macroscopic fluctuations of simple transport models, Phys. Rev. Lett. 99, 150602 (2007); Mapping out-of-equilibrium into equilibrium in one-dimensional transport models, J. Phys. A: Math. Theor. 41, 505001 (2008).
- [43] E. Bettelheim, N. R. Smith, and B. Meerson, Inverse scattering method solves the problem of full statistics of nonstationary heat transfer in the Kipnis-Marchioro-Presutti model, Phys. Rev. Lett. **128**, 130602 (2022); Full statistics of nonstationary heat transfer in the Kipnis-Marchioro-Presutti model, J. Stat. Mech. **2022**, 093103 (2022).
- [44] T. Agranov, S. Ro, Y. Kafri, and V. Lecomte, Macroscopic fluctuation theory and current fluctuations in active lattice gases, SciPost Phys. 14, 045 (2023).
- [45] E. Bettelheim and B. Meerson, Complete integrability of the problem of full statistics of nonstationary mass transfer in the simple inclusion process (2024), arXiv:2403.19536v2.
- [46] T. M. Liggett, Ergodic theorems for the asymmetric simple exclusion process, Trans. Am. Math. Soc. 213, 237–261 (1975).
- [47] S. Grosskinsky, Phase transitions in nonequilibrium stochastic particle systems with local conservation laws, Ph.D. thesis, Technical University of Munich (2004).
- [48] S. Saha and T. Sadhu, Current fluctuations in a semiinfinite line, J. Stat. Mech. 2023, 073207 (2023).
- [49] L. Williams and T. Sasamoto, Combinatorics of the asymmetric exclusion process on a semi-infinite lattice (2012), arXiv:1204.1114.
- [50] C. A. Tracy and H. Widom, The asymmetric simple exclusion process with an open boundary, J. Math. Phys. 54, 103301 (2013).
- [51] H. G. Duhart, P. Mörters, and J. Zimmer, The semiinfinite asymmetric exclusion process: Large deviations via matrix products, Potential Anal. 48, 301–323 (2018).
- [52] See Supplementary Material for details.
- [53] V. Lecomte, A. Imparato, and F. van Wijland, Current fluctuations in systems with diffusive dynamics, in and out of equilibrium, Prog. Theor. Phys. Suppl. 184, 276–289 (2010).
- [54] K. Johansson, Shape fluctuations and random matrices, Comm. Math. Phys. 209, 437–476 (2000).
- [55] C. A. Tracy and H. Widom, A fredholm determinant representation in ASEP, J. Stat. Phys. 132, 291–300 (2008).
- [56] C. A. Tracy and H. Widom, Total current fluctuations in the asymmetric simple exclusion process, J. Math. Phys. 50, 095204 (2009).
- [57] This perplexing similarity was brought to our attention by Gunter M. Schütz.
- [58] P. C. Martin, E. D. Siggia, and H. A. Rose, Statistical dy-

namics of classical systems, Phys. Rev. A 8, 423 (1973).

- [59] H. K. Janssen, On a Lagrangean for classical field dynamics and renormalization group calculations of dynamical critical properties, Z. Physik B 23, 377–380 (1976).
- [60] C. De Dominicis, Dynamics as a substitute for replicas in systems with quenched random impurities, Phys. Rev. B 18, 4913 (1978).
- [61] C. De Dominicis and L. Peliti, Field-theory renormalization and critical dynamics above T_c : Helium, antiferromagnets, and liquid-gas systems, Phys. Rev. B 18, 353 (1978).
- [62] M. J. Ablowitz and H. Segur, The inverse scattering transform: Semi-infinite interval, J. Math. Phys. 16, 1054–1056 (1975).
- [63] R. Baldasso, O. Menezes, A. Neumann, and R. R. Souza, Exclusion process with slow boundary, J. Stat. Phys. 167, 1112–1142 (2017).
- [64] T. Franco, P. Gonçalves, and A. Neumann, Nonequilibrium and stationary fluctuations of a slowed boundary symmetric exclusion, Stoch. Process. Their Appl. 129, 1413–1442 (2019).
- [65] P. Gonçalves, M. Jara, O. Menezes, and A. Neumann, Non-equilibrium and stationary fluctuations for the SSEP with slow boundary, Stoch. Process. Their Appl. 130, 4326–4357 (2020).
- [66] C. Erignoux, P. Gonçalves, and G. Nahum, Hydrodynamics for SSEP with non-reversible slow boundary dynamics: Part I, the critical regime and beyond, J. Stat. Phys. 181, 1433–1469 (2020).
- [67] C. Erignoux, P. Gonçalves, and G. Nahum, Hydrodynamics for SSEP with non-reversible slow boundary dynamics: Part II, below the critical regime, ALEA, Lat. Am. J. Probab. Math. Stat. 17, 791–823 (2020).
- [68] T. Franco, P. Gonçalves, and A. Neumann, Hydrodynamical behavior of symmetric exclusion with slow bonds, Ann. Inst. H. Poincaré Probab. Statist. 49, 402–427 (2013).
- [69] D. Erhard, T. Franco, P. Gonçalves, A. Neumann, and M. Tavares, Non-equilibrium fluctuations for the SSEP with a slow bond, Ann. Inst. H. Poincaré Probab. Statist. 56, 1099–1128 (2020).
- [70] C. Kipnis, C. Marchioro, and E. Presutti, Heat flow in an exactly solvable model, J. Stat. Phys. 27, 65–74 (1982).
- [71] G. Schütz and S. Sandow, Non-Abelian symmetries of stochastic processes: Derivation of correlation functions for random-vertex models and disordered-interactingparticle systems, Phys. Rev. E 49, 2726 (1994).
- [72] C. Giardinà, J. Kurchan, and F. Redig, Duality and exact correlations for a model of heat conduction, J. Math. Phys. 48, 033301 (2007).
- [73] P. L. Krapivsky and D. Stefanovic, Lattice gases with a point source, J. Stat. Mech. 2014, P09003 (2014).
- [74] P. L. Krapivsky, K. Mallick, and D. Sels, Free fermions with a localized source, J. Stat. Mech. 2019, 113108 (2019).
- [75] P. L. Krapivsky, K. Mallick, and D. Sels, Free bosons with a localized source, J. Stat. Mech. 2020, 063101 (2020).
- [76] J. L. Skinner, Kinetic Ising model for polymer dynamics: Applications to dielectric relaxation and dynamic depolarized light scattering, J. Chem. Phys. **79**, 1955–1964 (1983).
- [77] V. Gupta, S. K. Nandi, and M. Barma, Size-stretched exponential relaxation in a model with arrested states, Phys. Rev. E 102, 022103 (2020).

- [78] S. Mukherjee, P. Pareek, M. Barma, and S. K. Nandi, Stretched exponential to power-law: crossover of relaxation in a kinetically constrained model, J. Stat. Mech. 2024, 023205 (2024).
- [79] This work was initiated from a seminar by Mustansir Barma, who encouraged us to solve the problem.
- [80] E. Akkermans, T. Bodineau, B. Derrida, and O. Shpielberg, Universal current fluctuations in the symmetric exclusion process and other diffusive systems, EPL 103, 20001 (2013).
- [81] D. Bernard and T. Jin, Open quantum symmetric simple exclusion process, Phys. Rev. Lett. 123, 080601 (2019).
- [82] D. Bernard, Can the macroscopic fluctuation theory be quantized?, J. Phys. A: Math. Theor. 54, 433001 (2021).
- [83] B. Doyon, Lecture notes on generalised hydrodynamics, SciPost Phys. Lect. Notes , 18 (2020).
- [84] A. Bastianello, B. Bertini, B. Doyon, and R. Vasseur, Introduction to the special issue on emergent hydrodynamics in integrable many-body systems, J. Stat. Mech. 2022, 014001 (2022).

Supplementary Material: Large deviations of current for the symmetric simple exclusion process on a semi-infinite line and on an infinite line with a slow bond

Kapil Sharma,^{1, *} Soumyabrata Saha,^{1, †} Sandeep Jangid,^{1, ‡} and Tridib Sadhu^{1, §}

¹Department of Theoretical Physics, Tata Institute of Fundamental Research, Homi Bhabha Road, Mumbai 400005, India

(Dated: May 14, 2024)

In this Supplement, we present additional details complementing the *Letter*. We discuss details about our numerical simulation providing results on the finite-size effects. Subsequently, we explicitly write the MFT-Action for the current generating function of the SSEP on the semi-infinite line with slow coupling to the reservoir. Additional details about a symmetry of the least-Action leading to the simplified parameter dependence of the scgf are discussed. Certain steps of our calculations quoted in the *Letter* are presented in the later parts of the Supplement.

CONTENTS

Numerical details: The cloning algorithm	S-1
An Action formulation for the semi-infinite line SSEP coupled to a reservoir	S-1
Proof of eq. (22) using eq. (21) of the <i>Letter</i>	S-2
Proof of eq. (24) of the <i>Letter</i> using a rotational symmetry of the Action	S-3
Semi-infinite line SSEP slowly coupled to a reservoir	S-4
Infinite line SSEP with a slow bond Recovering the $fast$ infinite line results	S-5 S-5
Stretched exponential decay in spin-auto-correlation	S-5
References	S-6

NUMERICAL DETAILS: THE CLONING ALGORITHM

Our numerical results in Fig. 2 of the Letter are based on a continuous time cloning algorithm. The cloning algorithm for discrete-time Markov processes was introduced in [1], and subsequently extended for continuous-time processes in [2]. We suitably adapted the algorithm in [2, 3] for obtaining the scgf of current in the semi-infinite line SSEP. Our simulation was done on a finite lattice of L sites, initially populated with average density $\rho_b = 1/2$, with a reflecting boundary at site L, and a reservoir of density $\rho_a = 1/2$ connected at site 1 (See Fig. 1 in the Letter). The net current Q_T is measured at the left boundary over a period $T \ll L^2$ such that the effect of the reflecting boundary is negligible. The finite size effects due to the boundary, the measurement time, and the clone size are shown in Fig. S-1. Further improvements of numerical result require advanced computational resources, which are not currently available to us.

AN ACTION FORMULATION FOR THE SEMI-INFINITE LINE SSEP COUPLED TO A RESERVOIR

Here, we present the explicit expression of the MFT-Action for the semi-infinite line SSEP in presence of slow coupling $\gamma = \Gamma/\sqrt{T}$ with the reservoir (see model definition in Fig. 1 in *Letter*). The generating function of the time-integrated current Q_T across the boundary reservoir is given as a path integral [4]

$$\left\langle \mathrm{e}^{\lambda Q_T} \right\rangle = \int \mathcal{D}\left[\rho, \widehat{\rho}\right] \,\mathrm{e}^{-\sqrt{T}\,S_\lambda\left[\rho, \widehat{\rho}\right]}$$
(S-1)



FIG. S-1. The solid red line represents the theoretical value of the scgf $\mu_{\rm si}(\lambda, \rho_a, \rho_b)$ (eq. (2b) in the *Letter*) for the semi-infinite line SSEP with $\lambda = 1.0$ and $\rho_a = \rho_b = 1/2$. Results in (a) are for clone size $N_c = 5 \times 10^4$ and time T = 300, showing dependence on system length *L*. Results in (b) are for T = 250 and L = 200, showing dependence on the clone size N_c . Results in (c) are for L = 200 and $N_c = 10^5$, showing convergence to theoretical result at large *T*.

where the Action is

$$S_{\lambda}\left[\rho,\widehat{\rho}\right] = -\lambda \int_{0}^{\infty} \mathrm{d}x \left[\rho(x,1) - \rho(x,0)\right] + \mathcal{F}[\rho(x,0)] + \int_{0}^{1} \mathrm{d}t \left[\int_{0}^{\infty} \mathrm{d}x \left(\widehat{\rho}(x,t) \partial_{t}\rho(x,t)\right) - H\left[\rho,\widehat{\rho}\right]\right].$$
(S-2)

Here, the first term is due to Q_T in eq. (1) in the *Letter*, and

$$\mathcal{F}[\rho] = \int_0^\infty \mathrm{d}x \int_{\rho_b}^\rho \mathrm{d}r \, \frac{2\left(\rho - r\right)}{\sigma(r)} \tag{S-3}$$

with $\sigma(\rho) = 2 \rho (1 - \rho)$, is the contribution [5] from the probability of $\rho(x, 0)$ in the initial state of the semi-infinite line SSEP populated with Bernoulli measure of average density ρ_b . The effective Hamiltonian

$$H\left[\rho,\widehat{\rho}\right] = H_{\text{bdry}}\left[\rho(0,t),\widehat{\rho}(0,t)\right] + \int_{0}^{\infty} \mathrm{d}x \left[\frac{\sigma(\rho)}{2}\partial_{x}\widehat{\rho}(x,t) - \partial_{x}\rho(x,t)\right] \partial_{x}\widehat{\rho}(x,t) \tag{S-4}$$

where

$$H_{\text{bdry}}\left[\rho,\widehat{\rho}\right] = \Gamma\left[\left(e^{\widehat{\rho}} - 1\right)\rho_a\left(1 - \rho\right) + \left(e^{-\widehat{\rho}} - 1\right)\rho\left(1 - \rho_a\right)\right]$$
(S-5)

is the contribution [4] from fluctuations at the left boundary of the lattice. In (S-2) there is no condition imposed for the hydrodynamic fields at the left boundary. The boundary condition eq. (15) in the *Letter* emerges in the fast-coupling $\Gamma \to \infty$ limit, where any small boundary fluctuations bear excessive cost, leading to $\hat{\rho}(0,t) = 0$ for which $H_{\text{bdry}} = 0$. This is the MFT-formulation in eq. (11) of the *Letter* with λ set to 0.

PROOF OF EQ. (22) USING EQ. (21) OF THE LETTER

In the fast coupling limit $\Gamma \to \infty$, minimization of the Action (S-2) corresponds to the optimal profiles $(\rho, \hat{\rho}) \equiv (q_{\rm si}, p_{\rm si})$ having the Euler-Lagrange equations and boundary conditions mentioned in eq. (13) and eq. (15) of the *Letter*. Substituting $\partial_t q(x,t)$ from eq. (13a) of the *Letter* in the expression of $S_{\lambda}[q_{\rm si}, p_{\rm si}]$ from (S-2), and subsequently using the spatial boundary conditions of $q_{\rm si}$ and $p_{\rm si}$ after performing an integration by parts, the least-Action for the semi-infinite line SSEP with fugacity λ and $(\rho_a, \rho_b) \equiv (1/2, 0)$ becomes [4]

$$\mu_{\rm si}\left(\lambda, \frac{1}{2}, 0\right) = \lambda \int_0^\infty dx \left[q_{\rm si}(x, 1) - q_{\rm si}(x, 0)\right] - \int_0^1 dt \int_0^\infty dx \left[q_{\rm si} \left(1 - q_{\rm si}\right) \left(\partial_x p_{\rm si}\right)^2\right],\tag{S-6}$$

while the corresponding least-Action for the infinite line SSEP with fugacity 2λ and $(\rho_a, \rho_b) \equiv (1,0)$ in terms of the least-Action path is [5]

$$\mu_{\inf}(2\lambda, 1, 0) = 2\lambda \int_0^\infty dx \left[q_{\inf}(x, 1) - q_{\inf}(x, 0) \right] - \int_0^1 dt \int_{-\infty}^\infty dx \left[q_{\inf} \left(1 - q_{\inf} \right) \left(\partial_x p_{\inf} \right)^2 \right].$$
(S-7)

The primary differences between the two expressions (S-6, S-7) are in the fugacity and the domain of x-integration.

For the density pair $(\rho_a, \rho_b) \equiv (1, 0)$ in the infinite lattice problem, using the parity symmetry

$$q_{\inf}(x,t) = 1 - q_{\inf}(-x,t)$$
 and $p_{\inf}(x,t) = 2\lambda - p_{\inf}(-x,t)$ (S-8)

of the least-Action path, the generating function in (S-7) reduces to

$$\mu_{\inf}(2\lambda, 1, 0) = 2 \left\{ \lambda \int_0^\infty dx \left[q_{\inf}(x, 1) - q_{\inf}(x, 0) \right] - \int_0^1 dt \int_0^\infty dx \left[q_{\inf} \left(1 - q_{\inf} \right) \left(\partial_x p_{\inf} \right)^2 \right] \right\}$$
(S-9)

The expression in (S-9) is similar to (S-6), and subsequently, using the relations

$$q_{\rm si}(x,t) = q_{\rm inf}(x,t) \quad \text{and} \quad p_{\rm si}(x,t) = p_{\rm inf}(x,t) - \lambda \tag{S-10}$$

from eq. (21) of the *Letter*, it becomes apparent that the two least-Actions (S-6, S-9) are related by

$$\mu_{\rm si}\left(\lambda, \frac{1}{2}, 0\right) = \frac{1}{2}\,\mu_{\rm inf}(2\lambda, 1, 0),\tag{S-11}$$

which is the eq. (22) of the *Letter*.

PROOF OF EQ. (24) OF THE LETTER USING A ROTATIONAL SYMMETRY OF THE ACTION

The discussion in this Section follows from [5, 6]. The least-Action in eq. (12) of the *Letter* for the annealed case (fluctuating initial state) has an underlying rotational symmetry that follows from a direct correspondence to the Heisenberg spin chain [5, 7]. As a consequence of the symmetry, the least-Action corresponding to parameters $(\lambda, \rho_a, \rho_b)$ and $(\lambda', \rho'_a, \rho'_b)$ are the same as long as the parameters are related by $\omega(\lambda, \rho_a, \rho_b) = \omega(\lambda', \rho'_a, \rho'_b)$, with the function ω defined in eq. (2c) of the *Letter*. To see this explicitly, consider the mapping to $\rho'_a = \rho'_b = 1/2$ with the corresponding λ' related by

$$\omega(\lambda, \rho_a, \rho_b) = \omega\left(\lambda', \frac{1}{2}, \frac{1}{2}\right) = \left(\sinh\frac{\lambda'}{2}\right)^2.$$
(S-12)

A re-parametrization of ρ_a and ρ_b in terms of u and v, following

$$\rho_a = \frac{e^v \cosh u - 1}{e^\lambda - 1} \quad \text{and} \quad \rho_b = \frac{e^{-v} \cosh u - 1}{e^{-\lambda} - 1} \tag{S-13}$$

gives a simple solution $\lambda' = 2u$ of (S-12).

In terms of these re-parametrized variables, the least-Action path $(q_{\rm si}, p_{\rm si})$ in eq. (12) of the *Letter* corresponding to parameters $(\lambda, \rho_a, \rho_b)$ is related to the least-Action path $(q'_{\rm si}, p'_{\rm si})$ of the same problem, but corresponding to $(\lambda', 1/2, 1/2)$ by the transformation [5]

$$q_{\rm si} = \frac{1}{\sinh u \sinh \frac{\lambda}{2}} \left[e^{p'_{\rm si} - u} \sinh \frac{\lambda + u - v}{2} - \sinh \frac{\lambda - u - v}{2} \right] \left[q'_{\rm si} e^{u - p'_{\rm si}} \sinh \frac{u + v}{2} - (1 - q'_{\rm si}) \sinh \frac{u - v}{2} \right], \quad (S-14a)$$

$$p_{\rm si} = \log\left[1 + \frac{e^u \left(e^{\lambda} - 1\right) \left(e^{p'_{\rm si}} - 1\right)}{e^{p'_{\rm si}} \left(e^u - e^v\right) + e^u \left(e^{u+v} - 1\right)}\right].$$
(S-14b)

One way to see this is by noting that the transformed fields (S-14a, S-14b) keep the bulk Hamiltonian H[q, p] (S-4) invariant, and thus satisfy the same Euler-Lagrange equations. This has already been shown in [5] for the infinite-line problem and the analysis for the semi-infinite-line problem is similar. The important differences between the two problems are in the boundary conditions. For the semi-infinite-line problem, we explicitly verify that the transformation preserves the corresponding structure of the temporal boundary conditions (eq. (14) of the *Letter*).

$$p'_{\rm si}(x,0) = \lambda' + \int_{1/2}^{q'_{\rm si}(x,0)} \frac{\mathrm{d}r}{r(1-r)} \implies p_{\rm si}(x,0) = \lambda + \int_{\rho_b}^{q_{\rm si}(x,0)} \frac{\mathrm{d}r}{r(1-r)} , \quad \text{and} \tag{S-15a}$$

$$p'_{\rm si}(x,1) = \lambda' \implies p_{\rm si}(x,1) = \lambda$$
 (S-15b)

as well as the spatial boundary conditions

$$q'_{\rm si}(0,t) = \frac{1}{2}$$
, $p'_{\rm si}(0,t) = 0 \implies q_{\rm si}(0,t) = \rho_a$, $p_{\rm si}(0,t) = 0$, and (S-15c)

$$q'_{\rm si}(\infty,t) = \frac{1}{2} , \ p'_{\rm si}(\infty,t) = \lambda' \implies q_{\rm si}(\infty,t) = \rho_b , \ p_{\rm si}(\infty,t) = \lambda.$$
(S-15d)

The invariance of the Euler-Lagrange equations along with (S-15) culminate into

$$\mu(\lambda, \rho_a, \rho_b) = \mu\left(\lambda', \frac{1}{2}, \frac{1}{2}\right).$$
(S-16)

Using (S-12) to express λ' in terms of $\omega(\lambda, \rho_a, \rho_b)$ in (S-16), we arrive at

$$\mu(\lambda, \rho_a, \rho_b) = \mu\left(2\operatorname{arcsinh}\sqrt{\omega(\lambda, \rho_a, \rho_b)}, \frac{1}{2}, \frac{1}{2}\right),\tag{S-17}$$

thus proving the ω dependence of the scaf for the semi-infinite line SSEP.

SEMI-INFINITE LINE SSEP SLOWLY COUPLED TO A RESERVOIR

For slow coupling $\gamma = \Gamma/\sqrt{T}$ with the reservoir, contributions from the fluctuations at the boundary are relevant, as evidenced in the Hamiltonian (S-4). Effect of these boundary fluctuations can be incorporated in the current fluctuations using the additivity principle [8], whose basic premise is to treat the entire system as composed of two subsystems: the bond linking the reservoir with the semi-infinite lattice, and the lattice itself. The main idea is then to assume that for current fluctuations at large times, the two subsystems are independent of each other, except their dependence through the density at their common point, which is adjusted to maximize the probability of current

$$P(Q_T) \simeq \max_{\rho_0} P_{\text{bond}}(Q_T, \rho_a, \rho_0) P_{\text{si}}(Q_T, \rho_0, \rho_b)$$
(S-18)

for large T. In this description, the single slow bond is coupled at the two ends with reservoirs ρ_a and ρ_0 , and the semi-infinite lattice is coupled to a reservoir of density ρ_0 at its left end. For long times T, $P_{\text{bond}}(Q_T = j\sqrt{T}) \sim e^{-\sqrt{T}\phi_{\text{bond}}(j,\rho_a,\rho_0)}$, where $\phi_{\text{bond}}(j,\rho_a,\rho_0)$ is the Legendre-Fenchel transformation of $\Gamma \omega(\lambda,\rho_a,\rho_0)$ [8] with the function ω defined in eq. (2c) of the *Letter*. Using similar asymptotics (eq. (7) of the *Letter*) for the semi-infinite lattice, (S-18) gives the large deviation asymptotics $P(Q_T = j\sqrt{T}) \sim e^{-\sqrt{T}\phi_{\text{si}}^{\text{slow}}(j,\rho_a,\rho_b)}$ with the ldf

$$\phi_{\rm si}^{\rm slow}(j,\rho_a,\rho_b) = \min_{\rho_0} \left[\phi_{\rm bond}(j,\rho_a,\rho_0) + \phi_{\rm si}(j,\rho_a,\rho_b) \right]. \tag{S-19}$$

For the corresponding Legendre-Fenchel transformations, (S-19) gives [8]

$$\mu_{\rm si}^{\rm slow}(\lambda,\rho_a,\rho_b) = \max_{\rho(0)} \min_{\lambda_0} \left[\Gamma \,\omega(\lambda_0,\rho_a,\rho(0)) + R_{\rm si} \big(\omega(\lambda-\lambda_0,\rho(0),\rho_b) \big) \right],\tag{S-20}$$

where $R_{\rm si}$ is defined in eq. (2b) of the *Letter*.

The expression (S-20) further simplifies using an identity (see eq. (29) of [8]), which leads to

$$\mu_{\rm si}^{\rm slow}(\lambda,\rho_a,\rho_b) = \min_{z} \left[\Gamma \sinh^2\left(z-u\right) + R_{\rm si}\left(\sinh^2 z\right) \right],\tag{S-21}$$

with $\sinh^2 u = \omega(\lambda, \rho_a, \rho_b)$ as reported in eq. (25) of the *Letter*.

The variational problem in (S-21) could be explicitly solved using the expression for $R_{\rm si}$ in eq. (2b) of the *Letter*, leading to a parametric expression

$$\mu_{\rm si}^{\rm slow}(\lambda,\rho_a,\rho_b) = \Gamma \sinh^2\left(z-u\right) + \int_{-\infty}^{\infty} \frac{\mathrm{d}k}{2\pi} \log\left[1+\sinh^2 2z \,\mathrm{e}^{-k^2}\right] \tag{S-22a}$$

where z is given by the solution of the equation

$$\sinh 2(u-z) = \frac{\sinh 4z}{\Gamma \pi} \int_{-\infty}^{\infty} \frac{\mathrm{d}k}{\sinh^2 2z + \mathrm{e}^{k^2}}$$
(S-22b)

with $\sinh^2 u = \omega(\lambda, \rho_a, \rho_b).$

INFINITE LINE SSEP WITH A SLOW BOND

Following a similar construction as discussed in the previous section, the scgf of current for an infinite line SSEP with a single slow bond can be obtained by treating the entire system as composed of three subsystems: two semiinfinite systems with unit hopping rates coupled to each other by a single slow bond with hopping rate Γ/\sqrt{T} . This leads to a formula analogous to (S-20) for the scgf

$$\mu_{\inf}^{\text{slow}}(\lambda,\rho_a,\rho_b) = \max_{\rho_0,\rho_1} \min_{\lambda_0,\lambda_1} \Big[R_{\text{si}}\big(\omega(\lambda_0,\rho_a,\rho_0)\big) + \Gamma\,\omega(\lambda_1-\lambda_0,\rho_0,\rho_1) + R_{\text{si}}\big(\omega(\lambda-\lambda_1,\rho_1,\rho_b)\big) \Big],\tag{S-23}$$

where $\rho_{0(1)}$ is the density of the common site between the left (right) semi-infinite lattice and the slow bond. The analysis is very similar to the discussion in [8] for a related problem.

The expression (S-23) further simplifies using eq. (29) of [8] leading to

$$\mu_{\inf}^{\text{slow}}(\lambda,\rho_a,\rho_b) = \min_{z_a,z_b} \left[R_{\text{si}}(\sinh^2 z_a) + \Gamma \sinh^2 (z_a + z_b - u) + R_{\text{si}}(\sinh^2 z_b) \right]$$
(S-24)

with $\sinh^2 u = \omega(\lambda, \rho_a, \rho_b)$. This result is quoted in eq. (26) of the *Letter*.

Recovering the *fast* infinite line results

A self-consistency check of (S-24) is to recover the well-known result [9, 10] for the infinite line SSEP in the fast coupling limit $\Gamma \to \infty$. In this limit, the second term in (S-24) dominates and the minimization over z_a and z_b leads to the solutions $z_a = z_b = u/2$. Putting this back in the expression for the scgf and using the explicit expression for the scgf of a semi-infinite line SSEP (eq. (2b) in the *Letter*), we obtain

$$\mu_{\inf}(\lambda, \rho_a, \rho_b) = \int_{-\infty}^{\infty} \frac{\mathrm{d}k}{\pi} \log\left[1 + \omega(\lambda, \rho_a, \rho_b) \,\mathrm{e}^{-k^2}\right] \tag{S-25}$$

where we have used $\sinh^2 u = \omega(\lambda, \rho_a, \rho_b)$. This is the scgf for the infinite line SSEP.

STRETCHED EXPONENTIAL DECAY IN SPIN-AUTO-CORRELATION

For the energy-conserving spin-flip dynamics on a one-dimensional lattice, defined in [11], each site i at a given time t is assigned with an Ising spin $S_i(t) = \pm 1$. The dynamics is such that a spin can flip only if its two nearest neighbor's spins have opposite signs, therefore conserving the total energy (Fig. S-2). Due to this constraint, the number of domain walls remains conserved. In fact, the domain wall dynamics is exactly same as the movement of particles in SSEP on an infinite lattice.

This correspondence with SSEP relates the spin-auto-correlation to the particle current in SSEP. To see this we note that whenever a domain wall crosses through a site i (see Fig. S-2), the associated Ising spin, S_i , changes its sign. If Q_t is the net rightward flow of domain walls through the site i in time t, then

$$S_i(0) S_i(t) = \begin{cases} +1 & \text{for even } Q_t, \\ 1 & \text{for even } Q_t \end{cases}$$
(S-26a)

This implies the auto-correlation

$$\langle S_i(0) S_i(t) \rangle = P_t(\text{even } Q_T) - P_t(\text{odd } Q_T), \tag{S-27}$$

with the probability $P_t(\text{even } Q_T)$ of Q_t being even and $P_t(\text{odd } Q_T)$ for odd.

The difference of the two probabilities in (S-27) is related to the generating function of current $\langle e^{\lambda Q_t} \rangle$ in SSEP. For $\lambda = i\pi$, noting that Q_t only takes integer values, we see that

$$\langle e^{i\pi Q_t} \rangle = \sum_{Q_t} P(Q_t) e^{i\pi Q_t} = P_t(\text{even } Q_T) - P_t(\text{odd } Q_T),$$
 (S-28)



FIG. S-2. Dynamics of spins following a kinetic constraint of fixed energy: a spin can flip with unit rate only if its nearest neighbor's spins are opposite in sign. For example, the fifth spin from left has left neighbor down and right neighbor up, so it is allowed to flip, making the domain wall move leftward by one lattice unit. The movement of domain walls is analogous to the dynamics of particles in a SSEP.

From (S-27) and (S-28), it is evident that at large times, the spin-auto-correlation follows the asymptotics of the current scgf of SSEP. Using the results for the latter from [9] we see that for a large time t,

$$\langle S_i(0) S_i(t) \rangle \sim e^{\sqrt{t}\mu_{\inf}(i\pi,\rho,\rho)} = e^{-\sqrt{t/\tau}}$$
(S-29)

with $\tau = [\mu(i\pi, \rho, \rho)]^{-2}$. Here, the density ρ is the average density of domain walls in the spin dynamics. The result (S-29) is consistent with the bounds derived in [11]. Note that the correspondence with the current scgf of SSEP holds in the presence of slow regions in the the domain wall dynamics, or when the leftmost spin is coupled with a reservoir, relating to our corresponding results of current scgf for SSEP.

- [†] soumyabrata.saha@tifr.res.in
- ‡ sandeep.jangid@tifr.res.in
- [§] tridib@theory.tifr.res.in
- C. Giardinà, J. Kurchan, and L. Peliti, Direct evaluation of large-deviation functions, *Phys. Rev. Lett.* 96, 120603 (2006).
 V. Lecomte and J. Tailleur, A numerical approach to large deviations in continuous time, *J. Stat. Mech.* 2007, P03004
- (2007).
 [3] C. Pérez-Espigares and P. I. Hurtado, Sampling rare events across dynamical phase transitions, *Chaos* 29, 083106 (2019).
- [4] S. Saha and T. Sadhu, Current fluctuations in a semi-infinite line, J. Stat. Mech. **2023**, 073207 (2023).
- [5] B. Derrida and A. Gerschenfeld, Current fluctuations in one dimensional diffusive systems with a step initial density profile, J. Stat. Phys. 137, 978–1000 (2009).
- [6] [6] V. Lecomte, A. Imparato, and F. van Wijland, Current fluctuations in systems with diffusive dynamics, in and out of equilibrium, Prog. Theor. Phys. Suppl. 184, 276–289 (2010).
- [7] J. Tailleur, J. Kurchan, and V. Lecomte, Mapping nonequilibrium onto equilibrium: The macroscopic fluctuations of simple transport models, *Phys. Rev. Lett.* **99**, 150602 (2007); Mapping out-of-equilibrium into equilibrium in one-dimensional transport models, *J. Phys. A: Math. Theor.* **41**, 505001 (2008).
- [8] B. Derrida, O. Hirschberg, and T. Sadhu, Large deviations in the symmetric simple exclusion process with slow boundaries, J. Stat. Phys. 182, 15 (2021).
- B. Derrida and A. Gerschenfeld, Current fluctuations of the one dimensional symmetric simple exclusion process with step initial condition, J. Stat. Phys. 136, 1–15 (2009).
- [10] K. Mallick, H. Moriya, and T. Sasamoto, Exact solution of the macroscopic fluctuation theory for the symmetric exclusion process, *Phys. Rev. Lett.* **129**, 040601 (2022); Exact solutions to macroscopic fluctuation theory through classical integrable systems (2024), arXiv:2404.16434.
- [11] H. Spohn, Stretched exponential decay in a kinetic Ising model with dynamical constraint, Commun. Math. Phys. 125, 3–12 (1989).

^{*} kapil.sharma@tifr.res.in