ROGERS-RAMANUJAN IDENTITIES IN STATISTICAL MECHANICS

GEOFFREY B CAMPBELL

ABSTRACT. We describe the story of the Rogers-Ramanujan identities; being known for 85 years and having about 130 pure mathematics proofs, suddenly entering physics when Rodney Baxter solved the Hard Hexagon Model in Statistical Mechanics in 1980. We next cover the accompanying proofs by George E Andrews of other related Baxter identities arisen of Rogers-Ramanujan type, leading into a new flourishing partnership of Physics and Mathematics. Our narrative goes into the subsequent 44 years, explaining the progress in physics and mathematical analysis. Finally we show some related crossovers with regard to the *Elliptic qgamma function* and some *Vector Partition* generating functional equations; the latter of which may be new. The present paper is essentially chapter 11 of the author's 32 chapter book [23] to appear in June 2024.

1. Baxter's Hard Hexagon Model solved exactly by Rogers-Ramanujan identities

The years 1980 and 1981 were amazing in the theory of partitions; with significance arisen from a remarkable discovery in theoretical physics. In 1980 Baxter [10] found real-world physical application of the Rogers-Ramanujan identities coming directly from his solving the Hard Hexagon Model in Statistical Mechanics. This opened up a new world of research in both theoretical physics and partition-theoretic mathematics that is still evolving, over forty years later. This discovery has given rise to thousands of mathematics and physics research papers and promises continued relevance into future decades.

Some of this excitement can be gleaned from the important paper by Andrews in 1981 (see [3]) from quoting his opening paragraph here.

Think About It...

"In 1980, Baxter found his beautiful solution to the hard-hexagon model of statistical mechanics. His treatment of this model is naturally divided into six regimes that depend on values taken by various parameters associated with the model. Then in truly astounding fashion it turns out that eight Rogers-Ramanujan type identities, all essentially known to Rogers ([33], [34]), are the fundamental keys for finding infinite product representations of the related statistical mechanics partition functions in **regimes I**, **III**, **and IV**."

- George E. Andrews, Proceedings of the National Academy of Sciences, 1981

²⁰¹⁰ Mathematics Subject Classification. Primary: 11P84; Secondary: 37K60, 82B23.

Key words and phrases. Partition identities, identities of Rogers-Ramanujan type; Lattice dynamics, integrable lattice equations; Exactly solvable models; Bethe ansatz.

1.1. A few words about the types of lattice models studied here. Baxter formulates his approach based on the *hard hexagon* diagram (see next page Figure 1) which has constraints on the particles such that they can only be in certain places at certain moments.

The hard hexagon model is a two-dimensional lattice model of a gas of hard (i.e. non-overlapping) molecules. In it, particles are placed on the sites of the triangular lattice so that no two particles are together or adjacent. A typical allowed arrangement of particles is shown in Figure 1. If we regard each particle as the centre of a hexagon covering the six adjacent faces (such hexagons are shown shaded in the figure), then the rule only allows hexagons that do not overlap: hence the name of the model. For a lattice of N sites, the grand-partition function is

(1.1)
$$Z(z) = \sum_{n=0}^{n/3} z^n g(n, N)$$

where g(n, N) is the allowed number of ways of placing *n* particles on the lattice, and the sum is over all possible values of *n*. (Since no more than $\frac{1}{3}$ of the sites can be occupied, *n* takes values from 0 to $\frac{N}{3}$.)



A typical arrangement of particles (black circles) on the triangular lattice, such that no two particles are together or adjacent. The six faces round each particle are shaded: they form non-overlapping (ie. "hard") hexagons.

FIGURE 1. The particle arrangement for the Hard Hexagon Model.

Clearly, from the mathematical 'non-physicist' perspective, that this schematic diagram could lead to enumeration of well defined types of integer partitions was (back in 1980 at least) seen as a major departure from the more typical Ferrers Graphs, Durfee Squares and Plane Partition cube stack representations adopted in the literature by Andrews and others over the contemporary mathematical landscapes. As a mathematician already versed in the ideas of our earlier chapters, it becomes natural to ask how say, Bijection equivalences, *q*-series partition generating functions, and such, may become the same world as the statistical mechanics models involving assigned values of *spin charges* along a vertex or edge of a lattice of a certain structure, shape and form. We put such questions aside for now, as we want to present Baxter's extraordinary findings here. However, these questions of equivalences of theories and analyses are worthy of deeper understanding it seems. In seeking to calculate the free energy in the hard hexagon model as depicted in figure 1, Baxter was led to consider a range of logical options or *Regimes*, which are here depicted as Baxter described in 1980.

The three sub-lattices of the triangular lattice: sub-lattice 1 consists of all sites of type 1, and similarly for sub-lattices 2 and 3. Adjacent sites lie on different sublattices. a close-packed arrangement of particles (black circles) is shown: all sites of one sub-lattice (in this case sub-lattice 1) are occupied, the rest are empty. (See figure 2)



FIGURE 2. Baxter's three sub-lattices of the triangular lattice.

1.2. What Baxter found in 1980. So, continuing our discussion of equation (1.1), we want to calculate Z, or rather the partition-function per site of the infinite lattice

(1.2)
$$\kappa = \lim_{N \to \infty} Z^{\frac{1}{N}}$$

as a function of the positive real variable z. This z is known as the 'activity'. This problem can be put into what is called 'spin'-type language by associating with each site i a variable σ_i . However, instead of the usual approach in statistical mechanics models of letting σ_i take values 1 and -1, we take the values 0 and 1: if the site (lattice point) is empty, then $\sigma_i = 0$; if it is full then $\sigma_i = 1$. Thus σ_i is the number of particles at site i: the 'occupation number'. Then (1.1) can be written as

(1.3)
$$Z = \sum_{\sigma} z^{(\sigma_1 + \sigma_2 + \dots + \sigma_N)} \prod_{i,j} (1 - \sigma_i \sigma_j),$$

where the product is over all edges (i, j) of the triangular lattice, and the sum is over all values (0 and 1) of all the occupation numbers $\sigma_1, \sigma_2, ..., \sigma_N$.

We expect this model to undergo a phase transition from an homogeneous fluid state at low activity z to an inhomogeneous solid state at high activity z. To see this, divide the lattice into three sub-lattices 1, 2, 3, so that no two sites of the same type

GEOFFREY B CAMPBELL

are adjacent. Then there are three possible close-packed configurations of particles on the lattice: either all sites of type 1 are occupied, or all sites of type 2, or all sites of type 3. Suppose we fix the boundary sites as in the first possibility, i.e. all boundary sites of type 1 are full, and all other boundary sites are empty. Then for an infinite lattice the second and third possibilities give a negligible contribution to the sum-over-states in (1.3). Clearly, sites on different sub-lattices are no longer equivalent. Let ρ_r be the local density at a site of type r, given by,

(1.4)
$$\rho_r = \langle \sigma_l \rangle = Z^{-1} \sum_{\sigma} \sigma_l z^{(\sigma_1 + \sigma_2 + \dots + \sigma_N)} \prod_{i,j} (1 - \sigma_i \sigma_j),$$

where l is a site of type r.

When z is infinite, the system is close-packed with all sites of type 1 occupied, so $\rho_1 = 1$, $\rho_2 = \rho_3 = 0$. We can expand each ρ_r in inverse powers of z by considering successive perturbations of the close-packed state. For a site l deep inside a large lattice, this gives

(1.5)
$$\rho_1 = 1 - z^{-1} - 5z^{-2} - 34z^{-3} - 267z^{-4} - 2037z^{-5} - \dots$$
$$\rho_2 = \rho_3 = z^{-2} + 9z^{-3} + 80z^{-4} + 965z^{-5} + \dots$$

The system is therefore not homogeneous, since ρ_1, ρ_2, ρ_3 , are not all equal. This contrasts with the low-activity situation: starting from the state with all sites empty and successively introducing particles, we obtain

(1.6)
$$\rho_1 = \rho_2 = \rho_3 = z^1 - 7z^2 - 58z^3 - 519z^4 + 4856z^5 - \dots$$

To all orders in this expansion it is true that $\rho_1 = \rho_2 = \rho_3$.

The system is therefore inhomogeneous for sufficiently large z, and homogeneous for sufficiently small z if the series converge. There must be a critical value z_c , of z above which the system ceases to be homogeneous. Since the homogeneous phase is typical of a fluid, and the ordered inhomogeneous phase is typical of a solid, the model can be said to undergo a fluid to solid transition at $z = z_c$.

1.3. Baxter's 1980 numerical preliminaries. Baxter then applied the Corner Transfer Matrix approach to the matrix eigenvalues a_1, \ldots, a_{10} , for the hard hexagon model with z = 1. The values were approximate, being calculated from finite truncations of the triangular lattice analogue of the matrix equation related to this. The eigenvalues occur in groups of comparable magnitude, and it is sensible to include all members of a group. For this reason the truncations used were 2×2 , 3×3 , 5×5 , 7×7 and 10×10 . Each a_i was given for successively larger truncations, and clearly each tended rapidly to a limit. This limit is its exact value for the infinite-dimensional corner transfer matrix. Refer to Baxter [12, Chapter 13] for how the Corner Transfer Matrix context applies.

Baxter was able to determine the required coefficients up to 30 terms of the power series *exact solution*. Equipped with knowledge from his famous eight-vertex model exact solution from a few years earlier, Baxter was able to put the hard hexagon proposed solution expansion into the form

(1.7)
$$z = -x \prod_{n=1}^{\infty} (1 - x^n)^{c_n}.$$



The six regimes in the (L, M) plane, as listed at right. Shaded areas are unphysical, since $z = (1 - e^{-L}) (1 - e^{-M})/(e^{L+M} - e^{L} - e^{M})$ and $\Delta = z^{-\frac{1}{2}}(1 - z e^{L+M})$ for $\Delta = \Delta_1$ gives z therein to be negative. Regimes I, III, V are disordered, II and VI have triangular ordering, IV has square ordering. The system is critical on the (I, II), (III, IV) and (V, VI) boundaries, where $|\Delta| = \Delta_c$ the values of u on the (L, M) axes are indicated.

FIGURE 3. Baxter's six Regimes for the Hard Hexagon Model. The six regimes in the (L, M) plane. Shaded areas are not physical. Regimes I, III and V are disordered, and regimes II and VI have triangular ordering, while Regime IV has square ordering.

With this in mind he discovered that

Baxter then was able to infer that

(1.9)
$$z = -x \left(\frac{H(x)}{G(x)}\right)^{5},$$

where

(1.10)
$$G(x) = \prod_{n=1}^{\infty} \frac{1}{(1-x^{5n-4})(1-x^{5n-1})},$$

(1.11)
$$H(x) = \prod_{n=1}^{\infty} \frac{1}{(1-x^{5n-3})(1-x^{5n-2})}.$$

The mathematical partition theorist will will recognise the products G(x) and H(x) as from the Rogers-Ramanujan identities.

So, Baxter followed through this Corner Transfer Matrix approach to calculate the coefficients and determine a range of identities for each of the Hard Hexagon Regime types of allowable spin configurations. What he derived, is what Andrews subsequently was able to prove on his visit to Baxter in Australia the following year. So, we have set the scene for Andrews proofs in 1981.

1.4. What Andrews found in 1981. So, after Baxter's solution to the Hard Hexagon Model in 1980, the opening to a new world of research in both theoretical

physics and partition-theoretic mathematics was about to happen. George E. Andrews visited Baxter in 1981, and proved the identities arisen from the Regimes of Baxter's Hard Hexagon exact solutions.

In a landmark paper, Andrews [3] goes on then to say that Baxter found the following identities occurring for the six regimes, conspicuously missing (but later returning to) the more complicated Regime II, as follows:

Regime I

(1.12)
$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q)_n} = \frac{1}{\Pi(q, q^4; q^5)};$$

(1.13)
$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q)_n} = \frac{1}{\Pi(q^2, q^3; q^5)};$$

Regime III

(1.14)
$$\sum_{n=0}^{\infty} \frac{q^{\frac{n(3n-1)}{2}}}{(q)_n (q;q^2)_n} = \frac{\Pi(q^4, q^6, q^{10}; q^{10})}{(q)_{\infty}};$$

(1.15)
$$\sum_{n=0}^{\infty} \frac{q^{\frac{3n(n+1)}{2}}}{(q)_n(q;q^2)_{n+1}} = \frac{\Pi(q^2,q^8,q^{10};q^{10})}{(q)_{\infty}};$$

Regime IV

(1.16)
$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q)_{2n+1}} = \frac{\Pi(q^3, q^7, q^{10}; q^{10})\Pi(q^4, q^{16}; q^{20})}{(q)_{\infty}};$$

(1.17)
$$\sum_{n=0}^{\infty} \frac{q^{n(n+1)}}{(q)_{2n}} = \frac{\Pi(q, q^9, q^{10}; q^{10}) \Pi(q^8, q^{12}; q^{20})}{(q)_{\infty}};$$

(1.18)
$$\sum_{n=0}^{\infty} \frac{q^{n^2}}{(q)_{2n}} = \frac{1}{\Pi(q^4, q^{16}; q^{20})(q; q^2)_{\infty}};$$

(1.19)
$$\sum_{n=1}^{\infty} \frac{q^{n^2}}{(q)_{2n-1}} = \frac{q}{\Pi(q^8, q^{12}; q^{20})(q; q^2)_{\infty}}.$$

These use the standard notation of Slater [38],

(1.20)
$$(a)_n = (a;q)_n = \prod_{j=0}^{n-1} (1 - aq^j);$$

(1.21)
$$(a)_{\infty} = (a;q)_{\infty} = \prod_{j=0}^{\infty} (1 - aq^j);$$

 $\mathbf{6}$

(1.22)
$$\Pi(a_1, a_2, ..., a_r; q) = (a_1)_{\infty} (a_2)_{\infty} ... (a_r)_{\infty}.$$

These results either are given explicitly by Rogers ([33], [34]) or are immediate consequences of his work: (1.12) is equation 1 on page 328 of [33]; (1.13) is equation 2 on page 329 of [33]. (1.12) and (1.13) are of course the now famous Rogers-Ramanujan identities.

(1.14) is equation 2, line 3, on page 330 of [34]; (1.15) is equation 2, line 2, on page 330 of [34]; (1.16) is implicit in the identity of equations 2 and 3 on page 330 of [33] when $\lambda = 1$ and q is replaced by -q (explicitly given by Slater [38], equation 94); (1.17) is equation 13 on page 332 of [33]; (1.18) is the second equation of page 331 of [33]; (1.19) is equation 3, line 2, on page 330 of [34].

For **Regime II**, however, it turns out that one must consider the following rather complicated one-dimensional partition function:

(1.23)
$$F_k(\sigma_1) = \lim_{m \to \infty} \sum_{\sigma_2, \sigma_3, \dots, \sigma_m} q^{i(\sigma_i \sigma_{i+2} - \sigma_{i+1} + \overline{\sigma}_{i+1})}.$$

in which the summation runs over all possible (m-1)-tuples $(\sigma_2, \sigma_3, ..., \sigma_m)$ subject to the fairly complicated and stringent conditions:

$$\begin{cases} \sigma_{i} = 0 \text{ or } 1, & 1 \leq j \leq m; \\ \sigma_{i} + \sigma_{i+1} \leq 1, & 1 \leq j \leq m-1; \\ \sigma_{m} = \sigma_{m+1} = 0, & 1; \\ \overline{\sigma}_{i} = 1, & i \equiv k \pmod{3}; \\ \overline{\sigma}_{i} = 0, & i \neq k \pmod{3}. \end{cases}$$

Baxter obtains recurrence relations for refinements of these functions $F_k(\sigma_1)$; however, the techniques that he applies successfully to solve the recurrence relations in the other three regimes fail here. For this reason he is unable to find counterparts of the infinite series in (1.12) to (1.19). By direct expansion he obtains overwhelming evidence to conjecture that each of $F_1(0), F_1(1), F_2(0), F_2(1), F_3(0)$, and $F_3(1)$ are identical with elegant combinations of infinite products.

Next we shall give double series expansions for the $F_k(\sigma_1)$ that establish all six of Baxter's conjectures. Apart from their contribution to Baxter's solution of the hard-hexagon model, these results are also surprising mathematically. They are not apparently limiting cases of known basic hypergeometric series identities; this is in contradistinction to the fact that the place of equations (1.12) to (1.19) in the hierarchy of basic hypergeometric series was back then well known (see Slater [38], Bailey [7] and [8]). Here later, we shall describe the results and techniques required to establish these theorems.

2. HARD HEXAGON MODEL REGIME II IDENTITIES FROM BAXTER

So, the 1980 conjectures by Baxter for Regime II were rigorously proven by Andrews in 1981 (see [3]), although, as Andrews says, the results could have been known to, or derived by, Rogers [34] or Schur [36]) in 1917, or by Slater [38] originally in 1952. However, these results were not explicitly given back then, and Baxter's derivation was after all, innovative and a significant step forward in the theory. So we state the relevant identities in the following six equations.

Regime II

$$F_1(0) = \sum_{n=0}^{\infty} \sum_{0 \le r \le \frac{3n+1}{2}} \frac{q^{\frac{3n(n+1)}{2}-r}}{(q^2;q^2)_r(q)_{3n-2r+1}} = \frac{\Pi(q^4,q^{11},q^{15};q^{15}) + q\Pi(q,q^{14},q^{15};q^{15})}{(q)_{\infty}};$$

(2.2)

$$F_1(1) = \sum_{n=0}^{\infty} \sum_{0 \le r \le \frac{3n}{2}} \frac{q^{\frac{3n(n+1)}{2}-r}}{(q^2;q^2)_r(q)_{3n-2r}} = \frac{\Pi(q^7,q^8,q^{15};q^{15}) + q\Pi(q^2,q^{13},q^{15};q^{15})}{(q)_{\infty}};$$

(2.3)
$$F_2(0) = \sum_{n=1}^{\infty} \sum_{0 \le r \le \frac{3n-1}{2}} \frac{q^{\frac{n(3n-1)}{2}-r}}{(q^2;q^2)_r(q)_{3n-2r-1}} = \frac{\Pi(q^6,q^9,q^{15};q^{15})}{(q)_{\infty}}$$

$$= 1 + q + 2q^{2} + 3q^{3} + 5q^{4} + 7q^{5} + 10q^{6} + 14q^{7} + 20q^{8} + 26q^{9} + 36q^{10} + 47q^{11} + 63q^{12} + 81q^{13} + 106q^{14} + 135q^{15} + 174q^{16} + 219q^{17} + 278q^{18} + 347q^{19} + 436q^{20} + O(q^{21});$$

(2.4)
$$F_2(1) = \sum_{n=0}^{\infty} \sum_{0 \le r \le \frac{3n+1}{2}} \frac{q^{\frac{n(3n+5)}{2}+1-r}}{(q^2;q^2)_r(q)_{3n-2r+1}} = \frac{q\Pi(q^3,q^{12},q^{15};q^{15})}{(q)_{\infty}}$$

$$= q + q^{2} + 2q^{3} + 2q^{4} + 4q^{5} + 5q^{6} + 8q^{7} + 10q^{8} + 15q^{9} + 19q^{10} + 27q^{11} + 34q^{12} + 46q^{13} + 58q^{14} + 77q^{15} + 96q^{16} + 125q^{17} + 155q^{18} + 198q^{19} + 244q^{20} + O(q^{21});$$

(2.5)
$$F_3(0) = F_2(0) = \sum_{n=0}^{\infty} \sum_{0 \le r \le \frac{3n}{2}} \frac{q^{\frac{n(3n+1)}{2}-r}}{(q^2;q^2)_r(q)_{3n-2r}} = \frac{\Pi(q^6,q^9,q^{15};q^{15})}{(q)_{\infty}};$$

(2.6)
$$F_3(1) = F_2(1) = \sum_{n=1}^{\infty} \sum_{0 \le r \le \frac{3n-1}{2}} \frac{q^{\frac{n(3n+1)}{2}-r}}{(q^2;q^2)_r(q)_{3n-2r-1}} = \frac{q\Pi(q^3,q^{12},q^{15};q^{15})}{(q)_{\infty}}.$$

So, Baxter had conjectured the identity of each of the $F_k(\sigma_i)$ with the corresponding infinite products given above. His approach involved firstly, methods for the treatment of the expressions given in (1.22) so that the double series representations given above can be found. Secondly, a set of transformations was required to allow identification of the double series with the appropriate infinite product expression.

3. Outline of proofs of Baxter's Regime II Conjectures

Our methodology differs from that of Baxter immediately. Baxter's development of series-product identities relied on the taking of the limit as m tends to ∞ in (1.22). We instead find representations for the partition functions arising in **Regime III** with m remaining fixed and finite. We then make use of the significant fact that when m is finite we can map from **Regime III** to **Regime II** by the transformation $q \mapsto 1/q$. Our solution of **Regime III** (on the way proving (1.1) to (1.16) relies on the two following polynomial identities:

$$(3.1)$$

$$\sum_{n,r\geq 0} q^{\frac{n(3n+1)}{2}} \begin{bmatrix} N-2n-2r \\ n \end{bmatrix}_q \begin{bmatrix} r+n \\ r \end{bmatrix}_{q^2} q^r = \sum_{\lambda=-\infty}^{\infty} (-1)^{\lambda} q^{\lambda(5\lambda+1)} \begin{bmatrix} N \\ \left\lfloor \frac{n-5\lambda}{2} \right\rfloor_q;$$

(3.2)

$$\sum_{n,r\geq 0} q^{\frac{n(3n+1)}{2}} \begin{bmatrix} N-2n-2r-1\\n \end{bmatrix}_q \begin{bmatrix} r+n\\r \end{bmatrix}_{q^2} q^r = \sum_{\lambda=-\infty}^{\infty} (-1)^{\lambda} q^{\lambda(5\lambda-3)} \begin{bmatrix} N\\ \left\lfloor \frac{n-5\lambda}{2} \right\rfloor + 1 \end{bmatrix}_q;$$

in which

(3.3)
$$\begin{bmatrix} N \\ M \end{bmatrix}_{q} = \begin{cases} \frac{(1-q^{N})(1-q^{N-1})\cdots(1-q^{N-M+1})}{(1-q^{M})(1-q^{M-1})\cdots(1-q)}, & \text{for } M \ge 0; \\ 0, & \text{for } M < 0, N \ge 0 \end{cases}$$

and [x] = the largest integer not exceeding x. If in (3.1) we let $N \to \infty$, the first result required for **Regime III**, that being (3.3), is obtained. Similarly, if $N \to \infty$ in (4.1), we obtain (1.14).

To obtain (2.1) to (2.6), we replace N by 3N + a $(a = 0, \pm 1)$ in (3.1) and (3.2), and then replace q by 1/q. Next multiply by the minimal power of q necessary to produce polynomials in q, and then let $N \to \infty$. This process produces the identities of series and products described in these (2.5) to (2.6), and the relationship between **Regimes II** and **III** that follows from the replacement of q by 1/q resulting in the identity with the various $F_k(\sigma_l)$.

4. TRANSFORMS BETWEEN BAXTER'S REGIME II AND REGIME III

The results described here belie the fact that the Rogers-Ramanujan type identities for **regime II** of the hard-hexagon model are now rigorously established. On the other hand, there are numerous interesting long range questions more of interest in the theory of partitions and q-series that have been extensively explored under three approaches:

Approach (i): Suppose the Rogers-Ramanujan partition ideal [see Andrews [4, chapter 8] for detailed discussion of partition ideals] is replaced by another classical partition ideal; what happens in **Regimes II**, **III**, and **IV** appropriately modified?

Approach (ii): The $q \rightarrow 1/q$ duality of **Regimes II** and **III** also exists between **Regimes I** and **IV**. In fact, the relevant polynomial identities for this latter relationship are

GEOFFREY B CAMPBELL

(4.1)
$$\sum_{j\geq 0} q^{j^2} \begin{bmatrix} N-j \\ j \end{bmatrix}_q = \sum_{\lambda=-\infty}^{\infty} (-1)^{\lambda} q^{\frac{\lambda(5\lambda+1)}{2}} \begin{bmatrix} N \\ \left\lfloor \frac{N-5\lambda}{2} \right\rfloor_q;$$

(4.2)
$$\sum_{j\geq 0} q^{j(j+1)} \begin{bmatrix} N-j \\ j \end{bmatrix}_q = \sum_{\lambda=-\infty}^{\infty} (-1)^{\lambda} q^{\frac{\lambda(5\lambda-3)}{2}} \begin{bmatrix} N \\ \left\lfloor \frac{N+1-5\lambda}{2} \right\rfloor + 1 \end{bmatrix}_q.$$

These identities were completely stated in Andrews [2] and have their origin in the work of Schur [37]. The arguments used to obtain (2.1) to (2.6) from (3.1) and (3.2) may now be turned on (4.1) and (4.2) to obtain (1.16) to (1.19), a relationship previously unnoticed. Thus a *duality theory* between various sets of identities of the Rogers-Ramanujan type presents for exploration.

Approach (iii): The analytic duality described above has a corresponding manifestation in the partition-theoretic interpretations of the various identities considered. Thus the well-known combinatorial interpretations of (1.12) and (1.13) are *dual* to the combinatorial interpretations of (1.16) to (1.19). Since the 1980s this duality has been developed in two distinct streams, via statistical mechanics Integrable Systems and in the theory of mathematical integer partitions.

5. The partition function of the Hard Hexagon Model

The hard hexagon model occurs within the framework of the grand canonical ensemble, where the total number of particles (the *hexagons*) is allowed to vary naturally, and is fixed by a chemical potential. In the hard hexagon model, all valid states have zero energy, and so the only important thermodynamic control variable is the ratio of chemical potential to temperature $\frac{\mu}{\kappa T}$. The exponential of this ratio, $z = \exp\left(\frac{\mu}{\kappa T}\right)$ is called the activity and larger values correspond roughly to denser configurations.

For a triangular lattice with N sites, the grand partition function is

(5.1)
$$Z(z) = \sum_{n} z^{n} g(n, N) - 1 + Nz + \frac{1}{2} N(N-7)z^{2} + \cdots$$

where g(n, N) is the number of ways of placing n particles on distinct lattice sites such that no two are adjacent. The function κ is defined by

(5.2)
$$\kappa(z) = \lim Z(z)^{1/N} = 1 + z - 3z^2 + \cdots$$

so that $\log(\kappa)$ is the free energy per unit site. Solving the hard hexagon model means (roughly) finding an exact expression for κ as a function of z.

The **mean density** ρ is given for small z by

(5.3)
$$\rho = z \frac{d \log(\kappa)}{dz} = z - 7z^2 + 58z^3 - 519z^4 + 4856z^5 - \cdots$$

The vertices of the lattice fall into three classes numbered 1, 2, and 3, given by the 3 different ways to fill space with hard hexagons. There are 3 local densities ρ_1, ρ_2, ρ_3 , corresponding to the 3 classes of sites. When the activity is large the system approximates one of these 3 packings, so the local densities differ, but when the activity is below a critical point the three local densities are the same. The critical point separating the low-activity homogeneous phase from the high-activity ordered phase is

$$z_c = \frac{(11+5\sqrt{5})}{2} = \phi^5 = 11.09017\dots$$

with golden ratio $\phi = \frac{1+\sqrt{(5)}}{2}$. Above the critical point the local densities differ and in the phase where most hexagons are on sites of type 1 can be expanded as

$$\rho_1 = 1 - z^{-1} - 5z^{-2} - 34z^{-3} - 267z^{-4} - 2037z^{-5} - \cdots,$$

$$\rho_2 = z^{-2} - 9z^{-3} + 80z^{-4} + 965z^{-5} - \cdots.$$

5.1. Solution. The solution is given for small values of $z < z_c$ by

$$z = \frac{-xH(x)^5}{G(x)^5},$$

$$\kappa = \frac{H(x)^3Q(x^5)^2}{G(x)^2} \prod_{n \ge 1} \frac{(1 - x^{6n-4})(1 - x^{6n-3})^2(1 - x^{6n-2})}{(1 - x^{6n-5})(1 - x^{6n-1})(1 - x^{6n})^2},$$

$$\rho = \rho_1 = \rho_2 = \rho_3 = \frac{-xG(x)H(x^6)P(x^3)}{P(x)},$$

where

$$G(x) = \prod_{n \ge 1} \frac{1}{(1 - x^{5n-4})(1 - x^{5n-1})},$$

$$H(x) = \prod_{n \ge 1} \frac{1}{(1 - x^{5n-3})(1 - x^{5n-2})},$$

$$P(x) = \prod_{n \ge 1} (1 - x^{2n-1}) = \frac{Q(x)}{Q(x)^2},$$

$$Q(x) = \prod_{n \ge 1} (1 - x^n).$$

For large $z > z_c$ the solution (in the phase where most occupied sites have type 1) is given by

$$z = \frac{G(x)^5}{xH(x)^5},$$

$$\kappa = x^{\frac{1}{3}} \frac{G(x)^3 Q(x^5)^2}{H(x)^2} \prod_{n \ge 1} \frac{(1 - x^{3n-2})(1 - x^{3n-1})}{(1 - x^{3n})^2},$$

$$\rho_1 = \frac{H(x)Q(x)(G(x)Q(x) + x^2H(x^9)Q(x^9))}{Q(x^3)^2},$$

$$\rho_2 = \rho_3 = \frac{x^2H(x)Q(x)H(x^9)Q(x^9)}{Q(x^3)^2},$$

$$R = \rho_1 - \rho_2 = \frac{Q(x)Q(x^5)}{Q(x^3)^2}.$$

GEOFFREY B CAMPBELL

The functions G and H turn up in the Rogers-Ramanujan identities, and the function Q is the Euler function, which is closely related to the Dedekind eta function. If $x = e^{2\pi i \tau}$, then $q^{-1/60}G(x)$, $x^{11/60}H(x)$, $x^{-1/24}P(x)$, $z, \kappa, \rho, \rho_1, \rho_2$, and ρ_3 are modular functions of τ , while $x^{1/24}Q(x)$ is a modular form of weight 1/2. Since any two modular functions are related by an algebraic relation, this implies that the functions κ, z, R, ρ are all algebraic functions of each other of quite high degree.

6. Rogers-Ramanujan shift from Mathematics to Physics

In 1998 the physicists Alexander Berkovich and Barry M. McCoy from State University of New York wrote an expository paper [17] explaining how the story of the Rogers-Ramanujan identities transitioned from mathematics to physics. Put simply, in 1894 L.J. Rogers [33] proved the following identities for a = 0, 1 between infinite series and products valid for |q| < 1,

(6.1)
$$\sum_{n=0}^{\infty} \frac{q^{n(n+a)}}{(q)_n} = \prod_{n=1}^{\infty} \frac{1}{(1-q^{5n-1-a})(1-q^{5n-4+a})}$$
$$= \frac{1}{(q)_{\infty}} \sum_{m=-\infty}^{\infty} (q^{n(10n+1+2a)} - q^{(5n+2-a)(2n+1)}),$$

where $(q)_n = \prod_{j=0}^n (1-q^j)$.

A fascinating quote from Berkovich and McCoy on Rogers-Ramanujan identities:

Think About It...

"For about the first 85 years after their discovery interest in these identities and their generalizations was confined to mathematicians. Many ingenious proofs and relations applying combinatorics, basic hypergeometric functions and Lie algebras were discovered by MacMahon, Rogers, Schur, Ramanujan, Watson, Bailey, Slater, Gordon, Göllnitz, Andrews, Bressoud, Lepowsky and Wilson; so by 1980 there were over 130 isolated identities and several infinite families of identities known.

The entry of these identities into physics occurred in the early 1980s when Baxter [11], Andrews, Baxter and Forrester (see [3] and [27]), and the Kyoto group [24] encountered (6.1) and various generalizations in the computation of order parameters of certain lattice models of statistical mechanics.

A further glimpse of the relation to physics is seen in the development of conformal field theory by Belavin, Polyakov and Zamolodchikov [14] and the form of computation of characters of representations of Virasoro algebra by Kac [28], Feigin and Fuchs [25] and Rocha-Caridi [32]. The occurrence of (6.1) in this context led Kac [29] to suggest that every modular invariant representation of Virasoro should produce a Rogers-Ramanujan type identity."

- Alexander Berkovich and Barry M. McCoy, State University of New York, 1998 The full relation between physics and Rogers-Ramanujan identities is more extensive than we may think from these first indications. Starting in 1993 both Berkovich and McCoy (see for example [30, 31], [15, 16, 17, 18], and [5]) have fused the physical insight of solvable lattice models in statistical mechanics with the classical work of the first 85 years and the recent developments in conformal field theory to greatly enlarge the theory of Rogers-Ramanujan identities. We can summarize the results of this work and present some of the current results. Our point of view for the rest of this chapter will be influenced by the statistical mechanics, but we will try to indicate where alternative viewpoints exist. Hopefully some language differences between physicists and mathematicians can be understood.

The work of the last 30 years arising in physics problems has given a new context and point of view in the study of Rogers-Ramanujan identities. The emphasis is not the same as in the earlier mathematical investigations and thus it is good to discuss this before giving detailed results.

In mathematics, Rogers-Ramanujan identities always were seen as partitions of integers equivalence theorems or as generating function algebraic identities for integer partitions. This latter mostly took the form of q-series identities where one side was a sum of $(q)_n$ terms and the other side was an infinite product of $(q)_n$ terms. In physics the discussion still centered around partitions, but now defined by terms of statistical mechanical and conformal field theory applications. In physics nowadays in most cases where we have generalizations of the identities between the two sums, a product form is not known. So in physics, a Rogers-Ramanujan identity means the equality of the sums without further reference to possible product forms.

What is interesting about the two diverging approaches to this topic, is that underlying the mathematics and the physics is a universality that embraces both approaches. However, for now, the two approaches not just to Rogers-Ramanujan identities, but to the theory of partitions, are distinct, but equally valid.

The full range of Rogers-Ramanujan identities is by no means yet understood and it is anticipated that both in the mathematics and in the physics there is much still left to be discovered. The physics has grown into innumerable research areas due to the now well-established fact that the Yang-Baxter equation has long been recognised as the masterkey to integrability, providing the basis for exactly solved models which capture the fundamental physics of a number of realistic classical and quantum systems. The theory of partitions has therefore become an essential tool for physicists involved with exactly solvable models, of which there is now, including the hard-hexagon model, the Heisenberg spin chain, the transverse quantum Ising chain, a spin ladder model, the Lieb-Liniger Bose gas, the Gaudin-Yang Fermi gas and the two-site Bose-Hubbard model. For an eloquent review of this in the context of *condensed matter to ultracold atoms* see Batchelor and Foerster [9].

7. Partition generating functions in Physics are polynomial generalizations

The second insight which is also present in the very first papers on the connection of Rogers-Ramanujan identities with physics, (see Baxter [11], Andrews, Baxter and Forrester [6], and Forrester and Baxter [27]) is the fact that the physics will often lead to polynomial identities, with an order depending on an integer L, which yield infinite series identities as $L \to \infty$. The polynomial generalization of (6.1) is the identity first proven by Andrews [2] in 1970,

(7.1)
$$F_a(L,q) = B_a(L,q)$$

where

(7.2)
$$F_a(L,q) = \sum_{n=0}^{\infty} q^{n(n+a)} \begin{bmatrix} L-n-a \\ n \end{bmatrix}_q$$

and

(7.3)
$$B_a(L,q) = \sum_{n=-\infty}^{\infty} (-1)^n q^{n(5n+1+2a)/2} \begin{bmatrix} L \\ \lfloor \frac{1}{2}(L-5n-a) \rfloor \end{bmatrix}_q$$

where $\lfloor x \rfloor$ denotes the integer part of x and the Gaussian polynomials (q-binomial coefficients) are defined for integer m, n by

(7.4)
$$\begin{bmatrix} n \\ m \end{bmatrix}_q = \begin{cases} \frac{(q)_n}{(q)_m(q)_{n-m}}, & 0 \le m \le n; \\ 0, & \text{otherwise.} \end{cases}$$

The identity (7.1) is obtained by using

$$\lim_{n \to \infty} \left[\begin{array}{c} n \\ m \end{array} \right]_q = \frac{1}{(q)_m}.$$

So (7.1) is a generalization of the identity (6.1) which we will call a Rogers-Ramanujan identity.

There are many known generalizations of $F_a(L,q)$ that are written in terms of the following function due to Kedem et al. [31]

(7.5)
$$f = \sum_{\text{restrictions}} q^{\frac{1}{2}\mathbf{m}\mathbf{B}\mathbf{m} - \frac{1}{2}\mathbf{A}\mathbf{m}} \prod_{\alpha=1}^{n} \left[\begin{array}{c} ((1-\mathbf{B})\mathbf{m} + \frac{\mathbf{u}}{2})_{\alpha} \\ m_{\alpha} \end{array} \right]_{q}$$

where **m**, **u** and **A** are **n** dimensional vectors and **B** is an $n \times n$ dimensional matrix and the sum is over all values of the variables m_{α} possibly subject to some restrictions (such as being even or odd). In many cases the *q*-binomials are defined by (7.4) but there do occur cases with an extended definition

(7.6)
$$\begin{bmatrix} m+n \\ m \end{bmatrix}_q = \begin{cases} (q^{n+1})_m, & \text{for } m \ge 0, n \text{ integers;} \\ 0, & \text{otherwise.} \end{cases}$$

This brings (7.4) to cover negative n.

The function (7.5) is regarded as the partition function for a collection of n different species of free massless (right moving) fermions with a linear energy momentum relation $e(P_j^{\alpha})_{\alpha} = vP_j^{\alpha}$ where the momenta are quantized in units of $2\pi/M$ and are chosen from the sets

(7.7)
$$P_j^{\alpha} \in \left\{ P_{\min}^{\alpha}, P_{\min}^{\alpha} + \frac{2\pi}{M}, P_{\min}^{\alpha} + \frac{4\pi}{M}, P_{\min}^{\alpha} + \frac{6\pi}{M}, \dots, P_{\max}^{\alpha} \right\}$$

where

$$P_{\min}^{\alpha}(\mathbf{m}) = \frac{\pi}{M} \left[((\mathbf{B} - 1)\mathbf{m})_{\alpha} - \mathbf{A}_{\alpha} + 1 \right]$$

and

$$P_{\max}^{\alpha}(\mathbf{m}) = -P_{\min}^{\alpha}(\mathbf{m}) + \frac{\pi}{M} \left(\frac{\mathbf{u}}{2} - \mathbf{A}\right)$$

with the Fermi exclusion rule $P_j^{\alpha} \neq P_k^{\alpha}$ for $j \neq k$ and all $\alpha = 1, 2, \ldots, n$.

14

8. The Elliptic Q-Gamma Function

The elliptic gamma function is a generalization of the q-gamma function, which is itself the q-analog of the usual gamma function. It was first examined in detail by Ruijsenaars in 1997 as an integrable function (see [35]), and can be expressed in terms of the triplegamma function. It is defined by

(8.1)
$$\Gamma(z; p, q) = \prod_{m=0}^{\infty} \prod_{n=0}^{\infty} \frac{1 - p^{m+1} q^{n+1}/z}{1 - p^m q^n z}.$$

The following relation is easily seen to apply

(8.2)
$$\Gamma(z; p, q) \Gamma(pq/z; p, q) = 1.$$

Using the q-theta function $\theta(z;q) = \prod_{n=0}^{\infty} (1-q^n z)(1-q^{n+1}/z)$, it is well-known that

$$\begin{split} \Gamma(pz;p,q) &= \theta(z;q)\Gamma(z;p,q), \\ \Gamma(qz;p,q) &= \theta(z;p)\Gamma(z;p,q). \end{split}$$

At p = 0 we have $\Gamma(z; 0, q) (z, q)_{\infty} = 1$ using the infinite q-Pochhammer notation.

As with the usual gamma function, there is a duplication, a triplication and multiplication formula. We need firstly to define the function

(8.3)
$$\widetilde{\Gamma}(z;p,q) := \frac{(q,q)_{\infty}}{(p,p)_{\infty}} (\theta(q,p))^{1-z} \prod_{m=0}^{\infty} \prod_{n=0}^{\infty} \frac{1-p^{m+1}q^{n+1-z}}{1-p^m q^{n+z}}.$$

We follow the approach by Felder and Varchenko [26] who proved the

Theorem 8.1. If $r = q^n$ then

(8.4)
$$\widetilde{\Gamma}(nz; p, q) \widetilde{\Gamma}(\frac{1}{n}; p, r) \widetilde{\Gamma}(\frac{2}{n}; p, r) \cdots \widetilde{\Gamma}(\frac{n-1}{n}; p, r) = \left(\frac{\theta(r; p)}{\theta(q; p)}\right)^{nz-1} \widetilde{\Gamma}(z; p, q) \widetilde{\Gamma}(z + \frac{1}{n}; p, r) \cdots \widetilde{\Gamma}(z + \frac{n-1}{n}; p, r).$$

The two most significant corollaries of theorem 8.1 are as follows:

Corollary 8.1. The Duplication Formula.

(8.5)
$$\widetilde{\Gamma}(2z;p,q)\widetilde{\Gamma}(\frac{1}{2};p,q^2) = \left(\frac{\theta(q^2;p)}{\theta(q;p)}\right)^{2z-1} \widetilde{\Gamma}(z;p,q^2)\widetilde{\Gamma}(z+\frac{1}{2};p,q^2).$$

Corollary 8.2. The Triplication Formula. (8.6)

$$\widetilde{\Gamma}(3z;p,q)\widetilde{\Gamma}(\frac{1}{3};p,q^3)\widetilde{\Gamma}(\frac{2}{3};p,q^3) = \left(\frac{\theta(q^3;p)}{\theta(q;p)}\right)^{3z-1} \widetilde{\Gamma}(z;p,q^3)\widetilde{\Gamma}(z+\frac{1}{3};p,q^3)\widetilde{\Gamma}(z+\frac{2}{3};p,q^3)$$

The above theorem with the Duplication and Triplication formula cases may be worthwhile examining and comparing with the functional equations given in chapter 15 of the book by Campbell [23] soon to appear. For example, is there a connection applicable between the Multiplication Formula of the Elliptic Gamma Function and the functions cited, namely, **Definition 8.1.** Define the function $F_n(t)$ for all complex numbers a, t with |a|, |t| < 1, and for $n \ge 1$ with $|q_1|, |q_2|, ..., |q_n| < 1$ by the sequence of functions $F_n(t)$ with

(8.7)
$$F_0(-;a,t) = \frac{1-at}{1-t}$$

and for all of $|q_1|, |q_2|, |q_3|, ..., |q_n| < 1$, by

(8.8)
$$F_n(q_1, q_2, q_3, ..., q_n; a, t) = \prod_{\alpha_1, \alpha_2, \alpha_3, ..., \alpha_n \ge 0} \frac{1 - q_1^{\alpha_1} q_2^{\alpha_2} ... q_n^{\alpha_n} at}{1 - q_1^{\alpha_1} q_2^{\alpha_2} ... q_n^{\alpha_n} t} \equiv \sum_{k=1}^{\infty} {}_n A_k t^k$$

where $\alpha_1, \alpha_2, \alpha_3, ..., \alpha_n$ may be all zero.

We now state a functional equation theorem for $F_n(t)$, next highlighting the 2D, 3D and 4D cases as examples in a corollary.

Theorem 8.2. (see Campbell [23, Chapter 15]) If the left side of (8.8) is for the moment considered only as a function, $F_n(t)$, of t, then the following finite product holds,

(8.9)
$$\frac{1-at}{1-t} = \frac{F_n(t) \left(\prod_{S_2} F_n(q_{i_1}q_{i_2}t)\right) \left(\prod_{S_4} F_n(q_{i_1}q_{i_2}q_{i_3}q_{i_4}t)\right) \dots}{\left(\prod_{S_1} F_n(q_{i_1}t)\right) \left(\prod_{S_3} F_n(q_{i_1}q_{i_2}q_{i_3}t)\right) \left(\prod_{S_5} F_n(q_{i_1}q_{i_2}q_{i_3}q_{i_4}q_{i_5}t)\right) \dots}$$

the numerator products each being over the nth order even symmetric combinations whilst the denominator products are over the nth order odd symmetric combinations of the variables. We have used the notation $S_1, S_2, S_3, S_4, ...$ to denote the respective nth order symmetric function product of terms.

It is worth noting that we have used the abbreviation $F(t) = F_n(q_1, q_2, q_3, ..., q_n; a, t)$ and the right side of (8.9) is a finite product of such functions going up to the *n*th order symmetric function product of terms. It is also seen here that (8.9) is a functional equation for a *generalized right side of (8.8)* form of the *q*-binomial product. As an illustration we next give the n = 1 up to n = 4 examples of the functional equation (8.8), with $q_1 \rightarrow x, q_2 \rightarrow y, q_3 \rightarrow z$. With these variables, we assert

Corollary 8.3. (see Campbell [23, Chapter 15]) If for |w| < 1, |x| < 1, |y| < 1, |z| < 1, and all complex numbers a and t,

$$F_{1}(t) = F_{1}(z; a, t) = \prod_{j \ge 0} \frac{1 - z^{j} at}{1 - z^{j} t},$$

$$F_{2}(t) = F_{2}(y, z; a, t) = \prod_{j,k \ge 0} \frac{1 - y^{j} z^{k} at}{1 - y^{j} z^{k} t},$$

$$F_{3}(t) = F_{3}(x, y, z; a, t) = \prod_{j,k,l \ge 0} \frac{1 - x^{j} y^{k} z^{l} at}{1 - x^{j} y^{k} z^{l} t},$$

$$F_{4}(t) = F_{4}(w, x, y, z; a, t) = \prod_{j,k,l,m \ge 0} \frac{1 - w^{j} x^{k} y^{l} z^{m} at}{1 - w^{j} x^{k} y^{l} z^{m} t},$$

$$etc.$$

(8.10)

16

then

$$\begin{aligned} \frac{1-at}{1-t} &= \frac{F_1(t)}{F_1(zt)}, \\ \frac{1-at}{1-t} &= \frac{F_2(t)F_2(yzt)}{F_2(yt)F_2(zt)}, \\ \frac{1-at}{1-t} &= \frac{F_3(t)F_3(yzt)F_3(xyt)F_3(xzt)}{F_3(xt)F_3(yt)F_3(zt)F_3(xyzt)}, \\ \frac{1-at}{1-t} &= \frac{F_4(t)F_4(yzt)F_4(xyt)F_4(wxt)F_4(wyt)F_4(wyzt)F_4(wxyzt)}{F_4(wt)F_4(wt)F_4(wyt)F_4(wyzt)F_4(wyzt)}, \end{aligned}$$

```
(8.11) etc.
```

The above equations are suggestive of three as yet undeveloped theories:

- 1) The functional equations for generating functions imply recurrence relations for the nD vector partitions themselves. These are higher dimensional extensions of the Ramanujan integer partition congruences. See Bruinier [19], and Bruinier et al. [20] to [21] for unrestricted partition congruences.
- 2) An *n*D *m*-ary vector partition congruence theory; extending the binary partition congruences of Rödseth and Gupta. See Alkauskas [1] for an account of binary integer congruences.
- 3) The nD generating functions suggest their own versions of Duplication Formulas, Triplication formulas, and Multiplication formulas, similar to those stated here for the Elliptic Gamma function. This function is *integrable* and therefore related to existing theories in Statistical Mechanics, Yang-Baxter equations, Boltzmann weights and phase transitions for Lattice Models. (See the original 1997 account by Ruijsenaars [35], and more recently the 2016 physics in Bazhanov et al. [13] for example.

This narrative adjourns to future discussions on topics of functional equations, integrable functions, vector partition congruences and nD Lattice Models. The Elliptic Gamma Function is a 2D example of an nD function that may *play out* in a similar way in higher Euclidean spaces. The Elliptic Gamma Function seems useful for vector partition concepts not yet articulated. Also, our chapter 15 of Campbell [23, Chapter 15] on Vector Partition generating functions and functional equations may be related to solvable models associated with Yang-Baxter equations. This topic brings us deeply into the physics of Statistical Mechanics, but is beyond the present scope of our book. It may bring future researchers to consider cases of single or multiple spin degrees of freedom at each site of a higher dimensional lattice. The Yang-Baxter equations. For a 2023 contemporary appraisal of Yang-Baxter equations related to Brauer algebras see Cañada [22].

Think About It...

The 1982 book by Rodney Baxter *Exactly solved models in statistical* mechanics has over 7,180 citations as at early 2023.

These citations are probably split approximately evenly between physics and mathematics research articles and books; the breakup as follows: Highly Influential Citations = 919; Background Citations = 2,656; Methods Citations = 1,857; Results Citations = 125.

References

- ALKAUSKAS, G. Generalization of the Rödseth-Gupta Theorem on Binary Partitions. Lithuanian Mathematical Journal, April 2003.
- [2] ANDREWS, G.E. A polynomial identity which implies the Rogers-Ramanujan identities. Scr. Math., 28:297–305, 1970.
- [3] ANDREWS, G.E. The hard-hexagon model and Rogers-Ramanujan type identities. Proc Nat. Acad. Sci. USA, 78:5290–5292, 1981.
- [4] ANDREWS, G.E. and ASKEY, R. Enumeration of Partitions: The Role of Eulerian Series and q-Orthogonal Polynomials. In Martin Aigner, editor, *Higher Combinatorics*, pages 3–26, Dordrecht, 1977. Springer Netherlands.
- [5] ANDREWS, G.E. and BERKOVICH, A. A trinomial analogue of Bailey's lemma and N = 2 superconformal invariance. *Comm. Math. Phys.*, 192:245–260, 1998.
- [6] ANDREWS, G.E., BAXTER, R.J. and FORRESTER, P.J. Eightvertex SOS model and generalized Rogers-Ramanujan-type identities. J. Stat. Phys., 35:193–266, 1984.
- [7] BAILEY, W.N. Some identities in combinatory analysis. Proc. Lond. Math. Soc. Second Ser., 49:421–435, 1947.
- [8] BAILEY, W.N. Identities of the Rogers-Ramanujan type. Proc. Lond. Math. Soc. Second Ser., 50:1–10, 1949.
- [9] BATCHELOR, M.T. and FOERSTER, A. Yang-Baxter integrable models in experiments: From condensed matter to ultracold atoms. J. Phys. A: Math. Theor., 49(17), March 2016.
- [10] BAXTER, R.J. Hard hexagons: exact solution. Journal of Physics A: Mathematical and General, 13(3), 1980. L61-L70, Bibcode 1980JPhA...13L..61B.
- [11] BAXTER, R.J. Rogers-Ramanujan identities in the hard hexagon model. J. Stat. Phys., 26:427, 1981.
- [12] BAXTER, R.J. Exactly Solved Models in Statistical Mechanics. Academic Press, New York, 1982. Reprinted 1989 and 2007.
- [13] BAZHANOV, V.V., KELS, A.P. and SERGEEV, S.M. Quasi-classical expansion of the startriangle relation and integrable systems on quad-graphs. J. Phys. A: Math. Theor., 49:44pp., October 2016.
- [14] BELAVIN, A.A., POLYAKOV, A.M. and ZAMOLODCHIKOV, A.B. Infinite conformal symmetry in two-dimensional quantum field theory. *Nucl. Phys. B*, 241:333, 1984.
- [15] BERKOVICH, A. Fermionic counting of RSOS states and Virasoro character formulae for the unitary minimal series M(v, v + 1): Exact results. Nucl. Phys. B, 431(1-2):315-348, 1994.
- [16] BERKOVICH, A. and MCCOY, B.M. Continued fractions and fermionic representations for characters of M(p, p') minimal models. Lett. Math. Phys., 37(49), 1996.
- [17] BERKOVICH, A. and MCCOY, B.M. Rogers-Ramanujan Identities: A Century of Progress from Mathematics to Physics. *Documenta Mathematica*, Extra Volume ICM 1998 -III:163–172, 1998.
- [18] BERKOVICH, A., MCCOY, B.M. and SCHILLING, A. Rogers-Schur-Ramanujan type identities for M(p, p') minimal models of conformal field theory. *Comm. Math. Phys.*, 191(325), 1998.
- [19] BRUINIER, J.H. Borcherds products on O(2, l) and Chern classes of Heegner divisors, volume 1780 of Springer Lecture Notes in Mathematics. Springer-Verlag, 2002.
- [20] BRUINIER, J.H. and FUNKE, J. On two geometric theta lifts. Duke Math. J., 125:45–90, 2004.
- [21] BRUINIER, J.H., and ONO, K. Algebraic formulas for the coefficients of halfintegral weight harmonic weak Maass forms. Adv. Math., 246:198–219, 2013.
- [22] CANADAS, A.M.; BALLESTER-BOLINCHES, A.; and GAVIRIA, I.D.M. Solutions of the Yang-Baxter Equation Arising from Brauer Configuration Algebras. *Computation*, 11(1), 2023.
- [23] CAMPBELL, G.B. Vector Partitions, Visible Points, and Ramanujan Functions, CRC Press, Taylor and Francis Group, Boca Raton, London, New York, A Chapman & Hall Book, 2024. ISBN: 978-1-032-00366-5 (hbk), ISBN: 978-1-032- 00432-7 (pbk), ISBN: 978-1-003-17415-8 (ebk), DOI: 10.1201/9781003174158.

- [24] DATA, E., JIMBO, M., KUNIBA, A., MIWA, T. and OKADO, M. Exactly solvable SOS models: local height probabilities and theta function identities. *Nucl. Phys. B*, 290(231), 1987.
- [25] FEIGIN, B.L. and FUCHS, D.B. Verma modules over the Virasoro algebra. Funct. Anal. Appl., 17(3):241–242, 1983.
- [26] FELDER, G. and VARCHENKO, A. Multiplication Formulas for the Elliptic Gamma Function. arxiv preprint, Dec 2002. arXiv:math/0212155v1 [math.QA]. www.doi.org/10.48550/arXiv.math/0212155.
- [27] FORRESTER, P.J. and BAXTER, R.J. Further exact solutions of the eightvertex SOS model and generalizations of the Rogers-Ramanujan identities. J. Stat. Phys., 38(435), 1985.
- [28] KAC, V.G. Contravariant form for infinite-dimensional Lie algebras and superalgebras. Lect. Notes in Phys., 94:441, 1979.
- [29] KAC, V.G. Modular invariance in mathematics and physics. In Proceedings of the AMS Centennial Symposium, volume 337. American Mathematical Society, 1992.
- [30] KEDEM, R., KLASSEN, T.R., MCCOY, B.M. and MELZER, E. Fermionic quasi-particle representations for characters of $(G^{(1)})_1 \times (G^{(1)})_1/G^{(1)})_2$. *Phys. Lett. B*, 304:263–270, 1993.
- [31] KEDEM, R., KLASSEN, T.R., MCCOY, B.M. and MELZER, E. Fermionic sum representations for conformal field theory characters. *Phys. Lett. B*, 307:68–76, 1993.
- [32] ROCHA-CARIDI, A. Vertex Operators in Mathematics and Physics. Springer, Berlin, 1985. ed. J. Lepowsky, S. Mandelstam and I.M. Singer.
- [33] ROGERS, L.J. Second memoir on the expansion of certain infinite products. Proc. London Math. Soc., 25:318–343, 1894.
- [34] ROGERS, L.J. On Two Theorems of Combinatory Analysis and Some Allied Identities. Proc. Lond. Math. Soc. Second Ser., 16:315–336, 1917.
- [35] RUIJSENAARS, S.N.M. First order analytic difference equations and integrable quantum systems. J. Math. Phys., 38(2):1069–1146, February 1997.
- [36] SCHUR, I. Ein Beitrag zur additive Zahlentheorie und zue Theorie der Kettenbruche. S.-B. Preuss, Akad. Wies. Phys.-Math, Kl., pages 302–321, 1973. First printed 1917: Proceedings of the Royal Prussian Academy of Sciences in Berlin; Reprinted 1973: Ges. Abhandlungen, Vol. 2, pp. 117-136.
- [37] SCHUR, I. Zur additive Zahlentheorie und zue Theorie der Kettenbruche. S.-B. Preuss, Akad. Wies. Phys.-Math, Kl., pages 488–495, 1973. First printed 1917: Proceedings of the Royal Prussian Academy of Sciences in Berlin; Reprinted 1973: Ges. Abhandlungen, Vol. 3, pp. 43-50.
- [38] SLATER, L.J. Generalized Hypergeometric Series. Cambridge University Press, 1966.

MATHEMATICAL SCIENCES INSTITUTE, THE AUSTRALIAN NATIONAL UNIVERSITY, CANBERRA, ACT, 0200, AUSTRALIA

Email address: Geoffrey.Campbell@anu.edu.au