# The Pioneer anomaly as an effect of the dynamics of time

Antonio F. Rañada Facultad de Física, Universidad Complutense, E-28040 Madrid, Spain

1 July 2004

#### Abstract

A model is presented in which the Pioneer anomaly is not related to the motion of the spaceship, but is a consequence of the acceleration of the cosmological proper time  $\tau$  with respect to the coordinate parametric time t, i. e. of the value of  $d^2\tau/dt^2$  (> 0). Assuming flat spatial sections and that all the matter and energy is uniformly distributed throughout the universe, it is shown that this inequality is an effect of the background gravitational potential of the entire universe. According to this model, the light speed, while being constant if defined with respect to  $\tau$  (i. e. as  $d\ell/d\tau$ ) as is required in general relativity, would suffer an adiabatic secular acceleration,  $a_{\ell} = dc/dt > 0$ , if defined in terms of t (i. e. as  $d\ell/dt$ ). It turns out that such an adiabatic acceleration of light (according to the second definition), and a small acceleration of the Pioneer towards the Sun  $a_P$  could be mistaken the one for the other, because they do have the same fingerprint: a blue shift. However, this shift would be quite unrelated to any anomalous motion of the Pioneer, being just an observational effect of the acceleration of light with respect to time t, in such a way that  $a_P = a_\ell/2$ . A simple estimation predicts  $a_P \simeq 5.2 \times 10^{-10}$  m/s, just about 40 % smaller than the so-called Pioneer acceleration, which would correspond to the blueshift. The Pioneer anomaly turns out then to be an interesting case of the dynamics of time, its explanation involving the interplay between the two times  $\tau$  and t. The view presented here is the relativistic version of a previous Newtonian model.

PACS numbers: 04.80.Cc, 95.10.Eg, 95.55.Pe

Key words: Pioneer acceleration, Light speed, Dynamics of time.

#### Contents

- 1. Introduction
  - 1.1 The observation
  - 1.2 The purpose of this paper
  - 1.3 The dynamics of time and the two definitions of light speed
  - 1.4 Constant of light speed and observed light speed
  - 1.5 Equal fingerprints of two different phenomena
  - 1.6 Plan of the paper
- 2. On the speed of light and the coordinate time
- 3. The background gravitational potential and the acceleration of the non-proper light speed
- 4. Estimation of the adiabatic acceleration of the non-proper speed of light
  - 4.1 Time variation of the non-proper speed of light
- 4.2 Estimation of the non-proper accelerations of light and of the acceleration of the clocks
  - 5. The acceleration of cosmological time with respect to coordinate time 5.1 A toy model
  - 6. Summary and conclusions

Acknowledgements

References

#### 1 Introduction

1.1 The observation. Anderson et al reported in 1998 the observation of an anomalous acceleration  $a_{\rm P}$  in the Pioneer 10/11, Galileo and Ulysses spacecrafts, equal to  $a_{\rm P}=(8.74\pm1.33)\times10^{-10}{\rm m/s}^2$ , constant and directed towards the Sun [1]. More precisely, they observed a Doppler blue shift in the radio signals from these ships, which increases linearly in time, so that

$$\dot{\nu}/\nu = 2a_{\rm P}/c \,(= \text{const})\,,\tag{1}$$

where the factor 2 is because the Doppler effect refers to a two-way signal. Obviously, its simplest interpretation is that the ships were not following the predicted orbits but had an extra unmodelled acceleration towards the Sun, as if our star pulled a bit too much from them with a force independent of the distance. The effect is still unexplained and, intriguingly enough, does not show up in the planets [2]. Anderson et al said in 1998: "it is interesting to speculate on the possibility that the origin of the anomalous signal is new physics [1], and in 2002: "The veracity of the signal is now undisputed, although the source of the anomaly, some systematic or some not understood physics, is subject to debate" [3]. For an interesting argument showing that it may not necessarily be due to systematics, see [4].

1.2 The purpose of this paper. In some previous papers [5, 6, 7], a Newtonian model was presented that proposes an explanation for the Pioneer riddle, showing that it could be due to a universal and adiabatic acceleration of light, which would have the same observational fingerprint as the observed blue shift. That model is based on an application of the fourth Heisenberg relation to the sea of virtual electron-positron pairs which, combined with the expansion of the universe, would produce a progressive decrease of the optical density of the quantum vacuum, more precisely a progressive decrease of the permittivity and permeability of empty space, the consequence being a very small acceleration of light. The purpose of this paper is to propose a relativistic version of that Newtonian model. It must be emphasized that a near Newtonian situation will be always assumed in this paper, whenever the observed light speed will be mentioned.

1.3 The dynamics of time and the two definitions of the light speed. A primordial element of this model is the dynamics of time, which will be

understood here as the functional relation  $\tau = \tau(t)$  between the cosmological proper time  $\tau$  and the parametric coordinate time t. It is important, therefore, to emphasize that the light speed can be defined in general relativity in two different ways, (i) with respect to the cosmological proper time  $\tau$ ,  $c^* = \mathrm{d}\ell/\mathrm{d}\tau$  (= constant), and (ii) with respect to the coordinate time t,  $c = \mathrm{d}\ell/\mathrm{d}t = c(\mathbf{r},t)$ , where  $\mathrm{d}\ell$ ,  $\mathrm{d}\tau$  and  $\mathrm{d}t$  are elements of spatial distance, proper time and coordinate time along a null geodesic, respectively. These two speeds, both used in this paper, will be denoted as  $c^*$  and c, respectively, and will be called "proper light speed" and "non-proper speed of light speed". The derivative of c with respect to t will be called non-proper acceleration of light. The first is a universal constant of nature, the second is not but, quite on the contrary, depends generally on space and time  $c = c(\mathbf{r}, t)$ . Indeed, the element of interval can be written (assuming for simplicity that  $g_{0i} = 0$ )

$$ds^2 = c^{*2}d\tau^2 - d\ell^2, \qquad (2)$$

with  $d\tau = \sqrt{g_{00}} dt$  and  $d\ell^2 = g_{ij} dx^i dx^j$ , so that  $c^*$  is constant and c is

$$c = c(\mathbf{r}, t) = c^* \sqrt{g_{00}}. \tag{3}$$

Near the Newtonian limit,  $g_{00} \simeq 1 + 2\Phi/c^2$ , at first order,  $\Phi$  being the gravitational potential, so that

$$c = c(\mathbf{r}, t) = c^* [1 + \Phi(\mathbf{r}, t)/c^2], \qquad (4)$$

If  $c_0$  is the value of c at a reference terrestrial laboratory R, i.e.  $c_0 = c^*(1 + \Phi_R/c_0^2)$ ,

$$c = c(\mathbf{r}, t) = c_0[1 + \Phi(\mathbf{r}, t) - \Phi_R/c_0^2],$$
 (5)

The expression "light speed" means usually the constant  $c^*$  which, being a universal constant, is of the utmost importance. However, the non-proper light speed c is also used in some interesting cases. For instance, in the study of the bending of a light ray that grazes the Sun surface. It is observed that the bending angle is  $\phi = 1.75$ ". Let M and R be the mass and radius of the Sun. The interval around any star is given by the Schwarzschild metric, what implies that  $c = c(r) = c_{\infty}(1 - \eta R/r)$ , with  $c_{\infty} = c(\infty)$  and  $\eta = GM/c_{\infty}^2R \simeq 2.1 \times 10^{-6}$ . Einstein gave two formulae for this effect. The first (1907) is based only in the equivalence principle and gives  $\phi = 2\eta = 0.875$ ", just one half of the observed effect. The second (1916), in the frame of general relativity, gives the complete result  $\phi = 4\eta = 1.75$ ". The first one

can be obtained simply by considering the propagation of a wave light with the previous value of the light speed, in other words as the solution of the variational problem

$$\delta T = \delta \int_1^2 \frac{1 + \eta R/r}{c_\infty} \, \mathrm{d}\ell = 0 \,, \tag{6}$$

where  $\mathrm{d}\ell = \mathrm{d}x^2 + \mathrm{d}y^2 + \mathrm{d}z^2$  is the euclidean line element, *i.e.* as a consequence of the application of the Fermat principle to the non-proper light speed c. The complete effect is obtained by taking instead the non-euclidean spatial line element of the Schwarzschild geometry. Note that this implies that the problem is solved by assuming that the light propagates through with speed  $c = c_{\infty}(1 - \eta R/r)$  (taking into account the Riemannian character of the spatial metric). The same situation appears in the study of the delay of the radar echoes from the planets.

It will be assumed in this paper, for simplicity, that all the matter and energy in the universe are uniformly distributed. Since the universe is expanding, it is clear that c = c(t) and that there is a functional relation between the two times  $\tau = \tau(t)$ , determined by the background gravitational potential of all the universe. All this means that

$$c(t) = \frac{\mathrm{d}\ell}{\mathrm{d}\tau} \frac{\mathrm{d}\tau}{\mathrm{d}t} = c^* \frac{\mathrm{d}\tau}{\mathrm{d}t} \quad \text{and} \quad \frac{\mathrm{d}c(t)}{\mathrm{d}t} = c^* \frac{\mathrm{d}^2\tau}{\mathrm{d}t^2}. \tag{7}$$

Equation (7) states that the light speed c increases if the proper time  $\tau$  accelerates with respect to the coordinate time t, its time derivative being equal to the proper light speed  $c^*$  times the second derivative of  $\tau$  with respect to t. The notation  $a_{\ell} = \dot{c}(t) = \mathrm{d}c(t)/\mathrm{d}t$  will be used,  $a_{\ell}$  being termed the non-proper acceleration of light, or just the acceleration of light for short when there is no risk of confusion. It will be shown that c(t) must increase as a consequence of the interaction of light with the background gravitational field due to all the matter and energy in e universe, so that  $a_{\ell} > 0$ . This will give an explanation for the Pioneer anomaly.

It must be emphasized again that the proper speed of light  $c^*$  is constant in this model, as it must happen in general relativity. In fact,  $c^*$  it is a universal constant of nature. Expressions such as "acceleration of light" or "variation of light speed" are always applied here to the light speed c(t), not to  $c^*$ . In this sense, this model is not a variable light speed theory (of the class frequently known as VLS theories), but a model inside general relativity. A standard of velocity is still given by the constant  $c^*$ , which will allow

to measure the variation in time of the dimensional quantity c(t) by means of measurements of the dimensionless value of  $c(t)/c^*$ . The previous considerations are needed because the expression "light speed" means several different things and we must understand very carefully which one of its meanings we are using (see [8] for an interesting discussion).

1.4 Constant of light speed and observed light speed. This paper presents a relativistic argument which shows that the Pioneer anomaly could be a manifestation of the adiabatic increase of the non-proper light speed c. But it is often stated that it is impossible to measure an eventual variation in time of c, because it is a dimensional constant. This does not deny necessarily the existence of an acceleration of light. However, since the possibility can not be excluded that the standards of length or of time might be changing also, such a measurement would be ambiguous or meaningless. According to this view, widely shared by the metrologists, only non-dimensional constants can be measured as functions of time, as would be the case of the fine structure constant for instance.

However, this argument does not consider the fact that the quantity known usually as "the light speed" has two meanings: the observed speed of light and a constant of physics that appears in Einstein's equations (in fact, it has more than two meanings but these two are more important for the purpose of this work, see [8]). It is usually assumed as a matter of fact that these two faces are the same one, so that the values of the universal constant (a theoretical concept) and of the observed light speed (an empirical datum) are always identical. This seems to be the case, but physics is an experimental science and this must not be taken for granted necessarily. It can not be excluded a priori that the two values could be different at large scales. In that case, the observed light speed  $c_{\rm obs}$  could be a function of the constant light speed and the gravitational potential  $\Phi$  such as

$$c_{\text{obs}} = c^* \times g\left(\Phi/c^{*2}\right) \,, \tag{8}$$

at first order in potential, where  $c^*$  is the constant of nature and g is a dimensionless function. In that case and if  $\Phi$  depends on time, a time variation of the light could be detected meaningfully by measuring the dimensionless quantity  $c_{\text{obs}}/c^* = g(\Phi/c^{*2})$ . Even if the observed light speed changes in time, there would be still a standard of speed provided by the constant  $c^*$  (or any function of  $c^*$  which could be suitable).

1.5 Equal fingerprints of two different phenomena. It turns out (and this must be stressed) that an adiabatic increase of the coordinate time speed of light c (so that  $a_{\ell} = \dot{c} > 0$ , see section 1.3) has the same observational signature as a Doppler blue shift due to the approaching motion of the source, as was shown in [6, 7]. This can be understood easily, just by taking the wave equation with variable speed for the field, for instance  $[\nabla^2 - c^{-2}(t)\partial_t^2]\mathbf{E} = 0$ , with  $c(t) = c_0 + a_{\ell}t$  (the same result is obtained by using  $\mathbf{H}$  instead; note that the time in this equation is clearly the coordinate time). At first order in  $a_{\ell}$  the plane wave solutions are  $\mathbf{E} = \mathbf{E}_0 \exp\{-2\pi i[z/\lambda - (\nu_0 + \dot{\nu}t/2)t]\}$  with constant  $\lambda$  and

$$\dot{\nu}/\nu_0 = a_\ell/c \,, \tag{9}$$

which must be compared with (1) (the frequency is the time derivative of the phase, i.e.  $d[(\nu_0 + \dot{\nu}t/2)t]/dt = \nu_0 + \dot{\nu}t = \nu_0[1 + a_\ell t/c])$ . In other words, all the increase of the speed is used to increase the frequency, the wavelength remaining constant. This has two consequences: (i) an increase of non-proper light speed c with respect to the coordinate time t,  $a_\ell = 2a_P$ , would produce an extra blue shift in radio signals, linearly increasing in time and having the same observational effect as the unmodelled acceleration equal to  $a_P$  towards the Sun of the Pioneer and other spacecrafts, and (ii) the atomic clocks would be accelerating, since their periods would be decreasing. The blue shift would be due to the acceleration of light, not to the motion of the ships which would be following then and now the standard laws of gravitation without any extra pull from the Sun. This means that the Pioneer anomaly would be explained by finding a good reason for the coordinate time light speed c to increase.

1.6 Plan of the paper. In section 2, the speed of light defined in terms of the coordinate time c(t) will be considered, with basis on some developments by Einstein after proposing his equivalence principle and on the ideas of electric permittivity and magnetic permeability of gravitational origin. After that, it will be shown in section 3 that the background gravitational potential of all the universe implies an acceleration of light with respect to the coordinate time t, in the sense that  $a_{\ell} = \dot{c} > 0$  (see section 1.3). Section 4 contains a simple estimation of  $a_{\ell}$ , which turns out to be close, al least, to twice the Pioneer acceleration, the value that would solve the riddle. Section 5 analyzes the dynamics of time, i. e. the relation between  $\tau$  and t, as the basis for the explanation of the Pioneer anomaly: the speed of light is constant, if defined with respect to the cosmological proper time  $\tau$ , which in turn

accelerates with respect to the coordinate parametric time. The conclusions will be stated in Section 6.

## 2 On the speed of light and the coordinate time

Einstein's papers of the period 1907-1912, often considered just as matter for historians, are however of great interest for physicists, in particular because of his discussions on the variation of the light speed in a gravitational field [9, 10]. Since he had not yet proposed his general relativity, with the idea of proper time, he was obviously talking of the light speed  $c(t) = d\ell/dt$  (called non-proper light speed in section 1.3). In the last section of a review paper in 1907 [11], he introduces his principle of equivalence in a static situation, deducing from it that the light speed c must depend on the gravitational potential  $\Phi(\mathbf{r})$ , as well as his first formula for the bending of a light ray grazing a star. According to Pais, "the study of Maxwell equations in accelerated frames had taught him that the [observed] velocity of light is no longer a universal constant in the presence of gravitational fields" [9].

In 1911 he takes anew the question in a paper entitled "On the influence of gravitation on the propagation of light" [12], where he uses again his principle of equivalence. After a discussion on the synchronization of clocks, he concludes that "if we call the velocity of light at the origin of coordinates  $c_0$ , where we take  $\Phi = 0$ , then the velocity of light at a place with gravitational potential  $\Phi$  will be given as

$$c = c_0 \left( 1 + \Phi/c^2 \right)$$
." (10)

He thus confirms, with a more detailed analysis, the conclusion of his 1907 paper, stating also that eq. (10) is a first order approximation (although he does not tell which one, it is obvious that he refers to situations near the Newtonian limit). It can be written a bit more explicitly as

$$c(\mathbf{r},t) = c_0 \left\{ 1 + \Phi(\mathbf{r},t)/c^2(\mathbf{r},t) - \Phi_R/c_0^2 \right\},$$
 (11)

where  $\Phi_{\rm R}$  is a reference potential, at present time in a terrestrial laboratory R, where the observed light speed is  $c_0$ , i. e. the constant that appears in the tables. It follows from equations (10)-(11) that  $c = d\ell/dt$  must depend on  $\mathbf{r}$  and t, so that the deeper (more negative) is the potential, the smaller is

the light speed with respect to t and conversely (according to this Einstein formula, the light speed at the surface of the Sun, would be about 2 ppm lower than here at Earth).

In two papers in 1912 [13], he considers the light speed as a field in spacetime  $c(\mathbf{r},t)$  and states "a clock runs faster the greater the c of the location to which we bring it", a statement quite similar to consequence (ii) at the introduction of this work that the acceleration of light must cause or be related to an acceleration of atomic clocks, to be discussed in section 5. In a reply to a critical paper on relativity by M. Abraham, Einstein states "the constancy of the velocity of light can be maintained only insofar as one restricts oneself to spatio-temporal regions with constant gravitational potential. This is where, in my opinion, the limit of the principle of the constancy of the velocity of light — thought not of the principle of relativity — and therewith the limit of the validity of our current theory of relativity, lies" [14]. What Einstein says here is that the principle of relativity is not the same thing as the principle of constancy of light speed: the latter must not be taken as a necessary consequence of the former. Furthermore, he insists that the light velocity can depend on  $\Phi$ , as was the case with his eq. (10).

Two comments are important here: (i) as stated before, these statements by Einstein refer clearly to the light speed defined with respect to the coordinate time  $c = d\ell/dt$ , as explained in section 1.3, and (ii) they are still valid as first order approximations in general relativity. As Einstein himself found later, the light speed is constant and invariant if defined with respect to proper time  $c^* = d\ell/d\tau$  (= constant), but at that time he was mainly concerned with the effect of a gravitational field on Maxwell equations, in which the time derivatives are certainly with respect to the coordinate time t. Let us consider now this question (I will follow now the well known textbook The Classical Theory of Fields by Landau and Lifshitz, reference [15], section 90).

The electromagnetic tensor is defined in general relativity by means of a vector field such that  $F_{\mu\nu} = A_{\nu;\mu} - A_{\mu;\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$ . The electromagnetic vectors **E**, **D** and antisymmetric tensors  $B_{ij}$ ,  $H_{ij}$  are defined as follows

$$E_i = F_{0i} \,, \quad B_{ij} = F_{ij} \,, \quad D^i = -\sqrt{g_{00}} \, F^{0i} \,, \quad H^{ij} = \sqrt{g_{00}} \, F^{ij} \,,$$

the vectors **B**, **H** being the dual to the three-tensors  $B_{ij}$  and  $H_{ij}$ , i. e.  $B^i = -e^{ijk}B_{jk}/(2\sqrt{\gamma})$ ,  $H_i = -\sqrt{\gamma}\,e_{ijk}H^{jk}/2$ , where  $\gamma = \det(\gamma_{ij})$ ,  $\gamma_{ij} = -g_{ij}$  being the three-dimensional metric tensor (assuming for simplicity that  $g_{0i} = 0$ ).

It follows that

$$\mathbf{D} = \mathbf{E}/\sqrt{g_{00}}, \quad \mathbf{B} = \mathbf{H}/\sqrt{g_{00}},$$
 (12)

(see [15]). If the space is empty, i. e. without free charges or currents, the Maxwell equations can be written as

$$\nabla \cdot \mathbf{B} = 0, \quad \nabla \times \mathbf{E} = -\frac{1}{\sqrt{\gamma}} \partial_t (\sqrt{\gamma} \mathbf{B}).$$
 (13)

$$\nabla \cdot \mathbf{D} = 0, \quad \nabla \times \mathbf{H} = \frac{1}{\sqrt{\gamma}} \partial_t \left( \sqrt{\gamma} \, \mathbf{D} \right).$$
 (14)

In a static situation, these four equations have exactly the same form as in special relativity, since the factors  $\sqrt{\gamma}$  cancel. However, eq. (12) implies that the relative permittivity  $\epsilon_{\rm r}$  and permeability  $\mu_{\rm r}$  of empty space are different from 1, their common value being  $\epsilon_{\rm r} = \mu_{\rm r} = (g_{00})^{-1/2}$ . This is due to the geometry of spacetime. Near the Newtonian limit one has  $g_{00} = 1 + \Phi(\mathbf{r})/c^2 - \Phi_{\rm R}/c_0^2$ , where  $\Phi_{\rm R}$  is the potential at a reference laboratory where  $c = c_0$ , so that the empty space is like an inhomogeneous optical medium with  $\epsilon_{\rm r}(\mathbf{r}) = \mu_{\rm r}(\mathbf{r}) = 1 - [\Phi(\mathbf{r})/c^2 - \Phi_{\rm R}/c_0^2]$ . Since  $c = c_0/\sqrt{\epsilon_{\rm r}\mu_{\rm r}}$ , the light speed with respect to t is being given by Einstein eq. (11) at first order.

In this work, we will be interested mainly in the case of a potential depending only on time  $\Phi(t)$ , as is the case for the background potential of all the universe in the approximation that all its matter and energy are uniformly distributed. As will be seen later, the time derivatives of the potential (over  $c^2$ ) will be extremely small, of the order of the Hubble constant  $H_0 = 2.3 \times 10^{-18} \text{ s}^{-1}$ . The same applies, therefore, to the time derivatives of  $\gamma$ . It is easy then to deduce from eqs. (13)-(14) that the electromagnetic vector fields obey classical wave equations with light speed given by Einstein's equation (11), for frequencies such that  $\omega \gg H_0$ , i. e. for any practical purpose.

As a last remark to end this section, note that Einstein formula (11) can be written as

$$c(\mathbf{r},t) = c^* \left[ 1 + \Phi(\mathbf{r},t)/c^2(\mathbf{r},t) \right], \qquad (15)$$

where  $c^* = c_0[1 - \Phi_R/c_0^2]$  is the light speed for zero gravitational potential. It is also the light seed if defined with respect to proper time, as will be discussed later.

It seems clear, therefore, that, according to (10)-(11), the light speed with respect to coordinate time must be a function  $c(\mathbf{r}, t)$ , with both a space and

a time variation. The first depends on the distribution of matter and energy. The second must be dominated by a secular progressive increase, since the potential  $\Phi$  produced by all the matter and energy of the universe must be an increasing function of time because of the universal expansion.

### 3 The background gravitational potential and the acceleration of the non-proper light speed.

Let us take the element of interval in weak gravity [16, 17, 18, 19]

$$ds^{2} = e^{2\Phi_{loc}(\mathbf{r},t)/c^{2}} c^{*2} dt^{2} - d\ell^{2} \approx (1 + 2\Phi_{loc}/c^{2}) c^{*2} dt^{2} - d\ell^{2}, \qquad (16)$$

from which

$$c(\mathbf{r},t) = c^* [1 + \Phi_{\text{loc}}/c^2], \qquad (17)$$

at first order (compare with (15),  $\Phi_{\text{loc}}(\mathbf{r})$  being here the (weak) gravitational potential of nearby bodies, those that produce a non negligible acceleration  $g(\mathbf{r})$  at the observation point (the Solar System and the Galaxy, for observers near Earth). One has thus for the light speed at a generic point P along a ray,  $c(P) = c(R)[1 + (\Phi_{\text{loc}}(P) - \Phi_{\text{loc}}(R))/c^2]$ , R being a reference point.

Note that (17) is the same as Einstein formula. It implies that the difference between the non-proper light speed and the  $\tau$ -light speed is an effect of the gravitational potential since they are equal if  $\Phi = 0$ . The local variations of  $\Phi$  due to the inhomogeneities of the distribution of matter give local and small variations of  $c(\mathbf{r},t)$ , while its cosmological variation causes a secular increase which is seen as the blueshift, observed as the Pioneer effect.

It will be important in the following to know what kind of time is the variable t in (16). It is certainly a coordinate time, but this is not enough for our purpose, in view of the freedom to make changes of the spacetime coordinates. Since this element of interval is good near the classical limit, this t is a Newtonian time. In fact, the increment of the interval of a particle world line between the points  $P_1$  and  $P_2$  is

$$\Delta s = \int_{P_1}^{P_2} ds = \int_{t_1}^{t_2} c \left( 1 + \frac{2\Phi_{\text{loc}}}{c^2} - \frac{v^2}{c^2} \right)^{1/2} dt,$$

which, at first order in the potential, is equal to

$$\Delta s = c(t_2 - t_1) - \frac{1}{c} \int_{t_1}^{t_2} \left( \frac{v^2}{2} - \Phi_{\text{loc}} \right) dt$$
 (18)

As is seen, the condition for  $\Delta s$  to be stationary is equivalent to the Hamiltonian principle for a particle in the potential  $\Phi_{loc}$ . This shows that, for weak gravity, the time t in (16) is a Newtonian parametric time.

But we are not only submitted to the potential nearby bodies. Quite on the contrary, there is a background gravitational potential due to all the matter and and energy in the visible universe, which is assumed here to be uniformly distributed over flat surfaces t = constant, so that the background gravitational potential is a function of time  $\Phi_{\text{all}}(t)$ . That potential must be uniform and depend on time because of the expansion. We must therefore rewrite the interval (16) as

$$ds^2 = c^{*2} d\tau^2 - d\ell^2$$
, with  $d\tau = [1 + \Psi(t)]dt$ , (19)

at first order, where the dimensionless potential  $\Psi(t) = \Phi_{\rm all}(t)/c^2(t)$  the background potential of all the matter and energy. From now on, the notation  $\Psi = \Phi/c^2$  will be used.

It follows that

$$c = c(t) = c^*[1 + \Psi(t)] = c_0[1 + \Psi(t) - \Psi(t_0)], \qquad (20)$$

where  $t_0$  is the age of the universe, *i.e.* the present time, and  $c_0 = c(t_0) = c^*[1 + \Psi(t_0)]$ , at first order, is the present time value of the non-proper light speed at a terrestrial reference laboratory (*i.e.* the value in the tables). Note that this equation gives the variation of c(t) near  $t_0$  at first order. Taking now the t derivative of (20) at time  $t_0$ , one has  $\dot{c}(t_0) = c(t_0)\dot{\Psi}(t_0)$ . The same argument can be applied to the expression for c(t) near any other fixed time  $\tilde{t}$ , what implies that  $c(t) = c(\tilde{t}) \exp[\Psi(t) - \Psi(\tilde{t})] \, \forall t$ . In particular, taking  $\tilde{t} = t_0$ , one finds

$$c(t) = c_0 e^{[\Psi(t) - \Psi(t_0)]}. (21)$$

The shape of the function c(t) defined by (21) does not change if the reference time  $(t_0 \text{ or } \tilde{t})$  is changed because  $c(t_1)e^{-\Psi(t_1)} = c(t_2)e^{-\Psi(t_2)}$ . The interval can be written then as

$$ds^{2} = e^{2[\Psi(t) - \Psi(t_{0})]} c_{0}^{2} dt^{2} - d\ell^{2} = c^{*2} d\tau^{2} - d\ell^{2}, \text{ with } d\tau = e^{\Psi(t)} dt.$$
 (22)

Note that  $\tau$  is a well defined cosmological proper time. We see here that  $\tau$  accelerates with respect to t, its second derivative being not nil. This will be considered later with more detail.

The interval (19) is correct near the Newtonian limit, even if  $\Psi(t)$  is not small, since  $\nabla \Psi = 0$ . Indeed, if such a potential is included in (18), the Lagrangian will increase in a function of time, which does not have any effect on the equations of motion of the particle. This fact will be used in section 4. In any case,  $\Psi$  being space independent, it can be absorbed in a redefinition of the time. As a last remark in this section, note that the gravitational potential  $\Phi$  has been used even in cases in which its gradient can be neglected. For instance, by Ahluwalia and coworkers to study quantum aspects of gravity, eventual violations of the equivalence principle or gravitationally induced neutrino-oscillation phases, see [20, 21].

To summarize this section, the background gravitational potential of all the universe induces an acceleration of proper time with respect to coordinate time. Since the  $\tau$ -light speed is constant, the non-proper light speed necessarily accelerates.

### 4 Estimation of the adiabatic acceleration of the non-proper speed of light

**4.1 Time variation of the non-proper speed of light.** The effect of the expansion on the potential at a spacetime point will be considered now, an estimate being made later in section 4.2 of the non-proper acceleration of light  $a_{\ell} = \dot{c}(t_0)$  (7). The potential of all the universe at the terrestrial laboratory R can be written, with good approximation, as  $\Phi_{\rm all} = \Phi_{\rm loc}(R) + \Phi_{\rm av}(t)$ . The first term  $\Phi_{loc}(R)$  is the part due to the local inhomogeneities, i. e. the nearby bodies (the Solar System and the Milky Way). It is constant in time since these objects are not expanding. The second  $\Phi_{av}(t)$  is the space averaged potential due to all the mass and energy in the universe (except for the nearby bodies), assuming that they are uniformly distributed. Contrary to the first, it depends on time because of the expansion. The former has a non vanishing gradient but is small, the latter is space independent, but time dependent and much larger. The value of  $\Phi_{\rm loc}/c_0^2$  at R is the sum of the effects of the Earth, the Sun and the Milky Way, which are about  $-7 \times 10^{-10}$ ,  $-10^{-8}$ and  $-6 \times 10^{-7}$ , respectively, certainly with much smaller absolute values than  $\Phi_{\rm av}(t_0)$ , which of the order of  $-10^{-1}$  as will be seen below.

Taking the time derivative of eq. (20) or (21), the (observed) light speed

near present time  $t_0$  is equal to

$$c(t) = c_0[1 + a_t(t - t_0)] = c_0 + a_\ell(t - t_0), \tag{23}$$

the quantity  $a_t$  and the light acceleration  $a_\ell$  being

$$a_{t} = \dot{\Psi}(t_{0}), \qquad a_{\ell} = a_{t}c_{0} = \dot{\Psi}(t_{0})c_{0},$$
 (24)

Since the background gravitational potential of all the universe  $\Phi_{\rm av}(t)$  is increasing because of the expansion (the galaxies are separating and their interaction potential increasing) eqs. (20)-(24) show that a time increase of the non-proper speed of light c(t) must be expected also. In this sense, there is an acceleration of light  $a_{\ell} = \dot{c}$  (see section 1.3). Indeed, the arguments leading to (23) are clear and compelling. Unfortunately, a rigorous calculation of the quantities  $a_{\ell}$  and of  $a_{\rm t}$ , which would take into account all the eventual effects, is not easy. However, a simple, approximate and phenomenological estimation will be performed now. It is claimed that it is sensible and meaningful in spite of its simplicity (it may be convenient to stress again that all this is compatible with the constancy of the  $\tau$ -light speed  $c^* = d\ell/d\tau$ .)

**4.2 Estimation of the non-proper accelerations of light and of the acceleration of the clocks.** The inverse time  $a_t$  was in fact introduced by Anderson *et al* in reference [1], as the "clock acceleration", through the relation " $a_P = a_t c$ ". It will be shown later in section 5 that it is indeed the acceleration of the atomic clocks.

In the following a simple crude estimate of the values of the non-proper acceleration of light (section 1.3)  $a_{\ell} = \dot{c}(t_0)$  and the clocks acceleration will be made, taking (20) as starting point. Although it involves approximations and simplifications, it shows the main ideas of the model and gives an convincing representation of the phenomenon. Let  $\Omega_M$ ,  $\Omega_{\Lambda}$  be the corresponding present time relative densities of matter (ordinary plus dark) and dark energy corresponding to the cosmological constant  $\Lambda$ . We take a universe with k=0,  $\Omega_M=0.27$ ,  $\Omega_{\Lambda}=0.73$  and Hubble parameter  $H_0=71~{\rm km\cdot s^{-1}\cdot Mpc^{-1}}=2.3\times 10^{-18}~{\rm s^{-1}}$ . In order to determine the average potential  $\Phi_{\rm av}(t)$ , let  $\Phi_0(t_0)$  be the gravitational potential produced by the critical mass density distributed up to the present radius of the visible universe  $R_{\rm U}(t_0)=c_0/H_0=4,200~{\rm Mpc}$ ; taking into account the kinetic energy of the galaxies as a source of gravity, one has  $\Phi_0(t_0)/c_0^2=-\int_0^{c_0/H_0}c_0^{-2}G\rho_{\rm cr}4\pi r{\rm d}r/\sqrt{1-(H_0r/c_0)^2}=4\pi G\rho_{\rm cr}/H_0^2=-0.75$ . It must be

emphasized that, although this value of the potential might seem to be too large for this approximation to apply, there is no problem in fact since it is space independent and its time derivative is extremely small, as explained at the end of section 3. Its effect will be to accelerate adiabatically the proper time with respect to the coordinate time, as will be seen later.

Because, in this model, the light speed was smaller in the past, the radius of the visible universe is a function of time  $R_{\rm U}(t)$  that can be written as  $R_{\rm U}(t) = R_{\rm U}(t_0) \left[ \int_0^t c(t) {\rm d}t \, / \int_0^{t_0} c(t) {\rm d}t \right]$ . It turns out then that  $\Phi_0(t) = 4\pi G \rho_{\rm cr} R_{\rm U}^2(t)/c_0^2$ . Consequently, one has  $\dot{\Phi}_0(t_0) = \Phi_0(t_0) \times (2c_0/R_U)$ . Note that  $\Phi_0(t) \to 0$  when  $t \to 0$ .

The present time average potential is then  $\Phi_{\rm av}(t_0) = \Phi_0(\Omega_M - 2\Omega_\Lambda)$ . Because of the expansion of the universe, the gravitational potentials due to matter and dark energy equivalent to the cosmological constant vary in time as the inverse of the scale factor R(t) and as its square  $R^2(t)$ , respectively (with  $R(t) = (\Omega_M/\Omega_\Lambda)^{1/3} \sinh^{2/3} \left[ (3\Lambda)^{1/2} t/2 \right]$  for this model universe). This implies that the average background gravitational potential is given as

$$\Phi_{\rm av}(t) = \Phi_0(t_0) \left[ \frac{\int_0^t c(t) dt}{\int_0^{t_0} c(t) dt} \right]^2 \left[ \frac{\Omega_M}{R(t)} - 2\Omega_\Lambda R^2(t) \right]. \tag{25}$$

From eqs. (20)-(25) and a bit of simple algebra, the inverse time  $a_{\rm t}$  can be expressed as

$$a_{\rm t} = \frac{[1 - 3\Omega_{\Lambda}]\dot{\Phi}_0/c_0^2 - [1 + 3\Omega_{\Lambda}]H_0\Phi_0/c_0^2}{1 + 2(1 - 3\Omega_{\Lambda})\Phi_0/c_0^2} = H_0 \frac{(1 - 9\Omega_{\Lambda})\Phi_0/c_0^2}{1 + 2(1 - 3\Omega_{\Lambda})\Phi_0/c_0^2}.$$

Introducing in this equation the values of  $\Omega_M$ ,  $\Omega_{\Lambda}$ ,  $\Phi_0$  and  $\dot{\Phi}_0$ , the clock acceleration  $a_t$  and the light acceleration  $a_{\ell}$  are shown to take the values

$$a_{\rm t} \simeq 1.5 \, H_0 \,, \quad a_{\ell} = a_{\rm t} c_0 \simeq 10.4 \times 10^{-10} \,\,\mathrm{m/s}^2 \,.$$
 (26)

Since the anomaly would be explained if  $a_{\rm P} = a_{\ell}/2$  and the observed value is  $a_{\rm P} = (8.74 \pm 1.33) \times 10^{-10} {\rm m/s}^2$ , the predicted Pioneer acceleration only about 40 % off the observation. This is encouraging for such a simple estimate, but note that the main purpose of this work is less to get a precise fit of  $a_{\rm P}$  than to build an understanding of the phenomenon. It could not be otherwise, given the simplicity of the calculation. What matters is that the model predicts the existence of an adiabatic non-proper acceleration of light,

i.e. an increase of the non-proper light speed  $c=\mathrm{d}\ell/\mathrm{d}t$  implying a blue shift with a value close, at least, to the one observed in the Pioneer 10 and the spaceships. This is what would give an explanation of the riddle.

It must be underlined that such an acceleration of light is a simple consequence of Einstein formula (11), taking into account the expansion of the universe, since the potential  $\Phi_{\rm av}$  is increasing because of the expansion, so that c(t) must increase also (see section 1.3). This increase, in turn, produces a blue shift as shown in section 1.4, with the same observational signature as an extra attraction from the Sun. It could be argued, however, that the gravitational redshift would cancel the increase in c since the frequency decreases when photons climb up along an increasing potential. However, the well known expression for the gravitational redshift  $\Delta \nu/\nu_0 = -\Delta \Phi/c_0^2$  is not valid here, since it assumes static situations which is not the case in an expanding universe. To end this section, it can be said that Rosales has proposed an explanation for the Pioneer anomaly which, although different to the one considered here since he argues that it is a manifestation of the Berry phase, coincides in attributing the effect to the expansion of the universe [22].

## 5 The acceleration of cosmological time with respect to coordinate time

The main result of this paper is the predicted value of the Pioneer acceleration in section 4.2. This was made locally in time, in terms of some data at present time  $t_0$ , as the background gravitational potential of all the universe and its derivative with respect to the coordinate time t, assuming a near Newtonian situation. This section contains the development of some ideas based in a cosmological and global consideration of time.

As explained in section 2, Einstein asserted in 1912 that clocks run faster the higher is c (page 104 of the first paper of [13]; remember, his definition of the light speed was  $c(t) = d\ell/dt$ ). This means that the higher is c the larger is the quotient  $\Delta \tau/\Delta t$ , i.e. an interval of proper time over the corresponding interval of coordinate time. He was thinking then in clocks at different space points, but the same can be said clearly about clocks at different times: if he had known the universal expansion (unsuspected however at that time) he could have added "and, besides, clocks run faster as time goes on". The (cautious) claim of this and previous work is that the Pioneer acceleration is

an effect of the acceleration of light with respect to the coordinate time t (see section 1.3). These two statements are equivalent. Indeed, if the observed blue shift is the consequence of the acceleration of light, the basic units of time (i. e. the periods of electromagnetic waves) would be decreasing (see section 1.4), the atomic clocks going faster, and conversely. Otherwise stated, the Pioneer effect could be a manifestation of the acceleration of time, as measured by atomic clocks. In other words, of the acceleration of proper time with respect to coordinate time.

In (20) he proper time  $\tau$  was shown to be given as

$$d\tau = \sqrt{g_{00}} dt = e^{\Psi(t)} dt, \quad \tau(t) = \int_{t_i}^t e^{\Psi(t)} dt,$$
 (27)

 $t_i$  being an arbitrary initial time. As stated before, the time  $\tau$  accelerates with respect to t, the acceleration being obviously equal to

$$\frac{\mathrm{d}^2 \tau}{\mathrm{d}t^2} = \dot{\Psi}(t)e^{\Psi(t)} = a_t e^{\Psi(t)} \,, \tag{28}$$

because of the universal expansion.

However, a warning is necessary: although  $\tau$  is well defined and is the cosmological proper time (assuming a uniform distribution of mass and energy), it is not the time used in measurements with atomic clocks at Earth. Indeed, since the frequencies increase at the same rate as the light speed, the basic units of these clocks vary on time as 1/c(t). This means that, near present time  $t_0$ , the time of the atomic clocks, say  $\tau_{\rm at}(t;t_0)$ , verifies

$$d\tau_{at}(t;t_0) = e^{[\Psi(t) - \Psi(t_0)]} dt, \quad \frac{d^2 \tau_{at}(t;t_0)}{dt^2} \Big|_{t_0} = \dot{\Psi}(t_0) = a_t.$$
 (29)

As is seen,  $a_t$  is really the acceleration of the atomic clocks, as already stated by Anderson et al [1]. The speed of light, if measured or calculated with respect to this time, is constant and equal to

$$c[\tau_{\rm at}(t;t_0)] = \frac{\mathrm{d}\ell}{\mathrm{d}\tau_{\rm at}(t;t_0)} = c_0, \quad \text{while} \quad c(\tau) = \frac{\mathrm{d}\ell}{\mathrm{d}\tau} = c^*. \tag{30}$$

In other words, the measurements of the light speed with such atomic clocks gives the value of c(t) not of  $c^*$ . However,  $\tau_{\rm at}(t;t_0)$  is not a good universal time since its definition depends on  $t_0$  (the observation time); nevertheless

it is good near  $t_0$ . It verifies  $d\tau_{\rm at}(t;t_0)=e^{-\Psi(t_0)}d\tau(t)$ , and  $d^2\tau_{\rm at}/d\tau^2=0$ . This means that these two times,  $\tau_{\rm at}$  and  $\tau$ , do not accelerate with respect to one another, their time intervals being just proportional, the proportionality constant being  $e^{-\Psi(t_0)}$ . Furthermore  $d\tau_{\rm at}(t_0;t_0)=dt$ , what means that, at exactly the time  $t_0$ , the corresponding intervals of the times  $\tau_{\rm at}$  and t are equal. This clarifies the meaning of  $\tau_{\rm at}(t;t_0)$ : it is a redefinition of the cosmological proper time (by means of a multiplicative factor depending on  $t_0$ ), to ensure that the basic units of  $\tau_{\rm at}$  and t are equal at  $t_0$  (i. e. now).

Otherwise stated, the introduction of the factor  $e^{-\Psi(t_0)}$  makes sure that the atomic clocks which are used tick at time  $t_0$  at the same rate (if more exactly) as the mechanical classical clocks that measure parametric time t. This explains why a measure with the clocks with time  $\tau_{\rm at}(t;t_0)$  gives the value  $c(t_0)$ , not  $c^*$ . Since they are made to tick at the same rate as the coordinate time (as the barycenter dynamic time of the solar system, for instance) they give the same value for shorts time intervals near  $t_0$ . On the other hand a true measurement with atomic clocks would employ clocks that ticked at the same rate as the mechanical clocks at some initial time in the past without redefining their intervals, so that their "seconds", would be different now. The result of such a measurement would be the constant proper speed of light  $c^*$ .

Let us summarize which are the values of the light speed near present time  $t_0$ , as defined or measured with respect to the three times t,  $\tau$  and  $\tau_{\rm at}(t;t_0)$ , eqs (23) and (30),

$$c(t) = c_0[1 + a_t(t - t_0)], c(\tau) = c^*, c[\tau_{at}(t; t_0)] = c_0.$$
 (31)

As is seen, if c is defined: (i) with respect to the coordinate parametric time t, it increases linearly in t; (ii) with respect to the proper cosmological time  $\tau(t)$ , it is constant and equal to  $c^*$ ; and (iii) with respect to the time of the atomic clocks synchronized with t at time  $t_0$ ,  $\tau_{\rm at}(t;t_0)$ , it is constant and equal to  $c_0$ .

Assume, now, that we change  $t_0$  to  $t_0 + \Delta t_0$ ,  $\Delta t_0$  ( $\ll t_0$ ) being the time difference between two measurements of the light speed with atomic clocks (e. g. 1 year or 20 years). Taking into account that  $\Psi(t_0 + \Delta t_0) = \Psi(t_0) + a_t \Delta t_0$  and instead of (30), we will have for the light speed, at first order, if measured with respect to time  $\tau_{\rm at}(t; t_0 + \Delta t_0)$ ,

$$c[\tau_{\rm at}(t;t_0+\Delta t_0)] = \frac{\mathrm{d}\ell}{\mathrm{d}\tau_{\rm at}(t;t_0+\Delta t_0)} = \frac{1}{[1+\Psi(t)-\Psi(t_0)][1-a_t\Delta t_0]} \frac{\mathrm{d}\ell}{\mathrm{d}t}$$

$$= \frac{\mathrm{d}\ell}{\mathrm{d}t} (1 + a_{t} \Delta t_{0}) = c_{0} (1 + a_{t} \Delta t_{0}), \qquad (32)$$

instead of (30). This means that, in a measurement of the light speed with atomic clocks at times  $t_0$  and  $t_0 + \Delta t_0$  (if synchronized with coordinate time at the time of the measurement), one would find an acceleration of light with respect to t equal to  $a_{\ell} = a_t c_0$ , as in (24). This is the conclusion of this model: that the Pioneer anomaly is an effect of the dynamics of time [23, 24].

5.1 A toy model. This paper is about what happens now or in an interval around present time  $t_0$ , its main result being the prediction of an adiabatic acceleration of light  $a_{\ell}$  with respect to the coordinate time, eqs. (??)-(23), which would have the same observational signature as the Pioneer effect. However, this needs some clarification. What was shown is that the observed light speed is constant if measured with respect to the cosmological proper time  $\tau$ , while  $\tau$  accelerates with respect to the coordinate time t. As a consequence, light accelerates with respect to coordinate time t, so that  $\dot{c}(t) = a_{\ell} > 0$ . Moreover, a very simple approximate estimation gives for  $a_{\ell}$  a value just a 40% smaller than twice the Pioneer acceleration, the value that would explain exactly the effect (see section 1.3). Given the simplicity of the calculation, this seems encouraging.

It is tempting to explore as well the model towards the past, as a method to understand better the problem. The result of this exploration will be called now "a toy model" to emphasize that it is not taken necessarily as a rigorous theory. In spite of that, it may be useful to understand better the main ideas involved, from an intuitive point of view. To do that it suffices to take  $t_i = 0$  in (27), so that

$$\tau(t) = \int_0^t e^{\Psi(t)} dt, \qquad c = c_0 e^{[\Psi(t) - \Psi(t_0)]} = c^* e^{\Psi(t)}, \qquad (33)$$

with  $\Psi(t) = \Phi_{\rm av}(t)/c^2(t)$ ,  $\Phi_{\rm av}(t)$  being given in (25). Let us play with this toy model, in order to build an intuitive picture of the phenomenon. It must be underlined that, if the light speed were smaller in the past, it can be admitted that the universe is also smaller than what is now admitted. As an approximation, we will define the present value of the potential, introduced in section 4.2 as

$$\Phi_0(t_0)/c_0^2 = -\int_0^{R_{\rm U}} c_0^{-2} G \rho_{\rm cr} 4\pi r {\rm d}r / \sqrt{1 - (r/R_{\rm U})^2},$$

and change the value of  $R_{\rm U}$ . The corresponding Hubble parameter is  $H_0 = c_0/R_{\rm U}$ .

The most interesting results are summarized in two figures (the calculations are only approximate, no attempt having been made to refine the precision of the numerical values). In Figure 1, the cosmological proper time  $\tau$  (33) for  $R_{\rm U}(t_0)=3,000$  Mpc and  $R_{\rm U}(t_0)=3,400$  Mpc and the coordinate time t are plotted versus the coordinate time in units of the age of the universe. The line of the coordinate t is obviously a straight line. On the other hand, the acceleration of  $\tau$  with respect to t is seen in the curvature of the solid lines. Indeed, the other two curves verify  $d^2\tau/dt^2 < 0$  for  $t < t_{\rm c} \simeq 0.34\,t_0$ , and  $d^2\tau/dt^2 > 0$  for  $t > t_{\rm c}$ . Before  $t_{\rm c}$ ,  $\tau$  decelerates with respect to t, but it accelerates afterwards. The reason for that behavior is that the negative potential due to matter is dominant at the beginning, while near present time the positive potential due to the cosmological constant is more important (with the assumed value of  $\Omega_{\Lambda}$ ). The latter begins to take over the former at time  $t_{\rm c}$  because the potential  $\Psi(t)$  had a minimum then.

It happens that  $\tau(t_0) \simeq 1.20\,t_0$  (resp.  $1.27\,t_0$ ) for 3,000 Mpc (resp. 3,400 Mpc). In other words, the age of the universe, as measured by  $\tau$ , would be about  $1.20\,t_0$  (resp.  $1.27\,t_0$ ), while it is just  $t_0$  in terms of t. This would be due to the adiabatic acceleration  $a_t e^{\Psi(t)}$  of  $\tau$  with respect to t. Note that the difference between  $\tau$  and t was small until recent times.

In Figure 2, the non-proper light speed (33) is plotted against t for  $R_{\rm U}(t_0)=3,000~{\rm Mpc}$  (upper line) and  $R_{\rm U}(t_0)=3,400~{\rm Mpc}$  (lower line). As is shown, it decreases from t=0 until  $t\simeq 0.34\,t_0$ , where it has a minimum, increasing thereafter. It does not change much, however, before  $t\simeq 0.5\,t_0$ . Note that their values for t=0 turns out to be  $c^*\simeq 0.42\,c_0$  and  $c^*\simeq 0.31\,c_0$ , respectively. Also that (i) at  $t=\tau=0$ , both definitions of light speed give the same value  $c(t=0)=c(\tau=0)=c^*$ , and (ii) c(t) increases for t>0, while  $c(\tau)$  remains constant, so that  $c>c(\tau)$ , their difference increasing after the beginning.

It is seen in both figures that at about  $t = 0.35 t_0$  both the cosmological proper time  $\tau$  and the non-proper speed of light c begin to accelerate with respect to t, this effect being more clear after about  $t = 0.5 t_0$ . This is the transit from the slowing-down to the speeding-up, due to the effect of the dark energy (represented here by the cosmological constant).

There is a singularity at about  $R_{\rm U} \simeq 3,400$  Mpc. When that value is approached from below, the minimum of c(t) at  $t \simeq 0.35 t_0$  becomes more accused, the second derivative  ${\rm d}^2 c/{\rm d} t^2$  increasing, until the equation (33) fails




Figure 1: Cosmological proper time  $\tau$  (33) in the toy model (solid lines) and coordinate parametric time t (dashed line) versus parametric time, in units of the age of the universe  $t_0$ . The lower (resp. upper) solid line at  $t = t_0$  corresponds to  $R_{\rm U}(t_0) = 3,000$  Mpc (resp.  $R_{\rm U}(t_0) = 3,400$  Mpc)(explanation in the text).

to have a real solution for c around that value of t. This happens approximately when the Schwarzschild radius of the total mass of a distribution with the critical density inside a sphere of radius  $R_{\rm U}$  coincides with  $R_{\rm U}$ , although this is perhaps just a numerical coincidence.

#### 6 Summary and conclusions

In spite of its simplifications and approximations, the model presented here gives a promising explanation of the Pioneer anomaly, as a consequence of the interplay between the cosmological and proper time  $\tau$  and the coordinate and parametric time t. All through this work, it is assumed that (i) all the matter and energy in the universe are uniformly distributed and (ii) the near Newtonian approximation is acceptable. The idea of background gravitational potential of all the universe  $\Phi_{\rm av}(t)$  is important here. The conclusions of this paper are the following:

1. The light speed can be defined in two ways: (i) with respect to the




Figure 2: Non-proper light speed (33) in the toy model, in units of  $c_0$ , as function of the coordinate parametric time, in units of the age of the universe  $t_0$  (explanation in the text).

proper time  $\tau$ ,  $c^* = d\ell/d\tau$ , and (ii) with respect to the coordinate time t,  $c(t) = d\ell/dt$  (see section 1.3). With the first definition it is a universal constant  $c^*$ , with the second it is a function  $c(\mathbf{r},t)$ . They have been called here "proper speed of light" and "non-proper speed of light", respectively. Assuming a uniform distribution for all the matter and energy of the universe, the second definition gives a time dependent non-proper speed of light c(t), the variation of which is dominated by a secular adiabatic increase, due to the progressive augmentation of the background gravitational potential  $\Phi_{\rm av}(t)$  of all the universe as the galaxies separate. In this sense it is sensible to speak of an acceleration of light, even if the proper speed of light is constant. The present value of that acceleration would be  $a_{\ell} = \dot{c}(t_0) = a_t c_0$ ,  $a_t$  being the coordinate time derivative of the background potential of all the universe  $\Psi(t) = \Phi_{\rm av}(t)/c^2(t)$  and  $c_0$  the value of the light speed in the tables. A simple estimate predicts the value  $a_{\ell} = 10.4 \times 10^{-10} \text{ m/s}^2$ , what would mean an increase of c(t) of about 3.3 cm/s per year. Note, however, that this model is not a theory with variable light speed, since the proper speed of light is constant.

2. Such an adiabatic acceleration has the same observational signature, an extra blue shift, as an acceleration towards the Sun of a radio source such as the Pioneer. The agreement would be quantitatively exact if  $a_{\ell} \simeq 2a_{\rm P}$ , i. e. if the light acceleration were equal to twice the Pioneer acceleration (section 1.5). However,  $a_{\rm P}$  would be quite unrelated to any unmodelled motion, although it could be easily interpreted as an anomalous acceleration, even if the ship were following the exact trajectory predicted by the current theory of gravitation.

3. The phenomenon here reported is due to the relation between the cosmological proper time  $\tau$  and the coordinate time t. That relation is given as  $\tau = \int_0^t e^{\Psi(t)} \mathrm{d}t$ , so that so that  $\mathrm{d}^2\tau/\mathrm{d}t^2 = \dot{\Psi}\,e^{\Psi}$  (overdot means derivative with respect to t) (section 3 and 4). In other words,  $\tau$  accelerates with respect to t, since  $\mathrm{d}^2\tau/\mathrm{d}t^2 = \dot{\Psi}(t)e^{\Psi(t)} > 0$ . Consequently, while the light has constant speed with respect to  $\tau$ , it accelerates with respect to t. As all this indicates, the Pioneer phenomenon is a very interesting case of the dynamics of time [23, 24].

An important aspect of the problem refers to the detection of the blueshift. The time used in the Solar System is the barycenter dynamical time, which is originally a Newtonian parametric time, even if it is measured with atomic clocks in order to increase the precision. The blueshift can indeed be measured, as was the case, by using detectors with circuits calibrated in a macroscopic way. This is an indication in favor of this work. In any case, there is an interesting metrological problem.

To summarize: the conclusion of this paper is that the anomalous acceleration of the Pioneer 10/11 and the other two spacecrafts could be only apparent, not real, just an effect of the dynamics of time (i. e. the relation between he cosmological proper time and the coordinate parametric time) that shows up in the acceleration of light, if defined as the t derivative of  $c = d\ell/dt$  (the proper light speed  $c^* = d\ell/d\tau$  is constant as required in general relativity). This would cause the observed blue shift, which would be quite unrelated, however, to the motion of the space ships. Indeed, they would have followed the standard trajectories, as predicted by current gravitation theory. All this must be studied by the experts who know the details of the motion of the spacecrafts and of the metrological procedures involved in the observation. This should be done, since some of the consequences of the interplay between the two times t and  $\tau$  could be unexpected and surprising at the cosmological level.

Acknowledgements. I am indebted to A. Tiemblo for enlightening discussions on the dynamics of time, to J. L. Sebastián for information on microwaves and to G. Álvarez Galindo, C. Aroca, A. I. Gómez de Castro, E.

López and J. L. Rosales for discussions or encouragement.

#### References

- J. D. Anderson, Ph. A. Laing, E. L. Lau, A. S. Liu, M. Martin Nieto and S. G. Turyshev, Phys. Rev. Lett. 81, 2858 (1998).
- [2] J. D. Anderson, Ph. A. Laing, E. L. Lau, A. S. Liu, M. Martin Nieto and S. G. Turyshev, Phys. Rev. D 65, 082004 (2002).
- [3] J. D. Anderson, E. L. Lau, S. G. Turyshev, Ph. A. Laing and M. Martin Nieto, Mod. Phys. Lett. A 17, 875 (2002).
- [4] J. P. Mbelek and M. Michalski, Int. J. Mod. Phys. D 13, 865 (2004).
- [5] A. F. Rañada, astro-phys/0202224, Europhys. Lett. **61**, 174 (2003).
- [6] A. F. Rañada, gr-qc/0211052, Europhys. Lett. **63**, 653 (2003)
- [7] A. F. Rañada, Int J. Mod. Phys. D 12, 1755 (2003).
- [8] G. F. R. Ellis and J. Ph. Uzan, gr-qc/0305099 v2.
- [9] A. Pais, Subtle is the Lord. The science and the life of Albert Einstein (Oxford University Press, Oxford, 1984).
- [10] J. M. Sánchez Ron, El origen y desarrollo de la relatividad (Alianza Editorial, Madrid, 1985).
- [11] A. Einstein, Jahrbuch Radioaktivität und Electronik 4, 411 (1907); reprinted in *The collected papers of Albert Einstein*, English translation, Vol 2 (Princeton University Press, 1989), p. 252.
- [12] A. Einstein, Annalen der Physik 35, 898 (1911); reprinted in *The collected papers of Albert Einstein*, English translation, Vol 3 (Princeton University Press, 1993), p. 379.
- [13] A. Einstein, Annalen der Physik **38**, 355 (1912); **38**, 443 (1912); reprinted in *The collected papers of Albert Einstein*, English translation, Vol 4 (Princeton University Press, 1996), pp. 95 and 107.

- [14] A. Einstein, Ann. Physik 38, 1059 (1912); reprinted in *The collected papers of Albert Einstein* (1996), English translation, Vol 4 (Princeton University Press, Princeton, 1996), p. 130.
- [15] L. D. Landau and E. M. Lifshitz, *The Classical Theory of Fields*, 4th revised English edition (Pergamon Press, Oxford, 1975), chapter 10.
- [16] C. W. Misner, K. S. Thorne and J. A. Wheeler, *Gravitation* (Freeman, San Francisco, 1973).
- [17] M. V. Berry, *Principles of Cosmology and gravitation* (Adam Hler-Institute of Physics Publishing, Bristol, 1989).
- [18] C. M. Will, Theory and experiment in gravitational physics (Cambridge University Press, Cambridge, 1993), Rev. ed.
- [19] W. Rindler, *Relativity. Special, general and cosmological* (Oxford University Press, Oxford, 2001).
- [20] D. V. Ahluwalia, Gen. Rel. Grav. 29, 1491 (1997).
- [21] G. Z. Adunas, E. Rodríguez-Milla and D. V. Ahluwalia, Gen. Rel. Grav. 33, 183 (2001).
- [22] J. L. Rosales, gr-qc/0401014.
- [23] A. Tiemblo and R. Tresguerres, Gen. Rel. Grav. **34**, 31 (2002).
- [24] J. F. Barbero, A. Tiemblo and R. Tresguerres, Phys. Rev. D 57, 6104 (1998).