## HAMILTONIAN ISOTOPIES OF RELATIVELY EXACT LAGRANGIANS ARE ORIENTATION-PRESERVING

## JACK SMITH

ABSTRACT. Given a closed, orientable Lagrangian submanifold L in a symplectic manifold  $(X, \omega)$ , we show that if L is relatively exact then any Hamiltonian diffeomorphism preserving L setwise must preserve its orientation. In contrast to previous results in this direction, there are no spin hypotheses on L. Curiously, the proof uses only mod-2 coefficients in its singular and Floer cohomology rings.

Let  $(X, \omega)$  be a symplectic manifold which is compact or well-behaved (e.g. convex and cylindrical) at infinity. Recall that an isotopy  $\varphi_t$  of X is *Hamiltonian* if is generated by a time-dependent smooth function  $H_t : X \to \mathbb{R}$ , in the sense that the generating vector field  $V_t = \varphi_t^* \dot{\varphi}_t$  satisfies

$$\omega(V_t, -) = -\mathrm{d}H_t$$

The group  $\operatorname{Ham}(X)$  comprises the end-points  $\varphi_1$  of Hamiltonian isotopies  $\varphi_t$  starting from  $\varphi_0 = \operatorname{id}_X$ , and is an important subgroup of the symplectomorphism group  $\operatorname{Symp}(X)$ .

Let  $L \subset X$  be a closed, orientable Lagrangian submanifold, and define

$$\operatorname{Ham}(X, L) = \{ \varphi \in \operatorname{Ham}(X) : \varphi(L) = L \}.$$

Assume throughout that *L* is relatively exact, meaning that  $\omega$  vanishes on  $\pi_2(X, L)$ .

The purpose of this short note is to prove the following result.

**Theorem.** The action of Ham(X, L) on L is orientation-preserving.

**Example.** If L is homotopy equivalent to  $\mathbb{RP}^{2n+1}$  then  $\operatorname{Ham}(X, L)$  acts trivially on  $\operatorname{H}_*(L; \mathbb{Z})$ .

**Remark.** The *x*-axis in  $\mathbb{R}^2$ , with its standard symplectic form, can be rotated through angle  $\pi$  by a Hamiltonian isotopy, so compactness of *L* is essential in the theorem. Similarly, the equator in  $S^2$  can be turned over by a Hamiltonian isotopy, so relative exactness is also essential.

The action of Ham(X, L) on *L* induces a homomorphism

 $\operatorname{Ham}(X, L) \to \operatorname{GL}(\operatorname{H}_*(L; \mathbb{Z})),$ 

whose image is the (*Hamiltonian*) homological Lagrangian monodromy group  $\mathcal{H}_{L}^{\mathbb{Z}}$  of L. The theorem can be interpreted as saying that  $\mathcal{H}_{L}^{\mathbb{Z}}$  fixes the fundamental class [L], whilst in the example  $\mathcal{H}_{L}^{\mathbb{Z}}$  is trivial.

The group  $\mathcal{H}_{L}^{\mathbb{Z}}$  was introduced by M.-L. Yau in [4] to study the monotone Clifford and Chekanov tori in  $\mathbb{C}^2$ . It was extensively studied in the weakly exact setting, along with its analogue  $\mathcal{H}_{L}^{\mathbb{Z}/2}$  in GL(H<sub>\*</sub>( $L; \mathbb{Z}/2)$ ), by Hu–Lalonde–Leclercq in [1]. They showed that  $\mathcal{H}_{L}^{\mathbb{Z}/2}$  is trivial in this case, and that  $\mathcal{H}_{L}^{\mathbb{Z}}$  is also trivial if L is (relatively) spin and one restricts to those  $\varphi \in \text{Ham}(X, L)$  preserving the (relative) spin structure. Porcelli [2] subsequently gave a new proof of these results under weaker hypotheses, which extends to generalised homology theories. In particular, he showed that  $\mathcal{H}_{L}^{\mathbb{Z}}$  is trivial if L is spin, with no restriction on  $\varphi$ . Compared with these works, our result is therefore notable because of its lack of spin hypotheses on both L and  $\varphi$ . Note that the  $\mathbb{RP}^{2n+1}$  example is not spin for any even  $n \geq 2$ .

*Proof of the theorem.* We work throughout with  $\mathbb{Z}/2$  coefficients. First we recall some Floer theory. Suppose  $L_1$  and  $L_2$  are closed, orientable Lagrangian submanifolds in X for which the Floer cohomology  $HF(L_1, L_2)$  can be defined. This holds, for example, if they are relatively exact and Hamiltonian isotopic to each other. For each  $L_i$  a choice of orientation corresponds to a  $\mathbb{Z}/2$ -grading in the sense of [3], with respect to the canonical  $\mathbb{Z}/2$ -grading of X. Orienting both  $L_1$  and  $L_2$  then gives a  $\mathbb{Z}/2$ -grading on  $HF^*(L_1, L_2)$ . Reversing the orientation of  $L_1$  corresponds to shifting its grading, which we denote by  $L_1[1]$ , and there is then a natural identification  $HF^*(L_1[1], L_2) = HF^{*-1}(L_1, L_2)$ . Similarly for  $L_2$ , with  $HF^*(L_1, L_2[1]) = HF^{*+1}(L_1, L_2) = HF^{*-1}(L_1, L_2)$ . Given a third closed, oriented Lagrangian  $L_3$  for which the Floer cohomology with  $L_1$  and  $L_2$  can be defined, there is a Floer product

$$\mu^2$$
: HF\*( $L_2, L_3$ )  $\otimes$  HF\*( $L_1, L_2$ )  $\rightarrow$  HF\*( $L_1, L_3$ ).

This respects the  $\mathbb{Z}/2$ -gradings, and is associative in the obvious sense if we also consider a fourth Lagrangian. Each HF<sup>\*</sup>( $L_i$ ,  $L_i$ ) then becomes a  $\mathbb{Z}/2$ -graded ring, and this ring is unital. It is well-known that if  $L_i$  is relatively exact then the ring HF<sup>\*</sup>( $L_i$ ,  $L_i$ ) is given by H<sup>\*</sup>( $L_i$ ) with the grading collapsed modulo 2.

Now focus on our single closed, relatively exact Lagrangian *L*, on which we fix an orientation. A compactly-supported Hamiltonian isotopy  $\varphi_t$  induces a *continuation element* 

$$c(\varphi_t) \in \mathrm{HF}^0(\varphi_0(L), \varphi_1(L)).$$

This is functorial, in the sense that Floer products of continuation elements correspond to concatenations of Hamiltonian isotopies, and if  $\varphi_t$  is a constant isotopy then  $c(\varphi_t)$  is the unit. In particular this means that continuation elements are invertible with respect to the Floer product.

Suppose for contradiction that there exists a  $\varphi \in \text{Ham}(X, L)$  which reverses the orientation of L, and fix a Hamiltonian isotopy  $\varphi_t$  from  $\text{id}_X$  to  $\varphi$ . By cutting off the generating function  $H_t$  outside a compact set containing the sweepout  $\bigcup_t \varphi_t(L)$  of L, we may assume that the isotopy is compactly supported. It therefore has an associated continuation element  $c(\varphi_t) \in \text{HF}^0(L, \varphi(L))$ .

Since  $\varphi(L)$  is L with its orientation reversed, i.e.  $\varphi(L) = L[1]$ , we can view  $c(\varphi_t)$  as an element of  $HF^1(L, L)$ . By the above discussion, this element is invertible, and  $HF^*(L, L)$  is  $H^*(L)$  with grading collapsed mod 2, so we conclude that there exists an invertible element of odd degree in  $H^*(L)$ . But this is impossible: any odd-degree element lies in  $H^{>0}(L)$ , which is a proper ideal in  $H^*(L)$ , so it cannot be invertible. We obtain the desired contradiction, proving the theorem.

**Acknowledgements.** I am grateful to Noah Porcelli for useful correspondence and an anonymous referee for helpful comments.

## References

[1] S. HU, F. LALONDE, AND R. LECLERCQ, Homological Lagrangian monodromy, Geom. Topol., 15 (2011), pp. 1617–1650.

[2] N. PORCELLI, Families of relatively exact Lagrangians, free loop spaces and generalised homology, 2022.

[3] P. SEIDEL, Graded Lagrangian submanifolds, Bull. Soc. Math. France, 128 (2000), pp. 103–149.

[4] M.-L. YAU, Monodromy and isotopy of monotone Lagrangian tori, Math. Res. Lett., 16 (2009), pp. 531-541.

ST JOHN'S COLLEGE, CAMBRIDGE, CB2 1TP, UNITED KINGDOM

Email address: j.smith@dpmms.cam.ac.uk