Non-perturbative Quantum Gravity in Fock representations

T. Thiemann^{1*}

¹ Inst. for Quantum Gravity, FAU Erlangen – Nürnberg, Staudtstr. 7, 91058 Erlangen, Germany

May 3, 2024

Abstract

Perturbative quantum gravity starts from prescribing a background metric. That background metric is then used in order to carry out two separate steps: 1. One splits the non-perturbative metric into background and deviation from it (graviton) and expands the action in terms of the graviton which results in an ifinite series of unknown radius of convergence. 2. One constructs a Fock representation for the graviton and performs perturbative graviton quantum field theory on the fixed background as dictated by the perturbative action. The result is a non-renormalisable theory without predictive power.

It is therefore widely believed that a non-perturbative approach is mandatory in order to construct a fundamental, not only effective, predictive quantum field theory of the gravitational interaction. Since perturbation theory is by definition background dependent, the notions of background dependence (BD) and perturbation theory (PT) are often considered as symbiotic, as if they imply each other.

In the present work we point out that there is no such symbiosis, these two notions are in fact logically independent. In particular, one *can* use BD structures while while not using PT at all. Specifically, we construct BD Fock representations (step 2 above) for the full, *non-perturbative* metric rather than the graviton (*not* step 1 above) and therefore never perform a perturbative expansion. Despite the fact that the gravitational Lagrangean is a non-polynomial, not even analytic, function of the metric we show that e.g. the Hamiltonian constraint with any density weight can be defined as a quadratic form with dense form domain in such a representation.

1 Introduction

Today we have not yet constructed an interacting matter quantum field theory (QFT) in four dimensional Minkowski space that obeys the Wightman axioms [1]. Still, elementary particle physics has been spectacularly successful in using perturbation theory based on Fock representations for the free (i.e. non-interacting) QFT which can be constructed in any dimension. The chosen Fock representation is subordinate to a split of the non-perturbative Hamiltonian $H = H_0 + V$, which is a polynomial function of the fields, into a free piece H_0 which is a quadratic polynomial and the interaction V which is a polynomial of higher degree. While by construction H_0 is an operator on the free Fock space \mathcal{H}_0 with dense invariant domain \mathcal{D}_0 given by the span of Fock states, by Haag's theorem [2] the interaction term V is typically not an operator but only a quadratic form which is at the heart of the problem of giving meaning to powers of H as they appear in the would be unitary time evolution operator e^{itH} or the would be Moeller operator $e^{itH} e^{-itH_0}$ which is employed in the associated scattering matrix S. To define it as an operator would lead to a Wightman QFT and would require a representation subordinate to H and not to H_0 on a Hilbert space \mathcal{H} .

If one tries to construct the S-matrix nevertheless by Taylor expanding with respect to V as a quadratic form, one meets divergences that come from products of quadratic forms which is an ill-defined mathematical operation.

^{*}thomas.thiemann@gravity.fau.de

These divergences can be tamed using regulariation and removed by absorbing them into the monomial coefficients of the Hamiltonian. If this can be done order by order for a *finite* number of monomials one calls the theory perturbatively renormalisable. If this requires an infinite number of monomials one calls the theory perturbatively non-renormalisable. In the non-renormalisable case one calls the theory truncated to a finite number of monomials effective if one can argue that the neglected terms do not play any role up to a given energy resolution. Even in the renormalisable case there is no guarantee that the perturbation series converges in any sense. Importantly, one can do this also in curved spacetime (CST) i.e. for a general Lorentzian signature metric, not necessarily the Minkowski metric [3].

In quantum gravity one has tried to apply this programme to the gravitational field. A first difference with the matter field sector is that while the natural "background matter" field is the "zero" field, there is no such natural background for the metric. Therefore in a first step one splits the full metric g into a fixed background g_0 and a graviton deviation $h = g - g_0$. Then one expands the gravitational Lagrangean in powers of h at $g = g_0$. Since the gravitational Lagrangean is not polynomial, not even analytic, in g, the expansion is an infinite series with an unknown radius of convergence even classically. In a second step one collects the terms quadratic in hinto a free part (the analog of H_0) and the interacting rest (the analog of V). This must be accompanied by fixing the coordinate gauge freedom. One can then use the free part to construct a Fock representation for hadapted to it and q_0 . The rest of the programme then proceeds formally the same way as for matter QFT in the CST encoded by q_0 . Unfortunately the resulting theory is perturbatively non-renormalisable [4] in the following sense: the perturbative expansion of a Lagrangian in terms of h delivers n-th order monomial interaction terms L_n . One can now ask whether the counter terms triggered by L_n are found in the list $\{L_m\}_{m=0}^{\infty}$. If this is true for all $n \leq N$ and if this leaves no more than a finite number of free parameters then the theory would be renormalisable. However for a given L (say the Einstein-Hilbert Lagrangian) one finds even for relatively low orders n counter terms L'_m which belong to a different Lagrangian L' i.e. are not in the allowed list. This means that one needs to generalise the Lagrangian introducing more couplings and it is believed that this process never ends and results in a theory with infinitely many free parameters.

It is therefore widely believed that a fundamental QFT of gravity must be constructed using a non-perturbative approach (NPA). Examples for such programmes are, in alphabetical order, asymptotically safe quantum gravity (ASQG) [6], causal dynamical triangulations (CDT) [7], causal set theory (CST) [8], loop quantum gravity (LQG) [9] and string theory (ST) in the AdS/CFT incarnation [10]. See also [11] for a confrontation of these approaches in a single volume.

Since the perturbative approach to QG as sketched above uses a background metric g_0 as a starting point and since it has failed to construct a fundamental theory, many quantum geometers take background independence (BI) as a profound, almost holistic, guiding principle to construct a fundamental theory of QG. This would indeed be a necessary step, if the notions of background dependence (BD) and perturbative approach (PA) would be truly symbiotic, i.e. if indeed BD would imply a PA and vice versa. However, we would like to point our that *there is no such symbiosis*. Indeed, while one cannot A. perform perturbation theory without B. using a background metric, one can B. use a background metric without A. performing perturbation theory. Thus in terms of logical implication we have $A \Rightarrow B$ but $B \not\Rightarrow A$. That is, we may use a backround metric g_0 to perform various QFT constructions but without using g_0 to split the non-perturbative metric g into $g = g_0 + h$. Without such a split, the Lagrangean is not expanded in powers of of h and there is *no perturbation series at the classical level*.

In this paper we will use a background metric g_0 to construct a Fock representation, but not of the perturbation field $h = g - g_0$ but of the non-perturbative field g. In fact, even in perturbative QG one could have used a Fock representation for h based on a background metric g_1 which could be different from $g_0 = g - h$ although there it would be an unnecessary, even disadvantageous (because the quadratic part of the Hamiltonian is an operator rather than a quadratic form only if $g_1 = g_0$) blow-up of background structure. As a consequence the algebraic state (vacuum expectation value) underlying this Fock space structure is in particular not diffeomorphism invariant. This means that not a single Fock vector state, not even the vacuum (i.e. the vector annihilated by the annihilation operators) is diffeomorphism invariant. This appears to be in rather strong conflict in representing the gauge transformations of theory on the corresponding Hilbert space in the operator constraint approach and various operators such the Weyl operators that define the Fock representation and constraint operators that encode the quantum Einstein equations. However we remind of the phenomenon of symmetry breaking [12]: It is perfectly possible that an operator H is invariant under a unitary symmetry U while none of its ground states is (in that case the ground state is necessarily degenerate). With respect to the diffeomorphism group and the generators of diffeomorphisms (diffeomorphism constraints) it is perfectly possible that the generators are diffeomorphism co-variant while no state is invariant.

Such a background dependent representation is of any practical use only if one succeeds to formulate the quantum Einstein equations in mathematically rigorous way. In LQG one has gone rather far with a manifestly background independent representation of (a version tailored to non-abelian gauge theories of) the Weyl operators [13]: There is a unique such representation of Narnhofer-Thirring type [14] based on an inavariant state ω on the corresponding Weyl algebra [15]. The spatial diffeomorphism group consequently acts by unitary operators. However, one must pay a high price for this: The state ω is irregular, one cannot construct the spatial metric and its conjugate momentum simultaneously as operators. Similarly one cannot construct the generators of the spatial diffeomorphism group as operators. One can get away with this because one can at least obtain a representation of the spatial diffeomorphism group. However, the generators of the classical Einstein equations do not form a Poisson Lie algebra and thus Lie group techniques cannot be used to construct the exponentiated temporal generator, called the Hamiltonian constraint. One has to construct it directly in non-exponentiated form which requires regularisation and operator ordering choices and removal of the regulator based on spatial diffeomorphism invariance [16]. However, not all regularisation choices are washed out when the regulator is removed and leaves the Hamiltonian constraint with quantisation ambiguities. Furthermore, while the algebra of constraint operators closes with structure operators ordered to the right, these structure operators are not quantisations of the classical structure functions. Thus while there is no mathematcial anomaly, there is a physical one. These complications can be overcome in the Abelian truncation of vacuum gravity [17] but for a different Weyl algebra and with a different dense invariant domain of the constraint operators.

To avoid this anomaly in LQG, one can replace the Hamiltonian constraint by the spatially diffeomorphism invariant master constraint [18] or one can perform a reduced phase space quantisation [19] or one can try to define the constraints on a linear dual space of distributions without Hilbert space structure [20]. However, another annoying consequence of the irregularity of the state which penetrates to all three of these proposals is the fact the resulting Hilbert space is not separable. This again leads to quantisation ambiguities. It is these ambiguities and their removal that have motivated Hamiltonian renormalisation methods [21] and also the present work.

Fock representations are based on regular pure states on the Weyl algebra and thus we can define the spatial metric and its conjugate momentum as operator valued distributions. Moreover, the associated representation of the Weyl algebra is irreducible, the underlying vacuum expectation value state is pure, every vector is cyclic for it. As we will show, in a non-perturbative Fock representation one cannot easily define the constraints as *operators*, but one can define them as quadratic forms on a suitable dense form domain. There are no quantisation ambiguities, one *must* choose normal ordering. Moreover, although quadratic forms cannot be multiplied, in a recent contribution [31] we showed that for the generators of the spatial diffeomorphisms (spatial diffeomorphism constraints) one can define a *commutator* and check that it is free of physical anomalies. In this work we will show that given any representation, not necessaily in the Fock class, quadratic forms are sufficient to rigorously formulate and solve the Lorentzian or Euclidian quantum Einstein equations. Moreover, we will show that the commutator algebra can be meaningfully defined at least for Euclidian signature in background dependent Fock representations when the constraints are written in polynomial form. For the Lorentzian signature case this might also be possible by exactly the same technique but the computations, while straightforward, are much more complex and have not been finished yet.

As this works for a huge class of Fock representations one may ask which one to choose and whether this does not correspond to yet another source of ambiguity. However, note first of all that this choice of background structure B_0 consisting of a choice of background metric g_0 and a Fock state ω_0 based on it must also be made in the perturbative approach, this "background ambiguity" is in addition to the ambiguities that one encounters in the renormalisation process which requires to add an infinite number of counter terms corresponding to all possible curvature invariants. In the non-perturbative approach, if it can be accomplished, the counter term ambiguity is missing, which is an improvement. Next note that the background structure ambiguity B_0 is of the same quality as the representation ambiguity that one encounters in QFT in CST. The difference with QFT in CST is that there one neglects quantum gravity altogether and the background $g = g_0$ is considered as fixed. But even then there are infinitely many choices and one must downsize them by imposing additional physical requirements such

as the Hadamard condition on the 2-point function [22] which states that the short distance behaviour should be the same as that of the usual Fock representations of Minkowski space singled out by Poincaré invariance. In quantum gravity one can use either the operator constraint or the reduced phase space approach. In the operator constraint approach the choice of representation is likely not very important because what one is interested in is not the states of that representation but rather the solutions of the constraint equations which are distributions over a dense subspace in the representation space and that space of distributions could be independent of the choice of g_0 . Of course the space of solutions must be equipped with a Hilbert space structure by itself which opens a new representation problem, namely that of the gauge invariant observables. In a reduced phase space approach where one considers only representations of the observables from the outset, the choice of representation constructed from some g_0 enters only at that stage which is comparable to the problem of choice of representation in QFT in CST with the difference that now also quantum geometry fields have to be considered. One can now in principle again use the Hadamard condition to select suitable B_0 but with an additional input: The domain of the operators in question should be chosen in such a way that the quantum geometry fluctuations around g_0 are small. Such semiclassical, namely coherent, states do exist within Fock representations. This at least ties B_0 to the domain of the operators. Apart from that one needs further physical input in order to downsize the ambiguity B_0 , for instance one could ask that a maximal number of quadratic forms that enter the dynamics in fact turn into operators which is similar in nature to the Hadamard condition that ensures that the stress energy tensor is a well defined quadratic form. Finally we remind of Fell's theorem [23] which says that in principle one can approximate all states from a given one in the sense of matrix elements of an arbitrary but finite number of observables to arbitrary precision. Unfortunately the proof of that theorem is not constructive and therefore of little practical use.

We note that this background dependent but non-perturbative approach is much closer to the formalism of QFT in CST, which is believed to be framework to choose in the semiclassical limit of quantum gravity (vanishing geometry fluctuations), than when dealing with the background independent representation employed in LQG. For suitable matter the physical Hamiltonian of the reduced phase space formulation is spatially diffeomorphism invariant and one can promote the uniquness theorem [15] to the physical Hilbert space level. However, that physical Hilbert space is again non-separable and the representation is not regular with respect to either the metric or conjugate momentum, the physical Hamiltonian operator thus again suffers from quantisation ambiguities. In the Fock representation the physical Hilbert space is separable, the quantisation ambiguities are removed to a large extent but the physical Hamiltonian is now only a quadratic form, similar as in interacting QFT, and no longer an operator on the Fock space. Hence there are complementary advantages and disadvantages of Fock versus LQG representations. What makes the Fock representation attractive is that it is computationally much simpler and equips us with a huge aresenal or techniques.

Thus we see that the viability of Fock representations in quantum gravity depends on whether one is able to formulate either of i. the quantum constraints, ii. the master constraint, iii. the physical Hamiltonian as a densely defined quadratic form on the Fock space. Now for the spatial diffeomorphism constraints (SDC) this is easy to check they are quadratic polynomials in the fields. For the Hamiltonian constraint (HC) this is easy to check if one uses it in polynomial form which is possible by multiplying it by a sufficiently large power of the square root of the spatial metric upon which it becomes a polynomial as well. One can show that the SDC commutator can be defined in terms of quadratic forms and closes without anomaly [31]. For the HC this is less clear because the SDC and HC do not form a Lie algebra. If we replace the HC by a master constraint which is invariant under spatial diffeomorphisms then we do obtain a Lie algebra and the corresponding commutator algebra of quadratic forms has a chance to close. However, in order to be spatially diffeomorphism invariant, the master constraint must not be a polynomial! Rather it is a polynomial multiplied by inverse powers of the determinant of the spatial metric. Nevertheless it turns out that one can construct a densely defined quadratic form from the master constraint using a new technique which we present, to the best knowledge of the author, for the first time in this work. Finally, the physical Hamiltonian is obtained by solving the constraints for a canonical momentum conjugate to a scalar field which serves as a physical clock and thus is naturally a density of weight one and thus also not a polynomial. Therefore the same remarks as for the master constraint apply with additional complications originating from the fact that solving for the momenta involves choosing a square root. We present a partial solution to this problem which again defines a quadartic form.

The architecture of this work is as follows:

In section 2 we introduce the classical framework and define a class of Fock representations for quantum gravity both in the constraint quantisation and reduced phase space approach. We focus mostly on the geometry sector.

In section 3 we define the SDC and HC (in polynomial form) as quadratic forms on the Fock space. To define these, one makes use of mode structures on the one particle Hilbert space, mode cut-offs and limiting patterns for the cut-off removal. One exploits the freedom in the choice of such structures to tame the commutator of quadratic forms and take limits. We show that the hypersurface deformation algebra closes in this quadratic form sense without anomaly for Euclidean signature vacuum GR written in real valued self-dual connection variable in four spacetime dimensions. The only reason to restrict to Euclidian signature and four dimensions is that the real valued self-dual variables require that restriction but have the advantage of being polynomials of degree four only. The same technique can be applied in other dimensions and for Lorentzian signature but the computations are much harder as we encounter polynomials of degree ten or higher and have not been completed yet.

In section 4 we present the new technique to define any real power of the determinant of the spatial metric *at a point* as a densely defined quadratic form. The form domain must take into account that the quantum metric be non-degenerate if negative powers are involved. The same technique can be used to define geometric observables such as lengths, areas and volumes and the master constraint as quadratic forms in Fock representations. As for the physical Hamiltonian, which involves a square root, one can show that classically it has upper and lower bounds that do not involve a square root. One can then quantise these upper and lower bounds as quadratic forms on the Fock space which may serve as an approximation of the square root expression. We also mention some ideas of how to turn the quadratic forms into actual operators without using perturbative counter term methods.

In section 5 we summarise and conclude.

2 Classical Hamiltonian formulation

Our description will be brief, more details can be found in [9, 11].

2.1 Constraint formulation

A foliation of the spacetime D + 1 manifold $M \cong \mathbb{R} \times \sigma$ with a fixed D manifold σ into is encoded by lapse $N^0 := N$ and shift functions N^a , a = 1..D which define the timelike unit normal $n^\mu = N^{-1}[\delta^\mu_0 - N^a \delta^\mu_n]$, $\mu = 0,..,D$ of the t = const. hypersurfaces in the coordinates $x^0 = t \in \mathbb{R}$ and x^a on σ . One has the relation $g_{00} = -N^2 + q_{ab}N^aN^b, g_{0a} = q_{ab} N^b, g_{ab} = q_{ab}$ between N^μ and the components of the spacetime metric tensor $g_{\mu\nu}$. Let ω^A_μ , A = 1,..,d be a connection in a principal fibre bundle (we will shortly specialise it to the frame rotation bundle) with d-dimensional gauge group. The canonical Hamiltonian density is a linear combination of constraints C_μ , $\mu = 0,..,D$ and G_A , A = 1..d

$$H_{\rm can} = v^{\mu} P_{\mu} + v^A P_A + N^{\mu} C_{\mu} - \omega_0^A G_A \tag{2.1}$$

called secondary Hamiltonian constraint (HC) C_0 , spatial diffeomorphism constraint (SDC) C_a and Gauss constraint G_A respectively and of the primary constraints P_{μ} , P^A of momenta conjugate to N^{μ} , ω_0^A which are multiplied by the velocities $v^{\mu} = \dot{N}^{\mu}$, $v^A = \dot{\omega}^A$ that one cannot solve for in the Legendre transform of the Lagrangian. All constraints are first class.

The secondary constraints contain contributions from both geometry and matter. As nature contains fermions one is forced to consider D + 1-Bein fields e^I_{μ} , I = 0, .., D which constitute the spacetime metric $g_{\mu\nu} = \eta_{IJ}e^I_{\mu}e^J_{\nu}$ where η is the Minkowski metric. This implies in particular Gauss constraints corresponding to the D + 1 frame Lorentz group SO(1,D). One must remove the D boost generators among those Gauss constraints by imposing the time gauge $e^{\mu}_0 = n^{\mu}$ leaving D(D+1)/2 - D = D(D-1)/2 constraints for the frame rotation group SO(D) as the boost generators generate further secondary constraints on e^I_{μ} with respect to which they are second class. We will therefore understand that only those rotation generators are included in (2.1). Then the gravitational sector is entirely described by N^{μ} and the D-Bein e_a^j , j = 1, ..., D corresponding to $D + 1 + D^2 = (D + 1)^2 - D$ fields, together with the momenta P_{μ}, P_j^a conjugate to them and we have $q_{ab} = \delta_{jk} e_a^j e_b^k$. Since the constraints C_{μ}, G_A do not depend on N^{μ}, ω_0^A the primary constraints in fact have vanishing Poisson brackets with the secondary constraints. It follows that N^{μ}, ω_0^A are pure gauge and we can reduce the phase space by fixing $N^{\mu} = f^{\mu}, -\omega_0^A = f^A$ to be arbitrary Lagrange multiplier functions. Then the stability of that gauge enforces $v^{\mu} = v^A = 0$. Thus (2.1) simplifies to a linear combination of secondary constraints

$$H_{\rm can} = f^{\mu} C_{\mu} + f^A G_A \tag{2.2}$$

The geometry contributions to the constraints reads explicitly

$$C_{0} = [\det(q)]^{-1/2} [q_{ac} q_{bd} - [D-1]^{-1} q_{ab} q_{cd}] p^{ab} p^{bc} - [\det(q)]^{1/2} (R[q] + \Lambda)$$

$$C_{a} = P_{j}^{b} [e_{j}^{b}]_{,a} - [P_{j}^{b} e_{a}^{j}]_{,b}$$

$$G_{jk} = 2 P_{[j}^{b} \delta_{k]l}; e_{b}^{l}$$
(2.3)

where Λ is a cosmological constant and

$$p^{ab} := \frac{1}{2} P_j^{(a} e_k^{b)} \delta^{jk}, \ e_j^a \ e_a^k := \delta_j^k$$
(2.4)

R[q] is the Ricci scalar of q. As G_{jk} generates frame rotations, the quantities q_{ab} , P^{ab} are rotation invariant. Thus G_{jk} has vanishing Poisson brackets with C_0 but not with C_a . It is therefore convenient to decompose (indices j, k, l.. are moved with δ_{jk}, δ^{jk})

$$P_j^a = G_{jk} \ e^{ak} + 2 \ p^{ab} e_{bj} \tag{2.5}$$

and to rewrite ${\cal C}_a$ in manifestly Gauss invariant form

$$\hat{C}_a = C_a - [G_{jk}e^{bk}e^j_{b,a} - (G_{jk}e^{bk}e^j_a)_{,b}], = P^{bc} q_{bc,a} - 2(P^{bc} q_{ca})_{,b}$$
(2.6)

The non-vanishing Poisson brackets are (we set Newton's constant equal to unity)

$$\{P_j^a(x), e_b^k(y)\} = \delta(x, y) \ \delta_b^a \ \delta_j^k \tag{2.7}$$

In D = 3 it is possible to reformulate the theory in terms of SU(2) connections [25] by realising that

$$E_j^a := [\det(q)]^{1/2} e_j^a, \ A_a^j := \Gamma_a^j - [\det(q)]^{-1/2} q_{ab} P^{bj}$$
(2.8)

defines a canonical transformation $(P, e) \mapsto (E, A)$ where

$$e_{a,b}^{j} - \Gamma_{ba}^{c} e_{c}^{j} + \epsilon_{jlk} \Gamma[e]_{b}^{k} e_{a}^{l} := 0$$
(2.9)

defines the spin connection Γ_a^j of e from the Levi-Civita connection Γ_{ab}^c of q. In higher dimensions one can also arrive at a connection formulation [26] but one has to work with the hybrid connection of SO(D+1) and impose additional simplicity constraints. The non-vanishing Poisson brackets are then

$$\{E_j^a(x), A_b^k(y)\} = \delta(x, y) \ \delta_b^a \ \delta_j^k \tag{2.10}$$

In these variables the constraints are given in terms of the curvature $F = 2(dA + A \wedge A)$ of A by

$$\hat{C}_{0} = |\det(E)|^{-1/2} F_{ab}^{j} \epsilon_{jkl} E_{k}^{a} E_{l}^{b} - 2 |\det(E)|^{1/2} R[q]$$

$$\hat{C}_{a} = F_{ab}^{j} E_{j}^{b}$$

$$G_{j} = E_{j,a}^{a} + \epsilon_{jkl} A_{a}^{k} E^{bl}$$
(2.11)

where $C_0 - \hat{C}_0$ differ by terms proportional to G_j and $q_{ab} = |\det(E)|E_a^j E_{bj}, \ E_j^a E_a^k = \delta_j^k$.

The HC C_0 or \hat{C}_0 as produced by the Legendre transform of the Lagrangian carries a natural density weight of unity however then it is evidently not polynomial in the fields P_j^a, e_a^j or P^{ab}, q_{ab} or E_j^a, A_a^j . For purposes of Fock quantisation it is convenient to consider its polynomial version. This is obtained as follows: In the P_j^a, e_a^j formulation we have $[\det(q)]^{1/2} = |\det(e)|$. The Ricci scalar is given explicitly by

$$R = q^{ac} \,\delta^{c}_{d} \,R_{abc} \,^{d} = q^{ac} \,\delta^{c}_{d} \,(-2\partial_{[a}\Gamma^{d}_{b]c} + 2\Gamma^{e}_{c[a} \,\Gamma^{d}_{b]e}),$$

$$\Gamma^{d}_{ab} = q^{dc} \,\Gamma_{cab}, \, 2\Gamma_{cab} = 2q_{c(a,b)} - q_{ab,c}$$

$$q^{ab} = \det(q)^{-1} \,[(D-1)!]^{-1} \,\epsilon^{ac_{1..c}} e^{bd_{1..d}} - q_{c_{1}d_{1}} \cdot q_{c_{D-1}d} d_{D-1}$$

$$\Gamma^{d}_{ab,e} = q^{dc}_{,e} \Gamma_{cab} + q^{dc} \Gamma_{cab,e} = -q^{dg} \,q_{gh,e} \,q^{hc} \Gamma_{cab} + q^{dc} \Gamma_{cab,e}$$
(2.12)

It follows that $\det(q)^2 \Gamma_{ca}^e \Gamma_{be}^d$ is a monomial of degree 2D in 2D-2 factors of q and two factors of ∂q while $\det(q)^2 \Gamma_{bc,a}^d$ is a monomial of degree 2D consisting of two terms, the first term consisting of 2D-2 factors of q and two factors of ∂q while the second term consists of (2D-1) factors of q and one factor of $\partial^2 q$. Due to the contraction of the Ricci tensor with the inverse metric we require one more factor of $\det(q)$. It follows that

$$\hat{C}_0 := \det(q)^{5/2} C_0 \tag{2.13}$$

is a polynomial in the (q, p) polarisation where the "kinetic term" containing two facors of p^{ab} is of degree 2(D+1) in q_{ab} while the Ricci term has degree 2D + D - 1 = 3D - 1 in q_{ab} . For D = 3 its top degree is ten.

For the (e, P) polarisation we make use of the fact that the Ricci tensor can be written as $R_{abjk} e^{aj} e^{bk}$ and $R_{jk} = 2 [d\Gamma + \Gamma \wedge \Gamma]_{jk}$ where $\det(e) \Gamma_{jk}$ is a homogeneous polynomial of degree D with one factor of ∂e and D-1 factors of e. Since $|\det(e)| = \sqrt{\det(q)}$, by the same reasoning it is now sufficient to set

$$\hat{C}_0 := \det(q)^{3/2} C_0 \tag{2.14}$$

whose kinetic term containing two factors of P_j^a has 2(D+1) factors of e while the Ricci term has 2D+2(D-1) = 4D-2 factors of e. Its top degree is also ten in D = 3 (in fact it is homogeneous).

Finally in the (A, E) polarisation which is only available in D = 3 we have $R = R_{abjk}E^{aj}E^{bk}\det(q)^{-1}$ and again note that $\det(E)\Gamma_{jk}$ is a homogeneous polynomial of degree 3 with one factor of ∂E and two factors of E. Since $\det(E) = \det(q)$ we must use (2.13) again which contains eight facors of E in the F term and is a polynomial of degree 6 + 2 = 8 in the curvature term. Its top degree is also ten.

To summarise in all three polariasations we can generate a polynomial form of the Hamiltonian constraint with minimal top degree ten in D = 3 but it has density weight six in the (q, p) and (A, E) polarisation while it has density weight four in the (e, P) polarisation.

2.2 Master constraint formulation

Consider the smeared constraints

$$K(f, u, m) := \int_{\sigma} d^{D}x \left[f C_{0} + u^{a} C_{a} + m^{A} G_{A} \right]$$
(2.15)

They obey the following Poisson bracket relations

$$\{K(f, u, m), K(g, v, n)\} = -K(u[g] - v[f], [u, v] + q^{-1}(f \, dg - g \, df), [m, n] + u[n] - v[m])$$
(2.16)

where [u, v] is the commutator of vector fields, [m, n] the commutator of matrices, df the exterior derivative of a function and u[f] = df[u] the contraction of a vector field with the exterior derivative. The r.h.s. of (2.16) is a linear combination of smeared constraints, but the smearing functions are no longer constant functions on the phase space because of the appearance of q^{ab} . Thus the algebra of smeared constraints is no Lie algebra, also not in the polynomial version.

This is often argued to cause trouble in the constraint quantisation using Fock spaces due to the following reason: As we will see, the polynomial constraints can be quantised as densely defined quadratic forms in a Fock representation and this is sufficient to state the quantum constraint equations and to solve them. This requires

to normal order the polynomial constraints. Furthermore, we show that there exists a mathematically meaningful procedure to define the commutator of quadratic forms even though the product of quadratic forms is ill-defined. The result of the commutator calculation, if it exists, is then again a quadratic form which is necessarily normal ordered. However, the normal ordering of an expression of the form, say in D = 3,

$$\{\hat{C}_0(f), \hat{C}_0(g)\} = -\int d^D x \, \det(q)^3 q^{ab} (fg_{,b} - f_{,b}g) \, C_a \tag{2.17}$$

is not be given by

$$-\int d^{D}x : C_{a} :: \det(q)^{3}q^{ab} : (fg_{,b} - f_{,b}g)$$
(2.18)

Thus, if l is a linear functional on the form domain \mathcal{D} of the constraints such that $l[C_a(u^a)\psi) = 0$ for all vector fields u and all $\psi \in \mathcal{D}$ then $l([C(u), C(v)]\psi) \neq 0$ even if [C(u), C(v)] is given by the normal ordering of (2.17), i.e. even if there is no anomaly in the constraint algebra. This, in contrast to the situation in which the constraints are promoted to operators with dense invariant domain \mathcal{D} is, however, not a contradiction because $C(v)\psi$ is no longer in \mathcal{D} so that $l(C(u)C(v)\psi) \neq 0$ (in fact this expression diverges). This will be explained in more detail in the next section.

We can avoid these complications using the master constraint method. It comes in a minimal and a maximal version. In the minimal version it is only applied to C_0

$$M := \int d^D x \ C_0 \ \det(q)^{-1/2} C_0 \tag{2.19}$$

To see that the single M classically encodes the same information as the infinite number of $C_0(f)$ we note that M = 0 iff $C_0(f) = 0$ for all f thanks to the positivity of the integrand of (2.19). Next for a function F on the phase space we have $\{F, \{F, M\}\}_{M=0} = 0$ iff $\{C(f), F\}_{M=0} = 0$ for all f i.e. M defines the same weak Dirac observables. Since C_0 is a density of weight one it follows that $\{G(m), M\} = \{C_a(u^a), M\} = 0$ and in the quantum theory one is reduced to check the algebra of the $G(M), C_a(u^a)$ for anomaly freeness and that they commute with M, the commutator [M, M] = 0 is now trivial. However, the price to pay is that M in contrast to $G(m), C_a(u^a)$ is not polynomial and thus it is unclear how to define M as a quadratic form on the Fock space. Surprisingly, this is indeed possible as we will see in later sections.

Now instead of defining a solution by $l(\hat{C}_0(f)\psi = 0$ for all f, ψ one instead imposes $l(M\psi) = 0$ for all ψ where M is normal ordered. Normal ordering keeps symmetry of a quadratic form but destroys positivity. Note however that in an ordering such that C_0 is a symmetric operator we would formally have

$$\langle \psi, M\psi \rangle = \int d^D x ||([\det(q)]^{-1/4}C_0)(x)\psi||^2$$
 (2.20)

which is formally solved by those ψ with $\hat{C}_0(f)\psi = 0$ for all f. Thus "modulo ordering corrections" the condition $l(M\psi) = 0$ appears to be reasonable.

In the maximal version we combine *all constraints* into a single master constraint. As then we do not need to check any algebra, we can use the constraints in polynomial form. That is one would set e.g.

$$M = \int d^{D}x \left[\det(q) \ q^{ab} \ C_{a} \ C_{b} + \det(q)^{5} \ C_{0}^{2} + \delta^{AB} \ G_{A}G_{B} \right]$$
(2.21)

and normal order.

2.3 Reduced phase space formulation

In this case one removes the constraints classically using gauge fixing. This has the usual global reservations such as Gribov copies. It is in general true that constraining and quantisation do not commute thus we consider this as an alternative route which is only classically locally equivalent to the constraint method.

As for the non-gravitational Gauss constraints we can use one of the standard gauges such as axial gauge or Coulomb gauge. For the gravitational Gauss constraint we can use the upper triangular gauge on the D-bein [27]. That is, one can solve the relation

$$q_{ab} = e_a^j \delta_{jk} e_b^k \tag{2.22}$$

uniquely and algebraically for $e_a^j = u_a^j$, $u_a^j = 0$, a > j; $[u_j^a]_{j=a} > 0$. Here j = a refers to a choice of coordinate system in which we label coordinates by a = 1, ..., D as well as the internal indices j = 1, ..., D and one can state sufficient conditions under which this is compatible with the atlas of σ . We consider $q_{ab} = u_a^j u_{bj}$ as a function of of u_a^j and solve the Gauss constraints $G_{jk} = 0$ algebraically for P_j^a , a > j. The reduced phase is then coordinatised by $Q_j^a := P_j^a, u_a^j, a \le j$. Equivalently we invert $q_{ab} = q_{ab}[u]$ for $u_a^j = u_a^j[q]$ and pull back the symplectic structure

$$\Theta = \int d^D x \, \sum_{a \le j} Q_j^a \, [\delta u_a^j] = \int P^{ab} \, [\delta q_{ab}]$$
(2.23)

where $P^{ab} = \frac{1}{2} P_j^{(a} e^{b)j}$ at $e_a^j = u_a^j$, $G_{jk} = 0$ (if the fermionic current is not vanishing one must add the fermionic contribution to Θ to see this [27]).

Thus one may choose to work either with q_{ab} , P^{ab} or u_a^j , Q_j^a . Next one reduces the constraints C_{μ} . This may be done in various ways depending on the matter content. To have a concrete example in mind we consider the case of D + 1 minimally coupled, massless scalar fields ϕ^I , I = 0, ..., D with conjugate momentum π_I which contribute

$$\pi_I \phi_{,a}^I, \quad \frac{1}{2} \left[\delta^{IJ} \pi_I \pi_J [\det(q)]^{-1/2} + \delta_{IJ} \phi_{,a}^I \phi_{,b}^J [\det(q)]^{1/2} \right]$$
(2.24)

to C_a, C_0 respectively. We fix the coordinate gauge freedom

$$G^{I} = \phi^{I} - L^{-1} \delta^{I}_{\mu} x^{\mu} = 0, \ \phi^{I}_{*} = L^{-1} x^{I}$$
(2.25)

where we have introduced a length scale L for dimensional reasons (ϕ^I is dimension free while coordinates have length dimension). We then solve the constraints for π_I which yields

$$\pi_a^* = -L C_a', \ \pi_0^* = \sqrt{-2[\det(q)]^{1/2}\tilde{C}}, \ \tilde{C} = C' + \frac{1}{2}(L^2 \ \delta^{ab}C_a'C_b'/\sqrt{\det(q)} \ + L^{-2}q^{ab}\delta_{ab}\sqrt{\det(q)})$$
(2.26)

where C', C'_a are HC and SDC without the scalar contribution and we have chosen the positive root for π_0^* . The stability of the gauge condition under gauge transformations

$$G_{,0}^{I} + \{C_{\mu}(f^{\mu}), G^{I}\} = -L^{-1}\delta_{0}^{I} + L^{-1} f^{a}\delta_{a}^{I} + f^{0} [\det(q)]^{-1/2}\pi_{I} = 0$$
(2.27)

gives at $\phi = \phi_*, \pi = \pi^*$ with $Q = \sqrt{\det(q)}$

$$f_*^0 = L^{-1} Q [\pi_0^*]^{-1}, \ f_*^a = -L \ f_*^0 \ \pi_a * \ Q^{-1} = -\pi_a^* / \pi_0^*$$
(2.28)

The physical Hamiltonian is defined for a phase space function F independent of the gauge degrees of freedom ϕ, π by

$$\{H, F\} = \{C_{\mu}(f^{\mu}), F\}_{\phi - \phi_* = \pi - \pi^* = f - f_* = 0}$$
(2.29)

It drives the dynamics of the "true degrees of freedom" i.e. the non-scalar fields and working out (2.29) we find that it is explicitly given by

$$H = -L^{-1} \int d^D x \, \pi_0^* \tag{2.30}$$

In quantising the reduced phase space we only quantise the non-gauge fixed matter fields and the gravitational degrees of freedom q_{ab} , P^{ab} and we are no longer interested in the constraints but seek to implement the Hamiltonian H at least as a quadratic form. To do this, the square root in (2.25) is problematic because we can quantise only integrals over $\tilde{C}^n Q^m$ with $Q := \det(q)^{1/2}$ and $n \in \mathbb{N}_0$, $m \in \mathbb{Z}_0$ as quadratic forms as we will see.

To deal with this problem we offer the following preliminary solution: We note that the expression $-\tilde{C}$ under the square root is constrained to be positive in the classical theory. Consider any function $\Lambda_0 > 0$ and write

$$-L^{-1}\sqrt{-2Q\tilde{C}} = -\sqrt{2} \ L^{-1}Q\sqrt{\Lambda_0} \ [1 - (1 + \frac{\tilde{C}}{\Lambda_0 Q})]^{1/2}$$
(2.31)

We have $y:=1+\frac{\tilde{C}}{Q\Lambda_0}\leq 1$ and for such y we have the estimate

$$1 - \frac{y}{2} - \frac{y^2}{2} \le \sqrt{1 - y} \le 1 - \frac{y}{2} \tag{2.32}$$

To see this note that for $y \le 1$ the claimed upper bound is positive so the claimed upper inequality is equivalent to $1 - y \le (1 - y/2)^2$ which is identically satisfied even for all y. The claimed lower bound can be written as $\frac{1}{2}(1 - y)(2 + y)$ which for $y \le 1$ is positive for $y \ge -2$. Thus for $-2 \le y \le 1$ the claimed lower inequality is equivalent to $(1 - y)^2 (2 + y)^2 \le 4 (1 - y)$ which holds for y = 1 and for y < 1 is eqivalent to $(1 - y) (2 + y)^2 - 4 = -y^2(3 + y) \le 0$ which indeed holds for $-2 \le y \le 1$. Thus the claimed lower bound holds when the lower bound is positive and thus also for all other y < 1 as $\sqrt{1 - y}$ is not negative. Of course, sharper bounds can be obtained by taking the higher order Taylor expansion of $\sqrt{1 - y}$ in terms of y into account.

Accordingly, we can approximate the square root by an interpolating value between upper and lower bound

$$\sqrt{1-y} = 1 - y/2 - k y^2/2, \ k \in [0,1]$$
 (2.33)

Accordingly we could approximate the Hamiltonian by

$$H_k = \sqrt{2\Lambda_0} L^{-1} \int d^D x \, Q \, \sqrt{2\Lambda_0} L^{-1} \left[y/2 + k \, y^2/2 - 1 \right]$$
(2.34)

and for any $0 \le k \le$ the approximant H_k is in between the classically allowed extrema. Alternatively we could truncate the Taylor expansion of $-\sqrt{1-y}$ at some order N.

While upper and lower polynomial bounds are good to know an exact expression that avoids the square root would be benefitial. Such an expression is given by

$$\sqrt{z} = z \int_{\mathbb{R}} \frac{ds}{\sqrt{2\pi}} e^{-\frac{s^2}{2}z}$$
(2.35)

which however is no longer polynomial. It would require to be able to quantise $Q \ z \ e^{-s^2 z/2}$, $z = -\frac{\tilde{C}}{Q}$ as a quadratic form which is not known to be possible.

In lack of a better option we treat k or the truncation order N as a phenomenological parameter that serves to approximate the exact expression. We therefore have to make sure that the k dependent or higher than order N terms indeed are subleading by providing a choice for Λ_0 . To that extent, we will see in section 4 that we may actually define $\frac{\tilde{C}}{Q}$ as a quadratic form in Fock representations with dense form domain defined by the span of the excitations of a coherent state Ω_Z where Z denotes a point in the classical phase space such that the corresponding D-metric is positive definite. We thus pick

$$\Lambda_0 := - < \Omega_Z, \frac{\tilde{C}}{Q} \Omega_Z > \tag{2.36}$$

and quantise (2.34) by normal ordering. If Ω_Z is a coherent state peaked on a spatially very homogeneous universe then Λ_0 is expected to be almost a constant on sufficiently large scales. It is expected that the fluctuation factors $\Delta := 1 + \tilde{C}/(\Lambda_0 Q) = 1 - \frac{\tilde{C}}{Q} [<\frac{\tilde{C}}{Q}>]^{-1}$ contribute subdominantly in : $Q \Delta^n$: where : . : denotes normal ordering. See the end of section 4.3 for more details.

In any case, the point of this discussion is that the objects

$$\tilde{C}^n Q^{n-1} \tag{2.37}$$

can be quantised non-perturbatively as quadratic forms on the Fock space. The expansion in powers of n is not to be considered as a perturbative expansion but rather is similar to the relativistic expansion $p_0 = -\sqrt{m^2 + \vec{p}^2} = m + \frac{\vec{p}^2}{2m} + ...$ of the mass shell condition $-p_0^2 + \vec{p}^2 + m^2 = 0$ with the difference that the square root is not expanded at m but rather 0. This is because we have not solved for the analog of p_0 which would be the negative gravitational mode but rather for the analog of $p_D = -\sqrt{-[-p_0^2 + \vec{p}_{\perp}^2 + m^2]}$ where $\vec{p} = (\vec{p}_{\perp}, p_D)$ and the analog of m is the cosmological constant term. This cannot be avoided: It would have been more natural to solve the constraint C = 0 for the negative conformal gravitational mode. But this is not possible algebraically as the gravitational momenta, including the conformal mode, appear in C_a with derivatives.

2.4 Fock representations

Consider an arbitary background metric g of Euclidian signature on σ . From it we can construct the Laplacian $\Delta = g^{ab} \nabla_a \nabla_b$ where ∇_a is the torsion free covariant derivative compatible with g. The scalar one particle Hilbert space is defined by $L_2 \equiv L_2(\sigma, \sqrt{\det(g)} \ d^D x)$ and Δ is either essentially self-adjoint or at least has self-adjoint extensions of which we pick one. Let κ be an invertible (in the sense of the spectral theorem) function of Δ and consider canonically conjugate tensor fields $q_{b_1..b_B}^{a_1..a_A}$, $p_{a_1..a_A}^{b_1..b_B}$ of dual type (A, B, w = 0) and (B, A, w = 1) respectively where w is the density weight. We consider the annihilators

$$A_{c_1...c_{A+B}} := 2^{-1/2} \left[\kappa \ g_{c_1a_1}..g_{c_Aa_A} \ q_{c_{A+1}..c_{A+B}}^{a_1..a_A} - i\kappa^{-1} \ \omega^{-1}g_{c_{A+1}b_1}..g_{c_{A+B}b_B} \ p_{c_1..c_A}^{b_1..b_B} \right]$$
(2.38)

where

$$\omega := [\det(g)]^{1/2} \tag{2.39}$$

is invoked such that A is a background tensor of type (0, A + B, 0).

The annihilators are smeared with density weight zero test functions $f^{a_1..a_{A+B}}$ of type (A+B,0,0) relative to the volume form $\omega d^D x$

$$A(f) := \langle f, A \rangle_{L_2} := \int d^D x \,\omega \, [f^{c_1 \dots c_{A+B}}]^* \, A_{a_1 \dots a_{A+B}} = 2^{-1/2} \, [\langle \kappa f, q \rangle - i \langle \omega^{-1} \kappa^{-1} f, p \rangle]$$
(2.40)

where indices are pulled with g. Then using

$$[p_{a_1..a_A}^{b_1..b_B}(x), \ q_{b_1'..b_B'}^{a_1'..a_A'}(x')] = i\delta_{a_1}^{a_1'}..\delta_{a_A}^{a_A'}\delta_{b_1'}^{b_1}..\delta_{b_B'}^{b_B} \ \delta(x,x')$$
(2.41)

where $\int d^D x \delta(x, y) f(y) := f(x)$ we find

$$[A(f), A(g)^*] = \frac{1}{2} \left[< \kappa f, \kappa^{-1}g > + < \kappa^{-1}f, \kappa g > \right] = < f, g >$$
(2.42)

where self-adjointness of $\kappa = \kappa(\Delta)$ was used: We have with a multi-index $\mu = (a_1..a_C)$ and recalling that ∇_a acts on densities of weight w by a correction $-w \Gamma^b_{ab} = -[\ln(\omega^w)]_{,a}$ so that $\nabla_a \tilde{v}^a = \partial_a \tilde{v}^a$ for vector densities of weight one

$$< f, \Delta g > = \int d^{D}x \,\omega \, f_{\mu}^{*} \nabla_{a} (\nabla^{a} g^{\mu}) = \int d^{D}x \, f_{m} u^{*} \nabla_{a} (\nabla^{a} \omega g^{\mu})$$

$$= \int d^{D}x \, [\nabla_{a} (f_{\mu}^{*} \nabla^{a} \omega g^{\mu}) - (\nabla_{a} f_{\mu}^{*}) \, (\nabla^{a} \omega g^{\mu})]$$

$$= -\int d^{D}x \, (\nabla_{a} \omega f_{\mu}^{*}) \, (\nabla^{a} g^{\mu})$$

$$= -\int d^{D}x \, [\nabla_{a} (\nabla^{a} \omega f_{\mu}^{*}) \, g^{\mu}) - (\nabla_{a} (\nabla_{a} \omega f_{\mu}^{*})) \, g^{\mu}]$$

$$= < \Delta f, g >$$

$$(2.43)$$

assuming that the integrations by parts result in no boundary terms.

Fermions are naturally densities of weight 1/2 and for each Weyl fermion component ρ_I we have ρ_I^* as its anti-conjugate

$$[\rho_I(x), \rho_{I'}^*(x')]_+ = \delta_{II'} \,\,\delta(x, x') \tag{2.44}$$

We thus define the annihilator to be $B_I = \omega^{-1/2} \rho_I$.

The joint Fock vacuum is then defined by $A_{\mu}(x)\Omega = B_{I}(x)\Omega = 0$.

3 Polynomial SDC and HC as quadratic forms on Fock space

We first define the constraints as quadratic forms on the Fock space. Then we make some general remarks on the solution of constraints which are merely quadratic forms but not operators as well as the possibility of a commutator algebra of quadratic forms. We then give details for the case at hand when a Fock structure is available.

3.1 Polynomial constraint quadratic forms on Fock space

Apart from algebraic work, there is not much to do. We have already shown that in section 2 that both the SDC and HC can be written in polynomial form by multiplying the HC by a sufficiently high power of $\sqrt{\det(q)}$. Now for any polynonomial

$$P(x) = P(q(x), p(x), [\partial q](x), [\partial p](x), \rho(x), \rho^*(x), [\partial \rho](x), [\partial \rho^*](x))$$
(3.1)

of the fields and its first spatial derivatives where (q, p) and (ρ, ρ^*) denotes the collection of all bosonic and fermionic conjugate pairs respectively, we decompose q, p into annihilation and creation operators by inverting (2.38)

$$q = 2^{-1/2} [\otimes^A g^{-1}] \otimes [\otimes^B 1_D] \kappa^{-1} \cdot [A + A^*], \quad p = i \ 2^{-1/2} \ \omega[\otimes^A 1_D] \otimes [\otimes^B g] \kappa \cdot [A - A^*]$$
(3.2)

while $\rho = B, \rho^* = B^*$ are already annihilation and creation operators. Then in every monomial we order all occurrences of A^* left of all occurrences of A and all occurrences of B^* left of all occurrences of B. For the bosonic and fermionic sector we multiply by +1 and -1 for any interchange of the A, A^* and of the B, B^* respectively in this process.

The reordered polynomial is denoted by : P(x) : and given a definition of annihilation and creation operators the ordering of : P(x) : is unique. For any $x \in \sigma$, : P(x) : is a densely defined quadratic form on the Fock space \mathcal{H} where the dense form domain \mathcal{D} is the finite linear span of the Fock states

$$A(f_1)^* \dots A(f_M)^* B(g_1)^* \dots B(g_N)^* \Omega$$
(3.3)

where $M, N \in \mathbb{N}_0$ are the bosonic and fermionic particle numbers respectively and f_k, g_l are test functions (typically from the Schwartz function class $S(\sigma)$ of smooth functions of rapid decrease at $\partial \sigma$ if σ is not bounded or has a boundary). To see this we decompose P into monomials Q with say K_B, K_F creators and L_B, L_F annihilators (the subscripts denote bosonic and fermionic respectively) and consider

$$\langle \psi, : Q(x) : \psi' \rangle_{\mathcal{H}}$$
 (3.4)

where ψ, ψ' are of the form (3.3) with particle numbers M, N and M', N' respectively and let all annihilators and creators respectively act on ψ', ψ respectively. The result is zero if either of $K_B > M, K_F > N, L_B > M', L_F > N'$ or of the form

$$<\psi_1,\psi_1'>h(x)$$
 (3.5)

where the particle number of ψ_1, ψ'_1 respectively is given by $(M - K_B, N - K_F)$, $(M' - L_B, N' - L_F)$ and h is a test function again (a product of $K_B + K_F + L_B + L_F$ test functions involved in ψ, ψ' or their derivatives convoluted with $g, g^{-1}, \kappa, \kappa^{-1}$).

Now the smeared, polynomial, normal ordered constraints are given by

$$: C(f) := \int d^{D}x f^{\mu}(x) : C_{\mu}(x) :$$
(3.6)

and thus are symmetric (i.e. $\langle \psi, : C(f) : \psi' \rangle^* = \langle \psi', : C(f) : \psi \rangle$) quadratic forms on the Fock space with dense form domain \mathcal{D} because for any test function h the integral $\int d^D x f^{\mu} h$ exists.

3.2 Solution of the constraints as quadratic forms, constraint algebra and solutions

The usual approach to constraint quantisation is as follows [28]: Suppose that some constraints Z_I , $I \in \mathcal{I}$ some index set are defined *as operators* on a dense invariant domain \mathcal{D} of the Hilbert space \mathcal{H} , that is, $Z_I \mathcal{D} \subset \mathcal{D}$. A linear functional l on \mathcal{D} is called a solution iff $l[Z_I \psi] = 0$ for all $I \in \mathcal{I}$, $\psi \in \mathcal{D}$. Note that $Z_I \psi \in \mathcal{D}$ so that ldefined on \mathcal{D} is defined on $Z_I \psi$.

Now let b_n be an orthonormal basis of \mathcal{H} constructed from elements of \mathcal{D} (in the separable case we obtain this by using the Gram-Schmidt process, in the non-separable case this involves the axiom of choice. Fock spaces are separable). Then for an operator Z_I densely defined on invariant \mathcal{D} we have $Z_I \psi \in \mathcal{D} \subset \mathcal{H}$, thus

$$Z_I \psi = \sum_n b_n \langle b_n, Z_I \psi \rangle, \qquad (3.7)$$

and

$$||Z_I\psi||^2 = \sum_n |\langle b_n, Z_I\psi \rangle|^2$$
(3.8)

converges. A solution l is now determined by the coefficients

$$l_n := l[b_n] \tag{3.9}$$

and can be written as the linear functional

$$l := \sum_{n} l_n \langle b_n, . \rangle_{\mathcal{H}}$$
(3.10)

It follows that l is a solution iff

$$\sum_{n} l_n \langle b_n, Z_I \psi \rangle = 0 \tag{3.11}$$

for all $I \in \mathcal{I}, \psi \in \mathcal{D}$ which is now an infinite system of linear equations on the coefficients l_n with no growth condition on l_n as n varies.

Now suppose that the constraints Z_I are not operators on \mathcal{D} but merely quadratic forms. Then $Z_I\psi$ is illdefined as a vector in the Hilbert space, we can only compute matrix elements $\langle \psi, Z_I\psi' \rangle$. In particular, while we may try to use (3.7) as a formal definition, formula (3.8) now diverges. However, the condition (3.11) is still meaningful! That is, to state (3.11) it is sufficient that Z_I is a quadratic form on \mathcal{D} and that we know the coefficients (3.9) of l with respect to an ONB $b_n \in \mathcal{D}$. We thus require l to be a linear functional not only on \mathcal{D} but also on the formal range of Z_I on \mathcal{D} defined by (3.7). We may consider $Z_I\psi$ as an anti-linear functional on \mathcal{D} by

$$\psi' \mapsto (Z\psi)[\psi'] := \langle \psi', Z_I \psi \rangle \tag{3.12}$$

which is well-defined and then l acts on $Z_I\psi$ by

$$l[Z_I\psi] := \sum_n \ l[b_n] \ (Z\psi)[b_n] \tag{3.13}$$

This can converge even absolutely if the $l[b_n]$ decay sufficiently much faster than the $(Z_I\psi)[b_n]$ as n varies.

Assuming that the solution space consists not only of the trivial solution $l \equiv 0$, pick $l_0 \neq 0$. Then we may try to define an inner product, at least in the separable case, on the space L of solutions by

$$< l, l'>_L := \lim_M \frac{\sum_{n \le M} l[b_n]^* l'[b_n]}{\sum_{n \le M} |l_0[b_n]|^2}$$
(3.14)

where we assumed w.l.g. that b_n is labelled by $n \in \mathbb{N}_0$.

Finally suppose that the constraints originate from functions on the phase space which we denote by the same symbol Z_I and that form a Poisson Lie algebra

$$\{Z_I, Z_J\} = f_{IJ} \stackrel{K}{} Z_K \tag{3.15}$$

with structure constants f_{IJ}^{K} . If the Z_{I} are operators with dense invariant domain \mathcal{D} then for $\psi \in \mathcal{D}$ we have $Z_{I}\psi \in \mathcal{D}$ so that $[Z_{I}, Z_{J}]\psi \in \mathcal{D}$ and thus we define the quantum constraint algebra to be anomaly free iff

$$[Z_I, Z_J]\psi = i f_{IJ} {}^K Z_K \psi$$
(3.16)

for all $I, J \in \mathcal{I}, \psi \in \mathcal{D}$. If the Z_I are merely quadratic forms then $Z_I \psi$ is meaningless and while $\langle \psi, Z_I \psi' \rangle$ is well defined, again $\langle \psi, Z_I Z_J \psi' \rangle$ is meaningless. However, let

$$P_M := \sum_{n \le M} b_n < b_n. >_{\mathcal{H}}$$

$$(3.17)$$

be the projection onto \mathcal{H}_M , the completion of the linear span of the b_n , $n \leq M$, and suppose that for each $M < \infty$ the numbers

$$\langle \psi, Z_I P_M Z_J \psi' \rangle := \sum_{n \le M} \langle \psi, Z_I b_n \rangle \langle b_n, Z_J \psi' \rangle$$
(3.18)

exist. Then due to cancellation effects the weak limit

$$\langle \psi, [Z_I, Z_J]\psi' \rangle := \lim_M \left[\langle \psi, Z_I P_M Z_J \psi' \rangle - \langle \psi, Z_J P_M Z_I \psi' \rangle \right]$$

$$(3.19)$$

may exist. In this case we have a *commutator algebra of quadratic forms* and we define it to be free of anomalies iff

$$\langle \psi, [Z_I, Z_J] \psi' \rangle = i f_{IJ} {}^K \langle \psi, Z_K \psi' \rangle$$

$$(3.20)$$

for all $I, J \in \mathcal{I}, \psi, \psi' \in \mathcal{D}$ i.e. in the sense of quadratic forms which generalises (3.16).

In case that the classical constraints Z_I close as in (3.15) but with non-constant structure functions f_{IJ} K the condition (3.16) (for the case that Z_I are operators with dense invariant domain \mathcal{D}) is ill-defined as it stands because now f_{IJ} K is itself operator valued so that ordering issues arise. For instance, if Z_I, Z_J, f_{IJ} K are symmetric operators then an ordering consistent with the adjointness relations would be

$$[Z_I, Z_J]\psi = \frac{i}{2} (f_{IJ} {}^K Z_K + Z_K f_{IJ} {}^K) \psi$$
(3.21)

Let now l be a solution of the constraints in the sense of $l[Z_I\psi] = 0$ for all $I \in \mathcal{I}, \psi \in \mathcal{D}$ then as $Z_I\psi \in \mathcal{D}$ we would have the identity

$$l[[Z_I, Z_J]\psi] = 0 = \frac{i}{2} l[(f_{IJ} \ ^K \ Z_K + Z_K \ f_{IJ} \ ^K) \ \psi]$$
(3.22)

for all I, J, ψ . If moreover $f_{IJ} \stackrel{K}{=} \psi \in \mathcal{D}$ for all I, J, K, ψ then

$$0 = \frac{i}{2} l[[f_{IJ} {}^{K}, Z_{K}] \psi]$$
(3.23)

for all I, J, ψ . Thus l vanishes not only on all the Z_I but also on all of the $Z_{IJ} = i[f_{IJ} K, Z_K]$ which correspond to the classical Poisson brackets $\{f_{IJ} K, Z_K\}$ which are not required to vanish in the classical theory. This can be iterated and can result in a space of solutions which is too small [29]. Accordingly, in case of structure functions it is not sufficient to have e.g. the property (3.21), one must make sure that the space of solutions is large enough. This may may be achieved for non-symmetric constraint operators [16] or non-symmetric structure functions or for the case that above assumption $f_{IJ} K \psi \in \mathcal{D}$ fails.

Now consider the case that Z_I are only quadratic forms. Then the above obstruction argument fails because we do not have the property $Z_I\psi \in \mathcal{D}$ so that we cannot conclude from $l[Z_I\psi] = 0$ that also $l[Z_I Z_J\psi] = 0$. Hence in this case there is no immediate conclusion about the size of the space of solutions to be drawn even if the Z_I are symmetric quadratic forms. This is because of (3.19) which says that while $\langle \psi, Z_I Z_J \psi' \rangle$, $\langle \psi, Z_J Z_I \psi' \rangle$ are ill defined (divergent) the difference between $\langle \psi, Z_I P_M Z_J \psi' \rangle$ and $\langle \psi, Z_J P_M Z_I \psi' \rangle$ converges weakly as $M \to \infty$. Thus while we have

$$l[Z_I\psi] = \lim_{M'} \sum_{n' \le M'} l_{n'} < b_{n'}, Z_I\psi >= 0$$
(3.24)

we have

$$l[[Z_I, Z_J]\psi] := \lim_{M'} \sum_{n' \le M'} l_{n'} < b_{n'}, [Z_I, Z_J]\psi > := \lim_{M'} \sum_{n' \le M'} l_{n'} \lim_{M} \{ < b_{n'}, Z_I P_M Z_J \psi > - < b_n, Z_J P_M Z_I \psi > \}$$
(3.25)

If $P_M Z_J \psi \in \mathcal{H}_M$ and $P_M Z_I \psi \in \mathcal{H}_M$ would converge strongly to some $\psi_J, \psi_I \in \mathcal{D}$ then

$$l[[Z_I, Z_J]\psi] = \lim_{M'} \sum_{n' \le M'} l_{n'} \{ \langle b_{n'}, Z_I\psi_J \rangle - \langle b_{n'}, Z_J\psi_I \rangle \} = l[Z_I\psi_J] - l[Z_J\psi_I] = 0$$
(3.26)

However this is not the case, otherwise Z_I, Z_J would be operators. Alternatively note that

$$l[[Z_I, Z_J]\psi] = \lim_{M'} \lim_{M} \sum_{n' \in \leq M'} l_{n'} \sum_{n \leq M} \{ < b_{n'}, Z_I \ b_n > < b_n, Z_J\psi > - < b_{n'}, Z_J \ b_n > < b_n, Z_I\psi > \}$$
(3.27)

If we could interchange the limits $M, M' \to \infty$ then (3.27) would indeed vanish. But such interchange of limits is typically only allowed when the double sum is absolutely convergent (Fubini type of theorem) which is not the case as the existence of $l[Z_I\psi] = 0$ is crucially sensitive on the phases of the complex numbers l_n .

Note that if $[Z_I, Z_J]$ exists as a quadratic form and equals some ordered version of the classical expression $f_{IJ} \stackrel{K}{} Z_K$, that quantum version will be normal ordered in the case of the Fock quantisation. However then it will certainly not be ordered as $Z_K f_{IJ} \stackrel{K}{}$ and thus $l[[Z_I, Z_J]\psi] \neq 0$ even if $f_{IJ} \stackrel{K}{}$ has \mathcal{D} as invariant domain.

This shows that the case of non-trivial structure functions comes with extra complications and motivates to recast the algebra of constraints into a form in which the structure functions are replaced by structure constants. Locally this is always possible because one can in principle solve the constraints for some of the momenta and in this form the Poisson algebra of constraints is Abelian. Another possibility is the minimal or maximal master constraint approach discussed in the previous section. In Abelianised version and the minimal master constraint version the constraints are no longer polynomial which therefore calls for new tools developed in section 4. The maximal master constraint version allows for a polynomial formulation and therefore fits into the present general discussion. The classical maximal master constraint is constructed from the Z_I as

$$M = \frac{1}{2} \sum_{I,J} Z_I Q^{IJ} Z_J$$
(3.28)

where $(Q^{IJ})_{I,J\in\mathcal{I}}$ is a positive definite matrix which may be a non-constant function on the classical phase space. One may certainly choose Q^{IJ}, Z_I as polynomials so that M is a polynomial again and thus may be formulated as quadratic form in Fock representations by normal ordering.

One may criticise that the master constraint (3.28) is "blind" for possible anomalies in the algebra or that it does not allow to check whether the classical algebra is represented in the quantum theory. However, consider in the classical theory phase space functions G^I , so called gauge fixing conditions, such that $\Delta_I \ ^J := \{Z_I, G^J\}$ is non-degenrate close to the constraint surface $Z_I = 0, \ I \in \mathcal{I}$. Then

$$\{M, G^J\} [\Delta^{-1}]_J {}^M Q_{MI} =: \{M, G^J\} \hat{Q}_{JI} = Z_I + O(Z^2), \ Q_{IM} Q^{MJ} = \delta_I {}^J$$
(3.29)

so that Z_I maybe recovered from M using G^I . Then

$$\{Z_I, Z_J\} = f_{IJ} {}^K Z_K = f_{IJ} {}^M \hat{Q}_{KM} \{M, G^K\} + O(Z^2) = \hat{Q}_{KI} \hat{Q}_{LJ} \{\{M, G^K\}, \{M, G^L\}\} + O(Z^2)$$
(3.30)

i.e.

$$\{\{M, G^I\}, \{M, G^J\}\} = \hat{f}^{IJ}_K \{M, G^K\} + O(Z^2), \ \hat{f}^{IJ}_K = f_{MN} \ ^L Q^{IM} \ Q^{JN} \ Q_{KL}, \tag{3.31}$$

so that the $Z^I := \{M, G^I\}$ also close with structure functions, albeit in general with non polynomial ones. Thus classically M is truly equivalent to the set $\{Z_I\}_{I \in \mathcal{I}}$ and in the quantum theory one could in principle try to consider the constraint algebra defined by the $[M, G^I]$.

Of course the whole point of the master constraint is to circumvent that constraint algebra with structure functions and one expects that unitarily inequivalent representations of the CCR and CAR supporting M as a quadratic form differ in the "gap number"

$$M_0 := \inf_{0 \neq \psi \in \mathcal{D}} \frac{\langle \psi, M\psi \rangle}{||\psi||}$$
(3.32)

For instance, in Fock representations the classical positivity is destroyed in the normal ordering process while M is a symmetric quadratic form. One therefore expects that in Fock representations the expectation value of M w.r.t. $\psi \in \mathcal{D}$ takes full range in \mathbb{R} , i.e. $M_0 = -\infty$ so that solutions $l[M\psi] = 0$ for all $\psi \in \mathcal{D}$ should exist. In that case a systematic procedure consists in solving the infinite system

$$K_m := \sum_n l_n M_{n,m} = 0; \quad M_{n,m} = \langle b_n, M b_m \rangle$$
 (3.33)

as follows: Let $(b_n)_{n=0}^{\infty}$ be a labelling of the ONB by $n \in \mathbb{N}_0$ and set $M_{n,m}^{(0)} := M_{n,m}, m, n \ge 0$. Find all m such that $M_{n,m}^{(0)} = 0 \ \forall n \in \mathbb{N}_0$ and drop those m from the set \mathbb{N}_0 resulting in R_0 . For the smallest $m_0 \in R_0$ find the smallest $n = n_0$ in $L_0 := \mathbb{N}_0$ such that $M_{n,m_0} \ne 0$. Solve $K_{m_0} =$ for $l_{n_0} = \hat{l}_{n_0}$ and substitute that solution into the $K_m, m_0 \ne m \in R_0$. Collecting coefficients one can write these K_m as a sum over $l_n, n \in L_1 = L_0 - \{n_0\}$ with coefficients $M_{n,m}^{(1)}$. Drop all m from $R_0 - \{m_0\}$ for which $M_{n,m}^{(1)} = 0 \ \forall n \in L_1$ resulting in R_1 . Now iterate, thereby producing sets $L_0 \supset L_1 \supset ... \supset L_N \supset ..., N \in \mathbb{N}_0, L_{N+1} = L_N - \{n_N\}$. The space of solutions $\{l_n\}_{n \in \mathbb{N}_0}$ will be parametrised by the free coefficients $l_n, n \notin \{n_N\}_{N=0}^{\infty}$.

3.3 Implementation in Fock representations

Parts of the exposition of the present section are also considered in [31]. We include here a shorter version for reasons of self-containedness and refer to [31] for more details. Note that the index set \mathcal{I} considered in this section labels a basis e_I in the one particle Hilbert space and therefore can be used to smear constraints Z as $Z_I := \langle e_I, Z \rangle$. In this case the meaning of \mathcal{I} is the same as in the previous section. In general however the meaning of \mathcal{I} in this and the previous section is logically independent.

The concrete implementation of the programme described in the previous section in Fock representations is sytematised by picking an orthonormal basis of real valued functions e_I with mode label $I \in \mathcal{I}$ of the scalar one particle Hilbert space $L_2 = L_2(\sigma, d^D x \omega)$ and a background D-Bein h_j^a , j = 1..D with inverse co-D-Bein h_a^j such that $g_{ab}h_j^ah_k^b = \delta_{jk}$. Then $e_{I;j_1..j_A}^{k_1..k_B}$ is an ONB of tensors of type (A, B, 0) with components

$$[e_{I;j_1..j_A}^{k_1..k_B}]_{b_1..b_B}^{a_1..a_A}(x) = e_I(x) \prod_{r=1}^A h_{j_r}^{a_r}(x) \prod_{s=1}^B h_{b_s}^{k_s}(x)$$
(3.34)

with respect to the one particle inner product

$$\langle t, \hat{t} \rangle := \int_{\sigma} d^{D}x \ \omega \ \overline{t_{b_{1}..b_{B}}^{a_{1}..a_{A}}} \ \prod_{r=1}^{A} g_{a_{r}a_{r}'} \ \prod_{s=1}^{B} g^{b_{s}b_{s}'} \ \hat{t}_{b_{1}'..b_{B}'}^{a_{1}'..a_{A}'}$$
(3.35)

We define the background scalars of density weight zero (indices j, k are moved with the Kronecker δ)

$$q_{j_1..j_{A+B}} := h_{a_1j_1}..h_{a_Aj_A} h_{j_{A+1}}^{b_1}..h_{j_{A+B}}^{b_B} q_{b_1..b_B}^{a_1..a_A}, \quad p_{j_1..j_{A+B}} := \omega^{-1} h_{j_1}^{a_1}..h_{j_A}^{a_A} h_{b_1j_{A+1}}..h_{b_Bj_{A+B}} p_{a_1..a_A}^{b_1..b_B}, \quad (3.36)$$

which have Poisson brackets

$$\{p_{j_1..j_{A+B}}(x), q_{k_1..k_{A+B}}(y)\} = \delta(x, y) \prod_{r=1}^{A+B} \delta_{j_r k_r}$$
(3.37)

We define the bosonic smeared annihilators

$$A_{I;j_1..j_{A+B}} := \langle e_I, A_{j_1..j_{A+B}} \rangle := 2^{-1/2} \langle e_I, [\kappa \cdot q_{j_1..j_{A+B}} - i\kappa^{-1} \cdot p_{j_1..j_{A+B}} \rangle_{L_2}$$
(3.38)

where $\kappa = \kappa(\Delta), \ \Delta = \omega^{-1} \partial_a \omega g^{ab} \partial_b$ and the fermionic smeared annihilators

$$B_{I\alpha} = \langle e_I, B_{\alpha} \rangle := \langle e_I, \omega^{-1/2} \rho_{\alpha} \rangle_{L_2}$$
 (3.39)

which annihilate the Fock vacuum Ω .

The Fock orthonormal basis is then defined by

$$b_n := \prod_{I,\mu} \frac{[C_{I\mu}^*]^{n_{I\mu}}}{\sqrt{n_{I\mu}!}} \,\Omega, \ ||n|| := \sum_{I,\mu} n_{I,\mu} < \infty$$
(3.40)

where μ stands for one of the multi-indices $j_1...j_{A+B}$ or α respectively in which $C_{I,\mu}$ means $A_{I,\mu}$ or $B_{I,\mu}$ respectively and the *occupation numbers* $n_{I\mu} \in \mathbb{N}_0$.

Let \mathcal{D} be the finite linear span of the b_n . A truncation structure on \mathcal{I} is a sequence of finite mode sets $S_M \subset \mathcal{I}, \ M \in \mathbb{N}_0$ with $S_0 = \{I_0\}, \ S_M \subset S_{M+1}, \ \cup_M S_M = \mathcal{I}$ and I_0 is a single element in \mathcal{I} . It defines $\mathcal{D}_M \subset \mathcal{D}$ to be the span of the b_n with $n_{I,\mu} = 0$ for $I \notin S_M$. Note that the truncation structure does not impose any restriction on the occupation numbers $n_{I,\mu}$ for $I \in S_M$ except that by definition of a Fock state $||n|| := \sum_{I,\mu} n_{I\mu} < \infty$. For instance, for $\sigma = \mathbb{R}^D$ and $g_{ab} = \delta_{ab}$ the e_I could be the Hermite functions $I = (I_1, ..., I_D) \in \mathbb{N}_0^D$ and $I \in S_M$ iff $|I| = I_1 + ... + I_D \leq M$ and $I_0 := 0$. Similarly, for $\sigma = T^D$ the e_I could be the momentum eigenfunctions with $I = (I_1, ..., I_D) \in \mathbb{Z}^D$ and $I \in S_M$ iff $|I| = |I_1| + ... + |I_D| \leq M$ and again $I_0 := 0$.

The truncation structure can be used in order to define the truncations : $P_M(x)$: of a normal ordered polynomial : P(x) : as follows: We write : P(x) : in terms of the $A_{j_1..j_{A+B}}, B_{\alpha}$ and their adjoints and then use the completeness relation

$$\delta(x,y) = \sum_{I} e_{I}(x) [e_{I}(y)]^{*} \omega(y), \quad \int d^{D}y \, \delta(x,y) \, f(y) = f(x)$$
(3.41)

to expand the annihililation and creation operators in terms of the smeared versions (3.38), ((3.37). For each monomial of order N we obtain N sums over the $I_1, .., I_N \in \mathcal{I}$. The truncation consists in the restriction of those sums to $I_1, .., I_N \in S_M$. In this way the infinite series is truncated to a normal ordered polynomial in the smeared creation and annihilation operators, for each monomial of order N we obtain at most $|S_M|^N$ terms where $|S_M|$ is the finite number of elements of S_M .

Note that since $S_M \to \mathcal{I}, \ M \to \infty$, for each ψ, ψ' we find $M \in \mathbb{N}_0$ such that $\psi, \psi' \in \mathcal{D}_M$. Therefore

$$<\psi, : P(x): \psi' > = <\psi, : P_M(x): \psi' >$$
 (3.42)

To see this consider : P(x) : expanded into smeared annihilation and creation operators without truncation of the infinite series involved. Then let all creation operators act to the left on $\langle \psi, . \rangle$ and all annihilation operators to the right on ψ' . Whenever the mode label I of such an operator is not in S_M we can commute it trough the creation operators acting on Ω that are involved in ψ, ψ' whereby we pick at most a sign and then Ω is annihilated. It follows

$$: P(x) := w - \lim_{M \to \infty} : P_M(x) :$$
 (3.43)

i.e. every normal ordered polynomial is the weak operator topology limit of its truncations which is the appropriate operator topology to choose for quadratic forms.

Now quadratic forms cannot be multiplied, i.e. their product is no longer a quadratic form which is the source of most complications in QFT and necessiates regularisation and renormalisation procedures when considering the S-matrix (an infinite series of smeared interaction terms which itself is not an operator, only a quadratic form). However the truncations : $P_M(x)$: are actually operators with dense invariant domain \mathcal{D} because they are polynomials in smeared creation and annihilation operators with coefficients that take values in the Schwarz functions. Thus taking products of the truncations results in well defines operators densely defined on \mathcal{D} whose limit in general does not exist, not even as a quadratic form. However, due to cancellations involved, the limit of commutators of truncations may exist as a quadratic form.

Thus, given smeared, normal ordered polynomial constraints and their truncations

$$C(u) := \int d^D x \ u^{\mu}(x) \ : C_{\mu}(x) :, \ C_M(u) := \int d^D x \ u^{\mu}(x) \ : C_{\mu,M}(x) :, \tag{3.44}$$

we define, if it exists

$$[C(u), C(v)] := w - \lim_{M \to \infty} [C_M(u), C_M(v)]$$
(3.45)

In [31] we have carried out the analysis explicitly for the case of the spatial diffeomorphism constraints and found that

$$[C[u], C[v]] = iC(-[u, v])$$
(3.46)

where [u, v] is the commutator of vector fields. That is, for any choice of Fock representation of the above type, no matter that it is heavily background dependent, the algebra of SDC quadratic forms not only exists but it is free of anomalies! This shows that background dependence of elements of the quantum theory (here the representation of the CCR and CAR) does not spoil the background independence of the algebra of spatial diffeomorphisms.

We now want to analyse whether this result can be extended to the full algebra of constraints for Lorentzian GR, written in polynomial form. We consider first the vacuum sector. We have seen that we can work with (P^{ab}, q_{ab}) and in D = 3 the HC becomes a polynmial of degree ten in density weight six while in the (A, E) formulation it becomes a polynomial of degree twenty with density weight fourteen. The higher the polynomial degree, the more terms will come out not normal ordered in the commutator calculation and the more potentially divergent and/or anomalous terms will appear upon reinstalling normal order. Thus, to get a first taste of the complications involved we consider Euclidian vacuum GR in 3+1 dimensions where we have available the connection formluation mentioned in section 2 involving GC, SDC and HC constraints G_j, C_a, C_0 which are only second or fourth order polynomials respectively in density weight two. We thus expect the complications involved to be less severe than for Lorentzian signature. The classical Poisson algebra of the polynomial constraints

$$C(r, u, f) = \int d^{D}x \{ r^{j} G_{j} + u^{a} C_{a} + f C_{0} \}, \ G_{j} = E^{a}_{j,a} + \epsilon_{jkl} A^{k}_{a} E^{a}_{m} \delta^{lm},$$

$$C_{a} = E^{b}_{j} A^{j}_{b,a} - (E^{b}_{j} A^{j}_{a})_{,b}, \ C_{0} = F^{j}_{ab} \epsilon_{jkl} \delta^{km} \delta^{ln} E^{a}_{m} E^{b}_{n}, \ F^{j}_{ab} = 2A^{j}_{[b,a]} + \delta^{jm} \epsilon_{mkl} A^{k}_{a} A^{l}_{b}$$
(3.47)

is given by

$$\{C(r, u, f), C(s, v, g)\} = -C(r \times s + u[s] - v[r] - 4 Q(\omega, A), [u, v] - 4 Q(\omega, .), u[g] - v[f])$$

$$Q^{ab} = E^a_j E^a_k \delta^{jk}, \ \omega_a = f \ g_{,a} - g \ f_{,a}, \ u[s]^j = u^a \ s^j_{,a}, \ u[f] = u^a \ f_{,a} - u^a_{,a} \ f$$

$$(3.48)$$

Note that s, f transform as scalars of density weight zero and minus one respectively under spatial diffeomorphisms. We carry out the above programme in the simplest topology, background metric and Fock representation $\sigma = \mathbb{R}^3, g_{ab} = \delta_{ab}, \kappa = 1$ and use the Hermite basis e_I , $I \in \mathbb{N}_0^3$, $|I| = I_1 + I_2 + I_3$ for concreteness to define the truncation structure. A possible triad is $h_a^j := \delta_a^j$ and $\omega = [\det(g)]^{1/2} = 1$. The annihilation operator is then given by

$$A_{mk} := 2^{-1/2} \{ h_m^a \ A_a^l \ \delta_{lk} - i \delta_{ml} h_a^l E_k^a \}$$
(3.49)

and we have the non-vanishing commutators

$$A_{mj}(x), A_{nk}^*(y)] = \delta_{mn} \,\delta_{jk} \,\delta(x, y) \quad \Leftrightarrow \quad [E_j^a(x), A_b^k(y)] = i\delta_b^a \,\delta_j^k \,\delta(x, y) \tag{3.50}$$

and the smeared operators are simply

$$A_{Imj} = \langle e_I, A_{jk} \rangle_{L_2}, \ [A_{Imj}, A^*_{Jnk}] = \delta_{IJ} \ \delta_{mn} \delta_{jk}$$
 (3.51)

The normal ordered constraints can be expanded as (we define the derivatives $f_{,j} := \delta_j^a f_{,a}$ and Einstein summation

over twice repeated lower case indices independent of index position being understood)

$$G(r) = -\frac{i}{2^{1/2}} \sum_{I} r_{Imi} [A_{Imi} - A_{Imi}^{*}] + \frac{i}{2} \epsilon_{ijk} \sum_{I,J} r_{IJk} [A_{Imi}^{*} (A_{Jmj} - A_{Jmj}^{*}) + (A_{Jmj} - A_{Jmj}^{*})A_{Imi}];$$

$$r_{Imi} := \langle e_{I}, r_{,m}^{i} \rangle, r_{IJk} := \langle r^{k}, e_{I}e_{J} \rangle$$

$$C(u) = \frac{i}{2} \sum_{I,J} u_{IJmn} \{ [A_{Imj} - A_{Imj}^{*}] A_{Jnj} + A_{Jnj}^{*} [A_{Imj} - A_{Imj}^{*}] \};$$

$$u_{IJmn} := \delta_{mn} \langle e_{I}, u_{p}e_{J,p} \rangle + \langle e_{I}e_{J}, u_{n,m} \rangle$$

$$-2 C_{0}(f) = 2^{1/2} \sum_{I,J,K} f_{IJnK} \epsilon_{ijk} : [A_{Imi} - A_{Imi}^{*}] [A_{Jnj} - A_{Jnj}^{*}]] [A_{Kmk} + A_{Kmk}^{*}] :$$

$$+ \frac{1}{2} \sum_{I,J,K,L} f_{IJnK} \epsilon_{rij} \epsilon_{rkl} : [A_{Imi} - A_{Imi}^{*}] [A_{Jnj} - [A_{Jnj}^{*}]] [A_{Knk} + A_{Knk}^{*}] [A_{Lmk} + A_{Kmk}^{*}] :$$

$$f_{IJnK} = \langle (e_{I}e_{J}f)_{,n}, e_{K} \rangle, f_{IJKL} = \langle e_{I}e_{J}e_{KeL}, f \rangle$$

$$(3.52)$$

These inner products are well defined as the Hermite basis consists of Schwartz functions.

The computations are organised by using the compound index notation

$$\alpha = (Imi), \ \beta = (Jnj), \ \gamma = (Kpk), \ \delta = (Lql)$$
(3.53)

and

$$r_{\alpha} = r_{Imi}, \ r_{\alpha\beta} = r_{IJk}\epsilon_{kij}\delta_{mn}, \ u_{\alpha\beta} = u_{IJmn}\delta_{ij}, \ f_{\alpha\beta\gamma} = f_{IJnK}\delta_{mp}\epsilon_{ijk}, \ f_{\alpha\beta\gamma\delta} = f_{IJKL} \ \epsilon_{rij} \ \epsilon_{rkl} \ \delta_{mp} \ \delta_{nq}$$
(3.54)

which allows the compact notation (summation over repeated compound indices understood).

$$G(r) = G_{1}(r) + G_{2}(r) = -i2^{-1/2} r_{\alpha} F_{\alpha} + \frac{i}{2} r_{\alpha\beta} : G_{\alpha} F_{\beta} :$$

$$C(u) = C_{2}(u) = \frac{i}{2} u_{\alpha\beta} : F_{\alpha} G_{\beta} :$$

$$-2C_{0}(f) = C_{0;3}(f) + C_{0;4}(f) = 2^{1/2} f_{\alpha\beta\gamma} : F_{\alpha} F_{\beta} G_{\gamma} : +\frac{1}{2} f_{\alpha\beta\gamma\delta} : F_{\alpha} F_{\beta} G_{\gamma} G_{\delta} :$$

$$F_{\alpha} := A_{\alpha} - A_{\alpha}^{*}, \ G_{\alpha} := A_{\alpha} + A_{\alpha}^{*}, \qquad (3.55)$$

Here : (.) : denotes normal ordering of (.) in the usual way, e.g.

$$: F_{\alpha} F_{\beta} G_{\gamma} := A_{\gamma}^{*} : F_{\alpha} F_{\beta} :+: F_{\alpha} F_{\beta} : A_{\gamma}$$

$$: F_{\alpha} F_{\beta} := F_{\alpha} A_{\beta} - A_{\beta}^{*} F_{\alpha}$$
(3.56)

In (3.55) the summation range of $\alpha, \beta, \gamma, \delta$ is unconstrained, that is, there is no restriction on the range of the mode label I in $\alpha = (Imi)$ while of course $m, i \in \{1, 2, 3\}$ take only a finite range. Therefore the expressions (3.55) are formal only.

We define the ordering on the compound indices

$$\alpha = (Imi) \le M \quad \Leftrightarrow \quad I \in S_M \tag{3.57}$$

Let now

$$G_{M_{1};M_{2},M_{3}}(r) = G_{1;M_{1}}(r) + G_{2;M_{1},M_{3}}(r) = -\frac{i}{2^{1/2}} \sum_{\alpha \leq M_{1}} r_{\alpha} F_{\alpha} + \frac{i}{2} \sum_{\alpha \leq M_{2},\beta \leq M_{3}} r_{\alpha\beta} : G_{\alpha} F_{\beta} :$$

$$C_{M_{1},M_{2}}(u) = C_{2;M_{1},M_{2}}(u) = \frac{i}{2} \sum_{\alpha \leq M_{1},\beta \leq M_{2}} u_{\alpha\beta} : F_{\alpha} G_{\beta} :$$

$$-2C_{0,M_{1},...,M_{7}}(f) = C_{0;3;M_{1},...,M_{3}}(f) + C_{0;4;M_{4},...,M_{7}}(f)$$

$$= 2^{1/2} \sum_{\alpha \leq M_{1},\beta \leq M_{2},\gamma \leq M_{3}} f_{\alpha\beta\gamma} : F_{\alpha} F_{\beta} G_{\gamma} :$$

$$+\frac{1}{2} \sum_{\alpha \leq M_{4},\beta \leq M_{5},\gamma \leq M_{6},\delta \leq M_{7}} f_{\alpha\beta\gamma\delta} : F_{\alpha} F_{\beta} G_{\gamma} G_{\delta} :$$
(3.58)

be the expansions of F, G in terms of the e_I , the sums truncated as displayed. Thus 1. each occurring sum has now finite range and 2. individual normal ordered monomials have individual finite ranges. We can even take one step further and instead of summing over the same $\alpha \leq M$ for A_{α} , A_{α}^* occurring in G_{α} or F_{α} we can sum over different labels. This results in the explicit expressions

$$\begin{split} G_{M_1,...,M_8}(r) &= G_{1;M_1}^0(r) + G_{1;M_2}^1(r) + G_{2;M_3,M_4}^0(r) + G_{2;M_5,M_6}^1(r) + G_{2;M_7,M_8}^2(r) \\ G_{1;M_1}^0(r) &= -\frac{i}{2^{1/2}} \sum_{\alpha \leq M_1} r_{\alpha} A_{\alpha}, \ G_{1;M_2}^1(r) = \frac{i}{2^{1/2}} \sum_{\alpha \leq M_2} r_{\alpha} A_{\alpha}^* \\ G_{2,M_3,M_4}^0 &= \frac{i}{2} \sum_{\alpha \leq M_3,\beta \leq M_4} r_{\alpha\beta} A_{\alpha} A_{\beta} \\ G_{2,M_5,M_6}^1 &= \frac{i}{2} \sum_{\alpha \leq M_5,\beta \leq M_6} [r_{\alpha\beta} - r_{\beta\alpha}] A_{\alpha}^* A_{\beta} \\ G_{2,M_7,M_8}^1 &= \frac{i}{2} \sum_{\alpha \leq M_7,\beta \leq M_8} r_{\alpha\beta} A_{\alpha}^* A_{\beta}^* \\ C_{M_1..M_6}(u) &= C_{2;M_1,M_2}^0(u) + C_{2;M_3,M_4}^1(u) + C_{2;M_5,M_6}^2(u) \\ C_{2;M_1,M_2}^0(u) &= \frac{i}{2} \sum_{\alpha \leq M_1,\beta \leq M_2} u_{\alpha\beta} A_{\alpha} A_{\beta}, \ C_{1;M_3,M_4}^1(u) = \frac{i}{2} \sum_{\alpha \leq M_3,\beta \leq M_4} [-u_{\alpha\beta} + u_{\beta\alpha}] A_{\alpha}^* A_{\beta} \\ C_{2;M_5,M_6}^2(u) &= \frac{i}{2} \sum_{\alpha \leq M_7,\beta \leq M_8} u_{\alpha\beta} A_{\alpha}^* A_{\beta}^*, \\ -2C_{0,M_1,...,M_{28}}(f) &= \sum_{k=0}^3 C_{0;3;M_{3k+1},M_{3k+2},M_{3k+3}}(f) + \sum_{k=0}^4 C_{0;4;M_{12+4k+1},M_{12+4k+2},M_{12+4k+3},M_{12+4k+4}}(f) \\ C_{0;3;M_1,M_2,M_3}^0(f) &= 2^{1/2} \sum_{\alpha \leq M_{1,\beta} \leq M_{2,\gamma} \leq M_3} f_{\alpha\beta\gamma} A_{\alpha} A_{\beta} A_{\gamma}, \ ... \\ C_{0;4;M_{13}..M_{16}}^0(f) &= \frac{1}{2} \sum_{\alpha \leq M_{13,\beta} \leq M_{14,\gamma} \leq M_{15,\delta} \leq M_{16}} f_{\alpha\beta\gamma\delta} A_{\alpha} A_{\beta} A_{\gamma} A_{\delta}^* A_{\delta}^* \\ C_{0;4;M_{13}..M_{16}}^0(f) &= \frac{1}{2} \sum_{\alpha \leq M_{13,\beta} \leq M_{14,\gamma} \leq M_{16,\delta} \leq M_{16}} f_{\alpha\beta\gamma\delta} A_{\alpha}^* A_{\beta}^* A_{\gamma}^* A_{\delta}^* \\ C_{0;4;M_{25}..M_{28}}^0(f) &= \frac{1}{2} \sum_{\alpha < M_{25,\beta} \leq M_{26,\gamma} \leq M_{26,\gamma} \leq M_{26}} f_{\alpha\beta\gamma\delta} A_{\alpha}^* A_{\beta}^* A_{\gamma}^* A_{\delta}^* \\ C_{0;4;M_{25}..M_{28}}^0(f) &= \frac{1}{2} \sum_{\alpha < M_{25,\beta} \leq M_{26,\gamma} \leq M_{26}} f_{\alpha\beta\gamma\delta} A_{\alpha}^* A_{\beta}^* A_{\gamma}^* A_{\delta}^* \\ C_{0;4;M_{25}..M_{28}}^0(f) &= \frac{1}{2} \sum_{\alpha < M_{25,\beta} \leq M_{26,\gamma} \leq M_{26,\gamma} \leq M_{26}} f_{\alpha\beta\gamma\delta} A_{\alpha}^* A_{\beta}^* A_{\gamma}^* A_{\delta}^* \\ C_{0;4;M_{25}..M_{28}}^0(f) &= \frac{1}{2} \sum_{\alpha < M_{25,\beta} \leq M_{26,\gamma} < M_{27,\gamma} \leq M_{26}} f_{\alpha\beta\gamma\delta} A_{\alpha}^* A_{\beta}^* A_{\gamma}^* A_{\delta}^* \\ C_{0;4;M_{25}..M_{28}}^0(f) &= \frac{1}{2} \sum_{\alpha < M_{25,\beta} \leq M_{26,\gamma} < M_{27,\gamma} \leq M_{26}} f_{\alpha\beta\gamma\delta} A_{\alpha}^* A_{\beta}^* A_{\gamma}^* A_{\delta}^* \\ C_{0;4;M_{25}...M_{28}}^0(f) &= \frac{1}{2} \sum_{\alpha < M_{25,\beta} < M_{26,\gamma} < M_{27,\gamma} < M_{25}} \\ C_{0;4;M_{25}...M_{28}}^0$$

Here the superscript k and subscript N e.g. in $G_{N;M_1..M_N}^k(r)$; N = 1, 2; k = 0, 1, .., N means that we are looking at a normal ordered monomial of order N with k creation and N - k annihilation operators. For each such monomial we have numerous ways of taking the $M_1, .., M_N \to \infty$.

Definition 3.1.

Let for $0 \leq k \leq N \in \mathbb{N}$

$$E_{N;M_1,..,M_N}^k := \sum_{\alpha_1 \le M_1,..,\alpha_N \le M_N} E_{\alpha_1..\alpha_N}^k A_{\alpha_1}^* .. A_{\alpha_k}^* A_{\alpha_{k+1}} .. A_{\alpha_N}$$
(3.60)

be an N-th order normal ordered monomial in k creation operators and N - k annihilation operators. i.

A limiting pattern P in $1 \leq l \leq N$ indices is an ordered partition of $\{1, ..., N\}$ into subsets $U_1, ..., U_l$ i.e. $\cup_{r=1}^l U_r = \{1, ..., N\}, U_r \cap U_s = \emptyset \forall r \neq s.$ *ii.*

A limiting process subordinate to a limiting pattern P is the coincidence of cut-offs $M^r := M_{s_1} = ... = M_{s_{L_r}}$ for $U_r = \{s_1, ..., s_{L_r}\}$ followed by a sequential limit: First take $M^1 \to \infty$ at finite $M^2, ..., M^l$, then take $M^2 \to \infty$ at finite $M^3, ..., M^l$, ... and finally take $M^l \to \infty$. This precise process is denoted by $P \to \infty$. iii.

The extreme cases are the total coincidence limit l = 1 with $M^1 = M_1 = ... = M_N \to \infty$ and one of the N! total individual limits l = N defined by a permutation π in N elements i.e. $M_{i_{\pi(k)}} \to \infty$ at finite $M_{\pi(k+1)}, ..., M_{\pi(N)}$ in the order k = 1, 2, ..., N.

iv.

The quadratic form limit $E_{N;P}^k$, if it exists, subordinate to a limiting pattern P is the weak operator topology limit

$$\langle \psi, E_P^k \psi' \rangle := \lim_{P \to \infty} \langle \psi, E_{N;M_1,\dots,M_N}^k \psi' \rangle :$$

$$(3.61)$$

for all $\psi, \psi' \in \mathcal{D}$.

v.

A K-th order normal ordered polynomial is a sum of terms of the form (3.60) with $0 \le N \le K$, $0 \le k \le N$ each carrying its own $M_1^{(N)}, ..., M_N^{(N)}$. A limiting pattern and process applied to the polynomial is the sum of the individual monomial limiting patterns and monomials.

Note that the resulting $E_{N;P}^k$ is defined independently of the choice of ψ, ψ' in (3.61), i.e. the limiting process does not depend on those, so that the resulting object is indeed a quadratic form, i.e. a sesqui-linear form with form domain \mathcal{D} .

Various generalisations of the limiting pattern are conceivable and should be kept in mind for potential future applications:

1.

We could refine the pattern and make it also dependent on all the finite indices m_k, i_k of the $\alpha_k = (I_k, m_k, i_k), k = 1, ..., N$.

2.

Instead of a sequential limit we could consider coincidence limits at different "rates", say $M^r = \exp(c_r m), m \to \infty, c_r \in \mathbb{R}_+$ for some of the r or a mixture of those with sequantial limits etc. 3.

An expression of the form (3.60) could be split into several of the same form with different coefficients $E_{\alpha_1,..,\alpha_N;s}^k$, s = 1,..,S and each of them could be treated individually with its own limiting pattern.

For the purpose of this paper, the above, finite number of patterns will be sufficient. Then we have

Proposition 3.1.

The objects (3.55) are quadratic forms on the Fock space with dense form domain \mathcal{D} obtained from (3.59) or (3.58) via the weak limits (3.61) subordinate to any limiting pattern P, i.e. all those limits corresponding to different P coincide.

Note that the monomials of different order N or of different number of creation operators k in (3.59) are subjected to the limiting process individually, e.g. the first two terms of order N = 1 in the Gauss constraint are treated in the sense of (3.61) independently of each other and independently of any of the three terms of order N = 2 in accordance with item v. of above definition. The same applies for the various N = 3 contributions among themselves, the various N = 4 contributions among themselves and mutually independently of each other in the Hamiltonian constraint. The expression (3.58) is a special case of (3.59) in which we identify the cut-offs of those labels α , β respectively that originate from the same F_{α} , G_{β} respectively.

Proof. We sandwich (3.59) between fixed vectors $\psi, \psi' \in \mathcal{D}$. Then we find $M_0 \in \mathbb{N}$ such that $\psi, \psi \in \mathcal{D}_{M_0}$, the finite linear span of Fock states satisfying $n_I = 0$ for $I \notin S_{M_0}$. For each monomial of order N = 1, 2, 3, 4that appears in (3.59) we let annihilators act to the right on ψ' and creators act to the left on ψ . Following the pattern, we start with the the $\alpha's$ corresponding to U_1 to take full range. As the annihilators respectively creators commute among each other respectively, we can take those annihilators and creators to the outmost right and left respectively. Whenever such α violates $\alpha \leq M_0$ the corresponding term in the sum vanishes. In this way all those α are confined to $\alpha \leq M_0$. We proceed the same way with U_2 etc. Remarks:

Α.

Note that we could have considered also labelling the truncated Gauss constraint by only two cut-offs M_1, M_2 where M_1 labels the first order monomial and M_1, M_2 label the second order monomial. The same applies to the truncated Hamiltonian constraint which could be labelled by only four cut-offs $M_1, ..., M_4$ where only the first three label the third order monomial. This corresponds to a certain coincidence limit of cut-off labels between quadratic forms of different monomials order and is much more restrictive than item v. of definition 3.1. Similarly the labelling (3.58) corresponds to a coincidence limit that is obtained by expanding the continuum objects $F = A - A^*, G = A + A^*$ with respect to the e_I thereby obtaining $F_\alpha = A_\alpha - A^*_\alpha, G_\alpha = A_\alpha + A^*_\alpha$ and then decomposing the quadratic form into the A_α, A^*_α so obtained while the more general scheme first decomposes the quadratic form into the continuum A, A^* and then expands each factor A, A^* individually with respect to the e_I in each monomial. This therefore corresponds to a coincidence limit between cut-off labels of quadratic forms of the same monomial order but different numbers of creation operators $F_\alpha = A_\alpha - A^*_\alpha, G_\alpha = A_\alpha + A^*_\alpha$. which is also much more restrictive than item v. of the same definition. While for the objects (3.59) these coincidence limits exists and can be shown to coincide with the above limit these coincidental limits offer less flexibility in the choice of limiting patterns which becomes vital in the commutator calculations below. B.

In [31] we have applied truncation structures and limiting patterns already to the spatial diffeomorphism constraint. There the mixture issue mentioned in A. with respect to N does not arise as this is a monomial N = 2constraint and we have applied the same total coincidence limit $M_1 = M_2 = M \rightarrow \infty$ individually for all three occurring normal ordered monomials of the form $A^*_{\alpha} A^*_{\beta}$, $A^*_{\alpha} A_{\beta}$, $A_{\alpha} A_{\beta}$ i.e. we have taken a coincidence limit with respect to the k = 0, 1, 2 labelling. What is more, when computing the commutator as defined below, we not only took $M_1 = M_2 = M$ for the first SDC quadratic form and $M'_1 = M'_2 = M'$ for the second SDC quadratic form but even M = M'. This is not necessary and just one of the possibilities of taking the limit, leading however to the same unambiguous result in this case.

When computing commutators of the above constructed quadratic forms we need the following

Lemma 3.1.

We have

$$\begin{bmatrix} A_{\alpha_{1}}^{*}..A_{\alpha_{k}}^{*} \ A_{\beta_{1}}..A_{\beta_{l}}, A_{\gamma_{1}}^{*}..A_{\gamma_{p}}^{*} \ A_{\delta_{1}}..A_{\delta_{q}} \end{bmatrix}$$

$$= A_{\alpha_{1}}^{*}..A_{\alpha_{k}}^{*} \ \begin{bmatrix} A_{\beta_{1}}..A_{\beta_{l}}, A_{\gamma_{1}}^{*}..A_{\gamma_{p}}^{*} \end{bmatrix} A_{\delta_{1}}..A_{\delta_{q}} - A_{\gamma_{1}}^{*}..A_{\gamma_{p}}^{*} \ \begin{bmatrix} A_{\delta_{1}}..A_{\delta_{q}}, A_{\alpha_{1}}^{*}..A_{\alpha_{k}}^{*} \end{bmatrix} A_{\beta_{1}}..A_{\beta_{l}}$$

$$(3.62)$$

and

$$[A_{\beta_1}..A_{\beta_l}, A^*_{\gamma_1}..A^*_{\gamma_p}] = ([A_{\gamma_1}..A_{\gamma_p}, A^*_{\beta_1}..A^*_{\beta_l}])^*$$
(3.63)

and for $k \leq l$

$$[A_{\alpha_1}..A_{\alpha_k}, A^*_{\beta_1}..A^*_{\beta_l}] = \sum_{n=1}^k \sum_{1 \le r_1 < .. < r_n \le k; 1 \le s_1 < .. < s_n \le l} [\sum_{\pi \in S_n} \prod_{m=1}^n \delta_{\alpha_{r_{\pi(m)}}\beta_{s_m}}] [\prod_{s \notin \{s_1,..,s_n\}} A^*_{\beta_s}] [\prod_{r \notin \{r_1,..,r_n\}} A_{\alpha_r}]$$
(3.64)

The proof is not difficult: (3.62) follows directly from the basic CCR, (3.63) directly from the definition of the adjoint which allows us to assume $k \leq l$ in (3.64) and (3.64) follows by induction over k. Lemma 3.1 gives the explicit formula for restoring normal order in commutators of normal ordered monomials.

Suppose now that we have quadratic forms

$$Q_j = \sum_{N=0}^{N_j} \sum_{k=0}^{N} [Q_j]_N^k, \ [Q_j]_N^k := [Q_j]_{N;\alpha_1..\alpha_N}^k A_{\alpha_1}^*..A_{\alpha_k}^* A_{\alpha_{k+1}}..A_{\alpha_N}$$
(3.65)

with j = 1, 2 which as in (3.55) are uniquely defined using any limiting pattern so that the formal infinite range sums are in fact finite in any matrix element calculation. To compute the quadratic form commutator $[Q_1, Q_2]$,

if it exists, we have three possibilities.

Possibility 1: Truncating after commuting

We use the the expressions (3.65) with all labels α unconstrained, compute the commutator $[Q_1, Q_2]$ term by term corresponding to the terms labelled by N, k and N', k' respectively with $N \leq N_1, k \leq N, N' \leq N_2, k' \leq N'$ respectively at fixed labels α and restore normal order using lemma 3.1. Then for each normal ordered monomial that is produced we individually truncate the sums over all occurring α 's in that normal ordered monomial and finally for each of those monomials investigate the possible limiting patterns which result in well defined quadratic forms.

Possibility 2: Truncating before commuting

We use expression (3.65) and truncate the sums over $\alpha_1, ..., \alpha_N$ in the monomial corresponding to N, k by $\alpha_l \leq M_{j,N,k,l}$; j = 1, 2; $N = 1, ..., N_j$ k = 0, ...N; $l = 1..., N_j$, then we compute the commutator of $[Q_{1,\{M_1\}}, Q_{2,\{M_2\}}]$ of the so truncated quadratic forms where $\{M_j\}$ stands for the collection of those $M_{j,N,k,l}$, restore normal order using lemma 3.1 and finally investigate the possible limiting patterns that result in a well defined quadratic form.

Possibility 3: Mixture

One applies possibility 1 in the sense that one defines

$$[Q_1, Q_2] := \sum_{N=0, N'=0}^{N_1, N_2} \sum_{k, k'=0}^{N, N'} [[Q_1]_N^k, [Q_2]_{N'}^{k'}]$$
(3.66)

and then applies possibility 2 to each term in the finite sum (3.66).

The difference between the first two possibilities is that in the first possibility we have more truncation lables than in the second possibility. This is because in the second possibility only the original truncation labels $\{M_1\}, \{M_2\}$ appear in all possible monomials that are produced by the normal ordering identity (3.64) while in the first possibility each term on the right hand side of (3.64) gets equipped with its own individual truncation labels. Thus the first possibility allows for more limiting patterns and thus more possibilities to generate a well defined quadratic form. For the same reason it potentially also generates more ambiguity as there may be more than one way to generate a well defined quadratic form. One may argue for the first possibility by taking the point of view that the non-truncated quadratic forms Q_1, Q_2 are well defined, thus the commutator should be based on those and the afterwards termwise truncation in the decomposition into normal ordered expressions is part of the regularisation procedure for the commutator just as it is for any quadratic form by definition 3.1 as it was also used to defined Q_1, Q_2 . One may argue for the second possibility by taking the point of view that products of quadratic forms are ill-defined and thus they should be truncated before computing the commutator. That each term $[Q_j]_N^k$ is equipped with its individual truncation labels $M_{j,N,k,l}$ is justified by the assumption that all limiting patterns on $\{M_j\}$ result in the same quadratic form Q_j . Finally the mixed point of view can be justified when each term $[Q_j]_N^k$ by itself is a well defined quadratic form so that each of them by itself is obtained via any of the limiting patterns. Thus (3.66) is a possible definition if each of the individual $[[Q_1]_N^k, [Q_2]_{N'}^{k'}]$ can be turned into a well defined quadratic form.

Since for each of these prescriptions and limiting patterns the unconstrained $[Q_1, Q_2]$ is formally obtained, all of these possibilities appear to be equally justified a priori. It is similar to the problem of giving meaning to a formal multiple sum series of complex numbers (in our case obtained as matrix elements) which is not absolutely convergent but may have several sequential conditionally convergent sums. Thus giving a concrete conditional summing prescription is *part of the definition* of such a series.

Example 3.1.

Let for $m, n \in \mathbb{N}_0$ $a_{m,4n} := [m+1+4n]^{-1}, \ a_{m,4n+1} := -[m+1+4(n+1)]^{-1}, \ a_{m,4n+2} := -[m+2+4n]^{-1}, \ a_{m,4n+3} := [m+2+4(n+1)]^{-1}$ (3.67) Then the series $s := \sum_{(m,n) \in \mathbb{N}_0^2} a_{m,n}$ is ill defined as it stands as it is not absolutely convergent. The row sums

$$\sum_{m} a_{m,n} = \pm \infty \tag{3.68}$$

are plain divergent. The column sums are conditionally convergent

$$\sum_{n} a_{m,n} := \sum_{n=0}^{\infty} \left\{ \frac{1}{m+1+4n} - \frac{1}{m+1+4(n+1)} \right\} - \sum_{n=0}^{\infty} \left\{ \frac{1}{m+2+4n} - \frac{1}{m+2+4(n+1)} \right\}$$
$$= 4\sum_{n=0}^{\infty} \frac{1}{(m+1+4n)(m+1+4(n+1))} - 4\sum_{n=0}^{\infty} \frac{1}{(m+2+4n)(m+2+4(n+1))}$$
$$= \frac{1}{m+1} - \frac{1}{m+2} = \frac{1}{(m+1)(m+2)}$$
(3.69)

and thus the sequential and conditional sum

$$s := \sum_{m} \sum_{n} a_{m,n} = 1$$
 (3.70)

converges in this limiting pattern.

Note that this example is smilar to but different from examples illustrating Riemann's rearrangement theorem [35] which states that a conditionally convergent series $\sum_N c_N$ of real numbers can be made to converge to any real number by using a bijection $b : \mathbb{N} \to \mathbb{N}$; $N \mapsto b(N)$ on the indices and considerung $\sum_N c_{b(N)}$ instead. While \mathbb{N}_0^2 is in bijection with \mathbb{N} (e.g. via Cantor's map [35] B) and we can thus consider above $a_{m,n} = a_{B(N)} = c_N$ as a series in c_N , summing rows or columns first does not correspond to $N \to \infty$ but rather to taking limits of subseries of c_N and then summing the limits of those subseries. The rearrangement theorem however can be applied to those subsequences which in the above example are also only conditionally convergent.

In what follows we explore the mixed possibility 3, i.e. we will treat linear and bi-linear terms in the Gauss constraint as well as the tri-linear and quadri-linear terms in the Hamiltonian constraint as individually defined quadratic forms and then consider for those truncation before commutation. Moreover, we will constrain the limiting patterns to be such that in normal ordered expressions of the form : $F_{\alpha_1}..F_{\alpha_k}G_{\alpha_{k+1}}..G_{\alpha_N}$: we truncate $\alpha_k \leq M_k$. Definition 3.1 allows to be more general, e.g. in the normal ordered decomposition

$$\sum_{\alpha,\beta} E_{\alpha\beta} : F_{\alpha}G_{\beta} := \sum_{\alpha,\beta} E_{\alpha\beta}A_{\alpha}A_{\beta} + \left[-E_{\alpha\beta} + E_{\beta\alpha}\right]A_{\alpha}^*A_{\beta} - E_{\alpha\beta}A_{\alpha}^*A_{\beta}^*$$
(3.71)

we could truncate α, β in the the first, second, third term respectively by M_1, M_2 and M_3, M_4 and M_5, M_6 respectively instead of synchronising $M_1 = M_3 = M_5, M_2 = M_4 = M_6$. This is exactly the difference between (3.58) and (3.59). Thus we choose to work with (3.58). The advantage will be a much reduced computational effort using the fact that $[F_{\alpha}, F_{\beta}] = [G_{\alpha}, G_{\beta}] = 0$.

Proposition 3.2.

Consider the truncated constraint ingredients

$$G_{1;M_1}(r), \ G_{2;M_1,M_2}(r), \ C_{2;M_1,M_2}(u), \ C_{0;3,M_1..M_3}(f), \ C_{0;4,M_1..M_4}(f)$$

$$(3.72)$$

defined in (3.58) and define the constraint commutators by

$$\begin{split} [G(r),G(s)] &:= \frac{1}{2} \left\{ [G_1(r),G_1(s)] - [G_1(s),G_1(r)] \right\} + \frac{1}{2} \left\{ [G_2(r),G_2(s)] - [G_2(s),G_2(r)] \right\} \\ &+ \left\{ [G_1(r),G_2(s)] - [G_1(s),G_2(r)] \right\} \\ [C(u),C(v)] &:= \frac{1}{2} \left\{ [C_2(u),C_2(v)] - [C_2(v),C_2(u)] \right\} \\ [C(u),G(r)] &:= \left\{ [C_2(u),G_1(r)] \right\} + \left\{ [C_2(u),G_2(r)] \right\} \\ [G(r),C_0(f)] &:= \left\{ [G_1(r),C_{0,3}(f)] \right\} + \left\{ [G_2(r),C_{0,3}(f)] \right\} + \left\{ [G_1(r),C_{0,4}(f)] \right\} + \left\{ [G_2(r),C_{0,4}(f)] \right\} \\ [C(u),C_0(f)] &:= \left\{ [C_2(u),C_{0,3}(f)] \right\} + \left\{ [C_2(u),C_{0,4}(f)] \right\} \\ [C_0(f),C_0(g)] &:= \frac{1}{2} \left\{ [C_{0,3}(f),C_{0,3}(g)] - [C_{0,3}(g),C_{0,3}(f)] \right\} + \frac{1}{2} \left\{ [C_{0,4}(f),C_{0,4}(g)] - [C_{0,4}(g),C_{0,4}(f)] \right\} \\ &+ \left\{ [C_{0,3}(f),C_{0,4}(g)] - [C_{0,3}(g),C_{0,4}(f)] \right\} \end{split}$$

$$(3.73)$$

Consider each curly bracket expression on the right hand side of (3.73) individually. Each such curly bracket involves either one or two commutators. If one commutator appears, we just replace the quadratic forms by their correspondent in the list (3.72). E.g. $[C_2(u), C_{0;3}(f)]$ is replaced by $[C_{2;M_1,M_2}(u), C_{0;3;M_3,...,M_5}(f)]$ with arbitrary finite cut-offs $M_1...M_5$. If two commutators appear we replace the quadratic forms also by the correspondent in the list (3.72) but we use the same cut-offs in both commutators. For example we replace $[C_{0;3}(f), C_{0;4}(g)] - f \leftrightarrow g$ by $[C_{0;3;M_1,...,M_3}(f), C_{0;4;M_4,...,M_7}(g)] - f \leftrightarrow g$ with arbitrary finite cut-offs $M_1, ..., M_7$.

Then for each such curly bracket there exists a limiting pattern such that 1. the limiting process results in a well defined quadratic form and 2. those resulting quadratic forms add up exactly to the normal ordered quadratic forms corresponding to the Fock quantisations of the right hand sides of the corresponding classical Poisson bracket calculations. That is

$$[:Z_1:,:Z_2:] = i : \{Z_1, Z_2\} :$$
(3.74)

where Z_1, Z_2 are any of the $G(r), C(u), C_0(f)$.

Before going through the straightforward but tedious and lengthy proof a few remarks are in order:

1. Sources of divergences in commutator calculations and mechanisms to avoid them

When we use (3.64) to restore normal order in a commutator we end up with normal ordered monomials that come with coefficients D that are products of the coefficients of the quadratic forms that we compute the commutator of, times products of Kronecker symbols. Thus such a term is of the form

$$\sum_{\alpha_1 \le M_1..\alpha_L \le M_L} D_{\alpha_1..\alpha_L} \prod_{k=1}^n \delta_{\alpha_{r_k},\alpha_{s_k}} : \prod_{t \in \{1,..,L\} - \{r_1,..,r_n,s_1,..,s_n\}} C_{\alpha_s \sigma_s} :$$
(3.75)

where $\sigma = \pm, C_{\alpha,+} = A_{\alpha}^*, C_{\alpha,-} = A_{\alpha}$ and $\{r_1, ..., r_n, s_1, ..., s_n\} \subset \{1, ..., L\}$ with $2n \le L$ and $n \ge 1$. Clearly the labels α_t , $t \notin \{r_1, ..., r_n, s_1, ..., s_n\}$ are automatically truncated when sandwiching (3.75) between Fock states. Thus it is the sum over the remaining labels that is in danger of diverging. It turns out that only terms with $n \ge 2$ are dangerous. This can be seen from the typical example $D_{\alpha_1..\alpha_4} = \langle e_{I_1}e_{I_2}, f \rangle \langle e_{I_3}e_{I_4}, g \rangle$ where f, g are smearing functions of the constraints.

If we now perform the sum over $I_1 = I_3$ then by the completeness relation we end up with $\langle e_{I_2} e_{I_4}, f g \rangle$ which is finite. If we perform a second sum over $I_2 = I_4$ then we end up with $\langle [\sum_I e_I^2], fg \rangle$ which diverges because $\sum_I e_I(x) e_J(y) = \delta(x, y)$ so that

$$\Lambda(x) := \lim_{M \to \infty} \sum_{I \in S_M} e_I(x)^2 \tag{3.76}$$

diverges being formally equal to $\delta(x, x)$. Such diverging sequences appear all over the place in the *product* of quadratic forms. However, the commutator maybe better behaved because before we sum over $I_2 = I_4$ there is

a second term $\langle e_{I_2} e_{I_4}, g f \rangle$ which comes from the fact that $[Q(f), Q(g)] = \frac{1}{2}([Q(f), Q(g)] - [Q(g), Q(f)])$ and these two terms would in fact cancel (so called ultra locality). This is the reason for why we considered the curly bracket expressions with two commutators and equally chosen cut-offs so that such a cancellation can occur. This also shows why a sequential limiting pattern is of advantage.

On the other hand, we may have a term with a derivative of a smearing function $D_{\alpha_1..\alpha_4} = \langle e_{I_1}e_{I_2}, f \rangle \langle e_{I_3}e_{I_4}, g_{,j} \rangle$. Performing the first sum still delivers a finite result $\langle e_{I_2} e_{I_4}, f g_{,j} \rangle$ but subtracting the term with f, g interchanged in this case does not vanish but yields $\langle e_{I_2} e_{I_4}, [f g_{,j} - f_{,j}g] \rangle$. Now performing the second sum again gives $\langle \sum_I e_I^2 [f g_{,j} - f_{,j}g] \rangle$ which diverges. Thus, to avoid such divergences one will try to find a limiting pattern which avoids derivatives of smearing functions f, g of the constraints. For instance in

$$\langle e_I f, e_{J,j} \rangle \langle e_K g, e_L \rangle \delta_{IK} \delta_{JL} - f \leftrightarrow g$$

$$(3.77)$$

carrying out the sum over I = K first gives zero by the same mechanism as above while carrying out the sum over J = L first gives (integrations by parts do not create boundary terms because the e_I are Schwartz functions for $\sigma = [R]^3$ or all functions are periodic for $\sigma = T^3$)

$$- < [e_I f]_{,j} e_K, g > \delta_{IK} - f \leftrightarrow g = < [e_I e_K, [g_{,j} f - f_{,j} g] > \delta_{IK}$$
(3.78)

and the sum over I = K is again divergent. This demonstrates that the existence or non existence of the commutator as a quadratic form can depend rather drastically on the limiting pattern which is reminiscent of the situation with double sum series mentioned in example 3.1.

Now the observation is that the constraints except for the linear term in the Gauss constraint which due to its linearity cannot produce more than one Kronecker symbol and thus does not suffer from the above danger, all constraints contain the E_j^a witout derivatives (if we use $F_{ab}^j E_j^a = C_a + A_a^j G_j$ instead of C_a). Thus a promising limiting pattern is such that one carries out the sum over those α that label the F_{α} before carrying out the sum over those α that label the G_{α} which do come with derivatives. This strategy turns out to be successful here and may carry over to the Lorentzian constraints in all dimensions formulated in terms of ADM variables P, q where only q carries (up to second order) derivatives but not P.

2. General strategy of the computation

Step 1: At finite cut-offs we simply decompose a term of the form

$$[:F_{\alpha_1}..F_{\alpha_p} \ G_{\beta_{p+1}}..G_{\beta_m} :, \ F_{\beta_1}..F_{\beta_q} \ G_{\beta_{q+1}}..G_{\beta_n}]$$
(3.79)

into annihilation and creation operators. This yields 2^{m+n} terms of the form

$$[A_{\gamma_1}^*..A_{\gamma_k}^* A_{\gamma_{k+1}}..A_{\gamma_m}, A_{\delta_1}^*..A_{\delta_l}^* A_{\delta_{l+1}}..A_{\delta_n}]$$
(3.80)

where $0 \le k \le m$, $0 \le l \le n$ and the γ, δ respectively are permutations of the α, β labels respectively.

Step 2: Then we apply lemma 3.1 to (3.78). For each $u = 1, ..., \min(m - k, l)$ this gives $\binom{m-k}{u} \binom{l}{u}$ and for each $u = 1, ..., \min(k, n - l)$ this gives $\binom{k}{u} \binom{n-l}{u}$ normal ordered products of k + l - u creation and m + n - k - l - u annihilation operators respectively with coefficients that are totally symmetric linear combinations of u! products of u Kronecker symbols.

Step 3: We now substitute $A_{\alpha} = [G_{\alpha} + F_{\alpha}]/2$, $A_{\alpha}^* = [G_{\alpha} - F_{\alpha}]/2$ within the normal ordering symbol which gives 2^{m+n-2u} terms at given u in step 2. Since within the normal ordered symbol we can permute all F, G operators, we can rearrange the commutator into normal ordered monomials of the F, G with coefficients which involve at least one and at most min(m, n) products of Kronecker symbols of the form $\delta_{\alpha_i,\beta_j}$, i = 1, ..., m; j = 1, ..., n. Note that for m = n, p = q there will be no term with n factors of Kronecker symbols. This is because, suppressing indices, in $[:F^pG^{m-p}:]:F^pG^{m-p}:]$ the contributions to δ^m are coming from

$$[A^m + (-1)^p (A^*)^m, A^m + (-1)^p (A^*)^m] = (-1)^p ([\operatorname{symm} \otimes^m \delta] - [\operatorname{symm} \otimes^m \delta]^T) = 0$$
(3.81)

where symm means total symmetrisation and T transposition.

Step 4: To estimate the number of terms that one has to manipulate in these first 3 steps we note that of the 2^{m+n} terms obtained from step 1 there are $\binom{k}{m}\binom{l}{n}$ terms of the type (3.81), hence we obtain

$$N_{m,n} = \sum_{k=0}^{m} \sum_{l=0}^{n} \binom{m}{k} \binom{n}{l} \left[\sum_{u=1}^{\min(m-k,l)} u! \binom{m-k}{u} \binom{l}{u} 2^{m+n-2u} + \sum_{u=1}^{\min(n-l,k)} u! \binom{n-l}{u} \binom{k}{u} 2^{m+n-2u} \right]$$
(3.82)

terms in total. To estimate the number of these terms we approximate the binomial distribution by the Poisson distribution $2^{-k} \binom{k}{u} \approx e^{-1/2} \frac{2^{-u}}{u!}$. This allows to estimate the sums over u by extending to $u = \infty$. Then

$$N_{m,n} \approx \sum_{k=0}^{m} \sum_{l=0}^{n} \binom{m}{k} \binom{n}{l} 2^{m+n} e^{-1} \left[\sum_{u=0}^{\infty} \frac{2^{-4u}}{u!}\right] [2^{m-k+l} + 2^{n-l+k}]$$

= $2^{m+n} e^{15/16} \sum_{k=0}^{m} \sum_{l=0}^{n} \binom{m}{k} \binom{n}{l} [2^{m-k+l} + 2^{n-l+k}]$ (3.83)

Once again substituting for the Poisson distribution allows to perform the sum over k, l which we extend to ∞

$$N_{m,n} \approx = 2^{m+n} e^{-31/16} \sum_{k=0}^{m} \sum_{l=0}^{n} [2^{m-k+l} + 2^{n-l+k}] \frac{2^{m-k}}{k!} \frac{2^{n-l}}{l!}$$

$$\approx 4^{m+n} e^{-31/16} \sum_{k=0}^{m} \sum_{l=0}^{n} [2^{-2k} + 2^{-2l}] \frac{1}{k! l!} = 24^{m+n} e^{-13/16}$$
(3.84)

Thus we obtain an order of 4^{m+n} operations.

With an eye towards the computation for Lorentzian signature in q, P variables with top polynomial degree m = n = 10 in 3 spatial dimensioons this yiels 10^{12} terms to consider. For the present Euclidian calculation we have top degree m = n = 4 which yields merely 10^5 terms to consider. Yet, the final expression in terms of normal ordered monomials of F, G will contain only a small number of linearly independent terms. This is due to the fact that the original expression (3.79) is totally symmetric in the four groups of indices corresponding to the p, q factors of F respectively and the m - p, n - q factors of G respectively. While in the present case a calculation by hand is still possible and was indeed done by the author, it is still a good idea to use computer algebra tools to verify the calculations. In the present case we wrote a simple, not yet speed optimised code using pthon's SymPy package for symbolic calculus which in principle can deal with any m, n. For m = n = 4 the computation takes a few seconds with a 2.3MHz Intel core i5 processor. Therefore the computation will definitely be doable in Lorentzian signature in (p, q) variables using parallel computing which on the author's machine would momentarily take half a year.

Step 5: The crucial step is now to show that there exists at least one limiting pattern such that the terms involving more than one factor of Kronecker δ factors yield a finite result in the limiting process subordinate to that pattern. Note that the classical Poisson bracket calculation gives precisely one factor of Kronecker δ and therefore this term returns the result of the classical calculation automatically normal ordered. The chosen pattern must be the same for the terms involving $1, 2, ..., \min(m, n) - \delta_{m,n}$ factors of Kronecker δ because we have chosen truncation before commuting but individually for each F, G monomial into which a constraint decomposes which we regard as quadratic forms in their own right.

Proof.:

Organisation:

We compute commutators in increasing order of complexity and aim to show that there exist limiting pattern such that (all constraints are normnal ordered by definition)

- 1. Gauss-Gauss: $[G(r), G(s)] = i G(-r \times s).$
- 2. Diffeomorphism-Diffeomorphism: [C(u), C(v)] = i C(-[u, v]).
- 3. Diffeomorphism-Gauss: [C(u), G(r)] = i G(-u[r]).
- 4. Gauss-Hamiltonian: $[G(r), C_0(f)] = 0.$
- 5. Diffeomorphism-Hamiltonian: $[C(u), C_0(f)] = i H(-u[f]).$
- 6. Hamiltonian-Hamiltonian

$$[C_0(f), C_0(g)] = i \int d^D x [4(fg_{,a} - f_{,a}g)] : Q^{ab} F^j_{bc} E^b_j :$$
(3.85)

The last relation is, like the others, exactly *i* times the normal ordering of the classical Poisson bracket expression and thus *non-anomalous*. However, while $Q^{ab}F^{j}_{bc}E^{b}_{j} = Q^{ab} [C_b + A^{j}_bG_j]$ is a linear combination of constraints with second order and third order monomial structure functions we have

$$: Q^{ab} F^{j}_{bc} E^{b}_{j} :\neq C_{b} Q^{ab} + G_{j} A^{j}_{a}$$
(3.86)

because normal ordering is in particular symmetric. This is because the expression on the right hand side of (3.86) is ill defined on the Fock space even as a quadratic form. There is no contradiction for a solution lof the constraints to disobey $l([C_0(f), C_0(g)]\psi) = 0$ even if $l[C(f)\psi] = 0$ for all $f, \psi \in \mathcal{D}$ as $C_0(f)\psi \notin \mathcal{D}$ as we have emphasised before.

To organise the calculation we follow the above five steps applied to the commutators in the order 1.-6. listed above. Now the Gauss constraint consists of $G = G_1 + G_2$ where G_1, G_2 are of type : F : : F G :respectively, the spatial diffeomorphism constraint is $C = C_2$ which is of type : F G : as well and finally the Hamiltonian constraint is $C_0 = C_{0;3} + C_{0;4}$ where $C_{0;3}$, $C_{0;4}$ are of type : $F \ F \ G \ :, \ : \ F \ F \ G \ G \ :$ respectively. Thus we have 4 different types of normal ordered monomials of : $F^p G^{m-p}$: to consider with (p,m) = (1,1), (1,2), (2,3), (2,4) and correspondingly 10 different commutators of type (3.79) corresponding to (q, n) = (1, 1), (1, 2), (2, 3), (2, 4) as well and it suffices to consider $m \le n$. This part of the calculation can be performed following steps 1-3 above and is listed in appendix A. It has been obtained both by hand and by using computer algebra. After that we consider the terms with at least two factors of Kronecker symbols starting with the highest number of factors. As we have explained above, two Kroneckers can only be obtained for $m = 2 < n \leq 4$, $3 \leq m \leq n \leq 4$ (2+3 terms) and three Kroneckers can only be obtained for m = 3 < n < 4, m = n = 4 (1+1 terms). All other commutators are safe of singularities and the precise limiting pattern will be immaterial. This means that the following comutators in total are unproblematic: Gauss – Gauss, Gauss – Diffeo, Diffeo – Diffeo. The fact that there is no anomalous term in the Diffeo – Diffeo commutator appears to be in conflict with the fact that in one spatial dimension there is the well known Virasoro central term in the constraint algebra generated by the spatial diffeomorphism D and Hamiltonian constraint C, say in parametrised field theory PFT [33], given the fact that the spatial diffeomorphism constraint is theory independent in the sense that it follows purely algebraically from the tensor type that one considers, the information about the concrete Lagrangian of the theory is not encoded in the spatial diffeomorphism constraint but rather in the Hamiltonian constraint. However, a closer look reveals the following: In PFT one considers instead of C, D (in one spatial dimension these are quadratic in the fields and their conjugate momenta) two Poisson commuting constraints $C_{\pm} = D \pm C$ with Poisson algebras isomorphic to that of D and the algebra of the C_{\pm} is anomalous by central terms. However, these two central terms cancel in the algebra of the $D = C_+ + C_-$ while they add in $C = C_+ - C_-$. Thus only C is anomalous in PFT, there is no conflict as far as "universal" (theory independent) part D of the algebra is concerned. The absence of the anomaly in Euclidian GR in 3+1 dimensions is also not in conflict with the presence of the anomaly in PFT in 1+1 dimensions because this concerns the "non-universal" (theory dependent) part C of the algebra which is very different in these two theories.

Sandwiching between Fock states

In what follows we will use all the time the following "sandwiching between Fock states" argument: All commutators and normal reorderings are performed in a controlled way at finite cut-offs. The end result of that step depends on Kronecker symbols and surviving monomials of creation and annihilation operators

in normal ordered form. We are seeking for weak limit of that object, i.e. we sandwich it between any Fock states ψ, ψ' and remove the cut-offs according to some limiting pattern to be derived, and which is independent of ψ, ψ' . Now for any ψ, ψ' we find some M_0 dependening on ψ, ψ' such that $\psi, \psi' \in \mathcal{D}_{M_0}$. As we take the cut-offs to infinity according to some pattern, eventually even the smallest cut-off of that pattern exceeds M_0 . At this point all labels α which label one of the surviving annihilation and creation operators is constrained by M_0 rather than its assigned cut-off. The sum with respect to that label is no longer dependent on it so that the removal of the pre-assigned cut-off is trivial and the details of the limiting pattern get reduced to the remaining labels. The remaining task is then to solve the Kronecker symbols, to determine a limiting pattern consistent with them and to carry out the remaining sums in the prescribed order. Thus if a Kronecker symbol $\delta_{\alpha\beta}$ is given with $\alpha \leq M_1, \beta \leq M_2$ and the pattern is such that $M_1 \leq M_2$ then we must solve for β rather than α . The end result will then depend on finite sums constrained by M_0 which then can be recognised as a quadratic form sandwiched between those ψ, ψ' and the constraint on the sums by M_0 can be taken away so that the final result is independent of ψ, ψ' .

1. Gauss-Gauss

We have to compute three curly brackets.

1A.

The first curly bracket is, see (A.1)

$$[G_{1;M_1}(r), G_{1;M_2}(s)] - [G_{1;M_1}(s), G_{1;M_2}(r)] = 0 - 0 = 0$$
(3.87)

It vanishes trivially for any M_1, M_2 because $G_{1;M}(r)$ just depends on the combinations F_{α} which mutually commute. Therefore the resulting commutator quadratic form, obtained for any limiting pattern, is given by

$$[G_1(r), G_1(s)] = 0 (3.88)$$

1B.

The second curly bracket is, see (A.2)

$$[G_{1;M_{1}}(r), G_{2;M_{2},M_{3}}(s)] - [G_{1;M_{1}}(s), G_{2;M_{2},M_{3}}(r)] = \frac{1}{2\sqrt{2}} \sum_{\alpha \leq M_{1}, \beta \leq M_{2}, \gamma \leq M_{3}} (r_{\alpha}s_{\beta\gamma} - r \leftrightarrow s) [F_{\alpha}, :G_{\beta} F_{\gamma} :]$$

$$= \frac{1}{\sqrt{2}} \sum_{\alpha \leq M_{1}, \beta \leq M_{2}, \gamma \leq M_{3}} (r_{\alpha}s_{\beta\gamma} - r \leftrightarrow s) \delta_{\alpha,\beta} F_{\gamma}$$
(3.89)

By the sandwiching argument, $\gamma \leq M_0$ gets locked. Pick e.g. the limiting pattern $M_1 \to \infty$ before M_2 and carry out the Kronecker symbol in summing over α resulting in

$$\sum_{\beta \le M_2, \gamma \le M_0} \left[(r_\beta s_{\beta\gamma} - r \leftrightarrow s) F_\gamma \right]$$
(3.90)

The order in which we now take $M_2, M_3 \to \infty$ is irrelevant. We now take $M_2 \to \infty$ and have with $\beta = (Imi), \gamma = (Jnj)$ (again summation over repeated lower case latin letters is understood)

$$\sum_{\beta} r_{\beta} s_{\beta\gamma} = \sum_{I} r_{Imi} s_{Imi,Jnj} = \sum_{I} \langle e_{I}, r_{,m}^{i} \rangle \langle s^{l} e_{J}, e_{I} \rangle \delta_{mn} \epsilon_{lij} = \langle e_{J}, r_{,n}^{i} s^{l} \rangle \epsilon_{lij}$$
(3.91)

Thus the final result is

$$\frac{1}{\sqrt{2}} \sum_{I \le M_0} [\langle e_I, r_{,m}^i s^l \rangle - r \leftrightarrow s] \epsilon_{lij} F_{Imj}$$

$$= \frac{1}{\sqrt{2}} \sum_{I \le M_0} \langle e_I, (r^i s^l \epsilon_{lij} \rangle)_{,m} F_{Imj} = i [-\frac{i}{\sqrt{2}} \sum_{I \le M_0} (-r \times s)_{Imi} F_{Imi}] \quad (3.92)$$

where in the last step we applied the product rule. Thus the second curly bracket with e.g. the limiting pattern M_1 before M_2, M_3 yields

$$[G_1(r), G_2(s)] = i \ G_1(-r \times s) \tag{3.93}$$

1C.

The third curly bracket is given by, see (A.5)

$$-\frac{1}{8} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} [r_{\alpha\beta} \ s_{\gamma\delta} - r \leftrightarrow s] [: G_{\alpha}F_{\beta} ::: G_{\gamma}F_{\delta} :]$$

$$= -\frac{1}{4} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} [r_{\alpha\beta} \ s_{\gamma\delta} - r \leftrightarrow s] [\delta_{\beta\gamma} :: F_{\delta} \ G_{\alpha} :-\delta_{\delta\alpha} :: F_{\beta} \ G_{\gamma} :]$$
(3.94)

Pick the pattern M_2, M_4 before $M_1 = M_3$ and solve the kroneckers for β, δ respectively, then upon sandwiching this becomes

$$-\frac{1}{4} \sum_{\alpha \le M_0, \gamma \le M_3, \delta \le M_0} [r_{\alpha \gamma} \ s_{\gamma \delta} - r \leftrightarrow s] : F_{\delta} \ G_{\alpha} :$$

$$+\frac{1}{4} \sum_{\alpha \le M_1, \beta \le M_0, \gamma \le M_0} [r_{\alpha \beta} \ s_{\gamma \alpha} - r \leftrightarrow s] : F_{\beta} \ G_{\gamma} :$$

$$= -\frac{1}{4} \sum_{\alpha \le M_0, \beta \le M_0, \gamma \le M_3} [r_{\alpha \gamma} \ s_{\gamma \beta} - r \leftrightarrow s] : G_{\alpha} \ F_{\beta} :$$

$$+\frac{1}{4} \sum_{\gamma \le M_1, \beta \le M_0, \alpha \le M_0} [r_{\gamma \beta} \ s_{\alpha \gamma} - r \leftrightarrow s] : G_{\alpha} \ F_{\beta} :$$

$$= -\frac{1}{2} \sum_{\alpha \le M_0, \beta \le M_0, \gamma \le M_3} [r_{\alpha \gamma} \ s_{\gamma \beta} - r \leftrightarrow s] : G_{\alpha} \ F_{\beta} :$$
(3.95)

We can take now the remaining cut-off away and have with $\alpha = (Imi), \beta = (Jmj), \gamma = (Kuk), \delta = (Lvl)$

$$\sum_{\gamma} r_{\alpha\gamma} s_{\gamma\beta} = \sum_{J} \langle r_{p} e_{I}, e_{J} \rangle \epsilon_{pij} \,\delta_{mn} \langle e_{J}, s_{q} e_{L} \rangle \epsilon_{qkl} \delta_{uv} \delta_{jk} \delta_{nu} = \langle r_{p} s_{q}, e_{I} e_{L} \rangle \,\delta_{mv} \,2\delta_{p[l} \delta_{q]i} \quad (3.96)$$

This results in

$$-\sum_{I,J \le M_{0}} \langle e_{I}e_{J}, (r_{p}s_{q} - r_{q}s_{p}) \rangle \delta_{p[l}\delta_{q]i} : G_{Imi} F_{Jml} := -\frac{1}{2} \sum_{I,J \le M_{0}} \langle e_{I}e_{J}, (r_{l}s_{i} - r_{i}s_{l}) \rangle : G_{Imi}^{*}F_{Jml} :$$

$$= \frac{1}{2} \sum_{I,J \le M_{0}} (r \times s)_{IJ}^{l}\epsilon_{lij}\delta_{mn} : G_{Imi} F_{Jnj} :$$

$$= i \left[\frac{i}{2} \sum_{\alpha\beta \le M_{0}} (-r \times s)_{\alpha\beta} : G_{\alpha} F_{\beta} :\right]$$
(3.97)

Thus the third curly bracket e.g. with the limiting pattern M_2, M_4 before $M_1 = M_3$

$$[G_2(r), G_2(s)] = i \ G_2(-r \times s) \tag{3.98}$$

It follows altogether

$$[G(r), G(s)] = i \ G(-r \times s) \tag{3.99}$$

2. Diffeomorphism-Diffeomorphism

In this case the only curly bracket to consider is, see (A.5)

$$-\frac{1}{8} \qquad \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} \left[u_{\alpha\beta} \ v_{\gamma\delta} - u \leftrightarrow v \right] \left[: F_{\alpha}G_{\beta} : : : F_{\gamma}G_{\delta} : \right]$$
(3.100)

$$= -\frac{1}{4} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} [u_{\alpha\beta} \ v_{\gamma\delta} - u \leftrightarrow v] (\delta_{\alpha\delta} : F_{\gamma}G_{\beta} : -\delta_{\gamma\beta} : F_{\alpha}G_{\delta} :)$$
(3.101)

In [31] we showed that it is possible to choose the coincidence limit $M_1 = ... = M_4 \to \infty$. We show here that we may choose more general patterns without changing the result.

We pick the pattern M_1, M_3 before $M_2 = M_4$ and use the sandwiching argument, so that we can solve the Kronecker symbols for γ, δ respectively and obtain

$$-\frac{1}{4} \sum_{\beta \le M_0, \gamma \le M_0, \delta \le M_2} [u_{\delta\beta} v_{\gamma\delta} - u \leftrightarrow v] : F_{\gamma} G_{\beta} : +\frac{1}{4} \sum_{\alpha \le M_0, \beta \le M_2, \delta \le M_0} [u_{\alpha\beta} v_{\beta\delta} - u \leftrightarrow v] : F_{\alpha} G_{\delta} :$$

$$= -\frac{1}{4} \sum_{\alpha \le M_0, \beta \le M_0, \gamma \le M_2} \{ [v_{\alpha\gamma} u_{\gamma\beta} - u \leftrightarrow v] - [u_{\alpha\gamma} v_{\gamma\beta} - u \leftrightarrow v] \} : F_{\alpha} G_{\beta} :$$

$$= \frac{1}{2} \sum_{\alpha \le M_0, \beta \le M_0, \gamma \le M_2} [u_{\alpha\gamma} v_{\gamma\beta} - u \leftrightarrow v] : F_{\alpha} G_{\beta} :$$
(3.102)

We can now take $M_2 \rightarrow \infty$ and carry out the sums over γ

$$\sum_{\gamma} \{ [u_{\alpha\gamma}v_{\gamma\beta} - u \leftrightarrow v \}$$

$$= \sum_{K} \{ [\delta_{mr} +] \delta_{ik} [\delta_{rn} +] \delta_{kj} - u \leftrightarrow v \}$$

$$= \sum_{K} \{ [-<[e_{I}u_{p}]_{,p}, e_{K} > \delta_{mr} +][\delta_{rn} +] \delta_{ij} - u \leftrightarrow v \}$$

$$= \{ (-[e_{I}u_{p}]_{,p}\delta_{mr} + e_{I}u_{r,m}), (v_{q}[e_{J}]_{,q}\delta_{rn} + e_{J}v_{n,r}) > \delta_{ij} - u \leftrightarrow v \}$$

$$= \delta_{ij}\{ \delta_{mn} + +$$

$$+ \}$$

$$= \{ \delta_{mn} + \} \delta^{ij}$$

$$(3.103)$$

It follows that the result of the calculation is

$$i \frac{i}{2} \sum_{\alpha \le M_0, \beta \le M_0} (-[u, v])_{\alpha\beta} : F_{\alpha} G_{\beta} :$$
 (3.104)

Thus with e.g. the limiting pattern $M_1 = M_3$ before $M_2 = M_4$ we have

$$[C(u), C(v)] = i C(-[u, v])$$
(3.105)

3. Diffeomorphism-Gauss

We have to consider two curly brackets.

3A.

The first curly bracket is, see (A.2)

$$\frac{1}{2\sqrt{2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3} u_{\alpha\beta} r_{\gamma} [: F_{\alpha} G_{\beta} :, F_{\gamma}]$$

$$= -\frac{1}{\sqrt{2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3} u_{\alpha\beta} r_{\gamma} \delta_{\beta\gamma} F_{\alpha}$$
(3.106)

By sandwiching, α gets locked and we may pick e.g. the pattern M_3 before M_1, M_2 . Then we solve the Kronecker for γ and (3.106) becomes

$$-\frac{1}{\sqrt{2}}\sum_{\alpha \le M_0, \beta \le M_2} u_{\alpha\beta} r_{\beta} F_{\alpha}$$
(3.107)

The cut-off M_2 can now be removed and we find with $\alpha = (Imi), \ \beta = (Jjn)$

$$\sum_{\beta} u_{\alpha\beta} r_{\beta} = \sum_{J} [- \langle [u_{p}e_{I}]_{,p}, e_{J} \rangle \delta_{mn} + \langle e_{I}u_{n,m}, e_{J} \rangle] \delta_{ij} \langle e_{J}, r_{,n}^{j} \rangle$$
$$= \langle e_{I}, u_{p}r_{,m,p}^{i} + u_{p,m}r_{,p}^{i} \rangle = \langle e_{I}, (u_{p}r_{,p}^{i})_{,m} \rangle = (u[r])_{\alpha}$$
(3.108)

It follows that (3.107) becomes

$$i[-\frac{i}{\sqrt{2}}\sum_{\alpha \le M_0} (-u[r])_{\alpha} F_{\alpha}]$$
 (3.109)

Thus for e.g. the limiting pattern M_3 before M_1, M_2 we find

$$[C_2(u), G_1(r)] = iG_1(-u[r])$$
(3.110)

3B.

The second curly bracket is given by, see (A.5)

$$-\frac{1}{4} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} u_{\alpha\beta} r_{\delta\gamma} [: F_{\alpha}G_{\beta} ::: F_{\gamma}G_{\delta} :]$$

$$= -\frac{1}{2} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} u_{\alpha\beta} r_{\delta\gamma} [\delta_{\alpha\delta} :: F_{\gamma}G_{\beta} :-\delta_{\gamma\beta} :: F_{\alpha}G_{\delta} :]$$
(3.111)

Picking again the pattern with M_1, M_3 before M_2, M_4 and sandwiching between Fock states this becomes

$$-\frac{1}{2} \sum_{\beta \le M_0, \gamma \le M_0, \delta \le M_4} u_{\delta\beta} r_{\delta\gamma} : F_{\gamma} G_{\beta} :$$

+
$$\frac{1}{2} \sum_{\alpha \le M_0, \beta \le M_2, \delta \le M_0} u_{\alpha\beta} r_{\delta\beta} : F_{\alpha} G_{\delta} :$$
 (3.112)

We now take $M_2 = M_4 \rightarrow \infty$ and obtain

$$-\frac{1}{2} \sum_{\alpha \le M_0, \beta \le M_0, \gamma} u_{\gamma \alpha} r_{\gamma \beta} : G_{\alpha} F_{\beta} :$$

$$+\frac{1}{2} \sum_{\alpha \le M_0, \beta \le M_0, \gamma} u_{\beta \gamma} r_{\alpha \gamma} : G_{\alpha} F_{\beta} :$$

$$= -\frac{1}{2} \sum_{\alpha \le M_0, \beta \le M_0, \gamma} [u_{\gamma \alpha} r_{\gamma \beta} - u_{\beta \gamma} r_{\alpha \gamma}] : G_{\alpha} F_{\beta} :$$
(3.113)

We have with $\alpha = (Imi), \beta = (Jnj), \gamma = (Kqk)$

$$\sum_{\gamma} [u_{\gamma\alpha} r_{\gamma\beta} - u_{\beta\gamma} r_{\alpha\gamma}]$$

$$= \sum_{K} \{ [\langle e_{K}, u_{p}e_{I,p} \rangle \delta_{qm} + \langle e_{K}e_{I}, u_{m,q} \rangle] \delta_{ki} \epsilon_{lkj} \langle e_{K}e_{J}, r_{l} \rangle \delta_{qn}$$

$$-[\langle e_{J}, u_{p}e_{K,p} \rangle \delta_{nq} + \langle e_{J}e_{K}, u_{q,n} \rangle] \delta_{jk} \epsilon_{lik} \langle e_{I}e_{K}, r_{l} \rangle \delta_{mq} \}$$

$$= \sum_{K} \{ [\langle e_{K}, u_{p}e_{I,p} \rangle \delta_{nm} + \langle e_{K}, e_{I}u_{m,n} \rangle] \epsilon_{lij} \langle e_{J}r_{l}, e_{K} \rangle$$

$$-[-\langle (e_{J}u_{p})_{,p}, e_{K} \rangle \delta_{nm} + \langle e_{J}u_{m,n}, e_{K} \rangle] \epsilon_{lij} \langle e_{K}, e_{I}r_{l} \rangle \}$$

$$= \epsilon_{lij} \{ [\langle e_{J}r_{l}, u_{p}e_{I,p} \rangle \delta_{nm} + \langle e_{J}r_{l}, e_{I}u_{m,n} \rangle] - [-\langle (e_{J}u_{p})_{,p}, e_{I}r_{l} \rangle \delta_{nm} + \langle e_{J}u_{m,n}, e_{I}r_{l} \rangle] \}$$

$$= -\epsilon_{lij} \langle e_{I}e_{J}, u_{p}r_{l,p} \rangle \delta_{mn} = (u[r])_{\alpha\beta} \qquad (3.114)$$

It follows the end result

$$i\frac{i}{2}\sum_{\alpha\leq M_0,\beta\leq M_0} (-u[r])_{\alpha\beta}: G_\alpha \ F_\beta:$$
(3.115)

Therefore for e.g. the pattern M_1, M_3 before $M_2 = M_4$

$$[C_2(u), G_2(r)] = iG_2(-u[r])$$
(3.116)

and altogether

$$[C(u), G(r)] = iG(-u[r])$$
(3.117)

4. Gauss-Hamiltonian

We have to compute four curly bracket terms.

4A.

The first curly bracket is, see (A.3)

$$\frac{i}{2} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} r_{\alpha} f_{\beta \gamma \delta} [F_{\alpha}, : F_{\beta} F_{\gamma} G_{\delta} :]$$

$$= i \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4} r_{\alpha} f_{\beta \gamma \delta} \delta_{\alpha \delta} : F_{\beta} F_{\gamma} :$$
(3.118)

By sandwiching, β , γ get locked. We choose e.g. M_1 before M_2, M_3, M_4 and solve the Kronecker for α . Then (3.118) becomes

$$i \sum_{\beta \le M_0, \gamma \le M_0, \delta \le M_4} r_{\delta} f_{\beta\gamma\delta} : F_{\beta}F_{\gamma} :$$
(3.119)

The sum over δ can now be performed unconstrained and we have for $\beta = (Imi), \gamma = (Jnj), \delta = (Kpk)$

$$\sum_{\gamma} \qquad r_{\delta} f_{\beta\gamma\delta} = \sum_{K} \langle e_{K}, r_{k,p} \rangle \langle [e_{I}e_{J}f]_{,n}, e_{K} \rangle \langle \epsilon_{ijk}\delta_{mp} \rangle$$
$$= \langle r_{k,m}, [e_{I}e_{J}f]_{,n} \rangle \langle \epsilon_{ijk} = -\langle r_{k}, [e_{I}e_{J}f]_{,n,m} \rangle \langle \epsilon_{ijk} =: [f_{r}]_{\beta\gamma} \qquad (3.120)$$

Since $[f_r]_{\beta\gamma}$ is antisymmetric while : $F_{\beta} F_{\gamma}$: is symmetric under exchange of γ, β and since we sum both β, γ over full range in S_{M_0} , expression (3.119) vanishes identically. It follows for e.g. M_1 before $M_2 = M_3, M_4$

$$[G_1(r), C_{0;3}(f)] = 0 (3.121)$$

4B.

The second curly bracket is, see (A.4)

$$\frac{i}{4\sqrt{2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4, \mu \le M_5} r_\mu f_{\alpha\beta\gamma\delta} [F_\mu, : F_\alpha F_\beta G_\gamma G_\delta :]$$

$$= \frac{i}{\sqrt{2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4, \mu \le M_5} r_\mu f_{\alpha\beta\gamma\delta} : F_\alpha F_\beta G_{(\gamma} : \delta_{\delta)\mu}$$
(3.122)

By sandwiching, α, β and γ or δ get locked. We pick the pattern M_5 before $M_1, M_2, M_3 = M_4$ and (3.122) becomes after solving the Kronecker for μ

$$\frac{i}{2\sqrt{2}} \sum_{\alpha \le M_0, \beta \le M_0, \gamma \le M_0, \delta \le M_4} r_{\delta} f_{\alpha\beta\gamma\delta} : F_{\alpha}F_{\beta}G_{\gamma} : + \frac{i}{2\sqrt{2}} \sum_{\alpha \le M_0, \beta \le M_0, \gamma \le M_3, \delta \le M_0} r_{\gamma} f_{\alpha\beta\gamma\delta} : F_{\alpha}F_{\beta}G_{\delta} : (3.123)$$

Taking the remaining cut-off dependence to ∞ we find

$$\frac{i}{2\sqrt{2}} \sum_{\alpha \le M_0, \beta \le M_0, \gamma \le M_0, \delta} r_{\delta} \left[f_{\alpha\beta\gamma\delta} + f_{\alpha\beta\delta\gamma} \right] : F_{\alpha}F_{\beta}G_{\gamma} :$$
(3.124)

and with $\alpha = (Imi), \ \beta = (Jnj), \ \gamma = (Kpk), \delta = (Lql)$

$$\sum_{\delta} r_{\delta} \left[f_{\alpha\beta\gamma\delta} + f_{\alpha\beta\delta\gamma} \right] = \sum_{L} \langle e_{L}, r_{l,q} \rangle \left[\langle e_{I}e_{J}e_{K}e_{L}, f \rangle \delta_{mp}\delta_{nq}\epsilon_{tij}\epsilon_{tkl} + \langle e_{I}e_{J}e_{L}e_{K}, f \rangle \delta_{mq}\delta_{np}\epsilon_{tij}\epsilon_{tlk} \right]$$
$$= \left[\langle e_{I}e_{J}e_{K}, fr_{l,n} \rangle \delta_{mp} - \langle e_{I}e_{J}e_{K}, fr_{l,m} \rangle \delta_{np} \right]\epsilon_{tij}\epsilon_{tkl} =: \left[f^{r} \right]_{\alpha\beta\gamma}$$
(3.125)

Accordingly with e.g. the pattern M_5 before $M_1, M_2, M_3 = M_4$

$$[G_1(r), C_{0;4}(f)] = \frac{i}{2\sqrt{2}} \sum_{\alpha, \beta, \gamma} [f^r]_{\alpha\beta\gamma} : F_\alpha F_\beta G_\gamma :$$
(3.126)

4C.

The third curly bracket is, see (A.6)

$$-\frac{i}{2 2^{1/2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \mu \le M_4, \rho \le M_5} r_{\rho\mu} f_{\alpha\beta\gamma} [: F_{\mu} G_{\rho} :: F_{\alpha} F_{\beta} G_{\gamma} :]$$

$$= -\frac{i}{2 2^{1/2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \mu \le M_4, \rho \le M_5} r_{\rho\mu} f_{\alpha\beta\gamma} (2\delta_{\mu\gamma} :: F_{\alpha}F_{\beta}G_{\rho} :-4\delta_{\rho(\alpha} :: F_{\beta)}F_{\mu}G_{\gamma} :+4\delta_{\rho(\alpha}\delta_{\beta)\mu}G_{\gamma})$$

$$(3.127)$$

4C.i

This is the first incident that a potential divergence can appear because there is a term involving the product of two Kronecker symbols which we treat first. We assume e.g. a pattern with $M_4 \ge M_1, M_2 \ge M_3, M_5$, then in the term with two Kronecker symbols the index γ is locked by sandwiching and we can solve the Kroneckers for μ and α or β . The result, using $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \mu = (Mur), \rho = (Nvs),$ is proportional to

$$\sum_{\substack{\rho \leq M_5, \gamma \leq M_0}} [\sum_{\beta \leq M_2} r_{\rho\beta} f_{\rho\beta\gamma} + \sum_{\alpha \leq M_1} r_{\rho\alpha} f_{\alpha\rho\gamma}]G_{\gamma}$$

$$= \sum_{\substack{N \leq M_5, K \leq M_0}} [\sum_{J \leq M_2} r_{NJt} \delta_{vn} \epsilon_{tsj} f_{NJnK} \delta_{vp} \epsilon_{sjk} + \sum_{I \leq M_1} r_{NIt} \delta_{vm} \epsilon_{tsi} f_{INvK} \delta_{mp} \epsilon_{isk}]G_{Kpk}$$

$$= 2 \sum_{\substack{N \leq M_5, K \leq M_0}} [\sum_{J \leq M_2} r_{NJk} f_{NJpK} - \sum_{I \leq M_1} r_{NIk} f_{INpK}]G_{Kpk}$$

$$= 0 \qquad (3.128)$$

if $M_1 = M_2$ since $f_{INpK} = \langle (e_I e_N f)_{,p}, e_K \rangle = f_{NIpK}$. Thus we will adopt $M_1 = M_2$ from now on. Note that this result crucially rests on properties of $r_{\alpha\beta}, f_{\alpha\beta\gamma}$.

4C.ii

The contribution to (3.127) from the single Kronecker symbol terms is

$$-\frac{i}{2^{1/2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \mu \le M_4, \rho \le M_5} r_{\rho\mu} f_{\alpha\beta\gamma} \left[\delta_{\mu\gamma} : F_{\alpha}F_{\beta}G_{\rho} : -2\delta_{\rho(\alpha)} : F_{\beta}F_{\mu}G_{\gamma} : \right]$$
(3.129)

By sandwiching and solving for μ in the first term and α or β in the second respectively according to above

pattern this becomes

$$-\frac{i}{2^{1/2}} \left\{ \sum_{\alpha,\beta,\rho \leq M_{0},\gamma \leq M_{3}} r_{\rho\gamma} f_{\alpha\beta\gamma} : F_{\alpha}F_{\beta}G_{\rho} : -\sum_{\beta,\gamma,\mu \leq M_{0},\rho \leq M_{5}} r_{\rho\mu} f_{\rho\beta\gamma} : F_{\beta}F_{\mu}G_{\gamma} : -\sum_{\alpha,\gamma,\mu \leq M_{0},\rho \leq M_{5}} r_{\rho\mu} f_{\alpha\rho\gamma} : F_{\alpha}F_{\mu}G_{\gamma} : \right\}$$

$$= -\frac{i}{2^{1/2}} \left\{ \sum_{\alpha,\beta,\gamma \leq M_{0},\rho \leq M_{3}} r_{\gamma\rho} f_{\alpha\beta\rho} : F_{\alpha}F_{\beta}G_{\gamma} : -\sum_{\alpha,\beta,\gamma \leq M_{0},\rho \leq M_{5}} r_{\rho\beta} f_{\rho\alpha\gamma} : F_{\alpha}F_{\beta}G_{\gamma} : -\sum_{\alpha,\gamma,\beta \leq M_{0},\rho \leq M_{5}} r_{\rho\beta} f_{\alpha\rho\gamma} : F_{\alpha}F_{\beta}G_{\gamma} : \right\}$$

$$= -\frac{i}{2^{1/2}} \left\{ \sum_{\alpha,\beta,\gamma \leq M_{0},\rho \leq M_{3}} r_{\gamma\rho} f_{\alpha\beta\rho} - \sum_{\alpha,\beta,\gamma \leq M_{0},\rho \leq M_{5}} r_{\rho\beta} [f_{\rho\alpha\gamma} + f_{\alpha\rho\gamma}] \right\} : F_{\alpha}F_{\beta}G_{\gamma} : \quad (3.130)$$

We now take the remaining cut-offs on ρ away and carry out the unconstrained sum over $\rho = (Lql)$. This gives for the first term in the curly bracket of (3.130)

$$\sum_{\rho} r_{\gamma\rho} f_{\alpha\beta\rho} = \sum_{L} r_{KL}^{t} \delta_{pq} \epsilon_{tkl} f_{IJnL} \delta_{mq} \epsilon_{ijl}$$

$$= \sum_{L} \langle r_{t}e_{K}, e_{L} \rangle \langle e_{L}, (e_{I}e_{J}f)_{,n} \rangle \delta_{mp} \epsilon_{tkl} \epsilon_{ijl}$$

$$= \langle r_{t}e_{K}, (e_{I}e_{J}f)_{,n} \rangle \delta_{mp} \epsilon_{tkl} \epsilon_{ijl}$$

$$\sum_{\rho} r_{\rho\beta} [f_{\rho\alpha\gamma} + f_{\alpha\rho\gamma}] = \sum_{L} r_{LJ}^{t} \epsilon_{tlj} \delta_{qn} [f_{LImK} \delta_{qp} \epsilon_{lik} + f_{ILqK} \delta_{mp} \epsilon_{ilk}]$$

$$= \sum_{L} r_{LJ}^{t} \epsilon_{tlj} [f_{LImK} \delta_{np} - f_{ILnK} \delta_{mp}] \epsilon_{lik}$$

$$= -\sum_{L} \langle r_{t}e_{J}, e_{L} \rangle \epsilon_{tlj} [\langle e_{L}, fe_{I}e_{K,m} \rangle \delta_{np} - \langle e_{L}, fe_{I}e_{K,n} \rangle \delta_{mp}] \epsilon_{lik}$$

$$= -\epsilon_{tlj} \epsilon_{lik} [\langle r_{t}e_{J}fe_{I}, e_{K,m} \rangle \delta_{np} - \langle r_{t}e_{J}fe_{I}, e_{K,n} \rangle \delta_{mp}]$$
(3.131)

We use that : $F_{\alpha}F_{\beta}G_{\gamma} :=: F_{\beta}F_{\alpha}G_{\gamma}$: and replace in (3.131) both equalities by half the sum of (3.131) and (3.131) with α, β interchanged which results in

$$1. = \frac{1}{2} [\langle r_t e_K, (e_I e_J f)_{,n} \rangle \delta_{mp} - \langle r_t e_K, (e_I e_J f)_{,m} \rangle \delta_{np}] \epsilon_{tkl} \epsilon_{ijl}$$

$$2. = \frac{1}{2} [\epsilon_{tlj} \epsilon_{lik} - \epsilon_{tli} \epsilon_{ljk}] [\langle r_t e_J f e_I, e_{K,n} \rangle \delta_{mp} - \langle r_t e_J f e_I, e_{K,m} \rangle \delta_{np}]$$

$$= \frac{1}{2} \epsilon_{ijs} \epsilon_{uvs} \epsilon_{luk} \epsilon_{lvt} [\langle r_t e_J f e_I, e_{K,n} \rangle \delta_{mp} - \langle r_t e_J f e_I, e_{K,m} \rangle \delta_{np}]$$

$$= -\frac{1}{2} \epsilon_{ijs} \epsilon_{tks} [\langle r_t e_J f e_I, e_{K,n} \rangle \delta_{mp} - \langle r_t e_J f e_I, e_{K,m} \rangle \delta_{np}]$$

$$(3.132)$$

for the first and second equality respectively. The first minus the second of those two expressions that is required in (3.130) is given by

$$\frac{1}{2} \epsilon_{tkl} \epsilon_{ijl} \{ \langle r_t e_K, (e_I e_J f)_{,n} \rangle \delta_{mp} - \langle r_t e_K, (e_I e_J f)_{,m} \rangle \delta_{np} + \langle r_t e_J f e_I, e_{K,n} \rangle \delta_{mp} - \langle r_t e_J f e_I, e_{K,m} \rangle \delta_{np} \}$$

$$= -\frac{1}{2} \epsilon_{tkl} \epsilon_{ijl} \{ \langle r_{t,n}, e_I e_J e_K f \rangle \delta_{mp} - \langle r_{t,m}, e_I e_J e_K f \rangle \delta_{np} \}$$

$$= \frac{1}{2} [f^r]_{\alpha\beta\gamma} \qquad (3.133)$$

where the expression (3.125) reappeared.

It follows e.g. for the pattern M_4 before $M_1 = M_2$ before M_3, M_5

$$[G_2(r), C_{0;3}(f)] = -\frac{i}{2\sqrt{2}} \sum_{\alpha\beta\gamma} [f^r]_{\alpha\beta\gamma} : F_{\alpha} F_{\beta} G_{\gamma} :$$
(3.134)

which is excally the negative of (3.126).

4D.

The fourth curly bracket is, see (A.7)

$$-\frac{i}{8} \sum_{\alpha \le M_1 \beta \le M_2, \gamma \le M_3, \delta \le M_4, \mu \le M_5, \rho \le M_6} r_{\rho\mu} f_{\alpha\beta\gamma\delta} [: F_{\mu}G_{\rho} :: F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} :]$$

$$[: F_{\mu}G_{\rho} :: F_{\alpha}F_{\beta}G_{\gamma}G\delta :] = 4 : F_{\alpha}F_{\beta}G_{\rho}G_{(\gamma} : \delta_{\delta)\mu} + 4\delta_{\mu(\gamma} \delta_{\delta)\rho} : F_{\alpha}F_{\beta} :$$

$$-4 \delta_{\rho(\alpha} : F_{\beta})F_{\mu}G_{\gamma}G_{\delta} :+ 4\delta_{\rho(\alpha} \delta_{\beta)\mu} : G_{\gamma}G_{\delta} :$$

$$(3.135)$$

4D.i

Consider the contributions from two Kronecker symbols. Consider the pattern $M_5 \ge M_1, M_2 \ge M_6 \ge M_3, M_4$. For the term $\propto: F_{\alpha}F_{\beta}$: we obtain using the sandwiching argumment $\alpha, \beta \le M_0$ and after solving for μ, ρ a coefficient proportional to

$$\sum_{\gamma \le M_3, \delta \le M_4} \left[r_{\gamma\delta} + r_{\delta\gamma} \right] f_{\alpha\beta\gamma\delta} = 0 \tag{3.136}$$

because $r_{\gamma\delta} = -r_{\delta\gamma}$. For the term $\propto: G_{\gamma}G_{\delta}$: we obtain using the sandwiching argumment $\gamma, \delta \leq M_0$ and after solving for μ and α or β a coefficient proportional to

$$\sum_{\rho \le M_6} \left[\sum_{\beta \le M_2} r_{\rho\beta} f_{\rho\beta\gamma\delta} + \sum_{\alpha \le M_1} r_{\rho\alpha} f_{\alpha\rho\gamma\delta} \right]$$
(3.137)

Using $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \delta = (Lql), \mu = (Mur), \rho = (Nvs)$ the term in square brackets becomes

$$\sum_{J \leq M_2} r_{NJ}^t \,\delta_{vn} \,\epsilon_{tsj} \,f_{NJKL} \,\delta_{vp} \,\delta_{nq} \,\epsilon_{rsj} \,\epsilon_{rkl} + \sum_{I \leq M_1} r_{NI}^t \,\delta_{vm} \,\epsilon_{tsi} \,f_{INKL} \,\delta_{mp} \,\delta_{vq} \,\epsilon_{ris} \,\epsilon_{rkl}$$

$$= 2\epsilon_{tkl} \,\delta_{pq} \left[\sum_{J \leq M_2} r_{NJ}^t f_{JNKL} - \sum_{I \leq M_1} r_{NI}^t f_{INKL}\right]$$
(3.138)

which vanishes if $M_1 = M_2$ due to total symmetry of f_{IJKL} . Thus we will adopt $M_1 = M_2$ from now on.

4D.ii

The single Kronecker symbol contributions to (3.135) are given by

$$-\frac{i}{4} \sum_{\alpha \le M_1 \beta \le M_2, \gamma \le M_3, \delta \le M_4, \mu \le M_5, \rho \le M_6} r_{\rho\mu} f_{\alpha\beta\gamma\delta} \left[2 : F_\alpha F_\beta G_\rho G_{(\gamma} : \delta_{\delta)\mu} - 2 \delta_{\rho(\alpha} : F_\beta) F_\mu G_\gamma G_\delta :\right]$$
(3.139)

We solve for μ and α or β according to the chosen pattern and obtain with the sandwiching argument and

renaming indices carefully keeping track of the corresponding index range

$$-\frac{i}{4} \sum_{\alpha,\beta,\rho \leq M_{0}} \left[\sum_{\gamma \leq M_{0},\delta \leq M_{4}} f_{\alpha\beta\gamma\delta} : F_{\alpha}F_{\beta}G_{\rho}G_{\gamma} : r_{\rho\delta} + \sum_{\gamma \leq M_{3},\delta \leq M_{0}} f_{\alpha\beta\gamma\delta} : F_{\alpha}F_{\beta}G_{\rho}G_{\delta} : r_{\rho\gamma} \right] \\ +\frac{i}{4} \sum_{\mu,\gamma,\delta \leq M_{0},\rho \leq M_{6}} r_{\rho\mu} \left[\sum_{\beta \leq M_{0}} f_{\rho\beta\gamma\delta} : F_{\beta}F_{\mu}G_{\gamma}G_{\delta} : + \sum_{\alpha \leq M_{0}} f_{\alpha\rho\gamma\delta} : F_{\alpha}F_{\mu}G_{\gamma}G_{\delta} : \right] \\ = -\frac{i}{4} \sum_{\alpha,\beta,\gamma,\rho \leq M_{0}} \left[\sum_{\delta \leq M_{4}} r_{\rho\delta} f_{\alpha\beta\gamma\delta} + \sum_{\delta \leq M_{3}} r_{\rho\delta} f_{\alpha\beta\delta\gamma} \right] : F_{\alpha}F_{\beta}G_{\gamma}G_{\rho} : \\ +\frac{i}{4} \sum_{\beta,\mu,\gamma,\delta \leq M_{0};\rho \leq M_{6}} r_{\rho\mu} \left[f_{\rho\beta\gamma\delta} + f_{\beta\rho\gamma\delta} \right] : F_{\beta}F_{\mu}G_{\gamma}G_{\delta} : \\ = -\frac{i}{4} \sum_{\alpha,\beta,\gamma,\delta \leq M_{0};\rho \leq M_{6}} r_{\rho\rho} f_{\alpha\beta\gamma\rho} + \sum_{\rho \leq M_{3}} r_{\delta\rho} f_{\alpha\beta\rho\gamma} \right] : F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} : \\ +\frac{i}{4} \sum_{\alpha,\beta,\gamma,\delta \leq M_{0};\rho \leq M_{6}} r_{\rho\beta} \left[f_{\rho\alpha\gamma\delta} + f_{\alpha\rho\gamma\delta} \right] : F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} :$$
(3.140)

We take the final cut-offs away and find

$$-\frac{i}{4} \sum_{\alpha,\beta,\gamma,\delta \le M_0;\rho} \left\{ r_{\delta\rho} \left[f_{\alpha\beta\gamma\rho} + f_{\alpha\beta\rho\gamma} \right] - r_{\rho\beta} \left[f_{\rho\alpha\gamma\delta} + f_{\alpha\rho\gamma\delta} \right] : F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} : \right\}$$
(3.141)

Note that $r_{\delta\rho} = -r_{\rho\delta}$ and $f_{\alpha\beta\gamma\delta} = f_{\gamma\delta\alpha\beta}$. We have with $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \delta = (Lql), \rho = (Mus)$

$$\sum_{\rho} r_{\rho\delta} f_{\alpha\beta\gamma\rho} = \sum_{M} r_{ML}^{t} \,\delta_{uq} \,\epsilon_{tsl} \,f_{IJKM} \,\delta_{mp} \,\delta_{nu} \,\epsilon_{wij} \,\epsilon_{wks}$$
$$= \epsilon_{tsl} \,< e_{I}e_{J}e_{K}e_{L}, fr^{t} > \,\delta_{mp} \,\delta_{nq} \,\epsilon_{wij} \,\epsilon_{wks} =: [f_{r}]_{IJKL}^{t} \,\delta_{mp} \,\delta_{nq} \,\epsilon_{tsl} \,\epsilon_{wij} \,\epsilon_{wks} \tag{3.142}$$

Thus

$$\sum_{\rho} r_{\rho\delta} f_{\alpha\beta\rho\gamma} = [f_r]^t_{IJKL} \delta_{mq} \delta_{np} \epsilon_{tsl} \epsilon_{wij} \epsilon_{wsk}$$

$$\sum_{\rho} r_{\rho\beta} f_{\rho\alpha\gamma\delta} = [f_r]^t_{IJKL} \delta_{mq} \delta_{np} \epsilon_{tsj} \epsilon_{wsi} \epsilon_{wkl}$$

$$\sum_{\rho} r_{\rho\beta} f_{\alpha\rho\gamma\delta} = [f_r]^t_{IJKL} \delta_{mp} \delta_{nq} \epsilon_{tsj} \epsilon_{wis} \epsilon_{wkl} \qquad (3.143)$$

Thus the coefficient of $\frac{i}{4}$: $F_{\alpha}F_{\beta}G_{\gamma}G_{\delta}$: in (3.141) is

$$-[f_r]^t_{IJKL}\{[\epsilon_{wij}\epsilon_{tsl}\epsilon_{wsk} + \epsilon_{wkl}\epsilon_{tsj}\epsilon_{wsi}] [\delta_{mp}\delta_{nq} - \delta_{np}\delta_{mq}]$$
(3.144)

As : $F_{\alpha}F_{\beta}G_{\gamma}G_{\delta}$: is symmetric separately under exchange of α, β and γ, δ we replace (3.144) by half the sum of (3.144) and (3.144) with $\alpha \leftrightarrow \beta, \gamma \leftrightarrow \delta$. We obtain

$$-\frac{1}{2}[f_r]_{IJKL}^t \{ [\epsilon_{wij}(\epsilon_{tsl}\epsilon_{wsk} - \epsilon_{tsk}\epsilon_{wsl}) + \epsilon_{wkl}(\epsilon_{tsj}\epsilon_{wsi} - \epsilon_{tsi}\epsilon_{wsj})] [\delta_{mp}\delta_{nq} - \delta_{np}\delta_{mq}] \\ -\frac{1}{2}[f_r]_{IJKL}^t \{ [\epsilon_{wij}\epsilon_{vkl}\epsilon_{vtw} + \epsilon_{wkl}\epsilon_{vij}\epsilon_{vtw}] \} \\ -\frac{1}{2}[f_r]_{IJKL}^t \epsilon_{wij}\epsilon_{vkl} [\epsilon_{vtw} + \epsilon_{wtv} = 0$$

$$(3.145)$$

It follows e.g. with M_5 before $M_1 = M_2$ before M_6 before $M_3 = M_4$

$$[G_2(r), C_{0;4}(f)] = 0 (3.146)$$

Altogether

$$[G(r), C_0(f)] = 0 (3.147)$$

5. Diffeomorphism - Hamiltonian

We have two curly brackets to consider:

5A.

The first curly bracket is

$$-\frac{i}{2 \ 2^{1/2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \mu \le M_4, \rho \le M_5} u_{\mu\rho} \ f_{\alpha\beta\gamma} \ [: F_{\mu}G_{\rho} :, : F_{\alpha}F_{\beta}G_{\gamma} :]$$
(3.148)

We can use the same results as in item 4C., in particular (3.127). We pick the same pattern $M_4 \ge M_1, M_2 \ge M_3, M_5$.

5A.i

We consider first the contribution from two Kronecker symbols to (3.148) which after solving for μ and α or β is proportional to with $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \rho = (Lql)$

$$\sum_{\rho \le M_5} \left[\sum_{\beta \le M_2} u_{\beta\rho} f_{\rho\beta\gamma} + \sum_{\alpha \le M_1} u_{\alpha\rho} f_{\alpha\rho\gamma} \right] = 0$$
(3.149)

because $u_{\beta\rho} \propto \delta_{jl}$, $f_{\rho\beta\gamma} \propto \epsilon_{ljk}$ even without assuming $M_1 = M_2$.

5A.ii

The contribution from the single Kronecker symbol to (3.148) is given by

$$-\frac{i}{2^{1/2}}\sum_{\alpha\leq M_1,\beta\leq M_2,\gamma\leq M_3,\mu M_4,\rho\leq M_5} u_{\mu\rho} f_{\alpha\beta\gamma} \left[\delta_{\mu\gamma} : F_{\alpha}F_{\beta}G_{\rho} : -2\delta_{\rho(\alpha)} : F_{\beta}F_{\mu}G_{\gamma} :\right]$$
(3.150)

We solve for μ or α, β and sandwich

$$-\frac{i}{2^{1/2}} \{ \sum_{\alpha,\beta,\rho \le M_0; \gamma \le M_3} u_{\gamma\rho} f_{\alpha\beta\gamma} : F_{\alpha}F_{\beta}G_{\rho} : -\sum_{\beta,\mu,\gamma \le M_0,\rho \le M_5} u_{\mu\rho} f_{\rho\beta\gamma} : F_{\beta}F_{\mu}G_{\gamma} : -\sum_{\alpha,\mu,\gamma \le M_0;\rho \le M_5} u_{\mu\rho} f_{\alpha\rho\gamma} : F_{\alpha}F_{\mu}G_{\gamma} :] \}$$

$$= -\frac{i}{2^{1/2}} \sum_{\alpha,\beta,\gamma \le M_0} \{ \sum_{\rho \le M_3} u_{\rho\gamma} f_{\alpha\beta\rho} - \sum_{\rho \le M_5} (u_{\alpha\rho} f_{\rho\beta\gamma} + u_{\beta\rho} f_{\alpha\rho\gamma}) \} : F_{\alpha}F_{\beta}G_{\gamma} : \qquad (3.151)$$

Taking the remaining cut-offs to infinity the sum over ρ is unconstrained and we compute with $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \rho = (Lql)$

$$\sum_{\rho} \{ u_{\rho\gamma} f_{\alpha\beta\rho} - u_{\alpha\rho} f_{\rho\beta\gamma} - u_{\beta\rho} f_{\alpha\rho\gamma} \}$$

$$= \sum_{L} \{ u_{LK}^{qp} \delta_{lk} f_{IJnL} \delta_{mq} \epsilon_{ijl} - u_{IL}^{mq} \delta_{il} f_{LJnK} \delta_{qp} \epsilon_{ljk} - u_{JL}^{nq} \delta_{jl} f_{ILqK} \delta_{mp} \epsilon_{ilk} \}$$

$$= \sum_{L} \{ u_{LK}^{mp} f_{IJnL} - u_{IL}^{mp} f_{LJnK} - u_{JL}^{nq} f_{ILqK} \delta_{mp} \} \epsilon_{ijk}$$
(3.152)

Consider the coefficient of $\delta_{mp}\epsilon_{ijk}$ from the first two terms together with the whole third term

$$\sum_{L} \{ \langle e_{L}, u_{r}e_{K,r} \rangle \langle (e_{I}e_{J}f)_{,n}, e_{L} \rangle - \langle e_{I}, u_{r}e_{L,r} \rangle \langle (e_{L}e_{J}f)_{,n}, e_{K} \rangle - [\langle e_{J}, u_{r}e_{L,r} \rangle \delta_{nq} + \langle e_{J}e_{L}, u_{q,n} \rangle] \langle (e_{I}e_{L}f)_{,q}, e_{K} \rangle \}$$

$$= \sum_{L} \{ \langle e_{L}, u_{r}e_{K,r} \rangle \langle (e_{I}e_{J}f)_{,n}, e_{L} \rangle - \langle (e_{I}u_{r})_{,r}, e_{L} \rangle \langle e_{L}, e_{J}fe_{K,n} \rangle - [\langle (e_{J}u_{r})_{,r}e_{L} \rangle \delta_{nq} - \langle e_{L}, e_{J}u_{q,n} \rangle] \langle e_{L}, fe_{I}e_{K,q} \rangle \}$$

$$= \langle u_{r}e_{K,r}, (e_{I}e_{J}f)_{,n} \rangle - \langle (e_{I}u_{r})_{,r}, e_{J}fe_{K,n} \rangle - \langle (e_{J}u_{r})_{,r}, fe_{I}e_{K,n} \rangle + \langle e_{J}u_{r,n}, fe_{I}e_{K,r} \rangle$$

$$= \langle e_{K,r}, u_{r} (e_{I}e_{J}f)_{,n} + (e_{I}e_{J}f)u_{r,n} \rangle - \langle e_{K,n}, f[(e_{I}u_{r})_{,r}, e_{J} + (e_{J}u_{r})_{,r}e_{I}] \rangle$$

$$= \langle e_{K,r}, (u_{r} e_{I}e_{J}f)_{,n} \rangle - \langle e_{K,n}, f[(e_{I}e_{J}u_{r})_{,r} + u_{r,r}e_{I}e_{J}] \rangle$$

$$= \langle e_{K,n}, (u_{r}f_{,r} - u_{r,r}f) e_{I}e_{J} \rangle = - \langle (u[f]e_{I}e_{J})_{,n}e_{K} \rangle = -[u[f]]_{IJnK}$$
(3.153)

where we used $< (.)_m, (.)'_n > = < (.)_n, (.)'_m >$ in between. The coefficient $\not \propto \delta_{mp} \epsilon_{ijk}$ of (3.152) is

$$\sum_{L} \{ \langle e_{L}e_{K}, u_{p,m} \rangle \langle (e_{I}e_{J}f)_{,n}, e_{L} \rangle - \langle e_{I}e_{L}, u_{p,m} \rangle \langle (e_{L}e_{J}f)_{,n}, e_{K} \rangle \}$$

$$= \sum_{L} \{ \langle e_{L}, e_{K}u_{p,m} \rangle \langle (e_{I}e_{J}f)_{,n}, e_{L} \rangle + \langle e_{I}e_{L}, u_{p,m} \rangle \langle e_{L}, e_{J}fe_{K,n} \rangle \}$$

$$= \langle e_{K}u_{p,m}, (e_{I}e_{J}f)_{,n} \rangle + \langle e_{I}, u_{p,m}, e_{J}fe_{K,n} \rangle$$

$$= -\langle e_{I}e_{J}e_{K}f, u_{p,m,n} \rangle$$
(3.154)

Together with the factor ϵ_{ijk} (3.154) is a coefficient $[f_u]_{\alpha\beta\gamma}$ which is skew under exchange of α, β while : $F_{\alpha}F_{\beta}G_{\gamma}$: is symmetric. It follows that the first curly bracket is

$$i \left[-\frac{1}{2^{1/2}} \sum_{\alpha,\beta,\gamma \le M_0} (-u[f])_{\alpha\beta\gamma} : F_{\alpha}F_{\beta}G_{\gamma} :\right]$$
(3.155)

Thus e.g. for the pattern M_4 before $M_1 = M_2$ before M_3, M_5 we get

$$[C_2(u), C_{0;3}(f)] = iC_{0;3}(-u[f])$$
(3.156)

5B.

The second curly bracket is given by

$$-\frac{i}{8} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4, \mu \le M_5, \rho \le M_6} u_{\mu\rho} f_{\alpha\beta\gamma\delta} [: F_{\mu}G_{\rho} ::: F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} :]$$
(3.157)

We can use the same results as in item 4D., in particular (3.135). We pick the same pattern $M_5 \ge M_1, M_2 \ge M_6 \ge M_3, M_4$.

5B.i

We consider first the contribution from two Kronecker symbols to (3.157) The coefficient of : $F_{\alpha}F_{\beta}$: after solving for μ, ρ (which is consistent with the pattern as α, β get locked when sandwiching) is proportional to

$$-2\sum_{\gamma\delta} \left[u_{\gamma\delta} + u_{\delta\gamma}\right] f_{\alpha\beta\gamma\delta} \tag{3.158}$$

which vanishes as $u_{\gamma\delta}, u_{\delta\gamma} \propto \delta_{kl}$ while $f_{\alpha\beta\gamma\delta} \propto \epsilon_{rkl}$. The coefficient of : $G_{\gamma}G_{\delta}$: after solving for μ and α or β is proportional to

$$2\sum_{\rho} \left[\sum_{\beta} u_{\beta\rho} f_{\rho\beta\gamma\delta} + \sum_{\alpha} u_{\alpha\rho} f_{\alpha\rho\gamma\delta}\right]$$
(3.159)

which vanishes as $u_{\alpha\rho} \propto \delta_{il}$ while $f_{\alpha\rho\gamma\delta}, f_{\rho\alpha\gamma\delta} \propto \epsilon_{ril}$.

5B.ii

The single Kronecker symbol contribution to (3.157) is given by

$$-\frac{i}{4} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4, \mu \le M_5, \rho \le M_6} u_{\mu\rho} f_{\alpha\beta\gamma\delta} \left[2 : F_{\alpha}F_{\beta}G_{\rho}G_{(\gamma} : \delta_{\delta)\mu} - 2 \delta_{\rho(\alpha} : F_{\beta)}F_{\mu}G_{\gamma}G_{\delta} :\right]$$
(3.160)

We solve for μ and α or β and get by sandwiching

$$-\frac{i}{4} \{ \sum_{\alpha,\beta,\rho,\gamma \leq M_{0};\delta \leq M_{4}} u_{\delta\rho} f_{\alpha\beta\gamma\delta} : F_{\alpha}F_{\beta}G_{\rho}G_{\gamma} : + \sum_{\alpha,\beta\rho,\delta \leq M_{0};\gamma \leq M_{3}} u_{\gamma\rho} f_{\alpha\beta\gamma\delta} : F_{\alpha}F_{\beta}G_{\rho}G_{\delta} : - \sum_{\beta,\mu,\gamma,\delta \leq M_{0};\rho \leq M_{6}} u_{\mu\rho} f_{\alpha\rho\gamma\delta} : F_{\alpha}F_{\mu}G_{\gamma}G_{\delta} : + \sum_{\alpha,\mu,\gamma,\delta \leq M_{0};\rho \leq M_{6}} u_{\mu\rho} f_{\alpha\rho\gamma\delta} : F_{\alpha}F_{\mu}G_{\gamma}G_{\delta} : + \sum_{\alpha,\beta\gamma,\delta \leq M_{0};\rho \leq M_{4}} u_{\rho\delta} f_{\alpha\beta\gamma\rho} : F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} : + \sum_{\alpha,\beta\gamma,\delta \leq M_{0};\rho \leq M_{3}} u_{\rho\gamma} f_{\alpha\beta\rho\delta} : F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} : - \sum_{\alpha,\beta,\gamma,\delta \leq M_{0};\rho \leq M_{6}} u_{\alpha\rho} f_{\alpha\rho\gamma\delta} : F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} : - \sum_{\alpha,\beta,\gamma,\delta \leq M_{0};\rho \leq M_{6}} u_{\beta\rho} f_{\alpha\rho\gamma\delta} : F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} : \}$$
(3.161)

Taking the remaining cut-offs to infinity the sum over ρ becomes unconstrained and the coefficient of $-\frac{i}{4}: F_{\alpha}F_{\beta}G_{\gamma}G_{\delta}:$ in (3.161) becomes

$$\sum_{\rho} \{ u_{\rho\delta} f_{\alpha\beta\gamma\rho} + u_{\rho\gamma} f_{\alpha\beta\rho\delta} - u_{\alpha\rho} f_{\rho\beta\gamma\delta} - u_{\beta\rho} f_{\alpha\rho\gamma\delta} \}$$

$$= \sum_{M} \{ u_{ML}^{uq} \delta_{sl} f_{IJKM} \delta_{mp} \delta_{nu} \epsilon_{tij} \epsilon_{tks} + u_{MK}^{up} \delta_{sk} f_{IJML} \delta_{mu} \delta_{nq} \epsilon_{tij} \epsilon_{tsl} - u_{IM}^{mu} \delta_{is} f_{MJKL} \delta_{up} \delta_{nq} \epsilon_{tsj} \epsilon_{tkl} - u_{JM}^{nu} \delta_{js} f_{IMKL} \delta_{pm} \delta_{uq} \epsilon_{tis} \epsilon_{tkl} \}$$

$$= \epsilon_{tij} \epsilon_{tkl} \sum_{M} \{ \delta_{mp} [u_{ML}^{nq} f_{IJKM} - u_{JM}^{nq} f_{IMKL}] + \delta_{nq} [u_{MK}^{mp} f_{IJML} - u_{IM}^{mp} f_{MJKL}] \}$$
(3.162)

The contributions proportional to $\delta_{mp}\delta_{nq}$ to (3.162) are

$$\sum_{L} \{ \langle e_{M}, u_{r}e_{L,r} \rangle \langle e_{I}e_{J}e_{K}e_{M}, f \rangle - \langle e_{J}, u_{r}e_{M,r} \rangle \langle e_{I}e_{K}e_{L}e_{M}, f \rangle + \langle e_{M}, u_{r}e_{K,r} \rangle \langle e_{I}e_{J}e_{L}e_{M}, f \rangle - \langle e_{I}, u_{r}e_{M,r} \rangle \langle e_{J}e_{K}e_{L}e_{M}, f \rangle \}$$

$$= \langle fu_{r}, e_{L,r}e_{I}e_{J}e_{K} \rangle + \langle (e_{J}u_{r})_{,r}e_{I}e_{K}e_{L}, f \rangle + \langle fu_{r}, e_{K,r}e_{I}e_{J}e_{L} \rangle + \langle (e_{I}u_{r})_{,r}e_{J}e_{K}e_{L}, f \rangle$$

$$= \langle f, (e_{I}e_{J}e_{K}e_{L}u_{r})_{,r} \rangle + \langle f, e_{I}e_{J}e_{K}e_{L}u_{r,r} \rangle +$$

$$= -\langle u[f], e_{I}e_{J}e_{K}e_{L} \rangle = -(u[f])_{IJKL}$$
(3.163)

The contributions $\not \propto \delta_{mp} \delta_{nq}$ to (3.162) are

$$\sum_{M} \{ \delta_{mp}[\langle e_{M}e_{L}, u_{q,n} \rangle \langle e_{I}e_{J}e_{K}e_{M}, f \rangle - \langle e_{J}e_{M}, u_{q,n} \rangle \langle e_{I}e_{M}e_{K}e_{L}, f \rangle] + \delta_{nq}[\langle e_{M}e_{K}, u_{p,m} \rangle \langle e_{I}e_{J}e_{M}e_{L}, f \rangle - \langle e_{I}e_{M}, u_{p,m} \rangle \langle e_{M}e_{J}e_{K}e_{L}, f \rangle] \} = \delta_{mp} \langle fe_{I}e_{J}e_{K}e_{L}, [u_{q,n} - u_{q,n}] \rangle + \delta_{nq} \langle fe_{I}e_{J}e_{K}e_{L}, [u_{p,m} - u_{p,m}] \rangle = 0$$
(3.164)

Since

$$-(u[f])_{IJKL}\delta_{mp}\delta_{nq}\epsilon_{tij}\epsilon_{tkl} = -(u[f])_{\alpha\beta\gamma\delta}$$
(3.165)

we find that (3.157) reduces to

$$i\left[-\frac{1}{4}\sum_{\alpha\beta\gamma\delta\leq M_0}\left(-u[f]\right)_{\alpha\beta\gamma\delta}:F_{\alpha}F_{\beta}G_{\gamma}G_{\delta}:\right]$$
(3.166)

Thus, e.g. for the pattern M_5 before M_1, M_2 before M_6 before M_3, M_4

$$[C_2(u), C_{0;4}(f)] = i C_{0;4}(-u[f])$$
(3.167)

Altogether

$$[C(u), C_0(f)] = i C_0(-u[f])$$
(3.168)

6. Hamiltonian - Hamiltonian

We have to compute three curly brackets:

6A.

The first curly bracket is, see (A.8)

$$\frac{1}{4} \sum_{\alpha \leq M_1, \beta \leq M_2, \gamma \leq M_3, \mu \leq M_4, \nu \leq M_5, \rho \leq M_6} [f_{\alpha\beta\gamma} g_{\mu\nu\rho} - f \leftrightarrow g] [: F_{\alpha}F_{\beta}G_{\gamma} : , : F_{\mu}F_{\nu}G_{\rho} :]$$

$$[: F_{\alpha}F_{\beta}G_{\gamma} : , : F_{\mu}F_{\nu}G_{\rho} :]$$

$$= 4 [\delta_{\rho(\alpha} : F_{\beta}) F_{\mu} F_{\nu} G_{\gamma} : -\delta_{\gamma(\mu} : F_{\nu}) F_{\alpha} F_{\beta} G_{\rho} :] + 8 [\delta_{\gamma(\mu} \delta_{\nu)(\alpha} : F_{\beta}) G_{\rho} : -\delta_{\rho(\alpha} \delta_{\beta)(\mu} : F_{\nu}) G_{\gamma} :]$$
(3.169)

6A.i.

We consider first the terms containing two Kronecker factors and pick the pattern M_1, M_2, M_4, M_5 before M_3, M_6 . Using sandwiching, this allows to solve for μ, ν at locked ρ and α or β and to solve for α, β at locked γ and μ or ν . Hence this term, without the $f \leftrightarrow g$ contribution is given by

$$\frac{1}{2} \sum_{\gamma \leq M_3} \{ \sum_{\beta,\rho \leq M_0; \alpha \leq M_1} f_{\alpha\beta\gamma} \left[g_{\gamma\alpha\rho} + g_{\alpha\gamma\rho} \right] : F_\beta G_\rho : + \sum_{\alpha,\rho \leq M_0; \beta \leq M_2} f_{\alpha\beta\gamma} \left[g_{\gamma\beta\rho} + g_{\beta\gamma\rho} \right] : F_\alpha G_\rho : \}$$

$$= -\frac{1}{2} \sum_{\rho \leq M_6} \{ \sum_{\nu,\gamma \leq M_0; \mu \leq M_4} g_{\mu\nu\rho} \left[f_{\rho\mu\gamma} + f_{\mu\rho\gamma} \right] : F_\nu G_\gamma : + \sum_{\mu,\gamma \leq M_0; \nu \leq M_5} g_{\mu\nu\rho} \left[f_{\rho\nu\gamma} + f_{\nu\rho\gamma} \right] : F_\mu G_\gamma : \}$$

$$= \frac{1}{2} \sum_{\gamma \leq M_3; \beta, \rho \leq M_0} \{ \left[\sum_{\alpha \leq M_1} f_{\alpha\beta\gamma} + \sum_{\alpha \leq M_2} f_{\beta\alpha\gamma} \right] \left[g_{\gamma\alpha\rho} + g_{\alpha\gamma\rho} \right] : F_\beta G_\rho : \}$$

$$-\frac{1}{2} \sum_{\rho \leq M_6; \nu, \gamma \leq M_0} \{ \left[\sum_{\mu \leq M_4} g_{\mu\nu\rho} + \sum_{\mu \leq M_5} g_{\nu\mu\rho} \right] \left[f_{\rho\mu\gamma} + f_{\mu\rho\gamma} \right] : F_\nu G_\gamma : \}$$
(3.170)

In contrast to previous considerations, this does not vanish by inspection due to symmetry properties of the coefficients or when picking symmetric ranges $M_1 = M_2 = M_4 = M_5$ and $M_3 = M_6$ and not even when subtracting $f \leftrightarrow g$. To see that it vanishes nonetheless requires to perform the sums in the prescribed order. We pick $M_1 = M_2 = M_4 = M_5$, $M_3 = M_6$ and relabel μ, ν, γ by α, β, ρ in the second term to simplify (3.169) which becomes including the $f \leftrightarrow g$ term

$$\frac{1}{2} \sum_{\gamma \le M_3; \beta, \rho \le M_0, \alpha \le M_1} : F_{\beta} G_{\rho} : \left(\left\{ [f_{\alpha\beta\gamma} + f_{\beta\alpha\gamma}] [g_{\gamma\alpha\rho} + g_{\alpha\gamma\rho}] - [g_{\alpha\beta\gamma} + g_{\beta\alpha\gamma}] [f_{\gamma\alpha\rho} + f_{\alpha\gamma\rho}] \right\} - f \leftrightarrow g \right)$$

$$= 2 \sum_{\gamma \le M_3; \beta, \rho \le M_0, \alpha \le M_1} : F_{\beta} G_{\rho} : \left([f_{\alpha\beta\gamma} + f_{\beta\alpha\gamma}] [g_{\gamma\alpha\rho} + g_{\alpha\gamma\rho}] - f \leftrightarrow g \right) \quad (3.171)$$

We remove the cut-off M_1 , perform the sum over α and get with $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \rho = (Kpk), \beta = (Kpk$

(Lql) for the term in the round bracket of (3.170) without the $f \leftrightarrow g$ term

$$\sum_{I} [\langle (e_{I}e_{J}f)_{,n}, e_{K} \rangle \ \delta_{mp}\epsilon_{ijk} - \langle (e_{J}e_{I}f)_{,m}, e_{K} \rangle \ \delta_{np}\epsilon_{ijk}] \times [\langle (e_{K}e_{I}g)_{,m}, e_{L} \rangle \ \delta_{pq}\epsilon_{kil} - \langle (e_{I}e_{K}g)_{,p}, e_{L} \rangle \ \delta_{mq}\epsilon_{kil}] \\ = 2 \ \delta_{jl} \sum_{I} [\langle e_{I}e_{J}f, e_{K,n} \rangle \ \delta_{mp} - \langle e_{J}e_{I}f, e_{K,m} \rangle \ \delta_{np}] \times [\langle e_{K}e_{I}g, e_{L,m} \rangle \ \delta_{pq} - \langle e_{I}e_{K}g, e_{L,p} \rangle \ \delta_{mq}] \\ = 2 \ \delta_{jl} \sum_{I} [\langle e_{I}e_{J}f, e_{K,n} \rangle \langle e_{K}e_{I}g, e_{L,p} \rangle \ \delta_{pq} - \langle e_{I}e_{J}f, e_{K,n} \rangle \langle e_{I}e_{K}g, e_{L,p} \rangle \ \delta_{pq} \\ - \langle e_{J}e_{I}f, e_{K,m} \rangle \ \delta_{np} \ \langle e_{K}e_{I}g, e_{L,m} \rangle \ \delta_{pq} + \langle e_{J}e_{I}f, e_{K,q} \rangle \ \delta_{np} \ \langle e_{I}e_{K}g, e_{L,p} \rangle] \\ = -2 \ \delta_{jl} \ \langle e_{J} \ e_{K} \ [e_{K,m} \ e_{L,m} \ \delta_{pq} - e_{K,q} \ e_{L,p}], \ fg > \ \delta_{np}$$

$$(3.172)$$

Since (3.172) is symmetric under $f \leftrightarrow g$, (3.171) vanishes.

If instead we choose the pattern $M_3 = M_6$ before $M_1 = M_2 = M_4 = M_5$ then we must solve the two Kroneckers either for γ and e.g. one of α, β or for ρ and e.g. one of μ, ν . With sandwiching and noticing that with the chosen synchronised ranges of the cut-offs in (3.170) the first term is minus the second term under exchange of (α, β, γ) with (μ, ν, ρ) we find

$$-\sum_{\alpha,\beta,\mu,\nu\leq M_{1},\gamma,\rho\leq M_{3}} [f_{\alpha\beta\gamma} \ g_{\mu\nu\rho} - f \leftrightarrow g] \left[\delta_{\rho\alpha}\delta_{\beta\mu} : F_{\nu}G_{\gamma} : +\delta_{\rho\beta}\delta_{\alpha\mu} : F_{\nu}G_{\gamma} : +\delta_{\rho\alpha}\delta_{\beta\nu} : F_{\mu}G_{\gamma} : +\delta_{\rho\beta}\delta_{\alpha\nu} : F_{\mu}G_{\gamma} : \right]$$

$$= -\{\sum_{\nu\gamma\leq M_{0};\alpha,\beta\leq M_{1}} [f_{\alpha\beta\gamma} \ g_{\beta\nu\alpha} - f \leftrightarrow g] : F_{\nu}G_{\gamma} : +\sum_{\nu\gamma\leq M_{0};\alpha,\beta\leq M_{1},\gamma} [f_{\alpha\beta\gamma} \ g_{\alpha\nu\beta} - f \leftrightarrow g] : F_{\nu}G_{\gamma} : +\sum_{\mu\gamma\leq M_{0};\alpha,\beta\leq M_{1}} [f_{\alpha\beta\gamma} \ g_{\mu\alpha\beta} - f \leftrightarrow g] : F_{\mu}G_{\gamma} : +\sum_{\mu,\alpha\leq M_{0};\alpha,\beta\leq M_{1}} [f_{\alpha\beta\gamma} \ g_{\mu\alpha\beta} - f \leftrightarrow g] : F_{\mu}G_{\gamma} : +\sum_{\mu,\alpha\leq M_{0};\alpha,\beta\leq M_{1}} [f_{\alpha\beta\gamma} \ g_{\mu\alpha\beta} - f \leftrightarrow g] : F_{\mu}G_{\gamma} :$$

$$= -\sum_{\mu,\gamma\leq M_{0};\alpha,\beta\leq M_{1}} f_{\alpha\beta\gamma} \ [g_{\beta\mu\alpha} + g_{\alpha\mu\beta} + g_{\mu\beta\alpha} + g_{\mu\alpha\beta} - f \leftrightarrow g] : F_{\mu}G_{\gamma} :$$

$$(3.173)$$

In this expression we no longer have the freedom to carry out one limit before the other. We have with $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \mu = (Lql)$

$$\sum_{\alpha,\beta \leq M_{1}} f_{\alpha\beta\gamma} \left[g_{\beta\nu\alpha} + g_{\alpha\nu\beta} + g_{\nu\beta\alpha} + g_{\nu\alpha\beta} - f \leftrightarrow g \right]$$

$$= \sum_{I,J \leq M_{1}} f_{IJnK} \delta_{mp} \epsilon_{ijk} \left[g_{JLqI} \delta_{nm} \epsilon_{jli} + g_{ILqJ} \delta_{mn} \epsilon_{ilj} + g_{LJnI} \delta_{qm} \epsilon_{lji} + g_{LImJ} \delta_{qn} \epsilon_{lij} - f \leftrightarrow g \right]$$

$$= 2\delta_{kl} \sum_{I,J \leq M_{1}} f_{IJnK} \left[g_{JLqI} \delta_{np} - g_{ILqJ} \delta_{np} - g_{LJnI} \delta_{qp} + g_{LIpJ} \delta_{qn} - f \leftrightarrow g \right]$$

$$= -2\delta_{kl} \sum_{I,J \leq M_{1}} < e_{I}e_{J}f, e_{K,n} > \left[- < (e_{J}e_{L}g)_{,n}, e_{I} > \delta_{qp} + < (e_{I}e_{L}g)_{,p}, e_{J} > \delta_{qn} - f \leftrightarrow g \right] (3.174)$$

where we used that $f_{IJnK} = f_{JInK}$ to interchange I, J in the second term at finite M_1 to see that it cancels against the first term which again does not vanish at any finite M_1 . Now we remove M_1 and can perform the sum over one of I or J using the completeness relation and obtain after relabelling the remaining of J, Iinto I

$$-2\delta_{kl}\sum_{I} \left[- < (e_{I}e_{L}g)_{,r}, fe_{I}e_{K,r} > \delta_{qp} + < (e_{I}e_{L}g)_{,p}, fe_{I}e_{K,q} > -f \leftrightarrow g\right]$$

$$2\delta_{kl} \left[< \Lambda e_{L}, [g_{,r}f - f_{,r}g]e_{K,r} > \delta_{qp} - < \Lambda e_{L}[g_{,p}, f - f_{,p}g]e_{K,q} > \right]$$
(3.175)

where we introduced the quantity

$$\Lambda(x) := \lim_{M \to \infty} \Lambda_M(x), \ \Lambda_M(x) = \sum_{I \le M} e_I(x)^2$$
(3.176)

As already mentioned, Λ is formally given by $\delta(x, x)$ which is divergent. For both $\sigma = T^3$, \mathbb{R}^3 we have

$$\Lambda_M = \left[\frac{\zeta_M(0)}{\pi}\right]^3, \ \zeta_M(z) := \sum_{n=1}^M n^{-z}$$
(3.177)

where $\zeta_M(z)$ is the truncated Riemann zeta function [32] whose limit converges for $\Re(z) > 1$. Thus $\Lambda_M(x)$ is in this case independent of x. To regularise $\zeta_M(0)$ we may use the analytic extension of the zeta function which gives $\zeta(0) = -\frac{1}{2}$. Given any such regularisation of Λ we would conclude that in the pattern $M_3 = M_6$ before $M_1 = M_2 = M_4 = M_5$ the contribution to the curly bracket from the double Kronecker symbol is given by

$$-2\sum_{K,L\leq M_0} \delta_{kl} \left[<\Lambda e_L, \omega_r e_{K,r} > \delta_{qp} - <\Lambda e_L \omega_p e_{K,q} > \right] : F_{Lql}; G_{Kpk} :, \ \omega_r := fg_{,r} - gf_{,r}$$
(3.178)

To interpret this result consider

$$\int d^{3}x \, \delta^{ab} \omega_{a} : (\partial_{b}A_{c}^{k} - \partial_{c}A_{b}^{k})E_{k}^{c} := \frac{i}{2} \sum_{K,L} \delta^{rp} \, \delta_{kl} [<\omega_{r}, e_{K,p}e_{L} >: G_{Kqk}F_{Lql} := -<\omega_{r}, e_{K,q}e_{L} >: G_{Kpk}F_{Lql} :]$$

$$= \frac{i}{2} \sum_{K,L} \delta^{rp} \, \delta_{kl} [\delta_{pg} -] : G_{Kpk}F_{Lql} :]$$
(3.179)

It follows that (3.178) is given by

$$4i \int d^D x \left[\Lambda \omega_a\right] \delta^{ab} : (\partial_b A^k_c - \partial_c A^k_b) E^c_k : \qquad (3.180)$$

6A.ii.

We now focus on the single Kronecker symbol contribution to (3.169). In both schemes that we discuss for the first curly bracket we see that the second contribution to the single Kronecker term is the same as the first with α, β, γ interchanged with μ, ν, ρ with a minus sign. Relabelling indices produces another minus sign from the antisymmetric combination of the f, g dependence. Thus we may drop the second contribution and instead multiply by a factor of two. Then μ, ν, γ are locked by sandwiching and we solve for α or β using the $M_1 = M_2$ before M_3 pattern while we solve for ρ in the M_3 before $M_1 = M_2$ pattern.

The first pattern gives

$$\sum_{\substack{\alpha,\mu,\beta,\nu \leq M_{1};\gamma,\rho \leq M_{3}}} [f_{\alpha\beta\gamma} \ g_{\mu\nu\rho} - f \leftrightarrow g] : F_{\mu}F_{\nu}[\delta_{\rho\alpha}F_{\beta} + \delta_{\rho\beta}F_{\alpha}]G_{\gamma} :$$

$$= \sum_{\substack{\mu,\nu,\beta,\gamma \leq M_{0};\rho \leq M_{3}}} [f_{\rho\beta\gamma} \ g_{\mu\nu\rho} - f \leftrightarrow g] : F_{\mu}F_{\nu}F_{\beta}G_{\gamma} : + \sum_{\substack{\mu,\nu\alpha,\gamma \leq M_{0};\rho \leq M_{3}}} [f_{\alpha\rho\gamma} \ g_{\mu\nu\rho} - f \leftrightarrow g] : F_{\mu}F_{\nu}F_{\alpha}G_{\gamma} :$$

$$= \sum_{\substack{\mu,\nu,\alpha,\gamma \leq M_{0};\rho \leq M_{3}}} [(f_{\rho\alpha\gamma} + f_{\alpha\rho\gamma}) \ g_{\mu\nu\rho} - f \leftrightarrow g] : F_{\mu}F_{\nu}F_{\alpha}G_{\gamma} : \qquad (3.181)$$

and the second

$$\sum_{\substack{\alpha,\mu,\beta,\nu\leq M_{1};\gamma,\rho\leq M_{3}}} [f_{\alpha\beta\gamma} \ g_{\mu\nu\rho} - f \leftrightarrow g] : F_{\mu}F_{\nu}[\delta_{\rho\alpha}F_{\beta} + \delta_{\rho\beta}F_{\alpha}]G_{\gamma} :$$

$$= \sum_{\substack{\mu,\nu,\beta,\gamma\leq M_{0};\alpha\leq M_{1}}} [f_{\alpha\beta\gamma} \ g_{\mu\nu\alpha} - f \leftrightarrow g] : F_{\mu}F_{\nu} \ F_{\beta} \ G_{\gamma} : + \sum_{\substack{\mu,\nu,\alpha,\gamma\leq M_{0};\beta\leq M_{1}}} [f_{\alpha\beta\gamma} \ g_{\mu\nu\beta} - f \leftrightarrow g] : F_{\mu}F_{\nu}F_{\alpha} \ G_{\gamma} :$$

$$= \sum_{\substack{\mu,\nu,\alpha,\gamma\leq M_{0};\beta\leq M_{1}}} [(f_{\alpha\beta\gamma} + f_{\beta\alpha\gamma}) \ g_{\mu\nu\beta} - f \leftrightarrow g] : F_{\mu}F_{\nu}F_{\alpha} \ G_{\gamma} :$$

$$= \sum_{\substack{\mu,\nu,\alpha,\gamma\leq M_{0};\rho\leq M_{1}}} [(f_{\alpha\rho\gamma} + f_{\rho\alpha\gamma}) \ g_{\mu\nu\rho} - f \leftrightarrow g] : F_{\mu}F_{\nu}F_{\alpha} \ G_{\gamma} :$$

$$(3.182)$$

where we have relabelled β by ρ in the last step. Thus except for the range, (3.181) and (3.182) are identical. This is in general true for terms with a single Kronecker symbol.

The remaining cut-off M_3 or M_1 can now be removed and we have with $\alpha = (Imi), \gamma = (Jnj), \mu = (Mur), \nu = (Nvs), \rho = (Kpk)$

$$\sum_{\rho} [(f_{\rho\alpha\gamma} + f_{\alpha\rho\gamma}) g_{\mu\nu\rho} - f \leftrightarrow g]$$

$$= \sum_{K} [f_{KImJ}\delta_{pn}\epsilon_{kij} + f_{IKpJ}\delta_{mn}\epsilon_{ikj} - f \leftrightarrow g] g_{MNvK}\delta_{up}\epsilon_{rsk}$$

$$= \sum_{K} [f_{KImJ}\delta_{un} - f_{IKuJ}\delta_{mn} - f \leftrightarrow g] \epsilon_{kij}g_{MNvK}\epsilon_{krs}$$

$$= -\sum_{K} [\langle e_{K}e_{I}f, e_{J,m} \rangle \delta_{un} - \langle e_{K}e_{I}f, e_{J,u} \rangle \delta_{mn} - f \leftrightarrow g] \langle (e_{M}e_{N}g)_{,v}, e_{K} \rangle \epsilon_{kij}\epsilon_{krs}$$

$$= -[\langle (e_{M}e_{N}g)_{,v}e_{I}f, e_{J,m} \rangle \delta_{un} - \langle (e_{M}e_{N}g)_{,v}e_{I}f, e_{J,u} \rangle \delta_{mn} - f \leftrightarrow g] \epsilon_{kij}\epsilon_{krs}$$

$$= -[\langle e_{M}e_{N}e_{I}\omega_{v}, e_{J,m} \rangle \delta_{un} - \langle e_{M}e_{N}e_{I}\omega_{v}, e_{J,u} \rangle \delta_{mn}] \epsilon_{kij}\epsilon_{krs} \qquad (3.183)$$

Thus the single Kronecker symbol contribution to the curly bracket is

$$-\sum_{I,J,M,N \le M_{0}} [\langle e_{M}e_{N}e_{I}\omega_{v}, e_{J,m} \rangle : F_{Mnr}F_{Nvs}F_{Imi}G_{Jnj} : - \langle e_{M}e_{N}e_{I}\omega_{v}, e_{J,u} \rangle : F_{Mur}F_{Nvs}F_{Ini}G_{Jnj} :]\epsilon_{kij}\epsilon_{krs} - \sum_{I,J,M,N \le M_{0}} \langle e_{M}e_{N}e_{I}\omega_{v}, e_{J,m} \rangle : (F_{Mnr}F_{Imi} - F_{Mmr}F_{Ini})F_{Nvs}G_{Jnj} : (\delta_{ir}\delta_{js} - \delta_{is}\delta_{jr})$$

$$= -\sum_{I,J,M,N \le M_{0}} \langle e_{M}e_{N}e_{I}\omega_{v}, e_{J,m} \rangle \{ : (F_{Mnr}F_{Imr} - F_{Inr}F_{Mmr})F_{Nvj}G_{Jnj} : - i : (F_{Mnj}F_{Imi} - F_{Mmj}F_{Ini})F_{Nvi}G_{Jnj} :\}$$

$$= \sum_{I,J,M,N \le M_{0}} \langle e_{M}e_{N}e_{I}\omega_{v}, e_{J,m} \rangle : [(F_{Mnj}F_{Imi} - F_{Mmj}F_{Ini})F_{Nvi}G_{Jnj} :]$$

$$(3.184)$$

where in the step before the last we used that the normal ordered factor in the first term is antisymmetric in I, M while the prefactor coefficient is symmetric.

To interpret this result consider

$$\int d^{3}x \,\omega_{a} \,\delta^{ij} : E_{i}^{a} E_{j}^{b} (\partial_{b} A_{c}^{k} - \partial_{c} A_{b}^{k}) E_{k}^{c} :$$

$$= -\frac{i}{4} \sum_{I,J,M,N} < \omega_{v} e_{N} e_{M} e_{I}, e_{J,m} > \{ : (F_{Mmi} F_{Inj} - F_{Mni} F_{Imj}) G_{Jnj} F_{Nvi} :$$

$$= -\frac{i}{4} \sum_{I,J,M,N} < \omega_{v} e_{N} e_{M} e_{I}, e_{J,m} > \{ : (F_{Imi} F_{Mnj} - F_{Ini} F_{Mmj}) G_{Jnj} F_{Nvi} : \qquad (3.185)$$

which is precisely -i/4 times (3.184).

We summarise

$$[C_{0;3}(f), C_{0;3}(g)] = 4i \int d^3x \left[f g_{,a} - g f_{,a} \right] : \left[E^a_i e^b_j \delta^{ij} + \Lambda \delta^{ab} \right] \left(2\partial_{[b} A^k_{c]} \right) E^c_k$$
(3.186)

where $\Lambda = 0$ e.g. for the limiting pattern with $M_1 = M_2 = M_4 = M_5$ before $M_3 = M_6$ and $\Lambda \neq 0$ is a regularisation of $\delta(0,0)$ e.g. for the pattern $M_3 = M_6$ before $M_1 = M_2 = M_4 = M_5$.

6B.

The second curly bracket is given by

$$\frac{1}{4\ 2^{1/2}} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \mu \le M_4, \nu \le M_5, \rho \le M_6, \sigma \le M_7} \left[f_{\alpha\beta\gamma} \ g_{\mu\nu\rho\sigma} - f \leftrightarrow g \right] \left[: F_{\alpha}F_{\beta}G_{\gamma} : , : F_{\mu} \ F_{\nu} \ G_{\rho} \ G_{\sigma} : \right] (3.187)$$

where the evaluation of the commutator is given in (A.9).

6B.i

This is the first time that we have to deal with a triple product of Kronecker symbols whose contribution is

$$[:F_{\alpha}F_{\beta}G_{\gamma}:,:F_{\mu}F_{\nu}G_{\rho}G_{\sigma}:]_{\delta^{3}} = -16 F_{(\mu}\delta_{\nu)(\alpha}\delta_{\beta)(\rho}\delta_{\sigma)\gamma} - 8 F_{(\mu}\delta_{\nu)\gamma}\delta_{\alpha(\rho}\delta_{\sigma)\beta}$$
(3.188)

Whatever scheme we choose, the Kronecker symbols identify the same finite range sub-indices m = n, i = jin $\delta_{\alpha,\beta}$ if $\alpha = (Imi), \beta = (Jnj)$. Thus if we can show that (3.187) resticted to (3.188) vanishes due to symmetries among the finite range sub-indices in one scheme at finite remaining cut-offs, it does so in any other because different schemes differ only in the order in which the surviving ininite range indices are summed to ∞ . For this purpose we consider the pattern M_4, M_5, M_6, M_7 before M_1, M_2, M_3 and solve for μ, ν, ρ, σ . After relabelling, sandwiching this results in a term proportional to

$$\sum_{\mu \leq M_0} F_{\mu} \sum_{\alpha \leq M_1, \beta \leq M_2, \ \gamma \leq M_3} \times \left\{ f_{\alpha\beta\gamma} \left[(g_{\mu\alpha\beta\gamma} + g_{\mu\alpha\gamma\beta} + g_{\alpha\mu\beta\gamma} + g_{\alpha\mu\gamma\beta}) + (g_{\mu\beta\alpha\gamma} + g_{\mu\beta\gamma\alpha} + g_{\beta\mu\alpha\gamma} + g_{\beta\mu\gamma\alpha}) + (g_{\mu\gamma\alpha\beta} + g_{\mu\gamma\beta\alpha} + g_{\gamma\mu\alpha\beta} + g_{\gamma\mu\beta\alpha}) \right] - \right\}$$

With $\alpha = (Imi)$, $\beta = (Jnj)$, $\gamma = (Kpk)$, $\mu = (Lql)$ we note that the three groups of four g's is proportional to $\epsilon_{tli} \epsilon_{tjk}$, $\epsilon_{tlj} \epsilon_{tik}$, $\epsilon_{tlk} \epsilon_{tij}$ respectively while $f_{\alpha\beta\gamma}$ is proportional to ϵ_{ijk} . Summing over i, j, k = 1, 2, 3 gives respectively $2 \epsilon_{tli} \delta_{ti}$, $-2\epsilon_{tlj} \delta_{tj}$, $2\epsilon_{tlk} \delta_{tk}$ which all vanish when summing over t.

6B.ii

The double Kronecker contribution to the commutator in (3.187) is given by

$$[:F_{\alpha}F_{\beta}G_{\gamma}:,:F_{\mu} F_{\nu} G_{\rho} G_{\sigma}:]_{\delta^{2}} = 8 :F_{\mu} F_{\nu} F_{(\alpha}: \delta_{\beta)(\rho} \delta_{\sigma)\gamma} + 8 :G_{\rho} G_{\sigma} F_{(\alpha}: \delta_{\beta)(\mu} \delta_{\nu)\gamma} - 16 :G_{\gamma} G_{(\rho} \delta_{\sigma)(\alpha} \delta_{\beta)(\mu} F_{\nu)}$$

$$(3.190)$$

We consistently keep the pattern M_4, M_5, M_6, M_7 before M_1, M_2, M_3 .

Then the first term in (3.190) gives using sandwiching and relabelling a contribution proportional to, when solving for ρ, σ

$$2\sum_{\mu,\nu,\alpha\leq M_0} : F_{\mu} F_{\nu} F_{\alpha}; \sum_{\gamma\leq M_3} \left(\left[\sum_{\beta\leq M_2} f_{\alpha\beta\gamma} + \sum_{\beta\leq M_1} f_{\beta\alpha\gamma} \right] \left[g_{\mu\nu\beta\gamma} + g_{\mu\nu\gamma\beta} \right] - f \leftrightarrow g \right)$$
(3.191)

With $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \ \mu = (Mur), \ \nu = (Nvs)$ we have explicitly

$$f_{\alpha\beta\gamma} = -\langle e_I e_J f, e_{K,n} \rangle \delta_{mp} \epsilon_{ijk}, \ f_{\beta\alpha\gamma} = \langle e_I e_J f, e_{K,m} \rangle \delta_{np} \epsilon_{ijk}, g_{\mu\nu\beta\gamma} = \langle e_M e_N e_J e_K, g \rangle \delta_{un} \delta_{vp} \epsilon_{trs} \epsilon_{tjk}, \ g_{\mu\nu\gamma\beta} = -\langle e_M e_N e_J e_K, g \rangle \delta_{up} \delta_{vn} \epsilon_{trs} \epsilon_{tjk},$$
(3.192)

If we remove M_1, M_2 at finite M_3 we can perform the sum over J in (3.191) and obtain an expression which depends only on fg and from which we subtract the same expression with $f \leftrightarrow g$. Thus (3.191) vanishes in this pattern. If on the other hand we remove M_3 at finite M_1, M_2 then we possibly obtain a contribution which renormalises the cosmological constant because that term depends on the derivatives of f, g.

To see whether this is the case we substitute (3.192) into (3.191) and obtain

$$4\sum_{\mu,\nu,\alpha\leq M_{0}} : F_{\mu} F_{\nu} F_{\alpha} : \epsilon_{irs} \sum_{K\leq M_{3}} (\left[\sum_{J\leq M_{2}} [f_{IJnK}\delta_{mp} - \sum_{J\leq M_{1}} f_{JImK}\delta_{np}] g_{MNJK} [\delta_{un}\delta_{vp} - \delta_{up}\delta_{vn}] - f \leftrightarrow g\right)$$

$$= 4\sum_{\mu,\nu,\alpha\leq M_{0}} : F_{\mu}F_{\nu}F_{\alpha} : \epsilon_{irs} \sum_{K\leq M_{3}} (\{\sum_{J\leq M_{2}} [\delta_{un}\delta_{vm} - \delta_{um}\delta_{vn}] - \sum_{J\leq M_{1}} [\delta_{un}\delta_{vn} - \delta_{un}\delta_{vn}]\}f_{IJnK}g_{MNJK} - f \leftrightarrow g)$$

$$= 4\sum_{\mu,\nu,\alpha\leq M_{0}} : F_{\mu} F_{\nu} F_{\alpha} : \epsilon_{irs} \sum_{K\leq M_{3}} \sum_{J\leq M_{2}} (g_{MNJK} [f_{IJuK} \delta_{vm} - f_{IJvK} \delta_{um}] - f \leftrightarrow g)$$
(3.193)

We remove M_3 and perform the sum over K unconstrained which gives

$$-4 \sum_{M,N,I \le M_0} : F_{Mur} F_{Nvs} F_{Imi} : \epsilon_{irs} \sum_{J \le M_2} \times (\langle e_J^2 e_M e_N e_I, [f_{,u} g - f g_{,u}] \rangle \delta_{vm} - \langle e_J^2 e_M e_N e_I, [f_{,v} g - f g_{,v}] \rangle \delta_{um})$$

$$= -8 \sum_{M,N,I \le M_0} : F_{Mur} F_{Nvs} F_{Imi} : \epsilon_{irs} \sum_{J \le M_2} \langle e_J^2 e_M e_N e_I, [f_{,u} g - f g_{,u}] \rangle \delta_{vm}$$

$$= 8 \sum_{M,N,I \le M_0} \sum_{J \le M_2} : F_{Mur} F_{Nvs} F_{Ivi} : \epsilon_{irs} \langle e_J^2 e_M e_N e_I, \omega_u \rangle$$

$$= 0 \qquad (3.194)$$

with $\omega_u = f g_{,u} - f_{,u} g$. Here we have relabelled M, u, r with N, v, s in the second step and used commutativity under the normal ordering symbol while in the last step we used that $\epsilon_{irs} < e_J^2 e_M e_N e_I, \omega_u >$ is antisymmetric under $(N, s) \leftrightarrow (I, i)$ while : $F_{Mur} F_{Nvs} F_{Ivi}$: is symmetric and that $I, N \leq M_0$ have the same range. Accordingly, there is no renormalisation of the cosmological constant term in this pattern.

The second term in (3.190) gives after sandwiching, relabelling and solving for μ, ν

$$2\sum_{\rho,\sigma,\alpha\leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{\gamma\leq M_3} \left(\left[\sum_{\beta\leq M_2} f_{\alpha\beta\gamma} + \sum_{\beta\leq M_1} f_{\beta\alpha\gamma} \right] \left[g_{\beta\gamma\rho\sigma} + g_{\gamma\beta\rho\sigma} \right] \right) - f \leftrightarrow g \right)$$
(3.195)

Using the expressions (3.192) we see by the same argument that in the pattern M_1, M_2 before M_3 the expression (3.195) vanishes. In the pattern M_3 before M_1, M_2 the contribution including the prefactors to the curly bracket is with $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \rho = (Mur), \sigma = (Nvs)$

$$\frac{1}{2 2^{1/2}} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{\gamma \leq M_3} \left(\left[\sum_{\beta \leq M_2} f_{\alpha\beta\gamma} + \sum_{\beta \leq M_1} f_{\beta\alpha\gamma} \right] \left[g_{\beta\gamma\rho\sigma} + g_{\gamma\beta\rho\sigma} \right] \right) - f \leftrightarrow g \right)$$

$$= \frac{1}{2 2^{1/2}} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \left\{ \left[\sum_{J \leq M_2} f_{IJnK} \delta_{mp} \epsilon_{ijk} - \sum_{J \leq M_1} f_{JImK} \delta_{np} \epsilon_{ijk} \right] \times \left[g_{KJMN} \delta_{pu} \delta_{nv} \epsilon_{ikj} \epsilon_{trs} - g_{JKMN} \delta_{nu} \delta_{pv} \epsilon_{tkj} \epsilon_{trs} \right] - f \leftrightarrow g \right\}$$

$$= -\frac{1}{2^{1/2}} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \epsilon_{irs} \left\{ \left[\sum_{J \leq M_2} f_{IJnK} \delta_{mp} - \sum_{J \leq M_1} f_{JImK} \delta_{np} \right] \times \left[g_{KJMN} \delta_{pu} \delta_{nv} - g_{JKMN} \delta_{nu} \delta_{pv} \right] - f \leftrightarrow g \right\}$$

$$= -\frac{1}{2^{1/2}} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \epsilon_{irs} \times \left\{ \sum_{J \leq M_2} f_{IJnK} g_{JKMN} \left[\delta_{nu} \delta_{nv} - \delta_{nu} \delta_{nv} \right] - f \leftrightarrow g \right\}$$

$$= -\frac{1}{2^{1/2}} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \epsilon_{irs} \left\{ \sum_{J \leq M_2} g_{JKMN} \left[\delta_{mu} f_{IJvK} - \delta_{mv} f_{IJuK} \right] - f \leftrightarrow g \right\}$$

$$= -2^{1/2} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \epsilon_{irs} \left\{ \sum_{J \leq M_2} g_{JKMN} \left[\delta_{mu} f_{IJvK} - \delta_{mv} f_{IJuK} \right] - f \leftrightarrow g \right\}$$

$$= -2^{1/2} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \epsilon_{irs} \left\{ \sum_{J \leq M_2} g_{JKMN} \left[\delta_{mu} f_{IJvK} - \delta_{mv} f_{IJuK} \right] - f \leftrightarrow g \right\}$$

$$= -2^{1/2} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \epsilon_{irs} \left\{ \sum_{J \leq M_2} g_{JKMN} \left[\delta_{mu} f_{IJvK} - \delta_{mv} f_{IJuK} \right] - f \leftrightarrow g \right\}$$

$$= -2^{1/2} \sum_{\rho,\sigma,\alpha \leq M_0} : G_{\rho} G_{\sigma} F_{\alpha}; \sum_{K \leq M_3} \epsilon_{irs} \left\{ \sum_{J \leq M_2} g_{JKMN} \left[\delta_{mu} f_{IJvK} - f \leftrightarrow g \right\}$$

$$(3.196)$$

where in the last step we interchanged Mur and Nvs and used commutativity under the normal ordering symbol.

We remove M_3 and sum unconstrained over K which gives

$$-2^{1/2} \sum_{M,N,I \leq M_0} : G_{Mur} \ G_{Nvs} \ F_{Imi}; \sum_{J \leq M_2} \sum_{K \leq M_3} \epsilon_{irs} \left\{ g_{JKMN} \ f_{IJvK} \ \delta_{mu} - f \leftrightarrow g \right\}$$

$$= -2^{1/2} \sum_{M,N,I \leq M_0} : G_{Mur} \ G_{Nvs} \ F_{Iui}; \sum_{J \leq M_2} \epsilon_{irs} \left\{ < e_J^2 \ e_M e_N e_I, [f_vg - f \ g_{,v}] > \right\}$$

$$= 2^{1/2} \sum_{M,N,I \leq M_0} : G_{Mur} \ G_{Nvs} \ F_{Iui}; \ \epsilon_{irs} < \Lambda_{M_2} \ e_M e_N e_I \omega_v >$$
(3.197)

where $\Lambda_M = \sum_{I \leq M} e_I^2$, $\omega_v = fg_{,v} - f_{,v}g$.

To interpret \overline{th} is term we consider

$$\int d^3x \,\Lambda\omega_a \delta^{ab} \,\epsilon_{ijk} \,: A^j_b \,A^k_c \,E^c_i := \frac{i}{2 \,2^{1/2}} \,\epsilon^{irs} \sum_{I,M,N} < \Lambda\omega_v e_I e_M e_N > G_{Nvr} G_{Mus} F_{Jui} \tag{3.198}$$

Thus (3.197) is 4i times (3.198).

The third term in (3.190) gives after sandwiching, relabelling and solving for μ or ν and ρ or σ

$$-2\sum_{\mu,\sigma,\gamma\leq M_{0}}:F_{\mu} G_{\sigma} G_{\gamma}:\sum_{\alpha\leq M_{1},\beta\leq M_{2}} \left(f_{\alpha\beta\gamma} \left[g_{\mu\alpha\beta\sigma}+g_{\mu\beta\alpha\sigma}+g_{\alpha\mu\beta\sigma}+g_{beta\mu\alpha\sigma}+g_{\mu\alpha\sigma\beta}+g_{\mu\beta\sigma\alpha}+g_{\alpha\mu\sigma\beta}+g_{beta\mu\sigma\alpha}\right]-f\leftrightarrow g\right)$$

$$(3.199)$$

This term vanishes, in both patterns M_1, M_2 before M_3 or M_3 before M_1, M_2 because M_3 is locked, by the same argument as before because we only get contributions depending on fg when e.g. removing M_1 before M_2 and then we subtract the same expression with f, g interchanged.

6B.iii

We now consider the single Kronecker symbol contribution to (3.187) which is

$$[:F_{\alpha}F_{\beta}G_{\gamma}:,:F_{\mu}F_{\nu}G_{\rho}G_{\sigma}:]_{\delta^{1}} = -4\delta_{\gamma(\mu}:F_{\nu})F_{\alpha}F_{\beta}G_{\rho}G_{\sigma}:+8:F_{(\alpha}\delta_{\beta)(\rho}G_{\sigma})F_{\mu}F_{\nu}G_{\gamma}:$$
(3.200)

Its contribution to (3.187) is obtained by sandwiching and solving the Kronecker for μ or ν in the first term and for ρ or σ in the second

$$-\frac{1}{2 2^{1/2}} \sum_{\alpha,\beta,\rho,\sigma,\nu \le M_0} : F_{\nu} F_{\alpha} F_{\beta} G_{\rho} G_{\sigma} : \left\{ \sum_{\gamma \le M_3} f_{\alpha\beta\gamma} \left[g_{\gamma\nu\rho\sigma} + g_{\nu\gamma\rho\sigma} \right] - f \leftrightarrow g \right] \\ + \sum_{\alpha,\sigma,\mu,\nu,\gamma \le M_0} : F_{\alpha} G_{\sigma} F_{\mu} F_{\nu} G_{\gamma} : \left\{ \left[\sum_{\beta \le M_2} f_{\alpha\beta\gamma} + \sum_{\beta \le M_1} f_{\beta\alpha\gamma} \right] \left[g_{\mu\nu\beta\sigma} + g_{\mu\nu\sigma\beta} \right] - f \leftrightarrow g \right\}$$
(3.201)

We can now remove the remaining cut-offs. Then we see that the terms with the remaining sums over β vanish because they produce just terms depending on fg and therefore subtraction of the same term with f, g interchanged leads to cancellation. We are thus left with with the first term in (3.201) given by

$$-\frac{1}{2^{1/2}} \sum_{\alpha,\beta,\rho,\sigma,\mu \le M_0;\gamma \le M_3} \{ f_{\alpha\beta\gamma} \left[g_{\gamma\mu\rho\sigma} + g_{\mu\gamma\rho\sigma} - f \leftrightarrow g \right\} : F_{\mu}F_{\alpha}F_{\beta}G_{\rho}G_{\sigma} :$$
(3.202)

Using $\alpha = (Imi), \beta = (Jnj), \gamma = (Kpk), \mu = (Mqr), \rho = (Sus), \sigma = (Tvt)$ we can perform the sum over γ unconstrained removing M_3 . We have

$$\sum_{\gamma} \{ f_{\alpha\beta\gamma} \left[g_{\gamma\mu\rho\sigma} + g_{\mu\gamma\rho\sigma} - f \leftrightarrow g \} \}$$

$$= \sum_{K} \{ f_{IJnK} \delta_{mp} \epsilon_{ijk} \left[g_{KMST} \delta_{pu} \delta_{qv} \epsilon_{lkr} \epsilon_{lst} + g_{MKST} \delta_{qu} \delta_{pv} \epsilon_{lrk} \epsilon_{lst} \right] - f \leftrightarrow g \}$$

$$= \sum_{K} \{ f_{IJnK} g_{KMST} \epsilon_{ijk} \epsilon_{lkr} \epsilon_{lst} \left[\delta_{mu} \delta_{qv} - \delta_{qu} \delta_{mv} \right] - f \leftrightarrow g \}$$

$$= \{ \langle (e_I e_J f)_{,n}, e_M e_S e_T g \rangle - f \leftrightarrow g \} \epsilon_{ijk} \epsilon_{lkr} \epsilon_{lst} \left[\delta_{mu} \delta_{qv} - \delta_{qu} \delta_{mv} \right]$$

$$= - \langle e_I e_J e_M e_S e_T, \omega_n \rangle \epsilon_{ijk} \epsilon_{lkr} \epsilon_{lst} \left[\delta_{mu} \delta_{qv} - \delta_{qu} \delta_{mv} \right]$$
(3.203)

Thus (3.202) becomes

$$\frac{1}{2^{1/2}} \sum_{I,J,M,S,T \leq M_{0}} \langle e_{I}e_{J}e_{M}e_{S}e_{T}, \omega_{n} \rangle \langle \epsilon_{ijk}\epsilon_{lkr}\epsilon_{lst} : F_{Mvr}F_{Imi}F_{Jnj}(G_{Sms}G_{Tvt} - G_{Svs}G_{Tmt}) :$$

$$= \frac{1}{2^{1/2}} \sum_{I,J,M,S,T \leq M_{0}} \langle e_{I}e_{J}e_{M}e_{S}e_{T}, \omega_{n} \rangle \langle \epsilon_{ijk}\epsilon_{lkr}\epsilon_{lst} : F_{Mvr}F_{Imi}F_{Jnj}(G_{Sms}G_{Tvt} - G_{Tvs}G_{Smt}) :$$

$$= \frac{1}{2^{1/2}} \sum_{I,J,M,S,T \leq M_{0}} \langle e_{I}e_{J}e_{M}e_{S}e_{T}, \omega_{n} \rangle \langle \epsilon_{ijk}\epsilon_{lkr}\epsilon_{lst} : F_{Mvr}F_{Imi}F_{Jnj}(G_{Sms}G_{Tvt} + G_{Tvt}G_{Sms}) :$$

$$= 2^{1/2} \sum_{I,J,M,S,T \leq M_{0}} \langle e_{I}e_{J}e_{M}e_{S}e_{T}, \omega_{n} \rangle \langle \delta_{ir}\delta_{jl} - \delta_{il}\delta_{jr} \rangle \epsilon_{lst} : F_{Mvr}F_{Imi}F_{Jnj}G_{Sms}G_{Tvt} :$$

$$= 2^{1/2} \sum_{I,J,M,S,T \leq M_{0}} \langle e_{I}e_{J}e_{M}e_{S}e_{T}, \omega_{n} \rangle \langle \epsilon_{lst} : (F_{Mvr}F_{Imr}F_{Jnl} - F_{Mvr}F_{Iml}F_{Jnr})G_{Sms}G_{Tvt} :$$

$$= 2^{1/2} \sum_{I,J,M,S,T \leq M_{0}} \langle e_{I}e_{J}e_{M}e_{S}e_{T}, \omega_{n} \rangle \langle \epsilon_{lst} : F_{Jnj}F_{Mvj}G_{Tvi}G_{Smk}F_{Iml} :$$
(3.204)

where in the last relation we used that $F_{Mvr}F_{Imr}$ is symmetric under exchange of m, v while $\epsilon_{lst}G_{Sms}G_{Tvt}$ is antisymmetric when both are contracted with the totally symmetric $\langle e_I e_J e_S e_T e_M, \omega_n \rangle$.

To interpret this expression consider

$$\int d^3x \,\omega_a : E^a_j \,E^b_j \,\epsilon_{lik} \,A^i_b \,A^k_c \,E^c_l := -\frac{i}{22^{1/2}} \sum_{I,J,M,S,T} < e_I e_J e_M e_S e_T, \omega_n > \epsilon_{lik} : F_{Jnj} F_{Mvj} G_{Tvi} G_{Smk} F_{Iml} :$$
(3.205)

hence (3.204) is 4i times (3.205).

Altogether we obtain

$$[C_{0;3}(f), C_{0;4}(g)] = 4i \int d^3x \,\omega_a \,: (E^a_i E^b_j \delta^{ij} + \Lambda \delta^{ab}) \epsilon_{ijk} A^i_b A^j_c E^c_k : \qquad (3.206)$$

e.g. in the pattern with M_4, M_5, M_6, M_7 before M_1, M_2, M_3 and M_1, M_2 before M_3 and M_1 before M_2 for $\Lambda = 0$ and M_3 before M_1, M_2 for $\Lambda \neq 0$ a regulator dependent constant.

6C.

The third curly bracket is given by

$$\frac{1}{32} \sum_{\alpha \le M_1, \beta \le M_2, \gamma \le M_3, \delta \le M_4, \mu \le M_5, \nu \le M_6, \rho \le M_7, \sigma \le M_8} [f_{\alpha\beta\gamma\sigma} g_{\mu\nu\rho\sigma} - f \leftrightarrow g] [: F_{\alpha}F_{\beta}G_{\gamma}G_{\delta} :, F_{\mu}F_{\nu}G_{\rho}G_{\sigma} :]$$

$$(3.207)$$

where the commutator is worked out in (A.10).

Since $f_{\alpha\beta\gamma\delta}$, $g_{\mu\nu\rho\sigma}$ do not depend on derivatives of f, g, (3.207) vanishes for any pattern such that after solving the Kronecker symbols we obtain a sum that depends on cut-offs which are to be taken away sequentially. Consider e.g. the pattern M_n before M_{n-1} with n = 2, ..., 8. Then we can solve all Kronecker symbols for $\mu, \nu \rho, \sigma$. In the terms with 1, 2, 3 Kronecker factors the normal ordered monomial carries 6, 4, 2 indices that are locked by sandwiching and 1, 2, 3 of the indices μ, ν, ρ, σ are equated with one of $\alpha, \beta, \gamma, \delta$ so that 1, 2, 3 indices are left over for summing. Since these are summed sequentially according to the chosen pattern, removal of the top cut-off leads to an expression that depends just on fg minus the same expression depending on gf. Thus for the pattern $M_8 > M_7 > ... > M_1$ we find

$$[C_{0;4}(f), C_{0;4}(g)] = 0 (3.208)$$

Altogether

$$[C_0(f), C_0(g)] = i \int d^3x \left[4(fg_{,a} - f_{,a}g)\right] : \left(E_i^a E_j^b \delta^{ij} + \Lambda \delta^{ab}\right) F_{bc}^k E_k^c :$$
(3.209)

The course of the proof has revealed the following result:

Corollary 3.1.

There exist limiting patterns for the commutator of two normal ordered Hamiltonian constraints such that with $\omega = f \, dg - g \, df$

$$[C_0(f), C_0(g)] = 4i \int d^3x \,\omega_a \,: (\delta^{mn} \, E^a_m \, E^b_n + \Lambda \, \delta^{ab}) \, F^j_{bc} \, E^c_j : \qquad (3.210)$$

where Λ is a regularised constant depending on the limiting pattern.

Remarks:

0.

The investigation illustrates the fact that different limiting patterns lead to different results. We did not consider the most general pattern but focussed on those that tend to avoid derivatives of test functions in potentially divergent contributions.

1.

Note that we work in Planck units, restoring SI units would reveal that the Λ term is multiplied by ℓ_P^2 . 2.

The constant $\Lambda = \lim_M \Lambda_M$, $\Lambda_M(x) = \sum_I e_I(x)^2$ is naively infinite, but can be given a finite value using ζ function regularisation. At finite M, the function Λ_M is in fact a smooth function of rapid decrease for $\sigma = \mathbb{R}^3$ while for $\sigma = T^3$ it is just $\propto (M+1)^3$ without x dependence. It would be interesting to study this in more detail to see whether alternative limiting patterns exist such that Λ comes out finite without additional regularisation. 3.

As expected, the Λ term carries explicit dependence on the chosen background metric $g_{ab} = \delta_{ab}$. It is interesting to see that all the other terms carry no background dependence despite the fact that the chosen Fock representation also depends on that background.

4.

We have performed the calculation in the simplest possible Fock representation corresponding to a white noise covariance in order to minimise the already substantial computational effort. It is conceivable that both proposition and corollary also hold in more general Fock representations. This has been confirmed for the spatial diffeomorphism subalgebra in any dimension and for all tensor types in [31].

5.

It would be easy to extend the analysis by a cosmological constant term without increasing the computational effort too much and using the same tools. This just requires to extend the appendix by five more commutators namely [: F^3 :, : F^3 :] and [: F^3 ; .: X :], X = F, FG, FFG, FFGG. The fact that we compute commutators of linear combinations of quadratic forms using the distributive law allows to reuse the above analysis, one just has to supplement it with the additional commutators and limiting patterns.

6.

The "anomalous" Λ term is in fact again proportional to a normal ordered constraint, so one may call this a "soft anomaly" in the sense that the anomaly just changes the structure functions but not the set of constraints. This is similar to what happens in LQG [16] but while here we find a "quantum correction" to the "classically expected" structure function, in LQG the expected part is missing and one has just a quantum correction. 7.

As already mentioned, while the computation is therefore as non-anomalous as it can possibly be since only normal ordered expressions are well defined quadratic forms on the Fock space, the presence of the structure functions $\omega_a E_k^a E_k^b$ for C_b and $\omega_a E_k^a E_k^b A_b^j$ for G_j implies that $l([C_0(f), C_0(g)] \psi) \neq 0$ even if l solves all three types of constraints. Once again, there is no contradiction because only the object $[C_0(f), C_0(g)]$ is a well defined quadratic form since $C_0(g)\psi, C_0(f)\psi$ respectively is not in the form domain of $C_0(f), C_0(g)$ respectively for any Fock state ψ so that there is no reason to expect that this quantity vanishes. In that sense the commutator calculation merely confirms that one has a quantum represention of the hypersurface deformation algebra and thus has chosen a valid quantisation of the constraints.

As a byproduct we have also confirmed that the constraint algebra of Euclidian $U(1)^3$ gravity (e.g. [34] and references therein) is in this sense properly represented in the same Fock representation because going through the details of the proof one sees that one just has to drop the FG term from the Gauss constraint and the FFGG from the Hamiltonian constraint and thus can drop the commutators corresponding to 1B., 1C., 3B., 4B., 4C., 4D., 5B, 6B., 6C. Both the proposition and the corollary remain literally intact.

Inspection of the individual parts of the computation also reveals that the respective quadratic form monomials have the expected commutators among themselves: For example the FG part of the Gauss constraint has vanishing commutator seperately with the FFGG part of the Hamiltonian constraint which is the quantum statement of the fact that this term of the Gauss constraint generates SU(2) rotations and this part of the Hamiltonian constraint is invariant under those. In general it is reassuring to see that all parts of the Poisson bracket calculation are refelected in the quantum commutator calculation, most importantly that ultra-local terms drop out and that one can use the crucial commutativity of F, G, which in the classical calculation is granted, also in the quantum computation under the normal ordering symbol.

10.

8.

In LQG one avoids the complicated "Lorentzian correction" to the Euclidian Hamiltonian constraint $C_0(f)$, which depends on the Ricci scalar of spin connection Γ of E, by using either a "Wick rotation" [17] with generator $\int d^3x \Gamma_a^j E_i^a$ or by using the commutator equivalent of the Poisson bracket

 $\{\{V, C_0(1)\}, A_a^j\}$ $\{\{V, C_0(1)\}, A_b^k\}\{V, A_c^k\}\epsilon^{abc}\epsilon_{jkl}$ where $V = \int d^3x \sqrt{|\det(E)|}$ is the total volume. Both Γ, V depend non-polynomially on E so that this approach leaves the framework of the present section which relies on polynomial expressions. Thus within the present section one would need to deal with the polynomial version of the constraint mentioned at the beginning of this section.

The task to deal with non-polynomial expressions naturally leads us to the next section.

4 Geometric measures, master constraint and reduced Hamiltonian as quadratic forms on Fock space

As already mentioned, the reduced phase space approach unavoidably leads to rather non-polynomial expressions typically involving integrals over square roots of density weight two scalars which themselves depend on polynomial expressions of all the fields except for the spatial metric q_{ab} which enters also with integer powers of $\sqrt{\det(q)}$. This happens because wehn solving the constraints explicitly one is forced to solve them for canonical momenta because these are the only variables which enter the constraints polynomially and (at least sufficient subset of them) without spatial derivatives so that an explicit, algebraic and local solution is possible. The gauge fixing conditions are then imposed on the conjugate configuration variables. Then the reduced Hamiltonian is a linear combination of those momenta that one solved the constraints for which therefore is a density of weight one. Thus the power of $\sqrt{\det(q)}$ that enters various terms of the reduced Hamiltonian under the square root makes sure that the overall density weight is always two.

It is therefore of considerable interest to explore whether $[\sqrt{\det(q(x))}]^N$ can be defined as a densely defined quadratic form on the Fock space for any interger $N \in \mathbb{Z}$. If it can, then it will be a normal ordered expression of the gravitational annihilation and creation operators. Then more complicated expressions that involve also polynomials of the other fields can also be defined as quadratic forms again by normal ordering. Even more, we are not only interested in the reduced Hamiltonian but also other quadratic forms that involve $\sqrt{\det(q)}$ such as length, area and volume functions in three spatial dimensions and the various forms of the non-polynomial but spatially diffeomorphism invariant master constraint.

In the next subsection we consider these geoemtric measures in the q, p formulation (ADM variables) in any spatial dimension D where we obtain the most systematic results. Then we consider the geometric measures in the e, P formulation also in any D and the A, E formulation in D = 3 respectively which require a case by case analysis. Finally we construct the reduced Hamiltonian as a quadratic form using these tools.

4.1 Metric densities as quadratic forms in (q, p) variables

Let S be a submanifold of σ of dimension $1 \leq d \leq D$ and $Y : U \subset \mathbb{R}^d \to S \subset \sigma; y \mapsto x = Y(y)$ be the corresponding embedding where y, x are coordinates on S, σ respectively. For d = D we can pick Y = id and y = x. The volume of S is then given by

$$\mathsf{Vol}[S] = \int_U d^d y \,\sqrt{\det(([Y^*q](y)))} \tag{4.1}$$

On the other hand for the reduced dynamics or the master constraint we are interested in $[\sqrt{\det(q(x))}]^N$, $N \in \mathbb{Z}$ which formally also concerns the case Y = id. Thus we capture all cases of interest if we consider

$$Q_N^Y(y) := \sqrt{\det([Y^*q](y))}]^N$$
(4.2)

We pick a smooth background metric g on σ with Euclidian signature. We consider a Fock representation using the following (background scalar) annihilation operator

$$A_{jk} := 2^{-1/2} [\kappa \cdot q_{jk} - i \kappa^{-1} \cdot (\omega^{-1} \cdot p_{jk})]$$
(4.3)

where h_a^j is an adapted background D-Bein $g_{ab} = \delta_{jk} h_a^j h_b^k$ with inverse h_j^a and we defined the background scalars $q_{jk} = h_j^a h_k^b q_{ab}$, $p^{jk} = h_a^j h_b^k p^{ab}$ of background density weight zero and one respectively which enjoy canonical brackets

$$\{p^{jk}(x), q_{mn}(y)\} = \delta^j_{(m} \delta^k_{n)} \delta(x, y)$$

$$\tag{4.4}$$

The D-Bein indices j, k, l. are moved with the Kronecker symbol. Here κ is an invertible operator function of the Laplacian $\Delta = g^{ab} \nabla_a^g \nabla_b^g$ mapping scalars to scalars and $\omega = \sqrt{\det(g)}$ is the canonical background scalar density. The one particle Hilbert space is $\mathfrak{h} = L_2(\sigma, \omega \ d^D x)$. The operator $-\Delta$ is positive symmetric on \mathfrak{h} and has self-adjoint extensions with respect to which κ is defined using the spectral theorem. The operator κ has a symmetric bi-scalar integral kernel $\kappa(x, y) = \kappa(y, x)$ which acts on scalars f via

$$[\kappa \cdot f](x) := \int d^D y \,\omega(y) \,\kappa(x,y) \,f(y) \tag{4.5}$$

Denoting the kernel of its inverse by $\kappa^{-1}(x,y)$ defined analogous to (4.5) we find

$$\int d^{D}z \,\omega(z) \,\kappa(x,z) \,\kappa^{-1}(z,y) = \delta(x,y) \,\omega(y)^{-1}, \ \int d^{D}y \,\delta(x,y) \,f(y) = f(x) \tag{4.6}$$

The annihilation operator smeared with scalar background test functions is (summation over repeated indices is understood)

$$\langle f, A \rangle := \langle f^{jk}, A_{jk} \rangle_{\mathfrak{h}} = 2^{-1/2} \left[\langle \kappa \cdot f^{jk}, q_{jk} \rangle_{\mathfrak{h}} - \langle \omega^{-1} \kappa^{-1} \cdot f^{jk}, p_{jk} \rangle_{\mathfrak{h}} \right]$$
 (4.7)

Note the canonical commutation relations

$$[\langle f, A \rangle, \langle f', A \rangle] = 0, \ [\langle f, A \rangle, \ (\langle h, A \rangle)^*] = \langle f, h \rangle \cdot 1, \ \langle f, h \rangle := \int d^D x \ \omega(x) \ \overline{f^{jk}(x)} \ f'_{jk}(x)$$
(4.8)

A coherent state on the corresponding Fock space is defined by D(D+1)/2 complex valued scalar fields $Z_{jk} = Z_{kj}$ such that

$$|Z||^2 := \langle Z, Z \rangle, \langle Z, Z' \rangle := \int d^D x \ \omega \ \overline{Z_{jk}} \ \delta^{jm} \delta^{kn} \ Z'_{kn} < \infty$$

$$(4.9)$$

and

$$\Omega_Z := e^{-\frac{||Z||^2}{2}} e^{[\langle Z, A \rangle]^*} \Omega$$
(4.10)

where Ω is the Fock vacuum defined by (4.3). From Z one reconstructs classical q, p by

$$q_{ab} = 2^{-1/2} h_a^j h_b^k \kappa^{-1} \cdot [Z_{jk} + \overline{Z_{jk}}], \ p^{ab} = 2^{-1/2} i h^{aj} h^{bk} \omega \kappa \cdot [Z_{jk} - \overline{Z_{jk}}],$$
(4.11)

Note that the classical metric q from (4.11) on which the coherent state is concentrated is completely unrelated to the fixed background metric g_{ab} that is used to define the Fock representation.

Given any Ω_Z the span of its excitations

$$[\langle f_1, A \rangle]^* ..[\langle f_N, A \rangle]^* \Omega_Z = e^{\langle Z, A \rangle^*} [\langle f_1, A \rangle]^* ..[\langle f_N, A \rangle]^* \Omega$$
(4.12)

defines a dense subspace of the Fock space as $f_1, ..., f_N$ varies. This follows from the fact that the algebraic Fock state $< \Omega, .\Omega >$ is pure [23], therefore its GNS representation of the Weyl algebra or quivalently the annihilation and creation algebra is irreducible and thus every vector state such as Ω_Z is cyclic.

Proposition 4.1. In any Fock representation, (4.2) can be defined non-perturbatively as a quadratic form on a dense form domain which corresponds to the excitations of any coherent state of that Fock representation which is concentrated on a positive definite spatial metric.

Proof.:

If N is a positive even integer then Q_N^Y is a polynomial in q_{ab} , therefore a well defined quadratic form is obtained by normal ordering it. Thus we can restrict attention to the case that N is a positive odd integer or a negative integer. If N is a positive odd integer we define $Q_N^Y :=: Q_{N+1}^Y Q_{-1}^Y$: so we are left with the case of negative integers. We can define Q_{-N}^Y , N > 0 in various ways, for example by $: [Q_{-1}^Y]^N$: or by independent expressions which we explain below. To define $Q_{-1}^Y(y)$ as a quadratic form we note the identity

$$Q_{-1}^{Y}(y) = \int_{\mathbb{R}^d} \frac{d^d z}{[2\pi]^{d/2}} e^{-\frac{1}{2} \sum_{I,J=1}^d z^I [Y^*q]_{IJ}(y) z^J}$$
(4.13)

which holds in the classical theory for any positive definite metric q_{ab} . We now use (4.11) to express the Gaussian exponent in terms of A_{jk}, A_{jk}^* , specifically

$$\frac{1}{2} z^{T} [Y^{*}q](y) z = \langle f_{Y}(y,z), A \rangle + \langle f_{Y}(y,z), A \rangle^{*}, \quad f_{Y}^{jk}(x;y,z) = \hat{f}_{Y}^{jk}(y,z) \kappa^{-1}(Y(y),x), \\
\hat{f}_{Y}^{jk}(y,z) = 2^{-3/2} z^{I} z^{J} Y_{,I}^{a}(y) Y_{,J}^{b}(y) h_{a}^{j}(Y(y)) h_{b}^{k}(Y(y))$$
(4.14)

and normal order the exponential to arrive at the quadratic form equivalent of (4.13)

$$Q_{-1}^{Y}(y) := \int_{\mathbb{R}^d} \frac{d^d z}{[2\pi]^{d/2}} \left[E_Y(z,y) \right]^* \left[E_Y(z,y) \right], \quad E_Y(y,z) = e^{-\langle f_Y(y,z),A \rangle}$$
(4.15)

One possibility to define $Q_{-N}^{Y}(y)$ is then to take the N-th power of (4.15) and to normal order

$$Q_{-N}^{Y}(y) := \int_{\mathbb{R}^{ND}} \frac{d^{dN}z}{[2\pi]^{dN/2}} \left(\prod_{k=1}^{N} \left[E_{Y}(z_{k}, y) \right]^{*} \right) \left(\prod_{k=1}^{N} \left[E_{Y}(z_{k}, y) \right] \right)$$
(4.16)

Another possibility is to make use of the classical idenity, for $N \ge d$

$$Q_{-N}^{Y}(y) = \frac{1}{c_d} \int_{\mathbb{R}^{d^2}} d^{d^2} z \, |\det(z_1, ..., z_d)|^{N-d} \, e^{-\frac{1}{2} \sum_{k=1}^d \sum_{I,J=1}^d z_k^I \, [Y^*q]_{IJ}(y) \, z_k^J} \tag{4.17}$$

where c_d is the value of the integral in (4.17) obtained for $[Y^*q]_{IJ} = \delta_{IJ}$ and $(z_1, ..., z_d)$ is the d x d matrix whose k - th column is given by the vector z_k . We can now define Q_{-N}^Y by normal ordering the exponentials in (4.17) after decomposing them via (4.14). The advantage of (4.17) over (4.16) is that for large N there are fewer integrals to compute.

Yet another option is to take combinations of both possibilities or even to multiply by $1 = Q_{2M}^Y(y) Q_{-2M}^Y$ until the desired overall negative power of $Q_1^Y(y)$ is reached, followed by normal ordering.

It remains to show that these quadratic forms indeed have (4.12) as dense form domain. For this we just need to use the identity

$$E_{Y}(z,y) < f_{1}, A >^{*} ... < f_{N}, A >^{*} \Omega_{Z}$$

$$= [< f_{1}, A >^{*} + < f_{Y}(z,y), f_{1} >] ..[< f_{N}, A >^{*} + < f_{Y}(z,y), f_{N} >] E_{Y}(z,y) \Omega_{Z}$$

$$< f_{Y}(y,z), f >= \hat{f}_{Y}^{jk}(y,z) [\kappa^{-1} \cdot f_{jk}](Y(y))$$

$$E_{Y}(z,y) \Omega_{Z} = e^{-} \Omega_{Z}$$
(4.18)

where we made use of the fact that coherent states diagonalise the annihilation operator. Since

$$< f_{Y}(y,z), Z > + < f_{Y}(y,z), Z >^{*} = \hat{f}^{jk}(y,z) \ \kappa^{-1} \cdot [Z_{jk} + \overline{Z_{jk}}](Y(y)) = 2^{1/2} \ \hat{f}^{jk}(y,z) \ q_{jk}(Y(y)) = \frac{1}{2} z^{I} z^{J} [Y^{*}q]_{IJ}(y)$$

$$\tag{4.19}$$

it follows that the matrix elements of (4.15) and its descendants are Gaussian integrals in $z^I z^J$ with Gaussian factor $\exp(-z^T [Y^*q](y)z/2)$, q the classical metric determined by Z via (4.11), and coefficients $Y^*[(g \otimes g)[\kappa^{-1}f_k])_{IJ}(y)$, k = 1, ..., N. The result of the Gaussian integral is a polynomial in the inverse m^{IJ} of the pull back metric $m_{IJ} = [Y^*q]_{IJ}$ times $\sqrt{\det(m)}^{-1}$. To see this we introduce the new integration variables $z^A = m_I^A z^I$ where m_I^A is a d-Bein $m_{IJ} = m_I^A h_J^B \delta_{AB}$, express $z^I = h_A^I z^B$ where $h_I^A h_B^I = \delta_B^A$ and finally use that Gaussian integrals of $z^{A_1} \dots z^{A_{2N}}$ are Wick polynomials in the Kronecker symbols $\delta^{A_k A_l}$, $k, l = 1, \dots, 2N$.

It is interesting to note that the dense subspace consisting of the excitations of the Fock vacuum *does not* provide a form domain for (4.2) because the Fock vacuum is a coherent state concentrated on a classical metric of signature (0,..,0). The same is true for any other signature, $s = (\sigma_1,..,\sigma_D)$, $\sigma_k \in \{0,\pm 1\}$, k = 1,..,D different from (1,..,1) because the integral of $e^{-\sigma z^2/2}$ diverges unless $\sigma = 1$. This is the non-degeneracy footprint that the classical theory leaves in the quantum theory.

Corollary 4.1.

The volume functionals (4.1) are quadratic forms with the same dense form domain as in proposition 4.1. In particular, their expectation value with respect to coherent states Ω_Z in the form domain equals the value of the classical functional evaluated on the metric determined by Z.

Proof.:

We define the quadratic form by

$$\begin{aligned} \mathsf{Vol}[S] &:= \int_{U} d^{d}y \; : \; Q_{2}^{Y}(y) \; Q_{-1}^{Y}(y) := \int d^{d}y \; \int \frac{d^{d}z}{[2\pi]^{d/2}} \; E_{Y}(z,y)^{*} : \det([Y^{*}q](y) : \; E_{Y}(y,z) \\ \det([Y^{*}q](y) \; = \; \frac{1}{d!} \epsilon^{I_{1}..I_{d}} \; \epsilon^{J_{1}..J_{d}} [Y^{*}q]_{I_{1}J_{1}} \dots [Y^{*}q]_{I_{d}J_{d}} \\ [Y^{*}q]_{IJ}(y) &= < F_{Y,IJ}(y), A > + < F_{Y,IJ}(y), A >^{*}, \; F_{Y,IJ}^{ij}(x;y) = \hat{F}_{Y,IJ}^{ij}(y) \; \kappa^{-1}(Y(y),x), \\ \hat{F}_{Y,IJ}^{ij}(y) &= h_{a}^{i}(Y(y)) \; h_{b}^{j}(Y(y)) \; Y_{,I}^{a}(y) \; Y_{,J}^{b}(y) \end{aligned}$$
(4.20)

In computing matrix elements of (4.20) with respect to the states (4.12) the prescription of (4.20) is to perform the z integral before the y integral. As all annihilators and creators act on the ket and bra respectively the action of $E_Y(y, z)$ can be taken over from proposition 4.1 while the action of the normal ordered determinant returns a polynomial in the $\langle F_{Y,IJ}(y), f_k \rangle$, $\langle F_{Y,IJ}, Z \rangle$. The z integral can be performed explicitly and returns and overall factor of $Q_{-1}^Y(y)$ for q = q(Z) the classical metric determined by Z via (4.11). For the state Ω_Z itself (no excitation) the expectation value followed by the z integral just returns $Q_1^Y(y)$ for q = q(Z) and the y integral therefore Vol[S] at q = q[Z].

4.2 Metric densities as quadratic forms in other variables

Our results will be confined to defining integer powers of $Q = \sqrt{\det(q)}$ in the (e, p) formulation in any D and the (A, E) formulation in D = 3. The reason for why it is more complicated to define Vol[S] in these variables will become clear shortly.

We have the relations

$$Q^{2} = \det(q) = \det(e)^{2} = |\det(E)|$$
(4.21)

therefore

$$Q^{-1} = c \int_{\mathbb{R}^D} d^D z \ F(e \cdot z) = [\det(E)]^4 \ c_3 \int_{\mathbb{R}^9} d^9 z \ |\det((z_1, z_2, z_3))|^{1/2} \ F_3(E \cdot z_1, E \cdot 2, E \cdot z_3), \tag{4.22}$$

where $[e \cdot z]^j = e_a^j z^a$, $[E \cdot z_I]^a = E_j^a z_I^j$, F, F_3 are functions on $\mathbb{R}^D, \mathbb{R}^9$ respectively such that the integrals evaluated at $e_a^j = \delta_a^j$ and $E_j^a = \delta_a^j$ equal the constants c^{-1}, c_3^{-1} respectively. We now write F, F_3 in terms of their Fourier transforms

$$F(e \cdot z) = \int \frac{d^D k}{[2\pi]^D} \hat{F}(k) \ e^{i \ k_j e_a^j z_a}, \ F_3(\{E \cdot z\}) = \int \frac{d^9 k}{[2\pi]^9} \ \hat{F}_3(k) e^{i \ k_a^I E_j^a z_I^j}, \tag{4.23}$$

Then we pick, similar as for (q, P) variables a Fock quantisation of (e, p) and (A, E) respectively using the background metric g based on the annihilators

$$A_{jk} = 2^{-1/2} [\kappa \cdot h_j^a e_{ak} - i\kappa^{-1} \cdot \omega^{-1} h_{aj} p_k^a]$$

$$A_{jk} = 2^{-1/2} [\kappa \cdot h_j^a A_{ak} - i\kappa^{-1} \cdot \omega^{-1} h_{aj} E_k^a]$$
(4.24)

(for $j \le a$ in upper triangular gauge when the gravitational Gauss constraint is gauge fixed) which enjoy canonical commutation relations. Finally one normal orders the exponentials (Weyl elements) in (4.23).

To see why it is more dificult to define Vo[S] in these variables unless S is D dimensional and 3-dimensional respectively (in which case we just use $Q = Q^2Q^{-1} = \det(e)^2Q^{-1}$ for (e, p) and $Q = Q^4Q^{-3} = \det(E)^2Q^{-3}$ for (A, E), use (4.22) and normal order) we note that for other dimensions e.g.

$$[Y^*q]_{IJ} = \delta_{ij} [Y^*e]^i_I [Y^*e]^j_J \tag{4.25}$$

cannot be written as the determinant of a submatrix of e_a^j and thus it cannot be the result of a z-integral which returns such a determinant as a result of the Jacobean when changing integration variables. Thus we must use different techniques. For instance in D = 3 we can use for d = 1 and d = 2 respectively

$$\sqrt{\delta_{ij}[Y^*e]^i[Y^*e]^j}^{-1} = \int_{\mathbb{R}^3} \frac{d^3z}{2\pi^2 ||z||^2} e^{i[Y^*e]^j z_j}$$
$$\sqrt{\delta^{ij}[Y^*E]_i[Y^*E]_j}^{-1} = \int_{\mathbb{R}^3} \frac{d^3z}{2\pi^2 ||z||^2} e^{i[Y^*E]_j z^j}$$
(4.26)

with $[Y^*E]_j = \frac{1}{2} \epsilon^{IJ} \epsilon_{abc} Y^a_{,J} Y^b_{,J} E^c_j$ which is the integral that expresses the Green function of the Laplace operator as the Fourier of $||z||^{-2}$ up to a factor. Then we can use the relation $E^a_j = Qe^a_j$, $e^a_j e^k_a = \delta^k_j$ to relate E, eand then use (4.22), (4.26). Then the length of a curve or the area of a surface can be obtained by multiplying (4.26) with the polynomials $||Y^*e||^2$, $||Y^*E||^2$ respectively, normal ordering and integrating over the domain of the embedding Y. Other cases can be treated similarly.

4.3 Master constraint as a quadratic form on Fock space

The polynomial version of the master constraint is simply obtained by normal ordering of its integrand. The integrand of the non-polynomial version can be written as a polynomial times Q^{-N} for sufficiently large N. We apply proposition 4.1 and normal order the resulting expression.

4.4 Reduced Hamiltonian as a quadratic form on Fock space

We now apply this theory to the reduced Hamiltonian which as we have seen in section 2 typically takes the form

$$H = -\int d^D x \sqrt{-2 Q \tilde{C}}$$
(4.27)

where $C = \frac{\pi_0^2}{2Q} + \tilde{C}$, $Q = \sqrt{\det(q)}$ is the full Hamiltonian constraint and \tilde{C} is the contribution to C from geometry and matter fields other than ϕ^0, π_0 with gauge fixing $\phi^0 = t$. We pick a dense form domain based on a coherent state Ω_Z as above and define the quantity

$$\Lambda_0 := - \langle \Omega_Z, \tilde{c} | \Omega_Z \rangle, \ \tilde{c} := \frac{\tilde{C}}{Q}$$

$$(4.28)$$

where the spatial scalar $\frac{\tilde{C}}{Q}$ is quantised as a quadratic form in Fock representations for all geometry and matter fields by the above methods using the fact that $\tilde{C}/Q = [\tilde{C} \ Q^N]/Q^{N+1}$ and that $\tilde{C}Q^N$ is a polynomial for sufficiently large N (pick the minimal such N) and using normal ordering. Assuming that $\Lambda_0 > 0$ is a positive function on σ and that $\Delta = 1 + \frac{\tilde{c}}{\Lambda_0}$ is a small quantity we use the approximant derived in section (2) depending on the interpolation parameter k or the truncation parameter N and normal order, e.g.

$$H_{k,Z} := \int d^{D}x \sqrt{2\Lambda_{0}} : Q[-1 + \Delta + k \Delta^{2}]:$$
(4.29)

with $0 \le k \le 1$ where : $Q \Delta^n$:, n = 1, 2 is defined by decomposing into the terms : $\tilde{C}^l Q^{1-l}$:, l = 0, 1, 2 which are treated by the above methods. The terms independent of k are

$$H_{1,Z} = \int d^D x \, \frac{1}{\sqrt{2\Lambda_0}} \left[: \tilde{C} : -\Lambda_0 : Q :\right]$$
(4.30)

We have made the dependence of the approximant on Z via $\Lambda_0 = \Lambda_0(Z)$ explicit. Similar remarks concern the treatment of the higher order powers Δ^n in the N-th order truncation of the Hamiltonian.

Note that $\langle \Omega_Z, \Delta \Omega_Z \rangle = 0$ by construction and thus one expects the fluctuation term $\langle \Omega_Z, [H_{k,Z} - H_{1,Z}]\Omega_Z \rangle$ to be subleading. If $\tilde{C} = \Lambda Q + \hat{C}$ where $\Lambda > 0$ is a "bare" cosmological constant term and \hat{C} the remainder then

$$: \tilde{C} : -\Lambda_0 : Q :=: \hat{C} : + [\Lambda - \Lambda_0] : Q :$$
(4.31)

where $\epsilon := \Lambda - \Lambda_0$ is the dressed "cosmological function" which is a constant on sufficiciently large scales for suitable Ω_Z . It can be positive since we have $\langle : \tilde{C} : \rangle = \langle : \hat{C} : \rangle + \Lambda \langle : Q : \rangle = -\Lambda_0(1 - \delta) \langle : Q : \rangle$ with $\delta = \langle : \Delta Q : \rangle / \langle : Q : \rangle$, i.e. for e.g. $\epsilon = \Lambda_0 \delta$ we have $\langle : \hat{C} : \rangle = -[(2 - \delta)\Lambda_0 + \epsilon] \langle : Q : \rangle = -2\Lambda_0 \langle : Q : \rangle$ is negative due to the negative gravitational contribution.

This sketches how the above methods can be used in order to write k approximation of the exact square root Hamitonian as a quadratic form in Fock representations for both geoemtry and matter based on the background metric g and using normal ordering and the expression for Q^N provided in proposition 4.1 for Y = id. One expects that the matrix elements are expecially simple if q(Z) = g i.e. when the coherent state metric equals the background metric.

4.5 Towards operators

As quadratic forms cannot be multiplied, it is not possible to evaluate e.g. $Vol[S]^2$ on the form domain. However introducing modes e_I as in section 3 we can define actual operators such as $Vol_M[S]$ which are obtained by decomposing the part of Vol[S] which is polynomial in annihilation and creation operators, i.e. the piece : Q_2^Y : in (4.20), into modes I and to restrict the sum by I < M. Note that this is different from $P_{M,Z}$ Vol[S] $P_{M,Z}$ where $P_{M,Z}$ projects on the (closure of the) subspace $\mathcal{D}_{M,Z}$ given by the span of excitations of Ω_Z by polynomials in the $< f, A >^*$ where $< e_I, f^{ij} >= 0, I > M$, because Ω_Z is not annihilated by < f, A > (rather it returns < f, Z >). Then $Vol_M[S]$ in fact preserves $\mathcal{D}_{M,Z}$ because the resticted polynomial dependence is now a polynomial in annihilation and creation operators $A_I^*, A_I, I < M$ and no longer an infinite series. This however does not allow to remove the cutoff in contrast to the commutator algebra where there are cancellations.

To actually remove the regulator requires either the counterterm method familiar from the perturbative construction of the S matrix in usual QFT or methods from constructive QFT [36], see also [31]. The latter idea can be sketched as follows (we drop the label k, Z for simplicity): Consider the truncation H_M as defined above and try to find an invertible "dressing operator" $T_M : \mathcal{D}_0 \to \mathcal{D}_0$ where \mathcal{D}_0 is the span (4.12) such that $H_M T_M = T_M h_M$ where h_M has a strong limit h in the topology of \mathcal{H}_0 , the Fock space completion of \mathcal{D}_0 . One defines new scalar products $\langle T_M \psi, T_M \psi' \rangle_M := \langle T_M \psi, T_M \psi' \rangle / \langle T_M \Omega_Z, T_M \Omega_Z \rangle$ and $\hat{\mathcal{D}}_0$ as the subspace of \mathcal{D}_0 such that these inner producs converge as $M \to \infty$ thereby defining a new maps $T : \hat{\mathcal{D}}_0 \to \mathcal{D}$ and a new Hilbert space \mathcal{H} as the completion of \mathcal{D} . Then HT = Th if h has range in $\hat{\mathcal{D}}_0$ which defines H densely on \mathcal{D} with range in \mathcal{H} . Another possibility is a variant of the Friedrichs extension technique [24]:

Suppose that one could show that the quadratic form defining the physical Hamiltonian H, which is automatically symmetric on the physical Hilbert space \mathcal{H} , is bounded from below on a dense subspace $V \subset \mathcal{H}$ of the physical Hilbert space and such that it is closable. That is, there exists $k \geq 0$ such that $||\psi||_k^2 := \langle \psi, H\psi \rangle_{\mathcal{H}} + k||\psi||_{\mathcal{H}}^2 \geq 0$ for all $\psi \in V$ and such that V is in fact complete in the norm $||.||_{k+1}$, that is, $(V, < ... >_{k+1})$ is a Hilbert space. Then we can define an operator H' on \mathcal{H} with dense domain $\mathcal{D}' = R_k(\mathcal{H})$ where $R_k(\psi) \in V$, $\psi \in \mathcal{H}$ is the Riesz representative of the $||.||_{k+1}$ continuous linear form $f \in V \mapsto \langle \psi, f \rangle_{\mathcal{H}}$. Then for $\hat{\psi}'_1, \hat{\psi}'_2 \in \mathcal{D}'$ the formula $\langle \psi'_1, H'\psi'_2 \rangle_{\mathcal{H}} := \langle \psi'_1, H\psi'_2 \rangle_{\mathcal{H}}$ defines the matrix elements of a self-adjoint operator bounded from below by the same bound [24] and we can use the spectral theorem to construct the scattering matrix corresponding to H'.

5 Conclusion

The main result of the present manuscript is to demonstrate that, although counter intuitive, background dependent Fock representations can be used in a non perturbative fashion in order to construct a non-perturbative theory of quantum gravity. In non-perturbative quantum gravity there is no natural split of the Hamiltonian (constraint) into a "free" and an "interacting" part. Even in the polynomial version of the Hamiltonian constraint there is no term which is of quadratic order in the fields and the reduced Hamiltonian depends highly non-polynomially on the fields anyway.

In perturbative quantum gravity one artificially decomposes the full metric q_{ab} into a classical background g_{ab} and a "graviton" fluctuation $h_{ab} = q_{ab} - h_{ab}$ and one expands e.g. the reduced Hamiltonian in powers of h_{ab} which is an infinite series. Then one resorts to the formalism of perturbative QFT of h_{ab} on the background defined by g_{ab} by taking the free part (quadratic part of the expansion) of the Hamiltonian as an input for the choice of the Fock representation of h_{ab} . This is quite different from what we have done here because we never perform the split $q_{ab} = g_{ab} + h_{ab}$, we quantise the full metric q_{ab} and just use the background g_{ab} to define a Fock representation of the full q_{ab} (e.g. in order to define a background Laplacian etc.). In our formalism the fields g_{ab}, q_{ab} are totally independent. While we also have to use an expansion of the reduced Hamiltonian in order to deal with the square root, that expansion is of a different nature than the one used in perturbative quantum gravity because it does not use the auxiliary, classical background field g_{ab} as zeroth order but rather the expectation value of the argument of the quare root with respect to a coherent state Ω_Z which is concentrated on a classical Euclidian signature metric q_{ab}^0 constructed from Z. This metric is again is completely unrelated to g_{ab} . The excitations of Ω_Z by smeared polynomials of the quantum metric q_{ab} define a dense form domain of the resulting Hamiltonian quadratic form. The expansion is in powers of the fluctutations of the full argument of the square root which itself is not a polynomial but that non-polynomial dependence on the quantum metric can be dealt with using suitable Fock space techniques developed in section 4 of the present paper.

The resulting reduced Hamiltonian is then defined as a densely defined quadratic form but is not an operator. To define an actual operator one must use non-pertuerbative constructive QFT methods. These use the notion of mode cut-offs and were successfully employed to construct interacting scalar fields in 3d Minkowski space [36]. We also used elements of this in order to construct the polynomial quantum constraints of a version of Euclidian quantum gravity as quadratic forms and even their commutator algebra which is possible because commutators contain differences of products of quadratic forms and while products of quadratic forms are ill-defined such differences of products, carefully defined using mode cut-offs, can define well defined new quadratic forms.

The results of this paper have many applications of which we mention a few.

Α.

In quantum cosmology and black hole perturbation theory one performs a split $q_{ab} = q_{ab}^0 + q_{ab}^1$ where q_{ab}^0 is the "symmetric" part of the metric with respect to the given symmetry (spatial homogeneity, spherical symmetry, axi-symmetry...) and q_{ab}^1 is the "non-symmetric" remainder. However, instead of treating q^0, q^1 as g, h of the perturbative approach to quantum gravity one can quantise the background q^0 as well in order to study cumulative backreaction effects. Thus one wants to keep track of how the symmetric part interacts with the non-symmetric part. This viewpoint is in between the totally perturbative approach in which $g = q^0$ is fixed and just $h = q^1$ is quantised and the totally non-perturbative approach in which the full $q = q^0 + q^1$ is quantised without making

any difference between q^0, q^1 . For this reason it is sometimes called the "hybrid" approach [37, 38]. The point is now that the perturbative expansion of reduced Hamiltonian is by construction a polynomial in q^1 but retains a non-polynomial dependence on q^0 . We can therefore apply the non-perturbative Fock quantisation developed in this paper to the q^0 sector while we apply the usual perturbative Fock quantisation to the q^1 sector. B.

A natural regularisation of all the quadratic forms constructed in this paper such as Vol[S], $H_{k,Z}$ is obtained by considering $H_{k,Z,M}$ which is obtained from $H_{k,Z}$ by decomposing the polynomial dependence of $H_{k,Z}$ in terms of annihilation and creation operators and modes I and to restrict the sum over modes by I < M with mode cut-off M on the ONB e_I of $L_2(\sqrt{\det(g)}d^Dx, \sigma)$. Then $H_{k,Z,M}$ preserves the span $\mathcal{D}_{M,Z}$ of excitations of a coherent state Ω_Z where the test functions f of excitations $< f, A >^*$ have the property $< f, e_I >= 0, I > M$. Here Z is a point in the classical phase space subject to the condition that the metric q[Z] that is encoded by it has Euclidian signature. This is a real space regularisation independent of Fourier transform techniques which in principle works on any σ . It provides a UV cut-off as $M \to \infty$ corresponds to infinite spatial resolution. Thus at finite M this provides

1. a well defined theory of quantum gravity in interaction with matter

2. using ordinary Fock space techniques combined with new non-perturbative techniques, in particular all Hilbert spaces are separable

3. all gauge invariance has been removed.

Thus the powerful machinery of Fock spaces, Feynman diagrammes etc. can be applied e.g. in constructing a scattering matrix. We expect the theory to be especially simple when the coherent state label q[Z] coincides with the background metric g used to construct the Fock representation.

To take the $M \to \infty$ limit and define actual operators one needs to invoke renormalisation similar to [36] or using Hamiltonian/Wilsonian renormalisation (see [21] and references therein) based on the family of theories $(\mathcal{H}_{M,Z}, H_{k,Z,M})$.

С.

If one takes the quadratic form $H_{k,Z}$ as it is (no cut-off) one can construct the scattering matrix from it by the usual methods of QFT (Gell-Mann Low formula) if one can usefully decompose $H_{k,Z}$ into free and interacting terms. However the free term cannot just be a quadratic term because it does not exist in this non perturbative approach. One expects however that for matrix elements between matter excitations of the vector Ω_Z the Hamiltonian is effectively a QFT on CST Hamiltonian for matter fields for which such a decomposition is possible. There will be corrections as compared to QFT in CST due to fluctuation effects of Ω_Z and due to its gravitational excitations.

D.

The commutator calculation has been performed only for Euclidian vacuum gravity (however with cut-off removed!) in four dimensions in the (A, E) polarisation. By the methods introduced in the present work, it is possible to repeat the calculation for both signatures and any matter content in all three polarisations discussed and even in any dimension if one works in the (q, p) or (e, P) polarisation. However the computational effort is significantly higher.

A Nomal ordered commutators of normal ordered monomials

In this section we display the results of the computation of commutators of normal ordered monomials of $F_{\alpha} = A_{\alpha} - A_{\alpha}^*$, $G_{\alpha} = A_{\alpha} + A_{\alpha}^*$ as a linear combination of such normal ordered monomials. That linear combination is unique given the annihilation and creation algebra A_{α}, A_{α}^* , $[A_{\alpha}, A_{\beta}^*] = \delta_{\alpha\beta} E$ where E is the unit algebra element and $\alpha, \beta, ...$ can be from any index set. Since normal ordered monomials of F, G are separately totally symmetric in the labels of F, G factors respectively, that symmetry must be displayed also y the r.h.s. of the calculation which is a good consistency check. A normal ordered monomial is defined by its order m and the number p = 0, ..., m of factors F. If one computes the commutator of monomials with data (m, p), (n, q) respectively, the r.h.s. is a linear combination of normal ordered monomials of order m + n - 2k, k = 1, ...[(m + n)/2] where [.] is the Gauss bracket unless m = n, p = q in which case $1 \le k \le m - 1$. Each such monomial comes with k factors of Kronecker δ . The various powers of 2 can be attributed to first the commutator $[F_{\alpha}, G_{\beta}] = 2\delta_{\alpha\beta} E$ and the symmetrisation operation $K_{\alpha}L_{\beta} + K_{\alpha}L_{\beta} =: 2 K_{(\alpha}L_{\beta)}$ for any objects K, L, i.e. there is one factor of 2 for each

Kronecker δ and for each symmetrisation in 2 indices. Note that a factor of 2 Kronecker δ symmetrised in one pair of indices is automatically symmetrised in the other pair which is why some of the terms apparently lack an expected symmetry. Clearly there are no Kronecker δ for pairs of indices carried by the same monomial on the left hand side.

Thus it appears that one can almost uniquely guess the result of the right hand side by these simple rules if one simply writes down all possible normal ordered terms $F^{p-r}G^{m-p-s}F^{q-s}G^{n-q-r}$ following from [: F^pG^{m-p} :, : F^qG^{n-q} :] for the terms involving k = r + s Kronecker factors and then symmetrises and normal orders, the intuition being that one needs to take commutators [F, G] with F, G respectively taken from : F^pG^{m-p} :, : F^qG^{n-q} : respectively (plus sign) or vice versa (minus sign). However, this is not the case for the terms involving more than one Kronecker factor: For instance the double Kronecker terms $+: G^2:$, $+: F^2:$ in the computation of [: FG:,: $F^2G^2:$] are unexpected, one would rather have expected a term : FG: by this intuition which however is absent.

When doing the computation by hand, at least for low order monomials, the computational effort can be reduced by not using lemma 3.1 but rather successively using identities such as

$$: F_{\alpha} G_{\beta} := A_{\beta}^{*} F_{\alpha} + F_{\alpha} A_{\beta} = \frac{1}{2} [(G_{\beta} - F_{\beta}) F_{\alpha} + F_{\alpha} (G_{\beta} + F_{\beta})] = \frac{1}{2} [G_{\beta} F_{\alpha} + F_{\alpha} G_{\beta}]$$
$$: F_{\alpha} G_{\beta} G_{\gamma} := A_{\gamma}^{*} : F_{\alpha} G_{\beta} : + :F_{\alpha} G_{\beta} : A_{\gamma}$$

etc. which has the advantage to express the normal ordered objects in terms of symmetrised objects just containing F, G so that [F, F] = [G, G] = [F, G] - 2E = 0 can be exploited in the commutator calculation. When the comutator has been calculated, which at this point contains only one Kronecker factor, one needs to rearrange the result into the terms in the above list that one has produced which requires further commutations producing the additional powers of Kronecker symbols.

Although straightforward, this becomes quickly algebraically very involved, the hand written computation involves an order of 20 pages. The end result still fits on a single page. For better readability we have written E^k instead of E for the term that involves k Kronecker symbols.

1. [F, F]

$$[:F_{\alpha}:,:F_{\mu}:] = E \cdot 0 \tag{A.1}$$

2.
$$[F, FG]$$

$$[:F_{\alpha}:,:F_{\mu} G_{\rho}:] = 2 E \cdot \delta_{\alpha \rho} :F_{\mu}:$$
(A.2)

3. [F, FFG]

$$[:F_{\alpha}:,:F_{\mu} F_{\nu} G_{\rho}:] = 2 E \cdot \delta_{\alpha \rho} :F_{\mu} F_{\nu}:$$
(A.3)

4. [*F*, *FFGG*]

$$[:F_{\alpha}:,:F_{\mu} F_{\nu} G_{\rho} G_{\sigma}:] = 4 E \cdot \delta_{\alpha(\rho}:G_{\sigma}) F_{\mu} F_{\nu}:$$
(A.4)

5. [FG, FG]

$$[:F_{\alpha} G_{\gamma}:,:F_{\mu} G_{\rho}:] = 2 E \cdot (\delta_{\alpha\rho} :F_{\mu}G_{\gamma}:-\delta_{\mu\beta} :F_{\alpha}G_{\rho}:)$$
(A.5)

 $\mathbf{6.} \ [FG, FFG]$

$$[:F_{\alpha} G_{\gamma}:,:F_{\mu} F_{\nu} G_{\rho}:]$$

$$= 2 \{ E \cdot (\delta_{\alpha\rho} : F_{\mu} F_{\nu} G_{\gamma}: -2 \delta_{\gamma(\mu} : F_{\nu)} F_{\alpha} G_{\rho}:) + E^{2} \cdot (2 \delta_{\alpha(\mu} \delta_{\nu)\gamma} : G_{\rho}:) \}$$
(A.6)

7. [FG, FFGG]

$$[:F_{\alpha} G_{\gamma}:,:F_{\mu} F_{\nu} G_{\rho} G_{\sigma}:]$$

$$= 4 \{E \cdot (\delta_{\alpha(\rho}:G_{\sigma}) F_{\mu} F_{\nu} G_{\gamma}: -\delta_{\gamma(\mu}:F_{\nu)} F_{\alpha} G_{\rho} G_{\sigma}:)$$

$$+E^{2} \cdot (\delta_{\alpha(\mu} \delta_{\nu)\gamma}:G_{\rho} G_{\sigma}: +\delta_{\alpha(\rho} \delta_{\sigma)\gamma}:F_{\mu} F_{\nu}:)\}$$
(A.7)

8. [FFG, FFG]

$$[:F_{\alpha} F_{\beta} G_{\gamma}:,:F_{\mu} F_{\nu} G_{\rho}:]$$

$$= 4 \{E \cdot (\delta_{\rho(\alpha} : F_{\beta}) F_{\mu} F_{\nu} G_{\gamma}: -\delta_{\gamma(\mu} : F_{\nu)} F_{\alpha} F_{\beta} G_{\rho}:)$$

$$+E^{2} \cdot (\delta_{\gamma(\mu} \delta_{\nu)(\alpha} : F_{\beta}) G_{\rho}: -2 \delta_{\rho(\alpha} \delta_{\beta)(\mu} : F_{\nu)} G_{\gamma}:)\}$$
(A.8)

9. [*FFG*, *FFGG*]

$$[:F_{\alpha} F_{\beta} G_{\gamma}:,:F_{\mu} F_{\nu} G_{\rho} G_{\sigma}:]$$

$$= 4 \{E \cdot (-\delta_{\gamma(\mu} :F_{\nu)} F_{\alpha} F_{\beta} G_{\rho} G_{\sigma}:+2:F_{(\alpha} \delta_{\beta)(\rho} G_{\sigma}) F_{\mu} F_{\nu} G_{\gamma}:)$$

$$+E^{2} \cdot (2:F_{\mu} F_{\nu} F_{(\alpha}: \delta_{\beta)(\rho} \delta_{\sigma)\gamma}+2:G_{\rho} G_{\sigma} F_{(\alpha}: \delta_{\beta)(\mu} \delta_{\nu)\gamma}-4:G_{\gamma} G_{(\rho} \delta_{\sigma)(\alpha} \delta_{\beta)(\mu} F_{\nu)}:)$$

$$+E^{3} \cdot (-4:F_{(\mu}: \delta_{\nu)(\alpha} \delta_{\beta)(\rho} \delta_{\sigma)\gamma}-2:F_{(\mu}: \delta_{\nu)\gamma} \delta_{\alpha(\rho} \delta_{\sigma)\beta})\}$$
(A.9)

10. [*FFGG*, *FFGG*]

$$[:F_{\alpha} F_{\beta} G_{\gamma} G_{\delta}:, :F_{\mu} F_{\nu} G_{\rho} G_{\sigma}:]$$

$$= 8 \{E \cdot (-:F_{\alpha} F_{\beta} G_{\rho} G_{\sigma} G_{(\gamma} \delta_{\delta)(\mu} F_{\nu)}: + :F_{\mu} F_{\nu} G_{\gamma} G_{\delta} G_{(\rho} \delta_{\sigma)(\alpha} F_{\beta)}:)$$

$$+ E^{2} \cdot (-2:F_{\alpha} F_{\beta} F_{(\mu} \delta_{\nu)(\gamma} \delta_{\delta)(\rho} G_{\sigma)}: -2:G_{\gamma} G_{\delta} F_{(\mu} \delta_{\nu)(\alpha} \delta_{\beta)(\rho} G_{\sigma)}:$$

$$+ 2:F_{\mu} F_{\nu} G_{(\gamma} \delta_{\delta)(\rho} \delta_{\sigma)(\alpha} F_{\beta)}: + 2:G_{\rho} G_{\sigma} G_{(\gamma} \delta_{\delta)(\mu} \delta_{\nu)(\alpha} F_{\beta)}:)$$

$$+ E^{3} \cdot (4:F_{(\alpha} \delta_{\beta)(\mu} \delta_{\nu)(\gamma} \delta_{\delta)(\rho} G_{\sigma)}: + 2:F_{(\alpha} \delta_{\beta)(\rho} G_{\sigma)}: \delta_{\mu(\gamma} \delta_{\delta)\nu}$$

$$- 4:F_{(\mu} \delta_{\nu)(\alpha} \delta_{\beta)(\rho} \delta_{\sigma)(\gamma} G_{\delta)}: - 2:F_{(\mu} \delta_{\nu)(\gamma} G_{\delta)}: \delta_{\alpha(\rho} \delta_{\sigma)\beta}) \}$$

$$(A.10)$$

References

- [1] R. F. Streater, A. S. Wightman. PCT, spin and statistics and all that. Benjamin Inc., New York, 1964.
- [2] R. Haag, "Local Quantum Physics", Springer Verlag, Berlin, 1984
- [3] S. Fulling. Aspects of Quantum Field Theory in Curved Spacetime. London Math. Society Student Texts, vol. 17, 1989.
- [4] M.H. Goroff, A. Sagnotti, Phys. Lett. B160 (1985) 81; Nucl. Phys. B266 (1986) 709
- [5] C. P. Burgess. Introduction to Effective Field Theory. Ann. Rev. Nucl. Part. Sci. 57 (2007) 329-362. arXiv:hep-th/0701053
- [6] R. Percacci. An introduction to covariant quantum gravity and asymptotic safety. World Scientific, Singapore, 2017.

M. Reuter, F. Saueressig. Quantum gravity and the functional renormalization group. Cambridge monographs on mathematical physics, Cambridge, 2019.

[7] R. Loll. Quantum gravity from causal dynamical triangulations: a review. Class. Quantum Grav. 37 013002.
 e-Print: 1905.08669 [hep-th]

J. Ambjorn, A. Goerlich, J. Jurkiewicz, R. Loll. Causal dynamical triangulations and the search for a theory of quantum gravity. In: Proceedings of MG13. R. T. Jantzen, K. Rosquist, R. Ruffini. 120-137. World scientific, Singapore, 2013. e-Print: 1305.6680 [gr-qc]

[8] S. Surya. The causal set approach to quantum gravity. Living reviews in relativity 22 (2019) 1. arXiv:1903.11544 [gr-qc]

F. Dowker. Introduction to causal sets and their phenomenology. Gen. Rel. Grav. 45 (2013) 9, 1651-1667

[9] C. Rovelli, "Quantum Gravity", Cambridge University Press, Cambridge, 2004. T. Thiemann, "Modern Canonical Quantum General Relativity", Cambridge University, Press, P

T. Thiemann, "Modern Canonical Quantum General Relativity", Cambridge University Press, Cambridge, 2007

J. Pullin, R. Gambini, "A first course in Loop Quantum Gravity", Oxford University Press, New York, 2011 C. Rovelli, F. Vidotto, "Covariant Loop Quantum Gravity", Cambridge University Press, Cambridge, 2015

- [10] H. Nstase. introduction to the AdS/CFT correspondence. Cambridge university Press, Cambridge, 2015.
- [11] K. Giesel, T. Thiemann. Hamiltonian theory: dynamics. In: Handbook of Quantum Gravity. C. Bambi, L. Modesto, I. Shapiro (eds.), Springer Verlag, Berlin, 2023.
- [12] T-P. Cheng, L.F. li. Gauge theory of elementary particles. Oxford University Press, Oxford, 1984.
- [13] A. Ashtekar, C.J. Isham, "Representations of the Holonomy Algebras of Gravity and Non-Abelean Gauge Theories", Class. Quantum Grav. 9 (1992) 1433, [hep-th/9202053]
 A. Ashtekar, J. Lewandowski, "Representation theory of analytic Holonomy C* algebras", in "Knots and Quantum Gravity", J. Baez (ed.), Oxford University Press, Oxford 1994
 A. Ashtekar, J. Lewandowski, "Projective Techniques and Functional Integration for Gauge Theories", J. Math. Phys. 36, 2170 (1995), [gr-qc/9411046]
- [14] H. Narnhofer, W.E. Thirring. Covariant QED without indefinite metric. Rev. Math. Phys. 4 (1992) spec01, 197-211
- [15] C. Fleischhack, "Representations of the Weyl algebra in quantum geometry", Commun. Math. Phys. 285 (2009) 67-140, [math-ph/0407006]

J. Lewandowski, A. Okolow, H. Sahlmann, T. Thiemann, "Uniqueness of diffeomorphism invariant states on holonomy-flux algebras" Commun. Math. Phys. **267** (2006) 703-733, [gr-qc/0504147]

- [16] T. Thiemann, "Anomaly-free Formulation of non-perturbative, four-dimensional Lorentzian Quantum Gravity", Physics Letters B380 (1996) 257-264, [gr-qc/9606088]
 T. Thiemann, "Quantum Spin Dynamics (QSD)", Class. Quantum Grav. 15 (1998) 839-73, [gr-qc/9606089]; "Quantum Spin Dynamics (QSD) : II. The Kernel of the Wheeler-DeWitt Constraint Operator", Class. Quantum Grav. 15 (1998) 875-905, [gr-qc/9606090]; "Quantum Spin Dynamics (QSD) : IV. 2+1 Euclidean Quantum Gravity as a model to test 3+1 Lorentzian Quantum Gravity", Class. Quantum Grav. 15 (1998) 1249-1280, [gr-qc/9705018]; "Quantum Spin Dynamics (QSD) : V. Quantum Gravity as the Natural Regulator of the Hamiltonian Constraint of Matter Quantum Field Theories", Class. Quantum Grav. 15 (1998) 1281-1314, [gr-qc/9705019]
- [17] T. Thiemann. Exact quantisation of $U(1)^3$ quantum gravity via exponentiation of the hypersurface deformation algebroid. Class. Quant. Grav. 40 (2023) 24, 245003. e-Print: 2207.08302 [gr-qc]

- [18] T. Thiemann, "Quantum spin dynamics. VIII. The Master constraint", Class. Quant. Grav. 23 (2006) 2249-2266, [gr-qc/0510011]
- [19] K. Giesel, T. Thiemann, "Scalar Material Reference Systems and Loop Quantum Gravity", Class. Quant. Grav. 32 (2015) 135015, [arXiv:1206.3807]
- [20] R. Gambini, J. Lewandowski, D. Marolf, J. Pullin On the consistency of the constraint algebra in spin network quantum gravity. Int. J. Mod. Phys. D 7 (1998) 97-109; e-Print: gr-qc/9710018 [gr-qc]
 M. Varadarajan. Euclidean LQG Dynamics: An Electric Shift in Perspective. Class. Quant. Grav. 38 (2021) 13, 135020. e-Print: 2101.03115 [gr-qc]
 M. Varadarajan. Anomaly free quantum dynamics for Euclidean LQG. e-Print: 2205.10779 [gr-qc]
- [21] T. Thiemann. Renormalization, wavelets, and the Dirichlet-Shannon kernels. Phys. Rev. D 108 (2023) 12, 125008. e-Print: 2207.08294 [hep-th]
- [22] R. Wald. Quantum field theory in curved spacetime and black hole thermodynamics. University of chicago Press, chicago, 1994.
- [23] O. Bratteli, D. W. Robinson, "Operator Algebras and Quantum Statistical Mechanics", vol. 1,2, Springer Verlag, Berlin, 1997
- [24] M. Reed, B. Simon, "Methods of modern mathematical physics", vol. I-IV, Academic Press, 1980
- [25] A. Ashtekar, "New Variables for Classical and Quantum Gravity" Phys. Rev. Lett. 57 (1986) 2244-2247
 J. F. G. Barbero, "A real polynomial formulation of general relativity in terms of connections", Phys. Rev. D49 (1994) 6935-6938
- [26] N. Bodendorfer, A. Thurn, T. Thiemann. Towards Loop Quantum Supergravity (LQSG). Phys. Lett. B 711 (2012) 205-211. e-Print: 1106.1103 [gr-qc]
- [27] T. Thiemann. Quantum gravity in the triangular gauge. e-Print: 2305.06724 [gr-qc]
 T. Lang, S. Schander. Quantum Geometrodynamics Revived I. Classical Constraint Algebra. e-Print: 2305.10097 [gr-qc]
 - II. Hilbert Space of Positive Definite Metrics. 2305.09650 [gr-qc]
- [28] D. Marolf, D. Giulini. On the generality of refined algebraic quantization. Class. Quant. Grav. 16 (1999) 2479-2488; e-Print: gr-qc/9812024 [gr-qc]
- [29] P. Hajicek, K. Kuchar. Constraint quantization of parametrized relativistic gauge systems in curved spacetimes. Phys. Rev. D 41, 1091
- [30] M. Henneaux, C. Teitelboim. Quantisation of Gauge Systems. Princeton University Press, Princeton, 1992
- [31] T. Thiemann. Observations on representations of the spatial diffeomorphism group and algebra in all dimensions.
- [32] S. W. Hawking. Zeta function regularization of path integrals in curved space time. Commun. Math. Phys. 55 (1977) 133–148

E. Elizalde, S. D. Odintsov, A. Romeo, A. A. Bytsenko, S. Zerbini. Zeta regularization techniques with applications. World Scientific, Singapore, 1994

[33] K. Kuchar. Dirac Constraint Quantization of a Parametrized Field Theory by Anomaly - Free Operator Representations of Space-time Diffeomorphisms. Phys. Rev. D 39 (1989) 2263-2280.
K. Kuchar. Parametrized Scalar Field on R X S(1): Dynamical Pictures, Space-time Diffeomorphisms, and Conformal Isometries. Phys. Rev. D 39 (1989) 1579-1593
T. Thiemann, E. A. Zwicknagel. Hamiltonian renormalization. VI. Parametrized field theory on the cylinder. Phys.Rev. D 108 (2023) 12, 125006. e-Print: 2207.08290 [gr-qc]

- [34] T. Thiemann. Exact quantisation of $U(1)^3$ quantum gravity via exponentiation of the hypersurface deformation algebroid. Class. Quant. Grav. **35** (2023) 24, 245003. e-Print: 2207.08302 [gr-qc]
- [35] H. Heuser. Lehrbuch der Analysis. Vol. 1. Springer Verlag, Berlin, 2013
- [36] J. Glimm. Boson fields with the : Φ^4 : interaction in three dimensions. Comm. Math. Phys. 10 (1968) 1-47.
- [37] Beatriz Elizaga Navascues, Mercedes Martin-Benito, Guillermo A. Mena Marugan. Hybrid models in loop quantum cosmology Int. J. Mod. Phys. D25 (2016), 1642007. e-Print: arXiv:1608.05947
 Laura Castello Gomar, Mercedes Martin-Benito, Guillermo A. Mena Marugan. Gauge-Invariant Perturbations in Hybrid Quantum Cosmology JCAP 1506 (2015), 045. e-Print: arXiv:1503.03907
 Laura Castello Gomar, Mercedes Martin-Benito, Guillermo A. Mena Marugan. Quantum corrections to the Mukhanov-Sasaki equations. Phys. Rev. D93 (2016), 104025. e-Print: arXiv:1603.08448

[38] T. Thiemann. Symmetry reduction, gauge reduction, backreaction and consistent higher order perturbation theory.

T. Thiemann. Quantum Field Theory of Black Hole Perturbations with Backreaction: I. General framework. J. Neuser, T. Thiemann. Quantum Field Theory of Black Hole Perturbations with Backreaction: II. Spherically symmetric 2nd order Einstein sector.

III. Spherically symmetric 2nd order Maxwell sector.

J. Neuser. Quantum Field Theory of Black Hole Perturbations with Backreaction: IV. Spherically symmetric 2nd order Einstein-Maxwell in generalised gauges.