

The heat equation with time-correlated random potential in $d = 2$: Edwards-Wilkinson fluctuations

Sotirios Kotitsas*

Abstract

We consider the stochastic PDE:

$$\partial_t u(t, x) = \frac{1}{2} \Delta u(t, x) + \beta u(t, x) V(t, x),$$

in dimension $d = 2$, where the potential V is the space and time mollification of the two-dimensional space-time white noise. We show that after renormalizing, the fluctuations of the solution converge to the Edwards-Wilkinson limit with an explicit effective variance and constant effective diffusivity. Our main tool is a Markov chain on the space of paths which we use to establish an extension of the Kallianpur-Robbins law [KR53] to a specific regenerative process.

Keywords. critical SPDEs, stochastic heat equation, Edwards-Wilkinson limit, Kallianpur-Robbins theorem, regenerative processes.

Contents

1	Introduction	2
1.1	Comments and related results	3
1.2	Future Directions	6
1.3	Idea of the proof and outline	7
1.4	Notation	13
2	Proof of Theorem 1.1	13
3	The Markov Chain on Ω_1	17
3.1	The construction	17
3.2	The coupling in $d = 2$	21
3.3	The Mixing property	23
3.4	Intersections of independent paths	26
4	Proof of Proposition 1.4	35
A	Estimates on the Total Path Increments	45
B	Kallianpur-Robbins Estimates	50
C	The normalization constant $\zeta_{t/\epsilon^2}^{(\epsilon)}$	55

*Email: Sotirios.Kotitsas@warwick.ac.uk, University of Warwick, UK

1 Introduction

We are interested in the following stochastic heat equation in dimension $d = 2$:

$$\partial_t u(t, x) = \frac{1}{2} \Delta u(t, x) + \beta u(t, x) V(t, x), \quad u(0, x) = 1, \quad (1.1)$$

where

$$V(t, x) := \int_{\mathbb{R}^{1+2}} \phi(t-s) \psi(x-y) d\xi(s, y). \quad (1.2)$$

Here ξ is the space-time white noise on $\mathbb{R} \times \mathbb{R}^2$ over a probability space $(\mathbf{\Omega}, \mathcal{F}, \mathbf{P})$, and $\phi : \mathbb{R} \rightarrow \mathbb{R}$, $\psi : \mathbb{R}^2 \rightarrow \mathbb{R}$ are positive smooth functions. We assume that ϕ is supported on $[0, 1]$ and that ψ is symmetric and is supported on $|x| \leq 1/2$. The coupling constant β is tuning the strength of the random potential V .

The potential V is Gaussian and stationary. We denote by R its correlation function:

$$R(s, y) := \mathbf{E} \left[V(s, y) V(0, 0) \right] = \phi \star \tilde{\phi}(s) \psi \star \psi(y),$$

where \star denotes the convolution of two functions, \mathbf{E} is the expectation with respect to the white noise ξ and $\tilde{\phi}(s) = \phi(-s)$. The function R is compactly supported and $R(s, \cdot) = 0$ for $|s| \geq 1$.

Our goal is to understand the large space-time behavior of the solution to (1.1) i.e. the behavior of the random variable $u_\epsilon(t, x) = u(\frac{t}{\epsilon^2}, \frac{x}{\epsilon})$, as $\epsilon \rightarrow 0$. More specifically, we aim to understand the fluctuations of $u_\epsilon(t, x)$ as a distribution: For $g \in C_c^\infty(\mathbb{R}^2)$ we aim to find the limiting law as $\epsilon \rightarrow 0$ of

$$\int_{\mathbb{R}^2} u_\epsilon(t, x) g(x) dx,$$

after centering and re-scaling.

Similar problems have been considered before (see the next subsection for more details). In [GRZ18] the authors considered the same problem in $d \geq 3$. They proved that the limiting law in this case is Gaussian with mean 0 and with an explicit variance. More specifically, they proved that the limiting law is given by

$$\int_{\mathbb{R}^2} \mathcal{U}(t, x) g(x) dx,$$

where \mathcal{U} solves the additive stochastic heat equation with an effective diffusivity $a_{eff}(\beta) \in \mathbb{R}^{2 \times 2}$ (= the space of 2×2 matrices with real coefficients) and effective variance $v_{eff}(\beta) > 0$:

$$\partial_t \mathcal{U} = \frac{1}{2} \nabla a_{eff}(\beta) \nabla \mathcal{U} + \beta v_{eff}(\beta) \xi, \quad \mathcal{U}(0, x) = 0. \quad (1.3)$$

The additive stochastic heat equation is sometimes called the Edwards-Wilkinson equation.

Here we extend the Edwards-Wilkinson limit for the equation (1.1) to dimension $d = 2$. The important difference is that $d = 2$ is the critical dimension (see below for more details). For this reason, we also tune the coupling constant β to go 0 as $\epsilon \rightarrow 0$. More specifically, we consider the following equation:

$$\partial_t u(t, x) = \frac{1}{2} \Delta u(t, x) + \frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} u(t, x) V(t, x), \quad u(0, x) = 1. \quad (1.4)$$

Set:

$$u_\epsilon(t, x) := u\left(\frac{t}{\epsilon^2}, \frac{x}{\epsilon}\right),$$

where u solves (1.4). Our main result is the following:

Theorem 1.1. *Let $g \in C_c(\mathbb{R}^2)$ and:*

$$\hat{\beta}_c(R) = \sqrt{2\pi/\|R\|_1},$$

where $\|R\|_1 = \int_{\mathbb{R}^{1+2}} R(s, y) dy ds$. There exists $\zeta^{(\epsilon)} : \mathbb{R}_{>0} \rightarrow \mathbb{R}_{>0}$ such that $\zeta_{t/\epsilon^2}^{(\epsilon)} \rightarrow +\infty$ as $\epsilon \rightarrow 0$, and such that for all $\hat{\beta} < \hat{\beta}_c(R)$

$$\int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} u_\epsilon(t, x) g(x) dx \rightarrow \int_{\mathbb{R}^2} g(x) dx,$$

in probability as $\epsilon \rightarrow 0$. Moreover,

$$\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx \Rightarrow \int_{\mathbb{R}^2} \mathcal{U}(t, x) g(x) dx \quad (1.5)$$

in distribution, where \mathcal{U} solves the additive heat equation (1.3) with effective variance $v_{eff}(\hat{\beta}) > 0$ given by the formula:

$$v_{eff}(\hat{\beta})^2 = \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2}\right)^{-1} \cdot \|R\|_1^2, \quad (1.6)$$

and with trivial effective diffusivity $a_{eff}(\hat{\beta}) = I_{2 \times 2}$, where $I_{2 \times 2}$ is the 2 by 2 identity matrix.

The choice of flat initial data is for simplicity and our results can be generalized to more general initial conditions by using similar arguments.

The (first-order) asymptotics of $\zeta_{t/\epsilon^2}^{(\epsilon)}$, as $\epsilon \rightarrow 0$, are:

$$\zeta_{t/\epsilon^2}^{(\epsilon)} = (C_1 + o(1)) \frac{t}{\epsilon^2 \log \epsilon^{-1}} + o(1),$$

where C_1 is an explicit constant and $o(1)$ indicates a function of ϵ that goes to 0 as $\epsilon \rightarrow 0$ (see **Appendix C**).

1.1 Comments and related results

Equation (1.1) is a regularization of the multiplicative stochastic heat equation (mSHE), formally written as

$$\partial_t u(t, x) = \frac{1}{2} \Delta u(t, x) + \beta u(t, x) \xi(t, x),$$

where $x \in \mathbb{R}^d$, and $\xi(t, x)$ is the space-time white noise i.e. a mean zero Gaussian distribution with covariance

$$\mathbf{E}[\xi(s, y) \xi(t, x)] = \delta(s - t) \delta(x - y),$$

where δ is the Dirac delta function. Again, $\beta > 0$ is the coupling constant. This equation has a long history and is related to the KPZ equation (via the Cole-Hopf transform) and to directed polymers (via the Feynman-Kac formula). In dimension 1 we can give a direct meaning to this equation, by using the notions of the Ito-Walsh integral and the mild solution (see for example [Dal+08]). However, the noise ξ becomes increasingly irregular as the dimension increases and this theory does not work when $d \geq 2$. In particular, the product $u(t, x) \xi(t, x)$ does not make sense in $d \geq 2$.

The mSHE is a singular SPDE, a term usually reserved for SPDEs that do not make direct sense, usually due to some non-linearity in the equation, and we have to interpret them in a specific way. This class contains many interesting equations from mathematical physics like the KPZ equation, the Φ^4 -model, the Allen-Cahn equation, and the stochastic Navier-Stokes equations. Making direct sense of these equations is a challenging mathematical problem.

Some progress in studying these types of equations has been made in recent years most famously with the theories of regularity structures [Hai13],[Hai14], paracontrolled distributions [GP16] and renormalization group [Kup14; Duc22]. These theories focus on sub-critical singular SPDEs, which, loosely speaking, are those SPDEs that in small scales the cause of the problem (e.g. the non-linearity) is formally small, and the equation should behave as a nicer SPDE (see [CW17] for this semi-formal definition). The property of being sub-critical usually depends on the dimension of the underlying space and the specific form of the equation. For example, mSHE is sub-critical in $d = 1$, critical for $d = 2$, and supercritical for $d \geq 3$. Indeed if we take $u^\lambda(t, x) = u(\lambda^2 t, \lambda x)$, where u formally solves the mSHE, by the scaling properties of the space-time white noise $u^\lambda(t, x)$ is equal in distribution to the solution of

$$\partial_t u(t, x) = \frac{1}{2} \Delta u(t, x) + \lambda^{\frac{2-d}{2}} \beta u(t, x) \xi(t, x).$$

which demonstrates exactly this principle as $\lambda \rightarrow 0$.

For $d \geq 2$ the usual way to make sense of this equation is by regularizing the noise and studying the corresponding solution as we remove the regularization. For example, we can mollify the noise in space by an approximation of the identity. We define

$$\xi_\epsilon(t, x) = \psi_\epsilon \star' \xi(t, x)$$

where \star' denotes convolution in space, $\psi \in C_c^\infty(\mathbb{R}^d)$ and $\psi_\epsilon(x) = \epsilon^{-d} \psi(x/\epsilon)$. We then consider the corresponding solution u^ϵ with initial data equal¹ to 1. This solution now exists (again see [Dal+08]) for every $\epsilon > 0$. The interest now is the various limits of this random variable as $\epsilon \rightarrow 0$.

In general, as $\epsilon \rightarrow 0$, we also need to tune the coupling constant to get nontrivial limits. More specifically we need to take $\beta_\epsilon = \hat{\beta} \cdot \epsilon^{(d-2)/2}$. With this choice $u^\epsilon(t, x)$ is equal in distribution to $u(\frac{t}{\epsilon^2}, \frac{x}{\epsilon})$ where u solves (1.1) with $\beta = \hat{\beta}$ and V white in time (i.e. $\phi = \delta$).

In $d = 1$ the solution u^ϵ converges in L^2 to the mSHE without mollification, which is well defined. In higher dimensions the situation is different. For $d \geq 3$ it has been shown that there is a phase transition in $\hat{\beta}$ (see for example where, for $\hat{\beta}$ under a specific threshold $\hat{\beta}_c$ (called the weak disorder regime), the solution $u^\epsilon(t, x)$ has a nontrivial positive limit in distribution for every (t, x) . For $\hat{\beta} \geq \hat{\beta}_c$ the solution converges to zero. There is even a smaller regime $\hat{\beta} < \hat{\beta}_{L^2}$, called the L^2 regime, where the solution $u^\epsilon(t, x)$ has a finite second moment as $\epsilon \rightarrow 0$. In $d = 2$ we need to introduce a further modification to the equation by tuning the coupling constant as $\beta_\epsilon = \hat{\beta} / \sqrt{\log \epsilon^{-1}}$. In that case, a similar phase transition has been identified (see [CSZ17]) where the solution $u^\epsilon(t, x)$, for fixed (t, x) , converges in distribution to a nontrivial limit² for $\hat{\beta}$ below a critical value (called the subcritical regime) and to zero otherwise. It should be noted that these critical values $\hat{\beta}_c$ depend on the mollifier that we used. For example, in $d = 2$, $\hat{\beta}_c = \sqrt{2\pi} / \|\psi\|_1$.

The solution u^ϵ can be studied as a random distribution, in particular after we test it against a test function. In this case, the works [CSZ17] for $d = 2$ and [GL19], [LZ22], [CNN22] for $d \geq 3$, showed that we have a limit theorem and Gaussian fluctuations in the subcritical/ L^2 regime

¹This is just for simplicity.

²In particular to a log-normal random variable.

respectively. Regarding the fluctuations, it was proved that we fall into the Edwards-Wilkinson universality class. This means that

$$\frac{\hat{\beta}}{\beta_\epsilon} \int_{\mathbb{R}^2} (u^\epsilon(t, x) - \mathbf{E}[u^\epsilon(t, x)])g(x)dx$$

converges in distribution to

$$\int_{\mathbb{R}^2} \mathcal{U}(t, x)g(x)dx,$$

where $\mathcal{U}(t, x)$ solves equation (1.3). In both cases, $d = 2$ or $d \geq 3$, the effective diffusivity is trivial, $a_{eff} = I_{2 \times 2}$. For $d \geq 3$ we point out that the Edwards-Wilkinson limit has been proved in the full L^2 regime in [LZ22], [CNN22] while in [GL19] it was proved for $\hat{\beta}$ small enough.

The regularization we study here is via mollification of the noise in both space and time. We focus on studying the large-scale behavior of the solution with the noise regularized at a fixed scale. The added difficulty to this approach is that, since we do not use the Ito-Walsh integral, we destroy the martingale structure of the solution and create time correlations that add up exponentially. This is the reason for the exponential correction term appearing in the limit law in **Theorem 1.1** (which appears $d \geq 3$ as well). This term appears because the mean of $u_\epsilon(t, x)$ blows up exponentially as $\epsilon \rightarrow 0$ and the term $e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}}$ is introduced exactly so that the mean remains bounded as $\epsilon \rightarrow 0$.

This regularization has been studied in other dimensions as well. In [GT19] the authors study the mSHE with space-time mollification has been done in $d = 1$, and show that the corresponding solution converges to the solution of the mSHE without mollification in every L^p space. There is also the more general result of [HP14] where the authors show that solutions of a general class of SPDEs with space-time mollification converge to the corresponding SPDEs without mollification after renormalization.

In $d \geq 3$ the authors of [GRZ18] were able to prove the Edwards-Wilkinson limit as described above. The interesting aspect of this result is that they get an added effective diffusivity term i.e. $a_{eff} \neq I_{d \times d}$. Moreover, in [Dun+21] the authors were able to study the pointwise statistics of the solution and express the effective diffusivity and effective variance in terms of objects from stochastic homogenization theory. We should note that the result of [GRZ18] was proved for $\hat{\beta}$ being sufficiently small.

In $d = 2$ there are no results, to our knowledge, that treat the mSHE with space-time mollification either at the level of the pointwise statistics or at the level of local fluctuations (i.e. after we test against a test function). In analogy with the case where we use only mollification in space, we scale the coupling constant as $\beta = \hat{\beta}/\sqrt{\log \epsilon^{-1}}$. Again, we expect that there is a subcritical regime. This is suggested by **Theorem 1.1** since for all $\hat{\beta} < \hat{\beta}_c(R)$ we have a nontrivial Central Limit Theorem (CLT) and the effective variance that we get blows up at $\hat{\beta} = \hat{\beta}_c(R)$. Given this observation, we can prove **Theorem 1.1** in the optimal range of $\hat{\beta}$.

We should also note that unlike in $d \geq 3$, we get a "trivial" effective diffusivity for the Edwards-Wilkinson limit. The reason for this becomes apparent from the proof. Loosely speaking, the effective diffusivity in [GRZ18] is the second moment of the total path increment of a path with law absolutely continuous with respect to the Wiener measure. More specifically, its law is an exponentially tilted measure with the Wiener measure as the reference measure. These paths appear due to the Feynman-Kac formula and the exponential tilting is due to time mollification (see below for more details). It so happens that the exponential tilting depends on the coupling constant $\beta_\epsilon = \hat{\beta}/\sqrt{\log \epsilon^{-1}}$. As $\epsilon \rightarrow 0$ the tilting disappears making the total path increment look like the total path increment of a Brownian motion, which gives us the trivial effective diffusivity.

Finally, we should compare **Theorem 1.1** with its analog in $d = 2$ when we use space mollification. In the latter case, let u_ψ solve the mSHE with only space mollification, using ψ as the mollifier, with coupling constant $\beta_\epsilon = \hat{\beta}/\sqrt{\log \epsilon^{-1}}$. From [CSZ17], the effective variance of the Edwards-Wilkinson limit is:

$$\mathcal{V}_{eff}^2(\hat{\beta}) = \left(1 - \frac{\hat{\beta}^2 \|\psi\|_1^2}{2\pi}\right)^{-1} \|\psi\|_1^2.$$

If we additionally mollify time using ϕ , as in (1.2), then **Theorem 1.1** gives an effective variance:

$$v_{eff}^2(\hat{\beta}) = \left(1 - \frac{\hat{\beta}^2 \|\psi\|_1^2 \|\phi\|_1^2}{2\pi}\right)^{-1} \cdot \|\psi\|_1^2 \|\phi\|_1^2.$$

This shows that, when ϕ is a probability density

$$\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx$$

converges to the same limit as

$$\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} (u_\psi(t/\epsilon^2, x/\epsilon) - \mathbf{E}[u_\psi(t/\epsilon^2, x/\epsilon)]) g(x) dx.$$

Observe that this does not happen in $d \geq 3$. Indeed, in $d \geq 3$ when V is white in time the Edwards-Wilkinson limit has effective diffusivity $a_{eff} = I_{d \times d}$, while when V is correlated in time, $a_{eff} \neq I_{d \times d}$.

1.2 Future Directions

The mSHE is related to the KPZ equation via the Cole-Hopf transform. If $u(t, x)$ solves (1.4) then

$$h(t, x) := \log u(t, x)$$

solves the equation

$$\partial_t h(t, x) = \frac{1}{2} \Delta h(t, x) + |\nabla h(t, x)|^2 + \frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} V(t, x), \quad h(0, x) = 0.$$

In [CSZ20] it is proved that when V is white in time the random variable $h(t/\epsilon^2, x/\epsilon)$ converges in distribution to a Gaussian random variable for all $\hat{\beta}$ in the subcritical regime. Moreover, the fluctuations converge to the Edwards-Wilkinson limit. This is also proved in [Gu18] for small enough $\hat{\beta}$. It is interesting to try to combine our methods with the methods of [CSZ20] and of [Gu18] to prove that when V is as in (1.2), $h(t/\epsilon^2, x/\epsilon)$ is asymptotically Gaussian, with fluctuations in the Edwards-Wilkinson class.

Finally, we mention that in $d = 2$ with V white in time and at the critical point, the mSHE has an interesting and nontrivial behavior. Loosely speaking, when $\hat{\beta} = \hat{\beta}_c$ and for $g \in C_c(\mathbb{R}^2)$ the random variable

$$\int_{\mathbb{R}^2} u_\epsilon(t, x) g(x) dx \tag{1.7}$$

converges in distribution as $\epsilon \rightarrow 0$ [CSZ21]. The limiting object is called the critical $2d$ Stochastic Heat Flow. The moments of the critical $2d$ Stochastic Heat Flow admit explicit expressions [GQT21] and it is known that it is not a Gaussian multiplicative chaos [CSZ23]. **Theorem 1.1** indicates that a critical point $\hat{\beta}_c$ exists when V is correlated in time. Therefore, it is interesting to consider (1.4) with V as in (1.2) at the critical point and establish convergence in distribution for (1.7). If this is possible it is also interesting to compare this limiting object with the critical Stochastic Heat Flow obtained when V is white in time.

1.3 Idea of the proof and outline

We use methods and arguments inspired from [GRZ18] to prove **Theorem 1.1**. More specifically, for any $g \in C_c(\mathbb{R}^2)$, we need with two formulas from [GRZ18] for the random variable:

$$\int_{\mathbb{R}^2} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)])g(x)dx. \quad (1.8)$$

These formulas are proved in [GRZ18] for $d \geq 3$ and their proof carries exactly in the case $d = 2$. We start with the Feynman-Kac representation of $u(t, x)$. Since $V(t, x)$ is a (random) smooth function we have

$$u(t, x) = \mathbb{E}_B \left[\exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} \int_0^t V(t-s, x+B_s) ds \right) \right], \quad (1.9)$$

where \mathbb{E}_B is the expectation with respect to Brownian motion starting at 0, independent from the noise. Because of the time correlations induced by the time mollification, the usual martingale structure of the solution is destroyed. In practice, this is what induces the exponential tilting of the Wiener measure. More precisely, on the space:

$$\Omega_T = \{\omega \in C([0, T]) \mid \omega(0) = 0\}$$

equipped with the usual Borel σ -algebra, we define the following probability measure:

$$d\hat{\mathbb{P}}_T^{(\epsilon)}(B) := \exp \left(\frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \int_{[0, T]^2} R(s-u, B(s) - B(u)) dsdu - \zeta_T^{(\epsilon)} \right) W_T(dB), \quad (1.10)$$

where $W_T(dB)$ is the usual Wiener measure on Ω_T and $\zeta_T^{(\epsilon)}$ is the normalization constant:

$$\zeta_T^{(\epsilon)} := \log \int_{\Omega_T} \exp \left(\frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \int_{[0, T]^2} R(s-u, B(s) - B(u)) dsdu \right) W_T(dB). \quad (1.11)$$

We denote by $\hat{\mathbb{E}}_{B, T}$ the corresponding expectation. This exponential tilting already appears at the level of the moments of $u(t, x)$, as the next calculation shows:

Proposition 1.1. *If R is the correlation function of V then, for every $n \in \mathbb{N}$, we have:*

$$e^{-n\zeta_t^{(\epsilon)}} \mathbf{E}[u(t, x)^n] = \hat{\mathbb{E}}_{B^1, \dots, B^n; t} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \sum_{1 \leq i < j \leq n} \int_{[0, t]^2} R(s-u, B^i(s) - B^j(u)) dsdu \right) \right],$$

where $\hat{\mathbb{E}}_{B^1, \dots, B^n; t}$ denotes the expectation under n independent paths sampled from the exponentially tilted measure $\hat{\mathbb{P}}_t^{(\epsilon)}$.

Proof. For simplicity, we prove this for $n = 2$. The general formula is proved similarly. By plugging into (1.9) the definition of V , (1.2), we get:

$$u(t, x) = \mathbb{E}_B \left[\exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} \int_0^t \int_{\mathbb{R}} \int_{\mathbb{R}^2} \phi(t-s-s_1) \psi(x+B(s)-x_1) d\xi(s_1, x_1) \right) \right].$$

Now we define:

$$\Phi_{t, x, B}(s_1, x_1) := \int_0^t \phi(t-s-s_1) \psi(x+B(s)-x_1) ds, \quad (1.12)$$

$$M_{t,x,B}(\infty) := \int_{\mathbb{R}^{1+2}} \Phi_{t,x,B}(s_1, x_1) d\xi(s_1, x_1). \quad (1.13)$$

For every Brownian path B , $M_{t,x,B}(\infty)$ is a mean Gaussian random variable with second moment equal to

$$\int_{\mathbb{R}^{1+2}} \Phi_{t,x,B}(s_1, x_1)^2 ds_1 dx_1 = \int_{[0,t]^2} R(s-u, B(s) - B(u)) ds du. \quad (1.14)$$

With this notation, we can write

$$u(t, x) = \mathbb{E}_B \left[\exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} M_{t,x,B}(\infty) \right) \right].$$

From (1.11) and (1.14)

$$\mathbf{E}[u(t, x)] = \exp(\zeta_t^{(\epsilon)}).$$

For the second moment, we have

$$\mathbf{E}[u(t, x)^2] = \mathbb{E}_{B^1} \mathbb{E}_{B^2} \left[\mathbf{E} \left[\exp \left(\frac{\beta}{\sqrt{\log \epsilon^{-1}}} (M_{t,x,B^1}(\infty) + M_{t,x,B^2}(\infty)) \right) \right] \right],$$

which is equal to

$$\mathbb{E}_{B^1} \mathbb{E}_{B^2} \left[\exp \left(\frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} (\mathbf{E}[M_{t,x,B^1}(\infty)] + \mathbf{E}[M_{t,x,B^2}(\infty)] + 2\mathbf{E}[M_{t,x,B^1}(\infty)M_{t,x,B^2}(\infty)]) \right) \right], \quad (1.15)$$

where $\langle M_{t,x,B^1}, M_{t,x,B^2} \rangle(\infty)$ is the limit as $r \rightarrow \infty$ of the quadratic covariation $\langle M_{t,x,B^1}, M_{t,x,B^2} \rangle(r)$ of the martingales M_{t,x,B^1} and M_{t,x,B^2} . A straightforward calculation shows that

$$\mathbf{E}[M_{t,x,B^1}(\infty)M_{t,x,B^2}(\infty)] = \int_{[0,t]^2} R(s-u, B^1(s) - B^2(u)) ds du.$$

Plugging this in (1.15) we get

$$e^{-2\zeta_t^{(\epsilon)}} \mathbf{E}[u(t, x)^2] = \hat{\mathbb{E}}_{B^1, B^2; t} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0,t]^2} R(s-u, B^1(s) - B^2(u)) ds du \right) \right].$$

□

The above calculation indicates that $e^{-\zeta_t^{(\epsilon)}}$ is the correct exponential correction needed to keep $\mathbf{E}[u_\epsilon(t, x)]$ bounded. Indeed, for all $\epsilon > 0$,

$$e^{-\zeta_t^{(\epsilon)}} \mathbf{E}[u_\epsilon(t, x)] = 1. \quad (1.16)$$

The proof of **Theorem 1.1** is split into two steps. The first step is to find the limiting variance of:

$$\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_t^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx. \quad (1.17)$$

Observe that the convergence of the variance of (1.17) proves that $\int_{\mathbb{R}^2} e^{-\zeta_t^{(\epsilon)}} u_\epsilon(t, x) g(x) dx \rightarrow \int_{\mathbb{R}^2} g(x) dx$ in \mathbf{P} -probability as $\epsilon \rightarrow 0$.

The second step is using mixing arguments to prove the central limit theorem (1.5). These arguments allow us to approximate (1.17) in L^2 by a sum of independent random variables. Then we conclude by applying Lindenberg's CLT.

As mentioned, we need two formulas from [GRZ18]. We define

$$M_{t,x,B}(r) := \int_{-\infty}^r \int_{\mathbb{R}^2} \Phi_{t,x,B}(s_1, x_1) d\xi(s_1, x_1). \quad (1.18)$$

Observe that $(M_{t,x,B}(r))_{r \in \mathbb{R}}$ is a square-integrable continuous martingale with quadratic variation equal to:

$$\langle M_{t,x,B} \rangle(r) = \int_{-\infty}^r \int_{\mathbb{R}^2} \Phi_{t,x,B}(s_1, x_1)^2 ds_1 dx_1. \quad (1.19)$$

The first formula we need is contained in the following proposition. The proof in $d = 2$ is the same as in [GRZ18] (see Lemma 2.1 from the same paper):

Proposition 1.2. *We have that:*

$$(u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)])e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} = \frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} \int_{-1}^{t/\epsilon^2} \int_{\mathbb{R}^2} \hat{\mathbb{E}}_{B,t/\epsilon^2} \left[\Phi_{t,x,B}^\epsilon(r, y) \exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} M_{t,x,B}^\epsilon(r) - \frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \langle M_{t,x,B}^\epsilon \rangle(r) \right) \right] d\xi(r, y),$$

where $\Phi_{t,x,B}^\epsilon = \Phi_{t/\epsilon^2, x/\epsilon, B}$ and $M_{t,x,B}^\epsilon = M_{t/\epsilon^2, x/\epsilon, B}$, with $\Phi_{t,x,B}$ and $M_{t,x,B}$ defined in (1.9) and (1.12) respectively.

This formula is proved by combining the Clark-Ocone formula³

$$u(t, x) - \mathbf{E}[u(t, x)] = \int_{-\infty}^t \int_{\mathbb{R}^2} \mathbf{E}[D_{r,y} u(t, x) | \mathcal{F}_r] d\xi(r, y),$$

with the Feynman-Kac formula and similar calculations as in the previous proof.

Proposition 1.2 gives us a stochastic integral representation for (1.17)

$$\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx = \hat{\beta} \int_{-1}^{t/\epsilon^2} \int_{\mathbb{R}^2} Z_t^\epsilon(r, y) d\xi(r, y),$$

where:

$$Z_t^\epsilon(r, y) := \int_{\mathbb{R}^2} g(x) \hat{\mathbb{E}}_{B,t/\epsilon^2} \left[\Phi_{t,x,B}^\epsilon(r, y) \exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} M_{t,x,B}^\epsilon(r) - \frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \langle M_{t,x,B}^\epsilon \rangle(r) \right) \right] dx.$$

We are going to use this formula in the mixing arguments mentioned above to prove **Theorem 1.1**. The mixing arguments are implemented by modifying the martingale $M_{t,x,B}^\epsilon$ so that the stochastic integral above can be split into a sum of independent random variables and a 'negligible' remainder. Then, as mentioned, Lindenberg's CLT and the information we will get from the limiting variance will allow us to conclude.

This stochastic integral representation of (1.17) is also useful for the calculation of its limiting variance, which is the more complicated step. This is the content of the final formula needed from [GRZ18]. First, we need to introduce some notation. We define:

$$I_\epsilon(x_1, x_2, y, s_1, s_2, r) := \prod_{i=1}^2 g(\epsilon x_i - \epsilon B^i((t-r)/\epsilon^2 - s_i) + y), \quad (1.20)$$

³ $D_{r,y} u(t, x)$ is the Malliavin derivative of $u(t, x)$ at (r, y) , see [Nua09]

$$R_\phi(t_1, t_2) := \int_0^\infty \phi(s - t_1)\phi(s - t_2)ds, \quad (1.21)$$

$$J_\epsilon(M_1, M_2) := \frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{-1}^{M_1} \int_{-1}^{M_2} R_\phi(u_1, u_2) \psi \star \psi(x_1 - x_2 + \Delta B_{(t-r)/\epsilon^2 - s_1, (t-r)/\epsilon^2 + u_1}^1 - \Delta B_{(t-r)/\epsilon^2 - s_2, (t-r)/\epsilon^2 + u_2}^2) du_1 du_2, \quad (1.22)$$

where for a path B , we define $\Delta B_{s,u} = B(s) - B(u)$.

Proposition 1.3. *For any $t_1 < t - \epsilon^2$ we have the following formula:*

$$\mathbf{Var} \left(\int_{-\infty}^{t_1/\epsilon^2} \int_{\mathbb{R}^2} Z_t^\epsilon(r, y) d\xi(r, y) \right) = \int_0^t \int_{\mathbb{R}^6} \int_{[0,1]^2} \hat{\mathbb{E}}_{B,t/\epsilon^2} [I_\epsilon e^{J_\epsilon(r/\epsilon^2, r/\epsilon^2)}] \prod_{i=1}^2 \phi(s_i) \psi(x_i) d\bar{s} d\bar{x} dy dr, \quad (1.23)$$

with $d\bar{s} = ds_1 ds_2$, $d\bar{x} = dx_1 dx_2$.

The calculation is the same as in $d \geq 3$ and involves using Ito's isometry for the stochastic integral appearing above and a series of changes of variables. It is done in detail in [GRZ18] (see **Lemma 2.3** in the paper). Also, a similar formula is true for a general $t_1 \leq t$, with some modifications. The modifications are the following:

- The domain of s_1, s_2 changes to $[0, (t-r)/\epsilon^2]^2$.
- The domain of u_1, u_2 that appear in the definition of J_ϵ changes to $[-(t-r)/\epsilon^2, r/\epsilon^2]^2$.

Since ϕ is supported in $[0, 1]$, $R_\phi(u_1, u_2)$ is zero if either u_i is less than -1 . This shows that these changes create a difference only when $t - r \leq \epsilon^2$.

We are going to use this formula to find the limiting variance of (1.17). To do this, we analyze the functional appearing inside the integral on the right-hand side of (1.23). In particular, we define:

$$\mathcal{F}_\epsilon(r, y, M_1, M_2) = \int_{\mathbb{R}^4} \int_{[0,1]^2} \hat{\mathbb{E}}_{B,t/\epsilon^2} [I_\epsilon e^{J_\epsilon(M_1, M_2)}] \prod_{i=1}^2 \phi(s_i) \psi(x_i) d\bar{s} d\bar{x}. \quad (1.24)$$

We seek to find a limit for this object as well as good bounds in y so we can apply dominated convergence and find the limit of the variance. This is done in the following proposition, which is proved in **Section 4**:

Proposition 1.4. *Let $0 < M_1(\epsilon), M_2(\epsilon) \leq r$ be such that $\log M_i(\epsilon)/\log \epsilon^{-1} \rightarrow 0$ and such that for all ϵ small enough either $M_1(\epsilon) = M_2(\epsilon)$ or $M_1(\epsilon) - M_2(\epsilon) \geq c > 0$ for some c . Then for any $r \in (0, t)$, $y \in \mathbb{R}^2$, $\hat{\beta} < \hat{\beta}_c(R)$ and $k > 0$ we have:*

$$|\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2)| \leq C(1 \wedge |y|^{-k}), \quad (1.25)$$

for some constant C , depending only on k . Furthermore, as $\epsilon \rightarrow 0$ we have:

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) \rightarrow v_{eff}^2(\hat{\beta}) p_{t-r} \star g(y)^2,$$

where $p_t(y) = \frac{1}{2\pi t} e^{-|x|^2/2t}$, is the two dimensional heat kernel, and $v_{eff}^2(\hat{\beta})$ as in **Theorem 1.1**.

Remark 1.1. We put the condition

$$M_1(\epsilon) - M_2(\epsilon) \geq c > 0,$$

for some c , only for convenience. In this case, the proof of **Proposition 1.4** boils down to the case $M_1(\epsilon) = M_2(\epsilon)$. Indeed, if $M_1(\epsilon) - M_2(\epsilon) \geq c > 0$ then, for all ϵ small enough, $M_1(\epsilon)/\epsilon^2 \geq M_2(\epsilon)/\epsilon^2 + 1$. Moreover, since ϕ is supported on $[0, 1]$, $R_\phi(u_1, u_2) = 0$ when $|u_1 - u_2| \geq 1$. Therefore, if $M_1(\epsilon) - M_2(\epsilon) \geq c > 0$, from (1.22):

$$J_\epsilon(M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) = J_\epsilon(M_2(\epsilon)/\epsilon^2 + 1, M_2(\epsilon)/\epsilon^2),$$

for all ϵ sufficiently small. From (1.24), for all ϵ sufficiently small

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) = \mathcal{F}_\epsilon(r, y, M_2(\epsilon)/\epsilon^2 + 1, M_2(\epsilon)/\epsilon^2).$$

Using this equality, we are going to argue in **Section 4** that

$$|\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) - \mathcal{F}_\epsilon(r, y, M_2(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2)| \rightarrow 0,$$

as $\epsilon \rightarrow 0$.

This proposition gives us the limit for the variance appearing in (1.23) for any $t_1 < t - \epsilon^2$. We will argue in **Section 4** that this can be extended to the case $t_1 = t$, thus giving us the limiting variance. This is done in **Corollary 4.1**. Observe that the limiting variance is then equal to

$$\hat{\beta}^2 v_{eff}^2(\hat{\beta}) \int_0^t \int_{\mathbb{R}^2} p_{t-r} \star g(y)^2 dy dr,$$

which is equal to

$$\mathbf{Var} \left(\int_{\mathbb{R}^2} \mathcal{U}(t, x) g(x) dx \right),$$

with $\mathcal{U}(t, x)$ as in **Theorem 1.1**.

In **Section 2** we prove **Theorem 1.1** assuming **Proposition 1.4** and **Corollary 4.1**. As mentioned, we use the mixing arguments appearing in [GRZ18]. However, their arguments can give the CLT only in a restricted region of $\hat{\beta}$. This is because they eventually have to verify Lindenberg's principle and they do this by essentially estimating the fourth moment of (1.8). This restricts the range of $\hat{\beta}$ since they have estimates only for the second moment. We avoid this problem by using an argument from [CSZ20] which uses Gaussian hypercontractivity and the fact that we are strictly inside the subcritical regime.

The rest of the paper is dedicated to the proof of **Proposition 1.4**. It is clear that to prove it we have to deal with functionals of continuous paths which are distributed according to $\hat{\mathbb{P}}_T^{(\epsilon)}$. This is the bulk of this paper. We deal with this in a similar way as in [GRZ18] by introducing a Markov chain on the space of continuous paths over $[0, 1]$ and express the measure $\hat{\mathbb{P}}_T^{(\epsilon)}$ as a transition probability of this Markov chain. The way we can guess this Markov chain is pretty simple: We could try to build a path distributed according to $\hat{\mathbb{P}}_T^{(\epsilon)}$ by gluing together T -many smaller paths over $[0, 1]$ distributed according to $\hat{\mathbb{P}}_1^{(\epsilon)}$. However, these paths will interact under $\hat{\mathbb{P}}_T^{(\epsilon)}$, which will lead us to an explicit probability kernel that captures this interaction. This can be used to define the Markov chain, thus giving us an explicit way to construct a path distributed according to $\hat{\mathbb{P}}_T^{(\epsilon)}$ (for more details see the next section).

In **Sections 3.1** and **3.2** we study the properties of this Markov chain. This Markov chain satisfies a Doeblin condition: the transition probability measure is bounded below by a small multiple of the Wiener measure. From standard arguments, via a coupling with a sequence of Bernoulli random variables, we have a positive probability of sampling independently from the Wiener measure at each step of the Markov chain. This coupling gives the Markov chain a regeneration structure, and more specifically the path built from the Markov chain is a regenerative process [Asm03]. This leads to a random walk encoding the total path increment of the path built from the Markov chain at a regeneration time, which plays an important role in our arguments. Moreover, the fact that we tuned the coupling constant as:

$$\beta_\epsilon = \frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}},$$

leads to some simplifications when compared to the case of $d \geq 3$, for example, a stronger mixing property for the Markov chain as $\epsilon \rightarrow 0$ (see **Lemma 3.1**).

The second moment calculations necessary to prove **Proposition 1.4** is our main departure from the techniques in [GRZ18]. The added difficulty in our case is perhaps best seen at the level of the moments of $u_\epsilon(t, x)$. From **Proposition 1.1** we have:

$$e^{-2\zeta_{t/\epsilon^2}^{(\epsilon)}} \mathbf{E}[u_\epsilon(t, x)^2] = \hat{\mathbb{E}}_{B^1, B^2; t/\epsilon^2} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, t/\epsilon^2]^2} R(s-u, B^1(s) - B^2(u)) ds du \right) \right].$$

The formula holds true when $d \geq 3$, only with $\hat{\beta}^2/\log \epsilon^{-1}$ replaced by an ϵ -independent constant. To find the limit of $e^{-2\zeta_{t/\epsilon^2}^{(\epsilon)}} \mathbf{E}[u_\epsilon(t, x)^2]$ as $\epsilon \rightarrow 0$, we need to consider the limiting distribution of:

$$\int_{[0, t/\epsilon^2]^2} R(s-u, \omega^1(s) - \omega^2(u)) ds du, \quad (1.26)$$

where ω^1, ω^2 are two independent paths built from the Markov chain introduced in **Section 3.1**.

In $d \geq 3$ this is relatively easy to do, since (1.26) converges, as $\epsilon \rightarrow 0$, to a random variable with exponential moments. This is proved in [GRZ18] (Corollary 4.4 in the same paper). This result is somewhat expected since $(\epsilon \omega_{s/\epsilon^2}^1)_{s \leq t}$ converges to a Brownian motion with an effective diffusivity (see Proposition 4.1 in [GRZ18] for the precise statement).

In $d = 2$ we expect $(\epsilon \omega_{s/\epsilon^2}^1)_{s \leq t}$ to behave similarly and to converge to a Brownian motion. Since in $d = 2$ the Brownian motion is recurrent (1.26) diverges almost surely as $\epsilon \rightarrow 0$. This is the reason for the logarithmic dependence of the coupling constant β . Indeed, in **Section 3.4**, we will prove that:

$$\frac{1}{\log \epsilon^{-1}} \int_{[0, t/\epsilon^2]^2} R(s-u, \omega^1(s) - \omega^2(u)) ds du \quad (1.27)$$

converges in distribution to a multiple of an exponential random variable of rate 1.

If R does not depend on time, and ω^1, ω^2 are independent Brownian motions, then this result is the Kallianpur-Robbins law for the Brownian motion [KR53]. Therefore, we can think of the result of **Section 3.4** as a 'non-directed' version of the Kallianpur-Robbins law for the regenerative process (ω^1, ω^2) . Naturally, to prove this, we use the regenerative structure of the process. This means that the process can be split into paths of random length called cycles, such that the cycles are independent and identically distributed. We exploit this structure to analyze the additive functional (1.27), by splitting the functional as a sum over the cycles and then averaging over the law of the cycles. This will give us an additive functional of the total path increments. As mentioned, the total path increment is a random walk in \mathbb{R}^2 . Then, to properly analyze this additive

functional of the total path increments, we will prove a 'non-directed' version of a result in [KR54] (the discrete version of the Kallianpur-Robbins law) (see **Lemma B.1**).

In **Section 4**, we prove **Proposition 1.4** by exploiting the mixing property of the Markov chain and the information we got in **Section 3.4** for the limiting distribution of (1.27). The arguments we use here allow us to prove **Proposition 1.4** in the full subcritical regime:

$$\hat{\beta} < \hat{\beta}_c(R).$$

As mentioned, this regime is optimal since the effective variance is infinite for $\hat{\beta} \geq \hat{\beta}_c(R)$. We can get an explicit form for the effective variance and the corresponding critical value because we know the limiting distribution of the additive functionals appearing in our calculations.

Finally, there are three appendices presenting respectively, some lemmas concerning the total path increments, the 'non-directed' version of the 'discrete' Kallianpur-Robbins law that was mentioned before, and a result giving us the (first order) asymptotics of $\zeta_{t/\epsilon^2}^{(\epsilon)}$.

1.4 Notation

- For μ a probability measure on a measurable space we write $X \sim \mu$ when any random variable, defined over any probability space, has law equal to μ . We write \mathbb{E}_μ for the expectation with respect to μ . Abusing notation, we will write \mathbb{E}_X for the expectation with respect to the law of X again defined over any probability space. Finally, $\mu_1 \times \mu_2$ denotes the product measure of μ_1 and μ_2 .
- $Geo(\gamma)$ denotes the geometric distribution with parameter γ and W_T the Wiener measure on $C([0, T])$.
- For $t \in \mathbb{R}$ we write $[t] \in \mathbb{Z}$ for the integer part of t .
- We use the notation $x \lesssim y$ to mean $x \leq C \cdot y$ for some constant C , irrelevant to the current argument. We also use the big-O notation to mean the same thing: $x = O(y)$ means $x \lesssim y$. Similarly, we say that $x_\epsilon = o(y_\epsilon)$ as $\epsilon \rightarrow 0$ if $x_\epsilon/y_\epsilon \rightarrow 0$ as $\epsilon \rightarrow 0$. Finally, we write $x_\epsilon \sim y_\epsilon$ to indicate that $x_\epsilon/y_\epsilon \rightarrow 1$ as $\epsilon \rightarrow 0$ and we say that x_ϵ is asymptotic to y_ϵ .
- We use bold symbols to emphasize that the symbol is a vector. For two vectors $\mathbf{y}, \mathbf{x} \in \mathbb{R}^d$ we denote by \mathbf{y}^* the transpose of \mathbf{y} and by $\mathbf{x} \cdot \mathbf{y} = \mathbf{x}^* \mathbf{y}$ their inner product. Finally, we denote by $|\mathbf{x}|$ the standard Euclidean norm of \mathbf{x} .

2 Proof of Theorem 1.1

In this section we prove **Theorem 1.1** assuming **Proposition 1.4** and **Corollary 4.1**. From the latter, we get that the variance of

$$\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx$$

converges to a real number as $\epsilon \rightarrow 0$. This observation, when combined with (1.16) and the Markov inequality, proves that

$$\int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} u_\epsilon(t, x) g(x) dx \rightarrow \int_{\mathbb{R}^2} g(x) dx,$$

in **P**-probability, as $\epsilon \rightarrow 0$.

To prove the central limit theorem in **Theorem 1.1** recall that from **Proposition 1.2**, we have:

$$\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx = \hat{\beta} \int_{-1}^{t/\epsilon^2} \int_{\mathbb{R}^2} Z_t^\epsilon(r, y) d\xi(r, y),$$

where:

$$Z_t^\epsilon(r, y) := \int_{\mathbb{R}^2} g(x) \hat{\mathbb{E}}_{B,t} \left[\Phi_{t,x,B}^\epsilon(r, y) \exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} M_{t,x,B}^\epsilon(r) - \frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \langle M_{t,x,B}^\epsilon \rangle(r) \right) \right] dx. \quad (2.1)$$

We will split $[-1, t/\epsilon^2]$ into 'short' and 'long' intervals and then modify the martingale $M_{t,x,B}^\epsilon(r)$ over the long intervals. This will be done so that the contributions from the short intervals are negligible and the contributions over the long intervals are independent. Then, **Proposition 1.4** and Lindenbergs's criterion, will give the full central limit theorem.

More specifically, for $0 < a < \lambda < 2$, we split $[-1, t/\epsilon^2]$ into successive intervals of length ϵ^{-a} (the short intervals) and of length $\epsilon^{-\lambda}$ (the long intervals):

$$[-1, t/\epsilon^2] = [-1, \epsilon^{-a}] \cup (\epsilon^{-a}, \epsilon^{-a} + \epsilon^{-\lambda}] \cup \dots \cup (t_\epsilon, t/\epsilon^2],$$

where $t/\epsilon^2 - t_\epsilon = O(\epsilon^{-\lambda})$. For technical reasons we have to choose a and λ to depend on ϵ . We choose

$$a = 2 - \log \log \epsilon^{-1} / \log \epsilon^{-1} \text{ and } \lambda = 2 - \log \log \epsilon^{-1} / 2 \log \epsilon^{-1}.$$

We denote by $(I_{a,j})$ the collection of all short intervals (where we also include the interval containing t/ϵ^2 to this collection) and by $(I_{\lambda,j})$ the collection of all long intervals. The modification of $M_{t,x,B}^\epsilon$ is defined as

$$\tilde{M}_{t,x,B}^\epsilon(r) := \int_{-\infty}^r \int_{\mathbb{R}^2} \left(\int_0^{r_\epsilon} \phi(t/\epsilon^2 - s_1 - s) \psi(x/\epsilon + B(s_1) - y) ds_1 \right) d\xi(s, y), \quad (2.2)$$

where $r_\epsilon = t/\epsilon^2 - r + 1/2\epsilon^a$ and $r \in I_{\lambda,j}$, for some j (note that in this case, $r \geq \epsilon^{-a}$, and therefore $r_\epsilon < t/\epsilon^2$). Moreover, we define

$$\langle \tilde{M}_{t,x,B}^\epsilon \rangle(r) = \int_{-\infty}^r \int_{\mathbb{R}^2} \left(\int_0^{r_\epsilon} \phi(t/\epsilon^2 - s_1 - s) \psi(x/\epsilon + B(s_1) - y) \right)^2 ds_1 dy ds.$$

Also for $r \in I_{\lambda,j}$ define $\tilde{Z}_t^\epsilon(r, y)$ by (2.1), with $\tilde{M}_{t,x,B}^\epsilon$ replacing $M_{t,x,B}^\epsilon$. Finally, define:

$$\mathcal{X}_j(\hat{\beta}) = \int_{I_{\lambda,j}} \int_{\mathbb{R}^2} \tilde{Z}_t^\epsilon(r, y) d\xi(r, y). \quad (2.3)$$

From [GRZ18], the random variables $\mathcal{X}_j(\hat{\beta})$ are independent. Indeed, the integrand in (2.2) vanishes when $s \leq r - \epsilon^{-a}$, and ϵ is small enough. That is because $t/\epsilon^2 - s_1 - s \geq 1/2\epsilon^a > 1$ when $s \leq r - \epsilon^{-a}$ and ϵ small enough and so $\phi(t/\epsilon^2 - s_1 - s) = 0$. Therefore, $\tilde{M}_{t,x,B}^\epsilon(r)$ depends on the underlying noise $\xi(s, y)$ only for $s \in (r - \epsilon^{-a}, r]$. This proves that $(\mathcal{X}_j(\hat{\beta}))_j$ are independent.

Moreover, the following are true:

- If I_λ is the union of all long intervals then

$$\int_{I_\lambda} \int_{\mathbb{R}^2} \mathbf{E}[|Z_t^\epsilon(r, y) - \tilde{Z}_t^\epsilon(r, y)|^2] dy dr \rightarrow 0, \quad (2.4)$$

as $\epsilon \rightarrow 0$. Indeed, from [GRZ18] (proof of Lemma 3.1) we have

$$\int_{I_\lambda} \int_{\mathbb{R}^2} \mathbf{E}[|Z_t^\epsilon|^2](r, y) dy dr = \int_0^t \int_{\mathbb{R}^2} \mathbf{1}_{\{r/\epsilon^2 \in I_\lambda\}} \mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2) dy dr, \quad (2.5)$$

$$\int_{I_\lambda} \int_{\mathbb{R}^2} \mathbf{E}[|\tilde{Z}_t^\epsilon|^2](r, y) dy dr = \int_0^t \int_{\mathbb{R}^2} \mathbf{1}_{\{r/\epsilon^2 \in I_\lambda\}} \mathcal{F}_\epsilon(r, y, 1/2\epsilon^a, 1/2\epsilon^a) dy dr, \quad (2.6)$$

$$\int_{I_\lambda} \int_{\mathbb{R}^2} \mathbf{E}[Z_t^\epsilon(r, y) \tilde{Z}_t^\epsilon(r, y)] dy dr = \int_0^t \int_{\mathbb{R}^2} \mathbf{1}_{\{r/\epsilon^2 \in I_\lambda\}} \mathcal{F}_\epsilon(r, y, r/\epsilon^2, 1/2\epsilon^a) dy dr. \quad (2.7)$$

Recalling that $a = 2 - \log \log \epsilon^{-1} / \log \epsilon^{-1}$, so that $\epsilon^{-a} = \epsilon^{-2} / \log \epsilon^{-1}$, and using **Proposition 1.4**, we can verify that (2.5), (2.6) and (2.7) have the same limit. This proves (2.4).

- Similarly, if I_a is the union of all short intervals

$$\int_{I_a} \int_{\mathbb{R}^2} \mathbf{E}[|Z_t^\epsilon(r, y)|^2] dy dr \rightarrow 0, \quad (2.8)$$

as $\epsilon \rightarrow 0$. Indeed, from [GRZ18] (proof of Lemma 3.2)

$$\int_{I_a} \int_{\mathbb{R}^2} \mathbf{E}[|Z_t^\epsilon(r, y)|^2] dy dr = \int_0^t \int_{\mathbb{R}^2} \mathbf{1}_{\{r/\epsilon^2 \in I_a\}} \mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2) dy dr.$$

Note that here \mathcal{F}_ϵ has the modifications described after **Proposition 1.3**. Nevertheless, **Proposition 1.4** still holds (as it is argued in **Corollary 4.1** for example). By the bound provided by **Proposition 1.4** and $|\{r \in [0, t] : r/\epsilon^2 \in I_a\}| \rightarrow 0$ as $\epsilon \rightarrow 0$ we see that (2.8) holds.

- Now we will prove that for all $\hat{\beta} < \hat{\beta}_c(R)$

$$\hat{\beta} \sum_j \mathcal{X}_j^\epsilon(\hat{\beta}) \Rightarrow \int_{\mathbb{R}^2} \mathcal{U}(t, x) g(x) dx. \quad (2.9)$$

where \Rightarrow means convergence in distribution. Again from [GRZ18] (proof of Lemma 3.3), we have

$$\hat{\beta}^2 \sum_j \mathbf{Var}[\mathcal{X}_j^\epsilon(\hat{\beta})] = \beta^2 \sum_j \int_0^t \int_{\mathbb{R}^2} \mathbf{1}_{\{r/\epsilon^2 \in I_{\lambda, j}\}} \mathcal{F}_\epsilon(r, y, 1/2\epsilon^a, 1/2\epsilon^a) dy dr.$$

From **Proposition 1.4** this converges to $\hat{\beta}^2 v_{eff}^2(\hat{\beta}) \int_0^t \int_{\mathbb{R}^2} p_{t-r} \star g(y)^2 dy dr$, which is equal to the variance of $\int_{\mathbb{R}^2} \mathcal{U}(t, x) g(x) dx$. Now, since \mathcal{X}_j^ϵ are independent, to prove the full central limit theorem we need to check Lindeberg's condition

$$\sum_j \mathbf{E}[|\mathcal{X}_j^\epsilon(\hat{\beta})|^2 \mathbf{1}_{\{|\mathcal{X}_j^\epsilon(\hat{\beta})| > \delta\}}] \rightarrow 0, \quad (2.10)$$

for any $\delta > 0$ as $\epsilon \rightarrow 0$. Choose $p > 1$ such that $(2p - 1)\hat{\beta} < \hat{\beta}_c(R)$. From the Holder and the Chebyshev inequality:

$$\sum_j \mathbf{E}[|\mathcal{X}_j^\epsilon(\hat{\beta})|^2 \mathbf{1}_{\{|\mathcal{X}_j^\epsilon(\hat{\beta})| > \delta\}}] \leq \frac{1}{\delta} \sum_j \mathbf{E}[|\mathcal{X}_j^\epsilon(\hat{\beta})|^{2p}]^{1/p} (\mathbf{E}[|\mathcal{X}_j^\epsilon(\hat{\beta})|^2])^{1-1/p}.$$

Recall that $\lambda = 2 - \log \log \epsilon^{-1} / 2 \log \epsilon^{-1}$. From the bound in **Proposition 1.4**, $\mathbf{E}[\mathcal{X}_j^\epsilon(\hat{\beta})^2] \lesssim \epsilon^{2-\lambda} \lesssim 1 / \log \epsilon^{-1}$ for all j . Moreover:

$$\sum_j \mathbf{E}[\|\mathcal{X}_j^\epsilon(\hat{\beta})\|^{2p}]^{1/p} = \sum_j \|\mathcal{X}_j^\epsilon(\hat{\beta})\|_{2p}^2 \lesssim \sum_j \|\mathcal{X}_j^\epsilon((2p-1)\hat{\beta})\|_2^2 \lesssim 1,$$

as $\epsilon \rightarrow 0$, where we used **Lemma 2.1** below and the fact that the sum in the right-most side of the above inequality converges as $\epsilon \rightarrow 0$ when $(2p-1)\hat{\beta} < \hat{\beta}_c(R)$. These two observations prove (2.10), which proves (2.9).

These three items prove the full central limit theorem appearing in **Theorem 1.1**. The only thing left to prove is the following:

Lemma 2.1. *For $p > 2$ such that $(p-1)\hat{\beta} < \hat{\beta}_c(R)$ we have:*

$$\|\mathcal{X}_j^\epsilon(\hat{\beta})\|_p \lesssim \|\mathcal{X}_j^\epsilon((p-1)\hat{\beta})\|_2$$

Proof. As we will see, this is just an instance of hypercontractivity for Wiener chaos. Recall the definition of $\mathcal{X}_j(\hat{\beta})$:

$$\mathcal{X}_j(\hat{\beta}) = \int_{I_{\lambda,j}} \int_{\mathbb{R}^2} \tilde{Z}_t^\epsilon(r, y) d\xi(r, y),$$

where

$$\tilde{Z}_t^\epsilon(r, y) = \int_{\mathbb{R}^2} g(x) \hat{\mathbb{E}}_{B,t} \left[\Phi_{t,x,B}^\epsilon(r, y) \exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} \tilde{M}_{t,x,B}^\epsilon(r) - \frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \langle \tilde{M}_{t,x,B}^\epsilon(r) \rangle \right) \right] dx.$$

Observe that the random variable $\tilde{M}_{t,x,B}^\epsilon$, for a fixed realization of the path B , is an integral of a function that is deterministic with respect to the noise. Hence, it is a mean zero Gaussian random variable, and $\langle \tilde{M}_{t,x,B}^\epsilon \rangle$ is its second moment. Therefore, the exponential in the above equation is a Wick exponential [Jan97]:

$$\tilde{Z}_t^\epsilon(r, y) = \int_{\mathbb{R}^2} g(x) \hat{\mathbb{E}}_{B,t} \left[\Phi_{t,x,B}^\epsilon(r, y) : \exp \left(\frac{\hat{\beta}}{\sqrt{\log \epsilon^{-1}}} \tilde{M}_{t,x,B}^\epsilon(r) \right) : \right] dx,$$

where $: \exp X :$ is the Wick exponential of the random variable X . We use this to find a chaos expansion for $\mathcal{X}_j^\epsilon(\hat{\beta})$. We have that:

$$\begin{aligned} \tilde{Z}_t^\epsilon(r, y) = & \\ \int_{\mathbb{R}^2} g(x) \hat{\mathbb{E}}_{B,t} \left[\Phi_{t,x,B}^\epsilon(r, y) \left(1 + \sum_{k=1}^{\infty} \frac{\hat{\beta}^k}{(\log \epsilon^{-1})^k k!} \int \cdots \int_{\Delta_k \times \mathbb{R}^{2k}} \left[\prod_{j=1}^k \tilde{\Phi}_{t,x,B}^\epsilon(r_i, y_i) \right] \prod_{j=1}^k d\xi(r_i, y_i) \right) \right] dx, \end{aligned}$$

where $\Delta_k = \{-\infty < r_1 < \dots < r_k < r\}$ and $\tilde{\Phi}^\epsilon(s, y) = \int_{\mathbb{R}^2} \int_0^{r_\epsilon} \phi(t/\epsilon^2 - s_1 - s) \psi(x/\epsilon + B_{s_1} - y) ds_1$ with r_ϵ defined after (2.2). Plugging that in the expression for $\mathcal{X}_j^\epsilon(\hat{\beta})$ yields

$$\mathcal{X}_j^\epsilon(\hat{\beta}) = \mathcal{W}_0^\epsilon + \sum_{k=1}^{\infty} \frac{\hat{\beta}^k}{(\log \epsilon^{-1})^k k!} \mathcal{W}_k^\epsilon,$$

where

$$\mathcal{W}_k^\epsilon = \int \cdots \int_{\Delta'_{k+1} \times \mathbb{R}^{2(k+1)}} \left[\int_{\mathbb{R}^2} g(x) \hat{\mathbb{E}}_{B,t} \left[\prod_{j=1}^k \tilde{\Phi}_{t,x,B}^\epsilon(r_j, y_j) \Phi_{t,x,B}^\epsilon(r_{k+1}, y_{k+1}) \right] dx \right] \prod_{j=1}^{k+1} d\xi(r_j, y_j),$$

with $\Delta'_{k+1} = \{-\infty < r_1 < \dots < r_{k+1} < \infty, r_{k+1} \in I_{\lambda,j}\}$. Clearly \mathcal{W}_k^ϵ lies in the $k+1$ homogeneous Wiener chaos (again see [Jan97]) for all $k \in \mathbb{N}_0$. By hypercontractivity for Wiener chaos [Jan97] we get that:

$$\|\mathcal{X}_j^\epsilon(\hat{\beta})\|_p \lesssim \left(\sum_{k=0}^{\infty} \frac{((p-1)\beta)^k}{(\log \epsilon^{-1})^k k!} \|\mathcal{W}_k^\epsilon\|_2^2 \right)^{1/2} = \|\mathcal{X}_j^\epsilon((p-1)\hat{\beta})\|_2.$$

□

3 The Markov Chain on Ω_1

3.1 The construction

As mentioned, to prove **Proposition 1.4** we have to be able to control expressions of the form:

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[f(B)],$$

as $\epsilon \rightarrow 0$. This is done by introducing a Markov chain on Ω_1 . We follow the steps of [GRZ18] to construct this Markov chain and detail its properties.

The goal is to express the measure $\hat{\mathbb{P}}_T^{(\epsilon)}$ as a transition probability of a well-chosen Markov chain. To achieve this, we make the following crucial observation:

Let B be distributed according to the measure $\hat{\mathbb{P}}_T^{(\epsilon)}$ and recall the definition of this measure:

$$\hat{\mathbb{P}}_T^{(\epsilon)}(dB) = \exp\left(\frac{\beta^2}{2 \log \epsilon^{-1}} \int_{[0,T]^2} R(s-u, B(s) - B(u)) ds du - \zeta_T^{(\epsilon)}\right) W_T(dB).$$

Since $R(s, \cdot) = 0$ for $|s| \geq 1$, two points $B(s), B(u)$ interact only when $|s - u| < 1$. This suggests that we can split the path B into segments of length 1 and only neighboring segments will interact with each other. Now, heuristically, this interaction term can be seen as a transition probability. Therefore, if we choose the first segment according to an appropriate distribution, we can build a Markov chain using this transition probability. So we will get a sequence of paths in Ω_1 . Then, by gluing all of these paths together, we will get a path on Ω_T which will be distributed according to $\hat{\mathbb{P}}_T^{(\epsilon)}$.

Strictly speaking, the distribution of this path will not be exactly $\hat{\mathbb{P}}_T^{(\epsilon)}$ since we must account for some edge effects: The endpoint T may not be a natural number and for some technical reasons we may need to choose the first segment to be of length $\tau < 1$. In practice, in all expectations that we will encounter, there will be an extra term accounting for the edge effects. This term can be ignored since in $d \geq 3$ it asymptotically decouples from all relevant random variables we are considering (see **Lemma A.1** in [GRZ18]) and in $d = 2$, as $\epsilon \rightarrow 0$, it goes to 1 uniformly.

Let us be more specific. We consider any probability space $(\mathcal{X}, \mathcal{F}, \mathbb{P})$ that is rich enough to carry the random variables that we are going to encounter and we write \mathbb{E} for the corresponding expectation. As mentioned above we seek to build a continuous path over $[0, T]$ starting from 0 by gluing together paths from Ω_1 . The initial segment will be defined over $[0, \tau]$ where $\tau \in (0, 1]$.

Then the final segment will be defined over $[0, T - \tau - N]$ where $N = [T - \tau]$. Every other segment will be defined over $[0, 1]$.

To formalize this 'gluing' of the paths we set for $k = 1, \dots, N$, $\tau_{k+1} = \tau_k + 1$ with $\tau_0 = 0, \tau_1 = \tau$ and $\tau_{N+2} = T$. Now choose paths $x_0 \in \Omega_\tau$, $x_k \in \Omega_1$ for $k = 1, \dots, N$ and $x_{N+1} \in \Omega_{T-\tau-N}$. We can patch these paths together over the intervals (τ_k, τ_{k+1}) and make a path in Ω_T in the following way:

$$B(s) = \begin{cases} x_0(s) & \text{if } s \in [0, \tau], \\ B(\tau + k - 1) + x_k(s - \tau - k + 1) & \text{if } s \in [\tau + k - 1, \tau + k], k = 1, \dots, N, \\ B(\tau + N) + x_k(s - \tau - N) & \text{if } s \in [\tau + N, T]. \end{cases} \quad (3.1)$$

It is easily checked that $B \in \Omega_T$. Again following [GRZ18] we will write

$$B = [x_0, \dots, x_{N+1}]$$

for a path in Ω_T built from $(x_k)_{k=0, \dots, N+1}$.

With this terminology, we explain how to decompose the measure $\hat{\mathbb{P}}_T$. We write

$$\int_{[0, T]^2} R(s - u, B(s) - B(u)) ds du = \sum_{k, m=1}^{N+1} \int_{[\tau_k, \tau_{k+1}]} \int_{[\tau_m, \tau_{m+1}]} R(s - u, B(s) - B(u)) ds du.$$

Since $R(s, \cdot) = 0$ when $|s| \geq 1$ the integral inside the sum is nonzero only when $|k - m| \leq 1$ and so

$$\begin{aligned} & \int_{[0, T]^2} R(s - u, B(s) - B(u)) ds du = \\ & \sum_{k=0}^{N+1} \int_{[\tau_k, \tau_{k+1}]} \int_{[\tau_k, \tau_{k+1}]} R(s - u, B(s) - B(u)) ds du + 2 \sum_{k=0}^N \int_{[\tau_k, \tau_{k+1}]} \int_{[\tau_{k+1}, \tau_{k+2}]} R(s - u, B(s) - B(u)) ds du. \end{aligned}$$

For $B = [x_0, \dots, x_{N+1}]$ we have

$$B(\tau_k + s) = B(\tau_k) + x_k(s), \quad s \in [0, 1], \quad (3.2)$$

and

$$B(\tau_{k+1} + s) = B(\tau_k) + x_k(1) + x_{k+1}(s), \quad s \in [0, 1]. \quad (3.3)$$

Therefore, from (3.2) and the change of variables $(s, u) \rightarrow (\tau_k + s, \tau_k + u)$

$$\int_{[\tau_k, \tau_{k+1}]} \int_{[\tau_k, \tau_{k+1}]} R(s - u, B(s) - B(u)) ds du = \int_{[0, 1]^2} R(s - u, x_k(s) - x_k(u)) ds du,$$

for $k = 1, \dots, N$. From (3.3) and the change of variables $(s, u) \rightarrow (\tau_{k+1} + s, \tau_k + u)$ we also get

$$\int_{[\tau_{k+1}, \tau_{k+2}]} R(s - u, B(s) - B(u)) ds du = \int_{[0, 1]^2} R(s - u, x_{k+1}(s) + x_k(1) - x_k(u)) ds du,$$

for $k = 1, \dots, N$. Similar identities hold when $k = 0$ and when $k = N + 1$.

For $\beta_\epsilon = \hat{\beta} / \sqrt{\log \epsilon^{-1}}$ the above identities lead us to the following definitions:

$$D_\tau^{(\epsilon)}(x, y) := \beta_\epsilon^2 \int_{[0, \tau]^2} R(s - u, y(s) - x(u)) ds du, \quad x, y \in \Omega_\tau,$$

$$D^{(\epsilon)}(x, y) := \beta_\epsilon^2 \int_{[0,1]^2} R(s-u, y(s) - x(u)) dsdu, \quad x, y \in \Omega_1, \quad (3.4)$$

and

$$I_{0,1}^{(\epsilon)}(x, y) := \beta_\epsilon^2 \int_{[0,\tau]} \int_{[0,1]} R(s-u, y(s) + x(1) - x(u)) dsdu, \quad x \in \Omega_\tau, \quad y \in \Omega_1,$$

$$I_{N,N+1}^{(\epsilon)}(x, y) := \beta_\epsilon^2 \int_{[0,1]} \int_{[0,T-\tau-N]} R(s-u, y(s) + x(1) - x(u)) dsdu, \quad x \in \Omega_1, \quad y \in \Omega_{T-\tau-N}, \quad (3.5)$$

$$I^{(\epsilon)}(x, y) := \beta_\epsilon^2 \int_{[0,1]^2} R(s-u, y(s) + x(1) - x(u)) dsdu, \quad x, y \in \Omega_1. \quad (3.6)$$

The terms $D_\tau^{(\epsilon)}, D^{(\epsilon)}$ capture the self-interactions of each segment and the terms $I_{0,1}^{(\epsilon)}, I^{(\epsilon)}, I_{N,N+1}^{(\epsilon)}$ capture the interactions between neighboring segments. By our previous observations and by writing $B = [x_0, \dots, x_{N+1}]$ we have that

$$\begin{aligned} & \beta_\epsilon^2 \int_{[0,T]^2} R(s-u, B(s) - B(u)) dsdu = \\ & D_\tau^{(\epsilon)}(x_0, x_0) + \sum_{k=1}^{N+1} D^{(\epsilon)}(x_k, x_k) + 2 \left(I_{0,1}^{(\epsilon)}(x_0, x_1) + \sum_{k=1}^{N-1} I^{(\epsilon)}(x_k, x_{k+1}) + I_{N,N+1}^{(\epsilon)}(x_N, x_{N+1}) \right). \end{aligned}$$

It is easily checked that by writing $B = [x_0, \dots, x_{N+1}]$, where x_0, \dots, x_{N+1} are distributed like standard Brownian motions, then B is distributed like a Brownian motion in Ω_T . By plugging the above formula to the definition of $\hat{\mathbb{P}}_T^{(\epsilon)}(dB)$ and using this observation the measure $\hat{\mathbb{P}}_T^{(\epsilon)}(dB)$ is proportional to

$$\hat{\mathbb{P}}_\tau^{(\epsilon)}(dx_0) e^{I_{0,1}^{(\epsilon)}(x_0, x_1)} \prod_{k=1}^{N-1} \hat{\mathbb{P}}_1^{(\epsilon)}(dx_k) e^{I^{(\epsilon)}(x_k, x_{k+1})} \hat{\mathbb{P}}_1^{(\epsilon)}(dx_N) e^{I_{T-\tau-N}^{(\epsilon)}(x_N, x_{N+1})} \hat{\mathbb{P}}_{T-\tau-N}^{(\epsilon)}(dx_{N+1}).$$

We want to interpret this as a transition probability of a Markov chain. For this reason we use the Doob-Krein-Rutman theorem [Sne+12] which implies that we can find $\rho^{(\epsilon)} > 0$ and $\Psi^{(\epsilon)}$ on Ω_1 solving the following eigenvalue problem:

$$\int_{\Omega_1} e^{I^{(\epsilon)}(x,y)} \Psi^{(\epsilon)}(y) \hat{\mathbb{P}}_1^{(\epsilon)}(dy) = \rho^{(\epsilon)} \Psi^{(\epsilon)}(x), \quad (3.7)$$

so that $\rho^{(\epsilon)}$ is the largest possible eigenvalue, $\Psi^{(\epsilon)}$ is bounded above and below by positive constants, is normalized to have total mass 1 with respect to $\hat{\mathbb{P}}_1^{(\epsilon)}$ and is the unique eigenvector associated with $\rho^{(\epsilon)}$. More specifically, we have the bound

$$e^{-\|I^{(\epsilon)}\|_\infty} \leq \rho^{(\epsilon)} \leq e^{\|I^{(\epsilon)}\|_\infty},$$

which we get by integrating the above eigenvalue equation over y . This implies that

$$e^{-2\|I^{(\epsilon)}\|_\infty} \leq \Psi^{(\epsilon)}(x) \leq e^{2\|I^{(\epsilon)}\|_\infty}.$$

Now we can define the transition probabilities

$$\hat{\pi}^{(\epsilon)}(x, dy) := \frac{e^{I^{(\epsilon)}(x,y)} \Psi^{(\epsilon)}(y) \hat{\mathbb{P}}_1^{(\epsilon)}(dy)}{\rho^{(\epsilon)} \Psi^{(\epsilon)}(x)}, \quad (3.8)$$

$$\hat{\pi}_{N,N+1}^{(\epsilon)}(x, dy) := \frac{e^{I_{N,N+1}^{(\epsilon)}(x,y)} \hat{\mathbb{P}}_{T-1-N}^{(\epsilon)}(dy)}{f_{N,N+1}^{(\epsilon)}(x)}, \quad (3.9)$$

$$\hat{\pi}_{0,1}^{(\epsilon)}(x, dy) := \frac{e^{I_{0,1}^{(\epsilon)}(x,y)} \hat{\mathbb{P}}_{\tau}^{(\epsilon)}(dy)}{f_{0,1}^{(\epsilon)}(x)}, \quad (3.10)$$

where $f_{N,N+1}^{(\epsilon)}(x), f_{0,1}^{(\epsilon)}(x)$ are the normalization constants. With this notation the measure $\hat{\mathbb{P}}_T^{(\epsilon)}$ is proportional to

$$f_{0,1}^{(\epsilon)}(x_0) \hat{\mathbb{P}}_{\tau}^{(\epsilon)}(dx_0) \hat{\pi}_{0,1}^{(\epsilon)}(x_0, dx_1) \prod_{k=1}^{N-1} \hat{\pi}^{(\epsilon)}(x_k, dx_{k+1}) \hat{\pi}_{N,N+1}^{(\epsilon)}(x_N, dx_{N+1}) \frac{f_{N,N+1}^{(\epsilon)}(x_N)}{\Psi^{(\epsilon)}(x_N)}.$$

The Markov chain we are looking for is built from the probability kernels that appear above. In particular, we sample:

- $X_0 \in \Omega_{\tau}$ according to the distribution $(\mathcal{Z}_{\tau}^{(\epsilon)})^{-1} f_{0,1}^{(\epsilon)}(X_0) \hat{\mathbb{P}}_{\tau}^{(\epsilon)}(dX_0)$, where $\mathcal{Z}_{\tau}^{(\epsilon)} = \int_{\Omega_{\tau}} f_{0,1}^{(\epsilon)}(x) \hat{\mathbb{P}}_{\tau}^{(\epsilon)}(dx)$.
- (X_1, \dots, X_N) according to $\hat{\pi}_{0,1}^{(\epsilon)}(X_0, dX_1) \left(\prod_{k=1}^{N-1} \hat{\pi}^{(\epsilon)}(X_k, dX_{k+1}) \right) \hat{\pi}_{N,N+1}^{(\epsilon)}(X_N, dX_{N+1})$.

and once we sample our points we can build a path $B = [X_0, \dots, X_{N+1}]$ on Ω_T by gluing these paths together according to (3.1). Then for any measurable function $F : \Omega_T \rightarrow \mathbb{R}$,

$$\hat{\mathbb{E}}_{B,T}[F(B)] = \mathbb{E} \left[F([X_0, \dots, X_{N+1}]) c_{\tau,T}^{(\epsilon)} \frac{f_{N,N+1}^{(\epsilon)}(X_N)}{\Psi^{(\epsilon)}(X_N)} \right],$$

where $c_{\tau,T}^{(\epsilon)}$ is the appropriate normalization constant, determined by the equation

$$\frac{1}{c_{\tau,T}^{(\epsilon)}} = \mathbb{E} \left[\frac{f_{N,N+1}^{(\epsilon)}(X_N)}{\Psi^{(\epsilon)}(X_N)} \right]. \quad (3.11)$$

The term

$$c_{\tau,T}^{(\epsilon)} \frac{f_{N,N+1}^{(\epsilon)}(X_N)}{\Psi^{(\epsilon)}(X_N)}$$

is the term accounting for the edge effects born from splitting the interval $[0, T]$ into an interval of length τ and then intervals of length 1. As we will see, this term does not play a major role in our calculations.

We will mainly have to deal with expressions of the form

$$\hat{\mathbb{E}}_{B,T}[F(B^1, B^2)],$$

where B^1, B^2 are two independent paths sampled according to (1.10). By following the same construction as above we can convert this expectation into the following expectation:

$$\mathbb{E} \left[F([X_0, \dots, X_{N+1}], [Y_0, \dots, Y_{N+1}]) c_{\tau,T}^2 \frac{f_{N,N+1}^{(\epsilon)}(X_N)}{\Psi^{(\epsilon)}(X_N)} \frac{f_{N,N+1}^{(\epsilon)}(Y_N)}{\Psi^{(\epsilon)}(Y_N)} \right],$$

where $[X_0, \dots, X_{N+1}]$ and $[Y_0, \dots, Y_{N+1}]$ are paths built from two independent trajectories $(X_i)_{i=1, \dots, N+1}$, $(Y_i)_{i=1, \dots, N+1}$ respectively, sampled as described above. Let

$$\mathcal{G}_\epsilon(x, y) := c_{\tau, T}^2 \frac{f_{N, N+1}^{(\epsilon)}(x)}{\Psi^{(\epsilon)}(x)} \frac{f_{N, N+1}^{(\epsilon)}(y)}{\Psi^{(\epsilon)}(y)} \quad (3.12)$$

be the term accounting for the 'edge effects'. Then we have the identity:

$$\hat{\mathbb{E}}_{B, T}[F(B^1, B^2)] = \mathbb{E} \left[F([X_0, \dots, X_{N+1}], [Y_0, \dots, Y_{N+1}]) \mathcal{G}_\epsilon(X_N, Y_N) \right]. \quad (3.13)$$

3.2 The coupling in $d = 2$

In the remainder of the paper the Markov chain on $\Omega_1 \times \Omega_1$ with transition probability kernel

$$\hat{\pi}^{(\epsilon)} := \hat{\pi}^{(\epsilon)} \times \hat{\pi}^{(\epsilon)}$$

plays an important role. We dedicate the next few sections to its properties. First, we will introduce a coupling of this Markov chain to a sequence of Bernoulli random variables based on Doeblin's inequality.

Recall the definition of the transition measure

$$\hat{\pi}^{(\epsilon)}(x, dy) = \frac{e^{I^{(\epsilon)}(x, y)} \Psi^{(\epsilon)}(y) \hat{\mathbb{P}}_1^{(\epsilon)}(dy)}{\rho^{(\epsilon)} \Psi^{(\epsilon)}(x)},$$

and the definition of $\hat{\mathbb{P}}_1^{(\epsilon)}$

$$\hat{\mathbb{P}}_1^{(\epsilon)}(dx) = \exp \left(\frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \int_{[0, 1]^2} R(s - u, x(s) - x(u)) ds du - \zeta_1^{(\epsilon)} \right) W_1(dx).$$

Observe that, as $\epsilon \rightarrow 0$, the measure $\hat{\mathbb{P}}_1^{(\epsilon)}$ converges to W_1 in total variation. That is because the function R is compactly supported and we can use the dominated convergence theorem.

Moreover, recall the definition of $I^{(\epsilon)}(x, y)$ and of the associated eigenvalue problem defining $\Psi^{(\epsilon)}, \rho^{(\epsilon)}$:

$$I^{(\epsilon)}(x, y) = \beta_\epsilon^2 \int_{[0, 1]^2} R(s - u, x(1) + y(s) - x(u)) ds du,$$

$$\int_{\Omega_1} e^{I^{(\epsilon)}(x, y)} \Psi^{(\epsilon)}(y) \hat{\mathbb{P}}_1^{(\epsilon)}(dy) = \rho^{(\epsilon)} \Psi^{(\epsilon)}(x),$$

where we also recall that $\int_{\Omega_1} \Psi^{(\epsilon)}(y) \hat{\mathbb{P}}_1^{(\epsilon)}(dy) = 1$. We observe that as $\epsilon \rightarrow 0$, $I^{(\epsilon)}(x, y) \rightarrow 0$ uniformly over x, y , since R is bounded. This implies that $\rho^{(\epsilon)} \rightarrow 1$ since

$$e^{-\|I^{(\epsilon)}\|_\infty} \leq \rho^{(\epsilon)} \leq e^{\|I^{(\epsilon)}\|_\infty}.$$

This in turn implies that $\Psi^{(\epsilon)}(x) \rightarrow 1$ uniformly in x since we have

$$e^{-2\|I^{(\epsilon)}\|_\infty} \leq \Psi^{(\epsilon)}(x) \leq e^{2\|I^{(\epsilon)}\|_\infty},$$

for all x . These observations imply that:

$$\frac{e^{I^{(\epsilon)}(x, y)} \Psi^{(\epsilon)}(y)}{\rho^{(\epsilon)} \Psi^{(\epsilon)}(x)} \rightarrow 1,$$

uniformly over x, y . This proves the following proposition:

Proposition 3.1. *The transition probability kernel $\hat{\pi}^{(\epsilon)}(x, dy)$ converges in total variation to $W_1(dy)$ uniformly over x . In other words, we have:*

$$\sup_{x \in \Omega_1} d_{TV}(\hat{\pi}^{(\epsilon)}(x, dy), W_1(dy)) \rightarrow 0,$$

as $\epsilon \rightarrow 0$, where d_{TV} is the total variation distance between two probability measures. Furthermore for any $0 < \gamma < 1$, there is an $\epsilon_0 = \epsilon_0(\gamma)$ such that for all $x \in \Omega_1$ and A a Borel set in Ω_1

$$\hat{\pi}^{(\epsilon)}(x, A) \geq \gamma W_1(A), \quad (3.14)$$

for all $\epsilon < \epsilon_0$.

Remark 3.1. *Let $\tau_\epsilon \in (0, 1)$, $T = T(\epsilon) \rightarrow \infty$, as $\epsilon \rightarrow 0$ and $N_\epsilon = [T - \tau]$. Similar observations as the ones above prove that \mathcal{G}_ϵ , defined in (3.12), converges to 1 uniformly as $\epsilon \rightarrow 0$. Indeed, we see that $I_{N, N+1}^{(\epsilon)} \rightarrow 0$ uniformly as $\epsilon \rightarrow 0$, where $I_{N, N+1}^{(\epsilon)}$ is defined in (3.5). By looking at (3.9) we see that $f_{N, N+1}^{(\epsilon)} \rightarrow 1$ uniformly as $\epsilon \rightarrow 0$. This, along with our previous observations prove that $c_{\tau, T}^{(\epsilon)} \rightarrow 1$ as $\epsilon \rightarrow 0$. These observations imply that $\mathcal{G}_\epsilon \rightarrow 1$, uniformly as $\epsilon \rightarrow 0$.*

Since $\hat{\pi}^{(\epsilon)} = \hat{\pi}^{(\epsilon)} \times \hat{\pi}^{(\epsilon)}$, this proposition implies that for any $\gamma \in (0, 1)$, there is an ϵ_0 small enough such that for all Borel sets $A \subset \Omega_1^2$, for all $z \in \Omega_1 \times \Omega_1$ and all $\epsilon < \epsilon_0$

$$\hat{\pi}^{(\epsilon)}(z, A) \geq \gamma(W_1 \times W_1)(A). \quad (3.15)$$

We can choose γ to be such that both (3.14) and (3.15) are true. From here on we will fix γ and ϵ_0 and we will always assume that $\epsilon < \epsilon_0$.

This last observation gives us a Doeblin condition for the Markov chain. Therefore, we can introduce a coupling of the Markov chain with a sequence of i.i.d. Bernoulli random variables $(\eta_j)_{j \in \mathbb{N}}$. This coupling goes as follows:

We can write

$$\hat{\pi}^{(\epsilon)}(z, dy) = \gamma(W_1 \times W_1)(dy) + (1 - \gamma) \frac{\hat{\pi}^{(\epsilon)}(z, dy) - \gamma(W_1 \times W_1)(dy)}{1 - \gamma}.$$

So at each step j , if $\eta_j = 1$ we draw our next step from the measure $(W_1 \times W_1)(dy)$, independently from all the previous steps, and if not we draw (X_j, Y_j) from $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1 \times W_1}{1 - \gamma}$. This gives us a regenerative structure of the Markov chain since in every 'success' we forget what happened in the previous steps and we begin the chain anew. In particular, this structure decomposes a 'trajectory' of the Markov chain into independent cycles.

Now let $((X_j, Y_j))_{j \geq 1} \subseteq \Omega_1 \times \Omega_1$ be a trajectory of $\hat{\pi}^{(\epsilon)}$ with an initial segment $(X_0, Y_0) \sim W_1 \times W_1$. We consider the pair of paths $(\omega_{X_0}, \omega_{Y_0})$ in $C([0, \infty))^2$, each constructed by gluing $(X_j)_{j \geq 0}$, $(Y_j)_{j \geq 0}$ respectively, according to (3.1). We define the notion of the total path increment between two regeneration times. First, define the associated regeneration times:

$$T_0 := 0, T_i := \inf\{j > T_{i-1} : \eta_j = 1\}. \quad (3.16)$$

The times between two regenerations, $T_{k+1} - T_k$, $k = 0, 1, \dots$ are i.i.d. and distributed as geometric random variables with parameter γ . The corresponding total path increment for ω_{X_0} and ω_{Y_0} between two regenerations is defined by the equations:

$$\mathbf{X}_j^{(\epsilon)} := \sum_{k=T_j}^{T_{j+1}-1} X_k(1) \text{ and } \mathbf{Y}_j^{(\epsilon)} := \sum_{k=T_j}^{T_{j+1}-1} Y_k(1). \quad (3.17)$$

By construction, the random variables $((\mathbf{X}_j^{(\epsilon)}, \mathbf{Y}_j^{(\epsilon)})_{j \geq 0}$ are i.i.d.. Also, from (3.1)

$$\omega_{X_0}(T_j) = \sum_{k=0}^{j-1} \mathbf{X}_k^{(\epsilon)}, \quad \omega_{Y_0}(T_j) = \sum_{k=0}^{j-1} \mathbf{Y}_k^{(\epsilon)}, \quad (3.18)$$

and

$$\mathcal{C}_j := \left(T_{j+1} - T_j, (\omega_{X_0}(T_j + s) - \omega_{X_0}(T_j), \omega_{Y_0}(T_j + s) - \omega_{Y_0}(T_j))_{s \leq T_{j+1} - T_j} \right) \quad (3.19)$$

is equal in distribution to

$$\left(\theta, (\tilde{\omega}_{B^1}(s), \tilde{\omega}_{B^2}(s))_{s \leq \theta} \right), \quad (3.20)$$

where:

- $\theta \sim \text{Geo}(\gamma)$.
- $(B^1, B^2) \sim W_1 \times W_1$ independent from θ .
- $(\tilde{\omega}_{B^1}, \tilde{\omega}_{B^2})$ is a pair of paths, each built according to (3.1) using segments sampled from $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1 \times W_1}{1 - \gamma}$ with (B^1, B^2) as initial steps. Also, $(\tilde{\omega}_{B^1}, \tilde{\omega}_{B^2})$ is independent from θ .

All of these random variables mentioned above are also independent from $(\omega_{X_0}(s), \omega_{Y_0}(s))_{s \leq T_j}$.

The random variables \mathcal{C}_j are called cycles. From our previous observations the collection of all cycles $(\mathcal{C}_j)_{j \geq 1}$ is a collection of i.i.d. random variables with (3.20) as the common underlying distribution. This proves that the process $(\omega_{X_0}(s), \omega_{Y_0}(s))_{s \geq 0}$ can be decomposed into i.i.d. cycles. In full formality, one can prove that the process $(\omega_{X_0}(s), \omega_{Y_0}(s))_{s \geq 0}$ is a regenerative process [Asm03; Tho00].

3.3 The Mixing property

Here we detail the mixing mechanism of the Markov chain and some of its basic consequences. This is based on the observation that there are constants A_ϵ, B_ϵ such that they converge to 1 as $\epsilon \rightarrow 0$ and such that for all Borel sets of Ω_1 and $x \in \Omega_1$,

$$A_\epsilon W_1(A) \leq \hat{\pi}^{(\epsilon)}(x, A) \leq B_\epsilon W_1(A). \quad (3.21)$$

Indeed, recall that from (3.8), $\hat{\pi}^{(\epsilon)}$ is absolutely continuous with respect to the Wiener measure

$$\hat{\pi}^{(\epsilon)}(x, dy) = \frac{e^{I^{(\epsilon)}(x,y)} \Psi^{(\epsilon)}(y)}{\rho^{(\epsilon)} \Psi^{(\epsilon)}(x)} \cdot e^{D_1^{(\epsilon)}(y,y)} W_1(dy),$$

with $D^{(\epsilon)}$ as in (3.4) and $I^{(\epsilon)}$ as in (3.6). Since

$$e^{-\|I^{(\epsilon)}\|_\infty} \leq \rho^{(\epsilon)} \leq e^{\|I^{(\epsilon)}\|_\infty},$$

and

$$e^{-2\|I^{(\epsilon)}\|_\infty} \leq \Psi^{(\epsilon)}(x) \leq e^{2\|I^{(\epsilon)}\|_\infty},$$

we see that there is a constant $C > 0$ such that

$$e^{-C/\log \epsilon^{-1}} W_1(A) \leq \hat{\pi}^{(\epsilon)}(x, A) \leq e^{C/\log \epsilon^{-1}} W_1(A),$$

for all Borel sets $A \subseteq \Omega_1$ and $x \in \Omega_1$.

A similar inequality is true for $\hat{\pi}^{(\epsilon)}$ and $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1 \times W_1}{1 - \gamma}$ (only with different constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$) since both measures are absolutely continuous with respect to the Wiener measure, with density converging to 1 uniformly as $\epsilon \rightarrow 0$. With this observation, we can prove the following proposition:

Lemma 3.1. *Let F be a nonnegative integrable function on $(\Omega_1 \times \Omega_1)^{p+1}$ and let $((X_0, Y_0), \dots, (X_p, Y_p))$ be a sequence of random variables generated by $\hat{\pi}^{(\epsilon)}$ or $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1 \times W_1}{1 - \gamma}$ with $(X_0, Y_0) \sim W_1 \times W_1$. Then there are constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$ such that they converge to 1 as $\epsilon \rightarrow 0$ and such that for any $1 \leq k \leq m$*

$$\begin{aligned} & \mathcal{A}_\epsilon \mathbb{E} \left[F((X_0, Y_0), \dots, (\tilde{X}_k, \tilde{Y}_k), \dots, (X_p, Y_p)) \right] \\ & \leq \mathbb{E} \left[F((X_0, Y_0), \dots, (X_p, Y_p)) \right] \leq \\ & \mathcal{B}_\epsilon \mathbb{E} \left[F((X_0, Y_0), \dots, (\tilde{X}_k, \tilde{Y}_k), \dots, (X_p, Y_p)) \right], \end{aligned} \quad (3.22)$$

where $(X_0, Y_0), \dots, (\tilde{X}_k, \tilde{Y}_k), \dots, (X_p, Y_p)$ is generated as before only at step k we sample, $(\tilde{X}_k, \tilde{Y}_k) \sim W_1 \times W_1$ independently from all previous steps.

Proof. First, assume that F is bounded. We write $Z_k = (X_k, Y_k)$ for $k \geq 0$. We prove the proposition in the case where $(Z_k)_k$ is generated using $\hat{\pi}^{(\epsilon)}$. We prove this by writing the expectation of F under the Markov chain explicitly:

$$\mathbb{E} \left[F(Z_0, \dots, Z_k) \right] = \int_{\Omega_1} W_1 \times W_1(dz_0) \dots \int_{\Omega_1} \hat{\pi}^{(\epsilon)}(z_{k-1}, dz_k) \dots \int_{\Omega_1} \hat{\pi}^{(\epsilon)}(z_{m-1}, dz_m) F(z_0, \dots, z_m).$$

We aim to replace $\hat{\pi}^{(\epsilon)}(z_{k-1}, dz_k)$ by $W_1(dz_k)$: From (3.21) there are constants \mathcal{A}'_ϵ and \mathcal{B}'_ϵ converging to 1 as $\epsilon \rightarrow 0$, such that

$$\mathcal{A}'_\epsilon W_1(A) \leq \hat{\pi}^{(\epsilon)}(x, A) \leq \mathcal{B}'_\epsilon W_1(A),$$

for all Borel sets $A \subseteq \Omega_1 \times \Omega_1$. This inequality extends to all simple positive simple functions. In particular, there are constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$ converging to 1 as $\epsilon \rightarrow 0$ such that

$$\mathcal{A}_\epsilon W_1 \times W_1(f) \leq \hat{\pi}^{(\epsilon)}(x, f) \leq \mathcal{B}_\epsilon W_1 \times W_1(f), \quad (3.23)$$

where, for a measure μ and a function f , $\mu(f)$ is the integral of f with respect to μ . By the definition of the Lebesgue integral, (3.23) extends to all positive measurable functions f .

Therefore, from (3.23) we get the inequality:

$$\mathbb{E} \left[F(Z_0, \dots, Z_m) \right] \leq \mathcal{B}_\epsilon \int_{\Omega_1} W_1 \times W_1(dz_0) \dots \int_{\Omega_1} W_1(dz_k) \dots \int_{\Omega_1} \hat{\pi}^{(\epsilon)}(z_{m-1}, dz_m) F(z_0, \dots, z_m),$$

which is the right-hand side of (3.22). Similarly, we prove the left-hand side as well. When $(Z_k)_{k \in \mathbb{N}_0}$ is generated by $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1 \times W_1}{1 - \gamma}$, (3.22) is proved in the same way.

Once we have (3.22) for all bounded measurable positive F we may argue via a cutoff argument and monotone convergence to get the full result. \square

So when we have a positive functional of the Markov chain and we want to calculate the limit $\epsilon \rightarrow 0$ of its expectation, we may replace a finite number of steps by the standard Brownian motions that are independent of all previous steps. As $\epsilon \rightarrow 0$ this will not change the limiting mean since both upper and lower bounds will match.

Remark 3.2. • Using this result we can prove the same double inequality (with different constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$, still converging to 1, as $\epsilon \rightarrow 0$) for a general $F \in L^1$ by splitting F into its positive and negative parts.

- This lemma can also be applied in the case where F is a function of a finite number of independent trajectories of $\hat{\pi}^{(\epsilon)}$ (or trajectories generated by $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1 \times W_1}{1 - \gamma}$ with $(X_0, Y_0) \sim W_1 \times W_1$) and we want to replace parts of these trajectories by Brownian motions. In this case we will get bounds similar to (3.22) (again with a different set of constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$, converging to 1, as $\epsilon \rightarrow 0$).
- Finally, this lemma generalizes in the case where we consider the edge effects and we sample the first and the final steps of the Markov chain using $\hat{\pi}_{0,1} \times \hat{\pi}_{0,1}$ and $\hat{\pi}_{p-1,p} \times \hat{\pi}_{p-1,p}$ respectively. Indeed, it is easy to see that $\hat{\pi}_{0,1}$ and $\hat{\pi}_{p-1,p}$ satisfy a version of (3.21) which allow us to replace (X_1, Y_1) or (X_p, Y_p) by Brownian motions and get the bounds presented above. If we want to replace any other step the proof of **Lemma 3.1** works. A similar argument works in the case where we start the Markov chain by $\hat{\mathbb{P}}_\tau^{(\epsilon)}(dx_0)$ for some $\tau < 1$ or the (normalized) measure $f_{0,1}^{(\epsilon)}(x_0)\hat{\mathbb{P}}_\tau^{(\epsilon)}(dx_0)$ and we want to replace the starting point by a Brownian motion.

By conditioning on the regeneration length $T_{k+1} - T_k$, **Lemma 3.1** has the following consequence for functionals of the total path increments:

Corollary 3.1. There are constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$, converging to 1, as $\epsilon \rightarrow 0$, such that for any non-negative $F \in L^1(\mathbb{R}^{2p})$:

$$\begin{aligned} & \sum_{N_1=1, \dots, N_p=1}^{\infty} \left(\prod_{i=1}^p \mathcal{A}_\epsilon^{N_i} \gamma (1 - \gamma)^{N_i - 1} \right) \mathbb{E} \left[F((\mathbf{X}^{(N_1)}, \mathbf{Y}^{(N_1)}), \dots, (\mathbf{X}^{(N_p)}, \mathbf{Y}^{(N_p)})) \right] \\ & \leq \mathbb{E} \left[F((\mathbf{X}_1^{(\epsilon)}, \mathbf{Y}_1^{(\epsilon)}), \dots, (\mathbf{X}_p^{(\epsilon)}, \mathbf{Y}_p^{(\epsilon)})) \right] \leq \\ & \sum_{N_1=1, \dots, N_p=1}^{\infty} \left(\prod_{i=1}^p \mathcal{B}_\epsilon^{N_i} \gamma (1 - \gamma)^{N_i - 1} \right) \mathbb{E} \left[F((\mathbf{X}^{(N_1)}, \mathbf{Y}^{(N_1)}), \dots, (\mathbf{X}^{(N_p)}, \mathbf{Y}^{(N_p)})) \right], \end{aligned} \quad (3.24)$$

where $(\mathbf{X}^{N_i})_{i=1, \dots, p}, (\mathbf{Y}^{N_i})_{i=1, \dots, p}$ are two independent sets of independent mean zero Gaussian random variables with covariance matrix $N_i I_{2 \times 2}$ respectively.

Proof. We prove this for $p = 1$, with the general proof following the same lines, since $((\mathbf{X}_i^{(\epsilon)}, \mathbf{Y}_i^{(\epsilon)}))_{i \in \mathbb{N}}$ are i.i.d.. Recall the definition of $(\mathbf{X}_i^{(\epsilon)}, \mathbf{Y}_i^{(\epsilon)})$ from (3.17). By construction $(\mathbf{X}_i^{(\epsilon)}, \mathbf{Y}_i^{(\epsilon)})$ is equal in distribution to

$$\left(\sum_{k=0}^{\theta-1} X_k(1), \sum_{k=0}^{\theta-1} Y_k(1) \right),$$

where:

- $(X_0, Y_0) \sim W_1 \times W_1(dx_0, dy_0)$ and $(X_{k+1}, Y_{k+1}) \sim \frac{\hat{\pi}^{(\epsilon)}((X_k, Y_k), dx_{k+1}, dy_{k+1}) - \gamma W_1 \times W_1(dx_{k+1}, dy_{k+1})}{1 - \gamma}$ for $k \in \mathbb{N}$.
- $\theta \sim Geo(\gamma)$, independent from $(X_k, Y_k)_{k \in \mathbb{N}_0}$.

By conditioning on θ , we have

$$\mathbb{E}[F(\mathbf{X}_1^{(\epsilon)}, \mathbf{Y}_1^{(\epsilon)})] = \sum_{N=1} \gamma(1-\gamma)^{N-1} \mathbb{E} \left[F \left(\sum_{k=0}^{N-1} X_k(1), \sum_{k=0}^{N-1} Y_k(1) \right) \right].$$

From (3.22) and Remark 3.2, applied to the function $\tilde{F} : (\Omega_1 \times \Omega_1)^N \rightarrow \mathbb{R}$,

$$\tilde{F}((x_1, y_1), \dots, (x_N, y_N)) = F \left(\sum_{k=0}^{N-1} x_k(1), \sum_{k=0}^{N-1} y_k(1) \right),$$

we have the upper bound

$$\mathbb{E}[F(\mathbf{X}_1^{(\epsilon)}, \mathbf{Y}_1^{(\epsilon)})] \leq \sum_{N=1} \gamma(1-\gamma)^{N-1} \mathcal{B}_\epsilon^N \mathbb{E} \left[F \left(\sum_{j=0}^{N-1} B_j^1(1), \sum_{j=0}^{N-1} B_j^2(1) \right) \right],$$

where (B_j^1, B_j^2) are pairs of i.i.d. standard Brownian motions, and $\mathcal{B}_\epsilon \rightarrow 1$ as $\epsilon \rightarrow 0$. This proves the right-hand side of (3.24). Similarly, we prove the left-hand side of (3.24) and this concludes the proof. \square

3.4 Intersections of independent paths

The goal of this section is to calculate the limiting distribution for the random variable:

$$\frac{1}{\log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^2} \int_0^{M(\epsilon)/\epsilon^2} R(s-u, \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du, \quad (3.25)$$

where R is the correlation function of the space-time mollified white noise, $\omega_{X_0}, \omega_{Y_0}$ are two paths built from $\hat{\pi}^{(\epsilon)}$, X_0, Y_0 are two independent Brownian motions in Ω_1 and $M(\epsilon)$ is such that $\log M(\epsilon)/\log \epsilon^{-1} \rightarrow 0$, as $\epsilon \rightarrow 0$. This will be done by calculating all moments of the above functional, which is done in the following theorem:

Theorem 3.1. *Let $X_0, Y_0 \in \Omega_1$ and $M(\epsilon)$ be as above and let $x \in \mathbb{R}^2$ be fixed. Then, as $\epsilon \rightarrow 0$:*

$$\mathbb{E} \left[\left(\frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^2} \int_0^{M(\epsilon)/\epsilon^2} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du \right)^p \right] \rightarrow m_p,$$

where $m_p = p!$ is the p th moment of an exponential random variable with rate 1. In other words the random variable:

$$\frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^2} \int_0^{M(\epsilon)/\epsilon^2} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du,$$

as $\epsilon \rightarrow 0$ converges to an exponential random variable with rate 1.

Proof. We split the integral in (3.25) over regeneration intervals. We define

$$N_\epsilon := \max\{i / T_i < M(\epsilon)/\epsilon^2\}.$$

Since R is bounded and $R(s, \cdot) = 0$ for $|s| \geq 1$ and since $T_{N_\epsilon+1} - T_{N_\epsilon} \sim \text{Geo}(\gamma)$

$$\frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^2} \int_0^{M(\epsilon)/\epsilon^2} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du =$$

$$\frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \sum_{k,m=0}^{N_\epsilon-1} \int_{T_k}^{T_{k+1}} \int_{T_m}^{T_{m+1}} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du + O\left(\frac{1}{\log \epsilon^{-1}}\right),$$

where the (random) error term goes to 0 almost surely and in L^p , for every $p \in \mathbb{N}$. In particular, as $\epsilon \rightarrow 0$, (3.25) and

$$\frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \sum_{k,m=0}^{N_\epsilon-1} \int_{T_k}^{T_{k+1}} \int_{T_m}^{T_{m+1}} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du \quad (3.26)$$

have the same limiting moments.

Therefore, we focus on (3.26). Observe that $T_i - T_{i-1} \sim \text{Geo}(\gamma)$ and are independent, which implies that $T_{\lfloor M_2(\epsilon)\epsilon^{-2} \rfloor} / M(\epsilon)\epsilon^{-2} \rightarrow 1/\gamma$ a.s.. This implies that $N_\epsilon / M_2(\epsilon)\epsilon^{-2} \rightarrow \gamma$ in probability. We define:

$$\mathcal{M}(n) := \frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \sum_{k,m=0}^{n-1} \int_{T_k}^{T_{k+1}} \int_{T_m}^{T_{m+1}} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du,$$

and

$$A_{\epsilon,\delta} = \{|N_\epsilon / (M(\epsilon)\epsilon^{-2}) - \gamma| \geq \delta\}.$$

From **Lemma 3.2**, proved below, we have $\mathbb{E}[\mathcal{M}(M(\epsilon)\epsilon^{-2})^p] \rightarrow p!$, as $\epsilon \rightarrow 0$, for any $M(\epsilon)$ that satisfies the assumption of the theorem. We have:

$$\mathbb{E}[\mathcal{M}(N_\epsilon)^p] \geq \mathbb{E}[\mathbf{1}_{A_{\epsilon,\delta}^c} \mathcal{M}(N_\epsilon)^p].$$

But on $A_{\epsilon,\delta}^c$ we have $N_\epsilon \geq (\gamma - \delta)(M(\epsilon)\epsilon^{-2})$ and therefore:

$$\mathbb{E}[\mathcal{M}(N_\epsilon)^p] \geq \mathbb{E}[\mathbf{1}_{A_{\epsilon,\delta}^c} \mathcal{M}((\gamma - \delta)M(\epsilon)\epsilon^{-2})^p] = \mathbb{E}[(\mathbf{1}_{A_{\epsilon,\delta}^c} - 1)\mathcal{M}((\gamma - \delta)M(\epsilon)\epsilon^{-2})^p] + \mathbb{E}[\mathcal{M}((\gamma - \delta)M(\epsilon)\epsilon^{-2})^p].$$

Again from **Lemma 3.2**, the second term of the right-hand side in the above inequality converges to $p!$, while the first term is estimated above by:

$$\mathbb{E}[(\mathbf{1}_{A_{\epsilon,\delta}^c} - 1)^2]^{1/2} \mathbb{E}[\mathcal{M}((\gamma - \delta)M(\epsilon)\epsilon^{-2})^{2p}]^{1/2},$$

which goes to 0 as $\epsilon \rightarrow 0$, since $\mathbb{P}(A_{\epsilon,\delta}^c) \rightarrow 0$ and $\mathbb{E}[\mathcal{M}((\gamma - \delta)M(\epsilon)\epsilon^{-2})^{2p}] \rightarrow (2p)!$, as $\epsilon \rightarrow 0$. Therefore

$$\liminf_{\epsilon \rightarrow 0} \mathbb{E}[\mathcal{M}(N_\epsilon)^p] \geq p!.$$

For the upper bound, we argue similarly:

$$\mathbb{E}[\mathcal{M}(N_\epsilon)^p] = \mathbb{E}[\mathbf{1}_{A_{\epsilon,\delta}^c} \mathcal{M}(N_\epsilon)^p] + \mathbb{E}[\mathbf{1}_{A_{\epsilon,\delta}} \mathcal{M}(N_\epsilon)^p].$$

The first term is bounded by above by:

$$\mathbb{E}[\mathcal{M}((\gamma + \delta)M(\epsilon)\epsilon^{-2})^p].$$

which goes to $p!$ as $\epsilon \rightarrow 0$. For the second term observe that $N_\epsilon \leq 4 \cdot M(\epsilon)\epsilon^{-2}$, by definition. Therefore, the second term is bounded above by

$$\mathbb{E}[\mathbf{1}_{A_{\epsilon,\delta}} \mathcal{M}(4 \cdot M(\epsilon)\epsilon^{-2})^p] \leq \mathbb{P}[A_{\epsilon,\delta}]^{1/2} \mathbb{E}[\mathcal{M}(4 \cdot M(\epsilon)\epsilon^{-2})^{2p}]^{1/2} \rightarrow 0,$$

as $\epsilon \rightarrow 0$. Hence

$$\mathbb{E}[\mathcal{M}(N_\epsilon)^p] \rightarrow p!,$$

as $\epsilon \rightarrow 0$. □

Lemma 3.2. *Let $M(\epsilon)$ be such that $\log M(\epsilon)/\log \epsilon^{-1} \rightarrow 0$. Then as $\epsilon \rightarrow 0$ the p th moment of:*

$$\frac{1}{\log \epsilon^{-1}} \sum_{k,m=0}^{M(\epsilon)\epsilon^{-2}} \int_{T_k}^{T_{k+1}} \int_{T_m}^{T_{m+1}} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du \quad (3.27)$$

converges to $(\|R\|_1/2\pi)^p p!$.

Proof. Since $R(s, \cdot) = 0$ for $|s| \geq 1$, the p th moment of (3.27) is given by:

$$\begin{aligned} \mathbb{E} \left[\left(\frac{1}{\log \epsilon^{-1}} \sum_{k=1}^{M(\epsilon)\epsilon^{-2}} d_{k,k-1} + d_{k,k} + d_{k,k+1} \right)^p \right] &= p! \sum_{1 \leq k_1 < k_2 < \dots < k_p \leq M(\epsilon)\epsilon^{-2}} \mathbb{E} \left[\prod_{i=1}^p \left(d_{k_i, k_i-1} + d_{k_i, k_i} + d_{k_i, k_i+1} \right) \right] = \\ &= p! \sum_{\delta \in \{-1, 0, +1\}^p} \sum_{1 \leq k_1 < k_2 < \dots < k_p \leq M(\epsilon)\epsilon^{-2}} \mathbb{E} \left[\prod_{i=1}^p d_{k_i, k_i + \delta_i} \right], \end{aligned} \quad (3.28)$$

where for $\delta \in \{-1, 0, 1\}$:

$$d_{k, k+\delta} := \frac{1}{\log \epsilon^{-1}} \int_{T_k}^{T_{k+1}} \int_{T_{k+\delta}}^{T_{k+\delta+1}} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du. \quad (3.29)$$

By the change of variables $(s, u) \rightarrow (s + T_{k_i + \delta_i}, u + T_{k_i})$ and by adding and subtracting the terms $\omega_{X_0}(T_{k_i + \delta_i})$ and $\omega_{Y_0}(T_{k_i})$, $d_{k_i, k_i + \delta_i}$ is equal to:

$$\begin{aligned} \int_0^{T_{k_i+1} - T_{k_i}} \int_0^{T_{k_i + \delta_i + 1} - T_{k_i + \delta_i}} R \left(s - u - (T_{k_i} - T_{k_i + \delta_i}), x + \omega_{X_0}(T_{k_i + \delta_i}) - \omega_{Y_0}(T_{k_i}) + \right. \\ \left. \omega_{X_0}(s + T_{k_i + \delta_i}) - \omega_{X_0}(T_{k_i + \delta_i}) - (\omega_{Y_0}(u + T_{k_i}) - \omega_{Y_0}(T_{k_i})) \right) ds du. \end{aligned}$$

Let us explain our strategy. We need to show that

$$\sum_{\delta \in \{-1, 0, +1\}^p} \sum_{1 \leq k_1 < k_2 < \dots < k_p \leq M(\epsilon)\epsilon^{-2}} \mathbb{E} \left[\prod_{i=1}^p d_{k_i, k_i + \delta_i} \right],$$

converges to 1, as $\epsilon \rightarrow 0$.

Recall that, from **Section 3.2**, the process $(\omega_{X_0}, \omega_{Y_0})$ (and therefore the processes ω_{X_0} and ω_{Y_0} , considered separately) can be split into i.i.d. 'cycles'

$$(\mathcal{C}_j)_{j \geq 0} = \left(T_{j+1} - T_j, (\omega_{X_0}(T_j + s) - \omega_{X_0}(T_j), \omega_{Y_0}(T_j + s) - \omega_{Y_0}(T_j))_{s \leq T_{j+1} - T_j} \right)_{j \geq 0} \quad (3.30)$$

Observe that d_{k_i, k_i} is a functional of the cycle \mathcal{C}_{k_i} and of the point that the cycle initiates i.e. of $\omega_{X_0}(T_{k_i})$ and $\omega_{Y_0}(T_{k_i})$. From the regenerative structure of $(\omega_{X_0}, \omega_{Y_0})$ we see that \mathcal{C}_{k_i} is independent from

$$(\omega_{X_0}(T_{k_i}), \omega_{Y_0}(T_{k_i}))$$

Therefore we can write the expectation of d_{k_i, k_i} as

$$\mathbb{E}[d_{k_i, k_i}] = \mathbb{E}[\mathcal{G}_\epsilon(\omega_{X_0}(T_{k_i}) - \omega_{Y_0}(T_{k_i}))],$$

with

$$\mathcal{G}_\epsilon(z) = \mathbb{E}_{\text{cycle law}} \left[\int_0^\theta \int_0^\theta R(s-u, x+z + \tilde{\omega}_{B^1}(s) - \tilde{\omega}_{B^2}(u)) ds du \right],$$

where $(\theta, (\tilde{\omega}_{B^1}(s), \tilde{\omega}_{B^2}(s))_{s \leq \theta})$ is as in (3.20). So, in theory, if we want to calculate the limit of (3.28) we can average over the law of the cycles first. This will give us a functional of $(\omega_{X_0}(T_{k_i}) - \omega_{Y_0}(T_{k_i}))_{i=1, \dots, p}$. From (3.18) these random variables are sums of i.i.d. random variables, which guarantees their 'good behavior' (see **Proposition A.2**). Then the resulting additive functional of $(\omega_{X_0}(T_{k_i}) - \omega_{Y_0}(T_{k_i}))_{i=1, \dots, p}$ can be handled by standard results from [KR54].

There are two obstacles to implementing this strategy:

P.1 First, it is obvious that we have to consider the terms $d_{k_i, k_i + \delta_i}$, for $\delta_i \in \{-1, 0, 1\}$. When $\delta_i \neq 0$, the term $d_{k_i, k_i + \delta_i}$ compares the $k_i + \delta_i$ -th cycle of ω_{X_0} to the k_i -th cycle of ω_{Y_0} . So when we average over the law of these two cycles, we will get a function of $\omega_{X_0}(T_{k_i + \delta_i}) - \omega_{Y_0}(T_{k_i})$. This will lead us to a 'non-directed' additive functional of $(\omega_{X_0}(T_{k_i}) - \omega_{Y_0}(T_{k_i}))_{i=1, \dots, p}$ (see (3.37) below). Nevertheless, this can be handled with **Lemma B.1**.

P.2 Secondly, we cannot average over all cycles in the product

$$\mathbb{E} \left[\prod_{i=1}^p d_{k_i, k_i + \delta_i} \right] \quad (3.31)$$

to get an expectation of a function of $(\omega_{X_0}(T_{k_i + \delta_i}) - \omega_{Y_0}(T_{k_i}))_{i=1, \dots, p}$. The reason for this becomes apparent even in the case of $p = 2$ for the expectation

$$\mathbb{E}[d_{k_1, k_1} d_{k_2, k_2}],$$

with $k_1 < k_2$. As mentioned, d_{k_i} , $i = 1, 2$ is a function of \mathcal{C}_{k_i} and of $\omega_{X_0}(T_{k_i}) - \omega_{Y_0}(T_{k_i})$. However, $\omega_{X_0}(T_{k_2}) - \omega_{Y_0}(T_{k_2})$ is not independent from \mathcal{C}_{k_1} . Indeed, consider the paths $(\omega_{X_0}(T_{k_i} + s) - \omega_{X_0}(T_{k_i}), \omega_{Y_0}(T_{k_i} + u) - \omega_{Y_0}(T_{k_i}))_{s, u \in [0, T_{k_{i+1}} - T_{k_i}]}$. Then for $s = u = T_{k_{i+1}} - T_{k_i}$ these paths take the value:

$$(\omega_{X_0}(T_{k_{i+1}}) - \omega_{X_0}(T_{k_i}), \omega_{Y_0}(T_{k_{i+1}}) - \omega_{Y_0}(T_{k_i})).$$

From (3.18), this is equal to $(\mathbf{X}_{k_1}^{(\epsilon)}, \mathbf{Y}_{k_1}^{(\epsilon)})$. Since $k_2 > k_1$, again from (3.18),

$$\mathbf{X}_{k_1}^{(\epsilon)} - \mathbf{Y}_{k_1}^{(\epsilon)}$$

is one of the summands of $\omega_{X_0}(T_{k_2}) - \omega_{Y_0}(T_{k_2})$. This problem becomes more complicated in the general case. In particular, the product in (3.31) is a function of

$$\mathbf{Z}_{k_i} := \left(T_{k_{i+1}} - T_{k_i}, T_{k_i + \delta_i + 1} - T_{k_i + \delta_i}, (\omega_{X_0}(s + T_{k_i + \delta_i}) - \omega_{X_0}(T_{k_i + \delta_i}))_{s \leq T_{k_i + \delta_i + 1} - T_{k_i + \delta_i}}, \right. \\ \left. (\omega_{Y_0}(u + T_{k_i}) - \omega_{Y_0}(T_{k_i}))_{u \leq T_{k_{i+1}} - T_{k_i}} \right), \quad (3.32)$$

for $i = 1, \dots, p$ and of

$$(\omega_{X_0}(T_{k_1 + \delta_1}) - \omega_{Y_0}(T_{k_1}), \dots, \omega_{X_0}(T_{k_p + \delta_p}) - \omega_{Y_0}(T_{k_p})). \quad (3.33)$$

For the same reasons as in the previous example, the cycles

$$\left(T_{k_i+\delta_i+1} - T_{k_i+\delta_i}, (\omega_{X_0}(s+T_{k_i+\delta_i}) - \omega_{X_0}(T_{k_i+\delta_i})), \omega_{Y_0}(s+T_{k_i+\delta_i}) - \omega_{Y_0}(T_{k_i+\delta_i})_{s \in [0, T_{k_i+\delta_i+1} - T_{k_i+\delta_i}]} \right)$$

and

$$\left(T_{k_i+1} - T_{k_i}, (\omega_{X_0}(s+T_{k_i}) - \omega_{X_0}(T_{k_i})), \omega_{Y_0}(s+T_{k_i}) - \omega_{Y_0}(T_{k_i})_{s \in [0, T_{k_i+1} - T_{k_i}]} \right),$$

are not independent from $\omega_{X_0}(T_{k_j+\delta_j}) - \omega_{Y_0}(T_{k_j})$ for $i < j$. However, the latter depends on these cycles only through $\mathbf{X}_{k_i}^{(\epsilon)} - \mathbf{Y}_{k_i}^{(\epsilon)}$ and $\mathbf{X}_{k_i+\delta_i}^{(\epsilon)} - \mathbf{Y}_{k_i+\delta_i}^{(\epsilon)}$.

Now, we explain how we deal with item **P.2**. The idea is to write (3.33) as $\mathcal{S}^\epsilon + \mathcal{Y}^\epsilon$, for appropriate, independent random variables, so that \mathcal{S}^ϵ is independent from the cycles (3.30) and equal in distribution to a shifted version of (3.33). This is accomplished by 'throwing out' the terms $\mathbf{X}_{k_i}^{(\epsilon)} - \mathbf{Y}_{k_i}^{(\epsilon)}$ and $\mathbf{X}_{k_i+\delta_i}^{(\epsilon)} - \mathbf{Y}_{k_i+\delta_i}^{(\epsilon)}$ from the sum (3.18), defining $\omega_{X_0}(T_{k_j+\delta_j}) - \omega_{Y_0}(T_{k_j})$, $1 \leq i \leq j \leq p$. With such an expression in hand, and by averaging over the joint law of \mathcal{Y}^ϵ and of (3.32) we can write (3.31) as an expectation of a function of a shifted version of (3.33). Then, as mentioned, the resulting additive functional can be handled with **Lemma B.1**.

First, we will calculate the limit of:

$$\sum_{\delta \in \{-1, 0, +1\}^p} \sum_{\substack{1 \leq k_1 < k_2 < \dots < k_p \leq M(\epsilon)\epsilon^{-2}, \\ k_{i+1} > k_i + 2}} \mathbb{E} \left[\prod_{i=1}^p d_{k_i, k_i + \delta_i} \right]. \quad (3.34)$$

For $i = 2, \dots, p$ we write:

$$\omega_{Y_0}(T_{k_i}) = \bar{\mathbf{S}}_{k_i} + \sum_{j=1}^{i-1} (\mathbf{Y}_{k_j}^{(\epsilon)} + \mathbf{Y}_{k_j+\delta_j}^{(\epsilon)}) + \mathbf{Y}_{k_{i-1}}^{(\epsilon)}.$$

We included $\mathbf{Y}_{k_{i-1}}^{(\epsilon)}$ as well since in the case where $\delta_i = -1$, $\omega_{Y_0}(T_{k_i})$ also depends on $T_{k_i} - T_{k_{i-1}}$. Similarly, we write:

$$\omega_{Y_0}(T_{k_i+\delta_i}) = \mathbf{S}_{k_i+\delta_i} + \sum_{j=1}^{i-1} (\mathbf{X}_{k_j}^{(\epsilon)} + \mathbf{X}_{k_j+\delta_j}^{(\epsilon)}) + \mathbf{X}_{k_{i-1}}^{(\epsilon)},$$

where $\mathbf{S}_{k_i}, \bar{\mathbf{S}}_{k_i}$ are defined by the equations above and are independent from $\mathbf{X}_{k_j}^{(\epsilon)}, \mathbf{X}_{k_j+\delta_j}^{(\epsilon)}, \mathbf{Y}_{k_j}^{(\epsilon)}, \mathbf{Y}_{k_j+\delta_j}^{(\epsilon)}$, $j = 0, \dots, i-1$ and from $\mathbf{X}_{k_{i-1}}^{(\epsilon)}, \mathbf{Y}_{k_{i-1}}^{(\epsilon)}$. From the above observations the vector:

$$(\mathbf{S}_{k_1+\delta_1} - \bar{\mathbf{S}}_{k_1}, \dots, \mathbf{S}_{k_p+\delta_p} - \bar{\mathbf{S}}_{k_p}),$$

is independent from $(\mathbf{Z}_{k_l})_{l=1, \dots, p}$ and it is equal in distribution to the vector:

$$(\omega_{X_0}(T_{k_1-1+\delta_1}) - \omega_{Y_0}(T_{k_1-1}), \dots, \omega_{X_0}(T_{k_p+\delta_p-2p}) - \omega_{Y_0}(T_{k_p-2p})).$$

Now we focus on writing explicitly the the joint probability distribution of:

$$\left(\mathbf{Z}_{k_i}; \sum_{j=1}^{i-1} (\mathbf{X}_{k_j}^{(\epsilon)} - \mathbf{Y}_{k_j}^{(\epsilon)} + \mathbf{X}_{k_j+\delta_j}^{(\epsilon)} - \mathbf{Y}_{k_j+\delta_j}^{(\epsilon)}) + \mathbf{X}_{k_{i-1}}^{(\epsilon)} - \mathbf{Y}_{k_{i-1}}^{(\epsilon)} \right)_{i=1, \dots, p}. \quad (3.35)$$

Looking at (3.35), we see that it depends on the cycles $((\mathcal{C}_{k_i-1}, \mathcal{C}_{k_i}, \mathcal{C}_{k_i+1}))_{i=1, \dots, p}$. Since we have the restriction of $k_{i+1} > k_i + 2$, the triplets $(\mathcal{C}_{k_i-1}, \mathcal{C}_{k_i}, \mathcal{C}_{k_i+1})$ and $(\mathcal{C}_{k_j-1}, \mathcal{C}_{k_j}, \mathcal{C}_{k_j+1})$ are comprised by distinct cycles for $i \neq j$, and are therefore independent. The distribution of $(\mathcal{C}_{k_i-1}, \mathcal{C}_{k_i}, \mathcal{C}_{k_i+1})$ is equal to the distribution of

$$\left(\left(\theta_{-1}^i, (\tilde{\omega}_{B_{-1}^i}(s), \tilde{\omega}_{b_{-1}^i}(s))_{s \leq \theta_{-1}^i} \right), \left(\theta_0^i, (\tilde{\omega}_{B_0^i}(s), \tilde{\omega}_{b_0^i}(s))_{s \leq \theta_0^i} \right), \left(\theta_1^i, (\tilde{\omega}_{B_1^i}(s), \tilde{\omega}_{b_1^i}(s))_{s \leq \theta_1^i} \right) \right),$$

where:

- $(B_{-1}^i, b_{-1}^i)_{i=1, \dots, p}$, $(B_0^i, b_0^i)_{i=1, \dots, p}$, $(B_1^i, b_1^i)_{i=1, \dots, p}$ are independent collections of independent Brownian motions in $\Omega_1 \times \Omega_1$.
- $(\theta_{-1}^i)_{i=1, \dots, p}$, $(\theta_0^i)_{i=1, \dots, p}$, $(\theta_1^i)_{i=1, \dots, p}$ are independent collections of independent geometric random variables with parameter γ .
- $((\tilde{\omega}_{B_{-1}^i}, \tilde{\omega}_{b_{-1}^i}))_{i=1, \dots, p}$, $((\tilde{\omega}_{B_0^i}, \tilde{\omega}_{b_0^i}))_{i=1, \dots, p}$, $((\tilde{\omega}_{B_1^i}, \tilde{\omega}_{b_1^i}))_{i=1, \dots, p}$ are collections of continuous paths built from the transition probability kernel $\frac{\tilde{\pi} - \gamma W_1 \times W_1}{1 - \gamma}$ with (B_{-1}^i, b_{-1}^i) , (B_0^i, b_0^i) , (B_1^i, b_1^i) as initial steps, respectively. These are independent collections of independent paths. Each of these paths is also independent from $(\theta_{-1}^i, \theta_0^i, \theta_1^i)_{i=1, \dots, p}$.

This proves that (3.35) is equal in distribution to:

$$\mathcal{Z}^{(\epsilon)} := \left(\theta_0^i, \theta_{\delta_i}^i, (\tilde{\omega}_{B_{\delta_i}^{i+\delta_i}}(s))_{s \leq \theta_{i+\delta_i}}, (\tilde{\omega}_{b_0^i}(u))_{u \leq \theta_i}; \mathcal{Y}_i^{(\epsilon)} \right)_{i=1, \dots, p}, \quad (3.36)$$

where

$$\mathcal{Y}_i^{(\epsilon)} = \sum_{j=1}^{i-1} (\tilde{\omega}_{B_0^j}(\theta_0^j) - \tilde{\omega}_{b_0^j}(\theta_0^j) + \tilde{\omega}_{B_{\delta_j}^j}(\theta_{\delta_j}^j) - \tilde{\omega}_{b_{\delta_j}^j}(\theta_{\delta_j}^j)) + \tilde{\omega}_{B_{-1}^i}(\theta_{-1}^i) - \tilde{\omega}_{b_{-1}^i}(\theta_{-1}^i).$$

Furthermore $\mathcal{Z}^{(\epsilon)}$ is independent from $(\mathbf{S}_{k_i} - \bar{\mathbf{S}}_{k_i})_{i=1, \dots, p}$. This implies that (3.34) is equal to

$$\sum_{\delta \in \{-1, 0, +1\}^p} \sum_{\substack{1 \leq k_1 < k_2 < \dots < k_p \leq M(\epsilon) \epsilon^{-2}, \\ k_{i+1} > k_i + 2}} \mathbb{E} \left[\mathcal{G}^{\epsilon, \delta} \left(\omega_{X_0}(T_{k_1 + \delta_1}) - \omega_{Y_0}(T_{k_1}), \dots, \omega_{X_0}(T_{k_p + \delta_p - p + 1}) - \omega_{Y_0}(T_{k_p - p + 1}) \right) \right], \quad (3.37)$$

where

$$\mathcal{G}^{\epsilon, \delta}(z_1, \dots, z_p) = \mathbb{E}_{\mathcal{Z}^{(\epsilon)}} \left[\prod_{i=1}^p \int_0^{\theta_0^i} \int_0^{\theta_{\delta_i}^i} R \left(s - u + \mathbf{1}_{\delta_i=1} \theta_0^i - \mathbf{1}_{\delta_i=-1} \theta_{-1}^i, x + z_i + \mathcal{Y}_i^{(\epsilon)} + \tilde{\omega}_{B_0^i}^i(s) - \tilde{\omega}_{b_{\delta_i}^i}^i(u) \right) ds du \right]. \quad (3.38)$$

We want to apply **Lemma B.1** to (3.37), so we seek to prove that $(\mathcal{G}^{\epsilon, \delta})_{\epsilon > 0}$ satisfies the assumptions of the lemma, namely that:

L.1 The L^1 norm of $\mathcal{G}^{\epsilon, \delta}$ does not depend on ϵ .

L.2 We have

$$\lim_{M \rightarrow \infty} \sup_{\epsilon > 0} \int_{|x| \geq M_0} \mathcal{G}^{\epsilon, \delta}(x) dx = 0.$$

Observe that

$$\|\mathcal{G}^{\epsilon, \delta}\|_1 = \|\psi \star \psi\|_1 \mathbb{E} \left[\prod_{i=1}^p \int_0^{\theta_0^i} \int_0^{\theta_{\delta_i}^i} \phi \star \phi(s - u + \mathbf{1}_{\delta_i=1} \theta_0^i - \mathbf{1}_{\delta_i=-1} \theta_{-1}^i) ds du \right],$$

which is indeed an ϵ -independent constant, so **L.1** is true. To prove item **L.2** we observe that, from **Proposition A.1** and **Corollary A.1**,

$$\mathbb{E} \left[\mathbf{1}_{\{|k_\epsilon| \geq M_0\}} R \left(s - u + \mathbf{1}_{\delta_i=1} \theta_0^i - \mathbf{1}_{\delta_i=-1} \theta_{-1}^i, x + \mathcal{Y}_i^{(\epsilon)} + \tilde{\omega}_{B_0^i}^i(s) - \tilde{\omega}_{b_{\delta_i}^i}^i(u) \right) \right] \rightarrow 0,$$

as $M_0 \rightarrow \infty$, uniformly in ϵ , where

$$k_\epsilon = |\mathcal{Y}_i^{(\epsilon)}| + \sup_{s \leq \theta_0^i} |\tilde{\omega}_{B_0^i}^i(s)| + \sup_{s \leq \theta_{\delta_i}^i} |\tilde{\omega}_{b_{\delta_i}^i}^i(s)|.$$

Arguing similarly as in **Remark B.1**, $(\mathcal{G}^{\epsilon, \delta})_{\epsilon > 0}$ satisfies **L.2**. Therefore, **Lemma B.1** implies that (3.37) is asymptotic to:

$$\left((\gamma/2\pi) \log \epsilon^{-1} \right)^p \sum_{\delta \in \{-1, 0, +1\}^p} \|\mathcal{G}^{\epsilon, \delta}\|_1, \quad (3.39)$$

as $\epsilon \rightarrow 0$.

We need to show that this expression is equal to $(\|R\|_1 \log \epsilon^{-1} / 2\pi)^p$. Since $R(t, x) = \phi \star \tilde{\phi}(t) \psi \star \psi(x)$ it suffices to show that

$$\sum_{\delta \in \{-1, 0, +1\}^p} \mathbb{E} \left[\prod_{i=1}^p \int_0^{\theta_0^i} \int_0^{\theta_{\delta_i}^i} \phi \star \phi(s - u + \mathbf{1}_{\delta_i=1} \theta_0^i - \mathbf{1}_{\delta_i=-1} \theta_{-1}^i) ds du \right] = \|\phi \star \tilde{\phi}\|_1 = \|\phi\|_1^2$$

By writing the above expectation explicitly we have to calculate

$$\sum_{\delta \in \{-1, 0, +1\}^p} \sum_{\mathbf{N} \in \mathbb{N}^{3p}} \prod_{i=1}^p \gamma^3 (1 - \gamma)^{N_{-1}^i + N_0^i + N_1^i - 3} \prod_{i=1}^p \int_0^{N_0^i} \int_0^{N_{\delta_i}^i} \phi \star \tilde{\phi}(s - u - \mathbf{1}_{\delta_i=1} N_0^i + \mathbf{1}_{\delta_i=-1} N_{-1}^i) ds du.$$

For each $\delta \in \{-1, 0, +1\}^p$ we partition the set of indices $\{1, \dots, p\}$ into the sets $A_+ = \{\delta_i = 1\}$, $A_- = \{\delta_i = -1\}$ and $A_0 = \{\delta_i = 0\}$. Also define:

$$L_0(N) := N \cdot \int_0^1 \int_0^1 \phi \star \tilde{\phi}(s - u) ds du + 2 \cdot (N - 1) \int_0^1 \int_0^1 \phi \star \tilde{\phi}(s + u) ds du,$$

$$L_- = L_+ := \int_0^1 \int_0^1 \phi \star \tilde{\phi}(s + u) ds du.$$

With these definitions, the above quantity can be written as

$$\sum_{\{A_0, A_-, A_+\} \vdash \{1, \dots, p\}} \sum_{\mathbf{N} \in \mathbb{N}^{3p}} \prod_{i=1}^p \gamma^3 (1 - \gamma)^{N_{-1}^i + N_0^i + N_1^i - 3} \prod_{i \in A_0} L_0(N_0^i) L_+^{|A_+|} L_-^{|A_-|}, \quad (3.40)$$

where $\{A_0, A_-, A_+\} \vdash \{1, \dots, p\}$ means that the sets A_0, A_-, A_+ partition $\{1, \dots, p\}$, and we sum over all such partitions. By summing over all \mathbf{N} we that (3.40) is equal to

$$\sum_{\{A_0, A_-, A_+\} \vdash \{1, \dots, p\}} L_\gamma^{|A_0|} L_+^{|A_+|} L_-^{|A_-|},$$

where

$$L_\gamma := \frac{1}{\gamma} \left(\int_0^1 \int_0^1 \phi \star \tilde{\phi}(s-u) dsdu + 2 \int_0^1 \int_0^1 \phi \star \tilde{\phi}(s+u) dsdu \right) - 2 \int_0^1 \int_0^1 \phi \star \tilde{\phi}(s+u) dsdu.$$

Therefore (3.40) is equal to

$$\left(L_\gamma + L_+ + L_- \right)^p = (A(R)/\gamma)^p,$$

where

$$A(R) := \left(\int_0^1 \int_0^1 \phi \star \tilde{\phi}(s-u) dsdu + 2 \int_0^1 \int_0^1 \phi \star \tilde{\phi}(s+u) dsdu \right) \|\psi\|_1^2.$$

This implies that (3.34) is asymptotic to

$$\left((\gamma/2\pi) \log \epsilon^{-1} \right)^p (A(R)/\gamma)^p = \left(\frac{A(R)}{2\pi \log \epsilon^{-1}} \right)^p.$$

We will show that $A(R) = \|R\|_1$. Since, $\|R\|_1 = \|\phi\|_1^2 \|\psi\|_1^2$ it is sufficient to prove that

$$\int_0^1 \int_0^1 \phi \star \tilde{\phi}(s-u) dsdu + 2 \int_0^1 \int_0^1 \phi \star \tilde{\phi}(s+u) dsdu = \|\phi\|_1^2.$$

By a change of variables $s \rightarrow s+u$, the first term on the left-hand side is equal to:

$$\int_0^1 \int_{-u}^{1-u} \phi \star \tilde{\phi}(s) dsdu,$$

while the second term is equal to:

$$\int_0^1 \int_u^1 \phi \star \tilde{\phi}(s) dsdu + \int_0^1 \int_{1-u}^1 \phi \star \tilde{\phi}(s) dsdu,$$

where we also used the fact that $\phi \star \tilde{\phi}(s) = 0$ for $s \geq 1$. By adding these terms we get:

$$\int_0^1 \left(\int_{-u}^{1-u} \phi \star \tilde{\phi}(s) dsdu + \int_u^1 \phi \star \tilde{\phi}(s) dsdu + \int_{1-u}^1 \phi \star \tilde{\phi}(s) ds \right) du.$$

In the second term inside the integral we use the fact that $\phi \star \tilde{\phi}$ is symmetric, and we get that the above integral is equal to:

$$\int_0^1 \left(\int_{-u}^{1-u} \phi \star \tilde{\phi}(s) dsdu + \int_{-u}^{-1} \phi \star \tilde{\phi}(s) dsdu + \int_{1-u}^1 \phi \star \tilde{\phi}(s) ds \right) du = \int_0^1 \int_{-1}^1 \phi \star \tilde{\phi}(s) dsdu = \|\phi\|_1^2,$$

and therefore $A(R) = \|R\|_1$.

This proves that (3.39) is equal to $(\|R\|_1 \log \epsilon^{-1}/2\pi)^p$ which implies that

$$\left(\frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \right)^p \sum_{\substack{1 \leq k_1 < k_2 < \dots < k_p \leq M(\epsilon)\epsilon^{-2}, \\ k_{i+1} > k_i + 2}} \mathbb{E} \left[\prod_{i=1}^p d_{k_i, k_i-1} d_{k_i, k_i} d_{k_i, k_i+1} \right].$$

converges to 1 as $\epsilon \rightarrow 0$. This result implies that

$$\left(\frac{2\pi}{\|R\|_1 \log \epsilon^{-1}} \right)^p \sum_{1 \leq k_1 < k_2 < \dots < k_p \leq M(\epsilon)\epsilon^{-2}} \mathbb{E} \left[\prod_{i=1}^p \left(d_{k_i, k_i-1} d_{k_i, k_i} d_{k_i, k_i+1} \right) \right] \quad (3.41)$$

converges to 1 as $\epsilon \rightarrow 0$. Indeed for $r = 1, \dots, p$ and for $1 \leq i_1 < i_2 < \dots < i_r \leq p$, we define A_{i_1, \dots, i_r} to be the set:

$$\{k_i \in \mathbb{N}, i = 1, \dots, p : 1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, k_{i_j} \leq k_{i_{j-1}} + 2, k_i > k_{i-1} + 2, i \neq i_1, \dots, i_r\}. \quad (3.42)$$

The set A_{i_1, \dots, i_r} specifies where the restriction $k_{i+1} > k_i + 2$ fails. We consider the sum

$$\sum_{k_1, \dots, k_p \in A_{i_1, \dots, i_r}} \mathbb{E} \left[\prod_{i=1}^p \left(d_{k_i, k_{i-1}} d_{k_i, k_i} d_{k_i, k_{i+1}} \right) \right].$$

Observe that if there is an i such that $k_{i_r+1} = k_{i_r} + 1$ or $k_{i_r+1} = k_{i_r} + 2$. By repeating the same arguments as above, we can prove that this sum is of order $(\log \epsilon^{-1})^{p-r} = o((\log \epsilon^{-1})^p)$ and is therefore negligible compared to (3.34). This completes the proof. \square

Corollary 3.2. *Let $X_0, Y_0, R, M(\epsilon)$ be as before. Then for every $\hat{\beta}^2 < \|R\|_1/2\pi$*

$$\sup_{x \in \mathbb{R}^2, \epsilon \in (0,1)} \mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^{-2}} \int_0^{M(\epsilon)/\epsilon^{-2}} R(s-u, x + \omega_{X_0}(s) - \omega_{Y_0}(u)) ds du \right) \right] < \infty.$$

Proof. Using similar arguments as in **Proposition 1.1** we have

$$e^{-2\zeta_{M(\epsilon)/\epsilon^2}^{(\epsilon)}} \mathbf{E}[u(M(\epsilon)/\epsilon^2, x)u(M(\epsilon)/\epsilon^2, 0)] = \hat{\mathbb{E}}_{B, M(\epsilon)/\epsilon^2} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^{-2}} \int_0^{M(\epsilon)/\epsilon^{-2}} R(s-u, x + B^1(s) - B^2(u)) ds du \right) \right], \quad (3.43)$$

where we recall that $u(t, x)$ solves (1.4).

Now observe that $\mathbf{E}[u(t, x)^2] = \mathbf{E}[u(t, 0)^2]$ for all (t, x) . The Cauchy-Schwartz inequality implies that, for all $\epsilon > 0$, (3.43) is maximized when $x = 0$. Indeed we have

$$e^{-2\zeta_{M(\epsilon)/\epsilon^2}^{(\epsilon)}} \mathbf{E}[u(M(\epsilon)/\epsilon^2, x)u(M(\epsilon)/\epsilon^2, 0)] \leq e^{-2\zeta_{M(\epsilon)/\epsilon^2}^{(\epsilon)}} \mathbf{E}[u(M(\epsilon)/\epsilon^2, 0)^2],$$

and the right-hand side is equal to

$$\hat{\mathbb{E}}_{B, M(\epsilon)/\epsilon^2} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^{-2}} \int_0^{M(\epsilon)/\epsilon^{-2}} R(s-u, B^1(s) - B^2(u)) ds du \right) \right]. \quad (3.44)$$

Therefore it is sufficient to bound (3.44) for all $\epsilon \in (0, 1)$. We will use **Theorem 3.1** to argue that (3.44) is bounded for all $\hat{\beta}^2 < \|R\|_1/2\pi$.

We write the expectation in (3.44) as an expectation involving paths built from the Markov chain. We choose $T = M(\epsilon)/\epsilon^2$, $\tau = 1$, $N = [M(\epsilon)/\epsilon^2 - 1]$. We sample independently two random vectors $(X_0^1, \dots, X_{N_\epsilon+1}^1), (X_0^2, \dots, X_{N_\epsilon+1}^2)$ as described in **Section 3.1** (recall that $X_0^1, X_0^2 \in \Omega_\tau$ and $X_{N_\epsilon+1}^1, X_{N_\epsilon+1}^2 \in \Omega_{M(\epsilon)/\epsilon^2 - \tau - N_\epsilon}$). Recall that $[X_0^i, \dots, X_{N_\epsilon+1}^i]$ denotes the path built from $(X_0^i, \dots, X_{N_\epsilon+1}^i)$ as in (3.1). From (3.13), the expectation in (3.43) is equal to

$$\mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, M(\epsilon)/\epsilon^2]^2} R(s-u, x + [X_0^1, \dots, X_{N_\epsilon+1}^1](s) - [X_0^2, \dots, X_{N_\epsilon+1}^2](u)) ds du \right) \mathcal{G}_\epsilon(X_N, Y_N) \right],$$

where \mathcal{G}_ϵ is defined by (3.12). From **Remark 3.1**, \mathcal{G}_ϵ converges to 1 uniformly as $\epsilon \rightarrow 0$. Furthermore, since R is bounded, we have

$$\mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, M(\epsilon)/\epsilon^2]^2} R(s-u, [X_0^1, \dots, X_{N_\epsilon+1}^1](s) - [X_0^2, \dots, X_{N_\epsilon+1}^2](u)) ds du \right) \mathcal{G}_\epsilon(X_{N_\epsilon}, Y_{N_\epsilon}) \right] \lesssim$$

$$\mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, [M(\epsilon)/\epsilon^2] - 1]^2} R(s - u, [X_0^1, \dots, X_{N_\epsilon+1}^1](s) - [X_0^2, \dots, X_{N_\epsilon+1}^2](u)) dsdu \right) \right].$$

Observe that the integral in the right-hand side is over $[0, [M(\epsilon)/\epsilon^2] - 1]^2$ instead of $[0, M(\epsilon)/\epsilon^2]^2$. This ensures that $X_{N_\epsilon+1}^1, X_{N_\epsilon+1}^2$ will not appear in the above expectation. Hence, we can consider the paths that appear in the integral above as built from $\hat{\pi}^{(\epsilon)}$. Therefore, they are almost like the paths in the statement of **Theorem 3.1**, with the only difference being the distribution of the initial segment. From **Lemma 3.1** we can replace X_0^1, X_0^2 by independent Brownian motions. More specifically, if $(\omega_{X_0}, \omega_{Y_0})$ is a pair of paths built from $\hat{\pi}^{(\epsilon)}$ with $(X_0, Y_0) \sim W_1 \times W_1$ as the initial condition, then

$$\begin{aligned} \mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, M(\epsilon)/\epsilon^2]^2} R(s - u, [X_0^1, \dots, X_{N_\epsilon+1}^1](s) - [X_0^2, \dots, X_{N_\epsilon+1}^2](u)) dsdu \right) \right] &\lesssim \\ \mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, [M(\epsilon)/\epsilon^2] - 1]^2} R(s - u, \omega_{X_0}(s) - \omega_{Y_0}(u)) dsdu \right) \right]. \end{aligned}$$

for all $\epsilon > 0$ and $x \in \mathbb{R}^2$. This proves that for all $\epsilon \in (0, 1)$

$$\begin{aligned} \hat{\mathbb{E}}_{B, M(\epsilon)/\epsilon^2} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{M(\epsilon)/\epsilon^{-2}} \int_0^{M(\epsilon)/\epsilon^{-2}} R(s - u, B^1(s) - B^2(u)) dsdu \right) \right] &\lesssim \\ \mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, [M(\epsilon)/\epsilon^2] - 1]^2} R(s - u, \omega_{X_0}(s) - \omega_{Y_0}(u)) dsdu \right) \right]. \end{aligned}$$

From **Theorem 3.1** we have

$$\sup_{\epsilon \in (0, 1)} \mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, [M(\epsilon)/\epsilon^2] - 1]^2} R(s - u, \omega_{X_0}(s) - \omega_{X_0}(u)) dsdu \right) \right] < \infty,$$

which concludes the proof. \square

Remark 3.3. Recall that $\sqrt{2\pi/\|R\|_1} = \hat{\beta}_c(R)$. **Theorem 3.1** implies that

$$\mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{[0, [M(\epsilon)/\epsilon^2]^2} R(s - u, x + \omega_{X_0}(s) - \omega_{X_0}(u)) dsdu \right) \right] \rightarrow \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2} \right)^{-1},$$

as $\epsilon \rightarrow 0$. Moreover, arguing similarly as in the proof of **Corollary 3.2** we can prove that

$$\mathbf{E}[u_\epsilon(t, x)^2] \rightarrow \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2} \right)^{-1},$$

as $\epsilon \rightarrow 0$.

4 Proof of Proposition 1.4

In this section, we will prove **Proposition 1.4**. Recall the formula for $\mathcal{F}_\epsilon(r, y, M_1(\epsilon), M_2(\epsilon))$:

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon), M_2(\epsilon)) = \int_{\mathbb{R}^4} \int_{[0, 1]^2} \hat{\mathbb{E}}_{B, t/\epsilon^2} [I_\epsilon e^{J_\epsilon(M_1(\epsilon), M_2(\epsilon))}] \prod_{i=1}^2 \phi(s_i) \psi(x_i) d\bar{s} d\bar{x}, \quad (4.1)$$

where

$$I_\epsilon(x_1, x_2, y, s_1, s_2, r) = \prod_{i=1}^2 g(\epsilon x_i - \epsilon B_{(t-r)/\epsilon^2 - s_i}^i + y) \quad (4.2)$$

and

$$J_\epsilon(M_1(\epsilon), M_2(\epsilon)) = \frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{-1}^{M_1(\epsilon)} \int_{-1}^{M_2(\epsilon)} R_\phi(u_1, u_2) \psi \star \psi(x_1 - x_2 + \Delta B_{(t-r)/\epsilon^2 - s_1, (t-r)/\epsilon^2 + u_1}^1 - \Delta B_{(t-r)/\epsilon^2 - s_2, (t-r)/\epsilon^2 + u_2}^2) du_1 du_2, \quad (4.3)$$

with R_ϕ as in (1.21). Finally, recall that $M_1(\epsilon), M_2(\epsilon)$ satisfy the following assumptions:

A.1 $0 \leq M_1(\epsilon), M_2(\epsilon) \leq r$.

A.2 For $i = 1, 2$, $\log M_i(\epsilon) / \log \epsilon^{-1} \rightarrow 0$, as $\epsilon \rightarrow 0$.

A.3 Either $M_1(\epsilon) = M_2(\epsilon)$ or for all ϵ small enough $M_1(\epsilon) - M_2(\epsilon) \geq c > 0$, for some c .

First, we argue that it is sufficient to prove **Proposition 1.4** in the case $M_1(\epsilon) = M_2(\epsilon)$. Indeed, recall from **Remark 1.1** that if $M_1(\epsilon) - M_2(\epsilon) \geq c > 0$, then for all ϵ small enough

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) = \mathcal{F}_\epsilon(r, y, M_2(\epsilon)/\epsilon^2 + 1, M_2(\epsilon)/\epsilon^2).$$

From (4.3) we get that

$$J_\epsilon(M_2(\epsilon)/\epsilon^2 + 1, M_2(\epsilon)/\epsilon^2) = \mathcal{R}_\epsilon + J_\epsilon(M_2(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2), \quad (4.4)$$

with

$$\mathcal{R}_\epsilon := \frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{M_2(\epsilon)/\epsilon^2}^{M_2(\epsilon)/\epsilon^2 + 1} \int_{-1}^{M_2(\epsilon)/\epsilon^2} R_\phi(u_1, u_2) \psi \star \psi(x_1 - x_2 + \Delta B_{(t-r)/\epsilon^2 - s_1, (t-r)/\epsilon^2 + u_1}^1 - \Delta B_{(t-r)/\epsilon^2 - s_2, (t-r)/\epsilon^2 + u_2}^2) du_1 du_2. \quad (4.5)$$

Recall that, since ϕ is supported on $[0, 1]$, $R_\phi(u_1, u_2) = 0$ when $|u_1 - u_2| \geq 1$. Therefore, \mathcal{R}_ϵ is equal to

$$\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{M_2(\epsilon)/\epsilon^2}^{M_2(\epsilon)/\epsilon^2 + 1} \int_{M_2(\epsilon)/\epsilon^2 - 1}^{M_2(\epsilon)/\epsilon^2} R_\phi(u_1, u_2) \psi \star \psi(x_1 - x_2 + \Delta B_{(t-r)/\epsilon^2 - s_1, (t-r)/\epsilon^2 + u_1}^1 - \Delta B_{(t-r)/\epsilon^2 - s_2, (t-r)/\epsilon^2 + u_2}^2) du_1 du_2.$$

Since R_ϕ and $\psi \star \psi$ are bounded, we conclude that this term (and therefore \mathcal{R}_ϵ) converges to 0 as $\epsilon \rightarrow 0$, uniformly over the paths B^1, B^2 and over r, y, s, x_1, x_2 . Now, assume for the moment that **Proposition 1.4** is true for $\mathcal{F}_\epsilon(r, y, M_2(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2)$. Then, from our observations about \mathcal{R}_ϵ , (4.1), (4.4) and from Holder's inequality

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) - \mathcal{F}_\epsilon(r, y, M_2(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) \rightarrow 0,$$

as $\epsilon \rightarrow 0$ and

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) \lesssim \mathcal{F}_\epsilon(r, y, M_2(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2),$$

for all ϵ . Hence, **Proposition 1.4** is true for $\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2)$ as well.

Therefore, we will focus on

$$\mathcal{F}_\epsilon(r, y, M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2),$$

where $M(\epsilon)$ satisfies the assumptions **A.1** and **A.2** described above. For this case, we prove **Proposition 1.4** in **Lemma 4.5** and **Lemma 4.4**. First, we will calculate the limit of $\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon \exp(J_\epsilon)]$. Our main strategy will be to try to decouple I_ϵ and J_ϵ . This means that asymptotically we should have

$$\hat{\mathbb{E}}_{B,t/\epsilon^2} \left[I_\epsilon \exp(J_\epsilon) \right] \approx \hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon] \hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(J_\epsilon)].$$

Then the expectation

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(J_\epsilon)]$$

should converge to a constant, giving us the effective variance and

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon]$$

should converge to $p_{t-r} \star g(y)^2$, where p_t is the two-dimensional heat kernel. This heuristic is made formal in **Lemma 4.1** and **Lemma 4.3**.

Lemma 4.1. *For all $\hat{\beta} < \hat{\beta}_c(R)$ we have that*

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon] \rightarrow p_{t-r} \star g(y)^2,$$

and

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2))] \rightarrow \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2} \right)^{-1},$$

as $\epsilon \rightarrow 0$.

Proof. We calculate the limit of $\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon]$ first. Observe that I_ϵ is a functional of two independent paths distributed according to the exponentially tilted measure (1.10). We will emphasize this dependence by writing it as $I_\epsilon(B^1, B^2)$. We replace the paths that appear in (4.2) with paths built from the Markov chain constructed in **Section 3.1**. More specifically set

$$T = t/\epsilon^2, \tau = (t-r)/\epsilon^2 - [(t-r)/\epsilon^2] \text{ and } N_\epsilon = [t/\epsilon^2 - \tau].$$

We sample independently two random vectors $(\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1)$, $(\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2)$ as described in **Section 3.1**, with $\tilde{X}_0^1, \tilde{X}_0^2 \in \Omega_\tau$ and $\tilde{X}_{N_\epsilon+1}^1, \tilde{X}_{N_\epsilon+1}^2 \in \Omega_{t/\epsilon^2 - \tau - N_\epsilon}$. Then from (3.13), we have that:

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon(B^1, B^2)] = \mathbb{E} \left[\mathcal{G}_\epsilon(X_{N_\epsilon}^1, X_{N_\epsilon}^2) I_\epsilon([\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1], [\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2]) \right],$$

with \mathcal{G}_ϵ as in (3.12). From **Remark 3.1** \mathcal{G}_ϵ converges to 1 uniformly as $\epsilon \rightarrow 0$. Therefore, it is sufficient to calculate:

$$\mathbb{E} \left[I_\epsilon([\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1], [\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2]) \right],$$

as $\epsilon \rightarrow 0$.

We see that the term in the above expectation is a functional of $(\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1)$ and $(\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2)$. We will apply **Lemma 3.1** and replace the steps of these two trajectories of the Markov chain at $[(t-r)/\epsilon^2]$ and $[(t-r)/\epsilon^2] + 1$ by independent Brownian motions that are also independent of the previous steps. More specifically, consider the sequences $(X_0^1, \dots, X_{N_\epsilon+1}^1)$ and $(X_0^2, \dots, X_{N_\epsilon+1}^2)$, which

are generated in the same way as $(\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1)$ and $(\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2)$ respectively, but at the steps $k = \lceil (t-r)/\epsilon^2 \rceil, \lceil (t-r)/\epsilon^2 \rceil + 1$ we sample $(X_k^i, X_k^i) \sim W_1 \times W_1$, $i = 1, 2$ independently from the previous steps (also $(X_k^i, X_k^i), (X_k^j, X_k^j)$ are independent for $i \neq j$). Let $\boldsymbol{\omega}_{X_0^1} = [X_0^1, \dots, X_{N_\epsilon+1}^1]$ and $\boldsymbol{\omega}_{X_0^2} = [X_0^2, \dots, X_{N_\epsilon+1}^2]$ be the corresponding paths built according to (3.1). Then, from **Lemma 3.1** and **Remark 3.2**, there are constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$ going to 1 as $\epsilon \rightarrow 0$ such that:

$$\mathcal{A}_\epsilon \mathbb{E} \left[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \right] \leq \mathbb{E} \left[I_\epsilon([\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1], [\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2]) \right] \leq \mathcal{B}_\epsilon \mathbb{E} \left[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \right].$$

Therefore, it is sufficient to calculate the limit of:

$$\mathbb{E} \left[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \right].$$

By adding and subtracting the term $\epsilon \boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1)$ in (4.2) we have:

$$I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) = \prod_{i=1}^2 g(\epsilon x_i - \epsilon \boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1) - \epsilon(\boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - s_i) - \boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1)) + y_i).$$

Recall that (ω_X, ω_Y) denotes a pair of paths built from $\hat{\boldsymbol{\pi}}^{(\epsilon)} = \hat{\boldsymbol{\pi}}^{(\epsilon)} \times \hat{\boldsymbol{\pi}}^{(\epsilon)}$, with $(X, Y) \sim W_1 \times W_1$ as an initial step. For $s_i \in [0, 1]$, from (3.1) and the fact that $(X_{\lceil (t-r)/\epsilon^2 \rceil}^i)_{i=1,2}$ are independent Brownian motions in Ω_1 , independent from the previous steps of the Markov chain, we get that

$$(\boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - s_i) - \boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1))_{i=1,2} =$$

$$(\boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1 + 1 - s_i) - \boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1))_{i=1,2} \stackrel{d}{=} (\omega_{X_{\lceil (t-r)/\epsilon^2 \rceil}^i}(1 - s_i))_{i=1,2},$$

where the last equality is in distribution. The right hand side for $s_i \in [0, 1]$ is equal to $(X_{\lceil (t-r)/\epsilon^2 \rceil}^i(1 - s_i))_{i=1,2}$ which are independent from $\boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1)$, $i = 1, 2$. Since $(X_{\lceil (t-r)/\epsilon^2 \rceil}^1, X_{\lceil (t-r)/\epsilon^2 \rceil}^2) \sim W_1 \times W_1$, $I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$ is equal in distribution to:

$$\prod_{i=1}^2 g(\epsilon x_i - \epsilon \boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1) - \epsilon \tilde{B}_{(1-s_i)}^i + y_i)$$

where $(\tilde{B}^i)_{i=1,2}$ are independent Brownian motions, independent from $(\boldsymbol{\omega}_{X_0^i}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1))_{i=1,2}$. By **Lemma 4.2**, proved below, and **Remark 4.10**,

$$(\epsilon \boldsymbol{\omega}_{X_0^1}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1), \epsilon \boldsymbol{\omega}_{X_0^2}(\lceil (t-r)/\epsilon^2 \rceil + \tau - 1))$$

converges in distribution, as $\epsilon \rightarrow 0$, to a centered Gaussian random variable with covariance $(t-r)I_{4 \times 4}$. Therefore

$$\mathbb{E}[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})] \rightarrow p_{t-r} \star g(y)^2,$$

as $\epsilon \rightarrow 0$ and therefore $\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon(B^1, B^2)] \rightarrow p_{t-r} \star g(y)^2$, as $\epsilon \rightarrow 0$.

Now we look at $\hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(J_\epsilon)]$. Let $A_\epsilon = [-1, M(\epsilon)/\epsilon^2]^2 \cap \left([0, [M(\epsilon)/\epsilon^2] - 1]^2 \right)^c$. We write

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2))] = \hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(\tilde{J}_\epsilon(B^1, B^2) + \mathcal{J}_\epsilon(B^1, B^2))]$$

where:

$$\tilde{J}_\epsilon(B^1, B^2) = \frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{[M(\epsilon)/\epsilon^2]-1} \int_0^{[M(\epsilon)/\epsilon^2]-1} R(u_1 - u_2, x_1 - x_2 + \Delta B_{(t-r)/\epsilon^1 - s_1, (t-r)/\epsilon^2 + u_1}^1 - \Delta B_{(t-r)/\epsilon^2 - s_2, (t-r)/\epsilon^2 + u_2}^2) du_1 du_2 \quad (4.6)$$

and

$$\mathcal{J}_\epsilon(B^1, B^2) = \frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_{A_\epsilon} R_\phi(u_1, u_2) \psi \star \psi(x_1 - x_2 + \Delta B_{(t-r)/\epsilon^1 - s_1, (t-r)/\epsilon^2 + u_1}^1 - \Delta B_{(t-r)/\epsilon^2 - s_2, (t-r)/\epsilon^2 + u_2}^2) du_1 du_2 \quad (4.7)$$

where we also used the fact that $R_\phi(u_1, u_2) = \phi \star \tilde{\phi}(u_1 - u_2)$ when $u_1, u_2 \geq 0$. By arguing similarly as we did for \mathcal{R}_ϵ , defined in (4.5), we can see that $\mathcal{J}_\epsilon(B^1, B^2) \rightarrow 0$, as $\epsilon \rightarrow 0$, uniformly over B^1, B^2 . Therefore to calculate $\hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2)]$ as $\epsilon \rightarrow 0$, it is sufficient to calculate

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(\tilde{J}_\epsilon(B^1, B^2))],$$

as $\epsilon \rightarrow 0$. Using **Lemma 3.1** and arguing similarly as we did in the calculation of $\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon]$, we see that it is enough to calculate the limit of:

$$\mathbb{E}[\exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}))]. \quad (4.8)$$

By adding and subtracting $\boldsymbol{\omega}_{X_0^1}([(t-r)/\epsilon^2] + \tau)$ and $\boldsymbol{\omega}_{X_0^2}([(t-r)/\epsilon^2] + \tau)$ in (4.6) the paths in the spatial term inside the integral can be written as:

$$\sum_{i=1}^2 \left(\boldsymbol{\omega}_{X_0^i}((t-r)/\epsilon^2 + u_1) - \boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau) \right) + \left(\boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau) - \boldsymbol{\omega}_{X_0^i}((t-r)/\epsilon^2 - s_i) \right) \quad (4.9)$$

We deal with the two terms in the above sum, separately. Observe that for $i = 1$ and for $i = 2$ the corresponding summands are independent, by construction. We prove that

I.1 For $i = 1, 2$ the term $(\boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau) - \boldsymbol{\omega}_{X_0^i}((t-r)/\epsilon^2 - s_i))_{s_i \in [0,1]}$ is equal in distribution to a Brownian motion $(\tilde{B}_{s_i}^i)_{s_i \in [0,1]}$.

I.2 For $i = 1, 2$ the term $(\boldsymbol{\omega}_{X_0^i}((t-r)/\epsilon^2 + u_1) - \boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau))_{u_i \in [0, [M(\epsilon)/\epsilon^2]-1]}$ is equal in distribution to $(\omega_{X_{[(t-r)/\epsilon^2]+1}^i}(u_i))_{u_i \in [0, [M(\epsilon)/\epsilon^2]-1]}$.

I.3 The terms

$$(\boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau) - \boldsymbol{\omega}_{X_0^i}((t-r)/\epsilon^2 - s_i))_{s_i \in [0,1]}$$

and

$$(\boldsymbol{\omega}_{X_0^i}((t-r)/\epsilon^2 + u_1) - \boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau))_{u_i \in [0, [M(\epsilon)/\epsilon^2]-1]}$$

are independent.

Observe that these three items, along with **Theorem 3.1** and **Proposition 3.2**, give us the desired limit for (4.8). Indeed, $\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$ is equal in distribution to

$$\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{[M(\epsilon)/\epsilon^2]-1} \int_0^{[M(\epsilon)/\epsilon^2]-1} R \left(u_1 - u_2, x_1 - x_2 + \sum_{i=1}^2 \tilde{B}_{s_i}^i + \omega_{X_{[(t-r)/\epsilon^2]+1}^1}(u_1) - \omega_{X_{[(t-r)/\epsilon^2]+1}^2}(u_2) \right) du_1 du_2,$$

where $(\tilde{B}_{s_1}^1)_{s_1 \in [0,1]}, (\tilde{B}_{s_2}^2)_{s_2 \in [0,1]}$ are two independent Brownian motions that are also independent from

$$(\omega_{X_{[(t-r)/\epsilon^2]+1}^1}(u_1), \omega_{X_{[(t-r)/\epsilon^2]+1}^2}(u_2))_{u_1, u_2 \in [0, [M(\epsilon)/\epsilon^2]-1]}.$$

Also, recall that $(X_{[(t-r)/\epsilon^2]+1}^1, X_{[(t-r)/\epsilon^2]+1}^2) \sim W_1 \times W_1$. Now, if $b^1, b^2 \in \Omega_1$ are two (generic) Brownian motions, this equality in distribution proves that

$$\mathbb{E} \left[\exp(\tilde{J}_\epsilon(\omega_{X_0^1}, \omega_{X_0^2})) \right] = \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \mathbb{E} \left[\exp \left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{[M(\epsilon)/\epsilon^2]} \int_0^{[M(\epsilon)/\epsilon^2]} R \left(u_1 - u_2, x_1 - x_2 + y_1 + y_2 + \omega_{b^1}(u_1) - \omega_{b^2}(u_2) \right) du_1 du_2 \right) \right] p_{s_1}(y_1) p_{s_2}(y_2) dy_1 dy_2. \quad (4.10)$$

Now we apply **Theorem 3.1** and **Proposition 3.2** and get that:

$$\mathbb{E} \left[\exp(\tilde{J}_\epsilon(\omega_{X_0^1}, \omega_{X_0^2})) \right] \rightarrow \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2} \right)^{-1},$$

as $\epsilon \rightarrow 0$. Finally, from our previous remarks, $\hat{\mathbb{E}}_{B,t/\epsilon^2}[\exp(J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2))]$ has the same limit.

It is therefore sufficient to prove items **I.1-I.3**. To prove the item **I.1** we add and subtract $\omega_{X_0^i}([(t-r)/\epsilon^2] + \tau - 1)$. We have that

$$\omega_{X_0^i}([(t-r)/\epsilon^2] + \tau) - \omega_{X_0^i}([(t-r)/\epsilon^2] + \tau - s_i) = \left(\omega_{X_0^i}([(t-r)/\epsilon^2] + \tau) - \omega_{X_0^i}([(t-r)/\epsilon^2] + \tau - 1) \right) - \left(\omega_{X_0^i}([(t-r)/\epsilon^2] + \tau - s_i) - \omega_{X_0^i}([(t-r)/\epsilon^2] + \tau - 1) \right).$$

Recall that $X_{[(t-r)/\epsilon^2]}^i$ is sampled from the Wiener measure W_1 , independently from the previous steps. From this observation and (3.1) the right-hand side, for $s_i \in [0, 1]$, is equal in distribution to:

$$\omega_{X_{[(t-r)/\epsilon^2]}^i}(1) - \omega_{X_{[(t-r)/\epsilon^2]}^i}(1 - s_i). \quad (4.11)$$

Now for $s_i \in [0, 1]$ this is equal to $X_{[(t-r)/\epsilon^2]}^i(1) - X_{[(t-r)/\epsilon^2]}^i(1 - s_i)$ which is distributed like a Brownian motion at $1 - (1 - s_i) = s_i$ independent from $X_{[(t-r)/\epsilon^2]}^i(1 - s_i)$. This proves item **I.1**.

Now we focus on

$$(\omega_{X_0^i}([(t-r)/\epsilon^2] + \tau + u_i) - \omega_{X_0^i}([(t-r)/\epsilon^2] + \tau))_{u_i \in [0, [M(\epsilon)/\epsilon^2]-1]} \quad (4.12)$$

Since $X_{[(t-r)/\epsilon^2]+1}^i$ is sampled from W_1 , independently from the previous steps, (4.12) is independent from $(\omega_{X_0^i}([(t-r)/\epsilon^2] + \tau) - \omega_{X_0^i}([(t-r)/\epsilon^2] + \tau - s_i))_{s_i \in [0,1]}$. This proves item **I.3**. Furthermore, since $M(\epsilon) \leq r$ we have

$$\frac{t-r}{\epsilon^2} + [M(\epsilon)/\epsilon^2] - 1 \leq \tau + N_\epsilon,$$

This means that for $u_i \in [0, [M(\epsilon)/\epsilon^2] - 1]$, the step $X_{N_\epsilon+1}^i$ (which is sampled using $\hat{\pi}_{N_\epsilon, N_\epsilon+1}^\epsilon$, defined in (3.9)) does not appear in the path

$$\omega_{X_0^i}([(t-r)/\epsilon^2] + \tau + u_i) - \omega_{X_0^i}([(t-r)/\epsilon^2] + \tau)_{u_i \in [0, [M(\epsilon)/\epsilon^2]-1]}.$$

Using (3.1) this implies that (4.12) is equal in distribution to $(\omega_{X_{[(t-r)/\epsilon^2]+1}^i}(u_i))_{u_i \in [0, [M(\epsilon)/\epsilon^2]-1]}$ (which is built using only $\hat{\pi}^{(\epsilon)}$). This proves item **I.2** and concludes the proof. \square

The lemma that we used in the calculation of the limit of $\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon]$ is the following:

Lemma 4.2. For $t > 0$ and for $(X_0, Y_0) \sim W_1 \times W_1$ and $(\omega_{X_0}, \omega_{Y_0})$ built from $\hat{\pi}^{(\epsilon)}$ we have that:

$$\left(\epsilon \omega_{X_0}([t/\epsilon^2]), \epsilon \omega_{X_0}([t/\epsilon^2]) \right)$$

converges in distribution to $(\mathcal{N}_1, \mathcal{N}_2)$ where $\mathcal{N}_i \in \mathbb{R}^2$ are two independent centered Gaussian random variables with covariance matrix $tI_{2 \times 2}$.

Proof. Since $(\omega_{X_0}, \omega_{Y_0})$ is a pair of independent paths each built from $\hat{\pi}^{(\epsilon)}$ it is enough to prove this for ω_{X_0} . Define:

$$N_t^\epsilon = \max\{i / T_i < [t/\epsilon^2]\}.$$

From similar arguments as in **Proposition A.1** and **A.2**, the random variable

$$\epsilon \sum_{k=0}^{[t/\epsilon^2]} \mathbf{X}_k^{(\epsilon)}$$

converges in distribution, as $\epsilon \rightarrow 0$, to a centered Gaussian random variable with covariance matrix $\frac{t}{\gamma} I_{2 \times 2}$. Furthermore, since $T_n/n \rightarrow 1/\gamma$ a.s as $n \rightarrow \infty$ we see that:

$$\epsilon \sum_{k=0}^{N_t^\epsilon - 1} \mathbf{X}_k^{(\epsilon)} \tag{4.13}$$

converges in distribution to a centered Gaussian random variable with covariance matrix $tI_{2 \times 2}$. Finally, from **Corollary A.1** it is easy to see that:

$$\epsilon |\omega_{X_0}([t/\epsilon^2]) - \sum_{k=0}^{N_t^\epsilon} \mathbf{X}_k^{(\epsilon)}| \lesssim \sup_{s \in [T_{N_t^\epsilon - 1}, T_{N_t^\epsilon}]} \epsilon |\omega_{X_0}([t/\epsilon^2]) - \omega_{X_0}(T_{N_t^\epsilon - 1})| \rightarrow 0, \tag{4.14}$$

a.s. as $\epsilon \rightarrow 0$. This completes the proof. \square

Remark 4.1. Observe that this implies that $\epsilon \omega_{X_0}([t/\epsilon^2])$ (where $X_0 \sim W_\tau$, $\tau < 1$) converges in distribution to a centered Gaussian random variable with covariance $tI_{2 \times 2}$. Indeed, observe that $\epsilon \omega_{X_0}(N_t^\epsilon)$ is equal to (4.13) where only $\mathbf{X}_0^{(\epsilon)}$ has a different distribution from $\mathbf{X}_k^{(\epsilon)}$, $k \geq 1$. Then by following the same arguments as in the proof above, we see that (4.14) is still valid for ω_{X_0} . So it converges to a Gaussian random variable with covariance matrix $tI_{2 \times 2}$. Finally if $(\omega_{X_0}, \omega_{Y_0})$ are as in the proof of **Lemma 4.1** they are independent and therefore $(\epsilon \omega_{X_0}([t/\epsilon^2]), \epsilon \omega_{X_0}([t/\epsilon^2]))$ converges in distribution to $(\mathcal{N}_1, \mathcal{N}_2)$.

Now we show that I_ϵ and $\exp(J_\epsilon)$ decouple asymptotically. This is done in the proof of the following lemma:

Lemma 4.3. For all $\hat{\beta} < \hat{\beta}_c(R)$ we have that

$$\hat{\mathbb{E}}_{B,t/\epsilon^2} \left[I_\epsilon \exp(J_\epsilon) \right] \rightarrow \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2} \right)^{-1} p_{t-r} \star g(y)^2,$$

as $\epsilon \rightarrow 0$.

Proof. Using the same notation as in the proof of **Lemma 4.1** we have that

$$\hat{\mathbb{E}}_{B,t/\epsilon^2} \left[I_\epsilon \exp(J_\epsilon) \right] = \hat{\mathbb{E}}_{B,t/\epsilon^2} \left[I_\epsilon(B^1, B^2) \exp(\tilde{J}_\epsilon(B^1, B^2) + \mathcal{J}_\epsilon(B^1, B^2)) \right],$$

where $\tilde{J}_\epsilon(B^1, B^2)$ and $\mathcal{J}_\epsilon(B^1, B^2)$ are defined in (4.6) and (4.7), respectively. Since $\mathcal{J}_\epsilon(B^1, B^2) \rightarrow 0$, as $\epsilon \rightarrow 0$, uniformly over the paths B^1, B^2 , it is sufficient to consider

$$\hat{\mathbb{E}}_{B,t/\epsilon^2} \left[I_\epsilon(B^1, B^2) \exp(\tilde{J}_\epsilon(B^1, B^2)) \right],$$

and calculate its limit as $\epsilon \rightarrow 0$.

From (3.13) this expectation is equal to:

$$\mathbb{E} \left[\mathcal{G}_\epsilon(\tilde{X}_{N_\epsilon}^1, \tilde{X}_{N_\epsilon}^2) I_\epsilon([\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1], [\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2]) \exp(\tilde{J}_\epsilon([\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1], [\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2])) \right],$$

where $(\tilde{X}_0^1, \dots, \tilde{X}_{N_\epsilon+1}^1)$ and $(\tilde{X}_0^2, \dots, \tilde{X}_{N_\epsilon+1}^2)$ are as in the proof of **Lemma 4.1**. Combining the fact that \mathcal{G}_ϵ converges to 1 uniformly and **Lemma 3.1** we see that there are constants $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon$ such that:

$$\begin{aligned} & \mathcal{A}_\epsilon \mathbb{E} \left[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})) \right] \\ & \leq \hat{\mathbb{E}}_{B,t/\epsilon^2} \left[I_\epsilon \exp(J_\epsilon) \right] \leq \\ & \mathcal{B}_\epsilon \mathbb{E} \left[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})) \right], \end{aligned} \quad (4.15)$$

where $\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}$ are as in the proof of **Lemma 4.1** and $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon \rightarrow 1$, as $\epsilon \rightarrow 0$.

The terms $I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$ and $\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$ are independent. More specifically, the term (4.9), that appears in $\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$, is independent from $I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$. Indeed, observe that $\boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau + u_1) - \boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau)$, $i = 1, 2$ is independent from the paths $\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}$ at all times before $[(t-r)/\epsilon^2] + \tau$, and specifically from all times that appear in $I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$. The other terms $\boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau) - \boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau - s_i)$ as mentioned in the comments after (4.11) are independent from $\boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau - 1)$ and from $\boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau - s_i) - \boldsymbol{\omega}_{X_0^i}([(t-r)/\epsilon^2] + \tau - 1)$ and therefore from $I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$ as well. Therefore, $I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})$ and $\exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}))$ are indeed independent.

Finally, from the proof of **Lemma 4.1**

$$\mathbb{E} \left[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \right] \rightarrow p_{t-r} \star g(y)^2$$

and

$$\mathbb{E} \left[\exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2})) \right] \rightarrow \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2} \right)^{-1},$$

as $\epsilon \rightarrow 0$. This, combined with (4.15), concludes the proof. \square

Now **Lemma 4.1** and **Lemma 4.3** combined gives us the following:

Lemma 4.4. As $\epsilon \rightarrow 0$ and for all $\hat{\beta} < \hat{\beta}_c(R)$:

$$\mathcal{F}_\epsilon(r, y, M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2) \rightarrow v_{eff}^2(\hat{\beta})p_{t-r} \star g(y)^2,$$

where

$$v_{eff}^2(\hat{\beta}) = \left(1 - \frac{\hat{\beta}^2}{\hat{\beta}_c(R)^2}\right)^{-1} \cdot \|R\|_1.$$

Proof. Recall the definition of $\mathcal{F}_\epsilon(r, y, M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2)$ in (4.1) and of $J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2; x_1, x_2, s_1, s_2)$ in (4.3), where we emphasize the dependence of J_ϵ on x_1, x_2, s_1, s_2 . From **Lemma 4.3**, we see that if

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon e^{J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2; x_1, x_2, s_1, s_2)}]$$

is bounded for all sufficiently small ϵ and uniformly in x_1, x_2, s_1, s_2 then, from the dominated convergence theorem⁴, $\mathcal{F}_\epsilon(r, y) \rightarrow v_{eff}^2(\hat{\beta})p_{t-r} \star g(y)^2$ as $\epsilon \rightarrow 0$. We will prove that

$$\sup_{\epsilon \in (0,1)} \hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon e^{J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2; x_1, x_2, s_1, s_2)}] \lesssim 1, \quad (4.16)$$

uniformly in x_1, x_2, s_1, s_2 .

From (4.15) we have

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon e^{J_\epsilon(M(\epsilon)/\epsilon^2, M(\epsilon)/\epsilon^2; x_1, x_2, s_1, s_2)}] \lesssim \mathbb{E}\left[I_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}; x_1, x_2, s_1, s_2))\right].$$

Since g has compact support we see that:

$$\mathbb{E}\left[\tilde{I}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}) \exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}; x_1, x_2, s_1, s_2))\right] \lesssim \mathbb{E}\left[\exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}; x_1, x_2, s_1, s_2))\right].$$

Recall that, from (4.10)

$$\begin{aligned} & \mathbb{E}\left[\exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}; x_1, x_2, s_1, s_2))\right] = \\ & \int_{\mathbb{R}^2} \int_{\mathbb{R}^2} \mathbb{E}\left[\exp\left(\frac{\beta^2}{\log \epsilon^{-1}} \int_0^{[M(\epsilon)/\epsilon^2]} \int_0^{[M(\epsilon)/\epsilon^2]} R\left(u_1 - u_2, x_1 - x_2 + y_1 + y_2 + \omega_{b^1}(u_1) - \omega_{b^2}(u_2)\right) du_1 du_2\right)\right] p_{s_1}(y_1) p_{s_2}(y_2) dy_1 dy_2, \end{aligned} \quad (4.17)$$

where b^1, b^2 are two independent Brownian motions in Ω_1 .

Consider the term

$$\mathbb{E}\left[\exp\left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{[M(\epsilon)/\epsilon^2]} \int_0^{[M(\epsilon)/\epsilon^2]} R\left(u_1 - u_2, x_1 - x_2 + y_1 + y_2 + \omega_{b^1}(u_1) - \omega_{b^2}(u_2)\right) du_1 du_2\right)\right].$$

Then from **Proposition 3.2** we have

$$\sup_{\epsilon \in (0,1)} \mathbb{E}\left[\exp\left(\frac{\hat{\beta}^2}{\log \epsilon^{-1}} \int_0^{[M(\epsilon)/\epsilon^2]} \int_0^{[M(\epsilon)/\epsilon^2]} R\left(u_1 - u_2, x_1 - x_2 + y_1 + y_2 + \omega_{b^1}(u_1) - \omega_{b^2}(u_2)\right) du_1 du_2\right)\right] \lesssim 1,$$

⁴Recall that ϕ and ψ are compactly supported.

uniformly over x_1, x_2, y_1, y_2 . From (4.17) we get that

$$\sup_{\epsilon \in (0,1)} \mathbb{E}[\exp(\tilde{J}_\epsilon(\boldsymbol{\omega}_{X_0^1}, \boldsymbol{\omega}_{X_0^2}; x_1, x_2, y_1, y_2, s_1, s_2))] \lesssim 1,$$

uniformly over $x_1, x_2, y_1, y_2, s_1, s_2$, which in turn proves the uniform bound (4.16). This concludes the proof. \square

Finally, we prove the uniform bound (1.25):

Lemma 4.5. *Let $\hat{\beta} < \hat{\beta}_c(R)$ and $k > 0$. Then*

$$|\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2)| \lesssim (1 \wedge |y|^{-k}),$$

where the implied constant depends only on k .

Proof. From the proof of **Lemma 4.4** and the observations at the start of this section, it is easy to infer that for any $p > 1$ such that $p\hat{\beta} < \hat{\beta}_c(R)$

$$\sup_{\epsilon \in (0,1)} \hat{\mathbb{E}}_{B,t/\epsilon^2}[e^{pJ_\epsilon((M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2; x_1, x_2, s_1, s_2))}] \lesssim 1,$$

Hence, from Hölder's inequality

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) \lesssim \int_{\mathbb{R}^4} \int_{[0,1]^2} \hat{\mathbb{E}}_{B,t/\epsilon^2}[I_\epsilon^q]^{1/q} \prod_{i=1}^2 \phi(s_i) \psi(x_i) d\bar{s} d\bar{x},$$

where $1/p + 1/q = 1$. This implies that:

$$\mathcal{F}_\epsilon(r, y, M_1(\epsilon)/\epsilon^2, M_2(\epsilon)/\epsilon^2) \lesssim \int_{\mathbb{R}^2} \int_{[0,1]} \hat{\mathbb{E}}_{B,t/\epsilon^2}[g(\epsilon x - \epsilon B_{(t-r)/\epsilon^2-s} + y)^q]^{1/q} \phi(s) \psi(x) ds dx,$$

since g is compactly supported. Observe that for all k :

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[\mathbf{1}_{|\epsilon B_{(t-r)/\epsilon^2-s}| > M}] \lesssim \frac{1}{M^{2k}},$$

where the implied constant depends on k . The proof of this inequality is the same as in [GRZ18] (look at **Lemma 5.3** of the same paper) only we use **Corollary A.1** whenever they use **Lemma A.2**. With this inequality and since g and compactly supported:

$$\hat{\mathbb{E}}_{B,t/\epsilon^2}[g(\epsilon x - \epsilon B_{(t-r)/\epsilon^2-s} + y)^q]^{1/q} \lesssim 1 \wedge \frac{1}{|y|^k},$$

which implies (1.25). \square

As mentioned, **Lemma 4.4** and **Lemma 4.5** prove **Proposition 1.4**. We also have the following corollary:

Corollary 4.1. *For all $\hat{\beta} < \hat{\beta}_c(R)$ we have that*

$$\mathbf{Var}\left(\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx\right)$$

converges, as $\epsilon \rightarrow 0$, to

$$\mathbf{Var}\left(\int_{\mathbb{R}^2} \mathcal{U}(t, x) g(x) dx\right),$$

where \mathcal{U} is the solution to the Edwards-Wilkinson equation (1.3), with effective diffusivity equal to $I_{2 \times 2}$ and effective variance v_{eff} defined as in (1.6).

Proof. We have

$$\hat{\beta}^2 v_{eff}^2(\hat{\beta}) \int_0^t \int_{\mathbb{R}^2} p_{t-r} \star g(y)^2 dy dr = \mathbf{Var} \left(\int_{\mathbb{R}^2} \mathcal{U}(t, x) g(x) dx \right).$$

We want to apply **Proposition 1.4** to the formula for the variance that we got from **Proposition 1.3**. As mentioned in the remark after **Proposition 1.3** that formula has some modifications. Recall that s_1, s_2 in (4.1) will lie in $[0, (t-r)/\epsilon^2]$ and u_1, u_2 will lie in $[-(t-r)/\epsilon^2, M_1(\epsilon)]$, $[-(t-r)/\epsilon^2, M_2(\epsilon)]$. Let $\mathcal{F}_\epsilon(r, y, M_1(\epsilon), M_2(\epsilon))$ be defined by (4.1) but with these modifications. Then we have that

$$\mathbf{Var} \left(\sqrt{\log \epsilon^{-1}} \int_{\mathbb{R}^2} e^{-\zeta_{t/\epsilon^2}^{(\epsilon)}} (u_\epsilon(t, x) - \mathbf{E}[u_\epsilon(t, x)]) g(x) dx \right) = \hat{\beta} \int_0^t \int_{\mathbb{R}^2} \mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2) dy dr. \quad (4.18)$$

As argued in **Section 2** the differences between $\mathcal{F}_\epsilon(r, y, M_1(\epsilon), M_2(\epsilon))$ and $\mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2)$ come up only when $t-r \leq \epsilon^2$. Therefore, for fixed r, y

$$\mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2) = \mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2),$$

for all ϵ small enough. This proves that $\mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2) \rightarrow v_{eff}^2(\hat{\beta}) p_{t-r} \star g(y)^2$, as $\epsilon \rightarrow 0$. From this observation and (4.18), we see that to prove the corollary we need to bound $\mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2)$ by an integrable function so we can apply the dominated convergence theorem.

More specifically, we argue that $\mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2)$ satisfies the uniform bound (1.25). Indeed, observe that if $\mathbf{J}_\epsilon(r/\epsilon^2, r/\epsilon^2; x_1, x_2, s_1, s_2)$ is defined by (4.3) but with the range of u_1, u_2 changed to $[-(t-r)/\epsilon^2, r/\epsilon^2]$ then, since $r \in (0, t)$ we have

$$\mathbf{J}_\epsilon(r/\epsilon^2, r/\epsilon^2; x_1, x_2, s_1, s_2) \leq J'_\epsilon(t/\epsilon^2, t/\epsilon^2; x_1, x_2, s_1, s_2),$$

for all ϵ , where $J'_\epsilon(t/\epsilon^2, t/\epsilon^2; x_1, x_2, s_1, s_2)$ is also defined by (4.3) but with the range of u_1, u_2 changed to $[-t/\epsilon^2, t/\epsilon^2]$. For all ϵ small enough $J'_\epsilon(t/\epsilon^2, t/\epsilon^2; x_1, x_2, s_1, s_2) = J_\epsilon(t/\epsilon^2, t/\epsilon^2; x_1, x_2, s_1, s_2)$, since $R_\phi(u_1, u_2) = 0$ when $u_1 \leq -1$ or $u_2 \leq -1$. This proves that for $p > 1$ such that $p\hat{\beta} < \hat{\beta}_c(R)$ we have

$$\sup_{\epsilon \in (0,1)} \hat{\mathbb{E}}_{B, t/\epsilon^2} [e^{p\mathbf{J}_\epsilon(r/\epsilon^2, r/\epsilon^2; x_1, x_2, s_1, s_2)}] \lesssim \sup_{\epsilon \in (0,1)} \hat{\mathbb{E}}_{B, t/\epsilon^2} [e^{pJ_\epsilon(t/\epsilon^2, t/\epsilon^2; x_1, x_2, s_1, s_2)}] \lesssim 1,$$

uniformly over x_1, x_2, s_1, s_2 and r . From this estimate, and since g, ϕ are compactly supported, we get that

$$\mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2) \lesssim \int_{\mathbb{R}^2} \int_{[0, (t-r)/\epsilon^2]^2} \hat{\mathbb{E}}_{B, t/\epsilon^2} [g(\epsilon x - \epsilon B_{(t-r)/\epsilon^2 - s}^i + y)^q]^{1/q} \phi(s) \psi(x) ds dx,$$

where $1/p + 1/q = 1$. Now, by arguing in the same way as we did in the proof of **Lemma 4.5** we can prove that the uniform bound (1.25) is satisfied by $\mathcal{F}_\epsilon(r, y, r/\epsilon^2, r/\epsilon^2)$, for all ϵ small enough. This concludes the proof. \square

A Estimates on the Total Path Increments

In this appendix, we study the properties of the path increments between regeneration times. The main result is a local central limit theorem for sums of the total path increments that makes it possible to estimate the limiting distribution of additive functionals of $\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)$. The latter is done in the next appendix.

Proposition A.1. *The random variable $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ has zero mean. Moreover, as $\epsilon \rightarrow 0$, $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ converges in total variation to $\sum_{j=0}^{\theta-1} \mathcal{N}_j$ where:*

- $(\mathcal{N}_j)_{j \in \mathbb{N}_0}$ are i.i.d. Gaussian random variables with covariance matrix $2 \cdot I_{2 \times 2}$.
- $\theta \sim \text{Geo}(\gamma)$ and is independent from the $(\mathcal{N}_j)_{j \in \mathbb{N}}$.

Moreover, $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ has exponential moments uniformly in ϵ , i.e. for all $\boldsymbol{\lambda} \in \mathbb{R}^2$ small enough

$$\sup_{\epsilon > 0} \mathbb{E}[\exp(\boldsymbol{\lambda} \cdot (\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}))] \lesssim 1. \quad (\text{A.1})$$

Finally, the covariance matrix of $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ is bounded in ϵ and converges to $\frac{2}{\gamma} I_{2 \times 2}$ as $\epsilon \rightarrow 0$.

Proof. By symmetry, $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ has zero mean. In fact, it is easy to see that $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ is a symmetric random variable.

First, we show that $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ converges to a Gaussian. Inequality (3.24) implies that

$$\sum_{N=1}^{\infty} \gamma(1-\gamma)^{N-1} \mathcal{A}_\epsilon^N \mathbb{P}\left(\sum_{j=0}^{N-1} \mathcal{N}_j \in A\right) \leq \mathbb{P}(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)} \in A) \leq \sum_{N=1}^{\infty} \gamma(1-\gamma)^{N-1} \mathcal{B}_\epsilon^N \mathbb{P}\left(\sum_{j=0}^{N-1} \mathcal{N}_j \in A\right), \quad (\text{A.2})$$

where \mathcal{N}_j are i.i.d. Gaussian random variables with covariance matrix $2 \cdot I_{2 \times 2}$, and where $\mathcal{A}_\epsilon, \mathcal{B}_\epsilon \rightarrow 1$ as $\epsilon \rightarrow 0$. From dominated convergence

$$\lim_{\epsilon \rightarrow 0} \mathbb{P}\left(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)} \in A\right) = \mathbb{P}\left(\sum_{j=0}^{\theta-1} \mathcal{N}_j \in A\right).$$

We prove that this convergence happens uniformly over A . Observe that (A.2) implies that for $\mathcal{C}_\epsilon(N) = (\mathcal{B}_\epsilon^N - 1) \vee (1 - \mathcal{A}_\epsilon^N)$ ⁵ and for a generic $\theta \sim \text{Geo}(\gamma)$, independent from $(\mathcal{N}_j)_{j \in \mathbb{N}}$, we have

$$\left| \mathbb{P}(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)} \in A) - \mathbb{P}\left(\sum_{j=0}^{\theta-1} \mathcal{N}_j \in A\right) \right| \leq \sum_{N=1}^{\infty} \gamma(1-\gamma)^{N-1} \mathcal{C}_\epsilon(N) \mathbb{P}\left(\sum_{j=0}^{N-1} \mathcal{N}_j \in A\right).$$

For all Borel sets $A \subseteq \mathbb{R}^2$, the right-hand side of the above inequality is bounded above by

$$\sum_{N=1}^{\infty} \gamma(1-\gamma)^{N-1} \mathcal{C}_\epsilon(N),$$

which converges to 0 as $\epsilon \rightarrow 0$. Therefore, $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ converges in total variation to $\sum_{j=0}^{\theta-1} \mathcal{N}_j$.

We use (3.24) to show (A.1). We choose $F(\mathbf{x}) = \exp(\boldsymbol{\lambda} \cdot \mathbf{x}/c)$. Then, from (3.24), there is a constant \mathcal{B}_ϵ :

$$\mathbb{E}[F(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)})] \leq \sum_{N=1}^{\infty} \mathcal{B}_\epsilon^N \gamma(1-\gamma)^{N-1} \mathbb{E}_{\mathbf{X}^{(N)}, \mathbf{Y}^{(N)}}[F(\mathbf{X}^{(N)} - \mathbf{Y}^{(N)})],$$

where $(\mathbf{X}^{(N)}, \mathbf{Y}^{(N)})$ are as in (3.24). Since $\mathbf{X}^{(N)}, \mathbf{Y}^{(N)}$ are Gaussian, we get:

$$\mathbb{E}[F(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)})] \leq \sum_{N=1}^{\infty} \mathcal{B}_\epsilon^N \gamma(1-\gamma)^{N-1} e^{N|\boldsymbol{\lambda}|^2},$$

which is finite for all $\epsilon > 0$, for $\boldsymbol{\lambda}$ small enough (recall that $\mathcal{B}_\epsilon \rightarrow 1$ as $\epsilon \rightarrow 0$).

The last part of the proposition follows from a similar calculation as above, and tightness. \square

⁵Observe that $\mathcal{A}_\epsilon \leq 1 \leq \mathcal{B}_\epsilon$.

Remark A.1. Inequality (A.2) implies that $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ has a density with respect to the Lebesgue measure. This observation, combined with **Proposition A.1**, implies that the characteristic function of $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ converges uniformly to the corresponding characteristic function of $\sum_{j=0}^{\theta-1} \mathcal{N}_j$. This implies that for all ϵ small enough the characteristic function of $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ is integrable and, therefore the density of $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ is in $L^\infty(\mathbb{R}^2)$. It is also easy to see that all mixed moments of $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ are bounded in ϵ .

The observations in **Remark A.1** lead us to the following central limit theorem (and its local version):

Proposition A.2. Let $(X_0, Y_0) \sim W_1 \times W_1$. Then as $k \rightarrow \infty$ and $\epsilon \rightarrow 0$, the random variable

$$\frac{1}{\sqrt{k}}(\omega_{X_0}(T_k) - \omega_{Y_0}(T_k))$$

converges in distribution to a normal random variable in \mathbb{R}^2 with covariance matrix $\frac{2}{\gamma}I_{2 \times 2}$. Furthermore, if $F_k^{(\epsilon)}$ is the density of $\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)$ with respect to the Lebesgue measure, then

$$\sup_{\mathbf{x} \in \mathbb{R}^2} |kF_k^{(\epsilon)}(\mathbf{x}) - \frac{\gamma}{4\pi} \exp(-\gamma|\mathbf{x}|^2/4k)| \rightarrow 0, \quad (\text{A.3})$$

as $k \rightarrow \infty$ and $\epsilon \rightarrow 0$.

Proof. Recall that $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ has zero mean. The central limit theorem is proved via a standard calculation of the characteristic function. In particular, from (3.18) we have

$$\phi_{\frac{1}{\sqrt{k}}(\omega_{X_0}(T_k) - \omega_{Y_0}(T_k))}(\boldsymbol{\eta}) = (\phi_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\boldsymbol{\eta}/\sqrt{k}))^k,$$

where ϕ_X denotes the characteristic function of the random variable X .

As we observed in **Remark A.1**, all mixed moments of the vector $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ are bounded in ϵ . Therefore, Taylor's expansion yields

$$\phi_{\frac{1}{\sqrt{k}}(\omega_{X_0}(T_k) - \omega_{Y_0}(T_k))}(\boldsymbol{\eta}) = \left(1 - \frac{\boldsymbol{\eta}^* \cdot \mathbb{E}[(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)})(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)})^*] \cdot \boldsymbol{\eta}}{2k} + \mathbf{R}_{\epsilon,k}(\boldsymbol{\eta}) \right)^k,$$

where the error term $\mathbf{R}_{\epsilon,k}(\boldsymbol{\eta})$ satisfies the bound

$$\sup_{\epsilon \in (0,1)} |\mathbf{R}_{\epsilon,k}(\boldsymbol{\eta})| = o(1/k)|\boldsymbol{\eta}|^2.$$

From **Proposition A.1**

$$\mathbb{E}[(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)})(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)})^*] \rightarrow 2/\gamma I_{2 \times 2},$$

as $\epsilon \rightarrow 0$. Therefore:

$$\phi_{\frac{1}{\sqrt{k}}(\omega_{X_0}(T_k) - \omega_{Y_0}(T_k))}(\boldsymbol{\eta}) \rightarrow \exp(-|\boldsymbol{\eta}|^2/\gamma), \quad (\text{A.4})$$

as $k \rightarrow \infty$, $\epsilon \rightarrow 0$.

If $F_k^{(\epsilon)}(\mathbf{x})$ is the probability density of $\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)$, then by Fourier inversion,

$$F_k^{(\epsilon)}(\mathbf{x}) = \frac{1}{(2\pi)^2} \int_{\mathbb{R}^2} \exp(-i\mathbf{x} \cdot \boldsymbol{\eta}) \phi_{\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)}(\boldsymbol{\eta}) d\boldsymbol{\eta}. \quad (\text{A.5})$$

To prove (A.3), we go through the following steps:

F.1 $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ has a density $f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}} \in L^2(\mathbb{R}^2)$. There is an $\epsilon'_0 > 0$, such that

$$\sup_{\epsilon \leq \epsilon'_0} \int_{\mathbb{R}^2} f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\mathbf{x})^2 d\mathbf{x} \lesssim 1. \quad (\text{A.6})$$

F.2 There is an ϵ''_0 and an A such that

$$F_k^{(\epsilon)}(\mathbf{x}) = \frac{1}{(2\pi)^2 k} \int_{|\boldsymbol{\eta}| \leq A\sqrt{k}} \exp(-i(\mathbf{x} \cdot \boldsymbol{\eta})/\sqrt{k}) \phi_{\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)}(\boldsymbol{\eta}/\sqrt{k}) d\boldsymbol{\eta} + \mathcal{R}_{\epsilon,k}, \quad (\text{A.7})$$

with the error term $\mathcal{R}_{\epsilon,k}$ satisfying the bound

$$\sup_{\epsilon \leq \epsilon''_0} |\mathcal{R}_{\epsilon,k}| \lesssim 2^{-k}.$$

F.3 As $k \rightarrow \infty$ and $\epsilon \rightarrow 0$

$$\int_{|\boldsymbol{\eta}| \leq A\sqrt{k}} |\phi_{\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)}(\boldsymbol{\eta}/\sqrt{k}) - \exp(-|\boldsymbol{\eta}|^2/\gamma)| d\boldsymbol{\eta} \rightarrow 0. \quad (\text{A.8})$$

F.4 As $k \rightarrow \infty$,

$$\frac{1}{(2\pi)^2} \int_{|\boldsymbol{\eta}| \leq A\sqrt{k}} \exp(-i(\mathbf{x} \cdot \boldsymbol{\eta})/\sqrt{k}) \exp(-|\boldsymbol{\eta}|^2/\gamma) d\boldsymbol{\eta} = \frac{\gamma}{4\pi} \exp(-\gamma|\mathbf{x}|^2/4k) + o(1). \quad (\text{A.9})$$

Equations (A.7)-(A.9) prove that:

$$F_k^{(\epsilon)}(\mathbf{x}) = \frac{\gamma}{4\pi k} \exp(-\gamma|\mathbf{x}|^2/4k) + \mathcal{R}'_{\epsilon,k} \cdot 1/k,$$

where the error term $\mathcal{R}'_{\epsilon,k}$ goes to 0, as $k \rightarrow \infty$ and $\epsilon \rightarrow 0$. This proves (A.3).

To prove (A.6), recall that from **Remark A.1** $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ has a density with respect to the Lebesgue measure, which we denote by $f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}$. From (3.24)

$$\begin{aligned} \int_{\mathbb{R}^2} f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\mathbf{x})^2 d\mathbf{x} &= \mathbb{E}_{\mathbf{X}_1^{(\epsilon)}, \mathbf{Y}_1^{(\epsilon)}} [f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)})] \leq \\ &= \sum_{N=1}^{\infty} B_\epsilon^N \gamma (1-\gamma)^{N-1} \mathbb{E}_{\mathbf{X}^{(N)}, \mathbf{Y}^{(N)}} [f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\mathbf{X}^{(N)} - \mathbf{Y}^{(N)})], \end{aligned}$$

where $(\mathbf{X}^{(N)}, \mathbf{Y}^{(N)})$ are as in (3.24) and $B_\epsilon \rightarrow 1$ as $\epsilon \rightarrow 0$. Since the random variables $\mathbf{X}^{(N)}, \mathbf{Y}^{(N)}$ are Gaussian, we have:

$$\int_{\mathbb{R}^2} f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\mathbf{x})^2 d\mathbf{x} \lesssim \sum_{N=1}^{\infty} \frac{1}{N} B_\epsilon^N \gamma (1-\gamma)^{N-1} \int_{\mathbb{R}^2} f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\mathbf{x}) d\mathbf{x} \lesssim 1,$$

for all $\epsilon \leq \epsilon'_0$, where ϵ'_0 is such that $\sup_{\epsilon \leq \epsilon'_0} B_\epsilon < \frac{1}{(1+\gamma)(1-\gamma)}$. This proves (A.6).

To prove (A.7) we do a change of variables $\boldsymbol{\eta} \rightarrow \boldsymbol{\eta}/\sqrt{k}$ in the integral in (A.5). Then, we split the integral over the regions $|\boldsymbol{\eta}| > A\sqrt{k}$ and $|\boldsymbol{\eta}| \leq A\sqrt{k}$. The proof of (A.7) reduces to proving that there is an A and an ϵ''_0 such that:

$$\sup_{\epsilon \leq \epsilon''_0} \left| \frac{1}{k} \int_{|\boldsymbol{\eta}| > A\sqrt{k}} \phi_{\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)}(\boldsymbol{\eta}/\sqrt{k}) d\boldsymbol{\eta} \right| \lesssim 2^{-k}. \quad (\text{A.10})$$

We have that:

$$\frac{1}{k} \int_{|\boldsymbol{\eta}| > A\sqrt{k}} |\phi_{\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)}(\boldsymbol{\eta}/\sqrt{k})| d\boldsymbol{\eta} \leq \sup_{|\boldsymbol{\eta}| > A} |\phi_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\boldsymbol{\eta})|^{k-2} \frac{1}{k} \int_{\mathbb{R}^2} |\phi_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\boldsymbol{\eta}/\sqrt{k})|^2 d\boldsymbol{\eta}. \quad (\text{A.11})$$

The second term in the right-hand side of (A.11), after a change of variables $\boldsymbol{\eta} \rightarrow \sqrt{k}\boldsymbol{\eta}$ and Plancierel's theorem, is equal to $(2\pi)^2 \int_{\mathbb{R}^2} f_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\mathbf{x})^2 d\mathbf{x}$ which, from (A.6), is bounded in $\epsilon \leq \epsilon'_0$.

The first term on the right-hand side of (A.11) is exponentially small. Indeed, from **Proposition A.1**, for all $\boldsymbol{\eta}$

$$\phi_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\boldsymbol{\eta}) \rightarrow \phi_{\mathbf{Y}}(\boldsymbol{\eta}),$$

where $\mathbf{Y} = \sum_{i=0}^{\theta-1} \mathcal{N}_i$, with θ and $(\mathcal{N}_i)_{i \in \mathbb{N}_0}$ as in **Proposition A.1**. This convergence happens uniformly in $\boldsymbol{\eta}$, since $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$, \mathbf{Y} are absolutely continuous with respect to the Lebesgue measure and $\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}$ converges to \mathbf{Y} in total variation, as $\epsilon \rightarrow 0$. A straightforward calculation shows that:

$$\phi_{\mathbf{Y}}(\boldsymbol{\eta}) = \frac{\gamma}{e^{|\boldsymbol{\eta}|^2/2} - (1 - \gamma)}.$$

Therefore, there is an A such that $\sup_{|\boldsymbol{\eta}| > A} |\phi_{\mathbf{Y}}(\boldsymbol{\eta})| \leq 1/4$. This implies that, there is an $\epsilon''_0 \leq \epsilon'_0$ such that for all $\epsilon \leq \epsilon''_0$

$$\sup_{|\boldsymbol{\eta}| \geq A} |\phi_{\mathbf{X}_1^{(\epsilon)} - \mathbf{Y}_1^{(\epsilon)}}(\boldsymbol{\eta})| \leq 1/2,$$

which proves (A.10).

The proof of (A.8) goes along the same lines as the proofs of local limit theorems found in [LL10]. We skip the details. Finally, Gaussian integration proves (A.9) and this concludes the proof. \square

Finally, we are going to need the following corollary. It is the analog of **Lemma A.2** from [GRZ18]:

Corollary A.1. *Let $(X_k, Y_k)_{k \in \mathbb{N}_0}$ be a sequence of random variables generated by the transition probability $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1 \times W_1}{1 - \gamma}$, where $(X_0, Y_0) \sim \hat{\mathbb{P}}_1^{(\epsilon)}(dx_0) \times \hat{\mathbb{P}}_1^{(\epsilon)}(dy_0)$. Then $\mathbb{E}[X_k(1)] = 0$. Moreover, there is a $c_1 > 0$ such that*

$$\mathbb{P}[\sup_{s \in [0,1]} |X_k(s)| > t] \lesssim e^{-c_1 t^2}. \quad (\text{A.12})$$

Additionally, if $\theta \sim \text{Geo}(\gamma)$ is independent from $(X_k)_{k \in \mathbb{N}}$, then for some $c_2 > 0$:

$$\mathbb{P}\left[\sum_{k=1}^{\theta} \sup_{s \in [0,1]} |X_k(s)| > t\right] \lesssim e^{-c_2 t}. \quad (\text{A.13})$$

Therefore, $\sum_{k=1}^{\theta} \sup_{s \in [0,1]} |X_k(s)|$ has exponential tails, uniformly in ϵ . Finally

$$\mathbb{P}[\sup_{s \leq \theta} |B(s)| > t] \lesssim e^{-c_3 t}, \quad (\text{A.14})$$

for some $c_3 > 0$, where $B \in C([0, \infty))$ is the path built from $(X_k, Y_k)_{k \in \mathbb{N}_0}$.

Remark A.2. *Observe that the marginal law of $(X_k)_{k \in \mathbb{N}_0}$ is the same as the law of $(Z_k)_{\mathbb{N}_0}$, where this sequence is generated by $\frac{\hat{\pi}^{(\epsilon)} - \gamma W_1}{1 - \gamma}$ with $Z_0 \sim \hat{\mathbb{P}}_1^{(\epsilon)}(dz_0)$.*

Proof. The proof of $\mathbb{E}[X_k(1)] = 0$ is the same as in [GRZ18] (look at the proof of Lemma A.2).

Inequality (A.12) is a straightforward application of **Lemma 3.1** with $F(x) = \sup_{s \in [0,1]} |x(s)|$ and the corresponding bound

$$\mathbb{P}\left[\sup_{s \in [0,1]} |B(s)| > t\right] \lesssim e^{-c_1 t^2},$$

where B is a Brownian motion.

For (A.13), we condition on θ and apply **Lemma 3.1** to $F(x) = \sum_{k=1}^N \sup_{s \in [0,1]} |x_i(s)|$. If B^1, \dots, B^N are independent Brownian motions then

$$\mathbb{P}\left[\sum_{k=1}^{\theta} \sup_{s \in [0,1]} |X_k(s)| > t\right] \lesssim \sum_{N=1}^{\infty} \gamma(1-\gamma)^{N-1} \mathbb{P}\left[\sum_{k=1}^N \sup_{s \in [0,1]} |B^k(s)| > t\right].$$

We have that

$$\mathbb{P}\left[\sum_{k=1}^N \sup_{s \in [0,1]} |B^k(s)| > t\right] \leq e^{-ct} \mathbb{E}\left[\exp\left(c \sum_{k=1}^N \sup_{s \in [0,1]} |B^k(s)|\right)\right].$$

Therefore,

$$\mathbb{P}\left[\sum_{k=1}^N \sup_{s \in [0,1]} |X_k(s)| > t\right] \lesssim \sum_{N=1}^{\infty} \gamma(1-\gamma)^{N-1} e^{-ct} C^N,$$

for $C = \mathbb{E}[\exp(c \sup_{s \in [0,1]} |B^1(s)|)]$. We can conclude by taking c small enough.

Finally, (A.14) is proved similarly. \square

B Kallianpur-Robbins Estimates

Let $(X_0, Y_0) \sim W_1 \times W_1$ and $S_n = \omega_{X_0}(T_n) - \omega_{Y_0}(T_n)$. Recall that, from **Proposition A.2**,

$$F_n^{(\epsilon)}(x) = \frac{\gamma}{4\pi k} e^{-\gamma|x|^2/4k} + \mathcal{R}_{\epsilon,k} \cdot 1/k, \quad (\text{B.1})$$

for all $x \in \mathbb{R}$, where $\mathcal{R}_{\epsilon,k} \rightarrow 0$ as $k \rightarrow \infty$, $\epsilon \rightarrow 0$. These precise asymptotics for the density $F_n^{(\epsilon)}$ allow us to prove results on the occupation times of S_n .

We will make use of the following elementary estimates. Here, $M(\epsilon)$ is such that $\log M(\epsilon) / \log \epsilon^{-1} \rightarrow 0$ as $\epsilon \rightarrow 0$. We have

$$\frac{1}{(\log \epsilon^{-1})^p} \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{1}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})} \rightarrow 1, \quad (\text{B.2})$$

as $\epsilon \rightarrow 0$ and for $j = 1, \dots, p$

$$\frac{1}{(\log \epsilon^{-1})^p} \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{\mathcal{R}_{\epsilon, k_j - k_{j-1}} (1 + \mathcal{R}_{\epsilon, k_j})}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})} = o(1), \quad (\text{B.3})$$

as $\epsilon \rightarrow 0$, where $\mathcal{R}_{\epsilon,k}$ is as in (B.1). Similarly, for $j = 1, \dots, p$

$$\frac{1}{(\log \epsilon^{-1})^p} \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{(k_j - k_{j-1})^{-1}}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})} = o(1), \quad (\text{B.4})$$

as $\epsilon \rightarrow 0$.

The proof of **Proposition B.1** is similar to the proofs of the results in [KR54].

Proposition B.1. Let $M(\epsilon)$ be such that $\log M(\epsilon)/\log \epsilon^{-1} \rightarrow 0$ as $\epsilon \rightarrow 0$. Also let $(H_\epsilon)_{\epsilon>0} \subseteq L^1(\mathbb{R}^{2p})$ be a family of positive and bounded functions, such that for all ϵ , $\|H_\epsilon\|_1 = A$, for some constant A and such that

$$\lim_{M \rightarrow \infty} \sup_{\epsilon > 0} \int_{|x| \geq M} H_\epsilon(\mathbf{x}) d\mathbf{x} \rightarrow 0. \quad (\text{B.5})$$

For $S_n = \omega_{X_0}(T_n) - \omega_{Y_0}(T_n)$,

$$\left(\frac{4\pi}{\gamma \log \epsilon^{-1}} \right)^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \mathbb{E}[H_\epsilon(S_{k_1}, \dots, S_{k_p})] \rightarrow A, \quad (\text{B.6})$$

as $\epsilon \rightarrow 0$.

Proof. To prove (B.6) we write the expectation inside the sum explicitly. We have

$$\begin{aligned} & \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \mathbb{E}[H_\epsilon(S_{k_1}, \dots, S_{k_p})] = \\ & \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \int_{\mathbb{R}^{2p}} H_\epsilon(x_1, x_2, \dots, x_p) F_{k_1}^{(\epsilon)}(x_1) F_{k_2 - k_1}^{(\epsilon)}(x_2 - x_1) \dots F_{k_p - k_{p-1}}^{(\epsilon)}(x_p - x_{p-1}) dx_1 \dots dx_p. \end{aligned} \quad (\text{B.7})$$

We successively replace $F_{k_i - k_{i-1}}^{(\epsilon)}$, $i = 1, \dots, p$, by its asymptotic expansion (B.1). Adopting the notation in [KR54], (B.7) is equal to

$$\left(\frac{\gamma}{4\pi} \right)^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{\int_{\mathbb{R}^{2p}} H_\epsilon(x_1, x_2, \dots, x_p) e^{-Q_{k_1, \dots, k_p}(x_1, \dots, x_p)} dx_1 \dots dx_p}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})} + \mathcal{E}_\epsilon, \quad (\text{B.8})$$

where

$$Q_{k_1, \dots, k_p}(x_1, \dots, x_p) := \gamma \sum_{i=1}^p \frac{|x_i - x_{i-1}|^2}{4(k_i - k_{i-1})}, \quad (\text{B.9})$$

with $x_0 = 0$, $k_0 = 0$. The error term \mathcal{E}_ϵ is equal to

$$\sum_{j=1}^p \left(\frac{\gamma}{4\pi} \right)^j \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \mathcal{R}_{\epsilon, k_j - k_{j-1}} \int_{\mathbb{R}^{2p}} H_\epsilon(x_1, x_2, \dots, x_p) \frac{e^{-Q_{k_1, \dots, k_j}(x_1, \dots, x_j)}}{\prod_{i=1}^j (k_i - k_{i-1})} \left(\prod_{i>j}^p F_{k_i - k_{i-1}}^{(\epsilon)}(x_i - x_{i-1}) \right) d\mathbf{x}. \quad (\text{B.10})$$

We need to estimate both of the terms in (B.8). First, we deal with the error term (B.10). Observe that (B.1) implies that $F_n^{(\epsilon)}(x) \leq (1 + \mathcal{R}_{\epsilon, n})/n$. Using this bound, we see that the error term is

$$\leq \sum_{j=1}^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \int_{\mathbb{R}^{2p}} \mathcal{R}_{\epsilon, k_j - k_{j-1}} (1 + \mathcal{R}_{\epsilon, k_j}) \frac{H_\epsilon(x_1, x_2, \dots, x_p)}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})} dx_1 \dots dx_p.$$

Since for all ϵ , $\|H_\epsilon\|_1 = A$, this sum is equal to

$$A \sum_{j=1}^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{\mathcal{R}_{\epsilon, k_j - k_{j-1}} (1 + \mathcal{R}_{\epsilon, k_j})}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})}.$$

From (B.3), this term is $o((\log \epsilon^{-1})^p)$ as $\epsilon \rightarrow 0$. Therefore, $\mathcal{E}_\epsilon = o((\log \epsilon^{-1})^p)$.

Now, we focus on the first term in (B.8):

$$\left(\frac{\gamma}{4\pi}\right)^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{\int_{\mathbb{R}^{2p}} H_\epsilon(x_1, x_2, \dots, x_p) e^{-Q_{k_1, \dots, k_p}(x_1, \dots, x_p)} dx_1 \dots dx_p}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})}. \quad (\text{B.11})$$

We write (B.11) as

$$J_{p, \epsilon} + \left(\frac{\gamma A}{4\pi}\right)^p \sum_{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}} \frac{1}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})},$$

where

$$J_{p, \epsilon} = \left(\frac{\gamma}{4\pi}\right)^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{\int_{\mathbb{R}^{2p}} H_\epsilon(x_1, x_2, \dots, x_p) (e^{-Q_{k_1, \dots, k_p}(x_1, \dots, x_p)} - 1) dx_1 \dots dx_p}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})}. \quad (\text{B.12})$$

We aim to prove that $(\log \epsilon^{-1})^{-p} J_{p, \epsilon} \rightarrow 0$, as $\epsilon \rightarrow 0$. This will prove that (B.11) is asymptotic to $(\frac{\gamma A}{4\pi})^p$ as $\epsilon \rightarrow 0$, which proves (B.6).

From our assumption on (B.5), for any δ , there is a M_0 large enough such that

$$\sup_{\epsilon > 0} \int_{|\mathbf{x}| \geq M_0} H_\epsilon(x_1, x_2, \dots, x_p) d\mathbf{x} \leq \delta.$$

Therefore,

$$\left(\frac{\gamma}{4\pi}\right)^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{\int_{|\mathbf{x}| \geq M_0} H_\epsilon(x_1, x_2, \dots, x_p) |e^{-Q_{k_1, \dots, k_p}(x_1, \dots, x_p)} - 1| d\mathbf{x}}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})} \quad (\text{B.13})$$

is bounded above by

$$2\delta \left(\frac{\gamma}{4\pi}\right)^p \sum_{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}} \frac{1}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})}.$$

By dividing this term by $(\log \epsilon^{-1})^p$ and sending $\epsilon \rightarrow 0$, we see that it converges to $2\delta(\gamma/4\pi)^p$. Since δ is a arbitrary, (B.13) is negligible and the main contribution to (B.12) comes from

$$J''_{p, \epsilon} := \left(\frac{\gamma}{4\pi}\right)^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \frac{\int_{|\mathbf{x}| \leq M_0} H_\epsilon(x_1, x_2, \dots, x_p) (e^{-Q_{k_1, \dots, k_p}(x_1, \dots, x_p)} - 1) dx_1 \dots dx_p}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})}.$$

Observe that for all $|\mathbf{x}| \leq M_0$ we have

$$|e^{-Q_{k_1, \dots, k_p}(x_1, \dots, x_p)} - 1| \lesssim Q_{k_1, \dots, k_p}(x_1, \dots, x_p) \lesssim \sum_{i=1}^p \frac{1}{k_i - k_{i-1}}.$$

Therefore,

$$J_{p, \epsilon}'' \lesssim A \left(\frac{\gamma}{4\pi} \right)^p \sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \sum_{i=1}^p \frac{1}{k_1(k_2 - k_1) \dots (k_p - k_{p-1})} \cdot \frac{1}{k_i - k_{i-1}}.$$

From (B.4), the sum in the right-hand side is $o((\log \epsilon^{-1})^p)$ as $\epsilon \rightarrow 0$, and therefore $J_{p, \epsilon} = o((\log \epsilon^{-1})^p)$. This concludes the proof of (B.6). \square

Remark B.1. Let $(G_\epsilon)_{\epsilon \in (0,1)} \subseteq L^1(\mathbb{R}^{2p})$ satisfy the assumptions of **Proposition B.1** and define

$$\mathcal{G}_\epsilon(z) = \mathbb{E}_{Z^{(\epsilon)}}[G_\epsilon(\mathbf{z} + Z^{(\epsilon)})],$$

where $(Z^{(\epsilon)})_{\epsilon \in (0,1)}$ is a collection of tight random variables in \mathbb{R}^{2p} . The family $(\mathcal{G}_\epsilon)_{\epsilon \in (0,1)}$ satisfies the assumptions of **Proposition B.1**. To prove this we only need to prove (B.5). Since $(Z^{(\epsilon)})_{\epsilon \in (0,1)}$ is tight we have

$$\sup_{\epsilon \in (0,1)} \mathbb{P}[|Z^{(\epsilon)}| \geq M] \rightarrow 0, \quad (\text{B.14})$$

as $M \rightarrow \infty$. We also have

$$\int_{|z| \geq M} \mathcal{G}_\epsilon(\mathbf{z}) d\mathbf{z} \leq \mathbb{E} \left[\mathbf{1}_{\{|Z^{(\epsilon)}| \geq M/2\}} \int_{|z| \geq M - |Z^{(\epsilon)}|} G_\epsilon(\mathbf{z}) d\mathbf{z} \right] + \mathbb{E} \left[\mathbf{1}_{\{|Z^{(\epsilon)}| \leq M/2\}} \int_{|z| \geq M - |Z^{(\epsilon)}|} G_\epsilon(\mathbf{z}) d\mathbf{z} \right].$$

Therefore, since $(G_\epsilon)_{\epsilon > 0}$ satisfies the assumptions of **Proposition B.1** and from (B.14)

$$\int_{|z| \geq M} \mathcal{G}_\epsilon(\mathbf{z}) d\mathbf{z} \leq \mathbb{P}[|Z^{(\epsilon)}| \geq M/2] \int_{\mathbb{R}^{2p}} G_\epsilon(\mathbf{z}) d\mathbf{z} + \int_{|z| \geq M/2} G_\epsilon(\mathbf{z}) d\mathbf{z} \rightarrow 0,$$

as $M \rightarrow \infty$, uniformly in ϵ .

Finally, we prove the 'non-directed' version of **Proposition B.1**.

Lemma B.1. Let $(G_\epsilon)_{\epsilon \in (0,1)}$ be a collection of functions as in **Lemma B.1**. Also, let $\delta \in \{+1, -1, 0\}^p$ and $M(\epsilon)$ be such that $\log M(\epsilon) / \log \epsilon^{-1} \rightarrow 0$ as $\epsilon \rightarrow 0$. Then the mean of:

$$\sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} G_\epsilon(\omega_{X_0}(T_{k_1 + \delta_1}) - \omega_{Y_0}(T_{k_1}), \dots, \omega_{X_0}(T_{k_p + \delta_p}) - \omega_{Y_0}(T_{k_p})) \quad (\text{B.15})$$

is asymptotic to:

$$\left((\gamma/4\pi) \log \epsilon^{-1} \right)^p A,$$

as $\epsilon \rightarrow 0$, where $\|G_\epsilon\|_1 = A$.

Proof. We aim to use **Proposition B.1** to calculate the asymptotic of (B.15). The issue is the appearance of δ . We try to bring the functional in (B.15) to a functional involving only the random walk $\omega_{X_0}(T_k) - \omega_{Y_0}(T_k)$. We will exploit the fact that both $\omega_{X_0}(T_k)$ and $\omega_{Y_0}(T_k)$, as defined in (3.18), are sums of independent random variables. We will write the random vector appearing in (B.15) as:

$$(\omega_{X_0}(T_{k_1+\delta_1}) - \omega_{Y_0}(T_{k_1}), \dots, \omega_{X_0}(T_{k_p+\delta_p}) - \omega_{Y_0}(T_{k_p})) = (\mathbf{S}_{k_1}^{(\epsilon)}, \dots, \mathbf{S}_{k_p}^{(\epsilon)}) + \mathbf{Z}^{(\epsilon)}, \quad (\text{B.16})$$

where the vectors $(\mathbf{S}_{k_1}^{(\epsilon)}, \dots, \mathbf{S}_{k_p}^{(\epsilon)})$ and $\mathbf{Z}^{(\epsilon)}$ are independent. The distribution of the first vector will "look like" the distribution of $(\omega_{X_0}(T_{k_1}) - \omega_{Y_0}(T_{k_1}), \dots, \omega_{X_0}(T_{k_p}) - \omega_{Y_0}(T_{k_p}))$. Moreover, the distribution of $\mathbf{Z}^{(\epsilon)}$ will be independent from k_1, \dots, k_p . Then we average with respect to $\mathbf{Z}^{(\epsilon)}$ so that (B.15) is equal to

$$\sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \mathbb{E}[H(\mathbf{S}_{k_1}^{(\epsilon)}, \dots, \mathbf{S}_{k_p}^{(\epsilon)})],$$

for an appropriate H . Then, we will be able to apply **Proposition B.1** and conclude.

Let us make this more precise. From (3.18), we write for each $i = 1, \dots, p$:

$$\omega_{X_0}(T_{k_i+\delta_i}) - \omega_{Y_0}(T_{k_i}) = \mathbf{S}_{k_i} + \sum_{1 \leq j < i} (\mathbf{X}_{k_j-1}^{(\epsilon)} - \mathbf{Y}_{k_j-1}^{(\epsilon)} + \mathbf{X}_{k_j}^{(\epsilon)} - \mathbf{Y}_{k_j}^{(\epsilon)}) - \mathbf{Y}_{k_i-1}^{(\epsilon)} + \sum_{j=k_i-1}^{k_i+\delta_i-1} \mathbf{X}_j^{(\epsilon)},$$

where we make the convention that the last sum is 0 when $\delta_i = -1$. From (3.18), $\mathbf{S}_{k_i}^{(\epsilon)}$ is a sum of independent random variables. Moreover, \mathbf{S}_{k_i} is equal in distribution to $\omega_{X_0}(T_{k_i-2i+1}) - \omega_{Y_0}(T_{k_i-2i+1})$ and furthermore, the vector:

$$(\mathbf{S}_{k_1}, \dots, \mathbf{S}_{k_p})$$

is equal in distribution to

$$\left(\omega_{X_0}(T_{k_1-1}) - \omega_{Y_0}(T_{k_1-1}), \omega_{X_0}(T_{k_2-3}) - \omega_{Y_0}(T_{k_2-3}), \dots, \omega_{X_0}(T_{k_p-2p+1}) - \omega_{Y_0}(T_{k_p-2p+1}) \right)$$

and is independent of:

$$(\mathbf{X}_{k_1-1}^{(\epsilon)} - \mathbf{Y}_{k_1-1}^{(\epsilon)}, \mathbf{X}_{k_1}^{(\epsilon)} - \mathbf{Y}_{k_1}^{(\epsilon)}, \dots, \mathbf{X}_{k_p-1}^{(\epsilon)} - \mathbf{Y}_{k_p-1}^{(\epsilon)}, \mathbf{X}_{k_p}^{(\epsilon)} - \mathbf{Y}_{k_p}^{(\epsilon)}),$$

since these random variables do not appear inside the sum of any of the \mathbf{S}_{k_i} .

Now, define the vector $\mathbf{Z}^{(\epsilon)} \in \mathbb{R}^{2p}$, where the components $\mathbf{Z}^{(\epsilon)}(i) \in \mathbb{R}^2$ are given by:

$$\mathbf{Z}^{(\epsilon)}(i) := \sum_{1 \leq j < i} (\mathbf{X}_{k_j-1}^{(\epsilon)} - \mathbf{Y}_{k_j-1}^{(\epsilon)} + \mathbf{X}_{k_j}^{(\epsilon)} - \mathbf{Y}_{k_j}^{(\epsilon)}) - \mathbf{Y}_{k_i-1}^{(\epsilon)} + \sum_{j=k_i-1}^{k_i+\delta_i-1} \mathbf{X}_j^{(\epsilon)}.$$

By construction $\mathbf{Z}^{(\epsilon)}$ is independent from $(\mathbf{S}_{k_1}^{(\epsilon)}, \dots, \mathbf{S}_{k_p}^{(\epsilon)})$.

To prove that the distribution of $\mathbf{Z}^{(\epsilon)} = (\mathbf{Z}^{(\epsilon)}(1), \dots, \mathbf{Z}^{(\epsilon)}(p))$ is independent of k_1, \dots, k_p recall that the random variables $((\mathbf{X}_{k_i-1}^{(\epsilon)}, \mathbf{Y}_{k_i-1}^{(\epsilon)}), (\mathbf{X}_{k_i}^{(\epsilon)}, \mathbf{Y}_{k_i}^{(\epsilon)}))_{i=1, \dots, p}$ are i.i.d.⁶. Therefore, the vector

$$\left((\mathbf{X}_{k_1-1}^{(\epsilon)}, \mathbf{Y}_{k_1-1}^{(\epsilon)}), (\mathbf{X}_{k_1}^{(\epsilon)}, \mathbf{Y}_{k_1}^{(\epsilon)}), \dots, (\mathbf{X}_{k_p-1}^{(\epsilon)}, \mathbf{Y}_{k_p-1}^{(\epsilon)}), (\mathbf{X}_{k_p}^{(\epsilon)}, \mathbf{Y}_{k_p}^{(\epsilon)}) \right)$$

⁶Here, we made use of the restriction $k_i > k_{i-1} + 1$, so that $k_j < k_i - 1$ for all $j < i$.

is equal in distribution to

$$\left((\mathbf{X}_1^{(\epsilon)}, \mathbf{Y}_1^{(\epsilon)}), (\mathbf{X}_2^{(\epsilon)}, \mathbf{Y}_2^{(\epsilon)}), \dots, (\mathbf{X}_{2p-1}^{(\epsilon)}, \mathbf{Y}_{2p-1}^{(\epsilon)}), (\mathbf{X}_{2p}^{(\epsilon)}, \mathbf{Y}_{2p}^{(\epsilon)}) \right).$$

This means that the distribution of $\mathbf{Z}^{(\epsilon)}$ is equal to the distribution of $Z^{(\epsilon)} = (Z_1^{(\epsilon)}, \dots, Z_p^{(\epsilon)})$ where

$$Z_i^{(\epsilon)} := \sum_{1 \leq j < i} (\mathbf{X}_{2j-1}^{(\epsilon)} - \mathbf{Y}_{2j-1}^{(\epsilon)} + \mathbf{X}_{2j}^{(\epsilon)} - \mathbf{Y}_{2j}^{(\epsilon)}) - \mathbf{Y}_{2i-1}^{(\epsilon)} + \sum_{j=2i-1}^{2i+\delta_i-1} \mathbf{X}_j^{(\epsilon)}.$$

Therefore, the distribution of $\mathbf{Z}^{(\epsilon)}$ is independent from k_1, \dots, k_p . These observations prove (B.16). In particular, (B.15) is equal to

$$\sum_{\substack{1 \leq k_1 < \dots < k_p \leq [M(\epsilon)\epsilon^{-1}], \\ k_{i+1} > k_i + 2}} \mathbb{E} \left[\mathcal{G}^{(\epsilon)} \left(\omega_{X_0}(T_{k_1-1}) - \omega_{Y_0}(T_{k_1-1}), \dots, \omega_{X_0}(T_{k_p-2p+1}) - \omega_{Y_0}(T_{k_p-2p+1}) \right) \right], \quad (\text{B.17})$$

where $\mathcal{G}^{(\epsilon)} : \mathbb{R}^{2p} \rightarrow \mathbb{R}$ is given by:

$$\mathcal{G}^{(\epsilon)}(\mathbf{z}) := \mathbb{E}_{Z^{(\epsilon)}} \left[G(\mathbf{z} + Z^{(\epsilon)}) \right]. \quad (\text{B.18})$$

From **Proposition A.1** and **Remark B.1**, $(\mathcal{G}^{(\epsilon)})_{\epsilon \in (0,1)}$ satisfies the assumptions of **Proposition B.1** with $\|\mathcal{G}^{(\epsilon)}\|_1 = A$. We deduce that (B.17) is asymptotic to

$$\left((\gamma/4\pi) \log \epsilon^{-1} \right)^p A,$$

as $\epsilon \rightarrow 0$. Since (B.15) and (B.17) are equal, we get

$$\sum_{\substack{1 \leq k_1 < \dots < k_p \leq M(\epsilon)\epsilon^{-1}, \\ k_{i+1} > k_i + 2}} \mathbb{E} [G(\omega_{X_0}(T_{k_1+\delta_1}) - \omega_{Y_0}(T_{k_1}), \dots, \omega_{X_0}(T_{k_p+\delta_p}) - \omega_{Y_0}(T_{k_p}))] \sim A(\gamma/4\pi)^p (\log \epsilon^{-1})^p,$$

as $\epsilon \rightarrow 0$. □

C The normalization constant $\zeta_{t/\epsilon^2}^{(\epsilon)}$

In this appendix, we study the asymptotic behavior of $\zeta_{t/\epsilon^2}^{(\epsilon)}$. This is given by the following lemma:

Lemma C.1. *There is a constant $C_1 > 0$ such that*

$$\zeta_{t/\epsilon^2}^{(\epsilon)} = (C_1 + o(1)) \frac{t}{\epsilon^2 \log \epsilon^{-1}} + o(1), \quad (\text{C.1})$$

as $\epsilon \rightarrow 0$. The error terms depend only on R .

Proof. First, we assume that $t/\epsilon^2 = N_\epsilon \in \mathbb{N}$ and we set $\tau = 1$. Repeating the construction of the Markov chain, as in **Section 3.1**, while also keeping track of the normalization constants, yields

$$\hat{\mathbb{P}}_{N_\epsilon}^{(\epsilon)}(d\omega) = \Psi^{(\epsilon)}(x_0) \hat{\mathbb{P}}_1^{(\epsilon)}(dx_0) \prod_{k=0}^{N_\epsilon-2} \hat{\pi}^{(\epsilon)}(x_k, dx_{k+1}) \Psi^{(\epsilon)}(x_{N_\epsilon-1})^{-1} (\rho^{(\epsilon)})^{N_\epsilon-1} e^{N_\epsilon \zeta_1^{(\epsilon)} - \zeta_{N_\epsilon}^{(\epsilon)}}$$

with $\Psi^{(\epsilon)}$ and $\rho^{(\epsilon)}$ defined by the eigenvalue problem (3.7). Recall the normalization $\int_{\Omega_1} \Psi^{(\epsilon)}(x) \hat{\mathbb{P}}^{(\epsilon)}(dx) = 1$. Since $\hat{\mathbb{P}}_{N_\epsilon}^{(\epsilon)}$ is a probability measure

$$\mathbb{E}[\Psi^{(\epsilon)}(X_{N_\epsilon-1})^{-1}] = (\rho^{(\epsilon)})^{1-N_\epsilon} e^{-N_\epsilon \zeta_1^{(\epsilon)} + \zeta_{N_\epsilon}}$$

From the observations at the start of **Section 3.2**, $\Psi^{(\epsilon)}(x) \rightarrow 1$ uniformly in x , as $\epsilon \rightarrow 0$. Therefore,

$$\zeta_{N_\epsilon} = N_\epsilon \zeta_1^{(\epsilon)} - (N_\epsilon - 1) \log \rho^{(\epsilon)} + o(1). \quad (\text{C.2})$$

We estimate the terms $\zeta_1^{(\epsilon)}$ and $\log \rho^{(\epsilon)}$. For $\zeta_1^{(\epsilon)}$, recall its definition in (1.11)

$$\zeta_1^{(\epsilon)} = \log \int_{\Omega_1} \exp\left(\frac{\hat{\beta}^2}{2 \log \epsilon^{-1}} \int_{[0,1]^2} R(s-u, B(s) - B(u)) dsdu\right) W_1(dB).$$

By a Taylor expansion, we see that as $\epsilon \rightarrow 0$

$$\zeta_1^{(\epsilon)} = \frac{C}{\log \epsilon^{-1}} + o\left(\frac{1}{\log \epsilon^{-1}}\right), \quad (\text{C.3})$$

where

$$C = \mathbb{E}_B \left[\int_{[0,1]^2} R(s-u, B(s) - B(u)) dsdu \right].$$

Now for $\log \rho^{(\epsilon)}$, we look at the eigenvalue equation (3.7)

$$\int_{\Omega_1} e^{I^{(\epsilon)}(x,y)} \Psi^{(\epsilon)}(y) \hat{\mathbb{P}}_1^{(\epsilon)}(dy) = \rho^{(\epsilon)} \Psi^{(\epsilon)}(x),$$

with $I^{(\epsilon)}$ defined in (3.6). From the observations at the start of **Section 3.2**, $\hat{\mathbb{P}}_1^{(\epsilon)}$ converges to the Wiener measure in total variation as $\epsilon \rightarrow 0$. Moreover, $I^{(\epsilon)}(x,y) \rightarrow 0$ as $\epsilon \rightarrow 0$, uniformly in $x, y \in \Omega_1$. These two observations, combined with a Taylor expansion of the above exponential, show that

$$\rho^{(\epsilon)} - 1 = \frac{\tilde{C}}{\log \epsilon^{-1}} + o\left(\frac{1}{\log \epsilon^{-1}}\right),$$

as $\epsilon \rightarrow 0$, where

$$\tilde{C} = \int_{\Omega_1} \int_{\Omega_1} I(x,y) W_1(dx) W_1(dy), \quad (\text{C.4})$$

with

$$I(x,y) := \int_{[0,1]^2} R(s-u, y(s) + x(1) - x(u)) dsdu, \quad x, y \in \Omega_1.$$

This proves that

$$\log \rho^{(\epsilon)} = \frac{\tilde{C}}{\log \epsilon^{-1}} + o\left(\frac{1}{\log \epsilon^{-1}}\right), \quad (\text{C.5})$$

Now, plugging (C.5) and (C.3) to (C.2) yields (C.1) with:

$$C_1 = \mathbb{E}_B \left[\int_{[0,1]^2} R(s-u, B(s) - B(u)) dsdu \right] + \mathbb{E}_{B^1} \mathbb{E}_{B^2} \left[\int_{[0,1]^2} R(s-u, B^2(s) + B^1(1) - B^1(u)) dsdu \right].$$

When $t/\epsilon^2 \in \mathbb{R}_{>0}$ we have the same asymptotic, since

$$\zeta_{t/\epsilon^2}^{(\epsilon)} = \zeta_{[t/\epsilon^2]}^{(\epsilon)} + o(1)$$

and $[t/\epsilon^2]/(t/\epsilon^2) \rightarrow 1$ as $\epsilon \rightarrow 0$. □

Acknowledgments. *The author thanks their supervisor, Nikos Zygouras, for suggesting the problem and for many helpful discussions and suggestions. The author is supported by the Warwick Mathematics Institute Centre for Doctoral Training and gratefully acknowledges funding from the University of Warwick.*

References

- [Asm03] S. Asmussen. Applied Probability and Queues. Applications of mathematics : stochastic modelling and applied probability. Springer, 2003. ISBN: 9780387002118.
- [CSZ17] Francesco Caravenna, Rongfeng Sun, and Nikos Zygouras. Universality in marginally relevant disordered systems. In: *The Annals of Applied Probability* 27.5 (2017), pp. 3050–3112. ISSN: 10505164.
- [CSZ20] Francesco Caravenna, Rongfeng Sun, and Nikos Zygouras. The two-dimensional KPZ equation in the entire subcritical regime. In: *The Annals of Probability* 48.3 (2020), pp. 1086–1127. DOI: [10.1214/19-AOP1383](https://doi.org/10.1214/19-AOP1383).
- [CSZ21] Francesco Caravenna, Rongfeng Sun, and Nikos Zygouras. The critical 2d Stochastic Heat Flow. In: *Inventiones mathematicae* 233 (2021), pp. 325–460.
- [CSZ23] Francesco Caravenna, Rongfeng Sun, and Nikos Zygouras. The critical 2d stochastic heat flow is not a Gaussian multiplicative chaos. In: *The Annals of Probability* 51.6 (Nov. 2023). ISSN: 0091-1798. DOI: [10.1214/23-aop1648](https://doi.org/10.1214/23-aop1648).
- [CW17] Ajay Chandra and Hendrik Weber. Stochastic PDEs, Regularity structures, and interacting particle systems. en. In: *Annales de la Faculté des sciences de Toulouse : Mathématiques* Ser. 6, 26.4 (2017), pp. 847–909. DOI: [10.5802/afst.1555](https://doi.org/10.5802/afst.1555). URL: <http://www.numdam.org/articles/10.5802/afst.1555/>.
- [CNN22] Clément Cosco, Shuta Nakajima, and Makoto Nakashima. Law of large numbers and fluctuations in the sub-critical and L2 regions for SHE and KPZ equation in dimension $d \geq 3$. eng. In: *Stochastic processes and their applications* 151 (2022), pp. 127–173. ISSN: 0304-4149.
- [Dal+08] Robert C. Dalang, Davar Khoshnevisan, Carl Mueller, David Nualart, Yimin Xiao, and Firas Rassoul-Agha. A Minicourse on Stochastic Partial Differential Equations. In: 2008.
- [Duc22] Paweł Duch. Flow equation approach to singular stochastic PDEs. 2022. arXiv: [2109.11380](https://arxiv.org/abs/2109.11380).
- [Dun+21] Alexander Dunlap, Yu Gu, Lenya Ryzhik, and Ofer Zeitouni. The random heat equation in dimensions three and higher: the homogenization viewpoint. In: *Archive for Rational Mechanics and Analysis* 242.2 (2021), pp. 827–873.
- [Gu18] Yu Gu. Gaussian fluctuations from the 2D KPZ equation. In: *Stochastics and Partial Differential Equations: Analysis and Computations* 8 (2018), pp. 150–185. URL: <https://api.semanticscholar.org/CorpusID:115136843>.
- [GL19] Yu Gu and Jiawei Li. Fluctuations of a Nonlinear Stochastic Heat Equation in Dimensions Three and Higher. In: *SIAM J. Math. Anal.* 52 (2019), pp. 5422–5440. URL: <https://api.semanticscholar.org/CorpusID:207768470>.
- [GQT21] Yu Gu, Jeremy Quastel, and Li-Cheng Tsai. Moments of the 2D SHE at criticality. In: *Probability and Mathematical Physics* 2 (Mar. 2021), pp. 179–219. DOI: [10.2140/pmp.2021.2.179](https://doi.org/10.2140/pmp.2021.2.179).

- [GRZ18] Yu Gu, Lenya Ryzhik, and Ofer Zeitouni. The Edwards–Wilkinson Limit of the Random Heat Equation in Dimensions Three and Higher. In: *Communications in Mathematical Physics* 363.2 (July 2018), pp. 351–388. DOI: [10.1007/s00220-018-3202-0](https://doi.org/10.1007/s00220-018-3202-0). URL: <https://doi.org/10.1007%2Fs00220-018-3202-0>.
- [GT19] Yu Gu and Li-Cheng Tsai. Another look into the Wong-Zakai Theorem for Stochastic Heat Equation. In: *The Annals of Applied Probability* 29.5 (2019), pp. 3037–3061. ISSN: 10505164, 21688737. URL: <https://www.jstor.org/stable/26846324>.
- [GP16] Massimiliano Gubinelli and Nicolas Perkowski. KPZ Reloaded. In: *Communications in Mathematical Physics* 349.1 (Nov. 2016), pp. 165–269. DOI: [10.1007/s00220-016-2788-3](https://doi.org/10.1007/s00220-016-2788-3). URL: <https://doi.org/10.1007%2Fs00220-016-2788-3>.
- [Hai14] M. Hairer. A theory of regularity structures. In: *Inventiones mathematicae* 198.2 (Mar. 2014), pp. 269–504. DOI: [10.1007/s00222-014-0505-4](https://doi.org/10.1007/s00222-014-0505-4).
- [Hai13] Martin Hairer. Solving the KPZ equation. In: *Annals of Mathematics* 178.2 (2013), pp. 559–664. ISSN: 0003486X.
- [HP14] Martin Hairer and Etienne Pardoux. A Wong-Zakai theorem for stochastic PDEs. In: *Journal of the Mathematical Society of Japan* 67 (Sept. 2014). DOI: [10.2969/jmsj/06741551](https://doi.org/10.2969/jmsj/06741551).
- [Jan97] Svante Janson. Gaussian Hilbert Spaces. Cambridge Tracts in Mathematics. Cambridge University Press, 1997. DOI: [10.1017/CB09780511526169](https://doi.org/10.1017/CB09780511526169).
- [KR53] G. Kallianpur and H. Robbins. Ergodic Property of the Brownian Motion Process. In: *Proceedings of the National Academy of Sciences of the United States of America* 39.6 (1953), pp. 525–533. ISSN: 00278424. URL: <http://www.jstor.org/stable/88704>.
- [KR54] Gopinath Kallianpur and Herbert E. Robbins. The sequence of sums of independent random variables. In: *Duke Mathematical Journal* 21 (1954), pp. 285–307.
- [Kup14] Antti Kupiainen. Renormalization Group and Stochastic PDE’s. In: *Annales Henri Poincaré* 17 (Oct. 2014). DOI: [10.1007/s00023-015-0408-y](https://doi.org/10.1007/s00023-015-0408-y).
- [LL10] G.F. Lawler and V. Limic. Random Walk: A Modern Introduction. Cambridge Studies in Advanced Mathematics. Cambridge University Press, 2010. ISBN: 9781139488761. URL: <https://books.google.co.uk/books?id=UBQdwAZDeOEC>.
- [LZ22] Dimitris Lygkonis and Nikos Zygouras. Edwards-Wilkinson fluctuations for the directed polymer in the full L^2 -regime for dimensions $d \geq 3$. In: *Annales de l’Institut Henri Poincaré, Probabilités et statistiques* 58.1 (2022), pp. 65–104. DOI: [10.1214/21-AIHP1173](https://doi.org/10.1214/21-AIHP1173). URL: <https://doi.org/10.1214/21-AIHP1173>.
- [Nua09] D. Nualart. Malliavin Calculus and Its Applications. Regional conference series in mathematics. Conference Board of the Mathematical Sciences, 2009. ISBN: 9780821847794. URL: <https://books.google.co.uk/books?id=8HaNAwAAQBAJ>.
- [Sne+12] I.N. Sneddon, R. Dautray, P. Benilan, J.L. Lions, M. Cessenat, A. Gervat, A. Kavenoky, and H. Lanchon. Mathematical Analysis and Numerical Methods for Science and Technology: Volume 1 Physical Origins and Classical Methods. Springer Berlin Heidelberg, 2012. ISBN: 9783642615276. URL: <https://books.google.gr/books?id=mUzuCAAAQBAJ>.
- [Tho00] H. Thorisson. Coupling, Stationarity, and Regeneration. Probability and Its Applications. Springer New York, 2000. ISBN: 9780387987798.