## SPACELIKE INITIAL DATA FOR BLACK HOLE STABILITY

ALLEN JUNTAO FANG, JÉRÉMIE SZEFTEL, AND ARTHUR TOUATI

ABSTRACT. We construct initial data suitable for the Kerr stability conjecture, that is, solutions to the constraint equations on a spacelike hypersurface with boundary entering the black hole horizon that are arbitrarily decaying perturbations of a Kerr initial data set. This results from a more general perturbative construction on any asymptotically flat initial data set with the topology of  $\mathbb{R}^3 \setminus \{r < 1\}$ enjoying some analyticity near and at the boundary. In particular, we design a suitable mixed boundary condition for the elliptic operator of the conformal method in order to exclude the Killing initial data sets (KIDS).

#### 1. INTRODUCTION

The Kerr family of stationary black holes is generally believed to describe the black holes observed in astrophysics. An important question regarding whether or not this belief is substantiated is the stability property of this family. In this article, we construct initial data for the Kerr stability problem.

1.1. The initial value problem in general relativity. The Einstein vacuum equations (EVE) are the defining equations of Einstein's theory of general relativity under the assumption of vacuum and are given by

$$\mathbf{Ric}(\mathbf{g}) = 0,\tag{1.1}$$

where **g** is a Lorentzian metric defined on a 4-dimensional manifold  $\mathcal{M}$ , and **Ric** denotes the Ricci tensor. As first shown in the seminal papers [6, 7], there is a well-defined Cauchy problem for (1.1), where one solves for a maximal globally hyperbolic development ( $\mathcal{M}$ , **g**) of a given initial data triplet ( $\Sigma$ , g, k) consisting of a 3-dimensional Riemannian manifold ( $\Sigma$ , g) and a symmetric two tensor k satisfying the constraint equations. The constraint equations themselves are given by

$$R(g) + (\mathrm{tr}_g k)^2 - |k|_g^2 = 0,$$
  
div\_g k - dtr\_g k = 0, (1.2)

where R(g) is the scalar curvature of g.

1.2. The Kerr stability conjecture. Studying the stability properties of black hole solutions to EVE can be formulated in the context of the above Cauchy problem for EVE. We provide below a rough statement of the Kerr stability conjecture, and refer the reader for example to the introduction of [17, 21] for a more detailed discussion.

**Conjecture 1.1** (Kerr stability conjecture). If  $(\Sigma, g, k)$  is an initial data triplet such that (g, k) is an appropriately small perturbation of  $(g_{m,a}, k_{m,a})$  the initial data of a subextremal Kerr black hole  $\mathbf{g}_{m,a}$  (i.e. |a| < m), then the evolution of  $(\Sigma, g, k)$  under (1.1),  $(\mathcal{M}, \mathbf{g})$ , has a domain of outer communication that converges in the appropriate sense to a nearby member of the Kerr family  $(\mathcal{M}_{m_f,a_f}, \mathbf{g}_{m_f,a_f})$ .

In the context of black hole stability, there are generally three regions of interest in a black hole spacetime, separated by null cones arising from the initial data, as indicated by the three regions I, II, and III in Figure 1. We discuss briefly the stability results in each of these regions.



FIGURE 1. Penrose diagram of the future of initial perturbations of Kerr.

Region I is a compact region of spacetime, and thus, is a local existence regime. In region II, the full subextremal Kerr family was shown to be stable by [3, 28]. In region III, stability has been shown<sup>1</sup> for slowly-rotating Kerr black holes (i.e.  $|a| \ll m$ ) in [17–19, 21, 27], using initialization<sup>2</sup> from regions I and II, thus proving Conjecture 1.1 in the slowly-rotating case.

1.3. First version of the main result. Our goal is to construct (g, k) such that  $(\Sigma, g, k)$  is an initial data triplet satisfying the constraint equations (1.2) and is an arbitrarily decaying perturbation of the initial data for a given Kerr black hole in order to satisfy the requirements in [3, 28]. We remark that this will also induce data that suffices for [12, 20, 21]. We give here a rough version of our main result, see Theorem 3.3 for the precise statement.

**Theorem 1.2** (Rough statement). Let  $(\Sigma, g_{m,a}, k_{m,a})$  be the spacelike initial data set induced by the Kerr metric  $\mathbf{g}_{m,a}$ , and let  $q \in \mathbb{N}^*$ ,  $0 < \delta < 1$  and  $\varepsilon > 0$  small enough. There exists a codimension  $4q^2$  family of perturbations  $(g_{m,a} + \tilde{g}, k_{m,a} + \tilde{k})$  of  $(g_{m,a}, k_{m,a})$  such that

$$|\tilde{g}| \sim \varepsilon (1+r)^{-q-\delta}, \quad \left|\tilde{k}\right| \sim \varepsilon (1+r)^{-q-\delta-1},$$
(1.3)

and  $\left(\Sigma, g_{m,a} + \tilde{g}, k_{m,a} + \tilde{k}\right)$  satisfies the constraint equations (1.2).

Theorem 1.2 provides initial data for the works mentioned in Section 1.2:

- In region II, [3] requires  $q \ge 3$  and [28] requires  $q \ge 1$ , with  $0 < \delta < 1$  in both cases.
- In region III, [20, 21] require q = 1 and [12] requires q = 2, with  $\delta = \left(\frac{1}{2}\right)_+$  in both cases.

1.4. Construction of initial data sets. There is a rich literature on the constraint equations (1.2) and we refer the reader for example to [4] for a detailed review. The main two approaches to produce solutions to these equations are the conformal method and the gluing method. First introduced in [23], the conformal method proposes a conformal ansatz for (g, k) and transforms the quasi-linear underdetermined

<sup>&</sup>lt;sup>1</sup>See also [12, 20] for the stability of Schwarzschild (i.e. a = 0) for restrictive initial data.

<sup>&</sup>lt;sup>2</sup>This initialization takes place on the null cones of Figure 1 where the constraint equations take the form of non-linear ODEs, as opposed to (1.2) which are non-linear elliptic PDEs.

system (1.2) into a semi-linear elliptic system for a scalar function and a vector field. This approach has been successful in constructing many interesting solutions to the constraint equations, including those with large mean curvature or with apparent horizons [13, 25]. However these results fall short of constructing perturbations of a given black hole satisfying (1.3). This is due to the fact that these constructions are typically done within the range of invertibility of the relevant elliptic operators resulting in perturbations that decay like  $r^{-\delta}$  for  $0 < \delta < 1$ , falling short of the decay rates assumed in the aforementioned stability results, see the discussion after Theorem 1.2.

The gluing method, first introduced in [10] and later extended in [9, 11], consists of gluing any asymptotically flat solution of (1.2) to an exact Kerr initial data set outside a far away annulus. However, critically, these constructions, viewed as a perturbation of the background black hole metric, are compactly supported, whereas we aim to construct non-compactly supported initial data.

**Remark 1.3.** Gluing results use the Kerr initial data, viewed as a 10-parameter family, to saturate the 10-dimensional space of linear obstructions coming from the symmetries of Minkowski. Theorem 1.2 exhibits infinitely many 10-parameter families with the same asymptotics at main order. As a result, one should be able to glue asymptotically flat initial data to arbitrarily decaying perturbations of Kerr.

The authors have previously constructed and parametrized initial data perturbations of Minkowski space with arbitrary decay [14]. While the overall methodology is similar, there are two key differences between the current paper and [14]. We first recall that in [14] the Killing initial data sets (KIDS) presented a fundamental, linear obstacle to the construction. Since black hole spacetimes have fewer Killing vectorfields than Minkowski, there are fewer KIDS to deal with in the present paper than in [14]. In addition, motivated by the stability of Minkowski, initial data were constructed in [14] for a manifold without boundary. On the other hand, in the present paper, initial data are constructed for a manifold with boundary, reflecting the presence of an event horizon. The presence of the boundary adds flexibility, as the elliptic problem to be solved does not itself impose boundary conditions. This will play a crucial role in the ensuing proof, where we take advantage of this freedom.

1.5. Sketch of proof. We now give a brief sketch of the proof of Theorem 1.2. Rather than considering initial data  $(\Sigma, g, k)$  it will be convenient to instead consider initial data triplets  $(\Sigma, g, \pi)$ , where  $\pi = k - (\operatorname{tr}_g k)g$  will be referred to as the *reduced second fundamental form* of  $\Sigma$ .

Using the conformal method, we consider as an ansatz  $(g, \pi)$  of the form

$$(g,\pi) = \left( (1+\check{u})^4 \left( g_{m,a} + \check{g} \right), (1+\check{u})^2 \left( \pi_{m,a} + \check{\pi} + L_{g_{m,a}} \check{X} \right) \right), \tag{1.4}$$

where  $L_{g_{m,a}}$  is a modified Lie derivative (see (3.2)), and where  $(\check{g}, \check{\pi})$  are the "seed" of the construction and enjoy smallness and decay satisfying

$$\sup_{k=0,1} \left( r^k |\nabla^k \check{g}| + r^{k+1} |\nabla^k \check{\pi}| \right) \le \frac{\varepsilon}{(1+r)^{q+\delta}}.$$
(1.5)

The constraint equations (1.2) for  $(g, \pi)$  then become the following system of PDEs on  $\Sigma$ 

$$P\left(\check{u},\check{X}\right) = D\Phi[g_{m,a},\pi_{m,a}](\check{g},\check{\pi}) + \text{non-linear terms},$$
(1.6)

where P is an elliptic operator depending only on  $(g_{m,a}, \pi_{m,a})$ , and  $D\Phi[g_{m,a}, \pi_{m,a}]$  denotes the linearization of the constraint operator at  $(g_{m,a}, \pi_{m,a})$ . Theorem 1.2 then asks whether there exists a solution  $(\check{u}, \check{X})$  to (1.6) satisfying

$$|\breve{u}| + r \left| \nabla \breve{X} \right| \lesssim \frac{\varepsilon}{(1+r)^{q+\delta}}.$$
(1.7)

We remark that applying the conformal method does not impose boundary conditions on  $\partial \Sigma$  for the solutions  $(\check{u}, \check{X})$  to (1.6), which will play a critical role as mentioned above.

To address this question, we study the behavior of the elliptic operator P on weighted Sobolev spaces<sup>3</sup>  $H^2_{-q-\delta}(\Sigma)$ . Using qualitative properties of elliptic operators to understand the decay of their solutions

<sup>&</sup>lt;sup>3</sup>Recall that  $u \in H^2_{\eta}(\Sigma)$  implies  $|u| \lesssim r^{\eta}$  at infinity, see Definition 2.2.

is a classical method, and we illustrate the basic idea behind with toy model of the scalar Laplacian on Euclidean space before addressing the full problem at hand. The mapping properties of the Laplacian

$$\Delta: H^2_{\delta}(\mathbb{R}^3) \longrightarrow L^2_{\delta-2}(\mathbb{R}^3) \tag{1.8}$$

are well studied [2, 26], and in particular, it is known that the Laplacian as defined in (1.8) is a Fredholm operator, if  $\delta \notin \mathbb{Z}$ . This implies, via the Fredholm alternative, that provided h is orthogonal to the kernel of  $\Delta^*$ , then  $\Delta^{-1}h$  is well defined. Thus, given some  $h \in L^2_{\delta-2}(\mathbb{R}^3)$ , finding a perturbation  $\tilde{h}$  such that  $\Delta^{-1}(h+\tilde{h}) \in H^2_{\delta}(\mathbb{R}^3)$  reduces to solving for  $\tilde{h}$  such that

$$\left\langle h + \tilde{h}, w \right\rangle_{L^2(\mathbb{R}^3)} = 0, \quad \text{for all } w \in \ker(\Delta^*).$$
 (1.9)

In this case of the Laplacian, the kernel of  $\Delta^*$  can be explicitly described and it consists just of harmonic polynomials. Thus, a suitable perturbation  $\tilde{h}$  can be explicitly solved for. In particular, this toy model emphasizes the fact that a full characterization of the cokernel of the elliptic operator under consideration is crucial.

We now return to giving a sketch of the proof in our setting. It can be verified that P is also a Fredholm operator acting between weighted Sobolev spaces. However, characterizing the kernel of  $P^*$  is more delicate. We characterize the kernel of  $P^*$  as perturbations of the harmonic polynomials. The critical tool will be a good choice of mixed boundary condition on  $\partial \Sigma$  of the form

$$(B_{\nu} + F)\left(\breve{u}, \breve{X}\right) = 0, \tag{1.10}$$

such that the operator  $(P, B_{\nu} + F)$  acting on  $H^2_{-\delta}(\Sigma)$  for  $0 < \delta < 1$  will be injective, and where  $B_{\nu}$  is a Neumann type boundary operator defined on  $\partial \Sigma$ . We will also require that F vanishes identically on an open subset of  $\partial \Sigma$  and is the identity on another open subset. This condition on F is not necessary to characterize the kernel of  $P^*$  but will play a key role in eliminating certain obstructions in what follows.

To ensure the orthogonality conditions necessary for P to be invertible, we will modify the ansatz (1.4) to include compactly supported symmetric 2-tensors  $(\check{g}, \check{\pi})$  such that the elliptic system reads

$$P\left(\check{u},\check{X}\right) = D\Phi[g_{m,a},\pi_{m,a}]\left((\check{g},\check{\pi}) + (\check{g},\check{\pi})\right) + \text{non-linear terms.}$$
(1.11)

The correctors  $(\check{g}, \check{\pi})$  will then be chosen in a  $4q^2$  dimensional space such as to ensure the following analogue of (1.9):

$$\langle (\breve{g}, \breve{\pi}), D\Phi[g_{m,a}, \pi_{m,a}]^*(\mathbb{W}) \rangle = - \langle D\Phi[g_{m,a}, \pi_{m,a}](\breve{g}, \breve{\pi}), \mathbb{W} \rangle + \text{non-linear terms}$$
(1.12)

for all  $\mathbb{W}$  in the cokernel of  $(P, B_{\nu} + F)$ . The linearization of the system in (1.12) is solvable if  $D\Phi[g_{m,a}, \pi_{m,a}]^*$  acts injectively on the cokernel of  $(P, B_{\nu} + F)$  restricted to the support of  $(\breve{g}, \breve{\pi})$ . Now, since the Kerr initial data  $(g_{m,a}, \pi_{m,a})$  can be proved to be analytic near and at the boundary, elements of  $D\Phi[g_{m,a}, \pi_{m,a}]^*$  can also be proved to be analytic near and at the boundary. Therefore, they cannot satisfy the boundary condition (1.10) since we assumed that F vanishes in an open set of  $\partial\Sigma$  and is the identity on another one. This shows how  $D\Phi[g_{m,a}, \pi_{m,a}]^*$  acts injectively on the cokernel of  $(P, B_{\nu} + F)$ .

The above procedure illustrates how a careful choice of F allows us to simultaneously describe the mapping properties of P and also to ensure that we can appropriately solve for the correctors  $(\check{g}, \check{\pi})$  by eliminating the KIDS, i.e. the elements of ker $(D\Phi[g_{m,a}, \pi_{m,a}]^*)$ , which are well-known linear obstructions to solving (1.12). A straightforward fixed point argument then concludes the resolution of the non-linear system (1.11).

**Remark 1.4.** In fact, the only properties of the Kerr initial data set we use are its analyticity near and at the boundary, and the fact that it solves (1.2). For this reason, we prove a more general statement on asymptotically flat solutions of (1.2) which are analytic near and at the boundary, see Theorem 3.1, from which the result for Kerr is then deduced, see Theorem 3.3.

1.6. Outline of the article. We conclude this introduction by giving an outline of the article:

• In Section 2, we start with geometric and analytic preliminaries.

- In Section 3, we give a precise statement of our main result Theorem 3.1, from which we deduce Theorem 1.2.
- In Section 4, we study the properties of the elliptic operator P at the heart of our construction.
- Finally, in Section 5 we prove Theorem 3.1.

Acknowledgments. The first author acknowledges support from NSF award DMS-2303241 and support through Germany's Excellence Strategy EXC 2044 390685587, Mathematics Münster: Dynamics–Geometry–Structure.

## 2. Preliminaries

2.1. Notations and geometry. Throughout the paper, we will use Einstein summation notation, with roman indices referring to spatial indices  $\{1, 2, 3\}$ .

**Definition 2.1.** A triplet  $(\Sigma, \bar{g}, \bar{\pi})$ , where  $\Sigma$  is a 3-manifold with boundary,  $\bar{g}$  is a Riemannian metric on  $\Sigma$  and  $\bar{\pi}$  is a symmetric 2-tensor is called asymptotically flat if:

- (i)  $\Sigma$  is smooth, diffeomorphic to  $\mathbb{R}^3 \setminus B$  where B is an open ball in  $\mathbb{R}^3$  and there exists a global coordinate system  $(x^1, x^2, x^3)$  on  $\Sigma$  such that  $\partial \Sigma = \{r = r_0\}$  for some  $r_0 > 0$  where  $r := \sqrt{(x^1)^2 + (x^2)^2 + (x^3)^2}$ ,
- (ii)  $(\bar{g}, \bar{\pi})$  is smooth and asymptotically flat, i.e. there exists constants  $C_k$  such that for every  $k \in \mathbb{N}$  we have

$$\sup_{\Sigma} \left( r^{k+1} \left| \nabla^k (\bar{g}_{ij} - \delta_{ij}) \right| + r^{k+2} \left| \nabla^k \bar{\pi}_{ij} \right| \right) \le C_k,$$

where  $\nabla = (\partial_{x^1}, \partial_{x^2}, \partial_{x^3}).$ 

We will also refer to  $\Sigma$  satisfying the above conditions as an asymptotically flat manifold with boundary.

2.2. Function spaces. On  $(\Sigma, \bar{g}, \bar{\pi})$  an asymptotically flat manifold with boundary, as defined in Definition 2.1, we use the standard  $L^p(\Sigma)$  spaces for  $1 \leq p \leq \infty$ , where the integration is defined with respect to the volume form of  $\bar{g}$ . Moreover, the boundary  $\partial \Sigma$  is a compact 2-manifold and we will use the standard  $H^s(\partial \Sigma)$  spaces for  $s \in \mathbb{R}$ . We now define function spaces with weights at infinity.

**Definition 2.2.** Let  $(\Sigma, \overline{g}, \overline{\pi})$  be an asymptotically flat manifold with boundary as defined in Definition 2.1 and let  $k \in \mathbb{N}$  and  $\delta \in \mathbb{R}$ .

(i) We define the weighted Sobolev spaces  $H^k_{\delta}(\Sigma)$  to be the set of distribution on  $\Sigma$  for which the following norm is finite

$$\|u\|_{H^k_{\delta}(\Sigma)} \coloneqq \sum_{|\alpha| \le k} \left\| (1+r)^{-\delta - \frac{3}{2} + |\alpha|} \nabla^{\alpha} u \right\|_{L^2(\Sigma)}$$

We denote  $L^2_{\delta}(\Sigma) = H^0_{\delta}(\Sigma)$ .

(ii) We define the weighted Hölder spaces  $C^k_{\delta}(\Sigma)$  to be the set of distribution on  $\Sigma$  for which the following norm is finite

$$\|u\|_{C^k_{\delta}(\Sigma)} := \sum_{|\alpha| \le k} \left\| (1+r)^{-\delta + |\alpha|} \nabla^{\alpha} u \right\|_{L^{\infty}(\Sigma)}.$$

(iii) We extend these definitions to tensors of any type by summing over components in the global coordinates system  $(x^1, x^2, x^3)$  of Definition 2.1.

In the following lemma, we gather standard properties of these spaces. Proofs of these properties can be found in [5].

**Lemma 2.3.** Let  $k, k_1, k_2 \in \mathbb{N}$  and  $\delta, \delta_1, \delta_2 \in \mathbb{R}$ .

- (i) If  $k \geq 1$  and  $\delta' < \delta$  then the inclusion  $H^k_{\delta'}(\Sigma) \subset H^{k-1}_{\delta}(\Sigma)$  is compact. (ii) If  $k \geq 2$  then  $H^k_{\delta}(\Sigma) \subset C^0_{\delta}(\Sigma)$  and this inclusion is continuous.
- (iii) If  $k \leq \min(k_1, k_2)$ ,  $k < k_1 + k_2 \frac{3}{2}$  and  $\delta > \delta_1 + \delta_2$  then multiplication is a continuous map from  $H^{k_1}_{\delta_1}(\Sigma) \times H^{k_2}_{\delta_2}(\Sigma)$  to  $H^k_{\delta}(\Sigma)$ .

2.3. The constraint operator. Using the reduced second fundamental form  $\pi$ , the constraint equations (1.2) can be rewritten as

$$R(g) + \frac{1}{2} (\mathrm{tr}_g \pi)^2 - |\pi|_g^2 = 0,$$
  
$$\mathrm{div}_g \pi = 0.$$
 (2.1)

If g is a Riemannian metric on  $\Sigma$  and  $\pi$  is a symmetric 2-tensor on  $\Sigma$ , we define

$$\mathcal{H}(g,\pi) := R(g) + \frac{1}{2} (\mathrm{tr}_g \pi)^2 - |\pi|_g^2,$$
$$\mathcal{M}(g,\pi) := \mathrm{div}_g \pi.$$

For these operators, we will use the following notation for their expansion around some given  $(q, \pi)$ :

$$\mathcal{H}(g+h,\pi+\varpi) - \mathcal{H}(g,\pi) = D\mathcal{H}[g,\pi](h,\varpi) + Q_{\mathcal{H}}[g,\pi](h,\varpi), \tag{2.2}$$

$$\mathcal{M}(g+h,\pi+\varpi) - \mathcal{M}(g,\pi) = D\mathcal{M}[g,\pi](h,\varpi) + Q_{\mathcal{M}}[g,\pi](h,\varpi),$$
(2.3)

where  $D\mathcal{H}[g,\pi]$  and  $D\mathcal{M}[g,\pi]$  are the linearization of  $\mathcal{H}$  and  $\mathcal{M}$  and where  $Q_{\mathcal{H}}[g,\pi](h,\varpi)$  and  $Q_{\mathcal{H}}[g,\pi](h,\varpi)$ contain all quadratic and higher order terms in  $(h, \varpi)$ . Note that in this article, and in particular in the following lemma, we do not distinguish between coefficients of a metric g and coefficients of its inverse and thus simply write g in the schematic notation for error terms.

**Lemma 2.4.** Let g be a Riemannian metric on  $\Sigma$  and  $\pi$  a symmetric 2-tensor on  $\Sigma$ . We have

$$D\mathcal{H}[g,\pi](h,\varpi) = -\Delta_g \operatorname{tr}_g h + D^i D^j h_{ij} - h^{ij} \operatorname{Ric}(g)_{ij}$$

$$+ \operatorname{tr}_g \pi \left( \operatorname{tr}_g \varpi - \pi^{ij} h_{ij} \right) - 2\pi^{ij} \varpi_{ij} + 2h^{ij} g^{k\ell} \pi_{ik} \pi_{j\ell},$$

$$D\mathcal{M}[g,\pi](h,\varpi)_i = \operatorname{div}_g \varpi_i - h^{k\ell} D_k \pi_{\ell i} - \pi_i^j g^{k\ell} \left( D_k h_{\ell j} - \frac{1}{2} D_j h_{k\ell} \right) - \frac{1}{2} \pi^{k\ell} D_i h_{k\ell},$$

$$(2.4)$$

where D is the covariant derivative of q and all inverse are with respect to q. Moreover the terms  $Q_{\mathcal{H}}[g,\pi](h,\varpi)$  and  $Q_{\mathcal{M}}[g,\pi](h,\varpi)$  have the following schematic form

$$Q_{\mathcal{H}}[g,\pi](h,\varpi) = h\nabla^2 h + (g\nabla g)(h\nabla h) + g^2 \left((\nabla h)^2 + \varpi^2\right) + \left((\nabla g)^2 + \pi^2\right)h^2 + g\pi h\varpi + gh(\nabla h)^2 + h^2\nabla g\nabla h + h^2 \left((\nabla h)^2 + \varpi^2\right) + \pi h^2\varpi,$$
(2.6)

$$Q_{\mathcal{M}}[g,\pi](h,\varpi) = g^2 h \nabla \varpi + (g \nabla g) h \varpi + g \pi h \nabla h + h^2 \pi \nabla g + g h \varpi \nabla h + h^2 \varpi \nabla g + h^2 \pi \nabla h + h^2 \varpi \nabla h.$$
(2.7)

*Proof.* See [15] for (2.4) and (2.5). For (2.6) and (2.7), we use the schematic expressions

$$\begin{aligned} \mathcal{H}(g,\pi) &= g \nabla^2 g + (g \nabla g)^2 + (g \pi)^2 \\ \mathcal{M}(g,\pi) &= g \left( \nabla \pi + g \pi \nabla g \right). \end{aligned}$$

Isolating nonlinear terms in these expressions concludes the proof of the lemma.

**Definition 2.5.** The constraint operator is defined to be

$$\Phi(g,\pi) := \left(-\mathcal{H}(g,\pi), 2\mathcal{M}(g,\pi)\right). \tag{2.8}$$

Its linearization around some given  $(g,\pi)$  is denoted by  $D\Phi[g,\pi]$  and is given by

$$D\Phi[g,\pi](h,\varpi) = \left(-D\mathcal{H}[g,\pi](h,\varpi), 2D\mathcal{M}[g,\pi](h,\varpi)\right).$$
(2.9)

Its formal adjoint  $D\Phi[g,\pi]^*$  for the  $L^2(\Sigma)$  scalar product induced by the metric g will play a major role in our construction since elements of its kernel will be obstructions to solvability. From the formulas

(2.4), (2.5) and (2.8) we find that (f, X) belongs to ker $(D\Phi[g, \pi]^*)$  if and only if

$$\operatorname{Hess}_{g}(f) = f\left(\operatorname{Ric}(g)_{ij} + \frac{3}{4}\pi_{ij}\operatorname{tr}_{g}\pi - 2\pi_{ik}\pi_{j}^{k} + \frac{1}{4}|\pi|_{g}^{2}g_{ij}\right) + \pi_{kj}D_{i}X^{k} + \pi_{ki}D_{j}X^{k} + D_{X}\pi_{ij} - \frac{1}{2}X(\operatorname{tr}_{g}\pi)g_{ij},$$
(2.10)

$$\mathcal{L}_X g_{ij} = f\left(\pi_{ij} - \frac{1}{2} \mathrm{tr}_g \pi g_{ij}\right), \qquad (2.11)$$

where  $\mathcal{L}_X g$  is the Lie derivative with respect to X.

### 3. Statement of the main results

3.1. The main theorem. We now state the main result of this article, which handles any asymptotically flat solution of the constraint equations with analyticity near and at the boundary, see Remark 1.4. We will then apply it to Kerr in Section 3.2.

**Theorem 3.1.** Let  $(\Sigma, \bar{g}, \bar{\pi})$  be an asymptotically flat initial data as defined in Definition 2.1. Assume that there exists  $r_1 > r_0$  such that if  $\Omega := \Sigma \cap \{r_0 \le r < r_1\}$ , then the initial data set  $(\Omega, \bar{g}_{|\Omega}, \bar{\pi}_{|\Omega})$  is analytic<sup>4</sup>. Let  $q \in \mathbb{N}^*$  and  $0 < \delta < 1$ . There exists  $\varepsilon_0 > 0$  only depending on  $(\Sigma, \bar{g}, \bar{\pi})$ , q and  $\delta$  such that if  $0 < \varepsilon \le \varepsilon_0$  then the following holds: for all  $(\check{g}, \check{\pi})$  such that

$$\|\check{g}\|_{H^2_{-q-\delta}(\Sigma)} + \|\check{\pi}\|_{H^1_{-q-\delta-1}(\Sigma)} \le \varepsilon, \tag{3.1}$$

there exists a solution  $(g,\pi)$  to the constraint equations (2.1) on  $\Sigma$  of the form

$$g = u^4(\bar{g} + \check{g} + \check{g}),$$
  
$$\pi = u^2(\bar{\pi} + \check{\pi} + \check{\pi} + L_{\bar{g} + \check{g} + \check{g}}X),$$

where

$$L_g X := \mathcal{L}_X g - (\operatorname{div}_g X) g, \tag{3.2}$$

the scalar function u and the vector field X satisfy

 $\|u-1\|_{H^2_{-q-\delta}(\Sigma)} + \|X\|_{H^2_{-q-\delta}(\Sigma)} \le C\varepsilon,$ 

and  $(\check{g}, \check{\pi})$  belong to a vector space of dimension  $4q^2$  only depending on  $(\Sigma, \bar{g}, \bar{\pi})$  and composed of pairs of smooth symmetric 2-tensors compactly supported in  $\Sigma \cap \{r_0 < r < r_1\}$  and moreover

$$\|\breve{g}\|_{W^{2,\infty}(\Sigma)} + \|\breve{\pi}\|_{W^{1,\infty}(\Sigma)} \le C\varepsilon,$$

where C > 0 only depends on  $(\Sigma, \bar{g}, \bar{\pi})$ .

Theorem 3.1 will be proved in Section 5 relying on the properties of the linearized operator exhibited in Section 4.

3.2. Spacelike initial data for Kerr stability. In this section we present the main application of Theorem 3.1, i.e. the construction of initial data for the Kerr stability conjecture. Let m and a be two real numbers representing the mass and angular momentum of the Kerr black hole respectively. Let  $\mathcal{M}_{m,a} := \mathbb{R} \times (r_+, +\infty) \times (0, \pi) \times \mathbb{S}^1$  with  $r_+ = m + \sqrt{m^2 - a^2}$ . Then, the Kerr metric in Boyer-Lindquist coordinates  $(t, r, \theta, \phi)$  is given by

$$\mathbf{g}_{m,a} = -\frac{|q|^2 \Delta}{\Sigma^2} \mathrm{d}t^2 + \frac{\Sigma^2 \sin^2 \theta}{|q|^2} \left( \mathrm{d}\phi - \frac{2amr}{\Sigma^2} \mathrm{d}t \right)^2 + \frac{|q|^2}{\Delta} \mathrm{d}r^2 + |q|^2 \mathrm{d}\theta^2, \tag{3.3}$$

where

$$\Delta := r^2 + a^2 - 2mr, \quad q := r + ia\cos\theta, \quad \Sigma^2 := (r^2 + a^2)^2 - a^2\sin^2\theta\Delta,$$

which is well-defined for<sup>5</sup>  $(t, r, \theta, \phi) \in \mathcal{M}_{m,a}$ .

<sup>&</sup>lt;sup>4</sup>By this we mean analytic in  $\{r_0 < r < r_1\}$  and on the sphere  $\{r = r_0\}$ .

<sup>&</sup>lt;sup>5</sup>As usual, the coordinate singularities at the north and south poles of the spheres can be resolved by considering for example  $(x_{\pm}, y_{\pm}) = (\sin\theta\cos\phi, \sin\theta\cos\phi)$  as coordinates on neighborhoods of the north and south pole respectively.

It is well known that the singularity of the Kerr metric at the event horizon  $\{r = r_+\}$  is a coordinate singularity, and that the metric can in fact be smoothly extended past the event horizon. We can use this fact to construct a spacelike hypersurface of Kerr that enters into the black hole interior.

**Lemma 3.2.** There exist coordinates  $\tau$  and  $\varphi$  such that

(i) in the coordinates  $(\tau, r, \theta, \varphi)$  the Kerr metric  $\mathbf{g}_{m,a}$  is smooth on

$$\mathbb{R} \times [r_+(1-\delta_*),+\infty) \times (0,\pi) \times \mathbb{S}^1$$

and analytic in particular on

$$\mathbb{R} \times [r_+(1-\delta_*), 3m) \times (0, \pi) \times \mathbb{S}^1$$

for any  $0 < \delta_* < 1$ ,

*(ii) the hypersurface* 

$$\Sigma_{m,a} := \{\tau = 0\} \times [r_+(1-\delta_*), +\infty) \times (0,\pi) \times \mathbb{S}^1$$

is uniformly spacelike.

*Proof.* We recall the expression of the canonical ingoing principal null frame for the metric  $\mathbf{g}_{m,a}$ , see for example Section 2.7.2 in [21]:

$$e_{3} := \frac{r^{2} + a^{2}}{\Delta} \partial_{t} - \partial_{r} + \frac{a}{\Delta} \partial_{\phi}, \quad e_{4} := \frac{r^{2} + a^{2}}{|q|^{2}} \partial_{t} + \frac{\Delta}{|q|^{2}} \partial_{r} + \frac{a}{|q|^{2}} \partial_{\phi},$$

$$e_{1} := \frac{1}{|q|} \partial_{\theta}, \qquad e_{2} := \frac{a \sin \theta}{|q|} \partial_{t} + \frac{1}{|q| \sin \theta} \partial_{\phi}.$$
(3.4)

The new coordinates  $\tau$  and  $\varphi$  are defined in terms of the Boyer-Lindquist coordinates by

$$\tau = t + f(r), \quad \varphi = \phi + h(r), \tag{3.5}$$

where

$$h'(r) = \frac{a}{\Delta}, \quad f'(r) = \chi(r) \frac{r^2 + a^2}{\Delta} \sqrt{1 - \frac{2m^2\Delta}{(r^2 + a^2)^2}},$$

with f(4m) = 0 and where  $0 \le \chi \le 1$  is smooth and such that  $\chi = 1$  for r < 3m and  $\chi = 0$  for r > 4m. From (3.4) and (3.5) we first obtain

$$\begin{split} e_{3}(r) &= -1, \quad e_{4}(r) = \frac{\Delta}{|q|^{2}}, \quad e_{1}(r) = e_{2}(r) = 0, \\ e_{1}(\theta) &= \frac{1}{|q|}, \quad e_{3}(\theta) = e_{4}(\theta) = e_{2}(\theta) = 0, \\ e_{4}(\varphi) &= \frac{2a}{|q|^{2}}, \quad e_{2}(\varphi) = \frac{1}{|q|\sin\theta}, \quad e_{3}(\varphi) = e_{1}(\varphi) = 0, \\ e_{3}(\tau) &= \frac{r^{2} + a^{2}}{\Delta} \left( 1 - \chi(r)\sqrt{1 - \frac{2m^{2}\Delta}{(r^{2} + a^{2})^{2}}} \right), \quad e_{4}(\tau) = \frac{r^{2} + a^{2}}{|q|^{2}} \left( 1 + \chi(r)\sqrt{1 - \frac{2m^{2}\Delta}{(r^{2} + a^{2})^{2}}} \right), \\ e_{2}(\tau) &= \frac{a\sin\theta}{|q|}, \quad e_{1}(\tau) = 0. \end{split}$$

Together with the formula for the inverse metric

$$\mathbf{g}_{m,a}^{\alpha\beta} = -\frac{1}{2}e_3(x^{\alpha})e_4(x^{\beta}) - \frac{1}{2}e_3(x^{\beta})e_4(x^{\alpha}) + e_1(x^{\alpha})e_1(x^{\beta}) + e_2(x^{\alpha})e_2(x^{\beta})$$

from Lemma 4.1.3 in [21], this allows us to obtain the expression of the inverse Kerr metric in the new coordinates system  $(\tau, r, \theta, \varphi)$ :

$$\begin{aligned} \mathbf{g}_{m,a}^{rr} &= \frac{\Delta}{|q|^2}, \quad \mathbf{g}_{m,a}^{r\varphi} = \frac{a}{|q|^2}, \quad \mathbf{g}_{m,a}^{\varphi\varphi} = \frac{1}{|q|^2 \sin^2 \theta}, \quad \mathbf{g}_{m,a}^{\theta\theta} = \frac{1}{|q|^2}, \\ \mathbf{g}_{m,a}^{\tau\tau} &= \frac{1}{|q|^2} \left( (\chi^2 - 1) \frac{(r^2 + a^2)^2}{\Delta} - 2m^2 \chi^2 + a^2 \sin^2 \theta \right), \\ \mathbf{g}_{m,a}^{\tau r} &= \frac{r^2 + a^2}{|q|^2} \chi \sqrt{1 - \frac{2m^2 \Delta}{(r^2 + a^2)^2}}, \\ \mathbf{g}_{m,a}^{\tau\varphi} &= \frac{a}{|q|^2} \left( 1 + \frac{r^2 + a^2}{\Delta} \left( \chi \sqrt{1 - \frac{2m^2 \Delta}{(r^2 + a^2)^2}} - 1 \right) \right), \\ \mathbf{g}_{m,a}^{r\theta} &= \mathbf{g}_{m,a}^{\theta\varphi} = \mathbf{g}_{m,a}^{\tau\theta} = 0. \end{aligned}$$

We see on the above formulas that the inverse Kerr metric is smooth on the manifold  $\mathbb{R} \times [r_+(1 - \delta_*), +\infty) \times (0, \pi) \times \mathbb{S}^1$ . Moreover, if r < 3m then  $\chi = 1$  and the above formulas show that the inverse Kerr metric is actually analytic, which shows the first part of the lemma. We also easily check that there exists a constant  $c_{m,a} > 0$  only depending on m and a such that  $\mathbf{g}_{m,a}^{\tau\tau} \leq -c_{m,a}$  which shows the second part of the lemma.

Lemma 3.2 implies that the initial data set  $(\Sigma_{m,a}, g_{m,a}, \pi_{m,a})$  satisfies the assumptions of Theorem 3.1 and we immediately deduce the following statement (which is the precise version of Theorem 1.2).

**Theorem 3.3.** Let  $0 \leq |a| \leq m$  and  $(\Sigma_{m,a}, g_{m,a}, \pi_{m,a})$  be the hypersurface and initial data set constructed in Lemma 3.2. Let  $q \in \mathbb{N}^*$  and  $0 < \delta < 1$ . There exists  $\varepsilon_0 = \varepsilon_0(m, a, q, \delta) > 0$  such that if  $0 < \varepsilon \leq \varepsilon_0$ then the following holds: for all  $(\check{g}, \check{\pi})$  with

$$\|\check{g}\|_{H^2_{-a-\delta}(\Sigma_{m,a})} + \|\check{\pi}\|_{H^1_{-a-\delta}(\Sigma_{m,a})} \le \varepsilon,$$

there exists a solution  $(g,\pi)$  of the constraint equations (2.1) on  $\Sigma_{m,a}$  of the form

$$g = u^{4} \left( g_{m,a} + \check{g} + \check{g} \right),$$
  
$$\pi = u^{2} \left( \pi_{m,a} + \check{\pi} + \check{\pi} + L_{q_{m,a} + \check{q} + \check{q}} X \right),$$

where the correctors  $(\check{g}, \check{\pi})$  belong to a vector space of dimension  $4q^2$  only depending on (m, a) and composed of pairs of smooth symmetric 2-tensors compactly supported and where u, X and  $(\check{g}, \check{\pi})$  satisfy

$$\|u-1\|_{H^{2}_{-q-\delta}(\Sigma_{m,a})} + \|X\|_{H^{2}_{-q-\delta}(\Sigma_{m,a})} + \|\check{g}\|_{W^{2,\infty}(\Sigma_{m,a})} + \|\check{\pi}\|_{W^{1,\infty}(\Sigma_{m,a})} \lesssim \varepsilon.$$

**Remark 3.4.** We remark that Theorem 3.3 is true even in the extremal case where |a| = m. However, it is unclear if the data we construct for |a| = m are relevant to black hole stability, in particular seeing as extremal black holes are known to exhibit certain horizon linear instabilities [1, 16].

## 4. PROPERTIES OF THE LINEARIZED OPERATOR

Our first task is to develop a spectral theoretic framework for the following elliptic operator on  $\Sigma$  acting on pairs  $\mathbb{U} = (u, X)$  composed of a scalar function u and a vector field X:

$$P(\mathbb{U}) := \left(-8\Delta_{\bar{g}}u - 2\bar{\pi}^{ij}\mathcal{L}_X\bar{g}_{ij} + \mathrm{tr}_{\bar{g}}\bar{\pi}\mathrm{div}_{\bar{g}}X, -2\mathrm{div}_{\bar{g}}L_{\bar{g}}X_i - 8\bar{g}^{j\ell}\bar{\pi}_{i\ell}\partial_j u + 4\mathrm{tr}_{\bar{g}}\bar{\pi}\partial_i u\right).$$
(4.1)

This operator will naturally appear after we formulate the constraint equations following the conformal ansatz proposed in [11], see Section 5.1. For any  $\mathbb{U} = (u, X)$ , we will also define<sup>6</sup>

$$P^*(\mathbb{U}) \coloneqq \left(-8\Delta_{\bar{g}}u + 4\bar{\pi}^{ij}\mathcal{L}_X\bar{g}_{ij} - 4\operatorname{div}_{\bar{g}}(\operatorname{tr}_{\bar{g}}\bar{\pi}X), -2\operatorname{div}_{\bar{g}}L_{\bar{g}}X_i + 4\bar{\pi}^j_i\partial_j u - \partial_i(\operatorname{tr}_{\bar{g}}\bar{\pi}u)\right),$$
  

$$B_\nu(\mathbb{U}) \coloneqq \left(-8\partial_\nu u + \operatorname{tr}_{\bar{g}}\bar{\pi}\bar{g}(X,\nu) - 4\bar{\pi}(X,\nu), -2D_\nu X\right),$$
  

$$B_\nu^*(\mathbb{U}) \coloneqq \left(-8\partial_\nu u - 4\operatorname{tr}_{\bar{g}}\bar{\pi}\bar{g}(X,\nu) + 8\bar{\pi}(X,\nu), -2D_\nu X\right),$$

<sup>&</sup>lt;sup>6</sup>Directly integrating by parts  $\operatorname{div}_{\bar{g}}L_{\bar{g}}X_i$  would lead to the boundary term  $L_{\bar{g}}X(\nu, \cdot)$ . Here we instead first expand  $\operatorname{div}_{\bar{g}}L_{\bar{g}}X_i = \Delta_{\bar{g}}^{(1)}X_i + \operatorname{Ric}(\bar{g})(\partial_i, X)$  and then integrate by parts to obtain the boundary term  $D_{\nu}X$ .

with D being the covariant derivative associated with  $\bar{g}$  so that

$$\int_{\Sigma} \left( P(\mathbb{U}) \cdot \mathbb{V} - \mathbb{U} \cdot P^*(\mathbb{V}) \right) + \int_{\partial \Sigma} \left( B_{\nu}(\mathbb{U}) \cdot \mathbb{V} - \mathbb{U} \cdot B_{\nu}^*(\mathbb{V}) \right) = 0, \tag{4.2}$$

where  $\mathbb{U} \cdot \mathbb{V} = uv + \bar{g}(X, Y)$  for  $\mathbb{U} = (u, X)$  and  $\mathbb{V} = (v, Y)$ , and where  $\nu$  denotes the inward pointing unit normal to  $\partial \Sigma$  with respect to  $\bar{g}$ .

We will consider perturbations of the boundary operators  $B_{\nu}$  and  $B_{\nu}^*$ . For b a scalar function,  $\omega$  a 1-form, Z a vector field and  $\xi$  a (1,1)-tensor, we define  $F = (b, \omega, Z, \xi)$  so that F acts on the space of pairs  $\mathbb{U} = (u, X)$  of functions and vector fields through the formula

$$F(\mathbb{U}) = (bu + \omega(X), uZ + \xi(X))$$

We define the space of such smooth F restricted to  $\partial \Sigma$  by  $\mathcal{E}(\partial \Sigma)$ . If  $F \in \mathcal{E}(\partial \Sigma)$  we define its transpose  $F^* \in \mathcal{E}(\partial \Sigma)$  by the formula  $F^* = (b, Z^{\flat}, \omega^{\sharp}, \xi^{\natural})$ , where  $\xi^{\natural}$  is defined in coordinates by  $(\xi^{\natural})_i^{\ j} = \bar{g}^{j\ell} \xi_{\ell}^{\ k} \bar{g}_{ki}$ . This definition is such that

$$F(\mathbb{U}) \cdot \mathbb{V} = \mathbb{U} \cdot F^*(\mathbb{V}). \tag{4.3}$$

Also, we define  $\mathrm{Id} \in \mathcal{E}(\partial \Sigma)$  by  $\mathrm{Id} = (1, 0, 0, \delta)$  so that  $\mathrm{Id}(\mathbb{U}) = \mathbb{U}$ .

**Definition 4.1.** For  $F \in \mathcal{E}(\partial \Sigma)$  and  $\delta \in \mathbb{R}$  we define

$$P_{F,\delta} := \left( (P, B_{\nu} + F) : H^2_{\delta}(\Sigma) \longrightarrow L^2_{\delta-2}(\Sigma) \times H^{\frac{1}{2}}(\partial\Sigma) \right), \tag{4.4}$$

$$P_{F,\delta}^* := \left( (P^*, B_{\nu}^* + F^*) : H_{\delta}^2(\Sigma) \longrightarrow L_{\delta-2}^2(\Sigma) \times H^{\frac{1}{2}}(\partial\Sigma) \right).$$

$$(4.5)$$

Note that (4.2) and (4.3) imply

$$\langle P(\mathbb{U}), \mathbb{V} \rangle_{L^2(\Sigma)} + \langle (B_{\nu} + F)(\mathbb{U}), \mathbb{V} \rangle_{L^2(\partial \Sigma)} = \langle \mathbb{U}, P^*(\mathbb{V}) \rangle_{L^2(\Sigma)} + \langle \mathbb{U}, (B_{\nu}^* + F^*)(\mathbb{V}) \rangle_{L^2(\partial \Sigma)}.$$
(4.6)

4.1. **Basic properties.** We gather in the following proposition general facts about the operators (4.4) and (4.5), which follow from  $P_{f,\delta}$  and  $P_{f,\delta}^*$  being compact perturbations of the Laplace-Beltrami operator (acting on both scalar functions and vector fields) with Neumann boundary conditions.

**Proposition 4.2.** If  $\delta \notin \mathbb{Z}$  and  $F \in \mathcal{E}(\partial \Sigma)$ , then:

(i) There exists  $C_{F,\delta} > 0$  such that for all  $\mathbb{U} \in H^2_{\delta}(\Sigma)$  we have

$$\|\mathbb{U}\|_{H^{2}_{\delta}(\Sigma)} \leq C_{F,\delta} \left( \|P(\mathbb{U})\|_{L^{2}_{\delta-2}(\Sigma)} + \|(B_{\nu} + F)(\mathbb{U})\|_{H^{\frac{1}{2}}(\partial\Sigma)} + \|\mathbb{U}\|_{L^{2}_{\delta}(\Sigma)} \right),$$

$$\|\mathbb{U}\|_{H^{2}_{\delta}(\Sigma)} \leq C_{F,\delta} \left( \|P^{*}(\mathbb{U})\|_{L^{2}_{\delta-2}(\Sigma)} + \|(B^{*}_{\nu} + F^{*})(\mathbb{U})\|_{H^{\frac{1}{2}}(\partial\Sigma)} + \|\mathbb{U}\|_{L^{2}_{\delta}(\Sigma)} \right).$$

$$(4.7)$$

(ii) The operator  $P_{F,\delta}$  and  $P_{F,\delta}^*$  are Fredholm and if  $-1 < \delta < 0$  their index is zero.

*Proof.* We consider the rescaled Neumann Laplace-Beltrami operator of  $\overline{g}$  given by

$$\begin{split} \Delta_N : H^2_{\delta}(\Sigma) &\longrightarrow L^2_{\delta-2}(\Sigma) \times H^{\frac{1}{2}}(\partial\Sigma), \\ (u, X) &\longmapsto \left( \left( -8\Delta_{\bar{g}}u, -2\Delta_{\bar{g}}^{(1)}X \right), \left( -8\partial_{\nu}u, -2D_{\nu}X \right) \right), \end{split}$$

where  $\Delta_{\bar{g}}^{(1)}$  is the vectorial Laplace-Beltrami operators. Since  $\Delta_{\bar{g}}^{(1)}$  is a compact perturbation of  $\Delta_{\bar{g}}$  acting on components of vector fields in a given coordinate system, we deduce from Proposition 4.13 in [24] the following estimate

$$\|(u,X)\|_{H^{2}_{\delta}(\Sigma)} \lesssim \|\Delta_{N}(u,X)\|_{L^{2}_{\delta-2}(\Sigma) \times H^{\frac{1}{2}}(\partial\Sigma)} + \|(u,X)\|_{L^{2}_{\delta}(\Sigma)}.$$
(4.8)

Since both  $P_{F,\delta}$  and  $P_{F,\delta}^*$  are compact perturbations of  $\Delta_N$ , we deduce (4.7) from (4.8) using interpolation to control lower order terms. Using again the fact that  $P_{F,\delta}$  and  $P_{F,\delta}^*$  are compact perturbations of  $\Delta_N$ , the second part of the lemma follows from the fact that  $\Delta_N$  is Fredholm if  $\delta \notin \mathbb{Z}$  and is of index 0 if  $-1 < \delta < 0$ , see Propositions 1 and 6 in [25]. The adjoint  $(P_{F,\delta})^*$  of  $P_{F,\delta}$  is defined on the space  $L^2_{-1-\delta}(\Sigma) \times H^{-\frac{1}{2}}(\partial \Sigma)$  by the relation

$$\langle P(\mathbb{U}), \mathbb{V} \rangle_{L^2(\Sigma)} + \langle (B_{\nu} + F)(\mathbb{U}), \mathbb{h} \rangle_{L^2(\partial \Sigma)} = \langle \mathbb{U}, (P_{F,\delta})^*(\mathbb{V}, \mathbb{h}) \rangle_{L^2(\Sigma)}.$$
(4.9)

The adjoint  $(P_{F,\delta}^*)^*$  of  $P_{F,\delta}^*$  is defined similarly. In order to characterize the mapping properties of  $P_{F,\delta}$  and  $P_{F,\delta}^*$ , we will use the following lemma that characterizes the kernels of their adjoints.

**Lemma 4.3.** If  $\delta \notin \mathbb{Z}$  and  $F \in \mathcal{E}(\partial \Sigma)$  then

$$\ker\left(\left(P_{F,\delta}\right)^{*}\right) = \left\{\left(\mathbb{V}, \mathbb{V}_{\mid \partial \Sigma}\right) \mid \mathbb{V} \in \ker\left(P_{F,-1-\delta}^{*}\right)\right\},\tag{4.10}$$

$$\ker\left(\left(P_{F,\delta}^{*}\right)^{*}\right) = \left\{\left(\mathbb{V}, \mathbb{V}_{\mid \partial \Sigma}\right) \mid \mathbb{V} \in \ker\left(P_{F,-1-\delta}\right)\right\}.$$
(4.11)

*Proof.* We will only prove (4.10) since (4.11) follows by a similar argument. Let  $(\mathbb{V}, \mathbb{h}) \in L^2_{-1-\delta}(\Sigma) \times H^{-\frac{1}{2}}(\partial \Sigma)$ . From (4.6) and (4.9), we have that

$$\langle \mathbb{U}, (P_{F,\delta})^*(\mathbb{V}, \mathbb{h}) \rangle_{L^2(\Sigma)} = \langle \mathbb{U}, P^*(\mathbb{V}) \rangle_{L^2(\Sigma)} + \langle \mathbb{U}, (B^*_{\nu} + F^*)(\mathbb{V}) \rangle_{L^2(\partial\Sigma)} + \langle (B_{\nu} + F)(\mathbb{U}), \mathbb{h} - \mathbb{V} \rangle_{L^2(\partial\Sigma)}$$

for all  $\mathbb{U} \in H^2_{\delta}(\Sigma)$ . Then it is immediately clear that

$$\left\{ \left(\mathbb{V}, \mathbb{V}_{\mid \partial \Sigma}\right) \mid \mathbb{V} \in \ker\left(P_{F, -1-\delta}^*\right) \right\} \subset \ker\left(\left(P_{F, \delta}\right)^*\right).$$

We now prove the reverse inclusion. If  $(P_{F,\delta})^*(\mathbb{V},\mathbb{h}) = 0$ , then

$$\langle \mathbb{U}, P^*(\mathbb{V}) \rangle_{L^2(\Sigma)} + \langle \mathbb{U}, (B^*_{\nu} + F)(\mathbb{V}) \rangle_{L^2(\partial \Sigma)} + \langle (B_{\nu} + F)(\mathbb{U}), \mathbb{h} - \mathbb{V} \rangle_{L^2(\partial \Sigma)} = 0$$
(4.12)

for all  $\mathbb{U} \in H^2_{\delta}(\Sigma)$ . Taking  $\mathbb{U}$  compactly supported away from  $\partial \Sigma$ , the boundary terms disappear and we get  $P^*(\mathbb{V}) = 0$  on  $\Sigma$ . Since  $P^*(\mathbb{V}) = 0$ , then (4.12) becomes

$$\langle \mathbb{U}, (B_{\nu}^* + F^*)(\mathbb{V}) \rangle_{L^2(\partial \Sigma)} + \langle (B_{\nu} + F)(\mathbb{U}), \mathbb{h} - \mathbb{V} \rangle_{L^2(\partial \Sigma)} = 0,$$

and this should hold for every  $\mathbb{U} \in H^2_{\delta}(\Sigma)$ . In particular, it holds for every  $\mathbb{U} \in H^2_{\delta}(\Sigma)$  such that  $(B_{\nu} + F)(\mathbb{U}) = 0$  on  $\partial\Sigma$ , thus implying that  $(B^*_{\nu} + F^*)(\mathbb{V}) = 0$  on  $\partial\Sigma$ . Thus, using the elliptic regularity estimate (4.7) we infer from  $P^*(\mathbb{V}) = 0$  and  $(B^*_{\nu} + F^*)(\mathbb{V}) = 0$  that  $\mathbb{V} \in H^2_{-1-\delta}(\Sigma)$ . Therefore, for every  $\mathbb{U} \in H^2_{\delta}(\Sigma)$  we have

$$\langle (B_{\nu} + F)(\mathbb{U}), \mathbb{h} - \mathbb{V} \rangle_{L^{2}(\partial \Sigma)} = 0$$

which implies  $\mathbb{h} = \mathbb{V}$  on  $\partial \Sigma$ . Thus we have shown that

$$\operatorname{ker}\left(\left(P_{F,\delta}\right)^{*}\right) \subset \left\{\left(\mathbb{V},\mathbb{V}_{\mid \partial \Sigma}\right) \mid \mathbb{V} \in \operatorname{ker}\left(P_{F,-1-\delta}^{*}\right)\right\}$$

This concludes the proof of (4.10).

In the next lemma, we give a unique continuation result from the boundary for P and  $P^*$ .

**Lemma 4.4.** Let  $\Gamma$  be an open subset of  $\partial \Sigma$ . If  $\mathbb{U} \in H^2_{loc}(\Sigma)$  satisfies  $P(\mathbb{U}) = 0$  (resp.  $P^*(\mathbb{U}) = 0$ ) on  $\Sigma$  and  $B_{\nu}(\mathbb{U}) = \mathbb{U} = 0$  (resp.  $B^*_{\nu}(\mathbb{U}) = \mathbb{U} = 0$ ) on  $\Gamma$ , then  $\mathbb{U} = 0$  on  $\Sigma$ .

*Proof.* The proof follows along the same lines as the proof of Theorem 9.8 in [22], which is the analogue of Lemma 4.4 in the scalar case. Since P and  $P^*$  are diagonal operator at leading order, we can sum the four Carleman's estimates given by the proof of Theorem 9.8 in [22] and absorb the lower order coupling terms between the four components of  $\mathbb{U}$ .

4.2. A special boundary condition. In this section, we will construct a distinguished boundary condition so that  $P_{F,\delta}$  as defined in (4.1) has good mapping properties for any fixed  $\delta \in (0, 1)$ .

**Lemma 4.5.** Let  $F_0 \in \mathcal{E}(\partial \Sigma)$ . There exists  $\eta(F_0) > 0$  only depending on  $F_0$  such that for all  $F \in \mathcal{E}(\partial \Sigma)$  on  $\partial \Sigma$ ,

$$\|F - F_0\|_{H^{\frac{1}{2}}(\partial \Sigma)} \le \eta(F_0) \Longrightarrow \dim(\ker(P_{F,-\delta})) \le \dim(\ker(P_{F_0,-\delta})).$$

Proof. Lemma 4.5 follows immediately from Proposition 1.11 in [2] and the fact that

$$\|P_{F,-\delta} - P_{F_0,-\delta}\|_{\text{op}} \lesssim \|F - F_0\|_{H^{\frac{1}{2}}(\partial \Sigma)}$$

where  $\|\cdot\|_{op}$  refers to the standard operator norm.

In the next lemma, we show how to lower the dimension of the kernel in a constructive way.

**Lemma 4.6.** Let  $\mathcal{W}$  be an open subset of  $\partial \Sigma$  such that  $\partial \Sigma \setminus \mathcal{W}$  contains an open set. Let  $F_0 \in \mathcal{E}(\partial \Sigma)$  such that  $\dim(\ker(P_{F_0,-\delta})) > 0$ . Then there exists  $F_1 \in \mathcal{E}(\partial \Sigma)$  such that

$$\dim(\ker(P_{F_1,-\delta})) < \dim(\ker(P_{F_0,-\delta})), \quad and \quad (F_1 - F_0)|_{\mathcal{W}} = 0.$$
(4.13)

Proof. Using Proposition 4.2, the Fredholm index of  $P_{F_0,-\delta}$  is zero for  $\delta \in (0,1)$ . Thus, the assumption that dim $(\ker(P_{F_0,-\delta})) > 0$  implies that dim $(\ker((P_{F_0,-\delta})^*)) > 0$ . According to (4.10) this implies that dim $\left(\ker\left(P_{F_0,-1+\delta}^*\right)\right) > 0$ . Now consider  $(\mathbb{V}_0, \mathbb{V}_0^*) \in \ker(P_{F_0,-\delta}) \times \ker(P_{F_0,-1+\delta}^*)$  such that  $\mathbb{V}_0, \mathbb{V}_0^*$  are both non-zero. Since they satisfy  $(P, B_{\nu} + F_0)(\mathbb{V}_0) = 0$  and  $(P^*, B_{\nu}^* + F_0^*)(\mathbb{V}_0^*) = 0$ , unique continuation for the operators P and  $P^*$  (recall Lemma 4.4) implies that

$$\{\mathbb{V}_0 \neq 0\} \cap \{\mathbb{V}_0^* \neq 0\} \cap \partial \Sigma$$

is a dense open subset of  $\partial \Sigma$ . Thus, there exists  $H_0 \in \mathcal{E}(\partial \Sigma)$  such that  $H_{0|_{\mathcal{W}}} = 0$  and

$$\int_{\partial \Sigma} H_0(\mathbb{V}_0) \cdot \mathbb{V}_0^* = 1.$$
(4.14)

For  $\eta > 0$ , we define

$$F_{\eta} := F_0 + \eta \frac{H_0}{\|H_0\|_{H^{\frac{3}{2}}(\partial \Sigma)}}.$$

We will show that there exists some  $\eta$  sufficiently small such that dim  $(\ker(P_{F_{\eta},-\delta})) < \dim(\ker(P_{F_{0},-\delta}))$ . For  $\eta \leq \eta(F_{0})$ , Lemma 4.5 already implies that dim $(\ker(P_{F_{\eta},-\delta})) \leq \dim(\ker(P_{F_{0},-\delta}))$ . Now assume for the sake of contradiction that dim $(\ker(P_{F_{\eta},-\delta})) = \dim(\ker(P_{F_{0},-\delta}))$ . From Proposition 1.12 in [2] there exists  $0 < \eta'(F_{0}) \leq \eta(F_{0})$  and  $C(F_{0}) > 0$  such that if  $\eta \leq \eta'(F_{0})$  then

$$\left\| \mathbb{U} - \ker(P_{F_{\eta}, -\delta}) \right\|_{H^{2}_{-\delta}(\Sigma)} \le C(F_{0}) \left( \|P(\mathbb{U})\|_{L^{2}_{-\delta-2}(\Sigma)} + \|(B_{\nu} + F_{\eta})(\mathbb{U})\|_{H^{\frac{1}{2}}(\partial\Sigma)} \right),$$

for all  $\mathbb{U} \in H^2_{-\delta}(\Sigma)$ . We apply this to  $\mathbb{V}_0 \in \ker(P_{F_0,-\delta})$  as chosen above. Since  $\ker(P_{F_\eta,-\delta})$  is finitedimensional, there exists some  $\mathbb{U}_\eta \in \ker(P_{F_\eta,-\delta})$  such that

$$\|\mathbb{V}_0 - \mathbb{U}_\eta\|_{H^2_{-\delta}(\Sigma)} \lesssim C(F_0) \, \|\mathbb{V}_0\|_{H^2_{-\delta}(\Sigma)} \, \eta, \tag{4.15}$$

where we have used the continuous embeddings  $H^2_{-\delta}(\Sigma) \hookrightarrow H^{\frac{3}{2}}(\partial \Sigma) \hookrightarrow L^{\infty}(\partial \Sigma)$ . Then using (4.2), the fact that  $\mathbb{V}_0^* \in \ker(P_{F_0,-1+\delta}^*)$ , and that  $\mathbb{U}_\eta \in \ker(P_{F_\eta,-\delta})$ , we compute

$$\frac{\|H_0\|_{H^{\frac{3}{2}}(\partial\Sigma)}}{\eta} \int_{\Sigma} \left( P(\mathbb{U}_\eta) \cdot \mathbb{V}_0^* - \mathbb{U}_\eta \cdot P^*(\mathbb{V}_0^*) \right) = -\int_{\partial\Sigma} H_0(\mathbb{U}_\eta) \cdot \mathbb{V}_0^*$$

Since  $P(\mathbb{U}_{\eta}) = P^*(\mathbb{V}_0^*) = 0$  we have proved that

$$\int_{\partial \Sigma} H_0(\mathbb{U}_\eta) \cdot \mathbb{V}_0^* = 0$$

However (4.14) and (4.15) imply

$$\left| \int_{\partial \Sigma} H_0(\mathbb{U}_{\eta}) \cdot \mathbb{V}_0^* - 1 \right| \le CC(F_0) \, \|H_0\|_{H^{\frac{3}{2}}(\partial \Sigma)} \, \|\mathbb{V}_0^*\|_{H^{2}_{-\delta}(\Sigma)} \, \|\mathbb{V}_0\|_{H^{2}_{-\delta}(\Sigma)} \, \eta,$$

for some universal constant C > 0. Choosing

$$\eta_1(F_0) := \min\left(\eta'(F_0), \frac{1}{2CC(F_0) \|H_0\|_{H^{\frac{3}{2}}(\partial\Sigma)} \|\mathbb{V}_0^*\|_{H^2_{-\delta}(\Sigma)} \|\mathbb{V}_0\|_{H^2_{-\delta}(\Sigma)}}\right)$$

and taking  $0 < \eta \leq \eta_1(F_0)$  implies a contradiction. This proves that  $\dim(\ker(P_{F_{\eta},-\delta})) < \dim(\ker(P_{F_0,-\delta}))$ and concludes the proof of the lemma, after setting  $F_1 := F_{\eta}$ .

We can iterate the result of Lemma 4.6 to obtain the following corollary.

**Corollary 4.7.** Let  $\mathcal{U}$  and  $\mathcal{V}$  two open subsets of  $\partial \Sigma$  with  $\overline{\mathcal{U}} \cap \overline{\mathcal{V}} = \emptyset$ . There exists  $F_{\delta} \in \mathcal{E}(\partial \Sigma)$  such that  $F_{\delta|_{\mathcal{V}}} = \text{Id}, F_{\delta|_{\mathcal{U}}} = 0$  and

$$\ker(P_{F_{\delta},-\delta}) = \{0\},\tag{4.16}$$

$$\ker(P^*_{F_{\delta},-1+\delta}) = \{0\}. \tag{4.17}$$

*Proof.* First, note that (4.16) implies (4.17). Indeed, from (4.10), we have that dim  $\left(\ker\left(P_{F_{\delta},-1+\delta}^*\right)\right) = \dim\left(\ker\left(\left(P_{F_{\delta},-\delta}^*\right)^*\right)\right)$ . In view of the fact that  $P_{F_{\delta},-\delta}$  has Fredholm index zero from Proposition 4.2 this becomes

 $\dim\left(\ker\left(P_{F_{\delta},-1+\delta}^{*}\right)\right) = \dim\left(\ker\left(P_{F_{\delta},-\delta}\right)\right).$ 

Therefore, it only remains to construct  $F_{\delta}$  satisfying the requirements of the corollary and such that (4.16) holds. Since  $\overline{\mathcal{U}} \cap \overline{\mathcal{V}} = \emptyset$ , we may consider  $\chi_b$  a smooth function on  $\partial \Sigma$  such that  $\chi_{b|_{\mathcal{U}}} = 0$  and  $\chi_{b|_{\mathcal{V}}} = 1$  and set  $F_0 := \chi_b \text{Id}$ . Let  $n_0 := \dim(\ker(P_{F_0,-\delta}))$ . If  $n_0 = 0$ , the lemma is proved. If  $n_0 > 0$ , we construct iteratively a finite family of matrices  $(F_0, F_1, \ldots, F_{n_*})$  in the following way:

- (1) We apply Lemma 4.6 to  $F_0$  and  $\mathcal{W} := \mathcal{U} \cup \mathcal{V}$  (since  $\overline{\mathcal{U}} \cap \overline{\mathcal{V}} = \emptyset$ ,  $\partial \Sigma \setminus \mathcal{W}$  does indeed contain an open set), which gives the existence of  $F_1 \in \mathcal{E}(\partial \Sigma)$  with dim $(\ker(P_{F_1,-\delta})) < n_0$  and  $F_{1|_{\mathcal{W}}} = 0$ .
- (2) If  $(F_0, F_1, \ldots, F_j)$  is constructed, then we distinguish two cases:
  - (a) If dim $(\ker(P_{F_j,-\delta})) = 0$ , then set  $n_* = j$  and the iteration ends.
  - (b) Otherwise,  $\dim(\ker(P_{F_j,-\delta})) > 0$ , and Lemma 4.6 again implies the existence of  $F_{j+1} \in \mathcal{E}(\partial\Sigma)$  such that  $\dim(\ker(P_{F_{j+1},-\delta})) < \dim(\ker(P_{F_j,-\delta}))$  and  $(F_{j+1}-F_j)|_{W} = 0$ .

Since  $(\dim(\ker(P_{F_j,-\delta})))_j$  is a decreasing sequence of integers and since  $\dim(\ker(P_{F_0,-\delta})) = n_0$ , we have that  $n_* \leq n_0$ . We thus have a finite family  $(F_0, F_1, \ldots, F_{n_*})$  in  $\mathcal{E}(\partial \Sigma)$  with  $\ker(P_{F_{n_*},-\delta}) = \{0\}$ . Moreover, we can prove by induction that  $(F_{n_*} - F_0)_{|u \cup \nu} = 0$ . Since  $F_{0|\nu} = \operatorname{Id}$  and  $F_{0|\mu} = 0$ , this proves that  $F_{\delta} := F_{n_*}$  does satisfy the requirements of the lemma.

4.3. Adapted harmonic polynomials. We recall the definition of harmonic polynomials and vectorial harmonic polynomials in Euclidean space.

**Definition 4.8.** We identify the global coordinates of  $\mathbb{R}^3$  and  $\Sigma$  and consider the real spherical harmonics  $Y_{j,\ell}$  for  $j \ge 0$  and  $-j \le \ell \le j$  with an arbitrary normalization.

(i) For  $j \ge 1$  and  $-(j-1) \le \ell \le j-1$ , we define scalar functions

$$w_{j,\ell} := r^{j-1} Y_{j-1,\ell},$$

and if in addition k = 1, 2, 3, we define vector fields

$$W_{j,\ell,k} := w_{j,\ell} \partial_k.$$

(ii) For  $j \ge 1, -j \le \ell \le j$  and  $\alpha = 0, 1, 2, 3$ , we define pairs consisting of a scalar function and a vector field

$$\mathbb{W}_{j,\ell,\alpha}^{(e)} := \begin{cases} (w_{j,\ell}, 0), & \text{if } \alpha = 0, \\ (0, W_{j,\ell,\alpha}), & \text{if } \alpha = 1, 2, 3 \end{cases}$$

**Remark 4.9.** Note that for each  $q \in \mathbb{N}^*$  the family

$$\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right)_{1\leq j\leq q,-(j-1)\leq \ell\leq j-1,\alpha=0,1,2,3}$$

is a basis of homogeneous polynomials of degree  $1 \leq j \leq q$  moreover satisfying  $\Delta_e\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right) = 0$  where  $\Delta_e$  denotes here the Euclidean Laplacian.

Using the harmonic polynomials of Definition 4.8, we construct a family of harmonic polynomials adapted to  $P_{F,\delta}$ .

**Lemma 4.10.** Let  $j \ge 1$ ,  $-(j-1) \le \ell \le j-1$  and  $\alpha = 0, 1, 2, 3$ . There exists  $\widetilde{\mathbb{W}}_{j,\ell,\alpha} \in H^2_{j-2+\delta}(\Sigma)$  such that

$$\mathbb{W}_{j,\ell,\alpha} := \mathbb{W}_{j,\ell,\alpha}^{(e)} + \widetilde{\mathbb{W}}_{j,\ell,\alpha}$$

satisfies

$$\begin{cases} P^* \left( \mathbb{W}_{j,\ell,\alpha} \right) = 0, & \text{on } \Sigma, \\ \left( B_{\nu}^* + F_{\delta}^* \right) \left( \mathbb{W}_{j,\ell,\alpha} \right) = 0, & \text{on } \partial \Sigma. \end{cases}$$

*Proof.* It suffices to solve for  $\widetilde{\mathbb{W}}_{j,\ell,\alpha} \in H^2_{i-2+\delta}(\Sigma)$  such that

$$\begin{cases} P^*\left(\widetilde{\mathbb{W}}_{j,\ell,\alpha}\right) = -P^*\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right), & \text{on } \Sigma, \\ \left(B_{\nu}^* + F_{\delta}^*\right)\left(\widetilde{\mathbb{W}}_{j,\ell,\alpha}\right) = -\left(B_{\nu}^* + F_{\delta}^*\right)\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right), & \text{on } \partial\Sigma. \end{cases}$$

Now, observe that  $P^*$  is asymptotic to the Euclidean Laplacian in view of Definition 2.1. As a result, since  $\mathbb{W}_{j,\ell,\alpha}^{(e)} \in H_{j-1+\delta}^4(\Sigma)$ , we have that

$$P^*\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right) \in H^2_{j-4+\delta}(\Sigma)$$

since  $\Delta_e\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right) = 0$  in view of Remark 4.9. We can thus rewrite the above system for  $\widetilde{\mathbb{W}}_{j,\ell,\alpha}$  as

$$P_{F_{\delta},j-2+\delta}^{*}\left(\widetilde{\mathbb{W}}_{j,\ell,\alpha}\right) = -\left(P^{*}\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right), \left(B_{\nu}^{*}+F_{\delta}^{*}\right)\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right)\right)$$

Since  $j \ge 1$  we have  $\ker(P_{F_{\delta},1-j-\delta}) \subset \ker(P_{F_{\delta},-\delta}) = \{0\}$ , where we used (4.16). According to (4.11) this implies  $\ker\left(\left(P_{F_{\delta},j-2+\delta}^*\right)^*\right) = \{0\}$ , and hence that  $P_{F_{\delta},j-2+\delta}^*$  is surjective. Therefore we can always solve for  $\widetilde{W}_{j,\ell,\alpha}$ .

Using the adapted harmonic polynomials constructed in Lemma 4.10, we are now able to specify the orthogonality conditions necessary to invert  $P_{F_{\delta},-q-\delta}$ .

**Lemma 4.11.** Let  $q \in \mathbb{N}^*$ ,  $0 < \delta < 1$  and let  $\mathbb{Y} \in L^2_{-q-\delta-2}(\Sigma)$ . Then  $(\mathbb{Y}, 0) \in \operatorname{im}(P_{F_{\delta}, -q-\delta})$  if and only if

$$\left\langle \mathbb{W}_{j,\ell,\alpha},\mathbb{Y}\right\rangle_{L^{2}(\Sigma)}=0,\quad for \ all \ 1\leq j\leq q, \ -(j-1)\leq \ell\leq j-1 \ and \ \alpha=0,1,2,3$$

*Proof.* We start by proving that

$$\ker \left( P_{F_{\delta},q-1+\delta}^* \right) = \operatorname{span} \left( \mathbb{W}_{j,\ell,\alpha}, 1 \le j \le q, \ -(j-1) \le \ell \le j-1 \text{ and } \alpha = 0, 1, 2, 3 \right),$$
(4.18)

for all  $q \ge 1$ . The inclusion from right to left is immediate by definition of  $\mathbb{W}_{j,\ell,\alpha}$  in the Lemma 4.10 and the fact that  $j \le q$ . For the inclusion from left to right we proceed by induction on  $q \ge 1$ :

- For q = 1, let  $\mathbb{W} \in \ker\left(P_{F_{\delta},\delta}^*\right)$ . Since  $P^*$  is asymptotic to the Euclidean Laplacian  $\Delta$ , Theorem 1.17 in [2] and Remark 4.9 imply that there exists scalars  $\lambda^{\alpha}$  such that  $\mathbb{W} \lambda^{\alpha} \mathbb{W}_{1,0,\alpha}^{(e)} \in H^2_{-1+\delta}(\Sigma)$ . Lemma 4.10 thus implies that  $\mathbb{W} - \lambda^{\alpha} \mathbb{W}_{1,0,\alpha} \in \ker\left(P_{F_{\delta},-1+\delta}^*\right)$ . Thanks to (4.17), we obtain  $\mathbb{W} = \lambda^{\alpha} \mathbb{W}_{1,0,\alpha}$ . This proves (4.18) for q = 1.
- Assume that we have

$$\ker \left( P_{F_{\delta}, q-1+\delta}^* \right) = \operatorname{span} \left( \mathbb{W}_{j,\ell,\alpha}, 1 \le j \le q, \ -(j-1) \le \ell \le j-1 \text{ and } \alpha = 0, 1, 2, 3 \right)$$
(4.19)

for some  $q \ge 1$  and let  $\mathbb{W} \in \ker\left(P_{F_{\delta},q+\delta}^*\right)$ . From Theorem 1.17 in [2] and Remark 4.9 we again obtain the existence of scalars  $\lambda^{\ell,\alpha}$  such that  $\mathbb{W} - \lambda^{\ell,\alpha}\mathbb{W}_{q+1,\ell,\alpha}^{(e)} \in H_{q-1+\delta}^2(\Sigma)$ . Lemma 4.10 thus implies that  $\mathbb{W} - \lambda^{\ell,\alpha}\mathbb{W}_{q+1,\ell,\alpha} \in \ker\left(P_{F_{\delta},q-1+\delta}^*\right)$ . From (4.19) this gives

$$\mathbb{W} - \lambda^{\ell,\alpha} \mathbb{W}_{q+1,\ell,\alpha} \in \operatorname{span} \left( \mathbb{W}_{j,\ell,\alpha}, 1 \le j \le q, \, -(j-1) \le \ell \le j-1 \text{ and } \alpha = 0, 1, 2, 3 \right)$$

and thus

$$\mathbb{W} \in \text{span}(\mathbb{W}_{j,\ell,\alpha}, 1 \le j \le q+1, -(j-1) \le \ell \le j-1 \text{ and } \alpha = 0, 1, 2, 3)$$

This concludes the proof of (4.18). Now, by definition of the adjoint,  $(\mathbb{Y}, 0) \in \operatorname{im}(P_{F_{\delta}, -q-\delta})$  if and only if  $\langle \mathbb{Y}, \mathbb{V} \rangle_{L^{2}(\Sigma)} = 0$  for all  $\mathbb{V}$  such that  $(\mathbb{V}, \mathbb{h}) \in \ker((P_{F_{\delta}, -q-\delta})^{*})$  for some  $\mathbb{h}$ . Thanks to (4.10) we obtain

$$(\mathbb{Y},0) \in \operatorname{im}(P_{F_{\delta},-q-\delta}) \iff \langle \mathbb{Y},\mathbb{V} \rangle_{L^{2}(\Sigma)} = 0, \quad \text{for all } \mathbb{V} \in \ker\left(P_{F_{\delta},q-1+\delta}^{*}\right).$$

Using (4.18) concludes the proof of Lemma 4.11.

4.4. Conclusion. We summarize the properties of the operator P in (4.1) proved so far.

**Proposition 4.12.** Let  $q \in \mathbb{N}^*$  and  $0 < \delta < 1$ . There exists  $F_{\delta} \in \mathcal{E}(\partial \Sigma)$  and a family

 $(\mathbb{W}_{j,\ell,\alpha})_{1 \le j \le q, -(j-1) \le \ell \le j-1, \alpha = 0, 1, 2, 3}$ (4.20)

with the following properties

- (i) there exists  $\mathcal{U}$  and  $\mathcal{V}$  open subsets of  $\partial \Sigma$  such that  $F_{\delta|_{\mathcal{V}}} = \text{Id}$  and  $F_{\delta|_{\mathcal{U}}} = 0$ ,
- (ii) the family (4.20) is linearly independent and each  $\mathbb{W}_{j,\ell,\alpha}$  satisfies the estimate  $|\mathbb{W}_{j,\ell,\alpha}| \lesssim r^{j-1}$ and the system

$$\begin{cases} P^* \left( \mathbb{W}_{j,\ell,\alpha} \right) = 0, & \text{on } \Sigma, \\ \left( B^*_{\nu} + F^*_{\delta} \right) \left( \mathbb{W}_{j,\ell,\alpha} \right) = 0, & \text{on } \partial \Sigma, \end{cases}$$

(iii) if  $\mathbb{Y} \in L^2_{-a-\delta-2}(\Sigma)$  satisfies

$$\langle \mathbb{W}_{j,\ell,\alpha}, \mathbb{Y} \rangle_{L^2(\Sigma)} = 0, \quad \text{for all } 1 \le j \le q, \ -(j-1) \le \ell \le j-1 \text{ and } \alpha = 0, 1, 2, 3,$$
 (4.21)

then there exists a unique  $\mathbb{U} \in H^2_{-q-\delta}(\Sigma)$  with  $\|\mathbb{U}\|_{H^2_{-q-\delta}(\Sigma)} \lesssim \|\mathbb{Y}\|_{L^2_{-q-\delta-2}(\Sigma)}$  such that

$$\begin{cases} P(\mathbb{U}) = \mathbb{Y}, & on \Sigma, \\ (B_{\nu} + F_{\delta})(\mathbb{U}) = 0, & on \partial \Sigma. \end{cases}$$

*Proof.* We consider  $F_{\delta}$  as constructed in Corollary 4.7. The family (4.20) has been constructed in Lemma 4.10 and its linear independence follows from the linear independence of the family

$$\left(\mathbb{W}_{j,\ell,\alpha}^{(e)}\right)_{1\leq j\leq q,-(j-1)\leq \ell\leq j-1,\alpha=0,1,2,3}$$

and the asymptotics  $\mathbb{W}_{j,\ell,\alpha} - \mathbb{W}_{j,\ell,\alpha}^{(e)} = O\left(r^{j-2+\delta}\right)$  at infinity. If  $\mathbb{Y} \in L^2_{-q-\delta-2}(\Sigma)$  satisfies (4.21) then Lemma 4.11 implies that  $(\mathbb{Y}, 0) \in \operatorname{im}(P_{F_{\delta}, -q-\delta})$ , so that there exists  $\mathbb{U} \in H^2_{-q-\delta}(\Sigma)$  solving  $P_{F_{\delta}, -q-\delta}(\mathbb{U}) = (\mathbb{Y}, 0)$ . Uniqueness follows from the choice of  $F_{\delta}$  (Corollary 4.7) which implies that  $\ker(P_{F_{\delta}, -q-\delta}) \subset \ker(P_{F_{\delta}, -\delta}) = \{0\}$ . Finally, the estimate  $\|\mathbb{U}\|_{H^2_{-q-\delta}(\Sigma)} \lesssim \|\mathbb{Y}\|_{L^2_{-q-\delta-2}(\Sigma)}$  follows from (1.31) in [2].  $\Box$ 

## 5. Proof of the main theorem

This section is devoted to the proof of Theorem 3.1.

5.1. Conformal formulation of the constraint equations. Let  $(\Sigma, \bar{g}, \bar{\pi})$  an asymptotically flat initial data set satisfying the regularity assumptions of Theorem 3.1, and let  $(\check{g}, \check{\pi}) \in H^2_{-q-\delta}(\Sigma) \times H^1_{-q-\delta-1}(\Sigma)$  for some  $q \in \mathbb{N}^*$  and  $0 < \delta < 1$ . For  $\mathfrak{p} = (\check{g}, \check{\pi})$  compactly supported we define

$$(\hat{g}, \hat{\pi})(\mathfrak{p}) := (\bar{g} + \check{g}, \bar{\pi} + \check{\pi}) + \mathfrak{p}.$$

$$(5.1)$$

Following [11], we look for solutions of the constraint equations under the form

$$(g,\pi) = \left( (1+\check{u})^4 \hat{g}(\mathfrak{p}), (1+\check{u})^2 \left( \hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})} \check{X} \right) \right).$$
(5.2)

**Lemma 5.1.** If  $(g,\pi)$  is defined by (5.2) then the constraint equations  $\Phi(g,\pi) = 0$  rewrite as

$$P\left(\check{\mathbb{U}}\right) = D\Phi[\bar{g},\bar{\pi}]((\check{g},\check{\pi}) + \mathfrak{p}) + \mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right)$$
(5.3)

where the operator P is defined in (4.1),  $\check{\mathbb{U}} = (\check{u}, \check{X})$  and the remainder is defined by

$$\begin{aligned} \mathcal{R}\left(\mathfrak{p},\breve{\mathbb{U}}\right) &\coloneqq \left(-Q_{\mathcal{H}}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi}) \\ &-\check{u}\left(D\mathcal{H}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi})+Q_{\mathcal{H}}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi})\right) \\ &-\left(1+\check{u}\right)Q_{\mathcal{H}}[\hat{g}(\mathfrak{p}),\hat{\pi}(\mathfrak{p})]\left(0,L_{\hat{g}(\mathfrak{p})}\check{X}\right)-\check{u}\left(-2\hat{\pi}(\mathfrak{p})^{ij}\mathcal{L}_{\widetilde{X}}\hat{g}(\mathfrak{p})_{ij}+\mathrm{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\mathrm{div}_{\hat{g}(\mathfrak{p})}\check{X}\right) \\ &+8\left(\Delta_{\hat{g}(\mathfrak{p})}-\Delta_{\bar{g}}\right)\check{u}+2\left(\hat{\pi}(\mathfrak{p})^{ij}\mathcal{L}_{\widetilde{X}}\hat{g}(\mathfrak{p})_{ij}-\bar{\pi}^{ij}\mathcal{L}_{\widetilde{X}}\hat{g}_{ij}\right)-\left(\mathrm{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\mathrm{div}_{\hat{g}(\mathfrak{p})}-\mathrm{tr}_{\bar{g}}\bar{\pi}\mathrm{div}_{\bar{g}}\right)\check{X}, \\ &2Q_{\mathcal{M}}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi})_{i} \\ &+4(1+\check{u})^{-1}\hat{g}(\mathfrak{p})^{j\ell}\left(2L_{\hat{g}(\mathfrak{p})}\check{X}_{i\ell}\partial_{j}\check{u}-L_{\hat{g}(\mathfrak{p})}\check{X}_{j\ell}\partial_{i}\check{u}\right) \\ &+4((1+\check{u})^{-1}-1)\hat{g}(\mathfrak{p})^{j\ell}\left(2\hat{\pi}(\mathfrak{p})_{i\ell}\partial_{j}\check{u}-\hat{\pi}(\mathfrak{p})_{j\ell}\partial_{i}\check{u}\right) \\ &+2\left(\mathrm{div}_{\hat{g}(\mathfrak{p})}L_{\hat{g}(\mathfrak{p})}-\mathrm{div}_{\bar{g}}L_{\bar{g}}\right)\check{X}_{i}+8\left(\hat{g}(\mathfrak{p})^{j\ell}\hat{\pi}(\mathfrak{p})_{i\ell}-\bar{g}^{j\ell}\bar{\pi}_{i\ell}\right)\partial_{j}\check{u}-4\left(\mathrm{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})-\mathrm{tr}_{\bar{g}}\bar{\pi}\right)\partial_{i}\check{u}\right). \end{aligned}$$

*Proof.* In the proof of Theorem 1 in [11] we find the following formulas:

$$u^{5}\mathcal{H}\left(u^{4}\hat{g}(\mathfrak{p}), u^{2}\left(\hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)\right) = -8\Delta_{\hat{g}(\mathfrak{p})}u + u\mathcal{H}\left(\hat{g}(\mathfrak{p}), \hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)$$

and

$$\begin{split} u^{2}\mathcal{M}\left(u^{4}\hat{g}(\mathfrak{p}), u^{2}\left(\hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)\right)_{i} \\ &= \operatorname{div}_{\hat{g}(\mathfrak{p})}\left(\hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)_{i} + 2u^{-1}\hat{g}(\mathfrak{p})^{j\ell}\left(2\left(\hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)_{i\ell}\partial_{j}u - \left(\hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)_{j\ell}\partial_{i}u\right). \end{split}$$

On the one hand, thanks to (2.2) and (2.4), we can compute  $D\mathcal{H}[\hat{g}(\mathfrak{p}), \hat{\pi}(\mathfrak{p})](0, L_{\hat{g}}X)$  and rewrite the equation

$$\mathcal{H}\left(u^{4}\hat{g}(\mathfrak{p}), u^{2}\left(\hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)\right) = 0$$

 $\mathbf{as}$ 

$$-8\Delta_{\hat{g}(\mathfrak{p})}u - 2\hat{\pi}(\mathfrak{p})^{ij}\mathcal{L}_{X}\hat{g}(\mathfrak{p})_{ij} + \operatorname{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\operatorname{div}_{\hat{g}(\mathfrak{p})}X$$
  
$$= -\mathcal{H}(\hat{g}(\mathfrak{p}), \hat{\pi}(\mathfrak{p}))$$
  
$$- (u - 1)\mathcal{H}(\hat{g}(\mathfrak{p}), \hat{\pi}(\mathfrak{p})) - uQ_{\mathcal{H}}[\hat{g}(\mathfrak{p}), \hat{\pi}(\mathfrak{p})](0, L_{\hat{g}(\mathfrak{p})}X)$$
  
$$- (u - 1)\left(-2\hat{\pi}(\mathfrak{p})^{ij}\mathcal{L}_{X}\hat{g}(\mathfrak{p})_{ij} + \operatorname{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\operatorname{div}_{\hat{g}(\mathfrak{p})}X\right).$$

Using now the definition (5.1), (2.2) and  $\mathcal{H}(\bar{g}, \bar{\pi}) = 0$ , we rewrite this equation as

$$-8\Delta_{\bar{g}}u - 2\bar{\pi}^{ij}\mathcal{L}_X\bar{g}_{ij} + \operatorname{tr}_{\bar{g}}\bar{\pi}\operatorname{div}_{\bar{g}}X = -D\mathcal{H}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi}) - Q_{\mathcal{H}}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi}) - (u-1)\left(D\mathcal{H}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi}) + Q_{\mathcal{H}}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi})\right) - uQ_{\mathcal{H}}[\hat{g}(\mathfrak{p}),\hat{\pi}(\mathfrak{p})](0,L_{\hat{g}(\mathfrak{p})}X) - (u-1)\left(-2\hat{\pi}(\mathfrak{p})^{ij}\mathcal{L}_X\hat{g}(\mathfrak{p})_{ij} + \operatorname{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\operatorname{div}_{\hat{g}(\mathfrak{p})}X\right) + 8\Delta_{\hat{g}(\mathfrak{p})}u + 2\hat{\pi}(\mathfrak{p})^{ij}\mathcal{L}_X\hat{g}(\mathfrak{p})_{ij} - \operatorname{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\operatorname{div}_{\hat{g}(\mathfrak{p})}X - 8\Delta_{\bar{g}}u - 2\bar{\pi}^{ij}\mathcal{L}_X\bar{g}_{ij} + \operatorname{tr}_{\bar{g}}\bar{\pi}\operatorname{div}_{\bar{g}}X.$$

On the other hand, thanks to (2.3) and (2.5), as well as (5.1) and  $\operatorname{div}_{\bar{g}}\bar{\pi} = 0$ , we rewrite the equation

$$\mathcal{M}\left(u^4\hat{g}(\mathfrak{p}), u^2\left(\hat{\pi}(\mathfrak{p}) + L_{\hat{g}(\mathfrak{p})}X\right)\right)_i = 0$$

as

$$\begin{split} 2\mathrm{div}_{\bar{g}}L_{\bar{g}}X_{i} - 8\bar{g}^{j\ell}\bar{\pi}_{i\ell}\partial_{j}u + 4\mathrm{tr}_{\bar{g}}\bar{\pi}\partial_{i}u &= 2D\mathcal{M}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi})_{i} + 2Q_{\mathcal{M}}[\bar{g},\bar{\pi}](\check{g}+\check{g},\check{\pi}+\check{\pi})_{i} \\ &+ 8u^{-1}\hat{g}(\mathfrak{p})^{j\ell}L_{\hat{g}(\mathfrak{p})}X_{i\ell}\partial_{j}u - 4u^{-1}\mathrm{tr}_{\hat{g}(\mathfrak{p})}L_{\hat{g}(\mathfrak{p})}X_{\partial i}u \\ &+ 8(u^{-1}-1)\hat{g}(\mathfrak{p})^{j\ell}\hat{\pi}(\mathfrak{p})_{i\ell}\partial_{j}u - 4(u^{-1}-1)\mathrm{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\partial_{i}u \\ &+ 2\mathrm{div}_{\hat{g}(\mathfrak{p})}L_{\hat{g}(\mathfrak{p})}X_{i} + 8\hat{g}(\mathfrak{p})^{j\ell}\hat{\pi}(\mathfrak{p})_{i\ell}\partial_{j}u - 4\mathrm{tr}_{\hat{g}(\mathfrak{p})}\hat{\pi}(\mathfrak{p})\partial_{i}u \\ &- 2\mathrm{div}_{\bar{g}}L_{\bar{g}}X_{i} - 8\bar{g}^{j\ell}\bar{\pi}_{i\ell}\partial_{j}u + 4\mathrm{tr}_{\bar{g}}\bar{\pi}\partial_{i}u. \end{split}$$

Applying the above formula to  $u = 1 + \check{u}$  and  $X = \check{X}$  concludes the proof of the lemma.

The proof of Theorem 3.1 reduces to whether we can construct  $\mathfrak{p}$  such that (5.3) admits a solution  $\check{\mathbb{U}} \in H^2_{-q-\delta}(\Sigma)$ . Thanks to Proposition 4.12, we need  $\mathfrak{p}$  to ensure orthogonality conditions with the  $\mathbb{W}_{j,\ell,\alpha}$  for  $1 \leq j \leq q$ . Because of the remainder term in (5.3), the definition of  $\mathfrak{p}$  is non-linear and will follow from a fixed point argument performed in Section 5.3. More precisely for each  $\check{\mathbb{U}} \in H^2_{-q-\delta}(\Sigma)$  small enough we will construct and control  $\mathfrak{p}(\check{\mathbb{U}})$  so that the right-hand side of (5.3) is orthogonal to  $\mathbb{W}_{j,\ell,\alpha}$  for  $1 \leq j \leq q$ . We then solve the elliptic system in Section 5.4 and thus conclude the proof of Theorem 3.1.

5.2. Estimating the remainder. We start by estimating the remainder term in (5.3), i.e.  $\mathcal{R}(\mathfrak{p}, \check{\mathbb{U}})$ . For the sake of clarity, we introduce the schematic notations

 $\bar{\Gamma} = \{\nabla \bar{g}, \bar{\pi}\}, \quad \check{\Gamma} = \{\nabla \check{g}, \check{\pi}\}, \quad \check{\Gamma} = \{\nabla \check{g}, \check{\pi}\}, \quad \hat{\Gamma}(\mathfrak{p}) = \{\nabla \hat{g}(\mathfrak{p}), \hat{\pi}(\mathfrak{p})\}.$ 

Since  $(\check{g}, \check{\pi})$ ,  $\mathfrak{p}$  and  $\check{\mathbb{U}}$  will be small in appropriate spaces, we only write down quadratic terms when manipulating  $\mathcal{R}(\mathfrak{p}, \check{\mathbb{U}})$  since higher order terms are smaller and enjoy more decay and thus are easier to treat. Thanks to its expression given in Lemma 5.1 and the various schematic expressions of Lemma 2.4 we find

$$\mathcal{R}\left(\mathfrak{p},\breve{\mathbb{U}}\right) = \mathcal{R}_{1}\left(\mathfrak{p}\right) + \mathcal{R}_{2}\left(\mathfrak{p},\breve{\mathbb{U}}\right) + \mathcal{R}_{3}\left(\mathfrak{p},\breve{\mathbb{U}}\right) + \text{higher-order terms}$$

where we have schematically

$$\begin{split} \mathcal{R}_{1}\left(\mathfrak{p}\right) &= (\check{g}+\check{g})\nabla\left(\check{\Gamma}+\check{\Gamma}\right) + \bar{\Gamma}(\check{g}+\check{g})\left(\check{\Gamma}+\check{\Gamma}\right) + \left(\check{\Gamma}+\check{\Gamma}\right)^{2} + \bar{\Gamma}^{2}(\check{g}+\check{g})^{2},\\ \mathcal{R}_{2}\left(\mathfrak{p},\check{\mathbb{U}}\right) &= \hat{g}(\mathfrak{p})\left(\nabla\check{\mathbb{U}}\right)^{2} + \hat{\Gamma}(\mathfrak{p})\check{\mathbb{U}}\left(\nabla\check{\mathbb{U}}+\hat{\Gamma}(\mathfrak{p})\check{\mathbb{U}}\right),\\ \mathcal{R}_{3}\left(\mathfrak{p},\check{\mathbb{U}}\right) &= (\check{g}+\check{g})\nabla^{2}\check{\mathbb{U}} + \left((\check{g}+\check{g})\bar{\Gamma}+\check{\Gamma}+\check{\Gamma}\right)\nabla\check{\mathbb{U}}\\ &+ \left(\nabla\left(\check{\Gamma}+\check{\Gamma}\right) + (\check{g}+\check{g})\nabla\bar{\Gamma}+\bar{\Gamma}\left(\check{\Gamma}+\check{\Gamma}\right) + \bar{\Gamma}^{2}(\check{g}+\check{g})\right)\check{\mathbb{U}}. \end{split}$$

Note that we have omitted undifferentiated components of  $\bar{g}$  since they are O(1).

Lemma 5.2. Assume

$$\left\| \check{\mathbb{U}} \right\|_{H^2_{-q-\delta}(\Sigma)} \le C_0 \varepsilon, \tag{5.4}$$

$$\|\breve{g}\|_{W^{2,\infty}(\Sigma)} + \|\breve{\pi}\|_{W^{1,\infty}(\Sigma)} \le D_0\varepsilon,\tag{5.5}$$

for some  $C_0, D_0 > 0$ . If  $C_0$  and  $D_0$  are large enough and  $\varepsilon$  is small enough compared to  $C_0^{-1}$  and  $D_0^{-1}$  then

$$\left\| \mathcal{R}\left( \mathfrak{p}, \check{\mathbb{U}} \right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} \lesssim C(C_{0}, D_{0})\varepsilon^{2},$$
(5.6)

$$\left| \left\langle \mathcal{R}\left( \mathfrak{p}, \check{\mathbb{U}} \right), \mathbb{W}_{j,\ell,\alpha} \right\rangle_{L^{2}(\Sigma)} \right| \lesssim C(C_{0}, D_{0})\varepsilon^{2}.$$
(5.7)

*Proof.* Recall that  $\mathfrak{p} = (\check{g}, \check{\pi})$  is compactly supported and that its support will be fixed in the whole proof so that when estimating terms containing either  $\check{g}$  or  $\check{\pi}$  we can drop the weights at infinity. Moreover,

it suffices to estimate  $\mathcal{R}_1(\mathfrak{p})$ ,  $\mathcal{R}_2(\mathfrak{p}, \check{\mathbb{U}})$  and  $\mathcal{R}_3(\mathfrak{p}, \check{\mathbb{U}})$  since the higher order terms in  $\mathcal{R}(\mathfrak{p}, \check{\mathbb{U}})$  enjoy better decay and are smaller. We start with  $\mathcal{R}_1(\mathfrak{p})$ :

$$\begin{split} \|\mathcal{R}_{1}\left(\mathfrak{p}\right)\|_{L^{2}_{-q-\delta-2}(\Sigma)} &\lesssim \left\| (\breve{g}+\breve{g})\nabla\left(\breve{\Gamma}+\breve{\Gamma}\right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} + \left\| (\breve{g}+\breve{g})\left(\breve{\Gamma}+\breve{\Gamma}\right) \right\|_{L^{2}_{-q-\delta}(\Sigma)} \\ &+ \left\| \left(\breve{\Gamma}+\breve{\Gamma}\right)^{2} \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} + \left\| (\breve{g}+\breve{g})^{2} \right\|_{L^{2}_{-q-\delta+2}(\Sigma)}, \end{split}$$

where we used asymptotic flatness of  $(\bar{g}, \bar{\pi})$  from Definition 2.1, which implies  $|\bar{\Gamma}| \leq (1+r)^{-2}$ . Using now the continuous bilinear embeddings from Lemma 2.3 we obtain

$$\begin{aligned} \|\mathcal{R}_{1}\left(\mathfrak{p}\right)\|_{L^{2}_{-q-\delta-2}(\Sigma)} &\lesssim \|\check{g}+\check{g}\|_{H^{2}_{-q-\delta}(\Sigma)} \left(\left\|\nabla\left(\check{\Gamma}+\check{\Gamma}\right)\right\|_{L^{2}_{-q-\delta-2}(\Sigma)} + \left\|\check{\Gamma}+\check{\Gamma}\right\|_{H^{1}_{-q-\delta-1}(\Sigma)}\right) \\ &+ \left\|\check{\Gamma}+\check{\Gamma}\right\|_{H^{1}_{-q-\delta-1}(\Sigma)}^{2} + \left\|\check{g}+\check{g}\right\|_{H^{2}_{-q-\delta}(\Sigma)}^{2}. \end{aligned}$$

Using now (3.1) and (5.5) this gives

$$\left\|\mathcal{R}_{1}\left(\mathfrak{p}\right)\right\|_{L^{2}_{-q-\delta-2}(\Sigma)} \lesssim C(D_{0})\varepsilon^{2}.$$
(5.8)

We now estimate  $\mathcal{R}_2\left(\mathfrak{p}, \check{\mathbb{U}}\right)$ :

$$\left\|\mathcal{R}_{2}\left(\mathfrak{p},\check{\mathbb{U}}\right)\right\|_{L^{2}_{-q-\delta-2}(\Sigma)} \lesssim \left(1+C(D_{0})\varepsilon\right)\left(\left\|\left(\nabla\check{\mathbb{U}}\right)^{2}\right\|_{L^{2}_{-q-\delta-2}(\Sigma)}+\left\|\check{\mathbb{U}}\nabla\check{\mathbb{U}}\right\|_{L^{2}_{-q-\delta}(\Sigma)}+\left\|\check{\mathbb{U}}^{2}\right\|_{L^{2}_{-q-\delta+2}(\Sigma)}\right),$$

where we used (5.1), asymptotic flatness, (3.1) and (5.5) which imply  $\hat{g}(\mathfrak{p}) = 1 + O(r^{-1}) + O(\varepsilon C(D_0)r^{-q-\delta})$ and  $\hat{\Gamma}(\mathfrak{p}) = O(r^{-2}) + O(\varepsilon C(D_0)r^{-q-\delta-1})$ . The assumption (5.4) and the continuous bilinear embeddings from Lemma 2.3 then imply

$$\left\| \mathcal{R}_2\left( \mathfrak{p}, \check{\mathbb{U}} \right) \right\|_{L^2_{-q-\delta-2}(\Sigma)} \lesssim C(C_0, D_0) \varepsilon^2.$$
(5.9)

We finally estimate  $\mathcal{R}_3(\mathfrak{p}, \check{\mathbb{U}})$ :

$$\begin{split} \left\| \mathcal{R}_{3}\left(\mathfrak{p},\breve{\mathbb{U}}\right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} &\lesssim \left\| (\breve{g}+\breve{g})\nabla^{2}\breve{\mathbb{U}} \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} + \left\| (\breve{\Gamma}+\breve{\Gamma})\nabla\breve{\mathbb{U}} \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} + \left\| (\breve{g}+\breve{g})\nabla\breve{\mathbb{U}} \right\|_{L^{2}_{-q-\delta}(\Sigma)} \\ &+ \left\| \nabla \left(\breve{\Gamma}+\breve{\Gamma}\right)\breve{\mathbb{U}} \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} + \left\| (\breve{g}+\breve{g})\breve{\mathbb{U}} \right\|_{L^{2}_{-q-\delta+1}(\Sigma)} \\ &+ \left\| \left(\breve{\Gamma}+\breve{\Gamma}\right)\breve{\mathbb{U}} \right\|_{L^{2}_{-q-\delta}(\Sigma)} + \left\| (\breve{g}+\breve{g})\breve{\mathbb{U}} \right\|_{L^{2}_{-q-\delta+2}(\Sigma)}, \end{split}$$

where we used asymptotic flatness to write  $\overline{\Gamma} = O(r^{-2})$  and  $\nabla \overline{\Gamma} = O(r^{-3})$ . As above the continuous bilinear embeddings from Lemma 2.3 together with (3.1), (5.4) and (5.5) imply

$$\left\| \mathcal{R}_3\left( \mathfrak{p}, \check{\mathbb{U}} \right) \right\|_{L^2_{-q-\delta-2}(\Sigma)} \lesssim C(C_0, D_0) \varepsilon^2.$$
(5.10)

Putting (5.8), (5.9) and (5.10) together concludes the proof of (5.6). Finally, since  $j \leq q$  and  $\delta > 0$  imply  $\left| \langle \mathbb{V}, \mathbb{W}_{j,\ell,\alpha} \rangle_{L^2(\Sigma)} \right| \lesssim \|\mathbb{V}\|_{L^2_{-q-\delta-2}}$  (where we also used  $|\mathbb{W}_{j,\ell,\alpha}| \lesssim r^{j-1}$ ), (5.6) implies (5.7).

**Lemma 5.3.** Let  $\check{\mathbb{U}}, \check{\mathbb{U}}'$  satisfying (5.4) and  $\mathfrak{p}, \mathfrak{p}'$  satisfying (5.5). If  $C_0$  and  $D_0$  are large enough and  $\varepsilon$  is small enough compared to  $C_0^{-1}$  and  $D_0^{-1}$  then

$$\begin{aligned} \left\| \mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right) - \mathcal{R}\left(\mathfrak{p}',\check{\mathbb{U}}'\right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} & (5.11) \\ &\lesssim C(C_{0},D_{0})\varepsilon \left( \left\| \check{g} - \check{g}' \right\|_{W^{2,\infty}(\Sigma)} + \left\| \check{\pi} - \check{\pi}' \right\|_{W^{1,\infty}(\Sigma)} + \left\| \check{\mathbb{U}} - \check{\mathbb{U}}' \right\|_{H^{2}_{-q-\delta}(\Sigma)} \right), \\ & \left| \left\langle \mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right) - \mathcal{R}\left(\mathfrak{p}',\check{\mathbb{U}}'\right), \mathbb{W}_{j,\ell,\alpha} \right\rangle_{L^{2}(\Sigma)} \right| & (5.12) \\ &\lesssim C(C_{0},D_{0})\varepsilon \left( \left\| \check{g} - \check{g}' \right\|_{W^{2,\infty}(\Sigma)} + \left\| \check{\pi} - \check{\pi}' \right\|_{W^{1,\infty}(\Sigma)} + \left\| \check{\mathbb{U}} - \check{\mathbb{U}}' \right\|_{H^{2}_{-q-\delta}(\Sigma)} \right). \end{aligned}$$

*Proof.* We have

$$\begin{split} \mathcal{R}\left(\mathfrak{p},\breve{\mathbb{U}}\right) - \mathcal{R}\left(\mathfrak{p}',\breve{\mathbb{U}}'\right) &= \mathcal{R}_{1}\left(\mathfrak{p}\right) - \mathcal{R}_{1}\left(\mathfrak{p}'\right) + \mathcal{R}_{2}\left(\mathfrak{p},\breve{\mathbb{U}}\right) - \mathcal{R}_{2}\left(\mathfrak{p}',\breve{\mathbb{U}}'\right) \\ &+ \mathcal{R}_{3}\left(\mathfrak{p},\breve{\mathbb{U}}\right) - \mathcal{R}_{3}\left(\mathfrak{p}',\breve{\mathbb{U}}'\right) + \text{difference of higher order terms.} \end{split}$$

We have

$$\begin{aligned} \mathcal{R}_{1}\left(\mathfrak{p}\right) - \mathcal{R}_{1}\left(\mathfrak{p}'\right) &= (\breve{g} - \breve{g}')\nabla\left(\breve{\Gamma} + \breve{\Gamma}\right) + (\breve{g} + \breve{g}')\nabla\left(\breve{\Gamma} - \breve{\Gamma}'\right) \\ &+ \bar{\Gamma}\left((\breve{g} - \breve{g}')\left(\breve{\Gamma} + \breve{\Gamma}\right) + (\breve{g} + \breve{g}')\left(\breve{\Gamma} - \breve{\Gamma}'\right)\right) \\ &+ \left(\breve{\Gamma} - \breve{\Gamma}'\right)\left(\breve{\Gamma} + \breve{\Gamma} + \breve{\Gamma}'\right) + \bar{\Gamma}^{2}\left(\breve{g} - \breve{g}'\right)\left(\breve{g} + \breve{g} + \breve{g}'\right). \end{aligned}$$

By following the same lines as the proof of (5.8) we obtain

$$\|\mathcal{R}_{1}\left(\mathfrak{p}\right)-\mathcal{R}_{1}\left(\mathfrak{p}'\right)\|_{L^{2}_{-q-\delta-2}(\Sigma)} \lesssim C(C_{0})\varepsilon\left(\|\breve{g}-\breve{g}'\|_{W^{2,\infty}(\Sigma)}+\|\breve{\pi}-\breve{\pi}'\|_{W^{1,\infty}(\Sigma)}\right).$$
(5.13)

Next, we have

$$\begin{aligned} \mathcal{R}_{2}\left(\mathfrak{p},\breve{\mathbb{U}}\right) - \mathcal{R}_{2}\left(\mathfrak{p}',\breve{\mathbb{U}}'\right) &= \left(\breve{g} - \breve{g}'\right)\left(\nabla\breve{\mathbb{U}}\right)^{2} + \hat{g}(\mathfrak{p}')\nabla\left(\breve{\mathbb{U}} - \breve{\mathbb{U}}'\right)\nabla\left(\breve{\mathbb{U}} + \breve{\mathbb{U}}'\right) \\ &+ \left(\breve{\Gamma} - \breve{\Gamma}'\right)\breve{\mathbb{U}}\nabla\breve{\mathbb{U}} + \hat{\Gamma}(\mathfrak{p}')\left(\left(\breve{\mathbb{U}} - \breve{\mathbb{U}}'\right)\nabla\breve{\mathbb{U}} + \breve{\mathbb{U}}'\nabla\left(\breve{\mathbb{U}} - \breve{\mathbb{U}}'\right)\right) \\ &+ \left(\left(\breve{\Gamma} - \breve{\Gamma}'\right)\breve{\mathbb{U}} + \hat{\Gamma}(\mathfrak{p}')\left(\breve{\mathbb{U}} - \breve{\mathbb{U}}'\right)\right)\left(\hat{\Gamma}(\mathfrak{p})\breve{\mathbb{U}} + \hat{\Gamma}(\mathfrak{p}')\breve{\mathbb{U}}'\right). \end{aligned}$$

By following the same lines as the proof of (5.9) we obtain

$$\begin{aligned} \left\| \mathcal{R}_{2}\left(\mathfrak{p},\check{\mathbb{U}}\right) - \mathcal{R}_{2}\left(\mathfrak{p}',\check{\mathbb{U}}'\right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} &\lesssim C(C_{0},D_{0})\varepsilon \left\| \check{\mathbb{U}} - \check{\mathbb{U}}' \right\|_{H^{2}_{-q-\delta}(\Sigma)} \\ &+ C(C_{0},D_{0})\varepsilon^{2} \left( \left\| \check{g} - \check{g}' \right\|_{W^{2,\infty}(\Sigma)} + \left\| \check{\pi} - \check{\pi}' \right\|_{W^{1,\infty}(\Sigma)} \right). \end{aligned}$$

$$(5.14)$$

Finally, we have

$$\begin{split} \mathcal{R}_{3}\left(\mathfrak{p},\breve{\mathbb{U}}\right) - \mathcal{R}_{3}\left(\mathfrak{p}',\breve{\mathbb{U}}'\right) &= (\breve{g} - \breve{g}')\nabla^{2}\breve{\mathbb{U}} + (\breve{g} + \breve{g}')\nabla^{2}\left(\breve{\mathbb{U}} - \breve{\mathbb{U}}'\right) \\ &+ \left(\breve{\Gamma} - \breve{\Gamma} + (\breve{g} - \breve{g}')\bar{\Gamma}\right)\nabla\breve{\mathbb{U}} + \left(\breve{\Gamma} + \breve{\Gamma}' + (\breve{g} + \breve{g}')\bar{\Gamma}\right)\nabla\left(\breve{\mathbb{U}} - \breve{\mathbb{U}}'\right) \\ &+ \left(\nabla\left(\breve{\Gamma} - \breve{\Gamma}'\right) + (\breve{g} - \breve{g}')\nabla\bar{\Gamma} + \bar{\Gamma}\left(\breve{\Gamma} - \breve{\Gamma}'\right) + \bar{\Gamma}^{2}(\breve{g} - \breve{g}')\right)\breve{\mathbb{U}} \\ &+ \left(\nabla\left(\breve{\Gamma} + \breve{\Gamma}'\right) + (\breve{g} + \breve{g}')\nabla\bar{\Gamma} + \bar{\Gamma}\left(\breve{\Gamma} + \breve{\Gamma}'\right) + \bar{\Gamma}^{2}(\breve{g} + \breve{g}')\right)\left(\breve{\mathbb{U}} - \breve{\mathbb{U}}'\right). \end{split}$$

By following the same lines as the proof of (5.10) we obtain

$$\left\| \mathcal{R}_{3}\left(\mathfrak{p},\breve{\mathbb{U}}\right) - \mathcal{R}_{3}\left(\mathfrak{p}',\breve{\mathbb{U}}'\right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} \lesssim C(C_{0},D_{0})\varepsilon \left( \left\| \breve{g} - \breve{g}' \right\|_{W^{2,\infty}(\Sigma)} + \left\| \breve{\pi} - \breve{\pi}' \right\|_{W^{1,\infty}(\Sigma)} + \left\| \breve{\mathbb{U}} - \breve{\mathbb{U}}' \right\|_{H^{2}_{-q-\delta}(\Sigma)} \right).$$

$$(5.15)$$

By putting (5.13), (5.14) and (5.15) together we obtain (5.11). We conclude the proof of the lemma by noting that (5.11) implies (5.12), as (5.6) implies (5.7).

5.3. Solving for the corrector. In this section, we construct the corrector  $\mathfrak{p}$ , starting by deducing from the analyticity assumptions on  $(\Omega, \bar{g}_{|\Omega}, \bar{\pi}_{|\Omega})$  (see Theorem 3.1) that KIDS, i.e. elements of ker $(D\Phi[\bar{g}, \bar{\pi}]^*)$ , enjoy similar regularity.

**Lemma 5.4.** Any element of ker $(D\Phi[\bar{g}, \bar{\pi}]^*)$  is analytic on  $\{r_0 \leq r < r_1\}$ .

*Proof.* If  $(f, X) \in \ker(D\Phi[\bar{g}, \bar{\pi}]^*)$ , then (recall (2.10) and (2.11)) it satisfies a system of the form

$$\operatorname{Hess}_{\bar{g}}(f)^{ij} = \bar{A}^{ij}_{k\ell} D^k X^\ell + \bar{A}^{ij}_0 f + \bar{A}^{ij}_k X^k,$$
$$\mathcal{L}_X \bar{g}^{ij} = \bar{B}^{ij} f,$$

where the coefficients  $\bar{A}_{k\ell}^{ij}$ ,  $\bar{A}_0^{ij}$ ,  $\bar{A}_k^{ij}$  and  $\bar{B}^{ij}$  depends only on  $\bar{g}$  and  $\bar{\pi}$  and are analytic on  $\{r_0 \leq r < r_1\}$ . To simplify the exposition, in what follows we don't write the analytic coefficients of the equations satisfied by f and X. In particular the above equations rewrite schematically as

$$\operatorname{Hess}_{\bar{g}}(f) = DX + f + X, \tag{5.16}$$

$$\mathcal{L}_X \bar{g} = f. \tag{5.17}$$

Taking the trace of (5.17) implies  $\operatorname{div}_{\bar{g}}X = f$ . Taking also the divergence of (5.17) together with the identity

$$\operatorname{div}_{\bar{g}}\mathcal{L}_X \bar{g}_i = \Delta_{\bar{g}}^{(1)} X_i + \partial_i \operatorname{div}_{\bar{g}} X + Ric(\bar{g})(\partial_j, X)$$

implies that f and X solve a system of the form

$$\Delta_{\bar{g}}f = DX + f + X, \tag{5.18}$$

$$\Delta_{\bar{g}}^{(1)}X = Df + f + X, \tag{5.19}$$

where we also took the trace of (5.16) to get the first equation. Therefore, f and X solve an elliptic system with analytic coefficients and thus are analytic on  $\{r_0 < r < r_1\}$  thanks to standard elliptic regularity. It remains to prove that there are analytic on the boundary  $\partial \Sigma$ , and for that we will derive from (5.16)-(5.17) an elliptic system on  $\partial \Sigma$  with analytic coefficients.

For this, we introduce the standard notations from [8], i.e.  $\mathcal{D}$ , Hess and  $\mathcal{A}$  denote covariant derivative, Hessian and Laplace-Beltrami operator associated to the induced metric  $\mathbf{\vec{g}}$  on  $\partial \Sigma$  by  $\mathbf{\bar{g}}$ . We also decompose the vector field  $X = N\nu + \mathbf{\vec{X}}$  where  $\mathbf{\vec{X}} \in T\partial\Sigma$  and N is a scalar function. By assumption, every quantity related to  $\mathbf{\bar{g}}$  is analytic and for simplicity we don't write the analytic coefficients appearing in the following formulas such as the second fundamental form of  $\partial\Sigma$  as a submanifold of  $\Sigma$ . By plugging (5.18) into (5.16) we obtain a simpler form of the system for (f, X):

$$\operatorname{Hess}_{\bar{g}}(f) = DX + f + X,\tag{5.20}$$

$$\mathcal{L}_X \bar{g} = f. \tag{5.21}$$

We first contract (5.21) twice with  $\nu$  to obtain

$$\partial_{\nu}N = f + X. \tag{5.22}$$

Then, we contract (5.21) with one tangential vector field and  $\nu$  to get

$$\mathcal{D}_{\nu}X + \mathcal{D}N = f + N + X, \qquad (5.23)$$

which after taking the tangential divergence implies

Finally, we contract (5.21) with tangential vector fields, take the trace to obtain di/X = f and then take the tangential divergence of (5.21) to obtain the following elliptic equation on X

$$\mathbf{A}\mathbf{X} = \mathbf{P}^{\leq 1}(f + N + \mathbf{X}). \tag{5.25}$$

Using that  $[\Delta, \partial_{\nu}]$  involves  $\mathcal{D}\partial_{\nu}$  and  $\mathcal{D}^2$  we obtain from (5.25) the following

$$\Delta \mathcal{D}_{\nu} \mathcal{X} = \mathcal{D}^{\leq 1} (f + N + \mathcal{X} + \mathcal{D} \mathcal{X} + \partial_{\nu} f + \mathcal{D}_{\nu} \mathcal{X}), \qquad (5.26)$$

where we have used (5.22) to replace  $\partial_{\nu} N$ . We also commute (5.25) with D to obtain

$$\Delta \mathcal{D} X = \mathcal{D}^{\leq 1}(f + N + X) + \mathcal{D}^2(f + N + X).$$

Using (5.23) we replace  $\mathbb{D}^2 N$  by  $\mathbb{D}\mathbb{D}_{\nu} X$  and  $\mathbb{D} N$  to obtain

$$\Delta \mathcal{D} X = \mathcal{D}^{\leq 1}(f + N + X + \mathcal{D}_{\nu} X + \mathcal{D} X) + \mathcal{D}^{2} f.$$
(5.27)

We now look at the equation for f, i.e. (5.20). By contracting it in tangential directions we obtain

$$\operatorname{Hess} f = \partial_{\nu} f + D^{\leq 1} (f + N + X), \qquad (5.28)$$

where we have used (5.22) and (5.23) to replace  $\partial_{\nu}N$  and  $\not D_{\nu}X$ . Taking the tangential trace of (5.28) we obtain the equation for f:

$$\Delta f = \partial_{\nu} f + D^{\leq 1} (f + N + X).$$
(5.29)

Moreover, by using (5.28) we can also control  $D^2 f$  in (5.27) which becomes

$$\mathcal{A}\mathcal{D}\mathcal{X} = \partial_{\nu}f + \mathcal{D}^{\leq 1}(f + N + \mathcal{X} + \mathcal{D}_{\nu}\mathcal{X} + \mathcal{D}\mathcal{X}).$$
(5.30)

Also, we commute (5.29) with  $\partial_{\nu}$  to obtain an equation on  $\partial_{\nu} f$ , using again the fact that  $[\underline{A}, \partial_{\nu}]$  involves  $\underline{\mathcal{D}}\partial_{\nu}$  and  $\underline{\mathcal{D}}^2$ . We get:

$$\mathcal{A}\partial_{\nu}f = \partial_{\nu}^{2}f + \mathcal{D}^{\leq 1}(f + N + \mathcal{X} + \partial_{\nu}f + \mathcal{D}_{\nu}\mathcal{X})$$

$$(5.31)$$

where we have used (5.22) and (5.28) to replace  $\partial_{\nu} N$  and  $D^2 f$ . Finally, using the double contraction by  $\nu$  of (5.20) we can replace  $\partial_{\nu}^2$  in (5.31) to finally obtain

$$\Delta \partial_{\nu} f = D^{\leq 1} (f + N + X + \partial_{\nu} f + D_{\nu} X).$$
(5.32)

By using the schematic notation

 $V = \{f, N, \mathcal{X}, \partial_{\nu} f, \mathcal{D}_{\nu} \mathcal{X}, \mathcal{D} \mathcal{X}\},\$ 

we remark that combining equations (5.24), (5.25), (5.26), (5.29), (5.30) and (5.32) gives an elliptic system on  $\partial \Sigma$  of the form

 $\Delta V = \mathbb{D}^{\leq 1} V.$ 

In particular, the coefficients of this system are analytic by assumption and thus standard elliptic regularity implies that V is analytic on  $\partial \Sigma$ . This concludes the proof of Lemma 5.4.

We now define the space  $\mathcal{Z}_q$  of correctors.

**Definition 5.5.** We fix  $I \subset (r_0, r_1)$  an open interval in  $\mathbb{R}$ .

- (i) We define  $\chi_{corr} : \mathbb{R} \longrightarrow [0,1]$  to be a smooth function such that  $\chi_{corr}|_{\mathbb{R}\setminus I} = 0$  and  $\chi_{corr}|_{I} > 0$ .
- (ii) For  $j \ge 1$ ,  $-(j-1) \le \ell \le j-1$  and  $\alpha = 0, 1, 2, 3$  we define

$$Z_{j,\ell,\alpha} := \chi_{corr}(r) D\Phi[\bar{g},\bar{\pi}]^* (\mathbb{W}_{j,\ell,\alpha}).$$

(iii) For  $q \in \mathbb{N}^*$  we define the  $4q^2$  dimensional space of correctors

$$\mathcal{Z}_q := \operatorname{span}(Z_{j,\ell,\alpha}, 1 \le j \le q, \ -(j-1) \le \ell \le j-1 \ and \ \alpha = 0, 1, 2, 3),$$

endowed with the norm  $\|\cdot\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)}$ .

The following lemma will allow us to solve the equation for the corrector.

Lemma 5.6. Let  $q \in \mathbb{N}^*$ .

(i) The matrix

$$\left(\left\langle Z_{j,\ell,\alpha}, D\Phi[\bar{g},\bar{\pi}]^*\left(\mathbb{W}_{j',\ell',\alpha'}\right)\right\rangle_{L^2(\Sigma)}\right)_{1\leq j,j'\leq q,-(j-1)\leq \ell\leq j-1,-(j'-1)\leq \ell'\leq j'-1,0\leq \alpha,\alpha'\leq 3j'-1,0\leq \alpha,\alpha'<3j'-1,0\leq \alpha,\alpha'<3j'-1,0\leq \alpha,\alpha'<3j'-1,0\leq \alpha,\alpha'<3j'-1,0\leq \alpha,\alpha'<3j'-1,0\leq \alpha,\alpha'<3j'-1,0\leq \alpha,\alpha'<3j'-1,0\leq \alpha,\alpha'<3j'-1,0<\alpha,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-1,0<\beta,\alpha'<3j'-$$

is invertible.

(ii) Let  $\mu_{j,\ell,\alpha}$  be real numbers for  $1 \leq j \leq q$ ,  $-(j-1) \leq \ell \leq j-1$  and  $\alpha = 0, 1, 2, 3$ . There exists a unique  $\mathfrak{p} \in \mathbb{Z}_q$  such that

$$\langle \mathfrak{p}, D\Phi[\bar{g}, \bar{\pi}]^* (\mathbb{W}_{j,\ell,\alpha}) \rangle_{L^2(\Sigma)} = \mu_{j,\ell,\alpha},$$

and moreover p satisfies

$$\|\mathfrak{p}\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)}\lesssim \sup_{j,\ell,\alpha}|\mu_{j,\ell,\alpha}|.$$

*Proof.* We start with the first point of the lemma. Let  $\lambda^{j,\ell,\alpha}$  be real numbers defined such that

 $\left\langle \lambda^{j,\ell,\alpha} \chi_{corr}(r) D\Phi[\bar{g},\bar{\pi}]^* \left( \mathbb{W}_{j,\ell,\alpha} \right), D\Phi[\bar{g},\bar{\pi}]^* \left( \mathbb{W}_{j',\ell',\alpha'} \right) \right\rangle_{L^2(\Sigma)} = 0,$ 

for all  $1 \leq j' \leq q$ ,  $-(j'-1) \leq \ell' \leq j'-1$  and  $\alpha' = 0, 1, 2, 3$ . We multiply each equality by  $\lambda^{j',\ell',\alpha'}$  and sum over the indices  $j', \ell'$  and  $\alpha'$  to get

$$\left\langle \chi_{corr} D\Phi[\bar{g},\bar{\pi}]^* \left( \lambda^{j,\ell,\alpha} \mathbb{W}_{j,\ell,\alpha} \right), D\Phi[\bar{g},\bar{\pi}]^* \left( \lambda^{j,\ell,\alpha} \mathbb{W}_{j,\ell,\alpha} \right) \right\rangle_{L^2(\Sigma)} = 0,$$

Since  $\chi_{corr|_{\mathbb{R}\setminus I}} = 0$  and  $\chi_{corr|_{I}} > 0$  this implies

$$D\Phi[\bar{g},\bar{\pi}]^*\left(\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha}\right) = 0, \quad \text{on } \{r \in I\}.$$
(5.33)

Recall from Lemma 4.10 that  $P^*(\mathbb{W}_{j,\ell,\alpha}) = 0$  on  $\Sigma$  and by assumption (see Theorem 3.1) the coefficients of  $P^*$  are analytic in the region  $\{r_0 < r < r_1\}$ . Standard elliptic regularity thus implies that  $\mathbb{W}_{j,\ell,\alpha}$  is analytic in the region  $\{r_0 < r < r_1\}$ . Since the assumptions also imply that the coefficients of  $D\Phi[\bar{g},\bar{\pi}]^*$ are analytic in the region  $\{r_0 < r < r_1\}$ , we can extend (5.33) to this region, and by continuity to  $\{r_0 \leq r < r_1\}$ . We have thus proved

$$D\Phi[\bar{g},\bar{\pi}]^* \left(\lambda^{j,\ell,\alpha} \mathbb{W}_{j,\ell,\alpha}\right) = 0, \quad \text{on } \{r_0 \le r < r_1\}.$$
(5.34)

Now, Lemma 5.4 implies that  $\lambda^{j,\ell,\alpha} \mathbb{W}_{j,\ell,\alpha}$  is analytic on the boundary  $\partial \Sigma$ . Recall from Proposition 4.12 that  $(B_{\nu}^* + F_{\delta}^*)(\mathbb{W}_{j,\ell,\alpha}) = 0$  on  $\partial \Sigma$ . Restricting this to the open subset  $\mathcal{U}$  of  $\partial \Sigma$  where  $F_{\delta} = 0$  (which implies that  $F_{\delta}^* = 0$  on the same open subset) we obtain  $B_{\nu}^*(\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha})|_{\mathcal{U}} = 0$ . Since  $\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha}$  is analytic on the boundary, we extend this equality to  $B_{\nu}^*(\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha}) = 0$  on  $\partial \Sigma$ . Returning to the boundary conditions, this implies that  $F_{\delta}^*(\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha}) = 0$  on  $\partial \Sigma$ . Returning to the open subset  $\mathcal{V}$  of  $\partial \Sigma$  where  $F_{\delta} = \mathrm{Id}$  (which implies that  $F_{\delta}^* = \mathrm{Id}$  on the same open subset), we obtain  $\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha|_{\mathcal{V}}} = 0$  which again extends to  $\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha} = 0$  on  $\partial \Sigma$  thanks to analyticity. Together with Lemma 4.10 this implies that  $P^*(\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha}) = 0$  on  $\Sigma$ ,  $B_{\nu}^*(\lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha}) = \lambda^{j,\ell,\alpha}\mathbb{W}_{j,\ell,\alpha} = 0$  on  $\partial \Sigma$ . Lemma 4.4 thus implies that

$$\lambda^{j,\ell,\alpha} \mathbb{W}_{j,\ell,\alpha} = 0, \quad \text{on } \Sigma.$$
(5.35)

Since the family formed by the  $\mathbb{W}_{j,\ell,\alpha}$  is linearly independent on  $\Sigma$ , this implies that  $\lambda^{j,\ell,\alpha} = 0$  for all indices and concludes the proof of Lemma 5.6, since the second point follows directly from the first one.

**Remark 5.7.** The proof of Lemma 5.6 shows how the KIDS are obstructions to the construction of the corrector. In order to go from (5.34) to (5.35), we need to use a property distinguishing elements of  $\ker(D\Phi[\bar{g},\bar{\pi}]^*)$  and elements of  $\ker(P^*_{F_{\delta},q-1+\delta})$ . The key is to note that  $P^*(\mathbb{U}) = 0$  for all  $\mathbb{U} \in \ker(D\Phi[\bar{g},\bar{\pi}]^*)$ , which follows from the following alternative definition of P

$$P = D\Phi[\bar{g}, \bar{\pi}] \circ \Pi,$$

with  $\Pi(u, X) = (4u\bar{g}, 2u\bar{\pi} + L_{\bar{g}}X)$ . This implies that the only way to distinguish  $\ker(D\Phi[\bar{g}, \bar{\pi}]^*)$  from  $\ker(P^*_{F_{\delta}, g-1+\delta})$  is the boundary condition, and in particular the regularity at the boundary.

We now construct the corrector map. If  $\check{\mathbb{U}}$  is such that

$$\left\| \check{\mathbb{U}} \right\|_{H^2_{-q-\delta}(\Sigma)} \le C_0 \varepsilon, \tag{5.36}$$

we want the corrector  $\mathfrak{p} \in \mathcal{Z}_q$  to be such that the following orthogonality conditions hold

$$\left\langle D\Phi[\bar{g},\bar{\pi}]((\check{g},\check{\pi})+\mathfrak{p})+\mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right),\mathbb{W}_{j,\ell,\alpha}\right\rangle_{L^{2}(\Sigma)}=0,$$
(5.37)

for all  $1 \leq j \leq q$ ,  $-(j-1) \leq \ell \leq j-1$  and  $\alpha = 0, 1, 2, 3$ . According to Proposition 4.12 this will ensure the solvability of (5.3). Since each  $\mathfrak{p} \in \mathbb{Z}_q$  is compactly supported in  $\Sigma$  we can perform the following integration by part without picking up boundary terms

$$\langle D\Phi[\bar{g},\bar{\pi}](\mathfrak{p}),\mathbb{W}_{j,\ell,\alpha}\rangle_{L^{2}(\Sigma)} = \langle \mathfrak{p}, D\Phi[\bar{g},\bar{\pi}]^{*}(\mathbb{W}_{j,\ell,\alpha})\rangle_{L^{2}(\Sigma)},$$

so that (5.37) rewrites

$$\langle \mathfrak{p}, D\Phi[\bar{g}, \bar{\pi}]^*(\mathbb{W}_{j,\ell,\alpha}) \rangle_{L^2(\Sigma)} = - \langle D\Phi[\bar{g}, \bar{\pi}](\check{g}, \check{\pi}), \mathbb{W}_{j,\ell,\alpha} \rangle_{L^2(\Sigma)} - \left\langle \mathcal{R}\left(\mathfrak{p}, \check{\mathbb{U}}\right), \mathbb{W}_{j,\ell,\alpha} \right\rangle_{L^2(\Sigma)}.$$
(5.38)

**Definition 5.8.** Let  $D_0 > 0$ . The corrector map  $\Psi^{(c)}$  is defined by

$$\Psi^{(c)}: B_{\mathcal{Z}_q}(0, D_0 \varepsilon) \longrightarrow B_{\mathcal{Z}_q}(0, D_0 \varepsilon),$$
$$\mathfrak{p} \longmapsto \Psi^{(c)}(\mathfrak{p}),$$

such that  $\Psi^{(c)}(\mathfrak{p})$  solves

$$\left\langle \Psi^{(c)}(\mathfrak{p}), D\Phi[\bar{g},\bar{\pi}]^*(\mathbb{W}_{j,\ell,\alpha}) \right\rangle_{L^2(\Sigma)} = -\left\langle D\Phi[\bar{g},\bar{\pi}](\check{g},\check{\pi}), \mathbb{W}_{j,\ell,\alpha} \right\rangle_{L^2(\Sigma)} - \left\langle \mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right), \mathbb{W}_{j,\ell,\alpha} \right\rangle_{L^2(\Sigma)}$$
(5.39)

for all  $1 \le j \le q$ ,  $-(j-1) \le \ell \le j-1$  and  $\alpha = 0, 1, 2, 3$ .

**Lemma 5.9.** Let  $\check{\mathbb{U}}$  satisfying (5.36). If  $D_0$  is large enough and if  $\varepsilon$  is small enough compared to  $D_0^{-1}$  and  $C_0^{-1}$ , then the corrector map  $\Psi^{(c)}$  is a well-defined contraction.

*Proof.* Let  $\mathfrak{p} \in B_{\mathcal{Z}_q}(0, D_0 \varepsilon)$ . Thanks to Lemma 5.6 there exists a unique  $\Psi^{(c)}(\mathfrak{p}) \in \mathcal{Z}_q$  satisfying (5.39) for all  $1 \leq j \leq q, -(j-1) \leq \ell \leq j-1$  and  $\alpha = 0, 1, 2, 3$  satisfying moreover

$$\left\|\Psi^{(c)}(\mathfrak{p})\right\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)} \lesssim \sup_{j,\ell,\alpha} \left| \langle D\Phi[\bar{g},\bar{\pi}](\check{g},\check{\pi}), \mathbb{W}_{j,\ell,\alpha} \rangle_{L^{2}(\Sigma)} + \left\langle \mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right), \mathbb{W}_{j,\ell,\alpha} \right\rangle_{L^{2}(\Sigma)} \right|$$

We estimate both terms on the right-hand side. First since  $j \leq q$  and  $\delta > 0$  we have

$$\begin{aligned} |\langle D\Phi[\bar{g},\bar{\pi}](\check{g},\check{\pi}),\mathbb{W}_{j,\ell,\alpha}\rangle| &\lesssim \|D\Phi[\bar{g},\bar{\pi}](\check{g},\check{\pi})\|_{L^{2}_{-q-\delta-2}(\Sigma)} \\ &\lesssim \|\check{g}\|_{H^{2}_{-q-\delta}(\Sigma)} + \|\check{\pi}\|_{H^{1}_{-q-\delta-1}(\Sigma)} \\ &\leq \varepsilon \end{aligned}$$

where we have used (2.4), (2.5), asymptotic flatness and (3.1). For the second term in the estimate for  $\Psi^{(c)}(\mathfrak{p})$  we use (5.7). We obtain

$$\left\|\Psi^{(c)}(\mathfrak{p})\right\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)}\lesssim \varepsilon\left(1+C(C_0,D_0)\varepsilon\right).$$

Taking  $D_0$  large enough and  $\varepsilon$  small enough compared to  $D_0^{-1}$  and  $C_0^{-1}$  shows that  $\Psi^{(c)}(\mathfrak{p}) \in B_{\mathbb{Z}_q}(0, D_0\varepsilon)$ and thus that the corrector map  $\Psi^{(c)}$  is well-defined.

Next let  $\mathfrak{p}, \mathfrak{p}' \in B_{\mathbb{Z}_q}(0, D_0 \varepsilon)$ . Taking the difference of the relations defining  $\Psi^{(c)}(\mathfrak{p})$  and  $\Psi^{(c)}(\mathfrak{p}')$  and using Lemma 5.6 we obtain

$$\begin{split} \left\| \Psi^{(c)}(\mathfrak{p}) - \Psi^{(c)}(\mathfrak{p}') \right\|_{W^{2,\infty}(\Sigma) \times W^{1,\infty}(\Sigma)} &\lesssim \sup_{j,\ell,\alpha} \left| \left\langle \mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right) - \mathcal{R}\left(\mathfrak{p}',\check{\mathbb{U}}\right), \mathbb{W}_{j,\ell,\alpha} \right\rangle_{L^{2}(\Sigma)} \right. \\ &\lesssim C(C_{0},D_{0})\varepsilon \left\| \mathfrak{p} - \mathfrak{p}' \right\|_{W^{2,\infty}(\Sigma) \times W^{1,\infty}(\Sigma)}, \end{split}$$

where we have used (5.12). Taking  $\varepsilon$  small enough compared to  $D_0^{-1}$  and  $C_0^{-1}$  shows that  $\Psi^{(c)}$  is a contraction.

**Corollary 5.10.** Let  $\check{\mathbb{U}}$  satisfying (5.36). If  $\varepsilon$  is small enough compared to  $C_0^{-1}$ , there exists a unique  $\mathfrak{p}(\check{\mathbb{U}}) \in \mathcal{Z}_q$  such that (5.37) holds. Moreover we have

$$\left\| \mathfrak{p}\left(\check{\mathbb{U}}\right) \right\|_{W^{2,\infty}(\Sigma) \times W^{1,\infty}(\Sigma)} \lesssim \varepsilon$$
(5.40)

and

$$\left\|\mathfrak{p}\left(\check{\mathbb{U}}\right)-\mathfrak{p}\left(\check{\mathbb{U}}'\right)\right\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)} \lesssim C(C_0)\varepsilon \left\|\check{\mathbb{U}}-\check{\mathbb{U}}'\right\|_{H^2_{-q-\delta}(\Sigma)},\tag{5.41}$$

if  $\check{\mathbb{U}}$  and  $\check{\mathbb{U}}'$  both satisfy (5.36).

*Proof.* The proof of the first part of the corollary follows from Lemma 5.9 and the Banach fixed point theorem. For the proof of (5.41) we take the difference of the relations defining  $\mathfrak{p} := \mathfrak{p}\left(\check{\mathbb{U}}\right)$  and  $\mathfrak{p}' := \mathfrak{p}\left(\check{\mathbb{U}}'\right)$  together with Lemma 5.6 and (5.12) to obtain

$$\begin{split} \|\mathfrak{p}-\mathfrak{p}'\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)} &\lesssim \sup_{j,\ell,\alpha} \left| \left\langle \mathcal{R}\left(\mathfrak{p},\check{\mathbb{U}}\right) - \mathcal{R}\left(\mathfrak{p}',\check{\mathbb{U}}'\right),\mathbb{W}_{j,\ell,\alpha}\right\rangle_{L^{2}(\Sigma)} \right| \\ &\lesssim C(C_{0})\varepsilon \bigg( \left\|\mathfrak{p}-\mathfrak{p}'\right\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)} + \left\|\check{\mathbb{U}}-\check{\mathbb{U}}'\right\|_{H^{2}_{-q-\delta}(\Sigma)} \bigg). \end{split}$$

By taking  $\varepsilon$  small enough compared to  $C_0^{-1}$  we can absorb  $C(C_0)\varepsilon \|\mathfrak{p} - \mathfrak{p}'\|_{W^{2,\infty}(\Sigma)\times W^{1,\infty}(\Sigma)}$  by the LHS and obtain (5.41).

5.4. Solving the elliptic system. In this section we conclude the proof of Theorem 3.1 by solving the elliptic system for  $\check{\mathbb{U}}$  with the corrector  $\mathfrak{p}(\check{\mathbb{U}})$  constructed in Section 5.3. We want to solve

$$\begin{cases} P\left(\check{\mathbb{U}}\right) = D\Phi[\bar{g},\bar{\pi}]\left((\check{g},\check{\pi}) + \mathfrak{p}\left(\check{\mathbb{U}}\right)\right) + \mathcal{R}\left(\mathfrak{p}\left(\check{\mathbb{U}}\right),\check{\mathbb{U}}\right), \quad \text{on } \Sigma, \\ (B_{\nu} + F_{\delta})\left(\check{\mathbb{U}}\right) = 0, \quad \text{on } \partial\Sigma. \end{cases}$$
(5.42)

**Definition 5.11.** Let  $C_0 > 0$ . The solution map  $\Psi^{(s)}$  is defined by

$$\begin{split} \Psi^{(s)} : B_{H^2_{-q-\delta}(\Sigma)}(0,C_0\varepsilon) &\longrightarrow B_{H^2_{-q-\delta}(\Sigma)}(0,C_0\varepsilon) \\ & \check{\mathbb{U}} \longmapsto \Psi^{(s)}\left(\check{\mathbb{U}}\right) \end{split}$$

such that  $\Psi^{(s)}\left(\check{\mathbb{U}}\right)$  solves

$$\begin{cases}
P\left(\Psi^{(s)}\left(\check{\mathbb{U}}\right)\right) = D\Phi[\bar{g},\bar{\pi}]\left((\check{g},\check{\pi}) + \mathfrak{p}\left(\check{\mathbb{U}}\right)\right) + \mathcal{R}\left(\mathfrak{p}\left(\check{\mathbb{U}}\right),\check{\mathbb{U}}\right), \quad on \ \Sigma, \\
(B_{\nu} + F_{\delta})\left(\Psi^{(s)}\left(\check{\mathbb{U}}\right)\right) = 0, \quad on \ \partial\Sigma.
\end{cases}$$
(5.43)

**Lemma 5.12.** If  $C_0$  is large enough and  $\varepsilon$  is small enough compared to  $C_0^{-1}$ , the solution map  $\Psi^{(s)}$  is a well-defined contraction.

Proof. Let  $\check{\mathbb{U}} \in B_{H^2_{-q-\delta}(\Sigma)}(0, C_0\varepsilon)$ . First note that if  $\varepsilon$  is small enough compared to  $C_0^{-1}$  then Corollary 5.10 applies and the corrector  $\mathfrak{p}(\check{\mathbb{U}})$  exists and satisfy the bounds of Corollary 5.10. By construction, this corrector is such that

$$\left\langle D\Phi[\bar{g},\bar{\pi}]\left(\left(\check{g},\check{\pi}\right)+\mathfrak{p}\left(\check{\mathbb{U}}\right)\right)+\mathcal{R}\left(\mathfrak{p}\left(\check{\mathbb{U}}\right),\check{\mathbb{U}}\right),\mathbb{W}_{j,\ell,\alpha}\right\rangle_{L^{2}(\Sigma)}=0$$

for all  $1 \le j \le q$ ,  $-(j-1) \le \ell \le j-1$  and  $\alpha = 0, 1, 2, 3$ . Moreover, using (2.4), (2.5), asymptotic flatness, (3.1), (5.6) and (5.40) we obtain

$$\left\| D\Phi[\bar{g},\bar{\pi}]\left( \left(\check{g},\check{\pi}\right) + \mathfrak{p}\left(\check{\mathbb{U}}\right) \right) + \mathcal{R}\left( \mathfrak{p}\left(\check{\mathbb{U}}\right),\check{\mathbb{U}} \right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} \lesssim \varepsilon(1 + C(C_{0})\varepsilon).$$

Therefore, the third point in Proposition 4.12 implies that there exists a unique  $\Psi^{(s)}(\check{\mathbb{U}})$  solving (5.43) with the bound

$$\left\|\Psi^{(s)}\left(\check{\mathbb{U}}\right)\right\|_{H^{2}_{-q-\delta}(\Sigma)} \lesssim \varepsilon(1+C(C_{0})\varepsilon).$$

Choosing first  $C_0$  large enough and then  $\varepsilon$  small enough compared to  $C_0^{-1}$  implies that  $\Psi^{(s)}\left(\check{\mathbb{U}}\right) \in B_{H^2_{-q-\delta}(\Sigma)}(0, C_0\varepsilon)$  and thus that the solution map  $\Psi^{(s)}$  is well-defined.

Next let  $\check{\mathbb{U}}, \check{\mathbb{U}}' \in B_{H^2_{-q-\delta}(\Sigma)}(0, C_0 \varepsilon)$ . We substract the two systems defining  $\Psi^{(s)}(\check{\mathbb{U}})$  and  $\Psi^{(s)}(\check{\mathbb{U}}')$  together with Proposition 4.12 and (5.11) to obtain

$$\begin{split} \left\| \Psi^{(s)}\left(\check{\mathbb{U}}\right) - \Psi^{(s)}\left(\check{\mathbb{U}}'\right) \right\|_{H^{2}_{-q-\delta}(\Sigma)} &\lesssim \left\| D\Phi[\bar{g},\bar{\pi}]\left(\mathfrak{p}\left(\check{\mathbb{U}}\right) - \mathfrak{p}\left(\check{\mathbb{U}}'\right)\right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} \\ &+ \left\| \mathcal{R}\left(\mathfrak{p}\left(\check{\mathbb{U}}\right),\check{\mathbb{U}}\right) - \mathcal{R}\left(\mathfrak{p}\left(\check{\mathbb{U}}'\right),\check{\mathbb{U}}'\right) \right\|_{L^{2}_{-q-\delta-2}(\Sigma)} \\ &\lesssim \left(1 + C(C_{0})\varepsilon\right) \left\| \mathfrak{p}\left(\check{\mathbb{U}}\right) - \mathfrak{p}\left(\check{\mathbb{U}}'\right) \right\|_{W^{2,\infty}(\Sigma) \times W^{1,\infty}(\Sigma)} \\ &+ C(C_{0})\varepsilon \left\| \check{\mathbb{U}} - \check{\mathbb{U}}' \right\|_{H^{2}_{-q-\delta}(\Sigma)}. \end{split}$$

We now use (5.41) to obtain

$$\left|\Psi^{(s)}\left(\check{\mathbb{U}}\right)-\Psi^{(s)}\left(\check{\mathbb{U}}'\right)\right\|_{H^{2}_{-q-\delta}(\Sigma)} \lesssim C(C_{0})\varepsilon \left\|\check{\mathbb{U}}-\check{\mathbb{U}}'\right\|_{H^{2}_{-q-\delta}(\Sigma)}.$$

Taking  $\varepsilon$  small enough compared to  $C_0^{-1}$  shows that  $\Psi^{(s)}$  is a contraction.

In view of Lemma 5.12, the Banach fixed point theorem implies that if  $\varepsilon$  is small enough then there exists a unique solution  $\tilde{\mathbb{U}}$  to the system (5.3) with the bound

$$\left\| \check{\mathbb{U}} \right\|_{H^2_{-q-\delta}(\Sigma)} \lesssim \varepsilon.$$

This concludes the proof of Theorem 3.1.

# References

- Aretakis, Stefanos. Horizon Instability of Extremal Black Holes. Advances in Theoretical and Mathematical Physics 19 (3) (2015), pp. 507–530.
- Bartnik, Robert. The Mass of an Asymptotically Flat Manifold. Communications on Pure and Applied Mathematics 39 (5) (1986), pp. 661–693.
- [3] Caciotta, Giulio and Nicolò, Francesco. Non Linear Perturbations of Kerr Spacetime in External Regions and the Peeling Decay. Annales Henri Poincaré 11 (3) (2010), pp. 433–497.
- [4] Carlotto, Alessandro. The General Relativistic Constraint Equations. Living Reviews in Relativity 24 (1) (2021).
- [5] Choquet-Bruhat, Yvonne. General Relativity and the Einstein Equations. Oxford Mathematical Monographs. Oxford; New York: Oxford University Press, 2009. 785 pp.
- [6] Choquet-Bruhat, Yvonne. Théorème d'existence Pour Certains Systèmes d'équations Aux Dérivées Partielles Non Linéaires. Acta Mathematica 88 (1952), pp. 141–225.
- [7] Choquet-Bruhat, Yvonne and Geroch, Robert. Global Aspects of the Cauchy Problem in General Relativity. Communications in Mathematical Physics 14 (4) (1969), pp. 329–335.
- [8] Christodoulou, Demetrios and Klainerman, Sergiu. The Global Nonlinear Stability of the Minkowski Space. 1993.
- [9] Chrusciel, Piotr and Delay, Erwann. On Mapping Properties of the General Relativistic Constraints Operator in Weighted Function Spaces, with Applications. Mémoires de la Société Mathématique de France 1 (2003), pp. 1–103.
- [10] Corvino, Justin. Scalar Curvature Deformation and a Gluing Construction for the Einstein Constraint Equations: Communications in Mathematical Physics **214** (1) (2000), pp. 137–189.
- [11] Corvino, Justin and Schoen, Richard M. On the Asymptotics for the Vacuum Einstein Constraint Equations. Journal of Differential Geometry 73 (2) (2006), pp. 185–217.
- [12] Dafermos, Mihalis et al. The Non-Linear Stability of the Schwarzschild Family of Black Holes. arXiv: 2104.08222 [gr-qc, physics:math-ph]. preprint.
- [13] Dilts, James et al. Non-CMC Solutions of the Einstein Constraint Equations on Asymptotically Euclidean Manifolds. Classical and Quantum Gravity 31 (6) (2014).
- [14] Fang, Allen Juntao, Szeftel, Jérémie, and Touati, Arthur. *Initial Data for Minkowski Stability with Arbitrary Decay.* arXiv: 2401.14353 [gr-qc, physics:math-ph]. preprint.
- [15] Fischer, Arthur E., Marsden, Jerrold E., and Moncrief, Vincent. The Structure of the Space of Solutions of Einstein's Equations. I. One Killing Field. Annales de l'institut Henri Poincaré. Section A, Physique Théorique 33 (2) (1980), pp. 147–194.
- [16] Gajic, Dejan. Azimuthal Instabilities on Extremal Kerr. arXiv: 2302.06636 [gr-qc]. preprint.
- [17] Giorgi, Elena, Klainerman, Sergiu, and Szeftel, Jeremie. Wave Equations Estimates and the Nonlinear Stability of Slowly Rotating Kerr Black Holes. arXiv: 2205.14808. preprint.
- [18] Klainerman, Sergiu and Szeftel, Jérémie. Construction of GCM Spheres in Perturbations of Kerr. Annals of PDE 8 (2) (2022).
- [19] Klainerman, Sergiu and Szeftel, Jérémie. Effective Results on Uniformization and Intrinsic GCM Spheres in Perturbations of Kerr. Annals of PDE 8 (2) (2022).
- [20] Klainerman, Sergiu and Szeftel, Jérémie. Global Nonlinear Stability of Schwarzschild Spacetime under Polarized Perturbations. Annals of Mathematics Studies 210. Princeton University Press, 2020.
- [21] Klainerman, Sergiu and Szeftel, Jérémie. Kerr Stability for Small Angular Momentum. Pure and Applied Mathematics Quarterly 19 (3) (2023), pp. 791–1678.
- [22] Le Rousseau, Jérôme, Lebeau, Gilles, and Robbiano, Luc. Elliptic Carleman Estimates and Applications to Stabilization and Controllability, Volume II: General Boundary Conditions on Riemannian Manifolds. Vol. 98. Progress in Nonlinear Differential Equations and Their Applications. Cham: Springer International Publishing, 2022.

- [23] Lichnerowicz, André. L'intégration Des Équations de La Gravitation Relativiste et Le Problème Des n Corps. Journal de Mathématiques Pures et Appliquées **23** (1944), pp. 37–63.
- [24] Maxwell, David. Initial Data for Black Holes and Rough Spacetimes. University of Washington, 2004.
- [25] Maxwell, David. Solutions of the Einstein Constraint Equations with Apparent Horizon Boundaries. Communications in Mathematical Physics 253 (3) (2005), pp. 561–583.
- [26] McOwen, Robert C. The Behavior of the Laplacian on Weighted Sobolev Spaces. Communications on Pure and Applied Mathematics 32 (6) (1979), pp. 783–795.
- [27] Shen, Dawei. Construction of GCM Hypersurfaces in Perturbations of Kerr. Annals of PDE 9 (1) (2023).
- [28] Shen, Dawei. Kerr Stability in External Regions. accepted for publication in Annals of PDE (2023).

UNIVERSITÄT MÜNSTER, MÜNSTER, DEUTSCHLAND (ALLEN.JUNTAO.FANG@UNI-MUENSTER.DE)

CNRS & LABORATOIRE JACQUES-LOUIS LIONS, SORBONNE UNIVERSITÉ, PARIS, (JEREMIE-SZEFTEL@SORBONNE-UNIVERSITE.FR)

INSTITUT DES HAUTES ETUDES SCIENTIFIQUES, BURES-SUR-YVETTE, (TOUATI@IHES.FR)