

SHARP MAXIMAL FUNCTION ESTIMATES FOR MULTILINEAR PSEUDO-DIFFERENTIAL OPERATORS OF TYPE $(0, 0)$

BAE JUN PARK AND NAOHITO TOMITA

ABSTRACT. In this paper, we study sharp maximal function estimates for multilinear pseudo-differential operators. Our target is operators of type $(0, 0)$ for which a differentiation does not make any decay of the associated symbol. Analogous results for operators of type (ρ, ρ) , $0 < \rho < 1$, appeared in an earlier work of the authors [16], but a different approach is given for $\rho = 0$.

1. INTRODUCTION

Let l be a positive integer greater or equal to 1 in this paper. For $0 < r < \infty$ and locally r -th integrable functions f_1, \dots, f_l on \mathbb{R}^n , we define the l -sublinear Hardy-Littlewood maximal function by

$$\mathbf{M}_r(f_1, \dots, f_l)(x) := \sup_{Q:x \in Q} \left(\frac{1}{|Q|^l} \int_{Q \times \dots \times Q} \prod_{j=1}^l |f_j(u_j)|^r d\vec{u} \right)^{\frac{1}{r}}$$

where $d\vec{u} := du_1 \cdots du_l$ and the supremum is taken over all cubes containing x . For a locally r th integrable function f on \mathbb{R}^n , we also define the (inhomogeneous and sublinear) sharp maximal function by

$$\begin{aligned} \mathcal{M}_r^\sharp f(x) := & \sup_{Q:x \in Q, \ell(Q) \geq 1} \left(\frac{1}{|Q|} \int_Q |f(y)|^r dy \right)^{\frac{1}{r}} \\ & + \sup_{Q:x \in Q, \ell(Q) < 1} \inf_{c_Q \in \mathbb{C}} \left(\frac{1}{|Q|} \int_Q |f(y) - c_Q|^r dy \right)^{\frac{1}{r}} \end{aligned}$$

where the supremum ranges over all cubes in \mathbb{R}^n containing x whose side-length is ≥ 1 in the first one, and is < 1 in the second one. Then

$$(1.1) \quad \|f\|_{bmo(\mathbb{R}^n)} := \|\mathcal{M}_1^\sharp f\|_{L^\infty(\mathbb{R}^n)} \sim \|\mathcal{M}_r^\sharp f\|_{L^\infty(\mathbb{R}^n)}$$

for $1 < r < \infty$, where bmo denotes the localized version of the usual space of functions with bounded mean oscillation.

Given $m \in \mathbb{R}$ and $0 \leq \rho < 1$, the l -linear Hörmander symbol class $\mathbb{M}_l S_{\rho, \rho}^m(\mathbb{R}^n)$ consists of infinitely many differentiable functions $\sigma(x, \xi_1, \dots, \xi_l)$ on $(\mathbb{R}^n)^{l+1}$ satisfying the property

2020 *Mathematics Subject Classification.* Primary 47G30, 42B25, 35S05, 42B37.

Key words and phrases. Pseudo-differential operator, Multilinear operator, Sharp maximal function, Weighted norm inequality.

B. Park is supported in part by NRF grant 2022R1F1A1063637 and by POSCO Science Fellowship of POSCO TJ Park Foundation. B. Park is grateful for support by the Open KIAS Center at Korea Institute for Advanced Study. N. Tomita was supported by JSPS KAKENHI Grant Number 20K03700.

that for any multi-indices $\alpha, \beta_1, \dots, \beta_l \in \{0, 1, 2, \dots\}^n$ there exists a constant $C_{\alpha, \beta_1, \dots, \beta_l} > 0$ such that

$$|\partial_x^\alpha \partial_{\xi_1}^{\beta_1} \cdots \partial_{\xi_l}^{\beta_l} \sigma(x, \xi_1, \dots, \beta_l)| \leq C_{\alpha, \beta_1, \dots, \beta_l} (1 + |\xi_1| + \cdots + |\beta_l|)^{m - \rho(|\beta_1| + \cdots + |\beta_l| - |\alpha|)}.$$

Let $\mathcal{S}(\mathbb{R}^n)$ denote the Schwartz class on \mathbb{R}^n . For a symbol $\sigma \in \mathbb{M}_l S_{\rho, \rho}^m(\mathbb{R}^n)$, the corresponding l -linear pseudo-differential operator T_σ is defined as

$$T_\sigma(f_1, \dots, f_l)(x) := \int_{(\mathbb{R}^n)^l} \sigma(x, \xi_1, \dots, \xi_l) \prod_{j=1}^l \widehat{f_j}(\xi_j) e^{2\pi i \langle x, \xi_1 + \cdots + \xi_l \rangle} d\vec{\xi}$$

for $f_1, \dots, f_l \in \mathcal{S}(\mathbb{R}^n)$, where $\widehat{f}(\xi) := \int_{\mathbb{R}^n} f(x) e^{-2\pi i \langle x, \xi \rangle} dx$ is the Fourier transform of f , and $d\vec{\xi} := d\xi_1 \cdots d\xi_l$. Denote by $\text{Op}\mathbb{M}_l S_{\rho, \rho}^m(\mathbb{R}^n)$ the family of such operators with symbols in $\mathbb{M}_l S_{\rho, \rho}^m(\mathbb{R}^n)$.

Then we have the following sharp maximal function estimates which originated in Chanillo and Torchinsky [4] for linear operators. See also Naibo [13] for bilinear ones.

Theorem A. [16] *Let $1 < r \leq 2$, $0 < \rho < 1$, and $l \geq 1$. Suppose that $m \leq -\frac{nl}{r}(1 - \rho)$ and $\sigma \in \mathbb{M}_l S_{\rho, \rho}^m(\mathbb{R}^n)$.*

(1) *If $l = 1$, then*

$$\mathcal{M}_1^\sharp(T_\sigma f)(x) \lesssim M_r f(x), \quad x \in \mathbb{R}^n$$

for $f \in \mathcal{S}(\mathbb{R}^n)$.

(2) *If $l \geq 2$, then*

$$\mathcal{M}_{\frac{r}{l}}^\sharp(T_\sigma(f_1, \dots, f_l))(x) \lesssim M_r(f_1, \dots, f_l)(x), \quad x \in \mathbb{R}^n$$

for all $f_1, \dots, f_l \in \mathcal{S}(\mathbb{R}^n)$.

Here and in the sequel, the symbol $A \lesssim B$ indicates that $A \leq CB$ for some constant $C > 0$ independent of the variable quantities A and B , and $A \sim B$ if $A \lesssim B$ and $B \lesssim A$ hold simultaneously.

The key idea to prove Theorem A is the kernel estimates in [16, Lemmas 2.1 and 3.1]. In order to illustrate this, take a simple case when $\sigma(x, \xi) = \sigma(\xi)$ in $\mathbb{M}_1 S_{\rho, \rho}^m(\mathbb{R}^n)$. By applying (inhomogeneous) Littlewood-Paley decomposition, which will be actually discussed in the next section, we can write $T_\sigma f = \sum_{k \in \mathbb{N}_0} T_{\sigma_k} f$ and the kernel $K_k(x, y)$ of the operator T_{σ_k} satisfies the size estimate

$$(1.2) \quad |K_k(x, y)| \lesssim_M 2^{-k(\rho N - m - n)} \frac{1}{|x - y|^N}, \quad |x - y| \gtrsim 1$$

by using integration by parts N times. If $\rho > 0$, by taking N sufficiently large, we have enough exponential decay $2^{-k(\rho N - m - n)}$ (even though ρ is very small). Essentially similar properties are inherent in [16, Lemmas 2.1 and 3.1]. This is not that serious when k is small enough, but since we will finally take the infinite sum over $k \in \{0, 1, 2, \dots\}$, appropriate decay in k should be necessary to make it summable. However, when $\rho = 0$ the decay property is no longer available and this prevents Theorem A from being extended to $\rho = 0$.

In this paper, we will establish analogous pointwise estimates for $\rho = 0$, overcoming the obstacle mentioned above regarding (1.2). Our main result is

Theorem 1.1. Let $1 < r \leq 2$ and $l \geq 1$. Suppose that $m \leq -\frac{nl}{r}$ and $\sigma \in \mathbb{M}_l S_{0,0}^m(\mathbb{R}^n)$. Then we have

$$\mathcal{M}_{\frac{r}{l}}^\sharp(T_\sigma(f_1, \dots, f_l))(x) \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x), \quad x \in \mathbb{R}^n$$

for all $f_1, \dots, f_l \in \mathcal{S}(\mathbb{R}^n)$.

As an extension of Theorem A to $\rho = 0$, Theorem 1.1 will be proved in a seemingly similar structure. However, we would like to emphasize that if $\rho = 0$, sufficient exponential decay for k cannot be derived through multiple integration by parts, as seen in (1.2), so a different approach that can compensate for the critical issue is needed. For this one, we will adopt a nontrivial decomposition of a function f (e.g. (3.7) and (4.3) below).

Now we recall multi-variable extension of A_p weight classes, introduced by Lerner, Ombrosi, Pérez, Torres, and Trujillo-González [8].

Definition 1. Let $1 < p_1, \dots, p_l < \infty$ and $1/p = 1/p_1 + \dots + 1/p_l$. Then we define $A_{\vec{p}}$, $\vec{p} = (p_1, \dots, p_l)$, to be the class of l -tuples of weights $\vec{w} := (w_1, \dots, w_l)$ satisfying

$$\sup_Q \left[\left(\frac{1}{|Q|} \int_Q v_{\vec{w}}(x) dx \right)^{1/p} \prod_{j=1}^l \left(\frac{1}{|Q|} \int_Q (w_j(x))^{1-p'_j} dx \right)^{1/p_j} \right] < \infty$$

where p'_j denotes the Hölder conjugate of p_j and

$$v_{\vec{w}}(x) := \prod_{j=1}^l (w_j(x))^{p/p_j}.$$

If $l = 1$, the class $A_{\vec{p}}$ coincides with the classical Muckenhoupt A_p class. It is known in [8] that for $1 < p_1, \dots, p_l < \infty$ with $1/p = 1/p_1 + \dots + 1/p_l$,

$$(1.3) \quad \vec{w} = (w_1, \dots, w_l) \in A_{\vec{p}} \text{ if and only if } \|\mathbf{M}(f_1, \dots, f_l)\|_{L^p(v_{\vec{w}})} \lesssim \prod_{j=1}^l \|f_j\|_{L^{p_j}(w_j)}$$

for all locally integrable functions f_1, \dots, f_l , where $\mathbf{M} := \mathbf{M}_1$. As discussed in [16, Theorem 1.6], Theorem A, together with (1.3), deduces a weighted norm inequality for $T_\sigma \in \text{Op}\mathbb{M}_l S_{\rho,\rho}^{-\frac{nl}{r}(1-\rho)}(\mathbb{R}^n)$, involving $\vec{w} = (w_1, \dots, w_l) \in A_{\frac{p_1}{r}, \dots, \frac{p_l}{r}}$, and it can be naturally extended to $\rho = 0$ replacing Theorem A by Theorem 1.1.

Theorem 1.2. Let $l \geq 1$, $1 < r \leq 2$, and $r < p_1, \dots, p_l < \infty$ with $1/p = 1/p_1 + \dots + 1/p_l$. Suppose that $\sigma \in \mathbb{M}_l S_{0,0}^{-\frac{nl}{r}}(\mathbb{R}^n)$. If an l -tuple of weights $\vec{w} = (w_1, \dots, w_l)$ belongs to the class $A_{\frac{p_1}{r}, \dots, \frac{p_l}{r}}$, then the weighted norm inequality

$$\|T_\sigma(f_1, \dots, f_l)\|_{L^p(v_{\vec{w}})} \lesssim \prod_{j=1}^l \|f_j\|_{L^{p_j}(w_j)}$$

holds for $f_1, \dots, f_l \in \mathcal{S}(\mathbb{R}^n)$.

The proof is exactly same as that of [16, Theorem 1.6], simply employing Theorem 1.1 instead of Theorem A. So it is omitted here.

We also have the following end-point estimate as a corollary of Theorem 1.1.

Corollary 1.3. Suppose that $\sigma \in \mathbb{M}_l S_{0,0}^{-\frac{nl}{2}}(\mathbb{R}^n)$. Then we have

$$(1.4) \quad \|T_\sigma(f_1, \dots, f_l)\|_{BMO(\mathbb{R}^n)} \lesssim \prod_{j=1}^l \|f_j\|_{L^\infty(\mathbb{R}^n)}$$

for all $f_1, \dots, f_l \in \mathcal{S}(\mathbb{R}^n)$.

The inequality (1.4) follows from the BMO characterization

$$\|f\|_{BMO(\mathbb{R}^n)} \sim_p \sup_Q \inf_{c_Q} \left(\frac{1}{|Q|} \int_Q |f(x) - c_Q|^p dx \right)^{1/p}, \quad 0 < p < \infty$$

and Theorem 1.1. See [16, Corollary 1.5] for more details.

Notation. Let \mathbb{N} and \mathbb{Z} be the sets of all natural numbers and integers, respectively, and $\mathbb{N}_0 := \mathbb{N} \cup \{0\}$. For each set U in \mathbb{R}^n we denote by χ_U its characteristic function and by U^c its complement. For each cube Q in \mathbb{R}^n , let $\ell(Q)$ denote the side-length of Q .

2. PRELIMINARIES

2.1. Decomposition of Pseudo-differential operators. Let Φ be a Schwartz function on \mathbb{R}^{nl} such that its Fourier transform $\widehat{\Phi}$ is equal to 1 on the unit ball centered at the origin and is supported in the ball of radius 2. Let $\Psi \in \mathcal{S}(\mathbb{R}^{nl})$ satisfy $\widehat{\Psi}(\vec{\xi}) := \widehat{\Phi}(\vec{\xi}) - \widehat{\Phi}(2\vec{\xi})$ for $\vec{\xi} \in \mathbb{R}^{nl}$. For each $k \in \mathbb{Z}$, we define $\Psi_k(\vec{x}) := 2^{knl}\Psi(2^k\vec{x})$ for $\vec{x} \in \mathbb{R}^{nl}$. Then $\{\Phi\} \cup \{\Psi_k\}_{k \in \mathbb{N}}$ forms inhomogeneous Littlewood-Paley partition of unity. Note that

$$\text{supp}(\widehat{\Psi}_k) \subset \{\vec{\xi} \in \mathbb{R}^{nl} : 2^{k-1} \leq |\vec{\xi}| \leq 2^{k+1}\}, \quad k \in \mathbb{Z}$$

and

$$(2.1) \quad \widehat{\Phi}(\vec{\xi}) + \sum_{k \in \mathbb{N}} \widehat{\Psi}_k(\vec{\xi}) = 1.$$

Using (2.1), we decompose $\sigma \in \mathbb{M}_l S_{0,0}^m(\mathbb{R}^n)$ as

$$\sigma(x, \vec{\xi}) = \sigma(x, \vec{\xi})\widehat{\Phi}(\vec{\xi}) + \sum_{k \in \mathbb{N}} \sigma(x, \vec{\xi})\widehat{\Psi}_k(\vec{\xi}) =: \sigma_0(x, \vec{\xi}) + \sum_{k \in \mathbb{N}} \sigma_k(x, \vec{\xi}), \quad \vec{\xi} \in \mathbb{R}^{nl}$$

and then express

$$T_\sigma(f_1, \dots, f_l) = \sum_{k \in \mathbb{N}_0} T_{\sigma_k}(f_1, \dots, f_l)$$

where T_{σ_k} are the l -linear pseudo-differential operators associated with $\sigma_k \in \mathbb{M}_l S_{0,0}^m(\mathbb{R}^n)$. Here, we notice that for all multi-indices $\alpha, \beta_1, \dots, \beta_l \in (\mathbb{N}_0)^n$

$$|\partial_x^\alpha \partial_{\xi_1}^{\beta_1} \cdots \partial_{\xi_l}^{\beta_l} \sigma_k(x, \xi_1, \dots, \xi_l)| \lesssim_{\alpha, \beta_1, \dots, \beta_l} (1 + |\xi_1| + \cdots + |\xi_l|)^m$$

where the implicit constant in the inequality is independent of k . In such a case, we will say σ_k belongs to $\mathbb{M}_l S_{0,0}^m(\mathbb{R}^n)$ uniformly in k .

2.2. Generalized trace theorem for Sobolev spaces. For $s > 0$ and $N \in \mathbb{N}$, let $(I_N - \Delta_N)^{s/2}$ denote the Bessel potential acting on tempered distributions on \mathbb{R}^N , defined by

$$(I_N - \Delta_N)^{\frac{s}{2}} F := ((1 + 4\pi^2 |\cdot|^2)^{\frac{s}{2}} \widehat{F})^\vee.$$

The Sobolev space $L_s^2(\mathbb{R}^N)$ consists of all $F \in L^2(\mathbb{R}^N)$ satisfying

$$\|F\|_{L_s^2(\mathbb{R}^N)} := \|(I_N - \Delta_N)^{\frac{s}{2}} F\|_{L^2(\mathbb{R}^N)} = \left(\int_{\mathbb{R}^N} (1 + 4\pi^2 |\xi|^2)^s |\widehat{F}(\xi)|^2 d\xi \right)^{\frac{1}{2}} < \infty.$$

Lemma 2.1. [16, Lemma 5.1] *Let $l \geq 2$ and $s > 0$. Then*

$$\|\tilde{G}\|_{L_s^2(\mathbb{R}^n)} \lesssim_{s,r,n} \|G\|_{L_{s+\frac{(l-1)n}{2}}^2((\mathbb{R}^n)^l)}$$

for Schwartz functions G on $(\mathbb{R}^n)^l$, where $\tilde{G}(x) = G(x_1, \dots, x_l)|_{x_1=\dots=x_l=x}$ and $x, x_1, \dots, x_l \in \mathbb{R}^n$.

2.3. General l -linear estimates for $T_\sigma \in \mathbb{M}_l S_{0,0}^m(\mathbb{R}^n)$. Let h^p denote the local Hardy space, introduced by Goldberg [6]. The boundedness of (linear) pseudo-differential operators of type $(0,0)$ on h^p was given by Calderón and Vaillancourt [3] for $p = 2$, Coifman and Meyer [5] for $1 < p < \infty$, and Miyachi [10] and Päiväranta and Somersalo [14] for $0 < p \leq 1$. The multilinear setting was studied by Bényi and Torres [1], Michalowski, Rule, and Staubach [9], Bényi, Bernicot, Maldonado, Naibo, and Torres [2], Miyachi and Tomita [11], and finally Kato, Miyachi, and Tomita [7] proved the following.

Theorem B. *Let $l \geq 1$ and $0 < p_1, \dots, p_l \leq \infty$ with $1/p_1 + \dots + 1/p_l = 1/p$. Suppose that $m \in \mathbb{R}$ and $\sigma \in \mathbb{M}_l S_{0,0}^m(\mathbb{R}^n)$. Then*

$$m \leq m_0(\vec{p}) := -n \left[\sum_{j=1}^l \max \left(\frac{1}{p_j}, \frac{1}{2} \right) - \min \left(\frac{1}{p}, \frac{1}{2} \right) \right], \quad \vec{p} := (p_1, \dots, p_l)$$

if and only if

$$\|T_\sigma(f_1, \dots, f_l)\|_{Y^p(\mathbb{R}^n)} \lesssim \prod_{j=1}^l \|f_j\|_{Y^{p_j}(\mathbb{R}^n)}$$

for $f_1, \dots, f_l \in \mathcal{S}(\mathbb{R}^n)$, where

$$Y^p := \begin{cases} h^p & 0 < p < \infty \\ bmo & p = \infty \end{cases}.$$

Note that the inequality $\|f\|_{L^p} \lesssim \|f\|_{h^p}$ holds for locally integrable functions f , where $0 < p \leq 1$. Thus Theorem B deduces that if $1 < r \leq 2$ and $\sigma \in \mathbb{M}_l S_{0,0}^{-n(\frac{l}{r}-\frac{1}{2})}(\mathbb{R}^n)$, then

$$(2.2) \quad \|T_\sigma(f_1, \dots, f_l)\|_{L^{r/l}(\mathbb{R}^n)} \lesssim \prod_{j=1}^l \|f_j\|_{L^r(\mathbb{R}^n)}.$$

3. KEY ESTIMATES

We fix $k \in \mathbb{N}_0$. Let P_k be the concentric dilate of Q with $\ell(P_k) \geq 10\sqrt{n}\ell(Q)$ and χ_{P_k} denote its characteristic function. We write

$$f_j = f_j \chi_{P_k} + f_j \chi_{P_k^c} =: f_{j,k}^{(0)} + f_{j,k}^{(1)} \quad \text{for } j = 1, \dots, l$$

so that each $T_{\sigma_k}(f_1, \dots, f_l)$ can be decomposed as

$$T_{\sigma_k}(f_1, \dots, f_l) = \sum_{\mu_1, \dots, \mu_l \in \{0,1\}} T_{\sigma_k}(f_{1,k}^{(\mu_1)}, f_{2,k}^{(\mu_2)}, \dots, f_{l,k}^{(\mu_l)}).$$

Then a key estimate for the proof of Theorem 1.1 is

Proposition 3.1. *Let $1 < r \leq 2$, $l \geq 1$, $k \in \mathbb{N}_0$, and $\ell(Q) < 1$. Suppose that $\sigma \in \mathbb{M}_l S_{0,0}^{-\frac{nl}{r}}(\mathbb{R}^n)$ and $x \in Q$. Then we have*

$$(3.1) \quad \inf_{c_Q \in \mathbb{C}} \left(\frac{1}{|Q|} \int_Q |T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})(y) - c_Q|^{\frac{r}{l}} dy \right)^{\frac{l}{r}} \lesssim \ell(P_k)^{\frac{nl}{r}} \mathbf{M}_r(f_1, \dots, f_l)(x)$$

uniformly in k .

In order to prove Proposition 3.1, we first need the following lemma, which is actually an extension of [16, Lemma 5.4] to $\rho = 0$.

Lemma 3.2. *Let $1 < r \leq 2$ and $l \geq 1$. Suppose that $\sigma \in \mathbb{M}_l S_{0,0}^{-\frac{nl}{r}}(\mathbb{R}^n)$. Then we have*

$$\|T_{\sigma}(f_1, \dots, f_l)\|_{bmo(\mathbb{R}^n)} \lesssim \prod_{j=1}^l \|f_j\|_{L^r(\mathbb{R}^n)}$$

for $f_1, \dots, f_l \in \mathcal{S}(\mathbb{R}^n)$.

Proof. The linear case $l = 1$ has been already proved by Miyachi and Yabuta [12, Lemma 5.6] by using duality argument with $(h^1)^* = bmo$, but we provide another proof, using Sobolev embeddings into bmo . We notice that the operator

$$(I_n - \Delta_n)^{\frac{n}{4}} T_{\sigma}(I_n - \Delta_n)^{\frac{1}{2}n(\frac{1}{r} - \frac{1}{2})}$$

is a pseudo-differential operator associated with a symbol in $\mathbb{M}_1 S_{0,0}^0(\mathbb{R}^n)$. Then its $L^2(\mathbb{R}^n)$ boundedness, in view of Theorem B, deduces that

$$\begin{aligned} \|T_{\sigma}f\|_{L_{\frac{n}{2}}^2(\mathbb{R}^n)} &= \left\| \left((I_n - \Delta_n)^{\frac{n}{4}} T_{\sigma}(I_n - \Delta_n)^{\frac{1}{2}n(\frac{1}{r} - \frac{1}{2})} \right) (I_n - \Delta_n)^{-\frac{1}{2}n(\frac{1}{r} - \frac{1}{2})} f \right\|_{L^2(\mathbb{R}^n)} \\ &\lesssim \left\| (I_n - \Delta_n)^{-\frac{1}{2}n(\frac{1}{r} - \frac{1}{2})} f \right\|_{L^2(\mathbb{R}^n)} = \|f\|_{L_{-\frac{n}{2}(\frac{1}{r} - \frac{1}{2})}^2(\mathbb{R}^n)} \lesssim \|f\|_{L^r(\mathbb{R}^n)} \end{aligned}$$

for any Schwartz functions f on \mathbb{R}^n , where the last inequality follows from the Sobolev embedding theorem, but we note that $r > 1$. Therefore, by the embedding $L_{\frac{n}{2}}^2(\mathbb{R}^n) \hookrightarrow bmo(\mathbb{R}^n)$ (e.g. [15, Theorem 2.3]), we obtain

$$\|T_{\sigma}f\|_{bmo(\mathbb{R}^n)} \lesssim \|T_{\sigma}f\|_{L_{\frac{n}{2}}^2(\mathbb{R}^n)} \lesssim \|f\|_{L^r(\mathbb{R}^n)},$$

as desired.

Now we assume $l \geq 2$. In this case, we mimic the proof of [16, Lemma 5.4]. We first define the linear symbol Σ in $\mathbb{R}^{nl} \times \mathbb{R}^{nl}$ as

$$(3.2) \quad \Sigma(X, \Xi) := \sigma(x_1, \xi_1, \dots, \xi_l), \quad X := (x_1, \dots, x_l) \in (\mathbb{R}^n)^l, \quad \Xi := (\xi_1, \dots, \xi_l) \in (\mathbb{R}^n)^l.$$

Then it is obvious that the symbol Σ belongs to the linear Hörmander class $\mathbb{M}_1 S_{0,0}^{-\frac{nl}{r}}(\mathbb{R}^{nl})$. Moreover, the operator

$$(I_{nl} - \Delta_{nl})^{\frac{nl}{4}} T_\Sigma (I_{nl} - \Delta_{nl})^{\frac{1}{2}nl(\frac{1}{r}-\frac{1}{2})}$$

is a linear pseudo-differential operator in $\text{Op}\mathbb{M}_1 S_{0,0}^0(\mathbb{R}^{nl})$. Now, by applying its $L^2(\mathbb{R}^{nl})$ boundedness in Theorem B, we have

$$\begin{aligned} \|T_\Sigma(G)\|_{L^2_{\frac{nl}{2}}(\mathbb{R}^{nl})} &= \left\| \left((I_{nl} - \Delta_{nl})^{\frac{nl}{4}} T_\Sigma (I_{nl} - \Delta_{nl})^{\frac{1}{2}nl(\frac{1}{r}-\frac{1}{2})} \right) (I_{nl} - \Delta_{nl})^{-\frac{1}{2}nl(\frac{1}{r}-\frac{1}{2})} G \right\|_{L^2(\mathbb{R}^{nl})} \\ &\lesssim \left\| (I_{nl} - \Delta_{nl})^{-\frac{1}{2}nl(\frac{1}{r}-\frac{1}{2})} G \right\|_{L^2(\mathbb{R}^{nl})} = \|G\|_{L^2_{-nl(\frac{1}{r}-\frac{1}{2})}(\mathbb{R}^{nl})} \lesssim \|G\|_{L^r(\mathbb{R}^{nl})} \end{aligned}$$

for any Schwartz functions G on \mathbb{R}^{nl} , where the Sobolev embedding theorem is applied in the last inequality. Finally, since

$$T_\sigma(f_1, \dots, f_l)(x) = T_\Sigma(f_1 \otimes \dots \otimes f_l)(x_1, \dots, x_l)|_{x_1=\dots=x_l=x}, \quad x \in \mathbb{R}^n,$$

the embedding $L^2_{\frac{n}{2}}(\mathbb{R}^n) \hookrightarrow bmo(\mathbb{R}^n)$ and Lemma 2.1 with $s = \frac{n}{2}$ deduce

$$\begin{aligned} \|T_\sigma(f_1, \dots, f_l)\|_{bmo(\mathbb{R}^n)} &\lesssim \|T_\sigma(f_1, \dots, f_l)\|_{L^2_{\frac{n}{2}}(\mathbb{R}^n)} \\ &\lesssim \|T_\Sigma(f_1 \otimes \dots \otimes f_l)\|_{L^2_{\frac{nl}{2}}(\mathbb{R}^{nl})} \\ &\lesssim \|f_1 \otimes \dots \otimes f_l\|_{L^r(\mathbb{R}^{nl})} = \prod_{j=1}^l \|f_j\|_{L^r(\mathbb{R}^n)}. \end{aligned}$$

This completes the proof. \square

Now we prove Propositions 3.1.

Proof of Proposition 3.1. In view of (1.1), together with Hölder's inequality if $l \geq 2$, the left-hand side of (3.1) is no more than

$$\|T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})\|_{bmo(\mathbb{R}^n)}$$

and this is, via Lemma 3.2, bounded by a constant times

$$\begin{aligned} \prod_{j=1}^l \|f_j\|_{L^r(P_k)} &= \left(\int_{P_k \times \dots \times P_k} |f_1(u_1) \cdots f_l(u_l)|^r du_1 \cdots du_l \right)^{\frac{1}{r}} \\ (3.3) \quad &\lesssim \ell(P_k)^{\frac{nl}{r}} \mathbf{M}_r(f_1, \dots, f_l)(x) \end{aligned}$$

as $x \in Q \subset P_k$. Then (3.1) follows. \square

If one of μ_j is equal to 1, we benefit from the following proposition.

Proposition 3.3. [16, Propositions 2.3 and 5.3] *Let $1 \leq r \leq 2$, $l \geq 1$, $k \in \mathbb{N}_0$, and $x, y \in Q$. Suppose that $\sigma \in \mathbb{M}_l S_{0,0}^{-\frac{nl}{r}}(\mathbb{R}^n)$. If each μ_j is 0 or 1, excluding $\mu_1 = \dots = \mu_l = 0$, then we have*

$$(3.4) \quad |T_{\sigma_k}(f_{1,k}^{(\mu_1)}, \dots, f_{l,k}^{(\mu_l)})(y)| \lesssim_N \ell(P_k)^{-(N-\frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x)$$

and

$$|T_{\sigma_k}(f_{1,k}^{(\mu_1)}, \dots, f_{l,k}^{(\mu_l)})(y) - T_{\sigma_k}(f_{1,k}^{(\mu_1)}, \dots, f_{l,k}^{(\mu_l)})(x)|$$

$$(3.5) \quad \lesssim_N 2^k \ell(Q) \ell(P_k)^{-(N - \frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x)$$

for any $N > \frac{nl}{r}$.

Another key estimate is

Proposition 3.4. *Let $1 < r \leq 2$, $l \geq 1$, $x \in Q$, and $\ell(Q) < 1$. Suppose that $\sigma \in \mathbb{M}_l S_{0,0}^{-\frac{nl}{r}}(\mathbb{R}^n)$. Then we have*

$$(3.6) \quad \left(\frac{1}{|Q|} \int_Q \left| \sum_{k:2^k \ell(Q) \geq 1} T_{\sigma_k}(f_1, \dots, f_l)(y) \right|^{\frac{r}{l}} dy \right)^{\frac{l}{r}} \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x).$$

Proof. We note that $2^k \ell(Q) \geq 1$ implies $k \geq 1$. Therefore, we don't need to consider the case $k = 0$. We choose $0 < \epsilon < \frac{r}{2l} (\leq 1)$ and let $\mathcal{Q}_{k,\epsilon}^*$ be the concentric dilate of Q with $\ell(\mathcal{Q}_{k,\epsilon}^*) = 10\sqrt{nl}(2^k \ell(Q))^\epsilon (\geq 10\sqrt{nl} > 10\sqrt{nl} \ell(Q))$. Then we write

$$(3.7) \quad f_j = f_j \chi_{\mathcal{Q}_{k,\epsilon}^*} + f_j \chi_{(\mathcal{Q}_{k,\epsilon}^*)^c} =: f_{j,k}^{(0)} + f_{j,k}^{(1)} \quad \text{for } j = 1, \dots, l$$

so that the left-hand side of (3.6) is less than the sum of

$$\mathcal{J}_1 := \left(\sum_{k:2^k \ell(Q) \geq 1} \left(\frac{1}{|Q|} \int_Q \left| T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})(y) \right|^{\frac{r}{l}} dy \right)^{\min(1, \frac{l}{r})} \right)^{\max(1, \frac{l}{r})}$$

and

$$\mathcal{J}_2 := \sum_{\substack{\mu_1, \dots, \mu_l \in \{0,1\} \\ (\mu_1, \dots, \mu_l) \neq (0, \dots, 0)}} \left(\sum_{k:2^k \ell(Q) \geq 1} \left(\frac{1}{|Q|} \int_Q \left| T_{\sigma_k}(f_{k,1}^{(\mu_1)}, \dots, f_{k,l}^{(\mu_l)})(y) \right|^{\frac{r}{l}} dy \right)^{\min(1, \frac{l}{r})} \right)^{\max(1, \frac{l}{r})}$$

where the subadditivities of $\|\cdot\|_{\ell^r}$ and $\|\cdot\|_{\ell^{\frac{r}{l}}}$ if $l \geq 2$ are applied.

We consider \mathcal{J}_2 first. By applying (3.4) with $N > \frac{nl}{r}$, if $(\mu_1, \dots, \mu_l) \in \{0, 1\}^l \setminus \{(0, \dots, 0)\}$,

$$\begin{aligned} & \left(\frac{1}{|Q|} \int_Q \left| T_{\sigma_k}(f_{k,1}^{(\mu_1)}, \dots, f_{k,l}^{(\mu_l)})(y) \right|^{\frac{r}{l}} dy \right)^{\frac{l}{r}} \\ & \lesssim \ell(\mathcal{Q}_{k,\epsilon}^*)^{-(N - \frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x) \sim (2^k \ell(Q))^{-\epsilon(N - \frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x) \end{aligned}$$

so that

$$\begin{aligned} \mathcal{J}_2 & \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x) \left(\sum_{k:2^k \ell(Q) \geq 1} (2^k \ell(Q))^{-\epsilon(\frac{rN}{l} - n) \min(1, \frac{l}{r})} \right)^{\max(1, \frac{l}{r})} \\ & \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x). \end{aligned}$$

Now it remains to estimate \mathcal{J}_1 and we will separately treat the cases $l = 1$ and $l \geq 2$. Let us first consider the case $l = 1$. We define $\gamma_k := \Psi_{k-1} + \Psi_k + \Psi_{k+1} \in \mathcal{S}(\mathbb{R}^n)$ so that

$$\Psi_k * \gamma_k = \Psi_k \quad \text{for } k \geq 1.$$

This yields

$$T_{\sigma_k} f_k^{(0)} = T_{\sigma_k}(\gamma_k * f_k^{(0)}).$$

By applying Hölder's inequality and observing $2^{\frac{kn}{r}}\sigma_k \in \mathbb{M}_1 S_{0,0}^0(\mathbb{R}^n)$ uniformly in k , the L^2 boundedness for $2^{\frac{kn}{r}}T_{\sigma_k}$ in Theorem B deduces

$$\begin{aligned} \left(\frac{1}{|Q|} \int_Q |T_{\sigma_k} f_k^{(0)}(y)|^r dy \right)^{\frac{1}{r}} &\leq \ell(Q)^{-\frac{n}{2}} \|T_{\sigma_k}(\gamma_k * f_k^{(0)})\|_{L^2(\mathbb{R}^n)} \\ &\lesssim \ell(Q)^{-\frac{n}{2}} 2^{-\frac{kn}{r}} \|\gamma_k * f_k^{(0)}\|_{L^2(\mathbb{R}^n)} \end{aligned}$$

and this is bounded by

$$\ell(Q)^{-\frac{n}{2}} 2^{-\frac{kn}{r}} \|\gamma_k\|_{L^q(\mathbb{R}^n)} \|f_k^{(0)}\|_{L^r(\mathbb{R}^n)} \sim (2^k \ell(Q))^{-\frac{n}{2}} \|f\|_{L^r(\mathcal{Q}_{k,\epsilon}^*)} \lesssim (2^k \ell(Q))^{-n(\frac{1}{2}-\frac{\epsilon}{r})} M_r f(x)$$

where we applied Young's convolution inequality with $1 + \frac{1}{2} = \frac{1}{q} + \frac{1}{r}$. This finally proves

$$\mathcal{J}_1 \lesssim M_r f(x) \sum_{k: 2^k \ell(Q) \geq 1} (2^k \ell(Q))^{-n(\frac{1}{2}-\frac{\epsilon}{r})} \lesssim M_r f(x),$$

as desired.

Now assume $l \geq 2$. In this case, we will employ the trace theorem in Lemma 2.1 and the $L^2(\mathbb{R}^{nl})$ boundedness of a linear pseudo-differential operator belonging to $\mathbb{M}_1 S_{0,0}^0(\mathbb{R}^{nl})$. Since $2 < \frac{r}{\epsilon l}$, we may choose t such that

$$2 < t < \frac{r}{\epsilon l}.$$

Then by using Hölder's inequality with $\frac{r}{t} < t$ and the Sobolev embedding theorem,

$$\begin{aligned} \left(\frac{1}{|Q|} \int_Q |T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})(y)|^{\frac{r}{t}} dy \right)^{\frac{t}{r}} &\leq \frac{1}{|Q|^{\frac{1}{t}}} \|T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})\|_{L^t(\mathbb{R}^n)} \\ &\lesssim \ell(Q)^{-\frac{n}{t}} \|T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})\|_{L^2_{\frac{n}{2}-\frac{n}{t}}(\mathbb{R}^n)}. \end{aligned}$$

Let $\Gamma_k := \Psi_{k-1} + \Psi_k + \Psi_{k+1} \in \mathcal{S}(\mathbb{R}^{nl})$ so that $\Psi_k * \Gamma_k = \Psi_k$ for $k \geq 1$. Setting Σ as in (3.2) and $\Sigma_k := \Sigma \cdot \widehat{\Psi_k}$, we write

$$T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})(y) = T_{\Sigma_k} F_k(y, \dots, y), \quad y \in \mathbb{R}^n$$

where $T_{\Sigma_k} \in \text{Op}\mathbb{M}_1 S_{0,0}^{-\frac{nl}{r}}(\mathbb{R}^{nl})$ and $F_k := \Gamma_k * (f_{1,k}^{(0)} \otimes \dots \otimes f_{l,k}^{(0)})$ is a function defined on \mathbb{R}^{nl} . Therefore, applying Lemma 2.1 with $\frac{n}{2} - \frac{n}{t} > 0$,

$$\begin{aligned} \|T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})\|_{L^2_{\frac{n}{2}-\frac{n}{t}}(\mathbb{R}^n)} &\lesssim \|T_{\Sigma_k} F_k\|_{L^2_{\frac{nl}{2}-\frac{n}{t}}(\mathbb{R}^{nl})} \\ &= \left\| \left((I_{nl} - \Delta_{nl})^{\frac{1}{2}(\frac{nl}{2}-\frac{n}{t})} T_{\Sigma_k} \right) F_k \right\|_{L^2(\mathbb{R}^{nl})}. \end{aligned}$$

Observing

$$2^{kn(\frac{l}{r}+\frac{1}{t}-\frac{l}{2})} (I_{nl} - \Delta_{nl})^{\frac{1}{2}(\frac{nl}{2}-\frac{n}{t})} T_{\Sigma_k} \in \text{Op}\mathbb{M}_1 S_{0,0}^0(\mathbb{R}^{nl}) \quad \text{uniformly in } k,$$

its $L^2(\mathbb{R}^{nl})$ boundedness yields that the L^2 norm in the last expression is bounded by a constant times

$$2^{-kn(\frac{l}{r}+\frac{1}{t}-\frac{l}{2})} \|F_k\|_{L^2(\mathbb{R}^{nl})} \lesssim 2^{-kn(\frac{l}{r}+\frac{1}{t}-\frac{l}{2})} \|\Gamma_k\|_{L^q(\mathbb{R}^{nl})} \|f_{1,k}^{(0)} \otimes \dots \otimes f_{l,k}^{(0)}\|_{L^r(\mathbb{R}^{nl})}$$

$$\sim 2^{-kn(\frac{l}{r} + \frac{1}{t} - \frac{l}{2})} 2^{knl(1 - \frac{1}{q})} \prod_{j=1}^l \|f_j\|_{L^r(Q_{k,\epsilon}^*)} = 2^{-\frac{kn}{t}} \prod_{j=1}^l \|f_j\|_{L^r(Q_{k,\epsilon}^*)}$$

where the inequality follows from Young's inequality with $1 + \frac{1}{2} = \frac{1}{q} + \frac{1}{r}$. Using (3.3), the last displayed expression is controlled by a constant multiple of

$$2^{-\frac{kn}{t}} (2^k \ell(Q))^{\frac{\epsilon nl}{r}} \mathbf{M}_r(f_1, \dots, f_l)(x).$$

Finally, we have proved that

$$\left(\frac{1}{|Q|} \int_Q |T_{\sigma_k}(f_{1,k}^{(0)}, \dots, f_{l,k}^{(0)})(y)|^{\frac{r}{t}} dy \right)^{\frac{l}{r}} \lesssim (2^k \ell(Q))^{-n(\frac{1}{t} - \frac{\epsilon l}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x)$$

and this clearly implies that

$$\mathcal{J}_1 \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x) \left(\sum_{k: 2^k \ell(Q) \geq 1} (2^k \ell(Q))^{-n(\frac{r}{tl} - \epsilon)} \right)^{\frac{l}{r}} \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x).$$

This completes the proof. \square

4. PROOF OF THEOREM 1.1

Fixing a cube Q in \mathbb{R}^n and a point $x \in Q$, our focus shifts to demonstrating

$$(4.1) \quad \left(\frac{1}{|Q|} \int_Q |T_\sigma(f_1, \dots, f_l)(y)|^{\frac{r}{t}} dy \right)^{\frac{l}{r}} \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x) \quad \text{if } \ell(Q) \geq 1$$

and

$$(4.2) \quad \inf_{c_Q \in \mathbb{C}} \left(\frac{1}{|Q|} \int_Q |T_\sigma(f_1, \dots, f_l)(y) - c_Q|^{\frac{r}{t}} dy \right)^{\frac{l}{r}} \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x) \quad \text{if } \ell(Q) < 1$$

uniformly in Q and x .

4.1. Proof of (4.1). Assume $\ell(Q) \geq 1$. Let $0 < \epsilon < \frac{r}{2l}$. For each $k \in \mathbb{N}_0$, let $Q_{k,\epsilon}^*$ denote the concentric dilate of Q with $\ell(Q_{k,\epsilon}^*) = 10\sqrt{n}l2^k\ell(Q)$. Then we write

$$(4.3) \quad f_j = f_j \chi_{Q_{k,\epsilon}^*} + f_j \chi_{(Q_{k,\epsilon}^*)^c} =: \mathbf{f}_{j,k}^{(0)} + \mathbf{f}_{j,k}^{(1)} \quad \text{for } j = 1, \dots, l$$

so that $T_\sigma(f_1, \dots, f_l)$ can be decomposed as

$$T_\sigma(f_1, \dots, f_l) = \sum_{k \in \mathbb{N}_0} \sum_{\mu_1, \dots, \mu_l \in \{0, 1\}} T_{\sigma_k}(\mathbf{f}_{1,k}^{(\mu_1)}, \dots, \mathbf{f}_{l,k}^{(\mu_l)}).$$

Therefore, the left-hand side of (4.1) is controlled by a constant times

$$\sum_{\mu_1, \dots, \mu_l \in \{0, 1\}} \left(\sum_{k \in \mathbb{N}_0} \left(\frac{1}{|Q|} \int_Q |T_{\sigma_k}(\mathbf{f}_{1,k}^{(\mu_1)}, \dots, \mathbf{f}_{l,k}^{(\mu_l)})(y)|^{\frac{r}{t}} dy \right)^{\min(1, \frac{l}{r})} \right)^{\max(1, \frac{l}{r})}.$$

Now we claim that for each $k \in \mathbb{N}_0$,

$$(4.4) \quad \left(\frac{1}{|Q|} \int_Q |T_{\sigma_k}(\mathbf{f}_{1,k}^{(0)}, \dots, \mathbf{f}_{l,k}^{(0)})(y)|^{\frac{r}{t}} dy \right)^{\frac{l}{r}} \lesssim 2^{-kn(\frac{1}{2} - \frac{\epsilon l}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x)$$

and for $N > \frac{nl}{r}$

$$(4.5) \quad \left(\frac{1}{|Q|} \int_Q |T_{\sigma_k}(\mathbf{f}_{1,k}^{(\mu_1)}, \dots, \mathbf{f}_{l,k}^{(\mu_l)})(y)|^{\frac{r}{l}} dy \right)^{\frac{l}{r}} \lesssim 2^{-\epsilon k(N - \frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x)$$

if at least one of μ_j is equal to 1. Then (4.1) follows obviously.

By observing

$$2^{\frac{kn}{2}} \sigma_k \in \mathbb{M}_l S_{0,0}^{-n(\frac{l}{r} - \frac{1}{2})}(\mathbb{R}^n) \quad \text{uniformly in } k$$

and applying (2.2), the left-hand side of (4.4) is bounded by

$$\frac{1}{|Q|^{\frac{l}{r}}} \|T_{\sigma_k}(\mathbf{f}_{1,k}^{(0)}, \dots, \mathbf{f}_{l,k}^{(0)})\|_{L^{\frac{r}{l}}(\mathbb{R}^n)} \lesssim 2^{-\frac{kn}{2}} \ell(Q)^{-\frac{nl}{r}} \prod_{j=1}^l \|f_j\|_{L^r(Q_{k,\epsilon}^*)}.$$

Using the argument in (3.3) with $P_k = Q_{k,\epsilon}^*$, we have

$$\prod_{j=1}^l \|f_j\|_{L^r(Q_{k,\epsilon}^*)} \lesssim \ell(Q_{k,\epsilon}^*)^{\frac{nl}{r}} \mathbf{M}_r(f_1, \dots, f_l)(x) \sim (2^{\epsilon k} \ell(Q))^{\frac{nl}{r}} \mathbf{M}_r(f_1, \dots, f_l)(x)$$

and this yields (4.4).

Now assume that one of μ_j is 1. By employing (3.4), for $N > \frac{nl}{r}$ and $y \in Q$, we have

$$\begin{aligned} |T_{\sigma_k}(\mathbf{f}_{1,k}^{(\mu_1)}, \dots, \mathbf{f}_{l,k}^{(\mu_l)})(y)| &\lesssim (\ell(Q_{k,\epsilon}^*))^{-(N - \frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x) \\ &\sim (2^{\epsilon k} \ell(Q))^{-(N - \frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x) \\ &\leq 2^{-\epsilon k(N - \frac{nl}{r})} \mathbf{M}_r(f_1, \dots, f_l)(x) \end{aligned}$$

where the last inequality holds due to $\ell(Q) \geq 1$. This completes the proof of (4.5).

4.2. Proof of (4.2). Assume $\ell(Q) < 1$. Let P_Q be the concentric dilate of Q with $\ell(P_Q) = 10\sqrt{nl} (> 10\sqrt{nl}\ell(Q))$. We write

$$f_j = f_j \chi_{P_Q} + f_j \chi_{(P_Q)^c} =: \mathbf{f}_j^{(0)} + \mathbf{f}_j^{(1)} \quad \text{for } j = 1, \dots, l.$$

Then

$$\begin{aligned} &\left(\frac{1}{|Q|} \int_Q |T_\sigma(f_1, \dots, f_l)(y) - c_Q|^{\frac{r}{l}} dy \right)^{\frac{l}{r}} \\ &\lesssim \left(\frac{1}{|Q|} \int_Q |T_\sigma(\mathbf{f}_1^{(0)}, \dots, \mathbf{f}_l^{(0)})(y) - d_Q|^{\frac{r}{l}} dy \right)^{\frac{l}{r}} \\ &\quad + \sum_{\substack{\mu_1, \dots, \mu_l \in \{0,1\} \\ (\mu_1, \dots, \mu_l) \neq (0, \dots, 0)}} \left(\frac{1}{|Q|} \int_Q |T_\sigma(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(y) - \sum_{k: 2^k \ell(Q) < 1} T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(x)|^{\frac{r}{l}} dy \right)^{\frac{l}{r}} \end{aligned}$$

where

$$d_Q := c_Q - \sum_{\substack{\mu_1, \dots, \mu_l \in \{0,1\} \\ (\mu_1, \dots, \mu_l) \neq (0, \dots, 0)}} \sum_{k: 2^k \ell(Q) < 1} T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(x).$$

Therefore, in order to establish (4.2), we need to estimate

$$\mathcal{I}_1 := \inf_{d_Q \in \mathbb{C}} \left(\frac{1}{|Q|} \int_Q \left| T_\sigma(\mathbf{f}_1^{(0)}, \dots, \mathbf{f}_l^{(0)})(y) - d_Q \right|^{\frac{l}{r}} dy \right)^{\frac{l}{r}}$$

and for $(\mu_1, \dots, \mu_l) \in \{0, 1\}^l \setminus \{(0, \dots, 0)\}$

$$\mathcal{I}_2^{(\mu_1, \dots, \mu_l)} := \left(\frac{1}{|Q|} \int_Q \left| T_\sigma(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(y) - \sum_{k: 2^k \ell(Q) < 1} T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(x) \right|^{\frac{l}{r}} dy \right)^{\frac{l}{r}}.$$

Using Proposition 3.1, we have

$$\mathcal{I}_1 \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x).$$

For the other terms, we assume $(\mu_1, \dots, \mu_l) \in \{0, 1\}^l \setminus \{(0, \dots, 0)\}$ and write

$$\begin{aligned} & \left| T_\sigma(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(y) - \sum_{k: 2^k \ell(Q) < 1} T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(x) \right| \\ & \leq \sum_{k: 2^k \ell(Q) < 1} \left| T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(y) - T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(x) \right| \\ & \quad + \sum_{k: 2^k \ell(Q) \geq 1} \left| T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(y) \right| \\ & =: \mathcal{J}_1^{(\mu_1, \dots, \mu_l)}(y) + \mathcal{J}_2^{(\mu_1, \dots, \mu_l)}(y). \end{aligned}$$

By applying (3.5), if $y \in Q$, then we have

$$\left| T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(y) - T_{\sigma_k}(\mathbf{f}_1^{(\mu_1)}, \dots, \mathbf{f}_l^{(\mu_l)})(x) \right| \lesssim 2^k \ell(Q) \mathbf{M}_r(f_1, \dots, f_l)(x)$$

and thus

$$\mathcal{J}_1^{(\mu_1, \dots, \mu_l)}(y) \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x) \sum_{k: 2^k \ell(Q) < 1} 2^k \ell(Q) \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x).$$

This yields that

$$\left(\frac{1}{|Q|} \int_Q \left(\mathcal{J}_1^{(\mu_1, \dots, \mu_l)}(y) \right)^{\frac{l}{r}} dy \right)^{\frac{l}{r}} \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x).$$

Furthermore, Proposition 3.4, together with the property

$$|\mathbf{f}_j^{(\mu_j)}(y)| \leq |f_j(y)| \quad j = 1, \dots, l,$$

yields

$$\left(\frac{1}{|Q|} \int_Q \left(\mathcal{J}_2^{(\mu_1, \dots, \mu_l)}(y) \right)^{\frac{l}{r}} dy \right)^{\frac{l}{r}} \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x),$$

which proves

$$\mathcal{I}_2^{(\mu_1, \dots, \mu_l)} \lesssim \mathbf{M}_r(f_1, \dots, f_l)(x)$$

for $(\mu_1, \dots, \mu_l) \in \{0, 1\}^l \setminus \{(0, \dots, 0)\}$. This completes the proof.

REFERENCES

- [1] Á. Bényi and R. Torres, *Almost orthogonality and a class of bounded bilinear pseudodifferential operators*, Math. Res. Lett. **11** (2004), 1-11.
- [2] Á. Bényi, F. Bernicot, D. Maldonado, V. Naibo, and R. Torres, *On the Hörmander classes of bilinear pseudodifferential operators II*, Indiana Univ. Math. J. **62** (2013), 1733-1764.
- [3] A. P. Calderón and R. Vaillancourt, *A class of bounded pseudo-differential operators*, Proc. Nat. Acad. Sci. U.S.A. **69** (1972) 1185-1187.
- [4] S. Chanillo and A. Torchinsky, *Sharp function and weighted L^p estimates for a class of pseudo-differential operators*, Ark. Mat. **24** (1985), 1-25.
- [5] R. R. Coifman and Y. Meyer, *Au delà des opérateurs pseudo-différentiels*, Astérisque **57** (1978), 1-185.
- [6] D. Goldberg, *A local version of real Hardy spaces*, Duke Math. J. **46** (1979) 27-42.
- [7] T. Kato, A. Miyachi, and N. Tomita, *Boundedness of multilinear pseudo-differential operators with symbols in the Hörmander class $S_{0,0}$* , J. Funct. Anal. **282** (2022) Paper No. 109329, 28 pp.
- [8] A. Lerner, S. Ombrosi, C. Pérez, R. Torres, and R. Trujillo-González, *New maximal functions and multiple weights for the multilinear Calderón-Zygmund theory*, Adv. Math. **220** (2009), 1222-1264.
- [9] N. Michalowski, D. Rule, and W. Staubach, *Multilinear pseudodifferential operators beyond Calderón-Zygmund theory*, J. Math. Anal. Appl., **414**, (2014), 149-165.
- [10] A. Miyachi, *Estimates for pseudo-differential operators of class $S_{0,0}$* , Math. Nachr. **133** (1987), 135-154.
- [11] A. Miyachi and N. Tomita, *Calderón-Vaillancourt-type theorem for bilinear operators*, Indiana Univ. Math. J. **62** (2013), 1165-1201.
- [12] A. Miyachi and K. Yabuta, *Sharp function estimates for pseudo-differential operators of class $S_{\rho,\delta}^m$* , Bull. Fac. Sci. Ibaraki Univ. Ser. A **19** (1987), 15-30.
- [13] V. Naibo, *On the $L^\infty \times L^\infty \rightarrow BMO$ mapping property for certain bilinear pseudodifferential operators*, Proc. Amer. Math. Soc. **143** (2015), 5323 - 5336.
- [14] L. Päivärinta and E. Somersalo, *A generalization of the Calderón-Vaillancourt theorem to L^p and h^p* , Math. Nachr. **138** (1988) 145-156.
- [15] B. Park, *Some maximal inequalities on Triebel-Lizorkin spaces for $p = \infty$* , Math. Nachr. **292** (2019) 1137-1150.
- [16] B. Park and N. Tomita, *Sharp maximal function estimates for linear and multilinear pseudo-differential operators*, submitted.

B. PARK, DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 16419, REPUBLIC OF KOREA

Email address: bspark43@skku.edu

N. TOMITA, DEPARTMENT OF MATHEMATICS, GRADUATE SCHOOL OF SCIENCE, OSAKA UNIVERSITY, TOYONAKA, OSAKA 560-0043, JAPAN

Email address: tomita@math.sci.osaka-u.ac.jp