

Topology of restricted phase space thermodynamics in Kerr-Sen-Ads black holes

Bidyut Hazarika^{1*} and Prabwal Phukon^{1,2†}

1. *Department of Physics, Dibrugarh University, Dibrugarh, Assam, 786004.*

2. *Theoretical Physics Division, Centre for Atmospheric Studies,
Dibrugarh University, Dibrugarh, Assam, 786004.*

ABSTRACT

In this study, we investigate the thermodynamic topology of the Kerr-Sen-Ads black hole in restricted phase space. In the restricted phase space, a new parameter, central charge C , and its conjugate parameter μ are introduced, omitting the well-known PdV term in the first law of black hole thermodynamics. We study the local and global topology of the black hole by considering the black hole solution as topological defects in the free energy landscape. We compute the winding number and the total topological number at the thermodynamic defects. For our analysis, we have considered five ensembles of Kerr-Sen-Ads black holes in restricted phase space: fixed (Q, J, C) , fixed (ϕ, J, C) , fixed (Q, Ω, C) , fixed (Q, J, μ) , and fixed (ϕ, Ω, C) , where Q is the electric charge, J is the angular momentum, C is the central charge, ϕ is the electric potential conjugate to charge, Ω is the angular frequency conjugate to J , and finally, μ is the chemical potential. In the fixed (Q, J, C) , fixed (ϕ, J, C) , and fixed (Q, J, μ) ensembles, we find a topological charge of $+1$. In the fixed (Q, Ω, C) and fixed (ϕ, Ω, C) ensembles, depending on the values of the thermodynamic parameters, we find topological charges of -1 , 0 , and $+1$. Interestingly, in ensembles where we find the topological charge to be 0 , we observe both Hawking-Page and Davies type phase transitions. We show that both types of these phase transitions can be studied using a common vector field, and the topological charges associated with Davies type and Hawking-Page phase transitions are -1 and $+1$, respectively.

I. INTRODUCTION:

The study of black hole thermodynamics has made substantial strides since its inception in the 1970s [5–8]. Further studies have lead a multitude of intriguing findings which contribute to a better understanding of this area [9–17]. One captivating aspect of black hole thermodynamics is the study of black holes phase transitions. Black holes exhibit various forms of these transitions, including the Davies-type phase transition [18], the Hawking–Page phase transition [19], extremal phase transitions [20–28], van der Waals-type phase transitions [29–36] etc. The study of black hole phase transition behavior has been given a new dimension through the introduction of extended phase-space thermodynamics [37–49]. This framework introduces a novel pair of variables (P, V) into the domain of thermodynamic parameters, where pressure can be expressed in terms of the negative cosmological constant as :

$$P = \frac{-\Lambda}{8\pi G}$$

Subsequently, applying AdS/CFT correspondence [50], Visser introduced a new formalism [51] (which is an extension of extended thermodynamic phase space), by adding CFT central charge C and its conjugate parameter μ to the first law of black hole thermodynamics. Moreover in Visser’s formalism the pair (P, V) , is transformed into a new pair $(\mathcal{P}, \mathcal{V})$. Here the CFT volume $\mathcal{V} L^{d-2}$ and the corresponding pressure term \mathcal{P} is related through the CFT equation of state $E = (d - 2)\mathcal{P}\mathcal{V}$ where L is the Ads radius. d is the spacetime dimension in the bulk. So, for a charged rotating black hole the first law of black hole thermodynamics takes the form :

$$dE = TdS - \mathcal{P}d\mathcal{V} + \tilde{\phi}\tilde{Q} + \Omega J + \mu C$$

where, \tilde{Q} and $\tilde{\phi}$ are the rescaled electric charge and its conjugate electric potential respectively.

A restricted version of Visser’s formalism known as the restricted phase space thermodynamics (RPST) for Ads black holes was proposed [52]. These formalism has been applied to a number of black holes [53–63]. Studying black hole thermodynamics in RPST formalism has the potential to provide better understanding of AdS/CFT correspondence. Apart from extended phase space and RPST formalism, holographic thermodynamics [51, 64–66], geometric

* *rs.bidyuthazarika@dibru.ac.in*

† *prabwal@dibru.ac.in*

interpretation of phase transitions[67–76] are a few of the techniques used to study critical phenomenon in black hole thermodynamics.

A novel avenue in the context of understanding critical phenomena in black hole thermodynamics is the incorporation of topology in black hole thermodynamics[79, 80]. In one of these approaches, to investigate the local and global thermodynamic topology, black hole solutions are considered as topological defects within their thermodynamic phase space[80]. At these defects, a fundamental quantity known as the topological charge can be assigned. The topological charge signifies the distinct topological class present for a particular black hole solution. The study of thermodynamic topology has been extended to a range of black hole solutions found within various gravitational theories [81–130]. The following paragraph provides a brief overview of the off-shell method and outlines the procedure for computing the topological charge.

The expression for the off-shell free energy of a black hole with arbitrary mass is given by:

$$\mathcal{F} = E - \frac{S}{\tau} \quad (1)$$

Here, E and S are the energy and entropy of the black hole respectively. τ is the time scale parameter which equals to :

$$\tau = \frac{1}{T} \quad (2)$$

Here, T is the equilibrium temperature at the surface of the cavity that encloses the black hole. The time parameter τ is set freely to vary instead of the mass. Utilizing the generalized free energy, a vector field is constructed as follows :

$$\phi = (\phi^S, \phi^\Theta) = \left(\frac{\partial \mathcal{F}}{\partial S}, -\cot \Theta \csc \Theta \right) \quad (3)$$

Where $\theta = \pi/2$ and $\tau = 1/T$ represent a zero point of the vector field ϕ . In the θ component, a particular combination of trigonometric functions is chosen because at $\theta = \frac{\pi}{2}$, the vector field diverges, allowing one to easily identify the zero point of the S component by simply examining the vectors along the $\theta = \frac{\pi}{2}$ line. While constructing the vector field the Duan's ϕ mapping current theory has been taken into consideration [77, 78]. Using Duan's theory, the topological current can be written as

$$j^\mu = \frac{1}{2\pi} \epsilon^{\mu\nu\rho} \epsilon_{ab} \partial_\nu n^a \partial_\rho n^b \quad \mu, \nu, \rho = 0, 1, 2 \quad (4)$$

Here $\partial_\nu = \frac{\partial}{\partial x^\nu}$. It is important to note that the topological current is conserved or

$$\partial_\mu j^\mu = 0 \quad (5)$$

The normalized vector is calculated as :

$$n^a = \frac{\phi^a}{\|\phi\|}; \quad a = 1, 2 \quad \text{with} \quad \phi^1 = \phi^S, \quad \phi^2 = \phi^\Theta. \quad (6)$$

The two conditions that has to be satisfied by n^a are :

$$n^a n^a = 1 \quad \text{and} \quad n^a \partial_\nu n^a = 0. \quad (7)$$

To calculate the topological charge we use the residue method [115]. This method involves first determining the winding number (w_i) for each pole using the residue theorem and by adding the winding numbers for each pole, the topological charge can be computed. Initially, we find a solution for τ by solving the equation:

$$\frac{\partial \mathcal{F}}{\partial r_+} = 0 \quad (8)$$

Our aim is to create a two-dimensional space where we can introduce a real contour. This contour will encircle the thermodynamic imperfections described by equation 8. Thus the resulting solution, which depends on the horizon radius r_+ , is then expressed as a function of a complex variable z and denoted as $\mathcal{G}(z)$. Subsequently, a rational complex function $\mathcal{R}(z)$ is constructed as follows:

$$\mathcal{R}(z) = \frac{1}{\tau - \mathcal{G}(z)} \quad (9)$$

The complex function $R(z)$ is analytic except for a limited number of specific points $z_1, z_2, z_3, \dots, z_m$. To enclosed these points, a real closed contour C is introduced where the complex function is analytic within and on the contour. This is followed by employing the residue theorem to compute the integral of function $R(z)$ over the path C_i surrounding z_i or

$$\oint \frac{R(z)}{2\pi i} dz = \sum_{k=1}^m Res[R(z_k)] \quad (10)$$

Finally, local topology can be studied by computing the winding number at thermodynamic defects. We do that by just taking normalized residues at the poles of $\mathcal{R}(z)$

$$w_i = \frac{Res\mathcal{R}(z_i)}{|Res\mathcal{R}(z_i)|} = Sign[Res\mathcal{R}(z_i)] \quad (11)$$

Again, the Cauchy-Goursat theorem says that the integral around contour C , winding around the singular points, is equal to the integrals of $R(z)$ along C_i which z_i is enclosed. Applying this concept we can either consider a bigger contour which encompasses all thermodynamic defects or we can simply add the individual winding numbers w_i , computed around small individual contours. Hence the total winding number or topological charge, W , is then computed as

$$W = \sum_i w_i$$

In this study, we investigate the thermodynamic topology of Kerr-Sen-Ads black hole in restricted phase space. We consider the black hole solution as topological defects in the thermodynamic phase space. To study the local and the global topology of the black hole, we calculate the winding number and the total topological charge at the thermodynamic defects. In our analysis, we have considered five ensemble of Kerr-Sen-Ads black hole in restricted phase space. These are: fixed (Q, J, C) , fixed (ϕ, J, C) , fixed (Q, Ω, C) , fixed (Q, J, μ) and fixed (ϕ, Ω, C) where Q is the electric charge, J is the angular momentum, C is the central charge, ϕ is the electric potential conjugate to charge, Ω is the angular frequency conjugate to J and finally μ is the chemical potential. The selection of an ensemble frequently emerges as a decisive element influencing the thermodynamic characteristics and phase transitions exhibited by black holes[131–136]. For all these ensembles we study their thermodynamic topologies by computing the topological charge.

This paper is organized into the following sections: in Section **II**, we briefly review the Kerr-Sen-Ads black hole in restricted phase space. We analyse the thermodynamic topology of fixed (Q, J, C) ensemble in subsection **II.I.**, fixed (ϕ, J, C) in subsection **II.II.**, fixed (Q, Ω, C) in subsection **II.III.**, fixed (Q, J, μ) in subsection **II.IV.** and finally in subsection **II.V**, we analyze the same for fixed (ϕ, Ω, C) ensemble. The conclusions are presented in Section **III**.

II. KERR-SEN ADS BLACK HOLE IN RESTRICTED PHASE SPACE

The Kerr-Sen Ads black hole is the charged rotating black hole solution to the low-energy limit of heterotic string theory discovered by Sen [137]. The Kerr-Sen black holes in four dimensions have been examined from various perspectives[59, 138–146]. The thermodynamics of phase space in the extended phase space was studied in [139] and the thermodynamic of Kerr-Sen Ads black hole was studied in [59]. In this section a brief review of the Kerr-Sen-Ads black hole in restricted phase space is provided[59].

The metric for Kerr-Sen-Ads black hole in Boyer-Lindquist coordinate is given by :

$$ds^2 = -\frac{\Delta_r}{\rho^2} \left[dt - \frac{a \sin^2 \theta d\varphi}{\Xi} \right]^2 + \frac{\rho^2}{\Delta_r} dr^2 + \frac{\rho^2}{\Delta_\theta} d\theta^2 + \frac{\Delta_\theta \sin^2 \theta}{\rho^2} \left[a dt - \frac{r^2 + a^2}{\Xi} d\varphi \right]^2 \quad (12)$$

where

$$\Delta_r = (r^2 + 2br + a^2) \left(1 + \frac{r^2 + 2br}{l^2} \right) - 2Gmr$$

$$\Xi = 1 - \frac{a^2}{l^2} \quad , \quad \rho^2 = r^2 + 2br + a^2 \cos^2 \theta,$$

$$\Delta_\theta = 1 - \frac{a^2}{l^2} \cos^2 \theta$$

here b is the diatonic charge which can be express in terms of black hole mass m and electric charge q as $b = q^2/2m$. The expressions for ADM mass M and the angular momentum J and charge Q are :

$$M = \frac{m}{\Xi^2} \quad , \quad J = \frac{am}{\Xi^2} \quad \text{and} \quad Q = \frac{q}{\Xi} \quad (13)$$

using the condition $\Delta_r(r = r_+) = 0$ we can write M and J as

$$M = \frac{1}{2\Xi^2 Gr_+} (r^2 + 2br + a^2) \left(1 + \frac{r^2 + 2br}{l^2} \right) \quad (14)$$

and

$$J = \frac{a}{2\Xi^2 Gr_+} (r^2 + 2br + a^2) \left(1 + \frac{r^2 + 2br}{l^2} \right) \quad (15)$$

where r_+ is the event horizon radius and l is the Ads radius. To introduce RPST formalism two new quantities are defined which is the central charge C and its conjugate parameter μ [51, 52, 58].

$$C = \frac{l^2}{G}, \quad \mu = \frac{M - TS + \Omega J + \hat{\phi} \hat{Q}}{C} \quad (16)$$

Here \hat{Q} and $\hat{\phi}$ is the rescaled charge and electric potential expressed as:

$$\hat{Q} = \frac{Ql}{\sqrt{G}}, \quad \hat{\phi} = \frac{\phi\sqrt{G}}{l} \quad (17)$$

Using these defined quantities we write the two most important equations in RPST formalism. First one is the modified first law of black hole thermodynamic which is incorporated with a new term μdC and made free from the usual $\mathcal{P}d\mathcal{V}$ term :

$$dM = TdS + \Omega dJ + \phi dQ + \mu dC \quad (18)$$

and followed by the Euler-like relation :

$$M = TS + \Omega J + \hat{\phi} \hat{Q} + \mu C \quad (19)$$

Now, rewritting a, G as follows:

$$a = \frac{J}{M} \quad G = \frac{l^2}{C} \quad (20)$$

we can expressed the horizon radius r_+ as [59]:

$$r_+ = \frac{1}{2M} \left(\sqrt{\frac{-4J^2(\pi C + S) + \pi C Q^4 + 4l^2 M^2 S}{\pi C}} - Q^2 \right) \quad (21)$$

Putting equation.20 and equation.21 into equation.13, the ADM mass is obtained as [59] :

$$M = \frac{\sqrt{\pi C + S} \sqrt{\pi C (4\pi^2 J^2 + S^2) + 2\pi^2 Q^2 S + S^3}}{2\pi^{3/2} l \sqrt{CS}} \quad (22)$$

rest of the thermodynamic quantities can be computed using the first law of thermodynamics. For details of the RPST formlism of Kerr-Sen-Ads black hole see [59]. Now we study the thermodynamic topology in five different ensemble in the following subsections.

II.1. Fixed (Q,J,C) Ensemble

For Kerr sen Ads black hole,mass M in this ensemble is :

$$M = \frac{\sqrt{\pi C + S} \sqrt{\pi C (4\pi^2 J^2 + S^2) + 2\pi^2 Q^2 S + S^3}}{2\pi^{3/2} l \sqrt{CS}} \quad (23)$$

Using equation 23, free energy is computed as :

$$\mathcal{F} = M - S/\tau = -\frac{2\pi^{3/2}lS\sqrt{CS} - \tau\sqrt{\pi C + S}\sqrt{4\pi^3CJ^2 + \pi CS^2 + 2\pi^2Q^2S + S^3}}{2\pi^{3/2}l\tau\sqrt{CS}} \quad (24)$$

The components of the vector field ϕ is obtained as :

$$\phi^r = -\frac{4\pi^4C^3J^2\tau - 4\pi C^2S^3\tau - \pi^2C^3S^2\tau + 4\pi^{3/2}l\sqrt{\pi C + S}(CS)^{3/2}\sqrt{4\pi^3CJ^2 + \pi CS^2 + 2\pi^2Q^2S + S^3} - 2\pi^2CQ^2S^2\tau - 3CS^4\tau}{4\pi^{3/2}l\tau(CS)^{3/2}\sqrt{\pi C + S}\sqrt{4\pi^3CJ^2 + \pi CS^2 + 2\pi^2Q^2S + S^3}} \quad (25)$$

and

$$\phi^\Theta = -\cot \Theta \csc \Theta \quad (26)$$

The zero points of ϕ^r is also obtained as:

$$\tau = \frac{4\pi^{3/2}l(CS)^{3/2}\sqrt{\pi C + S}\sqrt{4\pi^3CJ^2 + \pi CS^2 + 2\pi^2Q^2S + S^3}}{-4\pi^4C^3J^2 + 4\pi C^2S^3 + \pi^2C^3S^2 + 2\pi^2CQ^2S^2 + 3CS^4} \quad (27)$$

The entropy is plotted against τ at $C = 0.2, Q = 1.5, J = 0.02$ and $l = 1$ in Figure.1. Here we observe three black hole branches : a small, an intermediate, and a large black hole branch. We also see a generation point at $\tau = 1.84076, S = 2.02107$ and an annihilation point at $\tau = 1.27059, S = 0.07658$ which are shown as blue dot and black dot respectively.




FIG. 1: τ vs S plot for Kerr Sen Ads black hole at $C = 0.2, Q = 1.5, J = 0.02$ and $l = 1$. The *LBH* and *SBH* regions represent the large and small black hole branches respectively.

We now calculate the topological charge of the black hole using the residue method. The complex function $\mathcal{R}(z)$ mentioned in section I is given by :

$$\mathcal{R}(z) = -\frac{-4\pi^4C^2J^2 + \pi^2C^2z^2 + 4\pi Cz^3 + 2\pi^2Q^2z^2 + 3z^4}{4\pi^4C^2J^2\tau - \pi^2C^2\tau z^2 + 4\pi^{3/2}l z\sqrt{Cz}\sqrt{\pi C + z}\sqrt{4\pi^3CJ^2 + \pi Cz^2 + 2\pi^2Q^2z + z^3} - 4\pi C\tau z^3 - 2\pi^2Q^2\tau z^2 - 3\tau z^4} \quad (28)$$

The denominator of the complex function is considered as a polynomial function $\mathcal{A}(z)$

$$\mathcal{A}(z) = 4\pi^4C^2J^2\tau - \pi^2C^2\tau z^2 + 4\pi^{3/2}l z\sqrt{Cz}\sqrt{\pi C + z}\sqrt{4\pi^3CJ^2 + \pi Cz^2 + 2\pi^2Q^2z + z^3} - 4\pi C\tau z^3 - 2\pi^2Q^2\tau z^2 - 3\tau z^4 \quad (29)$$

To calculate the topological charge, we put $C = 0.2, Q = 1.5, J = 0.02$ and $l = 1$ in equation 29.




FIG. 2: Plot for polynomial function $\mathcal{A}(z)$ for $\tau = 2.2, 2.4$ and 2.6

Figure 2 represents the plot for $\mathcal{A}(z)$. This demonstrates that there is one pole for $\tau < 1.27059$, three poles for $1.27059 < \tau < 1.84075$, and one pole for $\tau > 1.84075$. Using $\tau = 1.2$ for the large black hole branch, the pole is located at $z = 8.16619$ as suggested by the blue solid line in figure 2. The winding number is $w = +1$ around the pole $z = 8.16619$. Using $\tau = 1.6$, the poles for the intermediate black hole branch are located at $z_1 = 0.0235863$, $z_2 = 0.676384$, and $z_3 = 4.39265$, which is represented by the solid green line. The winding numbers around these three poles are $w_1 = +1$, $w_2 = -1$, and $w_3 = +1$, respectively, based on the sign of the residue around them. Thus,

$$W = w_1 + w_2 + w_3 = 1 - 1 + 1 = 1$$

is the topological charge. Similarly, as represented by the solid red line, for a small black hole branch, taking $\tau = 2$, the pole is located at $z = 0.0186124$, and the winding number is determined to be $+1$ based on the sign of residue surrounding the pole.

We now analyze the effect of thermodynamic parameter on the topological charge. We start with analysing the effect of Ads radius l keeping $C = 0.2, Q = 1.5, J = 0.02$ constant. We observe no effect of change in l on the topological charge. The same is illustrated in figure 3. Here we have considered two values of l keeping $C = 0.2, Q = 1.5, J = 0.02$ constant. In figure 13a, $l = 0.1$ and in figure 13b, $l = 100$. For both cases, we find the number of black hole branches equal to three and topological charge to be one.

Our above analysis is followed by studying the dependence of topological charge on the central charge C keeping $l = 1, Q = 1.5, J = 0.02$ fixed. It is seen that topological charge does not depend on variation in central charge C . Although for a specific range of C the three phases of the black hole merges to a single branch. However it does not alter the topological number. In figure 4a three black hole branches and topological charge 1 is observed. In 4b, we have only a single black hole branches with a topological charge of 1. In figure 4a the number of branch is again three and topological charge is 1.

Next, we want to vary q while holding $l = 1, C = 1, J = 0.02$ constant in order to see how the change in charge affects the topological charge. In Figures 5a, 5b, and 5c, respectively, we have set $Q = 0.1, Q = 1.5$, and $Q = 6$. Three black hole branches are found in figure 5a, one black hole branch is found in picture 5b, and three black hole branches are found once again in figure 5c. In each of the three scenarios, the topological charge is 1. Therefore, it is noted that charge likewise has no effect on the topological number.

Lastly, we perform the same study in figures 6a, 6b, and 6c to examine how J affects topological charge while maintaining constant l, Q, C at $l = 1, Q = 1.5, C = 0.2$. J is equal to 0.02 in figure 6a, 0.1 in figure 6b, and 1 in figure 6c. Topological charge in each of the three scenarios is 1. Thus, it is concluded that while the number of black hole branches may change in response to changes in angular momentum J , the topological charge remains constant at $W = 1$.

In conclusion, our findings show that the Kerr-Sen-Ads black hole in a fixed (Q, J, C) ensemble, the topological charge is 1. The topological charge is independent of the thermodynamic parameters Q, J, C .




FIG. 3: τ vs S plots for Kerr-Sen-AdS black hole in restricted phase space in fixed (q, J, C) ensemble. Here τ is plotted against S for two different l values while keeping $C = 0.2, Q = 1.5, J = 0.02$ fixed. Figure (a) shows τ vs S plot at $l = 0.1$ and in figure (b) shows τ vs S plot at $l = 100$. W denotes topological charge.




FIG. 4: τ vs S plots for Kerr-Sen-AdS black hole in restricted phase space in fixed (Q, J, C) ensemble. Here τ is plotted against S for three different C values while keeping $l = 1, Q = 1.5, J = 0.02$ fixed. Figure (a) shows τ vs S plot at $C = 0.5$, figure (b) shows τ vs S plot at $C = 1$ and in figure (c) shows τ vs S plot at $C = 10$. W denotes topological charge.

II.2. Fixed (ϕ, J, C) Ensemble

The potential ϕ , angular momentum J and central charge C is kept fixed in fixed (ϕ, J, C) ensemble. The potential ϕ is given by :

$$\phi = \frac{\partial M}{\partial q} = \frac{\sqrt{\pi} Q S \sqrt{\pi C + S}}{l \sqrt{C S} \sqrt{\pi C (4\pi^2 J^2 + S^2) + 2\pi^2 Q^2 S + S^3}} \quad (30)$$

To get an expression for q we solve equation 30 and get the expression as :

$$Q = \frac{\sqrt{\pi} \sqrt{C} l \phi \sqrt{4\pi^3 C J^2 + \pi C S^2 + S^3}}{\sqrt{S} \sqrt{-2\pi^2 C l^2 \phi^2 + \pi C + S}} \quad (31)$$




FIG. 5: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (q, J, C) ensemble. Here τ is plotted against S for three different Q values while keeping $l = 1, C = 1, J = 0.02$ fixed. Figure (a) shows τ vs S plot at $Q = 0.1$, figure (b) shows τ vs S plot at $Q = 1.5$ and in figure (c) shows τ vs S plot at $Q = 6$. W denotes topological charge.




FIG. 6: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (q, J, C) ensemble. Here τ is plotted against S for three different J values while keeping $l = 1, C = 0.2, Q = 1.5$ fixed. Figure (a) shows τ vs S plot at $J = 0.02$, figure (b) shows τ vs S plot at $J = 0.1$ and in figure (c) shows τ vs S plot at $J = 1$. W denotes topological charge.

Now the new mass (M_ϕ) in this ensemble is calculated using equation 31 as follows :

$$\begin{aligned}
 M_\phi &= M - q\phi \\
 &= \frac{\sqrt{\pi C + S} \sqrt{\frac{2\pi^3 C l^2 \phi^2 (4\pi^3 C J^2 + \pi C S^2 + S^3)}{-2\pi^2 C l^2 \phi^2 + \pi C + S}} + \pi C (4\pi^2 J^2 + S^2) + S^3}{2\pi^{3/2} l \sqrt{C S}} - \frac{\sqrt{\pi} \sqrt{C} l \phi^2 \sqrt{4\pi^3 C J^2 + \pi C S^2 + S^3}}{\sqrt{S} \sqrt{-2\pi^2 C l^2 \phi^2 + \pi C + S}} \quad (32)
 \end{aligned}$$

The off-shell free energy is computed using:

$$\mathcal{F} = M_\phi - S/\tau$$

We compute the expressions for ϕ^S and τ using the same method as in the preceding part.

In Figure.7, the entropy S vs τ is plotted at $C = 1, \phi = 1.5, J = 0.02$, and $l = 0.1$. Here also three black hole branches are visible. We see two points that are : a generation point at $\tau = 0.35285, S = 0.9876$ and an annihilation point at $\tau = 0.342307, S = 0.4037$, shown as blue and black dots, respectively

To analyze the effect of thermodynamic parameter on the topological charge we vary one particular parameter




FIG. 7: τ vs S plot for Kerr Sen Ads black hole at $C = 1, \phi = 1.5, J = 0.02$ and $l = 0.1$. The *LBH* and *SBH* regions represent the large and small black hole branches respectively.

keeping other parameter fixed. For this ensemble we observe no effect of change in any thermodynamic parameter on the topological charge. The same is illustrated in figure 8,9,10 and 11.

In figure 8, central charge C is varied keeping $l = 0.1, \phi = 1.5, J = 0.02$ fixed. For a specific range of C the three phases of the black hole is visible and it merges to a single branch for other values, through out the process the topological charge remains constant at 1. For $C = 10$, in figure 8a three black hole branches and topological charge 1 is observed. In 4b, we have only a single black hole branches with a topological charge of 1 when $C = 0.1$.




FIG. 8: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (ϕ, J, C) ensemble. Here τ is plotted against S for two different C values while keeping $l = 0.1, \phi = 1.5, J = 0.02$ fixed. Figure (a) shows τ vs S plot at $C = 10$, figure (b) shows τ vs S plot at $C = 0.1$. W denotes topological charge.

In figure 9, effect of variation of ϕ topological charge is shown while keeping $l = 0.1, C = 1, J = 0.02$ constant. In Figures 9a, 9b and 9c respectively, we have set $\phi = 0.5, \phi = 1$, and $\phi = 5$. Three black hole branches are found in figure 9b on the other hand only one black hole branch is found in figure 9a and figure 9c. In each of the three scenarios, the topological charge is 1. Therefore, it is noted that potential ϕ has no effect on the topological number.




FIG. 9: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (ϕ, J, C) ensemble. Here τ is plotted against S for three different ϕ values while keeping $l = 0.1, C = 1, J = 0.02$ fixed. Figure (a) shows τ vs S plot at $\phi = 0.5$, figure (b) shows τ vs S plot at $\phi = 1$ and in figure (c) shows τ vs S plot at $\phi = 5$. W denotes topological charge.

Lastly, in figures 10a, 10b, and 10c τ vs S is plotted to check how J affects topological charge while keeping constant l, ϕ, C at $l = 1, \phi = 1.5, C = 1$. J is equal to 0.02 in figure 10a, 0.1 in figure 10b, and 1 in figure 10c. In all three cases, topological charge is 1. Consequently, it is inferred that the topological charge at $W = 1$ remains constant, however the number of black hole branches may vary in response to variations in angular momentum J .




FIG. 10: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (ϕ, J, C) ensemble. Here τ is plotted against S for three different J values while keeping $l = 1, C = 0.2, \phi = 1.5$ fixed. Figure (a) shows τ vs S plot at $J = 0.02$, figure (b) shows τ vs S plot at $J = 0.1$ and in figure (c) shows τ vs S plot at $J = 1$. W denotes topological charge.

In conclusion, our findings show that the Kerr-Sen-Ads black hole in a fixed (Q, J, C) ensemble, the topological charge is 1. The topological charge is independent of the thermodynamic parameters Q, J, C .

II.3. Fixed (Q,Ω,C) Ensemble

In this ensemble, the angular frequency Ω , charge Q and central charge C is kept fixed. The angular frequency Ω is given by :

$$\Omega = \frac{2\pi^{3/2}CJ\sqrt{\pi C + S}}{l\sqrt{CS}\sqrt{\pi C(4\pi^2J^2 + S^2) + 2\pi^2Q^2S + S^3}} \quad (33)$$

By solving equation 33, we get the expression for J as :

$$J = \frac{lS\Omega\sqrt{\pi CS + 2\pi^2Q^2 + S^2}}{\pi^{3/2}\sqrt{C}\sqrt{4\pi C - 4l^2S\Omega^2 + 4S}} \quad (34)$$

The newly mass(M_Ω) in this ensemble :

$$M_\Omega = M - J\Omega$$

$$= -\frac{l^2S\Omega^2\sqrt{CS}\sqrt{\pi CS + 2\pi^2Q^2 + S^2} - \sqrt{C}\sqrt{\pi C + S}\sqrt{\pi C - l^2S\Omega^2} + S\sqrt{\frac{S(\pi C + S)(\pi CS + 2\pi^2Q^2 + S^2)}{\pi C - l^2S\Omega^2 + S}}}{2\pi^{3/2}\sqrt{C}l\sqrt{CS}\sqrt{\pi C - l^2S\Omega^2} + S} \quad (35)$$

The off-shell free energy is computed using:

$$\mathcal{F} = M_\Omega - S/\tau$$

We compute the expressions for ϕ^S and τ using the same method as in the preceding part.

In Figure.11, the entropy S vs τ is plotted at $C = 10, Q = 1.5, \Omega = 0.02$, and $l = 0.1$. We observe two black hole branches with a generation point at $(\tau, S) = (0.37635, 8.2427)$ represented by the black dot. We observe a large black hole branch shown as red solid line and a small black hole branch shown as black solid line. The topological number of the smaller black hole branch is -1 and that of the large black hole branch is $+1$. The total topological number is found to be $-1 + 1 = 0$.

Taking a cue from figure.11, study of Hawking-Page transition and Davies type of phase transition also can be studied




FIG. 11: τ vs S plot for Kerr Sen Ads black hole at $C = 1, \phi = 1.5, J = 0.02$ and $l = 0.1$. The red and the black colored solid represent the large and small black hole branches respectively.

by constructing a noval vector field given by [123]

$$\phi = \left(\frac{1}{S} \frac{\partial F^2}{\partial S}, -\cot\theta csc\theta \right) \quad (36)$$

Here F is the on-shell free energy with the condition $\tau = T = \frac{\partial M}{\partial S}$. From the vectorplot shown in figure.12 By solving




FIG. 12: A portion of the vector field ϕ on $S - \theta$ plane. The black dot of figure (a) represents Hawking page phase transition point and the blue dot in figure (b) represents Davies point.

$F = 0$ and $C_\Omega = T \frac{\partial S}{\partial T} = 0$, the Hawking Page phase transition point is found to be $S = 33.3908$ and the Davies point if found to be $S = 8.2427$ which exactly matches with the critical points of the vector field in equation 45. The black dot ($S = 8.2427$) in figure 12a is a Hawking page transition point and blue dot ($S = 33.3908$) is a Davies point.

The topological charge can be calculated by constructing a contour C around each zero point (C_0 and C_1 as shown in figure 12) which is parametrized as:

$$\begin{cases} S = a \cos \nu + r_0 \\ \theta = b \sin \nu + \frac{\pi}{2} \end{cases} \quad (37)$$

where $\nu \in (0, 2\pi)$. To calculate the deflection Ω of the vector field n along the contour C_1 and C_2 , we choose $a = b = 0.2$ and $r_0 =$ the zero point itself enclosed by the contours. We use the formula [77, 78]:

$$\Omega(\nu) = \int_0^\nu \epsilon_{12} n^1 \partial_\nu n^2 d\nu \quad (38)$$

where the unit vectors (n^1, n^2) are given by:

$$n^1 = \frac{\phi^S}{\sqrt{(\phi^S)^2 + (\phi^\Theta)^2}} \quad \text{and} \quad n^2 = \frac{\phi^\Theta}{\sqrt{(\phi^S)^2 + (\phi^\Theta)^2}}$$

Finally, the winding numbers w and topological charge W can be calculated as follows:

$$\begin{cases} w = \frac{1}{2\pi} \Omega(2\pi) \\ W = \sum_i w_i \end{cases} \quad (39)$$

In cases where the parameter region does not encompass any zero points, the topological charge equals to 0. Following the above procedure, we plot deflection Ω vs θ plot in figure.13. From the figure, the topological charge of the Hawking page phase transition (blue colored solid line) is +1 and that of Davis point (black colored solid line) is found to be -1.

To analyze the effect of thermodynamic parameter on the topological charge we vary one particular parameter keeping other parameter fixed. In figure 14, the value of central charge C is varied keeping $l = 0.1, Q = 1.5, \Omega = 0.02$ fixed. For a different value of C the topological charge is found to be 0 or 1. For $C = 0.01$, in figure 14a there are two black hole branches and topological charge is 0. In 14b, we have only a single black hole branch with a topological charge of 1 when $C = 1$. In 14c again we observe two black hole branch with a topological charge of 0 when $C = 15$.




FIG. 13: Kerr Sen AdS black hole in fixed (ϕ, J, C) ensemble : Figure (a) represents topological number calculation for Davies point around the C_0 contour in Figure.2 and Figure (b) represents the topological number calculation for Hawking-Page phase transition point around the C_1 contour




FIG. 14: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (Q, Ω, C) ensemble. Here τ is plotted against S for two different C values while keeping $l = 0.1, Q = 1.5, \Omega = 0.02$ fixed. Figure (a) shows τ vs S plot at $C = 0.01$, figure (b) shows τ vs S plot at $C = 0.1$, figure (c) shows the same at $C = 15$ W denotes topological charge.

In figure 15, effect of variation of Q on topological charge is shown while keeping $l = 0.1, C = 1, J = 0.02$ constant. In Figures 15a, 15b and 15c respectively, we have set $Q = 0.05, Q = 0.5$, and $Q = 3.5$. For $Q = 0.05$ and $Q = 3.5$, two black hole branches are found in figure 15a and figure.15c with topological charge 0. On the other hand only one black hole branch is found with topological charge 1 in figure 15b when $Q = 0.5$ is kept fixed.

Lastly, in figures 16a, 16b, and 16c τ vs S is plotted to check how Ω affects topological charge while keeping constant l, Q, C at $l = 1, Q = 0.05, C = 1$. Ω is equal to 0.01 in figure 16a, 5 in figure 16b, and 12 in figure 16c. For $\Omega = 5$ and $\Omega = 0.05$, the topological charge is 0 and for $\Omega = 12$, the topological charge is found to be -1 . In conclusion, our findings show that the Kerr-Sen-Ads black hole in a fixed (Q, Ω, C) ensemble, the topological charge is $-1, 0$ or 1 depending upon the values of the thermodynamic parameters Q, Ω, C .

II.4. Fixed (Q, J, μ) Ensemble

In this ensemble, the chemical potential μ , charge Q and angular momentum J is kept fixed. The chemical potential μ is given by :

$$\mu = \frac{S (\pi^2 C^2 (4\pi^2 J^2 + S^2) - S^2 (2\pi^2 Q^2 + S^2))}{4\pi^{3/2} l (CS)^{3/2} \sqrt{\pi C + S} \sqrt{\pi C (4\pi^2 J^2 + S^2) + 2\pi^2 Q^2 S + S^3}} \quad (40)$$




FIG. 15: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (ϕ, J, C) ensemble. Here τ is plotted against S for three different ϕ values while keeping $l = 0.1, C = 1, J = 0.02$ fixed. Figure (a) shows τ vs S plot at $\phi = 0.5$, figure (b) shows τ vs S plot at $\phi = 1$ and in figure (c) shows τ vs S plot at $\phi = 5$. W denotes topological charge.




FIG. 16: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (Q, Ω, C) ensemble. Here τ is plotted against S for three different Ω values while keeping $l = 1, C = 0.2, Q = 0.05$ fixed. Figure (a) shows τ vs S plot at $\Omega = 0.05$, figure (b) shows τ vs S plot at $\Omega = 5$ and in figure (c) shows τ vs S plot at $\Omega = 12$. W denotes topological charge.

By solving equation 40, we get the expression for C and substituting C in the following expression, newly mass (M_μ) in this ensemble as:

$$M_\mu = M - C\mu \quad (41)$$

The off-shell free energy is computed using:

$$\mathcal{F} = M_\mu - S/\tau$$

We compute the expressions for ϕ^S and τ using the same method as in the preceding part.

It is seen that, for this ensemble the topological charge is always 1 irrespective of the values of the thermodynamic parameters μ, Q and J . Moreover, only one branch is observed in the entropy S vs τ plot for all allowed values of the thermodynamic parameters. For example, in figure.17, the entropy S vs τ is plotted for $\mu = 1, Q = 1.5, J = 1.2$, and $l = 1$. We observe a single black hole branch with topological charge equals to 1.




FIG. 17: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (Q, J, μ) ensemble. Here τ is plotted against S while keeping $l = 1, \mu = 1, Q = 1.5$ and $J = 1.2$ fixed. The topological charge is found to be 1.

II.5. Fixed (ϕ, Ω, C) ensemble

The next ensemble we consider is the fixed (Ω, ϕ, C) ensemble. In this ensemble angular frequency Ω and electric potential ϕ are kept fixed. First, We substitute J from equation 33 in the expression for mass in equation 23 as follows:

$$M_J = \frac{\sqrt{S(\pi C + S)^2 (\pi C S + 2\pi^2 Q^2 + S^2)}}{2\pi^{3/2} l \sqrt{C S (\pi C - l^2 S \Omega^2 + S)}} \quad (42)$$

Now equation 42 becomes independent of variable J . Next we find out ϕ from equation 42 as follows:

$$\phi = \frac{\partial M_J}{\partial Q} = \frac{\sqrt{\pi Q} \sqrt{S(\pi C + S)^2 (\pi C S + 2\pi^2 Q^2 + S^2)}}{l \sqrt{C S (\pi C S + 2\pi^2 Q^2 + S^2) (\pi C - l^2 S \Omega^2 + S)}} \quad (43)$$

Accordingly, Q is given by equation 43 as:

$$q = \frac{\sqrt{C l} \sqrt{S} \phi \sqrt{\pi C + S} \sqrt{\pi C - l^2 S \Omega^2 + S}}{\sqrt{\pi} \sqrt{-\pi^2 C^2 (2l^2 \phi^2 - 1) + 2\pi C S (l^4 \Omega^2 \phi^2 - l^2 \phi^2 + 1) + S^2}}$$

Finally, we substitute Q in the following expression for the modified mass in fixed (Ω, Φ) ensemble, which is written as :

$$\tilde{M} = M - q\phi - J\Omega \quad (44)$$

Now \tilde{M} is a function of ϕ, Ω, C and l . Using equation 44, \mathcal{F} , ϕ^S and τ are constructed following standard procedure as shown in the previous sections.

We plot τ vs S curve for different values of C as shown in figure 18 keeping $\Omega = 1.2, \phi = 3.5, l = 0.1$ constant. In figure 18a and figure 18b, with $C = 0.05$ and $C = 0.5$ respectively, one black hole branch with topological charge $W = 1$ is observed. At $C = 10$, in figure 18c two black hole branches and topological charge 0 are found.

Next we investigate the topological charge with reference to a variation in the potential ϕ while keeping $C = 10, \Omega = 12, l = 0.1$ fixed. In figure 19a and 19b we set $\phi = 1$ and $\phi = 10$ respectively where we find one black hole branch and topological charge $W = -1$. In the latter case, where $\phi = 15$, we find two black hole branches with topological charge 0.

Finally, in figures 20, we fix $\phi = 0.01, C = 10, l = 0.1$ and vary Ω . In figure 20a and figure 20b, setting $\Omega = 5$ and $\Omega = 10$ respectively, we get two black hole branches and topological charge $W = 0$. In figure 20c, with $\Omega = 15$ a single




FIG. 18: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (ϕ, Ω, C) ensemble. Here τ is plotted against S for three different C values while keeping $l = 0.1, \phi = 3.5, \Omega = 1.2$ fixed. Figure (a) shows τ vs S plot at $C = 0.05$, figure (b) shows τ vs S plot at $C = 0.5$ and in figure (c) shows the same at $C = 10$. W denotes topological charge.




FIG. 19: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (ϕ, Ω, C) ensemble. Here τ is plotted against S for three different ϕ values while keeping $l = 0.1, C = 10, \Omega = 12$ fixed. Figure (a) shows τ vs S plot at $\phi = 1$, figure (b) shows τ vs S plot at $\phi = 10$ and in figure (c) shows the same at $\phi = 15$. W denotes topological charge.

black hole branch with topological charge $W = -1$ is observed.

So to summarize, the topological charge in fixed (ϕ, Ω, C) ensemble is $-1, 0$ or $+1$ depending on all the thermodynamic parameters C, Ω and ϕ .

Like fixed (Q, Ω, C) ensemble, here also we find a class of black hole, where topological charge is found to be zero. Here also both Hawking-Page transition and Davies type of phase transition are observed. For example, taking $C = 10, \phi = 0.01, \Omega = 5$ and $l = 0.1$ we observe two black hole branch as shown in figure.21 Using the following equation we construct the vector field

$$\phi = \left(\frac{1}{S} \frac{\partial F^2}{\partial S}, -\cot\theta \csc\theta \right) \quad (45)$$

The zero points of the vector field is found to be $(12.0763, \frac{\pi}{2})$ and $(36.276, \frac{\pi}{2})$. The vectorplot is shown in figure.22 The black dot ($S = 12.0763$) in figure 22a is a Davies transition point and blue dot ($S = 36.276$) is a Hawking-Page phase transition point. We have confirmed that, these zero points exactly matches with the critical points of the Hawking-page and Davies type phase transition. The topological charge is calculated by constructing a contour C




FIG. 20: τ vs S plots for Kerr-Sen-Ads black hole in restricted phase space in fixed (ϕ, Ω, C) ensemble. Here τ is plotted against S for three different Ω values while keeping $l = 0.1, C = 1, \phi = 0.01$ fixed. Figure (a) shows τ vs S plot at $\Omega = 5$, figure (b) shows τ vs S plot at $\Omega = 10$ and in figure (c) shows the same at $\Omega = 15$. W denotes topological charge.




FIG. 21: τ vs S plot for Kerr Sen Ads black hole at $C = 10, \phi = 0.01, \Omega = 5$ and $l = 0.1$. The red and the black colored solid represent the large and small black hole branches respectively.

around each zero point (C_0 and C_1 as shown in figure 22). As expected, the topological charge of the Hawking page phase transition is $+1$ and that of Davis point is found to be -1 as shown in figure.23




FIG. 22: A portion of the vector field ϕ on $S - \theta$ plane. The black dot of figure (a) represents Hawking page phase transition point and the blue dot in figure (b) represents Davies point.




FIG. 23: Kerr Sen AdS black hole in fixed (ϕ, Ω, C) ensemble : Figure (a) represents topological number calculation for Davies point around the C_0 contour in Figure.2 and Figure (b) represents the topological number calculation for Hawking-Page phase transition point around the C_1 contour

III. CONCLUSION

In this study, thermodynamic topology of the Kerr-Sen-Ads black hole in restricted phase space is analyzed. For our analysis, we have considered five ensembles of Kerr-Sen-Ads black holes in restricted phase space: fixed (Q, J, C) , fixed (ϕ, J, C) , fixed (Q, Ω, C) , fixed (Q, J, μ) , and fixed (ϕ, Ω, C) . We studied the local and global topology in each ensemble by computing the topological number. In the fixed (Q, J, C) , fixed (ϕ, J, C) , and fixed (Q, J, μ) ensembles, we found a topological charge of $+1$. In the fixed (Q, Ω, C) and fixed (ϕ, Ω, C) ensembles, we found topological charges of -1 , 0 , and $+1$. The results are summarized below.

Ensembles	Topological Charge	Annihilation Point	Generation Point
Fixed (Q, J, C)	1	1 or 0	1 or 0
Fixed (ϕ, J, C)	1	1 or 0	1 or 0
Fixed (Q, Ω, C)	-1, 0 or 1	0	1 or 0
Fixed (Q, J, μ)	1	0	0
Fixed (ϕ, Ω, C)	-1, 0 or 1	0	1 or 0

Notably, within ensembles where the topological charge is determined to be 0, occurrence of both Hawking-Page and Davies-type phase transitions is observed. We have shown that these distinct types of phase transitions can be comprehensively investigated employing a common vector field. Furthermore, both types of phase transitions can be distinguished by the associated topological charges. The topological charge linked with Davies-type and Hawking-Page phase transitions are found to be -1 and $+1$, respectively.

It is important to mention that three more possible ensembles exist in this system: fixed (ϕ, J, μ) , fixed (Q, Ω, μ) , and fixed (ϕ, Ω, μ) . Mathematically, we found it very challenging to conduct our analysis in these ensembles. We will endeavor to address this issue in our future work. It will be interesting to study the thermodynamic topology of various black hole systems in more challenging ensembles, which exist in other phase space formalisms or different modified theories of gravity.

IV. ACKNOWLEDGMENTS

BH would like to thank DST-INSPIRE, Ministry of Science and Technology fellowship program, Govt. of India for awarding the DST/INSPIRE Fellowship[IF220255] for financial support.

-
- [1] S. W. Wei and Y. X. Liu, Topology of black hole thermodynamics, Phys. Rev. D **105**, no.10, 104003 (2022) doi:10.1103/PhysRevD.105.104003 [arXiv:2112.01706 [gr-qc]].
 - [2] S. W. Wei, Y. X. Liu and R. B. Mann, Black Hole Solutions as Topological Thermodynamic Defects, Phys. Rev. Lett. **129**, no.19, 191101 (2022) doi:10.1103/PhysRevLett.129.191101 [arXiv:2208.01932 [gr-qc]].
 - [3] P. K. Yerra, C. Bhamidipati and S. Mukherji, Phys. Rev. D **106**, no.6, 064059 (2022) doi:10.1103/PhysRevD.106.064059 [arXiv:2208.06388 [hep-th]].
 - [4] K. Bhattacharya, K. Bamba and D. Singleton, Topological interpretation of extremal and Davies-type phase transitions of black holes, [arXiv:2402.18791 [gr-qc]].
 - [5] J. D. Bekenstein, Black holes and entropy, Phys. Rev. D **7**, 2333-2346 (1973) doi:10.1103/PhysRevD.7.2333
 - [6] S. W. Hawking, Black hole explosions, Nature **248**, 30-31 (1974) doi:10.1038/248030a0
 - [7] S. W. Hawking, Particle Creation by Black Holes, Commun. Math. Phys. **43**, 199-220 (1975) [erratum: Commun. Math. Phys. **46**, 206 (1976)] doi:10.1007/BF02345020
 - [8] J. M. Bardeen, B. Carter and S. W. Hawking, The Four laws of black hole mechanics, Commun. Math. Phys. **31**, 161-170 (1973) doi:10.1007/BF01645742
 - [9] R. M. Wald, Entropy and black-hole thermodynamics, Phys. Rev. D **20**, 1271-1282 (1979) doi:10.1103/PhysRevD.20.1271
 - [10] Jacob D Bekenstein. Black-hole thermodynamics, Physics Today, 33(1):24–31, 1980.
 - [11] R. M. Wald, The thermodynamics of black holes, Living Rev. Rel. **4**, 6 (2001) doi:10.12942/lrr-2001-6 [arXiv:gr-qc/9912119 [gr-qc]].
 - [12] S. Carlip, Black Hole Thermodynamics, Int. J. Mod. Phys. D **23**, 1430023 (2014) doi:10.1142/S0218271814300237 [arXiv:1410.1486 [gr-qc]].
 - [13] A.C.Wall, A Survey of Black Hole Thermodynamics, [arXiv:1804.10610 [gr-qc]].

- [14] P.Candelas and D.W.Sciama, Irreversible Thermodynamics of Black Holes, *Phys. Rev. Lett.* **38**, 1372-1375 (1977) doi:10.1103/PhysRevLett.38.1372
- [15] A. Chamblin, R. Emparan, C. V. Johnson and R. C. Myers, Holography, thermodynamics and fluctuations of charged AdS black holes, *Phys. Rev. D* **60**, 104026 (1999) doi:10.1103/PhysRevD.60.104026 [arXiv:hep-th/9904197 [hep-th]].
- [16] A. Chamblin, R. Emparan, C. V. Johnson and R. C. Myers, Charged AdS black holes and catastrophic holography, *Phys. Rev. D* **60**, 064018 (1999) doi:10.1103/PhysRevD.60.064018 [arXiv:hep-th/9902170 [hep-th]].
- [17] S. S. Gubser, I. R. Klebanov and A. M. Polyakov, Gauge theory correlators from noncritical string theory, *Phys. Lett. B* **428**, 105-114 (1998) doi:10.1016/S0370-2693(98)00377-3 [arXiv:hep-th/9802109 [hep-th]].
- [18] P. C. W. Davies, Thermodynamic Phase Transitions of Kerr-Newman Black Holes in De Sitter Space, *Class. Quant. Grav.* **6**, 1909 (1989) doi:10.1088/0264-9381/6/12/018
- [19] S. W. Hawking and D. N. Page, Thermodynamics of Black Holes in anti-De Sitter Space, *Commun. Math. Phys.* **87**, 577 (1983) doi:10.1007/BF01208266
- [20] A. Curir, Rotating black holes as dissipative spin-thermodynamical systems, *General Relativity and Gravitation*, **13**, 417, (1981) doi:10.1007/BF00756588
- [21] Anna Curir, Black hole emissions and phase transitions, *General Relativity and Gravitation*, **13**, 1177, (1981) doi:10.1007/BF00759866
- [22] D. Pavon and J. M. Rubi, Nonequilibrium Thermodynamic Fluctuations of Black Holes, *Phys. Rev. D* **37**, 2052-2058 (1988) doi:10.1103/PhysRevD.37.2052
- [23] D. Pavon, Phase transition in Reissner-Nordstrom black holes, *Phys. Rev. D* **43**, 2495-2497 (1991) doi:10.1103/PhysRevD.43.2495
- [24] O. Kaburaki, Critical behavior of extremal Kerr-Newman black holes, *Gen. Rel. Grav.* **28**, 843 (1996)
- [25] R. G. Cai, Z. J. Lu and Y. Z. Zhang, Critical behavior in (2+1)-dimensional black holes, *Phys. Rev. D* **55**, 853-860 (1997) doi:10.1103/PhysRevD.55.853 [arXiv:gr-qc/9702032 [gr-qc]].
- [26] R. G. Cai and J. H. Cho, Thermodynamic curvature of the BTZ black hole, *Phys. Rev. D* **60**, 067502 (1999) doi:10.1103/PhysRevD.60.067502 [arXiv:hep-th/9803261 [hep-th]].
- [27] Y. H. Wei, Thermodynamic critical and geometrical properties of charged BTZ black hole, *Phys. Rev. D* **80**, 024029 (2009) doi:10.1103/PhysRevD.80.024029
- [28] K. Bhattacharya, S. Dey, B. R. Majhi and S. Samanta, General framework to study the extremal phase transition of black holes, *Phys. Rev. D* **99**, no.12, 124047 (2019) doi:10.1103/PhysRevD.99.124047 [arXiv:1903.03434 [gr-qc]].
- [29] D. Kastor, S. Ray and J. Traschen, Enthalpy and the Mechanics of AdS Black Holes, *Class. Quant. Grav.* **26**, 195011 (2009) doi:10.1088/0264-9381/26/19/195011 [arXiv:0904.2765 [hep-th]].
- [30] B. P. Dolan, The cosmological constant and the black hole equation of state, *Class. Quant. Grav.* **28**, 125020 (2011) doi:10.1088/0264-9381/28/12/125020 [arXiv:1008.5023 [gr-qc]].
- [31] B. P. Dolan, Pressure and volume in the first law of black hole thermodynamics, *Class. Quant. Grav.* **28**, 235017 (2011) doi:10.1088/0264-9381/28/23/235017 [arXiv:1106.6260 [gr-qc]].
- [32] B. P. Dolan, Compressibility of rotating black holes, *Phys. Rev. D* **84**, 127503 (2011) doi:10.1103/PhysRevD.84.127503 [arXiv:1109.0198 [gr-qc]].
- [33] B. P. Dolan, Where Is the PdV in the First Law of Black Hole Thermodynamics?, doi:10.5772/52455 [arXiv:1209.1272 [gr-qc]].
- [34] D. Kubiznak and R. B. Mann, P-V criticality of charged AdS black holes, *JHEP* **07**, 033 (2012) doi:10.1007/JHEP07(2012)033 [arXiv:1205.0559 [hep-th]].
- [35] D. Kubiznak, R. B. Mann and M. Teo, Black hole chemistry: thermodynamics with Lambda, *Class. Quant. Grav.* **34**, no.6, 063001 (2017) doi:10.1088/1361-6382/aa5c69 [arXiv:1608.06147 [hep-th]].
- [36] K. Bhattacharya, B. R. Majhi and S. Samanta, Van der Waals criticality in AdS black holes: a phenomenological study, *Phys. Rev. D* **96**, no.8, 084037 (2017) doi:10.1103/PhysRevD.96.084037 [arXiv:1709.02650 [gr-qc]].
- [37] David Kubiznak and Robert B. Mann. *P-v criticality of charged AdS black holes*. *Journal of High Energy Physics*, **2012**(7),2012.
- [38] Natacha Altamirano, David Kubiznak, and Robert B. Mann. *Reentrant phase transitions in rotating anti-de sitter black holes*. *Physical Review D*, **88**(10),2013.
- [39] Natacha Altamirano, David Kubiznak, Robert B Mann, and Zeinab Sherkatghanad. *Kerr-ads analogue of triple point and solid/liquid/gas phase transition*. *Classical and Quantum Gravity*, **31**(4):042001, 2014.
- [40] Shao-Wen Wei and Yu-Xiao Liu. *Triple points and phase diagrams in the extended phase space of charged gauss-bonnet black holes in ads space.*, *Phys. Rev. D*, **90**:044057,2014.
- [41] Antonia M. Frassino, David Kubiznak, Robert B. Mann, and Fil Simovic, *Multiple reentrant phase transitions and triple points in lovelock thermodynamics.*, *Journal of High Energy Physics*, **2014**(9),2014.
- [42] Rong-Gen Cai, Li-Ming Cao, Li Li, and Run-Qiu Yang. *P-v criticality in the extended phase space of gauss-bonnet black holes in AdS space*. *Journal of High Energy Physics*, **2013**(9), 2013
- [43] Hao Xu, Wei Xu, and Liu Zhao. *Extended phase space thermodynamics for third-order lovelock black holes in diverse dimensions.*, *The European Physical Journal C*, **74**(9),2014.
- [44] Brian P Dolan, Anna Kostouki, David Kubiznak, and Robert B Mann. *Isolated critical point from lovelock gravity*. *Classical and Quantum Gravity*, **31**(24):242001, 2014
- [45] Robie A. Hennigar, W. G. Brenna, and Robert B. Mann. *P-v criticality in quasitopological gravity.*, *Journal of High Energy Physics*, **2015**(7), 2015.
- [46] Robie Hennigar and Robert Mann. *Reentrant phase transitions and van der waals behaviour for hairy black holes*. *Entropy***17**(12):8056–8072, 2015.

- [47] Robie A. Hennigar, Robert B. Mann, and Erickson Tjoa. *Superfluid black holes*. Physical Review Letters **118**(2),2017.
- [48] De-Cheng Zou,Ruihong Yue, and Ming Zhang. *Reentrant phase transitions of higher-dimensional ads black holes in drgt massive gravity*. The European Physical Journal C, **77**(4),2017.
- [49] Gogoi, Naba Jyoti and Phukon, Prabwal. *Thermodynamic geometry of 5D R-charged black holes in extended thermodynamic space*. Phys. Rev. D, **103**(126008),2021. doi:10.1103/PhysRevD.103.126008.
- [50] J. M. Maldacena, The Large N limit of superconformal field theories and supergravity, Adv. Theor. Math. Phys. **2**, 231-252 (1998) doi:10.4310/ATMP.1998.v2.n2.a1 [arXiv:hep-th/9711200 [hep-th]].
- [51] M. R. Visser, Holographic thermodynamics requires a chemical potential for color, Phys. Rev. D **105**, no.10, 106014 (2022)
- [52] G. Zeyuan and L. Zhao, *Restricted phase space thermodynamics for AdS black holes via holography*, Class. Quant. Grav. **39** (2022) no.7, 075019 doi:10.1088/1361-6382/ac566c [arXiv:2112.02386 [gr-qc]].
- [53] J. Sadeghi, M. Shokri, S. Gashti Noori and M. R. Alipour, *RPS thermodynamics of Taub–NUT AdS black holes in the presence of central charge and the weak gravity conjecture*. Gen. Rel. Grav. **54** (2022)
- [54] Y. Ladghami, B. Asfour, A. Bouali, A. Errahmani and T. Ouali, *4D-EGB black holes in RPS thermodynamics*, Phys. Dark Univ. **41** (2023),
- [55] X. Kong, Z. Zhang and L. Zhao, *Restricted phase space thermodynamics of charged AdS black holes in conformal gravity*, Chin. Phys. C **47** (2023)
- [56] M. R. Alipour, J. Sadeghi and M. Shokri, *WGC and WCCC of black holes with quintessence and cloud strings in RPS space*, Nucl. Phys. B **990** (2023)
- [57] T. Wang and L. Zhao, *Black hole thermodynamics is extensive with variable Newton constant*, Phys. Lett. B **827** (2022)
- [58] Z. Gao, X. Kong and L. Zhao, *Thermodynamics of Kerr-AdS black holes in the restricted phase space*, Eur. Phys. J. C **82** (2022)
- [59] Ali, Md and Ghosh, Sushant and Wang, Anzhong, *Thermodynamics of Kerr-Sen-AdS black holes in the restricted phase space*,Phys. Rev. D, **108**,2023
- [60] M. B. Awal and P. Phukon, *Restricted Phase Space Thermodynamics of NED-AdS Black Holes* [arXiv:2404.03261 [hep-th]].
- [61] Y. Ladghami, B. Asfour, A. Bouali, A. Errahmani and T. Ouali, *Barrow entropy and AdS black holes in RPS thermodynamics*, Phys. Dark Univ. **44** (2024), 101470 doi:10.1016/j.dark.2024.101470 [arXiv:2403.08991 [hep-th]].
- [62] Y. Ladghami, B. Asfour, A. Bouali, A. Errahmani and T. Ouali, *4D-EGB black holes in RPS thermodynamics*, Phys. Dark Univ. **41** (2023), 101261 doi:10.1016/j.dark.2023.101261 [arXiv:2303.00604 [hep-th]].
- [63] Y. Z. Du, H. F. Li, Y. Zhang, X. N. Zhou and J. X. Zhao, *Restricted Phase Space Thermodynamics of Einstein-Power-Yang–Mills AdS Black Hole*, Entropy **25** (2023) no.4, 687 doi:10.3390/e25040687 [arXiv:2210.02006 [hep-th]].
- [64] S. Dutta and G. S. Punia, *String theory corrections to holographic black hole chemistry*, Phys. Rev. D **106**, no.2, 026003 (2022)
- [65] T. F. Gong, J. Jiang and M. Zhang, *Holographic thermodynamics of rotating black holes*, JHEP **06**, 105 (2023)
- [66] W. Cong, D. Kubiznak, R. B. Mann and M. R. Visser, *Holographic CFT phase transitions and criticality for charged AdS black holes*, JHEP **08**, 174 (2022)
- [67] H. Quevedo and A. Sanchez, Geometric description of BTZ black holes thermodynamics, Phys. Rev. D **79**, 024012 (2009) doi:10.1103/PhysRevD.79.024012 [arXiv:0811.2524 [gr-qc]].
- [68] M. Akbar, H. Quevedo, K. Saifullah, A. Sanchez and S. Taj, Thermodynamic Geometry Of Charged Rotating BTZ Black Holes, Phys. Rev. D **83**, 084031 (2011) doi:10.1103/PhysRevD.83.084031 [arXiv:1101.2722 [gr-qc]].
- [69] S. H. Hendi and R. Naderi, Geometrothermodynamics of black holes in Lovelock gravity with a nonlinear electrodynamics, Phys. Rev. D **91**, no.2, 024007 (2015) doi:10.1103/PhysRevD.91.024007 [arXiv:1510.06269 [hep-th]].
- [70] T. Sarkar, G. Sengupta and B. Nath Tiwari, On the thermodynamic geometry of BTZ black holes, JHEP **11**, 015 (2006) doi:10.1088/1126-6708/2006/11/015 [arXiv:hep-th/0606084 [hep-th]].
- [71] S. H. Hendi, A. Sheykhi, S. Panahiyani and B. Eslam Panah, Phase transition and thermodynamic geometry of Einstein-Maxwell-dilaton black holes, Phys. Rev. D **92**, no.6, 064028 (2015) doi:10.1103/PhysRevD.92.064028 [arXiv:1509.08593 [hep-th]].
- [72] R. Banerjee, B. R. Majhi and S. Samanta, Thermogeometric phase transition in a unified framework, Phys. Lett. B **767**, 25-28 (2017) doi:10.1016/j.physletb.2017.01.040 [arXiv:1611.06701 [gr-qc]].
- [73] K. Bhattacharya and B. R. Majhi, Thermogeometric description of the van der Waals like phase transition in AdS black holes, Phys. Rev. D **95**, no.10, 104024 (2017) doi:10.1103/PhysRevD.95.104024 [arXiv:1702.07174 [gr-qc]].
- [74] K. Bhattacharya and B. R. Majhi, Thermogeometric study of van der Waals like phase transition in black holes: an alternative approach, Phys. Lett. B **802**, 135224 (2020) doi:10.1016/j.physletb.2020.135224 [arXiv:1903.10370 [gr-qc]].
- [75] N. J. Gogoi, G. K. Mahanta and P. Phukon, Geodesics in geometrothermodynamics (GTD) type II geometry of 4D asymptotically anti-de-Sitter black holes, Eur. Phys. J. Plus **138**, no.4, 345 (2023) doi:10.1140/epjp/s13360-023-03938-x
- [76] P. Kumar, S. Mahapatra, P. Phukon and T. Sarkar, Geodesics in Information Geometry : Classical and Quantum Phase Transitions, Phys. Rev. E **86**, 051117 (2012) doi:10.1103/PhysRevE.86.051117 [arXiv:1210.7135 [cond-mat.stat-mech]].
- [77] Y. S. Duan, *The structure of the topological current*,SLAC-PUB-3301,1984.
- [78] Yi-Shi Duan and Mo-Lin Ge. *SU(2) Gauge Theory and Electrodynamics with N Magnetic Monopoles*,Sci.Sin **9**,1072,1979
- [79] Shao-Wen Wei and Yu-Xiao Liu. *Topology of black hole thermodynamics*. Phys. Rev. D **105**:104003, 2022.
- [80] Shao-Wen Wei, Yu-Xiao Liu, and Robert B. Mann. *Black hole solutions as topological thermodynamic defects*. Phys. Rev. Lett.**129**:191101,2022.
- [81] C. Fairros, *Topological Interpretation Black Hole Phase Transition in Gauss-Bonnet Gravity*, [arXiv:2311.04050 [gr-qc]].
- [82] M. Rizwan and K. Jusufi, *Topological classes of thermodynamics of black holes in perfect fluid dark matter background* Eur. Phys. J. C **83**, no.10, 944 (2023) doi:10.1140/epjc/s10052-023-12126-1 [arXiv:2310.15182 [gr-qc]]

- [83] A. Mehmood and M. U. Shahzad, *Thermodynamic Topological Classifications of Well-Known Black Holes*, [arXiv:2310.09907 [hep-th]]
- [84] C. W. Tong, B. H. Wang and J. R. Sun, *Topology of black hole thermodynamics via Rényi statistics*, [arXiv:2310.09602 [gr-qc]].
- [85] M. U. Shahzad, A. Mehmood, S. Sharif and A. Övgün, *Criticality and topological classes of neutral Gauss–Bonnet AdS black holes in 5D*, *Annals Phys.* **458**, 169486 (2023) doi:10.1016/j.aop.2023.169486.
- [86] Z. Q. Chen and S. W. Wei, *Thermodynamics, Ruppeiner geometry, and topology of Born-Infeld black hole in asymptotic flat spacetime*, *Nucl. Phys. B* **996**, 116369 (2023) doi:10.1016/j.nuclphysb.2023.116369.
- [87] C. Liu, R. Li, K. Zhang and J. Wang, *Generalized free energy and dynamical state transition of the dyonic AdS black hole in the grand canonical ensemble*, [arXiv:2309.13931 [gr-qc]].
- [88] F. Barzi, H. El Moumni and K. Masmr, *Rényi Topology of Charged-flat Black Hole: Hawking-Page and Van-der-Waals Phase Transitions*, [arXiv:2309.14069 [hep-th]].
- [89] Y. Z. Du, H. F. Li, Y. B. Ma and Q. Gu, *Topology and phase transition for EPYM AdS black hole in thermal potential*, [arXiv:2309.00224 [hep-th]].
- [90] F. Demami, H. El Moumni, K. Masmr and S. Mazzou, *Thermodynamics and phase transition structure of charged black holes in $f(R)$ gravity background from Rényi statistics perception*, *Nucl. Phys. B* **994**, 116316 (2023) doi:10.1016/j.nuclphysb.2023.116316
- [91] R. Li, C. Liu, K. Zhang and J. Wang, *Topology of the landscape and dominant kinetic path for the thermodynamic phase transition of the charged Gauss-Bonnet-AdS black holes*, *Phys. Rev. D* **108**, no.4, 044003 (2023) doi:10.1103/PhysRevD.108.044003 [arXiv:2302.06201 [gr-qc]].
- [92] D. Wu, *Topological classes of thermodynamics of the four-dimensional static accelerating black holes*, *Phys. Rev. D* **108**, no.8, 084041 (2023) doi:10.1103/PhysRevD.108.084041 [arXiv:2307.02030 [hep-th]]
- [93] Y. S. Wang, Z. M. Xu and B. Wu, *Thermodynamic phase transition and winding number for the third-order Lovelock black hole*, [arXiv:2307.01569 [gr-qc]].
- [94] D. Chen, Y. He and J. Tao, *Topological classes of higher-dimensional black holes in massive gravity*, *Eur. Phys. J. C* **83**, no.9, 872 (2023) doi:10.1140/epjc/s10052-023-11983-0 [arXiv:2306.13286 [gr-qc]].
- [95] J. Sadeghi, S. Noori Gashti, M. R. Alipour and M. A. S. Afshar, *Bardeen black hole thermodynamics from topological perspective*, *Annals Phys.* **455**, 169391 (2023) doi:10.1016/j.aop.2023.169391 [arXiv:2306.05692 [hep-th]].
- [96] D. Wu, *Consistent thermodynamics and topological classes for the four-dimensional Lorentzian charged Taub-NUT spacetimes*, *Eur. Phys. J. C* **83**, no.7, 589 (2023) doi:10.1140/epjc/s10052-023-11782-7 [arXiv:2306.02324 [gr-qc]].
- [97] T. N. Hung and C. H. Nam, *Topology in thermodynamics of regular black strings with Kaluza–Klein reduction*, *Eur. Phys. J. C* **83**, no.7, 582 (2023) doi:10.1140/epjc/s10052-023-11768-5 [arXiv:2305.15910 [gr-qc]].
- [98] M. Y. Zhang, H. Chen, H. Hassanabadi, Z. W. Long and H. Yang, *Topology of nonlinearly charged black hole chemistry via massive gravity*, *Eur. Phys. J. C* **83**, no.8, 773 (2023) doi:10.1140/epjc/s10052-023-11933-w [arXiv:2305.15674 [gr-qc]].
- [99] N. J. Gogoi and P. Phukon, *Topology of thermodynamics in R-charged black holes*, *Phys. Rev. D* **107**, no.10, 106009 (2023) doi:10.1103/PhysRevD.107.106009
- [100] Z. M. Xu, Y. S. Wang, B. Wu and W. L. Yang, *Thermodynamic phase transition of the AdS black holes from the perspective of the complex analysis*, [arXiv:2305.05916 [gr-qc]].
- [101] M. R. Alipour, M. A. S. Afshar, S. Noori Gashti and J. Sadeghi, *Topological classification and black hole thermodynamics*, *Phys. Dark Univ.* **42**, 101361 (2023) doi:10.1016/j.dark.2023.101361 [arXiv:2305.05595 [gr-qc]].
- [102] P. K. Yerra, C. Bhamidipati and S. Mukherji, *Topology of critical points in boundary matrix duals*, [arXiv:2304.14988 [hep-th]].
- [103] N. J. Gogoi and P. Phukon, *Thermodynamic topology of 4D dyonic AdS black holes in different ensembles*, *Phys. Rev. D* **108**, no.6, 066016 (2023) doi:10.1103/PhysRevD.108.066016 [arXiv:2304.05695 [hep-th]].
- [104] C. Fairoos and T. Sharqui, *Topological nature of black hole solutions in dRGT massive gravity*, *Int. J. Mod. Phys. A* **38**, no.25, 2350133 (2023) doi:10.1142/S0217751X23501336 [arXiv:2304.02889 [gr-qc]].
- [105] R. Li and J. Wang, *Generalized free energy landscapes of charged Gauss-Bonnet-AdS black holes in diverse dimensions*, *Phys. Rev. D* **108**, no.4, 044057 (2023) doi:10.1103/PhysRevD.108.044057 [arXiv:2304.03425 [gr-qc]].
- [106] M. Zhang and J. Jiang, *Bulk-boundary thermodynamic equivalence: a topology viewpoint*, *JHEP* **06**, 115 (2023) doi:10.1007/JHEP06(2023)115 [arXiv:2303.17515 [hep-th]].
- [107] Y. Du and X. Zhang, *Topological classes of black holes in de-Sitter spacetime*, *Eur. Phys. J. C* **83**, no.10, 927 (2023) doi:10.1140/epjc/s10052-023-12114-5 [arXiv:2303.13105 [gr-qc]].
- [108] S. W. Wei, Y. P. Zhang, Y. X. Liu and R. B. Mann, *Static spheres around spherically symmetric black hole spacetime*, *Phys. Rev. Res.* **5**, no.4, 043050 (2023) doi:10.1103/PhysRevResearch.5.043050 [arXiv:2303.06814 [gr-qc]].
- [109] Y. Du and X. Zhang, *Topological classes of BTZ black holes*, [arXiv:2302.11189 [gr-qc]].
- [110] Q. Yu, Q. Xu and J. Tao, *Thermodynamics and microstructures of Euler–Heisenberg black hole in a cavity*, *Commun. Theor. Phys.* **75**, no.9, 095402 (2023) doi:10.1088/1572-9494/ace4b3 [arXiv:2302.09821 [gr-qc]].
- [111] N. Chatzifotis, P. Dorlis, N. E. Mavromatos and E. Papanonopoulos, *Thermal stability of hairy black holes*, *Phys. Rev. D* **107**, no.8, 084053 (2023) doi:10.1103/PhysRevD.107.084053 [arXiv:2302.03980 [gr-qc]].
- [112] D. Wu, *Classifying topology of consistent thermodynamics of the four-dimensional neutral Lorentzian NUT-charged spacetimes*, *Eur. Phys. J. C* **83**, no.5, 365 (2023) doi:10.1140/epjc/s10052-023-11561-4 [arXiv:2302.01100 [gr-qc]].
- [113] D. Wu and S. Q. Wu, *Topological classes of thermodynamics of rotating AdS black holes*, *Phys. Rev. D* **107**, no.8, 084002 (2023) doi:10.1103/PhysRevD.107.084002 [arXiv:2301.03002 [hep-th]].
- [114] D. Wu, *Topological classes of rotating black holes*, *Phys. Rev. D* **107**, no.2, 024024 (2023) doi:10.1103/PhysRevD.107.024024 [arXiv:2211.15151 [gr-qc]].

- [115] C. Fang, J. Jiang and M. Zhang, *Revisiting thermodynamic topologies of black holes*, JHEP **01**, 102 (2023) doi:10.1007/JHEP01(2023)102 [arXiv:2211.15534 [gr-qc]].
- [116] Z. Y. Fan, *Topological interpretation for phase transitions of black holes* Phys. Rev. D **107**, no.4, 044026 (2023) doi:10.1103/PhysRevD.107.044026 [arXiv:2211.12957 [gr-qc]].
- [117] C. Liu and J. Wang, *Topological natures of the Gauss-Bonnet black hole in AdS space*, Phys. Rev. D **107**, no.6, 064023 (2023) doi:10.1103/PhysRevD.107.064023 [arXiv:2211.05524 [gr-qc]].
- [118] N. C. Bai, L. Li and J. Tao, *Topology of black hole thermodynamics in Lovelock gravity* Phys. Rev. D **107**, no.6, 064015 (2023) doi:10.1103/PhysRevD.107.064015 [arXiv:2208.10177 [gr-qc]].
- [119] P. K. Yerra, C. Bhamidipati and S. Mukherji, *Topology of critical points and Hawking-Page transition*, Phys. Rev. D **106**, no.6, 064059 (2022) doi:10.1103/PhysRevD.106.064059 [arXiv:2208.06388 [hep-th]].
- [120] J. Sadeghi, M. A. S. Afshar, S. Noori Gashti and M. R. Alipour, *Thermodynamic topology of black holes from bulk-boundary, extended, and restricted phase space perspectives*, Annals Phys. **460** (2024), 169569 doi:10.1016/j.aop.2023.169569 [arXiv:2312.04325 [hep-th]].
- [121] N. J. Gogoi and P. Phukon, *Thermodynamic topology of 4D Euler-Heisenberg-AdS black hole in different ensembles*, [arXiv:2312.13577 [hep-th]].
- [122] B. Hazarika and P. Phukon, *Thermodynamic Topology of $D = 4, 5$ Horava Lifshitz Black Hole in Two Ensembles*, [arXiv:2312.06324 [hep-th]].
- [123] B. Hazarika, N. J. Gogoi and P. Phukon, *Revisiting thermodynamic topology of Hawking-Page and Davies type phase transitions*, [arXiv:2404.02526 [hep-th]].
- [124] B. Hazarika and P. Phukon, *Thermodynamic topology of black Holes in f(R) gravity*, Progress of Theoretical and Experimental Physics **4**(2024), <https://doi.org/10.1093/ptep/ptae035> [arXiv:2401.16756 [hep-th]].
- [125] M. Y. Zhang, H. Chen, H. Hassanabadi, Z. W. Long and H. Yang, *Thermodynamic topology of Kerr-Sen black holes via Rényi statistics*, [arXiv:2312.12814 [gr-qc]].
- [126] D. Chen, Y. He, J. Tao and W. Yang, *Topology of Hořava-Lifshitz black holes in different ensembles*, [arXiv:2311.11606 [hep-th]].
- [127] K. Bhattacharya, K. Bamba and D. Singleton, *Topological interpretation of extremal and Davies-type phase transitions of black holes*, [arXiv:2402.18791 [gr-qc]].
- [128] X. D. Zhu, D. Wu and D. Wen, *Topological classes of thermodynamics of the rotating charged AdS black holes in gauged supergravities*, [arXiv:2402.15531 [hep-th]].
- [129] H. Chen, M. Y. Zhang, H. Hassanabadi and Z. W. Long, *Thermodynamic topology of Phantom AdS Black Holes in Massive Gravity*, [arXiv:2404.08243 [gr-qc]].
- [130] D. Wu, S. Y. Gu, X. D. Zhu, Q. Q. Jiang and S. Z. Yang, *Topological classes of thermodynamics of the static multi-charge AdS black holes in gauged supergravities*, [arXiv:2402.00106 [hep-th]].
- [131] S. Mahapatra, P. Phukon and T. Sarkar, *On Black Hole Entropy Corrections in the Grand Canonical Ensemble*, Phys. Rev. D **84**, 044041 (2011)
- [132] N. J. Gogoi, G. K. Mahanta and P. Phukon, *Geodesics in geometrothermodynamics (GTD) type II geometry of 4D asymptotically anti-de-Sitter black holes*, Eur. Phys. J. Plus **138**
- [133] C. S. Peca and J. P. S. Lemos, *Thermodynamics of Reissner-Nordstrom anti-de Sitter black holes in the grand canonical ensemble* Phys. Rev. D **59**, 124007 (1999)
- [134] T. V. Fernandes and J. P. S. Lemos, *Grand canonical ensemble of a d-dimensional Reissner-Nordström black hole in a cavity* Phys. Rev. D **108**, no.8, 084053 (2023)
- [135] Z. Wang, H. Ren, J. Chen and Y. Wang, *Thermodynamics and phase transition of Bardeen black hole via Rényi statistics in grand canonical ensemble and canonical ensemble*, Eur. Phys. J. C **83**, no.6, 527 (2023)
- [136] C. Liu, R. Li, K. Zhang and J. Wang, *Generalized free energy and dynamical state transition of the dyonic AdS black hole in the grand canonical ensemble*, JHEP **11**, 068 (2023)
- [137] A. Sen, *Rotating charged black hole solution in heterotic string theory*, Phys. Rev. Lett. **69**, 1006-1009 (1992) doi:10.1103/PhysRevLett.69.1006 [arXiv:hep-th/9204046 [hep-th]].
- [138] M. Zhang and J. Jiang, *Strong cosmic censorship in near-extremal Kerr-Sen-de Sitter spacetime*, Eur. Phys. J. C **81**, no.11, 967 (2021) doi:10.1140/epjc/s10052-021-09753-x [arXiv:2110.09077 [gr-qc]].
- [139] M. Sharif and Q. Ama-Tul-Mughani, *P-V criticality and phase transition of the Kerr-Sen-AdS Black Hole*, Eur. Phys. J. Plus **136**, no.3, 284 (2021) doi:10.1140/epjp/s13360-021-01270-w
- [140] T. Roy, A. Sardar and U. Debnath, *Thermodynamic overview and heat engine efficiency of Kerr-Sen-AdS black hole*, Int. J. Geom. Meth. Mod. Phys. **20**, no.08, 2350136 (2023) doi:10.1142/S0219887823501360
- [141] M. F. A. R. Sakti and P. Burikham, *Dual CFT on Nariai limit for Kerr-Sen-dS black holes*, Eur. Phys. J. C **84**, no.2, 182 (2024) doi:10.1140/epjc/s10052-024-12537-8 [arXiv:2307.04929 [hep-th]].
- [142] H. M. Siahhan, *Pair production of scalars around near-extremal Kerr-Sen black holes*, Eur. Phys. J. C **80**, no.5, 387 (2020) doi:10.1140/epjc/s10052-020-7920-z [arXiv:1907.09372 [gr-qc]].
- [143] D. Wu, P. Wu, H. Yu and S. Q. Wu, *Are ultraspinning Kerr-Sen-AdS₄ black holes always superentropic?*, Phys. Rev. D **102**, no.4, 044007 (2020) doi:10.1103/PhysRevD.102.044007 [arXiv:2007.02224 [gr-qc]].
- [144] D. Wu, S. Q. Wu, P. Wu and H. Yu, *Aspects of the dyonic Kerr-Sen-AdS₄ black hole and its ultraspinning version*, Phys. Rev. D **103**, no.4, 044014 (2021) doi:10.1103/PhysRevD.103.044014 [arXiv:2010.13518 [gr-qc]].
- [145] A. Haldar, R. Biswas and B. Ghosh, *Extension of thermodynamics of the Kerr-Sen black hole under general uncertainty principle along with first-order corrected entropy*, Mod. Phys. Lett. A **37**, no.21, 2250131 (2022) doi:10.1142/S0217732322501310

- [146] M. F. A. R. Sakti and P. Burikham, *Dual CFT on a dyonic Kerr-Sen black hole and its gauged and ultraspinning counterparts*, Phys. Rev. D **106**, no.10, 106006 (2022) doi:10.1103/PhysRevD.106.106006 [arXiv:2206.10868 [hep-th]].