### The Birkhoff completion of finite lattices

Mohammad Abdulla<sup>1,2</sup>, Johannes Hirth<sup>1,2</sup>, Gerd Stumme<sup>1,2</sup>

<sup>1</sup> Knowledge & Data Engineering Group, University of Kassel, Germany
<sup>2</sup> Interdisciplinary Research Center for Information System Design,
University of Kassel, Germany

**Abstract.** We introduce the Birkhoff completion as the smallest distributive lattice in which a given finite lattice can be embedded as semilattice. We discuss its relationship to implicational theories, in particular to R. Wille's simply-implicational theories. By an example, we show how the Birkhoff completion can be used as a tool for ordinal data science.

**Keywords:** Formal Concept Analysis  $\cdot$  Distributive Lattices  $\cdot$  Implicational Theories

#### 1 Introduction

Formal Concept Analysis (FCA) is a popular means for data analysis, especially when at least one feature is not of nominal or interval scale type. It is known since the beginning that the resulting concept lattice can take the form of any finite<sup>3</sup> lattice. In real-world applications, though, surprisingly often the concept lattice carries more structure: being a distributive lattice which eventually is "cut off" at the bottom.

The likely reason for this unexpected emergence of distributivity is related to the fact that a concept lattice is distributive if and only if the implicational logic of its attributes can be described completely by implications that have only one attribute in their premise. The distributivity thus results from the prevalence of implications with one-element premises. According to Wille (2003) [10] this is caused "by the nature of human thought namely that everyday thinking predominantly uses logical inferences with one element premises."

If a concept lattice is distributive, this comes with several advantages. Besides a simpler — and easier to communicate — implicational logic, one benefits from the fact that the concept lattice can be decomposed into chains, which support an easier interpretation of the data and a natural grid-like visualisation by an embedding into the products of these chains. Furthermore the lattice is isomorphic to the order ideals of the ordered set of its meet-irreducible elements, which provides an interesting structure for supporting the navigation in the concept lattice.

<sup>&</sup>lt;sup>3</sup> Throughout this paper we assume that our data — the formal context — are finite. If we would also allow for infinite contexts then the concept lattice can take the form of any complete lattice.

But what if the concept lattice of formal context is not (completely) distributive? There are different possible explanations. The first is of course that the data we are analysing have a more complex logical structure. Another reason might be that the logic of our field of interest has a simple implicational structure but that the formal context which we have at hand is not perfectly representing it. It might for instance miss some of the objects or attributes or contain errors.

In this paper, we present a construction that may "repair" a non-distributive concept lattice whenever we have reason to assume that our context is not covering all potential objects of our domain of interest. The underlying construction is not specific to FCA, and all its building blocks — in particular Birkhoff's representation theorem for distributive lattices [2] — are known in lattice theory for a very long time. However, no-one has yet turned it into a construction for embedding a non-distributive lattice into a minimal distributive lattice.<sup>4</sup> Our proposed Birkhoff completion is therefore not only of interest to FCA but can be seen as a generic lattice-theoretical construction. We will therefore also introduce the Birkhoff completion in purely lattice-theoretical terms in Section 3. References to related work are not collected in a separate section, but mentioned whenever appropriate.

All methods presented in this paper are implemented and provided by the conexp-clj FCA research framework [5].

#### 2 Basics

In the following we recall some notions from order and lattice theory [1,2], and introduce some notations used in this work.

Let P and Q be two partially ordered sets. We say that P and Q are order-isomorphic, denoted by  $P \cong Q$ , if there exists an injective map  $\varphi$  from P onto Q such that  $x \leq y$  in P if and only if  $\varphi(x) \leq \varphi(y)$  in Q. Such a map is called an order-isomorphism. Associated with any ordered set is the corresponding familiy of down-sets, which plays a major role in our paper. Let P be an ordered set and  $Q \subseteq P$ . We say that Q is a down-set (or (order) ideal) if, whenever  $x \in Q$ ,  $y \in P$  and  $y \leq x$ , we have  $y \in Q$ . Dually, Q is an up-set (or (order) filter) if, whenever  $x \in Q$ ,  $y \in P$  and  $x \leq y$ , we have  $y \in Q$ . The down-set of  $x \in L$  is  $x := \{a \in L \mid a \leq x\}$ , and the up-set of  $x \in L$  is  $x := \{a \in L \mid a \leq x\}$ . We denote the family of the order ideals of P by  $x \in C$ 0, and the family of its order filters by  $x \in C$ 1. They are both lattices, under the inclusion order.

A partially ordered set L is called a *lattice* if, for any two elements  $x, y \in L$ , their infimum  $x \wedge y$  and supremum  $x \vee y$  exist in L. In a lattice, infimum and supremum are also called *meet* and *join*, resp. A join-semilattice is a partially ordered set that has a join for any nonempty finite subset. Dually, a meet-semilattice is a partially ordered set which has a meet for any nonempty finite

<sup>&</sup>lt;sup>4</sup> For a reader with background in lattice theory or general algebra it will be immediately clear that such an embedding will only work if we do not require the embedding to preserve all lattice operations. We will discuss this in detail in Section 3.




Fig. 1. The nondistributive lattices  $M_3$  and  $N_5$ 

subset. An element x of a (finite) lattice L is called meet-irreducible if  $\bigvee X = x$  implies  $x \in X$  for all  $X \subseteq L$ . Dually, x is called join-irreducible if  $\bigwedge X = x$  implies  $x \in X$  for all  $X \subseteq L$ . The set of all meet-irreducible elements of L is denoted by  $\mathcal{M}(L)$ , and the set of all its join-irreducibles is denoted by  $\mathcal{J}(L)$ . We call  $S \neq \emptyset$  a sublattice of L if  $(a, b \in S \Rightarrow a \lor b \in S)$  and  $a \land b \in S)$  holds.

Let L and K be lattices. A map  $f:L\to K$  is said to be a lattice homomorphism iff for all  $a,b\in L$ ,  $f(a\vee b)=f(a)\vee f(b)$  and  $f(a\wedge b)=f(a)\wedge f(b)$ . A bijective lattice homomorphism is a lattice isomorphism. If  $f:L\to K$  is a one-to-one homomorphism, then the sublattice f(L) of K is isomorphic to L and we refer to f as an embedding (of L into K).

A map  $\phi: L \to K$  between two (finite) lattices L and K is called join-preserving if  $\phi(\bigvee X) = \bigvee \phi(X)$  holds for all  $X \subseteq L$ . Dually, $\phi$  is meet preserving if  $\phi(\bigwedge X) = \bigwedge \phi(X)$  holds for all  $X \subseteq L$ . Distributivity in a lattice governs how the meet and join operations interact, ensuring consistency in lattice transformations.

**Definition 1.** A lattice L is said to be distributive if for all  $x, y, z \in L$ :

$$x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$$

**Theorem 1.** A lattice L is distributive if and only if there exists no sublattice  $A \subseteq L$  isomorphic to either  $M_3$  or  $N_5$ .

# 3 Fixing non distributivity based on Birkhoff's Representation Theorem

Our approach for fixing non-distributivity is based on Birkhoff's Representation Theorem [2], which states that for any finite distributive lattice L, the lattice  $\mathcal{I}(\mathcal{J}(L))$  of down-sets of the join-irreducible elements of L is isomorphic to L. It's important to note that we focus on this variation of the theorem tailored for finite lattices, as our work primarily involves finite structures.

Theorem 2 (Birkhoff's Representation Theorem).

4

If L is a finite<sup>5</sup> distributive lattice, then the map  $\eta: L \to \mathcal{I}(J(L))$  defined by  $\eta(a) = \{x \in \mathcal{J}(L) \mid x \leq a\}$  is an isomorphism of L onto  $(\mathcal{I}(J(L)), \subseteq)$ . Conversely, for every finite ordered set, the closure system of all order ideals is a distributive lattice D in which

$$\mathcal{J}(D) = \{ \downarrow x \, | \, x \in P \}$$

is supremum-dense.

We refer to Ganter and Wille (1999) [4] for the proof.

Although this theorem is usually based on the down-sets of the join-irreducible elements of L, the same also holds for the up-sets of the meet-irreducible elements of L: We have  $(\mathcal{I}(\mathcal{J}(L)), \subseteq) \cong (\mathcal{F}(\mathcal{M}(L)), \supseteq)$  for distributive lattices because in that case  $(\mathcal{J}(L), \subseteq) \cong (\mathcal{M}(L), \subseteq)$  holds (see Exercise 5.7 in Davey and Priestley, 2002 [1]) and because the complement of an order filter of an ordered set is always an order ideal and vice versa. As we want to link our work with observations about the implications between attributes in a formal context in Section 4, we prefer to continue with the equivalent dual representation:

Theorem 3 (Dual version of Birkhoff's Representation Theorem). If L is a finite distributive lattice, then the map  $\eta: L \to \mathcal{F}(\mathcal{M}(L))$  defined by  $\eta(a) = \{x \in \mathcal{M}(L) \mid x \geq a\}$  is an isomorphism of L onto  $(\mathcal{F}(\mathcal{M}(L)), \supseteq)$ . Conversely, for every finite ordered set P, the closure system of all order filters is a distributive lattice in which

$$\mathcal{M}(D) = \{ \uparrow x \mid x \in P \}$$


is infimum-dense.

**Corollary 1.** Let L be a finite lattice. Then the following statements are equivalent:


- (i) L is distributive.
- (ii)  $L \cong (\mathcal{I}(\mathcal{J}(L)), \subseteq)$ .
- (iii)  $L \cong (\mathcal{F}(\mathcal{M}(L)), \supseteq)$ .
- (iv) L is isomorphic to the lattice of all down-sets of some finite ordered set.
- (v) L is isomorphic to the lattice of all up-sets of some finite ordered set.

This implies that starting from any finite lattice L, one obtains two distributive lattices  $\mathcal{I}(\mathcal{J}(L))$  and  $\mathcal{F}(\mathcal{M}(L))$ , which are isomorphic to L if and only if L is distributive. This gives rise to the following constructions.

<sup>&</sup>lt;sup>5</sup> This theorem holds — like many of the statements in this paper — also for specific infinite lattices, the so-called doubly founded lattices. As our focus is on applications in data analysis, we consider only the finite case in the sequel, which is easier to handle.



**Fig. 2.**  $M_3$  (left), its down-set BC (middle) and its up-set BC(Right).



**Fig. 3.**  $N_5$  (left), its down-set BC (middle) and its up-set BC(Right).

**Definition 2.** Let  $(L, \leq)$  be a finite lattice. The down-set Birkhoff completion of L is the lattice  $(\mathcal{I}(\mathcal{J}(L)), \subseteq)$ . The up-set Birkhoff completion of L is the lattice  $(\mathcal{F}(\mathcal{M}(L)), \supseteq)$ . When we simply talk about the Birkhoff completion, we refer to the latter, and denote it by BC(L).

Remark 1. In terms of FCA, these two constructions are isomorphic to the two concept lattices  $\underline{\mathfrak{B}}(\mathcal{J}(L),\mathcal{J}(L),\not\geq)$  and  $\underline{\mathfrak{B}}(\mathcal{M}(L),\mathcal{M}(L),\not\geq)$ . In fact, the order ideals of  $\mathcal{J}(L)$  are exactly the concept extents of  $\underline{\mathfrak{B}}(\mathcal{J}(L),\mathcal{J}(L),\not\geq)$ , and the order filters of  $\mathcal{F}(L)$  are exactly the concept extents of  $\underline{\mathfrak{B}}(\mathcal{M}(L),\mathcal{M}(L),\not\geq)$  [9]. By abuse of notation, we will therefore also refer to these concept lattices as down-set Birkhoff completion and (up-set) Birkhoff completion, resp.

The Birkhoff completion BC(L) can be understood as a way to extend the original lattice L in a somewhat minimal way to a distributive lattice while preserving its essential structure.<sup>6</sup>

<sup>&</sup>lt;sup>6</sup> This is in a certain sense complementary to the construction of the free distributive completion of a partial complete lattice, which was introduced in Stumme (1998) [6].

**Theorem 4.** Let L be a finite lattice.

- 1. BC(L) is distributive.
- 2. The map

$$\iota : L \to BC(L) : x \mapsto \uparrow x \cap \mathcal{M}(L)$$

 $is\ a\ join-semilattice\ embedding.$ 

3. For any distributive lattice  $\hat{L}$  for which exists a join-semilattice embedding  $\varphi \colon L \to \hat{L}$ , there exists a join-semilattice embedding  $\varepsilon \colon BC(L) \to \hat{L}$  such that  $\varphi = \varepsilon \circ \iota$  holds.

*Proof.* 1. This follows directly from Theorem 3, since  $BC(L) = (\mathcal{F}(\mathcal{M}(L)), \supseteq)$ .

- 2. First, we note that for  $a, b \in L$  the equality  $\uparrow (a \lor b) = \uparrow a \cap \uparrow b$  holds. This implies that  $\iota(a \lor b) = \iota(a) \cap \iota(b)$ . Thus,  $\iota$  is joins preserving. In a finite lattice every element  $x \in L$  can be uniquely identified by the set of all meetirreducible elements  $m \in \mathcal{M}(L)$  for which  $m \ge x$ , i.e.,  $x = \bigwedge \{m \in \mathcal{M}(L) \mid m \ge x\}$ . Thus, is for  $x, y \in L$  with  $x \ne y$  the image  $\iota(x) \ne \iota(y)$ . Concluding,  $\iota$  is injective and therefore a join-semilattice embedding.
- 3. Let  $\kappa: BC(L) \to L$  with  $\kappa(A) := \{x \in L \mid \iota(x) \subseteq A\}$  and  $\varepsilon: BC(L) \to \hat{L}$  with  $\varepsilon(A) := \bigwedge_{\hat{L}} \{\varphi(x) \mid x \in \kappa(A)\}$  be two maps. For  $x, y \in L$ , we find that  $\iota(y) \subseteq \iota(x) \iff y \geq x$ . Combined with the fact that  $\varphi$  is an orderembedding we follow that  $(\varepsilon \circ \iota)(x) = \varphi(x)$ .

Next, we want to show that  $\varepsilon$  is join-preserving, i.e.,  $\varepsilon(A \vee_{BC(L)} B) = \varepsilon(A) \vee_{\hat{L}} \varepsilon(B)$ . For this, we first transform both sides of the equation to then show their equivalence:

$$\begin{split} \varepsilon(A) \vee_{\hat{L}} \varepsilon(B) &= \bigwedge_{\hat{L}} \{ \varphi(x) \mid x \in \kappa(A) \} \vee_{\hat{L}} \bigwedge_{\hat{L}} \{ \varphi(y) \mid y \in \kappa(B) \} \\ \text{(distributivity of } \hat{L}) &= \bigwedge_{\hat{L}} \{ \varphi(x) \vee_{\hat{L}} \varphi(y) \mid x \in \kappa(A), y \in \kappa(B) \} \\ (\varphi \text{ is join preserving}) &= \bigwedge_{\hat{L}} \{ \varphi(x \vee y) \mid x \in \kappa(A), y \in \kappa(B) \} \\ &= \bigwedge_{\hat{L}} \{ \varphi(z) \mid z = x \vee y, x \in \kappa(A), y \in \kappa(B) \}, \\ \varepsilon(A \vee_{BC(L)} B) &= \bigwedge_{\hat{L}} \{ \varphi(x) \mid x \in \kappa(A \vee_{BC(L)} B) \} \\ &= \bigwedge_{\hat{L}} \{ \varphi(z) \mid z \in \kappa(A \cap B) \} \\ &= \bigwedge_{\hat{L}} \{ \varphi(z) \mid z \in \kappa(A) \cap \kappa(B) \}. \end{split}$$

We proof their equality by showing for a  $z \in L$  that  $z \in \kappa(A) \cap \kappa(B)$  iff there exist  $x \in \kappa(A), y \in \kappa(B)$  with  $z = x \vee_L y$ .  $[\Rightarrow]$  This direction holds




Fig. 4. A lattice for which there exists an order embedding into a distributive lattice that has fewer elements than the up-set or down-set Birkhoff completion.


for x=z and y=z.  $[\Leftarrow]$  We can follow from  $z=x\vee_L y$  that  $z\geq x,y$ . From the fact that  $x\in\kappa(A),\,y\in\kappa(B)$ , we can follow that  $\iota(z)\subseteq\iota(x)\subseteq A$  and that  $\iota(z)\subseteq\iota(y)\subseteq B$ . Thus,  $z\in\kappa(A)$  and  $z\in\kappa(B)$ . Concluding,  $\varepsilon(A)\vee_{\hat{L}}\varepsilon(B)=\varepsilon(A\vee_{BC(L)}B)$  and  $\varepsilon$  is join-preserving.

It remains to be shown that  $\varepsilon$  is injective. Let  $A, B \in BC(L)$  with  $A \neq B$ . From this we can follow WLOG that there exists an  $x \in A$  with  $x \notin B$  and that there is no  $y \in B$  with  $y \leq x$  (otherwise x would have been in B). This implies that  $x \notin \kappa(B)$ . From the fact that  $\varphi$  is injective, we can follow that there is no  $y \in \kappa(B)$  with  $\varphi(x) = \varphi(y)$ . Moreover, since  $\varphi$  is an order embedding, we can follow that there is no  $y \in B$  such  $\varphi(y) \leq \varphi(x)$ . Hence,  $\varepsilon(B) \wedge_{\hat{L}} \varphi(x) \neq \varepsilon(B)$ , but  $\varepsilon(A) \wedge_{\hat{L}} \varphi(x) = \varepsilon(A)$ . Concluding,  $\varepsilon$  is injective and therefore a join-semilattice embedding.


Of course we cannot expect that we can always find a lattice embedding of L into BC(L) — because then BC(L) would have a non-distributive sublattice. Note that if we require the embedding to be an order embedding only, the embedding could be even smaller, as Figure 4 shows.

We conclude this section by observing that, in general, the up-set and the down-set Birkhoff completions of a lattice are not isomorphic. For example, consider Figures 5 and 6. However, as the example shows, both types of completions are closely related: The down-set Birkhoff completion of a finite lattice L is isomorphic to the dual of the (up-set) Birkhoff completion of the dual of L:

**Lemma 1.** Let L be a finite lattice. Then  $(\mathcal{I}(\mathcal{J}(L),\subseteq)) \cong (\mathcal{F}(\mathcal{M}(L^{\partial}),\supseteq))^{\partial}$ .



**Fig. 5.** L (left), its down-set BC (middle) and its up-set BC (right).



**Fig. 6.**  $L^{\partial}$  (left), its down-set BC (middle) and its up-set BC (right).

Proof. 
$$(\mathcal{J}(L), \subseteq) \cong (\mathcal{M}(L^{\partial}), \supseteq) \Longrightarrow (\mathcal{I}(\mathcal{J}(L)), \subseteq) \cong (\mathcal{F}(\mathcal{M}(L^{\partial}), \subseteq)) \cong (\mathcal{F}(\mathcal{M}(L^{\partial}), \supseteq))^{\partial}.$$

Note that the up-set and the down-set Birkhoff completions of a lattice L are of course isomorphic whenever L is distributive — because in that case both are also isomorphic to L itself. By duality, they are also isomorphic when L is isomorphic to its dual, as shown in Figure 2.

#### 4 The Birkhoff completion of a formal context

We will now transfer the construction introduced in the last section to the case when we do not start with a lattice, but with a formal context.

We start by recalling from Section 3 that for a finite lattice L its Birkhoff completion can be understood as the concept lattice  $\mathfrak{B}(\mathcal{M}(L), \mathcal{M}(L), \not\geq)$ . In the sequel, it will be convenient to differentiate, for any element of  $\mathcal{M}(L)$ , if it is considered in a specific situation as element of the object set or of the

attribute set of the context  $(\mathcal{M}(L), \mathcal{M}(L), \geq)$ . Therefore, we will frequently write this context as  $(\overline{\mathcal{M}(L)}, \mathcal{M}(L), \geq)$ , where  $\overline{\mathcal{M}(L)}$  contains a copy  $\overline{m}$  of each  $m \in \mathcal{M}(L)$ . The motivation for this notation of negating m is that, in the Birkhoff completion, for  $m \in L$ , the concept generated by the object  $\overline{m}$  is the largest concept that does not have the attribute m in its intent.

We can now define the Birkhoff completion of a formal context  $\mathbb{K} := (G, M, I)$  as extension of the set G of objects by the additional objects  $\overline{m} \in \mathcal{M}(M)$ . The incidence relation I will be extended by  $\not\geq$  for the new objects. It will be convenient, for  $m, n \in M$ , to write  $m \geq_{\mathbb{K}} n$  iff  $\{m\}' \supseteq \{n\}'$  in  $\mathbb{K}$  (which is equivalent to n implying m in the context  $\mathbb{K}$  ( $\mathbb{K} \models n \to m$ ) and to  $m \in \{n\}''$ ).

**Definition 3.** Let  $\mathbb{K} := (G, M, I)$  be a finite formal context. The Birkhoff completion of  $\mathbb{K}$  is

$$BC(\mathbb{K}) := (G \cup \overline{\mathcal{M}(M)}, M, I \cup \{(\overline{m}, n) \in \overline{\mathcal{M}(M)} \times M \mid m \not\geq_{\mathbb{K}} n\})$$

with  $\overline{\mathcal{M}(M)} := \{ \overline{m} \mid m \in \mathcal{M}(M) \}$  where  $\mathcal{M}(M)$  is the set of all meet-irreducible attributes of  $\mathbb{K}$ .

The following theorem shows that this is indeed a reasonable definition.

**Theorem 5.** Let  $\mathbb{K}$  be a finite formal context. Then  $\mathfrak{B}(BC(\mathbb{K})) \cong BC(\mathfrak{B}(\mathbb{K}))$ .

*Proof.* Let  $\mathbb{K} := (G, M, I)$  be a finite formal context.

$$BC(\underline{\mathfrak{B}}(\mathbb{K})) = \underline{\mathfrak{B}}(\mathcal{M}(\underline{\mathfrak{B}}(\mathbb{K})), \mathcal{M}(\underline{\mathfrak{B}}(\mathbb{K})), \geq) \tag{1}$$

$$\cong \underline{\mathfrak{B}}\left( \overline{\mathcal{M}(M)}, \mathcal{M}(M), \{(\overline{m}, n) \in \overline{\mathcal{M}(M)} \times \mathcal{M}(M) \mid m \not\geq_{\mathbb{K}} n \} \right)$$
 (2)

$$\cong \underline{\mathfrak{B}} \left( \overline{\mathcal{M}(M)} , M , \{(\overline{m}, n) \in \overline{\mathcal{M}(M)} \times M \mid m \not\geq_{\mathbb{K}} n \} \right)$$
 (3)

$$\cong \ \underline{\mathfrak{B}}\left(G\cup \overline{\mathcal{M}(M)}, \quad M \quad , I\cup \{(\overline{m},n)\in \overline{\mathcal{M}(M)}\times M\mid m\not\geq_{\mathbb{K}} n\}\right) \qquad (4)$$

$$= \mathfrak{B}\left(BC(\mathbb{K})\right) \tag{5}$$

- $(1) \cong (2)$ : The meet-irreducible elements of  $\mathfrak{B}(\mathbb{K})$  are exactly the attribute concepts of the attributes in  $\mathcal{M}(M)$ , and  $\mu(m) \not\geq \mu(n) \iff m \not\geq_{\mathbb{K}} nm$ .
- $(2)\cong (3)$ : We show that any attribute  $n\in M\setminus \mathcal{M}(M)$  is reducible in the context of line 3: Let  $B:=\{m\in \mathcal{M}(M)\mid m\geq_{\mathbb{K}} n\}$ . We show that  $B^\dagger=n^\dagger$  holds in the context of line 3, where  $\cdot^\dagger$  is the derivation operator in this context: First we observe that B is an order filter in  $(\mathcal{M}(M),\leq)$ , its complement  $C:=\mathcal{M}(M)\setminus B$  is an order ideal in  $(\mathcal{M}(M),\leq)$ , and it holds  $C=B^\dagger$ . On the other hand, we also have  $n^\dagger=\{\overline{m}\in \mathcal{M}(M)\mid m\not\geq_{\mathbb{K}} n\}=\mathcal{M}(M)\setminus \{\overline{m}\in \mathcal{M}(M)\mid m\geq_{\mathbb{K}} n\}=\mathcal{M}(M)\setminus B=C$ .
- (3)  $\cong$  (4): We use  $\cdot'$  as symbol for the derivation operator in  $\mathbb{K}$  and  $\cdot!$  for the derivation operator in  $BC(\mathbb{K})$  (which is the context shown in line 4). Let  $g \in G$ . We show that g is reducible in  $BC(\mathbb{K})$  by showing g! = A! with  $\overline{A} := \{\overline{m} \in \overline{\mathcal{M}(M)} \mid \neg(g,m) \in I\}$ :  $A! = \bigcap_{\overline{m} \in \overline{A}} \overline{m}! = \{n \in M \mid \forall \overline{m} \in \overline{A}: m \not\geq_{\mathbb{K}} n\} = \{n \in M \mid \forall m \in \mathcal{M}(M): m \in \mathcal{M}$

$$(4) \cong (5)$$
: By definition.

Example 1. The following example illustrates the construction. Figure 7 shows the complex usage of terms of the administrative geography of the United Kingdom [7], resulting from a long (and often bloody) historical process. The concept lattice in Figure 8 is based on the formal context which has the five countries (England, Scotland, Wales, Northern Ireland and Ireland) and the three crown dependencies (Isle of Man, Jersey and Guernsey) in the British Isles as objects, and the geographical features (marked green in Figure 7) 'British Isles', 'Ireland (Island)' and 'Great Britain' as well as the legal distinctions (blue) 'British Islands' and 'United Kingdom' as attributes. We added the geographical feature 'Channel Islands' as further attribute, as it is discussed as such in [7].

We note that the lattice is not completely distributive, as three of its sublattices are isomporphic to  $N_5$ . The up-set Birkhoff completion of the concept lattice is shown in Figure 9. Following the construction of Definition 3, it contains an additional, artificial (and potentially interesting) object for each of the five attributes. The down-set Birkhoff completion is shown in Figure 10. It contains an additional, artificial (and potentially interesting) attribute for each of the eight objects. In Section 6, we will discuss what we can learn from these completions about the administrative geography of the UK.




Fig. 7. Administrative geography of the United Kingdom [8].

#### 5 An implicational approach to the Birkhoff completion

It is long known that a finite concept lattice is distributive if and only if its canonical direct basis contains only implications that have one-element premises. After having recalled the basic notions, we will show in this section, that this




Fig. 8. Concept lattice about the administrative geography of the United Kingdom.




Fig. 9. Up-set Birkhoff completion of the concept lattice in Figure 8.




Fig. 10. Down-set Birkhoff completion of the concept lattice in Figure 8.

leads in a straightforward manner to another equivalent approach to defining the Birkhoff completion.

**Definition 4 (Proper Premise [4]).** Let (G, M, I) be a context. An attribute set  $A \subseteq M$  is a **proper premise** if

$$A^{\bullet} \coloneqq A'' \setminus \left(A \cup \bigcup_{n \in A} (A \ \{n\})''\right)$$

is not empty.

**Definition 5.** The canonical direct basis of a finite context  $\mathbb{K}$  is defined as the set  $\mathcal{CDB}(\mathbb{K})$  of all implications  $A \to A^{\bullet}$ , where A is a proper premise.

Based on the following theorem, we can infer that there is correspondence between distributive lattices and those that are isomorphic to concept lattices whose contexts proper premises are singletons.

**Theorem 6 (Theorem 41 (8) of [4]).** Let  $\mathbb{K}$  be a finite, reduced context. Then  $\mathfrak{B}(\mathbb{K})$  is distributive iff every proper premise of (G, M, I) is a singleton.

Thus the reduced contexts of concept lattices that are not distributive have proper premises that are of size 2 or larger. This motivates the following definition.

**Definition 6.** Let  $\mathbb{K}$  be a finite, reduced context and  $A \to A^{\bullet}$  an implication of its canonical direct basis. We call  $A \to A^{\bullet}$  a distributive implication if |A| = 1 and a non-distributive implication if  $|A| \ge 2$ .

With Theorem 5 (1  $\cong$  4) we showed that the Birkhoff completion of a concept lattice can be achieved by computing a finer closure system  $\operatorname{Int}(BC(\mathbb{K}))$  of  $\operatorname{Int}(\mathbb{K})$  on the set M, i.e.,  $\operatorname{Int}(\mathbb{K}) \subseteq \operatorname{Int}(BC(\mathbb{K}))$ . With the following proposition, we study this completion in terms of implications. For that, we make use of the dualism between closure systems and their implicational theories, i.e.,  $\operatorname{Int}(\mathbb{K}) \subseteq \operatorname{Int}(BC(\mathbb{K}))$  implies for the set of all attribute implications  $\operatorname{Th}(BC(\mathbb{K})) \subseteq \operatorname{Th}(\mathbb{K})$  [3].

**Lemma 2.** Let  $\mathbb{K}$  be a finite, reduced context,  $\mathcal{A}$  the set of its proper premises, and  $BC(\mathcal{A}) := \{A \in \mathcal{A} \mid |A| = 1\}$ . For all  $A \in BC(\mathcal{A})$  holds  $A^{\bullet} = A'' \setminus A$  independent of being computed in  $\mathbb{K}$  or in  $BC(\mathbb{K})$ . Furthermore  $BC(\mathcal{A})$  is the set of all proper premises of  $BC(\mathbb{K})$ .

*Proof.* As in the proof of Theorem 5, we use ' as symbol for the derivation operator in  $\mathbb{K}$  and '! for the derivation operator in  $BC(\mathbb{K})$ . We start by observing that  $BC(\mathbb{K})$  is attribute reduced since  $\mathbb{K}$  is attribute reduced and  $\mathrm{Int}(\mathbb{K}) \subseteq \mathrm{Int}(BC(\mathbb{K}))$ . As both  $\mathbb{K}$  and  $BC(\mathbb{K})$  are attribute reduced, we have in particular  $\emptyset'' = \emptyset = \emptyset$ !! Let now  $A = \{n\} \in BC(\mathcal{A})$ . With

$$A^{!!} = (A^!)^! = (A' \cup \{ \overline{m} \in \overline{\mathcal{M}(M)} \mid m \not \geq_{\mathbb{K}} n \})^! = (A')^! \cap \{ \overline{m} \in \overline{\mathcal{M}(M)} \mid m \not \geq_{\mathbb{K}} n \}^!$$

$$= A'' \cap \{ k \in M \mid m \not \geq_{\mathbb{K}} k \text{ for all } m \text{ with } m \not \geq_{\mathbb{K}} n \} = A'' \cap \{ k \in M \mid k \geq_{\mathbb{K}} n \} = A''$$

we obtain

$$\begin{split} A^{\bullet} \ \ &(\text{computed in } BC(\mathbb{K})) = A^{!!} \setminus (A \cup \bigcup_{n \in A} A \setminus \{n\}^{!!}) \\ &= A^{!!} \setminus (A \cup \bigcup_{n \in A} \emptyset^{!!}) = A^{!!} \setminus A = \qquad A'' \setminus A \qquad = A'' \setminus (A \cup \bigcup_{n \in A} \emptyset'') \\ &= A'' \setminus (A \cup \bigcup_{n \in A} A \setminus \{n\}'') = A^{\bullet} \ \ &(\text{computed in } \mathbb{K}) \ . \end{split}$$

It follows immediately that A is a proper premise of  $BC(\mathbb{K})$  because it is a proper premise of  $\mathbb{K}$  and thus  $A^{\bullet}$  is not empty.

It remains to show that all proper premises of  $BC(\mathbb{K})$  are contained in  $BC(\mathcal{A})$ . Let A be a proper premise of  $BC(\mathbb{K})$ . We can conclude from  $Int(\mathbb{K}) \subseteq Int(BC(\mathbb{K}))$  that  $A^{I_{\mathbb{K}}I_{\mathbb{K}}} \neq A$ . Moreover, since  $\mathbb{K}$  is reduced, we can infer that  $\emptyset \in Int(\mathbb{K})$ . Thus  $A^{\bullet}$  (computed in  $\mathbb{K}$ ) is not empty and A is a proper premise of  $\mathbb{K}$ . Additionally, we have |A| = 1 because of the distributivity of  $\mathfrak{B}(BC(\mathbb{K}))$ .  $\square$ 

**Corollary 2.** For a finite reduced context  $\mathbb{K}$  is  $BC(\mathcal{CDB}(\mathbb{K})) = \mathcal{CDB}(BC(\mathbb{K}))$  with

$$BC(\mathcal{CDB}(\mathbb{K})) := \{A \to A^{\bullet} \mid A \to A^{\bullet} \in \mathcal{CDB}(\mathbb{K}) \text{ and } |A| = 1\}$$
.

Note that for all proper premises A of  $\mathbb{K}$  with  $|A| \geq 2$ , A is not a proper premise of  $BC(\mathbb{K})$ , and  $A^{\bullet}$  evaluated in  $BC(\mathbb{K})$  will be a strict subset of  $A^{\bullet}$  evaluated in  $\mathbb{K}$ .

With Lemma 2 and Corollary 2, we have not only shown a relation between a concept lattice, its Birkhoff completion and their canonical direct bases, but we can also derive a procedure for the computation of the Birkhoff completion of the concept lattice. That is, compute the closure system of attribute sets that are closed with respect to the implicational theory  $BC(\mathcal{CDB}(\mathbb{K}))$ .

The condition that  $\mathbb{K}$  has to be reduced seems to be a constraint for the application of this proposition. However for a context that is not reduced, we can first reduce it, then compute the completion with respect to proper premises and lastly re-add the reducible attributes to the completion.

The canonical direct basis of the concept lattice (Figure 7) about the administrative geography of the UK is given in Figure 11. According to Lemma 2, the canonical direct base of the Birkhoff completion of the would thus consist of implications (1) to (3) — which can easily be verified in Figure 9.

```
\{UK\} \rightarrow \{British Islands\}
                                                                                                    (1)
                                    \{GB\} \rightarrow \{UK, British Islands\}
                                                                                                    (2)
                     \{Channel Islands\} \rightarrow \{British Islands\}
                                                                                                    (3)
 \{Ireland (Island), British Islands\} \rightarrow \{UK\}
                                                                                                    (4)
               \{UK, Channel Islands\} \rightarrow \{Ireland (Island), British Islands, GB\}
                                                                                                    (5)
               \{Channel Islands, GB\} \rightarrow \{UK, Ireland (Island), British Islands\}
                                                                                                    (6)
\{Ireland (Island), Channel Islands\} \rightarrow \{UK, British Islands, GB\}
                                                                                                    (7)
               \{Ireland (Island), GB\} \rightarrow \{UK, British Islands, Channel Islands\}
                                                                                                    (8)
```

Fig. 11. Canonical direct basis of the administrative geography of the UK.

## 6 Application: analysis of the administrative geography of the UK

We will now study by an example how the Birkhoff completion of the context interacts with the completion by implications; and what can be learnt from the Birkhoff completion. To this end, we consider again the administrative geopgraphy of the UK.

In the top left of the up-set BC in Figure 9, we can see that  $\overline{\text{British Islands}}$  is generating the same concept as Ireland (State). This indicates that the legal

distinction 'British Islands' is defined by exclusion – it contains all countries of the British Isles except the state of Ireland.

The other four new objects are all found at the lower part of the concept lattice, and all generate new concepts. These objects invalidate the implications with two-element premises of Figure 11: {Ireland (Island), British Islands}  $\rightarrow$  {UK} is invalidated by  $\overline{\text{UK}}$ , {UK, Channel Islands}  $\rightarrow$  {GB} by  $\overline{\text{GB}}$ , and {Ireland (Island), GB}  $\rightarrow$  {Channel Islands} by  $\overline{\text{Channel Islands}}$ .

Let us now assume for a moment, that geographical features are given by nature and not subject to modification, while legal distinctions might change over the years. We would thus assume that Ireland (Island), GB and the Channel Islands will always remain geographically disjoint. A new country having all attributes of  $\overline{\text{Channel Islands}}$  will therefore never exist. On the other hand, in the future there may arise new countries on the British Isles that are affiliated to different combinations of legal distinctions — or existing countries might change their affiliations. Therefore we cannot exclude the potential existence of countries having the attributes of  $\overline{\text{UK}}$ , of  $\overline{\text{GB}}$  or of  $\overline{\text{Ireland}(\text{Island})}$ .

We conclude this study of UK's administrative geography by a look at the down-set Birkhoff completion in Figure 10. First, we observe that all implications of the canonical direct base (Figure 11) still hold. Note that this is not contradicting Theorem 6 even though the down-set completion is distributive, because the original attributes (except 'British Islands') are not reduced any more.

As in the up-set completion we can observe that 'British Islands' is the complement of 'Ireland (State)', because Ireland (State) is attached to the same concept as 'British Islands'. The seven other new attibutes generate three new concepts. These concepts can be understood in two ways. The first is as disjunction of the attributes below. The top left concept, for instance, could be understood as 'Ireland (Island) or Channel Islands'. The second way is to consider them as negation, but note that we are now negating objects. While one is tempted to consider Wales, Scotland England) at the top left as '¬ Great Britain', the attribute Northern Island shows that it is not that simple: Because of the way it was constructed this attribute is incident with all objects that are neither 'Northern Island' nor other objects (such as 'Ireland (State)') which are implied (as object implication) by 'Northern Island'. Either way, we can understand these newly generated concepts as proposals for new concepts which are waiting for a denomination.

#### 7 Conclusion

Motivated by the fact that distributivity in concept lattices arises from the fact that everyday thinking predominantly uses simple implications, we have introduced the Birkhoff completion of a finite lattice L as the minimal distributive lattice in which L can be embeddeded as sub-semilattice. We have presented a

<sup>&</sup>lt;sup>7</sup> For sake of better readability, we will, in the sequel, subsume the crown dependencies under the term 'country'.

corresponding completion for formal contexts and a corresponding restriction of the implicational basis, and have shown their equivalences.

Depending on the characteristics of the dataset, either the up-set BC or the down-set BC might be more beneficial. If all objects under consideration are known from the beginning — like the countries in the British Isles — the down-set BC might focus our view on potentially new attributes for their description, while the up-set BC is only of interest, if we do not want to exclude the possibility of new objects.

We have defined the up-set Birkhoff completion of a lattice as standard rather than the down-set completion because of its dualism with the Birkhoff completion of the canonical direct base for the attribute implications. The same dualism holds of course also the down-set completion together with the object implications, but the latter are far less promeninent in everyday thinking.

Note that the Birkhoff completions are by no means the only ways to turn a non-distributive lattice into a distributive one. Other obvious ways are the deletion of objects and/or attributes, or the modification of the incidence relation of the formal context. Our future work will study the effects of these options more closely. When one is aiming at simplifying a given dataset so that all its implications are distributive, one could choose any of these options. Which choice is more adequate depends on the characteristics of the described objects.

#### References

- B.A.Davey, H.A.Priestley: Introduction to Lattices and Order. Cambridge University Press (2002)
- 2. Birkhoff, G.: Lattice theory. In: Colloquium Publications, vol. 25. American Mathematical Society, 3. edn. (1967)
- 3. Caspard, N., Monjardet, B.: The lattices of closure systems, closure operators, and implicational systems on a finite set: a survey. Discrete Applied Mathematics 127(2), 241–269 (2003). https://doi.org/https://doi.org/10.1016/S0166-218X(02) 00209-3, https://www.sciencedirect.com/science/article/pii/S0166218X02002093, ordinal and Symbolic Data Analysis (OSDA '98), Univ. of Massachusetts, Amherst, Sept. 28-30, 1998.
- 4. Ganter, B., Wille, R.: Formal Concept Analysis Mathematical Foundations. Springer (1999)
- Hanika, T., Hirth, J.: Conexp-clj a research tool for fca. In: Cristea, D., Ber, F.L., Missaoui, R., Kwuida, L., Sertkaya, B. (eds.) ICFCA (Supplements). CEUR Workshop Proceedings, vol. 2378, pp. 70–75. CEUR-WS.org (2019)
- Stumme, G.: Free distributive completions of partial complete lattices. In: Order 14, 179–189 (1998), http://www.kde.cs.uni-kassel.de/stumme/papers/2000/SIGKDD\_Explorations00.ps
- 7. Wikipedia: Administrative geography of the united kingdom (2024), https://en.wikipedia.org/wiki/Administrative geography of the United Kingdom
- 8. Wikipedia: British isles euler diagram (2024), https://en.wikipedia.org/wiki/File: British Isles Euler diagram 15.svg
- 9. Wille, R.: Tensorial decomposition of concept lattices. Order 2, 81–95 (1985)
- 10. Wille, R.: Truncated distributive lattices: Conceptual structures of simple-implicational theories. Order **20**(3), 229–238 (2003)